

Theoretical aspects of chiral separation in capillary elec

Journal of Chromatography A

603, 235-241

DOI: 10.1016/0021-9673(92)85366-2

Citation Report

#	ARTICLE	IF	CITATIONS
1	Theoretical aspects of chiral separation in capillary electrophoresis. Journal of Chromatography A, 1992, 609, 363-367.	3.7	230
2	Separation of the enantiomers of some racemic nonsteroidal aromatase inhibitors and barbiturates by capillary electrophoresis. Chirality, 1993, 5, 516-526.	2.6	53
3	Theory of chiral separation in capillary electrophoresis. Journal of Chromatography A, 1993, 636, 57-62.	3.7	172
4	Differential binding of tioconazole enantiomers to hydroxypropyl- β -cyclodextrin studied by capillary electrophoresis. Journal of Chromatography A, 1993, 636, 149-152.	3.7	128
5	Validation of a capillary electrophoresis method for the enantiomeric purity testing of fluparoxan. Journal of Chromatography A, 1993, 645, 193-196.	3.7	64
6	Theoretical aspects of chiral separation in capillary electrophoresis. Journal of Chromatography A, 1993, 635, 113-118.	3.7	164
7	Chiral separation of basic drugs using cyclodextrins as chiral pseudo-stationary phases in capillary electrophoresis. Journal of Chromatography A, 1993, 648, 267-274.	3.7	116
8	Quantitative aspects of the application of capillary electrophoresis to the analysis of pharmaceuticals and drug related impurities. Journal of Chromatography A, 1993, 646, 245-257.	3.7	99
9	Charged and uncharged cyclodextrins as chiral selectors in capillary electrophoresis. Chromatographia, 1993, 37, 475-481.	1.3	198
10	Simultaneous chiral separation of leucovorin and its major metabolite 5-methyl-tetrahydrofolate by capillary electrophoresis using cyclodextrins as chiral selectors: Estimation of the formation constant and mobility of the solute-cyclodextrin complexes. Chromatographia, 1993, 35, 419-429.	1.3	124
11	Chiral separation of basic drugs using cyclodextrin-modified capillary zone electrophoresis. Analytical Chemistry, 1993, 65, 885-893.	6.5	159
12	Fully Automated Analysis of Amino Acid Enantiomers by Derivatization and Chiral Separation on a Capillary Electrophoresis Instrument. Journal of Liquid Chromatography and Related Technologies, 1994, 17, 1883-1897.	1.0	26
13	Chiral separations of amino acids by capillary electrophoresis and high-performance liquid chromatography employing chiral crown ethers. Journal of Chromatography A, 1994, 685, 321-329.	3.7	45
14	Separation of enantiomers by capillary electrophoretic techniques. Journal of Chromatography A, 1994, 666, 295-319.	3.7	274
15	Capillary zone electrophoresis for separation of drug enantiomers using cyclodextrins as chiral selectors. Journal of Chromatography A, 1994, 666, 337-350.	3.7	87
16	Chiral separations of basic and acidic compounds in modified capillaries using cyclodextrin-modified capillary zone electrophoresis. Journal of Chromatography A, 1994, 666, 351-365.	3.7	56
17	Enantioselective separations using capillary electrophoresis. Chirality, 1994, 6, 25-40.	2.6	92
18	Investigation of enantioselective ligand-protein binding and displacement interactions using capillary electrophoresis. Chirality, 1994, 6, 230-238.	2.6	45

#	ARTICLE	IF	CITATIONS
19	Chiral separations by capillary electrophoresis: Present state of the art. <i>Electrophoresis</i> , 1994, 15, 755-762.	2.4	69
20	A theoretical approach to chiral capillary electrophoresis with some practical implications. <i>Electrophoresis</i> , 1994, 15, 774-778.	2.4	61
21	Separations of derivatized amino acid enantiomers by cyclodextrin-modified capillary electrophoresis: Mechanistic and molecular modeling studies. <i>Electrophoresis</i> , 1994, 15, 785-792.	2.4	62
22	Enantiomeric separation of salbutamol and related impurities using capillary electrophoresis. <i>Electrophoresis</i> , 1994, 15, 808-817.	2.4	57
23	Chiral separation of basic drugs by capillary zone electrophoresis with cyclodextrin additives. <i>Electrophoresis</i> , 1994, 15, 818-823.	2.4	117
24	Separation of enantiomers by affinity electrokinetic chromatography using avidin. <i>Electrophoresis</i> , 1994, 15, 848-853.	2.4	83
25	Enantiomeric separation of drugs by affinity electrokinetic chromatography using dextran sulfate. <i>Electrophoresis</i> , 1994, 15, 1335-1340.	2.4	48
26	Capillary zone electrophoresis in pharmaceutical and biomedical analysis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1994, 12, 579-611.	2.8	62
27	Stereoselective determination of verapamil and norverapamil by capillary electrophoresis. <i>Biomedical Applications</i> , 1994, 654, 121-127.	1.7	52
28	Cyclodextrins as enantioselective mobile phase modifiers for chiral capillary electrophoresis effects of pH and cyclodextrin concentration. <i>Biomedical Applications</i> , 1994, 657, 291-300.	1.7	34
29	Systematic approach to treatment of enantiomeric separations in capillary electrophoresis and liquid chromatography.. <i>Journal of Chromatography A</i> , 1994, 680, 147-155.	3.7	39
30	Practical aspects of chiral separations of pharmaceuticals by capillary electrophoresis I. Separation optimization. <i>Journal of Chromatography A</i> , 1994, 680, 157-162.	3.7	49
31	Chiral selectors with chelating properties in liquid chromatography: Fundamental reflections and selective review of recent developments. <i>Journal of Chromatography A</i> , 1994, 666, 55-76.	3.7	145
32	Chiral separation of drugs by capillary electrophoresis using β -cyclodextrin polymer. <i>Journal of Chromatography A</i> , 1994, 678, 333-342.	3.7	44
33	Chiral separation of drugs using cyclodextrins in capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 1994, 659, 449-457.	3.7	55
34	Optimization of a capillary electrophoresis method to determine the chiral purity of a drug. <i>Journal of Chromatography A</i> , 1994, 680, 609-621.	3.7	63
35	High-performance liquid chromatographic separation of cis- and trans isomers of proline-containing peptides. <i>Journal of Chromatography A</i> , 1994, 661, 7-12.	3.7	26
36	Effect of the degree of substitution of (2-hydroxy)propyl- β -cyclodextrin on the enantioseparation of organic acids by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1994, 678, 139-144.	3.7	56

#	ARTICLE	IF	CITATIONS
37	Plackett-Burman experimental design in chiral analysis using capillary electrophoresis. <i>Chromatographia</i> , 1994, 38, 723-729.	1.3	71
38	Factors affecting the separation of mandelic acid enantiomers by capillary electrophoresis. <i>Chromatographia</i> , 1994, 38, 730-736.	1.3	26
39	Capillary Electrophoresis with Chiral Selectors: Optimization of Separation and Determination of Thermodynamic Parameters for Binding of Tioconazole Enantiomers to Cyclodextrins. <i>Analytical Chemistry</i> , 1994, 66, 2866-2873.	6.5	144
40	Chiral Resolution of Cationic Drugs of Forensic Interest by Capillary Electrophoresis with Mixtures of Neutral and Anionic Cyclodextrins. <i>Analytical Chemistry</i> , 1994, 66, 4019-4026.	6.5	264
41	Sulfobutyl Ether .beta.-Cyclodextrin as a Chiral Discriminator for Use with Capillary Electrophoresis. <i>Analytical Chemistry</i> , 1994, 66, 4013-4018.	6.5	224
42	A Peak Resolution Model for the Capillary Electrophoretic Separation of the Enantiomers of Weak Acids with Hydroxypropyl .beta.-Cyclodextrin-Containing Background Electrolytes. <i>Analytical Chemistry</i> , 1994, 66, 619-627.	6.5	167
43	Chiral Separations Using Micellar Electrokinetic Capillary Chromatography and a Polymerized Chiral Micelle. <i>Analytical Chemistry</i> , 1994, 66, 3773-3776.	6.5	170
44	Separation of binaphthyl enantiomers by capillary zone electrophoresis and electrokinetic chromatography. <i>Journal of High Resolution Chromatography</i> , 1995, 18, 659-664.	1.4	32
45	Mechanisms of enantiomeric resolution in cyclodextrin-modified capillary electrophoretic separations of binaphthyl compounds. <i>Chirality</i> , 1995, 7, 401-408.	2.6	18
46	Rapid chiral separation methods development by cyclodextrin-mediated capillary electrophoresis for acidic and basic compounds. <i>Chirality</i> , 1995, 7, 409-414.	2.6	25
47	Determination of the enantiomeric purity of dihydroxy- and dimethoxy-2-aminotetralins by high-performance capillary electrophoresis with cyclodextrins as chiral selector. <i>Chirality</i> , 1995, 7, 459-468.	2.6	12
48	Chiral separation of naphthalene-2, 3-dialdehyde labelled peptides by cyclodextrin-modified electrokinetic chromatography. <i>Biomedical Chromatography</i> , 1995, 9, 295-301.	1.7	20
49	Chiral separations using heparin and dextran sulfate in capillary zone electrophoresis. <i>Analytica Chimica Acta</i> , 1995, 307, 185-191.	5.4	55
50	Viscous effects in capillary electrophoresis: Theory and experiment. <i>Electrophoresis</i> , 1995, 16, 2074-2085.	2.4	14
51	Chiral separation in capillary electrophoresis. <i>Electrophoresis</i> , 1995, 16, 2127-2131.	2.4	55
52	Electroosmotic control of chiral separation in capillary zone electrophoresis. <i>Electrophoresis</i> , 1995, 16, 2132-2136.	2.4	13
53	Protein chiral selectors in free-solution capillary electrophoresis and packed-capillary electrochromatography. <i>Journal of Chromatography A</i> , 1995, 694, 285-296.	3.7	150
54	Partial separation zone technique for the separation of enantiomers by affinity electrokinetic chromatography with proteins as chiral pseudo-stationary phases. <i>Journal of Chromatography A</i> , 1995, 694, 277-284.	3.7	170

#	ARTICLE	IF	CITATIONS
55	Extending the scope of chiral separation of basic compounds by cyclodextrin-mediated capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 1995, 692, 253-265.	3.7	81
56	Experimental evidence for the existence of duoselective (Type III) enantiomer separations in the capillary electrophoretic analysis of chiral weak acids. <i>Journal of Chromatography A</i> , 1995, 692, 319-325.	3.7	50
57	Enantioselective capillary electrophoresis of amino acid derivatives on cyclodextrin evaluation of structure-resolution relationships. <i>Journal of Chromatography A</i> , 1995, 697, 549-560.	3.7	64
58	Enantiomeric separation by capillary electrophoresis using a soluble neutral β^2 -cyclodextrin polymer. <i>Journal of Chromatography A</i> , 1995, 709, 89-98.	3.7	47
59	Systematic approach to treatment of enantiomeric separations in capillary electrophoresis and liquid chromatography II. A study of the enantiomeric separation of fluoxetine and norfluoxetine. <i>Journal of Chromatography A</i> , 1995, 700, 59-67.	3.7	56
60	Application of modified cyclodextrins in capillary electrophoresis for enantiomeric resolution of propranolol and analogues. <i>Journal of Chromatography A</i> , 1995, 705, 351-361.	3.7	29
61	Determination of β^2 -cyclodextrin inclusion complex constants for 3,4-dihydro-2-H-1-benzopyran enantiomers by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1995, 707, 311-326.	3.7	54
62	Enantiomeric purity determination of propranolol by cyclodextrin-modified capillary electrophoresis. <i>Journal of Chromatography A</i> , 1995, 717, 203-209.	3.7	91
63	Combined polymerized chiral micelle and β^3 -cyclodextrin for chiral separation in capillary electrophoresis. <i>Journal of Chromatography A</i> , 1995, 711, 297-304.	3.7	113
64	Determination of the enantiomeric purity of 5,6-dihydroxy-2-aminotetralin by high-performance capillary electrophoresis with crown ether as chiral selector. <i>Journal of Chromatography A</i> , 1995, 715, 143-149.	3.7	27
65	Use of organic modifiers to enhance chiral selectivity in capillary electrophoresis. <i>Amino Acids</i> , 1995, 8, 337-344.	2.7	14
66	Enantiomeric separation of denopamine by capillary electrophoresis and high-performance liquid chromatography using cyclodextrins. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1995, 13, 1483-1492.	2.8	19
67	Optical resolution of drugs by capillary electrophoretic techniques. <i>Journal of Chromatography A</i> , 1995, 694, 245-276.	3.7	274
68	Chiral separation of deprenyl and its major metabolites using cyclodextrin-modified capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 1995, 709, 157-162.	3.7	42
69	Capillary zone electrophoretic separation of the enantiomers of dipeptides based on host-guest complexation with a chiral crown ether. <i>Journal of Chromatography A</i> , 1995, 709, 81-88.	3.7	70
70	Resolution of the enantiomers of oxamniquine by capillary electrophoresis and high-performance liquid chromatography with cyclodextrins and heparin as chiral selectors. <i>Journal of Chromatography A</i> , 1995, 700, 51-58.	3.7	45
71	Direct and indirect chiral separation of amino acids by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1995, 704, 179-193.	3.7	103
72	Capillary zone electrophoretic determination of C2 ω —C18 linear saturated free fatty acids with indirect absorbance detection. <i>Journal of Chromatography A</i> , 1995, 708, 339-350.	3.7	72

#	ARTICLE	IF	CITATIONS
73	Chiral separation of amphetamines by high-performance capillary electrophoresis. Journal of Chromatography A, 1995, 717, 219-228.	3.7	92
74	Enantiomeric separation of amphetamine related drugs by capillary zone electrophoresis using native and derivatized β -cyclodextrin as chiral additives. Journal of Chromatography A, 1995, 710, 339-345.	3.7	57
75	Effect of urea addition on chiral separation of dansylamino acids by capillary zone electrophoresis with cyclodextrins. Journal of Chromatography A, 1995, 710, 331-337.	3.7	32
76	Buffer effects in the desionoselective/ionoselective/duoselective separation selectivity model-assisted optimization of the capillary electrophoretic separation of enantiomers I. Low-pH phosphate buffers. Journal of Chromatography A, 1995, 716, 197-205.	3.7	20
77	Enantiomeric separation of D/L-carnitine using HPLC and CZE after derivatization. Chromatographia, 1995, 40, 287-295.	1.3	52
78	Enantiomeric separation of trimetoquinol, denopamine and timepidium by capillary electrophoresis and HPLC and the application of capillary electrophoresis to the optical purity testing of the drugs. Chromatographia, 1995, 40, 638-644.	1.3	34
79	Capillary electrophoretic separation of basic drugs using surface-modified C8 capillaries and derivatized cyclodextrins as structural/chiral selectors. Chromatographia, 1995, 40, 697-704.	1.3	22
80	Effect of Micelles and Mixed Micelles on Efficiency and Selectivity of Antibiotic-Based Capillary Electrophoretic Enantioseparations. Analytical Chemistry, 1995, 67, 2088-2095.	6.5	121
81	Enantiomeric Separation of Drugs by Mucopolysaccharide-Mediated Electrokinetic Chromatography. Analytical Chemistry, 1995, 67, 2334-2341.	6.5	83
82	Effect of the Initial Potential Ramp on the Accuracy of Electrophoretic Mobilities in Capillary Electrophoresis. Analytical Chemistry, 1995, 67, 3079-3081.	6.5	29
83	Separation of PAHs by Capillary Electrophoresis with Laser-Induced Fluorescence Detection Using Mixtures of Neutral and Anionic β -Cyclodextrins. Analytical Chemistry, 1995, 67, 3004-3010.	6.5	75
84	Affinity Capillary Electrophoresis. Accounts of Chemical Research, 1995, 28, 461-468.	15.6	147
85	Chiral Separation and Detection of Terbutaline and Ephedrine by Capillary Electrophoresis Coupled with Ion Spray Mass Spectrometry. Analytical Chemistry, 1995, 67, 2054-2058.	6.5	134
86	Chiral Separations by Nonaqueous Capillary Electrophoresis. Analytical Chemistry, 1996, 68, 3460-3467.	6.5	173
87	Chiral separation of dansyl amino acids by capillary electrophoresis: Comparison of formamide and N-Methylformamide as background electrolytes. Chromatographia, 1996, 43, 242-246.	1.3	55
88	Separation of drug enantiomers by capillary electrophoresis: β -cyclodextrin as chiral solvating agent. Chromatographia, 1996, 42, 106-110.	1.3	25
89	Chiral separation of local anaesthetics with capillary electrophoresis. Chromatographia, 1996, 42, 263-268.	1.3	40
90	Determination of the binding of β -cyclodextrin derivatives to adamantane carboxylic acids using capillary electrophoresis. Chromatographia, 1996, 43, 659-662.	1.3	38

#	ARTICLE	IF	CITATIONS
91	Micellar electrokinetic capillary chromatography with in-situ charged micelles VI. Evaluation of novel chiral micelles consisting of steroidal-glycoside surfactant-borate complexes. Journal of Chromatography A, 1996, 724, 285-296.	3.7	63
92	Maximization of separation efficiency in capillary electrophoretic chiral separations by means of mobility-matching background electrolytes. Journal of Chromatography A, 1996, 730, 273-278.	3.7	44
93	Enantiomer separation of drugs by electrokinetic chromatography. Journal of Chromatography A, 1996, 735, 57-76.	3.7	131
94	Enantiomeric separation of amphetamine and phenylephrine by cyclodextrin-mediated capillary zone electrophoresis. Journal of Chromatography A, 1996, 735, 295-301.	3.7	23
95	Determination of the enantiomeric purity of dexfenfluramine by capillary electrophoresis: use of a Plackett-Burman design for the optimization of the separation. Journal of Chromatography A, 1996, 736, 281-289.	3.7	51
96	Chiral separation of dansyl-amino acids in a nonaqueous medium by capillary electrophoresis. Journal of Chromatography A, 1996, 737, 263-272.	3.7	86
97	Influence of the nature of the buffer on chiral separation in capillary electrophoresis. Journal of Chromatography A, 1996, 745, 61-71.	3.7	16
98	Cyclodextrins as buffer additives for the enantiomeric separation of pinacidil in capillary zone electrophoresis. Journal of Chromatography A, 1996, 749, 300-303.	3.7	7
99	Determination of the binding constant of salbutamol to unmodified and ethylated cyclodextrins by affinity capillary electrophoresis. Journal of Chromatography A, 1996, 735, 321-331.	3.7	48
100	Electrokinetic chromatography in suppressed electroosmotic flow environment: Use of a charged cyclodextrin for the separation of enantiomers and geometric isomers. Electrophoresis, 1996, 17, 1575-1583.	2.4	65
101	A practical procedure for the determination of association constants of the analyte-chiral selector equilibria by capillary zone electrophoresis. Electrophoresis, 1996, 17, 1921-1924.	2.4	18
102	Separation of neutral and basic enantiomers by cyclodextrin electrokinetic chromatography using anionic cyclodextrin derivatives as chiral pseudo-stationary phases. Journal of High Resolution Chromatography, 1996, 19, 421-433.	1.4	80
103	Chiral separation of methoxamine and lobeline using cyclodextrin as buffer additives in capillary zone electrophoresis. Journal of High Resolution Chromatography, 1996, 19, 478-480.	1.4	10
104	Use of cyclodextrins as chiral selector for the chiral separation of anticholinergic drugs such as anisodamine and glycopyrronium in capillary zone electrophoresis. Journal of High Resolution Chromatography, 1996, 19, 697-699.	1.4	8
105	Cyclodextrin-aided capillary electrophoretic separation and laser-induced fluorescence detection of polynuclear aromatic hydrocarbons (PAHs). , 1996, 9, 515-523.		9
106	Role of capillary electrophoresis methods in the drug development process. Chirality, 1996, 8, 108-121.	2.6	45
107	Capillary electrophoretic enantiomeric separations using the glycopeptide antibiotic, teicoplanin. Chirality, 1996, 8, 88-107.	2.6	98
108	Sulfated cyclodextrins for the chiral separations of catecholamines and related compounds in the reversed electrophoretic polarity mode. , 1996, 8, 316-324.		48

#	ARTICLE	IF	CITATIONS
109	Capillary electrophoresis of herbicides. III. Evaluation of octylmaltopyranoside chiral surfactant in the enantiomeric separation of phenoxy acid herbicides. , 1996, 8, 518-524.		35
110	Enantiomeric separations of basic pharmaceutical drugs by micellar electrokinetic chromatography using a chiral surfactant, N-dodecoxycarbonylvaline. Biomedical Applications, 1996, 683, 15-28.	1.7	50
111	Chiral separations of underivatized arylpropionic acids by capillary zone electrophoresis with various cyclodextrins Acidity and inclusion constant determinations. Journal of Chromatography A, 1996, 735, 311-320.	3.7	67
112	Examination of the origin, variation, and proper use of expressions for the estimation of association constants by capillary electrophoresis. Journal of Chromatography A, 1996, 721, 173-186.	3.7	273
113	Multiple-plug binding assays using affinity capillary electrophoresis electrophoresis. Journal of Chromatography A, 1996, 727, 291-299.	3.7	47
114	Enantiomeric separation of local anaesthetic drugs by micellar electrokinetic capillary chromatography with taurodeoxycholate as chiral selector. Journal of Chromatography A, 1996, 737, 301-313.	3.7	35
115	Direct chiral separation of amino acids derivatized with 2-(9-anthryl)ethyl chloroformate by capillary electrophoresis using cyclodextrins as chiral selectors Effect of organic modifiers on resolution and enantiomeric elution order. Journal of Chromatography A, 1996, 731, 283-292.	3.7	56
116	Separation of enatiomers of drugs by capillary electrophoresis III. β -cyclodextrin as chiral solvating agent. Journal of Chromatography A, 1996, 735, 333-343.	3.7	53
117	Comparative capillary electrophoretic and nuclear magnetic resonance studies of the chiral recognition of racemic metomidate with cyclodextrin hosts. Journal of Chromatography A, 1996, 732, 133-142.	3.7	69
118	Chiral separation of dioxypromethazine enantiomers by capillary electrophoresis using β -cyclodextrin as a chiral selector. Journal of Chromatography A, 1996, 732, 175-181.	3.7	24
119	Identification of chiral drug isomers by capillary electrophoresis. Journal of Chromatography A, 1996, 735, 77-121.	3.7	319
120	Capillary electrophoresis of cardiovascular drugs. Journal of Chromatography A, 1996, 735, 123-150.	3.7	23
121	Separation of the four stereoisomers of Z11556A, intermediate in the synthesis of triazole antimycotics, by capillary electrophoresis. Journal of Chromatography A, 1996, 741, 123-130.	3.7	8
122	Self-consistent framework for standardising mobilities in free solution capillary electrophoresis: applications to oligoglycines and oligoalanines. Journal of Chromatography A, 1996, 741, 99-113.	3.7	58
123	Chiral separation of muscarinic antagonists by capillary zone electrophoresis with cyclodextrin additives. Journal of Chromatography A, 1996, 741, 287-294.	3.7	9
124	Separation of positional and structural isomers by cyclodextrin-mediated capillary zone electrophoresis. Journal of Chromatography A, 1996, 745, 9-15.	3.7	27
125	Chiral separation in non-aqueous media by capillary electrophoresis using the ion-pair principle. Journal of Chromatography A, 1996, 745, 37-44.	3.7	113
126	Systematic approach to the treatment of enantiomeric separations in capillary electrophoresis and liquid chromatography III. A binding constant-retention factor relationship and effects of acetonitrile on the chiral separation of tioconazole. Journal of Chromatography A, 1996, 745, 25-35.	3.7	43

#	ARTICLE	IF	CITATIONS
127	Use of cyclodextrins in the capillary electrophoretic separation of erythro- and threo-mefloquine enantiomers. Journal of Chromatography A, 1996, 745, 17-23.	3.7	25
128	Determination of binding constants and the influence of methanol on the separation of drug enantiomers in cyclodextrin modified capillary electrophoresis. Journal of Chromatography A, 1996, 745, 181-187.	3.7	30
129	Utilization of in enantiomeric separation by micellar electrokinetic chromatography. Journal of Chromatography A, 1996, 749, 257-269.	3.7	22
130	Ergot alkaloids as chiral selectors in capillary electrophoresis determination of the separation mechanism. Journal of Chromatography A, 1996, 755, 251-259.	3.7	22
131	Migration behavior and separation of sulfonamides in capillary zone electrophoresis I. Influence of buffer pH and electrolyte modifier. Journal of Chromatography A, 1996, 755, 261-269.	3.7	37
132	Resolution improvement by use of carboxymethyl- β -cyclodextrin as chiral additive for the enantiomeric separation of basic drugs by capillary electrophoresis. Journal of Pharmaceutical and Biomedical Analysis, 1996, 14, 1107-1114.	2.8	52
133	Cyclodextrins and enantiomeric separations of drugs by liquid chromatography and capillary electrophoresis: basic principles and new developments. Biomedical Applications, 1996, 687, 303-336.	1.7	153
134	Determination of drug-cyclodextrin binding constants by capillary zone electrophoresis. Journal of Inclusion Phenomena and Macrocyclic Chemistry, 1996, 25, 253-256.	1.6	7
135	Chiral Separation of Reduced Haloperidol by Capillary Zone Electrophoresis with Heptakis (2,6-Di-o-Methyl)- β -cyclodextrin. Journal of Liquid Chromatography and Related Technologies, 1996, 19, 1567-1577.	1.0	9
136	Development of a Free Solution Capillary Electrophoresis Method for the Separation and Quantitation of D- and L-Phenylalanine Methyl Ester.. Analytical Letters, 1996, 29, 991-1001.	1.8	3
137	Determination of Association Constants in Cyclodextrin or Vancomycin-Modified Micellar Capillary Electrophoresis. Journal of Liquid Chromatography and Related Technologies, 1997, 20, 203-216.	1.0	15
138	Separation of the Enantiomers of Propranolol by Incorporation of Molecularly Imprinted Polymer Particles as Chiral Selectors in Capillary Electrophoresis. Analytical Communications, 1997, 34, 119-122.	2.2	59
139	Determination of Accurate Electroosmotic Mobility and Analyte Effective Mobility Values in the Presence of Charged Interacting Agents in Capillary Electrophoresis. Analytical Chemistry, 1997, 69, 4445-4451.	6.5	59
140	Enantiomeric Separations by Use of Calixarene Electrokinetic Chromatography. Analytical Chemistry, 1997, 69, 3239-3242.	6.5	70
141	Enantioseparation of 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate-derivatized amino acids by capillary zone electrophoresis using native and substituted β -cyclodextrins as chiral additives II. Evaluation of complexation constants. Journal of Chromatography A, 1997, 759, 167-175.	3.7	8
142	Reversed determination of the formation constants of 1-allyl terguride with mandelic acid optical isomers using capillary electrophoresis. Journal of Chromatography A, 1997, 772, 179-184.	3.7	9
143	Enantiomeric separation of the novel growth hormone secretagogue MK-0677 by capillary zone electrophoresis. Journal of Chromatography A, 1997, 773, 311-320.	3.7	22
144	Electrokinetic chromatography without electroosmotic flow. Journal of Chromatography A, 1997, 792, 125-141.	3.7	37

#	ARTICLE	IF	CITATIONS
145	Selectivity in capillary electrophoresis: the use of proteins. Journal of Chromatography A, 1997, 792, 349-369.	3.7	74
146	Neutral cyclodextrins as chiral agents for enantiomeric separations of chromanes in capillary electrophoresis. Chromatographia, 1997, 44, 313-319.	1.3	8
147	Separation of the enantiomers of basic drugs by affinity capillary electrophoresis using a partial filling technique and β -1-acid glycoprotein as chiral selector. Chromatographia, 1997, 44, 119-128.	1.3	73
148	Charged native β -cyclodextrin as a pseudostationary phase in electrokinetic chromatography. Fresenius' Journal of Analytical Chemistry, 1997, 357, 32-36.	1.5	9
149	Guidelines in selecting ligand concentrations for the determination of binding constants by affinity capillary electrophoresis. Biomedical Applications, 1997, 697, 77-88.	1.7	50
150	Simultaneous determination of the elimination profiles of the individual enantiomers of racemic isoproterenol using capillary electrophoresis and microdialysis sampling. Journal of Pharmaceutical and Biomedical Analysis, 1997, 15, 621-629.	2.8	29
151	A chiral capillary electrophoresis method for ropivacaine hydrochloride in pharmaceutical formulations: Validation and comparison with chiral liquid chromatography. Journal of Pharmaceutical and Biomedical Analysis, 1997, 15, 1051-1061.	2.8	37
152	Selected applications of capillaries with dynamic or permanent anodal electroosmotic flow in chiral separations by capillary electrophoresis. Journal of Pharmaceutical and Biomedical Analysis, 1997, 15, 1577-1584.	2.8	36
153	Use of $^1\text{H-NMR}$ spectroscopy to determine the enantioselective mechanism of neutral and anionic cyclodextrins in capillary electrophoresis. Journal of Pharmaceutical and Biomedical Analysis, 1997, 15, 1603-1619.	2.8	41
154	Quantitative description of migration behavior of porphyrins based on the dynamic complexation model in a nonaqueous capillary electrophoresis system. Electrophoresis, 1997, 18, 82-91.	2.4	52
155	Stereoselective analysis of herbicides by capillary electrophoresis using sulfobutyl ether β -cyclodextrin as chiral selector. Electrophoresis, 1997, 18, 227-234.	2.4	42
156	Quantitative description of analyte migration behavior based on dynamic complexation in capillary electrophoresis with one or more additives. Electrophoresis, 1997, 18, 706-716.	2.4	65
157	Chiral separations by capillary zone electrophoresis: Present state of the art. Electrophoresis, 1997, 18, 843-852.	2.4	69
158	Capillary zone electrophoresis at subzero temperatures II: Chiral separation of biogenic amines. Electrophoresis, 1997, 18, 873-883.	2.4	26
159	Three approaches to enantiomer separation of β -adrenergic antagonists by capillary electrochromatography. Electrophoresis, 1997, 18, 884-890.	2.4	129
160	Use of a zwitterionic cyclodextrin as a chiral agent for the separation of enantiomers by capillary electrophoresis. Electrophoresis, 1997, 18, 891-896.	2.4	29
161	Histamine-modified β -cyclodextrins for the enantiomeric separation of dansyl-amino acids in capillary electrophoresis. Electrophoresis, 1997, 18, 905-911.	2.4	65
162	Determination of association constants of dansyl-amino acids and β -cyclodextrin in N-methylformamide by capillary electrophoresis. Electrophoresis, 1997, 18, 919-923.	2.4	36

#	ARTICLE	IF	CITATIONS
163	Separation of enantiomers of drugs by capillary electrophoresis. Part 4: Hydroxypropyl-gamamacyclodextrin as chiral solvating agent. Electrophoresis, 1997, 18, 924-930.	2.4	39
164	Central composite design in the chiral analysis of amphetamines by capillary electrophoresis. Electrophoresis, 1997, 18, 931-937.	2.4	70
165	Chiral interactions in capillary zone electrophoresis: Computer simulation and comparison with experiment. Electrophoresis, 1997, 18, 938-942.	2.4	13
166	Determination of the enantiomeric purity of N-propionyl-6,7-dimethoxy-2-aminotetralin by cyclodextrin-modified micellar electrokinetic chromatography. Electrophoresis, 1997, 18, 996-1001.	2.4	15
167	Enantiomer separation of disopyramide with capillary electrophoresis using various cyclodextrins. Electrophoresis, 1997, 18, 1002-1006.	2.4	20
168	Enantioseparation of local anaesthetic drugs by capillary zone electrophoresis with cyclodextrins as chiral selectors using a partial filling technique. Electrophoresis, 1997, 18, 1019-1025.	2.4	32
169	Methods for the estimation of binding constants by capillary electrophoresis. Electrophoresis, 1997, 18, 2194-2202.	2.4	153
170	Capillary electrophoretic separations of enantiomers using cyclodextrin-containing background electrolytes. Electrophoresis, 1997, 18, 2305-2310.	2.4	88
171	Redefining the separation factor: A potential pathway to a unified separation science. Electrophoresis, 1997, 18, 2928-2934.	2.4	23
172	Resolution of the diastereomers of a large synthetic peptide by capillary electrophoresis using nonionic surfactants. Electrophoresis, 1997, 18, 2944-2949.	2.4	9
173	Cyclodextrin-modified micellar electrokinetic chromatography for separation of norgestrel enantiomers. Journal of High Resolution Chromatography, 1997, 20, 159-164.	1.4	2
174	Enantiomeric separation of amphetamine and phenylephrine by capillary zone electrophoresis with hydroxypropyl- β -cyclodextrin as chiral additive. Journal of High Resolution Chromatography, 1997, 20, 298-299.	1.4	2
175	Chiral Separations of Enantiomeric Pharmaceuticals by Capillary Electrophoresis Using Sulphobutyl Ether β -Cyclodextrin as Isomer Selector. Biomedical Chromatography, 1997, 11, 193-199.	1.7	29
176	Separation of Chiral Amino Acids by Micellar Electrokinetic Chromatography with Derivatized Cyclodextrins. , 1997, 11, 230-235.		26
177	Phosphated cyclodextrins as new acidic chiral additives for capillary electrophoresis. Journal of Separation Science, 1997, 9, 581-589.	1.0	30
178	Capillary electrophoresis of herbicides II. Evaluation of alkylglucoside chiral surfactants in the enantiomeric separation of phenoxy acid herbicides. Journal of Chromatography A, 1997, 757, 263-273.	3.7	52
179	Effect of β -cyclodextrin acetylation on the resolution of phenethylamines with capillary electrophoresis and nuclear magnetic resonance spectroscopy. Journal of Chromatography A, 1997, 758, 277-292.	3.7	44
180	Chiral separation of monoterpenes using mixtures of sulfated β -cyclodextrins and β -cyclodextrin as chiral additives in the reversed-polarity capillary electrophoresis mode. Journal of Chromatography A, 1997, 759, 149-155.	3.7	66

#	ARTICLE	IF	CITATIONS
181	Enantioseparation of 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate-derivatized-amino acids by capillary zone electrophoresis using native and substituted β -cyclodextrins as chiral additives I. Discussion of optimum separation conditions. Journal of Chromatography A, 1997, 759, 157-165.	3.7	13
182	Chiral resolution of diols by capillary electrophoresis using borate-cyclodextrin complexation. Journal of Chromatography A, 1997, 761, 269-275.	3.7	42
183	Migration order reversal of enantiomers in capillary electrophoretic separation. Journal of Chromatography A, 1997, 764, 151-156.	3.7	20
184	Chiral separation of neutral species by capillary electrophoresis evaluation of a theoretical model. Journal of Chromatography A, 1997, 761, 249-257.	3.7	49
185	Mobility measurements on dansylated amino acids. Journal of Chromatography A, 1997, 768, 153-159.	3.7	28
186	Chiral recognition and racemic resolution of drug enantiomers by electrophoresis based on host-guest complexation. Journal of Chromatography A, 1997, 763, 261-269.	3.7	15
187	Vacancy affinity capillary electrophoresis, a new method for measuring association constants. Journal of Chromatography A, 1997, 775, 313-326.	3.7	55
188	Enantioseparation using selected polysaccharides as chiral buffer additives in capillary electrophoresis. Journal of Chromatography A, 1997, 773, 331-338.	3.7	47
189	Principles and limitations of methods available for the determination of binding constants with affinity capillary electrophoresis. Journal of Chromatography A, 1997, 777, 329-353.	3.7	107
190	Dry look at the CHARM (charged resolving agent migration) model of enantiomer separations by capillary electrophoresis. Journal of Chromatography A, 1997, 777, 295-309.	3.7	173
191	Prediction of the migration behavior of analytes in capillary electrophoresis based on three fundamental parameters. Journal of Chromatography A, 1997, 781, 23-34.	3.7	25
192	Enantiomeric separation of a tetrapeptide with cyclodextrin extension of the model for chiral capillary electrophoresis by complex formation of one enantiomer molecule with more than one chiral selector molecules. Journal of Chromatography A, 1997, 782, 271-279.	3.7	39
193	Achiral selectivity in cyclodextrin-modified capillary electrophoresis. Journal of Chromatography A, 1997, 792, 431-444.	3.7	48
194	Selectivity in capillary electrophoresis in the presence of micelles, chiral selectors and non-aqueous media. Journal of Chromatography A, 1997, 792, 13-35.	3.7	105
195	Separation selectivity in chiral capillary electrophoresis with charged selectors. Journal of Chromatography A, 1997, 792, 269-295.	3.7	251
196	Enantioselectivity in chiral capillary electrophoresis with polysaccharides. Journal of Chromatography A, 1997, 792, 327-347.	3.7	85
197	Application of cyclodextrin-modified micellar electrokinetic chromatography to the separations of selected neutral pesticides and their enantiomers. Journal of Chromatography A, 1997, 792, 419-429.	3.7	73
198	Controlling enantioselectivity in chiral capillary electrophoresis with inclusion complexation. Journal of Chromatography A, 1997, 792, 227-267.	3.7	225

#	ARTICLE	IF	CITATIONS
199	Chiral separation principles in capillary electrophoresis. <i>Journal of Chromatography A</i> , 1997, 792, 179-225.	3.7	237
200	Study on the chiral separation of basic drugs by capillary zone electrophoresis using β -CD and derivatized β -CDs as chiral selectors. <i>Analytica Chimica Acta</i> , 1998, 359, 39-46.	5.4	25
201	Chiral separations of indan, tetralin and benzosuberan derivatives by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1998, 793, 135-143.	3.7	14
202	Chiral separation of 3-phenyl-3-(2-pyridyl)propylamines, and analogous guanidines and guanidine-N-carboxylic acid esters with high-performance liquid chromatography and capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 1998, 793, 77-90.	3.7	11
203	Separation of enantiomers of drugs by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1998, 793, 153-164.	3.7	46
204	Enantiomeric separation of propranolol and selected metabolites by using capillary electrophoresis with hydroxypropyl- β -cyclodextrin as chiral selector. <i>Journal of Chromatography A</i> , 1998, 793, 357-364.	3.7	49
205	Cationic β -cyclodextrin derivative for chiral separations. <i>Journal of Chromatography A</i> , 1998, 803, 261-271.	3.7	65
206	New derivatives of cyclodextrins as chiral selectors for the capillary electrophoretic separation of dichlorprop enantiomers. <i>Journal of Chromatography A</i> , 1998, 810, 193-200.	3.7	16
207	Simultaneous separation of o-, m-, p-fluoro-dl-phenylalanine and o-, m-, p-dl-tyrosine by ligand-exchange micellar electrokinetic capillary chromatography. <i>Journal of Chromatography A</i> , 1998, 813, 369-378.	3.7	61
208	Chiral separations by capillary zone electrophoresis with the use of cyanoethylated- β -cyclodextrin as chiral selector. <i>Journal of Chromatography A</i> , 1998, 817, 91-104.	3.7	23
209	Effect of various cyclodextrin derivatives on the resolution of fencamfamine isomers with capillary electrophoresis and nuclear magnetic resonance. <i>Journal of Chromatography A</i> , 1998, 818, 239-249.	3.7	18
210	Fast chiral separation of amino acid derivatives and acidic drugs by co-electroosmotic flow capillary electrophoresis with vancomycin as chiral selector. <i>Journal of Chromatography A</i> , 1998, 825, 81-87.	3.7	43
211	Effect of additives on chiral selectivities by β -cyclodextrin in capillary electrophoresis based on the phenomenon of solvatochromism. <i>Journal of Chromatography A</i> , 1998, 829, 301-307.	3.7	6
212	Chiral separation of racemic mexiletine hydrochloride using cyclodextrins as chiral additive by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1998, 795, 394-398.	3.7	9
213	Comparative capillary electrophoresis and NMR studies of enantioseparation of dimethindene with cyclodextrins. <i>Journal of Chromatography A</i> , 1998, 798, 315-323.	3.7	86
214	Chiral resolution of functionalized piperidine enantiomers by capillary electrophoresis with native, alkylated and anionic β -cyclodextrins. <i>Journal of Chromatography A</i> , 1998, 796, 375-383.	3.7	20
215	Use of chiral and achiral ion-pairing reagents in combination with cyclodextrins in capillary electrophoresis. <i>Journal of Chromatography A</i> , 1998, 798, 281-288.	3.7	34
216	Use of cationic cyclodextrin for enantioseparation by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1998, 798, 275-280.	3.7	33

#	ARTICLE	IF	CITATIONS
217	Mixed-mode capillary electrokinetic separation of positional explosive isomers using sodium dodecyl sulfate and negative- β -cyclodextrin derivatives. <i>Journal of Chromatography A</i> , 1998, 811, 225-232.	3.7	28
218	Nonaqueous capillary electrophoretic separation of enantiomers using the single-isomer heptakis(2,3-diacetyl-6-sulfato)- β -cyclodextrin as chiral resolving agent. <i>Journal of Chromatography A</i> , 1998, 816, 233-241.	3.7	82
219	Affinity capillary electrophoresis: A physical-organic tool for studying interactions in biomolecular recognition. <i>Electrophoresis</i> , 1998, 19, 367-382.	2.4	178
220	Dynamic complexation of solutes in capillary electrophoresis. <i>Electrophoresis</i> , 1998, 19, 383-387.	2.4	15
221	The effects of a mixture of charged and neutral additives on analyte migration behavior in capillary electrophoresis. <i>Electrophoresis</i> , 1998, 19, 388-396.	2.4	24
222	Enantioseparation of anaesthetic drugs by capillary zone electrophoresis using cyclodextrin-containing background electrolytes. <i>Electrophoresis</i> , 1998, 19, 731-737.	2.4	18
223	Capillary electrophoresis chiral separations of basic compounds using cationic cyclodextrin. <i>Electrophoresis</i> , 1998, 19, 2095-2100.	2.4	30
224	Capillary electrophoresis, nuclear magnetic resonance and mass spectrometry studies of opposite chiral recognition of chlorpheniramine enantiomers with various cyclodextrins. <i>Electrophoresis</i> , 1998, 19, 2101-2108.	2.4	50
225	Separation of chiral compounds by capillary electrophoresis. <i>Electrophoresis</i> , 1998, 19, 2798-2833.	2.4	144
226	Use of cyclodextrins in capillary electrophoresis: Resolution of tramadol enantiomers. <i>Electrophoresis</i> , 1998, 19, 2883-2889.	2.4	24
227	Separation of Enantiomers of Drugs by Capillary Electrophoresis, Part 6. Hydroxypropyl- β -cyclodextrin as Chiral Solvating Agent. <i>Journal of High Resolution Chromatography</i> , 1998, 21, 215-224.	1.4	31
228	Chiral Separation of Bencynonate and Econazole by Cyclodextrin-Modified Capillary Zone Electrophoresis. <i>Journal of High Resolution Chromatography</i> , 1998, 21, 421-423.	1.4	10
229	Enantiomeric Separation of β -Amino Alcohols by Capillary Electrophoresis Using DM- β -CD as Chiral Selector. <i>Journal of High Resolution Chromatography</i> , 1998, 21, 440-444.	1.4	10
230	Influence of the Buffer Organic Cation on the Chiral Separation of Some Basic Drugs by Capillary Zone Electrophoresis. <i>Journal of High Resolution Chromatography</i> , 1998, 21, 445-449.	1.4	11
231	Use of heptakis-(2,3-O-dimethyl-6-sulfato)- β -cyclodextrin, a new single-isomer chiral resolving agent for the separation of the enantiomers of neutral compounds by capillary electrophoresis. <i>Journal of Separation Science</i> , 1998, 10, 293-299.	1.0	24
232	Chiral separation of chloroquine and pemoline by capillary zone electrophoresis with sulfobutyl ether β -cyclodextrin as buffer additive. <i>Chromatographia</i> , 1998, 48, 310-313.	1.3	14
233	The effect of conductivity tuning in chiral separations by CE; Using hydroxypropyl- β -cyclodextrin in combination with tetraalkylammonium ions. <i>Chromatographia</i> , 1998, 48, 415-421.	1.3	15
234	Enantiomeric separation of tramadol hydrochloride and its metabolites by cyclodextrin-mediated capillary zone electrophoresis. <i>Biomedical Applications</i> , 1998, 707, 287-294.	1.7	32

#	ARTICLE	IF	CITATIONS
235	Determination of chiral pharmaceutical compounds, terbutaline, ketamine and propranolol, by on-line capillary electrophoresis-electrospray ionization mass spectrometry. Biomedical Applications, 1998, 714, 69-75.	1.7	55
236	Use of mobility ratios to estimate binding constants of ligands to proteins in affinity capillary electrophoresis. Biomedical Applications, 1998, 715, 203-210.	1.7	61
237	Quantitative relationship between retention in HPLC and CE via binding constants and selector concentrations. TrAC - Trends in Analytical Chemistry, 1998, 17, 148-157.	11.4	8
238	Separation of enantiomers of drugs by capillary electrophoresis Part 8. β -Cyclodextrin as chiral solvating agent. Talanta, 1998, 46, 743-749.	5.5	20
239	Capillary Electrophoresis with Electrochemical Detection for Chiral Separation of Optical Isomers. Analytical Chemistry, 1998, 70, 4030-4035.	6.5	16
240	Control of relative migration of small inorganic and organic anions with cyclodextrins in capillary electrophoresis (CE). Canadian Journal of Chemistry, 1998, 76, 194-198.	1.1	8
241	Enantiomeric Separations of Terbutaline by CE with a Sulfated β -Cyclodextrin Chiral Selector: A Quantitative Binding Study. Analytical Chemistry, 1998, 70, 5166-5171.	6.5	101
242	Monte Carlo Simulation of Error Propagation in the Determination of Binding Constants from Rectangular Hyperbolae. 1. Ligand Concentration Range and Binding Constant. Journal of Physical Chemistry A, 1998, 102, 8063-8071.	2.5	68
243	Enantioselective Separations. Journal of Chromatography Library, 1998, 60, 197-256.	0.1	3
244	Enantiomeric resolution of galanthamine and related drugs used in anti-Alzheimer therapy by means of capillary zone electrophoresis employing derivatized cyclodextrin selectors. Biomedical Applications, 1999, 730, 167-175.	1.7	28
245	Non-aqueous capillary electrophoresis chiral separations with sulfated β -cyclodextrin. Biomedical Applications, 1999, 731, 187-197.	1.7	54
246	Recent trends in enantioseparations using capillary electromigration techniques. TrAC - Trends in Analytical Chemistry, 1999, 18, 485-498.	11.4	65
247	Modelling of conditions for the enantiomeric separation of β -adrenergic sympathicomimetics by capillary electrophoresis using cyclodextrins as chiral selectors in a polyethylene glycol gel. Journal of Pharmaceutical and Biomedical Analysis, 1999, 19, 529-537.	2.8	19
248	Cyclodextrin-mediated micellar electrokinetic chromatography and matrix-assisted laser desorption/ionization time-of-flight mass spectrometry for the enantiomer separation of racemorphan in human urine. Journal of Chromatography A, 1999, 853, 439-447.	3.7	24
249	Rapid development of the enantiomeric separation of β -blockers by capillary electrophoresis using an experimental design approach. Journal of Chromatography A, 1999, 855, 681-693.	3.7	52
250	Separation of the enantiomers of four chiral drugs by neutral cyclodextrin-mediated capillary zone electrophoresis. Chromatographia, 1999, 49, 411-414.	1.3	9
251	Principle and applications of the partial filling technique in capillary electrophoresis. Chromatographia, 1999, 50, 497-506.	1.3	86
252	Chiral separation and determination of the optical purity of thiamphenicol-related chiral compound by capillary zone electrophoresis with cyclodextrins as chiral selectors. Chromatographia, 1999, 50, 317-320.	1.3	13

#	ARTICLE	IF	CITATIONS
253	Separation of chiral basic drugs with sulfobutyl- β -cyclodextrin in capillary electrophoresis. Chromatographia, 1999, 50, 363-368.	1.3	14
254	Riboflavin binding proteins as chiral selectors in HPLC and CE. Pharmaceutical Science & Technology Today, 1999, 2, 352-364.	0.7	8
255	Capillary Electrophoresis as an Efficient Tool in the Determination of Low Association Constants between Polyhydroxy Compound and Samarium Cation. Journal of High Resolution Chromatography, 1999, 22, 595-598.	1.4	2
256	Estimation of the pH-independent binding constants of alanylphenylalanine and leucylphenylalanine stereoisomers with β -cyclodextrin in the presence of urea. Electrophoresis, 1999, 20, 171-179.	2.4	60
257	Determination of association constants between enantiomers of orciprenaline and methyl- β -cyclodextrin as chiral selector by capillary zone electrophoresis using a partial filling technique. Electrophoresis, 1999, 20, 180-188.	2.4	31
258	Enantioseparation of tropa alkaloids by means of anionic cyclodextrin-modified capillary electrophoresis. Electrophoresis, 1999, 20, 1555-1560.	2.4	25
259	Separation of drugs by capillary electrophoresis, Part 10. Permethyl-alpha-cyclodextrin as chiral solvating agent. Electrophoresis, 1999, 20, 1878-1889.	2.4	20
260	Capillary electrophoresis enantioseparation of drugs using β -cyclodextrin polymer: Intramolecular synergistic effect. Electrophoresis, 1999, 20, 1900-1903.	2.4	6
261	Optical isomer separation of potential analgesic drug candidates by using capillary electrophoresis. Electrophoresis, 1999, 20, 2432-2437.	2.4	8
262	Considerations concerning interaction characterization of oligopeptide mixtures with vancomycin using affinity capillary electrophoresis-electrospray mass spectrometry. Electrophoresis, 1999, 20, 2462-2474.	2.4	33
263	Selector-selectand interactions in chiral capillary electrophoresis. Electrophoresis, 1999, 20, 2592-2604.	2.4	41
264	Comparison of sulfobutylether- and sulfated- β -cyclodextrins as additives for the chiral separation of basic spirobenzopyrans by capillary electrophoresis. Electrophoresis, 1999, 20, 2630-2637.	2.4	17
265	Application of sulfobutylether- β -cyclodextrin with specific degrees of substitution for the enantioseparation of pharmaceutical mixtures by capillary electrophoresis. Electrophoresis, 1999, 20, 2638-2649.	2.4	42
266	Binding constant dependency of amphetamines with various commercial methylated β -cyclodextrins. Electrophoresis, 1999, 20, 2670-2679.	2.4	30
267	Designed combination of chiral selectors for adjustment of enantioseparation selectivity in capillary electrophoresis. Electrophoresis, 1999, 20, 2691-2697.	2.4	88
268	Resolution of ephedrine derivatives by means of neutral and sulfatedheptakis(2,3-di-O-acetyl) β -cyclodextrins using capillary electrophoresis and nuclear magnetic resonance spectroscopy. Electrophoresis, 1999, 20, 2698-2704.	2.4	44
269	pKa shift-associated effects in enantioseparations by cyclodextrin-mediated capillary zone electrophoresis. Electrophoresis, 1999, 20, 2715-2722.	2.4	91
270	Equilibrium binding model of bile salt-mediated chiral micellar electrokinetic capillary chromatography. Electrophoresis, 1999, 20, 2754-2760.	2.4	9

#	ARTICLE	IF	CITATIONS
271	Enantioseparations by capillary electrophoresis using chiral glycosidic surfactants. I. Evaluation of cyclohexyl-pentyl- β -D-maltoside surfactant. Electrophoresis, 1999, 20, 2766-2771.	2.4	9
272	Selectivity in capillary electrokinetic separations. Electrophoresis, 1999, 20, 2989-3010.	2.4	38
273	Identification, quantitation, and characterization of biomolecules by capillary electrophoretic analysis of binding interactions. Electrophoresis, 1999, 20, 3122-3133.	2.4	148
274	Enantiomeric separation of glycyl dipeptides by capillary electrophoresis with cyclodextrins as chiral selectors. Electrophoresis, 1999, 20, 3417-3423.	2.4	21
275	Enantiomeric separation of fluoxetine and norfluoxetine in plasma and serum samples with high detection sensitivity capillary electrophoresis. Electrophoresis, 1999, 20, 3432-3438.	2.4	67
276	Enantiomeric resolution of some nonsteroidal antiinflammatory and anticoagulant drugs using β -cyclodextrins by capillary electrophoresis. Chirality, 1999, 11, 56-62.	2.6	33
277	Resolution of enantiomeric N-oxides by capillary electrophoresis using cyclodextrins as chiral selectors. Chirality, 1999, 11, 409-415.	2.6	13
278	Capillary electrophoresis chiral separation of basic pharmaceutical enantiomers with different charges using sulfated β -cyclodextrin. Journal of Separation Science, 1999, 11, 11-21.	1.0	31
279	Enantiomer separations of phenylephrine and synephrine by capillary electrochromatography on bare silica stationary phase using hydroxypropyl- β -cyclodextrin as a mobile phase additive. Journal of Separation Science, 1999, 11, 263-269.	1.0	30
280	Separation behavior of amino acid enantiomers in ligand exchange micellar electrokinetic chromatography. Journal of Separation Science, 1999, 11, 534-540.	1.0	29
281	Direct chiral resolution of aliphatic α -hydroxy acids using 2-hydroxypropyl- β -cyclodextrin in capillary electrophoresis. Analyst, The, 1999, 124, 55-59.	3.5	27
282	Symmetry Breaking during the Formation of β -Cyclodextrin~Imidazole Inclusion Compounds: A Capillary Electrophoresis Study. Analytical Chemistry, 1999, 71, 2046-2052.	6.5	36
283	Self-Association of Rifamycin B: A Possible Effects on Molecular Recognition. Journal of Physical Chemistry B, 1999, 103, 4338-4341.	2.6	6
284	Monte Carlo Simulation of Error Propagation in the Determination of Binding Constants from Rectangular Hyperbolae. 2. Effect of the Maximum-Response Range. Journal of Physical Chemistry A, 1999, 103, 197-202.	2.5	60
285	Fast Analytical-Scale Separations by Capillary Electrophoresis and Liquid Chromatography. Chemical Reviews, 1999, 99, 3081-3132.	47.7	68
286	Reversal Behaviors of the Enantiomer Migration Order and the Stereo-selectivity of Cu(II) Complex with Amino Acid Enantiomers in Ligand Exchange-Micellar Electrokinetic Chromatography.. Analytical Sciences, 2000, 16, 131-137.	1.6	41
288	High-speed chiral separations on microchip electrophoresis devices. Electrophoresis, 2000, 21, 211-219.	2.4	104
289	Capillary electrophoresis behavior of water-soluble anionic porphyrins in the presence of β -cyclodextrin and its O-methylated derivatives. Electrophoresis, 2000, 21, 619-626.	2.4	14

#	ARTICLE	IF	CITATIONS
290	Cellulases from the fungi <i>Phanerochaete chrysosporium</i> and <i>Trichoderma reesei</i> as chiral selectors in capillary electrophoresis: Applications with displacer plugs and sample preconcentration. <i>Electrophoresis</i> , 2000, 21, 1587-1596.	2.4	19
291	Simultaneous chiral separation of triadimefon and triadimenol by sulfated β -cyclodextrin-mediated capillary electrophoresis. <i>Electrophoresis</i> , 2000, 21, 1611-1619.	2.4	39
292	Experimental verification of a predicted, hitherto unseen separation selectivity pattern in the nonaqueous capillary electrophoretic separation of weak base enantiomers by octakis(2,3-diacetyl-6-sulfato)- β -cyclodextrin. <i>Electrophoresis</i> , 2000, 21, 2016-2024.	2.4	45
293	Enantiomeric separation of alanyl and leucyl dipeptides by capillary electrophoresis with cyclodextrins as chiral selectors. <i>Electrophoresis</i> , 2000, 21, 2397-2404.	2.4	21
294	Recent innovations in the use of charged cyclodextrins in capillary electrophoresis for chiral separations in pharmaceutical analysis. <i>Electrophoresis</i> , 2000, 21, 3220-3239.	2.4	156
295	Fast enantiomeric separation of uniconazole and diniconazole by electrokinetic chromatography using an anionic cyclodextrin: Application to the determination of analyte-selector apparent binding constants for enantiomers. <i>Electrophoresis</i> , 2000, 21, 3240-3248.	2.4	23
296	Experimental verification of a predicted, previously unseen separation selectivity pattern in the capillary electrophoretic separation of noncharged enantiomers by octakis(2,3-diacetyl-6-sulfato)- β -cyclodextrin. <i>Electrophoresis</i> , 2000, 21, 3249-3256.	2.4	16
297	Competitive binding of a Tris run buffer with chiral crown ether in chiral capillary electrophoresis. <i>Electrophoresis</i> , 2000, 21, 3618-3624.	2.4	17
298	Separation of Enantiomers of Drugs by Capillary Electrophoresis with Permethy- γ -Cyclodextrin as Chiral Solvating Agent. <i>Journal of High Resolution Chromatography</i> , 2000, 23, 413-429.	1.4	26
299	Enantiomeric separations by use of polymeric surfactant electrokinetic chromatography. <i>Journal of Chromatography A</i> , 2000, 875, 179-206.	3.7	70
300	Enantiomeric separations of drugs using mixtures of charged and neutral cyclodextrins. <i>Journal of Chromatography A</i> , 2000, 875, 123-134.	3.7	170
301	Enantiomeric separations by nonaqueous capillary electrophoresis. <i>Journal of Chromatography A</i> , 2000, 875, 277-293.	3.7	92
302	Calculation of stability constants for the chiral selector-enantiomer interactions from electrophoretic mobilities. <i>Journal of Chromatography A</i> , 2000, 875, 431-445.	3.7	50
303	Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2000, 875, 207-233.	3.7	45
304	Optimization of the cyclodextrin-assisted capillary electrophoresis separation of the enantiomers of phenoxyacid herbicides. <i>Journal of Chromatography A</i> , 2000, 875, 423-430.	3.7	28
305	tert-Butylcarbamoylequinine as chiral ion-pair agent in non-aqueous enantioselective capillary electrophoresis applying the partial filling technique. <i>Journal of Chromatography A</i> , 2000, 892, 509-521.	3.7	42
306	New pseudo-stationary phases for electrokinetic capillary chromatography. <i>Journal of Chromatography A</i> , 2000, 876, 169-182.	3.7	5
307	Analysis of polymyxin B sulfate by capillary zone electrophoresis with cyclodextrin as additive. <i>Journal of Chromatography A</i> , 2000, 879, 211-218.	3.7	35

#	ARTICLE	IF	CITATIONS
308	Fast development of separation methods for the chiral analysis of amino acid derivatives using capillary electrophoresis and experimental designs. <i>Journal of Chromatography A</i> , 2000, 883, 249-265.	3.7	42
309	Development and evaluation of a linear regression method for the prediction of maximal chiral separation of basic drug racemates by cyclodextrin-mediated capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 2000, 903, 245-254.	3.7	24
310	Direct chiral resolution of lactic acid in food products by capillary electrophoresis. <i>Journal of Chromatography A</i> , 2000, 875, 371-377.	3.7	47
311	Chiral capillary electrophoretic analysis of the enantiomeric purity of a pharmaceutical compound using sulfated β -cyclodextrin. <i>Journal of Chromatography A</i> , 2000, 875, 389-401.	3.7	47
312	Separation of drug enantiomers by capillary electrophoresis in the presence of neutral cyclodextrins. <i>Journal of Chromatography A</i> , 2000, 875, 135-161.	3.7	71
313	Design of experiments for capillary electrophoretic enantioresolution of salbutamol using dermatan sulfate. <i>Journal of Chromatography A</i> , 2000, 875, 411-422.	3.7	58
314	Enantioselective determination by capillary electrophoresis with cyclodextrins as chiral selectors. <i>Journal of Chromatography A</i> , 2000, 875, 89-122.	3.7	444
315	Enantiomeric separation of selenoaminoacid derivatives by cyclodextrin-modified micellar electrokinetic chromatography using a mixed micellar system of sodium dodecyl sulphate and taurodeoxycholic acid. <i>Analytica Chimica Acta</i> , 2000, 416, 1-7.	5.4	34
316	New derivative of β -cyclodextrin as chiral selectors for the capillary electrophoretic separation of chiral drugs. <i>Analytica Chimica Acta</i> , 2000, 421, 27-34.	5.4	26
317	Advanced statistical evaluation of complex formation constant from electrophoretic data. <i>Analytica Chimica Acta</i> , 2000, 407, 327-336.	5.4	10
318	Dependence of chiral separations on the amount of cyclodextrins as selectors, employing the partial filling technique in capillary zone electrophoresis. <i>Chromatographia</i> , 2000, 51, 226-230.	1.3	13
319	Selectivity difference between carboxyethyl- β -cyclodextrin and carboxymethyl- β -cyclodextrin in enantioseparations of drugs by capillary electrophoresis. <i>Chromatographia</i> , 2000, 51, 634-637.	1.3	2
320	Macrocyclic antibiotics as effective chiral selectors for enantiomeric resolution by liquid chromatography and capillary electrophoresis. <i>Chromatographia</i> , 2000, 52, 679-691.	1.3	108
321	Permethyl monoamino β -cyclodextrin a new chiral selective agent for capillary electrophoresis. <i>Chromatographia</i> , 2000, 53, 166-172.	1.3	17
322	CHIRAL SEPARATIONS Capillary Electrophoresis. , 2000, , 2287-2294.		1
323	Capillary Zone Electrophoresis. , 2000, , 139-208.		12
324	Capillary Zone Electrophoresis. , 2000, , 73-138.		7
325	Chiral Separations in Capillary Electrophoresis. <i>Chemical Reviews</i> , 2000, 100, 3715-3754.	47.7	168

#	ARTICLE	IF	CITATIONS
326	6.3. Capillary electrophoretic (CE) methods. Progress in Pharmaceutical and Biomedical Analysis, 2000, 4, 540-552.	0.1	0
327	Enantioseparation Of Aromatic Dipeptides Using Carboxymethyl- β -Cyclodextrin Polymer As Chiral Selector By Capillary Electrophoresis. Analytical Letters, 2000, 33, 1147-1165.	1.8	8
328	Recent advances in enantiomer separations by affinity capillary electrophoresis using proteins and peptides. Journal of Proteomics, 2001, 48, 103-116.	2.4	72
329	Quantitative determination of clenbuterol, salbutamol and tulobuterol enantiomers by capillary electrophoresis. Fresenius' Journal of Analytical Chemistry, 2001, 369, 212-219.	1.5	25
330	Enantioseparation of chiral phenothiazine derivatives in capillary electrophoresis using cyclodextrin type chiral selectors. Chromatographia, 2001, 53, S290-S295.	1.3	34
331	Chiral separation of newly synthesized arylpropionic acids by capillary electrophoresis using cyclodextrins or a glycopeptide antibiotic as chiral selectors. Chromatographia, 2001, 54, 87-92.	1.3	10
332	Modelling enantiomer separation by CE. Chromatographia, 2001, 54, S24-S41.	1.3	0
333	Comparison of binding of tetraphenylborate and tetraphenylphosphonium ions to cyclodextrins studied by capillary electrophoresis. Electrophoresis, 2001, 22, 117-122.	2.4	13
334	Chiral analysis of baclofen by β -cyclodextrin-modified capillary electrophoresis and laser-induced fluorescence detection. Electrophoresis, 2001, 22, 123-127.	2.4	39
335	Influence of neutral cyclodextrin concentration on plate numbers in capillary electrophoresis. Electrophoresis, 2001, 22, 1957-1973.	2.4	6
336	Modeling and optimization of the chiral selectivity of basic analytes in chiral capillary electrophoresis with negatively charged cyclodextrins using electrochemical detection. Electrophoresis, 2001, 22, 2025-2031.	2.4	15
337	Fundamental aspects of chiral separations by capillary electrophoresis. Electrophoresis, 2001, 22, 3079-3106.	2.4	215
338	Recent developments in chiral capillary electrophoresis and applications of this technique to pharmaceutical and biomedical analysis. Electrophoresis, 2001, 22, 3107-3130.	2.4	191
339	pH-Dependence of complexation constants and complex mobility in capillary electrophoresis separations of dipeptide enantiomers. Electrophoresis, 2001, 22, 3163-3170.	2.4	49
340	Rapid screening for chiral separations by short-end injection capillary electrophoresis using highly sulfated cyclodextrins as chiral selectors. Electrophoresis, 2001, 22, 3203-3215.	2.4	88
341	Chiral separation of pyrethroic acids with single isomer permethyl monoamino β -cyclodextrin selector. Electrophoresis, 2001, 22, 3232-3236.	2.4	22
342	Enantioseparation of chiral thiobarbiturates using cyclodextrin-modified capillary electrophoresis. Electrophoresis, 2001, 22, 3237-3242.	2.4	8
343	Experimental designs to investigate capillary electrophoresis-electrospray ionization-mass spectrometry enantioseparation with the partial-filling technique. Electrophoresis, 2001, 22, 3316-3326.	2.4	74

#	ARTICLE	IF	CITATIONS
344	Cyclodextrin-assisted capillary electrophoretic resolution of 1,1'-bi-2-naphthol atropisomers. <i>Electrophoresis</i> , 2001, 22, 3578-3582.	2.4	6
345	Interaction between 18-crown-6-tetracarboxylic acid and positional substituents of enantiomers and simultaneous separation of positional enantiomers of methyl-DL-tryptophans by capillary electrophoresis. <i>Electrophoresis</i> , 2001, 22, 2136-2142.	2.4	11
346	Electrophoretic chiral separation of pharmaceutical compounds with multiple stereogenic centers in charged cyclodextrin media. <i>Journal of Separation Science</i> , 2001, 24, 378-384.	2.5	15
347	Chiral separation of promethazine by capillary electrophoresis with end-column amperometric detection. <i>Journal of Separation Science</i> , 2001, 24, 658-662.	2.5	12
348	Evaluation of enantioselective nonaqueous ion-pair capillary electrophoresis as screening assay in the development of new ion exchange type chiral stationary phases. <i>Journal of Separation Science</i> , 2001, 24, 706-716.	2.5	21
349	High-performance chiral separation of fourteen triazole fungicides by sulfated β -cyclodextrin-mediated capillary electrophoresis. <i>Journal of Chromatography A</i> , 2001, 912, 171-179.	3.7	112
350	Determination of apparent binding constants of drugs by capillary electrophoresis using β -cyclodextrin as ligand and three different linear plotting methods. <i>Journal of Chromatography A</i> , 2001, 924, 483-491.	3.7	31
351	Comparison of monomeric and polymeric amino acid based surfactants for chiral separations. <i>Journal of Chromatography A</i> , 2001, 922, 329-338.	3.7	46
352	Non-aqueous capillary electrophoretic separation of enantiomeric amines with (α)-2,3:4,6-di-O-isopropylidene-2-keto-L-gulonic acid as chiral counter ion. <i>Journal of Chromatography A</i> , 2001, 922, 303-311.	3.7	39
353	Mechanistic study of enantiomeric recognition of primary amino compounds using an achiral crown ether with cyclodextrin by capillary electrophoresis and nuclear magnetic resonance. <i>Journal of Chromatography A</i> , 2001, 923, 229-239.	3.7	18
354	Chiral separation of six diastereomeric flavanone-7-O-glycosides by capillary electrophoresis and analysis of lemon juice. <i>Journal of Chromatography A</i> , 2001, 925, 279-289.	3.7	48
355	Rapid chiral on-chip separation with simplified amperometric detection. <i>Journal of Chromatography A</i> , 2001, 928, 225-232.	3.7	65
356	Enantioseparations in capillary electromigration techniques: recent developments and future trends. <i>Journal of Chromatography A</i> , 2001, 906, 309-363.	3.7	331
357	Enantioseparation of flobufen with cyclodextrins studied by capillary electrophoresis and NMR. <i>Pharmaceutical Research</i> , 2001, 18, 886-893.	3.5	8
358	Optimization Strategies and Modeling of Separations of Dansyl-Amino Acids by Cyclodextrin-Modified Capillary Electrophoresis. <i>Journal of Chromatographic Science</i> , 2001, 39, 411-419.	1.4	10
359	ENANTIOMERIC SEPARATION OF A DRUG SUBSTANCE USING CAPILLARY ELECTROPHORESIS WITH SULFATED- β -CYCLODEXTRIN. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2002, 25, 2999-3015.	1.0	4
360	Enantioseparation of Vinclozolin by β -Cyclodextrin-Modified Micellar Electrokinetic Chromatography. <i>Journal of Agricultural and Food Chemistry</i> , 2002, 50, 1312-1317.	5.2	27
361	Accurately describing weak analyte-additive interactions by capillary electrophoresis. <i>Electrophoresis</i> , 2002, 23, 880-888.	2.4	21

#	ARTICLE	IF	CITATIONS
362	Drug-plasma protein binding assay by electrokinetic chromatography-frontal analysis. Electrophoresis, 2002, 23, 951-955.	2.4	29
363	Chiral capillary electrophoretic determination of the enantiomeric purity of tetrahydronaphthalenic derivatives, melatoninergic ligands, using highly sulfated β -cyclodextrins. Electrophoresis, 2002, 23, 1121-1128.	2.4	20
364	Chiral recognition of functionalized cyclodextrins in capillary electrophoresis. Electrophoresis, 2002, 23, 1308-1313.	2.4	19
365	Modelling, optimisation and control of selectivity in the separation of aromatic bases by electrokinetic chromatography using a neutral cyclodextrin as a pseudostationary phase. Electrophoresis, 2002, 23, 1844.	2.4	8
366	Separation of reboxetine enantiomers by means of capillary electrophoresis. Electrophoresis, 2002, 23, 1870.	2.4	24
367	Electrolyte and additive effects on enantiomer separation of peptides by nonaqueous ion-pair capillary electrophoresis using tert.-butylcarbamoylquinine as chiral counterion. Electrophoresis, 2002, 23, 1887.	2.4	38
368	Combinatorial enantiomeric separation of diverse compounds using capillary array electrophoresis. Electrophoresis, 2002, 23, 2996-3005.	2.4	35
369	Enantioselective analysis of thiobencarb sulfoxide produced by metabolism of thiobencarb by hydroxypropyl- β -cyclodextrin modified micellar electrokinetic chromatography. Journal of Separation Science, 2002, 25, 1055-1062.	2.5	18
370	Enantiomer migration order in chiral capillary electrophoresis. Electrophoresis, 2002, 23, 4022-4035.	2.4	132
371	Optimized separation of pharmacologically active xanthenes from <i>Securidaca inappendiculata</i> by capillary electrophoresis. Chromatographia, 2002, 55, 217-223.	1.3	16
372	Knowledge-based system for method development of chiral separations with capillary electrophoresis using highly-sulphated cyclodextrins. Journal of Pharmaceutical and Biomedical Analysis, 2002, 27, 515-529.	2.8	52
373	Use of negatively charged cyclodextrins for the simultaneous enantioseparation of selected anesthetic drugs by capillary electrophoresis-mass spectrometry. Journal of Pharmaceutical and Biomedical Analysis, 2002, 27, 615-626.	2.8	45
374	Selected fundamental aspects of chiral electromigration techniques and their application to pharmaceutical and biomedical analysis. Journal of Pharmaceutical and Biomedical Analysis, 2002, 27, 373-399.	2.8	128
375	Chiral separation of methoxamine and lobeline in capillary zone electrophoresis using ethylbenzene-deactivated fused-silica capillary columns and cyclodextrins as buffer additives. Journal of Pharmaceutical and Biomedical Analysis, 2002, 29, 999-1003.	2.8	14
376	Estimation of binding constants by capillary electrophoresis. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2002, 768, 81-92.	2.3	198
377	Chiral separation of amino acids derivatized with fluoresceine-5-isothiocyanate by capillary electrophoresis and laser-induced fluorescence detection using mixed selectors of β -cyclodextrin and sodium taurocholate. Journal of Chromatography A, 2002, 955, 133-140.	3.7	76
378	Quantitative approach for the screening of cyclodextrins by nuclear magnetic resonance spectroscopy in support of chiral separations in liquid chromatography and capillary electrophoresis. Journal of Chromatography A, 2002, 961, 257-276.	3.7	30
379	Separation and migration behavior of structurally related phenothiazines in cyclodextrin-modified capillary zone electrophoresis. Journal of Chromatography A, 2002, 979, 399-408.	3.7	38

#	ARTICLE	IF	CITATIONS
380	Chiral separation of methadone, 2-ethylidene- 1,5-dimethyl-3,3-diphenylpyrrolidine (EDDP) and 2-ethyl-5-methyl-3,3-diphenyl-1-pyrroline (EMDP) by capillary electrophoresis using cyclodextrin derivatives. <i>Electrophoresis</i> , 2003, 24, 2106-2110.	2.4	16
381	Salt dependence on vancomycin-herbicide enantiomer binding: Capillary electrophoresis study and theoretical approach. <i>Electrophoresis</i> , 2003, 24, 1620-1626.	2.4	11
382	Direct chiral resolution of tartaric acid by ion-pair capillary electrophoresis using an aqueous background electrolyte with (1R,2R)-($\alpha\epsilon^6$)-1,2-diaminocyclohexane as a chiral counterion. <i>Electrophoresis</i> , 2003, 24, 2711-2715.	2.4	11
383	pH-dependent cyclodextrin capillary electrophoresis resolution of atropisomers. <i>Electrophoresis</i> , 2003, 24, 2456-2461.	2.4	12
384	Large-volume sample stacking combined with separation by 2-hydroxypropyl- β -cyclodextrin for analysis of isoxazolylpenicillins by capillary electrophoresis. <i>Electrophoresis</i> , 2003, 24, 3089-3096.	2.4	21
385	Complexation trends and binding constants between dextrin oligomers and small molecules as measured through affinity capillary electrophoresis. <i>Electrophoresis</i> , 2003, 24, 2914-2923.	2.4	8
386	Recent advances in enantioseparations of peptides by capillary electrophoresis. <i>Electrophoresis</i> , 2003, 24, 4063-4077.	2.4	51
387	Applications of affinity interactions in capillary electrophoresis. <i>Electrophoresis</i> , 2003, 24, 3879-3891.	2.4	83
388	Migration order of dipeptide and tripeptide enantiomers in the presence of single isomer and randomly sulfated cyclodextrins as a function of pH. <i>Electrophoresis</i> , 2003, 24, 1069-1076.	2.4	48
389	A high-throughput on-line microdialysis-capillary assay for D-serine. <i>Electrophoresis</i> , 2003, 24, 1227-1235.	2.4	52
390	Separation of isomers of dienolic acids by electromigration techniques. <i>Journal of Separation Science</i> , 2003, 26, 1253-1258.	2.5	7
391	Enantioseparation of dansyl amino acids on terguride-based chiral selectors. Part I: Capillary electrophoretic separation. <i>Journal of Separation Science</i> , 2003, 26, 851-856.	2.5	11
392	Separation of diastereomers of flavanone-7-O-glycosides by capillary electrophoresis using sulfobutyl ether- β -cyclodextrin as the selector. <i>Journal of Separation Science</i> , 2003, 26, 844-850.	2.5	30
393	Enantiomeric separation of aminoglutethimide by capillary electrophoresis using native cyclodextrins in single and dual systems. <i>Journal of Separation Science</i> , 2003, 26, 536-542.	2.5	13
394	Chiral separation of 2,3-allenoic acid by capillary zone electrophoresis using cyclodextrin derivatives. <i>Chirality</i> , 2003, 15, 201-205.	2.6	5
395	Comprehensive strategy for chiral separations using sulfated cyclodextrins in capillary electrophoresis. <i>Chirality</i> , 2003, 15, 709-723.	2.6	107
396	Mechanistic study of enantiomeric recognition with native β -cyclodextrin by capillary electrophoresis, reversed-phase liquid chromatography, nuclear magnetic resonance spectroscopy, electrospray mass spectrometry and circular dichroism techniques. <i>Journal of Chromatography A</i> , 2003, 987, 409-420.	3.7	37
397	Determination of free l- and d-alanine in hydrolysed protein fertilisers by capillary electrophoresis. <i>Journal of Chromatography A</i> , 2003, 985, 463-469.	3.7	17

#	ARTICLE	IF	CITATIONS
398	Exploration of the electrophoretic behaviour of borane cluster anions and of the capability of capillary electrophoresis to separate them chirally. <i>Journal of Chromatography A</i> , 2003, 984, 121-134.	3.7	17
399	Robustness testing of chiral separations by capillary electrophoresis using highly-sulfated cyclodextrins. <i>Journal of Chromatography A</i> , 2003, 1007, 165-177.	3.7	40
400	Chiral separation of dioxopromethazine in eye drops by CZE with charged cyclodextrin. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2003, 33, 157-164.	2.8	17
401	Simultaneous enantioselective separation of azelastine and three of its metabolites for the investigation of the enantiomeric metabolism in rats. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2003, 793, 389-404.	2.3	21
402	Enantioseparation using cyclosophoraoses as a novel chiral additive in capillary electrophoresis. <i>Carbohydrate Research</i> , 2003, 338, 1143-1146.	2.3	39
403	Indirect capillary electrophoresis with 8-anilino-1-naphthalenesulfonic acid as a fluorescence probe for determining the apparent stability constant of an inclusion complex formed between a cyclodextrin and a solute. <i>Journal of Chromatography A</i> , 2003, 987, 485-492.	3.7	5
404	Chiral separation of amines with N-benzoxycarbonylglycyl-L-proline as selector in non-aqueous capillary electrophoresis using methanol and 1,2-dichloroethane in the background electrolyte. <i>Journal of Chromatography A</i> , 2003, 984, 261-271.	3.7	29
405	Determination of the binding constant for the inclusion complex between procaine hydrochloride and β -cyclodextrin by capillary electrophoresis. <i>Talanta</i> , 2003, 59, 493-499.	5.5	24
406	Capillary Electrophoretic Behaviors of Pharmacologically Active Xanthenes from <i>Securidaca inappendiculata</i> with α -Cyclodextrin as a Buffer Additive. <i>Journal of Chromatographic Science</i> , 2003, 41, 182-186.	1.4	7
407	Use of cyclodextrins for separation of enantiomers. <i>Russian Chemical Reviews</i> , 2003, 72, 1035-1054.	6.5	32
408	Chiral Separation by Capillary Electrophoresis Using Polysaccharides. , 2004, 243, 343-354.		3
409	Chiral Separation of Deprenyl-N-Oxide Isomers by Capillary Electrophoresis Using Various Cyclodextrin Derivatives. <i>Journal of Chromatographic Science</i> , 2004, 42, 21-26.	1.4	10
411	Determination of equilibrium constants from chromatographic and electrophoretic measurements. <i>Journal of Chromatography A</i> , 2004, 1037, 15-28.	3.7	50
412	Separation and migration behavior of positional and structural naphthalenesulfonate isomers by cyclodextrin-mediated capillary electrophoresis. <i>Journal of Chromatography A</i> , 2004, 1033, 167-172.	3.7	20
413	Enantioseparation of four cis and trans diastereomers of 2 α ,3 α -didehydro-2 α ,3 α -dideoxythymidine analogs, by high-performance liquid chromatography and capillary electrophoresis. <i>Journal of Chromatography A</i> , 2004, 1034, 161-167.	3.7	17
414	Enatiomeric analysis of simendan by CE with β -CD as chiral selector compared with CMPA-HPLC. <i>Biomedical Chromatography</i> , 2004, 18, 866-871.	1.7	8
415	Enantioseparation of new nucleoside analogs, related to d4T and acyclovir, by chiral capillary electrophoresis using highly sulfated β -cyclodextrins. <i>Electrophoresis</i> , 2004, 25, 444-453.	2.4	15
416	Optimisation of selectivity in the separation of aromatic amino acid enantiomers using sulfated γ -cyclodextrin and dextran sulfate as pseudostationary phases. <i>Electrophoresis</i> , 2004, 25, 270-276.	2.4	16

#	ARTICLE	IF	CITATIONS
417	Chiral capillary electrophoretic resolution of baclofen, gabaergic ligand, using highly sulfated cyclodextrins. Electrophoresis, 2004, 25, 1111-1119.	2.4	18
418	Direct enantioseparation of catechin and epicatechin in tea drinks by 6-O- β -D-glucosyl- β -cyclodextrin-modified micellar electrokinetic chromatography. Electrophoresis, 2004, 25, 2892-2898.	2.4	44
419	Chiral separation of N-imidazole derivatives, aromatase inhibitors, by cyclodextrin-capillary zone electrophoresis. Mechanism of enantioselective recognition. Electrophoresis, 2004, 25, 2735-2744.	2.4	25
420	Influence of methanol on the enantioresolution of antihistamines with carboxymethyl- β -cyclodextrin in capillary electrophoresis. Electrophoresis, 2004, 25, 2838-2847.	2.4	20
421	Development and validation of a capillary electrophoresis method for the enantiomeric purity determination of SLV307, a basic potential antipsychotic compound. Electrophoresis, 2004, 25, 2854-2859.	2.4	10
422	Influence of (hydroxy)alkylamino substituents on enantioseparation ability of single-isomer amino- β -cyclodextrin derivatives in chiral capillary electrophoresis. Electrophoresis, 2004, 25, 2675-2686.	2.4	45
423	Ultrashort partial-filling technique in capillary electrophoresis for infinite resolution of tramadol enantiomers and its metabolites with highly sulfated cyclodextrins. Electrophoresis, 2004, 25, 2761-2771.	2.4	33
424	Differential effects of organic modifiers on the enantioseparation of dimetindene maleate with carboxymethyl- β -cyclodextrin in capillary electrophoresis. Journal of Separation Science, 2004, 27, 21-27.	2.5	14
425	Selectivity control in the separation of aromatic amino acid enantiomers with sulphated β -cyclodextrin. Journal of Chromatography A, 2004, 1031, 179-186.	3.7	9
426	Direct separation, identification and quantification of epimers 22R and 22S of budesonide by capillary gas chromatography on a short analytical column with Rtx [®] -5 stationary phase. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2004, 803, 191-200.	2.3	7
427	Evaluation of newly synthesized derivative of cyclodextrin for the capillary electrophoretic separation. Journal of Chromatography A, 2004, 1059, 181-189.	3.7	21
428	Enantiomeric separations of some acidic compounds with cationic cyclodextrin by capillary electrophoresis. Analytica Chimica Acta, 2004, 517, 95-101.	5.4	20
429	Enantiomeric analysis of rivastigmine in pharmaceuticals by cyclodextrin-modified capillary zone electrophoresis. Analytica Chimica Acta, 2004, 525, 43-51.	5.4	20
430	Effect of Different Substituted Position of Phenylcarbamoyl- β -CD on the Enantioseparation of Drugs by Capillary Electrophoresis. Analytical Letters, 2004, 37, 293-306.	1.8	1
431	Combined approach using capillary electrophoresis and NMR spectroscopy for an understanding of enantioselective recognition mechanisms by cyclodextrins. Chemical Society Reviews, 2004, 33, 337.	38.1	166
432	Analysis of Ion-Association Reaction in Aqueous Solution and Its Utilization by Capillary Zone Electrophoresis. Analytical Sciences, 2004, 20, 255-265.	1.6	22
433	2-O-(2-Hydroxybutyl)-.BETA.-cyclodextrin as a Chiral Selector for the Capillary Electrophoretic Separation of Chiral Drugs. Analytical Sciences, 2005, 21, 959-962.	1.6	18
434	Simultaneous Chiral Analysis of Methamphetamine and Related Compounds by Capillary Electrophoresis/Mass Spectrometry Using Anionic Cyclodextrin. Analytical Sciences, 2005, 21, 15-19.	1.6	44

#	ARTICLE	IF	CITATIONS
435	Rapid Enantioseparation of 1-Aminoindan by Microchip Electrophoresis with Linear-Imaging UV Detection. <i>Analytical Sciences</i> , 2005, 21, 61-65.	1.6	43
436	Enantioselective capillary electrophoretic separation of tryptophane- and tyrosine-methylesters in a dual system with a tetra-oxadiazia-crown-ether derivative and a cyclodextrin. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2005, 38, 601-608.	2.8	33
437	Chiral separation of tamsulosin by capillary electrophoresis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2005, 39, 691-696.	2.8	30
438	Enantioseparation of dansyl amino acids by a novel permanently positively charged single-isomer cyclodextrin: Mono-6-N-allylammonium-6-deoxy- β -cyclodextrin chloride by capillary electrophoresis. <i>Analytica Chimica Acta</i> , 2005, 546, 119-125.	5.4	28
439	Enantiomeric separation of 8 hydroxy, 10 carboxylic and 6 dansyl amino acids by mono(6-amino-6-deoxy)- β -cyclodextrin in capillary electrophoresis. <i>Analytica Chimica Acta</i> , 2005, 554, 156-162.	5.4	28
440	Affinity capillary electrophoresis on microchips. <i>Journal of Chromatography A</i> , 2005, 1063, 217-225.	3.7	42
441	Chiral separations by capillary electromigration techniques in nonaqueous media. <i>Journal of Chromatography A</i> , 2005, 1068, 3-30.	3.7	79
442	Comparison of vancomycin-based stationary phases with different chiral selector coverage for enantioselective separation of selected drugs in high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2005, 1088, 94-103.	3.7	55
443	Synthesis and application of mono-6-ammonium-6-deoxy- β -cyclodextrin chloride as chiral selector for capillary electrophoresis. <i>Journal of Chromatography A</i> , 2005, 1094, 187-191.	3.7	35
444	Separation and determination of norepinephrine, epinephrine and isoprenaline enantiomers by capillary electrophoresis in pharmaceutical formulation and human serum. <i>Journal of Chromatography A</i> , 2005, 1098, 166-171.	3.7	129
445	Analysis of Bioactive Triterpenes from <i>Rubus chingii</i> by Cyclodextrin-Modified Capillary Electrophoresis. <i>Chromatographia</i> , 2005, 62, 145-150.	1.3	20
446	Approach to method development and validation in capillary electrophoresis for enantiomeric purity testing of active basic pharmaceutical ingredients. <i>Electrophoresis</i> , 2005, 26, 2330-2341.	2.4	25
447	The formation of cucurbit[n]uril (n = 6, 7) complexes with amino compounds in aqueous formic acid studied by capillary electrophoresis. <i>Electrophoresis</i> , 2005, 26, 2214-2224.	2.4	26
448	Enantiomeric analysis of baclofen analogs by capillary zone electrophoresis, using highly sulfated cyclodextrins: Inclusion ionization constant pKa determination. <i>Electrophoresis</i> , 2005, 26, 2974-2983.	2.4	20
449	A family of single-isomer positively charged cyclodextrins as chiral selectors for capillary electrophoresis: Mono-6A-butylammonium-6A-deoxy- β -cyclodextrin tosylate. <i>Electrophoresis</i> , 2005, 26, 3125-3133.	2.4	35
450	Chiral separation in capillary electrophoresis using dual neutral cyclodextrins: Theoretical models of electrophoretic mobility difference and separation selectivity. <i>Electrophoresis</i> , 2005, 26, 3814-3823.	2.4	29
451	Enantioseparation of chiral N-imidazole derivatives by electrokinetic chromatography using highly sulfated cyclodextrins: Mechanism of enantioselective recognition. <i>Electrophoresis</i> , 2005, 26, 3824-3832.	2.4	15
452	Aptamer-oligonucleotide binding studied by capillary electrophoresis: Cation effect and separation efficiency. <i>Electrophoresis</i> , 2005, 26, 3247-3255.	2.4	25

#	ARTICLE	IF	CITATIONS
453	Synthesis and application of single-isomer 6-mono(alkylimidazolium)- β -cyclodextrins as chiral selectors in chiral capillary electrophoresis. <i>Electrophoresis</i> , 2005, 26, 3839-3848.	2.4	55
454	Enantioseparation in capillary electrophoresis using 2-O-(2-hydroxybutyl)- β -CD as a chiral selector. <i>Electrophoresis</i> , 2005, 26, 3890-3896.	2.4	14
455	Application of cyclodextrin-mediated capillary electrophoresis to determine the apparent binding constants and thermodynamic parameters of the alkylnaphthalene derivatives. <i>Electrophoresis</i> , 2005, 26, 3528-3537.	2.4	16
456	Influence of steric hindrance on enantioseparation of Dns-amino acids and pesticides on terguride based chiral selectors in capillary electrophoresis. <i>Journal of Separation Science</i> , 2005, 28, 673-677.	2.5	8
457	Enantioselective analysis of cetirizine in pharmaceuticals by cyclodextrin-mediated capillary electrophoresis. <i>Journal of Separation Science</i> , 2005, 28, 1278-1284.	2.5	28
458	18 Chiral separations. <i>Separation Science and Technology</i> , 2005, 6, 447-498.	0.2	9
459	Application of cyclodextrin-mediated capillary electrophoresis to simultaneously determine the apparent binding constants and thermodynamic parameters of naphthalenesulfonate derivatives. <i>Talanta</i> , 2006, 68, 1252-1258.	5.5	12
460	Principles of Enantiomer Separations in Electrokinetic Chromatography. , 0, , 179-206.		1
462	Chiral Capillary Electrophoresis of Agrochemicals in Real Samples. <i>Journal of Health Science</i> , 2006, 52, 489-494.	0.9	17
463	Chiral Capillary Electrophoresis of Amphetamine-Type Stimulants. <i>Journal of Health Science</i> , 2006, 52, 649-654.	0.9	11
464	Chiral Separation of Synthetic Vicinal Diol Compounds by Capillary Zone Electrophoresis with Borate Buffer and .BETA.-Cyclodextrin as Buffer Additive. <i>Analytical Sciences</i> , 2006, 22, 747-751.	1.6	5
465	Enantioselective separation in capillary electrophoresis using a novel mono-6A-propylammonium- β -cyclodextrin as chiral selector. <i>Analytica Chimica Acta</i> , 2006, 555, 63-67.	5.4	32
466	Determination of five anthraquinones in medicinal plants by capillary zone electrophoresis with β -cyclodextrin addition. <i>Journal of Chromatography A</i> , 2006, 1123, 134-137.	3.7	39
467	Chiral separation of four fluoroquinolone compounds using capillary electrophoresis with hydroxypropyl- β -cyclodextrin as chiral selector. <i>Journal of Chromatography A</i> , 2006, 1130, 296-301.	3.7	64
468	Enantioseparation of cis and trans nucleosides, aromatic analogues of stavudine, by capillary electrophoresis and high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2006, 1132, 141-147.	3.7	10
469	Enantiomeric separation of synthetic 2,3-dihydroxy-3-phenylpropionate compounds by β -cyclodextrin-modified capillary electrophoresis. <i>Journal of Chromatography A</i> , 2006, 1108, 258-262.	3.7	9
470	NMR studies of chiral recognition mechanisms: interaction of enantiomers of N-imidazole derivatives with cyclodextrin hosts. Correlation with the CD-EKC studies. <i>Tetrahedron: Asymmetry</i> , 2006, 17, 975-983.	1.8	12
471	Twenty Years of Separation of Cis-Trans (Z)-E Isomers. <i>Separation and Purification Reviews</i> , 2006, 35, 193-221.	5.5	10

#	ARTICLE	IF	CITATIONS
472	Determination of Ephedrine, Pseudo-Ephedrine and Caffeine in a Dietary Product by Capillary Electrophoresis. <i>Chromatographia</i> , 2006, 63, 143-148.	1.3	40
473	An observation of unusual temperature effects for enantioselective CZE employing highly sulfated- β -cyclodextrin. <i>Electrophoresis</i> , 2006, 27, 859-864.	2.4	9
474	Chiral separations by capillary electrophoresis: Recent developments and applications. <i>Electrophoresis</i> , 2006, 27, 2880-2895.	2.4	102
475	Enantiomeric separation of acidic compounds using single-isomer amino cyclodextrin derivatives in nonaqueous capillary electrophoresis. <i>Electrophoresis</i> , 2006, 27, 3434-3442.	2.4	31
476	Enantiomeric separation of some demethylated analogues of clofibric acid by capillary zone electrophoresis and nano-liquid chromatography. <i>Electrophoresis</i> , 2006, 27, 1227-1236.	2.4	17
477	Optimization of capillary electrophoretic enantioseparation for basic drugs with native β -CD as a chiral selector. <i>Electrophoresis</i> , 2006, 27, 2367-2375.	2.4	16
478	Chiral capillary electrophoretic determination of the enantiomeric purity of homocamptothecin derivatives, promising antitumor topoisomerase I inhibitors, using highly sulfated CDs and fluorescence detection. <i>Electrophoresis</i> , 2006, 27, 4717-4729.	2.4	10
479	Effect of alkali metal hydroxides on the enantioseparation of amines using di-O-isopropylidene-keto-L-gulonic acid as the selector in NACE. <i>Electrophoresis</i> , 2006, 27, 4469-4479.	2.4	10
480	Chiral separations by capillary electrophoresis. , 2006, , 241-275.		7
481	Enantiomeric Separation of Synthetic Amino Acids Using Capillary Zone Electrophoresis. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 30, 1421-1436.	1.0	13
482	Separation of 2-HPA by Capillary Electrophoresis. , 2007, , .		0
483	Enantioseparation of Isoxanthohumol in Beer by Hydroxypropyl- β -cyclodextrin-Modified Micellar Electrokinetic Chromatography. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 6547-6552.	5.2	13
484	Enantiomeric Separation of a Series of β -Lactams Using Capillary Zone Electrophoresis. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 30, 1709-1721.	1.0	12
485	Determination of Ofloxacin Enantiomers in Pharmaceutical Formulations by Capillary Electrophoresis. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 31, 348-360.	1.0	20
486	Investigation of β -CD-derivatized erythromycin as chiral selector in CE. <i>Electrophoresis</i> , 2007, 28, 2566-2572.	2.4	14
487	Chiral resolution of melatonineric ligands by EKC using highly sulfated CDs. <i>Electrophoresis</i> , 2007, 28, 3915-3921.	2.4	5
488	Stereoisomeric separation of pharmaceutical compounds using CE with a chiral crown ether. <i>Electrophoresis</i> , 2007, 28, 2658-2666.	2.4	31
489	Review of aqueous chiral electrokinetic chromatography (EKC) with an emphasis on chiral microemulsion EKC. <i>Electrophoresis</i> , 2007, 28, 2503-2526.	2.4	39

#	ARTICLE	IF	CITATIONS
490	Synthesis and NMR characterization of β -alanine-bridged hemispherodextrin, a very efficient chiral selector in EKC. <i>Electrophoresis</i> , 2007, 28, 2580-2588.	2.4	13
491	Chiral separability of hydrophobic boron cluster anions with native cyclodextrins in water-methanol background electrolytes. <i>Electrophoresis</i> , 2007, 28, 3639-3649.	2.4	12
492	Enantioselective analysis of the antipsychotic 9-hydroxyrisperidone, main metabolite of risperidone, by chiral capillary EKC using dual CDs. <i>Electrophoresis</i> , 2007, 28, 2683-2692.	2.4	19
493	Temperature effects on the enantioselectivity of basic analytes in capillary EKC using sulfated β -CDs as chiral selectors. <i>Electrophoresis</i> , 2007, 28, 3753-3758.	2.4	14
494	Peak splitting in the CE separation of enantiomers caused by organic solvents in the sample. <i>Electrophoresis</i> , 2007, 28, 3625-3638.	2.4	4
495	Electromigration of a heteroconjugated imidazole-acetate complex in ACN. <i>Electrophoresis</i> , 2007, 28, 3590-3599.	2.4	10
496	Simultaneous enantioseparation and sensitivity enhancement of basic drugs using large-volume sample stacking. <i>Electrophoresis</i> , 2007, 28, 3542-3552.	2.4	24
497	Complexation of tauro- and glyco-conjugated bile salts with three neutral β -CDs studied by ACE. <i>Electrophoresis</i> , 2007, 28, 3745-3752.	2.4	28
498	Chiral separation of dansyl amino acids in capillary electrophoresis using mono-(3-methyl-imidazolium)- β -cyclodextrin chloride as selector. <i>Journal of Separation Science</i> , 2007, 30, 1343-1349.	2.5	48
499	Ketopinic acid and diisopropylideneketogulonic acid as chiral ion-pair selectors in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2007, 1141, 287-294.	3.7	29
500	Influence of substituents on selectivity and efficiency of chiral separations of anions containing single nido-7,8-dicarbaundecaborane cluster with β -cyclodextrin. <i>Journal of Chromatography A</i> , 2007, 1143, 143-152.	3.7	8
501	Enantioseparations by using capillary electrophoretic techniques. <i>Journal of Chromatography A</i> , 2007, 1168, 45-70.	3.7	234
502	Effect of alkylimidazolium substituents on enantioseparation ability of single-isomer alkylimidazolium- β -cyclodextrin derivatives in capillary electrophoresis. <i>Analytica Chimica Acta</i> , 2007, 585, 227-233.	5.4	40
503	Kapillarelektrophorese, eine attraktive Methode zur Enantiomerenanalytik. <i>Pharmazie in Unserer Zeit</i> , 1999, 28, 186-196.	0.0	5
504	Characterization of the complexation of tauro- and glyco-conjugated bile salts with β -cyclodextrin and 2-hydroxypropyl- β -cyclodextrin using affinity capillary electrophoresis. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2008, 61, 161-169.	1.6	20
505	Cyclodextrins in capillary electrophoresis enantioseparations – Recent developments and applications. <i>Journal of Separation Science</i> , 2008, 31, 1991-2011.	2.5	146
506	Simultaneous enantioseparation of four β -agonists by capillary electrophoresis with cyclodextrin additives. Study of the enantioselective mechanism. <i>Journal of Separation Science</i> , 2008, 31, 3749-3754.	2.5	16
507	Monosubstituted positively charged cyclodextrins: Synthesis and applications in chiral separation. <i>Journal of Separation Science</i> , 2008, 31, 3246-3256.	2.5	57

#	ARTICLE	IF	CITATIONS
508	Synthesis of chiral vicinal diols and analysis of them by capillary zone electrophoresis. Chirality, 2008, 20, 75-83.	2.6	12
509	Recent advances in amino acid analysis by CE. Electrophoresis, 2008, 29, 207-223.	2.4	43
510	The role of cyclodextrins in chiral capillary electrophoresis. Electrophoresis, 2008, 29, 1701-1712.	2.4	116
511	Study of interaction between drug enantiomers and human serum albumin by flow injection capillary electrophoresis frontal analysis. Electrophoresis, 2008, 29, 2876-2883.	2.4	21
512	Enantiomeric separation of some flavanones using shinorhizobial linear octasaccharides in CE. Electrophoresis, 2008, 29, 4284-4290.	2.4	20
513	Enantiomeric separation of glycidyl tosylate by CE: Application to the study of catalytic asymmetric epoxidation of allyl alcohol. Electrophoresis, 2008, 29, 4575-4582.	2.4	5
515	Model of CE enantioseparation systems with a mixture of chiral selectors. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2008, 875, 30-34.	2.3	46
516	Stereoselective determination of trihexyphenidyl using carboxymethyl- β -cyclodextrin by capillary electrophoresis with field-amplified sample stacking. Microchemical Journal, 2008, 89, 34-41.	4.5	20
517	Possibilities of column coupling electrophoresis provided with a fiber-based diode array detection in enantioselective analysis of drugs in pharmaceutical and clinical samples. Journal of Chromatography A, 2008, 1179, 9-16.	3.7	21
518	Study of the complexation of risperidone and 9-hydroxyrisperidone with cyclodextrin hosts using affinity capillary electrophoresis and ^1H NMR spectroscopy. Journal of Chromatography A, 2008, 1215, 185-193.	3.7	38
519	18 Coupling CE and microchip-based devices with mass spectrometry. Separation Science and Technology, 2008, 9, 477-521.	0.2	9
520	Sensitivity improvement by using contactless conductivity rather than indirect UV detection for the determination of enantiomeric purity of amines by CE. Electrophoresis, 2009, 30, 487-498.	2.4	26
521	Enantioseparation of nicotine alkaloids in cigarettes by CE using sulfated β -CD as a chiral selector and a capillary coated with amino groups. Electrophoresis, 2009, 30, 349-356.	2.4	26
522	Chiral separations by CE employing CDs. Electrophoresis, 2009, 30, S203-10.	2.4	118
523	Affinity in Electrophoresis. Electrophoresis, 2009, 30, S229-39.	2.4	57
524	About the role of enantioselective selector-selectand interactions and the mobilities of diastereomeric associates in enantiomer separations using CE. Electrophoresis, 2009, 30, 2803-2811.	2.4	66
525	Use of coated capillaries for the electrophoretic separation of stereoisomers of a growth hormone secretagogue. Electrophoresis, 2009, 30, 3772-3779.	2.4	41
526	Separation of enantiomers with charged chiral selectors in CE. Electrophoresis, 2009, 30, S211-21.	2.4	116

#	ARTICLE	IF	CITATIONS
527	Advanced CE for chiral analysis of drugs, metabolites, and biomarkers in biological samples. Electrophoresis, 2009, 30, 2773-2802.	2.4	80
528	Enantioselective separation of chiral vicinal diols in capillary electrophoresis using a mono- α -aminoethylamino- β -cyclodextrin as a chiral selector. Journal of Separation Science, 2009, 32, 125-134.	2.5	11
529	Enantioseparation of chiral benzimidazole derivatives by electrokinetic chromatography using sulfated cyclodextrins. Journal of Separation Science, 2009, 32, 1907-1915.	2.5	11
530	Use of correction factors in mobility shift affinity capillary electrophoresis for weak analyte-ligand interactions. Journal of Separation Science, 2009, 32, 1712-1721.	2.5	27
531	Enantioselective analysis of primaquine and its impurity quinocide by capillary electrophoresis. Biomedical Chromatography, 2009, 23, 295-301.	1.7	10
532	6-O-(2-hydroxybutyl)- β -CD as a chiral selector for nonaqueous capillary electrophoretic separation of chiral drugs. Frontiers of Chemistry in China: Selected Publications From Chinese Universities, 2009, 4, 396-401.	0.4	3
533	A study of the interaction between enantiomers of zolmitriptan and hydroxypropyl- β -cyclodextrin by capillary electrophoresis. Analytical and Bioanalytical Chemistry, 2009, 393, 313-320.	3.7	18
534	Assessment of the complexation degree of camptothecin derivatives and cyclodextrins using spectroscopic and separative methodologies. Tetrahedron: Asymmetry, 2009, 20, 2482-2489.	1.8	13
535	The 3-amino-derivative of β -cyclodextrin as chiral selector of Dns-amino acids in electrokinetic chromatography. Journal of Chromatography A, 2009, 1216, 3678-3686.	3.7	17
536	Enantiomeric Resolution of Benzoxazolinonic Aminoalcohols, Potential Adrenergic Ligands, by LC and CE. Chromatographia, 2009, 69, 465-472.	1.3	4
537	Enantioselective Separation of Basic Drugs by CE with Polygalacturonic Acid as a Novel Chiral Selector. Chromatographia, 2009, 69, 1315-1320.	1.3	18
538	Enantioseparation of Dioxopromethazine Hydrochloride in Urine with Liquid-Liquid Extraction by CE-ECL Detection. Chromatographia, 2009, 70, 1291-1293.	1.3	12
539	Effect of β -Cyclodextrin on Drug Distribution Studied by Electrochemistry at Interfaces between Immiscible Electrolyte Solutions. Journal of Physical Chemistry B, 2009, 113, 7263-7269.	2.6	12
540	Enantioseparation of binaphthol and its monoderivatives by cyclodextrin-modified capillary zone electrophoresis: A mathematical approach. Talanta, 2009, 78, 631-637.	5.5	24
541	Simultaneous determination of R-(+)-, S-(-)-baclofen and impurity A by electrokinetic chromatography. Journal of Pharmaceutical and Biomedical Analysis, 2010, 51, 541-548.	2.8	19
542	Separation of vinca alkaloid enantiomers by capillary electrophoresis applying cyclodextrin derivatives and characterization of cyclodextrin complexes by nuclear magnetic resonance spectroscopy. Journal of Pharmaceutical and Biomedical Analysis, 2010, 53, 1258-1266.	2.8	29
543	Enantiomeric separation of epinephrine and salbutamol by micellar electrokinetic chromatography using β -cyclodextrin as chiral additive. Chinese Journal of Chemistry, 2010, 22, 845-848.	4.9	0
544	Glycogen: A novel branched polysaccharide chiral selector in CE. Electrophoresis, 2010, 31, 1044-1050.	2.4	34

#	ARTICLE	IF	CITATIONS
545	Enhanced selectivity in CZE multi- β -cyclodextrin chiral selector enantioseparation systems: Proposed separation mechanism. <i>Electrophoresis</i> , 2010, 31, 1435-1441.	2.4	54
546	Determination of the binding constants of modafinil enantiomers with sulfated β -cyclodextrin chiral selector by capillary electrophoresis using three different linear plotting methods. <i>Electrophoresis</i> , 2010, 31, 2957-2963.	2.4	17
548	Synthesis and application of a novel single isomer mono- β -deoxy- β -CD((2 <i>S</i> ,3 <i>S</i>)-(+)-2,3-epoxy-1,4-bis(isopropylideneamino)- β -cyclodextrin) as chiral selector in capillary electrophoresis. <i>Chirality</i> , 2010, 22, 914-921.	2.6	29
549	Capillary electrophoretic and nuclear magnetic resonance studies on the opposite affinity pattern of propranolol enantiomers towards various cyclodextrins. <i>Journal of Separation Science</i> , 2010, 33, 1617-1624.	2.5	52
550	Separation and characterization of modified pregabalins in terms of cyclodextrin complexation, using capillary electrophoresis and nuclear magnetic resonance. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2010, 51, 842-852.	2.8	51
551	Study of a new chiral selector: Sodium arsenyl-(l)-(+)-tartrate for capillary electrophoresis. <i>Journal of Chromatography A</i> , 2010, 1217, 1139-1148.	3.7	10
552	Application of charged single isomer derivatives of cyclodextrins in capillary electrophoresis for chiral analysis. <i>Journal of Chromatography A</i> , 2010, 1217, 953-967.	3.7	88
553	Chiral Recognition and Enantioseparation Mechanisms in Capillary Electrokinetic Chromatography. , 2010, , 97-152.		1
555	Chiral Separations. <i>Annual Review of Analytical Chemistry</i> , 2010, 3, 341-363.	5.4	109
556	Chiral Recognition in Separation Methods. , 2010, , .		81
557	Microchip Electrophoresis Using Linear-Imaging UV Detector. <i>Bunseki Kagaku</i> , 2011, 60, 725-734.	0.2	0
558	Complexation of triptolide and its succinate derivative with cyclodextrins: Affinity capillary electrophoresis, isothermal titration calorimetry and ¹ H NMR studies. <i>Journal of Chromatography A</i> , 2011, 1218, 8708-8714.	3.7	9
559	Enantioseparation of the New Antifungal Drug Iodiconazole and Structurally Related Triadimenol Analogues by CE with Neutral Cyclodextrin Additives. <i>Chromatographia</i> , 2011, 73, 1009-1014.	1.3	13
560	Chiral electromigration techniques in pharmaceutical and biomedical analysis. <i>Bioanalytical Reviews</i> , 2011, 3, 95-114.	0.2	11
561	Complexation of tauro- and glyco-conjugated bile salts with β -cyclodextrin and hydroxypropyl- β -cyclodextrin studied by affinity capillary electrophoresis and molecular modelling. <i>Journal of Separation Science</i> , 2011, 34, 3221-3230.	2.5	17
562	Accuracy and sensitivity of the determination of rate constants of interconversion in achiral and chiral environments by dynamic enantioselective electrophoresis. <i>Electrophoresis</i> , 2011, 32, 595-603.	2.4	6
563	Chiral separation and quantitation of cetirizine and hydroxyzine by maltodextrin-mediated CE in human plasma: Effect of zwitterionic property of cetirizine on enantioseparation. <i>Electrophoresis</i> , 2011, 32, 764-771.	2.4	31
564	Evaluation of the interaction between sitagliptin and cyclodextrin derivatives by capillary electrophoresis and nuclear magnetic resonance spectroscopy. <i>Electrophoresis</i> , 2011, 32, 2648-2654.	2.4	23

#	ARTICLE	IF	CITATIONS
565	Emerging subject for chiral separation science: Cluster boron compounds. <i>Chirality</i> , 2011, 23, 307-319.	2.6	25
566	Fundamental aspects of chiral electromigration techniques and application in pharmaceutical and biomedical analysis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2011, 55, 688-701.	2.8	90
567	Molecular Modeling Study of Chiral Separation and Recognition Mechanism of Î²-Adrenergic Antagonists by Capillary Electrophoresis. <i>International Journal of Molecular Sciences</i> , 2012, 13, 710-725.	4.1	51
568	8.10 Chromatographic Separations and Analysis: Cyclodextrin Mediated HPLC, GC and CE Enantiomeric Separations. , 2012, , 177-199.		12
569	Comparison of separation performances of novel Î²-cyclodextrin-based chiral stationary phases in high-performance liquid chromatographic enantioseparation. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2012, 70, 71-76.	2.8	15
570	Chiral separation of amino acids derivatised with fluorescein isothiocyanate by single isomer derivatives 3-monodeoxy-3-monoamino-Î²- and Î³-cyclodextrins: the effect of the cavity size. <i>Journal of Chromatography A</i> , 2012, 1269, 360-365.	3.7	23
571	Monte Carlo simulation of the enantioseparation process. <i>Physica A: Statistical Mechanics and Its Applications</i> , 2012, 391, 4389-4396.	2.6	5
572	Direct enantioseparation of lipoic acid in dietary supplements by capillary electrophoresis using trimethylâ€¢cyclodextrin as a chiral selector. <i>Electrophoresis</i> , 2012, 33, 2441-2445.	2.4	26
573	Simulation of the effects of complexâ€¢formation equilibria in electrophoresis: III. Simultaneous effects of chiral selector concentration and background electrolyte pH. <i>Electrophoresis</i> , 2012, 33, 3012-3020.	2.4	22
574	Affinity capillary electrophoresis method for investigation of bile salts complexation with sulfobutyl etherâ€¢cyclodextrin. <i>Journal of Separation Science</i> , 2012, 35, 2764-2772.	2.5	11
575	Chiral electrokinetic chromatographyâ€¢electrospray ionizationâ€¢mass spectrometry using a double junction interface. <i>Electrophoresis</i> , 2012, 33, 546-551.	2.4	9
576	Dynamic highâ€¢resolution computer simulation of electrophoretic enantiomer separations with neutral cyclodextrins as chiral selectors. <i>Electrophoresis</i> , 2012, 33, 958-969.	2.4	31
577	Determination of stability constants of complexes of neutral analytes with charged cyclodextrins by affinity capillary electrophoresis. <i>Electrophoresis</i> , 2012, 33, 1032-1039.	2.4	34
578	Simulation of the effects of complexâ€¢formation equilibria in electrophoresis: II. Experimental verification. <i>Electrophoresis</i> , 2012, 33, 948-957.	2.4	43
579	Chiral recognition of imperanene enantiomers by various cyclodextrins: A capillary electrophoresis and ¹H NMR spectroscopy study. <i>Electrophoresis</i> , 2012, 33, 1458-1464.	2.4	30
580	Extraction of Phenylalanine Enantiomers by Aqueous Two Phase Systems Containing Combinatorial Chiral Selector. <i>Chinese Journal of Chemistry</i> , 2012, 30, 965-969.	4.9	13
581	Effect of electrolyzed reduced water on chiral theanine and polyphenols in tea. <i>Food Chemistry</i> , 2012, 134, 1761-1766.	8.2	8
582	Enantiomeric separation of (<i>R,S</i>)-tolterodine and (<i>R,S</i>)-methoxytolterodine with negatively charged cyclodextrins by capillary electrophoresis. <i>Journal of Separation Science</i> , 2013, 36, 1561-1567.	2.5	18

#	ARTICLE	IF	CITATIONS
583	Simultaneous determination of flavonoid analogs in <i>Scutellariae Barbatae Herba</i> by β -cyclodextrin and acetonitrile modified capillary zone electrophoresis. <i>Talanta</i> , 2013, 105, 393-402.	5.5	50
584	Modified Cyclodextrins for Chiral Separation. , 2013, , .		22
585	Complexation of Buffer Constituents with Neutral Complexation Agents: Part II. Practical Impact in Capillary Zone Electrophoresis. <i>Analytical Chemistry</i> , 2013, 85, 8526-8534.	6.5	30
586	Differentiation of Enantiomers I. <i>Topics in Current Chemistry</i> , 2013, , .	4.0	15
587	Applicability and limitations of affinity capillary electrophoresis and vacancy affinity capillary electrophoresis methods for determination of complexation constants. <i>Electrophoresis</i> , 2013, 34, 761-767.	2.4	54
588	Chiral selectors in $\langle \text{scp} \rangle \text{CE} \langle / \text{scp} \rangle$: Recent developments and applications. <i>Electrophoresis</i> , 2013, 34, 178-204.	2.4	98
589	Separation selectivity patterns of fully charged achiral compounds in capillary electrophoresis with a neutral cyclodextrin. <i>Journal of Separation Science</i> , 2013, 36, 325-333.	2.5	0
590	Recent advances in electrodriven enantioseparations. <i>Journal of Separation Science</i> , 2013, 36, 52-74.	2.5	109
591	Synthesis and application of a chiral ionic liquid functionalized β -cyclodextrin as a chiral selector in capillary electrophoresis. <i>Biomedical Chromatography</i> , 2013, 27, 1027-1033.	1.7	27
592	Evaluation of the enantioselectivity of carbon nanoparticles α -modified chiral separation systems using dextrin as chiral selector by capillary electrokinetic chromatography. <i>Electrophoresis</i> , 2013, 34, 1901-1907.	2.4	26
593	Differentiation of Enantiomers by Capillary Electrophoresis. <i>Topics in Current Chemistry</i> , 2013, 340, 209-275.	4.0	24
594	Native β -Cyclodextrins and Their Neutral Derivatives for Capillary Electrophoresis. , 2013, , 145-186.		0
595	Cationic Cyclodextrins for Capillary Electrophoresis. , 2013, , 225-258.		0
596	Application of Polymeric Surfactants in Chiral Micellar Electrokinetic Chromatography (CMEKC) and CMEKC Coupled to Mass Spectrometry. <i>Methods in Molecular Biology</i> , 2013, 970, 319-348.	0.9	8
597	Electrically-enhanced microextraction combined with maltodextrin-modified capillary electrophoresis for quantification of tolterodine enantiomers in biological samples. <i>Microchemical Journal</i> , 2013, 106, 186-193.	4.5	46
598	Evaluation of electroosmotic markers in aqueous and nonaqueous capillary electrophoresis. <i>Electrophoresis</i> , 2013, 34, 3252-3259.	2.4	20
599	Enantioseparation of Tr \ddot{A} ger's Base Derivatives by Capillary Electrophoresis Using Cyclodextrins as Chiral Selectors. <i>Chirality</i> , 2013, 25, 379-383.	2.6	6
600	New single isomer negatively charged β - $\langle \text{scp} \rangle$ -cyclodextrin $\langle / \text{scp} \rangle$ derivatives as chiral selectors in $\langle \text{scp} \rangle$ capillary electrophoresis $\langle / \text{scp} \rangle$. <i>Electrophoresis</i> , 2013, 34, 1232-1240.	2.4	14

#	ARTICLE	IF	CITATIONS
601	Physicochemical profiling of drug candidates using capillary-based techniques. <i>Journal of Drug Delivery Science and Technology</i> , 2013, 23, 333-345.	3.0	8
602	Determination of effective mobilities of EOF markers in BGE containing sulfated β -cyclodextrin by a two-detector method. <i>Electrophoresis</i> , 2013, 34, 768-776.	2.4	19
603	Recent Trends in Chiral Separations by Nano Liquid Chromatography and Nano Capillary Electrophoresis. <i>Current Chromatography</i> , 2014, 1, 81-89.	0.3	19
604	Evaluation of the chiral separation ability of single isomer cationic β -cyclodextrins in capillary electrophoresis. <i>Electrophoresis</i> , 2014, 35, 2744-2751.	2.4	17
605	Hydrophilic interaction electrokinetic chromatography using bio-based nanofillers. <i>Electrophoresis</i> , 2014, 35, 2229-2236.	2.4	3
606	Characterizing the interaction between enantiomers of eight psychoactive drugs and highly sulfated β -cyclodextrin by counter-current capillary electrophoresis. <i>Biomedical Chromatography</i> , 2014, 28, 120-126.	1.7	11
607	Velocity gap mode of capillary electrophoresis developed for high-resolution chiral separations. <i>Electrophoresis</i> , 2014, 35, 2778-2784.	2.4	7
608	Cyclodextrin-mediated enantioseparation of phenylalanine amide derivatives and amino alcohols by capillary electrophoresis—Role of complexation constants and complex mobilities. <i>Electrophoresis</i> , 2014, 35, 2848-2854.	2.4	16
609	Separation and Migration Behavior of Dichlorophenols in β -Cyclodextrin-Modified Capillary Zone Electrophoresis. <i>Journal of the Chinese Chemical Society</i> , 2014, 61, 1025-1030.	1.4	0
610	Separation efficiency of dual-selector systems in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2014, 1330, 82-88.	3.7	21
611	Effect of competing binding modes on retention in chromatography and capillary electrophoresis. A theoretical consideration. <i>Journal of Separation Science</i> , 2014, 37, 390-392.	2.5	3
612	Application of cyclodextrins in chiral capillary electrophoresis. <i>Electrophoresis</i> , 2014, 35, 2701-2721.	2.4	141
613	Antagonists of the P_{2X7} receptor: Mechanism of enantioselective recognition using highly sulfated and sulfobutylether cyclodextrins by capillary electrokinetic chromatography. <i>Electrophoresis</i> , 2014, 35, 2892-2899.	2.4	9
614	Enantioseparation of meptazinol and its three intermediate enantiomers by capillary electrophoresis using a new cationic β -cyclodextrin derivative in single and dual cyclodextrin systems. <i>Biomedical Chromatography</i> , 2014, 28, 868-874.	1.7	17
615	Penicillin G as a novel chiral selector in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2014, 1326, 134-138.	3.7	16
616	Dynamic computer simulation of electrophoretic enantiomer migration order and separation in presence of a neutral cyclodextrin. <i>Electrophoresis</i> , 2014, 35, 2833-2841.	2.4	22
617	Determination of thermodynamic values of acidic dissociation constants and complexation constants of profens and their utilization for optimization of separation conditions by Simul 5 Complex. <i>Journal of Chromatography A</i> , 2014, 1364, 276-288.	3.7	27
618	Twenty years of development of dual and multi-selector models in capillary electrophoresis: A review. <i>Electrophoresis</i> , 2014, 35, 2688-2700.	2.4	40

#	ARTICLE	IF	CITATIONS
619	Maltodextrins as Chiral Selectors in CE: Molecular Structure Effect of Basic Chiral Compounds on the Enantioseparation. <i>Chirality</i> , 2014, 26, 620-628.	2.6	12
620	Visceral mesh modified with cyclodextrin for the local sustained delivery of ropivacaine. <i>International Journal of Pharmaceutics</i> , 2014, 476, 149-159.	5.2	25
621	Advances in chiral separations of small peptides by capillary electrophoresis and chromatography. <i>Journal of Separation Science</i> , 2014, 37, 2447-2466.	2.5	121
622	Novel methoxypropylimidazolium β^2 -cyclodextrin for improved enantioseparation of amino acids. <i>Talanta</i> , 2014, 128, 460-465.	5.5	17
624	Separation of Folinic Acid Diastereomers in Capillary Electrophoresis Using a New Cationic β^2 -Cyclodextrin Derivative. <i>PLoS ONE</i> , 2015, 10, e0120216.	2.5	7
625	Chiral Resolution Capabilities of DNA Oligonucleotides. <i>Analytical Chemistry</i> , 2015, 87, 5491-5495.	6.5	42
626	Synthesis of 2,3-O-dibenzyl-6-O-sulfobutyl- β and β^2 cyclodextrins: new chiral surfactants for capillary electrophoresis. <i>Tetrahedron Letters</i> , 2015, 56, 4451-4454.	1.4	13
627	Generalized model of electromigration with 1:1 (analyte:selector) complexation stoichiometry: Part II. Application to dual systems and experimental verification. <i>Journal of Chromatography A</i> , 2015, 1384, 147-154.	3.7	12
628	Study of the enantioseparation capability of chiral dual system based on chondroitin sulfate C in CE. <i>Electrophoresis</i> , 2015, 36, 607-614.	2.4	9
629	Methoxypropylamino β^2 -cyclodextrin clicked AC regioisomer for enantioseparations in capillary electrophoresis. <i>Analytica Chimica Acta</i> , 2015, 868, 73-79.	5.4	15
630	Development of a capillary electrophoretic method for determination of plasma clearance of iohexol in dogs and cats. <i>Biomedical Chromatography</i> , 2015, 29, 504-513.	1.7	7
631	Chiral Separations in Capillary Electrophoresis. , 2015, , .		2
632	Development of Electrophoretic Methods for Simultaneous Determination of Enantiomeric Ratio and Composition of Diastereomeric Salt Mixtures. <i>Chromatographia</i> , 2015, 78, 881-888.	1.3	6
633	A dispersive liquid-liquid microextraction and chiral separation of carvedilol in human plasma using capillary electrophoresis. <i>Bioanalysis</i> , 2015, 7, 1107-1117.	1.5	27
634	Chiral selectors in CE: Recent developments and applications (2012â€mid 2014). <i>Electrophoresis</i> , 2015, 36, 101-123.	2.4	67
635	Evaluation of sulfated maltodextrin as a novel anionic chiral selector for the enantioseparation of basic chiral drugs by capillary electrophoresis. <i>Electrophoresis</i> , 2015, 36, 305-311.	2.4	27
636	Influence of substituent position and cavity size of the regioisomers of monocarboxymethyl- β , β^2 , and β^3 -cyclodextrins on the apparent stability constants of their complexes with both enantiomers of Tröger's base. <i>Journal of Separation Science</i> , 2016, 39, 980-985.	2.5	15
637	The empirical comparison of cyclofructans and cyclodextrins as chiral selectors in capillary electrophoretic separation of atropisomers of <i>1,1'-binaphthalene-2,2'-diyl hydrogen phosphate</i> . <i>Journal of Separation Science</i> , 2016, 39, 973-979.		10

#	ARTICLE	IF	CITATIONS
638	Development and validation of a cyclodextrin- β -modified capillary electrophoresis method for the enantiomeric separation of vildagliptin enantiomers. <i>Electrophoresis</i> , 2016, 37, 1318-1325.	2.4	23
639	Cyclodextrin-assisted enantioseparation of warfarin and 10-hydroxywarfarin by capillary electrophoresis studied from the analytical and thermodynamic points of view. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016, 126, 60-65.	2.8	8
640	Advances in the Use of Cyclodextrins as Chiral Selectors in Capillary Electrokinetic Chromatography: Fundamentals and Applications. <i>Chromatographia</i> , 2016, 79, 1403-1435.	1.3	101
641	Single-isomer carboxymethyl- β -cyclodextrin as chiral resolving agent for capillary electrophoresis. <i>Journal of Chromatography A</i> , 2016, 1467, 445-453.	3.7	34
642	Affinity capillary electrophoresis: the theory of electromigration. <i>Analytical and Bioanalytical Chemistry</i> , 2016, 408, 8623-8641.	3.7	60
643	Three chiral ionic liquids as additives for enantioseparation in capillary electrophoresis and their comparison with conventional modifiers. <i>Journal of Chromatography A</i> , 2016, 1462, 146-152.	3.7	43
644	Enantiomer Separations by Capillary Electrophoresis. <i>Methods in Molecular Biology</i> , 2016, 1483, 277-299.	0.9	5
645	Chiral Separation of Uncharged Pomalidomide Enantiomers Using Carboxymethyl- β -Cyclodextrin: A Validated Capillary Electrophoretic Method. <i>Chirality</i> , 2016, 28, 199-203.	2.6	19
646	Estimation of apparent binding constant of complexes of selected acyclic nucleoside phosphonates with β -cyclodextrin by affinity capillary electrophoresis. <i>Electrophoresis</i> , 2016, 37, 239-247.	2.4	33
647	Physicochemical characterisation and cyclodextrin complexation of erlotinib. <i>Supramolecular Chemistry</i> , 2016, 28, 656-664.	1.2	14
648	Chiral separation of asenapine enantiomers by capillary electrophoresis and characterization of cyclodextrin complexes by NMR spectroscopy, mass spectrometry and molecular modeling. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016, 117, 398-404.	2.8	47
649	Characterization of complexes between phenethylamine enantiomers and β -cyclodextrin derivatives by capillary electrophoresis—Determination of binding constants and complex mobilities. <i>Electrophoresis</i> , 2017, 38, 1188-1200.	2.4	12
650	Validated capillary electrophoretic method for the enantiomeric quality control of <i>R</i> -praziquantel. <i>Electrophoresis</i> , 2017, 38, 1886-1894.	2.4	14
651	Use of various β -cyclodextrin derivatives as chiral selectors for the enantiomeric separation of ofloxacin and its five related substances by capillary electrophoresis. <i>Journal of Separation Science</i> , 2017, 40, 1784-1795.	2.5	21
652	Sequence requirements of oligonucleotide chiral selectors for the capillary electrophoresis resolution of low-affinity DNA binders. <i>Electrophoresis</i> , 2017, 38, 1383-1390.	2.4	14
653	Single and dual cyclodextrins systems for the enantiomeric and diastereoisomeric separations of structurally related dihydropyridone analogues. <i>Electrophoresis</i> , 2017, 38, 1922-1931.	2.4	10
654	Use of various β -cyclodextrin derivatives as chiral selectors for the enantiomeric separation of higenamine by capillary electrophoresis. <i>Microchemical Journal</i> , 2017, 134, 289-294.	4.5	13
655	Cyclodextrin complexation improves aqueous solubility of the antiepileptic drug, rufinamide: solution and solid state characterization of compound-cyclodextrin binary systems. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2017, 88, 43-52.	1.6	8

#	ARTICLE	IF	CITATIONS
656	Chiral separation of terbutaline and non-steroidal anti-inflammatory drugs by using a new lysine-bridged hemispherodextrin in capillary electrophoresis. Journal of Pharmaceutical and Biomedical Analysis, 2017, 145, 734-741.	2.8	14
657	Chiral surfaces: The many faces of chiral recognition. Current Opinion in Colloid and Interface Science, 2017, 32, 94-107.	7.4	31
658	Thermodynamic models to elucidate the enantioseparation of drugs with two stereogenic centers by micellar electrokinetic chromatography. Journal of Chromatography A, 2017, 1512, 133-142.	3.7	11
659	An analytical model for enantioseparation process in capillary electrophoresis. Physica A: Statistical Mechanics and Its Applications, 2017, 487, 153-163.	2.6	3
660	Enhancement of the conductivity detection signal in capillary electrophoresis systems using neutral cyclodextrins as sweeping agents. Electrophoresis, 2018, 39, 1390-1398.	2.4	9
661	Study of isomeric pentacyclic triterpene acids in traditional Chinese medicine of <i>Forsythiae Fructus</i> and their binding constants with β -cyclodextrin by capillary electrophoresis. Electrophoresis, 2018, 39, 1006-1013.	2.4	14
662	Into the theory of the partial filling affinity capillary electrophoresis and the determination of apparent stability constants of analyte-ligand complexes. Electrophoresis, 2018, 39, 742-751.	2.4	22
663	Equilibrium, structural and antibacterial characterization of moxifloxacin- β -cyclodextrin complex. Journal of Molecular Structure, 2018, 1166, 228-236.	3.6	30
664	Computer simulation and enantioselective capillary electrophoresis to characterize isomer mixtures of sulfated β -cyclodextrins. Electrophoresis, 2018, 39, 770-778.	2.4	12
665	Chiral separation of lenalidomide by liquid chromatography on polysaccharide-type stationary phases and by capillary electrophoresis using cyclodextrin selectors. Journal of Separation Science, 2018, 41, 1414-1423.	2.5	37
666	Molecular Modeling and Chiral Separation of Benzodiazepines by Capillary Electrophoresis Using Highly Sulfated Cyclodextrins. Journal of the Mexican Chemical Society, 2018, 62, .	0.6	4
667	Enantioseparations by Capillary Electromigration Techniques. , 2018, , 565-605.		2
668	Generalized model of the linear theory of electromigration and its application to electrokinetic chromatography: Theory and software PeakMaster 6-Next Generation. Electrophoresis, 2019, 40, 683-692.	2.4	24
669	Cyclodextrin-induced acidity modification of substituted cathinones studied by capillary electrophoresis supported by density functional theory calculations. Journal of Chromatography A, 2018, 1580, 142-151.	3.7	9
670	Enhanced enantioselectivity of native β -cyclodextrins by the synergy of chiral ionic liquids in capillary electrophoresis. Journal of Separation Science, 2018, 41, 4525-4532.	2.5	34
671	Contemporary theory of enantioseparations in capillary electrophoresis. Journal of Chromatography A, 2018, 1567, 2-25.	3.7	92
672	Fundamental aspects of field-amplified electrokinetic injection of cations for enantioselective capillary electrophoresis with sulfated cyclodextrins as selectors. Journal of Chromatography A, 2018, 1558, 85-95.	3.7	17
673	Preclinical evidence of enhanced analgesic activity of duloxetine complexed with succinyl- β -cyclodextrin: A comparative study with cyclodextrin complexes. International Journal of Pharmaceutics, 2019, 566, 391-399.	5.2	3

#	ARTICLE	IF	CITATIONS
674	Some thoughts about enantioseparations in capillary electrophoresis. <i>Electrophoresis</i> , 2019, 40, 2420-2437.	2.4	75
675	Separations of antifungal compounds in capillary electrophoresis with two anionic cyclodextrins. <i>Electrophoresis</i> , 2019, 40, 1986-1991.	2.4	2
676	Applicability assessment of the CASCade Two Dimensional SEDiment (CASC2Dâ€šSED) distributed hydrological model for flood forecasting across four typical medium and small watersheds in China. <i>Journal of Flood Risk Management</i> , 2019, 12, .	3.3	32
677	β -Cyclodextrin-copper (II) complex as chiral selector in capillary electrophoresis for the enantioseparation of β -blockers. <i>Journal of Chromatography A</i> , 2019, 1596, 233-240.	3.7	14
678	Synergistic enantioseparation systems with either cyclodextrins or cyclofructans and Lâ€šalanine <i>tert</i>-butyl ester lactate. <i>Electrophoresis</i> , 2019, 40, 539-546.	2.4	15
679	Determination of thermodynamic binding constants by affinity capillary electrophoresis. <i>Talanta</i> , 2019, 192, 448-454.	5.5	9
680	Clinical and pharmaceutical applications of affinity ligands in capillary electrophoresis: A review. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2020, 177, 112882.	2.8	40
681	Modeling-based optimization of the simultaneous enantiomeric separation of multicomponent mixtures of phenoxy acid herbicides using dual cyclodextrin systems by Capillary Electrophoresis. <i>Journal of Chromatography A</i> , 2020, 1610, 460552.	3.7	13
682	Investigation of the synergistic effect of chiral ionic liquids as additives in non-aqueous capillary electrophoresis for enantioseparation. <i>Journal of Chromatography A</i> , 2020, 1609, 460519.	3.7	36
683	Chiral covalent organic frameworks: design, synthesis and property. <i>Chemical Society Reviews</i> , 2020, 49, 6248-6272.	38.1	211
684	Binding constants determination of cyclodextrin inclusion complexes by affinity capillary electrophoresis. How to overcome the limitations induced by the UV-detector?. <i>Journal of Chromatography A</i> , 2020, 1623, 461209.	3.7	7
685	Use of Gamithromycin as a Chiral Selector in Capillary Electrophoresis. <i>Journal of Chromatography A</i> , 2020, 1624, 461099.	3.7	17
686	Investigation of the synergistic effect with chiral Dâ€špenicillamine functionalized gold nanoparticle as an additive for enantiomeric separation in capillary electrophoresis. <i>Electrophoresis</i> , 2020, 41, 1060-1066.	2.4	8
687	Single isomer cyclodextrins as chiral selectors in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2020, 1627, 461375.	3.7	47
688	Investigation of fusidic acid as a chiral selector in capillary electrophoresis. <i>Separation and Purification Technology</i> , 2020, 242, 116768.	7.9	17
689	Advances of capillary electrophoresis enantioseparations in pharmaceutical analysis (2017â€š2020). <i>Electrophoresis</i> , 2021, 42, 1709-1725.	2.4	51
690	Binding Constants Determination of Ofloxacin and Ornidazole Enantiomers with Sulfated β -Cyclodextrin as Single Ligand by Capillary Electrophoresis using Three Different Linear Plotting Methods. <i>Asian Journal of Chemistry</i> , 2021, 33, 1663-1670.	0.3	0
691	Bioanalysis of drugs and their metabolites by chiral electromigration techniques (2010â€š2020). <i>Electrophoresis</i> , 2021, 42, 1744-1760.	2.4	9

#	ARTICLE	IF	CITATIONS
692	Role of diaminonaphthalene based polymers as sensors in detection of biomolecules: A review. Results in Materials, 2021, 9, 100174.	1.8	5
693	Electrophoretic separation of multiple chiral center analyte with a three cyclodextrins mixture. Electrophoresis, 2021, 42, 1810-1817.	2.4	3
694	The Use of Dual Cyclodextrin Chiral Selector Systems in the Enantioseparation of Pharmaceuticals by Capillary Electrophoresis: An Overview. Molecules, 2021, 26, 2261.	3.8	19
695	Study on a novel buffer system to separate and quantify glycated hemoglobin A1c (HbA1c) and the mechanism between its key constituents and HbA1c. Microchemical Journal, 2021, 163, 105916.	4.5	1
696	Native and substituted cyclodextrins as chiral selectors for capillary electrophoresis enantioseparations: Structures, features, application, and molecular modeling. Electrophoresis, 2021, 42, 1676-1708.	2.4	52
697	Single Isomer N-Heterocyclic Cyclodextrin Derivatives as Chiral Selectors in Capillary Electrophoresis. Molecules, 2021, 26, 5271.	3.8	7
698	Functional materials in chiral capillary electrophoresis. Coordination Chemistry Reviews, 2021, 445, 214108.	18.8	25
699	Separation and Quantitation of Enantiomers. Chromatographia CE Series, 1998, , 70-100.	0.1	5
700	Applications of chiral capillary electrophoresis (cyclodextrin-capillary zone electrophoresis;) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 422 Td		
701	Pre-Concentration Approaches Combined with Capillary Electrophoresis in Bioanalysis of Chiral Cardiovascular Drugs. Pharmaceutical Sciences, 2015, 21, 229-243.	0.8	25
702	Modelling Enantiomer Separation by CE. Chromatographia CE Series, 2001, , 24-41.	0.1	0
703	Stereoselective Separations: Recent Advances in Capillary Electrophoresis and High-Performance Liquid Chromatography. Handbook of Experimental Pharmacology, 2003, , 45-75.	1.8	2
704	Chiral electromigration techniques in pharmaceutical and biomedical analysis. , 2013, , 225-244.		0
705	Quantitative Characterization of Derivatization Effects in Separations Techniques, I. Derivatization of Ibuprofen. Acta Polytechnica Hungarica, 2013, 10, .	2.9	0
707	3 Techniques used in the analysis of optically active compounds. 3.7 Electrophoresis. , 1995, , 243-278.		0
708	Determination of Drug-Cyclodextrin Binding Constants By Capillary Zone Electrophoresis. , 1996, , 675-678.		0
709	Electrolyte systems. , 1997, , 71-83.		0
710	Determination of enantiomeric purity of esomeprazole by capillary electrophoresis. Bulletin of Medical Sciences, 2020, 93, 93-101.	0.0	0

#	ARTICLE	IF	CITATIONS
711	Enantiomer Separation by Capillary Electrophoresis. , 0, , 333-368.		1
712	Separability of stereoisomers by electrokinetic chromatography in presence of a neutral selector â€“ Fundamental aspects assessed by computer simulation. Journal of Chromatography A, 2022, 1673, 463087.	3.7	3
713	Analytical methods for obtaining binding parameters of drugâ€“protein interactions: A review. Analytica Chimica Acta, 2022, 1219, 340012.	5.4	12
714	Application of Remote-Sensing-Based Hydraulic Model and Hydrological Model in Flood Simulation. Sustainability, 2022, 14, 8576.	3.2	4
715	Separation of planar chiral ferrocenes by capillary electrokinetic chromatography and liquid chromatography. Journal of Chromatography A, 2022, 1677, 463306.	3.7	6
716	Recognition in the Domain of Molecular Chirality: From Noncovalent Interactions to Separation of Enantiomers. Chemical Reviews, 2022, 122, 13235-13400.	47.7	77
717	Modelling the simultaneous chiral separation of a group of drugs by electrokinetic chromatography using mixtures of cyclodextrins. Journal of Chromatography A, 2022, 1681, 463444.	3.7	3
718	A feasibility study on the use of hydrophobic eutectic solvents as pseudo-stationary phases in capillary electrophoresis for chiral separations. Analytica Chimica Acta, 2023, 1239, 340693.	5.4	7
719	Chiral Separations of Pyrethroic Acids Using Cyclodextrin Selectors. Molecules, 2022, 27, 8718.	3.8	1
720	Cyclodextrins as chiral selectors in capillary electrophoresis: Recent trends in mechanistic studies. TrAC - Trends in Analytical Chemistry, 2023, 160, 116987.	11.4	17
723	Chiral separation by capillary electrokinetic chromatography with hydrophobic deep eutectic solvents as pseudo-stationary phases. Talanta, 2023, 260, 124556.	5.5	3
724	Leveraging the use of in vitro and computational methods to support the development of enabling oral drug products: An InPharma commentary. European Journal of Pharmaceutical Sciences, 2023, 188, 106505.	4.0	0
725	Strategies for capillary electrophoresis: Method development and validation for pharmaceutical and biological applicationsâ€“Updated and completely revised edition. Electrophoresis, 2023, 44, 1279-1341.	2.4	3
726	Deep eutectic solvents combined with beta-cyclodextrin derivatives for chiral separation of typical adrenergic receptor agonists by capillary electrophoresis with amperometric detection. Journal of Pharmaceutical and Biomedical Analysis, 2023, 236, 115748.	2.8	3
727	Relevance of the calculation of the diffusion coefficient in a capillary electrophoresis experiment. Electrophoresis, 2024, 45, 300-309.	2.4	0