

Theoretical aspects of chiral separation in capillary elec

Journal of Chromatography A

603, 235-241

DOI: 10.1016/0021-9673(92)85366-2

Citation Report

#	ARTICLE	IF	CITATIONS
1	Theoretical aspects of chiral separation in capillary electrophoresis. Journal of Chromatography A, 1992, 609, 363-367.	1.8	230
2	Separation of the enantiomers of some racemic nonsteroidal aromatase inhibitors and barbiturates by capillary electrophoresis. Chirality, 1993, 5, 516-526.	1.3	53
3	Theory of chiral separation in capillary electrophoresis. Journal of Chromatography A, 1993, 636, 57-62.	1.8	172
4	Differential binding of tioconazole enantiomers to hydroxypropyl- β -cyclodextrin studied by capillary electrophoresis. Journal of Chromatography A, 1993, 636, 149-152.	1.8	128
5	Validation of a capillary electrophoresis method for the enantiomeric purity testing of fluparoxan. Journal of Chromatography A, 1993, 645, 193-196.	1.8	64
6	Theoretical aspects of chiral separation in capillary electrophoresis. Journal of Chromatography A, 1993, 635, 113-118.	1.8	164
7	Chiral separation of basic drugs using cyclodextrins as chiral pseudo-stationary phases in capillary electrophoresis. Journal of Chromatography A, 1993, 648, 267-274.	1.8	116
8	Quantitative aspects of the application of capillary electrophoresis to the analysis of pharmaceuticals and drug related impurities. Journal of Chromatography A, 1993, 646, 245-257.	1.8	99
9	Charged and uncharged cyclodextrins as chiral selectors in capillary electrophoresis. Chromatographia, 1993, 37, 475-481.	0.7	198
10	Simultaneous chiral separation of leucovorin and its major metabolite 5-methyl-tetrahydrofolate by capillary electrophoresis using cyclodextrins as chiral selectors: Estimation of the formation constant and mobility of the solute-cyclodextrin complexes. Chromatographia, 1993, 35, 419-429.	0.7	124
11	Chiral separation of basic drugs using cyclodextrin-modified capillary zone electrophoresis. Analytical Chemistry, 1993, 65, 885-893.	3.2	159
12	Fully Automated Analysis of Amino Acid Enantiomers by Derivatization and Chiral Separation on a Capillary Electrophoresis Instrument. Journal of Liquid Chromatography and Related Technologies, 1994, 17, 1883-1897.	0.9	26
13	Chiral separations of amino acids by capillary electrophoresis and high-performance liquid chromatography employing chiral crown ethers. Journal of Chromatography A, 1994, 685, 321-329.	1.8	45
14	Separation of enantiomers by capillary electrophoretic techniques. Journal of Chromatography A, 1994, 666, 295-319.	1.8	274
15	Capillary zone electrophoresis for separation of drug enantiomers using cyclodextrins as chiral selectors. Journal of Chromatography A, 1994, 666, 337-350.	1.8	87
16	Chiral separations of basic and acidic compounds in modified capillaries using cyclodextrin-modified capillary zone electrophoresis. Journal of Chromatography A, 1994, 666, 351-365.	1.8	56
17	Enantioselective separations using capillary electrophoresis. Chirality, 1994, 6, 25-40.	1.3	92
18	Investigation of enantioselective ligand-protein binding and displacement interactions using capillary electrophoresis. Chirality, 1994, 6, 230-238.	1.3	45

#	ARTICLE	IF	CITATIONS
19	Chiral separations by capillary electrophoresis: Present state of the art. <i>Electrophoresis</i> , 1994, 15, 755-762.	1.3	69
20	A theoretical approach to chiral capillary electrophoresis with some practical implications. <i>Electrophoresis</i> , 1994, 15, 774-778.	1.3	61
21	Separations of derivatized amino acid enantiomers by cyclodextrin-modified capillary electrophoresis: Mechanistic and molecular modeling studies. <i>Electrophoresis</i> , 1994, 15, 785-792.	1.3	62
22	Enantiomeric separation of salbutamol and related impurities using capillary electrophoresis. <i>Electrophoresis</i> , 1994, 15, 808-817.	1.3	57
23	Chiral separation of basic drugs by capillary zone electrophoresis with cyclodextrin additives. <i>Electrophoresis</i> , 1994, 15, 818-823.	1.3	117
24	Separation of enantiomers by affinity electrokinetic chromatography using avidin. <i>Electrophoresis</i> , 1994, 15, 848-853.	1.3	83
25	Enantiomeric separation of drugs by affinity electrokinetic chromatography using dextran sulfate. <i>Electrophoresis</i> , 1994, 15, 1335-1340.	1.3	48
26	Capillary zone electrophoresis in pharmaceutical and biomedical analysis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1994, 12, 579-611.	1.4	62
27	Stereoselective determination of verapamil and norverapamil by capillary electrophoresis. <i>Biomedical Applications</i> , 1994, 654, 121-127.	1.7	52
28	Cyclodextrins as enantioselective mobile phase modifiers for chiral capillary electrophoresis effects of pH and cyclodextrin concentration. <i>Biomedical Applications</i> , 1994, 657, 291-300.	1.7	34
29	Systematic approach to treatment of enantiomeric separations in capillary electrophoresis and liquid chromatography.. <i>Journal of Chromatography A</i> , 1994, 680, 147-155.	1.8	39
30	Practical aspects of chiral separations of pharmaceuticals by capillary electrophoresis I. Separation optimization. <i>Journal of Chromatography A</i> , 1994, 680, 157-162.	1.8	49
31	Chiral selectors with chelating properties in liquid chromatography: Fundamental reflections and selective review of recent developments. <i>Journal of Chromatography A</i> , 1994, 666, 55-76.	1.8	145
32	Chiral separation of drugs by capillary electrophoresis using β -cyclodextrin polymer. <i>Journal of Chromatography A</i> , 1994, 678, 333-342.	1.8	44
33	Chiral separation of drugs using cyclodextrins in capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 1994, 659, 449-457.	1.8	55
34	Optimization of a capillary electrophoresis method to determine the chiral purity of a drug. <i>Journal of Chromatography A</i> , 1994, 680, 609-621.	1.8	63
35	High-performance liquid chromatographic separation of cis- and trans isomers of proline-containing peptides. <i>Journal of Chromatography A</i> , 1994, 661, 7-12.	1.8	26
36	Effect of the degree of substitution of (2-hydroxy)propyl- β -cyclodextrin on the enantioseparation of organic acids by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1994, 678, 139-144.	1.8	56

#	ARTICLE	IF	CITATIONS
37	Plackett-Burman experimental design in chiral analysis using capillary electrophoresis. <i>Chromatographia</i> , 1994, 38, 723-729.	0.7	71
38	Factors affecting the separation of mandelic acid enantiomers by capillary electrophoresis. <i>Chromatographia</i> , 1994, 38, 730-736.	0.7	26
39	Capillary Electrophoresis with Chiral Selectors: Optimization of Separation and Determination of Thermodynamic Parameters for Binding of Tioconazole Enantiomers to Cyclodextrins. <i>Analytical Chemistry</i> , 1994, 66, 2866-2873.	3.2	144
40	Chiral Resolution of Cationic Drugs of Forensic Interest by Capillary Electrophoresis with Mixtures of Neutral and Anionic Cyclodextrins. <i>Analytical Chemistry</i> , 1994, 66, 4019-4026.	3.2	264
41	Sulfobutyl Ether .beta.-Cyclodextrin as a Chiral Discriminator for Use with Capillary Electrophoresis. <i>Analytical Chemistry</i> , 1994, 66, 4013-4018.	3.2	224
42	A Peak Resolution Model for the Capillary Electrophoretic Separation of the Enantiomers of Weak Acids with Hydroxypropyl .beta.-Cyclodextrin-Containing Background Electrolytes. <i>Analytical Chemistry</i> , 1994, 66, 619-627.	3.2	167
43	Chiral Separations Using Micellar Electrokinetic Capillary Chromatography and a Polymerized Chiral Micelle. <i>Analytical Chemistry</i> , 1994, 66, 3773-3776.	3.2	170
44	Separation of binaphthyl enantiomers by capillary zone electrophoresis and electrokinetic chromatography. <i>Journal of High Resolution Chromatography</i> , 1995, 18, 659-664.	2.0	32
45	Mechanisms of enantiomeric resolution in cyclodextrin-modified capillary electrophoretic separations of binaphthyl compounds. <i>Chirality</i> , 1995, 7, 401-408.	1.3	18
46	Rapid chiral separation methods development by cyclodextrin-mediated capillary electrophoresis for acidic and basic compounds. <i>Chirality</i> , 1995, 7, 409-414.	1.3	25
47	Determination of the enantiomeric purity of dihydroxy- and dimethoxy-2-aminotetralins by high-performance capillary electrophoresis with cyclodextrins as chiral selector. <i>Chirality</i> , 1995, 7, 459-468.	1.3	12
48	Chiral separation of naphthalene-2, 3-dialdehyde labelled peptides by cyclodextrin-modified electrokinetic chromatography. <i>Biomedical Chromatography</i> , 1995, 9, 295-301.	0.8	20
49	Chiral separations using heparin and dextran sulfate in capillary zone electrophoresis. <i>Analytica Chimica Acta</i> , 1995, 307, 185-191.	2.6	55
50	Viscous effects in capillary electrophoresis: Theory and experiment. <i>Electrophoresis</i> , 1995, 16, 2074-2085.	1.3	14
51	Chiral separation in capillary electrophoresis. <i>Electrophoresis</i> , 1995, 16, 2127-2131.	1.3	55
52	Electroosmotic control of chiral separation in capillary zone electrophoresis. <i>Electrophoresis</i> , 1995, 16, 2132-2136.	1.3	13
53	Protein chiral selectors in free-solution capillary electrophoresis and packed-capillary electrochromatography. <i>Journal of Chromatography A</i> , 1995, 694, 285-296.	1.8	150
54	Partial separation zone technique for the separation of enantiomers by affinity electrokinetic chromatography with proteins as chiral pseudo-stationary phases. <i>Journal of Chromatography A</i> , 1995, 694, 277-284.	1.8	170

#	ARTICLE	IF	CITATIONS
55	Extending the scope of chiral separation of basic compounds by cyclodextrin-mediated capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 1995, 692, 253-265.	1.8	81
56	Experimental evidence for the existence of duoselective (Type III) enantiomer separations in the capillary electrophoretic analysis of chiral weak acids. <i>Journal of Chromatography A</i> , 1995, 692, 319-325.	1.8	50
57	Enantioselective capillary electrophoresis of amino acid derivatives on cyclodextrin evaluation of structure-resolution relationships. <i>Journal of Chromatography A</i> , 1995, 697, 549-560.	1.8	64
58	Enantiomeric separation by capillary electrophoresis using a soluble neutral β -cyclodextrin polymer. <i>Journal of Chromatography A</i> , 1995, 709, 89-98.	1.8	47
59	Systematic approach to treatment of enantiomeric separations in capillary electrophoresis and liquid chromatography II. A study of the enantiomeric separation of fluoxetine and norfluoxetine. <i>Journal of Chromatography A</i> , 1995, 700, 59-67.	1.8	56
60	Application of modified cyclodextrins in capillary electrophoresis for enantiomeric resolution of propranolol and analogues. <i>Journal of Chromatography A</i> , 1995, 705, 351-361.	1.8	29
61	Determination of β -cyclodextrin inclusion complex constants for 3,4-dihydro-2-H-1-benzopyran enantiomers by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1995, 707, 311-326.	1.8	54
62	Enantiomeric purity determination of propranolol by cyclodextrin-modified capillary electrophoresis. <i>Journal of Chromatography A</i> , 1995, 717, 203-209.	1.8	91
63	Combined polymerized chiral micelle and β -cyclodextrin for chiral separation in capillary electrophoresis. <i>Journal of Chromatography A</i> , 1995, 711, 297-304.	1.8	113
64	Determination of the enantiomeric purity of 5,6-dihydroxy-2-aminotetralin by high-performance capillary electrophoresis with crown ether as chiral selector. <i>Journal of Chromatography A</i> , 1995, 715, 143-149.	1.8	27
65	Use of organic modifiers to enhance chiral selectivity in capillary electrophoresis. <i>Amino Acids</i> , 1995, 8, 337-344.	1.2	14
66	Enantiomeric separation of denopamine by capillary electrophoresis and high-performance liquid chromatography using cyclodextrins. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1995, 13, 1483-1492.	1.4	19
67	Optical resolution of drugs by capillary electrophoretic techniques. <i>Journal of Chromatography A</i> , 1995, 694, 245-276.	1.8	274
68	Chiral separation of deprenyl and its major metabolites using cyclodextrin-modified capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 1995, 709, 157-162.	1.8	42
69	Capillary zone electrophoretic separation of the enantiomers of dipeptides based on host-guest complexation with a chiral crown ether. <i>Journal of Chromatography A</i> , 1995, 709, 81-88.	1.8	70
70	Resolution of the enantiomers of oxamniquine by capillary electrophoresis and high-performance liquid chromatography with cyclodextrins and heparin as chiral selectors. <i>Journal of Chromatography A</i> , 1995, 700, 51-58.	1.8	45
71	Direct and indirect chiral separation of amino acids by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1995, 704, 179-193.	1.8	103
72	Capillary zone electrophoretic determination of C ₂₁ –C ₁₈ linear saturated free fatty acids with indirect absorbance detection. <i>Journal of Chromatography A</i> , 1995, 708, 339-350.	1.8	72

#	ARTICLE	IF	CITATIONS
73	Chiral separation of amphetamines by high-performance capillary electrophoresis. <i>Journal of Chromatography A</i> , 1995, 717, 219-228.	1.8	92
74	Enantiomeric separation of amphetamine related drugs by capillary zone electrophoresis using native and derivatized β -cyclodextrin as chiral additives. <i>Journal of Chromatography A</i> , 1995, 710, 339-345.	1.8	57
75	Effect of urea addition on chiral separation of dansylamino acids by capillary zone electrophoresis with cyclodextrins. <i>Journal of Chromatography A</i> , 1995, 710, 331-337.	1.8	32
76	Buffer effects in the desionoselective/ionoselective/duoselective separation selectivity model-assisted optimization of the capillary electrophoretic separation of enantiomers I. Low-pH phosphate buffers. <i>Journal of Chromatography A</i> , 1995, 716, 197-205.	1.8	20
77	Enantiomeric separation of D/L-carnitine using HPLC and CZE after derivatization. <i>Chromatographia</i> , 1995, 40, 287-295.	0.7	52
78	Enantiomeric separation of trimetoquinol, denopamine and timepidium by capillary electrophoresis and HPLC and the application of capillary electrophoresis to the optical purity testing of the drugs. <i>Chromatographia</i> , 1995, 40, 638-644.	0.7	34
79	Capillary electrophoretic separation of basic drugs using surface-modified C8 capillaries and derivatized cyclodextrins as structural/chiral selectors. <i>Chromatographia</i> , 1995, 40, 697-704.	0.7	22
80	Effect of Micelles and Mixed Micelles on Efficiency and Selectivity of Antibiotic-Based Capillary Electrophoretic Enantioseparations. <i>Analytical Chemistry</i> , 1995, 67, 2088-2095.	3.2	121
81	Enantiomeric Separation of Drugs by Mucopolysaccharide-Mediated Electrokinetic Chromatography. <i>Analytical Chemistry</i> , 1995, 67, 2334-2341.	3.2	83
82	Effect of the Initial Potential Ramp on the Accuracy of Electrophoretic Mobilities in Capillary Electrophoresis. <i>Analytical Chemistry</i> , 1995, 67, 3079-3081.	3.2	29
83	Separation of PAHs by Capillary Electrophoresis with Laser-Induced Fluorescence Detection Using Mixtures of Neutral and Anionic β -Cyclodextrins. <i>Analytical Chemistry</i> , 1995, 67, 3004-3010.	3.2	75
84	Affinity Capillary Electrophoresis. <i>Accounts of Chemical Research</i> , 1995, 28, 461-468.	7.6	147
85	Chiral Separation and Detection of Terbutaline and Ephedrine by Capillary Electrophoresis Coupled with Ion Spray Mass Spectrometry. <i>Analytical Chemistry</i> , 1995, 67, 2054-2058.	3.2	134
86	Chiral Separations by Nonaqueous Capillary Electrophoresis. <i>Analytical Chemistry</i> , 1996, 68, 3460-3467.	3.2	173
87	Chiral separation of dansyl amino acids by capillary electrophoresis: Comparison of formamide and N-Methylformamide as background electrolytes. <i>Chromatographia</i> , 1996, 43, 242-246.	0.7	55
88	Separation of drug enantiomers by capillary electrophoresis: β -cyclodextrin as chiral solvating agent. <i>Chromatographia</i> , 1996, 42, 106-110.	0.7	25
89	Chiral separation of local anaesthetics with capillary electrophoresis. <i>Chromatographia</i> , 1996, 42, 263-268.	0.7	40
90	Determination of the binding of β -cyclodextrin derivatives to adamantane carboxylic acids using capillary electrophoresis. <i>Chromatographia</i> , 1996, 43, 659-662.	0.7	38

#	ARTICLE	IF	CITATIONS
91	Micellar electrokinetic capillary chromatography with in-situ charged micelles VI. Evaluation of novel chiral micelles consisting of steroidal-glycoside surfactant-borate complexes. <i>Journal of Chromatography A</i> , 1996, 724, 285-296.	1.8	63
92	Maximization of separation efficiency in capillary electrophoretic chiral separations by means of mobility-matching background electrolytes. <i>Journal of Chromatography A</i> , 1996, 730, 273-278.	1.8	44
93	Enantiomer separation of drugs by electrokinetic chromatography. <i>Journal of Chromatography A</i> , 1996, 735, 57-76.	1.8	131
94	Enantiomeric separation of amphetamine and phenylephrine by cyclodextrin-mediated capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 1996, 735, 295-301.	1.8	23
95	Determination of the enantiomeric purity of dexfenfluramine by capillary electrophoresis: use of a Plackett-Burman design for the optimization of the separation. <i>Journal of Chromatography A</i> , 1996, 736, 281-289.	1.8	51
96	Chiral separation of dansyl-amino acids in a nonaqueous medium by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1996, 737, 263-272.	1.8	86
97	Influence of the nature of the buffer on chiral separation in capillary electrophoresis. <i>Journal of Chromatography A</i> , 1996, 745, 61-71.	1.8	16
98	Cyclodextrins as buffer additives for the enantiomeric separation of pinacidil in capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 1996, 749, 300-303.	1.8	7
99	Determination of the binding constant of salbutamol to unmodified and ethylated cyclodextrins by affinity capillary electrophoresis. <i>Journal of Chromatography A</i> , 1996, 735, 321-331.	1.8	48
100	Electrokinetic chromatography in suppressed electroosmotic flow environment: Use of a charged cyclodextrin for the separation of enantiomers and geometric isomers. <i>Electrophoresis</i> , 1996, 17, 1575-1583.	1.3	65
101	A practical procedure for the determination of association constants of the analyte-chiral selector equilibria by capillary zone electrophoresis. <i>Electrophoresis</i> , 1996, 17, 1921-1924.	1.3	18
102	Separation of neutral and basic enantiomers by cyclodextrin electrokinetic chromatography using anionic cyclodextrin derivatives as chiral pseudo-stationary phases. <i>Journal of High Resolution Chromatography</i> , 1996, 19, 421-433.	2.0	80
103	Chiral separation of methoxamine and lobeline using cyclodextrin as buffer additives in capillary zone electrophoresis. <i>Journal of High Resolution Chromatography</i> , 1996, 19, 478-480.	2.0	10
104	Use of cyclodextrins as chiral selector for the chiral separation of anticholinergic drugs such as anisodamine and glycopyrronium in capillary zone electrophoresis. <i>Journal of High Resolution Chromatography</i> , 1996, 19, 697-699.	2.0	8
105	Cyclodextrin-aided capillary electrophoretic separation and laser-induced fluorescence detection of polynuclear aromatic hydrocarbons (PAHs). , 1996, 9, 515-523.		9
106	Role of capillary electrophoresis methods in the drug development process. <i>Chirality</i> , 1996, 8, 108-121.	1.3	45
107	Capillary electrophoretic enantiomeric separations using the glycopeptide antibiotic, teicoplanin. <i>Chirality</i> , 1996, 8, 88-107.	1.3	98
108	Sulfated cyclodextrins for the chiral separations of catecholamines and related compounds in the reversed electrophoretic polarity mode. , 1996, 8, 316-324.		48

#	ARTICLE	IF	CITATIONS
109	Capillary electrophoresis of herbicides. III. Evaluation of octylmaltopyranoside chiral surfactant in the enantiomeric separation of phenoxy acid herbicides. , 1996, 8, 518-524.		35
110	Enantiomeric separations of basic pharmaceutical drugs by micellar electrokinetic chromatography using a chiral surfactant, N-dodecoxy carbonylvaline. Biomedical Applications, 1996, 683, 15-28.	1.7	50
111	Chiral separations of underivatized arylpropionic acids by capillary zone electrophoresis with various cyclodextrins Acidity and inclusion constant determinations. Journal of Chromatography A, 1996, 735, 311-320.	1.8	67
112	Examination of the origin, variation, and proper use of expressions for the estimation of association constants by capillary electrophoresis. Journal of Chromatography A, 1996, 721, 173-186.	1.8	273
113	Multiple-plug binding assays using affinity capillary electrophoresis. Journal of Chromatography A, 1996, 727, 291-299.	1.8	47
114	Enantiomeric separation of local anaesthetic drugs by micellar electrokinetic capillary chromatography with taurodeoxycholate as chiral selector. Journal of Chromatography A, 1996, 737, 301-313.	1.8	35
115	Direct chiral separation of amino acids derivatized with 2-(9-anthryl)ethyl chloroformate by capillary electrophoresis using cyclodextrins as chiral selectors Effect of organic modifiers on resolution and enantiomeric elution order. Journal of Chromatography A, 1996, 731, 283-292.	1.8	56
116	Separation of enantiomers of drugs by capillary electrophoresis III. β -cyclodextrin as chiral solvating agent. Journal of Chromatography A, 1996, 735, 333-343.	1.8	53
117	Comparative capillary electrophoretic and nuclear magnetic resonance studies of the chiral recognition of racemic metomidate with cyclodextrin hosts. Journal of Chromatography A, 1996, 732, 133-142.	1.8	69
118	Chiral separation of dioxypromethazine enantiomers by capillary electrophoresis using β -cyclodextrin as a chiral selector. Journal of Chromatography A, 1996, 732, 175-181.	1.8	24
119	Identification of chiral drug isomers by capillary electrophoresis. Journal of Chromatography A, 1996, 735, 77-121.	1.8	319
120	Capillary electrophoresis of cardiovascular drugs. Journal of Chromatography A, 1996, 735, 123-150.	1.8	23
121	Separation of the four stereoisomers of Z11556A, intermediate in the synthesis of triazole antimycotics, by capillary electrophoresis. Journal of Chromatography A, 1996, 741, 123-130.	1.8	8
122	Self-consistent framework for standardising mobilities in free solution capillary electrophoresis: applications to oligoglycines and oligoalanines. Journal of Chromatography A, 1996, 741, 99-113.	1.8	58
123	Chiral separation of muscarinic antagonists by capillary zone electrophoresis with cyclodextrin additives. Journal of Chromatography A, 1996, 741, 287-294.	1.8	9
124	Separation of positional and structural isomers by cyclodextrin-mediated capillary zone electrophoresis. Journal of Chromatography A, 1996, 745, 9-15.	1.8	27
125	Chiral separation in non-aqueous media by capillary electrophoresis using the ion-pair principle. Journal of Chromatography A, 1996, 745, 37-44.	1.8	113
126	Systematic approach to the treatment of enantiomeric separations in capillary electrophoresis and liquid chromatography III. A binding constant-retention factor relationship and effects of acetonitrile on the chiral separation of tioconazole. Journal of Chromatography A, 1996, 745, 25-35.	1.8	43

#	ARTICLE	IF	CITATIONS
127	Use of cyclodextrins in the capillary electrophoretic separation of erythro- and threo-mefloquine enantiomers. <i>Journal of Chromatography A</i> , 1996, 745, 17-23.	1.8	25
128	Determination of binding constants and the influence of methanol on the separation of drug enantiomers in cyclodextrin modified capillary electrophoresis. <i>Journal of Chromatography A</i> , 1996, 745, 181-187.	1.8	30
129	Utilization of in enantiomeric separation by micellar electrokinetic chromatography. <i>Journal of Chromatography A</i> , 1996, 749, 257-269.	1.8	22
130	Ergot alkaloids as chiral selectors in capillary electrophoresis determination of the separation mechanism. <i>Journal of Chromatography A</i> , 1996, 755, 251-259.	1.8	22
131	Migration behavior and separation of sulfonamides in capillary zone electrophoresis I. Influence of buffer pH and electrolyte modifier. <i>Journal of Chromatography A</i> , 1996, 755, 261-269.	1.8	37
132	Resolution improvement by use of carboxymethyl- β -cyclodextrin as chiral additive for the enantiomeric separation of basic drugs by capillary electrophoresis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1996, 14, 1107-1114.	1.4	52
133	Cyclodextrins and enantiomeric separations of drugs by liquid chromatography and capillary electrophoresis: basic principles and new developments. <i>Biomedical Applications</i> , 1996, 687, 303-336.	1.7	153
134	Determination of drug-cyclodextrin binding constants by capillary zone electrophoresis. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 1996, 25, 253-256.	1.6	7
135	Chiral Separation of Reduced Haloperidol by Capillary Zone Electrophoresis with Heptakis (2,6-Di-o-Methyl)- β -cyclodextrin. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1996, 19, 1567-1577.	0.5	9
136	Development of a Free Solution Capillary Electrophoresis Method for the Separation and Quantitation of D-and L-Phenylalanine Methyl Ester.. <i>Analytical Letters</i> , 1996, 29, 991-1001.	1.0	3
137	Determination of Association Constants in Cyclodextrin or Vancomycin-Modified Micellar Capillary Electrophoresis. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1997, 20, 203-216.	0.5	15
138	Separation of the Enantiomers of Propranolol by Incorporation of Molecularly Imprinted Polymer Particles as Chiral Selectors in Capillary Electrophoresis. <i>Analytical Communications</i> , 1997, 34, 119-122.	2.2	59
139	Determination of Accurate Electroosmotic Mobility and Analyte Effective Mobility Values in the Presence of Charged Interacting Agents in Capillary Electrophoresis. <i>Analytical Chemistry</i> , 1997, 69, 4445-4451.	3.2	59
140	Enantiomeric Separations by Use of Calixarene Electrokinetic Chromatography. <i>Analytical Chemistry</i> , 1997, 69, 3239-3242.	3.2	70
141	Enantioseparation of 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate-derivatized amino acids by capillary zone electrophoresis using native and substituted β -cyclodextrins as chiral additives II. Evaluation of complexation constants. <i>Journal of Chromatography A</i> , 1997, 759, 167-175.	1.8	8
142	Reversed determination of the formation constants of 1-allyl terguride with mandelic acid optical isomers using capillary electrophoresis. <i>Journal of Chromatography A</i> , 1997, 772, 179-184.	1.8	9
143	Enantiomeric separation of the novel growth hormone secretagogue MK-0677 by capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 1997, 773, 311-320.	1.8	22
144	Electrokinetic chromatography without electroosmotic flow. <i>Journal of Chromatography A</i> , 1997, 792, 125-141.	1.8	37

#	ARTICLE	IF	CITATIONS
145	Selectivity in capillary electrophoresis: the use of proteins. <i>Journal of Chromatography A</i> , 1997, 792, 349-369.	1.8	74
146	Neutral cyclodextrins as chiral agents for enantiomeric separations of chromanes in capillary electrophoresis. <i>Chromatographia</i> , 1997, 44, 313-319.	0.7	8
147	Separation of the enantiomers of basic drugs by affinity capillary electrophoresis using a partial filling technique and β -1-acid glycoprotein as chiral selector. <i>Chromatographia</i> , 1997, 44, 119-128.	0.7	73
148	Charged native β -cyclodextrin as a pseudostationary phase in electrokinetic chromatography. <i>Fresenius' Journal of Analytical Chemistry</i> , 1997, 357, 32-36.	1.5	9
149	Guidelines in selecting ligand concentrations for the determination of binding constants by affinity capillary electrophoresis. <i>Biomedical Applications</i> , 1997, 697, 77-88.	1.7	50
150	Simultaneous determination of the elimination profiles of the individual enantiomers of racemic isoproterenol using capillary electrophoresis and microdialysis sampling. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1997, 15, 621-629.	1.4	29
151	A chiral capillary electrophoresis method for ropivacaine hydrochloride in pharmaceutical formulations: Validation and comparison with chiral liquid chromatography. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1997, 15, 1051-1061.	1.4	37
152	Selected applications of capillaries with dynamic or permanent anodal electroosmotic flow in chiral separations by capillary electrophoresis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1997, 15, 1577-1584.	1.4	36
153	Use of $^1\text{H-NMR}$ spectroscopy to determine the enantioselective mechanism of neutral and anionic cyclodextrins in capillary electrophoresis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1997, 15, 1603-1619.	1.4	41
154	Quantitative description of migration behavior of porphyrins based on the dynamic complexation model in a nonaqueous capillary electrophoresis system. <i>Electrophoresis</i> , 1997, 18, 82-91.	1.3	52
155	Stereoselective analysis of herbicides by capillary electrophoresis using sulfobutyl ether β -cyclodextrin as chiral selector. <i>Electrophoresis</i> , 1997, 18, 227-234.	1.3	42
156	Quantitative description of analyte migration behavior based on dynamic complexation in capillary electrophoresis with one or more additives. <i>Electrophoresis</i> , 1997, 18, 706-716.	1.3	65
157	Chiral separations by capillary zone electrophoresis: Present state of the art. <i>Electrophoresis</i> , 1997, 18, 843-852.	1.3	69
158	Capillary zone electrophoresis at subzero temperatures II: Chiral separation of biogenic amines. <i>Electrophoresis</i> , 1997, 18, 873-883.	1.3	26
159	Three approaches to enantiomer separation of β -adrenergic antagonists by capillary electrochromatography. <i>Electrophoresis</i> , 1997, 18, 884-890.	1.3	129
160	Use of a zwitterionic cyclodextrin as a chiral agent for the separation of enantiomers by capillary electrophoresis. <i>Electrophoresis</i> , 1997, 18, 891-896.	1.3	29
161	Histamine-modified β -cyclodextrins for the enantiomeric separation of dansyl-amino acids in capillary electrophoresis. <i>Electrophoresis</i> , 1997, 18, 905-911.	1.3	65
162	Determination of association constants of dansyl-amino acids and β -cyclodextrin in <i>N</i> -methylformamide by capillary electrophoresis. <i>Electrophoresis</i> , 1997, 18, 919-923.	1.3	36

#	ARTICLE	IF	CITATIONS
163	Separation of enantiomers of drugs by capillary electrophoresis. Part 4: Hydroxypropyl-gammacyclodextrin as chiral solvating agent. <i>Electrophoresis</i> , 1997, 18, 924-930.	1.3	39
164	Central composite design in the chiral analysis of amphetamines by capillary electrophoresis. <i>Electrophoresis</i> , 1997, 18, 931-937.	1.3	70
165	Chiral interactions in capillary zone electrophoresis: Computer simulation and comparison with experiment. <i>Electrophoresis</i> , 1997, 18, 938-942.	1.3	13
166	Determination of the enantiomeric purity of N-propionyl-6,7-dimethoxy-2-aminotetralin by cyclodextrin-modified micellar electrokinetic chromatography. <i>Electrophoresis</i> , 1997, 18, 996-1001.	1.3	15
167	Enantiomer separation of disopyramide with capillary electrophoresis using various cyclodextrins. <i>Electrophoresis</i> , 1997, 18, 1002-1006.	1.3	20
168	Enantioseparation of local anaesthetic drugs by capillary zone electrophoresis with cyclodextrins as chiral selectors using a partial filling technique. <i>Electrophoresis</i> , 1997, 18, 1019-1025.	1.3	32
169	Methods for the estimation of binding constants by capillary electrophoresis. <i>Electrophoresis</i> , 1997, 18, 2194-2202.	1.3	153
170	Capillary electrophoretic separations of enantiomers using cyclodextrin-containing background electrolytes. <i>Electrophoresis</i> , 1997, 18, 2305-2310.	1.3	88
171	Redefining the separation factor: A potential pathway to a unified separation science. <i>Electrophoresis</i> , 1997, 18, 2928-2934.	1.3	23
172	Resolution of the diastereomers of a large synthetic peptide by capillary electrophoresis using nonionic surfactants. <i>Electrophoresis</i> , 1997, 18, 2944-2949.	1.3	9
173	Cyclodextrin-modified micellar electrokinetic chromatography for separation of norgestrel enantiomers. <i>Journal of High Resolution Chromatography</i> , 1997, 20, 159-164.	2.0	2
174	Enantiomeric separation of amphetamine and phenylephrine by capillary zone electrophoresis with hydroxypropyl- β -cyclodextrin as chiral additive. <i>Journal of High Resolution Chromatography</i> , 1997, 20, 298-299.	2.0	2
175	Chiral Separations of Enantiomeric Pharmaceuticals by Capillary Electrophoresis Using Sulphobutyl Ether β -Cyclodextrin as Isomer Selector. <i>Biomedical Chromatography</i> , 1997, 11, 193-199.	0.8	29
176	Separation of Chiral Amino Acids by Micellar Electrokinetic Chromatography with Derivatized Cyclodextrins. , 1997, 11, 230-235.		26
177	Phosphated cyclodextrins as new acidic chiral additives for capillary electrophoresis. <i>Journal of Separation Science</i> , 1997, 9, 581-589.	1.0	30
178	Capillary electrophoresis of herbicides II. Evaluation of alkylglucoside chiral surfactants in the enantiomeric separation of phenoxy acid herbicides. <i>Journal of Chromatography A</i> , 1997, 757, 263-273.	1.8	52
179	Effect of β -cyclodextrin acetylation on the resolution of phenethylamines with capillary electrophoresis and nuclear magnetic resonance spectroscopy. <i>Journal of Chromatography A</i> , 1997, 758, 277-292.	1.8	44
180	Chiral separation of monoterpenes using mixtures of sulfated β -cyclodextrins and β -cyclodextrin as chiral additives in the reversed-polarity capillary electrophoresis mode. <i>Journal of Chromatography A</i> , 1997, 759, 149-155.	1.8	66

#	ARTICLE	IF	CITATIONS
181	Enantioseparation of 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate-derivatized-amino acids by capillary zone electrophoresis using native and substituted β -cyclodextrins as chiral additives I. Discussion of optimum separation conditions. <i>Journal of Chromatography A</i> , 1997, 759, 157-165.	1.8	13
182	Chiral resolution of diols by capillary electrophoresis using borate-cyclodextrin complexation. <i>Journal of Chromatography A</i> , 1997, 761, 269-275.	1.8	42
183	Migration order reversal of enantiomers in capillary electrophoretic separation. <i>Journal of Chromatography A</i> , 1997, 764, 151-156.	1.8	20
184	Chiral separation of neutral species by capillary electrophoresis evaluation of a theoretical model. <i>Journal of Chromatography A</i> , 1997, 761, 249-257.	1.8	49
185	Mobility measurements on dansylated amino acids. <i>Journal of Chromatography A</i> , 1997, 768, 153-159.	1.8	28
186	Chiral recognition and racemic resolution of drug enantiomers by electrophoresis based on host-guest complexation. <i>Journal of Chromatography A</i> , 1997, 763, 261-269.	1.8	15
187	Vacancy affinity capillary electrophoresis, a new method for measuring association constants. <i>Journal of Chromatography A</i> , 1997, 775, 313-326.	1.8	55
188	Enantioseparation using selected polysaccharides as chiral buffer additives in capillary electrophoresis. <i>Journal of Chromatography A</i> , 1997, 773, 331-338.	1.8	47
189	Principles and limitations of methods available for the determination of binding constants with affinity capillary electrophoresis. <i>Journal of Chromatography A</i> , 1997, 777, 329-353.	1.8	107
190	Dry look at the CHARM (charged resolving agent migration) model of enantiomer separations by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1997, 777, 295-309.	1.8	173
191	Prediction of the migration behavior of analytes in capillary electrophoresis based on three fundamental parameters. <i>Journal of Chromatography A</i> , 1997, 781, 23-34.	1.8	25
192	Enantiomeric separation of a tetrapeptide with cyclodextrin extension of the model for chiral capillary electrophoresis by complex formation of one enantiomer molecule with more than one chiral selector molecules. <i>Journal of Chromatography A</i> , 1997, 782, 271-279.	1.8	39
193	Achiral selectivity in cyclodextrin-modified capillary electrophoresis. <i>Journal of Chromatography A</i> , 1997, 792, 431-444.	1.8	48
194	Selectivity in capillary electrophoresis in the presence of micelles, chiral selectors and non-aqueous media. <i>Journal of Chromatography A</i> , 1997, 792, 13-35.	1.8	105
195	Separation selectivity in chiral capillary electrophoresis with charged selectors. <i>Journal of Chromatography A</i> , 1997, 792, 269-295.	1.8	251
196	Enantioselectivity in chiral capillary electrophoresis with polysaccharides. <i>Journal of Chromatography A</i> , 1997, 792, 327-347.	1.8	85
197	Application of cyclodextrin-modified micellar electrokinetic chromatography to the separations of selected neutral pesticides and their enantiomers. <i>Journal of Chromatography A</i> , 1997, 792, 419-429.	1.8	73
198	Controlling enantioselectivity in chiral capillary electrophoresis with inclusion complexation. <i>Journal of Chromatography A</i> , 1997, 792, 227-267.	1.8	225

#	ARTICLE	IF	CITATIONS
199	Chiral separation principles in capillary electrophoresis. <i>Journal of Chromatography A</i> , 1997, 792, 179-225.	1.8	237
200	Study on the chiral separation of basic drugs by capillary zone electrophoresis using β -CD and derivatized β -CDs as chiral selectors. <i>Analytica Chimica Acta</i> , 1998, 359, 39-46.	2.6	25
201	Chiral separations of indan, tetralin and benzosuberan derivatives by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1998, 793, 135-143.	1.8	14
202	Chiral separation of 3-phenyl-3-(2-pyridyl)propylamines, and analogous guanidines and guanidine-N-carboxylic acid esters with high-performance liquid chromatography and capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 1998, 793, 77-90.	1.8	11
203	Separation of enantiomers of drugs by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1998, 793, 153-164.	1.8	46
204	Enantiomeric separation of propranolol and selected metabolites by using capillary electrophoresis with hydroxypropyl- β -cyclodextrin as chiral selector. <i>Journal of Chromatography A</i> , 1998, 793, 357-364.	1.8	49
205	Cationic β -cyclodextrin derivative for chiral separations. <i>Journal of Chromatography A</i> , 1998, 803, 261-271.	1.8	65
206	New derivatives of cyclodextrins as chiral selectors for the capillary electrophoretic separation of dichlorprop enantiomers. <i>Journal of Chromatography A</i> , 1998, 810, 193-200.	1.8	16
207	Simultaneous separation of o-, m-, p-fluoro-dl-phenylalanine and o-, m-, p-dl-tyrosine by ligand-exchange micellar electrokinetic capillary chromatography. <i>Journal of Chromatography A</i> , 1998, 813, 369-378.	1.8	61
208	Chiral separations by capillary zone electrophoresis with the use of cyanoethylated- β -cyclodextrin as chiral selector. <i>Journal of Chromatography A</i> , 1998, 817, 91-104.	1.8	23
209	Effect of various cyclodextrin derivatives on the resolution of fencamfamine isomers with capillary electrophoresis and nuclear magnetic resonance. <i>Journal of Chromatography A</i> , 1998, 818, 239-249.	1.8	18
210	Fast chiral separation of amino acid derivatives and acidic drugs by co-electroosmotic flow capillary electrophoresis with vancomycin as chiral selector. <i>Journal of Chromatography A</i> , 1998, 825, 81-87.	1.8	43
211	Effect of additives on chiral selectivities by β -cyclodextrin in capillary electrophoresis based on the phenomenon of solvatochromism. <i>Journal of Chromatography A</i> , 1998, 829, 301-307.	1.8	6
212	Chiral separation of racemic mexiletine hydrochloride using cyclodextrins as chiral additive by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1998, 795, 394-398.	1.8	9
213	Comparative capillary electrophoresis and NMR studies of enantioseparation of dimethindene with cyclodextrins. <i>Journal of Chromatography A</i> , 1998, 798, 315-323.	1.8	86
214	Chiral resolution of functionalized piperidine enantiomers by capillary electrophoresis with native, alkylated and anionic β -cyclodextrins. <i>Journal of Chromatography A</i> , 1998, 796, 375-383.	1.8	20
215	Use of chiral and achiral ion-pairing reagents in combination with cyclodextrins in capillary electrophoresis. <i>Journal of Chromatography A</i> , 1998, 798, 281-288.	1.8	34
216	Use of cationic cyclodextrin for enantioseparation by capillary electrophoresis. <i>Journal of Chromatography A</i> , 1998, 798, 275-280.	1.8	33

#	ARTICLE	IF	CITATIONS
217	Mixed-mode capillary electrokinetic separation of positional explosive isomers using sodium dodecyl sulfate and negative- β -cyclodextrin derivatives. <i>Journal of Chromatography A</i> , 1998, 811, 225-232.	1.8	28
218	Nonaqueous capillary electrophoretic separation of enantiomers using the single-isomer heptakis(2,3-diacetyl-6-sulfato)- β -cyclodextrin as chiral resolving agent. <i>Journal of Chromatography A</i> , 1998, 816, 233-241.	1.8	82
219	Affinity capillary electrophoresis: A physical-organic tool for studying interactions in biomolecular recognition. <i>Electrophoresis</i> , 1998, 19, 367-382.	1.3	178
220	Dynamic complexation of solutes in capillary electrophoresis. <i>Electrophoresis</i> , 1998, 19, 383-387.	1.3	15
221	The effects of a mixture of charged and neutral additives on analyte migration behavior in capillary electrophoresis. <i>Electrophoresis</i> , 1998, 19, 388-396.	1.3	24
222	Enantioseparation of anaesthetic drugs by capillary zone electrophoresis using cyclodextrin-containing background electrolytes. <i>Electrophoresis</i> , 1998, 19, 731-737.	1.3	18
223	Capillary electrophoresis chiral separations of basic compounds using cationic cyclodextrin. <i>Electrophoresis</i> , 1998, 19, 2095-2100.	1.3	30
224	Capillary electrophoresis, nuclear magnetic resonance and mass spectrometry studies of opposite chiral recognition of chlorpheniramine enantiomers with various cyclodextrins. <i>Electrophoresis</i> , 1998, 19, 2101-2108.	1.3	50
225	Separation of chiral compounds by capillary electrophoresis. <i>Electrophoresis</i> , 1998, 19, 2798-2833.	1.3	144
226	Use of cyclodextrins in capillary electrophoresis: Resolution of tramadol enantiomers. <i>Electrophoresis</i> , 1998, 19, 2883-2889.	1.3	24
227	Separation of Enantiomers of Drugs by Capillary Electrophoresis, Part 6. Hydroxypropyl- β -cyclodextrin as Chiral Solvating Agent. <i>Journal of High Resolution Chromatography</i> , 1998, 21, 215-224.	2.0	31
228	Chiral Separation of Bencynonate and Econazole by Cyclodextrin-Modified Capillary Zone Electrophoresis. <i>Journal of High Resolution Chromatography</i> , 1998, 21, 421-423.	2.0	10
229	Enantiomeric Separation of β -Amino Alcohols by Capillary Electrophoresis Using DM- β -CD as Chiral Selector. <i>Journal of High Resolution Chromatography</i> , 1998, 21, 440-444.	2.0	10
230	Influence of the Buffer Organic Cation on the Chiral Separation of Some Basic Drugs by Capillary Zone Electrophoresis. <i>Journal of High Resolution Chromatography</i> , 1998, 21, 445-449.	2.0	11
231	Use of heptakis-(2,3-O-dimethyl-6-sulfato)- β -cyclodextrin, a new single-isomer chiral resolving agent for the separation of the enantiomers of neutral compounds by capillary electrophoresis. <i>Journal of Separation Science</i> , 1998, 10, 293-299.	1.0	24
232	Chiral separation of chloroquine and pemoline by capillary zone electrophoresis with sulfobutyl ether β -cyclodextrin as buffer additive. <i>Chromatographia</i> , 1998, 48, 310-313.	0.7	14
233	The effect of conductivity tuning in chiral separations by CE; Using hydroxypropyl- β -cyclodextrin in combination with tetraalkylammonium ions. <i>Chromatographia</i> , 1998, 48, 415-421.	0.7	15
234	Enantiomeric separation of tramadol hydrochloride and its metabolites by cyclodextrin-mediated capillary zone electrophoresis. <i>Biomedical Applications</i> , 1998, 707, 287-294.	1.7	32

#	ARTICLE	IF	CITATIONS
235	Determination of chiral pharmaceutical compounds, terbutaline, ketamine and propranolol, by on-line capillary electrophoresis-electrospray ionization mass spectrometry. <i>Biomedical Applications</i> , 1998, 714, 69-75.	1.7	55
236	Use of mobility ratios to estimate binding constants of ligands to proteins in affinity capillary electrophoresis. <i>Biomedical Applications</i> , 1998, 715, 203-210.	1.7	61
237	Quantitative relationship between retention in HPLC and CE via binding constants and selector concentrations. <i>TrAC - Trends in Analytical Chemistry</i> , 1998, 17, 148-157.	5.8	8
238	Separation of enantiomers of drugs by capillary electrophoresis Part 8. β -Cyclodextrin as chiral solvating agent. <i>Talanta</i> , 1998, 46, 743-749.	2.9	20
239	Capillary Electrophoresis with Electrochemical Detection for Chiral Separation of Optical Isomers. <i>Analytical Chemistry</i> , 1998, 70, 4030-4035.	3.2	16
240	Control of relative migration of small inorganic and organic anions with cyclodextrins in capillary electrophoresis (CE). <i>Canadian Journal of Chemistry</i> , 1998, 76, 194-198.	0.6	8
241	Enantiomeric Separations of Terbutaline by CE with a Sulfated β -Cyclodextrin Chiral Selector: A Quantitative Binding Study. <i>Analytical Chemistry</i> , 1998, 70, 5166-5171.	3.2	101
242	Monte Carlo Simulation of Error Propagation in the Determination of Binding Constants from Rectangular Hyperbolae. 1. Ligand Concentration Range and Binding Constant. <i>Journal of Physical Chemistry A</i> , 1998, 102, 8063-8071.	1.1	68
243	Enantioselective Separations. <i>Journal of Chromatography Library</i> , 1998, 60, 197-256.	0.1	3
244	Enantiomeric resolution of galanthamine and related drugs used in anti-Alzheimer therapy by means of capillary zone electrophoresis employing derivatized cyclodextrin selectors. <i>Biomedical Applications</i> , 1999, 730, 167-175.	1.7	28
245	Non-aqueous capillary electrophoresis chiral separations with sulfated β -cyclodextrin. <i>Biomedical Applications</i> , 1999, 731, 187-197.	1.7	54
246	Recent trends in enantioseparations using capillary electromigration techniques. <i>TrAC - Trends in Analytical Chemistry</i> , 1999, 18, 485-498.	5.8	65
247	Modelling of conditions for the enantiomeric separation of β -adrenergic sympathicomimetics by capillary electrophoresis using cyclodextrins as chiral selectors in a polyethylene glycol gel. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1999, 19, 529-537.	1.4	19
248	Cyclodextrin-mediated micellar electrokinetic chromatography and matrix-assisted laser desorption/ionization time-of-flight mass spectrometry for the enantiomer separation of racemorphan in human urine. <i>Journal of Chromatography A</i> , 1999, 853, 439-447.	1.8	24
249	Rapid development of the enantiomeric separation of β -blockers by capillary electrophoresis using an experimental design approach. <i>Journal of Chromatography A</i> , 1999, 855, 681-693.	1.8	52
250	Separation of the enantiomers of four chiral drugs by neutral cyclodextrin-mediated capillary zone electrophoresis. <i>Chromatographia</i> , 1999, 49, 411-414.	0.7	9
251	Principle and applications of the partial filling technique in capillary electrophoresis. <i>Chromatographia</i> , 1999, 50, 497-506.	0.7	86
252	Chiral separation and determination of the optical purity of thiamphenicol-related chiral compound by capillary zone electrophoresis with cyclodextrins as chiral selectors. <i>Chromatographia</i> , 1999, 50, 317-320.	0.7	13

#	ARTICLE	IF	CITATIONS
253	Separation of chiral basic drugs with sulfobutyl- β -cyclodextrin in capillary electrophoresis. <i>Chromatographia</i> , 1999, 50, 363-368.	0.7	14
254	Riboflavin binding proteins as chiral selectors in HPLC and CE. <i>Pharmaceutical Science & Technology Today</i> , 1999, 2, 352-364.	0.7	8
255	Capillary Electrophoresis as an Efficient Tool in the Determination of Low Association Constants between Polyhydroxy Compound and Samarium Cation. <i>Journal of High Resolution Chromatography</i> , 1999, 22, 595-598.	2.0	2
256	Estimation of the pH-independent binding constants of alanylphenylalanine and leucylphenylalanine stereoisomers with β -cyclodextrin in the presence of urea. <i>Electrophoresis</i> , 1999, 20, 171-179.	1.3	60
257	Determination of association constants between enantiomers of orciprenaline and methyl- β -cyclodextrin as chiral selector by capillary zone electrophoresis using a partial filling technique. <i>Electrophoresis</i> , 1999, 20, 180-188.	1.3	31
258	Enantioseparation of tropa alkaloids by means of anionic cyclodextrin-modified capillary electrophoresis. <i>Electrophoresis</i> , 1999, 20, 1555-1560.	1.3	25
259	Separation of drugs by capillary electrophoresis, Part 10. Permethyl-alpha-cyclodextrin as chiral solvating agent. <i>Electrophoresis</i> , 1999, 20, 1878-1889.	1.3	20
260	Capillary electrophoresis enantioseparation of drugs using β -cyclodextrin polymer: Intramolecular synergistic effect. <i>Electrophoresis</i> , 1999, 20, 1900-1903.	1.3	6
261	Optical isomer separation of potential analgesic drug candidates by using capillary electrophoresis. <i>Electrophoresis</i> , 1999, 20, 2432-2437.	1.3	8
262	Considerations concerning interaction characterization of oligopeptide mixtures with vancomycin using affinity capillary electrophoresis-electrospray mass spectrometry. <i>Electrophoresis</i> , 1999, 20, 2462-2474.	1.3	33
263	Selector-selectand interactions in chiral capillary electrophoresis. <i>Electrophoresis</i> , 1999, 20, 2592-2604.	1.3	41
264	Comparison of sulfobutylether- and sulfated- β -cyclodextrins as additives for the chiral separation of basic spirobenzopyrans by capillary electrophoresis. <i>Electrophoresis</i> , 1999, 20, 2630-2637.	1.3	17
265	Application of sulfobutylether- β -cyclodextrin with specific degrees of substitution for the enantioseparation of pharmaceutical mixtures by capillary electrophoresis. <i>Electrophoresis</i> , 1999, 20, 2638-2649.	1.3	42
266	Binding constant dependency of amphetamines with various commercial methylated β -cyclodextrins. <i>Electrophoresis</i> , 1999, 20, 2670-2679.	1.3	30
267	Designed combination of chiral selectors for adjustment of enantioseparation selectivity in capillary electrophoresis. <i>Electrophoresis</i> , 1999, 20, 2691-2697.	1.3	88
268	Resolution of ephedrine derivatives by means of neutral and sulfatedheptakis(2,3-di-O-acetyl) β -cyclodextrins using capillary electrophoresis and nuclear magnetic resonance spectroscopy. <i>Electrophoresis</i> , 1999, 20, 2698-2704.	1.3	44
269	pKa shift-associated effects in enantioseparations by cyclodextrin-mediated capillary zone electrophoresis. <i>Electrophoresis</i> , 1999, 20, 2715-2722.	1.3	91
270	Equilibrium binding model of bile salt-mediated chiral micellar electrokinetic capillary chromatography. <i>Electrophoresis</i> , 1999, 20, 2754-2760.	1.3	9

#	ARTICLE	IF	CITATIONS
271	Enantioseparations by capillary electrophoresis using chiral glycosidic surfactants. I. Evaluation of cyclohexyl-pentyl- β -D-maltoside surfactant. <i>Electrophoresis</i> , 1999, 20, 2766-2771.	1.3	9
272	Selectivity in capillary electrokinetic separations. <i>Electrophoresis</i> , 1999, 20, 2989-3010.	1.3	38
273	Identification, quantitation, and characterization of biomolecules by capillary electrophoretic analysis of binding interactions. <i>Electrophoresis</i> , 1999, 20, 3122-3133.	1.3	148
274	Enantiomeric separation of glycyI dipeptides by capillary electrophoresis with cyclodextrins as chiral selectors. <i>Electrophoresis</i> , 1999, 20, 3417-3423.	1.3	21
275	Enantiomeric separation of fluoxetine and norfluoxetine in plasma and serum samples with high detection sensitivity capillary electrophoresis. <i>Electrophoresis</i> , 1999, 20, 3432-3438.	1.3	67
276	Enantiomeric resolution of some nonsteroidal antiinflammatory and anticoagulant drugs using β -cyclodextrins by capillary electrophoresis. <i>Chirality</i> , 1999, 11, 56-62.	1.3	33
277	Resolution of enantiomeric N-oxides by capillary electrophoresis using cyclodextrins as chiral selectors. <i>Chirality</i> , 1999, 11, 409-415.	1.3	13
278	Capillary electrophoresis chiral separation of basic pharmaceutical enantiomers with different charges using sulfated β -cyclodextrin. <i>Journal of Separation Science</i> , 1999, 11, 11-21.	1.0	31
279	Enantiomer separations of phenylephrine and synephrine by capillary electrochromatography on bare silica stationary phase using hydroxypropyl- β -cyclodextrin as a mobile phase additive. <i>Journal of Separation Science</i> , 1999, 11, 263-269.	1.0	30
280	Separation behavior of amino acid enantiomers in ligand exchange micellar electrokinetic chromatography. <i>Journal of Separation Science</i> , 1999, 11, 534-540.	1.0	29
281	Direct chiral resolution of aliphatic β -hydroxy acids using 2-hydroxypropyl- β -cyclodextrin in capillary electrophoresis. <i>Analyst</i> , 1999, 124, 55-59.	1.7	27
282	Symmetry Breaking during the Formation of β -Cyclodextrin-Imidazole Inclusion Compounds: A Capillary Electrophoresis Study. <i>Analytical Chemistry</i> , 1999, 71, 2046-2052.	3.2	36
283	Self-Association of Rifamycin B: Possible Effects on Molecular Recognition. <i>Journal of Physical Chemistry B</i> , 1999, 103, 4338-4341.	1.2	6
284	Monte Carlo Simulation of Error Propagation in the Determination of Binding Constants from Rectangular Hyperbolae. 2. Effect of the Maximum-Response Range. <i>Journal of Physical Chemistry A</i> , 1999, 103, 197-202.	1.1	60
285	Fast Analytical-Scale Separations by Capillary Electrophoresis and Liquid Chromatography. <i>Chemical Reviews</i> , 1999, 99, 3081-3132.	23.0	68
286	Reversal Behaviors of the Enantiomer Migration Order and the Stereo-selectivity of Cu(II) Complex with Amino Acid Enantiomers in Ligand Exchange-Micellar Electrokinetic Chromatography. <i>Analytical Sciences</i> , 2000, 16, 131-137.	0.8	41
288	High-speed chiral separations on microchip electrophoresis devices. <i>Electrophoresis</i> , 2000, 21, 211-219.	1.3	104
289	Capillary electrophoresis behavior of water-soluble anionic porphyrins in the presence of β -cyclodextrin and its O-methylated derivatives. <i>Electrophoresis</i> , 2000, 21, 619-626.	1.3	14

#	ARTICLE	IF	CITATIONS
290	Cellulases from the fungi <i>Phanerochaete chrysosporium</i> and <i>Trichoderma reesei</i> as chiral selectors in capillary electrophoresis: Applications with displacer plugs and sample preconcentration. <i>Electrophoresis</i> , 2000, 21, 1587-1596.	1.3	19
291	Simultaneous chiral separation of triadimefon and triadimenol by sulfated β -cyclodextrin-mediated capillary electrophoresis. <i>Electrophoresis</i> , 2000, 21, 1611-1619.	1.3	39
292	Experimental verification of a predicted, hitherto unseen separation selectivity pattern in the nonaqueous capillary electrophoretic separation of weak base enantiomers by octakis(2,3-diacetyl-6-sulfato)- β -cyclodextrin. <i>Electrophoresis</i> , 2000, 21, 2016-2024.	1.3	45
293	Enantiomeric separation of alanyl and leucyl dipeptides by capillary electrophoresis with cyclodextrins as chiral selectors. <i>Electrophoresis</i> , 2000, 21, 2397-2404.	1.3	21
294	Recent innovations in the use of charged cyclodextrins in capillary electrophoresis for chiral separations in pharmaceutical analysis. <i>Electrophoresis</i> , 2000, 21, 3220-3239.	1.3	156
295	Fast enantiomeric separation of uniconazole and diniconazole by electrokinetic chromatography using an anionic cyclodextrin: Application to the determination of analyte-selector apparent binding constants for enantiomers. <i>Electrophoresis</i> , 2000, 21, 3240-3248.	1.3	23
296	Experimental verification of a predicted, previously unseen separation selectivity pattern in the capillary electrophoretic separation of noncharged enantiomers by octakis(2,3-diacetyl-6-sulfato)- β -cyclodextrin. <i>Electrophoresis</i> , 2000, 21, 3249-3256.	1.3	16
297	Competitive binding of a Tris run buffer with chiral crown ether in chiral capillary electrophoresis. <i>Electrophoresis</i> , 2000, 21, 3618-3624.	1.3	17
298	Separation of Enantiomers of Drugs by Capillary Electrophoresis with Permethy- γ -Cyclodextrin as Chiral Solvating Agent. <i>Journal of High Resolution Chromatography</i> , 2000, 23, 413-429.	2.0	26
299	Enantiomeric separations by use of polymeric surfactant electrokinetic chromatography. <i>Journal of Chromatography A</i> , 2000, 875, 179-206.	1.8	70
300	Enantiomeric separations of drugs using mixtures of charged and neutral cyclodextrins. <i>Journal of Chromatography A</i> , 2000, 875, 123-134.	1.8	170
301	Enantiomeric separations by nonaqueous capillary electrophoresis. <i>Journal of Chromatography A</i> , 2000, 875, 277-293.	1.8	92
302	Calculation of stability constants for the chiral selector-enantiomer interactions from electrophoretic mobilities. <i>Journal of Chromatography A</i> , 2000, 875, 431-445.	1.8	50
303	Chiral glycosidic surfactants for enantiomeric separation in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2000, 875, 207-233.	1.8	45
304	Optimization of the cyclodextrin-assisted capillary electrophoresis separation of the enantiomers of phenoxyacid herbicides. <i>Journal of Chromatography A</i> , 2000, 875, 423-430.	1.8	28
305	tert-Butylcarbamoylequinine as chiral ion-pair agent in non-aqueous enantioselective capillary electrophoresis applying the partial filling technique. <i>Journal of Chromatography A</i> , 2000, 892, 509-521.	1.8	42
306	New pseudo-stationary phases for electrokinetic capillary chromatography. <i>Journal of Chromatography A</i> , 2000, 876, 169-182.	1.8	5
307	Analysis of polymyxin B sulfate by capillary zone electrophoresis with cyclodextrin as additive. <i>Journal of Chromatography A</i> , 2000, 879, 211-218.	1.8	35

#	ARTICLE	IF	CITATIONS
308	Fast development of separation methods for the chiral analysis of amino acid derivatives using capillary electrophoresis and experimental designs. <i>Journal of Chromatography A</i> , 2000, 883, 249-265.	1.8	42
309	Development and evaluation of a linear regression method for the prediction of maximal chiral separation of basic drug racemates by cyclodextrin-mediated capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 2000, 903, 245-254.	1.8	24
310	Direct chiral resolution of lactic acid in food products by capillary electrophoresis. <i>Journal of Chromatography A</i> , 2000, 875, 371-377.	1.8	47
311	Chiral capillary electrophoretic analysis of the enantiomeric purity of a pharmaceutical compound using sulfated β -cyclodextrin. <i>Journal of Chromatography A</i> , 2000, 875, 389-401.	1.8	47
312	Separation of drug enantiomers by capillary electrophoresis in the presence of neutral cyclodextrins. <i>Journal of Chromatography A</i> , 2000, 875, 135-161.	1.8	71
313	Design of experiments for capillary electrophoretic enantioresolution of salbutamol using dermatan sulfate. <i>Journal of Chromatography A</i> , 2000, 875, 411-422.	1.8	58
314	Enantioselective determination by capillary electrophoresis with cyclodextrins as chiral selectors. <i>Journal of Chromatography A</i> , 2000, 875, 89-122.	1.8	444
315	Enantiomeric separation of selenoaminoacid derivatives by cyclodextrin-modified micellar electrokinetic chromatography using a mixed micellar system of sodium dodecyl sulphate and taurodeoxycholic acid. <i>Analytica Chimica Acta</i> , 2000, 416, 1-7.	2.6	34
316	New derivative of β -cyclodextrin as chiral selectors for the capillary electrophoretic separation of chiral drugs. <i>Analytica Chimica Acta</i> , 2000, 421, 27-34.	2.6	26
317	Advanced statistical evaluation of complex formation constant from electrophoretic data. <i>Analytica Chimica Acta</i> , 2000, 407, 327-336.	2.6	10
318	Dependence of chiral separations on the amount of cyclodextrins as selectors, employing the partial filling technique in capillary zone electrophoresis. <i>Chromatographia</i> , 2000, 51, 226-230.	0.7	13
319	Selectivity difference between carboxyethyl- β -cyclodextrin and carboxymethyl- β -cyclodextrin in enantioseparations of drugs by capillary electrophoresis. <i>Chromatographia</i> , 2000, 51, 634-637.	0.7	2
320	Macrocyclic antibiotics as effective chiral selectors for enantiomeric resolution by liquid chromatography and capillary electrophoresis. <i>Chromatographia</i> , 2000, 52, 679-691.	0.7	108
321	Permethyl monoamino β -cyclodextrin a new chiral selective agent for capillary electrophoresis. <i>Chromatographia</i> , 2000, 53, 166-172.	0.7	17
322	CHIRAL SEPARATIONS Capillary Electrophoresis. , 2000, , 2287-2294.		1
323	Capillary Zone Electrophoresis. , 2000, , 139-208.		12
324	Capillary Zone Electrophoresis. , 2000, , 73-138.		7
325	Chiral Separations in Capillary Electrophoresis. <i>Chemical Reviews</i> , 2000, 100, 3715-3754.	23.0	168

#	ARTICLE	IF	CITATIONS
326	6.3. Capillary electrophoretic (CE) methods. Progress in Pharmaceutical and Biomedical Analysis, 2000, 4, 540-552.	0.1	0
327	Enantioseparation Of Aromatic Dipeptides Using Carboxymethyl- β -Cyclodextrin Polymer As Chiral Selector By Capillary Electrophoresis. Analytical Letters, 2000, 33, 1147-1165.	1.0	8
328	Recent advances in enantiomer separations by affinity capillary electrophoresis using proteins and peptides. Journal of Proteomics, 2001, 48, 103-116.	2.4	72
329	Quantitative determination of clenbuterol, salbutamol and tulobuterol enantiomers by capillary electrophoresis. Fresenius' Journal of Analytical Chemistry, 2001, 369, 212-219.	1.5	25
330	Enantioseparation of chiral phenothiazine derivatives in capillary electrophoresis using cyclodextrin type chiral selectors. Chromatographia, 2001, 53, S290-S295.	0.7	34
331	Chiral separation of newly synthesized arylpropionic acids by capillary electrophoresis using cyclodextrins or a glycopeptide antibiotic as chiral selectors. Chromatographia, 2001, 54, 87-92.	0.7	10
333	Comparison of binding of tetraphenylborate and tetraphenylphosphonium ions to cyclodextrins studied by capillary electrophoresis. Electrophoresis, 2001, 22, 117-122.	1.3	13
334	Chiral analysis of baclofen by β -cyclodextrin-modified capillary electrophoresis and laser-induced fluorescence detection. Electrophoresis, 2001, 22, 123-127.	1.3	39
335	Influence of neutral cyclodextrin concentration on plate numbers in capillary electrophoresis. Electrophoresis, 2001, 22, 1957-1973.	1.3	6
336	Modeling and optimization of the chiral selectivity of basic analytes in chiral capillary electrophoresis with negatively charged cyclodextrins using electrochemical detection. Electrophoresis, 2001, 22, 2025-2031.	1.3	15
337	Fundamental aspects of chiral separations by capillary electrophoresis. Electrophoresis, 2001, 22, 3079-3106.	1.3	215
338	Recent developments in chiral capillary electrophoresis and applications of this technique to pharmaceutical and biomedical analysis. Electrophoresis, 2001, 22, 3107-3130.	1.3	191
339	pH-Dependence of complexation constants and complex mobility in capillary electrophoresis separations of dipeptide enantiomers. Electrophoresis, 2001, 22, 3163-3170.	1.3	49
340	Rapid screening for chiral separations by short-end injection capillary electrophoresis using highly sulfated cyclodextrins as chiral selectors. Electrophoresis, 2001, 22, 3203-3215.	1.3	88
341	Chiral separation of pyrethroic acids with single isomer permethyl monoamino β -cyclodextrin selector. Electrophoresis, 2001, 22, 3232-3236.	1.3	22
342	Enantioseparation of chiral thiobarbiturates using cyclodextrin-modified capillary electrophoresis. Electrophoresis, 2001, 22, 3237-3242.	1.3	8
343	Experimental designs to investigate capillary electrophoresis-electrospray ionization-mass spectrometry enantioseparation with the partial-filling technique. Electrophoresis, 2001, 22, 3316-3326.	1.3	74
344	Cyclodextrin-assisted capillary electrophoretic resolution of 1,1'-bi-2-naphthol atropisomers. Electrophoresis, 2001, 22, 3578-3582.	1.3	6

#	ARTICLE	IF	CITATIONS
345	Interaction between 18-crown-6-tetracarboxylic acid and positional substituents of enantiomers and simultaneous separation of positional enantiomers of methyl-DL-tryptophans by capillary electrophoresis. <i>Electrophoresis</i> , 2001, 22, 2136-2142.	1.3	11
346	Electrophoretic chiral separation of pharmaceutical compounds with multiple stereogenic centers in charged cyclodextrin media. <i>Journal of Separation Science</i> , 2001, 24, 378-384.	1.3	15
347	Chiral separation of promethazine by capillary electrophoresis with end-column amperometric detection. <i>Journal of Separation Science</i> , 2001, 24, 658-662.	1.3	12
348	Evaluation of enantioselective nonaqueous ion-pair capillary electrophoresis as screening assay in the development of new ion exchange type chiral stationary phases. <i>Journal of Separation Science</i> , 2001, 24, 706-716.	1.3	21
349	High-performance chiral separation of fourteen triazole fungicides by sulfated β -cyclodextrin-mediated capillary electrophoresis. <i>Journal of Chromatography A</i> , 2001, 912, 171-179.	1.8	112
350	Determination of apparent binding constants of drugs by capillary electrophoresis using β -cyclodextrin as ligand and three different linear plotting methods. <i>Journal of Chromatography A</i> , 2001, 924, 483-491.	1.8	31
351	Comparison of monomeric and polymeric amino acid based surfactants for chiral separations. <i>Journal of Chromatography A</i> , 2001, 922, 329-338.	1.8	46
352	Non-aqueous capillary electrophoretic separation of enantiomeric amines with (α)-2,3:4,6-di-O-isopropylidene-2-keto-L-gulonic acid as chiral counter ion. <i>Journal of Chromatography A</i> , 2001, 922, 303-311.	1.8	39
353	Mechanistic study of enantiomeric recognition of primary amino compounds using an achiral crown ether with cyclodextrin by capillary electrophoresis and nuclear magnetic resonance. <i>Journal of Chromatography A</i> , 2001, 923, 229-239.	1.8	18
354	Chiral separation of six diastereomeric flavanone-7-O-glycosides by capillary electrophoresis and analysis of lemon juice. <i>Journal of Chromatography A</i> , 2001, 925, 279-289.	1.8	48
355	Rapid chiral on-chip separation with simplified amperometric detection. <i>Journal of Chromatography A</i> , 2001, 928, 225-232.	1.8	65
356	Enantioseparations in capillary electromigration techniques: recent developments and future trends. <i>Journal of Chromatography A</i> , 2001, 906, 309-363.	1.8	331
357	Enantioseparation of flobufen with cyclodextrins studied by capillary electrophoresis and NMR. <i>Pharmaceutical Research</i> , 2001, 18, 886-893.	1.7	8
358	Optimization Strategies and Modeling of Separations of Dansyl-Amino Acids by Cyclodextrin-Modified Capillary Electrophoresis. <i>Journal of Chromatographic Science</i> , 2001, 39, 411-419.	0.7	10
359	ENANTIOMERIC SEPARATION OF A DRUG SUBSTANCE USING CAPILLARY ELECTROPHORESIS WITH SULFATED- β -CYCLODEXTRIN. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2002, 25, 2999-3015.	0.5	4
360	Enantioseparation of Vinclozolin by β -Cyclodextrin-Modified Micellar Electrokinetic Chromatography. <i>Journal of Agricultural and Food Chemistry</i> , 2002, 50, 1312-1317.	2.4	27
361	Accurately describing weak analyte-additive interactions by capillary electrophoresis. <i>Electrophoresis</i> , 2002, 23, 880-888.	1.3	21
362	Drug-plasma protein binding assay by electrokinetic chromatography-frontal analysis. <i>Electrophoresis</i> , 2002, 23, 951-955.	1.3	29

#	ARTICLE	IF	CITATIONS
363	Chiral capillary electrophoretic determination of the enantiomeric purity of tetrahydronaphthalenic derivatives, melatonergic ligands, using highly sulfated β -cyclodextrins. <i>Electrophoresis</i> , 2002, 23, 1121-1128.	1.3	20
364	Chiral recognition of functionalized cyclodextrins in capillary electrophoresis. <i>Electrophoresis</i> , 2002, 23, 1308-1313.	1.3	19
365	Modelling, optimisation and control of selectivity in the separation of aromatic bases by electrokinetic chromatography using a neutral cyclodextrin as a pseudostationary phase. <i>Electrophoresis</i> , 2002, 23, 1844.	1.3	8
366	Separation of reboxetine enantiomers by means of capillary electrophoresis. <i>Electrophoresis</i> , 2002, 23, 1870.	1.3	24
367	Electrolyte and additive effects on enantiomer separation of peptides by nonaqueous ion-pair capillary electrophoresis using tert.-butylcarbamoilquinine as chiral counterion. <i>Electrophoresis</i> , 2002, 23, 1887.	1.3	38
368	Combinatorial enantiomeric separation of diverse compounds using capillary array electrophoresis. <i>Electrophoresis</i> , 2002, 23, 2996-3005.	1.3	35
369	Enantioselective analysis of thiobencarb sulfoxide produced by metabolism of thiobencarb by hydroxypropyl- β -cyclodextrin modified micellar electrokinetic chromatography. <i>Journal of Separation Science</i> , 2002, 25, 1055-1062.	1.3	18
370	Enantiomer migration order in chiral capillary electrophoresis. <i>Electrophoresis</i> , 2002, 23, 4022-4035.	1.3	132
371	Optimized separation of pharmacologically active xanthenes from <i>Securidaca inappendiculata</i> by capillary electrophoresis. <i>Chromatographia</i> , 2002, 55, 217-223.	0.7	16
372	Knowledge-based system for method development of chiral separations with capillary electrophoresis using highly-sulphated cyclodextrins. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2002, 27, 515-529.	1.4	52
373	Use of negatively charged cyclodextrins for the simultaneous enantioseparation of selected anesthetic drugs by capillary electrophoresis- μ mass spectrometry. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2002, 27, 615-626.	1.4	45
374	Selected fundamental aspects of chiral electromigration techniques and their application to pharmaceutical and biomedical analysis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2002, 27, 373-399.	1.4	128
375	Chiral separation of methoxamine and lobeline in capillary zone electrophoresis using ethylbenzene-deactivated fused-silica capillary columns and cyclodextrins as buffer additives. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2002, 29, 999-1003.	1.4	14
376	Estimation of binding constants by capillary electrophoresis. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2002, 768, 81-92.	1.2	198
377	Chiral separation of amino acids derivatized with fluoresceine-5-isothiocyanate by capillary electrophoresis and laser-induced fluorescence detection using mixed selectors of β -cyclodextrin and sodium taurocholate. <i>Journal of Chromatography A</i> , 2002, 955, 133-140.	1.8	76
378	Quantitative approach for the screening of cyclodextrins by nuclear magnetic resonance spectroscopy in support of chiral separations in liquid chromatography and capillary electrophoresis. <i>Journal of Chromatography A</i> , 2002, 961, 257-276.	1.8	30
379	Separation and migration behavior of structurally related phenothiazines in cyclodextrin-modified capillary zone electrophoresis. <i>Journal of Chromatography A</i> , 2002, 979, 399-408.	1.8	38
380	Chiral separation of methadone, 2-ethylidene-1,5-dimethyl-3,3-diphenylpyrrolidine (EDDP) and 2-ethyl-5-methyl-3,3-diphenyl-1-pyrrolidine (EMDP) by capillary electrophoresis using cyclodextrin derivatives. <i>Electrophoresis</i> , 2003, 24, 2106-2110.	1.3	16

#	ARTICLE	IF	CITATIONS
381	Salt dependence on vancomycin-herbicide enantiomer binding: Capillary electrophoresis study and theoretical approach. <i>Electrophoresis</i> , 2003, 24, 1620-1626.	1.3	11
382	Direct chiral resolution of tartaric acid by ion-pair capillary electrophoresis using an aqueous background electrolyte with (1R,2R)-($\hat{\alpha}$)-1,2-diaminocyclohexane as a chiral counterion. <i>Electrophoresis</i> , 2003, 24, 2711-2715.	1.3	11
383	pH-dependent cyclodextrin capillary electrophoresis resolution of atropisomers. <i>Electrophoresis</i> , 2003, 24, 2456-2461.	1.3	12
384	Large-volume sample stacking combined with separation by 2-hydroxypropyl- $\hat{\beta}$ -cyclodextrin for analysis of isoxazolylpenicillins by capillary electrophoresis. <i>Electrophoresis</i> , 2003, 24, 3089-3096.	1.3	21
385	Complexation trends and binding constants between dextrin oligomers and small molecules as measured through affinity capillary electrophoresis. <i>Electrophoresis</i> , 2003, 24, 2914-2923.	1.3	8
386	Recent advances in enantioseparations of peptides by capillary electrophoresis. <i>Electrophoresis</i> , 2003, 24, 4063-4077.	1.3	51
387	Applications of affinity interactions in capillary electrophoresis. <i>Electrophoresis</i> , 2003, 24, 3879-3891.	1.3	83
388	Migration order of dipeptide and tripeptide enantiomers in the presence of single isomer and randomly sulfated cyclodextrins as a function of pH. <i>Electrophoresis</i> , 2003, 24, 1069-1076.	1.3	48
389	A high-throughput on-line microdialysis-capillary assay for D-serine. <i>Electrophoresis</i> , 2003, 24, 1227-1235.	1.3	52
390	Separation of isomers of dienolic acids by electromigration techniques. <i>Journal of Separation Science</i> , 2003, 26, 1253-1258.	1.3	7
391	Enantioseparation of dansyl amino acids on terguride-based chiral selectors. Part I: Capillary electrophoretic separation. <i>Journal of Separation Science</i> , 2003, 26, 851-856.	1.3	11
392	Separation of diastereomers of flavanone-7-O-glycosides by capillary electrophoresis using sulfobutyl ether- $\hat{\beta}$ -cyclodextrin as the selector. <i>Journal of Separation Science</i> , 2003, 26, 844-850.	1.3	30
393	Enantiomeric separation of aminoglutethimide by capillary electrophoresis using native cyclodextrins in single and dual systems. <i>Journal of Separation Science</i> , 2003, 26, 536-542.	1.3	13
394	Chiral separation of 2,3-allenoic acid by capillary zone electrophoresis using cyclodextrin derivatives. <i>Chirality</i> , 2003, 15, 201-205.	1.3	5
395	Comprehensive strategy for chiral separations using sulfated cyclodextrins in capillary electrophoresis. <i>Chirality</i> , 2003, 15, 709-723.	1.3	107
396	Mechanistic study of enantiomeric recognition with native $\hat{\beta}$ -cyclodextrin by capillary electrophoresis, reversed-phase liquid chromatography, nuclear magnetic resonance spectroscopy, electrospray mass spectrometry and circular dichroism techniques. <i>Journal of Chromatography A</i> , 2003, 987, 409-420.	1.8	37
397	Determination of free l- and d-alanine in hydrolysed protein fertilisers by capillary electrophoresis. <i>Journal of Chromatography A</i> , 2003, 985, 463-469.	1.8	17
398	Exploration of the electrophoretic behaviour of borane cluster anions and of the capability of capillary electrophoresis to separate them chirally. <i>Journal of Chromatography A</i> , 2003, 984, 121-134.	1.8	17

#	ARTICLE	IF	CITATIONS
399	Robustness testing of chiral separations by capillary electrophoresis using highly-sulfated cyclodextrins. <i>Journal of Chromatography A</i> , 2003, 1007, 165-177.	1.8	40
400	Chiral separation of dioxopromethazine in eye drops by CZE with charged cyclodextrin. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2003, 33, 157-164.	1.4	17
401	Simultaneous enantioselective separation of azelastine and three of its metabolites for the investigation of the enantiomeric metabolism in rats. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2003, 793, 389-404.	1.2	21
402	Enantioseparation using cyclophosphates as a novel chiral additive in capillary electrophoresis. <i>Carbohydrate Research</i> , 2003, 338, 1143-1146.	1.1	39
403	Indirect capillary electrophoresis with 8-anilino-1-naphthalenesulfonic acid as a fluorescence probe for determining the apparent stability constant of an inclusion complex formed between a cyclodextrin and a solute. <i>Journal of Chromatography A</i> , 2003, 987, 485-492.	1.8	5
404	Chiral separation of amines with N-benzoxycarbonylglycyl-L-proline as selector in non-aqueous capillary electrophoresis using methanol and 1,2-dichloroethane in the background electrolyte. <i>Journal of Chromatography A</i> , 2003, 984, 261-271.	1.8	29
405	Determination of the binding constant for the inclusion complex between procaine hydrochloride and β -cyclodextrin by capillary electrophoresis. <i>Talanta</i> , 2003, 59, 493-499.	2.9	24
406	Capillary Electrophoretic Behaviors of Pharmacologically Active Xanthenes from <i>Securidaca inappendiculata</i> with β -Cyclodextrin as a Buffer Additive. <i>Journal of Chromatographic Science</i> , 2003, 41, 182-186.	0.7	7
407	Use of cyclodextrins for separation of enantiomers. <i>Russian Chemical Reviews</i> , 2003, 72, 1035-1054.	2.5	32
408	Chiral Separation by Capillary Electrophoresis Using Polysaccharides. , 2004, 243, 343-354.		3
409	Chiral Separation of Deprenyl-N-Oxide Isomers by Capillary Electrophoresis Using Various Cyclodextrin Derivatives. <i>Journal of Chromatographic Science</i> , 2004, 42, 21-26.	0.7	10
411	Determination of equilibrium constants from chromatographic and electrophoretic measurements. <i>Journal of Chromatography A</i> , 2004, 1037, 15-28.	1.8	50
412	Separation and migration behavior of positional and structural naphthalenesulfonate isomers by cyclodextrin-mediated capillary electrophoresis. <i>Journal of Chromatography A</i> , 2004, 1033, 167-172.	1.8	20
413	Enantioseparation of four cis and trans diastereomers of 2',3'-didehydro-2',3'-dideoxythymidine analogs, by high-performance liquid chromatography and capillary electrophoresis. <i>Journal of Chromatography A</i> , 2004, 1034, 161-167.	1.8	17
414	Enantiomeric analysis of simendan by CE with β -CD as chiral selector compared with CMPA-HPLC. <i>Biomedical Chromatography</i> , 2004, 18, 866-871.	0.8	8
415	Enantioseparation of new nucleoside analogs, related to d4T and acyclovir, by chiral capillary electrophoresis using highly sulfated β -cyclodextrins. <i>Electrophoresis</i> , 2004, 25, 444-453.	1.3	15
416	Optimisation of selectivity in the separation of aromatic amino acid enantiomers using sulfated β -cyclodextrin and dextran sulfate as pseudostationary phases. <i>Electrophoresis</i> , 2004, 25, 270-276.	1.3	16
417	Chiral capillary electrophoretic resolution of baclofen, gabaergic ligand, using highly sulfated cyclodextrins. <i>Electrophoresis</i> , 2004, 25, 1111-1119.	1.3	18

#	ARTICLE	IF	CITATIONS
418	Direct enantioseparation of catechin and epicatechin in tea drinks by 6-O- β -D-glucosyl- β -cyclodextrin-modified micellar electrokinetic chromatography. <i>Electrophoresis</i> , 2004, 25, 2892-2898.	1.3	44
419	Chiral separation of N-imidazole derivatives, aromatase inhibitors, by cyclodextrin-capillary zone electrophoresis. Mechanism of enantioselective recognition. <i>Electrophoresis</i> , 2004, 25, 2735-2744.	1.3	25
420	Influence of methanol on the enantioresolution of antihistamines with carboxymethyl- β -cyclodextrin in capillary electrophoresis. <i>Electrophoresis</i> , 2004, 25, 2838-2847.	1.3	20
421	Development and validation of a capillary electrophoresis method for the enantiomeric purity determination of SLV307, a basic potential antipsychotic compound. <i>Electrophoresis</i> , 2004, 25, 2854-2859.	1.3	10
422	Influence of (hydroxy)alkylamino substituents on enantioseparation ability of single-isomer amino- β -cyclodextrin derivatives in chiral capillary electrophoresis. <i>Electrophoresis</i> , 2004, 25, 2675-2686.	1.3	45
423	Ultrashort partial-filling technique in capillary electrophoresis for infinite resolution of tramadol enantiomers and its metabolites with highly sulfated cyclodextrins. <i>Electrophoresis</i> , 2004, 25, 2761-2771.	1.3	33
424	Differential effects of organic modifiers on the enantioseparation of dimetindene maleate with carboxymethyl- β -cyclodextrin in capillary electrophoresis. <i>Journal of Separation Science</i> , 2004, 27, 21-27.	1.3	14
425	Selectivity control in the separation of aromatic amino acid enantiomers with sulphated β -cyclodextrin. <i>Journal of Chromatography A</i> , 2004, 1031, 179-186.	1.8	9
426	Direct separation, identification and quantification of epimers 22R and 22S of budesonide by capillary gas chromatography on a short analytical column with Rtx Å -5 stationary phase. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2004, 803, 191-200.	1.2	7
427	Evaluation of newly synthesized derivative of cyclodextrin for the capillary electrophoretic separation. <i>Journal of Chromatography A</i> , 2004, 1059, 181-189.	1.8	21
428	Enantiomeric separations of some acidic compounds with cationic cyclodextrin by capillary electrophoresis. <i>Analytica Chimica Acta</i> , 2004, 517, 95-101.	2.6	20
429	Enantiomeric analysis of rivastigmine in pharmaceuticals by cyclodextrin-modified capillary zone electrophoresis. <i>Analytica Chimica Acta</i> , 2004, 525, 43-51.	2.6	20
430	Effect of Different Substituted Position of Phenylcarbamoyl- β -CD on the Enantioseparation of Drugs by Capillary Electrophoresis. <i>Analytical Letters</i> , 2004, 37, 293-306.	1.0	1
431	Combined approach using capillary electrophoresis and NMR spectroscopy for an understanding of enantioselective recognition mechanisms by cyclodextrins. <i>Chemical Society Reviews</i> , 2004, 33, 337.	18.7	166
432	Analysis of Ion-Association Reaction in Aqueous Solution and Its Utilization by Capillary Zone Electrophoresis. <i>Analytical Sciences</i> , 2004, 20, 255-265.	0.8	22
433	2-O-(2-Hydroxybutyl)- β -cyclodextrin as a Chiral Selector for the Capillary Electrophoretic Separation of Chiral Drugs. <i>Analytical Sciences</i> , 2005, 21, 959-962.	0.8	18
434	Simultaneous Chiral Analysis of Methamphetamine and Related Compounds by Capillary Electrophoresis/Mass Spectrometry Using Anionic Cyclodextrin. <i>Analytical Sciences</i> , 2005, 21, 15-19.	0.8	44
435	Rapid Enantioseparation of 1-Aminoindan by Microchip Electrophoresis with Linear-Imaging UV Detection. <i>Analytical Sciences</i> , 2005, 21, 61-65.	0.8	43

#	ARTICLE	IF	CITATIONS
436	Enantioselective capillary electrophoretic separation of tryptophane- and tyrosine-methylesters in a dual system with a tetra-oxadiazacrown-ether derivative and a cyclodextrin. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2005, 38, 601-608.	1.4	33
437	Chiral separation of tamsulosin by capillary electrophoresis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2005, 39, 691-696.	1.4	30
438	Enantioseparation of dansyl amino acids by a novel permanently positively charged single-isomer cyclodextrin: Mono-6-N-allylammonium-6-deoxy- β -cyclodextrin chloride by capillary electrophoresis. <i>Analytica Chimica Acta</i> , 2005, 546, 119-125.	2.6	28
439	Enantiomeric separation of 8 hydroxy, 10 carboxylic and 6 dansyl amino acids by mono(6-amino-6-deoxy)- β -cyclodextrin in capillary electrophoresis. <i>Analytica Chimica Acta</i> , 2005, 554, 156-162.	2.6	28
440	Affinity capillary electrophoresis on microchips. <i>Journal of Chromatography A</i> , 2005, 1063, 217-225.	1.8	42
441	Chiral separations by capillary electromigration techniques in nonaqueous media. <i>Journal of Chromatography A</i> , 2005, 1068, 3-30.	1.8	79
442	Comparison of vancomycin-based stationary phases with different chiral selector coverage for enantioselective separation of selected drugs in high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2005, 1088, 94-103.	1.8	55
443	Synthesis and application of mono-6-ammonium-6-deoxy- β -cyclodextrin chloride as chiral selector for capillary electrophoresis. <i>Journal of Chromatography A</i> , 2005, 1094, 187-191.	1.8	35
444	Separation and determination of norepinephrine, epinephrine and isoprenaline enantiomers by capillary electrophoresis in pharmaceutical formulation and human serum. <i>Journal of Chromatography A</i> , 2005, 1098, 166-171.	1.8	129
445	Analysis of Bioactive Triterpenes from <i>Rubus chingii</i> by Cyclodextrin-Modified Capillary Electrophoresis. <i>Chromatographia</i> , 2005, 62, 145-150.	0.7	20
446	Approach to method development and validation in capillary electrophoresis for enantiomeric purity testing of active basic pharmaceutical ingredients. <i>Electrophoresis</i> , 2005, 26, 2330-2341.	1.3	25
447	The formation of cucurbit[n]uril (n = 6, 7) complexes with amino compounds in aqueous formic acid studied by capillary electrophoresis. <i>Electrophoresis</i> , 2005, 26, 2214-2224.	1.3	26
448	Enantiomeric analysis of baclofen analogs by capillary zone electrophoresis, using highly sulfated cyclodextrins: Inclusion ionization constant pKa determination. <i>Electrophoresis</i> , 2005, 26, 2974-2983.	1.3	20
449	A family of single-isomer positively charged cyclodextrins as chiral selectors for capillary electrophoresis: Mono-6A-butylammonium-6A-deoxy- β -cyclodextrin tosylate. <i>Electrophoresis</i> , 2005, 26, 3125-3133.	1.3	35
450	Chiral separation in capillary electrophoresis using dual neutral cyclodextrins: Theoretical models of electrophoretic mobility difference and separation selectivity. <i>Electrophoresis</i> , 2005, 26, 3814-3823.	1.3	29
451	Enantioseparation of chiralN-imidazole derivatives by electrokinetic chromatography using highly sulfated cyclodextrins: Mechanism of enantioselective recognition. <i>Electrophoresis</i> , 2005, 26, 3824-3832.	1.3	15
452	Aptamer-oligonucleotide binding studied by capillary electrophoresis: Cation effect and separation efficiency. <i>Electrophoresis</i> , 2005, 26, 3247-3255.	1.3	25
453	Synthesis and application of single-isomer 6-mono(alkylimidazolium)- β -cyclodextrins as chiral selectors in chiral capillary electrophoresis. <i>Electrophoresis</i> , 2005, 26, 3839-3848.	1.3	55

#	ARTICLE	IF	CITATIONS
454	Enantioseparation in capillary electrophoresis using 2-O-(2-hydroxybutyl)- β -CD as a chiral selector. <i>Electrophoresis</i> , 2005, 26, 3890-3896.	1.3	14
455	Application of cyclodextrin-mediated capillary electrophoresis to determine the apparent binding constants and thermodynamic parameters of the alkylnaphthalene derivatives. <i>Electrophoresis</i> , 2005, 26, 3528-3537.	1.3	16
456	Influence of steric hindrance on enantioseparation of Dns-amino acids and pesticides on terguride based chiral selectors in capillary electrophoresis. <i>Journal of Separation Science</i> , 2005, 28, 673-677.	1.3	8
457	Enantioselective analysis of cetirizine in pharmaceuticals by cyclodextrin-mediated capillary electrophoresis. <i>Journal of Separation Science</i> , 2005, 28, 1278-1284.	1.3	28
458	18 Chiral separations. <i>Separation Science and Technology</i> , 2005, 6, 447-498.	0.0	9
459	Application of cyclodextrin-mediated capillary electrophoresis to simultaneously determine the apparent binding constants and thermodynamic parameters of naphthalenesulfonate derivatives. <i>Talanta</i> , 2006, 68, 1252-1258.	2.9	12
460	Principles of Enantiomer Separations in Electrokinetic Chromatography. , 0, , 179-206.		1
462	Chiral Capillary Electrophoresis of Agrochemicals in Real Samples. <i>Journal of Health Science</i> , 2006, 52, 489-494.	0.9	17
463	Chiral Capillary Electrophoresis of Amphetamine-Type Stimulants. <i>Journal of Health Science</i> , 2006, 52, 649-654.	0.9	11
464	Chiral Separation of Synthetic Vicinal Diol Compounds by Capillary Zone Electrophoresis with Borate Buffer and .BETA.-Cyclodextrin as Buffer Additive. <i>Analytical Sciences</i> , 2006, 22, 747-751.	0.8	5
465	Enantioselective separation in capillary electrophoresis using a novel mono-6A-propylammonium- β -cyclodextrin as chiral selector. <i>Analytica Chimica Acta</i> , 2006, 555, 63-67.	2.6	32
466	Determination of five anthraquinones in medicinal plants by capillary zone electrophoresis with β -cyclodextrin addition. <i>Journal of Chromatography A</i> , 2006, 1123, 134-137.	1.8	39
467	Chiral separation of four fluoroquinolone compounds using capillary electrophoresis with hydroxypropyl- β -cyclodextrin as chiral selector. <i>Journal of Chromatography A</i> , 2006, 1130, 296-301.	1.8	64
468	Enantioseparation of cis and trans nucleosides, aromatic analogues of stavudine, by capillary electrophoresis and high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2006, 1132, 141-147.	1.8	10
469	Enantiomeric separation of synthetic 2,3-dihydroxy-3-phenylpropionate compounds by β -cyclodextrin-modified capillary electrophoresis. <i>Journal of Chromatography A</i> , 2006, 1108, 258-262.	1.8	9
470	NMR studies of chiral recognition mechanisms: interaction of enantiomers of N-imidazole derivatives with cyclodextrin hosts. Correlation with the CD-EKC studies. <i>Tetrahedron: Asymmetry</i> , 2006, 17, 975-983.	1.8	12
471	Twenty Years of Separation of Cis-Trans (Z)-E Isomers. <i>Separation and Purification Reviews</i> , 2006, 35, 193-221.	2.8	10
472	Determination of Ephedrine, Pseudo-Ephedrine and Caffeine in a Dietary Product by Capillary Electrophoresis. <i>Chromatographia</i> , 2006, 63, 143-148.	0.7	40

#	ARTICLE	IF	CITATIONS
473	An observation of unusual temperature effects for enantioselective CZE employing highly sulfated- β -cyclodextrin. <i>Electrophoresis</i> , 2006, 27, 859-864.	1.3	9
474	Chiral separations by capillary electrophoresis: Recent developments and applications. <i>Electrophoresis</i> , 2006, 27, 2880-2895.	1.3	102
475	Enantiomeric separation of acidic compounds using single-isomer amino cyclodextrin derivatives in nonaqueous capillary electrophoresis. <i>Electrophoresis</i> , 2006, 27, 3434-3442.	1.3	31
476	Enantiomeric separation of some demethylated analogues of clofibrac acid by capillary zone electrophoresis and nano-liquid chromatography. <i>Electrophoresis</i> , 2006, 27, 1227-1236.	1.3	17
477	Optimization of capillary electrophoretic enantioseparation for basic drugs with native β -CD as a chiral selector. <i>Electrophoresis</i> , 2006, 27, 2367-2375.	1.3	16
478	Chiral capillary electrophoretic determination of the enantiomeric purity of homocamptothecin derivatives, promising antitumor topoisomerase I inhibitors, using highly sulfated CDs and fluorescence detection. <i>Electrophoresis</i> , 2006, 27, 4717-4729.	1.3	10
479	Effect of alkali metal hydroxides on the enantioseparation of amines using di-O-isopropylidene-keto-L-gulonic acid as the selector in NACE. <i>Electrophoresis</i> , 2006, 27, 4469-4479.	1.3	10
480	Chiral separations by capillary electrophoresis. , 2006, , 241-275.		7
481	Enantiomeric Separation of Synthetic Amino Acids Using Capillary Zone Electrophoresis. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 30, 1421-1436.	0.5	13
482	Separation of 2-HPA by Capillary Electrophoresis. , 2007, , .		0
483	Enantioseparation of Isoxanthohumol in Beer by Hydroxypropyl- β -cyclodextrin-Modified Micellar Electrokinetic Chromatography. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 6547-6552.	2.4	13
484	Enantiomeric Separation of a Series of β -Lactams Using Capillary Zone Electrophoresis. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 30, 1709-1721.	0.5	12
485	Determination of Ofloxacin Enantiomers in Pharmaceutical Formulations by Capillary Electrophoresis. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 31, 348-360.	0.5	20
486	Investigation of β -CD-derivatized erythromycin as chiral selector in CE. <i>Electrophoresis</i> , 2007, 28, 2566-2572.	1.3	14
487	Chiral resolution of melatonineric ligands by EKC using highly sulfated CDs. <i>Electrophoresis</i> , 2007, 28, 3915-3921.	1.3	5
488	Stereoisomeric separation of pharmaceutical compounds using CE with a chiral crown ether. <i>Electrophoresis</i> , 2007, 28, 2658-2666.	1.3	31
489	Review of aqueous chiral electrokinetic chromatography (EKC) with an emphasis on chiral microemulsion EKC. <i>Electrophoresis</i> , 2007, 28, 2503-2526.	1.3	39
490	Synthesis and NMR characterization of β -alanine-bridged hemispherodextrin, a very efficient chiral selector in EKC. <i>Electrophoresis</i> , 2007, 28, 2580-2588.	1.3	13

#	ARTICLE	IF	CITATIONS
491	Chiral separability of hydrophobic boron cluster anions with native cyclodextrins in water-methanol background electrolytes. <i>Electrophoresis</i> , 2007, 28, 3639-3649.	1.3	12
492	Enantioselective analysis of the antipsychotic 9-hydroxyrisperidone, main metabolite of risperidone, by chiral capillary EKC using dual CDs. <i>Electrophoresis</i> , 2007, 28, 2683-2692.	1.3	19
493	Temperature effects on the enantioselectivity of basic analytes in capillary EKC using sulfated β -CDs as chiral selectors. <i>Electrophoresis</i> , 2007, 28, 3753-3758.	1.3	14
494	Peak splitting in the CE separation of enantiomers caused by organic solvents in the sample. <i>Electrophoresis</i> , 2007, 28, 3625-3638.	1.3	4
495	Electromigration of a heteroconjugated imidazole-acetate complex in ACN. <i>Electrophoresis</i> , 2007, 28, 3590-3599.	1.3	10
496	Simultaneous enantioseparation and sensitivity enhancement of basic drugs using large-volume sample stacking. <i>Electrophoresis</i> , 2007, 28, 3542-3552.	1.3	24
497	Complexation of tauro- and glyco-conjugated bile salts with three neutral β -CDs studied by ACE. <i>Electrophoresis</i> , 2007, 28, 3745-3752.	1.3	28
498	Chiral separation of dansyl amino acids in capillary electrophoresis using mono-(3-methyl-imidazolium)- β -cyclodextrin chloride as selector. <i>Journal of Separation Science</i> , 2007, 30, 1343-1349.	1.3	48
499	Ketopinic acid and diisopropylidene ketogulonic acid as chiral ion-pair selectors in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2007, 1141, 287-294.	1.8	29
500	Influence of substituents on selectivity and efficiency of chiral separations of anions containing single nido-7,8-dicarbaundecaborane cluster with β -cyclodextrin. <i>Journal of Chromatography A</i> , 2007, 1143, 143-152.	1.8	8
501	Enantioseparations by using capillary electrophoretic techniques. <i>Journal of Chromatography A</i> , 2007, 1168, 45-70.	1.8	234
502	Effect of alkylimidazolium substituents on enantioseparation ability of single-isomer alkylimidazolium- β -cyclodextrin derivatives in capillary electrophoresis. <i>Analytica Chimica Acta</i> , 2007, 585, 227-233.	2.6	40
503	Kapillarelektrophorese, eine attraktive Methode zur Enantiomerenanalytik. <i>Pharmazie in Unserer Zeit</i> , 1999, 28, 186-196.	0.0	5
504	Characterization of the complexation of tauro- and glyco-conjugated bile salts with β -cyclodextrin and 2-hydroxypropyl- β -cyclodextrin using affinity capillary electrophoresis. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2008, 61, 161-169.	1.6	20
505	Cyclodextrins in capillary electrophoresis enantioseparations – Recent developments and applications. <i>Journal of Separation Science</i> , 2008, 31, 1991-2011.	1.3	146
506	Simultaneous enantioseparation of four β -agonists by capillary electrophoresis with cyclodextrin additives. Study of the enantioselective mechanism. <i>Journal of Separation Science</i> , 2008, 31, 3749-3754.	1.3	16
507	Monosubstituted positively charged cyclodextrins: Synthesis and applications in chiral separation. <i>Journal of Separation Science</i> , 2008, 31, 3246-3256.	1.3	57
508	Synthesis of chiral vicinal diols and analysis of them by capillary zone electrophoresis. <i>Chirality</i> , 2008, 20, 75-83.	1.3	12

#	ARTICLE	IF	CITATIONS
509	Recent advances in amino acid analysis by CE. <i>Electrophoresis</i> , 2008, 29, 207-223.	1.3	43
510	The role of cyclodextrins in chiral capillary electrophoresis. <i>Electrophoresis</i> , 2008, 29, 1701-1712.	1.3	116
511	Study of interaction between drug enantiomers and human serum albumin by flow injection capillary electrophoresis frontal analysis. <i>Electrophoresis</i> , 2008, 29, 2876-2883.	1.3	21
512	Enantiomeric separation of some flavanones using shinorhizobial linear octasaccharides in CE. <i>Electrophoresis</i> , 2008, 29, 4284-4290.	1.3	20
513	Enantiomeric separation of glycidyl tosylate by CE: Application to the study of catalytic asymmetric epoxidation of allyl alcohol. <i>Electrophoresis</i> , 2008, 29, 4575-4582.	1.3	5
515	Model of CE enantioseparation systems with a mixture of chiral selectors. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2008, 875, 30-34.	1.2	46
516	Stereoselective determination of trihexyphenidyl using carboxymethyl- β -cyclodextrin by capillary electrophoresis with field-amplified sample stacking. <i>Microchemical Journal</i> , 2008, 89, 34-41.	2.3	20
517	Possibilities of column coupling electrophoresis provided with a fiber-based diode array detection in enantioselective analysis of drugs in pharmaceutical and clinical samples. <i>Journal of Chromatography A</i> , 2008, 1179, 9-16.	1.8	21
518	Study of the complexation of risperidone and 9-hydroxyrisperidone with cyclodextrin hosts using affinity capillary electrophoresis and ^1H NMR spectroscopy. <i>Journal of Chromatography A</i> , 2008, 1215, 185-193.	1.8	38
519	18 Coupling CE and microchip-based devices with mass spectrometry. <i>Separation Science and Technology</i> , 2008, 9, 477-521.	0.0	9
520	Sensitivity improvement by using contactless conductivity rather than indirect UV detection for the determination of enantiomeric purity of amines by CE. <i>Electrophoresis</i> , 2009, 30, 487-498.	1.3	26
521	Enantioseparation of nicotine alkaloids in cigarettes by CE using sulfated β -CD as a chiral selector and a capillary coated with amino groups. <i>Electrophoresis</i> , 2009, 30, 349-356.	1.3	26
522	Chiral separations by CE employing CDs. <i>Electrophoresis</i> , 2009, 30, S203-10.	1.3	118
523	Affinity in Electrophoresis. <i>Electrophoresis</i> , 2009, 30, S229-39.	1.3	57
524	About the role of enantioselective selector-selectand interactions and the mobilities of diastereomeric associates in enantiomer separations using CE. <i>Electrophoresis</i> , 2009, 30, 2803-2811.	1.3	66
525	Use of coated capillaries for the electrophoretic separation of stereoisomers of a growth hormone secretagogue. <i>Electrophoresis</i> , 2009, 30, 3772-3779.	1.3	41
526	Separation of enantiomers with charged chiral selectors in CE. <i>Electrophoresis</i> , 2009, 30, S211-21.	1.3	116
527	Advanced CE for chiral analysis of drugs, metabolites, and biomarkers in biological samples. <i>Electrophoresis</i> , 2009, 30, 2773-2802.	1.3	80

#	ARTICLE	IF	CITATIONS
528	Enantioselective separation of chiral vicinal diols in capillary electrophoresis using a mono- α -aminoethylamino- β -cyclodextrin as a chiral selector. <i>Journal of Separation Science</i> , 2009, 32, 125-134.	1.3	11
529	Enantioseparation of chiral benzimidazole derivatives by electrokinetic chromatography using sulfated cyclodextrins. <i>Journal of Separation Science</i> , 2009, 32, 1907-1915.	1.3	11
530	Use of correction factors in mobility shift affinity capillary electrophoresis for weak analyte-ligand interactions. <i>Journal of Separation Science</i> , 2009, 32, 1712-1721.	1.3	27
531	Enantioselective analysis of primaquine and its impurity quinocide by capillary electrophoresis. <i>Biomedical Chromatography</i> , 2009, 23, 295-301.	0.8	10
532	6-O-(2-hydroxybutyl)- β -CD as a chiral selector for nonaqueous capillary electrophoretic separation of chiral drugs. <i>Frontiers of Chemistry in China: Selected Publications From Chinese Universities</i> , 2009, 4, 396-401.	0.4	3
533	A study of the interaction between enantiomers of zolmitriptan and hydroxypropyl-beta-cyclodextrin by capillary electrophoresis. <i>Analytical and Bioanalytical Chemistry</i> , 2009, 393, 313-320.	1.9	18
534	Assessment of the complexation degree of camptothecin derivatives and cyclodextrins using spectroscopic and separative methodologies. <i>Tetrahedron: Asymmetry</i> , 2009, 20, 2482-2489.	1.8	13
535	The 3-amino-derivative of β -cyclodextrin as chiral selector of Dns-amino acids in electrokinetic chromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 3678-3686.	1.8	17
536	Enantiomeric Resolution of Benzoxazolinonic Aminoalcohols, Potential Adrenergic Ligands, by LC and CE. <i>Chromatographia</i> , 2009, 69, 465-472.	0.7	4
537	Enantioselective Separation of Basic Drugs by CE with Polygalacturonic Acid as a Novel Chiral Selector. <i>Chromatographia</i> , 2009, 69, 1315-1320.	0.7	18
538	Enantioseparation of Dioxopromethazine Hydrochloride in Urine with Liquid-Liquid Extraction by CE-ECL Detection. <i>Chromatographia</i> , 2009, 70, 1291-1293.	0.7	12
539	Effect of β -Cyclodextrin on Drug Distribution Studied by Electrochemistry at Interfaces between Immiscible Electrolyte Solutions. <i>Journal of Physical Chemistry B</i> , 2009, 113, 7263-7269.	1.2	12
540	Enantioseparation of binaphthol and its monoderivatives by cyclodextrin-modified capillary zone electrophoresis: A mathematical approach. <i>Talanta</i> , 2009, 78, 631-637.	2.9	24
541	Simultaneous determination of R-(+)-, S-(-)-baclofen and impurity A by electrokinetic chromatography. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2010, 51, 541-548.	1.4	19
542	Separation of vinca alkaloid enantiomers by capillary electrophoresis applying cyclodextrin derivatives and characterization of cyclodextrin complexes by nuclear magnetic resonance spectroscopy. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2010, 53, 1258-1266.	1.4	29
543	Enantiomeric separation of epinephrine and salbutamol by micellar electrokinetic chromatography using β -cyclodextrin as chiral additive. <i>Chinese Journal of Chemistry</i> , 2010, 22, 845-848.	2.6	0
544	Glycogen: A novel branched polysaccharide chiral selector in CE. <i>Electrophoresis</i> , 2010, 31, 1044-1050.	1.3	34
545	Enhanced selectivity in CZE multi-chiral selector enantioseparation systems: Proposed separation mechanism. <i>Electrophoresis</i> , 2010, 31, 1435-1441.	1.3	54

#	ARTICLE	IF	CITATIONS
546	Determination of the binding constants of modafinil enantiomers with sulfated β -cyclodextrin chiral selector by capillary electrophoresis using three different linear plotting methods. <i>Electrophoresis</i> , 2010, 31, 2957-2963.	1.3	17
548	Synthesis and application of a novel single isomer mono- β -deoxy- β -(2 <i>S</i> ,3 <i>S</i>)- α -(2,3- <i>O</i> -isopropylidene-1,4-tetramethylenediamine)- β -cyclodextrin as chiral selector in capillary electrophoresis. <i>Chirality</i> , 2010, 22, 914-921.		
549	Capillary electrophoretic and nuclear magnetic resonance studies on the opposite affinity pattern of propranolol enantiomers towards various cyclodextrins. <i>Journal of Separation Science</i> , 2010, 33, 1617-1624.	1.3	52
550	Separation and characterization of modified pregabalin in terms of cyclodextrin complexation, using capillary electrophoresis and nuclear magnetic resonance. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2010, 51, 842-852.	1.4	51
551	Study of a new chiral selector: Sodium arsenyl-(l)-(+)-tartrate for capillary electrophoresis. <i>Journal of Chromatography A</i> , 2010, 1217, 1139-1148.	1.8	10
552	Application of charged single isomer derivatives of cyclodextrins in capillary electrophoresis for chiral analysis. <i>Journal of Chromatography A</i> , 2010, 1217, 953-967.	1.8	88
553	Chiral Recognition and Enantioseparation Mechanisms in Capillary Electrokinetic Chromatography. , 2010, , 97-152.		1
555	Chiral Separations. <i>Annual Review of Analytical Chemistry</i> , 2010, 3, 341-363.	2.8	109
556	Chiral Recognition in Separation Methods. , 2010, , .		81
557	Microchip Electrophoresis Using Linear-Imaging UV Detector. <i>Bunseki Kagaku</i> , 2011, 60, 725-734.	0.1	0
558	Complexation of triptolide and its succinate derivative with cyclodextrins: Affinity capillary electrophoresis, isothermal titration calorimetry and ¹ H NMR studies. <i>Journal of Chromatography A</i> , 2011, 1218, 8708-8714.	1.8	9
559	Enantioseparation of the New Antifungal Drug Iodiconazole and Structurally Related Triadimenol Analogues by CE with Neutral Cyclodextrin Additives. <i>Chromatographia</i> , 2011, 73, 1009-1014.	0.7	13
560	Chiral electromigration techniques in pharmaceutical and biomedical analysis. <i>Bioanalytical Reviews</i> , 2011, 3, 95-114.	0.1	11
561	Complexation of tauro- and glyco- conjugated bile salts with β -cyclodextrin and hydroxypropyl- β -cyclodextrin studied by affinity capillary electrophoresis and molecular modelling. <i>Journal of Separation Science</i> , 2011, 34, 3221-3230.	1.3	17
562	Accuracy and sensitivity of the determination of rate constants of interconversion in achiral and chiral environments by dynamic enantioselective electrophoresis. <i>Electrophoresis</i> , 2011, 32, 595-603.	1.3	6
563	Chiral separation and quantitation of cetirizine and hydroxyzine by maltodextrin-mediated CE in human plasma: Effect of zwitterionic property of cetirizine on enantioseparation. <i>Electrophoresis</i> , 2011, 32, 764-771.	1.3	31
564	Evaluation of the interaction between sitagliptin and cyclodextrin derivatives by capillary electrophoresis and nuclear magnetic resonance spectroscopy. <i>Electrophoresis</i> , 2011, 32, 2648-2654.	1.3	23
565	Emerging subject for chiral separation science: Cluster boron compounds. <i>Chirality</i> , 2011, 23, 307-319.	1.3	25

#	ARTICLE	IF	CITATIONS
566	Fundamental aspects of chiral electromigration techniques and application in pharmaceutical and biomedical analysis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2011, 55, 688-701.	1.4	90
567	Molecular Modeling Study of Chiral Separation and Recognition Mechanism of β^2 -Adrenergic Antagonists by Capillary Electrophoresis. <i>International Journal of Molecular Sciences</i> , 2012, 13, 710-725.	1.8	51
568	8.10 Chromatographic Separations and Analysis: Cyclodextrin Mediated HPLC, GC and CE Enantiomeric Separations. , 2012, , 177-199.		12
569	Comparison of separation performances of novel β^2 -cyclodextrin-based chiral stationary phases in high-performance liquid chromatographic enantioseparation. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2012, 70, 71-76.	1.4	15
570	Chiral separation of amino acids derivatised with fluorescein isothiocyanate by single isomer derivatives 3-monodeoxy-3-monoamino- β^2 - and β^3 -cyclodextrins: the effect of the cavity size. <i>Journal of Chromatography A</i> , 2012, 1269, 360-365.	1.8	23
571	Monte Carlo simulation of the enantioseparation process. <i>Physica A: Statistical Mechanics and Its Applications</i> , 2012, 391, 4389-4396.	1.2	5
572	Direct enantioseparation of lipoic acid in dietary supplements by capillary electrophoresis using trimethyl- β^2 -cyclodextrin as a chiral selector. <i>Electrophoresis</i> , 2012, 33, 2441-2445.	1.3	26
573	Simulation of the effects of complex-formation equilibria in electrophoresis: III. Simultaneous effects of chiral selector concentration and background electrolyte pH. <i>Electrophoresis</i> , 2012, 33, 3012-3020.	1.3	22
574	Affinity capillary electrophoresis method for investigation of bile salts complexation with sulfobutyl ether- β^2 -cyclodextrin. <i>Journal of Separation Science</i> , 2012, 35, 2764-2772.	1.3	11
575	Chiral electrokinetic chromatography-electrospray ionization-mass spectrometry using a double junction interface. <i>Electrophoresis</i> , 2012, 33, 546-551.	1.3	9
576	Dynamic high-resolution computer simulation of electrophoretic enantiomer separations with neutral cyclodextrins as chiral selectors. <i>Electrophoresis</i> , 2012, 33, 958-969.	1.3	31
577	Determination of stability constants of complexes of neutral analytes with charged cyclodextrins by affinity capillary electrophoresis. <i>Electrophoresis</i> , 2012, 33, 1032-1039.	1.3	34
578	Simulation of the effects of complex-formation equilibria in electrophoresis: II. Experimental verification. <i>Electrophoresis</i> , 2012, 33, 948-957.	1.3	43
579	Chiral recognition of imperanene enantiomers by various cyclodextrins: A capillary electrophoresis and ¹ H-NMR spectroscopy study. <i>Electrophoresis</i> , 2012, 33, 1458-1464.	1.3	30
580	Extraction of Phenylalanine Enantiomers by Aqueous Two Phase Systems Containing Combinatorial Chiral Selector. <i>Chinese Journal of Chemistry</i> , 2012, 30, 965-969.	2.6	13
581	Effect of electrolyzed reduced water on chiral theanine and polyphenols in tea. <i>Food Chemistry</i> , 2012, 134, 1761-1766.	4.2	8
582	Enantiomeric separation of <i>R,S</i> -tolterodine and <i>R,S</i> -methoxytolterodine with negatively charged cyclodextrins by capillary electrophoresis. <i>Journal of Separation Science</i> , 2013, 36, 1561-1567.	1.3	18
583	Simultaneous determination of flavonoid analogs in <i>Scutellariae Barbatae Herba</i> by β^2 -cyclodextrin and acetonitrile modified capillary zone electrophoresis. <i>Talanta</i> , 2013, 105, 393-402.	2.9	50

#	ARTICLE	IF	CITATIONS
584	Modified Cyclodextrins for Chiral Separation. , 2013, , .		22
585	Complexation of Buffer Constituents with Neutral Complexation Agents: Part II. Practical Impact in Capillary Zone Electrophoresis. Analytical Chemistry, 2013, 85, 8526-8534.	3.2	30
586	Differentiation of Enantiomers I. Topics in Current Chemistry, 2013, , .	4.0	15
587	Applicability and limitations of affinity capillary electrophoresis and vacancy affinity capillary electrophoresis methods for determination of complexation constants. Electrophoresis, 2013, 34, 761-767.	1.3	54
588	Chiral selectors in <sc>CE</sc>: Recent developments and applications. Electrophoresis, 2013, 34, 178-204.	1.3	98
589	Separation selectivity patterns of fully charged achiral compounds in capillary electrophoresis with a neutral cyclodextrin. Journal of Separation Science, 2013, 36, 325-333.	1.3	0
590	Recent advances in electrodriven enantioseparations. Journal of Separation Science, 2013, 36, 52-74.	1.3	109
591	Synthesis and application of a chiral ionic liquid functionalized β -cyclodextrin as a chiral selector in capillary electrophoresis. Biomedical Chromatography, 2013, 27, 1027-1033.	0.8	27
592	Evaluation of the enantioselectivity of carbon nanoparticles β -modified chiral separation systems using dextrin as chiral selector by capillary electrokinetic chromatography. Electrophoresis, 2013, 34, 1901-1907.	1.3	26
593	Differentiation of Enantiomers by Capillary Electrophoresis. Topics in Current Chemistry, 2013, 340, 209-275.	4.0	24
594	Native β -Cyclodextrins and Their Neutral Derivatives for Capillary Electrophoresis. , 2013, , 145-186.		0
595	Cationic Cyclodextrins for Capillary Electrophoresis. , 2013, , 225-258.		0
596	Application of Polymeric Surfactants in Chiral Micellar Electrokinetic Chromatography (CMEKC) and CMEKC Coupled to Mass Spectrometry. Methods in Molecular Biology, 2013, 970, 319-348.	0.4	8
597	Electrically-enhanced microextraction combined with maltodextrin-modified capillary electrophoresis for quantification of tolterodine enantiomers in biological samples. Microchemical Journal, 2013, 106, 186-193.	2.3	46
598	Evaluation of electroosmotic markers in aqueous and nonaqueous capillary electrophoresis. Electrophoresis, 2013, 34, 3252-3259.	1.3	20
599	Enantioseparation of Tr β -Cyclodextrin's Base Derivatives by Capillary Electrophoresis Using Cyclodextrins as Chiral Selectors. Chirality, 2013, 25, 379-383.	1.3	6
600	New single isomer negatively charged β -cyclodextrin derivatives as chiral selectors in <sc>capillary electrophoresis</sc>. Electrophoresis, 2013, 34, 1232-1240.	1.3	14
601	Physicochemical profiling of drug candidates using capillary-based techniques. Journal of Drug Delivery Science and Technology, 2013, 23, 333-345.	1.4	8

#	ARTICLE	IF	CITATIONS
602	Determination of effective mobilities of EOF markers in BGE containing sulfated cyclodextrin by a two-detector method. <i>Electrophoresis</i> , 2013, 34, 768-776.	1.3	19
603	Recent Trends in Chiral Separations by Nano Liquid Chromatography and Nano Capillary Electrophoresis. <i>Current Chromatography</i> , 2014, 1, 81-89.	0.1	19
604	Evaluation of the chiral separation ability of single isomer cationic cyclodextrins in capillary electrophoresis. <i>Electrophoresis</i> , 2014, 35, 2744-2751.	1.3	17
605	Hydrophilic interaction electrokinetic chromatography using bio-based nanofillers. <i>Electrophoresis</i> , 2014, 35, 2229-2236.	1.3	3
606	Characterizing the interaction between enantiomers of eight psychoactive drugs and highly sulfated cyclodextrin by counter-current capillary electrophoresis. <i>Biomedical Chromatography</i> , 2014, 28, 120-126.	0.8	11
607	Velocity gap mode of capillary electrophoresis developed for high-resolution chiral separations. <i>Electrophoresis</i> , 2014, 35, 2778-2784.	1.3	7
608	Cyclodextrin-mediated enantioseparation of phenylalanine amide derivatives and amino alcohols by capillary electrophoresis—Role of complexation constants and complex mobilities. <i>Electrophoresis</i> , 2014, 35, 2848-2854.	1.3	16
609	Separation and Migration Behavior of Dichlorophenols in Cyclodextrin-Modified Capillary Zone Electrophoresis. <i>Journal of the Chinese Chemical Society</i> , 2014, 61, 1025-1030.	0.8	0
610	Separation efficiency of dual-selector systems in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2014, 1330, 82-88.	1.8	21
611	Effect of competing binding modes on retention in chromatography and capillary electrophoresis. A theoretical consideration. <i>Journal of Separation Science</i> , 2014, 37, 390-392.	1.3	3
612	Application of cyclodextrins in chiral capillary electrophoresis. <i>Electrophoresis</i> , 2014, 35, 2701-2721.	1.3	141
613	Antagonists of the P _{2X7} receptor: Mechanism of enantioselective recognition using highly sulfated and sulfobutylether cyclodextrins by capillary electrokinetic chromatography. <i>Electrophoresis</i> , 2014, 35, 2892-2899.	1.3	9
614	Enantioseparation of meptazinol and its three intermediate enantiomers by capillary electrophoresis using a new cationic cyclodextrin derivative in single and dual cyclodextrin systems. <i>Biomedical Chromatography</i> , 2014, 28, 868-874.	0.8	17
615	Penicillin G as a novel chiral selector in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2014, 1326, 134-138.	1.8	16
616	Dynamic computer simulation of electrophoretic enantiomer migration order and separation in presence of a neutral cyclodextrin. <i>Electrophoresis</i> , 2014, 35, 2833-2841.	1.3	22
617	Determination of thermodynamic values of acidic dissociation constants and complexation constants of profens and their utilization for optimization of separation conditions by Simul 5 Complex. <i>Journal of Chromatography A</i> , 2014, 1364, 276-288.	1.8	27
618	Twenty years of development of dual and multi-selector models in capillary electrophoresis: A review. <i>Electrophoresis</i> , 2014, 35, 2688-2700.	1.3	40
619	Maltodextrins as Chiral Selectors in CE: Molecular Structure Effect of Basic Chiral Compounds on the Enantioseparation. <i>Chirality</i> , 2014, 26, 620-628.	1.3	12

#	ARTICLE	IF	CITATIONS
620	Visceral mesh modified with cyclodextrin for the local sustained delivery of ropivacaine. International Journal of Pharmaceutics, 2014, 476, 149-159.	2.6	25
621	Advances in chiral separations of small peptides by capillary electrophoresis and chromatography. Journal of Separation Science, 2014, 37, 2447-2466.	1.3	121
622	Novel methoxypropylimidazolium β -cyclodextrin for improved enantioseparation of amino acids. Talanta, 2014, 128, 460-465.	2.9	17
624	Separation of Folinic Acid Diastereomers in Capillary Electrophoresis Using a New Cationic β -Cyclodextrin Derivative. PLoS ONE, 2015, 10, e0120216.	1.1	7
625	Chiral Resolution Capabilities of DNA Oligonucleotides. Analytical Chemistry, 2015, 87, 5491-5495.	3.2	42
626	Synthesis of 2,3-O-dibenzyl-6-O-sulfobutyl- α and β cyclodextrins: new chiral surfactants for capillary electrophoresis. Tetrahedron Letters, 2015, 56, 4451-4454.	0.7	13
627	Generalized model of electromigration with 1:1 (analyte:selector) complexation stoichiometry: Part II. Application to dual systems and experimental verification. Journal of Chromatography A, 2015, 1384, 147-154.	1.8	12
628	Study of the enantioseparation capability of chiral dual system based on chondroitin sulfate C in CE. Electrophoresis, 2015, 36, 607-614.	1.3	9
629	Methoxypropylamino β -cyclodextrin clicked AC regioisomer for enantioseparations in capillary electrophoresis. Analytica Chimica Acta, 2015, 868, 73-79.	2.6	15
630	Development of a capillary electrophoretic method for determination of plasma clearance of iohexol in dogs and cats. Biomedical Chromatography, 2015, 29, 504-513.	0.8	7
631	Chiral Separations in Capillary Electrophoresis. , 2015, , .		2
632	Development of Electrophoretic Methods for Simultaneous Determination of Enantiomeric Ratio and Composition of Diastereomeric Salt Mixtures. Chromatographia, 2015, 78, 881-888.	0.7	6
633	A dispersive liquid-liquid microextraction and chiral separation of carvedilol in human plasma using capillary electrophoresis. Bioanalysis, 2015, 7, 1107-1117.	0.6	27
634	Chiral selectors in CE: Recent developments and applications (2012-2014). Electrophoresis, 2015, 36, 101-123.	1.3	67
635	Evaluation of sulfated maltodextrin as a novel anionic chiral selector for the enantioseparation of basic chiral drugs by capillary electrophoresis. Electrophoresis, 2015, 36, 305-311.	1.3	27
636	Influence of substituent position and cavity size of the regioisomers of monocarboxymethyl- α , β , and γ -cyclodextrins on the apparent stability constants of their complexes with both enantiomers of Tröger's base. Journal of Separation Science, 2016, 39, 980-985.	1.3	15
637	The empirical comparison of cyclofructans and cyclodextrins as chiral selectors in capillary electrophoretic separation of atropisomers of 1,1'-binaphthalene-2,2'-diyl hydrogens phosphate. Journal of Separation Science, 2016, 39, 973-979.		10
638	Development and validation of a cyclodextrin-modified capillary electrophoresis method for the enantiomeric separation of vildagliptin enantiomers. Electrophoresis, 2016, 37, 1318-1325.	1.3	23

#	ARTICLE	IF	CITATIONS
639	Cyclodextrin-assisted enantioseparation of warfarin and 10-hydroxywarfarin by capillary electrophoresis studied from the analytical and thermodynamic points of view. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016, 126, 60-65.	1.4	8
640	Advances in the Use of Cyclodextrins as Chiral Selectors in Capillary Electrokinetic Chromatography: Fundamentals and Applications. <i>Chromatographia</i> , 2016, 79, 1403-1435.	0.7	101
641	Single-isomer carboxymethyl- β -cyclodextrin as chiral resolving agent for capillary electrophoresis. <i>Journal of Chromatography A</i> , 2016, 1467, 445-453.	1.8	34
642	Affinity capillary electrophoresis: the theory of electromigration. <i>Analytical and Bioanalytical Chemistry</i> , 2016, 408, 8623-8641.	1.9	60
643	Three chiral ionic liquids as additives for enantioseparation in capillary electrophoresis and their comparison with conventional modifiers. <i>Journal of Chromatography A</i> , 2016, 1462, 146-152.	1.8	43
644	Enantiomer Separations by Capillary Electrophoresis. <i>Methods in Molecular Biology</i> , 2016, 1483, 277-299.	0.4	5
645	Chiral Separation of Uncharged Pomalidomide Enantiomers Using Carboxymethyl- β -Cyclodextrin: A Validated Capillary Electrophoretic Method. <i>Chirality</i> , 2016, 28, 199-203.	1.3	19
646	Estimation of apparent binding constant of complexes of selected acyclic nucleoside phosphonates with β -cyclodextrin by affinity capillary electrophoresis. <i>Electrophoresis</i> , 2016, 37, 239-247.	1.3	33
647	Physicochemical characterisation and cyclodextrin complexation of erlotinib. <i>Supramolecular Chemistry</i> , 2016, 28, 656-664.	1.5	14
648	Chiral separation of asenapine enantiomers by capillary electrophoresis and characterization of cyclodextrin complexes by NMR spectroscopy, mass spectrometry and molecular modeling. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016, 117, 398-404.	1.4	47
649	Characterization of complexes between phenethylamine enantiomers and β -cyclodextrin derivatives by capillary electrophoresis—Determination of binding constants and complex mobilities. <i>Electrophoresis</i> , 2017, 38, 1188-1200.	1.3	12
650	Validated capillary electrophoretic method for the enantiomeric quality control of <i>R</i> -praziquantel. <i>Electrophoresis</i> , 2017, 38, 1886-1894.	1.3	14
651	Use of various β -cyclodextrin derivatives as chiral selectors for the enantiomeric separation of ofloxacin and its five related substances by capillary electrophoresis. <i>Journal of Separation Science</i> , 2017, 40, 1784-1795.	1.3	21
652	Sequence requirements of oligonucleotide chiral selectors for the capillary electrophoresis resolution of low-affinity DNA binders. <i>Electrophoresis</i> , 2017, 38, 1383-1390.	1.3	14
653	Single and dual cyclodextrins systems for the enantiomeric and diastereoisomeric separations of structurally related dihydropyridone analogues. <i>Electrophoresis</i> , 2017, 38, 1922-1931.	1.3	10
654	Use of various β -cyclodextrin derivatives as chiral selectors for the enantiomeric separation of higenamine by capillary electrophoresis. <i>Microchemical Journal</i> , 2017, 134, 289-294.	2.3	13
655	Cyclodextrin complexation improves aqueous solubility of the antiepileptic drug, rufinamide: solution and solid state characterization of compound-cyclodextrin binary systems. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2017, 88, 43-52.	0.9	8
656	Chiral separation of terbutaline and non-steroidal anti-inflammatory drugs by using a new lysine-bridged hemispherodextrin in capillary electrophoresis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2017, 145, 734-741.	1.4	14

#	ARTICLE	IF	CITATIONS
657	Chiral surfaces: The many faces of chiral recognition. <i>Current Opinion in Colloid and Interface Science</i> , 2017, 32, 94-107.	3.4	31
658	Thermodynamic models to elucidate the enantioseparation of drugs with two stereogenic centers by micellar electrokinetic chromatography. <i>Journal of Chromatography A</i> , 2017, 1512, 133-142.	1.8	11
659	An analytical model for enantioseparation process in capillary electrophoresis. <i>Physica A: Statistical Mechanics and Its Applications</i> , 2017, 487, 153-163.	1.2	3
660	Enhancement of the conductivity detection signal in capillary electrophoresis systems using neutral cyclodextrins as sweeping agents. <i>Electrophoresis</i> , 2018, 39, 1390-1398.	1.3	9
661	Study of isomeric pentacyclic triterpene acids in traditional Chinese medicine of <i>Forsythiae Fructus</i> and their binding constants with β -cyclodextrin by capillary electrophoresis. <i>Electrophoresis</i> , 2018, 39, 1006-1013.	1.3	14
662	Into the theory of the partial filling affinity capillary electrophoresis and the determination of apparent stability constants of analyte-glyand complexes. <i>Electrophoresis</i> , 2018, 39, 742-751.	1.3	22
663	Equilibrium, structural and antibacterial characterization of moxifloxacin- β -cyclodextrin complex. <i>Journal of Molecular Structure</i> , 2018, 1166, 228-236.	1.8	30
664	Computer simulation and enantioselective capillary electrophoresis to characterize isomer mixtures of sulfated β -cyclodextrins. <i>Electrophoresis</i> , 2018, 39, 770-778.	1.3	12
665	Chiral separation of lenalidomide by liquid chromatography on polysaccharide-type stationary phases and by capillary electrophoresis using cyclodextrin selectors. <i>Journal of Separation Science</i> , 2018, 41, 1414-1423.	1.3	37
666	Molecular Modeling and Chiral Separation of Benzodiazepines by Capillary Electrophoresis Using Highly Sulfated Cyclodextrins. <i>Journal of the Mexican Chemical Society</i> , 2018, 62, .	0.2	4
667	Enantioseparations by Capillary Electromigration Techniques. , 2018, , 565-605.		2
668	Generalized model of the linear theory of electromigration and its application to electrokinetic chromatography: Theory and software PeakMaster 6 Next Generation. <i>Electrophoresis</i> , 2019, 40, 683-692.	1.3	24
669	Cyclodextrin-induced acidity modification of substituted cathinones studied by capillary electrophoresis supported by density functional theory calculations. <i>Journal of Chromatography A</i> , 2018, 1580, 142-151.	1.8	9
670	Enhanced enantioselectivity of native β -cyclodextrins by the synergy of chiral ionic liquids in capillary electrophoresis. <i>Journal of Separation Science</i> , 2018, 41, 4525-4532.	1.3	34
671	Contemporary theory of enantioseparations in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2018, 1567, 2-25.	1.8	92
672	Fundamental aspects of field-amplified electrokinetic injection of cations for enantioselective capillary electrophoresis with sulfated cyclodextrins as selectors. <i>Journal of Chromatography A</i> , 2018, 1558, 85-95.	1.8	17
673	Preclinical evidence of enhanced analgesic activity of duloxetine complexed with succinyl- β -cyclodextrin: A comparative study with cyclodextrin complexes. <i>International Journal of Pharmaceutics</i> , 2019, 566, 391-399.	2.6	3
674	Some thoughts about enantioseparations in capillary electrophoresis. <i>Electrophoresis</i> , 2019, 40, 2420-2437.	1.3	75

#	ARTICLE	IF	CITATIONS
675	Separations of antifungal compounds in capillary electrophoresis with two anionic cyclodextrins. <i>Electrophoresis</i> , 2019, 40, 1986-1991.	1.3	2
676	Applicability assessment of the CASCade Two Dimensional SEDiment (CASC2Dâ€¦SED) distributed hydrological model for flood forecasting across four typical medium and small watersheds in China. <i>Journal of Flood Risk Management</i> , 2019, 12, .	1.6	32
677	β -Cyclodextrin-copper (II) complex as chiral selector in capillary electrophoresis for the enantioseparation of β -blockers. <i>Journal of Chromatography A</i> , 2019, 1596, 233-240.	1.8	14
678	Synergistic enantioseparation systems with either cyclodextrins or cyclofructans and Lâ€¦alanine <i><i>tert</i></i> butyl ester lactate. <i>Electrophoresis</i> , 2019, 40, 539-546.	1.3	15
679	Determination of thermodynamic binding constants by affinity capillary electrophoresis. <i>Talanta</i> , 2019, 192, 448-454.	2.9	9
680	Clinical and pharmaceutical applications of affinity ligands in capillary electrophoresis: A review. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2020, 177, 112882.	1.4	40
681	Modeling-based optimization of the simultaneous enantiomeric separation of multicomponent mixtures of phenoxy acid herbicides using dual cyclodextrin systems by Capillary Electrophoresis. <i>Journal of Chromatography A</i> , 2020, 1610, 460552.	1.8	13
682	Investigation of the synergistic effect of chiral ionic liquids as additives in non-aqueous capillary electrophoresis for enantioseparation. <i>Journal of Chromatography A</i> , 2020, 1609, 460519.	1.8	36
683	Chiral covalent organic frameworks: design, synthesis and property. <i>Chemical Society Reviews</i> , 2020, 49, 6248-6272.	18.7	211
684	Binding constants determination of cyclodextrin inclusion complexes by affinity capillary electrophoresis. How to overcome the limitations induced by the UV-detector?. <i>Journal of Chromatography A</i> , 2020, 1623, 461209.	1.8	7
685	Use of Gamithromycin as a Chiral Selector in Capillary Electrophoresis. <i>Journal of Chromatography A</i> , 2020, 1624, 461099.	1.8	17
686	Investigation of the synergistic effect with chiral Dâ€¦penicillamine functionalized gold nanoparticle as an additive for enantiomeric separation in capillary electrophoresis. <i>Electrophoresis</i> , 2020, 41, 1060-1066.	1.3	8
687	Single isomer cyclodextrins as chiral selectors in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2020, 1627, 461375.	1.8	47
688	Investigation of fusidic acid as a chiral selector in capillary electrophoresis. <i>Separation and Purification Technology</i> , 2020, 242, 116768.	3.9	17
689	Advances of capillary electrophoresis enantioseparations in pharmaceutical analysis (2017â€¦2020). <i>Electrophoresis</i> , 2021, 42, 1709-1725.	1.3	51
690	Binding Constants Determination of Ofloxacin and Ornidazole Enantiomers with Sulfated β -Cyclodextrin as Single Ligand by Capillary Electrophoresis using Three Different Linear Plotting Methods. <i>Asian Journal of Chemistry</i> , 2021, 33, 1663-1670.	0.1	0
691	Bioanalysis of drugs and their metabolites by chiral electromigration techniques (2010â€¦2020). <i>Electrophoresis</i> , 2021, 42, 1744-1760.	1.3	9
692	Role of diamionaphthalene based polymers as sensors in detection of biomolecules: A review. <i>Results in Materials</i> , 2021, 9, 100174.	0.9	5

#	ARTICLE	IF	CITATIONS
693	Electrophoretic separation of multiple chiral center analyte with a three cyclodextrins mixture. <i>Electrophoresis</i> , 2021, 42, 1810-1817.	1.3	3
694	The Use of Dual Cyclodextrin Chiral Selector Systems in the Enantioseparation of Pharmaceuticals by Capillary Electrophoresis: An Overview. <i>Molecules</i> , 2021, 26, 2261.	1.7	19
695	Study on a novel buffer system to separate and quantify glycosylated hemoglobin A1c (HbA1c) and the mechanism between its key constituents and HbA1c. <i>Microchemical Journal</i> , 2021, 163, 105916.	2.3	1
696	Native and substituted cyclodextrins as chiral selectors for capillary electrophoresis enantioseparations: Structures, features, application, and molecular modeling. <i>Electrophoresis</i> , 2021, 42, 1676-1708.	1.3	52
697	Single Isomer N-Heterocyclic Cyclodextrin Derivatives as Chiral Selectors in Capillary Electrophoresis. <i>Molecules</i> , 2021, 26, 5271.	1.7	7
698	Functional materials in chiral capillary electrophoresis. <i>Coordination Chemistry Reviews</i> , 2021, 445, 214108.	9.5	25
699	Separation and Quantitation of Enantiomers. <i>Chromatographia CE Series</i> , 1998, , 70-100.	0.1	5
700	Applications of chiral capillary electrophoresis (cyclodextrin-capillary zone electrophoresis;) Tj ETQq1 1 0.784314 rgBT /Overlqck 10 T		
701	Pre-Concentration Approaches Combined with Capillary Electrophoresis in Bioanalysis of Chiral Cardiovascular Drugs. <i>Pharmaceutical Sciences</i> , 2015, 21, 229-243.	0.8	25
702	Modelling Enantiomer Separation by CE. <i>Chromatographia CE Series</i> , 2001, , 24-41.	0.1	0
703	Stereoselective Separations: Recent Advances in Capillary Electrophoresis and High-Performance Liquid Chromatography. <i>Handbook of Experimental Pharmacology</i> , 2003, , 45-75.	0.9	2
704	Chiral electromigration techniques in pharmaceutical and biomedical analysis. , 2013, , 225-244.		0
707	3 Techniques used in the analysis of optically active compounds. <i>3.7 Electrophoresis</i> . , 1995, , 243-278.		0
708	Determination of Drug-Cyclodextrin Binding Constants By Capillary Zone Electrophoresis. , 1996, , 675-678.		0
709	Electrolyte systems. , 1997, , 71-83.		0
710	Determination of enantiomeric purity of esomeprazole by capillary electrophoresis. <i>Bulletin of Medical Sciences</i> , 2020, 93, 93-101.	0.0	0
711	Enantiomer Separation by Capillary Electrophoresis. , 0, , 333-368.		1
712	Separability of stereoisomers by electrokinetic chromatography in presence of a neutral selector â€“ Fundamental aspects assessed by computer simulation. <i>Journal of Chromatography A</i> , 2022, 1673, 463087.	1.8	3

#	ARTICLE	IF	CITATIONS
713	Analytical methods for obtaining binding parameters of drug-protein interactions: A review. <i>Analytica Chimica Acta</i> , 2022, 1219, 340012.	2.6	12
714	Application of Remote-Sensing-Based Hydraulic Model and Hydrological Model in Flood Simulation. <i>Sustainability</i> , 2022, 14, 8576.	1.6	4
715	Separation of planar chiral ferrocenes by capillary electrokinetic chromatography and liquid chromatography. <i>Journal of Chromatography A</i> , 2022, 1677, 463306.	1.8	6
716	Recognition in the Domain of Molecular Chirality: From Noncovalent Interactions to Separation of Enantiomers. <i>Chemical Reviews</i> , 2022, 122, 13235-13400.	23.0	77
717	Modelling the simultaneous chiral separation of a group of drugs by electrokinetic chromatography using mixtures of cyclodextrins. <i>Journal of Chromatography A</i> , 2022, 1681, 463444.	1.8	3
718	A feasibility study on the use of hydrophobic eutectic solvents as pseudo-stationary phases in capillary electrophoresis for chiral separations. <i>Analytica Chimica Acta</i> , 2023, 1239, 340693.	2.6	7
719	Chiral Separations of Pyrethroid Acids Using Cyclodextrin Selectors. <i>Molecules</i> , 2022, 27, 8718.	1.7	1
720	Cyclodextrins as chiral selectors in capillary electrophoresis: Recent trends in mechanistic studies. <i>TrAC - Trends in Analytical Chemistry</i> , 2023, 160, 116987.	5.8	17
723	Chiral separation by capillary electrokinetic chromatography with hydrophobic deep eutectic solvents as pseudo-stationary phases. <i>Talanta</i> , 2023, 260, 124556.	2.9	3