

# Actions of two novel prostaglandin analogs, SC-29169 and SC-58100, on human isolated urinary bladder

General Pharmacology

23, 805-809

DOI: [10.1016/0306-3623\(92\)90228-c](https://doi.org/10.1016/0306-3623(92)90228-c)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Non-adrenergic, non-cholinergic control of the urinary bladder. <i>World Journal of Urology</i> , 1994, 12, 233-44.	2.2	48
2	Response of the Isolated Guinea Pig Bladder to Exogenous and Endogenous Leukotrienes. <i>Journal of Urology</i> , 1994, 152, 1281-1286.	0.4	24
3	Effect of Anoxia on Urethral Response to Phenylephrine. <i>Journal of Urology</i> , 1995, 154, 1527-1531.	0.4	6
4	Misoprostol associated autonomic dysreflexia in a traumatic tetraplegic patient. <i>Spinal Cord</i> , 1996, 34, 121-122.	1.9	5
5	Pharmacological characterization of thromboxane and prostanoid receptors in human isolated urinary bladder. <i>British Journal of Pharmacology</i> , 1998, 124, 865-872.	5.4	69
6	Prostaglandin E2 release from isolated bladder strips in rats with spinal cord injury. <i>International Journal of Urology</i> , 2006, 13, 271-276.	1.0	43
7	The role of prostanoids in urinary bladder physiology. <i>Nature Reviews Urology</i> , 2012, 9, 283-290.	3.8	46
8	Prostanoid receptors mediating contraction in rat, macaque and human bladder smooth muscle in vitro. <i>European Journal of Pharmacology</i> , 2015, 769, 274-279.	3.5	11