

CITATION REPORT

List of articles citing

Climatic variations in the Argentine plains during the last 18,000 years

DOI: 10.1016/0031-0182(93)90013-9

Palaeogeography, Palaeoclimatology, Palaeoecology, 1993, 101, 209-220.

Source: <https://exaly.com/paper-pdf/24002579/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
213	A probable scenario for a dry climate in central Amazonia during the late Quaternary. 1994 , 21, 121-128		24
212	A climatic model for southwestern Amazonia in Last Glacial times. 1994 , 21, 163-169		38
211	A provisional aminostratigraphical framework for late Quaternary marine deposits in Buenos Aires province, Argentina. 1995 , 128, 85-104		9
210	Late Quaternary marginal marine deposits and palaeoenvironments from northeastern Buenos Aires Province, Argentina: A review. <i>Quaternary Science Reviews</i> , 1995 , 14, 223-254	3.9	89
209	The nature, distribution and formation of pans in arid zones. <i>Earth-Science Reviews</i> , 1995 , 38, 1-69	10.2	119
208	The Mactracea, Tellinacea and Myacea (Bivalvia) in the marine Holocene of northeastern Buenos Aires Province (Argentina, South America): indicators of environmental change. 1995 , 19, 297-332		5
207	Late Holocene Neoglacial episodes in southern South America and southern Africa: a comparison. 1995 , 5, 361-368		44
206	Southern Westerlies during the Holocene deduced from geomorphological and pedological studies in the Norte Chico, Northern Chile (27°-33°S). <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 1996 , 123, 107-119	2.9	103
205	Late Quaternary Vegetational and Climatic Changes in the Pampa Grassland of Argentina. 1996 , 45, 73-88		130
204	Late-Holocene glacier variations in the Cordillera Darwin, Tierra del Fuego, Chile. 1996 , 6, 353-358		44
203	Changes in the atmospheric CH ₄ gradient between Greenland and Antarctica during the Holocene. 1997 , 102, 15987-15997		178
202	Preliminary Vegetation Maps of the World since the Last Glacial Maximum: An Aid to Archaeological Understanding. 1997 , 24, 623-647		136
201	Hypothesis on the cause of extinction of the South American mastodonts. 1997 , 10, 29-38		25
200	Comment on A. Prieto's (1996) Paper "Late Quaternary Vegetational and Climatic Changes in the Pampa Grassland of Argentina" 1997 , 48, 251-252		2
199	Reply to the Letter to the Editor on "Late Quaternary Vegetational and Climatic Changes in the Pampa Grassland of Argentina" by M. Iriondo. 1997 , 48, 253-256		
198	Origin of Dissolved Groundwater Sulphate in Coastal Plain Sediments of the Rio de la Plata, Eastern Argentina. 1997 , 3, 305-328		15
197	Habitat islands, genetic diversity, and gene flow in a Patagonian rodent. <i>Molecular Ecology</i> , 1998 , 7, 667-78		70

196	A late-Holocene (. 1998 , 8, 473-479		48
195	Gliptodontes y Cazadores-Recolectores de la Region Pampeana (Argentina). 1998 , 9, 111-134		46
194	Variation in isotopic composition (C, O and Sr) of Holocene <i>Macra isabelleana</i> (Bivalvia) from the coast of Buenos Aires Province, Argentina. 1998 , 8, 613-621		19
193	Quaternary charophytes from Salina del Bebedero, Argentina: their relation with extant taxa and palaeolimnological significance. 1999 , 21, 307-323		18
192	Last Glacial Maximum and Hypsithermal in the Southern Hemisphere. 1999 , 62, 11-19		17
191	Climatic fluctuations in the Central Region of Argentina in the last 1000 years. 1999 , 62, 35-47		71
190	Predominance of arid climates indicated by mammals in the pampas of Argentina during the Late Pleistocene and Holocene. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 1999 , 147, 257-281	2.9	143
189	Taphonomic processes affecting late Quaternary molluscs along the coastal area of Buenos Aires Province (Argentina, Southwestern Atlantic). <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 1999 , 149, 283-304	2.9	30
188	Stable isotope analyses on the last 30 ka molluscan fauna from Pampa grassland, Bonaerense region, Argentina. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 1999 , 153, 289-308	2.9	48
187	DISPERSAL, VICARIANCE, AND CLOCKS: HISTORICAL BIOGEOGRAPHY AND SPECIATION IN A COSMOPOLITAN PASSERINE GENUS (ANTHUS: MOTACILLIDAE). 1999 , 53, 1536-1552		87
186	Dispersal, Vicariance, and Clocks: Historical Biogeography and Speciation in a Cosmopolitan Passerine Genus (Anthus: Motacillidae). 1999 , 53, 1536		85
185	Late Quaternary climate history of the Bolivian Altiplano. 2000 , 72, 37-51		97
184	Climatic events during the Late Pleistocene and Holocene in the Upper Parana River: Correlation with NE Argentina and South-Central Brazil. 2000 , 72, 73-85		77
183	Patagonian dust in Antarctica. 2000 , 68-71, 83-86		49
182	Pole-Equator-Pole Paleoclimates of the Americas: PEP 1 A Review*. 2000 , 103-123		1
181	The Late Pleistocene in Amazonia: A Paleoclimatic Approach. 2000 , 209-224		2
180	Variation in skull morphology and mastication in the fossil giant armadillos <i>Pampatherium</i> spp. and allied genera (Mammalia: Xenarthra: Pampatheriidae), with comments on their systematics and distribution. 2001 , 20, 743-754		41
179	Bone Diagenesis and Taphonomic History of the Paso Otero 1 Bone Bed, Pampas of Argentina. 2001 , 28, 1277-1290		24

178	Evidence for Late Quaternary aeolian activity in the Roraima Guyana Region. 2001 , 43, 63-80		21
177	Mid- to Late-Holocene pollen-based biome reconstructions for Colombia. <i>Quaternary Science Reviews</i> , 2001 , 20, 1289-1308	3.9	70
176	Testing models of diversification in mice in the <i>Abrothrix olivaceus/xanthorhinus</i> complex in Chile and Argentina. <i>Molecular Ecology</i> , 2001 , 10, 397-405	5.7	52
175	Cytogeography of the South American <i>Turnera sidoides</i> L. complex (Turneraceae, Leiocarpaceae). 2001 , 137, 189-196		6
174	Late Quaternary Eolian Records of the Americas and Their Paleoclimatic Significance. 2001 , 183-216		33
173	Feedback between aeolian dust, climate, and atmospheric CO ₂ in glacial time. 2002 , 17, 11-1-11-11		74
172	Stable isotope composition of <i>Littoridina australis</i> from the coast of Buenos Aires province, Argentina, during Holocene climatic fluctuations. 2002 , 35, 79-88		15
171	Pollen-based biome reconstructions for Colombia at 3000, 6000, 9000, 12 000, 15 000 and 18 000 14C yr ago: Late Quaternary tropical vegetation dynamics. 2002 , 17, 113-129		53
170	Putative Meteoritic Craters in Río Cuarto (Central Argentina) Interpreted as Eolian Structures. 2002 , 91, 9-24		7
169	Great white shark teeth used as pendants and possible tools by early-middle Holocene terrestrial mammal hunter-gatherers in the Eastern Pampas (Southern South America). 2003 , 13, 222-231		16
168	Late Pleistocene and Holocene palaeoenvironments in Golfo San Jorge, Patagonia: molluscan evidence. 2003 , 194, 3-30		35
167	Independent evolution of migration on the South American landscape in a long-distance temperate-tropical migratory bird, Swainson's flycatcher (<i>Myiarchus swainsoni</i>). 2003 , 30, 925-937		47
166	A NEW SPECIES OF LIOLAEMUS (SQUAMATA: LIOLAEMIDAE) FROM NORTHEASTERN ARGENTINA AND SOUTHERN PARAGUAY. 2003 , 59, 283-292		10
165	Isotopic constraints on the source of Argentinian loess with implications for atmospheric circulation and the provenance of Antarctic dust during recent glacial maxima. 2003 , 212, 181-196		92
164	Loess of southern South America. <i>Quaternary Science Reviews</i> , 2003 , 22, 1987-2006	3.9	226
163	Micromorphology and OSL dating of the basal part of the loess-paleosol sequence at La Mesada in Tucumán province, Northwest Argentina. 2003 , 106-107, 111-117		22
162	Redescription, shell variability and geographic distribution of <i>Plagiodontes dentatus</i> (Wood, 1828) (Gastropoda: Orthalicidae: Odontostominae) from Uruguay and Argentina. 2003 , 154, 1		13
161	Geographical zonation in the Neotropics of tree species characteristic of the Paraguay-Paraná Basin. 2004 , 31, 1489-1501		48

160	Mammalian biostratigraphy of Pleistocene fluvial deposits in northern Uruguay, South America. 2004 , 115, 347-357	35
159	Climate change: detection and attribution of trends from long-term geologic data. 2004 , 171, 433-450	20
158	Upper Pleistocene and Holocene paleosalinity and trophic state changes in relation to sea level variation in Rocha Lagoon, southern Uruguay. 2004 , 32, 117-135	32
157	Pedosedimentary development of part of a Late Quaternary loess-palaeosol sequence in northwest Argentina. 2004 , 19, 567-576	22
156	Quaternary of the Pantanal, west-central Brazil. 2004 , 114, 23-34	129
155	Floodplain construction in an anastomosed river. 2004 , 114, 55-65	47
154	Late Pleistocene vertebrates from northern Uruguay: tools for biostratigraphic, climatic and environmental reconstruction. 2004 , 114, 129-142	69
153	Sea-level fluctuations during the last 8600 years in the de la Plata river (Argentina). 2004 , 114, 155-165	96
152	Using Historical Climate Data to Evaluate Climate Trends: Issues of Statistical Inference. 2004 , 15, 1-10	2
151	Pantanal: a large South American wetland at a crossroads. 2005 , 24, 391-401	169
150	Diet and habitat of toxodont megaherbivores (Mammalia, Notoungulata) from the late Quaternary of South and Central America. 2005 , 64, 113-124	93
149	Relationships between Holocene sea-level variations, trophic development, and climatic change in Negra Lagoon, Southern Uruguay. 2005 , 33, 253-263	37
148	A New Fossil Furnariid from the Pleistocene of Uruguay, With Remarks on Nasal Type, Cranial Kinetics, and Relationships of the Extinct Genus Pseudoseisuropsis. 2005 , 107, 114-127	13
147	Vallecito Formation (Miocene): The evolution of an eolian system in an Andean foreland basin (northwestern Argentina). 2005 , 19, 343-357	25
146	Paleozoogeographic, biostratigraphic, and systematic aspects of the Genus Sclerocalyptus (Xenarthra, Glyptodontidae) of Argentina. 2005 , 20, 121-129	17
145	Human occupation of the eastern La Plata Basin and the adjacent littoral region during the mid-Holocene. 2005 , 132, 23-36	3
144	Mid-late Holocene cultural and environmental dynamics in Eastern Uruguay. 2005 , 132, 37-45	38
143	Was there a population replacement during the Late mid-Holocene in the southeastern Pampas of Argentina? Archaeological evidence and Paleoecological basis. 2005 , 132, 95-105	33

142	A New Fossil Furnariid from the Pleistocene of Uruguay, With Remarks on Nasal Type, Cranial Kinetics, and Relationships of the Extinct Genus <i>Pseudoseisuropsis</i> . 2005 , 107, 114		11
141	Rock magnetic study of fluvial Holocene soil from Buenos Aires province (Argentina). 2006 , 58, 1381-1387		1
140	Relationships between Trophic State, Paleosalinity and Climatic Changes during the First Holocene Marine Transgression in Rocha Lagoon, Southern Uruguay. 2006 , 35, 699-713		29
139	Paleolimnological evidence of Holocene climatic changes in Lake Blanca, southern Uruguay. 2006 , 36, 151-163		23
138	Biodiversity and its conservation in the Pantanal of Mato Grosso, Brazil. 2006 , 68, 278-309		329
137	The comparative biodiversity of seven globally important wetlands: a synthesis. 2006 , 68, 400-414		190
136	Vegetation and climate change Since 14,810 14C yr B.P. in southeastern uruguay and implications for the rise of early Formative societies. 2006 , 65, 20-32		62
135	A 2000-Year Global Temperature Reconstruction Based on Non-Treering Proxies. 2007 , 18, 1049-1058		54
134	Geomorphology and chronology of Late Quaternary dune fields of western Argentina. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2007 , 251, 300-320	2.9	63
133	Climate, Environment and Society in the Pacific During the Last Millennium. 2007 , 6, v-302		60
132	Proposiço formal do Alogrupo Alto Rio Paran (SP, PR e MS). 2007 , 7, 49-70		5
131	Molecular evidence for a species complex in the patagonian lizard <i>Liolaemus bibronii</i> and phylogeography of the closely related <i>Liolaemus gracilis</i> (Squamata: Liolaemini). 2007 , 43, 952-73		67
130	Mitochondrial sequence diversity of the southernmost extant New World monkey, <i>Alouatta caraya</i> . 2007 , 43, 202-15		18
129	Chloroplast DNA variation in the <i>Turnera sidoides</i> L. complex (Turneraceae): biogeographical implications. 2007 , 34, 427-436		35
128	Holocene landscape evolution along the Andean piedmont, Bolivian Chaco. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2008 , 260, 505-520	2.9	19
127	Late Quaternary paleosoldiment-sequences and landscape evolution along the Andean piedmont, Bolivian Chaco. 2008 , 98, 34-54		23
126	.		
125	Grazing Impacts in Vegetated Dune Fields: Predictions From Spatial Pattern Analysis. 2008 , 61, 194-203		20

- 124 Un Aporte Crítico a Partir de los Comentarios Sobre Montículos de la Cuenca de la Laguna Merín: Tiempo, Espacio Y Sociedad **2008**, 19, 325-335 3
- 123 The Pantanal of Mato Grosso: Linking Ecological Research, Actual Use and Management for Sustainable Development. 908-943 2
- 122 Late Holocene palaeoenvironments of the Nutria Mansa 1 archaeological site, Argentina. **2009**, 41, 273-296 14
- 121 Comments on the paper "Fibrous-clay mineral formation and soil evolution in Aridisols of northeastern Patagonia, Argentina" by P.J. Bouza, M. Simón, J. Aguilar, H. del Valle and M. Rostagno. **2009**, 150, 226-227
- 120 The Mar Chiquita Lake: An indicator of intraplate deformation in the central plain of Argentina. **2009**, 111, 111-122 14
- 119 The species of *Eutatus* (Mammalia, Xenarthra): Assessment, morphology and climate. **2009**, 210, 66-75 10
- 118 Historical climatic extremes as indicators for typical scenarios of Holocene climatic periods in the Pampean plain. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **2009**, 283, 107-119 2.9 16
- 117 Holocene intramontane lake development: A new model in the Jáchal River Valley, Andean Precordillera, San Juan, Argentina. **2009**, 28, 229-238 5
- 116 Geographic patterns of morphological variation in *Turnera sidoides* subsp. *pinnatifida* (Turneraceae). **2010**, 284, 231-253 17
- 115 Late Holocene climate and environment of the SE Pampa grasslands, Argentina, inferred from biological indicators in shallow, freshwater Lake Nahuel Rucú **2010**, 44, 761-775 38
- 114 Last millennium environmental changes and climate inferences in the Southeastern Atlantic forest, Brazil. **2010**, 82, 717-29 13
- 113 Radar remote sensing of wind-driven land degradation processes in northeastern Patagonia. **2010**, 39, 62-75 18
- 112 Quaternary deformation around the Palo Negro area, Pampa Norte, Argentina. **2010**, 29, 627-641 16
- 111 Influence of the post-Miocene tectonic activity on the geomorphology between Andes and Pampa Depresión in the area of Provincia de La Pampa, Argentina. **2010**, 121, 152-166 12
- 110 Quaternary mammalian faunas of the Pampean Region. **2010**, 212, 176-186 12
- 109 Population genetic structure of long-tailed pygmy rice rats (*Oligoryzomys longicaudatus*) from Argentina and Chile based on the mitochondrial control region. **2010**, 88, 23-35 14
- 108 A review of the Quaternary Tayassuidae (Mammalia, Artiodactyla) from the Tarija Valley, Bolivia. **2010**, 34, 7-20 13
- 107 The relationship between emergence of mound builders in SE Uruguay and climate change inferred from opal phytolith records. **2011**, 245, 62-73 24

106	Paleoenvironments in the Paso Otero locality during Late Pleistocene–Holocene (Pampean region, Argentina): An interdisciplinary approach. 2011 , 245, 37-47		25
105	Mamíferos del pleistoceno superior de Santiago del Estero (Argentina) y sus afinidades paleobiogeográficas. 2011 , 51, 83-100		8
104	Patterns of cytotype variation of <i>Turnera sidoides</i> subsp. <i>pinnatifida</i> (Turneraceae) in mountain ranges of central Argentina. 2011 , 124, 25-34		10
103	The role of historical sources in the functional representation of a river in the new world. The case of the Argentinian Paran�. 2011 , 3, 227-249		4
102	The high Andes, gene flow and a stable hybrid zone shape the genetic structure of a wide-ranging South American parrot. 2011 , 8, 16		27
101	Roedores y Marsupiales en Torno Al L�mite Pleistoceno/Holoceno en Catamarca, Argentina: Extinciones y Evoluci� Ambiental. 2011 , 48, 336		7
100	Late Quaternary megafans, fans and fluvio-aeolian interactions in the Bolivian Chaco, Tropical South America. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2012 , 356-357, 75-88	2.9	47
99	Quaternary fluvial systems of tropics: Major issues and status of research. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2012 , 356-357, 1-15	2.9	13
98	Pasture clearing from invasive woody plants in the Pantanal: a tool for sustainable management or environmental destruction?. 2012 , 20, 111-122		36
97	Effects of Quaternary climatic changes on the phylogeography and historical demography of the subterranean rodent <i>Ctenomys porteus</i> . 2012 , 286, 48-57		18
96	Phylogeography and palaeodistribution modelling in the Patagonian steppe: the case of <i>Mulinum spinosum</i> (Apiaceae). 2012 , 39, 1041-1057		27
95	A review of Tertiary climate changes in southern South America and the Antarctic Peninsula. Part 1: Oceanic conditions. 2012 , 247-248, 1-20		31
94	Geochemical evidence of the 8.2 ka event and other Holocene environmental changes recorded in paleolagoon sediments, southeastern Brazil. 2012 , 77, 31-43		14
93	Paleoenvironments and faunal extinctions: Analysis of the archaeological assemblages at the Paso Otero locality (Argentina) during the Late Pleistocene–Early Holocene. 2013 , 299, 53-63		14
92	Sponge spicules indicate Holocene environmental changes on the Nabileque River floodplain, southern Pantanal, Brazil. 2013 , 49, 171-183		29
91	Biogeographical history and diversification of <i>Petunia</i> and <i>Calibrachoa</i> (Solanaceae) in the Neotropical Pampas grassland. 2013 , 171, 140-153		55
90	Distribution of <i>Lyncodon patagonicus</i> (Carnivora, Mustelidae): changes from the Last Glacial Maximum to the present. 2013 , 94, 339-350		15
89	Current state of knowledge regarding South America wetlands and their future under global climate change. 2013 , 75, 113-131		158

88	Geomorphic expression of the southern Central Andes forebulge (37°S, Argentina). 2013 , 25, 361-367	15
87	Complex Geographical Distribution of Ploidy Levels in <i>Polylepis australis</i> (Rosaceae), an Endemic Tree Line Species in Argentina. 2014 , 175, 955-961	10
86	Land-use and land-cover effects on regional biodiversity distribution in a subtropical dry forest: a hierarchical integrative multi-taxa study. 2014 , 14, 1549-1561	51
85	Patterns in community assemblage and species richness of small mammals across an altitudinal gradient in semi-arid Patagonia, Argentina. 2014 , 106, 18-26	18
84	Floral variation and environmental heterogeneity in a tristylous clonal aquatic of the Pantanal wetlands of Brazil. 2014 , 114, 1637-49	16
83	Holocene (~4.5–1.7 cal. kyr BP) paleoenvironmental conditions in central Argentina inferred from entire-shell and intra-shell stable isotope composition of terrestrial gastropods. 2014 , 24, 1193-1205	26
82	The paradigm of paraglacial megafans of the San Juan river basin, Central Andes, Argentina. 2014 , 55, 166-172	9
81	Diversification in the South American Pampas: the genetic and morphological variation of the widespread <i>Petunia axillaris</i> complex (Solanaceae). <i>Molecular Ecology</i> , 2014 , 23, 374-89	5-7 35
80	G-banding patterns of the genus <i>Ctenomys</i> from Uruguay (Rodentia Ctenomyidae): high karyotype variation but chromosome arm conservation. 2014 , 67, 178-184	2
79	Palaeogeographic reconstruction of Minchin palaeolake system, South America: The influence of astronomical forcing. 2014 , 5, 249-259	8
78	Effects of past climate variability on fire and vegetation in the cerrado savanna of the Huanchaca Mesetta, NE Bolivia. 2015 , 11, 835-853	15
77	Regional Extinctions and Quaternary Shifts in the Geographic Range of <i>Lestodelphys halli</i> , the Southernmost Living Marsupial: Clues for Its Conservation. 2015 , 10, e0132130	15
76	Technological Organization in Hunting/Scavenging and Butchering Sites of Megamammals in the Pampa Grasslands (Argentina). 2015 , 1, 343-359	3
75	Ecosystem loss assessment following hydroelectric dam flooding: The case of Yacyreté, Argentina. 2015 , 1, 50-60	7
74	Chronological distribution and disturbance factors to evaluate population dynamics in Western Pampas, Argentina. 2015 , 356, 74-88	9
73	A model for the Guaraní expansion in the La Plata Basin and littoral zone of southern Brazil. 2015 , 356, 54-73	36
72	Why was the Highveld treeless? Looking laterally to the Pampas for global edaphic principles beyond biogeographical accidents. 2015 , 101, 98-106	8
71	Climate fluctuations as a cause of rarity in fairy armadillos. 2015 , 80, 452-458	4

70	Climate changes and solar cycles recorded at the Holocene Paraná Delta, and their impact on human population. <i>Scientific Reports</i> , 2015 , 5, 12851	4.9	11
69	Natural grasslands in the Chaco. A neglected ecosystem under threat by agriculture expansion and forest-oriented conservation policies. 2015 , 123, 40-46		44
68	Palaeoenvironmental reconstruction based on charophytes and sedimentology: Can the mid-Holocene Optimum be recognised in western Argentina?. 2015 , 120, 31-38		4
67	Delving into the variations in tree species composition and richness across South American subtropical Atlantic and Pampean forests. 2015 , 8, 242-260		114
66	Cordão Formation: loess deposits in the southern coastal plain of the state of Rio Grande do Sul, Brazil. 2016 , 88, 2143-2166		10
65	Anthropogenic geomorphic change as a potential generator of renewable geologic resources in the humid Pampa of Argentina. 2016 , 142, 177-189		1
64	Dunes, hills, waterholes, and saltpeter beds: Attractors for human populations in western Pampa, Argentina. 2016 , 422, 163-173		6
63	Influence of environmental heterogeneity on the distribution and persistence of a subterranean rodent in a highly unstable landscape. 2016 , 144, 711-722		9
62	A review of Late Quaternary inland dune systems of South America east of the Andes. 2016 , 410, 96-110		25
61	Paleoenvironmental insights into the Quaternary evolution of the southern Brazilian coast based on fossil and modern diatom assemblages. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2016 , 446, 108-124	2.9	16
60	The Quaternary of Uruguay at Regional Scale. 2016 , 63-67		
59	Phylogeography of <i>Cnesterodon decemmaculatus</i> (Cyprinodontiformes: poeciliidae) in Southern Pampas, Argentina: ancient versus recent patterns in freshwater fishes. 2016 , 99, 293-307		10
58	Climate teleconnections and indicators of coastal systems response. 2016 , 122, 64-76		16
57	The role of small carnivores in the movement of bones: implications for the Pampas archaeofaunal record, Argentina. 2016 , 8, 257-276		13
56	Agricultural production and bird conservation in complex landscapes of the dry Chaco. 2016 , 11, 188-202		10
55	Paleoenvironmental reconstruction of Quaternary valley-fill successions in summit paleosurfaces of southern Sierras Pampeanas (Córdoba Province, Argentina). 2017 , 442, 50-58		1
54	Late Quaternary Cuiabá megafan, Brazilian Pantanal: Channel patterns and paleoenvironmental changes. 2017 , 438, 108-125		17
53	Out of Amazonia: Late-Holocene climate change and the Tupiã-Guarani trans-continental expansion. 2017 , 27, 967-975		19

52	What happened in the South American Gran Chaco? Diversification of the endemic frog genus <i>Lepidobatrachus</i> Budgett, 1899 (Anura: Ceratophryidae). 2018 , 123, 123-136	10
51	Enhancing edaphoclimatic zoning by adding multivariate spatial statistics to regional data. 2018 , 310, 170-177	7
50	Late Holocene palynology of a saline lake in the Pantanal of Nhecolândia, Brazil. 2018 , 42, 457-465	5
49	Diverse yet endangered: pollen dispersal and mating system reveal inbreeding in a narrow endemic plant. 2019 , 12, 169-180	6
48	Impact of recycling and lateral sediment input on grain size fining trends—Implications for reconstructing tectonic and climate forcings in ancient sedimentary systems. 2019 , 31, 866-891	4
47	The glaciogenic origin of the Pleistocene calcareous dust in Argentina on the basis of field, mineralogical, textural, and geochemical analyses. 2019 , 91, 218-233	2
46	Pleistocene Mammal Communities and Their Extinction. 2019 , 289-302	
45	Early Holocene water well in the Pampas of Argentina: Human responses to water shortage events. 2019 , 29, 145-157	1
44	Late Quaternary vegetation history based on phytolith records in the eastern Chaco (Argentina). 2019 , 505, 21-33	6
43	Patterns of Resource Use and Isotopic Niche Overlap Among Guanaco (<i>Lama guanicoe</i>), Pampas Deer (<i>Ozotoceros bezoarticus</i>) and Marsh Deer (<i>Blastocerus dichotomus</i>) in the Pampas. Ecological, Paleoenvironmental and Archaeological Implications. 2020 , 25, 411-444	11
42	Late Quaternary climatic variability in northern Patagonia, Argentina, based on $\delta^{18}O$ of modern and fossil shells of <i>Amiantis purpurata</i> (Bivalvia, Veneridae). <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2020 , 560, 110012	2.9 0
41	Comparative evaluation of urban versus agricultural nitrate sources and sinks in an unconfined aquifer by isotopic and multivariate analyses. 2020 , 741, 140374	15
40	Origin, evolution, and sedimentary records of islands in large anabranching tropical rivers: The case of the Upper Paraná River, Brazil. 2020 , 358, 107118	10
39	The use of wood during prehispanic times in the Upper Paraná Delta revealed through analysis of ancient charcoal. 2021 , 30, 193-212	4
38	Late Pleistocene-Holocene paleoenvironments in the middle basin of the Salado river, province of Buenos Aires, Argentina. 2021 , 105, 103001	3
37	The mutual ecogeographical range and paleoclimatic reconstruction during the Late Pleistocene-Holocene in the Pampas (Argentina) using meso and microvertebrate fossils. 2021 , 31, 983-992	2
36	Study of dugout canoes from the coast of La Plata River and the islands of the Paraná Delta, Argentina. 1-25	
35	Species distribution modeling reveals the ecological niche of extinct megafauna from South America. 1-8	2

34	Pleistocene Mammals from Pampean Region (Argentina). Biostratigraphic, Biogeographic, and Environmental Implications. 2021 , 4, 15	0
33	From Patagonia to R�o de la Plata: Multistep long-distance littoral transport of Andean volcaniclastic sand along the Argentine passive margin.	4
32	Aquatic ferns (Salviniaceae) in the Holocene sediments from the Bermejo River, Eastern Chaco, Argentina. 2021 , 109, 103273	1
31	Transcontinental retroarc sediment routing controlled by subduction geometry and climate change (Central and Southern Andes, Argentina). 2021 , 33, 3406	3
30	Paleoenvironmental Reconstruction of the Destacamento R�o Salado Member, Canal de Las Escobas Formation (Middle Holocene), Northeastern Buenos Aires, Argentina: Sedimentological and Paleontological Evidence. 2021 , 305-336	2
29	Large Rivers and their Floodplains. 2020 , 55-101	2
28	Geomorphology. 2007 , 33-52	12
27	Biogeography of the Forests of the Paraguay-Paran� Basin. 2006 , 193-211	6
26	Travelling to the south: Phylogeographic spatial diffusion model in <i>Monttea aphylla</i> (Plantaginaceae), an endemic plant of the Monte Desert. 2017 , 12, e0178827	16
25	Phylogenetic niche conservatism and plant diversification in South American subtropical grasslands along multiple climatic dimensions. 2020 , 43, e20180291	3
24	Le r�le des sources historiques dans la compr�hension des dynamiques fluviales d�un grand fleuve de plaine. Le cas du Paran� argentin. 2013 , 19, 445-462	2
23	The effects of past climate variability on fire and vegetation in the cerr�o savanna ecosystem of the Huanchaca Mesetta, Noel Kempff Mercado National Park, NE Bolivia.	4
22	Evolution of Small Mammal Communities during the Late Quaternary in a Middle-Altitude Valley of Northwestern Argentina. 2019 , 56, 153	2
21	Neohelice (Decapoda: Brachyura: Varunidae) from the Mid-Holocene, Canal de las Escobas Formation, Northeastern Buenos Aires, Argentina. 2019 , 56, 243	7
20	OUP accepted manuscript.	0
19	Registro f�sil y distribuci�n de <i>Anadenathera</i> en Argentina desde el Mioceno hasta la actualidad. 2015 , 71, e031	0
18	Fish. 2019 , 57-64	
17	Site Formation Processes Since the Early Holocene in a Planosol Sequence from the Laguna Negra Basin. 2019 , 177-192	

16	Introduction. 2020 , 1-13		
15	Social Trajectories of Hunter-Gatherer Societies in Central Argentina: Exploration and Colonization of a Desert Landscape, La Pampa Province, Argentina. 2021 , 81-118		○
14	Phylogeographic Relationships among (Lepidoptera: Bombycidae) Populations and Their Relationships to Inferred from Mitochondrial Genomes.. 2022 , 11,		1
13	Evolutionary history and colonization patterns of the wing dimorphic grasshopper <i>Dichroplus vittatus</i> in two Argentinean biomes.. <i>Scientific Reports</i> , 2022 , 12, 2920	4.9	
12	Polyploidy, genome size variation and diversification in an autopolyploid complex: the case of <i>Turnera sidoides</i> (Passifloraceae, Turneroideae). <i>Systematics and Biodiversity</i> , 2022 , 20, 1-18	1.7	○
11	Neutral and outlier single nucleotide polymorphisms disentangle the evolutionary history of a coastal Solanaceae species.. <i>Molecular Ecology</i> , 2022 ,	5.7	
10	The Pampean region (Argentina) underwent larger variation in aridity than in temperature during the late Pleistocene: New evidence from the isotopic analysis of mammalian taxa. <i>Quaternary Science Reviews</i> , 2022 , 286, 107555	3.9	○
9	Detrital zircon provenance and transport pathways of Pleistocene-Holocene eolian sediment in the Pampean Plains, Argentina. <i>Bulletin of the Geological Society of America</i> ,	3.9	○
8	Andean retroarc-basin dune fields and Pampean Sand Sea (Argentina): Provenance and drainage changes driven by tectonics and climate. <i>Earth-Science Reviews</i> , 2022 , 231, 104077	10.2	○
7	EVOLUTIONARY DYNAMICS OF AUTOPOLYPLOIDS IN NATURAL POPULATIONS: THE CASE OF TURNERA SIDOIDES COMPLEX. 2022 , 33, 71-81		
6	The polygenetic floodplain of the Upper ParanRiver. 2022 , 103985		
5	Leguminosae endemic to the Chaco facing quaternary climate fluctuations. 36,		○
4	Conservation opportunities for threatened paleochannel grasslands in the South American Dry Chaco. 2023 , 71, 126306		○
3	Population genetic characteristics of the bumble bee <i>Bombus ardens ardens</i> (Hymenoptera: Apidae) in South Korea using novel microsatellite markers. 2023 , 26, 102071		○
2	A new species of the highly polytypic South American rodent <i>Ctenomys</i> increases the diversity of the magellanicus clade. 73, 289-312		○
1	Soils of Pampa Ga'cho, the Mixed Prairies of Southern Brazil. 2023 , 299-342		○