

Molecular evolution and phylogenetic implications of internal  
sequences of Ribosomal DNA in Winteraceae

American Journal of Botany

80, 1042-1055

DOI: [10.1002/j.1537-2197.1993.tb15332.x](https://doi.org/10.1002/j.1537-2197.1993.tb15332.x)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Chloroplast DNA systematics: a review of methods and data analysis. American Journal of Botany, 1994, 81, 1205-1224.	0.8	480
2	Radiation of the endemic genus <i>Dendroseris</i> (Asteraceae) on the Juan Fernandez Islands: evidence from sequences of the its regions of nuclear ribosomal DNA. American Journal of Botany, 1994, 81, 1494-1501.	0.8	161
3	A molecular phylogeny of Arceuthobium (Viscaceae) based on nuclear ribosomal DNA internal transcribed spacer sequences. American Journal of Botany, 1994, 81, 1149-1160.	0.8	102
4	Length variation in the internal transcribed spacers of ribosomal DNA in <i>Picea abies</i> and related species. Theoretical and Applied Genetics, 1994, 89-89, 969-974.	1.8	16
5	A conserved sequence in internal transcribed spacer 1 of plant nuclear rRNA genes. Plant Molecular Biology, 1994, 26, 775-778.	2.0	171
6	USE OF RIBOSOMAL DNA INTERNAL TRANSCRIBED SPACERS FOR PHYLOGENETIC STUDIES IN DIATOMS1. Journal of Phycology, 1994, 30, 507-512.	1.0	72
7	A Phylogenetic Analysis of Epilobium (Onagraceae) Based on Nuclear Ribosomal DNA Sequences. Systematic Botany, 1994, 19, 363.	0.2	195
8	NOTHOFAGUS AND PACIFIC BIOGEOGRAPHY. Cladistics, 1995, 11, 5-32.	1.5	106
9	Molecular phylogenetics of the subtribe Gentianinae (Gentianaceae) inferred from the sequences of internal transcribed spacers (ITS) of nuclear ribosomal DNA. Plant Systematics and Evolution, 1995, 196, 207-226.	0.3	76
10	Molecular phylogeny of the Pooideae (Poaceae) based on nuclear rDNA (ITS) sequences. Theoretical and Applied Genetics, 1995, 90, 389-398.	1.8	126
11	The its Region of Nuclear Ribosomal DNA: A Valuable Source of Evidence on Angiosperm Phylogeny. Annals of the Missouri Botanical Garden, 1995, 82, 247.	1.3	1,471
12	Phylogenetic relationships in tribe Carduoideae (Asteraceae) based on ITS sequences. American Journal of Botany, 1995, 82, 1056-1068.	0.8	78
13	Phylogenetic relationships in Maloideae (Rosaceae): evidence from sequences of the internal transcribed spacers of nuclear ribosomal DNA and its congruence with morphology. American Journal of Botany, 1995, 82, 903-918.	0.8	97
14	ITS Sequences and the Phylogeny of the Genus Robinsonia (Asteraceae). Systematic Botany, 1995, 20, 55.	0.2	124
15	Generic boundaries in the tribe Sileneae (Caryophyllaceae) as inferred from nuclear rDNA sequences. Taxon, 1995, 44, 525-542.	0.4	122
16	Phylogenetic relationships of the monogenomic species of the wheat tribe, Triticeae (Poaceae), inferred from nuclear rDNA (internal transcribed spacer) sequences. Genome, 1995, 38, 211-223.	0.9	211
17	Phylogeny of Agavaceae Based on ITS rDNA Sequence Variation. American Journal of Botany, 1996, 83, 1225.	0.8	57
18	The Origins of Arabidopsis suecica (Brassicaceae) as Indicated by Nuclear rDNA Sequences. Systematic Botany, 1996, 21, 559.	0.2	177

#	ARTICLE	IF	CITATIONS
19	Zea systematics: ribosomal ITS evidence. <i>Molecular Biology and Evolution</i> , 1996, 13, 612-622.	3.5	202
20	Infrageneric phylogeny of the genus <i>Gentiana</i> (Gentianaceae) inferred from nucleotide sequences of the internal transcribed spacers (ITS) of nuclear ribosomal DNA. <i>American Journal of Botany</i> , 1996, 83, 641-652.	0.8	82
21	Phylogeny of Agavaceae based on ITS rDNA sequence variation. <i>American Journal of Botany</i> , 1996, 83, 1225-1235.	0.8	52
22	Molecular divergence between disjunct taxa in eastern Asia and eastern North America. <i>American Journal of Botany</i> , 1996, 83, 1373-1378.	0.8	35
23	a molecular phylogeny of apiaceae subfamily Apioideae: evidence from nuclear ribosomal DNA internal transcribed spacer sequences. <i>American Journal of Botany</i> , 1996, 83, 234-251.	0.8	229
24	ANALYSIS OF ALEXANDRIUM (DINOPHYCEAE) SPECIES USING SEQUENCES OF THE 5.8S RIBOSOMAL DNA AND INTERNAL TRANSCRIBED SPACER REGIONS <sup>1</sup> . <i>Journal of Phycology</i> , 1996, 32, 424-432.	1.0	193
25	RAPD patterns, nrDNA ITS sequences and morphological patterns in <i>Silene</i> section <i>Sedoideae</i> (Caryophyllaceae). <i>Plant Systematics and Evolution</i> , 1996, 201, 93-116.	0.3	25
26	Length Variation in the Nuclear Ribosomal DNA Internal Transcribed Spacer Region of Non-Flowering Seed Plants. <i>Systematic Botany</i> , 1996, 21, 109.	0.2	112
27	Cladistic Relationships of <i>Kalmia</i> , <i>Leiophyllum</i> , and <i>Loiseleuria</i> (Phyllodoceae, Ericaceae) Based on rbcL and nrITS Data. <i>Systematic Botany</i> , 1996, 21, 17.	0.2	28
28	Discordance Between ITS and Chloroplast Topologies in the <i>Boykinia</i> Group (Saxifragaceae). <i>Systematic Botany</i> , 1996, 21, 169.	0.2	102
29	Phylogenetic Implications of rbcL and ITS Sequence Variation in the Berberidaceae. <i>Systematic Botany</i> , 1996, 21, 381.	0.2	37
30	Chloroplast DNA phylogeny, reticulate evolution, and biogeography of <i>Paeonia</i> (Paeoniaceae). <i>American Journal of Botany</i> , 1997, 84, 1120-1136.	0.8	1,289
31	Molecular evidence for a Mediterranean origin of the Macaronesian endemic genus <i>Argyranthemum</i> (Asteraceae). <i>American Journal of Botany</i> , 1997, 84, 1595-1613.	0.8	135
32	Persistent nuclear ribosomal DNA sequence polymorphism in the <i>Amelanchier</i> agamic complex (Rosaceae). <i>Molecular Biology and Evolution</i> , 1997, 14, 81-90.	3.5	251
33	Origin and relationships of <i>Saintpaulia</i> (Gesneriaceae) based on ribosomal DNA internal transcribed spacer (ITS) sequences. <i>American Journal of Botany</i> , 1997, 84, 956-965.	0.8	114
34	Phylogenetic and genomic relationships in the genus <i>Glycine</i> Willd. based on sequences from the ITS region of nuclear rDNA. <i>Genome</i> , 1997, 40, 57-68.	0.9	98
35	On the evolutionary origins of the cacti. <i>Taxon</i> , 1997, 46, 217-232.	0.4	123
36	A phylogenetic study of <i>Polygonum</i> sect. <i>Tovara</i> (Polygonaceae) based on ITS sequences of nuclear ribosomal DNA. <i>Journal of Plant Biology</i> , 1997, 40, 47-52.	0.9	13

#	ARTICLE	IF	CITATIONS
37	ITS sequences and speciation on far eastern Indigofera (Leguminosae). Journal of Plant Research, 1997, 110, 339-346.	1.2	13
38	Two Divergent Intragenomic rDNA ITS2 Types within a Monophyletic Lineage of the Fungus Fusarium Are Nonorthologous. Molecular Phylogenetics and Evolution, 1997, 7, 103-116.	1.2	1,815
39	Title is missing!. Genetic Resources and Crop Evolution, 1998, 45, 399-406.	0.8	9
40	Sequencing of the internal transcribed spacer region ITS1 as a molecular tool detecting variation in the Stylosanthes guianensis species complex. Theoretical and Applied Genetics, 1998, 96, 869-877.	1.8	23
41	ITS sequences from nuclear rDNA suggest unexpected phylogenetic relationships between Euro-Mediterranean, East Asiatic and North American taxa of Quercus (Fagaceae). Plant Systematics and Evolution, 1998, 211, 129-139.	0.3	35
42	Lengths and nucleotide sequences of the internal spacers of nuclear ribosomal DNA in gymnosperms and pteridophytes. Plant Systematics and Evolution, 1998, 213, 199-205.	0.3	25
43	Choosing an Approach and an Appropriate Gene for Phylogenetic Analysis. , 1998, , 1-42.		204
44	Shrinking the Violets: Phylogenetic Relationships of Infrageneric Groups in Viola (Violaceae) Based on Internal Transcribed Spacer DNA Sequences. Systematic Botany, 1998, 23, 439.	0.2	125
45	Phylogenetic Incongruence: Window into Genome History and Molecular Evolution. , 1998, , 265-296.		443
46	Molecular evidence for an African origin of the Hawaiian endemic Hesperomannia (Asteraceae). Proceedings of the National Academy of Sciences of the United States of America, 1998, 95, 15440-15445.	3.3	82
47	Biogeography and Floral Evolution of Baobabs Adansonia, Bombacaceae as Inferred From Multiple Data Sets. Systematic Biology, 1998, 47, 181-207.	2.7	276
48	The tortoise and the hare: choosing between noncoding plastome and nuclear Adh sequences for phylogeny reconstruction in a recently diverged plant group. American Journal of Botany, 1998, 85, 1301-1315.	0.8	423
49	Molecular phylogeny of the genus Hypochaeris using internal transcribed spacers of nuclear rDNA: inference for chromosomal evolution. Molecular Biology and Evolution, 1998, 15, 345-354.	3.5	58
50	Phylogenetic relationships among Fagopyrum species revealed by the nucleotide sequences of the ITS region of the nuclear rRNA gene.. Genes and Genetic Systems, 1998, 73, 201-210.	0.2	61
51	New Fagopyrum species revealed by morphological and molecular analyses.. Genes and Genetic Systems, 1998, 73, 85-94.	0.2	29
52	Phylogeny and biogeography of Aralia sect. Aralia (Araliaceae). American Journal of Botany, 1998, 85, 866-875.	0.8	76
53	Molecular systematics and phylogeography of the <i>Gibberella fujikuroi</i> species complex. Mycologia, 1998, 90, 465-493.	0.8	795
54	Molecular systematics of Clerodendrum (Lamiaceae): ITS sequences and total evidence. American Journal of Botany, 1999, 86, 98-107.	0.8	63

#	ARTICLE	IF	CITATIONS
55	Phylogenetic relationships in <i>Lupinus</i> (Fabaceae: Papilionoideae) based on internal transcribed spacer sequences (ITS) of nuclear ribosomal DNA. <i>American Journal of Botany</i> , 1999, 86, 590-607.	0.8	130
56	Molecular evidence for allopolyploid speciation and a single origin of the narrow endemic <i>Draba ladina</i> (Brassicaceae). <i>American Journal of Botany</i> , 1999, 86, 1282-1289.	0.8	80
57	Utility of rDNA ITS sequences in the systematics of <i>Teucrium</i> section <i>Polium</i> (Lamiaceae). <i>Plant Systematics and Evolution</i> , 1999, 215, 49-70.	0.3	30
58	Phylogenetic utility of the nuclear rDNA ITS region in subfamily <i>Xoroideae</i> (Rubiaceae): Comparisons with cpDNA <i>rbcL</i> sequence data. <i>Plant Systematics and Evolution</i> , 1999, 217, 119-135.	0.3	70
59	Polyloid speciation in <i>Hedera</i> (Araliaceae): Phylogenetic and biogeographic insights based on chromosome counts and ITS sequences. <i>Plant Systematics and Evolution</i> , 1999, 219, 165-179.	0.3	84
60	Morphological evolution and ecological diversification of the forest-dwelling poppies ( <i>Papaveraceae</i> ): <i>Tj ETQq1 1 0.784314 rgBT /Over</i> <i>Evolution</i> , 1999, 219, 181-197.	0.3	42
61	Phylogeny of <i>Weigela</i> and <i>Diervilla</i> (Caprifoliaceae) Based on Nuclear rDNA ITS Sequences: Biogeographic and Taxonomic Implications. <i>Journal of Plant Research</i> , 1999, 112, 331-341.	1.2	28
62	Relationships among Old and New World <i>Alliums</i> according to ITS DNA sequence analysis. <i>Theoretical and Applied Genetics</i> , 1999, 98, 422-433.	1.8	53
63	Molecular evidence for the phylogenetic position of <i>Hanabusaya asiatica</i> Nakai (Campanulaceae), an endemic species in Korea. <i>Journal of Plant Biology</i> , 1999, 42, 168-173.	0.9	8
64	Evidence on the Monophyly of <i>Astragalus</i> (Fabaceae) and its Major Subgroups Based on Nuclear Ribosomal DNA ITS and Chloroplast DNA <i>trnL</i> Intron Data. <i>Systematic Botany</i> , 1999, 24, 409.	0.2	164
65	High levels of intra- and inter-individual polymorphism in the rDNA ITS1 of <i>Caulerpa racemosa</i> (Chlorophyta). <i>European Journal of Phycology</i> , 2000, 35, 349-356.	0.9	64
66	Phylogeny of <i>Cyclamen</i> L. (Primulaceae): Evidence from morphology and sequence data from the internal transcribed spacers of nuclear ribosomal DNA. <i>Plant Systematics and Evolution</i> , 2000, 220, 147-160.	0.3	26
67	Phylogenetic relationships and genetic diversity among members of the <i>Festuca-Lolium</i> complex (Poaceae) based on ITS sequence data. <i>Plant Systematics and Evolution</i> , 2000, 224, 33-53.	0.3	68
68	Heterogeneity of the internal transcribed spacer 1 (ITS1) in <i>Tulipa</i> (Liliaceae). <i>Plant Systematics and Evolution</i> , 2000, 225, 29-41.	0.3	16
69	The internal transcribed spacer 2 region of the nuclear ribosomal DNA and the phylogeny of the moss family <i>Hylocomiaceae</i> . <i>Plant Systematics and Evolution</i> , 2000, 224, 127-137.	0.3	17
70	Genome evolution in polyploids. <i>Plant Molecular Biology</i> , 2000, 42, 225-249.	2.0	1,439
71	Gall mite molecular phylogeny and its relationship to the evolution of plant host specificity. <i>Experimental and Applied Acarology</i> , 2000, 24, 831-861.	0.7	34
72	Phylogeny and Phytogeography of <i>Eupatorium</i> (Eupatorieae, Asteraceae): Insights from Sequence Data of the nrDNA ITS Regions and cpDNA RFLP. <i>Journal of Plant Research</i> , 2000, 113, 79-89.	1.2	30

#	ARTICLE	IF	CITATIONS
73	Phylogenetic Relationships of Maples ( <i>Acer</i> L.; <i>Aceraceae</i> ) Implied by Nuclear Ribosomal ITS Sequences. <i>Journal of Plant Research</i> , 2000, 113, 193-202.	1.2	40
74	The classification and geography of the flowering plants: Dicotyledons of the class <i>Angiospermae</i> . <i>Botanical Review</i> , The, 2000, 66, 441-647.	1.7	136
75	Testing relative evolutionary rates and estimating divergence times among six genera of <i>Rhizophoraceae</i> using cpDNA and nrDNA sequences. <i>Science Bulletin</i> , 2000, 45, 1011-1015.	1.7	13
76	Diversification of the North American shrub genus <i>Ceanothus</i> ( <i>Rhamnaceae</i> ): conflicting phylogenies from nuclear ribosomal DNA and chloroplast DNA. <i>American Journal of Botany</i> , 2000, 87, 108-123.	0.8	83
77	High intraindividual variation in internal transcribed spacer sequences in <i>Aeschynanthus</i> ( <i>Gesneriaceae</i> ): implications for phylogenetics. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2000, 267, 1407-1415.	1.2	50
78	Genomic Origin and Organization of the Hybrid <i>Poa jemtlandica</i> ( <i>Poaceae</i> ) Verified by Genomic In Situ Hybridization and Chloroplast DNA Sequences. <i>Annals of Botany</i> , 2000, 85, 439-445.	1.4	33
79	Rapid Diversification of a Species-Rich Genus of Neotropical Rain Forest Trees. <i>Science</i> , 2001, 293, 2242-2245.	6.0	710
80	The phylogeny of ( <i>Gentianaceae</i> ) and its colonization of the southern hemisphere as revealed by nuclear and chloroplast DNA sequence variation. <i>Organisms Diversity and Evolution</i> , 2001, 1, 61-79.	0.7	154
81	Phylogenetic Patterns in Northern Hemisphere Plant Geography. <i>International Journal of Plant Sciences</i> , 2001, 162, S41-S52.	0.6	238
82	Phylogenetic Analysis of Sugarcane Rusts Based on Sequences of ITS, 5.8 S rDNA and D1/D2 Regions of LSU rDNA. <i>Journal of General Plant Pathology</i> , 2001, 67, 28-36.	0.6	53
83	Extensive Ribosomal DNA Genic Variation in the Columnar Cactus <i>Lophocereus</i> . <i>Journal of Molecular Evolution</i> , 2001, 53, 124-134.	0.8	73
84	Sources of error and confidence intervals in estimating the age of angiosperms from <i>rbcL</i> and 18S rDNA data. <i>American Journal of Botany</i> , 2001, 88, 1499-1516.	0.8	235
85	Phylogenetic Studies in <i>Streptocarpus</i> ( <i>Gesneriaceae</i> ): Reconstruction of Biogeographic History and Distribution Patterns. <i>Systematics and Geography of Plants</i> , 2001, 71, 545.	0.1	32
86	Phylogenetic relationships in family <i>Magnoliaceae</i> inferred from <i>ndhF</i> sequences. <i>American Journal of Botany</i> , 2001, 88, 717-728.	0.8	84
87	Divergent and reticulate species relationships in <i>Leucaena</i> ( <i>Fabaceae</i> ) inferred from multiple data sources: insights into polyploid origins and nrDNA polymorphism. <i>American Journal of Botany</i> , 2002, 89, 1057-1073.	0.8	87
88	HARDLY A RELICT: FREEZING AND THE EVOLUTION OF VESSELLESS WOOD IN <i>WINTERACEAE</i> . <i>Evolution; International Journal of Organic Evolution</i> , 2002, 56, 464.	1.1	1
89	PHYLOGENY OF <i>GAERTNERA</i> LAM. ( <i>RUBIACEAE</i> ) BASED ON MULTIPLE DNA MARKERS: EVIDENCE OF A RAPID RADIATION IN A WIDESPREAD, MORPHOLOGICALLY DIVERSE GENUS. <i>Evolution; International Journal of Organic Evolution</i> , 2002, 56, 42.	1.1	4
90	GENETIC DIFFERENTIATION BETWEEN NOBLE CRAYFISH, <i>ASTACUS ASTACUS</i> (L.), POPULATIONS DETECTED BY MICROSATELLITE LENGTH VARIATION IN THE RDNA ITS1 REGION. <i>Knowledge and Management of Aquatic Ecosystems: an International Journal on Aquatic Ecosystems</i> , 2002, , 691-706.	0.4	25

#	ARTICLE	IF	CITATIONS
91	Phylogenetic relationships in the Gesnerioideae (Gesneriaceae) based on nrDNA ITS and cpDNA <i>trnL</i> and <i>trnT</i> spacer region sequences. <i>American Journal of Botany</i> , 2002, 89, 296-311.	0.8	76
92	Molecular evidence for the taxonomic identity of Korean Adonis (Ranunculaceae). <i>Journal of Plant Research</i> , 2002, 115, 217-223.	1.2	15
93	Three nonorthologous ITS1 types are present in a polypore fungus <i>Trichaptum abietinum</i> . <i>Molecular Phylogenetics and Evolution</i> , 2002, 23, 112-122.	1.2	76
94	Evidence of multiple alleles of the nuclear 18S ribosomal RNA gene in sturgeon (Family: Acipenseridae). <i>Journal of Applied Ichthyology</i> , 2002, 18, 290-297.	0.3	28
95	Molecular phylogeny of genus <i>Vigna</i> subgenus <i>Ceratotropis</i> based on rDNA ITS and <i>atpB-rbcL</i> intergenic spacer of cpDNA sequences. <i>Genetica</i> , 2002, 114, 129-145.	0.5	68
96	A phylogeny of the Munnoziinae (Asteraceae, Liabeae): circumscription of <i>Munnozia</i> and a new placement of <i>M. perfoliata</i> . <i>Plant Systematics and Evolution</i> , 2003, 239, 171-185.	0.3	9
97	Unexpected high divergence in nrDNA ITS and extensive parallelism in floral morphology of <i>Pedicularis</i> (Orobanchaceae). <i>Plant Systematics and Evolution</i> , 2003, 240, 91-105.	0.3	44
98	Characterization of angiosperm nrDNA polymorphism, paralogy, and pseudogenes. <i>Molecular Phylogenetics and Evolution</i> , 2003, 29, 435-455.	1.2	384
99	Nuclear ribosomal DNA sequence polymorphism and hybridization in checker mallows ( <i>Sidalcea</i> ). <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 4</i>	1.2	45
100	Ribosomal ITS sequences and plant phylogenetic inference. <i>Molecular Phylogenetics and Evolution</i> , 2003, 29, 417-434.	1.2	1,592
101	Phylogenetic relationships among species of <i>Hypochaeris</i> (Asteraceae, Cichorieae) based on ITS, plastid <i>trnL</i> intron, <i>trnL</i> spacer, and <i>matK</i> sequences. <i>American Journal of Botany</i> , 2003, 90, 496-507.	0.8	61
102	Molecular Systematic Analysis Reveals Cryptic Tertiary Diversification of a Widespread Tropical Rain Forest Tree. <i>American Naturalist</i> , 2003, 162, 691-703.	1.0	186
103	Multiple Hybrid Formation in Natural Populations: Concerted Evolution of the Internal Transcribed Spacer of Nuclear Ribosomal DNA (ITS) in North American <i>Arabis divaricarpa</i> (Brassicaceae). <i>Molecular Biology and Evolution</i> , 2003, 20, 338-350.	3.5	212
104	A first molecular phylogenetic analysis of <i>Passiflora</i> (Passifloraceae). <i>American Journal of Botany</i> , 2003, 90, 1229-1238.	0.8	99
105	Phylogeny, Taxonomy, and Systematic Position of <i>Clethra</i> (Clethraceae, Ericales) with Notes on Biogeography: Evidence from Plastid and Nuclear DNA Sequences. <i>International Journal of Plant Sciences</i> , 2003, 164, 997-1006.	0.6	15
106	Biogeography of Discontinuously Distributed Hydrophytes: A Molecular Appraisal of Intercontinental Disjunctions. <i>International Journal of Plant Sciences</i> , 2003, 164, 917-932.	0.6	137
107	Phylogenetic analysis of ITS sequences suggests a Pliocene origin for the bipolar distribution of <i>Scleranthus</i> (Caryophyllaceae). <i>Australian Systematic Botany</i> , 2003, 16, 301.	0.3	37
108	Molecular genetic evidence for interspecific hybridization among endemic Hispaniolan <i>Bursera</i> (Burseraceae). <i>American Journal of Botany</i> , 2004, 91, 976-984.	0.8	35

#	ARTICLE	IF	CITATIONS
109	Recent Origin and Phylogenetic Utility of Divergent ITS Putative Pseudogenes: A Case Study from Naucleaeae (Rubiaceae). <i>Systematic Biology</i> , 2004, 53, 177-192.	2.7	106
110	Testing Hypotheses on Disjunctions Present in the Primarily Woody Boraginales: Ehretiaceae, Cordiaceae, and Heliotropiaceae, Inferred from ITS1 Sequence Data. <i>International Journal of Plant Sciences</i> , 2004, 165, S123-S135.	0.6	37
111	Phylogenetic relationships and biogeography of <i>Fuchsia</i> (Onagraceae) based on noncoding nuclear and chloroplast DNA data. <i>American Journal of Botany</i> , 2004, 91, 601-614.	0.8	93
112	Use of nuclear genes for phylogeny reconstruction in plants. <i>Australian Systematic Botany</i> , 2004, 17, 145.	0.3	367
113	Evolution and biogeography of the austral genus <i>Phyllocladus</i> (Podocarpaceae). <i>Journal of Biogeography</i> , 2004, 31, 1569-1577.	1.4	40
114	Taxonomic and phytogeographic implications from ITS phylogeny in <i>Berberis</i> (Berberidaceae). <i>Journal of Plant Research</i> , 2004, 117, 175-82.	1.2	54
115	The rDNA ITS Region in the Lessepsian Marine Angiosperm <i>Halophila stipulacea</i> (Forssk.) Aschers. (Hydrocharitaceae): Intragenomic Variability and Putative Pseudogenic Sequences. <i>Journal of Molecular Evolution</i> , 2004, 58, 115-121.	0.8	48
116	Evolution of GCYC, a Gesneriaceae homolog of CYCLOIDEA, within Gesnerioideae (Gesneriaceae). <i>Molecular Phylogenetics and Evolution</i> , 2004, 31, 765-779.	1.2	68
117	Evolution of multiple paralogous adenosine kinase genes in the moss genus <i>Hygroamblystegium</i> : phylogenetic implications. <i>Molecular Phylogenetics and Evolution</i> , 2004, 31, 505-516.	1.2	19
118	Floral development and molecular phylogeny support the generic status of <i>Tasmannia</i> (Winteraceae). <i>American Journal of Botany</i> , 2004, 91, 321-331.	0.8	30
119	Phylogeny, Molecular and Fossil Dating, and Biogeographic History of Annonaceae and Myristicaceae (Magnoliales). <i>International Journal of Plant Sciences</i> , 2004, 165, S55-S67.	0.6	77
120	Taxonomic and Phylogenetic Evaluation of the American "Thlaspi" Species: Identity and Relationship to the Eurasian Genus <i>Noccaea</i> (Brassicaceae). <i>Systematic Botany</i> , 2004, 29, 375-384.	0.2	54
121	Paraphyly in Tribe Onagreae: Insights into Phylogenetic Relationships of Onagraceae Based on Nuclear and Chloroplast Sequence Data. <i>Systematic Botany</i> , 2004, 29, 147-164.	0.2	74
122	Molecular Phylogeny and Biogeography of <i>Ribes</i> (Grossulariaceae), with an Emphasis on Gooseberries (subg. <i>Grossularia</i> ). <i>Systematic Botany</i> , 2004, 29, 77-96.	0.2	55
123	A Phylogenetic Analysis within Tribes Gloxinieae and Gesnerieae (Gesnerioideae: Gesneriaceae). <i>Systematic Botany</i> , 2004, 29, 947-958.	0.2	19
124	Phylogenetic Relationships in the Juncaceae Inferred from Nuclear Ribosomal DNA Internal Transcribed Spacer Sequence Data. <i>International Journal of Plant Sciences</i> , 2005, 166, 397-413.	0.6	22
125	INTRAGENOMIC NUCLEOTIDE POLYMORPHISM AMONG SMALL SUBUNIT (18S) RDNA PARALOGS IN THE DIATOM GENUS <i>SKELETONEMA</i> (BACILLARIOPHYTA)1. <i>Journal of Phycology</i> , 2005, 41, 1248-1257.	1.0	77
126	Phylogeny of <i>Aconitum</i> subgenus <i>Aconitum</i> (Ranunculaceae) inferred from ITS sequences. <i>Plant Systematics and Evolution</i> , 2005, 252, 11-25.	0.3	58



#	ARTICLE	IF	CITATIONS
127	Molecular phylogeny in Indian <i>Tinospora</i> species by DNA based molecular markers. <i>Plant Systematics and Evolution</i> , 2005, 256, 75-87.	0.3	15
128	Molecular phylogeny of <i>Phalaenopsis</i> Blume (Orchidaceae) based on the internal transcribed spacer of the nuclear ribosomal DNA. <i>Plant Systematics and Evolution</i> , 2005, 256, 1-16.	0.3	30
129	CLIMATE AND LIFE-HISTORY EVOLUTION IN EVENING PRIMROSES (OENOTHERA, ONAGRACEAE): A PHYLOGENETIC COMPARATIVE ANALYSIS. <i>Evolution; International Journal of Organic Evolution</i> , 2005, 59, 1914.	1.1	4
130	Rapid speciation and the evolution of hummingbird pollination in neotropical <i>Costus</i> subgenus <i>Costus</i> ( <i>Costaceae</i> ): evidence from nrDNA ITS and ETS sequences. <i>American Journal of Botany</i> , 2005, 92, 1899-1910.	0.8	204
131	Untangling <i>Gloxinieae</i> (Gesneriaceae). I. Phylogenetic patterns and generic boundaries inferred from nuclear, chloroplast, and morphological cladistic datasets. <i>Taxon</i> , 2005, 54, 389-410.	0.4	40
132	Molecular Phylogeny of <i>Anagallis</i> (Myrsinaceae) Based on ITS, trnL, and ndhF Sequence Data. <i>International Journal of Plant Sciences</i> , 2005, 166, 1019-1028.	0.6	22
133	Evolution by Reticulation: European <i>Dogroses</i> Originated by Multiple Hybridization Across the Genus <i>Rosa</i> . <i>Journal of Heredity</i> , 2005, 96, 4-14.	1.0	105
134	Evolution of the New Zealand mountain flora: Origins, diversification and dispersal. <i>Organisms Diversity and Evolution</i> , 2005, 5, 237-247.	0.7	127
135	Phylogeny of <i>Brassaiopsis</i> ( <i>Araliaceae</i> ) in Asia Based on Nuclear ITS and 5S-NTS DNA Sequences. <i>Systematic Botany</i> , 2005, 30, 872-886.	0.2	19
136	Phylogeny and Biogeography of Isophyllous Species of <i>Campanula</i> ( <i>Campanulaceae</i> ) in the Mediterranean Area. <i>Systematic Botany</i> , 2006, 31, 862-880.	0.2	79
137	A Nuclear Ribosomal DNA Phylogeny of <i>Acer</i> Inferred with Maximum Likelihood, Splits Graphs, and Motif Analysis of 606 Sequences. <i>Evolutionary Bioinformatics</i> , 2006, 2, 117693430600200.	0.6	51
138	Origin and diversification of <i>Thymelaea</i> ( <i>Thymelaeaceae</i> ): inferences from a phylogenetic study based on ITS (rDNA) sequences. <i>Plant Systematics and Evolution</i> , 2006, 257, 159-187.	0.3	17
139	A survey of nuclear ribosomal internal transcribed spacer substitution rates across angiosperms: an approximate molecular clock with life history effects. <i>BMC Evolutionary Biology</i> , 2006, 6, 36.	3.2	291
140	Phylogeny and Androecial Evolution in <i>Schisandraceae</i> , Inferred from Sequences of Nuclear Ribosomal DNA ITS and Chloroplast DNA trnL Regions. <i>International Journal of Plant Sciences</i> , 2006, 167, 539-550.	0.6	25
141	Phylogenetic relationships and generic boundaries in the <i>Episcieae</i> (Gesneriaceae) inferred from nuclear, chloroplast, and morphological data. <i>Taxon</i> , 2006, 55, 313-336.	0.4	46
142	The evolutionary history of <i>Sium</i> sensu lato ( <i>Apiaceae</i> ): dispersal, vicariance, and domestication as inferred from ITS rDNA phylogeny. <i>American Journal of Botany</i> , 2006, 93, 747-761.	0.8	45
143	Boreotropical migration explains hybridization between geographically distant lineages in the pantropical clade <i>Sideroxyleae</i> ( <i>Sapotaceae</i> ). <i>American Journal of Botany</i> , 2007, 94, 1491-1505.	0.8	52
144	Diversification of the Old World <i>Salsoleae</i> s.l. ( <i>Chenopodiaceae</i> ): Molecular Phylogenetic Analysis of Nuclear and Chloroplast Data Sets and a Revised Classification. <i>International Journal of Plant Sciences</i> , 2007, 168, 931-956.	0.6	145

#	ARTICLE	IF	CITATIONS
145	Measuring rDNA diversity in eukaryotic microbial systems: how intragenomic variation, pseudogenes, and PCR artifacts confound biodiversity estimates. <i>Molecular Ecology</i> , 2007, 16, 5326-5340.	2.0	267
146	Variation and similarities in pollen features in some basal angiosperms, with some taxonomic implications. <i>Plant Systematics and Evolution</i> , 2007, 263, 59-75.	0.3	14
147	Phylogeny of Bonatea (Orchidaceae: Habenariinae) based on molecular and morphological data. <i>Plant Systematics and Evolution</i> , 2007, 263, 253-268.	0.3	13
148	A Seed Related to Myristicaceae in the Early Eocene of Southern England. <i>Systematic Botany</i> , 2008, 33, 636-646.	0.2	19
149	Intra-individual and intra-species heterogeneity in nuclear rDNA ITS region of <i>Vigna</i> species from subgenus <i>Ceratotropis</i> . <i>Genetical Research</i> , 2008, 90, 299-316.	0.3	26
150	A phylogenetic analysis of Dipsacaceae based on four DNA regions. <i>Plant Systematics and Evolution</i> , 2009, 279, 69-86.	0.3	26
151	The Hawaiian Archipelago is a stepping stone for dispersal in the Pacific: an example from the plant genus <i>Melicope</i> (Rutaceae). <i>Journal of Biogeography</i> , 2009, 36, 230-241.	1.4	83
152	New Zealand phylogeography: evolution on a small continent. <i>Molecular Ecology</i> , 2009, 18, 3548-3580.	2.0	217
153	Phylogenetic relationships and generic delimitation in Inuleae subtribe Inulinae (Asteraceae) based on ITS and cpDNA sequence data. <i>Cladistics</i> , 2009, 25, 319-352.	1.5	39
154	Generic relationships and dating of lineages in Winteraceae based on nuclear (ITS) and plastid (rpS16) Tj ETQq1 1 0.784314 rgBT /Over	1.2	48
155	Molecular biology of aromatic plants and spices. A review.. <i>Flavour and Fragrance Journal</i> , 2010, 25, 272-281.	1.2	18
156	Plant or fungal sequences? An alternative optimized PCR protocol to avoid ITS (nrDNA) misamplification. <i>Brazilian Archives of Biology and Technology</i> , 2010, 53, 141-152.	0.5	12
157	Biogeographic Patterns of Diversification and the Origins of <i>Cleome</i> in the Neotropics (Cleomaceae). <i>Systematic Botany</i> , 2010, 35, 811-826.	0.2	77
158	Colonizing the American continent: Systematics of the genus <i>Arabis</i> in North America (Brassicaceae). <i>American Journal of Botany</i> , 2010, 97, 1040-1057.	0.8	35
159	<i>Shuaria</i> (Gesneriaceae), an Arborescent New Genus from the Cordillera del Cãndor and Amazonian Ecuador. <i>Systematic Botany</i> , 2010, 35, 662-674.	0.2	21
160	Evolution And Diversification Of Brook Weeds (Samolus, Samolaceae, Ericales). <i>International Journal of Plant Sciences</i> , 2011, 172, 250-266.	0.6	5
161	Independent Origin of Radial Floral Symmetry in the Gloxinieae (Gesnerioideae: Gesneriaceae) is Supported by the Rediscovery of <i>Phinaea pulchella</i> in Cuba. <i>Systematic Botany</i> , 2011, 36, 757-767.	0.2	7
162	Genetic and chemical components analysis of <i>Papaver setigerum</i> naturalized in Korea. <i>Forensic Science International</i> , 2012, 222, 387-393.	1.3	13

#	ARTICLE	IF	CITATIONS
163	Phylogenetic relationships among the North American cleomoids (Cleomaceae): A test of Iltis's reduction series. <i>American Journal of Botany</i> , 2013, 100, 2102-2111.	0.8	10
164	Molecular dating of Winteraceae reveals a complex biogeographical history involving both ancient Gondwanan vicariance and long-distance dispersal. <i>Journal of Biogeography</i> , 2014, 41, 894-904.	1.4	39
165	The Origin of Oranges: A Multi-locus Phylogeny of Rutaceae Subfamily Aurantioideae. <i>Systematic Botany</i> , 2016, 40, 1053-1062.	0.2	19
166	ITS non-concerted evolution and rampant hybridization in the legume genus <i>Lespedeza</i> (Fabaceae). <i>Scientific Reports</i> , 2017, 7, 40057.	1.6	43
167	Phylogenetic relationships among Egyptian species of Anthemideae (Asteraceae) using ITS markers. <i>Chromosome Botany</i> , 2017, 12, 1-6.	0.4	0
168	A Winteraceae pollen tetrad from the early Paleocene of western Greenland, and the fossil record of Winteraceae in Laurasia and Gondwana. <i>Journal of Biogeography</i> , 2018, 45, 567-581.	1.4	15
169	Herbarium collection-based phylogenetics of the ragweeds ( <i>Ambrosia</i> , Asteraceae). <i>Molecular Phylogenetics and Evolution</i> , 2018, 120, 335-341.	1.2	28
170	DNA barcoding the flowering plants from the tropical coral islands of Xisha (China). <i>Ecology and Evolution</i> , 2018, 8, 10587-10593.	0.8	8
171	Intergeneric relationships in the <i>Gorteria</i> clade of Arctotideae-Gorteriinae (Asteraceae), with description of a new genus, <i>Roessleria</i> . <i>South African Journal of Botany</i> , 2018, 118, 216-231.	1.2	0
172	The Eurasian steppe belt in time and space: Phylogeny and historical biogeography of the false flax ( <i>Camelina</i> Crantz, <i>Camelineae</i> , Brassicaceae). <i>Flora: Morphology, Distribution, Functional Ecology of Plants</i> , 2019, 260, 151477.	0.6	22
173	Molecular phylogeny and divergence times of <i>Onosma</i> ( <i>Boraginaceae</i> s.s.) based on nrDNA ITS and plastid <i>rpl32</i> and <i>trnL</i> (UAG) and <i>trnH-psbA</i> sequences. <i>Nordic Journal of Botany</i> , 2019, 37, e02060.	0.2	19
174	Diversification of New World Cleomaceae with emphasis on <i>Tarenaya</i> and the description of <i>Iltisiella</i> , a new genus. <i>Taxon</i> , 2020, 69, 321-336.	0.4	8
175	First South American Record of Winteroxylon, Eocene of Laguna del Hunco (Chubut, Patagonia, Argentina). <i>Journal of Paleontology</i> , 2020, 94, 185-197.	0.6	7
176	Molecular Systematics: 1994-1995. , 1997. , 470-499.		5
177	Nuclear and plastid DNA sequences reveal complex reticulate patterns in Australian water-lilies ( <i>Nymphaea</i> subgenus <i>Anecphyra</i> , <i>Nymphaeaceae</i> ). <i>Australian Systematic Botany</i> , 2008, 21, 229.	0.3	23
178	The Evolution of Ribosomal DNA Divergent Paralogues and Phylogenetic Implications. <i>Genetics</i> , 1997, 145, 821-832.	1.2	463
179	On the Evolutionary and Biogeographic History of <i>Saxifraga</i> sect. <i>Trachyphyllum</i> (Gaud.) Koch ( <i>Saxifragaceae</i> Juss.). <i>PLoS ONE</i> , 2013, 8, e69814.	1.1	25
180	Pollen morphology of extant Winteraceae: a study allowing SEM-based affiliation of its fossil representatives. <i>Acta Palaeobotanica</i> , 2017, 57, 339-396.	0.2	6

#	ARTICLE	IF	CITATIONS
181	The character of the New Zealand land snail fauna and communities: some evolutionary and ecological perspectives. <i>Records of the Western Australian Museum, Supplement</i> , 2005, 68, 53.	0.5	20
183	Phylogenetic Study of Korean <i>Chrysosplenium</i> Based on nrDNA ITS Sequences. <i>Korean Journal of Plant Resources</i> , 2011, 24, 358-369.	0.2	7
184	Molecular Phylogenetic Studies of Korean <i>Hydrocotyle</i> L.. <i>Korean Journal of Plant Resources</i> , 2012, 25, 490-497.	0.2	5
185	Ribosomal Internal Transcribed Spacer (ITS) DNA Variation in <i>Millepora</i> . <i>Journal of Marine Science: Research &amp; Development</i> , 2015, 06, .	0.4	0
186	Assessment of genetic diversity among different sugarcane genotypes using internal transcribed spacer (ITS) region of the ribosomal DNA (rDNA). <i>GSC Biological and Pharmaceutical Sciences</i> , 2018, 5, 017-025.	0.1	2
187	Identification of <i>Saposhinkoviae Radix</i> in concentrated Chinese medicine preparations by nested PCR and DNA sequencing methods. <i>Journal of Food and Drug Analysis</i> , 2005, 13, .	0.9	1
189	Extensive nrDNA ITS polymorphism in <i>Lycium</i> : Non-concerted evolution and the identification of pseudogenes. <i>Frontiers in Plant Science</i> , 0, 13, .	1.7	2
190	Phylogenetic inference and secondary structure predictions of Turkish genotypes of <i>Coriandrum sativum</i> (L.) based on ITS4 and ITS5 nrDNA sequences. <i>Plant Biotechnology Reports</i> , 0, , .	0.9	0
191	Evaluation of Genetic Variability among Three <i>Pistacia</i> Species Using Internal Transcribed Spacer 1 (ITS1) Marker. <i>Diversity</i> , 2022, 14, 1051.	0.7	2