

CITATION REPORT

List of articles citing

Quantitative assessment of tactile allodynia in the rat paw

DOI: 10.1016/0165-0270(94)90144-9
Journal of Neuroscience Methods, 1994, 53, 55-63.

Source: <https://exaly.com/paper-pdf/25107750/citation-report.pdf>

Version: 2024-04-29

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
2209	P2Y14 Receptor Antagonists Reverse Chronic Neuropathic Pain in a Mouse Model.		
2208	Design and Identification of a Novel, Functionally Subtype Selective GABAA Positive Allosteric Modulator (PF-06372865).		
2207	Cerebrospinal fluid beta-endorphin in models of hyperalgesia in the rat. 1995 , 59, 79-86		12
2206	The loss of antinociceptive efficacy of spinal morphine in rats with nerve ligation injury is prevented by reducing spinal afferent drive. 1995 , 199, 87-90		130
2205	Systemic and supraspinal, but not spinal, opiates suppress allodynia in a rat neuropathic pain model. 1995 , 199, 111-4		155
2204	Influence of selective alpha 2-adrenergic agents on mustard oil-induced central hyperalgesia in rats. 1995 , 281, 43-8		22
2203	POSTER COMMUNICATIONS. 1996 , 117, 130P-343P		6
2202	Antiallodynic effects of a CCKB antagonist in rats with nerve ligation injury: role of endogenous enkephalins. 1996 , 215, 161-4		32
2201	Effects of spinal cord stimulation on touch-evoked allodynia involve GABAergic mechanisms. An experimental study in the mononeuropathic rat. 1996 , 66, 287-95		141
2200	Characterization of time course of spinal amino acids, citrulline and PGE2 release after carrageenan/kaolin-induced knee joint inflammation: a chronic microdialysis study. 1996 , 67, 345-54		108
2199	Tactile allodynia and formalin hyperalgesia in streptozotocin-diabetic rats: effects of insulin, aldose reductase inhibition and lidocaine. 1996 , 68, 293-9		182
2198	Single intrathecal injections of dynorphin A or des-Tyr-dynorphins produce long-lasting allodynia in rats: blockade by MK-801 but not naloxone. 1996 , 68, 275-81		183
2197	Strain differences in adrenergic sensitivity of neuropathic pain behaviors in an experimental rat model. 1997 , 8, 3453-6		54
2196	A novel animal model to evaluate oxygen derived free radical damage in soft tissue. 1997 , 26, 363-72		36
2195	Regulation of expression of the sensory neuron-specific sodium channel SNS in inflammatory and neuropathic pain. 1997 , 10, 196-207		138
2194	Effect of subarachnoid gabapentin on tactile-evoked allodynia in a surgically induced neuropathic pain model in the rat. 1997 , 22, 249-56		139
2193	The effect of novel anti-epileptic drugs in rat experimental models of acute and chronic pain. 1997 , 324, 153-60		249

2192	Preserved acute pain and reduced neuropathic pain in mice lacking PKCgamma. 1997 , 278, 279-83	580
2191	Soft tissue repair capacity after oxygen-derived free radical-induced damage in one hindlimb of the rat. 1997 , 72, 60-9	18
2190	A metalloprotease-inhibitor reduces pain associated behavior in mice with experimental neuropathy. 1997 , 237, 45-8	70
2189	Adenosine receptor activation suppresses tactile hypersensitivity and potentiates spinal cord stimulation in mononeuropathic rats. 1997 , 223, 173-6	84
2188	The effects of mexiletine, desipramine and fluoxetine in rat models involving central sensitization. 1997 , 69, 161-9	94
2187	Electrophysiological characteristics of primary afferent fibers after systemic administration of anti-GD2 ganglioside antibody. 1997 , 69, 145-51	42
2186	Lumbar transplants of immortalized serotonergic neurons alleviate chronic neuropathic pain. 1997 , 72, 59-69	66
2185	Enhanced withdrawal responses to mechanical and thermal stimuli after bone injury. 1997 , 73, 325-337	16
2184	Supraspinal influence on hindlimb withdrawal thresholds and mustard oil-induced secondary allodynia in rats. 1997 , 42, 359-65	61
2183	Diminished inflammation and nociceptive pain with preservation of neuropathic pain in mice with a targeted mutation of the type I regulatory subunit of cAMP-dependent protein kinase. 1997 , 17, 7462-70	192
2182	Spinal pharmacology of tactile allodynia in diabetic rats. 1997 , 122, 1478-82	120
2181	ADVANCES IN PAIN RESEARCH. 1997 , 15, 229-234	1
2180	DIABETIC NEUROPATHY. 1997 , 15, 429-444	2
2179	Incidence of mononeuropathy in rats is influenced by pre-emptive alteration of spinal excitability. 1997 , 1, 53-9	12
2178	Primary afferent tachykinins are required to experience moderate to intense pain. 1998 , 392, 390-4	489
2177	Tactile hyperesthesia, altered epidermal innervation and plantar nerve injury in the hindfeet of rats housed on wire grates. 1998 , 788, 13-9	29
2176	ABT-594, a novel cholinergic channel modulator, is efficacious in nerve ligation and diabetic neuropathy models of neuropathic pain. 1998 , 801, 158-63	45
2175	Priming enhances endotoxin-induced thermal hyperalgesia and mechanical allodynia in rats. 1998 , 808, 13-22	25

2174	Systemic gabapentin and S(+)-3-isobutyl-gamma-aminobutyric acid block secondary hyperalgesia. 1998 , 810, 93-9		84
2173	Anti-allodynic actions of intravenous opioids in the nerve injured rat: potential utility of heroin and dihydroetorphine against neuropathic pain. 1998 , 357, 25-32		11
2172	An evaluation of the GABA uptake blocker tiagabine in animal models of neuropathic and nociceptive pain. 1998 , 44, 106-113		20
2171	Orphanin-FQ/nociceptin: lack of anti nociceptive, hyperalgesic or allodynic effects in acute thermal or mechanical tests following intracerebroventricular or intrathecal administration to mice or rats. 1998 , 2, 267-78		23
2170	Systemic morphine in the prevention of allodynia in the rat spinal nerve ligation model of neuropathic pain. 1998 , 2, 35-42		16
2169	Assessing mechanical allodynia in the rat paw with a new electronic algometer. <i>Journal of Neuroscience Methods</i> , 1998 , 84, 41-7	3	91
2168	A novel model of primary and secondary hyperalgesia after mild thermal injury in the rat. 1998 , 254, 25-8		58
2167	Weight bearing of the limb as a confounding factor in assessment of mechanical allodynia in the rat. 1998 , 74, 55-59		45
2166	The effect of thalidomide treatment on vascular pathology and hyperalgesia caused by chronic constriction injury of rat nerve. 1998 , 74, 83-91		145
2165	Excitotoxic spinal cord injury: behavioral and morphological characteristics of a central pain model. 1998 , 75, 141-155		193
2164	Partial sciatic nerve injury in the mouse as a model of neuropathic pain: behavioral and neuroanatomical correlates. 1998 , 76, 215-22		341
2163	Intrathecal administration of the mGluR compound, (S)-4CPG, attenuates hyperalgesia and allodynia associated with sciatic nerve constriction injury in rats. 1998 , 77, 59-66		60
2162	Hyperalgesia in experimental neuropathy is dependent on the TNF receptor 1. 1998 , 151, 138-42		230
2161	Broad-spectrum, non-opioid analgesic activity by selective modulation of neuronal nicotinic acetylcholine receptors. 1998 , 279, 77-81		325
2160	Central and Peripheral Analgesia Mediated by the Acetylcholinesterase-Inhibitor Neostigmine in the Rat Inflamed Knee Joint Model. 1998 , 86, 1027-1032		42
2159	The Anti-Allodynic Effects of Amitriptyline, Gabapentin, and Lidocaine in a Rat Model of Neuropathic Pain. 1998 , 87, 1360-1366		116
2158	The Anti-Allodynic Effects of Amitriptyline, Gabapentin, and Lidocaine in a Rat Model of Neuropathic Pain. 1998 , 87, 1360-1366		157
2157	The Effect of Electroconvulsive Treatment on Thermal Hyperalgesia and Mechanical Allodynia in a Rat Model of Peripheral Neuropathy. 1998 , 86, 584-587		6

2156	Gabapentin Reverses the Allodynia Produced by the Administration of Anti-GD2 Ganglioside, an Immunotherapeutic Drug. 1998 , 86, 111-116	27
2155	Central and peripheral analgesia mediated by the acetylcholinesterase-inhibitor neostigmine in the rat inflamed knee joint model. 1998 , 86, 1027-32	114
2154	In vivo antinociceptive activity of anti-rat mGluR1 and mGluR5 antibodies in rats. 1998 , 9, 731-5	80
2153	Gabapentin reverses the allodynia produced by the administration of anti-GD2 ganglioside, an immunotherapeutic drug. 1998 , 86, 111-6	40
2152	The effect of electroconvulsive treatment on thermal hyperalgesia and mechanical allodynia in a rat model of peripheral neuropathy. 1998 , 86, 584-7	9
2151	Hyperalgesia and allodynia induced by intrathecal (RS)-dihydroxyphenylglycine in rats. 1998 , 9, 1169-72	77
2150	Neuronal nitric oxide synthase mRNA upregulation in rat sensory neurons after spinal nerve ligation: lack of a role in allodynia development. 1999 , 19, 9201-8	79
2149	Mechanical and thermal hyperalgesia and ectopic neuronal discharge after chronic compression of dorsal root ganglia. 1999 , 82, 3347-58	238
2148	Receptor subtype mediating the adrenergic sensitivity of pain behavior and ectopic discharges in neuropathic Lewis rats. 1999 , 81, 2226-33	81
2147	Hyperalgesia caused by nerve transection: long-lasting block prevents early hyperalgesia in the receptive field of the surviving nerve. 1999 , 89, 1475-81	1
2146	Inflammatory Models of Pain and Hyperalgesia. 1999 , 40, 111-118	134
2145	Thermal and mechanical antinociceptive action of spinal vs peripherally administered clonidine in the rat inflamed knee joint model. 1999 , 83, 436-41	38
2144	Tierexperimentelle Untersuchungen bei neuropathischem Schmerz. 1999 , 13, 315-323	23
2143	Loss of antiallodynic and antinociceptive spinal/supraspinal morphine synergy in nerve-injured rats: restoration by MK-801 or dynorphin antiserum. 1999 , 831, 55-63	75
2142	A single intrathecal injection of GABA permanently reverses neuropathic pain after nerve injury. 1999 , 835, 334-9	77
2141	An adenosine kinase inhibitor attenuates tactile allodynia in a rat model of diabetic neuropathic pain. 1999 , 364, 141-6	32
2140	Preemptive intrathecal ketamine injection produces a long-lasting decrease in neuropathic pain behaviors in a rat model. 1999 , 24, 208-13	11
2139	Neutralizing antibodies to interleukin 1-receptor reduce pain associated behavior in mice with experimental neuropathy. 1999 , 270, 25-8	153

2138	Inhibition of spinal protein kinase C reduces nerve injury-induced tactile allodynia in neuropathic rats. 1999 , 276, 99-102	59
2137	Exogenous and endogenous adenosine enhance the spinal antiallodynic effects of morphine in a rat model of neuropathic pain. 1999 , 80, 31-6	74
2136	Heritability of nociception II. 'Types' of nociception revealed by genetic correlation analysis. 1999 , 80, 83-93	188
2135	Differential efficacy of intravenous lidocaine in alleviating ipsilateral versus contralateral neuropathic pain in the rat. 1999 , 80, 521-531	65
2134	Acute amitriptyline in a rat model of neuropathic pain: differential symptom and route effects. 1999 , 80, 643-653	125
2133	Spinal cannabinoids are anti-allodynic in rats with persistent inflammation. 1999 , 82, 199-205	127
2132	Inflammation-induced up-regulation of protein kinase Cgamma immunoreactivity in rat spinal cord correlates with enhanced nociceptive processing. 1999 , 88, 1267-74	111
2131	Sensory thresholds and the antinociceptive effects of GABA receptor agonists in mice lacking the beta3 subunit of the GABA(A) receptor. 2000 , 95, 795-806	55
2130	An improved method for assessing mechanical allodynia in the rat. 1999 , 67, 711-6	186
2129	Transplants of neuronal cells bioengineered to synthesize GABA alleviate chronic neuropathic pain. 1999 , 8, 87-101	91
2128	Intrathecal adenosine interacts with a spinal noradrenergic system to produce antinociception in nerve-injured rats. 1999 , 91, 1072-9	60
2127	Experimental Arthritis in the Rat Does Not Alter the Analgesic Potency of Intrathecal or Intraarticular Morphine. 1999 , 89, 403-408	4
2126	Experimental arthritis in the rat does not alter the analgesic potency of intrathecal or intraarticular morphine. 1999 , 89, 403-8	6
2125	Hyperalgesia Caused by Nerve Transection: Long-Lasting Block Prevents Early Hyperalgesia in the Receptive Field of the Surviving Nerve. 1999 , 89, 1475	18
2124	Mechanical allodynia in rats is blocked by a Ca ²⁺ permeable AMPA receptor antagonist. 1999 , 10, 3523-6	57
2123	Blockade of neuropathic pain by antisense targeting of tetrodotoxin-resistant sodium channels in sensory neurons. 2000 , 314, 201-13	26
2122	Initial characterization of the transplant of immortalized chromaffin cells for the attenuation of chronic neuropathic pain. 2000 , 9, 637-56	29
2121	The role of heme oxygenase in neuropathic and incisional pain. 2000 , 90, 677-82	30

2120	Antiallodynic effect of intrathecal gabapentin and its interaction with clonidine in a rat model of postoperative pain. 2000 , 92, 1126-31		86
2119	Injury to dorsal root ganglia alters innervation of spinal cord dorsal horn lamina involved in nociception. 2000 , 25, 537-42		34
2118	Effects of radolmidine, a novel alpha2 -adrenergic agonist compared with dexmedetomidine in different pain models in the rat. 2000 , 93, 473-81		50
2117	The effects of KRN5500, a spicamycin derivative, on neuropathic and nociceptive pain models in rats. 2000 , 91, 955-9		9
2116	An Analysis of Drug Interaction Between Morphine and Neostigmine in Rats with Nerve-Ligation Injury. 2000 , 90, 421		0
2115	An analysis of drug interaction between morphine and neostigmine in rats with nerve-ligation injury. 2000 , 90, 421-6		8
2114	Epineurial application of TNF elicits an acute mechanical hyperalgesia in the awake rat. 2000 , 5, 96-100		163
2113	An operant assay of thermal pain in conscious, unrestrained rats. <i>Journal of Neuroscience Methods</i> , 2000 , 97, 19-29	3	89
2112	Responses of rats to noxious mechanical stimulation of their tails during tail reperfusion following transient ischaemia. <i>Journal of Neuroscience Methods</i> , 2000 , 103, 173-80	3	11
2111	ABT-627, an endothelin ET(A) receptor-selective antagonist, attenuates tactile allodynia in a diabetic rat model of neuropathic pain. 2000 , 388, 29-35		82
2110	GM1 ganglioside restores abnormal responses to acute thermal and mechanical stimuli in aged rats. 2000 , 858, 380-5		16
2109	Thermal hyperalgesia and mechanical allodynia produced by intrathecal administration of the human immunodeficiency virus-1 (HIV-1) envelope glycoprotein, gp120. 2000 , 861, 105-16		220
2108	Low dose of tetrodotoxin reduces neuropathic pain behaviors in an animal model. 2000 , 871, 98-103		101
2107	Description of a short-term Taxol-induced nociceptive neuropathy in rats. 2000 , 887, 239-49		144
2106	Effect of nocistatin in pain modulation. 2000 , 45, 716-720		
2105	Key role for the epsilon isoform of protein kinase C in painful alcoholic neuropathy in the rat. 2000 , 20, 8614-9		110
2104	Anti-hyperalgesic effect of electroacupuncture in a model of post-incisional pain in rats. 2000 , 33, 957-60		14
2103	Mechanical allodynia caused by intraplantar injection of P2X receptor agonist in rats: involvement of heteromeric P2X2/3 receptor signaling in capsaicin-insensitive primary afferent neurons. 2000 , 20, RC90		156

2102	Antagonism of the melanocortin system reduces cold and mechanical allodynia in mononeuropathic rats. 2000 , 20, 8131-7	78
2101	Perfusion of the mechanically compressed lumbar ganglion with lidocaine reduces mechanical hyperalgesia and allodynia in the rat. 2000 , 84, 798-805	40
2100	Sympathetic nervous system: contribution to chronic pain. 2000 , 129, 451-468	50
2099	An analysis of changes in sensory thresholds to mild tactile and cold stimuli after experimental spinal cord injury in the rat. 2000 , 14, 287-300	77
2098	Altered pain responses in mice lacking alpha 1E subunit of the voltage-dependent Ca ²⁺ channel. 2000 , 97, 6132-7	204
2097	Extraterritorial neuropathic pain correlates with multisegmental elevation of spinal dynorphin in nerve-injured rats. 2000 , 86, 185-94	158
2096	Lumbar transplant of neurons genetically modified to secrete brain-derived neurotrophic factor attenuates allodynia and hyperalgesia after sciatic nerve constriction. 2000 , 86, 195-210	93
2095	Tactile allodynia in the absence of C-fiber activation: altered firing properties of DRG neurons following spinal nerve injury. 2000 , 85, 503-521	326
2094	Effects of purinergic and adrenergic antagonists in a rat model of painful peripheral neuropathy. 2000 , 87, 171-179	37
2093	Differential effects of spinal (R)-ketoprofen and (S)-ketoprofen against signs of neuropathic pain and tonic nociception: evidence for a novel mechanism of action of (R)-ketoprofen against tactile allodynia. 2000 , 87, 193-199	46
2092	Dorsal root section elicits signs of neuropathic pain rather than reversing them in rats with L5 spinal nerve injury. 2000 , 87, 213-219	62
2091	Possible role of inflammatory mediators in tactile hypersensitivity in rat models of mononeuropathy. 2000 , 88, 239-248	239
2090	Thalidomide treatment in chronic constrictive neuropathy decreases endoneurial tumor necrosis factor-alpha, increases interleukin-10 and has long-term effects on spinal cord dorsal horn met-enkephalin. 2000 , 88, 267-275	132
2089	Effects of intrathecal administration of ziconotide, a selective neuronal N-type calcium channel blocker, on mechanical allodynia and heat hyperalgesia in a rat model of postoperative pain. 2000 , 84, 151-8	82
2088	Spontaneous activity of axotomized afferent neurons after L5 spinal nerve injury in rats. 2000 , 84, 309-18	162
2087	Intrathecal lithium reduces neuropathic pain responses in a rat model of peripheral neuropathy. 2000 , 85, 59-64	23
2086	Spinal cyclooxygenase-2 is involved in development of allodynia after nerve injury in rats. 2000 , 97, 743-8	80
2085	Murine models of inflammatory, neuropathic and cancer pain each generates a unique set of neurochemical changes in the spinal cord and sensory neurons. 2000 , 98, 585-98	467

2084	Intrathecal S-nitroso-N-acetylpenicillamine and L-cysteine attenuate nerve injury-induced allodynia through noradrenergic activation in rats. 2000 , 101, 759-65	28
2083	Mediation of spinal nerve injury induced tactile allodynia by descending facilitatory pathways in the dorsolateral funiculus in rats. 2000 , 290, 129-32	76
2082	Assessment of allodynia and hyperalgesia after cisplatin administration to rats. 2000 , 291, 73-6	49
2081	Brainstem noradrenergic control of nociception is abnormal in the spontaneously hypertensive rat. 2000 , 291, 139-42	31
2080	Peripheral and spinal antihyperalgesic activity of SIB-1757, a metabotropic glutamate receptor (mGluR(5)) antagonist, in experimental neuropathic pain in rats. 2000 , 292, 115-8	97
2079	Enantioselective relief of neuropathic pain by systemic mexiletine in the rat. 2000 , 1, 128-137	5
2078	Early antinociception delays edema but does not reduce the magnitude of persistent pain in the formalin test. 2000 , 1, 218-28	14
2077	Allodynia and hyperalgesia produced by specific inhibition of spinal c-fos expression: lack of correlation with dynorphin content. 2001 , 2, 241-9	4
2076	The effect of lumbar sympathectomy in the spinal nerve ligation model of neuropathic pain. 2001 , 2, 270-8	19
2075	Systemic administration of CNI-1493, a p38 mitogen-activated protein kinase inhibitor, blocks intrathecal human immunodeficiency virus-1 gp120-induced enhanced pain states in rats. 2001 , 2, 326-33	36
2074	Nerve injury-induced tactile allodynia is mediated via ascending spinal dorsal column projections. 2001 , 90, 105-11	100
2073	Antiallodynic effect of intrathecally administered 5-HT(2) agonists in rats with nerve ligation. 2001 , 90, 173-9	71
2072	Antihyperalgesic effects of the muscarinic receptor ligand vedaclidine in models involving central sensitization in rats. 2001 , 93, 221-227	9
2071	Differential effects of NMDA and group I mGluR antagonists on both nociception and spinal cord protein kinase C translocation in the formalin test and a model of neuropathic pain in rats. 2001 , 94, 17-29	83
2070	Non-noxious A fiber afferent input enhances capsaicin-induced mechanical hyperalgesia in the rat. 2001 , 94, 169-175	20
2069	A new model of sciatic inflammatory neuritis (SIN): induction of unilateral and bilateral mechanical allodynia following acute unilateral peri-sciatic immune activation in rats. 2001 , 94, 231-244	243
2068	P75-expressing elements are necessary for anti-allodynic effects of spinal clonidine and neostigmine. 2001 , 102, 681-6	8
2067	Cytotoxic targeting of isolectin IB4-binding sensory neurons. 2001 , 108, 143-55	110

2066	Role of protein kinase Cepsilon and protein kinase A in a model of paclitaxel-induced painful peripheral neuropathy in the rat. 2001 , 108, 507-15	136
2065	The effect of lumbar sympathectomy on increased tactile sensitivity in spinal nerve ligated rats. 2001 , 298, 99-102	15
2064	Changes in brain content of nociceptin/orphanin FQ and endomorphin 2 in a rat model of neuropathic pain. 2001 , 311, 13-6	27
2063	A murine model of opioid-induced hyperalgesia. 2001 , 86, 56-62	118
2062	The paired homeodomain protein DRG11 is required for the projection of cutaneous sensory afferent fibers to the dorsal spinal cord. 2001 , 31, 59-73	119
2061	Intrathecal HIV-1 envelope glycoprotein gp120 induces enhanced pain states mediated by spinal cord proinflammatory cytokines. 2001 , 21, 2808-19	275
2060	Chemokines and glycoprotein120 produce pain hypersensitivity by directly exciting primary nociceptive neurons. 2001 , 21, 5027-35	411
2059	Upregulation of dorsal root ganglion (alpha)2(delta) calcium channel subunit and its correlation with allodynia in spinal nerve-injured rats. 2001 , 21, 1868-75	491
2058	Pronociceptive actions of dynorphin maintain chronic neuropathic pain. 2001 , 21, 1779-86	244
2057	Hypoalgesia and hyperalgesia with inherited hypertension in the rat. 2001 , 280, R345-54	30
2056	Inhibition of neuropathic pain by selective ablation of brainstem medullary cells expressing the mu-opioid receptor. 2001 , 21, 5281-8	223
2055	The effects of peripheral administration of a novel selective antagonist for prostaglandin E receptor subtype EP(1), ONO-8711, in a rat model of postoperative pain. 2001 , 92, 233-8	42
2054	The interaction between intrathecal neostigmine and GABA receptor agonists in rats with nerve ligation injury. 2001 , 93, 1297-303	19
2053	Stereospecific effect of pregabalin on ectopic afferent discharges and neuropathic pain induced by sciatic nerve ligation in rats. 2001 , 95, 1473-9	64
2052	Role of endogenous ATP at the incision area in a rat model of postoperative pain. 2001 , 12, 1701-4	38
2051	Allosteric adenosine modulation to reduce allodynia. 2001 , 95, 416-20	31
2050	Pain models display differential sensitivity to Ca ²⁺ -permeable non-NMDA glutamate receptor antagonists. 2001 , 95, 965-73	55
2049	Chronic blockade of melanocortin receptors alleviates allodynia in rats with neuropathic pain. 2001 , 93, 1572-7, table of contents	35

2048	[Animal models and peripheral nociception tests for the study of neuropathic pain]. 2001 , 118, 89-95	3
2047	Tonic descending facilitation from the rostral ventromedial medulla mediates opioid-induced abnormal pain and antinociceptive tolerance. 2001 , 21, 279-86	275
2046	A rat model of postthoracotomy pain: behavioural and spinal cord NK-1 receptor assessment. 2001 , 48, 665-76	23
2045	Assessment of neurotoxicity following repeated cremophor/ethanol injections in rats. 2001 , 3, 301-6	26
2044	Model of neuropathic intermittent claudication in the rat: methodology and application. <i>Journal of Neuroscience Methods</i> , 2001 , 104, 191-8	3 35
2043	Sciatic inflammatory neuritis (SIN): behavioral allodynia is paralleled by peri-sciatic proinflammatory cytokine and superoxide production. 2001 , 6, 111-29	62
2042	Sex hormones regulate the contribution of PKCepsilon and PKA signalling in inflammatory pain in the rat. 2001 , 13, 2227-33	89
2041	Suppression of inflammatory and neuropathic pain symptoms in mice lacking the N-type Ca ²⁺ channel. 2001 , 20, 2349-56	259
2040	Genetic enhancement of inflammatory pain by forebrain NR2B overexpression. 2001 , 4, 164-9	271
2039	Knockdown of spinal metabotropic glutamate receptor 1 (mGluR(1)) alleviates pain and restores opioid efficacy after nerve injury in rats. 2001 , 132, 354-67	96
2038	Contribution of alpha(2) receptor subtypes to nerve injury-induced pain and its regulation by dexmedetomidine. 2001 , 132, 1827-36	73
2037	Spinal N-acetyl-alpha-linked acidic dipeptidase (NAALADase) inhibition attenuates mechanical allodynia induced by paw carrageenan injection in the rat. 2001 , 909, 138-44	57
2036	Chronic administration of amitriptyline and caffeine in a rat model of neuropathic pain: multiple interactions. 2001 , 430, 211-8	54
2035	Opioid-induced hyperalgesia and incisional pain. 2001 , 93, 204-9	99
2034	Injury type-specific calcium channel alpha 2 delta-1 subunit up-regulation in rat neuropathic pain models correlates with antiallodynic effects of gabapentin. 2002 , 303, 1199-205	321
2033	Effect of 2-(phosphono-methyl)-pentanedioic acid on allodynia and afferent ectopic discharges in a rat model of neuropathic pain. 2002 , 300, 662-7	48
2032	Spinal and peripheral mu opioids and the development of secondary tactile allodynia after thermal injury. 2002 , 94, 968-74, table of contents	24
2031	Transgenic over-expression of galanin in injured primary sensory neurons. 2002 , 13, 2129-32	23

2030	Hyperalgesia During Opioid Abstinence: Mediation by Glutamate and Substance P. 2002 , 95, 979-984	22
2029	Changes in expression of voltage-dependent ion channel subunits in dorsal root ganglia of rats with radicular injury and pain. 2002 , 27, 1517-24; discussion 1525	45
2028	Central nociceptive role of prostacyclin (IP) receptor induced by peripheral inflammation. 2002 , 13, 93-6	42
2027	Postdelivery of alfentanil and ketamine has no effect on intradermal capsaicin-induced pain and hyperalgesia. 2002 , 18, 373-9	19
2026	Hyperalgesia during opioid abstinence: mediation by glutamate and substance p. 2002 , 95, 979-84, table of contents	41
2025	Pharmacology of spinal glutamatergic receptors in post-thermal injury-evoked tactile allodynia and thermal hyperalgesia. 2002 , 96, 617-26	42
2024	Time-dependent descending facilitation from the rostral ventromedial medulla maintains, but does not initiate, neuropathic pain. 2002 , 22, 5129-36	280
2023	Concentration-effect relationships for intravenous alfentanil and ketamine infusions in human volunteers: effects on acute thresholds and capsaicin-evoked hyperpathia. 2002 , 42, 70-80	49
2022	Immortalized chromaffin cells disimmortalized with Cre/lox site-directed recombination for use in cell therapy for pain after partial nerve injury. 2002 , 175, 49-60	15
2021	Effect of lumbar 5 ventral root transection on pain behaviors: a novel rat model for neuropathic pain without axotomy of primary sensory neurons. 2002 , 175, 23-34	78
2020	Behavioural assessment of dimethylsulfoxide neurotoxicity in rats. 2002 , 132, 117-21	29
2019	Effects of ablation of N- and R-type Ca(2+) channels on pain transmission. 2002 , 43, 1-7	78
2018	Inhibition of neuropathic pain by decreased expression of the tetrodotoxin-resistant sodium channel, NaV1.8. 2002 , 95, 143-52	319
2017	Therapeutic administration of nitric oxide synthase inhibitors reverses hyperalgesia but not inflammation in a rat model of polyarthritis. 2002 , 95, 215-223	18
2016	Attenuation of hyperalgesia in a rat model of neuropathic pain after intrathecal pre- or post-treatment with a neurokinin-1 antagonist. 2002 , 95, 277-285	69
2015	Mechanical hyperalgesia after an L5 ventral rhizotomy or an L5 ganglionectomy in the rat. 2002 , 96, 63-72	74
2014	Analgesic and anti-inflammatory effects of A-286501, a novel orally active adenosine kinase inhibitor. 2002 , 96, 107-18	43
2013	Lamotrigine in spinal cord injury pain: a randomized controlled trial. 2002 , 96, 375-383	188

2012	Heritability of nociception. III. Genetic relationships among commonly used assays of nociception and hypersensitivity. 2002 , 97, 75-86	155
2011	Spinal noradrenergic activation mediates allodynia reduction from an allosteric adenosine modulator in a rat model of neuropathic pain. 2002 , 97, 117-25	45
2010	A mouse model of neuropathic cancer pain. 2002 , 99, 167-74	69
2009	Efficacy of systemic morphine suggests a fundamental difference in the mechanisms that generate bone cancer vs inflammatory pain. 2002 , 99, 397-406	161
2008	Reduced basal release of serotonin from the ventrobasal thalamus of the rat in a model of neuropathic pain. 2002 , 99, 359-66	28
2007	Progressive tactile hypersensitivity after a peripheral nerve crush: non-noxious mechanical stimulus-induced neuropathic pain. 2002 , 100, 155-62	51
2006	'Knock-down' of spinal CB1 receptors produces abnormal pain and elevates spinal dynorphin content in mice. 2002 , 100, 203-9	31
2005	Differential screening-selected gene aberrative in neuroblastoma protein modulates inflammatory pain in the spinal dorsal horn. 2002 , 110, 579-86	6
2004	Different peripheral mechanisms mediate enhanced nociception in metabolic/toxic and traumatic painful peripheral neuropathies in the rat. 2002 , 111, 389-97	84
2003	Enhanced release of adenosine in rat hind paw following spinal nerve ligation: involvement of capsaicin-sensitive sensory afferents. 2002 , 114, 379-87	10
2002	Chronic neuropathic pain is accompanied by global changes in gene expression and shares pathobiology with neurodegenerative diseases. 2002 , 114, 529-46	216
2001	Role of primary afferent nerves in allodynia caused by diabetic neuropathy in rats. 2002 , 114, 291-9	190
2000	Neuropathic pain is associated with alterations of nitric oxide synthase immunoreactivity and catalytic activity in dorsal root ganglia and spinal dorsal horn. 2002 , 58, 161-71	66
1999	Involvement of adenosine in the anti-allodynic effect of amitriptyline in streptozotocin-induced diabetic rats. 2002 , 328, 129-32	77
1998	Altered synaptophysin expression in the rat spinal cord after chronic constriction injury of sciatic nerve. 2002 , 333, 155-8	24
1997	A water-soluble synthetic spicamycin derivative (San-Gly) decreases mechanical allodynia in a rodent model of neuropathic pain. 2002 , 330, 37-40	7
1996	Potential mechanisms of neuropathic pain in diabetes. 2002 , 50, 205-28	122
1995	Sustained morphine exposure induces a spinal dynorphin-dependent enhancement of excitatory transmitter release from primary afferent fibers. 2002 , 22, 6747-55	207

1994	Phosphorylation of extracellular signal-regulated kinase in primary afferent neurons by noxious stimuli and its involvement in peripheral sensitization. 2002 , 22, 7737-45	281
1993	Degeneration of myelinated efferent fibers induces spontaneous activity in uninjured C-fiber afferents. 2002 , 22, 7746-53	192
1992	Presurgical ketoprofen, but not morphine, dipyron, diclofenac or tenoxicam, preempts post-incisional mechanical allodynia in rats. 2002 , 35, 111-9	19
1991	Antinociceptive potency of aminoglycoside antibiotics and magnesium chloride: a comparative study on models of phasic and incisional pain in rats. 2002 , 35, 395-403	10
1990	The 5-HT ₃ subtype of serotonin receptor contributes to nociceptive processing via a novel subset of myelinated and unmyelinated nociceptors. 2002 , 22, 1010-9	303
1989	Selective mediation of nerve injury-induced tactile hypersensitivity by neuropeptide Y. 2002 , 22, 9858-67	87
1988	Dorsal horn-enriched genes identified by DNA microarray, in situ hybridization and immunohistochemistry. 2002 , 3, 11	26
1987	Dual effect of local application of nitric oxide donors in a model of incision pain in rats. 2002 , 441, 57-65	49
1986	Actions of intrathecal omega-conotoxins CVID, GVIA, MVIIA, and morphine in acute and neuropathic pain in the rat. 2002 , 451, 279-86	132
1985	Antisense oligonucleotide knockdown of mGluR1 alleviates hyperalgesia and allodynia associated with chronic inflammation. 2002 , 73, 401-10	39
1984	Antinociceptive effects following intrathecal pretreatment with selective metabotropic glutamate receptor compounds in a rat model of neuropathic pain. 2002 , 73, 411-8	91
1983	Group II mGluR receptor agonists are effective in persistent and neuropathic pain models in rats. 2002 , 73, 419-27	102
1982	Substance P release in the spinal dorsal horn following peripheral nerve injury. 2002 , 36, 252-6	17
1981	Antibody directed against GD(2) produces mechanical allodynia, but not thermal hyperalgesia when administered systemically or intrathecally despite its dependence on capsaicin sensitive afferents. 2002 , 930, 67-74	21
1980	Possible involvement of a muscarinic receptor in the anti-allodynic action of a 5-HT ₂ receptor agonist in rats with nerve ligation injury. 2002 , 932, 124-8	26
1979	Role for both spinal cord COX-1 and COX-2 in maintenance of mechanical hypersensitivity following peripheral nerve injury. 2002 , 937, 94-9	79
1978	Combined systemic administration of morphine and magnesium sulfate attenuates pain-related behavior in mononeuropathic rats. 2002 , 943, 101-4	24
1977	Changes of sodium channel expression in experimental painful diabetic neuropathy. 2002 , 52, 786-92	159

1976	Impaired sensory function in heterozygous P0 knockout mice is associated with nodal changes in sensory nerves. 2002 , 67, 167-73	21
1975	The effect of combined systemic administration of morphine and L-NAME, a nitric oxide synthase inhibitor, on behavioral signs of neuropathic pain in rats. 2002 , 30, 143-153	13
1974	Intrathecal carbachol and clonidine produce a synergistic antiallodynic effect in rats with a nerve ligation injury. 2002 , 49, 178-84	4
1973	Localization of N-type Ca ²⁺ channels in the rat spinal cord following chronic constrictive nerve injury. 2002 , 147, 456-63	112
1972	Alteration of dorsal root ganglion P2X3 receptor expression and function following spinal nerve ligation in the rat. 2002 , 147, 511-9	77
1971	Morphological and pharmacological evidence for the role of peripheral prostaglandins in the pathogenesis of neuropathic pain. 2002 , 15, 1037-47	101
1970	Amelioration of chronic neuropathic pain after partial nerve injury by adeno-associated viral (AAV) vector-mediated over-expression of BDNF in the rat spinal cord. 2002 , 9, 1387-95	118
1969	ATP induced three types of pain behaviors, including allodynia. 2003 , 59, 56-63	18
1968	Upregulation of FGF-2 in reactive spinal cord astrocytes following unilateral lumbar spinal nerve ligation. 2003 , 148, 366-76	47
1967	The role of voltage-gated sodium channels in neuropathic pain. 2003 , 13, 291-7	139
1966	Activation of spinal orexin-1 receptor produces anti-allodynic effect in the rat carrageenan test. 2003 , 481, 175-80	32
1965	Enhanced thermal antinociceptive potency and anti-allodynic effects of morphine following spinal administration of endotoxin. 2003 , 960, 209-18	17
1964	Involvement of NMDA-nitric oxide pathways in the development of tactile hypersensitivity evoked by the loose-ligation of inferior alveolar nerves in rats. 2003 , 963, 232-43	31
1963	Interactions of 5-HT ₂ receptor agonists with acetylcholine in spinal analgesic mechanisms in rats with neuropathic pain. 2003 , 965, 114-20	25
1962	Chronic constriction injury of sciatic nerve induces the up-regulation of descending inhibitory noradrenergic innervation to the lumbar dorsal horn of mice. 2003 , 970, 110-8	89
1961	Cannabinoid CB1 receptor inhibition of mechanically evoked responses of spinal neurones in control rats, but not in rats with hindpaw inflammation. 2003 , 474, 209-16	15
1960	Acetic acid conditioning stimulus induces long-lasting antinociception of somatic inflammatory pain. 2003 , 74, 841-9	21
1959	Animal and cellular models of chronic pain. 2003 , 55, 949-65	156

1958	The role of uninjured nerve in spinal nerve ligated rats points to an improved animal model of neuropathic pain. 2003 , 7, 473-9	22
1957	Coupling gene chip analyses and rat genetic variances in identifying potential target genes that may contribute to neuropathic allodynia development. 2003 , 87, 560-73	95
1956	Activation of peripheral cannabinoid CB1 receptors inhibits mechanically evoked responses of spinal neurons in noninflamed rats and rats with hindpaw inflammation. 2003 , 18, 2239-43	53
1955	Internal and external factors affecting the development of neuropathic pain in rodents. Is it all about pain?. 2003 , 3, 326-42	15
1954	Multiple actions of systemic artemin in experimental neuropathy. 2003 , 9, 1383-9	129
1953	Neurotrophic factors as novel therapeutics for neuropathic pain. 2003 , 2, 460-72	114
1952	Characterization of cyclophosphamide cystitis, a model of visceral and referred pain, in the mouse: species and strain differences. 2003 , 170, 1008-12	92
1951	Models of neuropathic pain in the rat. 2003 , Chapter 5, Unit5.32	29
1950	Heme oxygenase type 2 participates in the development of chronic inflammatory and neuropathic pain. 2003 , 4, 101-7	26
1949	Spared nerve injury model of neuropathic pain in the mouse: a behavioral and anatomic analysis. 2003 , 4, 465-70	203
1948	Mechanical sensation and pain thresholds in patients with chronic arthropathies. 2003 , 4, 203-11	93
1947	Peripheral norepinephrine exacerbates neuritis-induced hyperalgesia. 2003 , 4, 212-21	34
1946	Paw withdrawal thresholds and persistent hindlimb flexion in experimental mononeuropathies. 2003 , 4, 222-30	10
1945	Gene expression profiling of melanocortin system in neuropathic rats supports a role in nociception. 2003 , 118, 111-8	30
1944	Further characterization of high mobility group box 1 (HMGB1) as a proinflammatory cytokine: central nervous system effects. 2003 , 24, 254-65	110
1943	An animal model of nociceptive peripheral neuropathy following repeated cisplatin injections. 2003 , 182, 12-20	114
1942	Intrathecal nerve growth factor restores opioid effectiveness in an animal model of neuropathic pain. 2003 , 45, 543-52	44
1941	Intraplantar injection of anandamide inhibits mechanically-evoked responses of spinal neurones via activation of CB2 receptors in anaesthetised rats. 2003 , 45, 404-11	59

1940	Attenuation of mechanical hyperalgesia following spinal cord injury by administration of antibodies to nerve growth factor in the rat. 2003 , 336, 117-20	69
1939	Systemic agmatine attenuates tactile allodynia in two experimental neuropathic pain models in rats. 2003 , 339, 88-90	34
1938	Anti-mechanical allodynic effect of intrathecal and intracerebroventricular injection of orexin-A in the rat neuropathic pain model. 2003 , 347, 183-6	59
1937	Effect of systemic FR140423, a new analgesic compound, in a rat model of postoperative pain: contribution of delta-opioid receptors. 2003 , 350, 29-32	4
1936	Discovery and in vivo evaluation of new melanocortin-4 receptor-selective peptides. 2003 , 24, 271-80	19
1935	Anti-allodynic effect of NW-1029, a novel Na(+) channel blocker, in experimental animal models of inflammatory and neuropathic pain. 2003 , 102, 17-25	51
1934	Powerful antinociceptive effects of the cone snail venom-derived subtype-selective NMDA receptor antagonists conantokins G and T. 2003 , 101, 109-16	108
1933	Spinal nerve injury activates prostaglandin synthesis in the spinal cord that contributes to early maintenance of tactile allodynia. 2003 , 101, 139-47	31
1932	Transgene-mediated enkephalin release enhances the effect of morphine and evades tolerance to produce a sustained antiallodynic effect in neuropathic pain. 2003 , 102, 135-42	102
1931	Multiple phases of relief from experimental mechanical allodynia by systemic lidocaine: responses to early and late infusions. 2003 , 103, 21-9	49
1930	'CatWalk' automated quantitative gait analysis as a novel method to assess mechanical allodynia in the rat; a comparison with von Frey testing. 2003 , 102, 203-9	198
1929	Involvement of the anterior pretectal nucleus in the control of persistent pain: a behavioral and c-Fos expression study in the rat. 2003 , 103, 163-74	27
1928	Peripheral and central sites of action for A-85380 in the spinal nerve ligation model of neuropathic pain. 2003 , 103, 269-276	35
1927	Cyclooxygenase-1 in the spinal cord plays an important role in postoperative pain. 2003 , 104, 15-23	85
1926	A substance P receptor (NK1) antagonist can reverse vascular and nociceptive abnormalities in a rat model of complex regional pain syndrome type II. 2003 , 104, 75-84	65
1925	The role of calcitonin gene-related peptide (CGRP) in the generation and maintenance of mechanical allodynia and hyperalgesia in rats after intradermal injection of capsaicin. 2003 , 104, 201-8	91
1924	Intramuscular injection of tumor necrosis factor-alpha induces muscle hyperalgesia in rats. 2003 , 104, 579-588	123
1923	Reversal of experimental neuropathic pain by T-type calcium channel blockers. 2003 , 105, 159-68	161

1922	5-hydroxytryptamine1A receptor is involved in the bee venom induced inflammatory pain. 2003 , 106, 135-42	18
1921	A peripheral cannabinoid mechanism suppresses spinal fos protein expression and pain behavior in a rat model of inflammation. 2003 , 117, 659-70	91
1920	Glial cell line-derived neurotrophic factor contributes to delayed inflammatory hyperalgesia in adjuvant rat pain model. 2003 , 117, 503-12	47
1919	Prostaglandin E2 inhibits the potassium current in sensory neurons from hyperalgesic Kv1.1 knockout mice. 2003 , 119, 65-72	19
1918	Selective activation of cannabinoid CB(2) receptors suppresses spinal fos protein expression and pain behavior in a rat model of inflammation. 2003 , 119, 747-57	143
1917	Sexual dimorphism for protein kinase c epsilon signaling in a rat model of vincristine-induced painful peripheral neuropathy. 2003 , 119, 831-8	52
1916	Pro-nociceptive effects of neuromedin U in rat. 2003 , 120, 467-74	48
1915	Expression of auxiliary beta subunits of sodium channels in primary afferent neurons and the effect of nerve injury. 2003 , 121, 441-50	53
1914	Heme oxygenase type 2 modulates behavioral and molecular changes during chronic exposure to morphine. 2003 , 121, 999-1005	43
1913	Cyclooxygenase 2 in infiltrating inflammatory cells in injured nerve is universally up-regulated following various types of peripheral nerve injury. 2003 , 121, 691-704	81
1912	A new animal model of vincristine-induced nociceptive peripheral neuropathy. 2003 , 24, 797-805	86
1911	Impaired neuropathic pain responses in mice lacking the chemokine receptor CCR2. 2003 , 100, 7947-52	488
1910	Activation of CB2 cannabinoid receptors by AM1241 inhibits experimental neuropathic pain: pain inhibition by receptors not present in the CNS. 2003 , 100, 10529-33	415
1909	Transgenic overexpression of galanin in the dorsal root ganglia modulates pain-related behavior. 2003 , 100, 6180-5	64
1908	Sensory function above lesion level in spinal cord injury patients with and without pain. 2003 , 20, 71-6	37
1907	HSV-mediated gene transfer of the glial cell-derived neurotrophic factor provides an antiallodynic effect on neuropathic pain. 2003 , 8, 367-75	87
1906	Antiallodynic effects of intrathecal orexins in a rat model of postoperative pain. 2003 , 307, 1065-71	68
1905	Models of neuropathic pain in the rat. 2003 , Chapter 9, Unit 9.14	32

1904	Function of gamma-aminobutyric acid receptor/channel rho 1 subunits in spinal cord. 2003 , 278, 48321-9	35
1903	Sensory function in spinal cord injury patients with and without central pain. 2003 , 126, 57-70	172
1902	Intrathecal clonidine reduces hypersensitivity after nerve injury by a mechanism involving spinal m4 muscarinic receptors. 2003 , 96, 1403-1408	17
1901	Severity and duration of hyperalgesia in rat varies with type of nerve lesion. 2003 , 53, 1200-8; discussion 1208-9	12
1900	Cyclooxygenase-1 in the spinal cord is altered after peripheral nerve injury. 2003 , 99, 1175-9	57
1899	Effects of morphine on mechanical allodynia in a rat model of central neuropathic pain. 2003 , 14, 1017-1020	15
1898	Dexmedetomidine Fails to Cause Hyperalgesia After Cessation of Chronic Administration. 2003 , 96, 195-200	10
1897	Dexmedetomidine fails to cause hyperalgesia after cessation of chronic administration. 2003 , 96, 195-200, table of contents	24
1896	Magnesium chloride and ruthenium red attenuate the antiallodynic effect of intrathecal gabapentin in a rat model of postoperative pain. 2003 , 98, 1472-9	20
1895	A single intravenous injection of KRN5500 (antibiotic spicamycin) produces long-term decreases in multiple sensory hypersensitivities in neuropathic pain. 2003 , 97, 174-82, table of contents	9
1894	Effects of morphine on mechanical allodynia in a rat model of central neuropathic pain. 2003 , 14, 1017-20	24
1893	Redistribution of Na(V)1.8 in uninjured axons enables neuropathic pain. 2003 , 23, 158-66	312
1892	Neuronal hyperpolarization-activated pacemaker channels drive neuropathic pain. 2003 , 23, 1169-78	267
1891	Enhanced evoked excitatory transmitter release in experimental neuropathy requires descending facilitation. 2003 , 23, 8370-9	118
1890	Absence of thermal hyperalgesia in serotonin transporter-deficient mice. 2003 , 23, 708-15	103
1889	Resiniferatoxin induces paradoxical changes in thermal and mechanical sensitivities in rats: mechanism of action. 2003 , 23, 2911-9	120
1888	Spinal glia and proinflammatory cytokines mediate mirror-image neuropathic pain in rats. 2003 , 23, 1026-40	571
1887	Increased sensitivity of injured and adjacent uninjured rat primary sensory neurons to exogenous tumor necrosis factor-alpha after spinal nerve ligation. 2003 , 23, 3028-38	251

1886	Tumor necrosis factor-alpha induces mechanical allodynia after spinal nerve ligation by activation of p38 MAPK in primary sensory neurons. 2003 , 23, 2517-21	447
1885	A conditional deletion of the NR1 subunit of the NMDA receptor in adult spinal cord dorsal horn reduces NMDA currents and injury-induced pain. 2003 , 23, 5031-40	154
1884	Primary cultures of neonatal rat spinal cord. 2004 , 99, 203-13	11
1883	Isolation and culture of sensory neurons from the dorsal-root ganglia of embryonic or adult rats. 2004 , 99, 189-202	32
1882	Calcium-permeable alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid/kainate receptors mediate development, but not maintenance, of secondary allodynia evoked by first-degree burn in the rat. 2004 , 310, 223-9	38
1881	Calcitonin gene-related peptide receptor activation produces PKA- and PKC-dependent mechanical hyperalgesia and central sensitization. 2004 , 92, 2859-66	134
1880	Pituitary adenylate cyclase-activating polypeptide is required for the development of spinal sensitization and induction of neuropathic pain. 2004 , 24, 7283-91	87
1879	Altered sodium channel expression in second-order spinal sensory neurons contributes to pain after peripheral nerve injury. 2004 , 24, 4832-9	214
1878	Proteinase-activated receptor 2-mediated potentiation of transient receptor potential vanilloid subfamily 1 activity reveals a mechanism for proteinase-induced inflammatory pain. 2004 , 24, 4293-9	256
1877	A role for proinflammatory cytokines and fractalkine in analgesia, tolerance, and subsequent pain facilitation induced by chronic intrathecal morphine. 2004 , 24, 7353-65	346
1876	Sensory neuron-specific receptor activation elicits central and peripheral nociceptive effects in rats. 2004 , 101, 7175-80	92
1875	Three exercise paradigms differentially improve sensory recovery after spinal cord contusion in rats. 2004 , 127, 1403-14	240
1874	The candidate neuroprotective agent artemin induces autonomic neural dysplasia without preventing peripheral nerve dysfunction. 2004 , 32, 275-94	33
1873	Constitutive spinal cyclooxygenase-2 participates in the initiation of tissue injury-induced hyperalgesia. 2004 , 24, 2727-32	83
1872	Semaphorin3A inhibits nerve growth factor-induced sprouting of nociceptive afferents in adult rat spinal cord. 2004 , 24, 819-27	99
1871	Spinal dorsal horn calcium channel alpha2delta-1 subunit upregulation contributes to peripheral nerve injury-induced tactile allodynia. 2004 , 24, 8494-9	191
1870	Selective activation of cannabinoid CB2 receptors suppresses hyperalgesia evoked by intradermal capsaicin. 2004 , 308, 446-53	108
1869	Does Gabapentin Act as an Agonist at Native GABAB Receptors?. 2004 , 11, 346-355	0

1868	Integrin signaling in inflammatory and neuropathic pain in the rat. 2004 , 19, 634-42	89
1867	TRPV1 and CB(1) receptor-mediated effects of the endovanilloid/endocannabinoid N-arachidonoyl-dopamine on primary afferent fibre and spinal cord neuronal responses in the rat. 2004 , 20, 175-84	58
1866	Antinociceptive effects of N-acetylaspartylglutamate (NAAG) peptidase inhibitors ZJ-11, ZJ-17 and ZJ-43 in the rat formalin test and in the rat neuropathic pain model. 2004 , 20, 483-94	82
1865	Deletion of the CCK2 receptor gene reduces mechanical sensitivity and abolishes the development of hyperalgesia in mononeuropathic mice. 2004 , 20, 1577-86	25
1864	Central sprouting of uninjured small fiber afferents in the adult rat spinal cord following spinal nerve ligation. 2004 , 20, 1705-12	21
1863	Cannabinoid CB2 receptor activation inhibits mechanically evoked responses of wide dynamic range dorsal horn neurons in naïve rats and in rat models of inflammatory and neuropathic pain. 2004 , 20, 2311-20	138
1862	Evidence that exogenous and endogenous fractalkine can induce spinal nociceptive facilitation in rats. 2004 , 20, 2294-302	229
1861	Antinociception induced by stimulating the anterior pretectal nucleus in two models of pain in rats. 2004 , 31, 608-13	18
1860	Spinal NF-κB activation induces COX-2 upregulation and contributes to inflammatory pain hypersensitivity. 2004 , 19, 3375-81	185
1859	Diabetes-induced expression of activating transcription factor 3 in mouse primary sensory neurons. 2004 , 9, 242-54	29
1858	Reduction by gabapentin of K ⁺ -evoked release of [3H]-glutamate from the caudal trigeminal nucleus of the streptozotocin-treated rat. 2004 , 141, 574-9	31
1857	Fractalkine (CX3CL1) and fractalkine receptor (CX3CR1) distribution in spinal cord and dorsal root ganglia under basal and neuropathic pain conditions. 2004 , 20, 1150-60	307
1856	Nefopam potentiates morphine antinociception in allodynia and hyperalgesia in the rat. 2004 , 77, 695-703	20
1855	Morphine and endomorphin-1 differently influence pronociceptin/orphanin FQ system in neuropathic rats. 2004 , 78, 171-8	25
1854	Effect of K ⁺ channel modulators on the antiallodynic effect of gabapentin. 2004 , 484, 201-8	35
1853	Decrease in serotonin concentration in raphe magnus nucleus and attenuation of morphine analgesia in two mice models of neuropathic pain. 2004 , 484, 217-23	38
1852	Riboflavin reduces hyperalgesia and inflammation but not tactile allodynia in the rat. 2004 , 492, 35-40	19
1851	Assessing the role of metabotropic glutamate receptor 5 in multiple nociceptive modalities. 2004 , 506, 107-18	88

1850	Effect of a graded single constriction of the rat sciatic nerve on pain behavior and expression of immunoreactive NPY and NPY Y1 receptor in DRG neurons and spinal cord. 2004 , 1006, 87-99	42
1849	Role of prostaglandin receptor subtype EP1 in prostaglandin E2-induced nociceptive transmission in the rat spinal dorsal horn. 2004 , 1010, 62-8	34
1848	Antinociceptive effects of bethanechol or dimethylphenylpiperazinium in models of phasic or incisional pain in rats. 2004 , 1018, 272-82	12
1847	Sciatic nerve regeneration in mice and rats: recovery of sensory innervation is followed by a slowly retreating neuropathic pain-like syndrome. 2004 , 1027, 67-72	95
1846	The behavioral and neuroanatomical effects of IB4-saporin treatment in rat models of nociceptive and neuropathic pain. 2004 , 1029, 65-76	52
1845	Does gabapentin act as an agonist at native GABA(B) receptors?. 2004 , 11, 346-55	22
1844	Prevention of sensory disorders in diabetic Sprague-Dawley rats by aldose reductase inhibition or treatment with ciliary neurotrophic factor. 2004 , 47, 718-24	125
1843	The anti-hyperalgesic activity of retigabine is mediated by KCNQ potassium channel activation. 2004 , 369, 382-90	61
1842	Electrochemically synthesized polypyrrole and Cu-plated nylon/spandex for electrotherapeutic pad electrode. 2004 , 91, 4064-4071	52
1841	Activation of p38 mitogen-activated protein kinase in spinal hyperactive microglia contributes to pain hypersensitivity following peripheral nerve injury. 2004 , 45, 89-95	411
1840	Effects of spinal nerve ligation on immunohistochemically identified neurons in the L4 and L5 dorsal root ganglia of the rat. 2004 , 475, 575-89	78
1839	A preclinical post laminectomy rat model mimics the human post laminectomy syndrome. <i>Journal of Neuroscience Methods</i> , 2004 , 137, 283-9	3 22
1838	Mechanical and cold allodynia in a rat spinal cord contusion model. 2004 , 21, 25-31	27
1837	A rat pain model of vincristine-induced neuropathy. 2004 , 99, 91-8	21
1836	Modeling diabetic sensory neuropathy in rats. 2004 , 99, 55-65	57
1835	Functional genomic analysis in pain research using hybridization arrays. 2004 , 99, 239-53	
1834	Neurokinin-1 receptor gene expression in the mouse dorsal horn increases with neuropathic pain. 2004 , 5, 71-6	11
1833	Differential blockade of nerve injury-induced thermal and tactile hypersensitivity by systemically administered brain-penetrating and peripherally restricted local anesthetics. 2004 , 5, 281-9	22

1832	Spinal neural cyclooxygenase-2 mediates pain caused in a rat model of lumbar disk herniation. 2004 , 5, 385-91	31
1831	Spinal gap junctions: potential involvement in pain facilitation. 2004 , 5, 392-405	131
1830	Detection and assessment of xenobiotic-induced sensory neuropathy. 2005 , Chapter 11, Unit11.13	
1829	Pain Research. 2004 ,	2
1828	Attenuation of mechanical but not thermal hyperalgesia by electroacupuncture with the involvement of opioids in rat model of chronic inflammatory pain. 2004 , 63, 99-103	62
1827	Cyclooxygenase inhibition in nerve-injury- and TNF-induced hyperalgesia in the rat. 2004 , 185, 160-8	79
1826	Partial sciatic nerve transection causes redistribution of pain-related peptides and lowers withdrawal threshold. 2004 , 188, 444-51	15
1825	Tart cherry anthocyanins suppress inflammation-induced pain behavior in rat. 2004 , 153, 181-8	138
1824	Novel mechanism of enhanced nociception in a model of AIDS therapy-induced painful peripheral neuropathy in the rat. 2004 , 107, 147-58	96
1823	Substance P signaling contributes to the vascular and nociceptive abnormalities observed in a tibial fracture rat model of complex regional pain syndrome I. 2004 , 108, 95-107	144
1822	Antiallodynic effects of intrathecally administered 5-HT(2C) receptor agonists in rats with nerve injury. 2004 , 108, 163-9	52
1821	Snake venom phospholipase A2s (Asp49 and Lys49) induce mechanical allodynia upon peri-sciatic administration: involvement of spinal cord glia, proinflammatory cytokines and nitric oxide. 2004 , 108, 180-91	56
1820	Contribution of sensitized P2X receptors in inflamed tissue to the mechanical hypersensitivity revealed by phosphorylated ERK in DRG neurons. 2004 , 108, 258-266	76
1819	Evidence that CB-1 and CB-2 cannabinoid receptors mediate antinociception in neuropathic pain in the rat. 2004 , 109, 124-31	104
1818	HIV-1 gp120 stimulates proinflammatory cytokine-mediated pain facilitation via activation of nitric oxide synthase-I (nNOS). 2004 , 110, 517-530	80
1817	Attenuation of mechanical allodynia by clinically utilized drugs in a rat chemotherapy-induced neuropathic pain model. 2004 , 110, 56-63	94
1816	Peri-sciatic proinflammatory cytokines, reactive oxygen species, and complement induce mirror-image neuropathic pain in rats. 2004 , 110, 299-309	110
1815	Snake venom components enhance pain upon subcutaneous injection: an initial examination of spinal cord mediators. 2004 , 111, 65-76	46

1814	Reactive oxygen species (ROS) play an important role in a rat model of neuropathic pain. 2004 , 111, 116-24	325
1813	Chronic post-ischemia pain (CPIP): a novel animal model of complex regional pain syndrome-type I (CRPS-I; reflex sympathetic dystrophy) produced by prolonged hindpaw ischemia and reperfusion in the rat. 2004 , 112, 94-105	233
1812	In vivo evidence for apoptosis, but not inflammation in the hindlimb muscle of neuropathic rats. 2004 , 112, 121-30	24
1811	Effects of low-affinity NMDA receptor channel blockers in two rat models of chronic pain. 2004 , 47, 175-83	32
1810	Participation of brainstem nuclei in the pronociceptive effect of lesion or neural block of the anterior pretectal nucleus in a rat model of incisional pain. 2004 , 47, 117-27	11
1809	Mouse strains that lack spinal dynorphin upregulation after peripheral nerve injury do not develop neuropathic pain. 2004 , 123, 43-52	55
1808	Peripheral glutamate receptors contribute to mechanical hyperalgesia in a neuropathic pain model of the rat. 2004 , 128, 169-76	56
1807	The contribution of autophosphorylated alpha-calcium-calmodulin kinase II to injury-induced persistent pain. 2004 , 128, 889-98	49
1806	Proteomic identification of brainstem cytosolic proteins in a neuropathic pain model. 2004 , 128, 193-200	28
1805	Peri-sciatic administration of indomethacin early after nerve injury can attenuate the development of tactile allodynia in a rat model of L5 single spinal nerve injury. 2004 , 356, 37-40	10
1804	Melanocortin 4 receptor is expressed in the dorsal root ganglions and down-regulated in neuropathic rats. 2004 , 358, 79-82	36
1803	Involvement of peripherally released substance P and calcitonin gene-related peptide in mediating mechanical hyperalgesia in a traumatic neuropathy model of the rat. 2004 , 360, 129-32	56
1802	Formalin-induced spinal cord calcium/calmodulin-dependent protein kinase II alpha expression is modulated by heme oxygenase in mice. 2004 , 360, 61-4	25
1801	Cannabinoids blocks tactile allodynia in diabetic mice without attenuation of its antinociceptive effect. 2004 , 368, 82-6	56
1800	Ketamine potentiates the effect of electroacupuncture on mechanical allodynia in a rat model of neuropathic pain. 2004 , 368, 327-31	50
1799	The effect of WIN 55,212-2, a cannabinoid agonist, on tactile allodynia in diabetic rats. 2004 , 371, 167-70	65
1798	Paclitaxel-induced neuropathic hypersensitivity in mice: responses in 10 inbred mouse strains. 2004 , 74, 2593-604	102
1797	Reduced inflammatory pain in mice deficient in the differential screening-selected gene aberrative in neuroblastoma. 2004 , 25, 504-14	1

1796	Voltage-gated sodium channels and hyperalgesia. 2004 , 44, 371-97	277
1795	Absence of mechanical allodynia and Abeta-fiber sprouting after sciatic nerve injury in mice lacking membrane-type 5 matrix metalloproteinase. 2004 , 557, 125-8	53
1794	Efficacy of duloxetine, a potent and balanced serotonin-norepinephrine reuptake inhibitor in persistent pain models in rats. 2004 , 311, 576-84	241
1793	Intrathecal clonidine potentiates suppression of tactile hypersensitivity by spinal cord stimulation in a model of neuropathy. 2004 , 99, 135-139	47
1792	Characterization of a new animal model for evaluation of persistent postthoracotomy pain. 2004 , 99, 1453-1460	74
1791	Preoperative sciatic nerve block decreases mechanical allodynia more in young rats: is preemptive analgesia developmentally modulated?. 2004 , 99, 140-145	22
1790	Repeated dosing with oral allosteric modulator of adenosine A1 receptor produces tolerance in rats with neuropathic pain. 2004 , 100, 956-61	15
1789	Production of paradoxical sensory hypersensitivity by alpha 2-adrenoreceptor agonists. 2004 , 100, 1538-44	29
1788	Effect of cyclooxygenase-1 inhibition in postoperative pain is developmentally regulated. 2004 , 101, 1031-5	8
1787	Disability of development of tolerance to morphine and U-50,488H, a selective kappa-opioid receptor agonist, in neuropathic pain model mice. 2004 , 94, 305-12	18
1786	Supernatant of traumatized muscle induces inflammation and pain, but not microcirculatory perfusion failure and apoptotic cell death. 2005 , 24, 219-25	12
1785	Involvement of N-methyl-D-aspartate-type glutamate receptor epsilon1 and epsilon4 subunits in tonic inflammatory pain and neuropathic pain. 2005 , 16, 1667-70	12
1784	Transient cervical nerve root compression in the rat induces bilateral forepaw allodynia and spinal glial activation: mechanical factors in painful neck injuries. 2005 , 30, 1924-32	67
1783	Morphine can enhance the antiallodynic effect of intrathecal R-PIA in rats with nerve ligation injury. 2005 , 100, 461-468	8
1782	Spinal opioid receptor like1 receptor agonist, but not N-methyl-D-aspartic acid antagonist, reverses the secondary mechanical allodynia induced by intradermal injection of capsaicin in rats. 2005 , 100, 1087-1092	9
1781	The monoamine-mediated antiallodynic effects of intrathecally administered milnacipran, a serotonin noradrenaline reuptake inhibitor, in a rat model of neuropathic pain. 2005 , 100, 1406-1410	65
1780	Preoperative inhibition of cyclooxygenase-1 in the spinal cord reduces postoperative pain. 2005 , 100, 1390-1393	42
1779	Local anesthetics for the treatment of neuropathic pain: on the limits of meta-analysis. 2005 , 101, 1736-1737	8

1778	Enhanced wind-up of the C-fiber-mediated nociceptive flexor reflex movement following painful diabetic neuropathy in mice. 2005 , 97, 195-202		15
1777	Spinal muscarinic and nicotinic subtypes activated by clonidine in postincisional pain. 2005 , 103, 1253-8		11
1776	Systemic Meloxicam Reduces Tactile Allodynia Development After L5 Single Spinal Nerve Injury in Rats. 2005 , 30, 351-355		15
1775	Adenosine reduces glutamate release in rat spinal synaptosomes. 2005 , 103, 1060-5		19
1774	Medullary N-type and P/Q-type calcium channels contribute to neuropathy-induced allodynia. 2005 , 16, 563-6		22
1773	Pharmacological evaluation of the selective spinal nerve ligation model of neuropathic pain in the rat. <i>Journal of Neuroscience Methods</i> , 2005 , 144, 175-81	3	84
1772	Long-term changes of c-Fos expression in the rat spinal cord following chronic constriction injury. 2005 , 9, 345-54		16
1771	Anti-fibroblast growth factor-2 antibodies attenuate mechanical allodynia in a rat model of neuropathic pain. 2005 , 27, 315-24		20
1770	Evidence that gabapentin reduces neuropathic pain by inhibiting the spinal release of glutamate. 2005 , 94, 1131-9		124
1769	Controlling pathological pain by adenovirally driven spinal production of the anti-inflammatory cytokine, interleukin-10. 2005 , 21, 2136-48		183
1768	Ca ²⁺ /calmodulin-dependent protein kinase II in the spinal cord contributes to neuropathic pain in a rat model of mononeuropathy. 2005 , 21, 2467-74		40
1767	Inhibitory effects of CB1 and CB2 receptor agonists on responses of DRG neurons and dorsal horn neurons in neuropathic rats. 2005 , 22, 371-9		129
1766	Inhibition of peripheral vanilloid TRPV1 receptors reduces noxious heat-evoked responses of dorsal horn neurons in naïve, carrageenan-inflamed and neuropathic rats. 2005 , 22, 361-70		49
1765	Involvement of spinal cord nuclear factor kappaB activation in rat models of proinflammatory cytokine-mediated pain facilitation. 2005 , 22, 1977-86		117
1764	An initial investigation of spinal mechanisms underlying pain enhancement induced by fractalkine, a neuronally released chemokine. 2005 , 22, 2775-82		142
1763	Role of descending noradrenergic system and spinal alpha2-adrenergic receptors in the effects of gabapentin on thermal and mechanical nociception after partial nerve injury in the mouse. 2005 , 144, 703-14		110
1762	BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain. 2005 , 438, 1017-21		1420
1761	Activation of ERK/CREB pathway in spinal cord contributes to chronic constrictive injury-induced neuropathic pain in rats. 2005 , 26, 789-98		97

1760	Synergy between intrathecal omega-conotoxin CVID and dexmedetomidine to attenuate mechanical hypersensitivity in the rat. 2005 , 506, 221-7	7
1759	ABT-594 (a nicotinic acetylcholine agonist): anti-allodynia in a rat chemotherapy-induced pain model. 2005 , 509, 43-8	38
1758	Anti-allodynic and anti-hyperalgesic effects of nociceptin receptor antagonist, JTC-801, in rats after spinal nerve injury and inflammation. 2005 , 510, 223-8	25
1757	Anti-allodynic interactions between NMDA receptor channel blockers and morphine or clonidine in neuropathic rats. 2005 , 519, 80-5	11
1756	The antihyperalgesic effects of the T-type calcium channel blockers ethosuximide, trimethadione, and mibefradil. 2005 , 521, 79-85	46
1755	Spinal nerve ligation increases alpha2-adrenergic receptor G-protein coupling in the spinal cord. 2005 , 1038, 76-82	48
1754	Glutamate uptake is attenuated in spinal deep dorsal and ventral horn in the rat spinal nerve ligation model. 2005 , 1041, 38-47	61
1753	Effect of systemic and intrathecal gabapentin on allodynia in a new rat model of postherpetic neuralgia. 2005 , 1042, 108-13	35
1752	Anti-hyperalgesic effects of intrathecally administered neuropeptide W-23, and neuropeptide B, in tests of inflammatory pain in rats. 2005 , 1045, 97-106	33
1751	Increased hyperalgesia after tissue injury and faster recovery of allodynia after nerve injury in the GalR1 knockout mice. 2005 , 39, 217-21	16
1750	Knock down of the alpha 5 nicotinic acetylcholine receptor in spinal nerve-ligated rats alleviates mechanical allodynia. 2005 , 80, 135-43	26
1749	Bone cancer pain: from model to mechanism to therapy. 2005 , 29, S32-46	110
1748	Gene transfer of glutamic acid decarboxylase reduces neuropathic pain. 2005 , 57, 914-8	83
1747	CCL2 and CXCL1 trigger calcitonin gene-related peptide release by exciting primary nociceptive neurons. 2005 , 82, 51-62	109
1746	End-to-side nerve grafting of the tibial nerve to bridge a neuroma-in-continuity. 2005 , 25, 155-64; discussion 164-6	14
1745	Centrally mediated antihyperalgesic and antiallodynic effects of zonisamide following partial nerve injury in the mouse. 2005 , 372, 107-14	20
1744	The antinociceptive and anxiolytic-like effects of the metabotropic glutamate receptor 5 (mGluR5) antagonists, MPEP and MTEP, and the mGluR1 antagonist, LY456236, in rodents: a comparison of efficacy and side-effect profiles. 2005 , 179, 207-17	137
1743	Effects of intrathecally administered NaV1.8 antisense oligonucleotide on the expression of sodium channel mRNA in dorsal root ganglion. 2005 , 25, 696-9	4

1742	Expression of pERK in the Neurons and Microglia in the Spinal Cord of the Rat with Neuropathic Pain. 2005 , 1, 19	1
1741	Hyperexcitability of axotomized and neighboring unaxotomized sensory neurons is reduced days after perineural clonidine at the site of injury. 2005 , 94, 3159-67	29
1740	The Effect of ATP-sensitive Potassium Channel on R-PIA Induced Mechanical Allodynia in a Peripheral Neuropathic Rat. 2005 , 18, 107	2
1739	Distinct roles of group III metabotropic glutamate receptors in control of nociception and dorsal horn neurons in normal and nerve-injured Rats. 2005 , 312, 120-6	62
1738	Changes in electrophysiological properties and sodium channel Nav1.3 expression in thalamic neurons after spinal cord injury. 2005 , 128, 2359-71	142
1737	Transgenic expression of a dominant-negative ASIC3 subunit leads to increased sensitivity to mechanical and inflammatory stimuli. 2005 , 25, 9893-901	105
1736	Effect of morphine on deep dorsal horn projection neurons depends on spinal GABAergic and glycinergic tone: implications for reduced opioid effect in neuropathic pain. 2005 , 315, 696-703	26
1735	Plasticity of the Central Nervous System and Chronic Pain. 2005 , 1, 27-36	
1734	Lack of the central nervous system- and neural crest-expressed forkhead gene Foxs1 affects motor function and body weight. 2005 , 25, 5616-25	40
1733	Sciatic Inflammatory Neuritis. 2005 , 1, 3-11	1
1732	Upregulation of forebrain NMDA NR2B receptors contributes to behavioral sensitization after inflammation. 2005 , 25, 11107-16	185
1731	Upregulation of the voltage-gated sodium channel beta2 subunit in neuropathic pain models: characterization of expression in injured and non-injured primary sensory neurons. 2005 , 25, 10970-80	97
1730	Altered behavioral responses to noxious stimuli and fear in glutamate receptor 5 (GluR5)- or GluR6-deficient mice. 2005 , 25, 977-84	93
1729	Comparison of the antinociceptive profiles of gabapentin and 3-methylgabapentin in rat models of acute and persistent pain: implications for mechanism of action. 2005 , 313, 1209-16	58
1728	Efficacy of duloxetine, a potent and balanced serotonergic and noradrenergic reuptake inhibitor, in inflammatory and acute pain models in rodents. 2005 , 312, 726-32	123
1727	Nonviral gene transfer of human hepatocyte growth factor improves streptozotocin-induced diabetic neuropathy in rats. 2005 , 54, 846-54	43
1726	Activin induces tactile allodynia and increases calcitonin gene-related peptide after peripheral inflammation. 2005 , 25, 9227-35	50
1725	Cholecystokinin in the rostral ventromedial medulla mediates opioid-induced hyperalgesia and antinociceptive tolerance. 2005 , 25, 409-16	164

1724	A blocking antibody to nerve growth factor attenuates skeletal pain induced by prostate tumor cells growing in bone. 2005 , 65, 9426-35	178
1723	Alpha2-adrenoceptor stimulation transforms immune responses in neuritis and blocks neuritis-induced pain. 2005 , 25, 8988-94	59
1722	Contribution of the tetrodotoxin-resistant voltage-gated sodium channel NaV1.9 to sensory transmission and nociceptive behavior. 2005 , 102, 9382-7	217
1721	Spinal L-type calcium channel blockade abolishes opioid-induced sensory hypersensitivity and antinociceptive tolerance. 2005 , 101, 1730-1735	31
1720	Loss of neurons from laminae I-III of the spinal dorsal horn is not required for development of tactile allodynia in the spared nerve injury model of neuropathic pain. 2005 , 25, 6658-66	115
1719	Systemic meloxicam reduces tactile allodynia development after L5 single spinal nerve injury in rats. 2005 , 30, 351-5	7
1718	Regulation of two rat mas-related genes in a model of neuropathic pain. 2005 , 142, 58-64	10
1717	Skin denervation, neuropathology, and neuropathic pain in a laser-induced focal neuropathy. 2005 , 18, 40-53	32
1716	Spinal blockade of TNF blocks spinal nerve ligation-induced increases in spinal P-p38. 2005 , 379, 209-13	109
1715	Effect of the cannabinoid ajulemic acid on rat models of neuropathic and inflammatory pain. 2005 , 382, 231-5	20
1714	Experiments with nitric oxide synthase inhibitors in spinal nerve ligated rats provide no evidence of a role for nitric oxide in neuropathic mechanical allodynia. 2005 , 385, 179-83	20
1713	Functional alteration of inhibitory influences on spinal motor output in painful diabetic neuropathy in rats. 2005 , 389, 152-6	4
1712	The effects of intraspinal L-NOARG or SIN-1 on the control by descending pathways of incisional pain in rats. 2005 , 76, 1939-51	9
1711	Endogenous and exogenous melanocortin antagonists induce anti-allodynic effects in a model of rat neuropathic pain. 2005 , 157, 55-62	27
1710	Choosing the correct functional assay: a comprehensive assessment of functional tests in the rat. 2005 , 163, 143-58	88
1709	TRPV1 desensitisation and endogenous vanilloid involvement in the enhanced analgesia induced by capsaicin in inflamed tissues. 2005 , 67, 476-81	22
1708	Ectopic discharges from injured nerve fibers are highly correlated with tactile allodynia only in early, but not late, stage in rats with spinal nerve ligation. 2005 , 191, 128-36	56
1707	Stepwise motor and all-or-none sensory recovery is associated with nonlinear sparing after incremental spinal cord injury in rats. 2005 , 191, 251-65	111

1706	Nociceptor-specific gene deletion using heterozygous NaV1.8-Cre recombinase mice. 2005 , 113, 27-36	157
1705	Spinal noradrenaline transporter inhibition by reboxetine and Xen2174 reduces tactile hypersensitivity after surgery in rats. 2005 , 113, 271-276	39
1704	Intrathecal morphine and ketorolac analgesia after surgery: comparison of spontaneous and elicited responses in rats. 2005 , 113, 376-385	27
1703	Medium and large injured dorsal root ganglion cells increase TRPV-1, accompanied by increased alpha2C-adrenoceptor co-expression and functional inhibition by clonidine. 2005 , 113, 386-394	82
1702	Minocycline attenuates mechanical allodynia and proinflammatory cytokine expression in rat models of pain facilitation. 2005 , 115, 71-83	529
1701	Peripheral inflammation selectively increases TRPV1 function in IB4-positive sensory neurons from adult mouse. 2005 , 115, 37-49	115
1700	Anti-NGF therapy profoundly reduces bone cancer pain and the accompanying increase in markers of peripheral and central sensitization. 2005 , 115, 128-41	236
1699	Spinal CK2 regulates nociceptive signaling in models of inflammatory pain. 2005 , 115, 182-90	22
1698	Enhancement of NMDA receptor phosphorylation of the spinal dorsal horn and nucleus gracilis neurons in neuropathic rats. 2005 , 116, 62-72	99
1697	Neuropathic pain: early spontaneous afferent activity is the trigger. 2005 , 116, 243-256	148
1696	Intraneural injection of interleukin-1beta and tumor necrosis factor-alpha into rat sciatic nerve at physiological doses induces signs of neuropathic pain. 2005 , 116, 257-263	230
1695	Heat and mechanical hyperalgesia in mice model of cancer pain. 2005 , 117, 19-29	59
1694	Relationship between sodium channel NaV1.3 expression and neuropathic pain behavior in rats. 2005 , 117, 145-53	132
1693	Activated PKA and PKC, but not CaMKIIalpha, are required for AMPA/Kainate-mediated pain behavior in the thermal stimulus model. 2005 , 117, 259-270	41
1692	Anti-allodynic efficacy of the chi-conopeptide, Xen2174, in rats with neuropathic pain. 2005 , 118, 112-24	70
1691	Mechanisms underlying enhanced P2X receptor-mediated responses in the neuropathic pain state. 2005 , 119, 38-48	97
1690	Assessment of nociception in acrylamide-induced neuropathy in rats. 2005 , 119, 104-112	17
1689	Basal and carrageenan-induced pain behavior in Sprague-Dawley, Lewis and Fischer rats. 2005 , 85, 177-86	25

1688	mGlu and NMDA receptor contributions to capsaicin-induced thermal and mechanical hypersensitivity. 2005 , 48, 325-32	55
1687	Comparison of responses of ventral posterolateral and posterior complex thalamic neurons in naive rats and rats with hindpaw inflammation: mu-opioid receptor mediated inhibitions. 2005 , 48, 607-16	14
1686	Analgesic effects of the selective group II (mGlu2/3) metabotropic glutamate receptor agonists LY379268 and LY389795 in persistent and inflammatory pain models after acute and repeated dosing. 2005 , 49 Suppl 1, 206-18	72
1685	Increased sensitivity to acute and persistent pain in neuron-specific endothelin-1 knockout mice. 2005 , 130, 349-58	26
1684	alpha2-adrenoceptors inhibit the intracellular Ca ²⁺ response to electrical stimulation in normal and injured sensory neurons, with increased inhibition of calcitonin gene-related peptide expressing neurons after injury. 2005 , 131, 189-97	29
1683	A new role for the renin-angiotensin system in the rat periaqueductal gray matter: angiotensin receptor-mediated modulation of nociception. 2005 , 132, 453-63	52
1682	Imbalance between the expression of NK1 and GABAB receptors in nociceptive spinal neurons during secondary hyperalgesia: a c-Fos study in the monoarthritic rat. 2005 , 132, 905-16	12
1681	Increased responsiveness of rat dorsal horn neurons in vivo following prolonged intrathecal exposure to interferon-gamma. 2005 , 135, 969-77	28
1680	Identification and biological evaluation of 4-(3-trifluoromethylpyridin-2-yl)piperazine-1-carboxylic acid (5-trifluoromethylpyridin-2-yl)amide, a high affinity TRPV1 (VR1) vanilloid receptor antagonist. 2005 , 48, 1857-72	152
1679	Topical high molecular weight hyaluronan reduces radicular pain post laminectomy in a rat model. 2005 , 5, 494-502	26
1678	Activation of the spinal cord complement cascade might contribute to mechanical allodynia induced by three animal models of spinal sensitization. 2005 , 6, 174-83	44
1677	Hyperpolarization-activated, cation-nonspecific, cyclic nucleotide-modulated channel blockade alleviates mechanical allodynia and suppresses ectopic discharge in spinal nerve ligated rats. 2005 , 6, 417-24	60
1676	A behavioral model of neuropathic pain induced by ligation of the common peroneal nerve in mice. 2005 , 6, 747-56	59
1675	Neonatal chronic hind paw inflammation alters sensitization to intradermal capsaicin in adult rats: a behavioral and immunocytochemical study. 2005 , 6, 798-808	42
1674	Up-and-Down Method. 2005 ,	
1673	Formalin injection causes a coordinated spinal cord CO/NO-cGMP signaling system response. 2005 , 1, 33	7
1672	Neuropathic pain develops normally in mice lacking both Na(v)1.7 and Na(v)1.8. 2005 , 1, 24	146
1671	Spared nerve injury rats exhibit thermal hyperalgesia on an automated operant dynamic thermal escape task. 2005 , 1, 18	33

1670	Controlling neuropathic pain by adeno-associated virus driven production of the anti-inflammatory cytokine, interleukin-10. 2005 , 1, 9	148
1669	Sodium channel expression in the ventral posterolateral nucleus of the thalamus after peripheral nerve injury. 2006 , 2, 27	55
1668	HSV-mediated expression of interleukin-4 in dorsal root ganglion neurons reduces neuropathic pain. 2006 , 2, 6	121
1667	Inflammatory and neuropathic pain animals exhibit distinct responses to innocuous thermal and motoric challenges. 2006 , 2, 1	15
1666	Activated microglia contribute to the maintenance of chronic pain after spinal cord injury. 2006 , 26, 4308-17	488
1665	Spontaneous pain, both neuropathic and inflammatory, is related to frequency of spontaneous firing in intact C-fiber nociceptors. 2006 , 26, 1281-92	324
1664	Animal Models of Nociception. 221-235	10
1663	Sciatic chronic constriction injury produces cell-type-specific changes in the electrophysiological properties of rat substantia gelatinosa neurons. 2006 , 96, 579-90	111
1662	Calibrated forceps: a sensitive and reliable tool for pain and analgesia studies. 2006 , 7, 32-9	55
1661	Covariance among age, spinal p38 MAP kinase activation and allodynia. 2006 , 7, 337-45	7
1660	Differential blockade of nerve injury-induced shift in weight bearing and thermal and tactile hypersensitivity by milnacipran. 2006 , 7, 513-20	37
1659	Spinal cord glia and interleukin-1 do not appear to mediate persistent allodynia induced by intramuscular acidic saline in rats. 2006 , 7, 757-67	37
1658	Spinal glial activation contributes to postoperative mechanical hypersensitivity in the rat. 2006 , 7, 816-22	99
1657	Reduced expression of endothelin B receptors and mechanical hyperalgesia in experimental chronic diabetes. 2006 , 201, 399-406	18
1656	Fractalkine and minocycline alter neuronal activity in the spinal cord dorsal horn. 2006 , 580, 4306-10	62
1655	TRPA1 mediates the inflammatory actions of environmental irritants and proalgesic agents. 2006 , 124, 1269-82	1447
1654	Aging, peripheral nerve injury and nociception: effects of the antioxidant 16-desmethyltirilazad. 2006 , 166, 159-65	17
1653	Interleukin-1alpha gene-deficient mice show reduced nociceptive sensitivity in models of inflammatory and neuropathic pain but not post-operative pain. 2006 , 167, 355-64	96

1652	Allodynia and hyperalgesia suppression by a novel analgesic in experimental neuropathic pain. 2006 , 350, 358-63	10
1651	Nociceptor-derived brain-derived neurotrophic factor regulates acute and inflammatory but not neuropathic pain. 2006 , 31, 539-48	127
1650	Nerve growth factor mediates peripheral mechanical hypersensitivity that accompanies experimental cystitis in mice. 2006 , 392, 193-7	60
1649	Expression of IL-1beta in supraspinal brain regions in rats with neuropathic pain. 2006 , 407, 176-81	85
1648	Intracellular messengers involved in spontaneous pain, heat hyperalgesia, and mechanical allodynia induced by intrathecal dihydroxyphenylglycine. 2006 , 409, 224-9	3
1647	Therapeutic efficacy of prosaposin-derived peptide on different models of allodynia. 2006 , 121, 14-21	18
1646	Glucocorticoid inhibition of vascular abnormalities in a tibia fracture rat model of complex regional pain syndrome type I. 2006 , 121, 158-67	44
1645	Chronic pain and genetic background interact and influence opioid analgesia, tolerance, and physical dependence. 2006 , 121, 232-240	58
1644	Analgesic effect of vitamin E is mediated by reducing central sensitization in neuropathic pain. 2006 , 122, 53-62	93
1643	Peripheral 5-HT _{2A} receptor antagonism attenuates primary thermal hyperalgesia and secondary mechanical allodynia after thermal injury in rats. 2006 , 122, 130-6	57
1642	Cancer-related bone pain is attenuated by a systemically available delta-opioid receptor agonist. 2006 , 122, 174-81	52
1641	Involvement of the TTX-resistant sodium channel Nav 1.8 in inflammatory and neuropathic, but not post-operative, pain states. 2006 , 123, 75-82	132
1640	The role of tumor necrosis factor-alpha in the neuropathic pain induced by Lumbar 5 ventral root transection in rat. 2006 , 123, 306-321	171
1639	Spinal cord dynorphin expression increases, but does not drive microglial prostaglandin production or mechanical hypersensitivity after incisional surgery in rats. 2006 , 125, 43-52	20
1638	(L)-Phenylglycine, but not necessarily other alpha2delta subunit voltage-gated calcium channel ligands, attenuates neuropathic pain in rats. 2006 , 125, 136-42	36
1637	Clonidine maintains intrathecal self-administration in rats following spinal nerve ligation. 2006 , 125, 257-263	34
1636	Screening for pain phenotypes: analysis of three congenic mouse strains on a battery of nine nociceptive assays. 2006 , 126, 24-34	61
1635	Capsaicin-evoked brain activation and central sensitization in anaesthetised rats: a functional magnetic resonance imaging study. 2006 , 126, 35-45	30

1634	Mechanical sensory threshold testing using nylon monofilaments: the pain field's "tin standard". 2006 , 124, 13-7	53
1633	Endogenous beta-endorphin induces thermal analgesia at the initial stages of a murine osteosarcoma. 2006 , 27, 2778-85	29
1632	Limbic and HPA axis function in an animal model of chronic neuropathic pain. 2006 , 88, 67-76	106
1631	Development and expression of neuropathic pain in CB1 knockout mice. 2006 , 50, 111-22	37
1630	Fluvoxamine, a selective serotonin reuptake inhibitor, exerts its antiallodynic effects on neuropathic pain in mice via 5-HT _{2A/2C} receptors. 2006 , 51, 866-72	43
1629	Effects of NAAG peptidase inhibitor 2-PMPA in model chronic pain - relation to brain concentration. 2006 , 51, 1163-71	43
1628	Analgesic effects of serotonergic, noradrenergic or dual reuptake inhibitors in the carrageenan test in rats: evidence for synergism between serotonergic and noradrenergic reuptake inhibition. 2006 , 51, 1172-80	33
1627	Effects of intrathecal and i.c.v. administration of neuropeptide W-23 and neuropeptide B on the mechanical allodynia induced by partial sciatic nerve ligation in rats. 2006 , 137, 265-73	10
1626	Spinal nerve ligation does not alter the expression or function of GABA(B) receptors in spinal cord and dorsal root ganglia of the rat. 2006 , 138, 1277-87	41
1625	Rescue and regeneration of injured peripheral nerve axons by intrathecal insulin. 2006 , 139, 429-49	71
1624	The roles of different subtypes of opioid receptors in mediating the nucleus submedius opioid-evoked antiallodynia in a neuropathic pain model of rats. 2006 , 138, 1319-27	11
1623	Absence of Reelin results in altered nociception and aberrant neuronal positioning in the dorsal spinal cord. 2006 , 139, 1385-96	37
1622	Inhibitory M ₂ muscarinic receptors are upregulated in both axotomized and intact small diameter dorsal root ganglion cells after peripheral nerve injury. 2006 , 140, 259-68	33
1621	Role of astrocytic S100beta in behavioral hypersensitivity in rodent models of neuropathic pain. 2006 , 140, 1003-10	51
1620	Descending facilitation from the rostral ventromedial medulla maintains nerve injury-induced central sensitization. 2006 , 140, 1311-20	116
1619	Secondary hyperalgesia in the monoarthritic rat is mediated by GABA _B and NK1 receptors of spinal dorsal horn neurons: a behavior and c-fos study. 2006 , 141, 2087-95	22
1618	Comparison of antinociceptive actions of standard analgesics in attenuating capsaicin and nerve-injury-induced mechanical hypersensitivity. 2006 , 143, 587-96	78
1617	Potentiation of Morphine's Antinociception by Group II and Group III Metabotropic Glutamate Receptors Agonists on a Rat Incisional Pain. 2006 , 19, 131	1

1616	Does the tibial and sural nerve transection model represent sympathetically independent pain?. 2006 , 47, 847-51	15
1615	EConotoxins GVIA, MVIIA and CVID: SAR and Clinical Potential. 2006 , 4, 193-214	20
1614	. 2006 ,	8
1613	Mechanisms of the antinociceptive action of gabapentin. 2006 , 100, 471-86	185
1612	Adverse effects of gabapentin and lack of anti-allodynic efficacy of amitriptyline in the streptozotocin model of painful diabetic neuropathy. 2006 , 14, 42-51	14
1611	Injection of nuclear factor-kappa B decoy into the sciatic nerve suppresses mechanical allodynia and thermal hyperalgesia in a rat inflammatory pain model. 2006 , 31, 2904-8	19
1610	Age-dependent responses to nerve injury-induced mechanical allodynia. 2006 , 104, 344-50	32
1609	Blockade of the complement C5a receptor reduces incisional allodynia, edema, and cytokine expression. 2006 , 104, 1274-82	59
1608	A genetic analysis of opioid-induced hyperalgesia in mice. 2006 , 104, 1054-62	93
1607	Perineural clonidine reduces p38 mitogen-activated protein kinase activation in sensory neurons. 2006 , 17, 1313-7	12
1606	Vector-mediated gene transfer to express inhibitory neurotransmitters in dorsal root ganglion reduces pain in a rodent model of lumbar radiculopathy. 2006 , 31, 1555-8	15
1605	B1 and TRPV-1 receptor genes and their relationship to hyperalgesia following spinal cord injury. 2006 , 31, 2778-82	45
1604	Up-regulation of acid-sensing ion channel 3 in dorsal root ganglion neurons following application of nucleus pulposus on nerve root in rats. 2006 , 31, 2048-52	38
1603	Chronic activation of spinal adenosine A1 receptors results in hypersensitivity. 2006 , 17, 1619-22	5
1602	Neuropathic pain is enhanced in delta-opioid receptor knockout mice. 2006 , 23, 830-4	93
1601	Sensory neuronal phenotype in galanin receptor 2 knockout mice: focus on dorsal root ganglion neurone development and pain behaviour. 2006 , 23, 627-36	48
1600	CB2 receptor-mediated antihyperalgesia: possible direct involvement of neural mechanisms. 2006 , 23, 1530-8	222
1599	Spatial and temporal relationship between monocyte chemoattractant protein-1 expression and spinal glial activation following peripheral nerve injury. 2006 , 97, 772-83	266

1598	The structure and mode of action of different botulinum toxins. 2006 , 13 Suppl 4, 1-9	173
1597	Altered pain behavior and regeneration after nerve injury in TNF receptor deficient mice. 2006 , 11, 294-303	45
1596	Intrathecal administration of Cav3.2 and Cav3.3 antisense oligonucleotide reverses tactile allodynia and thermal hyperalgesia in rats following chronic compression of dorsal root of ganglion. 2006 , 27, 1547-52	39
1595	Dynorphin A activates bradykinin receptors to maintain neuropathic pain. 2006 , 9, 1534-40	121
1594	Actions of the FAAH inhibitor URB597 in neuropathic and inflammatory chronic pain models. 2006 , 147, 281-8	219
1593	Inhibition of fatty acid amide hydrolase produces analgesia by multiple mechanisms. 2006 , 148, 102-13	143
1592	Spinal alpha(2)-adrenergic and muscarinic receptors and the NO release cascade mediate supraspinally produced effectiveness of gabapentin at decreasing mechanical hypersensitivity in mice after partial nerve injury. 2006 , 148, 233-44	51
1591	Physiological properties of spinal lamina II GABAergic neurons in mice following peripheral nerve injury. 2006 , 577, 869-78	57
1590	An animal model of chronic inflammatory pain: pharmacological and temporal differentiation from acute models. 2006 , 10, 537-49	62
1589	Animal models of pain for drug discovery. 2006 , 1, 323-34	19
1588	Effects of the RANKL inhibitor, osteoprotegerin, on the pain and histopathology of bone cancer in rats. 2006 , 23, 167-75	25
1587	Dynorphin A, kappa opioid receptors and the antinociceptive efficacy of asimadoline in streptozotocin-induced diabetic rats. 2006 , 49, 2775-85	25
1586	Benfotiamine relieves inflammatory and neuropathic pain in rats. 2006 , 530, 48-53	48
1585	The nitric oxide-cyclic GMP-protein kinase G-K ⁺ channel pathway participates in the antiallodynic effect of spinal gabapentin. 2006 , 531, 87-95	38
1584	Dissociation between post-surgical pain behaviors and spinal Fos-like immunoreactivity in the rat. 2006 , 531, 108-17	14
1583	mu- but not delta- and kappa-opioid receptors in the ventrolateral orbital cortex mediate opioid-induced antiallodynia in a rat neuropathic pain model. 2006 , 1076, 68-77	19
1582	Pharmacological evidence for the participation of NO-cyclic GMP-PKG-K ⁺ channel pathway in the antiallodynic action of resveratrol. 2006 , 84, 535-42	34
1581	Molecular mechanisms of neuropathic pain-phenotypic switch and initiation mechanisms. 2006 , 109, 57-77	193

1580	Reciprocal regulation of nuclear factor kappa B and its inhibitor ZAS3 after peripheral nerve injury. 2006 , 7, 4	17
1579	Behavioral assessment of neuropathic pain in preclinical models. 2006 , 67, 302-307	11
1578	Modulation of neurotrophic growth factors as a therapeutic strategy for neuropathic pain. 2006 , 67, 389-403	3
1577	Heterozygous P0 deficiency protects mice from vincristine-induced polyneuropathy. 2006 , 84, 37-46	26
1576	Thiamine and cyanocobalamin relieve neuropathic pain in rats: synergy with dexamethasone. 2006 , 77, 53-62	28
1575	Chapter 14 Descending excitatory systems. 2006 , 81, 193-210	9
1574	Antiallodynic and antihyperalgesic effects of selective competitive GLUK5 (GluR5) ionotropic glutamate receptor antagonists in the capsaicin and carrageenan models in rats. 2006 , 319, 396-404	32
1573	Analgesic effects of fatty acid amide hydrolase inhibition in a rat model of neuropathic pain. 2006 , 26, 13318-27	141
1572	muO-conotoxin MrVIB selectively blocks Nav1.8 sensory neuron specific sodium channels and chronic pain behavior without motor deficits. 2006 , 103, 17030-5	161
1571	TRPV1 receptors in the CNS play a key role in broad-spectrum analgesia of TRPV1 antagonists. 2006 , 26, 9385-93	237
1570	Impaired nociception and inflammatory pain sensation in mice lacking the prokineticin receptor PKR1: focus on interaction between PKR1 and the capsaicin receptor TRPV1 in pain behavior. 2006 , 26, 6716-27	112
1569	A peptide c-Jun N-terminal kinase (JNK) inhibitor blocks mechanical allodynia after spinal nerve ligation: respective roles of JNK activation in primary sensory neurons and spinal astrocytes for neuropathic pain development and maintenance. 2006 , 26, 3551-60	419
1568	The glial modulatory drug AV411 attenuates mechanical allodynia in rat models of neuropathic pain. 2006 , 2, 279-91	95
1567	Block of peripheral nerve sodium channels selectively inhibits features of neuropathic pain in rats. 2006 , 69, 823-32	51
1566	Alleviation of neuropathic pain by intrathecal injection of antisense oligonucleotides to p65 subunit of NF-kappaB. 2006 , 97, 553-8	58
1565	Differential and synergistic effects of selective norepinephrine and serotonin reuptake inhibitors in rodent models of pain. 2007 , 320, 1178-85	61
1564	Characterization of the antinociceptive actions of bicipadine in models of acute, persistent, and chronic pain. 2007 , 321, 1208-25	49
1563	Organ cross talk modulates pelvic pain. 2007 , 293, R1191-8	88

1562	Acute and Neuropathic Pain. 2007 , 327-349	
1561	High-fat diet induced neuropathy of pre-diabetes and obesity: effects of "healthy" diet and aldose reductase inhibition. 2007 , 56, 2598-608	191
1560	Are alpha9alpha10 nicotinic acetylcholine receptors a pain target for alpha-conotoxins?. 2007 , 72, 1406-10	96
1559	Modulation of thalamic nociceptive processing after spinal cord injury through remote activation of thalamic microglia by cysteine cysteine chemokine ligand 21. 2007 , 27, 8893-902	168
1558	Upregulation of substance P in low-threshold myelinated afferents is not required for tactile allodynia in the chronic constriction injury and spinal nerve ligation models. 2007 , 27, 2035-44	33
1557	Antibodies to nerve growth factor reverse established tactile allodynia in rodent models of neuropathic pain without tolerance. 2007 , 322, 282-7	86
1556	Pain assessment following experimental maxillofacial surgical procedure in sheep. 2007 , 41, 345-52	8
1555	Expression of CCR2 in both resident and bone marrow-derived microglia plays a critical role in neuropathic pain. 2007 , 27, 12396-406	325
1554	Gliosis alters expression and uptake of spinal glial amino acid transporters in a mouse neuropathic pain model. 2007 , 3, 141-53	45
1553	Antinociceptive pharmacology of N-(4-chlorobenzyl)-N'-(4-hydroxy-3-iodo-5-methoxybenzyl) thiourea, a high-affinity competitive antagonist of the transient receptor potential vanilloid 1 receptor. 2007 , 321, 791-8	62
1552	Design and evaluation of small interfering RNAs that target expression of the N-methyl-D-aspartate receptor NR1 subunit gene in the spinal cord dorsal horn. 2007 , 322, 982-8	33
1551	Extracellular signal-regulated kinase-regulated microglia-neuron signaling by prostaglandin E2 contributes to pain after spinal cord injury. 2007 , 27, 2357-68	165
1550	Encyclopedia of Pain. 2007 , 1424-1427	1
1549	Reduced expression of A-type potassium channels in primary sensory neurons induces mechanical hypersensitivity. 2007 , 27, 9855-65	113
1548	Upregulation of acid-sensing ion channel ASIC1a in spinal dorsal horn neurons contributes to inflammatory pain hypersensitivity. 2007 , 27, 11139-48	122
1547	Decreased nociceptive sensitization in mice lacking the fragile X mental retardation protein: role of mGluR1/5 and mTOR. 2007 , 27, 13958-67	161
1546	Control of chronic pain by the ubiquitin proteasome system in the spinal cord. 2007 , 27, 8226-37	28
1545	Inflammation-induced reduction of spontaneous activity by adjuvant: A novel model to study the effect of analgesics in rats. 2007 , 320, 194-201	86

1544	Tolerance to the antinociceptive and antiexudative effects of morphine in a murine model of peripheral inflammation. 2007 , 322, 360-8	28
1543	The supraspinally mediated analgesic effects of zonisamide in mice after peripheral nerve injury are independent of the descending monoaminergic system. 2007 , 104, 335-40	4
1542	[Experimental methods in pain using neuropathic and inflammatory animal models]. 2007 , 130, 39-44	4
1541	Analgesia induced by perineural clonidine is enhanced in persistent neuritis. 2007 , 18, 67-71	10
1540	Ischemic preconditioning prevents skeletal muscle tissue injury, but not nerve lesion upon tourniquet-induced ischemia. 2007 , 63, 788-97	27
1539	Does facet joint inflammation induce radiculopathy?: an investigation using a rat model of lumbar facet joint inflammation. 2007 , 32, 406-12	41
1538	Comparison of the mGluR1 antagonist A-841720 in rat models of pain and cognition. 2007 , 18, 273-81	16
1537	Anti-TNF-alpha antibody reduces pain-behavioral changes induced by epidural application of nucleus pulposus in a rat model depending on the timing of administration. 2007 , 32, 413-6	56
1536	Glial phosphorylated p38 MAP kinase mediates pain in a rat model of lumbar disc herniation and induces motor dysfunction in a rat model of lumbar spinal canal stenosis. 2007 , 32, 159-67	54
1535	Altered response to formalin by L5 spinal nerve ligation in rats: a behavioral and molecular study. 2007 , 104, 936-43	5
1534	Osteopontin expression and function within the dorsal root ganglion. 2007 , 18, 153-7	15
1533	Histaminergic Involvement in Neuropathic Pain Produced by Partial Ligation of the Sciatic Nerve in Rats. 2007 , 32, 124-129	21
1532	Rho kinase inhibitor improves motor dysfunction and hypoalgesia in a rat model of lumbar spinal canal stenosis. 2007 , 32, 2070-5	17
1531	Increase of 200-kDa neurofilament-immunoreactive afferents in the substantia gelatinosa in allodynic rats induced by compression of the dorsal root ganglion. 2007 , 32, 1265-71	18
1530	Opioid self-administration in the nerve-injured rat: relevance of antiallodynic effects to drug consumption and effects of intrathecal analgesics. 2007 , 106, 312-22	86
1529	Chronic Intrathecal Infusion of Minocycline Prevents the Development of Spinal-Nerve Ligation-Induced Pain in Rats. 2007 , 32, 209-216	
1528	Anti-allodynic effects of peripheral delta opioid receptors in neuropathic pain. 2007 , 127, 84-93	97
1527	Activation of cyclin-dependent kinase 5 (Cdk5) in primary sensory and dorsal horn neurons by peripheral inflammation contributes to heat hyperalgesia. 2007 , 127, 109-20	59

1526	NMDA receptor-independent synaptic plasticity in the central amygdala in the rat model of neuropathic pain. 2007 , 127, 161-72	172
1525	Catechol-O-methyltransferase inhibition increases pain sensitivity through activation of both beta2- and beta3-adrenergic receptors. 2007 , 128, 199-208	207
1524	Spinal NK-1 receptor expressing neurons mediate opioid-induced hyperalgesia and antinociceptive tolerance via activation of descending pathways. 2007 , 129, 35-45	83
1523	Prolonged maintenance of capsaicin-induced hyperalgesia by brief daily vibration stimuli. 2007 , 129, 93-101	10
1522	Anatomical and functional analysis of aquaporin 1, a water channel in primary afferent neurons. 2007 , 131, 8-20	69
1521	Neuropeptide Y acts at Y1 receptors in the rostral ventral medulla to inhibit neuropathic pain. 2007 , 131, 83-95	36
1520	Reactive oxygen species (ROS) are involved in enhancement of NMDA-receptor phosphorylation in animal models of pain. 2007 , 131, 262-271	180
1519	Modulation of persistent nociceptive inputs in the anterior pretectal nucleus of the rat. 2007 , 132, 42-52	19
1518	Oral and spinal melatonin reduces tactile allodynia in rats via activation of MT2 and opioid receptors. 2007 , 132, 273-280	66
1517	The role of reactive oxygen species in capsaicin-induced mechanical hyperalgesia and in the activities of dorsal horn neurons. 2007 , 133, 9-17	129
1516	Engineering an endomorphin-2 gene for use in neuropathic pain therapy. 2007 , 133, 29-38	53
1515	D-cycloserine reduces neuropathic pain behavior through limbic NMDA-mediated circuitry. 2007 , 132, 108-23	85
1514	Morphine treatment accelerates sarcoma-induced bone pain, bone loss, and spontaneous fracture in a murine model of bone cancer. 2007 , 132, 154-68	82
1513	Attenuated cold sensitivity in TRPM8 null mice. 2007 , 54, 379-86	569
1512	Tumor necrosis factor-alpha induces long-term potentiation of C-fiber evoked field potentials in spinal dorsal horn in rats with nerve injury: the role of NF-kappa B, JNK and p38 MAPK. 2007 , 52, 708-15	111
1511	Peripheral kinin B(1) and B(2) receptor-operated mechanisms are implicated in neuropathic nociception induced by spinal nerve ligation in rats. 2007 , 53, 48-57	28
1510	Pregabalin, S-(+)-3-isobutylgaba, activates the descending noradrenergic system to alleviate neuropathic pain in the mouse partial sciatic nerve ligation model. 2007 , 53, 842-53	70
1509	Role of peripheral hyperpolarization-activated cyclic nucleotide-modulated channel pacemaker channels in acute and chronic pain models in the rat. 2007 , 144, 1477-85	85

1508	Roles of different subtypes of opioid receptors in mediating the ventrolateral orbital cortex opioid-induced inhibition of mirror-neuropathic pain in the rat. 2007 , 144, 1486-94	19
1507	Further characterization of a rat model of varicella zoster virus-associated pain: Relationship between mechanical hypersensitivity and anxiety-related behavior, and the influence of analgesic drugs. 2007 , 144, 1495-508	106
1506	Small interfering RNA-mediated selective knockdown of Na(V)1.8 tetrodotoxin-resistant sodium channel reverses mechanical allodynia in neuropathic rats. 2007 , 146, 812-21	99
1505	Continuous intra-arterial application of substance P induces signs and symptoms of experimental complex regional pain syndrome (CRPS) such as edema, inflammation and mechanical pain but no thermal pain. 2007 , 148, 757-65	22
1504	Mice overexpressing chemokine ligand 2 (CCL2) in astrocytes display enhanced nociceptive responses. 2007 , 149, 706-14	100
1503	Plasticity in intact A delta- and C-fibers contributes to cold hypersensitivity in neuropathic rats. 2007 , 150, 182-93	35
1502	Effects of carrageenan and morphine on acute inflammation and pain in Lewis and Fischer rats. 2007 , 21, 68-78	24
1501	Clonidine reduces hypersensitivity and alters the balance of pro- and anti-inflammatory leukocytes after local injection at the site of inflammatory neuritis. 2007 , 21, 569-80	28
1500	Interleukin-6 mediates low-threshold mechanical allodynia induced by intrathecal HIV-1 envelope glycoprotein gp120. 2007 , 21, 660-7	49
1499	Intrathecal interleukin-10 gene therapy attenuates paclitaxel-induced mechanical allodynia and proinflammatory cytokine expression in dorsal root ganglia in rats. 2007 , 21, 686-98	220
1498	Role of the CX3CR1/p38 MAPK pathway in spinal microglia for the development of neuropathic pain following nerve injury-induced cleavage of fractalkine. 2007 , 21, 642-51	272
1497	Enhanced 3,5-dihydroxyphenylglycine-induced sustained nociceptive behaviors in rats with neuropathy or chronic inflammation. 2007 , 184, 150-6	6
1496	Increased TRPA1, TRPM8, and TRPV2 expression in dorsal root ganglia by nerve injury. 2007 , 358, 1058-64	108
1495	Expression profiles of BDNF splice variants in cultured DRG neurons stimulated with NGF. 2007 , 362, 682-8	19
1494	Intravenous paclitaxel administration in the rat induces a peripheral sensory neuropathy characterized by macrophage infiltration and injury to sensory neurons and their supporting cells. 2007 , 203, 42-54	203
1493	Antinociceptive effect of cannabinoid agonist WIN 55,212-2 in rats with a spinal cord injury. 2007 , 204, 454-7	31
1492	Activation of transcription factor c-jun in dorsal root ganglia induces VIP and NPY upregulation and contributes to the pathogenesis of neuropathic pain. 2007 , 204, 467-72	19
1491	At-level neuropathic pain is induced by lumbosacral ventral root avulsion injury and ameliorated by root reimplantation into the spinal cord. 2007 , 204, 273-82	26

1490	The role of uninjured C-afferents and injured afferents in the generation of mechanical hypersensitivity after partial peripheral nerve injury in the rat. 2007 , 204, 288-98	29
1489	p38 activation in uninjured primary afferent neurons and in spinal microglia contributes to the development of neuropathic pain induced by selective motor fiber injury. 2007 , 204, 355-65	89
1488	Peri-sciatic administration of recombinant rat TNF-alpha induces mechanical allodynia via upregulation of TNF-alpha in dorsal root ganglia and in spinal dorsal horn: the role of NF-kappa B pathway. 2007 , 205, 471-84	82
1487	The neurocytokine, interleukin-6, corrects nerve dysfunction in experimental diabetes. 2007 , 207, 23-9	19
1486	Activation of phosphatidylinositol 3-kinase and protein kinase B/Akt in dorsal root ganglia and spinal cord contributes to the neuropathic pain induced by spinal nerve ligation in rats. 2007 , 206, 269-79	99
1485	Long-term synaptic plasticity in the spinal dorsal horn and its modulation by electroacupuncture in rats with neuropathic pain. 2007 , 208, 323-32	100
1484	Antinociceptive and anti-inflammatory properties from the bulbs of <i>Cipura paludosa</i> Aubl. 2007 , 112, 19-25	28
1483	Role of spinal voltage-dependent calcium channel alpha 2 delta-1 subunit in the expression of a neuropathic pain-like state in mice. 2007 , 80, 2015-24	13
1482	Antihyperalgesic effects induced by the IL-1 receptor antagonist anakinra and increased IL-1beta levels in inflamed and osteosarcoma-bearing mice. 2007 , 81, 673-82	53
1481	Antihyperalgesic effects of loperamide in a model of rat neuropathic pain are mediated by peripheral delta-opioid receptors. 2007 , 411, 143-6	42
1480	Optimizing the transplant dose of a human neuronal cell line graft to treat SCI pain in the rat. 2007 , 414, 121-5	15
1479	The peripheral role of group I metabotropic glutamate receptors on nociceptive behaviors in rats with knee joint inflammation. 2007 , 416, 123-7	22
1478	Pharmacological validation of a model of cystitis pain in the mouse. 2007 , 421, 250-2	44
1477	FK506 reduces the severity of cutaneous hypersensitivity in rats with a spinal cord contusion. 2007 , 58, 95-9	14
1476	Involvement of substance P and calcitonin gene-related peptide in development and maintenance of neuropathic pain from spinal nerve injury model of rat. 2007 , 58, 245-9	78
1475	Encyclopedia of Pain. 2007 , 98-98	
1474	Morphine reduces local cytokine expression and neutrophil infiltration after incision. 2007 , 3, 28	87
1473	Transformation of the output of spinal lamina I neurons after nerve injury and microglia stimulation underlying neuropathic pain. 2007 , 3, 27	189

1472	Actions of N-arachidonyl-glycine in a rat inflammatory pain model. 2007 , 3, 24	55
1471	Prostaglandin E2 potentiation of P2X3 receptor mediated currents in dorsal root ganglion neurons. 2007 , 3, 22	53
1470	FM1-43 is a permeant blocker of mechanosensitive ion channels in sensory neurons and inhibits behavioural responses to mechanical stimuli. 2007 , 3, 1	49
1469	4-Hydroxynonenal, an endogenous aldehyde, causes pain and neurogenic inflammation through activation of the irritant receptor TRPA1. 2007 , 104, 13519-24	584
1468	A critical role of toll-like receptor 2 in nerve injury-induced spinal cord glial cell activation and pain hypersensitivity. 2007 , 282, 14975-83	225
1467	Social and environmental enrichment improves sensory and motor recovery after severe contusive spinal cord injury in the rat. 2007 , 24, 1761-72	64
1466	Development of orally bioavailable and CNS penetrant biphenylaminocyclopropane carboxamide bradykinin B1 receptor antagonists. 2007 , 50, 272-82	47
1465	Pathophysiology of pain. 2007 , 91, 1-12	49
1464	Chronic intrathecal infusion of minocycline prevents the development of spinal-nerve ligation-induced pain in rats. 2007 , 32, 209-16	42
1463	Histaminergic involvement in neuropathic pain produced by partial ligation of the sciatic nerve in rats. 2007 , 32, 124-9	22
1462	Subarachnoid transplant of a human neuronal cell line attenuates chronic allodynia and hyperalgesia after excitotoxic spinal cord injury in the rat. 2007 , 8, 33-50	46
1461	Walker 256 tumor-bearing rats as a model to study cancer pain. 2007 , 8, 412-21	49
1460	Age- and sex-specific nociceptive response to endothelin-1. 2007 , 8, 657-66	20
1459	Lumbar sympathectomy attenuates cold allodynia but not mechanical allodynia and hyperalgesia in rats with spared nerve injury. 2007 , 8, 931-7	23
1458	Analgesic, Anti-inflammatory, and Anti-pyretic Activity. 2007 , 983-1116	2
1457	The Combined Antiallodynic Effect of Gabapentin and Milnacipran in a Rat Neuropathic Pain Model. 2007 , 20, 8	3
1456	Antiallodynic effect of pregabalin in rat models of sympathetically maintained and sympathetic independent neuropathic pain. 2007 , 48, 41-7	28
1455	The Effect of Treatment with Intrathecal Ginsenosides in a Rat Model of Postoperative Pain. 2007 , 20, 100	12

1454	Modification of antiallodynic and antinociceptive effects of morphine by peripheral and central action of fluoxetine in a neuropathic mice model. 2007 , 58, 369-79		12
1453	Comparison of pre- versus post-incision administration of intraplantar indomethacin and MK886 in a rat model of postoperative pain. 2007 , 40, 1141-7		2
1452	Evaluation of NR2B peptide as subunit vaccines against experimental neuropathic pain. 2007 , 120, 643-647		2
1451	Modification of classical neurochemical markers in identified primary afferent neurons with A β -, A δ -, and C-fibers after chronic constriction injury in mice. 2007 , 502, 325-36		120
1450	Spinal AMPA receptor inhibition attenuates mechanical allodynia and neuronal hyperexcitability following spinal cord injury in rats. 2007 , 85, 2352-9		31
1449	Discovery of a novel class of benzazepinone Na(v)1.7 blockers: potential treatments for neuropathic pain. 2007 , 17, 4630-4		62
1448	Benzazepinone Nav1.7 blockers: potential treatments for neuropathic pain. 2007 , 17, 6172-7		39
1447	Synergistic interactions between the dual serotonergic, noradrenergic reuptake inhibitor duloxetine and the non-steroidal anti-inflammatory drug ibuprofen in inflammatory pain in rodents. 2007 , 11, 208-15		23
1446	Peripheral endothelin A receptor antagonism attenuates carcinoma-induced pain. 2007 , 11, 406-14		53
1445	Behavioral and sensory changes after direct ischemia-reperfusion injury in rats. 2007 , 11, 677-84		15
1444	Spinal administration of a delta opioid receptor agonist attenuates hyperalgesia and allodynia in a rat model of neuropathic pain. 2007 , 11, 685-93		54
1443	Differences in inflammatory pain in nNOS-, iNOS- and eNOS-deficient mice. 2007 , 11, 810-8		77
1442	The CatWalk method: a detailed analysis of behavioral changes after acute inflammatory pain in the rat. <i>Journal of Neuroscience Methods</i> , 2007 , 163, 9-16	3	92
1441	Validation of an automated punctate mechanical stimuli delivery system designed for fMRI studies in rodents. <i>Journal of Neuroscience Methods</i> , 2007 , 163, 31-7	3	6
1440	GDNF-chitosan blended nerve guides: a functional study. 2007 , 1, 360-7		49
1439	Gene transfer to interfere with TNF α signaling in neuropathic pain. 2007 , 14, 1010-6		92
1438	Prevention of neuropathic pain in an animal model of spare nerve injury following oral immunization with recombinant adenovirus serotype 5-mediated NR2B gene transfer. 2007 , 14, 1681-7		12
1437	Activation of peripheral cannabinoid CB1 and CB2 receptors suppresses the maintenance of inflammatory nociception: a comparative analysis. 2007 , 150, 153-63		80

1436	Intraplantar PGE2 causes nociceptive behaviour and mechanical allodynia: the role of prostanoid E receptors and protein kinases. 2007 , 150, 727-37	70
1435	The kinin B1 receptor antagonist SSR240612 reverses tactile and cold allodynia in an experimental rat model of insulin resistance. 2007 , 152, 280-7	25
1434	Actions of the endocannabinoid transport inhibitor AM404 in neuropathic and inflammatory pain models. 2007 , 34, 1186-90	33
1433	Local administration of N-acetylaspartylglutamate (NAAG) peptidase inhibitors is analgesic in peripheral pain in rats. 2007 , 25, 147-58	45
1432	Episodic dural stimulation in awake rats: a model for recurrent headache. 2007 , 47, 1026-36	155
1431	A comparison of the glutamate release inhibition and anti-allodynic effects of gabapentin, lamotrigine, and riluzole in a model of neuropathic pain. 2007 , 100, 1289-99	85
1430	Markedly attenuated acute and chronic pain responses in mice lacking adenylyl cyclase-5. 2007 , 6, 120-7	27
1429	Neuronal nitric oxide synthase is upregulated in a subset of primary sensory afferents after nerve injury which are necessary for analgesia from alpha2-adrenoceptor stimulation. 2007 , 1127, 52-8	13
1428	Involvement of cellular prion protein in the nociceptive response in mice. 2007 , 1151, 84-90	8
1427	Behavioral characterization and effect of clinical drugs in a rat model of pain following spinal cord compression. 2007 , 1185, 117-28	41
1426	Neurochemical evidence that supraspinally administered gabapentin activates the descending noradrenergic system after peripheral nerve injury. 2007 , 556, 69-74	44
1425	Altered antinociceptive efficacy of tramadol over time in rats with painful peripheral neuropathy. 2007 , 559, 32-7	24
1424	Acute and chronic administration of the cannabinoid receptor agonist CP 55,940 attenuates tumor-evoked hyperalgesia. 2007 , 558, 73-87	44
1423	Anti-nociceptive and anti-allodynic effects of a high affinity NOP hexapeptide [Ac-RY(3-Cl)YRWR-NH ₂] (Syn 1020) in rodents. 2007 , 560, 29-35	17
1422	Treatment with the xanthine oxidase inhibitor, allopurinol, improves nerve and vascular function in diabetic rats. 2007 , 561, 63-71	64
1421	Minocycline and pentoxifylline attenuate allodynia and hyperalgesia and potentiate the effects of morphine in rat and mouse models of neuropathic pain. 2007 , 560, 142-9	180
1420	Antinociceptive efficacy of lacosamide in rat models for tumor- and chemotherapy-induced cancer pain. 2007 , 565, 98-104	42
1419	Possible involvement of spinal noradrenergic mechanisms in the antiallodynic effect of intrathecally administered 5-HT _{2C} receptor agonists in the rats with peripheral nerve injury. 2007 , 567, 89-94	26

1418	Subcutaneous, intrathecal and periaqueductal grey administration of asimadoline and ICI-204448 reduces tactile allodynia in the rat. 2007 , 573, 75-83	23
1417	Altered hippocampal long-term potentiation after peripheral nerve injury in mice. 2007 , 574, 127-32	50
1416	Melatonin reduces formalin-induced nociception and tactile allodynia in diabetic rats. 2007 , 577, 203-10	54
1415	Pain mechanisms and their implication for the management of pain in farm and companion animals. 2007 , 174, 227-39	68
1414	Involvement of kinin B1 receptor and oxidative stress in sensory abnormalities and arterial hypertension in an experimental rat model of insulin resistance. 2007 , 41, 375-87	36
1413	Neutrophils invade lumbar dorsal root ganglia after chronic constriction injury of the sciatic nerve. 2007 , 184, 164-71	61
1412	Dissociation of spinal microglia morphological activation and peripheral inflammation in inflammatory pain models. 2007 , 192, 40-8	58
1411	The antipsychotic drug, fluphenazine, effectively reverses mechanical allodynia in rat models of neuropathic pain. 2008 , 195, 559-68	15
1410	Microglial activation in different models of peripheral nerve injury of the rat. 2007 , 38, 245-51	13
1409	Depletion of vesicular zinc in dorsal horn of spinal cord causes increased neuropathic pain in mice. 2008 , 21, 151-8	18
1408	Local loperamide inhibits thermal hyperalgesia but not mechanical allodynia induced by intratibial inoculation of melanoma cells in mice. 2008 , 28, 981-90	13
1407	Distinct responses of DREAM to electroacupuncture stimulation with different frequencies during physiological and inflammatory conditions in rats. 2008 , 33, 2070-7	9
1406	Mu-opioid receptor in the nucleus submedius: involvement in opioid-induced inhibition of mirror-image allodynia in a rat model of neuropathic pain. 2008 , 33, 2134-41	8
1405	Interleukin-1 beta of Red nucleus involved in the development of allodynia in spared nerve injury rats. 2008 , 188, 379-84	33
1404	Urinary bladder function and somatic sensitivity in vasoactive intestinal polypeptide (VIP)-/- mice. 2008 , 36, 175-87	39
1403	Electrospun micro- and nanofiber tubes for functional nervous regeneration in sciatic nerve transections. 2008 , 8, 39	247
1402	Nuclear factor-kappaB decoy amelioration of spinal cord injury-induced inflammation and behavior outcomes. 2008 , 86, 566-80	54
1401	CNS-infiltrating CD4+ T lymphocytes contribute to murine spinal nerve transection-induced neuropathic pain. 2008 , 38, 448-58	159

1400	Lyn tyrosine kinase is required for P2X(4) receptor upregulation and neuropathic pain after peripheral nerve injury. 2008 , 56, 50-8	84
1399	Activation of dorsal horn microglia contributes to diabetes-induced tactile allodynia via extracellular signal-regulated protein kinase signaling. 2008 , 56, 378-86	133
1398	Temporal changes of astrocyte activation and glutamate transporter-1 expression in the spinal cord after spinal nerve ligation-induced neuropathic pain. 2008 , 291, 513-8	38
1397	Early behavioral and histological outcomes following a novel traumatic partial nerve lesion. <i>Journal of Neuroscience Methods</i> , 2008 , 172, 236-44	3 5
1396	Effect of peripheral endothelin-1 concentration on carcinoma-induced pain in mice. 2008 , 12, 293-300	62
1395	Imidazopyridines: a novel class of hNav1.7 channel blockers. 2008 , 18, 1696-701	30
1394	Central action of prosaptide TX14(A) against gp120-induced allodynia in rats. 2008 , 12, 76-81	6
1393	Neuregulin 1 is a pronociceptive cytokine that is regulated by progesterone in the spinal cord: implications for sex specific pain modulation. 2008 , 12, 94-103	23
1392	Expression of c-Fos in the parabrachial nucleus following peripheral nerve injury in rats. 2008 , 12, 172-9	7
1391	Sciatic nerve cuffing in mice: a model of sustained neuropathic pain. 2008 , 12, 591-9	96
1390	Morphine-induced depression of spinal excitation is not altered following acute disruption of GABA(A) or GABA(B) receptor activity. 2008 , 12, 677-85	5
1389	Reduced thermal hyperalgesia and enhanced peripheral nerve injury after hind paw inflammation in mice lacking the serotonin-transporter. 2008 , 12, 790-7	24
1388	Chronic, but not acute, tricyclic antidepressant treatment alleviates neuropathic allodynia after sciatic nerve cuffing in mice. 2008 , 12, 1008-17	62
1387	HC-030031, a TRPA1 selective antagonist, attenuates inflammatory- and neuropathy-induced mechanical hypersensitivity. 2008 , 4, 48	313
1386	Design and synthesis of a novel series of N-alkyl isatin acylhydrazone derivatives that act as selective cannabinoid receptor 2 agonists for the treatment of neuropathic pain. 2008 , 51, 4932-47	48
1385	Inhibition of fatty acid amide hydrolase produces PPAR-alpha-mediated analgesia in a rat model of inflammatory pain. 2008 , 155, 1297-306	71
1384	MDA7: a novel selective agonist for CB2 receptors that prevents allodynia in rat neuropathic pain models. 2008 , 155, 1104-16	57
1383	Spinal cord injury-induced attenuation of GABAergic inhibition in spinal dorsal horn circuits is associated with down-regulation of the chloride transporter KCC2 in rat. 2008 , 586, 5701-15	100

1382	Differential effects of cannabinoid receptor agonists on regional brain activity using pharmacological MRI. 2008 , 153, 367-79	74
1381	In vitro and in vivo characterization of A-796260: a selective cannabinoid CB2 receptor agonist exhibiting analgesic activity in rodent pain models. 2008 , 153, 390-401	115
1380	Gabapentin produces PKA-dependent pre-synaptic inhibition of GABAergic synaptic transmission in LC neurons following partial nerve injury in mice. 2008 , 105, 933-42	43
1379	Intra-neural administration of fractalkine attenuates neuropathic pain-related behaviour. 2008 , 106, 640-9	34
1378	JAK/STAT3 pathway is activated in spinal cord microglia after peripheral nerve injury and contributes to neuropathic pain development in rat. 2008 , 107, 50-60	164
1377	Contribution of the Reelin signaling pathways to nociceptive processing. 2008 , 27, 523-37	16
1376	Evidence for a novel functional role of cannabinoid CB(2) receptors in the thalamus of neuropathic rats. 2008 , 27, 1722-30	70
1375	Non-stereoselective reversal of neuropathic pain by naloxone and naltrexone: involvement of toll-like receptor 4 (TLR4). 2008 , 28, 20-9	297
1374	Expression changes of multiple brain-derived neurotrophic factor transcripts in selective spinal nerve ligation model and complete Freund's adjuvant model. 2008 , 1206, 13-9	21
1373	Effects of selective spinal nerve ligation on acetic acid-induced nociceptive responses and ASIC3 immunoreactivity in the rat dorsal root ganglion. 2008 , 1219, 26-31	30
1372	The activation of extracellular signal-regulated protein kinase 5 in spinal cord and dorsal root ganglia contributes to inflammatory pain. 2008 , 1215, 76-86	23
1371	Comparison of icilin- and cold-evoked responses of spinal neurones, and their modulation of mechanical activity, in a model of neuropathic pain. 2008 , 1215, 87-96	26
1370	A comparison of spinal Iba1 and GFAP expression in rodent models of acute and chronic pain. 2008 , 1219, 116-26	143
1369	The quantitative assessment of functional impairment and its correlation to infarct volume in rats with transient middle cerebral artery occlusion. 2008 , 1230, 303-9	14
1368	Rats with chronic post-ischemia pain exhibit an analgesic sensitivity profile similar to human patients with complex regional pain syndrome--type I. 2008 , 583, 97-102	22
1367	Secondary hyperalgesia in the rat first degree burn model is independent of spinal cyclooxygenase and nitric oxide synthase. 2008 , 587, 118-23	5
1366	Ameliorative effects of amiloride and pralidoxime in chronic constriction injury and vincristine induced painful neuropathy in rats. 2008 , 587, 104-11	86
1365	Blockade of sensory abnormalities and kinin B(1) receptor expression by N-acetyl-L-cysteine and ramipril in a rat model of insulin resistance. 2008 , 589, 66-72	39

1364	Pyritinol reduces nociception and oxidative stress in diabetic rats. 2008 , 590, 170-6	11
1363	Crotalphine induces potent antinociception in neuropathic pain by acting at peripheral opioid receptors. 2008 , 594, 84-92	36
1362	Multiplicative interactions to enhance gabapentin to treat neuropathic pain. 2008 , 598, 21-6	28
1361	Effects of norketamine enantiomers in rodent models of persistent pain. 2008 , 90, 676-85	44
1360	Role of p38 mitogen activated protein kinase in a model of osteosarcoma-induced pain. 2008 , 90, 664-75	26
1359	Interaction between morphine and norketamine enantiomers in rodent models of nociception. 2008 , 90, 769-77	16
1358	Spinal nerve ligation reduces nitric oxide synthase activity and expression: effect of resveratrol. 2008 , 90, 742-7	20
1357	Y1 receptor knockout increases nociception and prevents the anti-allodynic actions of NPY. 2008 , 24, 885-91	21
1356	Contralateral neuropathic pain following a surgical model of unilateral nerve injury in rats. 2008 , 33, 211-6	22
1355	Modulation of mechanical and thermal nociceptive sensitivity in the laboratory mouse by behavioral state. 2008 , 9, 174-84	62
1354	Sympathetic vasoconstrictor antagonism and vasodilatation relieve mechanical allodynia in rats with chronic postischemia pain. 2008 , 9, 423-33	32
1353	Animal Models and Neuropathic Pain. 2008 , 857-864	1
1352	New frontiers in assessing pain and analgesia in laboratory animals. 2008 , 3, 1099-108	11
1351	A model of cystitis pain in the mouse. 2008 , Chapter 5, Unit 5.52	5
1350	Gene transfer of GLT-1, a glial glutamate transporter, into the spinal cord by recombinant adenovirus attenuates inflammatory and neuropathic pain in rats. 2008 , 4, 65	70
1349	The role of TRPV1 in different subtypes of dorsal root ganglion neurons in rat chronic inflammatory nociception induced by complete Freund's adjuvant. 2008 , 4, 61	127
1348	Brain derived neurotrophic factor (BDNF) contributes to the pain hypersensitivity following surgical incision in the rats. 2008 , 4, 27	71
1347	Chronic morphine administration enhances nociceptive sensitivity and local cytokine production after incision. 2008 , 4, 7	54

1346	Chronic Pain and Microglia: The Role of ATP. 2008 , 55-67	13
1345	Investigation of TRPV1 loss-of-function phenotypes in transgenic shRNA expressing and knockout mice. 2008 , 37, 579-89	61
1344	MMPs initiate Schwann cell-mediated MBP degradation and mechanical nociception after nerve damage. 2008 , 39, 619-27	86
1343	Ameliorative effects of <i>Ocimum sanctum</i> in sciatic nerve transection-induced neuropathy in rats. 2008 , 120, 56-62	59
1342	A novel human foamy virus mediated gene transfer of GAD67 reduces neuropathic pain following spinal cord injury. 2008 , 432, 13-8	35
1341	Local application of capsaicin alleviates mechanical hyperalgesia after spinal nerve transection. 2008 , 433, 199-204	14
1340	Differential galanin upregulation in dorsal root ganglia and spinal cord after graded single ligature nerve constriction of the rat sciatic nerve. 2008 , 35, 94-100	24
1339	Role of kinin B1 and B2 receptors in a rat model of neuropathic pain. 2008 , 8, 188-96	30
1338	Actions of N-arachidonyl-glycine in a rat neuropathic pain model. 2008 , 54, 189-93	75
1337	Down-regulation of NR2B receptors partially contributes to analgesic effects of Gentiopicroside in persistent inflammatory pain. 2008 , 54, 1175-81	55
1336	Depression of C fibre-evoked spinal field potentials by the spinal delta opioid receptor is enhanced in the spinal nerve ligation model of neuropathic pain: involvement of the mu-subtype. 2008 , 55, 1376-82	7
1335	Reduced potassium-chloride co-transporter expression in spinal cord dorsal horn neurons contributes to inflammatory pain hypersensitivity in rats. 2008 , 152, 502-10	85
1334	Pharmacological characterization and gene expression profiling of an L5/L6 spinal nerve ligation model for neuropathic pain in mice. 2008 , 153, 492-500	42
1333	Activation of microglia and p38 mitogen-activated protein kinase in the dorsal column nucleus contributes to tactile allodynia following peripheral nerve injury. 2008 , 153, 1245-55	74
1332	Local erythropoietin signaling enhances regeneration in peripheral axons. 2008 , 154, 767-83	55
1331	Group II metabotropic glutamate receptor activation attenuates peripheral sensitization in inflammatory states. 2008 , 154, 754-66	29
1330	Intact Adelta-fibers up-regulate transient receptor potential A1 and contribute to cold hypersensitivity in neuropathic rats. 2008 , 154, 1054-66	56
1329	Fatty acid amide hydrolase inhibition enhances the anti-allodynic actions of endocannabinoids in a model of acute pain adapted for the mouse. 2008 , 154, 1554-61	22

1328	GABAA receptor modulation in dorsal root ganglia in vivo affects chronic pain after nerve injury. 2008 , 154, 1539-53	50
1327	Spinal nerve ligation-induced activation of nuclear factor kappaB is facilitated by prostaglandins in the affected spinal cord and is a critical step in the development of mechanical allodynia. 2008 , 155, 902-13	13
1326	Anatomical localization and expression pattern for the NMDA-2D receptor subunit in a rat model of neuropathic pain. 2008 , 155, 492-502	9
1325	Inflammatory pain-induced signaling events following a conditional deletion of the N-methyl-D-aspartate receptor in spinal cord dorsal horn. 2008 , 155, 948-58	37
1324	Tactile allodynia can occur in the spared nerve injury model in the rat without selective loss of GABA or GABA(A) receptors from synapses in laminae I-II of the ipsilateral spinal dorsal horn. 2008 , 156, 193-202	73
1323	Analgesic effects of intrathecally administered 26RFa, an intrinsic agonist for GPR103, on formalin test and carrageenan test in rats. 2008 , 157, 214-22	28
1322	Selective stimulation of either tumor necrosis factor receptor differentially induces pain behavior in vivo and ectopic activity in sensory neurons in vitro. 2008 , 157, 414-23	72
1321	Tramadol and several anticonvulsants synergize in attenuating nerve injury-induced allodynia. 2008 , 134, 254-262	28
1320	Characterization of cell proliferation in rat spinal cord following peripheral nerve injury and the relationship with neuropathic pain. 2008 , 135, 37-47	149
1319	Inflammation-induced changes in rostral ventromedial medulla mu and kappa opioid receptor mediated antinociception. 2008 , 136, 320-330	25
1318	Brain derived nerve growth factor induces spinal noradrenergic fiber sprouting and enhances clonidine analgesia following nerve injury in rats. 2008 , 136, 348-355	72
1317	Intrathecal neuropeptide Y reduces behavioral and molecular markers of inflammatory or neuropathic pain. 2008 , 137, 352-365	61
1316	Axonal accumulation of hyperpolarization-activated cyclic nucleotide-gated cation channels contributes to mechanical allodynia after peripheral nerve injury in rat. 2008 , 137, 495-506	82
1315	Interaction between protein kinase D1 and transient receptor potential V1 in primary sensory neurons is involved in heat hypersensitivity. 2008 , 137, 574-588	16
1314	Norepinephrine-induced nociception and vasoconstrictor hypersensitivity in rats with chronic post-ischemia pain. 2008 , 137, 640-651	48
1313	Spread of excitation across modality borders in spinal dorsal horn of neuropathic rats. 2008 , 135, 300-310	55
1312	Peripherally acting mu-opioid receptor agonist attenuates neuropathic pain in rats after L5 spinal nerve injury. 2008 , 138, 318-329	91
1311	Species and strain differences in rodent sciatic nerve anatomy: implications for studies of neuropathic pain. 2008 , 136, 188-201	192

1310	Propentofylline attenuates allodynia, glial activation and modulates GABAergic tone after spinal cord injury in the rat. 2008 , 138, 410-422	108
1309	Endogenous kappa-opioid receptor systems inhibit hyperalgesia associated with localized peripheral inflammation. 2008 , 138, 423-439	26
1308	Thermal nociception and TRPV1 function are attenuated in mice lacking the nucleotide receptor P2Y2. 2008 , 138, 484-496	67
1307	Interleukin-1 alpha has antiallodynic and antihyperalgesic activities in a rat neuropathic pain model. 2008 , 138, 587-597	83
1306	NK-1 receptors in the rostral ventromedial medulla contribute to hyperalgesia produced by intraplantar injection of capsaicin. 2008 , 139, 34-46	22
1305	Glutamate receptor ligands attenuate allodynia and hyperalgesia and potentiate morphine effects in a mouse model of neuropathic pain. 2008 , 139, 117-126	98
1304	Lack of TRPV1 inhibits cystitis-induced increased mechanical sensitivity in mice. 2008 , 139, 158-167	84
1303	Injury discharges regulate calcium channel alpha-2-delta-1 subunit upregulation in the dorsal horn that contributes to initiation of neuropathic pain. 2008 , 139, 358-366	61
1302	Profound reduction of somatic and visceral pain in mice by intrathecal administration of the anti-migraine drug, sumatriptan. 2008 , 139, 533-540	46
1301	Activation of peripheral ephrinBs/EphBs signaling induces hyperalgesia through a MAPKs-mediated mechanism in mice. 2008 , 139, 617-631	26
1300	Allodynia and hyperalgesia in diabetic rats are mediated by GABA and depletion of spinal potassium-chloride co-transporters. 2008 , 140, 48-57	109
1299	A 2A adenosine receptor regulates glia proliferation and pain after peripheral nerve injury. 2008 , 140, 95-103	53
1298	Analgesic effect in rodents of native and recombinant Ph alpha 1beta toxin, a high-voltage-activated calcium channel blocker isolated from armed spider venom. 2008 , 140, 115-126	76
1297	The persistence of a long-term negative affective state following the induction of either acute or chronic pain. 2008 , 140, 436-445	60
1296	Antinociceptive effect of the C-terminus of murine S100A9 protein on experimental neuropathic pain. 2008 , 29, 1806-14	11
1295	Pbx3 is required for normal locomotion and dorsal horn development. 2008 , 314, 23-39	19
1294	Tumor necrosis factor- α of Red nucleus involved in the development of neuropathic allodynia. 2008 , 77, 233-6	29
1293	Remote activation of microglia and pro-inflammatory cytokines predict the onset and severity of below-level neuropathic pain after spinal cord injury in rats. 2008 , 212, 337-47	184

1292	BDNF induces late-phase LTP of C-fiber evoked field potentials in rat spinal dorsal horn. 2008 , 212, 507-14	71
1291	Conditional deletion of the NMDA-NR1 receptor subunit gene in the central nucleus of the amygdala inhibits naloxone-induced conditioned place aversion in morphine-dependent mice. 2008 , 213, 57-70	28
1290	Proinflammatory cytokines oppose opioid-induced acute and chronic analgesia. 2008 , 22, 1178-89	237
1289	Characterization of a model of chronic orofacial hyperalgesia in the rat: contribution of NA(V) 1.8. 2008 , 9, 522-31	29
1288	Intrathecal rosiglitazone acts at peroxisome proliferator-activated receptor-gamma to rapidly inhibit neuropathic pain in rats. 2008 , 9, 639-49	79
1287	Oxycodone plus ultra-low-dose naltrexone attenuates neuropathic pain and associated mu-opioid receptor-Gs coupling. 2008 , 9, 700-13	59
1286	Spinal dynorphin and bradykinin receptors maintain inflammatory hyperalgesia. 2008 , 9, 1096-105	30
1285	Interferon-gamma is a critical modulator of CB(2) cannabinoid receptor signaling during neuropathic pain. 2008 , 28, 12136-45	106
1284	Functional gait evaluation of collagen chitosan nerve guides for sciatic nerve repair. 2008 , 14, 365-70	18
1283	Deficiency in complement C1q improves histological and functional locomotor outcome after spinal cord injury. 2008 , 28, 13876-88	56
1282	Crucial role of CB(2) cannabinoid receptor in the regulation of central immune responses during neuropathic pain. 2008 , 28, 12125-35	145
1281	Sulfasalazine blocks the development of tactile allodynia in diabetic rats. 2008 , 57, 2801-8	28
1280	Percutaneous pulsed radiofrequency reduces mechanical allodynia in a neuropathic pain model. 2008 , 107, 1406-11	24
1279	Evaluation tools and animal models of peripheral neuropathies. 2008 , 5, 72-108	11
1278	A new nerve growth factor-mimetic peptide active on neuropathic pain in rats. 2008 , 28, 2698-709	89
1277	Pain Testing in the Laboratory Mouse. 2008 , 549-560	5
1276	Sensory neuron targeting by self-complementary AAV8 via lumbar puncture for chronic pain. 2008 , 105, 1055-60	108
1275	Neuropathic pain memory is maintained by Rac1-regulated dendritic spine remodeling after spinal cord injury. 2008 , 28, 13173-83	90

1274	P2Y12 receptors in spinal microglia are required for neuropathic pain after peripheral nerve injury. 2008 , 28, 4949-56	214
1273	Activation of the neuronal extracellular signal-regulated kinase 2 in the spinal cord dorsal horn is required for complete Freund's adjuvant-induced pain hypersensitivity. 2008 , 28, 14087-96	44
1272	Disruption of cAMP and prostaglandin E2 transport by multidrug resistance protein 4 deficiency alters cAMP-mediated signaling and nociceptive response. 2008 , 73, 243-51	90
1271	Reversal of chronic inflammatory pain by acute inhibition of Ca ²⁺ /calmodulin-dependent protein kinase II. 2008 , 325, 267-75	68
1270	Experimental autoimmune prostatitis induces chronic pelvic pain. 2008 , 294, R1268-75	67
1269	Capsular ligament involvement in the development of mechanical hyperalgesia after facet joint loading: behavioral and inflammatory outcomes in a rodent model of pain. 2008 , 25, 1383-93	57
1268	Contralateral Neuropathic Pain Following a Surgical Model of Unilateral Nerve Injury in Rats. 2008 , 33, 211-216	43
1267	Zonisamide suppresses pain symptoms of formalin-induced inflammatory and streptozotocin-induced diabetic neuropathy. 2008 , 107, 213-20	23
1266	Growth factors as therapeutics for diabetic neuropathy. 2008 , 9, 47-59	60
1265	Effect of etanercept, a tumor necrosis factor-alpha inhibitor, on neuropathic pain in the rat chronic constriction injury model. 2008 , 33, 227-34	47
1264	Antiallodynic and antihyperalgesic effect of milnacipran in mice with spinal nerve ligation. 2008 , 106, 1309-15, table of contents	33
1263	Low-dose systemic bupivacaine prevents the development of allodynia after thoracotomy in rats. 2008 , 107, 1587-91	17
1262	The antiallodynic effect of Neurotropin is mediated via activation of descending pain inhibitory systems in rats with spinal nerve ligation. 2008 , 107, 1064-9	47
1261	Do corticosteroids produce additional benefit in nerve root infiltration for lumbar disc herniation?. 2008 , 33, 743-7	54
1260	Contralateral neuropathic pain and neuropathology in dorsal root ganglion and spinal cord following hemilateral nerve injury in rats. 2008 , 33, 1344-51	67
1259	Direct application of the TNF-alpha inhibitor, etanercept, does not affect CGRP expression and phenotypic change of DRG neurons following application of nucleus pulposus onto injured sciatic nerves in rats. 2008 , 33, 2403-8	18
1258	The Expression of the Ca ⁺⁺ Channel α 1 Subunit and TRPM8 in the Dorsal Root Ganglion of Sympathetically Maintained Pain and Sympathetic Independent Pain Rat Models. 2008 , 21, 11	1
1257	Effect of the Preoperative Intercostal Nerve Block in a Rat Model of Postthoracotomy Pain. 2008 , 21, 106	1

1256	Spinal Metabotropic Glutamate Receptors (mGluRs) are Involved in the Melittin-induced Nociception in Rats. 2008 , 12, 237-43	1
1255	The role of alpha 6 integrin in prostate cancer migration and bone pain in a novel xenograft model. 2008 , 3, e3535	41
1254	The interaction of gabapentin and N6-(2-phenylisopropyl)-adenosine R-(-)-isomer (R-PIA) on mechanical allodynia in rats with a spinal nerve ligation. 2008 , 23, 678-84	8
1253	Cannabinoid modulation of cutaneous Adelta nociceptors during inflammation. 2008 , 100, 2794-806	26
1252	Long-term Follow-up of Cutaneous Hypersensitivity in Rats with a Spinal Cord Contusion. 2008 , 12, 299-306	14
1251	The Role of the Peripheral Chemokine, CCL3, in Hyperalgesia following Peripheral Nerve Injury in the Rat. 2008 , 21, 187	
1250	[Effect of dehydroepiandrosterone on affected and unaffected hindlimb muscles in rats with neuropathic pain induced by unilateral peripheral nerve injury]. 2009 , 39, 632-40	1
1249	The Antiallodynic Effects of Intrathecal Zaprinas in Rats with Chronic Constriction Injury of the Sciatic Nerve. 2009 , 22, 16	1
1248	In vivo delta opioid receptor internalization controls behavioral effects of agonists. 2009 , 4, e5425	141
1247	Crosstalk between spinal astrocytes and neurons in nerve injury-induced neuropathic pain. 2009 , 4, e6973	76
1246	A Selective Role for alpha3 Subunit Glycine Receptors in Inflammatory Pain. 2009 , 2, 14	29
1245	(-)-Linalool attenuates allodynia in neuropathic pain induced by spinal nerve ligation in c57/bl6 mice. 2009 , 85, 221-35	26
1244	Acute inhibition of Ca ²⁺ /calmodulin-dependent protein kinase II reverses experimental neuropathic pain in mice. 2009 , 330, 650-9	50
1243	Proteomic analysis uncovers novel actions of the neurosecretory protein VGF in nociceptive processing. 2009 , 29, 13377-88	45
1242	Enduring reversal of neuropathic pain by a single intrathecal injection of adenosine 2A receptor agonists: a novel therapy for neuropathic pain. 2009 , 29, 14015-25	82
1241	Activation of TRPV1 in the spinal cord by oxidized linoleic acid metabolites contributes to inflammatory hyperalgesia. 2009 , 106, 18820-4	161
1240	Molecular mechanisms of lipoic acid modulation of T-type calcium channels in pain pathway. 2009 , 29, 9500-9	48
1239	Models and mechanisms of hyperalgesia and allodynia. 2009 , 89, 707-58	819

1238	Morphological and functional reorganization of rat medial prefrontal cortex in neuropathic pain. 2009 , 106, 2423-8	291
1237	Membrane-delimited coupling of TRPV1 and mGluR5 on presynaptic terminals of nociceptive neurons. 2009 , 29, 10000-9	57
1236	Deletion of the glutamate receptor 5 subunit of kainate receptors affects the development of morphine tolerance. 2009 , 328, 579-87	13
1235	Activation of extracellular signal-regulated protein kinases 5 in the spinal cord contributes to the neuropathic pain behaviors induced by CCI in rats. 2009 , 31, 1037-43	5
1234	Ifenprodil induced antinociception and decreased the expression of NR2B subunits in the dorsal horn after chronic dorsal root ganglia compression in rats. 2009 , 108, 1015-20	27
1233	Nonviral retrograde gene transfer of human hepatocyte growth factor improves neuropathic pain-related phenomena in rats. 2009 , 17, 42-50	23
1232	Neuropathic pain is maintained by brainstem neurons co-expressing opioid and cholecystokinin receptors. 2009 , 132, 778-87	73
1231	Involvement of JAK-STAT signaling/function after cyclophosphamide-induced bladder inflammation in female rats. 2009 , 297, F1038-44	26
1230	Vascular and neural dysfunctions in obese Zucker rats: effect of AVE7688. 2009 , 2009, 912327	14
1229	Intranasal insulin ameliorates experimental diabetic neuropathy. 2009 , 58, 934-45	63
1228	Blockade of endocannabinoid-degrading enzymes attenuates neuropathic pain. 2009 , 330, 902-10	250
1227	A role of SCN9A in human epilepsies, as a cause of febrile seizures and as a potential modifier of Dravet syndrome. 2009 , 5, e1000649	172
1226	Variable threshold of trigeminal cold-thermosensitive neurons is determined by a balance between TRPM8 and Kv1 potassium channels. 2009 , 29, 3120-31	146
1225	Selective T-type calcium channel blockade alleviates hyperalgesia in ob/ob mice. 2009 , 58, 2656-65	99
1224	Retrograde viral vector-mediated inhibition of pontospinal noradrenergic neurons causes hyperalgesia in rats. 2009 , 29, 12855-64	33
1223	The co-inhibitory molecule PD-1 modulates disease severity in a model for an inherited, demyelinating neuropathy. 2009 , 33, 96-103	13
1222	Increased pain and neurogenic inflammation in mice deficient of neutral endopeptidase. 2009 , 35, 177-83	28
1221	Beta2-adrenoceptors are essential for desipramine, venlafaxine or reboxetine action in neuropathic pain. 2009 , 33, 386-94	63

1220	Biogenic amine depletion causes chronic muscular pain and tactile allodynia accompanied by depression: A putative animal model of fibromyalgia. 2009 , 146, 26-33	138
1219	Tocotrienol ameliorates behavioral and biochemical alterations in the rat model of alcoholic neuropathy. 2009 , 145, 129-35	36
1218	Prefrontal cortex and spinal cord mediated anti-neuropathy and analgesia induced by sarcosine, a glycine-T1 transporter inhibitor. 2009 , 145, 176-83	23
1217	Peripheral and central sensitization in remote spinal cord regions contribute to central neuropathic pain after spinal cord injury. 2009 , 147, 265-76	159
1216	Vanilloid receptor TRPV1-mediated phosphorylation of ERK in murine adjuvant arthritis. 2009 , 17, 244-51	33
1215	Intra-articular magnesium sulfate (MgSO ₄) reduces experimental osteoarthritis and nociception: association with attenuation of N-methyl-D-aspartate (NMDA) receptor subunit 1 phosphorylation and apoptosis in rat chondrocytes. 2009 , 17, 1485-93	53
1214	Erythropoietin overrides the triggering effect of DNA platination products in a mouse model of cisplatin-induced neuropathy. 2009 , 10, 77	19
1213	Mammalian somatosensory mechanotransduction. 2009 , 19, 362-9	80
1212	Expression of monocyte chemoattractant protein-1 in rat dorsal root ganglia and spinal cord in experimental models of neuropathic pain. 2009 , 1251, 103-11	48
1211	Thalamic neuron hyperexcitability and enlarged receptive fields in the STZ model of diabetic pain. 2009 , 1268, 154-161	55
1210	Differential mediation of descending pain facilitation and inhibition by spinal 5HT-3 and 5HT-7 receptors. 2009 , 1280, 52-9	154
1209	Maternal separation decreases the stability of mushroom spines in adult mice somatosensory cortex. 2009 , 1294, 45-51	26
1208	Peripheral antinociceptive effects of mu- and delta-opioid receptor agonists in NOS2 and NOS1 knockout mice during chronic inflammatory pain. 2009 , 602, 41-9	31
1207	Activity of new NOP receptor ligands in a rat peripheral mononeuropathy model: potentiation of morphine anti-allodynic activity by NOP receptor antagonists. 2009 , 610, 49-54	13
1206	Gabapentin and pregabalin ameliorate mechanical hypersensitivity after spinal cord injury in mice. 2009 , 609, 65-8	22
1205	B vitamins alleviate indices of neuropathic pain in diabetic rats. 2009 , 612, 41-7	45
1204	Effect of 5-trifluoromethyl-4,5-dihydro-1H-pyrazoles on chronic inflammatory pain model in rats. 2009 , 616, 91-100	44
1203	Differential activation of spinal microglial and astroglial cells in a mouse model of peripheral neuropathic pain. 2009 , 623, 65-72	132

1202	2,3-Dihydro-1-benzofuran derivatives as a series of potent selective cannabinoid receptor 2 agonists: design, synthesis, and binding mode prediction through ligand-steered modeling. 2009 , 4, 1615-29	57
1201	Synergistic antiallodynic interaction of the metamizol-gabapentin combination. 2009 , 70, 386-394	2
1200	Critical role of microglial CD40 in the maintenance of mechanical hypersensitivity in a murine model of neuropathic pain. 2009 , 39, 3562-9	26
1199	Spinal p38 mitogen-activated protein kinase mediates allodynia induced by first-degree burn in the rat. 2009 , 87, 948-55	30
1198	Nuclear factor-kappa B decoy suppresses nerve injury and improves mechanical allodynia and thermal hyperalgesia in a rat lumbar disc herniation model. 2009 , 18, 1001-7	36
1197	Diabetic painful and insensate neuropathy: pathogenesis and potential treatments. 2009 , 6, 638-47	199
1196	Analgesic and Antipyretic Activities of a Novel Tetrapeptide in Rats. 2009 , 15, 293-301	3
1195	Design and assessment of a potent sodium channel blocking derivative of mexiletine for minimizing experimental neuropathic pain in several rat models. 2009 , 34, 1816-23	7
1194	Nerve growth factor of red nucleus involvement in pain induced by spared nerve injury of the rat sciatic nerve. 2009 , 34, 1612-8	27
1193	Effects of chronic inflammation and morphine tolerance on the expression of phospho-ERK 1/2 and phospho-P38 in the injured tissue. 2009 , 379, 315-23	9
1192	Proteome of synaptosome-associated proteins in spinal cord dorsal horn after peripheral nerve injury. 2009 , 9, 1241-53	38
1191	Characteristics and physiological role of hyperpolarization activated currents in mouse cold thermoreceptors. 2009 , 587, 1961-76	49
1190	Sensitization of lamina I spinoparabrachial neurons parallels heat hyperalgesia in the chronic constriction injury model of neuropathic pain. 2009 , 587, 2005-17	22
1189	Involvement of platelet-activating factor in ultraviolet B-induced hyperalgesia. 2009 , 129, 167-74	13
1188	Suppression of complete Freund's adjuvant-induced adjuvant arthritis by cobra toxin. 2009 , 30, 219-27	54
1187	Reactive oxygen species in rats with chronic post-ischemia pain. 2009 , 53, 648-56	36
1186	Spinal NK-1 receptor-expressing neurons and descending pathways support fentanyl-induced pain hypersensitivity in a rat model of postoperative pain. 2009 , 29, 727-37	56
1185	Glial activation in the rostroventromedial medulla promotes descending facilitation to mediate inflammatory hypersensitivity. 2009 , 30, 229-41	66

1184	Reduction of allodynia by intrathecal transplantation of microencapsulated porcine chromaffin cells. 2009 , 33, 240-9		13
1183	Modulation of neuropathic pain by a glial-derived factor. 2009 , 10, 1229-36		23
1182	Peripheral and central sites of action for the non-selective cannabinoid agonist WIN 55,212-2 in a rat model of post-operative pain. 2009 , 157, 645-55		18
1181	Intrathecal injection of the neurosteroid, DHEAS, produces mechanical allodynia in mice: involvement of spinal sigma-1 and GABA receptors. 2009 , 157, 666-73		27
1180	NCX 2057, a novel NO-releasing derivative of ferulic acid, suppresses inflammatory and nociceptive responses in in vitro and in vivo models. 2009 , 158, 569-79		23
1179	Inhibiting astrocytic activation: a novel analgesic mechanism of ketamine at the spinal level?. 2009 , 109, 1691-700		20
1178	Plasticity and emerging role of BKCa channels in nociceptive control in neuropathic pain. 2009 , 110, 352-62		73
1177	Chronic constriction injury induces aquaporin-2 expression in the dorsal root ganglia of rats. 2009 , 215, 498-505		23
1176	Effects of transgene-mediated endomorphin-2 in inflammatory pain. 2009 , 13, 380-6		46
1175	Comparison of dorsal root ganglion gene expression in rat models of traumatic and HIV-associated neuropathic pain. 2009 , 13, 387-98		67
1174	Behavioural and histological observations of sensory impairment caused by tight ligation of the trigeminal nerve in mice. <i>Journal of Neuroscience Methods</i> , 2009 , 181, 67-72	3	22
1173	Development of a mechanical stimulator and force measurement system for the assessment of nociceptive thresholds in pigs. <i>Journal of Neuroscience Methods</i> , 2009 , 182, 64-70	3	19
1172	Discovery of a novel class of isoxazoline voltage gated sodium channel blockers. 2009 , 19, 5329-33		12
1171	Discovery of isoxazole voltage gated sodium channel blockers for treatment of chronic pain. 2009 , 19, 5334-8		14
1170	Neuropathic pain: models and mechanisms. 2009 , 15, 1711-6		55
1169	Pain: Mechanisms and Measurement. 2009 ,		
1168	Antinociceptive synergistic effect of spinal mGluR2/3 antagonist and glial cells inhibitor on peripheral inflammation-induced mechanical hypersensitivity. 2009 , 79, 219-23		21
1167	Role of the spinal cord NR2B-containing NMDA receptors in the development of neuropathic pain. 2009 , 215, 298-307		118

1166	L1 cell adhesion molecule is essential for the maintenance of hyperalgesia after spinal cord injury. 2009 , 216, 22-34	26
1165	Selective inhibition of JNK with a peptide inhibitor attenuates pain hypersensitivity and tumor growth in a mouse skin cancer pain model. 2009 , 219, 146-55	49
1164	Attenuation of morphine tolerance by minocycline and pentoxifylline in naive and neuropathic mice. 2009 , 23, 75-84	130
1163	Stress exacerbates neuropathic pain via glucocorticoid and NMDA receptor activation. 2009 , 23, 851-60	105
1162	Transgene-mediated enkephalin expression attenuates signs of naloxone-precipitated morphine withdrawal in rats with neuropathic pain. 2009 , 197, 84-9	18
1161	Housing supplementation decreases the magnitude of inflammation-induced nociception in rats. 2009 , 197, 230-3	25
1160	The CatWalk method: assessment of mechanical allodynia in experimental chronic pain. 2009 , 198, 477-80	35
1159	Early exercise in spinal cord injured rats induces allodynia through TrkB signaling. 2009 , 381, 339-44	17
1158	Antinociceptive effects of the marine snail peptides conantokin-G and conotoxin MVIIA alone and in combination in rat models of pain. 2009 , 56, 556-63	43
1157	Pharmacological assessments of nitric oxide synthase isoforms and downstream diversity of NO signaling in the maintenance of thermal and mechanical hypersensitivity after peripheral nerve injury in mice. 2009 , 56, 702-8	100
1156	Antinociceptive effects of chronic administration of uncompetitive NMDA receptor antagonists in a rat model of diabetic neuropathic pain. 2009 , 57, 121-6	63
1155	Antinociceptive effects of the non-selective cannabinoid receptor agonist CP 55,940 are absent in CB1(-/-) and not CB2(-/-) mice in models of acute and persistent pain. 2009 , 57, 235-41	37
1154	The contributing role of CD14 in toll-like receptor 4 dependent neuropathic pain. 2009 , 158, 896-903	89
1153	Effects of distal nerve injuries on dorsal-horn neurons and glia: relationships between lesion size and mechanical hyperalgesia. 2009 , 158, 904-14	25
1152	High force reaching task induces widespread inflammation, increased spinal cord neurochemicals and neuropathic pain. 2009 , 158, 922-31	54
1151	Noradrenergic neurons in the locus coeruleus contribute to neuropathic pain. 2009 , 160, 174-85	82
1150	Distribution and tumor necrosis factor-alpha isoform binding specificity of locally administered etanercept into injured and uninjured rat sciatic nerve. 2009 , 160, 492-500	23
1149	Remote astrocytic and microglial activation modulates neuronal hyperexcitability and below-level neuropathic pain after spinal injury in rat. 2009 , 161, 895-903	114

1148	Cognitive impairment of prefrontal-dependent decision-making in rats after the onset of chronic pain. 2009 , 161, 671-9	84
1147	MKK3, an upstream activator of p38, contributes to formalin phase 2 and late allodynia in mice. 2009 , 162, 462-71	25
1146	Changes in expression of sensory organ-specific microRNAs in rat dorsal root ganglia in association with mechanical hypersensitivity induced by spinal nerve ligation. 2009 , 164, 711-23	125
1145	Evidence for a role of heat shock protein-90 in toll like receptor 4 mediated pain enhancement in rats. 2009 , 164, 1821-32	57
1144	The pro-nociceptive effects of remifentanyl or surgical injury in mice are associated with a decrease in delta-opioid receptor mRNA levels: Prevention of the nociceptive response by on-site delivery of enkephalins. 2009 , 141, 88-96	41
1143	Profiling of dynamically changed gene expression in dorsal root ganglia post peripheral nerve injury and a critical role of injury-induced glial fibrillary acidic protein in maintenance of pain behaviors [corrected]. 2009 , 143, 114-22	79
1142	Peripheral inflammation increased the synaptic expression of NMDA receptors in spinal dorsal horn. 2009 , 144, 162-9	50
1141	Differential regulation of TRP channels in a rat model of neuropathic pain. 2009 , 144, 187-99	88
1140	Electroacupuncture suppresses capsaicin-induced secondary hyperalgesia through an endogenous spinal opioid mechanism. 2009 , 145, 332-340	47
1139	Role of substance P signaling in enhanced nociceptive sensitization and local cytokine production after incision. 2009 , 145, 341-349	77
1138	Long-lasting neonatal inflammation enhances pain responses to subsequent inflammation, but not peripheral nerve injury in adult rats. 2009 , 27, 215-22	11
1137	Pharmacological characteristics of the hind paw weight bearing difference induced by chronic constriction injury of the sciatic nerve in rats. 2009 , 84, 622-6	34
1136	Peripheral and spinal antihyperalgesic activity of najanalgesin isolated from <i>Naja naja atra</i> in a rat experimental model of neuropathic pain. 2009 , 460, 191-5	12
1135	Role of spinal cholecystokinin in neuropathic pain after spinal cord hemisection in rats. 2009 , 462, 303-7	16
1134	Sustained attention deficits in rats with chronic inflammatory pain. 2009 , 463, 98-102	49
1133	Decreased opioid analgesia in weanling rats exposed to endothelin-1 during infancy. 2009 , 466, 144-8	5
1132	Evaluation of side effects through selective ablation of the mu opioid receptor expressing descending nociceptive facilitatory neurons in the rostral ventromedial medulla with dermorphin-saporin. 2009 , 30, 1096-106	14
1131	Sigma-1 receptors regulate activity-induced spinal sensitization and neuropathic pain after peripheral nerve injury. 2009 , 145, 294-303	123

1130	JNK-induced MCP-1 production in spinal cord astrocytes contributes to central sensitization and neuropathic pain. 2009 , 29, 4096-108	417
1129	Discovery of selective nonpeptidergic neuropeptide FF2 receptor agonists. 2009 , 52, 6511-4	24
1128	Tyrosine phosphorylation of the N-Methyl-D-Aspartate receptor 2B subunit in spinal cord contributes to remifentanyl-induced postoperative hyperalgesia: the preventive effect of ketamine. 2009 , 5, 76	79
1127	Large A-fiber activity is required for microglial proliferation and p38 MAPK activation in the spinal cord: different effects of resiniferatoxin and bupivacaine on spinal microglial changes after spared nerve injury. 2009 , 5, 53	77
1126	Behavioral phenotypes of mice lacking purinergic P2X4 receptors in acute and chronic pain assays. 2009 , 5, 28	134
1125	Formalin-induced behavioural hypersensitivity and neuronal hyperexcitability are mediated by rapid protein synthesis at the spinal level. 2009 , 5, 27	64
1124	Intrathecal delivery of PDGF produces tactile allodynia through its receptors in spinal microglia. 2009 , 5, 23	22
1123	Antidepressants inhibit P2X4 receptor function: a possible involvement in neuropathic pain relief. 2009 , 5, 20	86
1122	Behavioural and electrophysiological characterisation of experimentally induced osteoarthritis and neuropathy in C57Bl/6 mice. 2009 , 5, 18	59
1121	Transforming growth factor-beta1 impairs neuropathic pain through pleiotropic effects. 2009 , 5, 16	95
1120	MRI structural brain changes associated with sensory and emotional function in a rat model of long-term neuropathic pain. 2009 , 47, 1007-14	194
1119	Activation of p-38alpha MAPK contributes to neuronal hyperexcitability in caudal regions remote from spinal cord injury. 2009 , 220, 154-61	28
1118	TRPV1 receptor in expression of opioid-induced hyperalgesia. 2009 , 10, 243-52	55
1117	Additive antinociceptive effects of the selective Nav1.8 blocker A-803467 and selective TRPV1 antagonists in rat inflammatory and neuropathic pain models. 2009 , 10, 306-15	35
1116	siRNA-mediated knockdown of the NR1 subunit gene of the NMDA receptor attenuates formalin-induced pain behaviors in adult rats. 2009 , 10, 380-90	40
1115	Joint distraction magnitude is associated with different behavioral outcomes and substance P levels for cervical facet joint loading in the rat. 2009 , 10, 436-45	54
1114	Role of NFkappaB in an animal model of complex regional pain syndrome-type I (CRPS-I). 2009 , 10, 1161-9	50
1113	6-Methoxy-N-alkyl isatin acylhydrazone derivatives as a novel series of potent selective cannabinoid receptor 2 inverse agonists: design, synthesis, and binding mode prediction. 2009 , 52, 433-44	64

1112	The effects of punctured nucleus pulposus on lumbar radicular pain in rats: a behavioral and immunohistochemical study. 2009 , 11, 492-500	17
1111	Ranolazine attenuates behavioral signs of neuropathic pain. 2009 , 20, 755-8	27
1110	Biochemical and biological properties of 4-(3-phenyl-[1,2,4] thiadiazol-5-yl)-piperazine-1-carboxylic acid phenylamide, a mechanism-based inhibitor of fatty acid amide hydrolase. 2009 , 108, 316-29	58
1109	The effect of amitriptyline on ectopic discharge of primary afferent fibers in the L5 dorsal root in a rat model of neuropathic pain. 2009 , 108, 1671-9	9
1108	A peripherally acting Na(v)1.7 sodium channel blocker reverses hyperalgesia and allodynia on rat models of inflammatory and neuropathic pain. 2009 , 109, 951-8	49
1107	From mouse to man: the 5-HT3 receptor modulates physical dependence on opioid narcotics. 2009 , 19, 193-205	52
1106	Use of transduced adipose tissue stromal cells as biologic minipumps to deliver levodopa for the treatment of neuropathic pain: possibilities and limitations. 2009 , 18, 1341-58	7
1105	Demonstration of antiallodynic effects of the cyclooxygenase-2 inhibitor meloxicam on established diabetic neuropathic pain in mice. 2009 , 110, 213-7	25
1104	Protein kinase A-dependence of the supraspinally mediated analgesic effects of gabapentin on thermal and mechanical hypersensitivity. 2009 , 110, 223-6	9
1103	Peripheral neuropathy in rats exposed to dichloroacetate. 2009 , 68, 985-93	39
1102	Nerve growth factor mediates mechanical allodynia in a mouse model of type 2 diabetes. 2009 , 68, 1229-43	72
1101	The effects of Ginkgo biloba extract EGb 761 on mechanical and cold allodynia in a rat model of neuropathic pain. 2009 , 108, 1958-63	35
1100	Complete Freund's adjuvant-induced intervertebral discitis as an animal model for discogenic low back pain. 2009 , 109, 1287-96	33
1099	The reactions of glial cells and endoneurial macrophages in the dorsal root ganglion and their contribution to pain-related behavior after application of nucleus pulposus onto the nerve root in rats. 2010 , 35, 264-71	40
1098	Spinal cord stimulation-induced analgesia: electrical stimulation of dorsal column and dorsal roots attenuates dorsal horn neuronal excitability in neuropathic rats. 2010 , 113, 1392-405	133
1097	Local anesthetic sciatic nerve block and nerve fiber damage in diabetic rats. 2010 , 35, 343-50	57
1096	The effects of resiniferatoxin in an experimental rat thoracotomy model. 2010 , 110, 228-32	8
1095	The effect of pulsed radiofrequency current on mechanical allodynia induced with resiniferatoxin in rats. 2010 , 111, 784-90	48

1094	Anti-nociceptive effect of bovine milk-derived lactoferrin in a rat lumbar disc herniation model. 2010, 35, 1663-7	8
1093	The reactions of glial cells and endoneurial macrophages in the dorsal root ganglion and their contribution to pain-related behavior after application of nucleus pulposus onto the nerve root in rats. 2010, 35, 10-7	14
1092	Pharmacological evaluation of analgesic effects of the cholecystokinin2 receptor antagonist Z-360 in mouse models of formalin- and cancer-induced pain. 2010, 33, 244-8	9
1091	Mexiletine reverses oxaliplatin-induced neuropathic pain in rats. 2010, 112, 473-6	35
1090	Contribution of the chemokine (C-C motif) ligand 2 (CCL2) to mechanical hypersensitivity after surgical incision in rats. 2010, 112, 1250-8	38
1089	Microinjection of morphine into thalamic nucleus submedius depresses bee venom-induced inflammatory pain in the rat. 2010, 60, 1355-1363	12
1088	Antinociceptive and anti-inflammatory properties of 7-hydroxycoumarin in experimental animal models: potential therapeutic for the control of inflammatory chronic pain. 2010, 62, 205-13	52
1087	Bilateral hyperexcitability of thalamic VPL neurons following unilateral spinal injury in rats. 2010, 60, 59-66	28
1086	Attenuation of neuropathy-induced allodynia following intraplantar injection of pregabalin. 2010, 57, 664-71	19
1085	Endogenous enkephalin does not contribute to the cerebral anti-hyperalgesic action of gabapentin. 2010, 53, 1428-32	1
1084	ADAMTS-5 deficient mice do not develop mechanical allodynia associated with osteoarthritis following medial meniscal destabilization. 2010, 18, 572-80	99
1083	Glucosamine sulfate reduces experimental osteoarthritis and nociception in rats: association with changes of mitogen-activated protein kinase in chondrocytes. 2010, 18, 1192-202	58
1082	Limited BDNF contributes to the failure of injury to skin afferents to produce a neuropathic pain condition. 2010, 148, 148-157	30
1081	N-Methyl-D-aspartate receptor (NMDAR) independent maintenance of inflammatory pain. 2010, 148, 237-246	45
1080	Acetazolamide and midazolam act synergistically to inhibit neuropathic pain. 2010, 148, 302-308	95
1079	The c-Jun N-terminal kinase 1 (JNK1) in spinal astrocytes is required for the maintenance of bilateral mechanical allodynia under a persistent inflammatory pain condition. 2010, 148, 309-319	122
1078	Aberrant sensory responses are dependent on lesion severity after spinal cord contusion injury in mice. 2010, 148, 328-342	35
1077	Anti-GD(2) with an FC point mutation reduces complement fixation and decreases antibody-induced allodynia. 2010, 149, 135-142	93

1076	Peripheral inflammation induces tumor necrosis factor dependent AMPA receptor trafficking and Akt phosphorylation in spinal cord in addition to pain behavior. 2010 , 149, 243-253	130
1075	Pharmacological activation of 5-HT7 receptors reduces nerve injury-induced mechanical and thermal hypersensitivity. 2010 , 149, 483-494	67
1074	Microglial/macrophage GRK2 determines duration of peripheral IL-1beta-induced hyperalgesia: contribution of spinal cord CX3CR1, p38 and IL-1 signaling. 2010 , 150, 550-560	79
1073	Spinal hyperpolarization-activated cyclic nucleotide-gated cation channels at primary afferent terminals contribute to chronic pain. 2010 , 151, 87-96	47
1072	Role of peripheral endothelin receptors in an animal model of complex regional pain syndrome type 1 (CRPS-I). 2010 , 151, 174-183	38
1071	A peripherally restricted cannabinoid receptor agonist produces robust anti-nociceptive effects in rodent models of inflammatory and neuropathic pain. 2010 , 151, 337-344	73
1070	Characterization of the acute and persistent pain state present in K/BxN serum transfer arthritis. 2010 , 151, 394-403	94
1069	Neuronal hyperexcitability in the dorsal horn after painful facet joint injury. 2010 , 151, 414-421	51
1068	Spinal-, brainstem- and cerebrally mediated responses at- and below-level of a spinal cord contusion in rats: evaluation of pain-like behavior. 2010 , 151, 670-679	69
1067	The analgesic and toxic effects of nornicotine enantiomers alone and in interaction with morphine in rodent models of acute and persistent pain. 2010 , 94, 352-62	15
1066	H4 receptor antagonism exhibits anti-nociceptive effects in inflammatory and neuropathic pain models in rats. 2010 , 95, 41-50	74
1065	Phosphatidylinositol 3-kinase mediates pain behaviors induced by activation of peripheral ephrinBs/EphBs signaling in mice. 2010 , 95, 315-24	28
1064	Potentialiation of morphine antiallodynic efficacy by ACPT-III, a group III metabotropic glutamate receptor agonist, in rat spinal nerve ligation-induced neuropathic pain. 2010 , 96, 108-13	8
1063	Antinociceptive effect of ambroxol in rats with neuropathic spinal cord injury pain. 2010 , 97, 249-55	34
1062	Changes in Abeta non-nociceptive primary sensory neurons in a rat model of osteoarthritis pain. 2010 , 6, 37	28
1061	Sildenafil and glyceryl trinitrate reduce tactile allodynia in streptozotocin-injected rats. 2010 , 631, 17-23	6
1060	Regulation of spinal neuroimmune responses by prolonged morphine treatment in a rat model of cancer induced bone pain. 2010 , 1326, 162-73	30
1059	Decrease in the descending inhibitory 5-HT system in rats with spinal nerve ligation. 2010 , 1330, 45-60	39

1058	Tibia tumor-induced cancer pain involves spinal p38 mitogen-activated protein kinase activation via TLR4-dependent mechanisms. 2010 , 1346, 213-23	45	
1057	Lack of analgesic efficacy of spinal ondansetron on thermal and mechanical hypersensitivity following spinal nerve ligation in the rat. 2010 , 1352, 83-93	28	
1056	Chemical or surgical sympathectomy prevents mechanical hyperalgesia induced by intraplantar injection of bee venom in rats. 2010 , 1353, 86-93	13	
1055	Antinociceptive effects of histamine H3 receptor antagonist in the preclinical models of pain in rats and the involvement of central noradrenergic systems. 2010 , 1354, 74-84	32	
1054	Differential effects of adenosine A1 receptor on pain-related behavior in normal and nerve-injured rats. 2010 , 1361, 23-30	36	
1053	Inhibition of NF-kappaB prevents mechanical allodynia induced by spinal ventral root transection and suppresses the re-expression of Nav1.3 in DRG neurons in vivo and in vitro. 2010 , 1363, 151-8	36	
1052	Activation of astrocytes in the spinal cord contributes to the development of bilateral allodynia after peripheral nerve injury in rats. 2010 , 1363, 72-80	42	
1051	Sciatic nerve injury in adult rats causes distinct changes in the central projections of sensory neurons expressing different glial cell line-derived neurotrophic factor family receptors. 2010 , 518, 3024-45	23	
1050	Nerve injury-activated microglia engulf myelinated axons in a P2Y12 signaling-dependent manner in the dorsal horn. 2010 , 58, 1838-46	54	
1049	Spinal injection of TNF- α -activated astrocytes produces persistent pain symptom mechanical allodynia by releasing monocyte chemoattractant protein-1. 2010 , 58, 1871-80	89	
1048	Alteration of sensory neurons and spinal response to an experimental osteoarthritis pain model. 2010 , 62, 2995-3005	127	
1047	Altered expression of sodium channel distribution in the dorsal root ganglion after gradual elongation of rat sciatic nerves. 2010 , 28, 481-6	8	
1046	Noradrenaline stimulates ATP release from DRG neurons by targeting beta(3) adrenoceptors as a factor of neuropathic pain. 2010 , 224, 345-51	25	
1045	A novel animal model of graded neuropathic pain: utility to investigate mechanisms of population heterogeneity. <i>Journal of Neuroscience Methods</i> , 2010 , 193, 47-53	3	35
1044	Anti-allodynic effect of the flavonoid myricetin in a rat model of neuropathic pain: Involvement of p38 and protein kinase C mediated modulation of Ca ²⁺ channels. 2010 , 14, 992-8	32	
1043	Synthesis and bradykinin inhibitory activity of novel non-peptide compounds, and evaluation of in vivo analgesic activity. 2010 , 18, 2327-2336	4	
1042	Synthesis and T-type calcium channel blocking activity of novel diphenylpiperazine compounds, and evaluation of in vivo analgesic activity. 2010 , 18, 5938-44	16	
1041	Substituted biaryl oxazoles, imidazoles, and thiazoles as sodium channel blockers. 2010 , 20, 5536-40	13	

1040	Substituted biaryl pyrazoles as sodium channel blockers. 2010 , 20, 5480-3	12
1039	Discovery of a novel class of biphenyl pyrazole sodium channel blockers for treatment of neuropathic pain. 2010 , 20, 7479-82	17
1038	Modality of hyperalgesia tested, not type of nerve damage, predicts pharmacological sensitivity in rat models of neuropathic pain. 2010 , 14, 503-9	18
1037	The role of N-methyl-D-aspartate receptor subunit NR2B in spinal cord in cancer pain. 2010 , 14, 496-502	54
1036	Sumatriptan alleviates nitroglycerin-induced mechanical and thermal allodynia in mice. 2010 , 30, 170-8	144
1035	Spinal and peripheral analgesic effects of the CB2 cannabinoid receptor agonist AM1241 in two models of bone cancer-induced pain. 2010 , 160, 561-73	67
1034	Effects of COX-2 inhibition on spinal nociception: the role of endocannabinoids. 2010 , 160, 669-76	29
1033	Spinal glial TLR4-mediated nociception and production of prostaglandin E(2) and TNF. 2010 , 160, 1754-64	77
1032	Sustained morphine-mediated pain sensitization and antinociceptive tolerance are blocked by intrathecal treatment with Raf-1-selective siRNA. 2010 , 161, 51-64	13
1031	Reversal of acid-induced and inflammatory pain by the selective ASIC3 inhibitor, APETx2. 2010 , 161, 950-60	92
1030	Evidence that pregabalin reduces neuropathic pain by inhibiting the spinal release of glutamate. 2010 , 113, 552-61	72
1029	Inflammatory hyperalgesia induces essential bioactive lipid production in the spinal cord. 2010 , 114, 981-93	44
1028	Reduced inflammatory and neuropathic pain and decreased spinal microglial response in fractalkine receptor (CX3CR1) knockout mice. 2010 , 114, 1143-57	86
1027	Light touch induces ERK activation in superficial dorsal horn neurons after inflammation: involvement of spinal astrocytes and JNK signaling in touch-evoked central sensitization and mechanical allodynia. 2010 , 115, 505-14	52
1026	High-throughput behavioral phenotyping in the expanded panel of BXD recombinant inbred strains. 2010 , 9, 129-59	130
1025	Coding of facial expressions of pain in the laboratory mouse. 2010 , 7, 447-9	743
1024	Endocannabinoid regulation of spinal nociceptive processing in a model of neuropathic pain. 2010 , 31, 1414-22	23
1023	Effects of interleukin-6 treatment on neurovascular function, nerve perfusion and vascular endothelium in diabetic rats. 2010 , 12, 689-99	26

1022	Pharmacological characterization of a novel cannabinoid ligand, MDA19, for treatment of neuropathic pain. 2010 , 111, 99-109	30
1021	Mediators and Mechanisms of Inflammatory Pain. 217-233	
1020	Pungent general anesthetics activate transient receptor potential-A1 to produce hyperalgesia and neurogenic bronchoconstriction. 2010 , 112, 1452-63	54
1019	Antinociceptive Effect of Memantine and Morphine on Vincristine-induced Peripheral Neuropathy in Rats. 2010 , 23, 179-85	23
1018	MDL-28170 has no analgesic effect on CCI induced neuropathic pain in mice. 2010 , 15, 3038-47	6
1017	Evidence for a role of endocannabinoids, astrocytes and p38 phosphorylation in the resolution of postoperative pain. 2010 , 5, e10891	52
1016	Are spinal GABAergic elements related to the manifestation of neuropathic pain in rat?. 2010 , 14, 59-69	21
1015	Intrathecal siRNA against Toll-like receptor 4 reduces nociception in a rat model of neuropathic pain. 2010 , 7, 251-9	72
1014	Effect of perioperative perineural injection of dexamethasone and bupivacaine on a rat spared nerve injury model. 2010 , 23, 166-71	12
1013	Effect of sildenafil on neuropathic pain and hemodynamics in rats. 2010 , 51, 82-7	16
1012	Validation of a preclinical spinal safety model: effects of intrathecal morphine in the neonatal rat. 2010 , 113, 183-99	35
1011	Participation of neuronal nitric oxide synthase in experimental neuropathic pain induced by sciatic nerve transection. 2010 , 43, 367-76	30
1010	Antiallodynic Effect of Thalidomide and Morphine on Rat Spinal Nerve Ligation-induced Neuropathic Pain. 2010 , 23, 172-8	14
1009	Analgesic effects of a substituted N-triazole oxindole (TROX-1), a state-dependent, voltage-gated calcium channel 2 blocker. 2010 , 334, 545-55	79
1008	The role of nitric oxide in the local antiallodynic and antihyperalgesic effects and expression of delta-opioid and cannabinoid-2 receptors during neuropathic pain in mice. 2010 , 334, 887-96	52
1007	Ligand-directed trafficking of the μ opioid receptor in vivo: two paths toward analgesic tolerance. 2010 , 30, 16459-68	109
1006	IL-6- and NGF-induced rapid control of protein synthesis and nociceptive plasticity via convergent signaling to the eIF4F complex. 2010 , 30, 15113-23	150
1005	SOCS3-mediated blockade of JAK/STAT3 signaling pathway reveals its major contribution to spinal cord neuroinflammation and mechanical allodynia after peripheral nerve injury. 2010 , 30, 5754-66	139

1004	Genetic targeting of ERK1 suggests a predominant role for ERK2 in murine pain models. 2010 , 30, 11537-47	36
1003	Monoamine-dependent, opioid-independent antihypersensitivity effects of intrathecally administered milnacipran, a serotonin noradrenaline reuptake inhibitor, in a postoperative pain model in rats. 2010 , 334, 1059-65	33
1002	Surgical incision induces anxiety-like behavior and amygdala sensitization: effects of morphine and gabapentin. 2010 , 2010, 705874	16
1001	Overexpression of NGF in mouse urothelium leads to neuronal hyperinnervation, pelvic sensitivity, and changes in urinary bladder function. 2010 , 298, R534-47	119
1000	Metabotropic glutamate receptor-5 and protein kinase C-epsilon increase in dorsal root ganglion neurons and spinal glial activation in an adolescent rat model of painful neck injury. 2010 , 27, 2261-71	26
999	Animal models of acute and chronic inflammatory and nociceptive pain. 2010 , 617, 41-55	13
998	Sensory stimulation prior to spinal cord injury induces post-injury dysesthesia in mice. 2010 , 27, 777-87	10
997	Increased spinal dynorphin levels and phospho-extracellular signal-regulated kinases 1 and 2 and c-Fos immunoreactivity after surgery under remifentanil anesthesia in mice. 2010 , 77, 185-94	29
996	Analgesic (omega)-conotoxins CVIE and CVIF selectively and voltage-dependently block recombinant and native N-type calcium channels. 2010 , 77, 139-48	50
995	P2X7-dependent release of interleukin-1beta and nociception in the spinal cord following lipopolysaccharide. 2010 , 30, 573-82	236
994	Chronic spontaneous activity generated in the somata of primary nociceptors is associated with pain-related behavior after spinal cord injury. 2010 , 30, 14870-82	104
993	NADPH oxidase 2-derived reactive oxygen species in spinal cord microglia contribute to peripheral nerve injury-induced neuropathic pain. 2010 , 107, 14851-6	170
992	The roles of T-type calcium channel in the development of neuropathic pain following chronic compression of rat dorsal root ganglia. 2010 , 85, 295-300	53
991	Evaluating intra-articular drug delivery for the treatment of osteoarthritis in a rat model. 2010 , 16, 81-92	46
990	Triptan-induced enhancement of neuronal nitric oxide synthase in trigeminal ganglion dural afferents underlies increased responsiveness to potential migraine triggers. 2010 , 133, 2475-88	79
989	Identification and functional characterization of a stable, centrally active derivative of the neurotensin (8-13) fragment as a potential first-in-class analgesic. 2010 , 53, 4623-32	27
988	In vitro and in vivo effects on neural crest stem cell differentiation by conditional activation of Runx1 short isoform and its effect on neuropathic pain behavior. 2010 , 115, 56-64	7
987	Key role for spinal dorsal horn microglial kinin B1 receptor in early diabetic pain neuropathy. 2010 , 7, 36	64

986	Loss of GABAergic interneurons in laminae I-III of the spinal cord dorsal horn contributes to reduced GABAergic tone and neuropathic pain after spinal cord injury. 2010 , 27, 729-37	126
985	Pain Control on Demand Based on Pulsed Radio-Frequency Stimulation of the Dorsal Root Ganglion Using a Batteryless Implantable CMOS SoC. 2010 , 4, 350-9	30
984	A partial L5 spinal nerve ligation induces a limited prolongation of mechanical allodynia in rats: an efficient model for studying mechanisms of neuropathic pain. 2010 , 471, 43-7	22
983	Plasticity in the synaptic number associated with neuropathic pain in the rat spinal dorsal horn: A stereological study. 2010 , 486, 24-8	16
982	Prenatal exposure to methamphetamine alters the mechanical withdrawal threshold and tonic hyperalgesia in the offspring. 2010 , 31, 432-8	13
981	A cannabinoid 2 receptor agonist attenuates bone cancer-induced pain and bone loss. 2010 , 86, 646-53	65
980	Intrathecal gabapentin and clonidine synergistically inhibit allodynia in spinal nerve-ligated rats. 2010 , 87, 565-71	19
979	Levodopa analgesia in experimental neuropathic pain. 2010 , 83, 304-9	33
978	Spinal high-mobility group box 1 contributes to mechanical allodynia in a rat model of bone cancer pain. 2010 , 395, 572-6	47
977	VGLUT2-dependent sensory neurons in the TRPV1 population regulate pain and itch. 2010 , 68, 529-42	166
976	Comparison of mechanical allodynia and the affective component of inflammatory pain in rats. 2010 , 58, 537-43	45
975	Cannabinoid receptor-mediated antinociception with acetaminophen drug combinations in rats with neuropathic spinal cord injury pain. 2010 , 58, 758-66	40
974	Pain is a salient "stressor" that is mediated by corticotropin-releasing factor-1 receptors. 2010 , 59, 160-6	36
973	Peripheral contributions to the mechanical hyperalgesia following a lumbar 5 spinal nerve lesion in rats. 2010 , 165, 221-32	13
972	Evidence that intrathecal morphine-3-glucuronide may cause pain enhancement via toll-like receptor 4/MD-2 and interleukin-1beta. 2010 , 165, 569-83	129
971	Topography and time course of changes in spinal neuropeptide Y immunoreactivity after spared nerve injury. 2010 , 165, 914-22	12
970	Effects of neurokinin-1 receptor agonism and antagonism in the rostral ventromedial medulla of rats with acute or persistent inflammatory nociception. 2010 , 165, 902-13	31
969	Possible involvement of toll-like receptor 4/myeloid differentiation factor-2 activity of opioid inactive isomers causes spinal proinflammation and related behavioral consequences. 2010 , 167, 880-93	97

968	Evoked pain behavior and spinal glia activation is dependent on tumor necrosis factor receptor 1 and 2 in a mouse model of bone cancer pain. 2010 , 169, 463-74	46
967	Subtype-specific changes in 5-HT receptor-mediated modulation of C fibre-evoked spinal field potentials are triggered by peripheral nerve injury. 2010 , 168, 831-41	50
966	Morphine tolerance attenuates the resolution of postoperative pain and enhances spinal microglial p38 and extracellular receptor kinase phosphorylation. 2010 , 169, 843-54	50
965	Mammalian target of rapamycin in spinal cord neurons mediates hypersensitivity induced by peripheral inflammation. 2010 , 169, 1392-402	69
964	Spinal upregulation of glutamate transporter GLT-1 by ceftriaxone: therapeutic efficacy in a range of experimental nervous system disorders. 2010 , 169, 1888-900	98
963	The role of dopamine receptors in ventrolateral orbital cortex-evoked anti-nociception in a rat model of neuropathic pain. 2010 , 169, 1872-80	28
962	Reserpine causes biphasic nociceptive sensitivity alteration in conjunction with brain biogenic amine tones in rats. 2010 , 169, 1860-71	27
961	Enhanced nociceptive responding in two rat models of depression is associated with alterations in monoamine levels in discrete brain regions. 2010 , 171, 1300-13	58
960	Contribution of the spinal cord BDNF to the development of neuropathic pain by activation of the NR2B-containing NMDA receptors in rats with spinal nerve ligation. 2010 , 222, 256-66	113
959	Combining ketamine with astrocytic inhibitor as a potential analgesic strategy for neuropathic pain ketamine, astrocytic inhibitor and pain. 2010 , 6, 50	21
958	Deficiency of the negative immune regulator B7-H1 enhances inflammation and neuropathic pain after chronic constriction injury of mouse sciatic nerve. 2010 , 222, 153-60	22
957	Validity of acute and chronic tactile sensory testing after spinal cord injury in rats. 2010 , 225, 366-76	48
956	Unilateral focal burn injury is followed by long-lasting bilateral allodynia and neuronal hyperexcitability in spinal cord dorsal horn. 2010 , 11, 119-30	66
955	Learned avoidance from noxious mechanical simulation but not threshold semmes weinstein filament stimulation after nerve injury in rats. 2010 , 11, 280-6	52
954	A novel method to quantify histochemical changes throughout the mediolateral axis of the substantia gelatinosa after spared nerve injury: characterization with TRPV1 and substance P. 2010 , 11, 388-98	20
953	Characterization of Fasudil in preclinical models of pain. 2010 , 11, 941-9	14
952	Pain intensity and duration can be enhanced by prior challenge: initial evidence suggestive of a role of microglial priming. 2010 , 11, 1004-14	77
951	Minocycline attenuates mechanical allodynia and central sensitization following peripheral second-degree burn injury. 2010 , 11, 1146-54	56

950	The antinociceptive effect of (-)-linalool in models of chronic inflammatory and neuropathic hypersensitivity in mice. 2010 , 11, 1222-9	90
949	Fatty acid amide hydrolase and monoacylglycerol lipase inhibitors produce anti-allodynic effects in mice through distinct cannabinoid receptor mechanisms. 2010 , 11, 1420-8	95
948	Synthesis, structure-activity relationship, and pharmacological profile of analogs of the ASIC-3 inhibitor A-317567. 2010 , 1, 19-24	23
947	Analgesic effect of highly reversible α -conotoxin FVIA on N type Ca^{2+} channels. 2010 , 6, 97	33
946	Abnormal nociception and opiate sensitivity of STOP null mice exhibiting elevated levels of the endogenous alkaloid morphine. 2010 , 6, 96	7
945	Distinctive response of CNS glial cells in oro-facial pain associated with injury, infection and inflammation. 2010 , 6, 79	44
944	Hypolocomotion, asymmetrically directed behaviors (licking, lifting, flinching, and shaking) and dynamic weight bearing (gait) changes are not measures of neuropathic pain in mice. 2010 , 6, 34	79
943	p38 mediates mechanical allodynia in a mouse model of type 2 diabetes. 2010 , 6, 28	49
942	Down-regulation of Toll-like receptor 4 gene expression by short interfering RNA attenuates bone cancer pain in a rat model. 2010 , 6, 2	102
941	Analgesia. 2010 ,	6
940	Locomotor activity in a novel environment as a test of inflammatory pain in rats. 2010 , 617, 67-78	15
939	Sodium channel blockers for neuropathic pain. 2010 , 20, 755-79	36
938	Differential effects of nociceptin/orphanin FQ (NOP) receptor agonists in acute versus chronic pain: studies with bifunctional NOP/ μ receptor agonists in the sciatic nerve ligation chronic pain model in mice. 2011 , 339, 687-93	48
937	Involvement of substance P in peripheral neuropathy induced by paclitaxel but not oxaliplatin. 2011 , 337, 226-35	40
936	Antitussive effect of naringin on experimentally induced cough in Guinea pigs. 2011 , 77, 16-21	43
935	Peripheral nerve injury leads to working memory deficits and dysfunction of the hippocampus by upregulation of TNF- α in rodents. 2011 , 36, 979-92	160
934	Behavioral Measures of Pain Thresholds. 2011 , 1, 383-412	28
933	Anti-arthritic activity of standardised extract of <i>Phyllanthus amarus</i> in Freund's complete adjuvant induced arthritis. 2011 , 1, 185-190	27

932	Coexisting mechanical hypersensitivity and anxiety in a rat model of spinal cord injury and the effect of pregabalin, morphine, and midazolam treatment. 2018 , 2, 139-145	7
931	Effects of the excitatory amino acid transporter subtype 2 (EAAT-2) inducer ceftriaxone on different pain modalities in rat. 2018 , 2, 132-136	10
930	Single-walled carbon nanotubes chemically functionalized with polyethylene glycol promote tissue repair in a rat model of spinal cord injury. 2011 , 28, 2349-62	98
929	Animal Models of Pain. 2011 ,	2
928	Spinal protein kinase M μ underlies the maintenance mechanism of persistent nociceptive sensitization. 2011 , 31, 6646-53	102
927	Impaired sensory nerve function and axon morphology in mice with diabetic neuropathy. 2011 , 106, 905-14	43
926	The rat intervertebral disk degeneration pain model: relationships between biological and structural alterations and pain. 2011 , 13, R165	37
925	Kinematic and dynamic gait compensations in a rat model of lumbar radiculopathy and the effects of tumor necrosis factor-alpha antagonism. 2011 , 13, R137	16
924	N-Acetyl-l-Cysteine Attenuates Ischemia/Reperfusion Injury-Induced Allodynia and N-Methyl-d-Aspartate Receptor Activation in Rats. 2011 , 72, 216-27	5
923	Sustained blockade of neurotrophin receptors TrkA, TrkB and TrkC reduces non-malignant skeletal pain but not the maintenance of sensory and sympathetic nerve fibers. 2011 , 48, 389-98	50
922	Electroacupuncture attenuates mechanical and warm allodynia through suppression of spinal glial activation in a rat model of neuropathic pain. 2011 , 86, 403-11	52
921	Protective effect of Acorus calamus L. in rat model of vincristine induced painful neuropathy: an evidence of anti-inflammatory and anti-oxidative activity. 2011 , 49, 2557-63	61
920	Dynamic temporal and spatial regulation of mu opioid receptor expression in primary afferent neurons following spinal nerve injury. 2011 , 15, 669-75	51
919	Brain-derived neurotrophic factor contributes to spinal long-term potentiation and mechanical hypersensitivity by activation of spinal microglia in rat. 2011 , 25, 322-34	97
918	Adoptive transfer of peripheral immune cells potentiates allodynia in a graded chronic constriction injury model of neuropathic pain. 2011 , 25, 503-13	21
917	TNF α is involved in neuropathic pain induced by nucleoside reverse transcriptase inhibitor in rats. 2011 , 25, 1668-76	29
916	Somatosensory and sensorimotor consequences associated with the heterozygous disruption of the autism candidate gene, Gabrb3. 2011 , 216, 36-45	59
915	Decoy strategy targeting the brain-derived neurotrophic factor exon I to attenuate tactile allodynia in the neuropathic pain model of rats. 2011 , 408, 139-44	14

914	Antinociceptive properties of bergenin. 2011 , 74, 2062-8	43
913	Substance P in the cerebrospinal fluid-contacting nucleus contributes to morphine physical dependence in rats. 2011 , 488, 188-92	13
912	Antinociceptive effect of intrathecal cannabinoid receptor agonist WIN 55,212-2 in a rat bone tumor pain model. 2011 , 493, 67-71	23
911	Analgesic effect of intrathecally γ -aminobutyric acid transporter-1 inhibitor NO-711 administrating on neuropathic pain in rats. 2011 , 494, 6-9	9
910	Up-regulation of spinal glutamate transporters contributes to anti-hypersensitive effects of valproate in rats after peripheral nerve injury. 2011 , 502, 52-5	36
909	Transient tactile allodynia following intrathecal puncture in mouse: contributions of Toll-like receptor signaling. 2011 , 504, 215-8	8
908	Post-injury administration of minocycline: an effective treatment for nerve-injury induced neuropathic pain. 2011 , 70, 305-12	62
907	Anti-allodynic effects of intrathecally and intracerebroventricularly administered 26RFa, an intrinsic agonist for GRP103, in the rat partial sciatic nerve ligation model. 2011 , 32, 1262-9	11
906	GABAergic disinhibition induced pain hypersensitivity by upregulating NMDA receptor functions in spinal dorsal horn. 2011 , 60, 921-9	29
905	Lacosamide has protective disease modifying properties in experimental vincristine neuropathy. 2011 , 61, 600-7	25
904	Selective knockdown of NMDA receptors in primary afferent neurons decreases pain during phase 2 of the formalin test. 2011 , 172, 474-82	35
903	Spinal activation of delta opioid receptors alleviates cancer-related bone pain. 2011 , 183, 221-9	27
902	Upregulation of neuronal nitric oxide synthase in the periphery promotes pain hypersensitivity after peripheral nerve injury. 2011 , 190, 367-78	28
901	Expression of monocyte chemoattractant protein-1 and its induction by tumor necrosis factor receptor 1 in sensory neurons in the ventral rhizotomy model of neuropathic pain. 2011 , 190, 354-66	10
900	Phosphodiesterase-5 is a therapeutic target for peripheral neuropathy in diabetic mice. 2011 , 193, 399-410	69
899	Activating transcription factor 4, a mediator of the integrated stress response, is increased in the dorsal root ganglia following painful facet joint distraction. 2011 , 193, 377-86	24
898	Manual acupuncture inhibits mechanical hypersensitivity induced by spinal nerve ligation in rats. 2011 , 193, 370-6	25
897	Bipolar spinal cord stimulation attenuates mechanical hypersensitivity at an intensity that activates a small portion of A-fiber afferents in spinal nerve-injured rats. 2011 , 199, 470-80	42

896	Brain-derived neurotrophic factor-activated astrocytes produce mechanical allodynia in neuropathic pain. 2011 , 199, 452-60		26
895	Development of potent μ and κ opioid agonists with high lipophilicity. 2011 , 54, 382-6		40
894	TNF- α enhances the currents of voltage gated sodium channels in uninjured dorsal root ganglion neurons following motor nerve injury. 2011 , 227, 279-86		61
893	Development of mechanical and thermal nociceptive threshold testing devices in unrestrained birds (broiler chickens). <i>Journal of Neuroscience Methods</i> , 2011 , 201, 220-7	3	19
892	A tropomyosine receptor kinase inhibitor blocks spinal neuroplasticity essential for the anti-hypersensitivity effects of gabapentin and clonidine in rats with peripheral nerve injury. 2011 , 12, 94-100		10
891	Does chronic pain alter the normal interaction between cardiovascular and pain regulatory systems? Pain modulation in the hypertensive-monoarthritic rat. 2011 , 12, 194-204		10
890	Radiotelemetric and symptomatic evaluation of pain in the rat after laparotomy: long-term benefits of perioperative ropivacaine care. 2011 , 12, 246-56		10
889	Rewarding electrical brain stimulation in rats after peripheral nerve injury: decreased facilitation by commonly abused prescription opioids. 2011 , 115, 1271-80		32
888	[Effects of nitric oxide synthase inhibitor on hindlimb muscles in rats with neuropathic pain induced by unilateral peripheral nerve injury]. 2011 , 41, 520-7		3
887	Effects of low and high molecular weight hyaluronic acids on peridural fibrosis and inflammation in lumbar laminectomized rats. 2011 , 24, 191-8		5
886	The expression of corticotropin-releasing factor and its receptors in the spinal cord and dorsal root ganglion in a rat model of neuropathic pain. 2011 , 44, 60-8		10
885	A comparison of pain-related behavior following local application of nucleus pulposus and/or mechanical compression on the dorsal root ganglion. 2011 , 57, 46-53		7
884	Animal Pain Models and Behavior Tests. 2011 , 31, 103		2
883	Neuropathic pain as a process: reversal of chronification in an animal model. 2011 , 4, 315-23		11
882	The selective neuronal nitric oxide synthase inhibitor 7-nitroindazole has acute analgesic but not cumulative effects in a rat model of peripheral neuropathy. 2011 , 4, 85-90		2
881	Progesterone prevents development of neuropathic pain in a rat model: Timing and duration of treatment are critical. 2011 , 4, 91-101		28
880	Subarachnoid Transplant of the Human Neuronal hNT2.19 Serotonergic Cell Line Attenuates Behavioral Hypersensitivity without Affecting Motor Dysfunction after Severe Contusive Spinal Cord Injury. 2011 , 2011, 891605		9
879	Design, synthesis and biological activity evaluation of arylpiperazine derivatives for the treatment of neuropathic pain. 2011 , 16, 5785-806		6

878	Effects of glyceic regulation on chronic postischemia pain. 2011 , 115, 614-25	8
877	IL-4 deficiency is associated with mechanical hypersensitivity in mice. 2011 , 6, e28205	43
876	Acid solution is a suitable medium for introducing QX-314 into nociceptors through TRPV1 channels to produce sensory-specific analgesic effects. 2011 , 6, e29395	21
875	A single subanesthetic dose of ketamine relieves depression-like behaviors induced by neuropathic pain in rats. 2011 , 115, 812-21	126
874	Regular exercise reverses sensory hypersensitivity in a rat neuropathic pain model: role of endogenous opioids. 2011 , 114, 940-8	193
873	The macrophage-mediated effects of the peroxisome proliferator-activated receptor-gamma agonist rosiglitazone attenuate tactile allodynia in the early phase of neuropathic pain development. 2011 , 113, 398-404	40
872	The intrathecally administered kappa-2 opioid agonist GR89696 and interleukin-10 attenuate bone cancer-induced pain through synergistic interaction. 2011 , 113, 934-40	12
871	Cannabidiol prevents the development of cold and mechanical allodynia in paclitaxel-treated female C57Bl6 mice. 2011 , 113, 947-50	68
870	Lidocaine suppresses the increased extracellular signal-regulated kinase/cyclic AMP response element-binding protein pathway and pro-inflammatory cytokines in a neuropathic pain model of rats. 2011 , 28, 106-11	13
869	Intrathecal lemnalol, a natural marine compound obtained from Formosan soft coral, attenuates nociceptive responses and the activity of spinal glial cells in neuropathic rats. 2011 , 22, 739-50	29
868	Lack of beneficial effect for preemptive analgesia in postoperative pain control: verifying the efficacy of preemptive analgesia with N-methyl-D-aspartate receptor antagonists in a modified animal model of postoperative pain. 2011 , 112, 710-8	5
867	Lidocaine attenuates the development of diabetic-induced tactile allodynia by inhibiting microglial activation. 2011 , 113, 941-6	28
866	Spinal metabotropic glutamate receptor 4 is involved in neuropathic pain. 2011 , 22, 244-8	24
865	Behavioral signs of chronic back pain in the SPARC-null mouse. 2011 , 36, 95-102	44
864	Changes in spinal cord expression of fractalkine and its receptor in a rat model of disc herniation by autologous nucleus pulposus. 2011 , 36, E753-60	21
863	Changes in expression of mRNA for interleukin-8 and effects of interleukin-8 receptor inhibitor in the spinal dorsal horn in a rat model of lumbar disc herniation. 2011 , 36, 2139-46	43
862	Transient receptor potential vanilloid 1 mediates pain in mice with severe sickle cell disease. 2011 , 118, 3376-83	107
861	Intrathecal administration of AS1928370, a transient receptor potential vanilloid 1 antagonist, attenuates mechanical allodynia in a mouse model of neuropathic pain. 2011 , 34, 1105-8	27

860	Dynamics of Circadian Thalamocortical Flow of Information during a Peripheral Neuropathic Pain Condition. 2011 , 5, 43	16
859	Blockade of persistent sodium currents contributes to the riluzole-induced inhibition of spontaneous activity and oscillations in injured DRG neurons. 2011 , 6, e18681	54
858	The spared nerve injury (SNI) model of induced mechanical allodynia in mice. 2011 ,	38
857	5-hydroxytryptamine type 3 receptor modulates opioid-induced hyperalgesia and tolerance in mice. 2011 , 114, 1180-9	54
856	Intrathecal herpes simplex virus type 1 amplicon vector-mediated human proenkephalin reduces chronic constriction injury-induced neuropathic pain in rats. 2011 , 4, 529-33	13
855	Instability of spatial encoding by CA1 hippocampal place cells after peripheral nerve injury. 2011 , 33, 2255-64	17
854	Forced-exercise delays neuropathic pain in experimental diabetes: effects on voltage-activated calcium channels. 2011 , 118, 224-36	50
853	Central and peripheral sites of action for CB ₁ receptor mediated analgesic activity in chronic inflammatory and neuropathic pain models in rats. 2011 , 162, 428-40	97
852	Predictive validity of behavioural animal models for chronic pain. 2011 , 164, 1195-206	112
851	A spinal mechanism of action for duloxetine in a rat model of painful diabetic neuropathy. 2011 , 164, 159-69	42
850	Longitudinal study of painful diabetic neuropathy in the Zucker diabetic fatty rat model of type 2 diabetes: impaired basal G-protein activity appears to underpin marked morphine hyposensitivity at 6 months. 2011 , 12, 437-50	21
849	Insulin implants prevent the temporal development of mechanical allodynia and opioid hyposensitivity for 24-wks in streptozotocin (STZ)-diabetic Wistar rats. 2011 , 12, 782-93	11
848	Botulinum toxin decreases hyperalgesia and inhibits P2X3 receptor over-expression in sensory neurons induced by ventral root transection in rats. 2011 , 12, 1385-94	34
847	Effect of the cation-chloride cotransporter inhibitor furosemide in a rat model of postoperative pain. 2011 , 12, 1427-34	4
846	Neuronal CCL21 up-regulates microglia P2X4 expression and initiates neuropathic pain development. 2011 , 30, 1864-73	117
845	Assessing nociceptive sensitivity in mouse models of inflammatory and neuropathic trigeminal pain. <i>Journal of Neuroscience Methods</i> , 2011 , 201, 46-54	3 40
844	A novel mechanism of inhibition of high-voltage activated calcium channels by β -conotoxins contributes to relief of nerve injury-induced neuropathic pain. 2011 , 152, 259-266	69
843	Pain tests provoke modality-specific cardiovascular responses in awake, unrestrained rats. 2011 , 152, 274-284	12

842	Cognitive-emotional sensitization contributes to wind-up-like pain in phantom limb pain patients. 2011 , 152, 157-162	77
841	Behavioral indices of ongoing pain are largely unchanged in male mice with tissue or nerve injury-induced mechanical hypersensitivity. 2011 , 152, 990-1000	125
840	Calcium channel alpha-2-delta-1 protein upregulation in dorsal spinal cord mediates spinal cord injury-induced neuropathic pain states. 2011 , 152, 649-655	96
839	Increased hippocampal glycine uptake and cognitive dysfunction after peripheral nerve injury. 2011 , 152, 809-817	59
838	Genetic ablation of delta opioid receptors in nociceptive sensory neurons increases chronic pain and abolishes opioid analgesia. 2011 , 152, 1238-1248	118
837	A novel slow-inactivation-specific ion channel modulator attenuates neuropathic pain. 2011 , 152, 833-843	59
836	Statins alleviate experimental nerve injury-induced neuropathic pain. 2011 , 152, 1033-1043	55
835	Neuropathic pain- and glial derived neurotrophic factor-associated regulation of cadherins in spinal circuits of the dorsal horn. 2011 , 152, 924-935	19
834	GDNF levels in the lower lip skin in a rat model of trigeminal neuropathic pain: implications for nonpeptidergic fiber reinnervation and parasympathetic sprouting. 2011 , 152, 1502-1510	35
833	Oxaliplatin elicits mechanical and cold allodynia in rodents via TRPA1 receptor stimulation. 2011 , 152, 1621-1631	220
832	Lesion of the rostral anterior cingulate cortex eliminates the aversiveness of spontaneous neuropathic pain following partial or complete axotomy. 2011 , 152, 1641-1648	149
831	GRK2 in sensory neurons regulates epinephrine-induced signalling and duration of mechanical hyperalgesia. 2011 , 152, 1649-1658	40
830	Antinociceptive effect of Brazilian armed spider venom toxin Tx3-3 in animal models of neuropathic pain. 2011 , 152, 2224-2232	45
829	Contribution of afferent pathways to nerve injury-induced spontaneous pain and evoked hypersensitivity. 2011 , 152, 1997-2005	74
828	Engagement of descending inhibition from the rostral ventromedial medulla protects against chronic neuropathic pain. 2011 , 152, 2701-2709	149
827	Genotype-selective phenotypic switch in primary afferent neurons contributes to neuropathic pain. 2011 , 152, 2413-2426	50
826	Chronic neuropathic pain-like behavior correlates with IL-1 β expression and disrupts cytokine interactions in the hippocampus. 2011 , 152, 2827-2835	90
825	Spinal TLR4 mediates the transition to a persistent mechanical hypersensitivity after the resolution of inflammation in serum-transferred arthritis. 2011 , 152, 2881-2891	102

824	Monosodium iodoacetate-induced osteoarthritis produces pain-depressed wheel running in rats: implications for preclinical behavioral assessment of chronic pain. 2011 , 98, 35-42	62
823	Blockade of 5-HT7 receptors reduces tactile allodynia in the rat. 2011 , 99, 591-7	31
822	Analgesic effects of the ethanolic extract from <i>Magnolia ovata</i> (Magnoliaceae) trunk bark and of N-acetylxylpine, a semi-synthetic analogue of xylpine. 2011 , 18, 143-7	4
821	The impact of prenatal stress on basal nociception and evoked responses to tail-docking and inflammatory challenge in juvenile pigs. 2011 , 104, 728-37	28
820	Delayed postoperative latent pain sensitization revealed by the systemic administration of opioid antagonists in mice. 2011 , 657, 89-96	50
819	Chromaffin cell transplant in spinal cord reduces secondary allodynia induced by formalin in the rat. Role of opioid receptors and β -adrenoceptors. 2011 , 668, 147-54	4
818	Antinociceptive effects induced through the stimulation of spinal cannabinoid type 2 receptors in chronically inflamed mice. 2011 , 668, 184-9	14
817	Involvement of mast cells in a mouse model of postoperative pain. 2011 , 672, 88-95	54
816	Rac1-regulated dendritic spine remodeling contributes to neuropathic pain after peripheral nerve injury. 2011 , 232, 222-33	55
815	Role of peripheral polyamines in the development of inflammatory pain. 2011 , 82, 269-77	17
814	Potentiation of analgesic efficacy but not side effects: co-administration of an α 4 β 2 neuronal nicotinic acetylcholine receptor agonist and its positive allosteric modulator in experimental models of pain in rats. 2011 , 82, 967-76	45
813	α 4 β 2 neuronal nicotinic receptor positive allosteric modulation: an approach for improving the therapeutic index of α 4 β 2 nAChR agonists in pain. 2011 , 82, 959-66	63
812	Identification and SAR of selective inducible nitric oxide synthase (iNOS) dimerization inhibitors. 2011 , 21, 6888-94	9
811	Pregabalin attenuates place escape/avoidance behavior in a rat model of spinal cord injury. 2011 , 1370, 129-35	42
810	Novel peptide ligands with dual acting pharmacophores designed for the pathophysiology of neuropathic pain. 2011 , 1395, 1-11	27
809	Activation of spinal and supraspinal cannabinoid-1 receptors leads to antinociception in a rat model of neuropathic spinal cord injury pain. 2011 , 1412, 44-54	44
808	Taltirelin, a thyrotropin-releasing hormone analog, alleviates mechanical allodynia through activation of descending monoaminergic neurons in persistent inflammatory pain. 2011 , 1414, 50-7	6
807	Efficacy of impropgan, a non-opioid analgesic, in neuropathic pain. 2011 , 1424, 32-7	1

806	Adenosine triphosphate-sensitive potassium channel blockers attenuate the antiallodynic effect of R-PIA in neuropathic rats. 2011 , 112, 1494-9	5
805	Intra-arterial tert-Butyl-hydroperoxide infusion induces an exacerbated sensory response in the rat hind limb and is associated with an impaired tissue oxygen uptake. 2011 , 34, 49-57	
804	Anti-hyperalgesic effect of CaMKII inhibitor is associated with downregulation of phosphorylated CREB in rat spinal cord. 2011 , 25, 87-92	18
803	Etanercept attenuates pain-related behavior following compression of the dorsal root ganglion in the rat. 2011 , 20, 1877-84	26
802	Systemic and spinal administration of etanercept, a tumor necrosis factor alpha inhibitor, blocks tactile allodynia in diabetic mice. 2011 , 48, 135-42	43
801	Electrophysiological properties of spinal wide dynamic range neurons in neuropathic pain rats following spinal nerve ligation. 2011 , 27, 1-8	14
800	Attenuating effect of hydroalcoholic extract of Acorus calamus in vincristine-induced painful neuropathy in rats. 2011 , 65, 480-7	32
799	Role of TRPM8 in dorsal root ganglion in nerve injury-induced chronic pain. 2011 , 12, 120	83
798	Attenuating effect of Acorus calamus extract in chronic constriction injury induced neuropathic pain in rats: an evidence of anti-oxidative, anti-inflammatory, neuroprotective and calcium inhibitory effects. 2011 , 11, 24	49
797	Enhanced neuroinflammation and pain hypersensitivity after peripheral nerve injury in rats expressing mutated superoxide dismutase 1. 2011 , 8, 33	33
796	Inhibition of spinal astrocytic c-Jun N-terminal kinase (JNK) activation correlates with the analgesic effects of ketamine in neuropathic pain. 2011 , 8, 6	37
795	Increased local concentration of complement C5a contributes to incisional pain in mice. 2011 , 8, 80	37
794	Temporal control of gene deletion in sensory ganglia using a tamoxifen-inducible Advillin-Cre-ERT2 recombinase mouse. 2011 , 7, 100	46
793	Differential expression of microRNAs in mouse pain models. 2011 , 7, 17	124
792	Analgesic tone conferred by constitutively active mu opioid receptors in mice lacking β -arrestin 2. 2011 , 7, 24	19
791	Peripheral effects of morphine and expression of μ opioid receptors in the dorsal root ganglia during neuropathic pain: nitric oxide signaling. 2011 , 7, 25	36
790	TRESK channel contribution to nociceptive sensory neurons excitability: modulation by nerve injury. 2011 , 7, 30	103
789	Activation of ATP-sensitive potassium channels antagonize nociceptive behavior and hyperexcitability of DRG neurons from rats. 2011 , 7, 35	29

788	Monosodium iodoacetate-induced joint pain is associated with increased phosphorylation of mitogen activated protein kinases in the rat spinal cord. 2011, 7, 39	41
787	Transient inflammation-induced ongoing pain is driven by TRPV1 sensitive afferents. 2011, 7, 4	72
786	Glial TNF α in the spinal cord regulates neuropathic pain induced by HIV gp120 application in rats. 2011, 7, 40	79
785	Central nervous system mast cells in peripheral inflammatory nociception. 2011, 7, 42	52
784	Cancer-induced bone pain sequentially activates the ERK/MAPK pathway in different cell types in the rat spinal cord. 2011, 7, 48	57
783	Reduced spinal microglial activation and neuropathic pain after nerve injury in mice lacking all three nitric oxide synthases. 2011, 7, 50	35
782	Mechanisms underlying purinergic P2X3 receptor-mediated mechanical allodynia induced in diabetic rats. 2011, 7, 60	52
781	Targeting adenosine monophosphate-activated protein kinase (AMPK) in preclinical models reveals a potential mechanism for the treatment of neuropathic pain. 2011, 7, 70	155
780	Genetic evidence for an essential role of neuronally expressed IL-6 signal transducer gp130 in the induction and maintenance of experimentally induced mechanical hypersensitivity in vivo and in vitro. 2011, 7, 73	30
779	Inhibition of p38-MAPK signaling pathway attenuates breast cancer induced bone pain and disease progression in a murine model of cancer-induced bone pain. 2011, 7, 81	27
778	Autophagy impairment in a mouse model of neuropathic pain. 2011, 7, 83	59
777	Genetic enhancement of behavioral itch responses in mice lacking phosphoinositide 3-kinase- α (PI3K α) 2011, 7, 96	7
776	PKM ζ is essential for spinal plasticity underlying the maintenance of persistent pain. 2011, 7, 99	87
775	<i>Emblica officinalis</i> corrects functional, biochemical and molecular deficits in experimental diabetic neuropathy by targeting the oxido-nitrosative stress mediated inflammatory cascade. 2011, 25, 1527-36	41
774	NR2B phosphorylation at tyrosine 1472 in spinal dorsal horn contributed to N-methyl-D-aspartate-induced pain hypersensitivity in mice. 2011, 89, 1869-76	30
773	Gait and behavior in an IL1 α -mediated model of rat knee arthritis and effects of an IL1 antagonist. 2011, 29, 694-703	19
772	Functional characterization of a chronic cyclophosphamide-induced overactive bladder model in mice. 2011, 30, 1659-65	56
771	A potent and selective indole N-type calcium channel (Ca(v)2.2) blocker for the treatment of pain. 2011, 21, 869-73	21

770	Insular cortex lesion diminishes neuropathic and inflammatory pain-like behaviours. 2011 , 15, 132-8		41
769	Amelioration of functional, biochemical and molecular deficits by epigallocatechin gallate in experimental model of alcoholic neuropathy. 2011 , 15, 286-92		20
768	Prevention of oxaliplatin-induced mechanical allodynia and neurodegeneration by neurotrophin in the rat model. 2011 , 15, 344-50		54
767	A novel rat forelimb model of neuropathic pain produced by partial injury of the median and ulnar nerves. 2011 , 15, 459-66		9
766	Postnatal expression of the homeobox gene <i>Prrxl1</i> (<i>Drg11</i>) is increased in inflammatory but not neuropathic pain. 2011 , 15, 477-81		5
765	Direct injection into the dorsal root ganglion: technical, behavioral, and histological observations. <i>Journal of Neuroscience Methods</i> , 2011 , 199, 43-55	3	61
764	Intrathecal PKA-selective siRNA treatment blocks sustained morphine-mediated pain sensitization and antinociceptive tolerance in rats. <i>Journal of Neuroscience Methods</i> , 2011 , 199, 62-8	3	14
763	Intrathecal lentiviral-mediated RNA interference targeting PKC α attenuates chronic constriction injury-induced neuropathic pain in rats. 2011 , 22, 465-75		40
762	Schwann cell proliferation and macrophage infiltration are evident at day 14 after painful cervical nerve root compression in the rat. 2011 , 28, 2429-38		26
761	Transplantation of human glial restricted progenitors and derived astrocytes into a contusion model of spinal cord injury. 2011 , 28, 579-94		57
760	Intervertebral disc degeneration-induced expression of pain-related molecules: glial cell-derived neurotrophic factor as a key factor. 2011 , 23, 329-34		20
759	Amelioration of neuropathic pain by novel transient receptor potential vanilloid 1 antagonist AS1928370 in rats without hyperthermic effect. 2011 , 336, 743-50		68
758	Repeated vulvovaginal fungal infections cause persistent pain in a mouse model of vulvodynia. 2011 , 3, 101ra91		91
757	Chronic intrathecal infusion of gabapentin prevents nerve ligation-induced pain in rats. 2011 , 106, 699-705		25
756	Spinal phosphoinositide 3-kinase-Akt-mammalian target of rapamycin signaling cascades in inflammation-induced hyperalgesia. 2011 , 31, 2113-24		104
755	Involvement of calcitonin gene-related peptide and CCL2 production in CD40-mediated behavioral hypersensitivity in a model of neuropathic pain. 2011 , 7, 117-28		15
754	Peripheral nerve injury alters blood-spinal cord barrier functional and molecular integrity through a selective inflammatory pathway. 2011 , 31, 10819-28		177
753	Caudal granular insular cortex is sufficient and necessary for the long-term maintenance of allodynic behavior in the rat attributable to mononeuropathy. 2011 , 31, 6317-28		42

752	JAK-STAT3 pathway regulates spinal astrocyte proliferation and neuropathic pain maintenance in rats. 2011 , 134, 1127-39	193
751	Hindlimb muscle atrophy occurs from peripheral nerve damage in a rat neuropathic pain model. 2011 , 13, 44-54	5
750	Neuronal loss in the rostral ventromedial medulla in a rat model of neuropathic pain. 2011 , 31, 17028-39	62
749	Spinal cord Toll-like receptor 4 mediates inflammatory and neuropathic hypersensitivity in male but not female mice. 2011 , 31, 15450-4	307
748	Microglial Ca(2+)-activated K(+) channels are possible molecular targets for the analgesic effects of S-ketamine on neuropathic pain. 2011 , 31, 17370-82	69
747	Axon regeneration can facilitate or suppress hindlimb function after olfactory ensheathing glia transplantation. 2011 , 31, 4298-310	50
746	CCL2 released from neuronal synaptic vesicles in the spinal cord is a major mediator of local inflammation and pain after peripheral nerve injury. 2011 , 31, 5865-75	147
745	Dynamic changes in the microRNA expression profile reveal multiple regulatory mechanisms in the spinal nerve ligation model of neuropathic pain. 2011 , 6, e17670	104
744	p38/MAPK inhibitor modulates the expression of dorsal horn GABA(B) receptors in the spinal nerve ligation model of neuropathic pain. 2011 , 18, 150-5	25
743	Neuropeptides contribute to peripheral nociceptive sensitization by regulating interleukin-1 β production in keratinocytes. 2011 , 113, 175-83	53
742	Tonic inhibition of chronic pain by neuropeptide Y. 2011 , 108, 7224-9	89
741	Inter-regional contribution of enhanced activity of the primary somatosensory cortex to the anterior cingulate cortex accelerates chronic pain behavior. 2011 , 31, 7631-6	81
740	Perspectives on: information and coding in mammalian sensory physiology: probing mammalian touch transduction. 2011 , 138, 291-301	15
739	Distinct mechanisms underlying pronociceptive effects of opioids. 2011 , 31, 16748-56	55
738	Antinociceptive effect of riluzole in rats with neuropathic spinal cord injury pain. 2011 , 28, 127-34	41
737	Reduced expression of SynGAP, a neuronal GTPase-activating protein, enhances capsaicin-induced peripheral sensitization. 2011 , 106, 309-18	6
736	Antinociceptive Activity of Trichilia catigua Hydroalcoholic Extract: New Evidence on Its Dopaminergic Effects. 2011 , 2011, 120820	21
735	CXCL5 mediates UVB irradiation-induced pain. 2011 , 3, 90a60	82

734	The animal model of spinal cord injury as an experimental pain model. 2011 , 2011, 939023	41
733	Evaluation of cutaneous allodynia following induction of cortical spreading depression in freely moving rats. 2011 , 31, 1090-100	38
732	A Comparison of the α /3/5 Selective Positive Allosteric Modulators L-838,417 and TPA023 in Preclinical Models of Inflammatory and Neuropathic Pain. 2011 , 2011, 608912	21
731	Retinoic acid reduces chemotherapy-induced neuropathy in an animal model and patients with lung cancer. 2011 , 77, 987-95	51
730	CCL2 and CCL3 are essential mediators of pelvic pain in experimental autoimmune prostatitis. 2012 , 303, R580-9	46
729	Modification of rat model of sciatica induced by lumbar disc herniation and the anti-inflammatory effect of osthole given by epidural catheterization. 2012 , 90, 251-63	11
728	The Inhibition of Spinal Astrocytic JAK2-STAT3 Pathway Activation Correlates with the Analgesic Effects of Triptolide in the Rat Neuropathic Pain Model. 2012 , 2012, 185167	32
727	Microinjection of HSV-1 amplicon vector-mediated human proenkephalin into the periaqueductal grey attenuates neuropathic pain in rats. 2012 , 122, 189-94	4
726	Sphingosine 1-phosphate receptor 2 antagonist JTE-013 increases the excitability of sensory neurons independently of the receptor. 2012 , 108, 1473-83	16
725	Microglial cathepsin B contributes to the initiation of peripheral inflammation-induced chronic pain. 2012 , 32, 11330-42	57
724	Anti-inflammatory mesenchymal stem cells (MSC2) attenuate symptoms of painful diabetic peripheral neuropathy. 2012 , 1, 557-65	58
723	Time-dependent cross talk between spinal serotonin 5-HT _{2A} receptor and mGluR1 subserves spinal hyperexcitability and neuropathic pain after nerve injury. 2012 , 32, 13568-81	27
722	P2Y ₁ purinoceptor inhibition reduces extracellular signal-regulated protein kinase 1/2 phosphorylation in spinal cord and dorsal root ganglia: implications for cancer-induced bone pain. 2012 , 44, 367-72	24
721	μ Opioid mechanisms for ADL5747 and ADL5859 effects in mice: analgesia, locomotion, and receptor internalization. 2012 , 342, 799-807	51
720	Effect of acute versus continuous glycemic control on duration of local anesthetic sciatic nerve block in diabetic rats. 2012 , 37, 595-600	15
719	TRPC1 contributes to light-touch sensation and mechanical responses in low-threshold cutaneous sensory neurons. 2012 , 107, 913-22	62
718	Up-regulation of Cav β subunit in primary sensory neurons increases voltage-activated Ca ²⁺ channel activity and nociceptive input in neuropathic pain. 2012 , 287, 6002-13	27
717	Distinct Nav1.7-dependent pain sensations require different sets of sensory and sympathetic neurons. 2012 , 3, 791	188

716	TRPC3 and TRPC6 are essential for normal mechanotransduction in subsets of sensory neurons and cochlear hair cells. 2012 , 2, 120068	106
715	Enhanced GABAergic activity in the mouse primary somatosensory cortex is insufficient to alleviate chronic pain behavior with reduced expression of neuronal potassium-chloride cotransporter. 2012 , 32, 16552-9	30
714	Peripheral inflammatory disease associated with centrally activated IL-1 system in humans and mice. 2012 , 109, 12728-33	105
713	Cyclin-dependent kinase 5 controls TRPV1 membrane trafficking and the heat sensitivity of nociceptors through KIF13B. 2012 , 32, 14709-21	65
712	N-methyl-D-aspartate receptor- and calpain-mediated proteolytic cleavage of K ⁺ -Cl ⁻ cotransporter-2 impairs spinal chloride homeostasis in neuropathic pain. 2012 , 287, 33853-64	101
711	Murine immunodeficiency virus-induced peripheral neuropathy and the associated cytokine responses. 2012 , 189, 3724-33	20
710	Differential association of receptor-Gβγ complexes with β-arrestin2 determines recycling bias and potential for tolerance of μ-opioid receptor agonists. 2012 , 32, 4827-40	53
709	Formulation, Optimization and In Vivo Evaluation of Fucoidan-Based Cream with Anti-Inflammatory Properties. 2021 , 19,	5
708	Relating spinal injury-induced neuropathic pain and spontaneous afferent activity to sleep and respiratory dysfunction.	
707	Antinociceptive and Antipruritic Effects of HSK21542, a Peripherally-Restricted Kappa Opioid Receptor Agonist, in Animal Models of Pain and Itch. 2021 , 12, 773204	0
706	Role of Adenosine Kinase in Sphingosine-1-Phosphate Receptor 1-Induced Mechano-Hypersensitivities. 2021 , 1	0
705	Effects of Ozone on Hippocampus BDNF and Fos Expressions in Rats with Chronic Compression of Dorsal Root Ganglia. 2021 , 2021, 5572915	1
704	Innovations and advances in modelling and measuring pain in animals. 2021 ,	10
703	Contextual control of conditioned pain tolerance and endogenous analgesic systems: Evidence for sex-based differences in endogenous opioid engagement.	
702	Suppression of P2X3 receptor-mediated currents by the activation of β-adrenergic receptors in rat dorsal root ganglion neurons. 2021 ,	1
701	β-helmenene relieves neuropathic pain in mice through the regulation on C-X-C motif chemokine receptor 3 and GABAA receptor. 2021 ,	1
700	Astaxanthin alleviates inflammatory pain by regulating the p38 mitogen-activated protein kinase and nuclear factor-erythroid factor 2-related factor/heme oxygenase-1 pathways in mice. 2021 ,	0
699	Overexpression of RAPGEF3 enhances the therapeutic effect of dezocine in treatment of neuropathic pain. 2021 , 44, e20200463	0

- 698 Inhibiting spinal secretory phospholipase A after painful nerve root injury attenuates established pain and spinal neuronal hyperexcitability by altering spinal glutamatergic signaling.. **2021**, 17, 17448069211066221 ⁰
- 697 Abnormal Somatosensory Behaviors Associated With a Gain-of-Function Mutation in TRPV3 Channels.. **2021**, 14, 790435 2
- 696 Glutamate in primary afferents is required for itch transmission.. **2021**, 2
- 695 Pain-like behavior in the collagen antibody-induced arthritis model is regulated by lysophosphatidic acid and activation of satellite glia cells.. **2022**, 101, 214-230 1
- 694 The glial cell's role in antinociceptive differential effects of oxytocin upon female and male rats.. **2022**, 0
- 693 Association Between the Accumulation of Pentosidine at the Sciatic Nerve and Cutaneous Hindpaw Hypersensitivity in a Rat Ovariectomy Model.. **2022**, 14, e21059
- 692 Targeted up-regulation of Drp1 in dorsal horn attenuates neuropathic pain hypersensitivity by increasing mitochondrial fission.. **2021**, 49, 102216 0
- 691 LncRNA MRAK159688 facilitates morphine tolerance by promoting REST-mediated inhibition of mu opioid receptor in rats.. **2022**, 206, 108938 0
- 690 Pharmacopuncture of Bauhinia variegata Nanoemulsion Formulation against Diabetic Peripheral Neuropathic Pain. **2020**, 23, 30-36 2
- 689 Kinin B1 receptor is involved in mechanical nociception in a fibromyalgia-like model in mice. 1, 1431
- 688 The Convergent Neuroscience of Affective Pain and Substance Use Disorder.. **2021**, 41, 14 0
- 687 EXPERIMENTAL STUDY OF NEUROPATHIC PAIN TREATMENT UNDER MULTIPLE SCLEROSIS MODELING. **2021**, 4, 141
- 686 A Guide to Preclinical Models of Zoster-Associated Pain and Postherpetic Neuralgia. **2021**, 0
- 685 Suppression of joint pain in transient receptor potential vanilloid 4 knockout rats with monoiodoacetate-induced osteoarthritis. **2021**, 6, e951 2
- 684 Thermal hyperalgesia and dynamic weight bearing share similar recovery dynamics in a sciatic nerve entrapment injury model.. **2021**, 10, 100079 0
- 683 Structure-based design of a chemical probe set for the 5-HT5A serotonin receptor.
- 682 Effect of Sensory Deprivation of Nasal Respiratory on Behavior of C57BL/6J Mice.. **2021**, 11, 0
- 681 Ovariectomy increases paclitaxel-induced mechanical hypersensitivity and reduces anti-inflammatory CD4+ T cells in the dorsal root ganglion of female mice.

- 680 Nerve decompression surgery suppresses TNF- α expression and T cell infiltration in a rat sciatic nerve chronic constriction injury model.. **2022**,
- 679 Blockage of HCN Channels Inhibits the Function of P2X Receptors in Rat Dorsal Root Ganglion Neurons.. **2022**, 47, 1083 ○
- 678 From Initiation to Maintenance: HIV-1 Gp120-induced Neuropathic Pain Exhibits Different Molecular Mechanisms in the Mouse Spinal Cord Via Bioinformatics Analysis Based on RNA Sequencing.. **2022**, 1
- 677 Small molecule targeting NaV1.7 via inhibition of the CRMP2-Ubc9 interaction reduces pain in chronic constriction injury (CCI) rats.. **2022**, 16, 1-8 ○
- 676 Mouse Models of Osteoarthritis: A Summary of Models and Outcomes Assessment.. **2022**, ○
- 675 Design and synthesis of pyrido[2,3-d]pyrimidine derivatives for a novel PAC1 receptor antagonist.. **2022**, 231, 114160 ○
- 674 A female-specific role for Calcitonin Gene-Related Peptide (CGRP) in rodent pain models.. **2022**, 2
- 673 Protectin DX Attenuates Lumbar Radicular Pain of Non-compressive Disc Herniation by Autophagy Flux Stimulation Adenosine Monophosphate-Activated Protein Kinase Signaling.. **2021**, 12, 784653
- 672 Transcranial Direct Current Stimulation Alleviates the Chronic Pain of Osteoarthritis by Modulating NMDA Receptors in Midbrain Periaqueductal Gray in Rats.. **2022**, 15, 203-214 ○
- 671 Effects of β -Carotin and Green Tea Powder Diets on Alleviating the Symptoms of Gouty Arthritis and Improving Gut Microbiota in C57BL/6 Mice.. **2022**, 13, 837182 ○
- 670 Nuclear Factor κ B-COX2 Pathway Activation in Non-myelinating Schwann Cells Is Necessary for the Maintenance of Neuropathic Pain .. **2021**, 15, 782275 ○
- 669 Pharmacological interactions between intrathecal pregabalin plus tianeptine or clopidogrel in a rat model of neuropathic pain.. **2022**, 35, 59-65 ○
- 668 Evaluation of calcium-sensitive adenylyl cyclase AC1 and AC8 mRNA expression in the anterior cingulate cortex of mice with spared nerve injury neuropathy.. **2022**, 11, 100081 1
- 667 Transcriptomic Profiling in Mice With CB1 receptor Deletion in Primary Sensory Neurons Suggests New Analgesic Targets for Neuropathic Pain.. **2021**, 12, 781237 1
- 666 Targeting the A adenosine receptor to prevent and reverse chemotherapy-induced neurotoxicities in mice.. **2022**, 10, 11 ○
- 665 Increased Brain-Derived Neurotrophic Factor Levels in Cerebrospinal Fluid During the Acute Phase in TBI-Induced Mechanical Allodynia in the Rat Model.. **2022**, 15, 229-239 1
- 664 Lysophosphatidyl-choline 16: 0 mediates chronic joint pain associated to rheumatic diseases through acid-sensing ion channel 3.. **2022**, 2
- 663 Analgesic Effect of SH003 and Trichosanthes kirilowii Maximowicz in Paclitaxel-Induced Neuropathic Pain in Mice. **2022**, 44, 718-730 ○

- 662 CGRP Administration into the Cerebellum Evokes Migraine-like Behaviors Predominately in Female Mice.
- 661 Inhibition of Spinal Interleukin-33 Attenuates Peripheral Inflammation and Hyperalgesia in Experimental Arthritis.. **2022**, 1 0
- 660 BoNT/A alleviates neuropathic pain in osteoarthritis by down-regulating the expression of P2X4R in spinal microglia.. **2021**, 0
- 659 GPR55 and GPR119 Receptors Contribute to the Processing of Neuropathic Pain in Rats.. **2022**, 15, 1
- 658 The possibility of alleviating chronic neuropathic pain and related behaviors by the direct suppression of the parabrachial nucleus.. **2021**, 95, 180-187 0
- 657 The Role of Transient Receptor Potential A1 and G Protein-Coupled Receptor 39 in Zinc-Mediated Acute and Chronic Itch in Mice.. **2021**, 14, 768731
- 656 Neuronal allodynic mechanisms of Slc7a5 (LAT1) in the spared nerve injury rodent model of neuropathic pain.. **2022**, 474, 397 1
- 655 Methods and protocols for translatable rodent models of postsurgical pain.. **2022**, 168, 249-276
- 654 Preconditioning by voluntary wheel running attenuates later neuropathic pain via Nrf2 antioxidant signaling in rats.. **2022**, 1
- 653 Reversal of neuropathic pain is associated with corticostriatal functional reorganization after nerve repair in the spared nerve injury model.. **2022**,
- 652 Inhibition of oligodendrocyte apoptosis in the prelimbic medial prefrontal cortex prevents fentanyl-induced hyperalgesia in rats.. **2022**, 0
- 651 SAHA Inhibits Somatic Hyperalgesia Induced by Stress Combined with Orofacial Inflammation Through Targeting Different Spinal 5-HT Receptor Subtypes.. **2022**, 1
- 650 In-Depth Characterization of Somatic and Orofacial Sensitive Dysfunctions and Interfering-Symptoms in a Relapsing-Remitting Experimental Autoimmune Encephalomyelitis Mouse Model.. **2021**, 12, 789432 0
- 649 Nicotinamide Riboside Alleviates Corneal and Somatic Hypersensitivity Induced by Paclitaxel in Male Rats.. **2022**, 63, 38 1
- 648 The Effect of Melatonin on Radicular Pain in a Rat Model of Lumbar Disc Herniation.. **2022**,
- 647 Characterization of pain-related behaviors, changes in bone microarchitecture and sensory innervation induced by chronic cadmium exposure in adult mice.. **2022**, 89, 99-99 0
- 646 Sex-dependent pain trajectories induced by prolactin require an inflammatory response for pain resolution.. **2022**, 0
- 645 Development of hepatic impairment aggravates chemotherapy-induced peripheral neuropathy following oxaliplatin treatment: Evidence from clinical and preclinical studies.. **2022**, 148, 315-325

644	The analgesic properties of Yu-Xue-Bi tablets in the inflammatory pain mice: By the inhibition of CCL3-mediated macrophage transmigration into the spinal cord.. 2022 , 289, 115051	0
643	Characterisation of the pathophysiology of neuropathy and sensory dysfunction in a mouse model of recessive dystrophic epidermolysis bullosa.. 2022 ,	
642	The efficacy of duloxetine depends on spinal cholinergic plasticity in neuropathic pain model rats.. 2022 , 12, 188-196	1
641	Joint Damage and Neuropathic Pain in Rats Treated With Lysophosphatidic Acid.. 2022 , 13, 811402	0
640	Schwann cell endosome CGRP signals elicit periorbital mechanical allodynia in mice.. 2022 , 13, 646	2
639	Alfaxalone improved in acute stress-induced tactile hypersensitivity and anxiety-like behavior in mice.. 2022 ,	1
638	A diencephalic circuit in rats for opioid analgesia but not positive reinforcement.. 2022 , 13, 764	1
637	The Surface Amine Group of Ultrasmall Magnetic Iron Oxide Nanoparticles Produce Analgesia in the Spinal Cord and Decrease Long-Term Potentiation.. 2022 , 14,	
636	Cdyl Deficiency Brakes Neuronal Excitability and Nociception through Promoting Kcnb1 Transcription in Peripheral Sensory Neurons.. 2022 , e2104317	1
635	Kindlin-2 preserves integrity of the articular cartilage to protect against osteoarthritis.	2
634	An Improved Model of Type 2 Diabetes with Effects on Glucose Tolerance, Neuropathy and Retinopathy with and without Obesity.. 2022 , 113740	0
633	Medullary kappa-opioid receptor neurons inhibit pain and itch through a descending circuit.	0
632	Spinal cord-wide structural disruption in type 2 diabetes rescued by exenatide "a glucagon-like peptide-1 analogue" via down-regulating inflammatory, oxidative stress and apoptotic signaling pathways.. 2022 , 121, 102079	0
631	Chronic electrical stimulation of peripheral nerves via deep-red light transduced by an implanted organic photocapacitor.. 2021 ,	10
630	Structures of the μ receptor enable docking for bioactive ligand discovery. 2021 ,	24
629	Cholinergic modulation is independent of T lymphocytes in a mouse model of neuropathic pain.. 2022 , 18, 17448069221076634	
628	Methods and protocols for chemotherapy-induced peripheral neuropathy (CIPN) mouse models using paclitaxel.. 2022 , 168, 277-298	0
627	Effect of Pregabalin and Gabapentin on Nociceptive Behaviors Induced by Spinal Nerve Ligation. 2022 , 10, 114-123	

626	Structure-Based Design of a Chemical Probe Set for the 5-HT Serotonin Receptor.. 2022 ,	3
625	Epidural Posterior Insular Stimulation Alleviates Neuropathic Pain Manifestations in Rats With Spared Nerve Injury Through Endogenous Opioid System.. 2022 ,	0
624	The Antinociceptive Potential of Leaf Extract, (-)-Epicatechin, and Rutin against Chronic Constriction Injury-Induced Neuropathic Pain in Rats.. 2022 , 11,	0
623	Identification of Altered Evoked and Non-Evoked Responses in a Heterologous Mouse Model of Endometriosis-Associated Pain.. 2022 , 10,	1
622	The circ_0002538/miR-138-5p/PLLP axis regulates Schwann cell migration and myelination in diabetic peripheral neuropathy.	0
621	Macrophages transfer mitochondria to sensory neurons to resolve inflammatory pain.. 2021 ,	6
620	Mechanical Conflict-Avoidance Assay to Measure Pain Behavior in Mice.. 2022 ,	0
619	Nrf2 Activation Mediates Antiallodynic Effect of Electroacupuncture on a Rat Model of Complex Regional Pain Syndrome Type-I through Reducing Local Oxidative Stress and Inflammation.. 2022 , 2022, 8035109	2
618	Intravenous administration of human mesenchymal stem cells derived from adipose tissue and umbilical cord improves neuropathic pain via suppression of neuronal damage and anti-inflammatory actions in rats.. 2022 , 17, e0262892	0
617	Exploring the Correlation Between the Regulation of Macrophages by Regulatory T Cells and Peripheral Neuropathic Pain.. 2022 , 16, 813751	0
616	Development of Dibenzothiazepine Derivatives as Multifunctional Compounds for Neuropathic Pain.. 2022 , 15,	
615	RGMA signal in Macrophages Induces Neutrophil-related Astrocytopathy in NMO.. 2022 ,	0
614	Migraine susceptibility is modulated by food triggers and analgesic overuse via sulfotransferase inhibition.. 2022 , 23, 36	1
613	Sirtuin 2 Alleviates Chronic Neuropathic Pain by Suppressing Ferroptosis in Rats.. 2022 , 13, 827016	1
612	Contextual control of conditioned pain tolerance and endogenous analgesic systems.. 2022 , 11,	1
611	Electroacupuncture Reduces Visceral Pain Via Cannabinoid CB2 Receptors in a Mouse Model of Inflammatory Bowel Disease.. 2022 , 13, 861799	0
610	Analysis of Potential Hub Genes for Neuropathic Pain Based on Differential Expression in Rat Models.. 2022 , 2022, 6571987	
609	Duration of reduction in enduring stress-induced hyperalgesia via FKBP51 inhibition depends on timing of administration relative to traumatic stress exposure.. 2022 ,	1

- 608 Site of Nerve Division Affects Pain-Related Behavior and Spinal Cord Glial Proliferation after C7 Neurotomy in a Rat Stroke Model.. **2022**, 2022, 7446482
- 607 Transient receptor potential melastatin 8 (TRPM8) is required for nitroglycerin and calcitonin gene-related peptide induced migraine-like pain behaviors in mice.. **2022**, 0
- 606 Secondary auditory cortex mediates a sensorimotor mechanism for action timing.. **2022**, 25, 330-344 1
- 605 High-Throughput Screening for Extracellular Inhibitors of the FLT3 Receptor Tyrosine Kinase Reveals Chemically Diverse and Druggable Negative Allosteric Modulators.. **2022**, 0
- 604 Unbiased proteomic analysis detects painful systemic inflammatory profile in the serum of nerve injured mice.
- 603 Oxytocin inhibits hindpaw hyperalgesia induced by orofacial inflammation combined with stress.. **2022**, 17448069221089591 0
- 602 Somatostatin venom analogs evolved by fish-hunting cone snails: From prey capture behavior to identifying drug leads.. **2022**, 8, eabk1410 2
- 601 Translocator Protein (TSPO) Alleviates Neuropathic Pain by Activating Spinal Autophagy and Nuclear SIRT1/PGC-1 β Signaling in a Rat L5 SNL Model.. **2022**, 15, 767-778 2
- 600 Peripheral Neuropathy Phenotyping in Rat Models of Type 2 Diabetes Mellitus: Evaluating Uptake of the Neurodiab Guidelines and Identifying Future Directions.. **2022**, 46, 198-221 0
- 599 Translating outcomes from the clinical setting to preclinical models: chronic pain and functionality in chronic musculoskeletal pain.. **2022**, 1
- 598 CD8+ T cell-derived IL-13 increases macrophage IL-10 to resolve neuropathic pain.. **2022**, 7, 1
- 597 Loss of Stathmin-2, a hallmark of TDP-43-associated ALS, causes motor neuropathy. 0
- 596 Metformin Relieves Bortezomib-Induced Neuropathic Pain by Regulating AMPKa2-Mediated Autophagy in the Spinal Dorsal Horn.. **2022**, 1 1
- 595 Spinal 5-HT_{2A} Receptor is Involved in Electroacupuncture Inhibition of Chronic Pain.. **2022**, 17448069221087583
- 594 Chronic facial inflammatory pain-induced anxiety is associated with bilateral deactivation of the rostral anterior cingulate cortex.. **2022**,
- 593 The cAMP Response Element- Binding Protein/Brain-Derived Neurotrophic Factor Pathway in Anterior Cingulate Cortex Regulates Neuropathic Pain and Anxiodepression Like Behaviors in Rats.. **2022**, 15, 831151 2
- 592 Optimization of a Pyrimidinone Series for Selective Inhibition of Ca/Calmodulin-Stimulated Adenylyl Cyclase 1 Activity for the Treatment of Chronic Pain.. **2022**, 0
- 591 Dezocine relieves the postoperative hyperalgesia in rats through suppressing the hyper-action of Akt1/GSK-3 β pathway.. **2022**, 1

- 590 Conditioned Medium From the Stem Cells of Human Exfoliated Deciduous Teeth Ameliorates Neuropathic Pain in a Partial Sciatic Nerve Ligation Model.. **2022**, 13, 745020
- 589 Pain and Pain Management in Sea Turtle and Herpetological Medicine: State of the Art.. **2022**, 12, 0
- 588 Antinociceptive activity of the *Caesalpinia eriostachys* Benth. ethanolic extract, fractions, and isolated compounds in mice.
- 587 Effect of Riluzole on the Expression of HCN2 in Dorsal Root Ganglion Neurons of Diabetic Neuropathic Pain Rats.. **2022**, 2022, 8313415 1
- 586 The Insulin receptor regulates the persistence of mechanical nociceptive sensitization in flies and mice.
- 585 A spinal microglia population involved in remitting and relapsing neuropathic pain.. **2022**, 376, 86-90 9
- 584 The T-Type Calcium Channel Cav3.2 in Somatostatin Interneurons in Spinal Dorsal Horn Participates in Mechanosensation and Mechanical Allodynia in Mice.. **2022**, 16, 875726
- 583 Altered Spinal Homeostasis and Maladaptive Plasticity in GFAP Null Mice Following Peripheral Nerve Injury.. **2022**, 11, 0
- 582 Methacrylic acid-based biomaterials promote peripheral innervation in the subcutaneous space of mice.
- 581 Involvement of Opioid Peptides in the Analgesic Effect of Spinal Cord Stimulation in a Rat Model of Neuropathic Pain.. **2022**, 1 0
- 580 The small molecule compound C65780 alleviates pain by stabilizing voltage-gated sodium channels in the inactivated and slowly-recovering state.. **2022**, 109057
- 579 An Injectable Hydrogel for Treatment of Chronic Neuropathic Pain.. **2022**, e2100529 0
- 578 Genome-Wide Expression Profiling by RNA-Sequencing in Spinal Cord Dorsal Horn of a Rat Chronic Postsurgical Pain Model to Explore Potential Mechanisms Involved in Chronic Pain.. **2022**, 15, 985-1001
- 577 Sex-dependent antiallodynic effect of α -adrenergic receptor agonist tizanidine in rats with experimental neuropathic pain.. **2022**, 174855 0
- 576 The BDNF-TrkB signaling pathway in the rostral anterior cingulate cortex is involved in the development of pain aversion in rats with bone cancer via NR2B and ERK-CREB signaling.. **2022**, 0
- 575 Effects of heme oxygenase 1 in the molecular changes and neuropathy associated with type 2 diabetes in mice.. **2022**, 199, 114987 0
- 574 Calcitonin gene-related peptide is not involved in neuropathic pain induced by partial sciatic nerve ligation in mice.. **2022**, 136615
- 573 TET1 overexpression attenuates paclitaxel-induced neuropathic pain through rescuing K1.1 expression in primary sensory neurons of male rats.. **2022**, 297, 120486 1

572	Dorsal horn neuronal sparing predicts the development of at-level mechanical allodynia following cervical spinal cord injury in mice.. 2022 , 352, 114048	0
571	Trifluoro-icaritin alleviates chronic inflammatory pain through $\alpha 7$ nAChR-mediated suppression of HMGB1/NF- κ B signaling in the spinal cord of rats.. 2022 , 183, 13-26	0
570	Fuzi decoction ameliorates pain and cartilage degeneration of osteoarthritic rats through PI3K-Akt signaling pathway and its clinical retrospective evidence.. 2022 , 100, 154071	0
569	MicroRNA-489-3p attenuates neuropathic allodynia by regulating oncoprotein DEK/TET1-dependent epigenetic modification in the dorsal horn.. 2022 , 210, 109028	
568	Acetaminophen and pregabalin attenuate central sensitization in rodent models of nociplastic widespread pain.. 2022 , 210, 109029	0
567	Antibody-induced pain-like behavior and bone erosion: links to subclinical inflammation, osteoclast activity and ASIC3-dependent sensitization.. 2021 ,	3
566	α -Antitrypsin derived SP16 peptide demonstrates efficacy in rodent models of acute and neuropathic pain. 2022 , 36, e22093	2
565	PACAP-PAC1 Receptor Inhibition is Effective in Models of Opioid-Induced Medication Overuse Headache.	
564	Single-Dose P2 X4R Single-Chain Fragment Variable Antibody Permanently Reverses Chronic Pain in Male Mice.. 2021 , 22,	1
563	The Causal Role of Magnesium Deficiency in the Neuroinflammation, Pain Hypersensitivity and Memory/Emotional Deficits in Ovariectomized and Aged Female Mice.. 2021 , 14, 6633-6656	0
562	MicroRNA-155-5p promotes neuroinflammation and central sensitization via inhibiting SIRT1 in a nitroglycerin-induced chronic migraine mouse model. 2021 , 18, 287	3
561	EFFICACY OF P-COUMARIC ACID IN CHRONIC CONSTRICTION INJURY INDUCED NEUROPATHIC PAIN IN RATS. 2021 , 58, 52-58	
560	Hydrogen Sulfide Inhibits Inflammatory Pain and Enhances the Analgesic Properties of Delta Opioid Receptors.. 2021 , 10,	1
559	Human IAPP is a driver of painful diabetic peripheral neuropathy.	
558	Scorpion Venom peptide, AGAP inhibits TRPV1 and potentiates the analgesic effect of lidocaine.. 2021 , 7, e08560	2
557	Anthrax toxins regulate pain signaling and can deliver molecular cargoes into ANTXR2 DRG sensory neurons.. 2021 ,	3
556	Theta-burst stimulation of primary afferents drives long-term potentiation in the spinal cord and persistent pain via α 1-bound NMDA receptors. 2021 ,	4
555	Gait analysis as a robust pain behavioural endpoint in the chronic phase of the monoiodoacetate-induced knee joint pain in the rat.. 2022 , 33, 23-31	0

- 554 Repeated Low-Dose Acrolein Triggers Irreversible Lamina Propria Edema in Urinary Bladder, Transient Voiding Behavior and Widening of Eyes to Mechanical Stimuli.. **2021**, 10,
- 553 Effect of chrysin on mechanical hyperalgesia in chronic constriction injury-induced neuropathic pain in rat model. **2020**, 10, 189 1
- 552 Anti-inflammatory potential of liraglutide, a glucagon-like peptide-1 receptor agonist, in rats with peripheral acute inflammation.. **2022**, 1
- 551 Sex-Specific B Cell and Anti-Myelin Autoantibody Response After Peripheral Nerve Injury.. **2022**, 16, 835800 0
- 550 The role of PTEN in primary sensory neurons in processing itch and thermal information in mice.. **2022**, 39, 110724 0
- 549 Receptor-Interacting Protein Kinase 3 Inhibition Relieves Mechanical Allodynia and Suppresses NLRP3 Inflammasome and NF- κ B in a Rat Model of Spinal Cord Injury.. **2022**, 15, 861312 0
- 548 Long-term male-specific chronic pain via telomere- and p53-mediated spinal cord cellular senescence.. **2022**, 132, 1
- 547 Hedgehog signaling plays a crucial role in hyperalgesia associated with neuropathic pain in mice.. **2022**,
- 546 Botulinum toxin type a combined with transcranial direct current stimulation reverses the chronic pain induced by osteoarthritis in rats.. **2022**, 0
- 545 Behavioral and receptor expression studies on the primary somatosensory cortex and anterior cingulate cortex oxytocin involvement in modulation of sensory and affective dimensions of neuropathic pain induced by partial sciatic nerve ligation in rats.. **2022**, 113818 0
- 544 Sertoli cell transplantation attenuates microglial activation and inhibits TRPC6 expression in neuropathic pain induced by spinal cord injury.. **2022**, 251, 113807 0
- 543 Stephalagine, an aporphinic alkaloid with therapeutic effects in acute gout arthritis in mice.. **2022**, 115291 1
- 542 Data_Sheet_1.PDF. **2020**,
- 541 Image_1.PDF. **2020**,
- 540 Image_2.PDF. **2020**,
- 539 Table_1.DOCX. **2020**,
- 538 Table_10.DOCX. **2020**,
- 537 Table_2.XLSX. **2020**,

536 Table_3.XLSX. 2020,

535 Table_4.XLSX. 2020,

534 Table_5.XLSX. 2020,

533 Table_6.XLSX. 2020,

532 Table_7.XLSX. 2020,

531 Table_8.XLSX. 2020,

530 Table_9.XLSX. 2020,

529 Image_1.PDF. 2020,

528 Table_1.PDF. 2020,

527 Table_2.PDF. 2020,

526 Table_3.PDF. 2020,

525 Table_4.PDF. 2020,

524 Image_1.tif. 2020,

523 Image_2.tif. 2020,

522 Table_1.xlsx. 2020,

521 Table_2.xlsx. 2020,

520 Table_3.xlsx. 2020,

519 Data_Sheet_1.pdf. 2020,

518 Image_1.TIF. 2018,

517 Image_2.TIF. 2018,

516 Image_3.TIF. 2018,

515 Image_4.tif. 2018,

514 Image_5.tif. 2018,

513 Image_6.TIF. 2018,

512 Image_7.TIF. 2018,

511 Image_8.TIF. 2018,

510 DataSheet_1.pdf. 2020,

509 Image_1.TIF. 2018,

508 DataSheet_1.pdf. 2020,

507 Table_1.docx. 2019,

506 Image_1.JPEG. 2018,

505 Image_2.jpg. 2018,

504 Image_3.jpg. 2018,

503 Presentation_1.PDF. 2018,

502 Presentation_1.pdf. 2018,

501 Table_1.XLSX. 2018,

500 Table_2.XLSX. 2018,

499 Table_3.XLSX. 2018,

498 Image_1.JPEG. 2018,

497 Image_1.TIF. 2018,

496 Table_1.XLSX. 2018,

495 Table_1.DOCX. 2018,

494 Data_Sheet_1.docx. 2019,

493 Data_Sheet_2.xlsx. 2019,

492 Image_1.TIF. 2019,

491 Data_Sheet_1.PDF. 2019,

490 DataSheet_1.pdf. 2020,

489 Data_Sheet_1.pdf. 2019,

488 Image_1.tif. 2020,

487 Image_2.tif. 2020,

486 Table_1.xlsx. 2019,

485 Presentation_1.PDF. 2018,

484 Data_Sheet_1.ZIP. 2020,

483 Data_Sheet_2.ZIP. 2020,

482 Image_1.tif. **2020**,

481 Image_2.tif. **2020**,

480 Image_3.tif. **2020**,

479 Image_4.tif. **2020**,

478 Table_1.docx. **2020**,

477 Data_Sheet_1.PDF. **2019**,

476 Data_Sheet_1.PDF. **2019**,

475 Image_1.pdf. **2018**,

474 NGF monoclonal antibody DS002 alleviates chemotherapy-induced peripheral neuropathy in rats.. **2022**,

2

473 Transient astrocytic mGluR5 expression drives synaptic plasticity and subsequent chronic pain in mice.. **2022**, 219,

1

472 The transcription factor CCAAT/enhancer-binding protein β in spinal microglia contributes to pre-operative stress-induced prolongation of postsurgical pain.. **2022**, 17448069221099360

0

471 Environmental enrichment mitigates the long-lasting sequelae of perinatal fentanyl exposure in mice.. **2022**,

0

470 Upregulation of Hevin Contributes to Postoperative Pain Hypersensitivity by Inducing Neurexin1 β /Neuroigin1-Mediated Synaptic Targeting of GluA1-Containing Ampa Receptors in Rat Dorsal Horn.

469 Using Corneal Confocal Microscopy to Identify Therapeutic Agents for Diabetic Neuropathy.. **2022**, 11,

468 Centrally expressed Cav3.2 T-type calcium channel is critical for the initiation and maintenance of neuropathic pain.

467 Resolvin D1 Alleviates Mechanical Allodynia via ALX/FPR2 Receptor Targeted Nod-like Receptor Protein 3/Extracellular Signal-Related Kinase Signaling in a Neuropathic Pain Model.. **2022**,

0

466 Chronic pain recruits hypothalamic dynorphin/kappa opioid receptor signalling to promote wakefulness and vigilance.. **2022**,

0

465 A ketogenic diet prevents methylglyoxal-evoked nociception by scavenging methylglyoxal.. **2022**,

0

464	Chronic pain-induced neuronal plasticity in the bed nucleus of the stria terminalis causes maladaptive anxiety.. 2022 , 8, eabj5586	1
463	CGRP Administration Into the Cerebellum Evokes Light Aversion, Tactile Hypersensitivity, and Nociceptive Squint in Mice.. 2022 , 3, 861598	0
462	Selective Targeting of Serotonin 5-HT1a and 5-HT3 Receptors Attenuates Acute and Long-Term Hypersensitivity Associated With Neonatal Procedural Pain.. 2022 , 3, 872587	
461	Alleviative effect of microRNA-497 on diabetic neuropathic pain in rats in relation to decreased USP15.. 2022 , 1	
460	Structural analysis of TrkA mutations in patients with congenital insensitivity to pain reveals PLC β as an analgesic drug target.. 2022 , 15, eabm6046	0
459	Cannabinoid CB2 receptors are upregulated via bivalent histone modifications and control primary afferent input to the spinal cord in neuropathic pain.. 2022 , 101999	1
458	The sex-specific role of sensory neuron LKB1 on metabolic stress-induced mechanical hypersensitivity and mitochondrial respiration.. 2022 ,	0
457	Targeted ubiquitination of sensory neuron calcium channels reduces the development of neuropathic pain.. 2022 , 119, e2118129119	1
456	Precision spinal gene delivery-induced functional switch in nociceptive neurons reverses neuropathic pain.. 2022 ,	0
455	S-ketamine promotes autophagy and alleviates neuropathic pain by inhibiting PI3K/Akt/mTOR signaling pathway. 1	
454	Bladder Oxidative Stress and HMGB1 Release Contribute to PAR4-Mediated Bladder Pain in Mice. 2022 , 16,	1
453	IL-17 exacerbates experimental autoimmune prostatitis via CXCL1/CXCL2-mediated neutrophil infiltration.. 2022 , e14455	0
452	Downregulation of Fat Mass and Obesity-Related Protein in the Anterior Cingulate Cortex Participates in Anxiety- and Depression-Like Behaviors Induced by Neuropathic Pain. 2022 , 16,	1
451	Grpr expression defines a population of superficial dorsal horn vertical cells that have a role in both itch and pain.. 2022 ,	0
450	Acute inflammatory response via neutrophil activation protects against the development of chronic pain.. 2022 , 14, eabj9954	1
449	Cytotoxicity and inflammation induced by <i>Philodryas patagoniensis</i> venom.. 2022 , 109356	1
448	Systemic, Intrathecal, and Intracerebroventricular Antihyperalgesic Effects of the Calcium Channel Blocker CTK 01512-2 Toxin in Persistent Pain Models.. 2022 ,	0
447	Unbiased proteomic analysis detects painful systemic inflammatory profile in the serum of nerve injured mice.. 2022 ,	2

- 446 Measuring Mouse Somatosensory Reflexive Behaviors with High-Speed Videography, Statistical Modeling, and Machine Learning. **2022**, 441-456
- 445 Use of Intraspinally Delivered Chemogenetic Receptor, PSAM-GlyR, to Probe the Behavioral Role of Spinal Dorsal Horn Neurons. **2022**, 417-439
- 444 Activation of locus coeruleus-spinal cord α 1-adrenergic neurons alleviates neuropathic pain in mice via reducing neuroinflammation from astrocytes and microglia in spinal dorsal horn. **2022**, 19, 2
- 443 Sex-Specific Transcriptomic Signatures in Brain Regions Critical for Neuropathic Pain-Induced Depression. **2022**, 15, 0
- 442 Altered prefrontal-striatal theta-band oscillatory dynamics underlie working memory deficits in neuropathic pain rats. 0
- 441 SUMOylation of Kir7.1 participates in neuropathic pain through regulating its membrane expression in spinal cord neurons.
- 440 Behavioral and inflammatory sex differences revealed by celecoxib nanotherapeutic treatment of peripheral neuroinflammation. **2022**, 12, 0
- 439 Longitudinal RNA Sequencing of Skin and DRG Neurons in Mice with Paclitaxel-Induced Peripheral Neuropathy. **2022**, 7, 72
- 438 Targeted muscle reinnervation prevents and reverses rat pain behaviors following nerve transection. **2022**, Publish Ahead of Print, 0
- 437 Inhibition of DAGL α as a therapeutic target for pain in sickle cell disease. 0
- 436 Dysregulation of Immune Response Mediators and Pain-Related Ion Channels Is Associated with Pain-like Behavior in the GLA KO Mouse Model of Fabry Disease. **2022**, 11, 1730 1
- 435 Microglia-mediated degradation of perineuronal nets promotes pain. 5
- 434 Low-Intensity 10 kHz Spinal Cord Stimulation Reduces Behavioral and Neural Hypersensitivity in a Rat Model of Painful Diabetic Neuropathy. Volume 15, 1503-1513 1
- 433 Medullary kappa-opioid receptor neurons inhibit pain and itch through a descending circuit. 0
- 432 Low-Molecular-Weight Perorally Active Nerve Growth Factor Mimetic Reduces Manifestations of Diabetic Neuropathy in Wistar Rats. **2022**, 173, 37-40
- 431 Slack Potassium Channels Modulate TRPA1-Mediated Nociception in Sensory Neurons. **2022**, 11, 1693 0
- 430 Mammalian Sterile 20-Like Kinase 1 Mediates Neuropathic Pain Associated with Its Effects on Regulating Mitophagy in Schwann Cells. **2022**, 2022, 1-17
- 429 Olfactory exposure to late-pregnant and lactating mice causes stress-induced analgesia in male mice. **2022**, 8,

- 428 Electroacupuncture Ameliorates Mechanical Allodynia of a Rat Model of CRPS-I via Suppressing NLRP3 Inflammasome Activation in Spinal Cord Dorsal Horn Neurons. 16,
- 427 Local effects of natural alkylamides from *Acmella oleracea* and synthetic isobutylalkyl amide on neuropathic and postoperative pain models in mice. **2022**, 105224 ○
- 426 Nalbuphine alleviates inflammation by down-regulating NF- κ B in an acute inflammatory visceral pain rat model. **2022**, 23,
- 425 Activation of Double-Stranded RNA-Activated Protein Kinase in the Dorsal Root Ganglia and Spinal Dorsal Horn Regulates Neuropathic Pain Following Peripheral Nerve Injury in Rats.
- 424 RSK1 promotes mammalian axon regeneration by inducing the synthesis of regeneration-related proteins. **2022**, 20, e3001653 ○
- 423 Objective and Quantitative Evaluation of Spontaneous Pain-Like Behaviors Using Dynamic Weight-Bearing System in Mouse Models of Postsurgical Pain. Volume 15, 1601-1612 ○
- 422 Differential effects of remifentanyl and sufentanyl anesthesia on post-operative pain and cognitive functions. **2022**, 108, 108888 ○
- 421 The Development of Mechanical Allodynia in Diabetic Rats Revealed by Single-Cell RNA-Seq. 15, ○
- 420 Developmentally determined intersectional genetic strategies to dissect adult somatosensory circuit function.
- 419 Long-lasting reflexive and nonreflexive pain responses in two mouse models of fibromyalgia-like condition. **2022**, 12, 2
- 418 Neuroinflammation in the anterior cingulate cortex: the potential supraspinal mechanism underlying the mirror-image pain following motor fiber injury. **2022**, 19, ○
- 417 Titrating the Translational Relevance of a Low-Level Repetitive Head Impact Model. 13,
- 416 Activation of Dopamine D2 Receptors Alleviates Neuronal Hyperexcitability in the Lateral Entorhinal Cortex via Inhibition of HCN Current in a Rat Model of Chronic Inflammatory Pain.
- 415 Spinal CCK1 receptors contribute to somatic pain hypersensitivity induced by malocclusion via a reciprocal neuron-glia signaling cascade. **2022**, ○
- 414 Sex-Related Pain Behavioral Differences following Unilateral NGF Injections in a Rat Model of Low Back Pain. **2022**, 11, 924 ○
- 413 Spinal TRPA1 Contributes to the Mechanical Hypersensitivity Effect Induced by Netrin-1. **2022**, 23, 6629 ○
- 412 Regulatory T-cells and IL-5 mediate pain outcomes in a preclinical model chronic muscle pain.. 174480692211106
- 411 UNC5B Overexpression Alleviates Peripheral Neuropathic Pain by Stimulating Netrin-1-Dependent Autophagic Flux in Schwann Cells. 1

- 410 Alterations of monoamine neurotransmitters, HPA-axis hormones, and inflammation cytokines in reserpine-induced hyperalgesia and depression comorbidity rat model. **2022**, 22, 1
- 409 [6]-Shogaol Attenuates Oxaliplatin-Induced Allodynia through Serotonergic Receptors and GABA in the Spinal Cord in Mice. **2022**, 15, 726 0
- 408 Complement Receptor 3 Pathway and NMDA Receptor 2B Subunit Involve Neuropathic Pain Associated with Spinal Cord Injury. Volume 15, 1813-1823
- 407 Stimulation of CGRP-expressing Neurons in the Medial Cerebellar Nucleus Induces Light and Touch Sensitivity in Mice. **2022**, 100098 0
- 406 Migraine and peripheral pain models show differential alterations in neuronal complexity.
- 405 A Mouse Model of Cancer Induced Bone Pain: From Pain to Movement. 16, 0
- 404 Low-dose interleukin-2 reverses chronic migraine-related sensitizations through peripheral interleukin-10 and transforming growth factor beta-1 signaling. **2022**, 12, 100096
- 403 Potent CaV3.2 channel inhibitors exert analgesic effects in acute and chronic pain models. **2022**, 153, 113310 0
- 402 Microglia and macrophages contribute to the development and maintenance of sciatica in lumbar disc herniation. **2022**, Publish Ahead of Print, 1
- 401 Intrathecal dexmedetomidine attenuates mechanical allodynia through the downregulation of brain-derived neurotrophic factor in a mild traumatic brain injury rat.
- 400 ACT001 Inhibits TLR4 Signaling by Targeting Co-Receptor MD2 and Attenuates Neuropathic Pain. 13, 0
- 399 Disruption of lactate metabolism in the peripheral nervous system leads to motor-selective deficits. 0
- 398 The Interaction between Carbon Monoxide and Hydrogen Sulfide during Chronic Joint Pain in Young Female Mice. **2022**, 11, 1271 0
- 397 Liquid Biopsy-Based Biomarkers of Inflammatory Nociception Identified in Male Rats. 13,
- 396 Shaoyao Gancuo Decoction Ameliorates Paclitaxel-Induced Peripheral Neuropathy via Suppressing TRPV1 and TLR4 Signaling Expression in Rats. Volume 16, 2067-2081 1
- 395 Loss of Stathmin-2, a hallmark of TDP-43-associated ALS, causes motor neuropathy. **2022**, 39, 111001 1
- 394 Electroacupuncture Attenuates Post-Inflammatory IBS-Associated Visceral and Somatic Hypersensitivity and Correlates With the Regulatory Mechanism of Epac1/Biezo2 Axis. 13, 0
- 393 Metabotropic Glutamate Receptor 5 in the Dysgranular Zone of Primary Somatosensory Cortex Mediates Neuropathic Pain in Rats. **2022**, 10, 1633

- 392 Ultrasound-Triggered on Demand Lidocaine Release Relieves Postoperative Pain. 10,
- 391 Sciatic nerve block downregulates the BDNF pathway to alleviate the neonatal incision-induced exaggeration of incisional pain via decreasing microglial activation. **2022**,
- 390 Cryptochrome Stabilization Ameliorates Chronic Pain.
- 389 Oral Cancer Cells Release Vesicles that Cause Pain. 2200073 1
- 388 Age-associated changes in knee osteoarthritis, pain-related behaviors, and dorsal root ganglia immunophenotyping of male and female mice.
- 387 Early life stress exposure increases susceptibility to high fat/high sucrose diet in female mice.
- 386 Antinociceptive and Analgesic Effects of (2R,6R)-hydroxynorketamine. JPET-AR-2022-001278 1
- 385 Effect of chronic lithium on mechanical sensitivity and trabecular bone loss induced by type-1 diabetes mellitus in mice.
- 384 Mechanisms of Neuropathic Pain and Pain-Relieving Effects of Exercise Therapy in a Rat Neuropathic Pain Model. Volume 15, 1925-1938
- 383 Neuropathic Pain Induces Interleukin-1 β -Sensitive Bimodal Glycinergic Activity in the Central Amygdala. **2022**, 23, 7356
- 382 Controlled activation of cortical astrocytes modulates neuropathic pain-like behaviour. **2022**, 13, 1
- 381 Selective block of sensory neuronal T-type/Cav3.2 activity mitigates neuropathic pain behavior in a rat model of osteoarthritis pain. **2022**, 24,
- 380 Hydrogen Sulfide Increases the Analgesic Effects of μ - and κ Opioid Receptors during Neuropathic Pain: Pathways Implicated. **2022**, 11, 1321 0
- 379 Enhancement of Mitochondrial Function by the Neurogenic Molecule NSI-189 Accompanies Reversal of Peripheral Neuropathy and Memory Impairment in a Rat Model of Type 2 Diabetes. **2022**, 2022, 1-13
- 378 Anti-Inflammatory and Analgesic Properties of the Cannabis Terpene Myrcene in Rat Adjuvant Monoarthritis. **2022**, 23, 7891 0
- 377 Endogenous κ Opioid μ Neuropeptide Y Y1 receptor synergy silences chronic postoperative pain.
- 376 Time course and localization of nerve growth factor expression and sensory nerve growth during progression of knee osteoarthritis in rats. **2022**, 0
- 375 IL-10 alleviates radicular pain by inhibiting TNF- α /p65 dependent Nav1.7 up-regulation in DRG neurons of rats. **2022**, 1791, 147997 0

- 374 SR 57227A, a serotonin type-3 receptor agonist, as a candidate analgesic agent targeting nociplastic pain. **2022**, 622, 143-148
- 373 Next generation behavioral sequencing for advancing pain quantification. **2022**, 76, 102598
- 372 Upregulation of Hevin contributes to postoperative pain hypersensitivity by inducing neurexin1 α /neuroligin1-mediated synaptic targeting of GluA1-containing AMPA receptors in rat dorsal horn. **2022**, 1792, 148004
- 371 Potentiation of morphine antinociception and inhibition of diabetic neuropathic pain by the multi-chemokine receptor antagonist peptide RAP-103. **2022**, 306, 120788 ○
- 370 Inhibition of the RNA Regulator HuR by SRI-42127 Attenuates Neuropathic Pain After Nerve Injury Through Suppression of Neuroinflammatory Responses. ○
- 369 Light-Induced Activation of a Specific Type-5 Metabotropic Glutamate Receptor Antagonist in the Ventrobasal Thalamus Causes Analgesia in a Mouse Model of Breakthrough Cancer Pain. **2022**, 23, 8018
- 368 Low-Intensity Focused Ultrasound Alleviates Chronic Neuropathic Pain-Induced Allodynia by Inhibiting Neuroplasticity in the Anterior Cingulate Cortex. **2022**, 2022, 1-11
- 367 Mouse Models of Surgical and Neuropathic Pain Produce Distinct Functional Alterations to Prodynorphin-Expressing Neurons in the Prelimbic Cortex.
- 366 Individual differences in oxycodone addiction-like behaviors in a large cohort of heterogeneous stock (HS) rats.
- 365 Sex-specific effects of neuropathic pain on long-term pain behavior and mortality in mice. **2022**, Publish Ahead of Print, ○
- 364 LncRNA-84277 is involved in chronic pain-related depressive behaviors through miR-128-3p/SIRT1 axis in central amygdala. 15,
- 363 Analgesic effect of nobiletin against neuropathic pain induced by the chronic constriction injury of the sciatic nerve in mice.
- 362 Activation of δ -containing GABAA receptors induces antinociception under physiological and pathological conditions. **2022**, Publish Ahead of Print,
- 361 No evidence for cognitive impairment in an experimental rat model of knee osteoarthritis and associated chronic pain.
- 360 Long-lasting visceral hypersensitivity in a model of DSS -induced colitis in rats.
- 359 Pharmacological and non-pharmacological therapeutic interventions for the treatment of spinal cord injury-induced pain. 3, ○
- 358 SEPT9 Upregulation in Satellite Glial Cells Associated with Diabetic Polyneuropathy in a Type 2 Diabetes-like Rat Model. **2022**, 23, 9372 ○
- 357 Supraspinal melatonin MT 2 receptor agonism alleviates pain via a neural circuit that recruits mu opioid receptors. 1

- 356 Activation of GDNF-ERK-Runx1 signaling contributes to P2X3R gene transcription and bone cancer pain. **2022**, 104936 1
- 355 Insight into the possible mechanism(s) involved in the antinociceptive and antineuropathic activity of *Descurainia sophia* L. Webb ex Prantl essential oil. **2022**, 115638
- 354 Wnt3a/YTHDF1 Regulated Oxaliplatin-Induced Neuropathic Pain Via TNF- α /IL-18 Expression in the Spinal Cord. 0
- 353 Antioxidant effect of grape seed extract corrects experimental autoimmune encephalomyelitis behavioral dysfunctions, demyelination, and glial activation. 13,
- 352 Sensory neuron dysfunction in orthotopic mouse models of colon cancer. **2022**, 19, 0
- 351 Resting-State Functional Connectivity Analyses: Brain Functional Reorganization in a Rat Model of Postherpetic Neuralgia. **2022**, 12, 1029
- 350 Analgesic Dorsal Root Ganglion Field Stimulation Blocks Both Afferent and Efferent Spontaneous Activity in Sensory Neurons of Rats with Monosodium Iodoacetate-induced Osteoarthritis. **2022**, 0
- 349 Transgenic Mice for the Translational Study of Neuropathic Pain and Dystonia. **2022**, 23, 8580
- 348 Intrathecal Co-administration of Morphine Facilitated the Anti-nociceptive of Bupivacaine in a Rat Model of Acute Postoperative Pain. **2022**,
- 347 Etv4 regulates nociception by controlling peptidergic sensory neuron development and peripheral tissue innervation. **2022**, 149,
- 346 Unbiased analysis of the dorsal root ganglion after peripheral nerve injury: no neuronal loss, no gliosis, but satellite glial cell plasticity. **2022**, Publish Ahead of Print, 0
- 345 A model of painful vaso-occlusive crisis in mice with sickle cell disease.
- 344 The RSK-RPS6 Axis Controls the Preconditioning Effect and Induces Spinal Cord Regeneration.
- 343 Analgesic Characteristics of NanoBEO Released by an Airless Dispenser for the Control of Agitation in Severe Dementia. **2022**, 27, 4987 1
- 342 Somatotopy of mouse spinothalamic innervation and the localisation of a noxious stimulus requires DCC expression by Phox2a neurons. JN-RM-1164-22
- 341 Methacrylic acid-based biomaterials promote peripheral innervation in the subcutaneous space of mice. **2022**, 121764
- 340 Development of a spontaneous pain indicator based on brain cellular calcium using deep learning. 1
- 339 Ligustrazine mitigates chronic venous disease-induced pain hyperalgesia through desensitization of inflammation-associated TRPA1 activity in DRG. **2022**, 115667

- 338 The adverse effects of repeated, intravenous morphine on recovery following spinal cord injury in young, male rats are blocked by a kappa opioid receptor antagonist.
- 337 MAO-B inhibitor, KDS2010, alleviates spinal nerve ligation-induced neuropathic pain in rats through competitively blocking the BDNF/TrkB/NR2B signaling. **2022**, 1
- 336 Inhibition of itch by neurokinin 1 receptor (Tacr1) -expressing ON cells in the rostral ventromedial medulla in mice. 11, 0
- 335 Effects of Neural Mobilization on Sensory Dysfunction and Peripheral Nerve Degeneration in Rats with Painful Diabetic Neuropathy.
- 334 Dexmedetomidine Alleviates Neuropathic Pain via the TRPC6-p38 MAPK Pathway in the Dorsal Root Ganglia of Rats. Volume 15, 2437-2448
- 333 Mechanical and Heat Hyperalgesia upon Withdrawal from Chronic Intermittent Ethanol Vapor depends on Sex, Exposure Duration and Blood Alcohol Concentration in Mice.
- 332 FGF 10 Inhibited Spinal Microglial Activation in Neuropathic Pain via PPAR- γ -TNF- α Signaling. **2022**, 500, 52-62
- 331 Cannabidiol attenuates hypersensitivity and oxidative stress after traumatic spinal cord injury in rats. **2022**, 788, 136855 0
- 330 Down-regulation of NR2B receptors contributes to the analgesic and antianxiety effects of enriched environment mediated by endocannabinoid system in the inflammatory pain mice. **2022**, 435, 114062
- 329 Effects of treatment with a carbon monoxide donor and an activator of heme oxygenase 1 on the nociceptive, apoptotic and/or oxidative alterations induced by persistent inflammatory pain in the central nervous system of mice. **2022**, 188, 169-178 0
- 328 Role of spinal RIP3 in inflammatory pain and electroacupuncture-mediated analgesic effect in mice. **2022**, 306, 120839 1
- 327 Peroxisome proliferator-activated receptor gamma agonist ELB00824 suppresses oxaliplatin-induced pain, neuronal hypersensitivity, and oxidative stress. **2022**, 218, 109233
- 326 Dimethyl Fumarate Ameliorates Paclitaxel-Induced Neuropathic Pain in Rats. **2022**, 0
- 325 SIRT3-Mediated CypD-K166 Deacetylation Alleviates Neuropathic Pain by Improving Mitochondrial Dysfunction and Inhibiting Oxidative Stress. **2022**, 2022, 1-17 1
- 324 Role of 5-HT_{2A} Receptor in Modulating Glutamatergic Activity in the Ventrolateral Orbital Cortex: Implication in Trigeminal Neuralgia. **2022**, 502, 107-116 0
- 323 Targeting 5-HT_{2A} receptors and Kv7 channels in PFC to attenuate chronic neuropathic pain in rats using a spared nerve injury model. **2022**, 789, 136864 0
- 322 The effects of low intensity focused ultrasound on neuronal activity in pain processing regions in a rodent model of common peroneal nerve injury. **2022**, 789, 136882 0
- 321 Glucocorticoid regulation of lactate release from spinal astrocytes contributes to the induction of spinal LTP of C-fiber-evoked field potentials and the development of mechanical allodynia. **2022**, 219, 109253 0

- 320 A Possible Mechanism for Development of Working Memory Impairment in Male Mice Subjected to Inflammatory Pain. **2022**, 503, 17-27 ○
- 319 Persistent sensory changes and sex differences in transgenic mice conditionally expressing HIV-1 Tat regulatory protein. **2022**, 358, 114226 ○
- 318 Discovery of pimozide derivatives as novel T-type calcium channel inhibitors with little binding affinity to dopamine D2 receptors for treatment of somatic and visceral pain. **2022**, 243, 114716 ○
- 317 Fluoxetine reverses hyperactivity of anterior cingulate cortex and attenuates chronic stress-induced hyperalgesia. **2022**, 220, 109259 ○
- 316 Benzo[b]thiophene-2-carboxamides as novel opioid receptor agonists with potent analgesic effect and reduced constipation. **2022**, 243, 114728 ○
- 315 Neuroprotective Effects of Virgin Coconut Oil Solubilized Curcumin in Diabetic Peripheral Neuropathy. **2022**, 17, 36-42 ○
- 314 Acute and Chronic Pain Preclinical Models to Study the Analgesic Properties of Melatonergic Compounds. **2022**, 453-461 ○
- 313 Lidocaine reduces pain behaviors by inhibiting the expression of Nav1.7 and Nav1.8 and diminishing sympathetic sprouting in SNI rats. **2022**, 18, 174480692211249 ○
- 312 Changes in the expression of endocannabinoid system components in an experimental model of chemotherapy-induced peripheral neuropathic pain: Evaluation of sex-related differences. **2023**, 359, 114232 ○
- 311 Inflammation-induced mitochondrial and metabolic disturbances in sensory neurons control the switch from acute to chronic pain. ○
- 310 The role of somatosensory innervation of adipose tissues. **2022**, 609, 569-574 2
- 309 Comparison of Pain-Like behaviors in two surgical incision animal models in C57BL/6J mice. **2022**, 12, 100103 ○
- 308 Spinal neurovascular coupling is preserved despite time dependent alterations of spinal cord blood flow responses in a rat model of chronic back pain: implications for functional spinal cord imaging.. **2022**, Publish Ahead of Print, ○
- 307 Chronic pain causes Tau-mediated hippocampal pathology and memory deficits. 1
- 306 Synchronous firing of dorsal horn neurons at the origin of dorsal root reflexes in naïve and paw-inflamed mice. 16, 1
- 305 Involvement of TLR2, TLR4, NLRP3, and IL-17 in pain induced by a novel Sprague-Dawley rat model of experimental autoimmune encephalomyelitis. 3, ○
- 304 Meteorin alleviates Paclitaxel-induced peripheral neuropathy in mice. ○
- 303 Dysregulation of dopamine neurotransmission in the nucleus accumbens in immobilization-induced hypersensitivity. 13, ○

302	DICAM in the Extracellular Vesicles from Astrocytes Attenuates Microglia Activation and Neuroinflammation. 2022 , 11, 2977	0
301	Functional Gait Assessment Using Manual, Semi-Automated and Deep Learning Approaches Following Standardized Models of Peripheral Nerve Injury in Mice. 2022 , 12, 1355	0
300	Pharmacological Interaction of Quercetin Derivatives of <i>Tilia americana</i> and Clinical Drugs in Experimental Fibromyalgia. 2022 , 12, 916	0
299	Identification of regulatory elements in primary sensory neurons involved in neuropathic pain.	0
298	Human PMSCs-derived small extracellular vesicles alleviate neuropathic pain through miR-26a-5p/Wnt5a in SNI mice model. 2022 , 19,	1
297	Anti-inflammatory and anti-allodynic activities in mice of the essential oil and desmethyloencecalin from <i>Zinnia grandiflora</i> Nuttall. 1-12	0
296	Lysine demethylation KDM5B downregulates SIRT3 -mediated mitochondrial glucose and lipid metabolism in diabetic neuropathy.	0
295	Macrophage Infiltration Initiates RIP3/MLKL-Dependent Necroptosis in Paclitaxel-Induced Neuropathic Pain. 2022 , 2022, 1-10	0
294	Targeting FoxO transcription factors with HDAC inhibitors for the treatment of osteoarthritis. <i>annrheumdis-2021-22126</i>	0
293	Sex-related differences in oxaliplatin-induced changes in the expression of transient receptor potential channels and their contribution to cold hypersensitivity. 2022 , 788, 136863	0
292	Simultaneous monitoring of mouse grip strength, force profile, and cumulative force profile distinguishes muscle physiology following surgical, pharmacologic and diet interventions. 2022 , 12,	1
291	Activation of VIP interneurons in the prefrontal cortex ameliorates neuropathic pain aversiveness. 2022 , 40, 111333	0
290	Activation of GPR183 by 7 β -25-Dihydroxycholesterol Induces Behavioral Hypersensitivity through Mitogen-Activated Protein Kinase and Nuclear Factor- κ B. 2022 , 383, 172-181	0
289	A parabrachial-to-amygdala circuit that determines hemispheric lateralization of somatosensory processing.	0
288	HDAC6 inhibition reverses cisplatin-induced mechanical hypersensitivity via tonic delta opioid receptor signaling. JN-RM-1182-22	0
287	Angiotensin type 2 receptor antagonism as a new target to manage gout.	1
286	Monoamine control of descending pain modulation after mild traumatic brain injury. 2022 , 12,	0
285	Regulation of neuropathic pain by microglial Orai1 channels.	1

284	Small synthetic hyaluronan disaccharide BIS014 mitigates neuropathic pain in mice. 2022,	0
283	Effect of basic fibroblast growth factor injection on peripheral nerve injury at the watershed zone in a rat model. 2022,	0
282	Uncompetitive, adduct-forming SARM1 inhibitors are neuroprotective in preclinical models of nerve injury and disease. 2022,	1
281	Keratinocyte PIEZO1 modulates cutaneous mechanosensation. 11,	1
280	A parabrachial-to-amygdala circuit that determines hemispheric lateralization of somatosensory processing. 2022,	1
279	Hydrogen-Rich Water as a Novel Therapeutic Strategy for the Affective Disorders Linked with Chronic Neuropathic Pain in Mice. 2022, 11, 1826	0
278	High-fat diet causes mechanical allodynia in the absence of injury or diabetic pathology. 2022, 12,	0
277	Pattern of Self-Administered Vapor Fentanyl Exposure Determines Long-term Behavior Consequences, in Mice with or without Neuropathic Pain.	0
276	Structure-based discovery of nonopioid analgesics acting through the β_2 A -adrenergic receptor. 2022, 377,	3
275	Pharmacogenetic inhibition of TrkB signaling in adult mice attenuates mechanical hypersensitivity and improves locomotor function after spinal cord injury. 16,	0
274	Combination of paeoniflorin and liquiritin alleviates neuropathic pain by lipid metabolism and calcium signaling coordination. 13,	0
273	Interactions Among Non-Coding RNAs and mRNAs in the Trigeminal Ganglion Associated with Neuropathic Pain. Volume 15, 2967-2988	1
272	What Did We Learn About Fracture Pain from Animal Models?. Volume 15, 2845-2856	0
271	Hypothalamic A11 Nuclei Regulate the Circadian Rhythm of Spinal Mechanonociception through Dopamine Receptors and Clock Gene Expression. 2022, 12, 1411	0
270	Cilostazol Alleviates NLRP3 Inflammasome-Induced Allodynia/Hyperalgesia in Murine Cerebral Cortex Following Transient Ischemia: Focus on TRPA1/Glutamate and Akt/Dopamine/BDNF/Nrf2 Trajectories.	0
269	Transcriptional profiling of TGF- β superfamily members in lumbar DRGs of rats following sciatic nerve axotomy and activin C inhibits neuropathic pain. 2022, 22,	0
268	Medial prefrontal cortex nitric oxide modulates neuropathic pain behavior through mu opioid receptors in rats. 2022, 35, 413-422	0
267	Ningmitai capsules have anti-inflammatory and pain-relieving effects in the chronic prostatitis/chronic pelvic pain syndrome mouse model through systemic immunity. 13,	0

- 266 Piezo2 voltage-block regulates mechanical pain sensitivity. ○
- 265 (2R,6R)-hydroxynorketamine (HNK) reverses mechanical hypersensitivity in a model of localized inflammatory pain. **2022**, 109276 ○
- 264 Analgesic Effectiveness and Dorsal Root Ganglia Protein Modulation of a Peripheral Adenosine Monophosphate Kinase Alpha Activator (O304) Following Lumbar Disk Puncture in the Mouse. Publish Ahead of Print, ○
- 263 Metformin promotes angiogenesis and functional recovery in aged mice after spinal cord injury by adenosine monophosphate-activated protein kinase/ endothelial nitric oxide synthase pathway. **2022**, 0 ○
- 262 Sex-distinct microglial activation and myeloid cell infiltration in the spinal cord after painful peripheral injury. **2022**, 12, 100106 ○
- 261 NLRP3-Mediated Piezo1 Upregulation in ACC Inhibitory Parvalbumin-Expressing Interneurons Is Involved in Pain Processing after Peripheral Nerve Injury. **2022**, 23, 13035 ○
- 260 TRPM8 contributes to sex dimorphism by promoting recovery of normal sensitivity in a mouse model of chronic migraine. **2022**, 13, ○
- 259 Upregulation of Nav1.6 Mediated by the p38 MAPK Pathway in the Dorsal Root Ganglia Contributes to Cancer-Induced Bone Pain in Rats. **2022**, 11, 3375 ○
- 258 TRPV1 participates in neuropathic pain after spinal cord injury by mediating the proliferation and activation of CX3CL1-positive glial cells in the spinal dorsal horn. ○
- 257 Neuroimaging uncovers neuronal and metabolic changes in pain modulatory brain areas in a rat model of chemotherapy-induced neuropathy [MEMRI and ex vivo spectroscopy studies. **2022**, ○
- 256 Cerium Oxide Nanoparticles Alleviate Neuropathic Pain by Modulating Macrophage Polarization in a Rat SCI Model. Volume 15, 3369-3380 ○
- 255 PATHOPHYSIOLOGICAL CHARACTERIZATION OF THE TOWNES MOUSE MODEL FOR SICKLE CELL DISEASE. **2022**, ○
- 254 Sensory Neuron-Specific Deletion of Tropomyosin Receptor Kinase A (TrkA) in Mice Abolishes Osteoarthritis (OA) Pain via NGF/TrkA Intervention of Peripheral Sensitization. **2022**, 23, 12076 ○
- 253 LncRNA XR_351665 contributes to chronic pain-induced depression by upregulating DNMT1 via sponging miR-152-3p. **2022**, ○
- 252 Unveiling Targets for Treating Postoperative Pain: The Role of the TNF- α /p38 MAPK/NF- κ B/Nav1.8 and Nav1.9 Pathways in the Mouse Model of Incisional Pain. **2022**, 23, 11630 ○
- 251 Spinal cord retinoic acid receptor signaling gates mechanical hypersensitivity in neuropathic pain. **2022**, ○
- 250 A novel implantable device for sensory and affective assessment of orofacial pain in rats. 9, ○
- 249 Green light exposure elicits anti-inflammation, endogenous opioid release and dampens synaptic potentiation to relieve post-surgical pain. **2022**, ○

- 248 Role of leukotriene B4 (LTB4)-LTB4 receptor 1 signaling in post-incisional nociceptive sensitization and local inflammation in mice. **2022**, 17, e0276135 ○
- 247 Synergistic action between a synthetic cannabinoid compound and tramadol in neuropathic pain rats. **2022**, 72, 509-527 ○
- 246 The natural product Argentatin C attenuates postoperative pain via inhibition of voltage-gated sodium and T-type voltage-gated calcium channels. ○
- 245 Synaptic plasticity in two cell types of central amygdala for regulation of emotion and pain. 16, ○
- 244 A randomized, double-blinded, placebo-controlled, crossover study of the HCN channel blocker ivabradine in a capsaicin-induced pain model in healthy volunteers. **2022**, 12, ○
- 243 Long-Term Blockade of Nociceptive Nav1.7 Channels Is Analgesic in Rat Models of Knee Arthritis. **2022**, 12, 1571 ○
- 242 Identification of sensory dysfunction and nervous structure changes in Fam134b knockout mice. 1-8 ○
- 241 Preoperative Acute Sleep Deprivation Causes Postoperative Pain Hypersensitivity and Abnormal Cerebral Function. ○
- 240 Oral Administration of Glutathione Trisulfide Increases Reactive Sulfur Levels in Dorsal Root Ganglion and Ameliorates Paclitaxel-Induced Peripheral Neuropathy in Mice. **2022**, 11, 2122 ○
- 239 Schwann cells promote sensory neuron excitability during development. ○
- 238 HDAC2 in primary sensory neurons constitutively restrains chronic pain by repressing β -casein expression and associated NMDA receptor activity. JN-RM-0735-22 ○
- 237 The RNA-binding protein HuR is integral to the function of nociceptors in mice and humans. JN-RM-1630-22 ○
- 236 Establishment of a bipedal rat model of lumbar facet joint osteoarthritis using intraarticular injection of urinary plasminogen activator. **2022**, 17, ○
- 235 Meteorin alleviates Paclitaxel-induced peripheral neuropathic pain in mice. **2022**, ○
- 234 Lipid Nanoparticle Delivery of Chemically Modified NGF R100W mRNA Alleviates Peripheral Neuropathy. 2202127 ○
- 233 Stabilizer-induced different in vivo behaviors for intramuscularly long-acting celecoxib nanocrystals. **2022**, 628, 122298 ○
- 232 Nociceptor neurons affect cancer immunosurveillance. ○
- 231 An NTS-CeA projection modulates depression-like behaviors in a mouse model of chronic pain. **2022**, 174, 105893 ○

- 230 Dexmedetomidine alleviates pain in MPTP-treated mice by activating the AMPK/mTOR/NF- κ B pathways in astrocytes. **2022**, 791, 136933 0
- 229 Employment of PEGylated ultra-deformable transferosomes for transdermal delivery of tapentadol with boosted bioavailability and analgesic activity in post-surgical pain **2022**, 628, 122274 3
- 228 Continuous release of mefloquine featured in electrospun fiber membranes alleviates epidural fibrosis and aids in sensory neurological function after lumbar laminectomy. **2022**, 17, 100469 0
- 227 Protocols for the social transfer of pain and analgesia in mice. **2022**, 3, 101756 0
- 226 Intrathecal minocycline does not block the adverse effects of repeated, intravenous morphine administration on recovery of function after SCI. **2023**, 359, 114255 0
- 225 Establishment and validation of a rodent model of endometriosis to evaluate the effect of new therapeutic strategies. **2022**, 8, 036-042 0
- 224 Methyl ferulic acid alleviates neuropathic pain by inhibiting Nox4-induced ferroptosis in dorsal root ganglia neurons in rats. 0
- 223 C-low threshold mechanoreceptor activation becomes sufficient to trigger affective pain in spinal cord-injured mice in association with increased respiratory rates. 0
- 222 Cancer cells co-opt nociceptive nerves to thrive in nutrient-poor environments and upon nutrient-starvation therapies. **2022**, 0
- 221 Brief opioid exposure paradoxically augments primary afferent input to spinal excitatory neurons via μ -opioid dependent presynaptic NMDA receptors. JN-RM-1704-22 1
- 220 Istradefylline protects from cisplatin-induced nephrotoxicity and peripheral neuropathy while preserving cisplatin antitumor effects. **2022**, 132, 0
- 219 Neuropathic injury drives a generalized negative affective state in mice. 0
- 218 Autophagy Dysfunction in a Diabetic Peripheral Neuropathy Model **Publish Ahead of Print**, 0
- 217 Fluorescence visualization of the neuropathic pain triad in trigeminal neuralgia. 0
- 216 Inducible co-stimulatory molecule (ICOS) alleviates Paclitaxel induced peripheral neuropathy via an IL-10-mediated mechanism in female mice. 0
- 215 The contribution of endocytosis to sensitization of nociceptors and synaptic transmission in nociceptive circuits. **2022**, Publish Ahead of Print, 0
- 214 Analgesic effects and arthritic changes following intra-articular injection of diclofenac etalhyaluronate in a rat knee osteoarthritis model. **2022**, 23, 0
- 213 Spinal dopaminergic D1 and D5 receptors contribute to reserpine-induced fibromyalgia-like pain in rats. **2022**, 148167 0

- 212 Bone marrow mesenchymal stem cell-derived exosomes shuttling miR-150-5p alleviates mechanical allodynia in rats by targeting NOTCH2 in microglia. **2022**, 28, ○
- 211 The circ_0002538/miR-138-5p/plasmolipin axis regulates Schwann cell migration and myelination in diabetic peripheral neuropathy. **2022**, 0 ○
- 210 A High-Fat High-Sugar Diet and Adversity Early in Life Modulate Pain Outcomes at the Behavioural and Molecular Level in Adolescent Rats: The Role of Sex. **2022**, ○
- 209 Quercetin Attenuates Nitroglycerin-Induced Migraine Headaches by Inhibiting Oxidative Stress and Inflammatory Mediators. **2022**, 14, 4871 ○
- 208 BHF177 Suppresses Diabetic Neuropathic Pain by Blocking PKC/CaMKII/ERK1/2/CREB Signaling Pathway through Activating GABAB Receptor. **2022**, 2022, 1-10 ○
- 207 C781, a β -arrestin biased antagonist at protease-activated receptor-2 (PAR2), displays in vivo efficacy against protease-induced pain in mice. **2022**, ○
- 206 TRPV4 is not the molecular sensor for bacterial lipopolysaccharides-induced calcium signaling. **2023**, 383, 104651 ○
- 205 Huc-MSCs-derived exosomes attenuate neuropathic pain by inhibiting activation of the TLR2/MyD88/NF- κ B signaling pathway in the spinal microglia by targeting Rsad2. **2023**, 114, 109505 ○
- 204 A hydrolysate of poly-trans-[(2-carboxyethyl)germasesquioxane] (Ge-132) suppresses Cav3.2-dependent pain by sequestering exogenous and endogenous sulfide. **2023**, 59, 102579 ○
- 203 Sural hypersensitivity after nerve transection depends on anatomical differences in the distal tibial nerve of mice and rats. **2023**, 246, 152038 1
- 202 Effectiveness of losartan on infrapatellar fat pad/synovial fibrosis and pain behavior in the monoiodoacetate-induced rat model of osteoarthritis pain. **2023**, 158, 114121 ○
- 201 Sex-specific differences in alcohol-induced pain sensitization. **2023**, 225, 109354 ○
- 200 Netrin-1 mediates nerve innervation and angiogenesis leading to discogenic pain. **2023**, 39, 21-33 ○
- 199 Injectable bioorthogonal hydrogel (BIOGEL) accelerates tissue regeneration in degenerated intervertebral discs. **2023**, 23, 551-562 1
- 198 Acacetin alleviates neuroinflammation and oxidative stress injury via the Nrf2/HO-1 pathway in a mouse model of spinal cord injury. **2022**, 13, 483-494 ○
- 197 LncRNA MEG3-TRPV1 signaling regulates chronic inflammatory pain in rats. **2022**, 18, 174480692211442 ○
- 196 Monoaminergic mediation of hyperalgesic and analgesic descending control of nociception in mice. **2022**, Publish Ahead of Print, ○
- 195 Pro-opiomelanocortin neurons in the nucleus of the solitary tract mediate endorphinergic endogenous analgesia in mice. **2022**, Publish Ahead of Print, ○

- 194 Clinically relevant mouse models of Charcot-Marie-Tooth type 2S. ○
- 193 Desulfovibrio confers resilience to the comorbidity of pain and anxiety in a mouse model of chronic inflammatory pain. ○
- 192 Neuron Navigator 1 Regulates Learning, Memory, and the Response to Multiple Potentially Addictive Drugs. ○
- 191 A Purine Derivative Containing an Organoselenium Group Protects Against Memory Impairment, Sensitivity to Nociception, Oxidative Damage, and Neuroinflammation in a Mouse Model of Alzheimer's Disease. ○
- 190 Centrally expressed Cav3.2 T-type calcium channel is critical for the initiation and maintenance of neuropathic pain. 11, ○
- 189 Syringaresinol Alleviates Oxaliplatin-Induced Neuropathic Pain Symptoms by Inhibiting the Inflammatory Responses of Spinal Microglia. **2022**, 27, 8138 ○
- 188 Reversal of Peripheral Neuropathic Pain by the Small-Molecule Natural Product Narirutin via Block of Nav1.7 Voltage-Gated Sodium Channel. **2022**, 23, 14842 ○
- 187 Age and sex drive differential behavioral and neuroimmune phenotypes during postoperative pain. **2022**, ○
- 186 Molecular Mechanics Simulations and Experimental Investigation of the Effect of Tadalafil on Various Inflammatory Pain Mediators. **2022**, 7, 43747-43758 ○
- 185 The anterior cingulate cortex contributes to the analgesic rather than the anxiolytic effects of duloxetine in chronic pain-induced anxiety. 16, ○
- 184 Activation of P2X4 receptors in midbrain cerebrospinal fluid-contacting nucleus leads to mechanical hyperalgesia in chronic constriction injury rats. ○
- 183 JMJD3 is a Key Regulator of Dopamine Synthesis in Ventral Tegmental Area Dopamine Neurons. ○
- 182 Effects of ethanol on mechanical allodynia and dynamic weight bearing in male and female mice with spared nerve injury. ○
- 181 Age-dependent and modality-specific changes in the phenotypic markers Nav1.8, ASIC3, P2X3 and TRPM8 in male rat primary sensory neurons during healthy aging. ○
- 180 C-low threshold mechanoreceptor activation becomes sufficient to trigger affective pain in spinal cord-injured mice in association with increased respiratory rates. 16, ○
- 179 The upregulation of NLRP3 inflammasome in dorsal root ganglion by ten-eleven translocation methylcytosine dioxygenase 2 (TET2) contributed to diabetic neuropathic pain in mice. **2022**, 19, ○
- 178 Sphaeranthus indicus Linn ameliorates streptozotocin-induced experimental diabetic neuropathy by targeting oxidative stress-mediated alterations. **2022**, 8, ○
- 177 Stathmin-2 loss leads to neurofilament-dependent axonal collapse driving motor and sensory denervation. ○

- 176 Spinal glycinergic currents are reduced in a rat model of neuropathic pain following partial nerve ligation but not chronic constriction injury.. ○
- 175 Evaluation of Peripheral Nerve Injury According to the Severity of Damage Using 18F-FDG PET/MRI in a Rat Model of Sciatic Nerve Injury. ○
- 174 Newly repopulated spinal cord microglia exhibit a unique transcriptome and coincide with sex-independent pain resolution. ○
- 173 Cyclovirobuxine D, a cardiovascular drug from traditional Chinese medicine, alleviates inflammatory and neuropathic pain mainly via inhibition of voltage-gated Cav3.2 channels. 13, ○
- 172 New Treatment for the Cognitive and Emotional Deficits Linked with Paclitaxel-Induced Peripheral Neuropathy in Mice. **2022**, 11, 2387 ○
- 171 Andrographolide Relieves Post-Operative Wound Pain but Affects Local Angiogenesis. **2022**, 15, 1586 ○
- 170 Sulfatide-selectin signaling in the spinal cord induces mechanical allodynia. ○
- 169 Intervention mechanism of electroacupuncture on the EP1-TRPV1 pathway in the dorsal root ganglion of rats in the transition from acute to chronic pain. **2022**, ○
- 168 A Novel Mouse Model of Chronic Primary Pain Conditions that Integrates Clinically Relevant Genetic and Environmental Factors. ○
- 167 TIAM1-mediated synaptic plasticity underlies comorbid depression-like and ketamine antidepressant-like actions in chronic pain. **2022**, 132, 1
- 166 Using the DNA methylation profile of the stress driver gene FKBP5 for chronic pain diagnosis. ○
- 165 Bestrophin-1 participates in neuropathic pain induced by spinal nerve transection but not spinal nerve ligation. **2022**, ○
- 164 Lidocaine alleviates inflammation and pruritus in atopic dermatitis by blocking different population of sensory neurons. ○
- 163 Spinal Astrocyte-Neuron Lactate Shuttle Contributes to the Pituitary Adenylate Cyclase-Activating Polypeptide/PAC1 Receptor-Induced Nociceptive Behaviors in Mice. **2022**, 12, 1859 ○
- 162 Synergistic Antinociceptive Interaction upon Co-Administration of Intravenous Ketamine and Lidocaine in a Thermal and Inflammatory Pain Model in Wistar Rats. 2200229 ○
- 161 Paclitaxel Inhibits KCNQ Channels in Primary Sensory Neurons to Initiate the Development of Painful Peripheral Neuropathy. **2022**, 11, 4067 ○
- 160 TMEM100 Regulates Neuropathic Pain by Reducing the Expression of Inflammatory Factors. ○
- 159 TRPV4 activation prevents lipopolysaccharide-induced painful bladder hypersensitivity in rats by regulating immune pathways. 13, ○

- 158 Sustained release of locally delivered celecoxib provides pain relief for osteoarthritis: a proof of concept in dog patients. **2022**, ○
- 157 Non-neuronal TRPA1 encodes mechanical allodynia associated with neurogenic inflammation and partial nerve injury in rats. ○
- 156 Radiocaine: An Imaging Marker of Neuropathic Injury. **2022**, 13, 3661-3667 ○
- 155 Musical therapy attenuates neuroma pain by modifying leptin expression. **2022**, 22, ○
- 154 TRPV1 and GABAB1 in the Cerebrospinal Fluid-Contacting Nucleus are Jointly Involved in Chronic Inflammatory Pain in Rats. Volume 15, 3931-3939 ○
- 153 Discovery of BAY-390, a Selective CNS Penetrant Chemical Probe as Transient Receptor Potential Ankyrin 1 (TRPA1) Antagonist. 1
- 152 The chronification mechanism of orofacial inflammatory pain: Facilitation by GPER1 and microglia in the rostral ventral medulla. 15, ○
- 151 Glucocorticoid nanoformulations relieve chronic pelvic pain syndrome and may alleviate depression in mice. ○
- 150 Betulinic acid analogs inhibit N- and T-type voltage-gated calcium channels to attenuate nerve-injury associated neuropathic and formalin models of pain. **2023**, 13, 100116 ○
- 149 Electroacupuncture ameliorates depression-like behaviors comorbid to chronic neuropathic pain via Tet1-mediated restoration of adult neurogenesis. 1
- 148 Ketolysis is Required for the Proper Development and Function of the Somatosensory Nervous System. ○
- 147 Anti-inflammatory and analgesic effect of Forsythiaside B on complete Freund's adjuvant-induced inflammatory pain in mice. **2023**, ○
- 146 CircFhit Modulates GABAergic Synaptic Transmission via Regulating the Parental Gene Fhit Expression in the Spinal Dorsal Horn in a Rat Model of Neuropathic Pain. ○
- 145 Netrin-3 Suppresses Diabetic Neuropathic Pain by Gating the Intra-epidermal Sprouting of Sensory Axons. ○
- 144 Temporal alterations of pituitary adenylate cyclase activating polypeptide and its receptors in a rat model induced by recurrent chemical stimulations: Relevant to chronic migraine. **2023**, 19, 174480692311521 ○
- 143 Somatostatin Neurons from Periaqueductal Gray to Medulla Facilitate Neuropathic Pain in Male Mice. **2023**, ○
- 142 The involvement of the noradrenergic system in the antinociceptive effect of cucurbitacin D on mice with paclitaxel-induced neuropathic pain. 13, ○
- 141 Transient Reflexive Pain Responses and Chronic Affective Nonreflexive Pain Responses Associated with Neuroinflammation Processes in Both Spinal and Supraspinal Structures in Spinal Cord-Injured Female Mice. **2023**, 24, 1761 ○

- 140 Up-Regulation of ProBDNF/p75NTR Signaling in Spinal Cord Drives Inflammatory Pain in Male Rats. Volume 16, 95-107 ○
- 139 PACAP-PAC1 Receptor Inhibition is Effective in Opioid Induced Hyperalgesia and Medication Overuse Headache Models. **2023**, 105950 ○
- 138 Hypertension contributes to exacerbated osteoarthritis pathophysiology in rats in a sex-dependent manner. **2023**, 25, 1
- 137 Attenuation of SCI-Induced Hypersensitivity by Intensive Locomotor Training and Recombinant GABAergic Cells. **2023**, 10, 84 ○
- 136 Prevention of Chemotherapy-Induced Peripheral Neuropathy by Inhibiting C-X-C Motif Chemokine Receptor 2. **2023**, 24, 1855 ○
- 135 AQP4 mitigates chronic neuropathic pain-induced cognitive impairment in mice. **2023**, 440, 114282 ○
- 134 Mesenchymal stem cell-derived extracellular vesicles carrying miR-99b-3p restrain microglial activation and neuropathic pain by stimulating autophagy. **2023**, 115, 109695 ○
- 133 Artificial Neural Networks Coupled with MALDI-TOF MS Serum Fingerprinting To Classify and Diagnose Pathological Pain Subtypes in Preclinical Models. **2023**, 14, 300-311 ○
- 132 FDG PET Imaging of the Pain Matrix in Neuropathic Pain Model Rats. **2023**, 11, 63 ○
- 131 Antihyperalgesic and Antiallodynic Effects of Amarisolide A and Salvia amarissima Ortega in Experimental Fibromyalgia-Type Pain. **2023**, 13, 59 ○
- 130 Preclinical Studies of Posttraumatic Headache and the Potential Therapeutics. **2023**, 12, 155 ○
- 129 The role of adenosine receptor ligands on inflammatory pain: possible modulation of TRPV1 receptor function. 1
- 128 miR-1306-3p directly activates P2X3 receptors in primary sensory neurons to induce visceral pain in rats. **2022**, Publish Ahead of Print, ○
- 127 Targeting the vascular endothelial growth factor A/neuropilin 1 axis for relief of neuropathic pain. **2022**, Publish Ahead of Print, ○
- 126 Modulation of the inflammatory response by pre-emptive administration of IMT504 reduces postoperative pain in rats and has opioid-sparing effects. **2023**, ○
- 125 Mast cell-derived BH4 is a critical mediator of postoperative pain. ○
- 124 Epidermal SIRT1 and BDNF modulate mechanical allodynia in mouse models of diabetic neuropathy. ○
- 123 Dermal macrophages set pain sensitivity by modulating the amount of tissue NGF through an SNX25/Nrf2 pathway. ○

122	4-methylumbelliferone enhances neuroplasticity in the central nervous system: potential oral treatment for SCI.	1
121	Identification of a human estrogen receptor β -tetrapeptidic fragment with dual antiproliferative and anti-nociceptive action. 2023 , 13,	1
120	Genome-wide association study suggests a critical contribution of the adaptive immune system to chronic post-surgical pain.	0
119	Rodent Animal Models of Endometriosis-Associated Pain: Unmet Needs and Resources Available for Improving Translational Research in Endometriosis. 2023 , 24, 2422	0
118	Salmonella biofilm and its importance in the pathogenesis. 2023 , 447-459	0
117	High fat/high sucrose diet worsens metabolic outcomes and widespread hypersensitivity following early life stress exposure in female mice.	0
116	A sleep-active basalocortical pathway crucial for generation and maintenance of chronic pain.	0
115	Complex I inhibitor of oxidative phosphorylation in advanced solid tumors and acute myeloid leukemia: phase I trials. 2023 , 29, 115-126	3
114	Evidence for Endogenous Opioid Dependence Related to Latent Sensitization in a Rat Model of Chronic Inflammatory Pain. 2023 , 24, 2812	0
113	Time-dependent and selective microglia-mediated removal of spinal synapses in neuropathic pain. 2023 , 42, 112010	0
112	Inducible co-stimulatory molecule (ICOS) alleviates paclitaxel-induced neuropathic pain via an IL-10-mediated mechanism in female mice. 2023 , 20,	0
111	The Role of IL-6 and TMEM100 in Lumbar Discogenic Pain and the Mechanism of the Glycine-Serine-Threonine Metabolic Axis: A Metabolomic and Molecular Biology Study. Volume 16, 437-461	0
110	Neuropeptide Y-expressing dorsal horn inhibitory interneurons gate spinal pain and itch signalling.	0
109	Caraway Oil as a Multimodal Therapy for Neuropathic Pain: Investigating the Mechanisms of Action in Rats with Chronic Constriction Injury. 2023 , 13, 2989	0
108	Stem cell-derived small extracellular vesicles containing miR-27b-3p attenuated osteoarthritis through inhibition of leukaemia inhibitory factor. 2023 ,	0
107	Kinin B1 and B2 receptors mediate cancer pain associated with both the tumor and oncology therapy using aromatase inhibitors. 2023 , 13,	1
106	Nanofat lysate ameliorates pain and cartilage degradation of osteoarthritis through activation of TGF- β 1/Smad2/3 signaling of chondrocytes. 14,	0
105	Multi-omics analysis revealed the role of CYP1A2 in the induction of mechanical allodynia in type 1 diabetes. 14,	0

- 104 No evidence for cognitive impairment in an experimental rat model of knee osteoarthritis and associated chronic pain. **2023**, ○
- 103 Rat model of attention-deficit hyperactivity disorder exhibits delayed recovery from acute incisional pain due to impaired descending noradrenergic inhibition. **2023**, 13, ○
- 102 Inhibition of spinal ferroptosis-like cell death alleviates hyperalgesia and spontaneous pain in a mouse model of bone cancer pain. **2023**, 102700 ○
- 101 Identification of an essential spinoparabrachial pathway for mechanical itch. **2023**, ○
- 100 Kinin B2 and B1 Receptors Activation Sensitize the TRPA1 Channel Contributing to Anastrozole-Induced Pain Symptoms. **2023**, 15, 1136 ○
- 99 Nerve injury-induced gut dysbiosis contributes to spinal cord TNF- α expression and nociceptive sensitization. **2023**, 110, 155-161 ○
- 98 GPR55 contributes to neutrophil recruitment and mechanical pain induction after spinal cord compression in mice. **2023**, 110, 276-287 ○
- 97 Trimetazidine alleviates paclitaxel-induced peripheral neuropathy through modulation of TLR4/p38/NF- κ B and klotho protein expression. **2023**, 376, 110446 ○
- 96 Schwann cell insulin-like growth factor receptor type-1 mediates metastatic bone cancer pain in mice. **2023**, 110, 348-364 ○
- 95 Persistent peripheral inflammation and pain induces immediate early gene activation in supraspinal nuclei in rats. **2023**, 446, 114395 ○
- 94 Phenelzine-based probes reveal Secernin-3 is involved in thermal nociception. **2023**, 125, 103842 ○
- 93 Role of the blood-spinal cord barrier: An adheren junction regulation mechanism that promotes chronic postsurgical pain. **2023**, 660, 65-72 ○
- 92 Spinal dopaminergic D1-and D2-like receptors have a sex-dependent effect in an experimental model of fibromyalgia. **2023**, 948, 175696 ○
- 91 Toxicological and pharmacological effects of *Eugenia brasiliensis* Lam. (Myrtaceae) leaves in mice. **2023**, 309, 116309 ○
- 90 Identification of key genes involved in neural regeneration and the repairing effect of BDNF-overexpressed BMSCs on spinal cord ischemia-reperfusion injury in rats. **2023**, 160, 114293 ○
- 89 Transcranial direct current stimulation attenuates the chronic pain of osteoarthritis in rats via the top-down modulation. ○
- 88 The role of the endocannabinoid 2-arachidonoylglycerol in the in vivo spinal oxytocin-induced antinociception in male rats. **2023**, 363, 114383 ○
- 87 Group II metabotropic glutamate receptor activation suppresses ATP currents in rat dorsal root ganglion neurons. **2023**, 227, 109443 ○

- 86 Satellite glial cells drive the transition from acute to chronic pain in a rat model of hyperalgesic priming. 16, ○
- 85 Red Ginger Extract Prevents the Development of Oxaliplatin-Induced Neuropathic Pain by Inhibiting the Spinal Noradrenergic System in Mice. **2023**, 11, 432 ○
- 84 NFATc2-dependent epigenetic downregulation of the TSC2/Beclin-1 pathway is involved in neuropathic pain induced by oxaliplatin. **2023**, 19, 174480692311582 ○
- 83 Antinociceptive Effects of Aaptamine, a Sponge Component, on Peripheral Neuropathy in Rats. **2023**, 21, 113 ○
- 82 Nanomedicines Reprogram Synovial Macrophages by Scavenging Nitric Oxide and Silencing CA9 in Progressive Osteoarthritis. **2023**, 10, ○
- 81 The role of ectopic P granules protein 5 homolog (EPG5) in DHPG -induced pain sensitization in mice. **2023**, 165, 196-210 ○
- 80 Electroacupuncture Alleviates CFA-Induced Inflammatory Pain via PD-L1/PD-1-SHP-1 Pathway. **2023**, 60, 2922-2936 ○
- 79 Synchronized activity of sensory neurons initiates cortical synchrony in a model of neuropathic pain. **2023**, 14, ○
- 78 A subset of antibodies targeting citrullinated proteins confers protection from rheumatoid arthritis. **2023**, 14, ○
- 77 Anxiety-related behaviors without observation of generalized pain in a mouse model of endometriosis. 17, ○
- 76 miR-26a-5p alleviates CFA-induced chronic inflammatory hyperalgesia through Wnt5a/CaMKII/NFAT signaling in mice. **2023**, 29, 1254-1271 ○
- 75 Critical Pronociceptive Role of Family 2 Voltage-Gated Calcium Channels in a Novel Mouse Model of HIV-Associated Sensory Neuropathy. **2023**, 60, 2954-2968 ○
- 74 Mouse models of surgical and neuropathic pain produce distinct functional alterations to prodynorphin expressing neurons in the prelimbic cortex. **2023**, 13, 100121 ○
- 73 Protease-Activated Receptor 2 (PAR2) expressed in sensory neurons contributes to signs of pain and neuropathy in paclitaxel treated mice. ○
- 72 Polyphyllin VI screened from Chonglou by cell membrane immobilized chromatography relieves inflammatory pain by inhibiting inflammation and normalizing the expression of P2X3 purinoceptor. 14, ○
- 71 Effect of atomoxetine on ADHD-pain hypersensitization comorbidity in 6-OHDA lesioned mice. **2023**, 75, 342-357 ○
- 70 CCL17 Blockade by CCL17mAb/GSK-J4 Ameliorates Hyperalgesia in a Rat Model of Postoperative Pain. **2023**, 46, 533-541 ○
- 69 The Constitutive Activity of Spinal 5-HT6 Receptors Contributes to Diabetic Neuropathic Pain in Rats. **2023**, 13, 364 ○

- 68 Resveratrol suppresses neuroinflammation to alleviate mechanical allodynia by inhibiting Janus kinase 2/signal transducer and activator of transcription 3 signaling pathway in a rat model of spinal cord injury. 16,
- 67 Sex Differences in Behavior and Learning Abilities in Adult Rats. **2023**, 13, 547
- 66 The thalamic reticular nucleus-lateral habenula circuit regulates depressive-like behaviors in chronic stress and chronic pain.
- 65 Pathological Characteristics of Monosodium Iodoacetate-Induced Osteoarthritis in Rats.
- 64 Gut microbiota promotes pain chronicity in Myosin1A deficient male mice.
- 63 Methyl Ferulic Acid Alleviates Neuropathic Pain by Inhibiting Nox4-induced Ferroptosis in Dorsal Root Ganglia Neurons in Rats. **2023**, 60, 3175-3189
- 62 Assessing Orofacial Pain Behaviors in Animal Models: A Review. **2023**, 13, 390
- 61 Paclitaxel A Valuable Tool for Inducing Visceral Pain in Preclinical Testing?. **2023**, 3, 108-119
- 60 PL02 alleviates joint Pain and Inflammation, and subchondral bone deterioration in the Monosodium iodoacetate-induced Osteoarthritis rat model.
- 59 Mechanical and Heat Hyperalgesia upon Withdrawal From Chronic Intermittent Ethanol Vapor Depends on Sex, Exposure Duration, and Blood Alcohol Concentration in Mice. **2023**,
- 58 Aspirin-triggered Resolvin D1 ameliorates activation of the NLRP3 inflammasome via induction of autophagy in a rat model of neuropathic pain. 14,
- 57 Comparison of in vivo behaviors of intramuscularly long-acting celecoxib nanosuspensions with different particle sizes for the postoperative pain treatment. **2023**, 636, 122793
- 56 Excess Growth Hormone Triggers Inflammation-Associated Arthropathy, Subchondral Bone Loss, and Arthralgia. **2023**,
- 55 Low-dose celecoxib-loaded PCL fibers reverse intervertebral disc degeneration by up-regulating CHSY3 expression. **2023**, 21,
- 54 Astrocytic pyruvate dehydrogenase kinase-lactic acid axis involvement in glia-neuron crosstalk contributes to morphine-induced hyperalgesia in mice. **2023**,
- 53 Pharmacokinetics of Orally Applied Cannabinoids and Medical Marijuana Extracts in Mouse Nervous Tissue and Plasma: Relevance for Pain Treatment. **2023**, 15, 853
- 52 Estrogen Mediates the Sexual Dimorphism of GT1b-Induced Central Pain Sensitization. **2023**, 12, 808
- 51 Kinin B1 and B2 Receptors Contribute to Cisplatin-Induced Painful Peripheral Neuropathy in Male Mice. **2023**, 15, 852

- 50 Environmentally stressed human nucleus pulposus cells trigger the onset of discogenic low back pain. ○
- 49 Structure-based discovery of cannabinoid-1 receptor agonists with reduced side effects. ○
- 48 Properties of Nav1.8Chr2-positive and Nav1.8Chr2-negative afferent mechanoreceptors in the hindpaw glabrous skin of mice. **2023**, 16, ○
- 47 Activation of the bile acid receptors TGR5 and FXR in the spinal dorsal horn alleviates neuropathic pain. ○
- 46 Spinal cannabinoid receptor 2 activation alleviates neuropathic pain by regulating microglia and suppressing P2X7 receptor. 16, ○
- 45 Role of Glutamatergic Projections from Lateral Habenula to Ventral Tegmental Area in Inflammatory Pain-Related Spatial Working Memory Deficits. **2023**, 11, 820 ○
- 44 Inhibition of Nonsense-Mediated Decay Induces Nociceptive Sensitization through Activation of the Integrated Stress Response. **2023**, 43, 2921-2933 ○
- 43 Peripheral transient receptor potential vanilloid type 4 hypersensitivity contributes to chronic sickle cell disease pain. **2023**, Publish Ahead of Print, ○
- 42 Sexual Dimorphism in Lesion Size and Sensorimotor Responses Following Spinal Cord Injury. 13, ○
- 41 Combined non-psychoactive Cannabis components cannabidiol and Δ^9 -caryophyllene reduce chronic pain via CB1 interaction in a rat spinal cord injury model. **2023**, 18, e0282920 ○
- 40 Systematic identification of potential key microRNAs and circRNAs in the dorsal root ganglia of mice with sciatic nerve injury. 16, ○
- 39 Human IAPP is a contributor to painful diabetic peripheral neuropathy. **2023**, 133, ○
- 38 Duloxetine and Amitriptyline Reduce Neuropathic Pain by Inhibiting Primary Sensory Input to Spinal Dorsal Horn Neurons via α - and β -Adrenergic Receptors. **2023**, 14, 1261-1277 ○
- 37 Proteomic Analysis of the Spinal Dorsal Horn in Mice with Neuropathic Pain After Exercise. Volume 16, 973-984 ○
- 36 Dopamine D2 receptors in the extended amygdala scale the optimization of seeking and avoidance behaviors. ○
- 35 Cerebellar nuclei neurons projecting to the lateral parabrachial nucleus modulate classical fear conditioning. **2023**, 42, 112291 ○
- 34 Upregulated spinal histone deacetylases induce nociceptive sensitization by inhibiting the GABA system in chronic constriction injury-induced neuropathy in rats. ○
- 33 Identification of brain-to-spinal circuits controlling the laterality and duration of mechanical allodynia in mice. **2023**, 42, 112300 ○

- 32 Effects of 1-methyl-1, 2, 3, 4-tetrahydroisoquinoline on a diabetic neuropathic pain model. 14, ○
- 31 Persistent Changes in Mechanical Nociception in Rats With Traumatic Brain Injury Involving Polytrauma. **2023**, ○
- 30 Voltage-gated calcium channel subunit $\alpha_1\text{H}$ in spinal dorsal horn neurons contributes to aberrant excitatory synaptic transmission and mechanical hypersensitivity after peripheral nerve injury. 16, ○
- 29 Sex differences in the role of parabrachial in nociception and pain in awake mice. ○
- 28 Electroacupuncture at Fengchi(GB20) and Yanglingquan(GB34) Ameliorates Paralgnesia through Microglia-Mediated Neuroinflammation in a Rat Model of Migraine. **2023**, 13, 541 ○
- 27 Through Its Powerful Antioxidative Properties, L-Theanine Ameliorates Vincristine-Induced Neuropathy in Rats. **2023**, 12, 803 ○
- 26 Topiramate affords neuroprotection in diabetic neuropathy model via downregulating spinal GFAP/inflammatory burden and improving neurofilament production. 1-15 ○
- 25 Terpenes from *Cannabis sativa* Induce Antinociception in Mouse Chronic Neuropathic Pain via Activation of Spinal Cord Adenosine A2A Receptors. ○
- 24 Suberoylanilide Hydroxamic Acid Ameliorates Pain Sensitization in Central Neuropathic Pain After Spinal Cord Injury via the HDAC5/NEDD4/SCN9A Axis. ○
- 23 Pilot Studies on the Novel Hypothesis that TSPO (Peripheral Benzodiazepine Receptor) Is Involved in Benzodiazepine- \bar{A} -Drug-Physical Dependence and/or Withdrawal. **2023**, 14, 72-83 ○
- 22 Cannabidiol in the dorsal hippocampus attenuates emotional and cognitive impairments related to neuropathic pain: role of prelimbic neocortex-hippocampal connections. ○
- 21 Evaluation of Photobiomodulation Therapy (117 and 90 seconds) on Pain, Regeneration, and Epigenetic Factors (HDAC 2, DNMT3a) Expression Following Spinal Cord Injury in A Rat Model. ○
- 20 Intrathecal non-viral interleukin-10 gene therapy ameliorates neuropathic pain as measured by both classical static allodynia and a novel supra-spinally mediated pain assay, the Two-Arm Rodent Somatosensory (TARS) task. **2023**, ○
- 19 Effect of Lycopene alone and along with Coenzyme-Q10 in Streptozotocin Induced Peripheral Neuropathy: Biochemical & Behavioural Study. ○
- 18 The effect of risperidone on behavioral reactions and gene expression of pro and anti-inflammatory cytokines in neuropathic pain model induced by chronic constriction injury of the sciatic nerve in rat. ○
- 17 New botulinum neurotoxin constructs for treatment of chronic pain. **2023**, 6, e202201631 ○
- 16 HSP90 inhibition in the mouse spinal cord enhances opioid signaling by suppressing an AMPK-mediated negative feedback loop. **2023**, 16, ○
- 15 Characterization of pain-, anxiety-, and cognition-related behaviors in the complete Freund's adjuvant model of chronic inflammatory pain in Wistar-Kyoto rats. 4, ○

- 14 Transient receptor potential channel vanilloid 1 (TRPV1) contributes to facial mechanical hypersensitivity in a mouse model of atopic asthma. **2023**, 100149 ○
- 13 Reduced pain sensitivity of episodic pain syndrome model mice carrying a Nav1.9 mutation by ANP-230, a novel sodium channel blocker. **2023**, 9, e15423 ○
- 12 Altered Brain Expression of DNA Methylation and Hydroxymethylation Epigenetic Enzymes in a Rat Model of Neuropathic Pain. **2023**, 24, 7305 ○
- 11 The RSK2-RPS6 axis promotes axonal regeneration in the peripheral and central nervous systems. **2023**, 21, e3002044 ○
- 10 Acanthoscurria gomesiana spider-derived Mygalin in the prelimbic prefrontal cortex modulates neuropathic pain and depression comorbid. ○
- 9 Cannabidiol Treatment Shows Therapeutic Efficacy in a Rodent Model of Social Transfer of Pain in Pair-Housed Male Mice. ○
- 8 Ameliorative Effects of Isoeugenol and Eugenol against Impaired Nerve Function and Inflammatory and Oxidative Mediators in Diabetic Neuropathic Rats. **2023**, 11, 1203 ○
- 7 Cingulate protein arginine methyltransferases 1 regulates peripheral hypersensitivity via fragile X messenger ribonucleoprotein. 16, ○
- 6 Mild Traumatic Brain Injury Induced Augmented Postsurgical Pain Is Driven by Central Serotonergic Pain-Facilitatory Signaling. Publish Ahead of Print, ○
- 5 Intra-Articular Delivery of an Indoleamine 2,3-Dioxygenase Galectin-3 Fusion Protein for Osteoarthritis Treatment in Male Lewis Rats. ○
- 4 Establishment of an animal model of sciatic nerve injury induced by local anesthetics. **2023**, 42, 096032712311733 ○
- 3 Periaqueductal grey and spinal cord pathology contribute to pain in Parkinson's disease. **2023**, 9, ○
- 2 Acetaldehyde via CGRP receptor and TRPA1 in Schwann cells mediates ethanol-evoked periorbital mechanical allodynia in mice: relevance for migraine. **2023**, 30, ○
- 1 Scaling up IL glutamatergic outputs to the amygdala alleviates opioid induced hyperalgesia in male rats. ○