

CITATION REPORT

List of articles citing

Premature translational termination triggers mRNA decapping

DOI: 10.1038/370578a0
Nature, 1994, 370, 578-81.

Source: <https://exaly.com/paper-pdf/25191944/citation-report.pdf>

Version: 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
346	Mutational analysis of the neurofibromatosis type 1 gene in childhood myelodysplastic syndromes using a protein truncation assay. 2002 , 68, 157-70		
345	Identification of an additional gene required for eukaryotic nonsense mRNA turnover. 1995 , 92, 10354-8		98
344	Decoying the cap- mRNA degradation system by a double-stranded RNA virus and poly(A)- mRNA surveillance by a yeast antiviral system. <i>Molecular and Cellular Biology</i> , 1995 , 15, 2763-71	4.8	116
343	Dystrophin, its gene, and the dystrophinopathies. 1995 , 33, 177-231		22
342	Initiation of translation can occur only in a restricted region of the CYC1 mRNA of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1995 , 15, 1021-33	4.8	27
341	Identification and characterization of a sequence motif involved in nonsense-mediated mRNA decay. <i>Molecular and Cellular Biology</i> , 1995 , 15, 2231-44	4.8	119
340	Synthetic lethality of <i>sep1</i> (<i>xrn1</i>) <i>ski2</i> and <i>sep1</i> (<i>xrn1</i>) <i>ski3</i> mutants of <i>Saccharomyces cerevisiae</i> is independent of killer virus and suggests a general role for these genes in translation control. <i>Molecular and Cellular Biology</i> , 1995 , 15, 2719-27	4.8	78
339	Translational regulation in response to changes in amino acid availability in <i>Neurospora crassa</i> . <i>Molecular and Cellular Biology</i> , 1995 , 15, 5235-45	4.8	56
338	The <i>ade6</i> gene of the fission yeast as a target for antisense and ribozyme RNA-mediated suppression. 1995 , 5, 295-305		9
337	Characterization of cis-acting sequences and decay intermediates involved in nonsense-mediated mRNA turnover. <i>Molecular and Cellular Biology</i> , 1995 , 15, 809-23	4.8	132
336	Turnover mechanisms of the stable yeast PGK1 mRNA. <i>Molecular and Cellular Biology</i> , 1995 , 15, 2145-56	4.8	281
335	Abundance and half-life of the distinct oat phytochrome A3 and A4 mRNAs. 1995 , 29, 367-77		3
334	A simple p53 functional assay for screening cell lines, blood, and tumors. 1995 , 92, 3963-7		389
333	Trans-acting factors in yeast pre-rRNA and pre-snoRNA processing. 1995 , 73, 803-12		41
332	The <i>Xenopus</i> intron-encoded U17 snoRNA is produced by exonucleolytic processing of its precursor in oocytes. 1995 , 23, 4670-6		31
331	Identification of a novel component of the nonsense-mediated mRNA decay pathway by use of an interacting protein screen. <i>Genes and Development</i> , 1995 , 9, 437-54	12.6	215
330	Multiple functions for the poly(A)-binding protein in mRNA decapping and deadenylation in yeast. <i>Genes and Development</i> , 1995 , 9, 2421-32	12.6	222

329	Purification and characterization of an estrogen-regulated <i>Xenopus</i> liver polysomal nuclease involved in the selective destabilization of albumin mRNA. 1995 , 270, 6108-18		55
328	Degradation of mRNA in eukaryotes. 1995 , 81, 179-83		584
327	Suppression of translation frameshift by upstream termination codon. 1996 , 221, 602-8		6
326	An essential component of the decapping enzyme required for normal rates of mRNA turnover. <i>Nature</i> , 1996 , 382, 642-6	50.4	285
325	From transcript to protein. 1996 , 85, 963-72		65
324	mRNA decapping activities and their biological roles. 1996 , 78, 1049-55		15
323	Nitrogen metabolite signalling involves the C-terminus and the GATA domain of the <i>Aspergillus</i> transcription factor AREA and the 3' untranslated region of its mRNA.. <i>EMBO Journal</i> , 1996 , 15, 2791-2801	13	115
322	Utilizing the GCN4 leader region to investigate the role of the sequence determinants in nonsense-mediated mRNA decay.. <i>EMBO Journal</i> , 1996 , 15, 2810-2819	13	32
321	Effects of nonsense mutations on nuclear and cytoplasmic adenine phosphoribosyltransferase RNA. <i>Molecular and Cellular Biology</i> , 1996 , 16, 4426-35	4.8	61
320	Genetic and biochemical characterization of mutations in the ATPase and helicase regions of the Upf1 protein. <i>Molecular and Cellular Biology</i> , 1996 , 16, 5477-90	4.8	178
319	Mutations in trans-acting factors affecting mRNA decapping in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996 , 16, 5830-8	4.8	152
318	Intracellular targeting and mRNA interactions of the eukaryotic translation initiation factor eIF4E in the yeast <i>Saccharomyces cerevisiae</i> . 1996 , 1308, 142-50		7
317	Purification and characterization of a 5' to 3' exoribonuclease from rabbit reticulocytes that degrades capped and uncapped RNAs. 1996 , 237, 171-9		5
316	Making sense of nonsense in yeast. 1996 , 21, 433-8		86
315	End-plate acetylcholine receptor deficiency due to nonsense mutations in the epsilon subunit. 1996 , 40, 810-7		150
314	Allele-specific measurement of low-density lipoprotein receptor transcript levels. 1996 , 8, 126-33		10
313	Control of mRNA stability in higher plants. 1996 , 32, 63-78		77
312	mRNA sequences influencing translation and the selection of AUG initiator codons in the yeast <i>Saccharomyces cerevisiae</i> . 1996 , 19, 1225-39		55

311	mRNA turnover in yeast promoted by the MATalpha1 instability element. 1996 , 24, 4304-12		30
310	Novel mutations detected in the TSC2 gene from both sporadic and familial TSC patients. 1996 , 5, 249-56		76
309	A mouse cytoplasmic exoribonuclease (mXRN1p) with preference for G4 tetraplex substrates. 1997 , 136, 761-73		282
308	Relationship between yeast polyribosomes and Upf proteins required for nonsense mRNA decay. 1997 , 272, 22163-72		88
307	Cloning and characterization of HUPF1, a human homolog of the <i>Saccharomyces cerevisiae</i> nonsense mRNA-reducing UPF1 protein. 1997 , 25, 814-21		128
306	Destabilization of human alpha-globin mRNA by translation anti-termination is controlled during erythroid differentiation and is paralleled by phased shortening of the poly(A) tail. 1997 , 272, 6607-13		53
305	Cleavage properties of an estrogen-regulated polysomal ribonuclease involved in the destabilization of albumin mRNA. 1997 , 25, 735-42		35
304	Disruption of ribosomal scanning on the 5' untranslated region, and not restriction of translational initiation per se, modulates the stability of nonaberrant mRNAs in the yeast <i>Saccharomyces cerevisiae</i> . 1997 , 272, 9131-40		41
303	Ssd1p of <i>Saccharomyces cerevisiae</i> associates with RNA. 1997 , 272, 16103-9		89
302	Rat1p and Xrn1p are functionally interchangeable exoribonucleases that are restricted to and required in the nucleus and cytoplasm, respectively. <i>Molecular and Cellular Biology</i> , 1997 , 17, 6122-30	4.8	160
301	The role of mRNA stability in the control of globin gene expression. 1997 , 57, 249-87		32
300	Messenger RNA deadenylation precedes decapping in mammalian cells. 1997 , 94, 5628-33		192
299	Life and death in the cytoplasm: messages from the 3' end. 1997 , 7, 220-32		296
298	Ankyrin Bugey: a de novo deletional frameshift variant in exon 6 of the ankyrin gene associated with spherocytosis. 1997 , 54, 242-8		15
297	The Arabidopsis AUX1 gene: a model system to study mRNA processing in plants. 1998 , 36, 463-71		32
296	A single amino acid substitution in yeast eIF-5A results in mRNA stabilization. <i>EMBO Journal</i> , 1998 , 17, 2914-25	13	154
295	Isolation and characterization of Dcp1p, the yeast mRNA decapping enzyme. <i>EMBO Journal</i> , 1998 , 17, 1487-96	13	129
294	mRNA surveillance mitigates genetic dominance in <i>Caenorhabditis elegans</i> . 1998 , 260, 176-84		54

293	CYCLE is a second bHLH-PAS clock protein essential for circadian rhythmicity and transcription of <i>Drosophila</i> period and timeless. 1998 , 93, 805-14		519
292	The 3' to 5' degradation of yeast mRNAs is a general mechanism for mRNA turnover that requires the SKI2 DEVH box protein and 3' to 5' exonucleases of the exosome complex. <i>EMBO Journal</i> , 1998 , 17, 1497-506	13	506
291	Mutational analysis of exoribonuclease I from <i>Saccharomyces cerevisiae</i> . 1998 , 26, 3707-16		41
290	Accelerated mRNA decay in conditional mutants of yeast mRNA capping enzyme. 1998 , 26, 2050-7		81
289	mRNA stabilization by poly(A) binding protein is independent of poly(A) and requires translation. <i>Genes and Development</i> , 1998 , 12, 3226-35	12.6	168
288	The surveillance complex interacts with the translation release factors to enhance termination and degrade aberrant mRNAs. <i>Genes and Development</i> , 1998 , 12, 1665-77	12.6	270
287	Telomere length regulation and telomeric chromatin require the nonsense-mediated mRNA decay pathway. <i>Molecular and Cellular Biology</i> , 1998 , 18, 6121-30	4.8	63
286	Capped mRNA degradation intermediates accumulate in the yeast <i>spb8-2</i> mutant. <i>Molecular and Cellular Biology</i> , 1998 , 18, 5062-72	4.8	130
285	Deadenylation-dependent and -independent decay pathways for alpha1-tubulin mRNA in <i>Chlamydomonas reinhardtii</i> . <i>Molecular and Cellular Biology</i> , 1998 , 18, 1498-505	4.8	22
284	Intron function in the nonsense-mediated decay of beta-globin mRNA: indications that pre-mRNA splicing in the nucleus can influence mRNA translation in the cytoplasm. <i>Rna</i> , 1998 , 4, 801-15	5.8	255
283	Mutations in VPS16 and MRT1 stabilize mRNAs by activating an inhibitor of the decapping enzyme. <i>Molecular and Cellular Biology</i> , 1999 , 19, 7568-76	4.8	31
282	Aberrant mRNAs with extended 3' UTRs are substrates for rapid degradation by mRNA surveillance. <i>Rna</i> , 1999 , 5, 1299-307	5.8	165
281	The cis acting sequences responsible for the differential decay of the unstable MFA2 and stable PGK1 transcripts in yeast include the context of the translational start codon. <i>Rna</i> , 1999 , 5, 420-33	5.8	67
280	Mutations in the MOF2/SUI1 gene affect both translation and nonsense-mediated mRNA decay. <i>Rna</i> , 1999 , 5, 794-804	5.8	29
279	mRNA surveillance in eukaryotes: kinetic proofreading of proper translation termination as assessed by mRNP domain organization?. <i>Rna</i> , 1999 , 5, 711-9	5.8	90
278	Overexpression of truncated Nmd3p inhibits protein synthesis in yeast. <i>Rna</i> , 1999 , 5, 1055-70	5.8	25
277	Nonsense mutations in the alcohol dehydrogenase gene of <i>Drosophila melanogaster</i> correlate with an abnormal 3' end processing of the corresponding pre-mRNA. <i>Rna</i> , 1999 , 5, 562-73	5.8	42
276	An internal open reading frame triggers nonsense-mediated decay of the yeast SPT10 mRNA. <i>EMBO Journal</i> , 1999 , 18, 6134-45	13	89

275	Evidence for a 3P5Pdecay pathway for c-myc mRNA in mammalian cells. 1999 , 274, 16174-9		32
274	Nonsense-mediated mRNA decay in health and disease. 1999 , 8, 1893-900		753
273	Mechanisms of mRNA surveillance in eukaryotes. 1999 , 33, 229-60		180
272	Recognition of yeast mRNAs as "nonsense containing" leads to both inhibition of mRNA translation and mRNA degradation: implications for the control of mRNA decapping. 1999 , 10, 3971-8		75
271	5Pto 3Pexoribonucleolytic activity is a normal component of chloroplast mRNA decay pathways. 1999 , 19, 521-31		60
270	The virion host shutoff function of herpes simplex virus degrades the 5Pend of a target mRNA before the 3Pend. 1999 , 264, 195-204		66
269	The plasminogen activator system: biology and regulation. 1999 , 56, 104-32		265
268	RNA surveillance. Unforeseen consequences for gene expression, inherited genetic disorders and cancer. 1999 , 15, 74-80		342
267	Should we kill the messenger? The role of the surveillance complex in translation termination and mRNA turnover. <i>BioEssays</i> , 1999 , 21, 685-96	4.1	81
266	Current perspectives on mRNA stability in plants: multiple levels and mechanisms of control. 1999 , 4, 429-438		116
265	The DCP2 protein is required for mRNA decapping in <i>Saccharomyces cerevisiae</i> and contains a functional MutT motif. <i>EMBO Journal</i> , 1999 , 18, 5411-22	13	264
264	Rothmund-thomson syndrome responsible gene, RECQL4: genomic structure and products. 1999 , 61, 268-76		139
263	Analysis of mRNA decay pathways in <i>Saccharomyces cerevisiae</i> . 1999 , 17, 3-10		8
262	Linking mRNA turnover and translation: assessing the polyribosomal association of mRNA decay factors and degradative intermediates. 1999 , 17, 28-37		52
261	Monitoring mRNA decapping activity. 1999 , 17, 46-51		49
260	SMG-2 is a phosphorylated protein required for mRNA surveillance in <i>Caenorhabditis elegans</i> and related to Upf1p of yeast. <i>Molecular and Cellular Biology</i> , 1999 , 19, 5943-51	4.8	187
259	An mRNA stability complex functions with poly(A)-binding protein to stabilize mRNA in vitro. <i>Molecular and Cellular Biology</i> , 1999 , 19, 4552-60	4.8	206
258	Assembly of the alpha-globin mRNA stability complex reflects binary interaction between the pyrimidine-rich 3Puntranslated region determinant and poly(C) binding protein alphaCP. <i>Molecular and Cellular Biology</i> , 1999 , 19, 4572-81	4.8	115

257	Novel Upf2p orthologues suggest a functional link between translation initiation and nonsense surveillance complexes. <i>Molecular and Cellular Biology</i> , 2000 , 20, 8944-57	4.8	134
256	Genomic structure, chromosomal localization and identification of mutations in the xeroderma pigmentosum variant (XPV) gene. 2000 , 19, 4721-8		50
255	PIN domains in nonsense-mediated mRNA decay and RNAi. 2000 , 10, R888-90		131
254	Nonsense-mediated decay of glutathione peroxidase 1 mRNA in the cytoplasm depends on intron position. <i>EMBO Journal</i> , 2000 , 19, 4734-44	13	72
253	The Puf3 protein is a transcript-specific regulator of mRNA degradation in yeast. <i>EMBO Journal</i> , 2000 , 19, 6602-11	13	209
252	Cap-dependent deadenylation of mRNA. <i>EMBO Journal</i> , 2000 , 19, 1079-86	13	152
251	A Sm-like protein complex that participates in mRNA degradation. <i>EMBO Journal</i> , 2000 , 19, 1661-71	13	296
250	Upf1p control of nonsense mRNA translation is regulated by Nmd2p and Upf3p. <i>Molecular and Cellular Biology</i> , 2000 , 20, 4591-603	4.8	93
249	Glucose-regulated turnover of mRNA and the influence of poly(A) tail length on half-life. 2000 , 275, 14155-66		22
248	mRNA decapping in yeast requires dissociation of the cap binding protein, eukaryotic translation initiation factor 4E. <i>Molecular and Cellular Biology</i> , 2000 , 20, 7933-42	4.8	143
247	Yeast exosome mutants accumulate 3Pextended polyadenylated forms of U4 small nuclear RNA and small nucleolar RNAs. <i>Molecular and Cellular Biology</i> , 2000 , 20, 441-52	4.8	301
246	<i>Saccharomyces cerevisiae</i> RAI1 (YGL246c) is homologous to human DOM3Z and encodes a protein that binds the nuclear exoribonuclease Rat1p. <i>Molecular and Cellular Biology</i> , 2000 , 20, 4006-15	4.8	101
245	A nuclear 3P5Pexonuclease involved in mRNA degradation interacts with Poly(A) polymerase and the hnRNA protein Npl3p. <i>Molecular and Cellular Biology</i> , 2000 , 20, 604-16	4.8	179
244	The role of nuclear cap binding protein Cbc1p of yeast in mRNA termination and degradation. <i>Molecular and Cellular Biology</i> , 2000 , 20, 2827-38	4.8	58
243	Rapid induction of nuclear transcripts and inhibition of intron decay in response to the polymerase II inhibitor DRB. 2000 , 299, 1179-91		25
242	Human Upf proteins target an mRNA for nonsense-mediated decay when bound downstream of a termination codon. 2000 , 103, 1121-31		456
241	Function of the ski4p (Csl4p) and Ski7p proteins in 3Pto-5Pdegradation of mRNA. <i>Molecular and Cellular Biology</i> , 2000 , 20, 8230-43	4.8	151
240	Mechanisms and control of mRNA decapping in <i>Saccharomyces cerevisiae</i> . 2000 , 69, 571-95		129

239	Viral and cellular mRNA capping: past and prospects. 2000 , 55, 135-84		291
238	Molecular identification of smg-4, required for mRNA surveillance in <i>C. elegans</i> . 2001 , 268, 153-64		26
237	Defects in the mRNA export factors Rat7p, Gle1p, Mex67p, and Rat8p cause hyperadenylation during 3P _{end} formation of nascent transcripts. <i>Rna</i> , 2001 , 7, 753-64	5.8	70
236	The DEAD box helicase, Dhh1p, functions in mRNA decapping and interacts with both the decapping and deadenylase complexes. <i>Rna</i> , 2001 , 7, 1717-27	5.8	268
235	Yeast mRNA decapping enzyme. 2001 , 342, 226-33		8
234	Splicing and 3P _{end} formation in the definition of nonsense-mediated decay-competent human beta-globin mRNPs. <i>EMBO Journal</i> , 2001 , 20, 532-40	13	60
233	A novel mRNA-decapping activity in HeLa cytoplasmic extracts is regulated by AU-rich elements. <i>EMBO Journal</i> , 2001 , 20, 1134-43	13	130
232	The Y14 protein communicates to the cytoplasm the position of exon-exon junctions. <i>EMBO Journal</i> , 2001 , 20, 2062-8	13	155
231	The cap-to-tail guide to mRNA turnover. 2001 , 2, 237-46		632
230	Nonsense-mediated decay of human HEXA mRNA. <i>Molecular and Cellular Biology</i> , 2001 , 21, 5512-9	4.8	74
229	Upf1p, Nmd2p, and Upf3p regulate the decapping and exonucleolytic degradation of both nonsense-containing mRNAs and wild-type mRNAs. <i>Molecular and Cellular Biology</i> , 2001 , 21, 1515-30	4.8	81
228	Characterization of a general stabilizer element that blocks deadenylation-dependent mRNA decay. 2001 , 276, 30995-1003		16
227	The mRNA cap structure stimulates rate of poly(A) removal and amplifies processivity of degradation. 2001 , 276, 27923-9		76
226	mRNA decay enzymes: decappers conserved between yeast and mammals. 2002 , 99, 12512-4		25
225	The hDcp2 protein is a mammalian mRNA decapping enzyme. 2002 , 99, 12663-8		266
224	Regulation of pathways of mRNA destabilization and stabilization. 2002 , 72, 129-64		53
223	Analyzing mRNA decay in <i>Saccharomyces cerevisiae</i> . 2002 , 351, 648-60		17
222	An mRNA surveillance mechanism that eliminates transcripts lacking termination codons. 2002 , 295, 2258-61		421

221	Identification of a human decapping complex associated with hUpf proteins in nonsense-mediated decay. <i>Molecular and Cellular Biology</i> , 2002 , 22, 8114-21	4.8	296
220	Regulation of c-myc mRNA decay by translational pausing in a coding region instability determinant. <i>Molecular and Cellular Biology</i> , 2002 , 22, 3959-69	4.8	115
219	Role of RNA surveillance proteins Upf1/CpaR, Upf2 and Upf3 in the translational regulation of yeast CPA1 gene. 2002 , 41, 224-31		12
218	The scavenger mRNA decapping enzyme DcpS is a member of the HIT family of pyrophosphatases. <i>EMBO Journal</i> , 2002 , 21, 4699-708	13	194
217	SMG-5, required for <i>C.elegans</i> nonsense-mediated mRNA decay, associates with SMG-2 and protein phosphatase 2A. <i>EMBO Journal</i> , 2003 , 22, 641-50	13	107
216	Glucocorticoid receptor-JNK interaction mediates inhibition of the JNK pathway by glucocorticoids. <i>EMBO Journal</i> , 2003 , 22, 6035-44	13	90
215	Genetic background affects relative nonsense mRNA accumulation in wild-type and upf mutant yeast strains. 2003 , 43, 171-7		31
214	Post-transcriptional control of gene expression: effectors of mRNA decay. 2003 , 49, 267-76		36
213	Interaction between Ski7p and Upf1p is required for nonsense-mediated 3Pto-5PmRNA decay in yeast. <i>EMBO Journal</i> , 2003 , 22, 3951-9	13	59
212	Computational modeling and experimental analysis of nonsense-mediated decay in yeast. 2003 , 113, 533-45		122
211	An NMD pathway in yeast involving accelerated deadenylation and exosome-mediated 3P->5P degradation. 2003 , 11, 1405-13		166
210	Nonsense-mediated mRNA decay in mammalian cells involves decapping, deadenylating, and exonucleolytic activities. 2003 , 12, 675-87		290
209	Genome-wide analysis of mRNAs regulated by the nonsense-mediated and 5Pto 3PmRNA decay pathways in yeast. 2003 , 12, 1439-52		331
208	Cytoplasmic degradation of splice-defective pre-mRNAs and intermediates. 2003 , 12, 1453-65		92
207	Molecular biology. A place to die, a place to sleep. 2003 , 300, 753-5		36
206	Rapid deadenylation triggered by a nonsense codon precedes decay of the RNA body in a mammalian cytoplasmic nonsense-mediated decay pathway. <i>Molecular and Cellular Biology</i> , 2003 , 23, 4805-13	4.8	102
205	Nonsense-containing mRNAs that accumulate in the absence of a functional nonsense-mediated mRNA decay pathway are destabilized rapidly upon its restitution. <i>Molecular and Cellular Biology</i> , 2003 , 23, 842-51	4.8	62
204	The Upf-dependent decay of wild-type PPR1 mRNA depends on its 5PUTR and first 92 ORF nucleotides. 2003 , 31, 3157-65		22

203	Degradation of normal mRNA in the nucleus of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2003 , 23, 5502-15	4.8	100
202	Complexes between the nonsense-mediated mRNA decay pathway factor human upf1 (up-frameshift protein 1) and essential nonsense-mediated mRNA decay factors in HeLa cells. 2003 , 373, 775-83		30
201	Leaky termination at premature stop codons antagonizes nonsense-mediated mRNA decay in <i>S. cerevisiae</i> . <i>Rna</i> , 2004 , 10, 691-703	5.8	127
200	Evidence against a direct role for the Upf proteins in frameshifting or nonsense codon readthrough. <i>Rna</i> , 2004 , 10, 1721-9	5.8	28
199	Premature termination codons enhance mRNA decapping in human cells. 2004 , 32, 488-94		57
198	Decapping reaction of mRNA requires Dcp1 in fission yeast: its characterization in different species from yeast to human. 2004 , 136, 805-12		21
197	Nonsense-mediated mRNA decay: splicing, translation and mRNP dynamics. 2004 , 5, 89-99		950
196	The enzymes and control of eukaryotic mRNA turnover. 2004 , 11, 121-7		637
195	Crystal structure of Dcp1p and its functional implications in mRNA decapping. 2004 , 11, 249-56		71
194	Nonsense-mediated messenger RNA decay is initiated by endonucleolytic cleavage in <i>Drosophila</i> . <i>Nature</i> , 2004 , 429, 575-8	50.4	190
193	An enemy within: fly reconnaissance deploys an endonuclease to destroy nonsense-containing mRNA. 2004 , 14, 594-7		6
192	Nonsense-mediated mRNA decay: terminating erroneous gene expression. 2004 , 16, 293-9		347
191	Messenger RNA decay in mammalian cells: the exonuclease perspective. 2004 , 41, 265-78		23
190	Eukaryotic mRNA decapping. 2004 , 73, 861-90		397
189	AtXRN4 degrades mRNA in <i>Arabidopsis</i> and its substrates include selected miRNA targets. 2004 , 15, 173-83		295
188	Regulation of mRNA decay: decapping goes solo. 2004 , 15, 1-2		20
187	SMG7 acts as a molecular link between mRNA surveillance and mRNA decay. 2004 , 16, 587-96		222
186	The role of SMG-1 in nonsense-mediated mRNA decay. 2005 , 1754, 305-15		64

185	Concerted action of poly(A) nucleases and decapping enzyme in mammalian mRNA turnover. 2005 , 12, 1054-63		336
184	Applying the brakes on gene expression. 2005 , 12, 1024-5		1
183	The yeast EDC1 mRNA undergoes deadenylation-independent decapping stimulated by Not2p, Not4p, and Not5p. <i>EMBO Journal</i> , 2005 , 24, 1033-45	13	72
182	RNA decapping inside and outside of processing bodies. 2005 , 17, 326-31		60
181	Nonsense-mediated mRNA decay: molecular insights and mechanistic variations across species. 2005 , 17, 316-25		366
180	A novel frameshift mutation (+G) at codons 15/16 in a beta0 thalassaemia gene results in a significant reduction of beta globin mRNA values. 2005 , 58, 923-6		11
179	Cap-binding protein 1-mediated and eukaryotic translation initiation factor 4E-mediated pioneer rounds of translation in yeast. 2005 , 102, 4258-63		45
178	Scavenger decapping activity facilitates 5Pto 3PmRNA decay. <i>Molecular and Cellular Biology</i> , 2005 , 25, 9764-72	4.8	32
177	Processing bodies require RNA for assembly and contain nontranslating mRNAs. <i>Rna</i> , 2005 , 11, 371-82	5.8	504
176	Novel growth hormone receptor mutation in a Chinese patient with Laron syndrome. 2005 , 18, 209-13		3
175	Gene set coregulated by the <i>Saccharomyces cerevisiae</i> nonsense-mediated mRNA decay pathway. 2005 , 4, 2066-77		18
174	Recruitment and activation of mRNA decay enzymes by two ARE-mediated decay activation domains in the proteins TTP and BRF-1. <i>Genes and Development</i> , 2005 , 19, 351-61	12.6	355
173	Domain interactions within the Ski2/3/8 complex and between the Ski complex and Ski7p. <i>Rna</i> , 2005 , 11, 1291-302	5.8	45
172	Crystal structures of human DcpS in ligand-free and m7GDP-bound forms suggest a dynamic mechanism for scavenger mRNA decapping. 2005 , 347, 707-18		45
171	Multiple processing body factors and the ARE binding protein TTP activate mRNA decapping. 2005 , 20, 905-15		365
170	Identification of the gene encoding the enzyme deficient in mucopolysaccharidosis IIIC (Sanfilippo disease type C). 2006 , 79, 738-44		61
169	Targeting of aberrant mRNAs to cytoplasmic processing bodies. 2006 , 125, 1095-109		244
168	TayBachs Disease. 2006 ,		

167	Early nonsense: mRNA decay solves a translational problem. 2006 , 7, 415-25		210
166	No mercy for messages that mess with the ribosome. 2006 , 13, 299-301		14
165	Synaptotagmin: fusogenic role for calcium sensor?. 2006 , 13, 301-3		30
164	Crystal structure and functional analysis of Dcp2p from <i>Schizosaccharomyces pombe</i> . 2006 , 13, 63-70		86
163	Endonucleolytic cleavage of eukaryotic mRNAs with stalls in translation elongation. <i>Nature</i> , 2006 , 440, 561-4	50.4	500
162	Impact of nonsense-mediated mRNA decay on the global expression profile of budding yeast. 2006 , 2, e203		104
161	Sodium-induced GCN4 expression controls the accumulation of the 5Pto 3PRNA degradation inhibitor, 3Pphosphoadenosine 5Pphosphate. 2006 , 281, 3276-82		15
160	Ebs1p, a negative regulator of gene expression controlled by the Upf proteins in the yeast <i>Saccharomyces cerevisiae</i> . 2006 , 5, 301-12		19
159	Sbp1p affects translational repression and decapping in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2006 , 26, 5120-30	4.8	49
158	Mutant LYS2 mRNAs retained and degraded in the nucleus of <i>Saccharomyces cerevisiae</i> . 2006 , 103, 10871-6		19
157	Conventional 3Pend formation is not required for NMD substrate recognition in <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 2006 , 12, 1441-5	5.8	15
156	Quality control of eukaryotic mRNA: safeguarding cells from abnormal mRNA function. <i>Genes and Development</i> , 2007 , 21, 1833-56	12.6	446
155	An erythroid-enriched endoribonuclease (ErEN) involved in alpha-globin mRNA turnover. 2007 , 14, 131-6		7
154	TTP and BRF proteins nucleate processing body formation to silence mRNAs with AU-rich elements. <i>Genes and Development</i> , 2007 , 21, 719-35	12.6	192
153	Association of yeast Upf1p with direct substrates of the NMD pathway. 2007 , 104, 20872-7		71
152	A genomic screen in yeast reveals novel aspects of nonstop mRNA metabolism. <i>Genetics</i> , 2007 , 177, 773-84		57
151	A specific role for the C-terminal region of the Poly(A)-binding protein in mRNA decay. 2007 , 35, 6017-28		43
150	The decapping activator Lsm1p-7p-Pat1p complex has the intrinsic ability to distinguish between oligoadenylated and polyadenylated RNAs. <i>Rna</i> , 2007 , 13, 998-1016	5.8	111

149	Saccharomyces cerevisiae Ebs1p is a putative ortholog of human Smg7 and promotes nonsense-mediated mRNA decay. 2007 , 35, 7688-97		58
148	Components of the Arabidopsis mRNA decapping complex are required for early seedling development. 2007 , 19, 1549-64		99
147	P bodies and the control of mRNA translation and degradation. 2007 , 25, 635-46		988
146	Initiation of RNA decay in Escherichia coli by 5Ppyrophosphate removal. 2007 , 27, 79-90		200
145	The nonsense-mediated decay RNA surveillance pathway. 2007 , 76, 51-74		938
144	MicroRNAs--micro in size but macro in function. 2008 , 275, 4929-44		112
143	Messenger RNA regulation: to translate or to degrade. <i>EMBO Journal</i> , 2008 , 27, 471-81	13	339
142	Plant stress granules and mRNA processing bodies are distinct from heat stress granules. 2008 , 56, 517-30		179
141	Nuclear pre-mRNA Processing in Plants. <i>Current Topics in Microbiology and Immunology</i> , 2008 ,	3-3	3
140	Nonsense-mediated mRNA decay in yeast does not require PAB1 or a poly(A) tail. 2008 , 29, 134-40		51
139	The control of mRNA decapping and P-body formation. 2008 , 32, 605-15		313
138	A cryptic unstable transcript mediates transcriptional trans-silencing of the Ty1 retrotransposon in <i>S. cerevisiae</i> . <i>Genes and Development</i> , 2008 , 22, 615-26	12.6	157
137	Deadenylation is prerequisite for P-body formation and mRNA decay in mammalian cells. 2008 , 182, 89-101		154
136	Deadenylation-independent stage-specific mRNA degradation in Leishmania. 2008 , 36, 1634-44		25
135	Chapter 1. Methods to study no-go mRNA decay in Saccharomyces cerevisiae. 2008 , 449, 3-21		
134	Analysis of cytoplasmic mRNA decay in Saccharomyces cerevisiae. 2008 , 448, 409-27		15
133	Control of mRNA stability by SAPKs. 2008 , 20, 159-170		3
132	Chapter 6. Qualitative and quantitative assessment of the activity of the yeast nonsense-mediated mRNA decay pathway. 2008 , 449, 127-47		18

131	LSM1 over-expression in <i>Saccharomyces cerevisiae</i> depletes U6 snRNA levels. 2009 , 37, 5529-36		22
130	Structural aspects of RNA helicases in eukaryotic mRNA decay. 2009 , 29, 339-49		3
129	Activation of decapping involves binding of the mRNA and facilitation of the post-binding steps by the <i>Lsm1-7-Pat1</i> complex. <i>Rna</i> , 2009 , 15, 1837-48	5.8	25
128	Execution of nonsense-mediated mRNA decay: what defines a substrate?. 2009 , 21, 394-402		219
127	mRNA Degradation Machinery in Plants. 2009 , 52, 114-124		50
126	SMG6 promotes endonucleolytic cleavage of nonsense mRNA in human cells. 2009 , 16, 49-55		292
125	Crystal structure of Ski8p, a WD-repeat protein with dual roles in mRNA metabolism and meiotic recombination. 2004 , 13, 2673-84		30
124	A convergence of rRNA and mRNA quality control pathways revealed by mechanistic analysis of nonfunctional rRNA decay. 2009 , 34, 440-50		138
123	Cutting the nonsense: the degradation of PTC-containing mRNAs. 2010 , 38, 1615-20		90
122	Regulation of mRNA decapping. 2010 , 1, 253-65		90
121	Nonsense-mediated mRNA decay in human cells: mechanistic insights, functions beyond quality control and the double-life of NMD factors. 2010 , 67, 677-700		246
120	Nonsense-mediated mRNA decapping occurs on polyribosomes in <i>Saccharomyces cerevisiae</i> . 2010 , 17, 244-7		63
119	All things must pass: contrasts and commonalities in eukaryotic and bacterial mRNA decay. 2010 , 11, 467-78		135
118	Regulation of mRNA Turnover. 2010 , 2311-2315		1
117	Translational competence of ribosomes released from a premature termination codon is modulated by NMD factors. <i>Rna</i> , 2010 , 16, 1832-47	5.8	35
116	Identification and analysis of the interaction between Edc3 and Dcp2 in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2010 , 30, 1446-56	4.8	53
115	5Pto 3PmRNA decay factors colocalize with Ty1 gag and human APOBEC3G and promote Ty1 retrotransposition. <i>Journal of Virology</i> , 2010 , 84, 5052-66	6.6	49
114	How and where are nonsense mRNAs degraded in mammalian cells?. 2010 , 7, 28-32		61

113	Interactions between Upf1 and the decapping factors Edc3 and Pat1 in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2011 , 6, e26547	3.7	19
112	Mechanisms of deadenylation-dependent decay. 2011 , 2, 167-83		204
111	trRNA mimicry in translation termination and beyond. 2011 , 2, 647-68		40
110	Differential utilization of decapping enzymes in mammalian mRNA decay pathways. <i>Rna</i> , 2011 , 17, 419-288	3.8	49
109	XRN1: A Major 5' to 3' Exoribonuclease in Eukaryotic Cells. 2012 , 31, 97-114		11
108	<i>Drosophila</i> mutants show NMD pathway activity is reduced, but not eliminated, in the absence of Smg6. <i>Rna</i> , 2012 , 18, 1475-86	5.8	13
107	Modulation of poly(A)-specific ribonuclease (PARN): current knowledge and perspectives. 2012 , 19, 4838-49		23
106	Analysis of nonsense-mediated mRNA decay in <i>Saccharomyces cerevisiae</i> . 2012 , Chapter 27, Unit 27.3		5
105	Translation drives mRNA quality control. 2012 , 19, 594-601		271
104	Ways and means of eukaryotic mRNA decay. 2012 , 1819, 593-603		50
103	Mechanism of the initiation of mRNA decay: role of eRF3 family G proteins. 2012 , 3, 743-57		23
102	P-bodies and stress granules: possible roles in the control of translation and mRNA degradation. 2012 , 4, a012286		433
101	RNA degradation in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2012 , 191, 671-702	4	260
100	Roles of target site location and sequence complementarity in trans-acting siRNA formation in <i>Arabidopsis</i> . 2012 , 69, 217-26		29
99	mRNP quality control goes regulatory. 2012 , 28, 70-7		38
98	mRNA quality control pathways in <i>Saccharomyces cerevisiae</i> . 2013 , 38, 615-40		8
97	The Hbs1-Dom34 protein complex functions in non-stop mRNA decay in mammalian cells. 2013 , 288, 17832-43		57
96	The Upf factor complex interacts with aberrant products derived from mRNAs containing a premature termination codon and facilitates their proteasomal degradation. 2013 , 288, 28630-40		22

95	Ten Years of Progress in GW/P Body Research. <i>Advances in Experimental Medicine and Biology</i> , 2013 ,	3.6	4
94	Nonsense-mediated mRNA decay: from mechanistic insights to impacts on human health. 2013 , 12, 25-36		28
93	Biogenesis, turnover, and mode of action of plant microRNAs. 2013 , 25, 2383-99		640
92	Structural and functional control of the eukaryotic mRNA decapping machinery. 2013 , 1829, 580-9		86
91	Deadenylation and P-bodies. <i>Advances in Experimental Medicine and Biology</i> , 2013 , 768, 183-95	3.6	16
90	The SMG5-SMG7 heterodimer directly recruits the CCR4-NOT deadenylase complex to mRNAs containing nonsense codons via interaction with POP2. <i>Genes and Development</i> , 2013 , 27, 2125-38	12.6	125
89	The role of disordered protein regions in the assembly of decapping complexes and RNP granules. <i>Genes and Development</i> , 2013 , 27, 2628-41	12.6	136
88	Multiple transcripts from a 3'UTR reporter vary in sensitivity to nonsense-mediated mRNA decay in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2013 , 8, e80981	3.7	8
87	The Not4 RING E3 Ligase: A Relevant Player in Cotranslational Quality Control. 2013 , 2013, 548359		13
86	A novel NF1 frame-shift mutation (c.702_703delGT) in a Chinese family with neurofibromatosis type 1. 2014 , 13, 5395-404		5
85	Human nonsense-mediated RNA decay initiates widely by endonucleolysis and targets snoRNA host genes. <i>Genes and Development</i> , 2014 , 28, 2498-517	12.6	119
84	Protecting the proteome: Eukaryotic cotranslational quality control pathways. 2014 , 204, 467-76		102
83	EXOSC8 mutations alter mRNA metabolism and cause hypomyelination with spinal muscular atrophy and cerebellar hypoplasia. 2014 , 5, 4287		80
82	Fungal RNA Biology. 2014 ,		3
81	Cbc2p, Upf3p and eIF4G are components of the DRN (Degradation of mRNA in the Nucleus) in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2014 , 14, 922-32	3.1	8
80	Yeast Edc3 targets RPS28B mRNA for decapping by binding to a 3'UTR untranslated region decay-inducing regulatory element. <i>Molecular and Cellular Biology</i> , 2014 , 34, 1438-51	4.8	18
79	Expression of Subtelomeric lncRNAs Links Telomeres Dynamics to RNA Decay in. <i>Non-coding RNA</i> , 2015 , 1, 94-126	7.1	7
78	The messenger RNA decapping and recapping pathway in <i>Trypanosoma</i> . 2015 , 112, 6967-72		27

77	Not4-dependent translational repression is important for cellular protein homeostasis in yeast. <i>EMBO Journal</i> , 2015 , 34, 1905-24	13	34
76	Nonsense-Mediated mRNA Decay: Degradation of Defective Transcripts Is Only Part of the Story. 2015 , 49, 339-66		167
75	Use of Cellular Decapping Activators by Positive-Strand RNA Viruses. 2016 , 8,		6
74	ATP hydrolysis by UPF1 is required for efficient translation termination at premature stop codons. 2016 , 7, 14021		28
73	Interrogating the degradation pathways of unstable mRNAs with XRN1-resistant sequences. 2016 , 7, 13691		24
72	Targeted mRNA Degradation. 2016 , 317-326		
71	RNA decay systems enhance reciprocal switching of sense and antisense transcripts in response to glucose starvation. 2016 , 21, 1276-1289		6
70	Nonsense-Mediated Decay Restricts LncRNA Levels in Yeast Unless Blocked by Double-Stranded RNA Structure. 2016 , 61, 379-392		67
69	The cytoplasmic mRNA degradation factor Pat1 is required for rRNA processing. 2016 , 13, 455-65		3
68	LncRNAs, lost in translation or licence to regulate?. 2017 , 63, 29-33		28
67	High-resolution profiling of NMD targets in yeast reveals translational fidelity as a basis for substrate selection. <i>Rna</i> , 2017 , 23, 735-748	5.8	51
66	Nonsense-mediated mRNA decay in Tetrahymena is EJC independent and requires a protozoa-specific nuclease. 2017 , 45, 6848-6863		14
65	Feline leukocyte adhesion (CD18) deficiency caused by a deletion in the integrin α (ITGB2) gene. 2017 , 46, 391-400		6
64	Ksp1-dependent phosphorylation of eIF4G modulates post-transcriptional regulation of specific mRNAs under glucose deprivation conditions. 2018 , 46, 3047-3060		8
63	RNA stability and metabolism in regulated cell death, aging and diseases. <i>FEMS Yeast Research</i> , 2018 , 18,	3.1	11
62	Birth, coming of age and death: The intriguing life of long noncoding RNAs. 2018 , 79, 143-152		10
61	Nonsense-mediated mRNA decay involves two distinct Upf1-bound complexes. <i>EMBO Journal</i> , 2018 , 37,	13	26
60	Human MARF1 is an endoribonuclease that interacts with the DCP1:2 decapping complex and degrades target mRNAs. 2018 , 46, 12008-12021		12

59	Nonsense mRNA suppression via nonstop decay. <i>ELife</i> , 2018 , 7,	8.9	31
58	The UPF1 interactome reveals interaction networks between RNA degradation and translation repression factors in Arabidopsis. 2018 , 96, 119-132		30
57	Destabilization of Eukaryote mRNAs by 5' Proximal Stop Codons Can Occur Independently of the Nonsense-Mediated mRNA Decay Pathway. 2019 , 8,		1
56	RNA degradomes reveal substrates and importance for dark and nitrogen stress responses of Arabidopsis XRN4. 2019 , 47, 9216-9230		16
55	Cauliflower mosaic virus transactivator protein (TAV) can suppress nonsense-mediated decay by targeting VARICOSE, a scaffold protein of the decapping complex. 2019 , 9, 7042		12
54	States of decay: The systems biology of mRNA stability. 2019 , 15, 48-57		2
53	In silico analysis of codon usage and rare codon clusters in the halophilic bacteria L-asparaginase. 2020 , 75, 151-160		1
52	When Long Noncoding Becomes Protein Coding. <i>Molecular and Cellular Biology</i> , 2020 , 40,	4.8	49
51	Nonsense-Mediated mRNA Decay: Pathologies and the Potential for Novel Therapeutics. 2020 , 12,		18
50	mRNA Editing, Processing and Quality Control in. <i>Genetics</i> , 2020 , 215, 531-568	4	9
49	Smart-ORF: a single-molecule method for accessing ribosome dynamics in both upstream and main open reading frames. 2021 , 49, e26		0
48	Nuclear SR-protein mediated mRNA quality control is continued in cytoplasmic nonsense-mediated decay. 2021 , 18, 1390-1407		2
47	A synonymous mutation results in exon skipping, loss of function and worse patient prognosis. 2021 , 24, 102173		4
46	The RNA-binding protein Puf5 contributes to buffering of mRNA upon chromatin-mediated changes in nascent transcription. <i>Journal of Cell Science</i> , 2021 , 134,	5.3	
45	From Yeast to Mammals, the Nonsense-Mediated mRNA Decay as a Master Regulator of Long Non-Coding RNAs Functional Trajectory. <i>Non-coding RNA</i> , 2021 , 7,	7.1	4
44	LISTERIN E3 Ubiquitin Ligase and Ribosome-Associated Quality Control (RQC) Mechanism. <i>Molecular Neurobiology</i> , 2021 , 58, 6593-6609	6.2	1
43	Functions of the cytoplasmic exosome. <i>Advances in Experimental Medicine and Biology</i> , 2011 , 702, 79-90	3.6	39
42	State of decay: an update on plant mRNA turnover. <i>Current Topics in Microbiology and Immunology</i> , 2008 , 326, 179-99	3.3	8

41	Control of mRNA stability in higher plants. 1996 , 63-77		1
40	Analysis of mutations in the yeast mRNA decapping enzyme. <i>Genetics</i> , 1999 , 151, 1273-85	4	64
39	Temperature-sensitive mutations in the <i>Saccharomyces cerevisiae</i> MRT4, GRC5, SLA2 and THS1 genes result in defects in mRNA turnover. <i>Genetics</i> , 1999 , 153, 35-47	4	49
38	Two related proteins, Edc1p and Edc2p, stimulate mRNA decapping in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2001 , 157, 27-37	4	77
37	The yeast cytoplasmic Lsm1/Pat1p complex protects mRNA 3' termini from partial degradation. <i>Genetics</i> , 2001 , 158, 1445-55	4	80
36	Curbing the nonsense: the activation and regulation of mRNA surveillance. <i>Genes and Development</i> , 2001 , 15, 2781-5	12.6	76
35	Nonsense-mediated mRNA decay: insights into mechanism from the cellular abundance of human Upf1, Upf2, Upf3, and Upf3X proteins. <i>Cold Spring Harbor Symposia on Quantitative Biology</i> , 2001 , 66, 313-20	3.9	18
34	Nonsense-mediated mRNA decay in <i>Saccharomyces cerevisiae</i> : a quality control mechanism that degrades transcripts harboring premature termination codons. <i>Cold Spring Harbor Symposia on Quantitative Biology</i> , 2001 , 66, 321-8	3.9	16
33	Translation but not the encoded sequence is essential for the efficient propagation of the defective interfering RNAs of the coronavirus mouse hepatitis virus. <i>Journal of Virology</i> , 1995 , 69, 3744-51	6.6	41
32	Replication and packaging of coronavirus infectious bronchitis virus defective RNAs lacking a long open reading frame. <i>Journal of Virology</i> , 1996 , 70, 8660-8	6.6	51
31	Posttranscriptional control of gene expression in yeast. <i>Microbiology and Molecular Biology Reviews</i> , 1998 , 62, 1492-553	13.2	173
30	mRNA stability in mammalian cells. <i>Microbiological Reviews</i> , 1995 , 59, 423-50		1083
29	Mechanisms and control of mRNA turnover in <i>Saccharomyces cerevisiae</i> . <i>Microbiological Reviews</i> , 1996 , 60, 233-49		221
28	A nonsense mutation in the erythrocyte band 3 gene associated with decreased mRNA accumulation in a kindred with dominant hereditary spherocytosis. <i>Journal of Clinical Investigation</i> , 1996 , 97, 373-80	15.9	28
27	A factor required for nonsense-mediated mRNA decay in yeast is exported from the nucleus to the cytoplasm by a nuclear export signal sequence. <i>Journal of Cell Science</i> , 1998 , 111, 3129-3143	5.3	47
26	mRNA surveillance: the perfect persist. <i>Journal of Cell Science</i> , 2002 , 115, 3033-3038	5.3	164
25	Multispecies reconstructions uncover widespread conservation, and lineage-specific elaborations in eukaryotic mRNA metabolism. <i>PLoS ONE</i> , 2018 , 13, e0192633	3.7	12
24	Isolation and characterization of a mutant defective in utilization of exogenous phosphatidylethanolamine in <i>Saccharomyces cerevisiae</i> . <i>Journal of General and Applied Microbiology</i> , 2007 , 53, 255-8	1.5	2

23	Transcriptome maps of general eukaryotic RNA degradation factors. <i>ELife</i> , 2019 , 8,	8.9	15
22	Nuclear mRNA Quality Control and Cytoplasmic NMD Are Linked by the Guard Proteins Gbp2 and Hrb1. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	0
21	Turnover of mRNA In Eukaryotic Cells. 2001 , 245-257		2
20	Post-Transcriptional Control of the GAP-43 mRNA by the ELAV-Like Protein HuD. <i>Growth Hormone</i> , 2002 , 157-174		
19	Transcription, Translation, and the Control of Gene Expression.		
18	Cytoplasmic mRNA Surveillance Pathways. 2014 , 195-216		
17	Messenger RNA Ribonucleases and mRNA Turnover in <i>Saccharomyces cerevisiae</i> . 1997 , 589-620		
16	Turnover of mRNA in Eukaryotic Cells. 1999 , 205-216		
15	Detection and Degradation of Nonsense-mediated mRNA Decay Substrates Involve Two Distinct Upf1-bound Complexes.		
14	The RNA-binding protein Puf5 buffers mRNA levels against chromatin-mediated changes in nascent transcription.		
13	In silico Evaluation of Substrate Binding Site and Rare Codons in the Structure of CYP152A1. <i>Current Proteomics</i> , 2020 , 17, 10-22	0.7	
12	Nitrogen metabolite signalling involves the C-terminus and the GATA domain of the <i>Aspergillus</i> transcription factor AREA and the 3' untranslated region of its mRNA. <i>EMBO Journal</i> , 1996 , 15, 2791-801 ¹³		56
11	Utilizing the GCN4 leader region to investigate the role of the sequence determinants in nonsense-mediated mRNA decay. <i>EMBO Journal</i> , 1996 , 15, 2810-9	13	26
10	Interaction between Nmd2p and Upf1p is required for activity but not for dominant-negative inhibition of the nonsense-mediated mRNA decay pathway in yeast. <i>Rna</i> , 1996 , 2, 153-70	5.8	52
9	When cells stop making sense: effects of nonsense codons on RNA metabolism in vertebrate cells. <i>Rna</i> , 1995 , 1, 453-65	5.8	393
8	Progressive polysomal association of Upf1, Upf2, and Upf3 as ribosomes traverse mRNA coding regions.		
7	Unusual SMG suspects recruit degradation enzymes in nonsense-mediated mRNA decay.. <i>BioEssays</i> , 2022 , e2100296	4.1	
6	An Evolutionarily Conserved AU-Rich Element in the 3' Untranslated Region of a Transcript Misannotated as a Long Noncoding RNA Regulates RNA Stability.. <i>Molecular and Cellular Biology</i> , 2022 , e0050521	4.8	

5	Post-transcriptional regulation during stress.. <i>FEMS Yeast Research</i> , 2022 ,	3.1	○
4	Eukaryotic mRNA decapping factors: molecular mechanisms and activity.		○
3	The Destabilization of IL-2 mRNA by a Premature Stop Codon and Its Differential Stabilization by Trans-Acting Inhibitors of Protein Synthesis Do Not Support a Role for Active Translation in mRNA Stability. 1999 , 163, 3321-3330	8	
2	Recoding of Nonsense Mutation as a Pharmacological Strategy. 2023 , 11, 659		○
1	Disruption of Dcp1 leads to a Dcp2-dependent aberrant ribosome profiles in <i>Aspergillus nidulans</i> .		○