

CITATION REPORT

List of articles citing

Deadenylation of the unstable mRNA encoded by the yeast MFA2 gene leads to decapping followed by 5'→3' digestion of the transcript

DOI: 10.1101/gad.8.7.855

Genes and Development, 1994, 8, 855-66.

Source: <https://exaly.com/paper-pdf/25441472/citation-report.pdf>

Version: 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
442	A regulatory role for the 5' and 3' untranslated regions in differential expression of hsp83 in <i>Leishmania</i> . <i>Nucleic Acids Research</i> , 1994 , 22, 2922-9	20.1	68
441	Selective degradation of early-response-gene mRNAs: functional analyses of sequence features of the AU-rich elements. 1994 , 14, 8471-82		236
440	Metabolic stability of mRNA in yeast--a potential target for modulating productivity?. 1994 , 12, 444-9		7
439	Mechanisms of mRNA degradation in eukaryotes. 1994 , 19, 336-40		212
438	Premature translational termination triggers mRNA decapping. 1994 , 370, 578-81		332
437	The levels of yeast gluconeogenic mRNAs respond to environmental factors. 1994 , 224, 473-81		43
436	Detection and characterization of a 3' untranslated region ribonucleoprotein complex associated with human alpha-globin mRNA stability. 1995 , 15, 1769-77		186
435	Erythroid cell-specific mRNA stability elements in the alpha 2-globin 3' untranslated region. 1995 , 15, 2457-65		158
434	Identification of an additional gene required for eukaryotic nonsense mRNA turnover. 1995 , 92, 10354-8		98
433	Polyadenylation helps regulate mRNA decay in <i>Escherichia coli</i> . 1995 , 92, 1807-11		224
432	Effects of mutations in the <i>Saccharomyces cerevisiae</i> RNA14, RNA15, and PAP1 genes on polyadenylation in vivo. 1995 , 15, 6979-86		50
431	Identification and characterization of a sequence motif involved in nonsense-mediated mRNA decay. 1995 , 15, 2231-44		119
430	Synthetic lethality of <i>sep1</i> (<i>xrn1</i>) <i>ski2</i> and <i>sep1</i> (<i>xrn1</i>) <i>ski3</i> mutants of <i>Saccharomyces cerevisiae</i> is independent of killer virus and suggests a general role for these genes in translation control. 1995 , 15, 2719-27		78
429	An efficient strategy to isolate full-length cDNAs based on an mRNA cap retention procedure (CAPture). 1995 , 15, 3363-71		71
428	Degradation of the soybean ribulose-1,5-bisphosphate carboxylase small-subunit mRNA, SRS4, initiates with endonucleolytic cleavage. 1995 , 15, 6641-52		23
427	Characterization of cis-acting sequences and decay intermediates involved in nonsense-mediated mRNA turnover. 1995 , 15, 809-23		132
426	Turnover mechanisms of the stable yeast PGK1 mRNA. 1995 , 15, 2145-56		281

425	The nonamer UUAUUUAUU is the key AU-rich sequence motif that mediates mRNA degradation. 1995 , 15, 2219-30		460
424	Poly(A) tail length control is caused by termination of processive synthesis. 1995 , 270, 2800-8		147
423	Trans-acting factors in yeast pre-rRNA and pre-snoRNA processing. 1995 , 73, 803-12		41
422	Exonucleolytic processing of small nucleolar RNAs from pre-mRNA introns. <i>Genes and Development</i> , 1995 , 9, 1411-24	12.6	127
421	5' exonuclease-2 of <i>Saccharomyces cerevisiae</i> . Purification and features of ribonuclease activity with comparison to 5' exonuclease-1. 1995 , 270, 16063-9		98
420	Multiple functions for the poly(A)-binding protein in mRNA decapping and deadenylation in yeast. <i>Genes and Development</i> , 1995 , 9, 2421-32	12.6	222
419	Purification and characterization of an estrogen-regulated <i>Xenopus</i> liver polysomal nuclease involved in the selective destabilization of albumin mRNA. 1995 , 270, 6108-18		55
418	Glucose-dependent turnover of the mRNAs encoding succinate dehydrogenase peptides in <i>Saccharomyces cerevisiae</i> : sequence elements in the 5' untranslated region of the <i>Ip</i> mRNA play a dominant role. 1995 , 6, 1125-43		66
417	Diversity of cytoplasmic functions for the 3' untranslated region of eukaryotic transcripts. 1995 , 7, 386-92		214
416	Degradation of mRNA in eukaryotes. 1995 , 81, 179-83		584
415	Degradation of maternal string mRNA is controlled by proteins encoded on maternally contributed transcripts. 1995 , 51, 217-26		12
414	Identification of functional domains in the Sep1 protein (= Kem1, Xrn1), which is required for transition through meiotic prophase in <i>Saccharomyces cerevisiae</i> . 1995 , 104, 215-22		19
413	An essential component of the decapping enzyme required for normal rates of mRNA turnover. 1996 , 382, 642-6		285
412	From transcript to protein. 1996 , 85, 963-72		65
411	mRNA decapping activities and their biological roles. 1996 , 78, 1049-55		15
410	Multiple steps in the regulation of transcription-factor level and activity. 1996 , 317 (Pt 2), 329-42		125
409	Mutations in trans-acting factors affecting mRNA decapping in <i>Saccharomyces cerevisiae</i> . 1996 , 16, 5830-8		152
408	Functional mapping of the translation-dependent instability element of yeast MAT α 1 mRNA. 1996 , 16, 3833-43		33

407	Degradation of CYC1 mRNA in the yeast <i>Saccharomyces cerevisiae</i> does not require translation. 1996 , 93, 8895-900		6
406	Purification and characterization of a 5' to 3' exoribonuclease from rabbit reticulocytes that degrades capped and uncapped RNAs. 1996 , 237, 171-9		5
405	Control of mRNA stability in higher plants. 1996 , 32, 63-78		77
404	mRNA turnover in yeast promoted by the MAT α 1 instability element. <i>Nucleic Acids Research</i> , 1996 , 24, 4304-12	20.1	30
403	The 3' end of yeast 5.8S rRNA is generated by an exonuclease processing mechanism. <i>Genes and Development</i> , 1996 , 10, 502-13	12.6	153
402	Regulated nuclear polyadenylation of <i>Xenopus</i> albumin pre-mRNA. <i>Nucleic Acids Research</i> , 1996 , 24, 4078-83	20.1	16
401	Fatty acid-responsive control of mRNA stability. Unsaturated fatty acid-induced degradation of the <i>Saccharomyces</i> OLE1 transcript. 1996 , 271, 25801-9		71
400	PAN3 encodes a subunit of the Pab1p-dependent poly(A) nuclease in <i>Saccharomyces cerevisiae</i> . 1996 , 16, 5744-53		140
399	A mouse cytoplasmic exoribonuclease (mXRN1p) with preference for G4 tetraplex substrates. 1997 , 136, 761-73		282
398	The Frequency and Degree of Cosuppression by Sense Chalcone Synthase Transgenes Are Dependent on Transgene Promoter Strength and Are Reduced by Premature Nonsense Codons in the Transgene Coding Sequence. 1997 , 9, 1357-1368		150
397	Translation initiation factors eIF-iso4G and eIF-4B interact with the poly(A)-binding protein and increase its RNA binding activity. 1997 , 272, 16247-55		210
396	Postfertilization deadenylation of mRNAs in <i>Xenopus laevis</i> embryos is sufficient to cause their degradation at the blastula stage. 1997 , 17, 209-18		68
395	Destabilization of human alpha-globin mRNA by translation anti-termination is controlled during erythroid differentiation and is paralleled by phased shortening of the poly(A) tail. 1997 , 272, 6607-13		53
394	Ribosome concentration contributes to discrimination against poly(A)- mRNA during translation initiation in <i>Saccharomyces cerevisiae</i> . 1997 , 272, 6004-10		29
393	Poly(A) tail shortening by a mammalian poly(A)-specific 3' exoribonuclease. 1997 , 272, 10448-56		175
392	Cleavage properties of an estrogen-regulated polysomal ribonuclease involved in the destabilization of albumin mRNA. <i>Nucleic Acids Research</i> , 1997 , 25, 735-42	20.1	35
391	The poly(A) tail inhibits the assembly of a 3' to 5' exonuclease in an in vitro RNA stability system. 1997 , 17, 398-406		91
390	Modulation of the fate of cytoplasmic mRNA by AU-rich elements: key sequence features controlling mRNA deadenylation and decay. 1997 , 17, 4611-21		315

389	Rok1p is a putative RNA helicase required for rRNA processing. 1997 , 17, 3398-407		92
388	Rat1p and Xrn1p are functionally interchangeable exoribonucleases that are restricted to and required in the nucleus and cytoplasm, respectively. 1997 , 17, 6122-30		160
387	The role of mRNA stability in the control of globin gene expression. 1997 , 57, 249-87		32
386	Messenger RNA deadenylylation precedes decapping in mammalian cells. 1997 , 94, 5628-33		192
385	A calcium-metalloribozyme with autodecapping and pyrophosphatase activities. 1997 , 36, 14107-19		14
384	Structural modifications of RNA influence the 5' exoribonucleolytic hydrolysis by XRN1 and HKE1 of <i>Saccharomyces cerevisiae</i> . 1997 , 235, 799-805		39
383	Life and death in the cytoplasm: messages from the 3' end. 1997 , 7, 220-32		296
382	Cloning and characterization of mouse Dhm2 cDNA, a functional homolog of budding yeast SEP1. 1997 , 191, 161-6		11
381	A viral sequence in the 3' untranslated region mimics a 5' cap in facilitating translation of uncapped mRNA. <i>EMBO Journal</i> , 1997 , 16, 4107-16	13	124
380	Lithium toxicity in yeast is due to the inhibition of RNA processing enzymes. <i>EMBO Journal</i> , 1997 , 16, 7184-95	13	201
379	A single amino acid substitution in yeast eIF-5A results in mRNA stabilization. <i>EMBO Journal</i> , 1998 , 17, 2914-25	13	154
378	The deadenylating nuclease (DAN) is involved in poly(A) tail removal during the meiotic maturation of <i>Xenopus</i> oocytes. <i>EMBO Journal</i> , 1998 , 17, 5427-37	13	206
377	Isolation and characterization of Dcp1p, the yeast mRNA decapping enzyme. <i>EMBO Journal</i> , 1998 , 17, 1487-96	13	129
376	A tale of two termini: a functional interaction between the termini of an mRNA is a prerequisite for efficient translation initiation. 1998 , 216, 1-11		240
375	8 Transcript Analysis. 1998 , 26, 119-139		
374	Identification and developmental expression of a 5' exoribonuclease from <i>Drosophila melanogaster</i> . 1998 , 79, 51-5		42
373	The 3' to 5' degradation of yeast mRNAs is a general mechanism for mRNA turnover that requires the SKI2 DEVH box protein and 3' to 5' exonucleases of the exosome complex. <i>EMBO Journal</i> , 1998 , 17, 1497-506	13	506
372	Mutational analysis of exoribonuclease I from <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1998 , 26, 3707-16	20.1	41

371	Accelerated mRNA decay in conditional mutants of yeast mRNA capping enzyme. <i>Nucleic Acids Research</i> , 1998 , 26, 2050-7	20.1	81
370	Control of translation initiation in animals. 1998 , 14, 399-458		447
369	A mutated human homologue to yeast Upf1 protein has a dominant-negative effect on the decay of nonsense-containing mRNAs in mammalian cells. 1998 , 95, 10009-14		157
368	mRNA stabilization by poly(A) binding protein is independent of poly(A) and requires translation. <i>Genes and Development</i> , 1998 , 12, 3226-35	12.6	168
367	AUUUA sequences direct mRNA deadenylation uncoupled from decay during <i>Xenopus</i> early development. 1998 , 18, 7537-45		91
366	Telomere length regulation and telomeric chromatin require the nonsense-mediated mRNA decay pathway. 1998 , 18, 6121-30		63
365	RNA polymerase I-promoted <i>HIS4</i> expression yields uncapped, polyadenylated mRNA that is unstable and inefficiently translated in <i>Saccharomyces cerevisiae</i> . 1998 , 18, 665-75		41
364	Poly(A) tail length control in <i>Saccharomyces cerevisiae</i> occurs by message-specific deadenylation. 1998 , 18, 6548-59		178
363	Capped mRNA degradation intermediates accumulate in the yeast <i>spb8-2</i> mutant. 1998 , 18, 5062-72		130
362	Processing of the intron-encoded U18 small nucleolar RNA in the yeast <i>Saccharomyces cerevisiae</i> relies on both exo- and endonucleolytic activities. 1998 , 18, 3376-83		57
361	Deadenylation-dependent and -independent decay pathways for alpha1-tubulin mRNA in <i>Chlamydomonas reinhardtii</i> . 1998 , 18, 1498-505		22
360	Processing of the precursors to small nucleolar RNAs and rRNAs requires common components. 1998 , 18, 1181-9		179
359	Granulocyte-Macrophage Colony-stimulating Factor. 1998 , 261-275		
358	Mutations in <i>VPS16</i> and <i>MRT1</i> stabilize mRNAs by activating an inhibitor of the decapping enzyme. 1999 , 19, 7568-76		31
357	Aberrant mRNAs with extended 3'UTRs are substrates for rapid degradation by mRNA surveillance. <i>Rna</i> , 1999 , 5, 1299-307	5.8	165
356	The cis acting sequences responsible for the differential decay of the unstable <i>MFA2</i> and stable <i>PGK1</i> transcripts in yeast include the context of the translational start codon. <i>Rna</i> , 1999 , 5, 420-33	5.8	67
355	Overexpression of truncated Nmd3p inhibits protein synthesis in yeast. <i>Rna</i> , 1999 , 5, 1055-70	5.8	25
354	Identification of cis-acting RNA leader elements required for chloroplast <i>psbD</i> gene expression in <i>Chlamydomonas</i> . 1999 , 11, 957-70		112

353	Mechanisms of mRNA surveillance in eukaryotes. 1999 , 33, 229-60	180
352	Recognition of yeast mRNAs as "nonsense containing" leads to both inhibition of mRNA translation and mRNA degradation: implications for the control of mRNA decapping. 1999 , 10, 3971-8	75
351	5' to 3' exoribonucleolytic activity is a normal component of chloroplast mRNA decay pathways. 1999 , 19, 521-31	60
350	The virion host shutoff function of herpes simplex virus degrades the 5' end of a target mRNA before the 3' end. 1999 , 264, 195-204	66
349	Decapping of stabilized, polyadenylated mRNA in yeast pab1 mutants. 1999 , 15, 687-702	32
348	Analysis of mRNA decay pathways in <i>Saccharomyces cerevisiae</i> . 1999 , 17, 3-10	8
347	Monitoring mRNA decapping activity. 1999 , 17, 46-51	49
346	Assays for analyzing exonucleases in vitro. 1999 , 17, 52-9	7
345	SMG-2 is a phosphorylated protein required for mRNA surveillance in <i>Caenorhabditis elegans</i> and related to Upf1p of yeast. 1999 , 19, 5943-51	187
344	An mRNA stability complex functions with poly(A)-binding protein to stabilize mRNA in vitro. 1999 , 19, 4552-60	206
343	Assembly of the alpha-globin mRNA stability complex reflects binary interaction between the pyrimidine-rich 3' untranslated region determinant and poly(C) binding protein alphaCP. 1999 , 19, 4572-81	115
342	Mutations in translation initiation factors lead to increased rates of deadenylation and decapping of mRNAs in <i>Saccharomyces cerevisiae</i> . 1999 , 19, 5247-56	202
341	Active-site mutations in the Xrn1p exoribonuclease of <i>Saccharomyces cerevisiae</i> reveal a specific role in meiosis. 1999 , 19, 5930-42	48
340	Poly(A)-binding protein I of <i>Leishmania</i> : functional analysis and localisation in trypanosomatid parasites. <i>Nucleic Acids Research</i> , 2000 , 28, 1211-20	20.1 32
339	Stability determinants in the chloroplast psbB/T/H mRNAs of <i>Chlamydomonas reinhardtii</i> . 2000 , 21, 469-82	85
338	Yeast Sm-like proteins function in mRNA decapping and decay. 2000 , 404, 515-8	340
337	Functions of Lsm proteins in mRNA degradation and splicing. 2000 , 12, 346-50	137
336	Size distribution of the urokinase mRNA decay intermediates in different tissues and cell lines. 2000 , 1517, 33-45	3

335	The Puf3 protein is a transcript-specific regulator of mRNA degradation in yeast. <i>EMBO Journal</i> , 2000 , 19, 6602-11	13	209
334	Cap-dependent deadenylation of mRNA. <i>EMBO Journal</i> , 2000 , 19, 1079-86	13	152
333	Upf1p control of nonsense mRNA translation is regulated by Nmd2p and Upf3p. 2000 , 20, 4591-603		93
332	Exoribonucleases and their multiple roles in RNA metabolism. 2001 , 66, 67-105		108
331	mRNA decapping in yeast requires dissociation of the cap binding protein, eukaryotic translation initiation factor 4E. 2000 , 20, 7933-42		143
330	<i>Saccharomyces cerevisiae</i> RAI1 (YGL246c) is homologous to human DOM3Z and encodes a protein that binds the nuclear exoribonuclease Rat1p. 2000 , 20, 4006-15		101
329	A nuclear 3'5' exonuclease involved in mRNA degradation interacts with Poly(A) polymerase and the hnRNA protein Npl3p. 2000 , 20, 604-16		179
328	Novel features of the XRN-family in Arabidopsis: evidence that AtXRN4, one of several orthologs of nuclear Xrn2p/Rat1p, functions in the cytoplasm. 2000 , 97, 13985-90		138
327	Deletion of the PAT1 gene affects translation initiation and suppresses a PAB1 gene deletion in yeast. 2000 , 20, 3538-49		45
326	Emerging regulatory mechanisms for fibrinolytic gene expression. 2000 , 14, 146-154		4
325	Interaction between a poly(A)-specific ribonuclease and the 5' cap influences mRNA deadenylation rates in vitro. 2000 , 5, 479-88		154
324	Identification of a regulated pathway for nuclear pre-mRNA turnover. 2000 , 102, 765-75		318
323	Mechanisms and control of mRNA decapping in <i>Saccharomyces cerevisiae</i> . 2000 , 69, 571-95		129
322	Nonsense-mediated mRNA decay in <i>Saccharomyces cerevisiae</i> . 2001 , 274, 15-25		113
321	Regulated ARE-mediated mRNA decay in <i>Saccharomyces cerevisiae</i> . 2001 , 7, 1191-200		106
320	New twists in understanding the fate of antisense oligodeoxynucleotide mRNA targets. 2001 , 8, 732-3		2
319	Targeting an mRNA for decapping: displacement of translation factors and association of the Lsm1p-7p complex on deadenylated yeast mRNAs. 2001 , 8, 1075-83		191
318	The transcription factor associated Ccr4 and Caf1 proteins are components of the major cytoplasmic mRNA deadenylase in <i>Saccharomyces cerevisiae</i> . 2001 , 104, 377-86		475

317	Functional link between the mammalian exosome and mRNA decapping. 2001 , 107, 751-62		212
316	The mechanism and regulation of deadenylation: identification and characterization of <i>Xenopus</i> PARN. <i>Rna</i> , 2001 , 7, 875-86	5.8	86
315	Full-length cDNAs: more than just reaching the ends. 2001 , 6, 57-80		47
314	Computational modeling of eukaryotic mRNA turnover. <i>Rna</i> , 2001 , 7, 1192-212	5.8	104
313	The DEAD box helicase, Dhh1p, functions in mRNA decapping and interacts with both the decapping and deadenylase complexes. <i>Rna</i> , 2001 , 7, 1717-27	5.8	268
312	Yeast mRNA decapping enzyme. 2001 , 342, 226-33		8
311	Analysis of XRN orthologs by complementation of yeast mutants and localization of XRN-GFP fusion proteins. 2001 , 342, 269-82		10
310	5'→3' exoribonuclease from rabbit reticulocytes. 2001 , 342, 282-92		
309	<i>Drosophila</i> 5'→3' exoribonuclease Pacman. 2001 , 342, 293-302		10
308	Ski7p G protein interacts with the exosome and the Ski complex for 3' to 5' mRNA decay in yeast. <i>EMBO Journal</i> , 2001 , 20, 4684-93	13	168
307	A novel mRNA-decapping activity in HeLa cytoplasmic extracts is regulated by AU-rich elements. <i>EMBO Journal</i> , 2001 , 20, 1134-43	13	130
306	The <i>dhp1(+)</i> gene, encoding a putative nuclear 5'→3' exoribonuclease, is required for proper chromosome segregation in fission yeast. <i>Nucleic Acids Research</i> , 2001 , 29, 1326-33	20.1	32
305	COL5A1 exon 14 splice acceptor mutation causes a functional null allele, haploinsufficiency of alpha 1(V) and abnormal heterotypic interstitial fibrils in Ehlers-Danlos syndrome II. 2001 , 276, 13356-64		27
304	5' exoribonuclease 1: Xrn1. 2001 , 342, 251-9		36
303	Upf1p, Nmd2p, and Upf3p regulate the decapping and exonucleolytic degradation of both nonsense-containing mRNAs and wild-type mRNAs. 2001 , 21, 1515-30		81
302	Diazaborine treatment of Baker's yeast results in stabilization of aberrant mRNAs. 2001 , 276, 36419-24		13
301	Characterization of a general stabilizer element that blocks deadenylation-dependent mRNA decay. 2001 , 276, 30995-1003		16
300	Absence of Dbp2p alters both nonsense-mediated mRNA decay and rRNA processing. 2001 , 21, 7366-79		76

299	Degradation of the unstable EP1 mRNA in <i>Trypanosoma brucei</i> involves initial destruction of the 3' untranslated region. <i>Nucleic Acids Research</i> , 2001 , 29, 4707-15	20.1	35
298	The yeast POP2 gene encodes a nuclease involved in mRNA deadenylation. <i>Nucleic Acids Research</i> , 2001 , 29, 2448-55	20.1	164
297	The target of rapamycin signaling pathway regulates mRNA turnover in the yeast <i>Saccharomyces cerevisiae</i> . 2001 , 12, 3428-38		49
296	The mRNA cap structure stimulates rate of poly(A) removal and amplifies processivity of degradation. 2001 , 276, 27923-9		76
295	mRNA decay enzymes: decappers conserved between yeast and mammals. 2002 , 99, 12512-4		25
294	Evidence for a stabilizer element in the untranslated regions of <i>Drosophila glutathione S-transferase D1</i> mRNA. 2002 , 277, 34700-7		6
293	The hDcp2 protein is a mammalian mRNA decapping enzyme. 2002 , 99, 12663-8		266
292	mRNA decay is rapidly induced after spore germination of <i>Saccharomyces cerevisiae</i> . 2002 , 277, 40505-12		25
291	Exosome-mediated recognition and degradation of mRNAs lacking a termination codon. 2002 , 295, 2262-4		430
290	Temperature-sensitive mutants of the exosome subunit Rrp43p show a deficiency in mRNA degradation and no longer interact with the exosome. <i>Nucleic Acids Research</i> , 2002 , 30, 4186-98	20.1	27
289	Regulation of pathways of mRNA destabilization and stabilization. 2002 , 72, 129-64		53
288	Analyzing mRNA decay in <i>Saccharomyces cerevisiae</i> . 2002 , 351, 648-60		17
287	An mRNA surveillance mechanism that eliminates transcripts lacking termination codons. 2002 , 295, 2258-61		421
286	The mammalian exosome mediates the efficient degradation of mRNAs that contain AU-rich elements. <i>EMBO Journal</i> , 2002 , 21, 165-74	13	288
285	Processing of 3' extended read-through transcripts by the exosome can generate functional mRNAs. 2002 , 9, 1285-96		138
284	mRNA degradation machines in eukaryotic cells. 2002 , 84, 821-37		111
283	Analysis of the products of mRNA decapping and 3' to 5' decay by denaturing gel electrophoresis. <i>Rna</i> , 2002 , 8, 959-65	5.8	8
282	Replacement of the yeast TRP4 3' untranslated region by a hammerhead ribozyme results in a stable and efficiently exported mRNA that lacks a poly(A) tail. <i>Rna</i> , 2002 , 8, 336-44	5.8	16

281	Messenger RNA degradation: beginning at the end. 2002 , 12, R285-7		70
280	Non-stop decay--a new mRNA surveillance pathway. 2002 , 24, 785-8		81
279	In vivo evidence for 5' to 3' exonuclease degradation of an unstable chloroplast mRNA. 1998 , 13, 85-96		101
278	KEM1 is involved in filamentous growth of <i>Saccharomyces cerevisiae</i> . 2002 , 216, 33-8		14
277	<i>Arabidopsis thaliana</i> atrab28: a nuclear targeted protein related to germination and toxic cation tolerance. 2002 , 50, 249-59		38
276	MRNA stability and the control of gene expression: implications for human disease. 2002 , 27, 957-80		156
275	The DEAD box protein Dhh1 stimulates the decapping enzyme Dcp1. <i>EMBO Journal</i> , 2002 , 21, 2788-97	13	137
274	Ccr4p is the catalytic subunit of a Ccr4p/Pop2p/Notp mRNA deadenylase complex in <i>Saccharomyces cerevisiae</i> . <i>EMBO Journal</i> , 2002 , 21, 1427-36	13	264
273	SMG-5, required for <i>C.elegans</i> nonsense-mediated mRNA decay, associates with SMG-2 and protein phosphatase 2A. <i>EMBO Journal</i> , 2003 , 22, 641-50	13	107
272	Glucocorticoid receptor-JNK interaction mediates inhibition of the JNK pathway by glucocorticoids. <i>EMBO Journal</i> , 2003 , 22, 6035-44	13	90
271	The CCR4-NOT complex plays diverse roles in mRNA metabolism. 2003 , 73, 221-50		122
270	Interaction between Ski7p and Upf1p is required for nonsense-mediated 3' to 5' mRNA decay in yeast. <i>EMBO Journal</i> , 2003 , 22, 3951-9	13	59
269	Possible role of 3' to 5' phosphoadenosine-5' phosphate phosphatase in the etiology and therapy of bipolar disorder. 2003 , 27, 723-7		7
268	Identification of human SEP1 as a glial cell line-derived neurotrophic factor-inducible protein and its expression in the nervous system. 2003 , 121, 899-906		2
267	Lsm Proteins are required for normal processing and stability of ribosomal RNAs. 2003 , 278, 2147-56		63
266	A 3' exonuclease that specifically interacts with the 3' end of histone mRNA. 2003 , 12, 295-305		97
265	Nonsense-mediated mRNA decay in mammalian cells involves decapping, deadenylating, and exonucleolytic activities. 2003 , 12, 675-87		290
264	Genome-wide analysis of mRNAs regulated by the nonsense-mediated and 5' to 3' mRNA decay pathways in yeast. 2003 , 12, 1439-52		331

263	Cytoplasmic degradation of splice-defective pre-mRNAs and intermediates. 2003 , 12, 1453-65		92
262	Decapping and decay of messenger RNA occur in cytoplasmic processing bodies. 2003 , 300, 805-8		987
261	Functional characterization of the mammalian mRNA decapping enzyme hDcp2. <i>Rna</i> , 2003 , 9, 1138-47	5.8	97
260	Nonsense-containing mRNAs that accumulate in the absence of a functional nonsense-mediated mRNA decay pathway are destabilized rapidly upon its restitution. 2003 , 23, 842-51		62
259	Analysis of recombinant yeast decapping enzyme. <i>Rna</i> , 2003 , 9, 231-8	5.8	135
258	Launching the yeast 23S RNA Narnavirus shows 5Qand 3Qcis-acting signals for replication. 2003 , 100, 2568-73		34
257	p38 Mitogen-activated protein kinase stabilizes mRNAs that contain cyclooxygenase-2 and tumor necrosis factor AU-rich elements by inhibiting deadenylation. 2003 , 278, 39470-6		82
256	A yeast homologue of Hsp70, Ssa1p, regulates turnover of the MFA2 transcript through its AU-rich 3Quntranslated region. 2003 , 23, 2623-32		28
255	The enhancer of decapping proteins, Edc1p and Edc2p, bind RNA and stimulate the activity of the decapping enzyme. <i>Rna</i> , 2003 , 9, 239-51	5.8	63
254	Premature termination codons enhance mRNA decapping in human cells. <i>Nucleic Acids Research</i> , 2004 , 32, 488-94	20.1	57
253	Positive and negative regulation of poly(A) nuclease. 2004 , 24, 5521-33		69
252	Identification of Edc3p as an enhancer of mRNA decapping in <i>Saccharomyces cerevisiae</i> . 2004 , 166, 729-39		122
251	Inhibition of 5Qto 3QmRNA degradation under stress conditions in <i>Saccharomyces cerevisiae</i> : from GCN4 to MET16. <i>Rna</i> , 2004 , 10, 458-68	5.8	16
250	Recruitment of the Puf3 protein to its mRNA target for regulation of mRNA decay in yeast. <i>Rna</i> , 2004 , 10, 1625-36	5.8	58
249	Crystal structure of Dcp1p and its functional implications in mRNA decapping. <i>Nature Structural and Molecular Biology</i> , 2004 , 11, 249-56	17.6	71
248	Bringing the role of mRNA decay in the control of gene expression into focus. 2004 , 20, 491-7		226
247	The association of Alu repeats with the generation of potential AU-rich elements (ARE) at 3Q untranslated regions. 2004 , 5, 97		21
246	Messenger RNA decay in mammalian cells: the exonuclease perspective. 2004 , 41, 265-78		23

245	Messenger RNA turnover in eukaryotes: pathways and enzymes. 2004 , 39, 197-216		271
244	Eukaryotic mRNA decapping. 2004 , 73, 861-90		397
243	AtPARN is an essential poly(A) ribonuclease in Arabidopsis. 2004 , 328, 95-102		58
242	Posttranscriptional regulation of the karyogamy gene by Kem1p/Xrn1p exoribonuclease and Rok1p RNA helicase of <i>Saccharomyces cerevisiae</i> . 2004 , 321, 1032-9		8
241	AtXRN4 degrades mRNA in Arabidopsis and its substrates include selected miRNA targets. 2004 , 15, 173-83		295
240	Nuclear pre-mRNA decapping and 5' degradation in yeast require the Lsm2-8p complex. 2004 , 24, 9646-57		53
239	Process or perish: quality control in mRNA biogenesis. <i>Nature Structural and Molecular Biology</i> , 2005 , 12, 482-8	17.6	104
238	The yeast EDC1 mRNA undergoes deadenylation-independent decapping stimulated by Not2p, Not4p, and Not5p. <i>EMBO Journal</i> , 2005 , 24, 1033-45	13	72
237	RNA decapping inside and outside of processing bodies. 2005 , 17, 326-31		60
236	Folding free energies of 5'UTRs impact post-transcriptional regulation on a genomic scale in yeast. 2005 , 1, e72		97
235	Context-dependent codon bias and messenger RNA longevity in the yeast transcriptome. 2005 , 22, 1403-11		31
234	<i>Caenorhabditis elegans</i> decapping proteins: localization and functional analysis of Dcp1, Dcp2, and Dcp5 during embryogenesis. 2005 , 16, 5880-90		55
233	Mutations in the <i>Saccharomyces cerevisiae</i> LSM1 gene that affect mRNA decapping and 3' end protection. 2005 , 170, 33-46		49
232	p38 mitogen-activated protein kinase/Hog1p regulates translation of the AU-rich-element-bearing MFA2 transcript. 2005 , 25, 9753-63		21
231	Regulation of protease and protease inhibitor gene expression: the role of the 3'UTR and lessons from the plasminogen activating system. 2005 , 80, 169-215		
230	Conservation of the deadenylase activity of proteins of the Caf1 family in human. <i>Rna</i> , 2005 , 11, 487-94	5.8	77
229	Multiple processing body factors and the ARE binding protein TTP activate mRNA decapping. 2005 , 20, 905-15		365
228	General translational repression by activators of mRNA decapping. 2005 , 122, 875-86		489

227	Differential inhibition of mRNA degradation pathways by novel cap analogs. 2006 , 281, 1857-67		62
226	Control of mRNA stability in eukaryotes. 2006 , 34, 30-34		88
225	ARE-mRNA degradation requires the 5' decay pathway. 2006 , 7, 72-7		188
224	Sodium-induced GCN4 expression controls the accumulation of the 5' to 3' RNA degradation inhibitor, 3' phosphoadenosine 5' phosphate. 2006 , 281, 3276-82		15
223	Cell and molecular biology of the exosome: how to make or break an RNA. 2006 , 251, 159-208		28
222	Translation-independent inhibition of mRNA deadenylation during stress in <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 2006 , 12, 1835-45	5.8	69
221	Tpa1p is part of an mRNP complex that influences translation termination, mRNA deadenylation, and mRNA turnover in <i>Saccharomyces cerevisiae</i> . 2006 , 26, 5237-48		43
220	Roles of a <i>Trypanosoma brucei</i> 5' to 3' exoribonuclease homolog in mRNA degradation. <i>Rna</i> , 2006 , 12, 2171-86	5.8	63
219	Sbp1p affects translational repression and decapping in <i>Saccharomyces cerevisiae</i> . 2006 , 26, 5120-30		49
218	Yeast transcripts cleaved by an internal ribozyme provide new insight into the role of the cap and poly(A) tail in translation and mRNA decay. <i>Rna</i> , 2006 , 12, 1323-37	5.8	79
217	PAB1 self-association precludes its binding to poly(A), thereby accelerating CCR4 deadenylation in vivo. 2007 , 27, 6243-53		33
216	Microtubule disruption stimulates P-body formation. <i>Rna</i> , 2007 , 13, 493-502	5.8	64
215	Identification of functional domains in <i>Arabidopsis thaliana</i> mRNA decapping enzyme (AtDcp2). <i>Nucleic Acids Research</i> , 2008 , 36, 203-16	20.1	24
214	Accumulation of polyadenylated mRNA, Pab1p, eIF4E, and eIF4G with P-bodies in <i>Saccharomyces cerevisiae</i> . 2007 , 18, 2592-602		117
213	A+U-rich instability elements differentially activate 5' and 3' mRNA decay. 2007 , 27, 2791-9		51
212	A specific role for the C-terminal region of the Poly(A)-binding protein in mRNA decay. <i>Nucleic Acids Research</i> , 2007 , 35, 6017-28	20.1	43
211	The decapping activator Lsm1p-7p-Pat1p complex has the intrinsic ability to distinguish between oligoadenylated and polyadenylated RNAs. <i>Rna</i> , 2007 , 13, 998-1016	5.8	111
210	Degradation of hsp70 and other mRNAs in <i>Drosophila</i> via the 5' decay pathway and its regulation by heat shock. 2007 , 282, 21818-28		34

209	5Cto-3Cexoribonuclease activity in bacteria: role of RNase J1 in rRNA maturation and 5Cstability of mRNA. 2007 , 129, 681-92		271
208	Initiation of RNA decay in Escherichia coli by 5Cpyrophosphate removal. 2007 , 27, 79-90		200
207	P bodies: at the crossroads of post-transcriptional pathways. 2007 , 8, 9-22		751
206	A single subunit, Dis3, is essentially responsible for yeast exosome core activity. <i>Nature Structural and Molecular Biology</i> , 2007 , 14, 15-22	17.6	346
205	Regulation and function of maternal mRNA destabilization during early Drosophila development. 2007 , 75, 482-506		18
204	cDNA-AFLP analysis reveals differential gene expression in response to salt stress in foxtail millet (<i>Setaria italica</i> L.). 2008 , 40, 241-51		61
203	The bacterial enzyme RppH triggers messenger RNA degradation by 5Cpyrophosphate removal. 2008 , 451, 355-8		322
202	Structural insights into the dual activity of RNase J. <i>Nature Structural and Molecular Biology</i> , 2008 , 15, 206-12	17.6	142
201	The 5Cto-3Cexoribonuclease pacman is required for epithelial sheet sealing in Drosophila and genetically interacts with the phosphatase puckered. 2008 , 100, 687-701		18
200	Roles of eukaryotic Lsm proteins in the regulation of mRNA function. 2009 , 272, 149-89		69
199	Methods to determine mRNA half-life in <i>Saccharomyces cerevisiae</i> . 2008 , 448, 267-84		25
198	The control of mRNA decapping and P-body formation. 2008 , 32, 605-15		313
197	Diverse aberrancies target yeast mRNAs to cytoplasmic mRNA surveillance pathways. 2008 , 1779, 550-7		7
196	Messenger RNA half-life measurements in mammalian cells. 2008 , 448, 335-57		145
195	Pat1 contains distinct functional domains that promote P-body assembly and activation of decapping. 2008 , 28, 1298-312		91
194	A novel class of mRNA-containing cytoplasmic granules are produced in response to UV-irradiation. 2008 , 19, 4980-92		27
193	The 3Cuntranslated region of sindbis virus represses deadenylation of viral transcripts in mosquito and Mammalian cells. 2008 , 82, 880-92		62
192	Upf1 potentially serves as a RING-related E3 ubiquitin ligase via its association with Upf3 in yeast. <i>Rna</i> , 2008 , 14, 1950-8	5.8	45

191	NMD: multitasking between mRNA surveillance and modulation of gene expression. 2008 , 62, 185-243		59
190	Novel low abundance and transient RNAs in yeast revealed by tiling microarrays and ultra high-throughput sequencing are not conserved across closely related yeast species. 2008 , 4, e1000299		29
189	lsm1 mutations impairing the ability of the Lsm1p-7p-Pat1p complex to preferentially bind to oligoadenylated RNA affect mRNA decay in vivo. <i>Rna</i> , 2008 , 14, 2149-58	5.8	19
188	Analysis of cytoplasmic mRNA decay in <i>Saccharomyces cerevisiae</i> . 2008 , 448, 409-27		15
187	<i>S. cerevisiae</i> Vts1p induces deadenylation-dependent transcript degradation and interacts with the Ccr4p-Pop2p-Not deadenylase complex. <i>Rna</i> , 2008 , 14, 1328-36	5.8	30
186	The DEAD-box RNA helicase Ded1p affects and accumulates in <i>Saccharomyces cerevisiae</i> P-bodies. 2008 , 19, 984-93		79
185	A kinetic assay to monitor RNA decapping under single- turnover conditions. 2008 , 448, 23-40		19
184	A role for Lsm1p in response to ultraviolet-radiation damage in <i>Saccharomyces cerevisiae</i> . 2008 , 170, 411-21		3
183	The 5' 3' exonuclease Pacman is required for normal male fertility and is dynamically localized in cytoplasmic particles in <i>Drosophila</i> testis cells. 2008 , 416, 327-35		25
182	The role of deadenylation in the degradation of unstable mRNAs in trypanosomes. <i>Nucleic Acids Research</i> , 2009 , 37, 5511-28	20.1	50
181	Stm1 modulates mRNA decay and Dhh1 function in <i>Saccharomyces cerevisiae</i> . 2009 , 181, 93-103		25
180	Cordycepin interferes with 3' end formation in yeast independently of its potential to terminate RNA chain elongation. <i>Rna</i> , 2009 , 15, 837-49	5.8	39
179	Activation of decapping involves binding of the mRNA and facilitation of the post-binding steps by the Lsm1-7-Pat1 complex. <i>Rna</i> , 2009 , 15, 1837-48	5.8	25
178	Polysomes, P bodies and stress granules: states and fates of eukaryotic mRNAs. 2009 , 21, 403-8		304
177	Decapping is preceded by 3' uridylation in a novel pathway of bulk mRNA turnover. <i>Nature Structural and Molecular Biology</i> , 2009 , 16, 616-23	17.6	163
176	The relationship of prions and translation. <i>Wiley Interdisciplinary Reviews RNA</i> , 2010 , 1, 81-9	9.3	8
175	Distinct roles for Caf1, Ccr4, Edc3 and CutA in the co-ordination of transcript deadenylation, decapping and P-body formation in <i>Aspergillus nidulans</i> . 2010 , 76, 503-16		24
174	All things must pass: contrasts and commonalities in eukaryotic and bacterial mRNA decay. 2010 , 11, 467-78		135

173	Regulation of mRNA Turnover. 2010 , 2311-2315		1
172	Human Pat1b connects deadenylation with mRNA decapping and controls the assembly of processing bodies. 2010 , 30, 4308-23		104
171	Dcp2 phosphorylation by Ste20 modulates stress granule assembly and mRNA decay in <i>Saccharomyces cerevisiae</i> . 2010 , 189, 813-27		71
170	Comparison of U2OS and Huh-7 cells for identifying host factors that affect hepatitis C virus RNA replication. 2010 , 91, 2238-48		28
169	CUCU modification of mRNA promotes decapping and transcript degradation in <i>Aspergillus nidulans</i> . 2010 , 30, 460-9		44
168	An RNA pseudoknot is required for production of yellow fever virus subgenomic RNA by the host nuclease XRN1. 2010 , 84, 11395-406		129
167	PUF3 acceleration of deadenylation in vivo can operate independently of CCR4 activity, possibly involving effects on the PAB1-mRNP structure. 2010 , 399, 562-75		36
166	An RNA pyrophosphohydrolase triggers 5' exonucleolytic degradation of mRNA in <i>Bacillus subtilis</i> . 2011 , 43, 940-9		89
165	A brief survey of mRNA surveillance. 2011 , 36, 585-92		29
164	Mechanisms of deadenylation-dependent decay. <i>Wiley Interdisciplinary Reviews RNA</i> , 2011 , 2, 167-83	9.3	204
163	Structural and functional insights into eukaryotic mRNA decapping. <i>Wiley Interdisciplinary Reviews RNA</i> , 2011 , 2, 193-208	9.3	37
162	Roles of Puf proteins in mRNA degradation and translation. <i>Wiley Interdisciplinary Reviews RNA</i> , 2011 , 2, 471-92	9.3	111
161	The role of the 5' exoribonuclease XRN1 in transcriptome-wide mRNA degradation. <i>Rna</i> , 2011 , 17, 2039-47	5.8	45
160	Processive RNA decay by the exosome: merits of a quantitative Bayesian sampling approach. 2011 , 8, 55-60		6
159	A quantitative assay for measuring mRNA decapping by splinted ligation reverse transcription polymerase chain reaction: qSL-RT-PCR. <i>Rna</i> , 2011 , 17, 535-43	5.8	19
158	Evidence that XRN4, an Arabidopsis homolog of exoribonuclease XRN1, preferentially impacts transcripts with certain sequences or in particular functional categories. <i>Rna</i> , 2011 , 17, 501-11	5.8	60
157	Masking the 5' terminal nucleotides of the hepatitis C virus genome by an unconventional microRNA-target RNA complex. 2011 , 108, 3193-8		230
156	Different nuclease requirements for exosome-mediated degradation of normal and nonstop mRNAs. 2011 , 108, 2366-71		58

155	Coordinated destruction of cellular messages in translation complexes by the gammaherpesvirus host shutoff factor and the mammalian exonuclease Xrn1. 2011 , 7, e1002339		65
154	Cytoplasmic mRNA 3' tagging in eukaryotes: does it spell the end?. 2012 , 40, 810-4		7
153	XRN1: A Major 5' to 3' Exoribonuclease in Eukaryotic Cells. 2012 , 31, 97-114		11
152	Biochemistry and Function of RNA Exosomes. 2012 , 31, 1-30		3
151	A noncoding RNA produced by arthropod-borne flaviviruses inhibits the cellular exoribonuclease XRN1 and alters host mRNA stability. <i>Rna</i> , 2012 , 18, 2029-40	5.8	150
150	Global analysis of mRNA decay intermediates in <i>Saccharomyces cerevisiae</i> . 2012 , 109, 11764-9		41
149	A direct interaction between DCP1 and XRN1 couples mRNA decapping to 5' exonucleolytic degradation. <i>Nature Structural and Molecular Biology</i> , 2012 , 19, 1324-31	17.6	111
148	mRNA decapping factors and the exonuclease Xrn2 function in widespread premature termination of RNA polymerase II transcription. 2012 , 46, 311-24		145
147	Ways and means of eukaryotic mRNA decay. 2012 , 1819, 593-603		50
146	P-bodies and stress granules: possible roles in the control of translation and mRNA degradation. 2012 , 4, a012286		433
145	RGG motif proteins: modulators of mRNA functional states. 2012 , 11, 2594-9		63
144	Mass spectrometric identification of proteins that interact through specific domains of the poly(A) binding protein. 2012 , 287, 711-730		18
143	mRNA 3' tagging is induced by nonsense-mediated decay and promotes ribosome dissociation. 2012 , 32, 2585-95		34
142	The eIF4E-binding protein Eap1p functions in Vts1p-mediated transcript decay. <i>PLoS ONE</i> , 2012 , 7, e47131	13.7	13
141	Rat1 and Xrn2: The Diverse Functions of the Nuclear Rat1/Xrn2 Exonuclease. 2012 , 31, 131-63		3
140	Degradation of mRNAs that lack a stop codon: a decade of nonstop progress. <i>Wiley Interdisciplinary Reviews RNA</i> , 2012 , 3, 649-60	9.3	96
139	RNA degradation in <i>Saccharomyces cerevisiae</i> . 2012 , 191, 671-702		260
138	The RRM1 domain of the poly(A)-binding protein from <i>Saccharomyces cerevisiae</i> is critical to control of mRNA deadenylation. 2013 , 288, 401-12		9

137	mRNAs containing the histone 3Cstem-loop are degraded primarily by decapping mediated by oligouridylation of the 3Cend. <i>Rna</i> , 2013 , 19, 1-16	5.8	41
136	Global analysis of eukaryotic mRNA degradation reveals Xrn1-dependent buffering of transcript levels. 2013 , 52, 52-62		152
135	The discovery and analysis of P Bodies. 2013 , 768, 23-43		67
134	Eukaryotic mRNA decay: methodologies, pathways, and links to other stages of gene expression. 2013 , 425, 3750-75		99
133	XRN 5C3Cexoribonucleases: structure, mechanisms and functions. 2013 , 1829, 590-603		210
132	Exonuclease hDIS3L2 specifies an exosome-independent 3C5Cdegradation pathway of human cytoplasmic mRNA. <i>EMBO Journal</i> , 2013 , 32, 1855-68	13	113
131	Maternally recruited DCP1A and DCP2 contribute to messenger RNA degradation during oocyte maturation and genome activation in mouse. 2013 , 88, 11		65
130	A genome-wide RNAi screen reveals that mRNA decapping restricts bunyaviral replication by limiting the pools of Dcp2-accessible targets for cap-snatching. <i>Genes and Development</i> , 2013 , 27, 1511-25	12.6	68
129	mRNA Stability. 2013 , 1-26		
128	Cytoplasmic 5C3Cexonuclease Xrn1p is also a genome-wide transcription factor in yeast. <i>Frontiers in Genetics</i> , 2014 , 5, 1	4.5	257
127	Lsm2 and Lsm3 bridge the interaction of the Lsm1-7 complex with Pat1 for decapping activation. 2014 , 24, 233-46		35
126	Human DDX6 effects miRNA-mediated gene silencing via direct binding to CNOT1. <i>Rna</i> , 2014 , 20, 1398-409	4.9	91
125	RppH-dependent pyrophosphohydrolysis of mRNAs is regulated by direct interaction with DapF in Escherichia coli. <i>Nucleic Acids Research</i> , 2014 , 42, 12746-57	20.1	23
124	RNA-RNA interactions and pre-mRNA mislocalization as drivers of group II intron loss from nuclear genomes. 2014 , 111, 6612-7		17
123	The activation of the decapping enzyme DCP2 by DCP1 occurs on the EDC4 scaffold and involves a conserved loop in DCP1. <i>Nucleic Acids Research</i> , 2014 , 42, 5217-33	20.1	64
122	Quality control: Is there quality control of localized mRNAs?. 2014 , 204, 863-8		18
121	Cloning of Hsp21 gene and its expression in Chinese shrimp Fenneropenaeus chinensis in response to WSSV challenge. 2014 , 55, 231-8		11
120	Regulatory mechanisms of RNA function: emerging roles of DNA repair enzymes. 2014 , 71, 2451-65		25

119	Interrelations between translation and general mRNA degradation in yeast. <i>Wiley Interdisciplinary Reviews RNA</i> , 2014 , 5, 747-63	9.3	58
118	Widespread Co-translational RNA Decay Reveals Ribosome Dynamics. 2015 , 161, 1400-12		171
117	Role of cytoplasmic deadenylation and mRNA decay factors in yeast apoptosis. 2015 , 15,		8
116	General and MicroRNA-Mediated mRNA Degradation Occurs on Ribosome Complexes in Drosophila Cells. 2015 , 35, 2309-20		35
115	Interrogating the degradation pathways of unstable mRNAs with XRN1-resistant sequences. 2016 , 7, 13691		24
114	A conserved virus-induced cytoplasmic TRAMP-like complex recruits the exosome to target viral RNA for degradation. <i>Genes and Development</i> , 2016 , 30, 1658-70	12.6	35
113	The decapping activator Edc3 and the Q/N-rich domain of Lsm4 function together to enhance mRNA stability and alter mRNA decay pathway dependence in <i>Saccharomyces cerevisiae</i> . 2016 , 5, 1388-1399		5
112	The DEAD-Box Protein Dhh1p Couples mRNA Decay and Translation by Monitoring Codon Optimality. 2016 , 167, 122-132.e9		148
111	Global Analysis of Truncated RNA Ends Reveals New Insights into Ribosome Stalling in Plants. 2016 , 28, 2398-2416		68
110	mRNA decapping in 3D. <i>Nature Structural and Molecular Biology</i> , 2016 , 23, 954-956	17.6	5
109	RNA voyeurism: A coming of age story. 2016 , 98, 10-17		11
108	Standing your ground to exoribonucleases: Function of Flavivirus long non-coding RNAs. 2016 , 212, 70-7		25
107	When mRNA translation meets decay. 2017 , 45, 339-351		27
106	Spatiotemporal Uncoupling of MicroRNA-Mediated Translational Repression and Target RNA Degradation Controls MicroRNP Recycling in Mammalian Cells. 2017 , 37,		23
105	Translation initiation events on structured eukaryotic mRNAs generate gene expression noise. <i>Nucleic Acids Research</i> , 2017 , 45, 6981-6992	20.1	12
104	Small molecule induced poly(A) single strand to self-structure conformational switching: evidence for the prominent role of H-bonding interactions. 2017 , 13, 1000-1009		6
103	Trick or TREAT: A Scary-Good New Approach for Single-Molecule mRNA Decay Analysis. 2017 , 68, 476-477		4
102	The organization and regulation of mRNA-protein complexes. <i>Wiley Interdisciplinary Reviews RNA</i> , 2017 , 8, e1369	9.3	40

101	Simultaneous detection of mRNA transcription and decay intermediates by dual colour single mRNA FISH on subcellular resolution. <i>Nucleic Acids Research</i> , 2017 , 45, e49	20.1	15
100	Attacked from All Sides: RNA Decay in Antiviral Defense. 2017 , 9,		42
99	Impact of Methods on the Measurement of mRNA Turnover. 2017 , 18,		30
98	The ApaH-like phosphatase TbALPH1 is the major mRNA decapping enzyme of trypanosomes. 2017 , 13, e1006456		22
97	The influence of microRNAs and poly(A) tail length on endogenous mRNA-protein complexes. 2017 , 18, 211		28
96	Nuclear mRNA degradation tunes the gain of the unfolded protein response in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2018 , 46, 1139-1156	20.1	7
95	Identification of phlebovirus and arenavirus RNA sequences that stall and repress the exoribonuclease XRN1. 2018 , 293, 285-295		23
94	Revisiting the Closed-Loop Model and the Nature of mRNA 5'3' Communication. 2018 , 72, 805-812		48
93	The DEAD-box RNA-binding protein DDX6 regulates parental RNA decay for cellular reprogramming to pluripotency. <i>PLoS ONE</i> , 2018 , 13, e0203708	3.7	8
92	A Plant Biologist's Toolbox to Study Translation. 2018 , 9, 873		16
91	A folded viral noncoding RNA blocks host cell exoribonucleases through a conformationally dynamic RNA structure. 2018 , 115, 6404-6409		49
90	Spt6 Association with RNA Polymerase II Directs mRNA Turnover During Transcription. 2018 , 70, 1054-1066.e420		
89	RNA degradomes reveal substrates and importance for dark and nitrogen stress responses of <i>Arabidopsis</i> XRN4. <i>Nucleic Acids Research</i> , 2019 , 47, 9216-9230	20.1	16
88	Bacterial ribonucleases and their roles in RNA metabolism. 2019 , 54, 242-300		57
87	Cauliflower mosaic virus transactivator protein (TAV) can suppress nonsense-mediated decay by targeting VARICOSE, a scaffold protein of the decapping complex. 2019 , 9, 7042		12
86	Detection and quantification of RNA decay intermediates using XRN1-resistant reporter transcripts. 2019 , 14, 1603-1633		12
85	Negative Regulation of the Mis17-Mis6 Centromere Complex by mRNA Decay Pathway and EKC/KEOPS Complex in. 2019 , 9, 1815-1823		
84	Asymmetric distribution of glucose transporter mRNA provides a growth advantage in yeast. <i>EMBO Journal</i> , 2019 , 38,	13	3

83	Colocalization of mA and G-Quadruplex-Forming Sequences in Viral RNA (HIV, Zika, Hepatitis B, and SV40) Suggests Topological Control of Adenosine -Methylation. 2019 , 5, 218-228		26
82	A recap of RNA recapping. <i>Wiley Interdisciplinary Reviews RNA</i> , 2019 , 10, e1504	9.3	27
81	Dom34 mediates targeting of exogenous RNA in the antiviral OAS/RNase L pathway. <i>Nucleic Acids Research</i> , 2019 , 47, 432-449	20.1	16
80	Engineered viral RNA decay intermediates to assess XRN1-mediated decay. 2019 , 155, 116-123		3
79	RNA-Seq approach for accurate characterization of splicing efficiency of yeast introns. 2020 , 176, 25-33		5
78	The Dynamics of Cytoplasmic mRNA Metabolism. 2020 , 77, 786-799.e10		33
77	The landscape of eukaryotic mRNPs. <i>Rna</i> , 2020 , 26, 229-239	5.8	29
76	Suppressors of mRNA Decapping Defects Restore Growth Without Major Effects on mRNA Decay Rates or Abundance. 2020 , 216, 1051-1069		2
75	Codon and amino acid content are associated with mRNA stability in mammalian cells. <i>PLoS ONE</i> , 2020 , 15, e0228730	3.7	19
74	Tissue-specific changes in the RNA structurome mediate salinity response in. <i>Rna</i> , 2020 , 26, 492-511	5.8	12
73	Fated for decay: RNA elements targeted by viral endonucleases. 2021 , 111, 119-125		2
72	Genome-scale molecular principles of mRNA half-life regulation in yeast. 2021 , 288, 3428-3447		0
71	RNA Metabolism Guided by RNA Modifications: The Role of SMUG1 in rRNA Quality Control. 2021 , 11,		6
70	discovery of RNA proximal proteins via proximity-dependent biotinylation. 2021 , 18, 2203-2217		3
69	A (dis)integrated stress response: Genetic diseases of eIF2 β regulators. <i>Wiley Interdisciplinary Reviews RNA</i> , 2021 , e1689	9.3	2
68	Global view on the metabolism of RNA poly(A) tails in yeast <i>Saccharomyces cerevisiae</i> . 2021 , 12, 4951		4
67	Roles of mRNA poly(A) tails in regulation of eukaryotic gene expression. 2021 ,		14
66	Functions of the cytoplasmic exosome. 2011 , 702, 79-90		39

65	mRNA Degradation and Decay. 2014 , 159-193	3
64	Cytoplasmic fate of eukaryotic mRNA: identification and characterization of AU-binding proteins. 1997 , 18, 141-72	18
63	Cap-independent translation and internal initiation of translation in eukaryotic cellular mRNA molecules. 1995 , 203, 155-77	26
62	Control of mRNA stability in higher plants. 1996 , 63-77	1
61	Purification of a human polyribosome-associated 3' to 5' exoribonuclease.. 1994 , 269, 31814-31821	44
60	Regulation of long non-coding RNAs and genome dynamics by the RNA surveillance machinery. 2020 , 21, 123-136	72
59	RNA polymerase III defects suppress a conditional-lethal poly(A) polymerase mutation in <i>Saccharomyces cerevisiae</i> . 1996 , 143, 1149-61	11
58	Analysis of mutations in the yeast mRNA decapping enzyme. 1999 , 151, 1273-85	64
57	Temperature-sensitive mutations in the <i>Saccharomyces cerevisiae</i> MRT4, GRC5, SLA2 and THS1 genes result in defects in mRNA turnover. 1999 , 153, 35-47	49
56	Inhibition of mRNA turnover in yeast by an <i>xrn1</i> mutation enhances the requirement for eIF4E binding to eIF4G and for proper capping of transcripts by Ceg1p. 2000 , 155, 31-42	13
55	Two related proteins, Edc1p and Edc2p, stimulate mRNA decapping in <i>Saccharomyces cerevisiae</i> . 2001 , 157, 27-37	77
54	The yeast cytoplasmic Lsm1/Pat1p complex protects mRNA 3' termini from partial degradation. 2001 , 158, 1445-55	80
53	The yeast ubiquitin protease, Ubp3p, promotes protein stability. 2002 , 162, 1079-89	18
52	Identification of Edc3p as an Enhancer of mRNA Decapping in <i>Saccharomyces cerevisiae</i> . 2004 , 166, 729-739	17
51	In vivo discovery of RNA proximal proteins in human cells via proximity-dependent biotinylation.	4
50	Non-invasive measurement of mRNA decay reveals translation initiation as the major determinant of mRNA stability.	3
49	The Dynamics of Cytoplasmic mRNA Metabolism.	3
48	ELAV proteins stabilize deadenylated intermediates in a novel in vitro mRNA deadenylation/degradation system. <i>Genes and Development</i> , 1999 , 13, 188-201	12.6 211

47	Nonsense-mediated mRNA decay in <i>Saccharomyces cerevisiae</i> : a quality control mechanism that degrades transcripts harboring premature termination codons. 2001 , 66, 321-8		16
46	The virion host shutoff protein of herpes simplex virus type 1: messenger ribonucleolytic activity in vitro. 1996 , 70, 2411-9		87
45	The 3' untranslated region of the B19 parvovirus capsid protein mRNAs inhibits its own mRNA translation in nonpermissive cells. 1997 , 71, 9482-9		41
44	Selective degradation of early-response-gene mRNAs: functional analyses of sequence features of the AU-rich elements. 1994 , 14, 8471-8482		94
43	Posttranscriptional control of gene expression in yeast. 1998 , 62, 1492-553		173
42	mRNA stability in mammalian cells. <i>Microbiological Reviews</i> , 1995 , 59, 423-50		1083
41	Mechanisms and control of mRNA turnover in <i>Saccharomyces cerevisiae</i> . <i>Microbiological Reviews</i> , 1996 , 60, 233-49		221
40	Double-stranded RNA viruses of <i>Saccharomyces cerevisiae</i> . <i>Microbiological Reviews</i> , 1996 , 60, 250-65		206
39	A factor required for nonsense-mediated mRNA decay in yeast is exported from the nucleus to the cytoplasm by a nuclear export signal sequence. <i>Journal of Cell Science</i> , 1998 , 111, 3129-3143	5.3	47
38	Linking π to π diverse and dynamic RNA-based mechanisms to regulate gene expression by 5' to 3' communication. <i>F1000Research</i> , 2016 , 5,	3.6	3
37	Phosphorylation of mRNA decapping protein Dcp1a by the ERK signaling pathway during early differentiation of 3T3-L1 preadipocytes. <i>PLoS ONE</i> , 2013 , 8, e61697	3.7	18
36	Effects of the yeast RNA-binding protein Whi3 on the half-life and abundance of CLN3 mRNA and other targets. <i>PLoS ONE</i> , 2013 , 8, e84630	3.7	29
35	SPT5 affects the rate of mRNA degradation and physically interacts with CCR4 but does not control mRNA deadenylation. <i>American Journal of Molecular Biology</i> , 2012 , 02, 11-20	0.2	1
34	Non-invasive measurement of mRNA decay reveals translation initiation as the major determinant of mRNA stability. <i>ELife</i> , 2018 , 7,	8.9	79
33	Turnover of mRNA In Eukaryotic Cells. 2001 , 245-257		2
32	mRNA polyadenylation. 2. Formation of poly(A) tails in yeast, plant, prokaryote and virus mRNAs. <i>Biopolymers and Cell</i> , 2001 , 17, 185-202	0.3	1
31	Messenger RNA Ribonucleases and mRNA Turnover in <i>Saccharomyces cerevisiae</i> . 1997 , 589-620		
30	RNA-Processing RNases in Mammalian Cells. 1997 , 553-587		

29	mRNA Decay Machinery in Plants: Approaches and Potential Components. 1998 , 125-133		
28	Turnover of mRNA in Eukaryotic Cells. 1999 , 205-216		
27	Codon Usage and Amino Acid Identity Are Major Determinants of MRNA Stability in Humans. <i>SSRN Electronic Journal</i> ,	1	1
26	A folded viral noncoding RNA blocks host cell exoribonucleases through programmed remodeling of RNA structure.		
25	Asymmetric Distribution of Glucose Transporter mRNA Provides Growth Advantage.		
24	Tissue-specific changes in the RNA structurome mediate salinity response in Arabidopsis.		
23	Suppressors of mRNA decapping defects isolated by experimental evolution ameliorate transcriptome disruption without restoring mRNA decay.		
22	Genome-scale conserved molecular principles of mRNA half-life regulation.		
21	Identification of the cis-elements mediating the autogenous control of ribosomal protein L2 mRNA stability in yeast. <i>EMBO Journal</i> , 1995 , 14, 4022-30	13	14
20	Cytoplasmic 3' poly(A) addition induces 5' cap ribose methylation: implications for translational control of maternal mRNA. <i>EMBO Journal</i> , 1995 , 14, 6301-10	13	37
19	Overexpression of poly(A) binding protein prevents maturation-specific deadenylation and translational inactivation in <i>Xenopus</i> oocytes. <i>EMBO Journal</i> , 1996 , 15, 900-9	13	45
18	Interaction between Nmd2p and Upf1p is required for activity but not for dominant-negative inhibition of the nonsense-mediated mRNA decay pathway in yeast. <i>Rna</i> , 1996 , 2, 153-70	5.8	52
17	A cis-acting element known to block 3' mRNA degradation enhances expression of polyA-minus mRNA in wild-type yeast cells and phenocopies a ski mutant. <i>Rna</i> , 2001 , 7, 1566-77	5.8	19
16	Transcription feedback dynamics in the wake of cytoplasmic degradation shutdown.		
15	Distinct ribosome states trigger diverse mRNA quality control pathways.		0
14	Shaping the Innate Immune Response Through Post-Transcriptional Regulation of Gene Expression Mediated by RNA-Binding Proteins.. <i>Frontiers in Immunology</i> , 2021 , 12, 796012	8.4	1
13	The RNA binding protein Nab2 genetically interacts with multiple RNA exosome cofactors to regulate target RNAs.		
12	Eukaryotic mRNA Decapping Activation.. <i>Frontiers in Genetics</i> , 2022 , 13, 832547	4.5	1

11	PABLO-QA: A sensitive assay for quantifying monophosphorylated RNA 5' ends.. <i>STAR Protocols</i> , 2022 , 3, 101190	1.4	
10	Decoupling of degradation from deadenylation reshapes poly(A) tail length in yeast meiosis. <i>Nature Structural and Molecular Biology</i> , 2021 , 28, 1038-1049	17.6	1
9	OUP accepted manuscript. <i>Nucleic Acids Research</i> ,	20.1	1
8	RNA binding proteins Smaug and Cup induce CCR4-NOT-dependent deadenylation of the nanos mRNA in a reconstituted system.		0
7	Keeping development on time: Insights into post-transcriptional mechanisms driving oscillatory gene expression during vertebrate segmentation. <i>Wiley Interdisciplinary Reviews RNA</i> ,	9.3	
6	Distinct elongation stalls during translation are linked with distinct pathways for mRNA degradation. 11,		1
5	Eukaryotic mRNA decapping factors: molecular mechanisms and activity.		0
4	Deadenylation rate is not a major determinant of RNA degradation in yeast.		0
3	RNA binding proteins Smaug and Cup induce CCR4-NOT-dependent deadenylation of the nanos mRNA in a reconstituted system.		0
2	Arabidopsis mRNA decay landscape shaped by XRN 5'-3' exoribonucleases.		0
1	Regulation of eukaryotic mRNA deadenylation and degradation by the Ccr4-Not complex. 11,		0