

Long-term hippocampal slices: A model system for investigating pathologic processes

Journal of Neuroscience Research

42, 294-305

DOI: [10.1002/jnr.490420303](https://doi.org/10.1002/jnr.490420303)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Stable maintenance of glutamate receptors and other synaptic components in long-term hippocampal slices. <i>Hippocampus</i> , 1995, 5, 425-439.	0.9	86
2	Depolarization with high-K ⁺ causes Ca ²⁺ -independent but partially Cl ⁻ -dependent glutamate release in rat hippocampal slice cultures. <i>Neuroscience Research</i> , 1996, 25, 399-402.	1.0	8
3	Organotypic slice cultures: a technique has come of age. <i>Trends in Neurosciences</i> , 1997, 20, 471-477.	4.2	782
4	Sprouting and functional recovery in co-cultures between old and young hippocampal organotypic slices. <i>Neuroscience</i> , 1997, 80, 1127-1136.	1.1	24
5	Reactive Astrocytosis from Excitotoxic Injury in Hippocampal Organ Culture Parallels that Seen in Vivo. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 1997, 17, 26-43.	2.4	63
6	Microelectrode arrays for electrophysiological monitoring of hippocampal organotypic slice cultures. <i>IEEE Transactions on Biomedical Engineering</i> , 1997, 44, 1159-1163.	2.5	75
7	Correction to "A Bioelectric Inverse Imaging Technique Based On Surface Laplacians". <i>IEEE Transactions on Biomedical Engineering</i> , 1997, 44, 1163-1163.	2.5	0
8	Microdialysis monitoring of extracellular glutamate combined with the simultaneous recording of evoked field potentials in hippocampal organotypic slice cultures. <i>Journal of Neuroscience Methods</i> , 1997, 74, 65-76.	1.3	20
9	Trimethyltin (TMT) neurotoxicity in organotypic rat hippocampal slice cultures. <i>Brain Research</i> , 1998, 783, 305-315.	1.1	43
10	Amyloid β protein is internalized selectively by hippocampal field CA1 and causes neurons to accumulate amyloidogenic carboxyterminal fragments of the amyloid precursor protein. , 1998, 397, 139-147.		81
11	Lysosomal Dysfunction Reduces Brain-Derived Neurotrophic Factor Expression. <i>Experimental Neurology</i> , 1998, 150, 128-135.	2.0	15
12	Brain damage due to episodic alcohol exposure in vivo and in vitro: furosemide neuroprotection implicates edema-based mechanism. <i>FASEB Journal</i> , 1998, 12, 221-230.	0.2	96
13	Calcium Waves Precede Electrophysiological Changes of Spreading Depression in Hippocampal Organ Cultures. <i>Journal of Neuroscience</i> , 1998, 18, 3416-3425.	1.7	91
14	Neuroprotective interaction effects of NMDA and AMPA receptor antagonists in an in vitro model of cerebral ischemia. <i>Brain Research</i> , 1999, 816, 299-308.	1.1	47
15	Microglial-neuronal interactions in synaptic damage and recovery. <i>Journal of Neuroscience Research</i> , 1999, 58, 191-201.	1.3	153
16	Anatomical and functional reconstruction of the nigrostriatal system in vitro: Selective innervation of the striatum by dopaminergic neurons. , 1999, 58, 553-566.		16
17	Markers for neuronal degeneration in organotypic slice cultures. <i>Brain Research Protocols</i> , 1999, 3, 278-290.	1.7	266
18	Role of apolipoprotein E and estrogen in mossy fiber sprouting in hippocampal slice cultures. <i>Neuroscience</i> , 1999, 91, 1009-1016.	1.1	69

#	ARTICLE	IF	CITATIONS
19	Coupling of Organotypic Brain Slice Cultures to Silicon-Based Arrays of Electrodes. <i>Methods</i> , 1999, 18, 160-172.	1.9	35
20	Integrin-type signaling has a distinct influence on NMDA-induced cytoskeletal disassembly. , 2000, 59, 827-832.		27
21	Metabolic activity of cultured rat brainstem, hippocampal and spinal cord slices. <i>Journal of Neuroscience Methods</i> , 2000, 99, 1-7.	1.3	34
22	Excitatory amino acid neurotoxicity and modulation of glutamate receptor expression in organotypic brain slice cultures. <i>Amino Acids</i> , 2000, 19, 7-21.	1.2	73
23	Dynamics of NMDAR-mediated neurotoxicity during chronic ethanol exposure and withdrawal. <i>Neuropharmacology</i> , 2000, 39, 218-226.	2.0	43
24	Interface Organotypic Hippocampal Slice Cultures. , 2001, , 13-27.		10
25	Effects of age and insulin-like growth factor-1 on neuron and synapse numbers in area CA3 of hippocampus. <i>Neuroscience</i> , 2001, 107, 231-238.	1.1	45
26	Intracellular Deposition, Microtubule Destabilization, and Transport Failure: An "Early" Pathogenic Cascade Leading to Synaptic Decline. <i>Journal of Neuropathology and Experimental Neurology</i> , 2002, 61, 640-650.	0.9	59
27	Survival Signaling and Selective Neuroprotection Through Glutamatergic Transmission. <i>Experimental Neurology</i> , 2002, 174, 37-47.	2.0	65
28	Gas1 Is Induced during and Participates in Excitotoxic Neuronal Death. <i>Molecular and Cellular Neurosciences</i> , 2002, 19, 417-429.	1.0	39
29	Reciprocal Interactions Between Microglia and Neurons: From Survival to Neuropathology. <i>Reviews in the Neurosciences</i> , 2002, 13, 221-42.	1.4	188
30	Blockade of NMDA-receptors or calcium-channels attenuates the ischaemia-evoked efflux of glutamate and phosphoethanolamine and depression of neuronal activity in rat organotypic hippocampal slice cultures. <i>Comptes Rendus - Biologies</i> , 2002, 325, 495-504.	0.1	10
31	Spreading depression: a teleological means for self-protection from brain ischemia. , 2002, , 142-158.		7
32	Peptidyl γ -keto amide inhibitor of calpain blocks excitotoxic damage without affecting signal transduction events. <i>Journal of Neuroscience Research</i> , 2002, 67, 787-794.	1.3	29
33	Transforming growth factor γ 1 prevents IL-1 γ -induced microglial activation, whereas TNF γ - and IL-6-stimulated activation are not antagonized. <i>Glia</i> , 2002, 40, 109-120.	2.5	78
34	Novel Cathepsin D Inhibitors Block the Formation of Hyperphosphorylated Tau Fragments in Hippocampus. <i>Journal of Neurochemistry</i> , 2002, 74, 1469-1477.	2.1	67
35	The neuropathogenic contributions of lysosomal dysfunction. <i>Journal of Neurochemistry</i> , 2002, 83, 481-489.	2.1	181
36	Cytosolic Proteolysis of τ , by Cathepsin D in Hippocampus Following Suppression of Cathepsins B and L. <i>Journal of Neurochemistry</i> , 1996, 67, 1846-1855.	2.1	97

#	ARTICLE	IF	CITATIONS
37	Evaluation of drugs acting at glutamate transporters in organotypic hippocampal cultures: new evidence on substrates and blockers in excitotoxicity. <i>Neurochemical Research</i> , 2002, 27, 5-13.	1.6	26
38	Organotypic Slice Cultures from Transgenic Mice as Disease Model Systems. <i>Journal of Molecular Neuroscience</i> , 2002, 19, 317-320.	1.1	32
39	Brain Damage Results in Down-Regulation of N-Acetylaspartate as a Neuronal Osmolyte. <i>NeuroMolecular Medicine</i> , 2003, 3, 95-104.	1.8	32
40	Acetylcholinesterase activation in organotypic rat hippocampal slice cultures deprived of oxygen and glucose. <i>Neuroscience Letters</i> , 2003, 348, 123-125.	1.0	13
41	Integrins regulate neuronal neurotrophin gene expression through effects on voltage-sensitive calcium channels. <i>Neuroscience</i> , 2003, 118, 925-940.	1.1	34
42	Repeated Exposures to Subthreshold Doses of Chlorpyrifos in Rats: Hippocampal Damage, Impaired Axonal Transport, and Deficits in Spatial Learning. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2003, 305, 375-384.	1.3	96
43	Lysosomal Activation Is a Compensatory Response Against Protein Accumulation and Associated Synaptopathogenesis—An Approach for Slowing Alzheimer Disease?. <i>Journal of Neuropathology and Experimental Neurology</i> , 2003, 62, 451-463.	0.9	140
44	Recombinant vesicular stomatitis virus vectors as oncolytic agents in the treatment of high-grade gliomas in an organotypic brain tissue slice—glioma coculture model. <i>Journal of Neurosurgery</i> , 2004, 100, 1049-1059.	0.9	34
45	Neutralization of Transthyretin Reverses the Neuroprotective Effects of Secreted Amyloid Precursor Protein (APP) in APPSw Mice Resulting in Tau Phosphorylation and Loss of Hippocampal Neurons: Support for the Amyloid Hypothesis. <i>Journal of Neuroscience</i> , 2004, 24, 7707-7717.	1.7	243
46	The pathogenic activation of calpain: a marker and mediator of cellular toxicity and disease states. <i>International Journal of Experimental Pathology</i> , 2004, 81, 323-339.	0.6	98
47	Multiplexed Cytokine Protein Expression Profiles from Spreading Depression in Hippocampal Organotypic Cultures. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2004, 24, 829-839.	2.4	54
48	Homeostatic plasticity in hippocampal slice cultures involves changes in voltage-gated Na ⁺ channel expression. <i>Brain Research</i> , 2004, 998, 155-163.	1.1	44
49	Functional validation of adult hippocampal organotypic cultures as an in vitro model of brain injury. <i>Brain Research</i> , 2004, 1001, 125-132.	1.1	81
50	Crush injury induces NMDA-receptor-dependent delayed nerve cell death in rat entorhinal—hippocampal slice cultures. <i>Brain Research</i> , 2004, 1025, 35-42.	1.1	5
51	Biphasic NF- κ B activation in the excitotoxic hippocampus. <i>Acta Neuropathologica</i> , 2004, 108, 173-82.	3.9	15
52	P/Q Ca ²⁺ channel blockade stops spreading depression and related pyramidal neuronal Ca ²⁺ rise in hippocampal organ culture. <i>Hippocampus</i> , 2004, 14, 356-367.	0.9	39
53	Repeated contact with subtoxic soman leads to synaptic vulnerability in hippocampus. <i>Journal of Neuroscience Research</i> , 2004, 77, 739-746.	1.3	18
54	The brain angiotensin system and extracellular matrix molecules in neural plasticity, learning, and memory. <i>Progress in Neurobiology</i> , 2004, 72, 263-293.	2.8	150

#	ARTICLE	IF	CITATIONS
55	Prevention of neuron and oligodendrocyte degeneration by interleukin-6 (IL-6) and IL-6 receptor/IL-6 fusion protein in organotypic hippocampal slices. <i>Molecular and Cellular Neurosciences</i> , 2004, 25, 301-311.	1.0	84
56	Proteases and Neuronal Plasticity. , 2005, , 235-270.		0
57	Blocking cannabinoid activation of FAK and ERK1/2 compromises synaptic integrity in hippocampus. <i>European Journal of Pharmacology</i> , 2005, 508, 47-56.	1.7	49
58	Cytoskeletal, synaptic, and nuclear protein changes associated with rat interface organotypic hippocampal slice culture development. <i>Developmental Brain Research</i> , 2005, 160, 275-286.	2.1	22
59	Cellular Responses to Protein Accumulation Involve Autophagy and Lysosomal Enzyme Activation. <i>Rejuvenation Research</i> , 2005, 8, 227-237.	0.9	44
61	Activity-Dependent Dendritic Arborization Mediated by CaM-Kinase I Activation and Enhanced CREB-Dependent Transcription of Wnt-2. <i>Neuron</i> , 2006, 50, 897-909.	3.8	413
62	Immune Modulation by Estrogens: Role in CNS HIV-1 Infection. <i>Endocrine</i> , 2006, 29, 289-298.	2.2	29
63	Oxidative Stress and Lysosomes: CNS-Related Consequences and Implications for Lysosomal Enhancement Strategies and Induction of Autophagy. <i>Antioxidants and Redox Signaling</i> , 2006, 8, 185-196.	2.5	86
64	Fibronectin and neuroprotective effect of granulocyte colony-stimulating factor in focal cerebral ischemia. <i>Brain Research</i> , 2006, 1098, 161-169.	1.1	56
65	3-Nitropropionic acid toxicity in hippocampus: Protection through N-methyl-D-aspartate receptor antagonism. <i>Hippocampus</i> , 2006, 16, 834-842.	0.9	33
66	Acid-sensing ion channel 1a is a postsynaptic proton receptor that affects the density of dendritic spines. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 16556-16561.	3.3	178
67	Brain Slices as Models for Neurodegenerative Disease and Screening Platforms to Identify Novel Therapeutics. <i>Current Neuropharmacology</i> , 2007, 5, 19-33.	1.4	200
68	Lysosomal Dysfunction Produces Distinct Alterations in Synaptic $\hat{1}\pm$ -Amino-3-Hydroxy-5-Methylisoxazolepropionic Acid and N-Methyl-D-Aspartate Receptor Currents in Hippocampus. <i>Journal of Neuropathology and Experimental Neurology</i> , 2007, 66, 779-788.	0.9	20
69	Inositol 1,4,5 trisphosphate receptor and chromogranin B are concentrated in different regions of the hippocampus. <i>Journal of Neuroscience Research</i> , 2007, 85, 2026-2036.	1.3	16
70	Neuroprotective activity of selective mGlu1 and mGlu5 antagonists in vitro and in vivo. <i>European Journal of Pharmacology</i> , 2007, 554, 18-29.	1.7	41
71	Microtubule-stabilizing agent prevents protein accumulation-induced loss of synaptic markers. <i>European Journal of Pharmacology</i> , 2007, 562, 20-27.	1.7	56
72	Hydrogen Peroxide Induces Lysosomal Protease Alterations in PC12 Cells. <i>Neurochemical Research</i> , 2007, 32, 1499-1510.	1.6	22
73	Gephyrin Alterations Due to Protein Accumulation Stress are Reduced by the Lysosomal Modulator Z-Phe-Ala-Diazomethylketone. <i>Journal of Molecular Neuroscience</i> , 2008, 34, 131-139.	1.1	10

#	ARTICLE	IF	CITATIONS
74	Targeting IKK2 by pharmacological inhibitor AS602868 prevents excitotoxic injury to neurons and oligodendrocytes. <i>Journal of Neural Transmission</i> , 2008, 115, 693-701.	1.4	11
75	Neutralization of the membrane protein Nogo-66 enhances growth and reactive sprouting in established organotypic hippocampal slice cultures. <i>European Journal of Neuroscience</i> , 2008, 28, 1808-1824.	1.2	35
76	Determination of I^{Ca} -Potential in Rat Organotypic Hippocampal Cultures. <i>Biophysical Journal</i> , 2008, 94, 4561-4569.	0.2	26
77	Activity-Dependent Synaptogenesis: Regulation by a CaM-Kinase Kinase/CaM-Kinase I/ I^{Ca} PIX Signaling Complex. <i>Neuron</i> , 2008, 57, 94-107.	3.8	200
78	mGlu1 \pm receptors are co-expressed with CB1 receptors in a subset of interneurons in the CA1 region of organotypic hippocampal slice cultures and adult rat brain. <i>Neuropharmacology</i> , 2008, 55, 428-439.	2.0	21
79	An activity-regulated microRNA controls dendritic plasticity by down-regulating p250GAP. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 9093-9098.	3.3	495
80	The neuroprotective effects of xenon and helium in an in vitro model of traumatic brain injury*. <i>Critical Care Medicine</i> , 2008, 36, 588-595.	0.4	150
81	Lysosomal Modulatory Drugs for a Broad Strategy Against Protein Accumulation Disorders. <i>Current Alzheimer Research</i> , 2009, 6, 438-445.	0.7	14
82	Propofol: neuroprotection in an in vitro model of traumatic brain injury. <i>Critical Care</i> , 2009, 13, R61.	2.5	53
83	Argon: Neuroprotection in in vitro models of cerebral ischemia and traumatic brain injury. <i>Critical Care</i> , 2009, 13, R206.	2.5	115
84	Organotypic Hippocampal Slice Cultures as a Model of Limbic Epileptogenesis. <i>NeuroMethods</i> , 2009, , 183-201.	0.2	5
85	Animal Models of Epilepsy. <i>NeuroMethods</i> , 2009, , .	0.2	4
86	A novel method of organotypic brain slice culture using connexin-specific antisense oligodeoxynucleotides to improve neuronal survival. <i>Brain Research</i> , 2010, 1353, 194-203.	1.1	22
87	Organotypic Cultures as Tool to Test Long-Term Effects of Chemicals on the Nervous System. <i>Current Medicinal Chemistry</i> , 2010, 17, 987-1001.	1.2	31
88	An activity-induced microRNA controls dendritic spine formation by regulating Rac1-PAK signaling. <i>Molecular and Cellular Neurosciences</i> , 2010, 43, 146-156.	1.0	242
89	Microglia and neuroprotection: From in vitro studies to therapeutic applications. <i>Progress in Neurobiology</i> , 2010, 92, 293-315.	2.8	226
90	Submicromolar $\text{A}\beta_{42}$ reduces hippocampal glutamate receptors and presynaptic markers in an aggregation-dependent manner. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2011, 1812, 1664-1674.	1.8	18
91	Protective Effects of Positive Lysosomal Modulation in Alzheimer's Disease Transgenic Mouse Models. <i>PLoS ONE</i> , 2011, 6, e20501.	1.1	77

#	ARTICLE	IF	CITATIONS
92	A New Generation Fatty Acid Amide Hydrolase Inhibitor Protects Against Kainate-Induced Excitotoxicity. <i>Journal of Molecular Neuroscience</i> , 2011, 43, 493-502.	1.1	45
93	Positive Lysosomal Modulation As a Unique Strategy to Treat Age-Related Protein Accumulation Diseases. <i>Rejuvenation Research</i> , 2012, 15, 189-197.	0.9	43
94	Equipotent Inhibition of Fatty Acid Amide Hydrolase and Monoacylglycerol Lipase – Dual Targets of the Endocannabinoid System to Protect against Seizure Pathology. <i>Neurotherapeutics</i> , 2012, 9, 801-813.	2.1	49
95	Impact of Microglial Activation on Astroglial Connexin Expression and Function in Brain Inflammation. , 2013, , 247-260.		4
96	Neuroprotection against Traumatic Brain Injury by Xenon, but Not Argon, Is Mediated by Inhibition at the N-Methyl-D-Aspartate Receptor Glycine Site. <i>Anesthesiology</i> , 2013, 119, 1137-1148.	1.3	105
97	A Genome-Wide Screen of CREB Occupancy Identifies the RhoA Inhibitors Par6C and Rnd3 as Regulators of BDNF-Induced Synaptogenesis. <i>PLoS ONE</i> , 2013, 8, e64658.	1.1	57
98	The Blood-Brain Barrier as a Target in Traumatic Brain Injury Treatment. <i>Archives of Medical Research</i> , 2014, 45, 698-710.	1.5	107
99	Different modes of herpes simplex virus type 1 spread in brain and skin tissues. <i>Journal of NeuroVirology</i> , 2014, 20, 18-27.	1.0	13
100	PARP-1 activation causes neuronal death in the hippocampal CA1 region by increasing the expression of Ca ²⁺ -permeable AMPA receptors. <i>Neurobiology of Disease</i> , 2014, 70, 43-52.	2.1	39
101	Leptin Induces Hippocampal Synaptogenesis via CREB-Regulated MicroRNA-132 Suppression of p250GAP. <i>Molecular Endocrinology</i> , 2014, 28, 1073-1087.	3.7	74
102	Forebrain microglia from wild-type but not adult 5xFAD mice prevent amyloid- β plaque formation in organotypic hippocampal slice cultures. <i>Scientific Reports</i> , 2015, 5, 14624.	1.6	82
103	Ethanol withdrawal is required to produce persisting N-methyl-d-aspartate receptor-dependent hippocampal cytotoxicity during chronic intermittent ethanol exposure. <i>Alcohol</i> , 2015, 49, 219-227.	0.8	27
104	Organotypic brain slice cultures: A review. <i>Neuroscience</i> , 2015, 305, 86-98.	1.1	324
105	Multiple Single-Unit Long-Term Tracking on Organotypic Hippocampal Slices Using High-Density Microelectrode Arrays. <i>Frontiers in Neuroscience</i> , 2016, 10, 537.	1.4	20
106	Conversion of Synthetic A β to In Vivo Active Seeds and Amyloid Plaque Formation in a Hippocampal Slice Culture Model. <i>Journal of Neuroscience</i> , 2016, 36, 5084-5093.	1.7	41
107	Blast waves from detonated military explosive reduce GluR1 and synaptophysin levels in hippocampal slice cultures. <i>Experimental Neurology</i> , 2016, 286, 107-115.	2.0	18
108	Blast Injury Science and Engineering. , 2016, , .		11
109	Organotypic Hippocampal Slices as Models for Stroke and Traumatic Brain Injury. <i>Molecular Neurobiology</i> , 2016, 53, 4226-4237.	1.9	43

#	ARTICLE	IF	CITATIONS
110	A new method allowing long-term potentiation recordings in hippocampal organotypic slices. <i>Brain and Behavior</i> , 2017, 7, e00692.	1.0	2
111	Organotypic Hippocampal Slice Cultures As a Model to Study Neuroprotection and Invasiveness of Tumor Cells. <i>Journal of Visualized Experiments</i> , 2017, , .	0.2	16
112	A β 242-mediated proteasome inhibition and associated tau pathology in hippocampus are governed by a lysosomal response involving cathepsin B: Evidence for protective crosstalk between protein clearance pathways. <i>PLoS ONE</i> , 2017, 12, e0182895.	1.1	18
113	Xenon Protects against Blast-Induced Traumatic Brain Injury in an <i>In Vitro</i> Model. <i>Journal of Neurotrauma</i> , 2018, 35, 1037-1044.	1.7	30
114	Preparation of organotypic brain slice cultures for the study of Alzheimer's disease. <i>F1000Research</i> , 2018, 7, 592.	0.8	14
115	Replenishment of Organotypic Hippocampal Slice Cultures with Neonatal or Adult Microglia. <i>Methods in Molecular Biology</i> , 2019, 2034, 127-147.	0.4	2
116	Noble gas neuroprotection: xenon and argon protect against hypoxic-ischaemic injury in rat hippocampus <i>In Vitro</i> via distinct mechanisms. <i>British Journal of Anaesthesia</i> , 2019, 123, 601-609.	1.5	35
117	Early Synaptic Alterations and Selective Adhesion Signaling in Hippocampal Dendritic Zones Following Organophosphate Exposure. <i>Scientific Reports</i> , 2019, 9, 6532.	1.6	13
118	rAAV-based brain slice culture models of Alzheimer's and Parkinson's disease inclusion pathologies. <i>Journal of Experimental Medicine</i> , 2019, 216, 539-555.	4.2	48
119	Organotypic brain slice cultures to model neurodegenerative proteinopathies. <i>Molecular Neurodegeneration</i> , 2019, 14, 45.	4.4	69
120	Amyloid- β Oligomers Induce Only Mild Changes to Inhibitory Bouton Dynamics. <i>Journal of Alzheimer's Disease Reports</i> , 2021, 5, 153-160.	1.2	6
122	Rat Hippocampal Slice Culture Models for the Evaluation of Neuroprotective Agents. <i>Methods in Molecular Biology</i> , 2012, 846, 343-354.	0.4	33
123	Modelling Blast Brain Injury. , 2016, , 173-182.		1
124	Leptin stimulates synaptogenesis in hippocampal neurons via KLF4 and SOCS3 inhibition of STAT3 signaling. <i>Molecular and Cellular Neurosciences</i> , 2020, 106, 103500.	1.0	17
125	Competitive Inhibition at the Glycine Site of the N-Methyl-D-Aspartate Receptor Mediates Xenon Neuroprotection against Hypoxia-Ischemia. <i>Anesthesiology</i> , 2010, 112, 614-622.	1.3	86
126	Preparation of organotypic brain slice cultures for the study of Alzheimer's disease. <i>F1000Research</i> , 2018, 7, 592.	0.8	31
127	Acute death of astrocytes in blast-exposed rat organotypic hippocampal slice cultures. <i>PLoS ONE</i> , 2017, 12, e0173167.	1.1	25
128	Effect of the bone marrow multipotent mesenchymal stromal cells to the neural tissue after ischemic injury <i>in vitro</i> . <i>Cell and Organ Transplantation</i> , 2014, 2, 74-78.	0.2	1

#	ARTICLE	IF	CITATIONS
129	Polo-like kinase 2 inhibition reduces serine-129 phosphorylation of physiological nuclear alpha-synuclein but not of the aggregated alpha-synuclein. PLoS ONE, 2021, 16, e0252635.	1.1	18
130	Assessment of Neuronal Damage in Brain Slice Cultures Using Machine Learning Based on Spatial Features. Frontiers in Neuroscience, 2021, 15, 740178.	1.4	1
135	Organotypic whole hemisphere brain slice models to study the effects of donor age and oxygen-glucose-deprivation on the extracellular properties of cortical and striatal tissue. Journal of Biological Engineering, 2022, 16, .	2.0	1
136	Lymphatic cells do not functionally integrate into 3D organotypic brain slice cultures, but aggregate around penetrating blood vessels. Experimental Brain Research, 0, , .	0.7	0
137	Modelling Blast Brain Injury. , 2022, , 315-325.		0