

Role of lipid peroxidation in H₂O₂-induced renal epithel

American Journal of Physiology - Renal Physiology

268, F30-F38

DOI: [10.1152/ajprenal.1995.268.1.f30](https://doi.org/10.1152/ajprenal.1995.268.1.f30)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Effects of cumene hydroperoxide on cellular cation composition in frog kidney proximal tubular cells. <i>Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology</i> , 2000, 126, 251-262.	0.8	0
2	Adenosine attenuates oxidant injury in human proximal tubular cells via A1 and A2a adenosine receptors. <i>American Journal of Physiology - Renal Physiology</i> , 2002, 282, F844-F852.	1.3	78
3	Oxidative stress and gene expression of antioxidant enzymes in the renal cortex of streptozotocin-induced diabetic rats. <i>Molecular and Cellular Biochemistry</i> , 2003, 243, 147-152.	1.4	129
4	Bioflavonoids attenuate renal proximal tubular cell injury during cold preservation in Euro-Collins and University of Wisconsin solutions. <i>Kidney International</i> , 2003, 63, 554-563.	2.6	40
5	Garlic supplementation prevents oxidative DNA damage in essential hypertension. <i>Molecular and Cellular Biochemistry</i> , 2005, 275, 85-94.	1.4	74
6	In vitro efficacy of three lazaroids in a model of acute chemical neuronal hypoxia. <i>Neuroscience Letters</i> , 2006, 407, 171-175.	1.0	4
7	Anti-inflammatory and antinecrotic effects of the volatile anesthetic sevoflurane in kidney proximal tubule cells. <i>American Journal of Physiology - Renal Physiology</i> , 2006, 291, F67-F78.	1.3	85
8	Evaluation of selected parameters of the antioxidative system in patients with type 2 diabetes in different periods of metabolic compensation. <i>Archivum Immunologiae Et Therapiae Experimentalis</i> , 2007, 55, 335-340.	1.0	12
9	Idiopathic recurrent calcium urolithiasis (IRCU): variation of fasting urinary protein is a window to pathophysiology or simple consequence of renal stones in situ?. <i>European Journal of Medical Research</i> , 2009, 14, 378.	0.9	3
10	Comparative and combined effect of chlorogenic acid and tetrahydrocurcumin on antioxidant disparities in chemical induced experimental diabetes. <i>Molecular and Cellular Biochemistry</i> , 2010, 341, 109-117.	1.4	50
11	Isoflurane Protects Human Kidney Proximal Tubule Cells against Necrosis via Sphingosine Kinase and Sphingosine-1-Phosphate Generation. <i>American Journal of Nephrology</i> , 2010, 31, 353-362.	1.4	51
12	Hydroxytyrosol glucuronides protect renal tubular epithelial cells against H2O2 induced oxidative damage. <i>Chemico-Biological Interactions</i> , 2011, 193, 232-239.	1.7	26
13	Protective effects of propofol against hydrogen peroxide-induced oxidative stress in human kidney proximal tubular cells. <i>Korean Journal of Anesthesiology</i> , 2012, 63, 441.	0.9	8
14	The Prostaglandin E ₂ Type 4 Receptor Participates in the Response to Acute Oxidant Stress in Airway Epithelial Cells. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2012, 341, 552-563.	1.3	13
15	Antioxidative effects of fermented sesame sauce against hydrogen peroxide-induced oxidative damage in LLC-PK1 porcine renal tubule cells. <i>Nutrition Research and Practice</i> , 2014, 8, 138.	0.7	12
16	Protective effects of methanolic extract from fruits of <i>Lycium ruthenicum</i> Murr on 2,2-azobis (2-amidinopropane) dihydrochloride-induced oxidative stress in LLC-PK1 cells. <i>Pharmacognosy Magazine</i> , 2014, 10, 522.	0.3	19
17	<i>Nitraria retusa</i> fruit prevents penconazole-induced kidney injury in adult rats through modulation of oxidative stress and histopathological changes. <i>Pharmaceutical Biology</i> , 2017, 55, 1061-1073.	1.3	35