

CITATION REPORT

List of articles citing

A phosphotyrosine interaction domain

DOI: 10.1016/0092-8674(95)90347-x
Cell, 1995, 80, 693-4.

Source: <https://exaly.com/paper-pdf/26490154/citation-report.pdf>

Version: 2024-04-24

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
190	The phosphotyrosine interaction domain of Shc binds an LXNPXY motif on the epidermal growth factor receptor. <i>Molecular and Cellular Biology</i> , 1995 , 15, 4403-9	4.8	116
189	Chapter 26. Tyrosine Kinase Induced Mitogenesis Breaking the Link With Cancer. 1995 , 30, 247-253		12
188	The WW domain of Yes-associated protein binds a proline-rich ligand that differs from the consensus established for Src homology 3-binding modules. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995 , 92, 7819-23	11.5	498
187	Structure and function of the phosphotyrosine binding (PTB) domain. 1995 , 64, 221-35		18
186	Recognition and specificity in protein tyrosine kinase-mediated signalling. 1995 , 20, 470-5		322
185	A new flavor in phosphotyrosine recognition. 1995 , 3, 421-4		18
184	SAM: a novel motif in yeast sterile and Drosophila polyhomeotic proteins. 1995 , 4, 1928-30		133
183	Role of IRS-2 in insulin and cytokine signalling. 1995 , 377, 173-7		773
182	Structure and ligand recognition of the phosphotyrosine binding domain of Shc. 1995 , 378, 584-92		338
181	L-O-(2-malonyl)tyrosine: a new phosphotyrosyl mimetic for the preparation of Src homology 2 domain inhibitory peptides. 1995 , 38, 4270-5		71
180	The Drosophila insulin receptor homolog: a gene essential for embryonic development encodes two receptor isoforms with different signaling potential.. <i>EMBO Journal</i> , 1995 , 14, 3373-3384	13	232
179	Signal transduction by the alpha 6 beta 4 integrin: distinct beta 4 subunit sites mediate recruitment of Shc/Grb2 and association with the cytoskeleton of hemidesmosomes.. <i>EMBO Journal</i> , 1995 , 14, 4470-4481	12	188
178	Coupling of the proto-oncogene product c-Cbl to the epidermal growth factor receptor. <i>Journal of Biological Chemistry</i> , 1995 , 270, 25332-5	5.4	86
177	The pleckstrin homology domain in insulin receptor substrate-1 sensitizes insulin signaling. <i>Journal of Biological Chemistry</i> , 1995 , 270, 11715-8	5.4	87
176	The phosphotyrosine interaction domain of SHC recognizes tyrosine-phosphorylated NPXY motif. <i>Journal of Biological Chemistry</i> , 1995 , 270, 14863-6	5.4	152
175	Distinct modes of interaction of SHC and insulin receptor substrate-1 with the insulin receptor NPEY region via non-SH2 domains. <i>Journal of Biological Chemistry</i> , 1995 , 270, 23258-62	5.4	67
174	Evidence for a differential interaction of SHC and the insulin receptor substrate-1 (IRS-1) with the insulin-like growth factor-I (IGF-I) receptor in the yeast two-hybrid system. <i>Journal of Biological Chemistry</i> , 1995 , 270, 23456-60	5.4	81

173	Cloning of a novel phosphoprotein regulated by colony-stimulating factor 1 shares a domain with the Drosophila disabled gene product. <i>Journal of Biological Chemistry</i> , 1995 , 270, 14184-91	5.4	119
172	The regions of the Fe65 protein homologous to the phosphotyrosine interaction/phosphotyrosine binding domain of Shc bind the intracellular domain of the Alzheimer β amyloid precursor protein. <i>Journal of Biological Chemistry</i> , 1995 , 270, 30853-6	5.4	243
171	Shc binding to nerve growth factor receptor is mediated by the phosphotyrosine interaction domain. <i>Journal of Biological Chemistry</i> , 1995 , 270, 15125-9	5.4	106
170	New frontiers in insulin receptor substrate signaling. 1995 , 6, 209-15		52
169	HIV nuclear import is governed by the phosphotyrosine-mediated binding of matrix to the core domain of integrase. <i>Cell</i> , 1995 , 83, 569-76	56.2	325
168	Characterization of a novel protein-binding module--the WW domain. <i>FEBS Letters</i> , 1995 , 369, 67-71	3.8	286
167	The PTB domain: a new protein module implicated in signal transduction. 1995 , 20, 277-80		210
166	4RO-[2-(2-fluoromalonyl)]-L-tyrosine: a phosphotyrosyl mimic for the preparation of signal transduction inhibitory peptides. 1996 , 39, 1021-7		98
165	Molecular cloning and characterization of Byp, a murine receptor-type tyrosine phosphatase similar to human DEP-1. <i>FEBS Letters</i> , 1996 , 378, 7-14	3.8	33
164	Association of the protein tyrosine phosphatase PTP1C with the protein tyrosine kinase c-Src in human platelets. <i>FEBS Letters</i> , 1996 , 383, 165-9	3.8	33
163	Protein fold recognition by mapping predicted secondary structures. <i>Journal of Molecular Biology</i> , 1996 , 259, 349-65	6.5	119
162	The intracellular cytoplasmic domain of the Alzheimer β disease amyloid precursor protein interacts with phosphotyrosine-binding domain proteins in the yeast two-hybrid system. <i>FEBS Letters</i> , 1996 , 397, 197-200	3.8	124
161	PH domains: diverse sequences with a common fold recruit signaling molecules to the cell surface. <i>Cell</i> , 1996 , 85, 621-4	56.2	459
160	Structure of the IRS-1 PTB domain bound to the juxtamembrane region of the insulin receptor. <i>Cell</i> , 1996 , 85, 695-705	56.2	278
159	How do protein kinases recognize their substrates?. 1996 , 1314, 191-225		362
158	Protein sequence motifs. 1996 , 6, 366-76		119
157	Numb antagonizes Notch signaling to specify sibling neuron cell fates. 1996 , 17, 21-6		367
156	Control of daughter cell fates during asymmetric division: interaction of Numb and Notch. 1996 , 17, 27-41		583

155	Asymmetric localization of a mammalian numb homolog during mouse cortical neurogenesis. 1996 , 17, 43-53		420
154	Binding of zinc finger protein ZPR1 to the epidermal growth factor receptor. <i>Science</i> , 1996 , 272, 1797-803	3.3	96
153	Association of a novel human FE65-like protein with the cytoplasmic domain of the beta-amyloid precursor protein. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996 , 93, 10832-7	11.5	153
152	Interaction between the phosphotyrosine binding domain of Shc and the insulin receptor is required for Shc phosphorylation by insulin in vivo. <i>Journal of Biological Chemistry</i> , 1996 , 271, 3959-62	5.4	55
151	The Drosophila Numb protein inhibits signaling of the Notch receptor during cell-cell interaction in sensory organ lineage. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996 , 93, 11925-32	11.5	262
150	Formin binding proteins bear WWP/WW domains that bind proline-rich peptides and functionally resemble SH3 domains.. <i>EMBO Journal</i> , 1996 , 15, 1045-1054	13	169
149	The phosphotyrosine interaction domains of X11 and FE65 bind to distinct sites on the YENPTY motif of amyloid precursor protein. <i>Molecular and Cellular Biology</i> , 1996 , 16, 6229-41	4.8	429
148	The Drosophila insulin receptor activates multiple signaling pathways but requires insulin receptor substrate proteins for DNA synthesis. <i>Molecular and Cellular Biology</i> , 1996 , 16, 2509-17	4.8	81
147	Structure and distribution of modules in extracellular proteins. 1996 , 29, 119-67		266
146	PDGF and FGF receptors in health and disease. 1996 , 179-228		3
145	The MEN II syndromes and the role of the ret proto-oncogene. 1996 , 70, 179-222		47
144	Signal-transducing protein phosphorylation cascades mediated by Ras/Rho proteins in the mammalian cell: the potential for multiplex signalling. 1996 , 318 (Pt 3), 729-47		447
143	SH2 domain protein interaction and possibilities for pharmacological intervention. <i>Cellular Signalling</i> , 1996 , 8, 75-86	4.9	23
142	Structure and function of the WW domain. 1996 , 65, 113-32		229
141	The PI/PTB domain: a new protein interaction domain involved in growth factor receptor signaling. 1996 , 128, 235-41		41
140	Mammalian NUMB is an evolutionarily conserved signaling adapter protein that specifies cell fate. 1996 , 6, 1134-45		142
139	WW domains. 1996 , 4, 495-9		85
138	pH-Dependent self-association of the Src homology 2 (SH2) domain of the Src homologous and collagen-like (SHC) protein. 1996 , 5, 405-13		9

137	Structure of the WW domain of a kinase-associated protein complexed with a proline-rich peptide. 1996 , 382, 646-9		379
136	Phosphotyrosine-independent binding of SHC to the NPLH sequence of murine protein-tyrosine phosphatase-PEST. Evidence for extended phosphotyrosine binding/phosphotyrosine interaction domain recognition specificity. <i>Journal of Biological Chemistry</i> , 1996 , 271, 8424-9	5.4	66
135	Annexin II is a novel player in insulin signal transduction. Possible association between annexin II phosphorylation and insulin receptor internalization. <i>Journal of Biological Chemistry</i> , 1996 , 271, 29489-96	5.4	100
134	cDNA cloning and chromosome mapping of the human Fe65 gene: interaction of the conserved cytoplasmic domains of the human beta-amyloid precursor protein and its homologues with the mouse Fe65 protein. 1996 , 5, 1589-98		95
133	Interaction between the insulin receptor and its downstream effectors. Use of individually expressed receptor domains for structure/function analysis. <i>Journal of Biological Chemistry</i> , 1996 , 271, 6998-7003	5.4	34
132	The pleckstrin homology domain is the principal link between the insulin receptor and IRS-1. <i>Journal of Biological Chemistry</i> , 1996 , 271, 24300-6	5.4	133
131	Identification of residues within the SHC phosphotyrosine binding/phosphotyrosine interaction domain crucial for phosphopeptide interaction. <i>Journal of Biological Chemistry</i> , 1996 , 271, 1813-6	5.4	35
130	Enhanced tumorigenic behavior of glioblastoma cells expressing a truncated epidermal growth factor receptor is mediated through the Ras-Shc-Grb2 pathway. <i>Journal of Biological Chemistry</i> , 1996 , 271, 25639-45	5.4	127
129	Thermodynamic studies of SHC phosphotyrosine interaction domain recognition of the NPXpY motif. <i>Journal of Biological Chemistry</i> , 1996 , 271, 4770-5	5.4	28
128	Binding specificity and mutational analysis of the phosphotyrosine binding domain of the brain-specific adaptor protein ShcC. <i>Journal of Biological Chemistry</i> , 1996 , 271, 11787-91	5.4	16
127	Affinity, specificity, and kinetics of the interaction of the SHC phosphotyrosine binding domain with asparagine-X-X-phosphotyrosine motifs of growth factor receptors. <i>Journal of Biological Chemistry</i> , 1996 , 271, 264-9	5.4	59
126	Systematic mapping of potential binding sites for Shc and Grb2 SH2 domains on insulin receptor substrate-1 and the receptors for insulin, epidermal growth factor, platelet-derived growth factor, and fibroblast growth factor. <i>Journal of Biological Chemistry</i> , 1996 , 271, 5603-9	5.4	51
125	Activation-induced association of a 145-kDa tyrosine-phosphorylated protein with Shc and Syk in B lymphocytes and macrophages. <i>Journal of Biological Chemistry</i> , 1996 , 271, 1145-52	5.4	73
124	Identification of six novel autophosphorylation sites on fibroblast growth factor receptor 1 and elucidation of their importance in receptor activation and signal transduction. <i>Molecular and Cellular Biology</i> , 1996 , 16, 977-89	4.8	334
123	The role of the Shc phosphotyrosine interaction/phosphotyrosine binding domain and tyrosine phosphorylation sites in polyoma middle T antigen-mediated cell transformation. <i>Journal of Biological Chemistry</i> , 1997 , 272, 20671-7	5.4	28
122	Mints, Munc18-interacting proteins in synaptic vesicle exocytosis. <i>Journal of Biological Chemistry</i> , 1997 , 272, 31459-64	5.4	262
121	The 66-kDa Shc isoform is a negative regulator of the epidermal growth factor-stimulated mitogen-activated protein kinase pathway. <i>Journal of Biological Chemistry</i> , 1997 , 272, 28042-9	5.4	112
120	The WW domain of neural protein FE65 interacts with proline-rich motifs in Mena, the mammalian homolog of <i>Drosophila</i> enabled. <i>Journal of Biological Chemistry</i> , 1997 , 272, 32869-77	5.4	191

119	14-3-3 protein binds to insulin receptor substrate-1, one of the binding sites of which is in the phosphotyrosine binding domain. <i>Journal of Biological Chemistry</i> , 1997 , 272, 25267-74	5.4	126
118	Characterization of the WW domain of human yes-associated protein and its polyproline-containing ligands. <i>Journal of Biological Chemistry</i> , 1997 , 272, 17070-7	5.4	147
117	Distinct tyrosine autophosphorylation sites negatively and positively modulate neu-mediated transformation. <i>Molecular and Cellular Biology</i> , 1997 , 17, 5410-25	4.8	124
116	DNA-binding protein Pur alpha and transcription factor YY1 function as transcription activators of the neuron-specific FE65 gene promoter. 1997 , 328 (Pt 1), 293-300		61
115	Chapter 6 Molecular basis of insulin action. 1997 , 161-202		
114	Tyrosine phosphorylation of transmembrane ligands for Eph receptors. <i>Science</i> , 1997 , 275, 1640-3	33.3	351
113	High-affinity binding of the Drosophila Numb phosphotyrosine-binding domain to peptides containing a Gly-Pro-(p)Tyr motif. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997 , 94, 7204-9	11.5	64
112	The N terminus of the Drosophila Numb protein directs membrane association and actin-dependent asymmetric localization. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997 , 94, 13005-10	11.5	90
111	A Tyrosine-Phosphorylated Protein of 140 kD Is Constitutively Associated With the Phosphotyrosine Binding Domain of Shc and the SH3 Domains of Grb2 in Acute Myeloid Leukemia Cells. <i>Blood</i> , 1997 , 89, 2024-2035	2.2	13
110	Sequence-specific recognition of the internalization motif of the Alzheimer β amyloid precursor protein by the X11 PTB domain. <i>EMBO Journal</i> , 1997 , 16, 6141-50	13	132
109	The role of the PH domain in the signal-dependent membrane targeting of Sos. <i>EMBO Journal</i> , 1997 , 16, 1351-9	13	107
108	FBP WW domains and the Abl SH3 domain bind to a specific class of proline-rich ligands. <i>EMBO Journal</i> , 1997 , 16, 2376-83	13	178
107	On the classification and evolution of protein modules. 1997 , 16, 545-51		29
106	Phosphatidylcholine signaling in response to CSF-1. 1997 , 46, 24-30		11
105	SH2 and PTB domain interactions in tyrosine kinase signal transduction. 1997 , 1, 227-34		76
104	Negative signaling in B cells: SHIP Grbs Shc. 1997 , 18, 424-7		74
103	PDZ domains: targeting signalling molecules to sub-membranous sites. 1997 , 19, 469-79		376
102	Constitutively tyrosine phosphorylated p52 Shc in breast cancer cells: correlation with ErbB2 and p66 Shc expression. 1998 , 49, 119-28		33

101	Insulin signal transduction through protein kinase cascades. 1998 , 182, 31-48		272
100	Insulin induces tyrosine phosphorylation of Shc and stimulates Shc/GRB2 association in insulin-sensitive tissues of the intact rat. 1998 , 8, 193-200		17
99	Disabled-2 (Dab2) is an SH3 domain-binding partner of Grb2. <i>Oncogene</i> , 1998 , 16, 1561-9	9.2	109
98	Activation loop tyrosines contribute varying roles to TrkB autophosphorylation and signal transduction. <i>Oncogene</i> , 1998 , 16, 1691-700	9.2	23
97	DOC-2, a candidate tumor suppressor gene in human epithelial ovarian cancer. <i>Oncogene</i> , 1998 , 16, 2381-92		145
96	DOC-2/hDab2, a candidate tumor suppressor gene involved in the development of gestational trophoblastic diseases. <i>Oncogene</i> , 1998 , 17, 419-24	9.2	79
95	Structure of a Numb PTB domain-peptide complex suggests a basis for diverse binding specificity. <i>Nature Structural Biology</i> , 1998 , 5, 1075-83		98
94	The human FE65 gene: genomic structure and an intronic biallelic polymorphism associated with sporadic dementia of the Alzheimer type. 1998 , 103, 295-303		65
93	Candidate adaptor protein CED-6 promotes the engulfment of apoptotic cells in <i>C. elegans</i> . <i>Cell</i> , 1998 , 93, 961-72	56.2	179
92	Fe65 and the protein network centered around the cytosolic domain of the Alzheimer β beta-amyloid precursor protein. <i>FEBS Letters</i> , 1998 , 434, 1-7	3.8	92
91	Modulators of Notch signaling. 1998 , 9, 609-17		68
90	Adjacent carboxyterminal tyrosine phosphorylation events identify functionally distinct ErbB2 receptor subsets: implications for molecular diagnostics. <i>Experimental Cell Research</i> , 1998 , 241, 467-75	4.2	10
89	A rho-associated protein kinase, ROK α , binds insulin receptor substrate-1 and modulates insulin signaling. <i>Journal of Biological Chemistry</i> , 1998 , 273, 4740-6	5.4	55
88	X11 interaction with beta-amyloid precursor protein modulates its cellular stabilization and reduces amyloid beta-protein secretion. <i>Journal of Biological Chemistry</i> , 1998 , 273, 22351-7	5.4	122
87	IB1, a JIP-1-related nuclear protein present in insulin-secreting cells. <i>Journal of Biological Chemistry</i> , 1998 , 273, 1843-6	5.4	123
86	Identification of an evolutionarily conserved heterotrimeric protein complex involved in protein targeting. <i>Journal of Biological Chemistry</i> , 1998 , 273, 31633-6	5.4	160
85	Role of tyrosine phosphorylation in ligand-induced regulation of transcytosis of the polymeric Ig receptor. <i>Molecular Biology of the Cell</i> , 1998 , 9, 1787-802	3.5	45
84	Functional analysis of the Numb phosphotyrosine-binding domain using site-directed mutagenesis. <i>Journal of Biological Chemistry</i> , 1998 , 273, 10381-8	5.4	36

83	The mammalian numb phosphotyrosine-binding domain. Characterization of binding specificity and identification of a novel PDZ domain-containing numb binding protein, LNX. <i>Journal of Biological Chemistry</i> , 1998 , 273, 9179-87	5.4	93
82	Insulin signal transduction through protein kinase cascades. 1998 , 31-48		9
81	A role for the insulin-interleukin (IL)-4 receptor motif of the IL-4 receptor alpha-chain in regulating activation of the insulin receptor substrate 2 and signal transducer and activator of transcription 6 pathways. Analysis by mutagenesis. <i>Journal of Biological Chemistry</i> , 1998 , 273, 9898-905	5.4	22
80	Interaction of cytosolic adaptor proteins with neuronal apolipoprotein E receptors and the amyloid precursor protein. <i>Journal of Biological Chemistry</i> , 1998 , 273, 33556-60	5.4	446
79	Phosphorylation of Enabled by the Drosophila Abelson tyrosine kinase regulates the in vivo function and protein-protein interactions of Enabled. <i>Molecular and Cellular Biology</i> , 1998 , 18, 152-60	4.8	72
78	Numb-associated kinase interacts with the phosphotyrosine binding domain of Numb and antagonizes the function of Numb in vivo. <i>Molecular and Cellular Biology</i> , 1998 , 18, 598-607	4.8	65
77	Disabled is a putative adaptor protein that functions during signaling by the sevenless receptor tyrosine kinase. <i>Molecular and Cellular Biology</i> , 1998 , 18, 4844-54	4.8	26
76	SMART, a simple modular architecture research tool: identification of signaling domains. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998 , 95, 5857-64	11.5	3017
75	Disabled-1 binds to the cytoplasmic domain of amyloid precursor-like protein 1. <i>Journal of Neuroscience</i> , 1999 , 19, 7507-15	6.6	160
74	Interaction of c-Jun amino-terminal kinase interacting protein-1 with p190 rhoGEF and its localization in differentiated neurons. <i>Journal of Biological Chemistry</i> , 1999 , 274, 35113-8	5.4	105
73	Crystal structure of the pleckstrin homology-phosphotyrosine binding (PH-PTB) targeting region of insulin receptor substrate 1. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999 , 96, 8378-83	11.5	85
72	Different tyrosine autophosphorylation requirements in fibroblast growth factor receptor-1 mediate urokinase-type plasminogen activator induction and mitogenesis. <i>Molecular Biology of the Cell</i> , 1999 , 10, 23-33	3.5	26
71	Characterization of four mammalian numb protein isoforms. Identification of cytoplasmic and membrane-associated variants of the phosphotyrosine binding domain. <i>Journal of Biological Chemistry</i> , 1999 , 274, 33097-104	5.4	144
70	Regulation of beta-amyloid secretion by FE65, an amyloid protein precursor-binding protein. <i>Journal of Biological Chemistry</i> , 1999 , 274, 7952-7	5.4	170
69	The function of PTB domain proteins. <i>Kidney International</i> , 1999 , 56, 1230-7	9.9	55
68	The surprisingly flexible PTB domain. <i>Nature Structural Biology</i> , 1999 , 6, 7-10		15
67	Disabled-2 inactivation is an early step in ovarian tumorigenicity. <i>Oncogene</i> , 1999 , 18, 3104-13	9.2	117
66	A WW domain-containing yes-associated protein (YAP) is a novel transcriptional co-activator. <i>EMBO Journal</i> , 1999 , 18, 2551-62	13	436

65	Raf-like Ras/Rap-binding domains in RGS12- and still-life-like signalling proteins. <i>Journal of Molecular Medicine</i> , 1999 , 77, 695-8	5.5	54
64	The insulin receptor tyrosine kinase domain in a chimaeric epidermal growth factor-insulin receptor generates Ca ²⁺ signals through the PLC-gamma1 pathway. <i>BBA - Proteins and Proteomics</i> , 1999 , 1431, 421-32		4
63	The disabled 1 phosphotyrosine-binding domain binds to the internalization signals of transmembrane glycoproteins and to phospholipids. <i>Molecular and Cellular Biology</i> , 1999 , 19, 5179-88	4.8	338
62	Suppression of epidermal growth factor-induced phospholipase C activation associated with actin rearrangement in rat hepatocytes in primary culture. <i>Hepatology</i> , 2000 , 32, 947-57	11.2	11
61	Multiple modes of peptide recognition by the PTB domain of the cell fate determinant Numb. <i>EMBO Journal</i> , 2000 , 19, 1505-15	13	77
60	Peptidomimetic SH2 Domain Antagonists for Targeting Signal Transduction. <i>Topics in Current Chemistry</i> , 2000 , 17-59		21
59	Overexpression, purification, and characterization of glutaminase-interacting protein, a PDZ-domain protein from human brain. <i>Protein Expression and Purification</i> , 2001 , 23, 411-8	2	13
58	ConSurf: an algorithmic tool for the identification of functional regions in proteins by surface mapping of phylogenetic information. <i>Journal of Molecular Biology</i> , 2001 , 307, 447-63	6.5	380
57	IRS-1 tyrosine phosphorylation reflects insulin-induced metabolic and mitogenic responses in 3T3-L1 pre-adipocytes. <i>Archives of Physiology and Biochemistry</i> , 2001 , 109, 52-62	2.2	5
56	Autosomal recessive hypercholesterolemia caused by mutations in a putative LDL receptor adaptor protein. <i>Science</i> , 2001 , 292, 1394-8	33.3	448
55	c-Jun N-terminal kinase (JNK)-interacting protein-1b/islet-brain-1 scaffolds Alzheimer's amyloid precursor protein with JNK. <i>Journal of Neuroscience</i> , 2001 , 21, 6597-607	6.6	169
54	Surfing the insulin signaling web. <i>European Journal of Clinical Investigation</i> , 2001 , 31, 966-77	4.6	63
53	hFE65L influences amyloid precursor protein maturation and secretion. <i>Journal of Neurochemistry</i> , 1999 , 73, 985-93	6	65
52	Disabled-2 colocalizes with the LDLR in clathrin-coated pits and interacts with AP-2. <i>Traffic</i> , 2001 , 2, 111-23	3.7	214
51	Notch signaling in mammary gland tumorigenesis. <i>Journal of Mammary Gland Biology and Neoplasia</i> , 2001 , 6, 23-36	2.4	87
50	Doc-2/hDab2 expression is up-regulated in primary pancreatic cancer but reduced in metastasis. <i>Laboratory Investigation</i> , 2001 , 81, 863-73	5.9	23
49	Transgenic expression of numb inhibits notch signaling in immature thymocytes but does not alter T cell fate specification. <i>Journal of Immunology</i> , 2002 , 168, 3173-80	5.3	43
48	The direct recruitment of BLNK to immunoglobulin alpha couples the B-cell antigen receptor to distal signaling pathways. <i>Molecular and Cellular Biology</i> , 2002 , 22, 2524-35	4.8	109

47	Autosomal recessive hypercholesterolaemia in Sardinia, Italy, and mutations in ARH: a clinical and molecular genetic analysis. <i>Lancet, The</i> , 2002 , 359, 841-7	4.0	129
46	Lipoprotein receptors in the nervous system. <i>Annual Review of Biochemistry</i> , 2002 , 71, 405-34	29.1	346
45	The endocytic protein alpha-Adaptin is required for numb-mediated asymmetric cell division in <i>Drosophila</i> . <i>Developmental Cell</i> , 2002 , 3, 221-31	10.2	313
44	A role for APP in motility and transcription?. <i>Trends in Pharmacological Sciences</i> , 2002 , 23, 203-5; discussion 205-6	13.2	4
43	An adipocentric view of signaling and intracellular trafficking. <i>Diabetes/Metabolism Research and Reviews</i> , 2002 , 18, 345-56	7.5	130
42	Down-regulation of DOC-2 in colorectal cancer points to its role as a tumor suppressor in this malignancy. <i>Diseases of the Colon and Rectum</i> , 2002 , 45, 1242-8	3.1	36
41	SH2 and PTB domains in tyrosine kinase signaling. <i>Science Signaling</i> , 2003 , 2003, RE12	8.8	189
40	Crystal structures of the Dab homology domains of mouse disabled 1 and 2. <i>Journal of Biological Chemistry</i> , 2003 , 278, 36572-81	5.4	101
39	Cytoplasmic tail adaptors of Alzheimer's amyloid-beta protein precursor. <i>Molecular Neurobiology</i> , 2004 , 30, 185-200	6.2	7
38	Polarized distribution of mRNAs encoding a putative LDL receptor adaptor protein, xARH (autosomal recessive hypercholesterolemia) in <i>Xenopus</i> oocytes. <i>Mechanisms of Development</i> , 2004 , 121, 1249-58	1.7	6
37	Generating neuronal diversity in the <i>Drosophila</i> central nervous system: a view from the ganglion mother cells. <i>Developmental Dynamics</i> , 2005 , 232, 609-16	2.9	30
36	A novel EB-1/AIDA-1 isoform, AIDA-1c, interacts with the Cajal body protein coilin. <i>BMC Cell Biology</i> , 2005 , 6, 23		16
35	Phosphorylation of Numb family proteins. Possible involvement of Ca ²⁺ /calmodulin-dependent protein kinases. <i>Journal of Biological Chemistry</i> , 2005 , 280, 35108-18	5.4	42
34	Endocytosis of megalin by visceral endoderm cells requires the Dab2 adaptor protein. <i>Journal of Cell Science</i> , 2005 , 118, 5345-55	5.3	81
33	Phosphorylation of Numb regulates its interaction with the clathrin-associated adaptor AP-2. <i>FEBS Letters</i> , 2006 , 580, 5797-801	3.8	19
32	Membrane and juxtamembrane targeting by PH and PTB domains. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2006 , 1761, 850-67	5	125
31	Modulation of lipoprotein receptor functions by intracellular adaptor proteins. <i>Cellular Signalling</i> , 2006 , 18, 1560-71	4.9	87
30	Expression of mDab1 promotes the stability and processing of amyloid precursor protein and this effect is counteracted by X11alpha. <i>Neurobiology of Aging</i> , 2007 , 28, 377-88	5.6	38

29	Regulation of receptor tyrosine kinase signaling by GRKs and beta-arrestins. <i>Annual Review of Physiology</i> , 2007 , 69, 561-77	23.1	68
28	Symmetric division and expression of its regulatory gene Numb in human cervical squamous carcinoma cells. <i>Pathobiology</i> , 2009 , 76, 149-54	3.6	19
27	The APP-interacting protein FE65 is required for hippocampus-dependent learning and long-term potentiation. <i>Learning and Memory</i> , 2009 , 16, 537-44	2.8	20
26	Differential requirement for Dab2 in the development of embryonic and extra-embryonic tissues. <i>BMC Developmental Biology</i> , 2013 , 13, 39	3.1	31
25	Endocytosis and Physiology: Insights from Disabled-2 Deficient Mice. <i>Frontiers in Cell and Developmental Biology</i> , 2016 , 4, 129	5.7	20
24	NUMB regulates the endocytosis and activity of the anaplastic lymphoma kinase in an isoform-specific manner. <i>Journal of Molecular Cell Biology</i> , 2019 , 11, 994-1005	6.3	6
23	Endocytic Adaptors in Cardiovascular Disease. <i>Frontiers in Cell and Developmental Biology</i> , 2020 , 8, 6241597	5.7	2
22	Dab2 gene variant is associated with increased coronary artery disease risk in Chinese Han population. <i>Medicine (United States)</i> , 2020 , 99, e20924	1.8	3
21	Proteins Implicated In Alzheimer Disease. <i>Advances in Experimental Medicine and Biology</i> , 1998 , 161-180	3.6	9
20	App Localization and Trafficking in the Central Nervous System. <i>Advances in Behavioral Biology</i> , 1998 , 487-494		1
19	Tyrosine Phosphorylation in Oxidative Stress. 1997 , 181-199		5
18	Signal transduction by the receptor tyrosine kinase Ret. <i>Recent Results in Cancer Research</i> , 1998 , 154, 271-81	1.5	45
17	Function of PTB domains. <i>Current Topics in Microbiology and Immunology</i> , 1998 , 228, 23-38	3.3	28
16	WW (WWP) domains: from structure to function. <i>Current Topics in Microbiology and Immunology</i> , 1998 , 228, 115-33	3.3	33
15	WW domain-mediated interactions reveal a spliceosome-associated protein that binds a third class of proline-rich motif: the proline glycine and methionine-rich motif. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998 , 95, 10602-7	11.5	124
14	SH2 and PTB Domains in Tyrosine Kinase Signaling. <i>Science Signaling</i> , 2003 , 2003, re12-re12	8.8	26
13	Purification and Molecular Cloning of SH2- and SH3-Containing Inositol Polyphosphate-5-Phosphatase, Which Is Involved in the Signaling Pathway of Granulocyte-Macrophage Colony-Stimulating Factor, Erythropoietin, and Bcr-Abl. <i>Blood</i> , 1997 , 89, 2745-2756	2.2	34
12	Hormonal induction and roles of Disabled-2 in lactation and involution. <i>PLoS ONE</i> , 2014 , 9, e110737	3.7	6

11	Pleckstrin Homology (PH) Domains.		
10	PTB Domains.		
9	FcRI-Induced Protein Tyrosine Phosphorylation. <i>Molecular Biology Intelligence Unit</i> , 1997 , 33-54		2
8	Signaling Components and Pathways. 1998 , 41-95		
7	Design and Use of Synthetic Peptides as Biological Models. 1998 , 207-257		
6	Asymmetric Cell Division and Fate Specification in the Drosophila Central Nervous System. 1999 , 273-285		
5	The Drosophila insulin receptor homolog: a gene essential for embryonic development encodes two receptor isoforms with different signaling potential. <i>EMBO Journal</i> , 1995 , 14, 3373-84	13	96
4	Signal transduction by the alpha 6 beta 4 integrin: distinct beta 4 subunit sites mediate recruitment of Shc/Grb2 and association with the cytoskeleton of hemidesmosomes. <i>EMBO Journal</i> , 1995 , 14, 4470-81	13	75
3	Formin binding proteins bear WWP/WW domains that bind proline-rich peptides and functionally resemble SH3 domains. <i>EMBO Journal</i> , 1996 , 15, 1045-54	13	56
2	Insulin signal transduction through protein kinase cascades. <i>Molecular and Cellular Biochemistry</i> , 1998 , 182, 31-48	4.2	79
1	Pleckstrin Homology Domains.		0