

CITATION REPORT

List of articles citing

Hypertension in the hemodialysis population: a survey of 649 patients

DOI: 10.1016/0272-6386(95)90492-1
American Journal of Kidney Diseases, 1995, 26, 461-8.

Source: <https://exaly.com/paper-pdf/26663442/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| # | Paper | IF | Citations |
|-----|--|-----|-----------|
| 152 | Hypertension in the hemodialysis population: any relation to one-year survival?. <i>American Journal of Kidney Diseases</i> , 1996 , 28, 737-40 | 7.4 | 78 |
| 151 | Hypertension in the dialysis population: no easy answers. 1996 , 19, 693-694 | | 4 |
| 150 | Predictors of stroke in patients receiving chronic hemodialysis. 1996 , 50, 1672-5 | | 93 |
| 149 | Use of the medical differential diagnosis to achieve optimal end-stage renal disease outcomes. 1997 , 4, 97-111 | | 1 |
| 148 | NKF-DOQI clinical practice guidelines for hemodialysis adequacy. National Kidney Foundation. <i>American Journal of Kidney Diseases</i> , 1997 , 30, S15-66 | 7.4 | 105 |
| 147 | Ambulatory blood pressure monitoring in dialysis patients and estimation of mean interdialytic blood pressure. <i>American Journal of Kidney Diseases</i> , 1997 , 29, 678-84 | 7.4 | 67 |
| 146 | Reply. <i>American Journal of Kidney Diseases</i> , 1997 , 29, 811-812 | 7.4 | 3 |
| 145 | Dialysate sodium delivery can alter chronic blood pressure management. <i>American Journal of Kidney Diseases</i> , 1997 , 29, 383-91 | 7.4 | 73 |
| 144 | Hypertension in hemodialysis patients: more questions than answers. <i>American Journal of Kidney Diseases</i> , 1997 , 30, 577-8 | 7.4 | 5 |
| 143 | The development of concepts of volume control. 1998 , 4, 267-273 | | |
| 142 | Evaluation of the Losartan in Hemodialysis (ELHE) Study. 1998 , 68, S125-9 | | 34 |
| 141 | Drug therapy in haemodialysis patients. Special considerations in the elderly. 1998 , 12, 441-59 | | 19 |
| 140 | Evaluating volume status in hemodialysis patients. 1998 , 5, 64-74 | | 23 |
| 139 | Dialysis schedule-related fluid state and cardiovascular effects. <i>Nephrology Dialysis Transplantation</i> , 1998 , 13 Suppl 6, 91-3 | 4.3 | 7 |
| 138 | Impaired renal adaptation to stress in the elderly with isolated systolic hypertension. 1999 , 34, 1106-11 | | 15 |
| 137 | Supervised atenolol therapy in the management of hemodialysis hypertension. 1999 , 55, 1528-35 | | 75 |
| 136 | Role of angiotensin-converting enzyme inhibitors and angiotensin-receptor blockers in the prevention of progression of renal disease. 1999 , 1, 431-5 | | 2 |

| | | | |
|-----|---|-----|-----|
| 135 | Factors associated with inadequate blood pressure control in hypertensive hemodialysis patients. <i>American Journal of Kidney Diseases</i> , 1999 , 33, 498-506 | 7.4 | 137 |
| 134 | Hypertension in the hemodialysis population? High time for answers. <i>American Journal of Kidney Diseases</i> , 1999 , 33, 592-4 | 7.4 | 28 |
| 133 | Role of home blood pressure monitoring in hemodialysis patients. <i>American Journal of Kidney Diseases</i> , 1999 , 33, 682-7 | 7.4 | 53 |
| 132 | Dialysis Therapy in the United States: A Historical Perspective. <i>Home Hemodialysis International International Symposium on Daily Home Hemodialysis</i> , 1999 , 3, 9-12 | | 7 |
| 131 | Prevalence of hypertension in patients on peritoneal dialysis: results of an Italian multicentre study. <i>Nephrology Dialysis Transplantation</i> , 1999 , 14, 1536-40 | 4.3 | 80 |
| 130 | Is Hypertension Beneficial to the Conventional Hemodialysis Patient?. <i>Hemodialysis International</i> , 2000 , 4, 59-61 | 1.7 | |
| 129 | Hypertension in chronic renal failure and end-stage renal disease patients treated with haemodialysis or peritoneal dialysis. <i>Nephrology Dialysis Transplantation</i> , 2000 , 15 Suppl 5, 105-10 | 4.3 | 21 |
| 128 | Nocturnal blood pressure and 24-hour pulse pressure are potent indicators of mortality in hemodialysis patients. 2000 , 57, 2485-91 | | 184 |
| 127 | Inadequate diagnosis and therapy of arterial hypertension as causes of left ventricular hypertrophy in uremic dialysis patients. 2000 , 58, 260-8 | | 40 |
| 126 | Changes in renal autacoids and hemodynamics associated with aging and isolated systolic hypertension. 2000 , 62, 117-33 | | 5 |
| 125 | Treatment of hypertension in the hemodialysis patient: beneficial or not?. 2000 , 2, 441-4 | | 5 |
| 124 | Prevalence of hospitalization and prognosis of patients on chronic dialysis. 2000 , 4, 236-240 | | 7 |
| 123 | Stroke in Patients Undergoing Dialysis. 2000 , 23, 77-82 | | 7 |
| 122 | Risk for posttransplant diabetes mellitus with current immunosuppressive medications. <i>American Journal of Kidney Diseases</i> , 2000 , 35, 562 | 7.4 | 5 |
| 121 | Reply. <i>American Journal of Kidney Diseases</i> , 2000 , 35, 562 | 7.4 | 0 |
| 120 | Hypertension in the hemodialysis patient. <i>American Journal of Kidney Diseases</i> , 2000 , 35, 563 | 7.4 | |
| 119 | Interdialytic weight gain, compliance with dialysis regimen, and age are independent predictors of blood pressure in hemodialysis patients. <i>American Journal of Kidney Diseases</i> , 2000 , 35, 257-65 | 7.4 | 109 |
| 118 | Influence of atrial fibrillation on the morbido-mortality of patients on hemodialysis. 2000 , 140, 886-90 | | 126 |

| | | | |
|-----|--|-----|-----|
| 117 | I. NKF-K/DOQI Clinical Practice Guidelines for Hemodialysis Adequacy: update 2000. <i>American Journal of Kidney Diseases</i> , 2001 , 37, S7-S64 | 7.4 | 102 |
| 116 | Risk factors for hypertension in chronic hemodialysis patients: baseline data from the HEMO study. 2001 , 21, 280-8 | | 91 |
| 115 | Hypertension in hemodialysis patients. 2001 , 3, 496-502 | | 22 |
| 114 | Prediction of hypertension in chronic hemodialysis patients. 2001 , 60, 1982-9 | | 79 |
| 113 | Lisinopril therapy for hemodialysis hypertension: hemodynamic and endocrine responses. <i>American Journal of Kidney Diseases</i> , 2001 , 38, 1245-50 | 7.4 | 53 |
| 112 | Controversy on optimal blood pressure on haemodialysis: normotensive blood pressure values are essential for survival. <i>Nephrology Dialysis Transplantation</i> , 2001 , 16, 469-74 | 4.3 | 19 |
| 111 | Sympathetic nervous system and chronic renal failure. 2001 , 23, 69-75 | | 16 |
| 110 | The pharmacokinetics and pharmacodynamics of angiotensin-receptor blockers in end-stage renal disease. 2002 , 3, 247-54 | | 19 |
| 109 | Sympathetic overactivity as a cause of hypertension in chronic renal failure. <i>Journal of Hypertension</i> , 2002 , 20, 3-9 | 1.9 | 100 |
| 108 | Long-term control of hypertension in dialysis patients by low dose atenolol. 2002 , 25, 269-75 | | 4 |
| 107 | Hemodialysis-associated hypertension: pathophysiology and therapy. <i>American Journal of Kidney Diseases</i> , 2002 , 39, 227-44 | 7.4 | 166 |
| 106 | A comparison of standardized versus "usual" blood pressure measurements in hemodialysis patients. <i>American Journal of Kidney Diseases</i> , 2002 , 39, 1226-30 | 7.4 | 84 |
| 105 | Sympathetic nervous system function in renal hypertension. 2002 , 4, 229-36 | | 20 |
| 104 | Pulse pressure and risk of total mortality and cardiovascular events in patients on chronic hemodialysis. 2002 , 61, 717-26 | | 130 |
| 103 | A new technique for establishing dry weight in hemodialysis patients via whole body bioimpedance. 2002 , 61, 2250-8 | | 142 |
| 102 | Assessment of blood pressure in hemodialysis patients. <i>Seminars in Dialysis</i> , 2002 , 15, 299-304 | 2.5 | 38 |
| 101 | Effect of hypertension before beginning dialysis on survival of hemodialysis patients. <i>American Journal of Kidney Diseases</i> , 2003 , 41, 814-21 | 7.4 | 27 |
| 100 | Extracellular volume changes and blood pressure levels in hemodialysis patients. <i>Hemodialysis International</i> , 2003 , 7, 135-42 | 1.7 | 5 |

| | | | |
|----|---|-----|-----|
| 99 | Traditional cardiac risk factors in individuals with chronic kidney disease. <i>Seminars in Dialysis</i> , 2003 , 16, 118-27 | 2.5 | 71 |
| 98 | Prevalence of vascular risk factors and vascular disease in predialysis chronic renal failure. 2003 , 8, 274-9 | | 9 |
| 97 | Prevalence, treatment, and control of hypertension in chronic hemodialysis patients in the United States. 2003 , 115, 291-7 | | 346 |
| 96 | Patterns of antihypertensive medication use in hemodialysis patients. 2004 , 61, 1473-8 | | 14 |
| 95 | Blood pressure in haemodialysis patients: the importance of the relationship between the renin-angiotensin-aldosterone system, salt intake and extracellular volume. 2004 , 5, 14-22 | | 7 |
| 94 | Reference values for multifrequency bioimpedance analysis in dialysis patients. 2004 , 22, 301-6 | | 56 |
| 93 | Blood pressure measurement in hemodialysis patients. <i>Seminars in Dialysis</i> , 2004 , 17, 250-4 | 2.5 | 16 |
| 92 | Noncardiac consequences of hypertension in hemodialysis patients. <i>Seminars in Dialysis</i> , 2004 , 17, 304-6 | 2.5 | 11 |
| 91 | Improved blood pressure control with nocturnal hemodialysis: review of clinical observations and physiologic mechanisms. 2004 , 6, 140-4 | | 6 |
| 90 | Underweight rather than overweight is associated with higher prevalence of hypertension: BP vs BMI in haemodialysis population. <i>Nephrology Dialysis Transplantation</i> , 2004 , 19, 427-32 | 4.3 | 22 |
| 89 | Diurnal variation of blood pressure; reproducibility and association with left ventricular hypertrophy in hemodialysis patients. 2005 , 10, 25-32 | | 48 |
| 88 | Opinion: What Clinical Insights from the Early Days of Dialysis Are Being Overlooked Today?. <i>Seminars in Dialysis</i> , 2005 , 18, 19-21 | 2.5 | |
| 87 | Opinion: What Clinical Insights from the Early Days of Dialysis Are Being Overlooked Today?. <i>Seminars in Dialysis</i> , 2005 , 18, 16-18 | 2.5 | |
| 86 | Opinion: What Clinical Insights from the Early Days of Dialysis Are Being Overlooked Today?. <i>Seminars in Dialysis</i> , 2005 , 18, 15-16 | 2.5 | |
| 85 | What clinical insights from the early days of dialysis are being overlooked today?. <i>Seminars in Dialysis</i> , 2005 , 18, 13-21 | 2.5 | 3 |
| 84 | Opinion: What Clinical Insights from the Early Days of Dialysis Are Being Overlooked Today?. <i>Seminars in Dialysis</i> , 2005 , 18, 18-19 | 2.5 | 2 |
| 83 | Hypertension and survival in chronic hemodialysis patients--past lessons and future opportunities. 2005 , 67, 1-13 | | 127 |
| 82 | Interdialytic weight gain as a marker of blood pressure, nutrition, and survival in hemodialysis patients. 2005 , S63-8 | | 80 |

| | | | |
|----|--|-----|-----|
| 81 | Haemodialysis-unresponsive blood pressure: cardiovascular mortality predictor?. 2005 , 10, 438-41 | | 10 |
| 80 | Discordance of influence of hypertension on mortality and cardiovascular risk in hemodialysis patients. <i>American Journal of Kidney Diseases</i> , 2005 , 45, 112-8 | 7.4 | 31 |
| 79 | Hypertension in pediatric patients on long-term dialysis: a report of the North American Pediatric Renal Transplant Cooperative Study (NAPRTCS). <i>American Journal of Kidney Diseases</i> , 2005 , 45, 309-15 | 7.4 | 108 |
| 78 | Effect of spironolactone on blood pressure and the renin-angiotensin-aldosterone system in oligo-anuric hemodialysis patients. <i>American Journal of Kidney Diseases</i> , 2005 , 46, 94-101 | 7.4 | 86 |
| 77 | Blood pressure control before and after starting dialysis. 2005 , 99, c86-91 | | 6 |
| 76 | Blood pressure control in haemodialysis patients: an audit. 2006 , 11, 100-4 | | 11 |
| 75 | Effects of imidapril on left ventricular mass in chronic hemodialysis patients. 2006 , 29, 253-60 | | 29 |
| 74 | Development and validation of bioimpedance analysis prediction equations for dry weight in hemodialysis patients. 2007 , 2, 675-80 | | 29 |
| 73 | Aortic stiffness, left ventricular hypertrophy and weekly averaged blood pressure (WAB) in patients on haemodialysis. <i>Nephrology Dialysis Transplantation</i> , 2007 , 22, 1198-204 | 4.3 | 40 |
| 72 | ACE inhibitor-based, directly observed therapy for hypertension in hemodialysis patients. 2007 , 27, 522-9 | | 12 |
| 71 | Clinical use of laboratory tests for the identification of secondary forms of arterial hypertension. 2007 , 44, 1-85 | | 41 |
| 70 | Increased body mass index is not a reliable marker of good nutrition in hemodialysis patients. <i>Renal Failure</i> , 2007 , 29, 487-93 | 2.9 | 9 |
| 69 | Reducing Mortality in Dialysis Patients: Why Does Hypertension Continue to be Overlooked?. <i>Seminars in Dialysis</i> , 2007 , 10, 250-250 | 2.5 | 6 |
| 68 | Dialysis and Hypertension: Worthy of Investigation?. <i>Seminars in Dialysis</i> , 2007 , 11, 109-112 | 2.5 | |
| 67 | Reversing the risk factor paradox: is daily nocturnal hemodialysis the solution?. <i>Seminars in Dialysis</i> , 2007 , 20, 539-43 | 2.5 | 23 |
| 66 | Blood pressure control in pediatric hemodialysis: the Midwest Pediatric Nephrology Consortium Study. 2007 , 22, 547-53 | | 32 |
| 65 | Thoughts and Progress. <i>Artificial Organs</i> , 2008 , 21, 402-432 | 2.6 | |
| 64 | Antihypertensive prescription in pediatric dialysis: a practitioner survey by the Midwest Pediatric Nephrology Consortium study. <i>Hemodialysis International</i> , 2009 , 13, 307-15 | 1.7 | 12 |

| | | | |
|----|---|------|-----|
| 63 | Blood pressure measurements and carotid intima media thickness in hemodialysis patients. <i>Therapeutic Apheresis and Dialysis</i> , 2009 , 13, 288-93 | 1.9 | 9 |
| 62 | Blood pressure assessment in haemodialysis patients: comparison between pre-dialysis blood pressure and ambulatory blood pressure measurement. 2009 , 14, 283-90 | | 9 |
| 61 | Hypertension in Hemodialysis Patients. 2010 , 6, 44-54 | | 1 |
| 60 | Effects of efonidipine, an L- and T-type calcium channel blocker, on the renin-angiotensin-aldosterone system in chronic hemodialysis patients. 2010 , 51, 188-92 | | 8 |
| 59 | Why assistive technology is needed for probing of dry weight. 2011 , 31, 197-202 | | 17 |
| 58 | Impact of volume status on blood pressure and left ventricle structure in patients undergoing chronic hemodialysis. <i>Renal Failure</i> , 2011 , 33, 377-81 | 2.9 | 12 |
| 57 | Nutritional Intake in Adult Hemodialysis Patients. 2011 , 26, 45-56 | | 16 |
| 56 | Epidemiology of interdialytic ambulatory hypertension and the role of volume excess. 2011 , 34, 381-90 | | 57 |
| 55 | Hypertension and kidney disease: is renalase a new player or an innocent bystander?. <i>Journal of Hypertension</i> , 2012 , 30, 457-62 | 1.9 | 16 |
| 54 | Prognostic value of 48-hour ambulatory blood pressure measurement and cardiovascular mortality in hemodialysis patients. <i>Kidney and Blood Pressure Research</i> , 2012 , 35, 326-31 | 3.1 | 9 |
| 53 | Home systolic blood pressure on the morning of dialysis days has prognostic impact for hypertensive hemodialysis patients. 2012 , 16, 427-32 | | 3 |
| 52 | Blood pressure and blood volume: acute and chronic considerations in hemodialysis. <i>Seminars in Dialysis</i> , 2013 , 26, 62-72 | 2.5 | 9 |
| 51 | Long-term antihypertensive effects of aliskiren, a direct renin inhibitor, in chronic hemodialysis patients. <i>Therapeutic Apheresis and Dialysis</i> , 2013 , 17, 524-31 | 1.9 | 3 |
| 50 | Effects of nutritional parameters on nocturnal blood pressure in patients undergoing hemodialysis. <i>Renal Failure</i> , 2013 , 35, 946-50 | 2.9 | 6 |
| 49 | Which fluid space is affected by ultrafiltration during hemodiafiltration?. <i>Hemodialysis International</i> , 2014 , 18, 384-90 | 1.7 | 17 |
| 48 | Assessment and management of hypertension in patients on dialysis. <i>Journal of the American Society of Nephrology: JASN</i> , 2014 , 25, 1630-46 | 12.7 | 106 |
| 47 | Identification of impeding factors for dry weight achievement in end-stage renal disease after appropriate kidney graft function. <i>Artificial Organs</i> , 2014 , 38, 113-20 | 2.6 | 2 |
| 46 | Does pharmacotherapy improve cardiovascular outcomes in hemodialysis patients?. <i>Hemodialysis International</i> , 2015 , 19 Suppl 3, S40-50 | 1.7 | 1 |

| | | | |
|----|--|------|----|
| 45 | Relationship between different blood pressure measurements and left ventricular mass by cardiac magnetic resonance imaging in end-stage renal disease. <i>Journal of the American Society of Hypertension</i> , 2015 , 9, 275-84 | | 14 |
| 44 | Epidemiology, diagnosis and management of hypertension among patients on chronic dialysis. <i>Nature Reviews Nephrology</i> , 2016 , 12, 636-47 | 14.9 | 35 |
| 43 | Hypertension in the Hemodialysis Patient. <i>Advances in Experimental Medicine and Biology</i> , 2017 , 956, 327-340 | 3.6 | 3 |
| 42 | Hypertension in the Dialysis Patient. 2016 , 133-166 | | |
| 41 | Hypertension in dialysis patients: a consensus document by the European Renal and Cardiovascular Medicine (EURECA-m) working group of the European Renal Association-European Dialysis and Transplant Association (ERA-EDTA) and the Hypertension and the Kidney working group of the European Society of Hypertension (ESH). <i>Nephrology Dialysis Transplantation</i> , 2017 , 32, 620-640 | 4.3 | 79 |
| 40 | Hypertension in dialysis patients: a consensus document by the European Renal and Cardiovascular Medicine (EURECA-m) working group of the European Renal Association - European Dialysis and Transplant Association (ERA-EDTA) and the Hypertension and the Kidney working group of the European Society of Hypertension (ESH). <i>Journal of Hypertension</i> , 2017 , 35, 457-476 | 1.9 | 35 |
| 39 | The Combination of Beta Blockers and Renin-Angiotensin System Blockers Improves Survival in Incident Hemodialysis Patients: A Propensity-Matched Study. <i>Kidney International Reports</i> , 2017 , 2, 665-675 | 4.1 | 2 |
| 38 | N-Terminal Prohormone of Brain Natriuretic Peptide but not C-Terminal Pre-Pro Vasopressin (Copeptin) Level is Associated with the Response to Antihypertensive Therapy in Haemodialysis Patients. <i>Kidney and Blood Pressure Research</i> , 2017 , 42, 1013-1022 | 3.1 | 3 |
| 37 | Mechanisms and mediators of hypertension induced by erythropoietin and related molecules. <i>Nephrology Dialysis Transplantation</i> , 2018 , 33, 1690-1698 | 4.3 | 18 |
| 36 | Aldosterone antagonists for people with chronic kidney disease requiring dialysis. <i>The Cochrane Library</i> , 2018 , | 5.2 | 1 |
| 35 | Hypertension and Chronic Kidney Disease. 2018 , 311-320 | | |
| 34 | Prevalence and control of hypertension by 48-h ambulatory blood pressure monitoring in haemodialysis patients: a study by the European Cardiovascular and Renal Medicine (EURECA-m) working group of the ERA-EDTA. <i>Nephrology Dialysis Transplantation</i> , 2019 , 34, 1542-1548 | 4.3 | 12 |
| 33 | Systolic and diastolic hypertension among patients on hemodialysis: Musings on volume overload, arterial stiffness, and erythropoietin. <i>Seminars in Dialysis</i> , 2019 , 32, 507-512 | 2.5 | 2 |
| 32 | Development of an Artificial Intelligence Model to Guide the Management of Blood Pressure, Fluid Volume, and Dialysis Dose in End-Stage Kidney Disease Patients: Proof of Concept and First Clinical Assessment. <i>Kidney Diseases (Basel, Switzerland)</i> , 2019 , 5, 28-33 | 3.3 | 36 |
| 31 | Association of body mass index and uncontrolled blood pressure with cardiovascular mortality in peritoneal dialysis patients. <i>Journal of Human Hypertension</i> , 2019 , 33, 106-114 | 2.6 | 1 |
| 30 | Hypertension in patients on dialysis: diagnosis, mechanisms, and management. <i>Jornal Brasileiro De Nefrologia: Orgao Oficial De Sociedades Brasileira E Latino-Americana De Nefrologia</i> , 2019 , 41, 400-411 | 1.5 | 13 |
| 29 | Increased Weight Gain During the Long Interdialytic Period Is Associated with Minor Effects on Blood Pressure Control in Clinically Stable In-Centre Haemodialysis Patients. <i>Nephron</i> , 2019 , 141, 87-97 | 3.3 | 3 |
| 28 | Evaluation of management and factors associated with hypertension control in hemodialysis patients at a tertiary-care hospital in Pakistan. <i>Drugs and Therapy Perspectives</i> , 2020 , 36, 396-403 | 1.5 | 2 |

| | | | |
|----|--|-----|----|
| 27 | Prevalence and control of hypertension among patients on haemodialysis. <i>European Journal of Clinical Investigation</i> , 2020 , 50, e13292 | 4.6 | 1 |
| 26 | Blood pressure targets and pharmacotherapy for hypertensive patients on hemodialysis. <i>Expert Opinion on Pharmacotherapy</i> , 2020 , 21, 1219-1240 | 4 | 0 |
| 25 | Role of Artificial Intelligence in Kidney Disease. <i>International Journal of Medical Sciences</i> , 2020 , 17, 970-984 | 3.7 | 11 |
| 24 | Usefulness of ABPM and bioimpedance for the treatment and control of hypertension in patients on chronic haemodialysis. <i>Nefrologia</i> , 2021 , 41, 17-26 | 1.5 | 1 |
| 23 | Usefulness of ABPM and bioimpedance for the treatment and control of hypertension in patients on chronic haemodialysis. <i>Nefrologia</i> , 2021 , 41, 17-26 | 0.4 | |
| 22 | Aldosterone antagonists for people with chronic kidney disease requiring dialysis. <i>The Cochrane Library</i> , 2021 , 2, CD013109 | 5.2 | 3 |
| 21 | Pathophysiology of hypertension in renal failure. <i>Seminars in Nephrology</i> , 2002 , 22, 17-26 | 4.8 | 16 |
| 20 | Prevalence of hypertension among Iranian hemodialysis patients and associated risk factors: a nationwide multicenter study. <i>Pakistan Journal of Biological Sciences</i> , 2008 , 11, 910-4 | 0.8 | 3 |
| 19 | Hypertension in Renal Failure. 2007 , 693-705 | | |
| 18 | Hypertensive Emergencies. 2008 , 624-634 | | |
| 17 | Acute Dialysis with the GENIUS System. 2009 , 1299-1305 | | |
| 16 | Antihypertensive agents for dialysis patients. | | |
| 15 | Akute Probleme während der Dialysebehandlung. 2011 , 133-143 | | |
| 14 | Akute Probleme während der Dialysebehandlung. 2014 , 139-149 | | |
| 13 | Antihypertensive agents for dialysis patients. | | |
| 12 | Hypertension in Dialysis Patients: Clinical Epidemiology, Pathogenesis, Diagnosis, and Treatment. <i>Updates in Hypertension and Cardiovascular Protection</i> , 2018 , 383-417 | 0.1 | |
| 11 | Dialysis Therapy in the United States: A Historical Perspective. <i>Home Hemodialysis International International Symposium on Daily Home Hemodialysis</i> , 1999 , 3, 9-12 | | |
| 10 | Is Hypertension Beneficial to the Conventional Hemodialysis Patient?. <i>Hemodialysis International</i> , 2000 , 4, 59-61 | 1.7 | |

| | | | |
|---|--|-----|----|
| 9 | The management of hypertension in hemodialysis and CAPD patients. <i>Hippokratia</i> , 2007 , 11, 171-4 | 0.4 | 3 |
| 8 | A PDA-based dietary self-monitoring intervention to reduce sodium intake in an in-center hemodialysis patient. <i>Patient Preference and Adherence</i> , 2008 , 2, 177-84 | 2.4 | 12 |
| 7 | The association between blood pressure level and serum uric acid concentration in hemodialysis patients. <i>Journal of Nephropathology</i> , 2015 , 4, 85-90 | 0.6 | 5 |
| 6 | Effects of Mean Artery Pressure and Blood pH on Survival Rate of Patients with Acute Kidney Injury Combined with Acute Hypoxic Respiratory Failure: A Retrospective Study. <i>Medicina (Lithuania)</i> , 2021 , 57, | 3.1 | 1 |
| 5 | [Out-of-center blood pressure measurements in dialysis patients: Feasibility and comparison of methods].. <i>Nephrologie Et Therapeutique</i> , 2022 , | 0.6 | |
| 4 | Safety and Efficacy of Spironolactone in Dialysis-Dependent Patients: Meta-Analysis of Randomized Controlled Trials.. <i>Frontiers in Medicine</i> , 2022 , 9, 828189 | 4.9 | |
| 3 | The Relationship Between Pulse Pressure and Hypervolemia in Hemodialysis Patients. <i>Medical Journal of Western Black Sea</i> , 2021 , 5, 380-385 | 3.5 | |
| 2 | Hypertension in Dialysis Patients: Diagnostic Approaches and Evaluation of Epidemiology. 2022 , 12, 2961 | | 0 |
| 1 | Extended-Hours Hemodialysis without Dietary Restrictions Increases Body Mass Index and Normalizes Hypertension: A Case Report. 2023 , | | 0 |