

# CITATION REPORT

List of articles citing

Poly(rC) binding protein 2 binds to stem-loop IV of the poliovirus RNA 5' noncoding region: identification by automated liquid chromatography-tandem mass spectrometry

DOI: 10.1073/pnas.93.20.11115

Proceedings of the National Academy of Sciences of the United States of America, 1996, 93, 11115-20.

Source: <https://exaly.com/paper-pdf/26991746/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
184	RNA Determinants of Picornavirus Cap-Independent Translation Initiation. <b>1997</b> , 8, 242-255		49
183	RNA Signals in Enterovirus and Rhinovirus Genome Replication. <b>1997</b> , 8, 256-273		44
182	Variable flow liquid chromatography-tandem mass spectrometry and the comprehensive analysis of complex protein digest mixtures. <b>1997</b> , 8, 1059-1069		75
181	Switch from translation to RNA replication in a positive-stranded RNA virus. <i>Genes and Development</i> , <b>1998</b> , 12, 2293-304	12.6	391
180	Cellular factors in the transcription and replication of viral RNA genomes: a parallel to DNA-dependent RNA transcription. <i>Virology</i> , <b>1998</b> , 244, 1-12	3.6	240
179	Cytoplasmic regulatory functions of the KH-domain proteins hnRNPs K and E1/E2. <b>1998</b> , 23, 409-11		152
178	The identification of peptide modifications derived from gel-separated proteins using electrospray triple quadrupole and ion trap analyses. <b>1998</b> , 19, 989-97		62
177	Rapid protein identification using a microscale electrospray LC/MS system on an ion trap mass spectrometer. <b>1998</b> , 9, 194-201		124
176	MgATP binding to the nucleotide-binding domains of the eukaryotic cytoplasmic chaperonin induces conformational changes in the putative substrate-binding domains. <b>1998</b> , 7, 1524-30		24
175	Mass spectrometry. <b>1998</b> , 70, 647R-716R		185
174	Molecular aspects of poliovirus biology with a special focus on the interactions with nerve cells. <b>1998</b> , 4, 1-26		34
173	Genetics, Pathogenesis and Evolution of Picornaviruses. <b>1999</b> , 287-343		45
172	Polypyrimidine-tract binding protein (PTB) is necessary, but not sufficient, for efficient internal initiation of translation of human rhinovirus-2 RNA. <i>Rna</i> , <b>1999</b> , 5, 344-59	5.8	122
171	Differential utilization of poly(rC) binding protein 2 in translation directed by picornavirus IRES elements. <i>Rna</i> , <b>1999</b> , 5, 1570-85	5.8	120
170	Structure and function of a small RNA that selectively inhibits internal ribosome entry site-mediated translation. <i>Nucleic Acids Research</i> , <b>1999</b> , 27, 562-72	20.1	14
169	Properties of cloned and expressed human RNase H1. <i>Journal of Biological Chemistry</i> , <b>1999</b> , 274, 28270-8	5.4	125
168	The N-terminal K homology domain of the poly(rC)-binding protein is a major determinant for binding to the poliovirus 5' untranslated region and acts as an inhibitor of viral translation. <i>Journal of Biological Chemistry</i> , <b>1999</b> , 274, 38163-70	5.4	59

167	Intracellular determinants of picornavirus replication. <b>1999</b> , 7, 76-82		73
166	Paradoxes of the replication of picornaviral genomes. <i>Virus Research</i> , <b>1999</b> , 62, 129-47	6.4	96
165	Rhinovirus 2A proteinase mediated stimulation of rhinovirus RNA translation is additive to the stimulation effected by cellular RNA binding proteins. <i>Virus Research</i> , <b>1999</b> , 62, 119-28	6.4	16
164	eIF4 initiation factors: effectors of mRNA recruitment to ribosomes and regulators of translation. <b>1999</b> , 68, 913-63		1729
163	unr, a cellular cytoplasmic RNA-binding protein with five cold-shock domains, is required for internal initiation of translation of human rhinovirus RNA. <i>Genes and Development</i> , <b>1999</b> , 13, 437-48	12.6	199
162	The RNA encompassing the internal ribosome entry site in the poliovirus 5' untranslated region enhances the encapsidation of genomic RNA. <i>Virology</i> , <b>2000</b> , 273, 391-9	3.6	17
161	Internal ribosome entry sites of viral and cellular RNAs. <b>2000</b> , 34, 157-167		1
160	Identification of an RNA hairpin in poliovirus RNA that serves as the primary template in the in vitro uridylylation of VPg. <i>Journal of Virology</i> , <b>2000</b> , 74, 10359-70	6.6	244
159	A group B coxsackievirus/poliovirus 5' untranslated region chimera can act as an attenuated vaccine strain in mice. <i>Journal of Virology</i> , <b>2000</b> , 74, 4047-56	6.6	46
158	Genetic analysis of a poliovirus/hepatitis C virus (HCV) chimera: interaction between the poliovirus cloverleaf and a sequence in the HCV 5' untranslated region results in a replication phenotype. <i>Journal of Virology</i> , <b>2000</b> , 74, 6223-6	6.6	20
157	Interactions of viral protein 3CD and poly(rC) binding protein with the 5' untranslated region of the poliovirus genome. <i>Journal of Virology</i> , <b>2000</b> , 74, 2219-26	6.6	191
156	Identification of two novel mammalian genes establishes a subfamily of KH-domain RNA-binding proteins. <b>2000</b> , 67, 301-16		49
155	Translational control of viral gene expression in eukaryotes. <b>2000</b> , 64, 239-80		250
154	Internal ribosome entry sites in eukaryotic mRNA molecules. <i>Genes and Development</i> , <b>2001</b> , 15, 1593-612	2.6	759
153	Interaction of poly(rC)-binding protein 2 with the 5' terminal stem loop of the hepatitis C-virus genome. <i>Virus Research</i> , <b>2001</b> , 73, 67-79	6.4	50
152	Poly(rC) binding proteins mediate poliovirus mRNA stability. <i>Rna</i> , <b>2001</b> , 7, 1126-41	5.8	61
151	A novel protein-RNA binding assay: functional interactions of the foot-and-mouth disease virus internal ribosome entry site with cellular proteins. <i>Rna</i> , <b>2001</b> , 7, 114-22	5.8	30
150	Translational Control of the Picornavirus Phenotype. <b>2001</b> , 35, 591-599		5

149	Cell-specific proteins regulate viral RNA translation and virus-induced disease. <i>EMBO Journal</i> , <b>2001</b> , 20, 6899-908	13	90
148	5Qloverleaf in poliovirus RNA is a cis-acting replication element required for negative-strand synthesis. <i>EMBO Journal</i> , <b>2001</b> , 20, 1439-48	13	222
147	Translation of the human c-myc P0 tricistronic mRNA involves two independent internal ribosome entry sites. <b>2001</b> , 20, 4270-80		27
146	Internal ribosome entry site-mediated translation of a mammalian mRNA is regulated by amino acid availability. <i>Journal of Biological Chemistry</i> , <b>2001</b> , 276, 12285-91	5.4	124
145	Investigating the structure of human RNase H1 by site-directed mutagenesis. <i>Journal of Biological Chemistry</i> , <b>2001</b> , 276, 23547-53	5.4	57
144	A predicted secondary structural domain within the internal ribosome entry site of echovirus 12 mediates a cell-type-specific block to viral replication. <i>Journal of Virology</i> , <b>2001</b> , 75, 6472-81	6.6	22
143	Distinct poly(rC) binding protein KH domain determinants for poliovirus translation initiation and viral RNA replication. <i>Journal of Virology</i> , <b>2002</b> , 76, 12008-22	6.6	110
142	Long-term circulation of vaccine-derived poliovirus that causes paralytic disease. <i>Journal of Virology</i> , <b>2002</b> , 76, 6791-9	6.6	79
141	Cell proteins TIA-1 and TIAR interact with the 3Qstem-loop of the West Nile virus complementary minus-strand RNA and facilitate virus replication. <i>Journal of Virology</i> , <b>2002</b> , 76, 11989-2000	6.6	167
140	Translation of polioviral mRNA is inhibited by cleavage of polypyrimidine tract-binding proteins executed by polioviral 3C(pro). <i>Journal of Virology</i> , <b>2002</b> , 76, 2529-42	6.6	120
139	The poly(C)-binding proteins: a multiplicity of functions and a search for mechanisms. <i>Rna</i> , <b>2002</b> , 8, 265-788		335
138	Expression of poly(C)-binding proteins is differentially regulated by hypoxia and ischemia in cortical neurons. <b>2002</b> , 110, 191-8		29
137	Recombination of poliovirus RNA proceeds in mixed replication complexes originating from distinct replication start sites. <i>Journal of Virology</i> , <b>2002</b> , 76, 10960-71	6.6	39
136	Determinants of the recognition of enteroviral cloverleaf RNA by coxsackievirus B3 proteinase 3C. <i>Rna</i> , <b>2002</b> , 8, 188-201	5.8	45
135	A tenascin-C aptamer identified by tumor cell SELEX: systematic evolution of ligands by exponential enrichment. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2003</b> , 100, 15416-21	11.5	474
134	Activity of a type 1 picornavirus internal ribosomal entry site is determined by sequences within the 3Qnontranslated region. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2003</b> , 100, 15125-30	11.5	53
133	Identification of mRNAs associated with alphaCP2-containing RNP complexes. <b>2003</b> , 23, 7055-67		62
132	Long-range RNA-RNA interaction between the 5Qnontranslated region and the core-coding sequences of hepatitis C virus modulates the IRES-dependent translation. <i>Rna</i> , <b>2003</b> , 9, 599-606	5.8	63

131	Analysis of the cloverleaf element in a human rhinovirus type 14/poliovirus chimera: correlation of subdomain D structure, ternary protein complex formation and virus replication. <b>2003</b> , 84, 2203-2216		28
130	Functional dissection of a poliovirus cis-acting replication element [PV-cre(2C)]: analysis of single- and dual-cre viral genomes and proteins that bind specifically to PV-cre RNA. <i>Journal of Virology</i> , <b>2003</b> , 77, 5152-66	6.6	73
129	Molecular mechanisms of attenuation of the Sabin strain of poliovirus type 3. <i>Journal of Virology</i> , <b>2004</b> , 78, 11097-107	6.6	79
128	Specific recognition of the C-rich strand of human telomeric DNA and the RNA template of human telomerase by the first KH domain of human poly(C)-binding protein-2. <i>Journal of Biological Chemistry</i> , <b>2004</b> , 279, 48126-34	5.4	16
127	20S proteasome differentially alters translation of different mRNAs via the cleavage of eIF4F and eIF3. <b>2004</b> , 16, 575-86		68
126	Formation of an alphaCP1-KH3 complex with UC-rich RNA. <b>2005</b> , 34, 423-9		3
125	Hepatitis C virus internal ribosome entry site-dependent translation in <i>Saccharomyces cerevisiae</i> is independent of polypyrimidine tract-binding protein, poly(rC)-binding protein 2, and La protein. <i>Journal of Virology</i> , <b>2005</b> , 79, 10126-37	6.6	23
124	Functional interaction of heterogeneous nuclear ribonucleoprotein C with poliovirus RNA synthesis initiation complexes. <i>Journal of Virology</i> , <b>2005</b> , 79, 3254-66	6.6	77
123	The polypyrimidine tract binding protein is required for efficient picornavirus gene expression and propagation. <i>Journal of Virology</i> , <b>2005</b> , 79, 6172-9	6.6	37
122	Structure and RNA binding of the third KH domain of poly(C)-binding protein 1. <i>Nucleic Acids Research</i> , <b>2005</b> , 33, 1213-21	20.1	22
121	Crystal structure of the first KH domain of human poly(C)-binding protein-2 in complex with a C-rich strand of human telomeric DNA at 1.7 Å. <i>Journal of Biological Chemistry</i> , <b>2005</b> , 280, 38823-30	5.4	47
120	Amino acid changes in proteins 2B and 3A mediate rhinovirus type 39 growth in mouse cells. <i>Journal of Virology</i> , <b>2005</b> , 79, 5363-73	6.6	37
119	Two cellular proteins that interact with a stem loop in the simian hemorrhagic fever virus 3' UTR RNA. <i>Virus Research</i> , <b>2005</b> , 109, 109-24	6.4	14
118	End-to-end communication in the modulation of translation by mammalian RNA viruses. <i>Virus Research</i> , <b>2006</b> , 119, 43-51	6.4	50
117	Translational control by viral proteinases. <i>Virus Research</i> , <b>2006</b> , 119, 76-88	6.4	100
116	Internal initiation: IRES elements of picornaviruses and hepatitis c virus. <i>Virus Research</i> , <b>2006</b> , 119, 2-15	6.4	60
115	Polyadenylation of genomic RNA and initiation of antigenomic RNA in a positive-strand RNA virus are controlled by the same cis-element. <i>Nucleic Acids Research</i> , <b>2006</b> , 34, 2953-65	20.1	42
114	Replication of poliovirus requires binding of the poly(rC) binding protein to the cloverleaf as well as to the adjacent C-rich spacer sequence between the cloverleaf and the internal ribosomal entry site. <i>Journal of Virology</i> , <b>2007</b> , 81, 10017-28	6.6	62

113	Crystal structure of the third KH domain of human poly(C)-binding protein-2 in complex with a C-rich strand of human telomeric DNA at 1.6 Å resolution. <i>Nucleic Acids Research</i> , <b>2007</b> , 35, 2651-60	20.1	28
112	X-ray crystallographic and NMR studies of protein-protein and protein-nucleic acid interactions involving the KH domains from human poly(C)-binding protein-2. <i>Rna</i> , <b>2007</b> , 13, 1043-51	5.8	34
111	Cellular protein modification by poliovirus: the two faces of poly(rC)-binding protein. <i>Journal of Virology</i> , <b>2007</b> , 81, 8919-32	6.6	116
110	The myotonic dystrophy type 2 protein ZNF9 is part of an ITAF complex that promotes cap-independent translation. <b>2007</b> , 6, 1049-58		47
109	Epidemics to eradication: the modern history of poliomyelitis. <b>2007</b> , 4, 70		57
108	hnRNP E1 and E2 have distinct roles in modulating HIV-1 gene expression. <b>2007</b> , 4, 28		38
107	A nucleo-cytoplasmic SR protein functions in viral IRES-mediated translation initiation. <i>EMBO Journal</i> , <b>2007</b> , 26, 459-67	13	139
106	Protein-RNA tethering: the role of poly(C) binding protein 2 in poliovirus RNA replication. <i>Virology</i> , <b>2008</b> , 374, 280-91	3.6	22
105	The linker domain of poly(rC) binding protein 2 is a major determinant in poliovirus cap-independent translation. <i>Virology</i> , <b>2008</b> , 378, 243-53	3.6	33
104	Poly(rC)-binding protein 2 interacts with the oligo(rC) tract of coxsackievirus B3. <b>2008</b> , 366, 917-21		28
103	Interaction of poly(rC)-binding protein 2 domains KH1 and KH3 with coxsackievirus RNA. <b>2008</b> , 377, 500-503		9
102	Modulation of enteroviral proteinase cleavage of poly(A)-binding protein (PABP) by conformation and PABP-associated factors. <i>Virology</i> , <b>2008</b> , 375, 59-72	3.6	28
101	Structure of a construct of a human poly(C)-binding protein containing the first and second KH domains reveals insights into its regulatory mechanisms. <i>Journal of Biological Chemistry</i> , <b>2008</b> , 283, 28757-66	5.4	42
100	Poly(rC) binding proteins and the 5'UTR of uncapped poliovirus mRNA function during de novo assembly of polysomes. <i>Journal of Virology</i> , <b>2008</b> , 82, 5835-46	6.6	14
99	Genetic interactions between an essential 3' cis-acting RNA pseudoknot, replicase gene products, and the extreme 3' end of the mouse coronavirus genome. <i>Journal of Virology</i> , <b>2008</b> , 82, 1214-28	6.6	71
98	Cleavage of poly(A)-binding protein by poliovirus 3C proteinase inhibits viral internal ribosome entry site-mediated translation. <i>Journal of Virology</i> , <b>2008</b> , 82, 9389-99	6.6	48
97	Introduction: RNA viruses. 1-2		
96	Neurotropic picornaviruses. 3-25		1

95	Depletion of the poly(C)-binding proteins alphaCP1 and alphaCP2 from K562 cells leads to p53-independent induction of cyclin-dependent kinase inhibitor (CDKN1A) and G1 arrest. <i>Journal of Biological Chemistry</i> , <b>2009</b> , 284, 9039-49	5.4	50
94	SYNCRIP (synaptotagmin-binding, cytoplasmic RNA-interacting protein) is a host factor involved in hepatitis C virus RNA replication. <i>Virology</i> , <b>2009</b> , 386, 249-56	3.6	33
93	Altered interactions between stem-loop IV within the 5' noncoding region of coxsackievirus RNA and poly(rC) binding protein 2: effects on IRES-mediated translation and viral infectivity. <i>Virology</i> , <b>2009</b> , 389, 45-58	3.6	55
92	Viral and host proteins involved in picornavirus life cycle. <i>Journal of Biomedical Science</i> , <b>2009</b> , 16, 103	13.3	128
91	PCBP2 mediates degradation of the adaptor MAVS via the HECT ubiquitin ligase AIP4. <b>2009</b> , 10, 1300-8		233
90	Inhibition of hepatitis C virus IRES-mediated translation by oligonucleotides. <i>Virus Research</i> , <b>2009</b> , 146, 29-35	6.4	4
89	The 5' 5C-L-PCBP RNP complex, 3' poly(A) tail and 2A(pro) are required for optimal translation of poliovirus RNA. <i>Virology</i> , <b>2010</b> , 397, 14-22	3.6	18
88	Mechanistic consequences of hnRNP C binding to both RNA termini of poliovirus negative-strand RNA intermediates. <i>Journal of Virology</i> , <b>2010</b> , 84, 4229-42	6.6	49
87	Mechanistic intersections between picornavirus translation and RNA replication. <b>2011</b> , 80, 1-24		22
86	Diverse roles of host RNA binding proteins in RNA virus replication. <i>RNA Biology</i> , <b>2011</b> , 8, 305-15	4.8	101
85	Poliovirus-mediated disruption of cytoplasmic processing bodies. <i>Journal of Virology</i> , <b>2011</b> , 85, 64-75	6.6	86
84	SARS coronavirus nsp1 protein induces template-dependent endonucleolytic cleavage of mRNAs: viral mRNAs are resistant to nsp1-induced RNA cleavage. <i>PLoS Pathogens</i> , <b>2011</b> , 7, e1002433	7.6	224
83	Re-localization of cellular protein SRp20 during poliovirus infection: bridging a viral IRES to the host cell translation apparatus. <i>PLoS Pathogens</i> , <b>2011</b> , 7, e1002127	7.6	44
82	Viral subversion of host functions for picornavirus translation and RNA replication. <i>Future Virology</i> , <b>2012</b> , 7, 179-191	2.4	42
81	Picornavirus modification of a host mRNA decay protein. <i>MBio</i> , <b>2012</b> , 3, e00431-12	7.8	45
80	Contribution of the first K-homology domain of poly(C)-binding protein 1 to its affinity and specificity for C-rich oligonucleotides. <i>Nucleic Acids Research</i> , <b>2012</b> , 40, 5101-14	20.1	29
79	Genetic characterization of enterovirus 71 isolated from patients with severe disease by comparative analysis of complete genomes. <i>Journal of Medical Virology</i> , <b>2012</b> , 84, 931-9	19.7	42
78	The 40-80 nt region in the 5' NCR of genome is a critical target for inactivating poliovirus by chlorine dioxide. <i>Journal of Medical Virology</i> , <b>2012</b> , 84, 526-35	19.7	14



77	Thiouracil cross-linking mass spectrometry: a cell-based method to identify host factors involved in viral amplification. <i>Journal of Virology</i> , <b>2013</b> , 87, 8697-712	6.6	35
76	Viral proteinase requirements for the nucleocytoplasmic relocalization of cellular splicing factor SRp20 during picornavirus infections. <i>Journal of Virology</i> , <b>2013</b> , 87, 2390-400	6.6	31
75	EWSR1 binds the hepatitis C virus cis-acting replication element and is required for efficient viral replication. <i>Journal of Virology</i> , <b>2013</b> , 87, 6625-34	6.6	30
74	Norovirus genome circularization and efficient replication are facilitated by binding of PCBP2 and hnRNP A1. <i>Journal of Virology</i> , <b>2013</b> , 87, 11371-87	6.6	25
73	Novel function of the poly(c)-binding protein ECP2 as a transcriptional activator that binds to single-stranded DNA sequences. <i>International Journal of Molecular Medicine</i> , <b>2013</b> , 32, 1187-94	4.4	2
72	Optimization of chemiluminescent detection of mitochondrial RNA-protein interaction by nonradioactive mobility shift. <i>African Journal of Biotechnology</i> , <b>2013</b> , 12, 1915-1920	0.6	3
71	A cytoplasmic RNA virus generates functional viral small RNAs and regulates viral IRES activity in mammalian cells. <i>Nucleic Acids Research</i> , <b>2014</b> , 42, 12789-805	20.1	39
70	The mechanism of translation initiation on Type 1 picornavirus IRESs. <i>EMBO Journal</i> , <b>2014</b> , 33, 76-92	13	103
69	Differential cleavage of IRES trans-acting factors (ITAFs) in cells infected by human rhinovirus. <i>Virology</i> , <b>2014</b> , 449, 35-44	3.6	8
68	Mitogen-activated protein kinase-interacting kinase regulates mTOR/AKT signaling and controls the serine/arginine-rich protein kinase-responsive type 1 internal ribosome entry site-mediated translation and viral oncolysis. <i>Journal of Virology</i> , <b>2014</b> , 88, 13149-60	6.6	32
67	Induction of viral, 7-methyl-guanosine cap-independent translation and oncolysis by mitogen-activated protein kinase-interacting kinase-mediated effects on the serine/arginine-rich protein kinase. <i>Journal of Virology</i> , <b>2014</b> , 88, 13135-48	6.6	29
66	Inhibition of poliovirus-induced cleavage of cellular protein PCBP2 reduces the levels of viral RNA replication. <i>Journal of Virology</i> , <b>2014</b> , 88, 3192-201	6.6	31
65	Picornaviruses and nuclear functions: targeting a cellular compartment distinct from the replication site of a positive-strand RNA virus. <i>Frontiers in Microbiology</i> , <b>2015</b> , 6, 594	5.7	57
64	Initiation of protein-primed picornavirus RNA synthesis. <i>Virus Research</i> , <b>2015</b> , 206, 12-26	6.4	55
63	Iron Toxicity in the Retina Requires Alu RNA and the NLRP3 Inflammasome. <i>Cell Reports</i> , <b>2015</b> , 11, 1686-93.6	93.6	54
62	Cytotoxic and immunogenic mechanisms of recombinant oncolytic poliovirus. <i>Current Opinion in Virology</i> , <b>2015</b> , 13, 81-5	7.5	30
61	The role of Misshapen NCK-related kinase (MINK), a novel Ste20 family kinase, in the IRES-mediated protein translation of human enterovirus 71. <i>PLoS Pathogens</i> , <b>2015</b> , 11, e1004686	7.6	28
60	AtLa1 protein initiates IRES-dependent translation of WUSCHEL mRNA and regulates the stem cell homeostasis of Arabidopsis in response to environmental hazards. <i>Plant, Cell and Environment</i> , <b>2015</b> , 38, 2098-114	8.4	27



59	Nuclear proteins hijacked by mammalian cytoplasmic plus strand RNA viruses. <i>Virology</i> , <b>2015</b> , 479-480, 457-74	3.6	46
58	Mutational robustness and resilience of a replicative cis-element of RNA virus: Promiscuity, limitations, relevance. <i>RNA Biology</i> , <b>2015</b> , 12, 1338-54	4.8	8
57	Enterovirus Control of Translation and RNA Granule Stress Responses. <i>Viruses</i> , <b>2016</b> , 8, 93	6.2	23
56	Diverse Strategies Used by Picornaviruses to Escape Host RNA Decay Pathways. <i>Viruses</i> , <b>2016</b> , 8,	6.2	12
55	Poliovirus. <b>2016</b> , 1-26		
54	Divergent Requirement for a DNA Repair Enzyme during Enterovirus Infections. <i>MBio</i> , <b>2015</b> , 7, e01931-15	3.8	10
53	Structure of RNA Stem Loop B from the Picornavirus Replication Platform. <i>Biochemistry</i> , <b>2017</b> , 56, 2549-2557	3.5	5
52	Regulation of BC200 RNA-mediated translation inhibition by hnRNP E1 and E2. <i>FEBS Letters</i> , <b>2017</b> , 591, 393-405	3.8	17
51	Control of the negative IRES trans-acting factor KHSRP by ubiquitination. <i>Nucleic Acids Research</i> , <b>2017</b> , 45, 271-287	20.1	121
50	Role of MAPK/MNK1 signaling in virus replication. <i>Virus Research</i> , <b>2018</b> , 253, 48-61	6.4	56
49	Stimulation of the Internal Ribosome Entry Site (IRES)-Dependent Translation of Enterovirus 71 by DDX3X RNA Helicase and Viral 2A and 3C Proteases. <i>Frontiers in Microbiology</i> , <b>2018</b> , 9, 1324	5.7	18
48	Exploitation of nuclear functions by human rhinovirus, a cytoplasmic RNA virus. <i>PLoS Pathogens</i> , <b>2018</b> , 14, e1007277	7.6	8
47	Picornaviruses and RNA Metabolism: Local and Global Effects of Infection. <i>Journal of Virology</i> , <b>2019</b> , 93,	6.6	5
46	Strand-specific affinity of host factor hnRNP C1/C2 guides positive to negative-strand ratio in Coxsackievirus B3 infection. <i>RNA Biology</i> , <b>2019</b> , 16, 1286-1299	4.8	7
45	eIF4G2 balances its own mRNA translation via a PCBP2-based feedback loop. <i>Rna</i> , <b>2019</b> , 25, 757-767	5.8	4
44	Staufen1 Protein Participates Positively in the Viral RNA Replication of Enterovirus 71. <i>Viruses</i> , <b>2019</b> , 11,	6.2	8
43	Conformational flexibility in the enterovirus RNA replication platform. <i>Rna</i> , <b>2019</b> , 25, 376-387	5.8	5
42	Translation control of Enterovirus A71 gene expression. <i>Journal of Biomedical Science</i> , <b>2020</b> , 27, 22	13.3	2

41	PCBP1 and PCBP2 both bind heavily oxidized RNA but cause opposing outcomes, suppressing or increasing apoptosis under oxidative conditions. <i>Journal of Biological Chemistry</i> , <b>2020</b> , 295, 12247-12261	5.4	8
40	Structure of the PCBP2/stem-loop IV complex underlying translation initiation mediated by the poliovirus type I IRES. <i>Nucleic Acids Research</i> , <b>2020</b> , 48, 8006-8021	20.1	6
39	ANXA2 Facilitates Enterovirus 71 Infection by Interacting with 3D Polymerase and PI4KB to Assist the Assembly of Replication Organelles. <i>Virologica Sinica</i> , <b>2021</b> , 1	6.4	1
38	The Role of RNA-Binding Proteins in IRES-Dependent Translation. <i>Growth Hormone</i> , <b>2002</b> , 1-33		3
37	A cell cycle-dependent protein serves as a template-specific translation initiation factor. <i>Genes and Development</i> , <b>2000</b> , 14, 2028-2045	12.6	154
36	Picornavirus Genome: an Overview. 125-148		14
35	Proteins Involved in the Function of Picornavirus Internal Ribosomal Entry Sites. 171-183		5
34	Possible Unifying Mechanism of Picornavirus Genome Replication. 225-246		36
33	Processing of a cellular polypeptide by 3CD proteinase is required for poliovirus ribonucleoprotein complex formation. <i>Journal of Virology</i> , <b>1997</b> , 71, 578-85	6.6	34
32	Functional involvement of polypyrimidine tract-binding protein in translation initiation complexes with the internal ribosome entry site of foot-and-mouth disease virus. <i>Journal of Virology</i> , <b>1997</b> , 71, 8330-9	6.6	77
31	Requirement of poly(rC) binding protein 2 for translation of poliovirus RNA. <i>Journal of Virology</i> , <b>1997</b> , 71, 6243-6	6.6	246
30	trans-encapsidation of a poliovirus replicon by different picornavirus capsid proteins. <i>Journal of Virology</i> , <b>1998</b> , 72, 7972-7	6.6	24
29	Interaction of poly(rC) binding protein 2 with the 5' noncoding region of hepatitis A virus RNA and its effects on translation. <i>Journal of Virology</i> , <b>1998</b> , 72, 9668-75	6.6	66
28	A new internal ribosomal entry site 5' boundary is required for poliovirus translation initiation in a mouse system. <i>Journal of Virology</i> , <b>1998</b> , 72, 2398-405	6.6	12
27	A small yeast RNA blocks hepatitis C virus internal ribosome entry site (HCV IRES)-mediated translation and inhibits replication of a chimeric poliovirus under translational control of the HCV IRES element. <i>Journal of Virology</i> , <b>1998</b> , 72, 5638-47	6.6	35
26	Complete protein linkage map of poliovirus P3 proteins: interaction of polymerase 3Dpol with VPg and with genetic variants of 3AB. <i>Journal of Virology</i> , <b>1998</b> , 72, 6732-41	6.6	109
25	Hepatitis B virus RNA-binding proteins associated with cytokine-induced clearance of viral RNA from the liver of transgenic mice. <i>Journal of Virology</i> , <b>1999</b> , 73, 474-81	6.6	81
24	Absence of internal ribosome entry site-mediated tissue specificity in the translation of a bicistronic transgene. <i>Journal of Virology</i> , <b>1999</b> , 73, 2729-38	6.6	31

23	Multimerization of poly(rC) binding protein 2 is required for translation initiation mediated by a viral IRES. <i>Rna</i> , <b>2004</b> , 10, 1266-76	5.8	42
22	Giardavirus internal ribosome entry site has an apparently unique mechanism of initiating translation. <i>PLoS ONE</i> , <b>2009</b> , 4, e7435	3.7	6
21	Additive Promotion of Viral Internal Ribosome Entry Site-Mediated Translation by Far Upstream Element-Binding Protein 1 and an Enterovirus 71-Induced Cleavage Product. <i>PLoS Pathogens</i> , <b>2016</b> , 12, e1005959	7.6	28
20	Identifying the cellular location of brain cytoplasmic 200 RNA using an RNA-recognizing antibody. <i>BMB Reports</i> , <b>2017</b> , 50, 318-322	5.5	7
19	Poly-C Binding Proteins: Cellular Regulators of mRNA Fate and Function. <i>Growth Hormone</i> , <b>2002</b> , 53-69		
18	Informatics For Protein Identification by Tandem Mass Spectrometry; Focused on Two Most-widely Applied Algorithms, Mascot and SEQUEST. <i>Interdisciplinary Bio Central</i> , <b>2006</b> , 1, 88-93		
17	Positive-Strand RNA Virus Replication in Vertebrate Hosts.		
16	Interference with Cellular Gene Expression. 163-180		
15	Genome Replication I: the Players. 105-125		
14	Poliomyelitis. 337-351		
13	Host restriction factor A3G inhibits the replication of Enterovirus D68 through competitively binding 5QTR with PCBP1. <i>Journal of Virology</i> , <b>2021</b> , JVI0170821	6.6	1
12	A cell cycle-dependent protein serves as a template-specific translation initiation factor. <i>Genes and Development</i> , <b>2000</b> , 14, 2028-45	12.6	231
11	Oncolytic immunotherapy through tumor-specific translation and cytotoxicity of poliovirus. <i>Discovery Medicine</i> , <b>2015</b> , 19, 359-65	2.5	9
10	RNA-Binding Proteins as Regulators of Internal Initiation of Viral mRNA Translation.. <i>Viruses</i> , <b>2022</b> , 14,	6.2	0
9	Picornavirus translation strategies.. <i>FEBS Open Bio</i> , <b>2022</b> ,	2.7	1
8	Poly(C)-binding Protein 2 Regulates the p53 Expression via Interactions with the 5QTerminal Region of p53 mRNA.. <i>International Journal of Molecular Sciences</i> , <b>2021</b> , 22,	6.3	1
7	Image_1.JPEG. <b>2018</b> ,		
6	Image_2.JPEG. <b>2018</b> ,		

5 Image\_3.JPEG. **2018**,

4 PKR Binds Enterovirus IRESs, Displaces Host Translation Factors, and Impairs Viral Translation to Enable Innate Antiviral Signaling. *MBio*, 7.8 ○

3 Multiple functions of heterogeneous nuclear ribonucleoproteins in the positive single-stranded RNA virus life cycle. 13, 2

2 Elusive Trans-Acting Factors Which Operate with Type I (Poliovirus-like) IRES Elements. **2022**, 23, 15497 ○

1 RNA Structure in the 5' Untranslated Region of Enterovirus D68 Strains with Differing Neurovirulence Phenotypes. **2023**, 15, 295 ○