

Cold activation of complement in sera from patients with infection on interferon therapy

Journal of Gastroenterology and Hepatology (Australia)

11, 341-346

DOI: [10.1111/j.1440-1746.1996.tb01381.x](https://doi.org/10.1111/j.1440-1746.1996.tb01381.x)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Cold activation of complement enhancing with the duration of storage in sera from blood donors with hepatitis C virus RNA. <i>International Hepatology Communications</i> , 1996, 5, 89-96.	0.7	0
2	Hepatitis C Virus Infection in Patients With Psoriasis. <i>Archives of Dermatology</i> , 1996, 132, 1391.	1.7	7
3	Antibodies to hepatitis C virus and elevated transaminase levels in a town of hyperendemicity in Iwate, Japan. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 1997, 12, 67-72.	1.4	4
4	Changes in the acute phase complement component and IL-6 levels in patients with chronic hepatitis C receiving interferon α -2b. <i>Immunology Letters</i> , 2000, 72, 69-74.	1.1	14
5	Hepatitis C virus infection and myositis: a polymerase chain reaction study. <i>Acta Neuropathologica</i> , 2000, 99, 271-276.	3.9	17
6	In vitro cold activation of complement shown by an overestimation of total complement 4: a study in patients with hepatitis C virus infection. <i>Annals of Clinical Biochemistry</i> , 2001, 38, 687-693.	0.8	5
7	Loss of Circulating Hepatitis C Virus in Children Who Developed a Persistent Carrier State after Mother-to-Baby Transmission. <i>Pediatric Research</i> , 1997, 42, 263-267.	1.1	37