

TNFÎ± concentrations and mRNA expression are increased

Journal of Reproductive Immunology

32, 157-169

DOI: [10.1016/s0165-0378\(96\)00998-9](https://doi.org/10.1016/s0165-0378(96)00998-9)

Citation Report

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Cytokine Expression by First-Trimester Human Chorionic Villi. American Journal of Reproductive Immunology, 1998, 40, 309-318. | 1.2 | 36 |
| 2 | Circulating Levels of Immunoreactive Cytokines in Women with Preeclampsia. American Journal of Reproductive Immunology, 1998, 40, 102-111. | 1.2 | 365 |
| 3 | Placental mitochondria as a source of oxidative stress in pre-eclampsia. Placenta, 1998, 19, 581-586. | 0.7 | 242 |
| 4 | Plasma tumor necrosis factor- α soluble receptor p55 (sTNFp55) concentrations in eclamptic, preeclamptic and normotensive pregnant Zimbabwean women. Journal of Reproductive Immunology, 1998, 40, 159-173. | 0.8 | 42 |
| 5 | Maternal-Placental Interactions of Oxidative Stress and Antioxidants in Preeclampsia. Seminars in Reproductive Medicine, 1998, 16, 93-104. | 0.5 | 222 |
| 6 | Influence of human leukocyte antigen and tumour necrosis factor genes on the development of pre-eclampsia. Human Reproduction Update, 1999, 5, 94-102. | 5.2 | 34 |
| 7 | Interaction of the protein C/protein S anticoagulant system, the endothelium and pregnancy. Blood Reviews, 1999, 13, 127-146. | 2.8 | 29 |
| 8 | Expression of the placental cytokines tumor necrosis factor α , interleukin 1β , and interleukin 10 is increased in preeclampsia. American Journal of Obstetrics and Gynecology, 1999, 181, 915-920. | 0.7 | 210 |
| 9 | Maternal Second Trimester Serum Tumor Necrosis Factor- α -soluble Receptor p55 (sTNFp55) and Subsequent Risk of Preeclampsia. American Journal of Epidemiology, 1999, 149, 323-329. | 1.6 | 105 |
| 10 | Leukocytosis is Proportional to HELLP Syndrome Severity. Southern Medical Journal, 2000, 93, 768-771. | 0.3 | 13 |
| 11 | Activin A Exerts both Pro- and Anti-inflammatory Effects on Human Term Gestational Tissues. Placenta, 2000, 21, 38-43. | 0.7 | 91 |
| 12 | Tumor necrosis factor- α soluble receptor p55 (sTNFp55) and risk of preeclampsia in Peruvian women. Journal of Reproductive Immunology, 2000, 47, 49-63. | 0.8 | 24 |
| 13 | Induction of Placental Heme Oxygenase-1 Is Protective Against TNF α -induced Cytotoxicity and Promotes Vessel Relaxation. Molecular Medicine, 2000, 6, 391-409. | 1.9 | 141 |
| 14 | Modulation of indoleamine 2,3-dioxygenase by interferon-gamma in human placental chorionic villi. Molecular Human Reproduction, 2000, 6, 369-374. | 1.3 | 34 |
| 15 | Invasive Cytotrophoblasts Manifest Evidence of Oxidative Stress in Preeclampsia. American Journal of Pathology, 2000, 156, 321-331. | 1.9 | 245 |
| 16 | Postnatal androgenization induces premature aging of rat ovaries. Steroids, 2000, 65, 190-205. | 0.8 | 35 |
| 17 | Stress and Pregnancy Loss: Role of Immune Mediators, Hormones and Neurotransmitters. American Journal of Reproductive Immunology, 2001, 46, 117-123. | 1.2 | 91 |
| 18 | Expression of Inflammatory Cytokines in Placentas from Women with Preeclampsia. Journal of Clinical Endocrinology and Metabolism, 2001, 86, 2505-2512. | 1.8 | 292 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Role of psychoneuroimmunology in reproduction. <i>Der Gynakologe</i> , 2001, 34, 539-546. | 1.0 | 1 |
| 20 | Cytokine Levels in Midtrimester Amniotic Fluid in Normal Pregnancy and in the Prediction of Pre-eclampsia. <i>Scandinavian Journal of Immunology</i> , 2001, 53, 310-314. | 1.3 | 44 |
| 21 | Increased Superoxide Generation is Associated with Decreased Superoxide Dismutase Activity and mRNA Expression in Placental Trophoblast Cells in Pre-eclampsia. <i>Placenta</i> , 2001, 22, 206-212. | 0.7 | 206 |
| 22 | Effect of Hypoxia, Oxidative Stress and Lipopolysaccharides on the Release of Prostaglandins and Cytokines from Human Term Placental Explants. <i>Placenta</i> , 2001, 22, S45-S50. | 0.7 | 54 |
| 23 | Neutrophil Activation Induced by Placental Factors in Normal and Pre-eclamptic Pregnancies In Vitro. <i>Placenta</i> , 2001, 22, 560-565. | 0.7 | 45 |
| 24 | Lack of association of severe preeclampsia with maternal and fetal mutant alleles for tumor necrosis factor $\text{I}\alpha$ and lymphotoxin $\text{I}\alpha$ genes and plasma tumor necrosis factor $\text{I}\alpha$ levels. <i>American Journal of Obstetrics and Gynecology</i> , 2001, 184, 1273-1277. | 0.7 | 37 |
| 25 | Activin A and activin receptors in gestational tissue from preeclamptic pregnancies. <i>Journal of Endocrinology</i> , 2001, 171, 57-64. | 1.2 | 61 |
| 26 | Regulation of Proinflammatory Cytokines in Human Lung Epithelial Cells Infected with <i>Mycoplasma pneumoniae</i> . <i>Infection and Immunity</i> , 2002, 70, 3649-3655. | 1.0 | 111 |
| 27 | Searching for preeclampsia genes: the current position. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2002, 105, 94-113. | 0.5 | 138 |
| 28 | Cytokines of the Placenta and Extra-placental Membranes: Roles and Regulation During Human Pregnancy and Parturition. <i>Placenta</i> , 2002, 23, 257-273. | 0.7 | 297 |
| 29 | Adriana and Luisa Castellucci Award Lecture 2001 Hypoxia Inducible Factor-1: Oxygen Regulation of Trophoblast Differentiation in Normal and Pre-eclamptic Pregnancies – A Review. <i>Placenta</i> , 2002, 23, S47-S57. | 0.7 | 183 |
| 30 | Increased apoptosis in the syncytiotrophoblast in human term placentas complicated by either preeclampsia or intrauterine growth retardation. <i>American Journal of Obstetrics and Gynecology</i> , 2002, 186, 158-166. | 0.7 | 459 |
| 31 | Transforming growth factor-beta1 does not relate to hypertension in pre-eclampsia. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2002, 29, 968-971. | 0.9 | 22 |
| 32 | Pathophysiology of Preeclampsia: Linking Placental Ischemia/Hypoxia with Microvascular Dysfunction. <i>Microcirculation</i> , 2002, 9, 147-160. | 1.0 | 279 |
| 33 | Macrophage Inhibitory Cytokine-1 in Gestational Tissues and Maternal Serum in Normal and Pre-eclamptic Pregnancy. <i>Placenta</i> , 2003, 24, 100-106. | 0.7 | 77 |
| 34 | Protease Chymotrypsin Mediates the Endothelial Expression of P- and E-selectin, but not ICAM and VCAM, Induced by Placental Trophoblasts from Pre-eclamptic Pregnancies. <i>Placenta</i> , 2003, 24, 851-861. | 0.7 | 29 |
| 35 | Th1/Th2 balance in preeclampsia. <i>Journal of Reproductive Immunology</i> , 2003, 59, 161-173. | 0.8 | 269 |
| 36 | Placental Tissue Interleukin-10 Receptor Distribution in Pre-eclampsia. <i>American Journal of Reproductive Immunology</i> , 2003, 49, 377-381. | 1.2 | 16 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Intracellular glutathione and lipid peroxide availability and the secretion of vasoactive substances by human umbilical vein endothelial cells after incubation with TNF- α . <i>European Journal of Clinical Investigation</i> , 2003, 33, 176-182. | 1.7 | 20 |
| 38 | Functional proteomics of neurokinin B in the placenta indicates a novel role in regulating cytotrophoblast antioxidant defences. <i>Proteomics</i> , 2003, 3, 2044-2051. | 1.3 | 54 |
| 39 | Preeclamptic women are deficient of interleukin-10 as assessed by cytokine release of trophoblast cells in vitro. <i>Cytokine</i> , 2003, 23, 119-125. | 1.4 | 67 |
| 40 | Induction of Proinflammatory Cytokines in Human Lung Epithelial Cells during Chlamydia pneumoniae Infection. <i>Infection and Immunity</i> , 2003, 71, 614-620. | 1.0 | 58 |
| 41 | Adrenomedullin Is Decreased in Preeclampsia Because of Failed Response to Epidermal Growth Factor and Impaired Syncytialization. <i>Hypertension</i> , 2003, 42, 895-900. | 1.3 | 32 |
| 42 | Hypoxia-Inducible Factor-1 Transactivates Transforming Growth Factor- β 3 in Trophoblast. <i>Endocrinology</i> , 2004, 145, 4113-4118. | 1.4 | 84 |
| 43 | Oxidative stress in the placenta. <i>Histochemistry and Cell Biology</i> , 2004, 122, 369-382. | 0.8 | 664 |
| 45 | Secretion of Tumor Necrosis Factor- α from Human Placental Tissues Induced by Hypoxia-Reoxygenation Causes Endothelial Cell Activation in Vitro. <i>American Journal of Pathology</i> , 2004, 164, 1049-1061. | 1.9 | 190 |
| 46 | Prediction of Preeclampsia: Can It Be Achieved?. <i>Obstetrical and Gynecological Survey</i> , 2004, 59, 464-482. | 0.2 | 33 |
| 47 | Impact of genes related to immune tolerance and inflammation (tumour necrosis factor- α), Tj ETQq1 1 0.784314 rgBT /Overlock 10 TTS 2005, 23, 2187-2191. | 0.3 | 24 |
| 48 | Prevention of Preeclampsia With Antioxidants: Evidence From Randomized Trials. <i>Clinical Obstetrics and Gynecology</i> , 2005, 48, 416-429. | 0.6 | 27 |
| 49 | The Role of Placental Oxidative Stress and Lipid Peroxidation in Preeclampsia. <i>Obstetrical and Gynecological Survey</i> , 2005, 60, 807-816. | 0.2 | 133 |
| 50 | Neutrophils from pregnant women produce thromboxane and tumor necrosis factor- α in response to linoleic acid and oxidative stress. <i>American Journal of Obstetrics and Gynecology</i> , 2005, 193, 830-835. | 0.7 | 30 |
| 51 | Placental imbalance of Th1- and Th2-type cytokines in preeclampsia. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2005, 84, 788-793. | 1.3 | 48 |
| 52 | Glucocorticoids inhibit placental cytokines from cultured normal and preeclamptic placental explants. <i>Placenta</i> , 2005, 26, 654-660. | 0.7 | 26 |
| 53 | Serum and placental interleukin-18 are elevated in preeclampsia. <i>Journal of Reproductive Immunology</i> , 2005, 65, 77-87. | 0.8 | 37 |
| 54 | Expression of TNF α and its receptors R1 and R2 in human alveolar epithelial cells exposed to organic dust and the effects of 8-bromo-cAMP and protein kinase A modulation. <i>Inflammation Research</i> , 2005, 54, 281-288. | 1.6 | 19 |
| 55 | Cytokines, Hypoxia, and Preeclampsia. <i>Journal of the Society for Gynecologic Investigation</i> , 2005, 12, 385-387. | 1.9 | 7 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 56 | Interleukin-10 regulates arterial pressure in early primate pregnancy. <i>Cytokine</i> , 2005, 29, 176-185. | 1.4 | 38 |
| 57 | Regulation of Monocyte Chemoattractant Protein-1 Expression by Tumor Necrosis Factor- α and Interleukin-1 β in First Trimester Human Decidual Cells. <i>American Journal of Pathology</i> , 2006, 168, 445-452. | 1.9 | 121 |
| 58 | Hypoxia and Reoxygenation: a Possible Mechanism for Placental Oxidative Stress in Preeclampsia. <i>Taiwanese Journal of Obstetrics and Gynecology</i> , 2006, 45, 189-200. | 0.5 | 243 |
| 59 | Monocytes of preeclamptic women spontaneously synthesize pro-inflammatory cytokines. <i>Clinical Immunology</i> , 2006, 118, 268-275. | 1.4 | 117 |
| 60 | Immunological Analogy Between Allograft Rejection, Recurrent Abortion and Pre-Eclampsia – the Same Basic Mechanism?. <i>Human Immunology</i> , 2006, 67, 492-511. | 1.2 | 98 |
| 61 | Impact of genetic variation of tumor necrosis factor- α on gestational hypertension. <i>Chinese Medical Journal</i> , 2006, 119, 719-724. | 0.9 | 24 |
| 62 | Antihypertensive drugs clonidine, diazoxide, hydralazine and furosemide regulate the production of cytokines by placentas and peripheral blood mononuclear cells in normal pregnancy. <i>Journal of Hypertension</i> , 2006, 24, 915-922. | 0.3 | 45 |
| 63 | Total peroxy radical-trapping ability and anti-oxidant vitamins of the umbilical venous plasma and the placenta in pre-eclampsia. <i>Journal of Obstetrics and Gynaecology Research</i> , 2006, 32, 32-41. | 0.6 | 21 |
| 64 | Shared and disparate components of the pathophysiologies of fetal growth restriction and preeclampsia. <i>American Journal of Obstetrics and Gynecology</i> , 2006, 195, 40-49. | 0.7 | 382 |
| 65 | PAF Levels and PAF-AH Activities in Placentas from Normal and Preeclamptic Pregnancies. <i>Placenta</i> , 2006, 27, 744-749. | 0.7 | 12 |
| 66 | Tumour necrosis factor- α impairs chorionic gonadotrophin β -subunit expression and cell fusion of human villous cytotrophoblast. <i>Molecular Human Reproduction</i> , 2006, 12, 601-609. | 1.3 | 65 |
| 67 | Difference in Neutrophil Superoxide Generation During Pregnancy Between Preeclampsia and Essential Hypertension. <i>Hypertension</i> , 2007, 49, 1436-1441. | 1.3 | 26 |
| 68 | Inflammatory Changes in Preeclampsia: Current Understanding of the Maternal Innate and Adaptive Immune Response. <i>Obstetrical and Gynecological Survey</i> , 2007, 62, 191-201. | 0.2 | 73 |
| 69 | Inflammatory response in preeclampsia. <i>Molecular Aspects of Medicine</i> , 2007, 28, 210-219. | 2.7 | 76 |
| 70 | Increased Apoptosis in First Trimester Extravillous Trophoblasts from Pregnancies at Higher Risk of Developing Preeclampsia. <i>American Journal of Pathology</i> , 2007, 170, 1903-1909. | 1.9 | 88 |
| 71 | Anti-Hypertensive Drugs Alter Cytokine Production from Preeclamptic Placentas and Peripheral Blood Mononuclear Cells. <i>Hypertension in Pregnancy</i> , 2007, 26, 343-356. | 0.5 | 16 |
| 72 | Maternal serum soluble CD30 is increased in normal pregnancy, but decreased in preeclampsia and small for gestational age pregnancies. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2007, 20, 867-878. | 0.7 | 34 |
| 73 | CXCL10/IP-10: A missing link between inflammation and anti-angiogenesis in preeclampsia?. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2007, 20, 777-792. | 0.7 | 112 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 74 | Nicotine Inhibits Cytokine Production by Placenta Cells via NF κ B: Potential Role in Pregnancy-Induced Hypertension. <i>Molecular Medicine</i> , 2007, 13, 576-583. | 1.9 | 53 |
| 75 | Leptin, IL-10 and Inflammatory Markers (TNF- α , IL-6 and IL-8) in Pre-Eclamptic, Normotensive Pregnant and Healthy Non-Pregnant Women. <i>American Journal of Reproductive Immunology</i> , 2007, 58, 21-30. | 1.2 | 201 |
| 76 | Placental Oxidative Stress in a Rat Model of Preeclampsia. <i>Placenta</i> , 2007, 28, 52-58. | 0.7 | 37 |
| 77 | Role of placentally produced inflammatory and regulatory cytokines in pregnancy and the etiology of preeclampsia. <i>Seminars in Immunopathology</i> , 2007, 29, 151-162. | 2.8 | 167 |
| 78 | Elevated Plasma Levels of Interleukin 1 β , Tumour Necrosis Factor α and Monocyte Chemotactic Protein 1 Are Associated with Pregnancy Toxaemia in Ewes. <i>Veterinary Research Communications</i> , 2007, 31, 565-573. | 0.6 | 12 |
| 79 | The combination of ApoCIII, hepatic lipase and hormone sensitive lipase gene polymorphisms suggests an association with susceptibility to gestational hypertension. <i>Journal of Human Genetics</i> , 2007, 52, 244-254. | 1.1 | 20 |
| 80 | ORIGINAL ARTICLE: Association of Tumor Necrosis Factor α and Interleukin α 10 Gene Polymorphisms in Iranian Patients with Pre α eclampsia. <i>American Journal of Reproductive Immunology</i> , 2008, 60, 179-185. | 1.2 | 31 |
| 81 | Hydralazine reduces leukocyte migration through different mechanisms in spontaneously hypertensive and normotensive rats. <i>European Journal of Pharmacology</i> , 2008, 589, 206-214. | 1.7 | 20 |
| 82 | Preeclampsia: Impaired inflammatory response mediated by Toll-like receptors. <i>Journal of Reproductive Immunology</i> , 2008, 78, 80-83. | 0.8 | 26 |
| 83 | Altered Interleukin-6 Receptor, IL-6R and gp130, Production and Expression and Decreased SOCS-3 Expression in Placentas from Women with Pre-eclampsia. <i>Placenta</i> , 2008, 29, 1024-1028. | 0.7 | 38 |
| 84 | Low birth weight in response to salt restriction during pregnancy is not due to alterations in uterine-placental blood flow or the placental and peripheral renin α angiotensin system. <i>Physiology and Behavior</i> , 2008, 95, 145-151. | 1.0 | 38 |
| 85 | Association between tumor necrosis factor (TNF) α G-308A gene polymorphism and preeclampsia complicated by severe fetal growth restriction. <i>Clinica Chimica Acta</i> , 2008, 392, 52-57. | 0.5 | 60 |
| 86 | Pathophysiology of hypertension during preeclampsia: linking placental ischemia with endothelial dysfunction. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2008, 294, H541-H550. | 1.5 | 432 |
| 87 | Recent Progress Toward the Understanding of the Pathophysiology of Hypertension During Preeclampsia. <i>Hypertension</i> , 2008, 51, 982-988. | 1.3 | 134 |
| 88 | Exogenous Soluble VEGF Receptor-1 (sFlt-1) Regulates Th1/Th2 Cytokine Production from Normal Placental Explants via Intracellular Calcium. <i>Hypertension in Pregnancy</i> , 2009, 28, 448-456. | 0.5 | 8 |
| 89 | Human Tumour Necrosis Factor: Physiological and Pathological Roles in Placenta and Endometrium. <i>Placenta</i> , 2009, 30, 111-123. | 0.7 | 286 |
| 90 | Calcitriol inhibits TNF α -induced inflammatory cytokines in human trophoblasts. <i>Journal of Reproductive Immunology</i> , 2009, 81, 17-24. | 0.8 | 144 |
| 91 | Placental production and maternal serum and urine levels of inhibin A and activin A are modified by antihypertensive therapy in hypertensive disorders of pregnancy. <i>Clinical Endocrinology</i> , 2009, 70, 924-931. | 1.2 | 11 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 92 | ORIGINAL ARTICLE: Maternal Circulating TNF α Levels are Highly Correlated with IL-10 Levels, but not IL-6 and IL-8 Levels, in Women with Preeclampsia. American Journal of Reproductive Immunology, 2009, 62, 269-274. | 1.2 | 32 |
| 93 | The Phenotypic and Metabolic Characteristics of Neutrophils in Normal Pregnancy and Preeclampsia. Current Immunology Reviews, 2009, 5, 243-248. | 1.2 | 0 |
| 94 | Possible therapeutic effect of Magnesium sulfate in pre-eclampsia by the down-regulation of placental tumor necrosis factor-alpha secretion. European Cytokine Network, 2010, 21, 58-64. | 1.1 | 12 |
| 95 | Altered levels of insulin-like growth factor binding protein proteases in preeclampsia and intrauterine growth restriction. Prenatal Diagnosis, 2010, 30, 815-820. | 1.1 | 24 |
| 96 | ORIGINAL ARTICLE: Leptin Gene (TTTC) _n Microsatellite Polymorphism as well as Leptin Receptor R223Q and PPAR γ 2 P12A Substitutions are not Associated with Hypertensive Disorders in Pregnancy. American Journal of Reproductive Immunology, 2010, 63, 310-317. | 1.2 | 9 |
| 97 | Role of Peroxiredoxin III in the Pathogenesis of Pre-eclampsia as Evidenced in Mice. Oxidative Medicine and Cellular Longevity, 2010, 3, 71-73. | 1.9 | 12 |
| 98 | Leukocytes of pregnant women with small-for-gestational age neonates have a different phenotypic and metabolic activity from those of women with preeclampsia. Journal of Maternal-Fetal and Neonatal Medicine, 2010, 23, 476-487. | 0.7 | 40 |
| 99 | Angiotensin Receptor Agonistic Autoantibody-Mediated Tumor Necrosis Factor- α Induction Contributes to Increased Soluble Endoglin Production in Preeclampsia. Circulation, 2010, 121, 436-444. | 1.6 | 70 |
| 100 | The roles of cellular reactive oxygen species, oxidative stress and antioxidants in pregnancy outcomes. International Journal of Biochemistry and Cell Biology, 2010, 42, 1634-1650. | 1.2 | 565 |
| 101 | Endoplasmic reticulum stress in the pathogenesis of early-onset pre-eclampsia. Pregnancy Hypertension, 2011, 1, 72-78. | 0.6 | 107 |
| 102 | A meta-analysis of tumor necrosis factor-alpha, interleukin-6, and interleukin-10 in preeclampsia. Cytokine, 2011, 56, 550-559. | 1.4 | 98 |
| 103 | Immunologic Characteristics of Preeclampsia, a Comprehensive Review. American Journal of Reproductive Immunology, 2011, 65, 377-394. | 1.2 | 90 |
| 104 | TNF- α inhibits trophoblast integration into endothelial cellular networks. Placenta, 2011, 32, 241-246. | 0.7 | 66 |
| 105 | Protective effects of hydrogen-rich saline in preeclampsia rat model. Placenta, 2011, 32, 681-686. | 0.7 | 22 |
| 106 | ADAM17 regulates TNF α production by placental trophoblasts. Placenta, 2011, 32, 975-980. | 0.7 | 29 |
| 107 | Concentraciones de interleucina 10 en preeclámpticas y embarazadas normotensas sanas. Clinica E Investigacion En Ginecología Y Obstetricia, 2011, 38, 164-168. | 0.1 | 0 |
| 108 | Regulation of pregnancy-associated plasma protein A2 (PAPPA2) in a human placental trophoblast cell line (BeWo). Reproductive Biology and Endocrinology, 2011, 9, 48. | 1.4 | 38 |
| 109 | Maternal Schistosomiasis Japonica Is Associated with Maternal, Placental, and Fetal Inflammation. Infection and Immunity, 2011, 79, 1254-1261. | 1.0 | 55 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 110 | IL-6-induced pathophysiology during pre-eclampsia: potential therapeutic role for magnesium sulfate?. International Journal of Interferon, Cytokine and Mediator Research, 2011, 2011, 59. | 1.1 | 59 |
| 111 | Activation of NF- κ B in Placentas of Women with Preeclampsia. Hypertension in Pregnancy, 2012, 31, 243-251. | 0.5 | 103 |
| 112 | Trophoblastic progranulin expression is upregulated in cases of fetal growth restriction and preeclampsia. Journal of Perinatal Medicine, 2012, 40, 475-481. | 0.6 | 21 |
| 113 | Calcitriol inhibits interleukin-10 expression in cultured human trophoblasts under normal and inflammatory conditions. Cytokine, 2012, 57, 316-321. | 1.4 | 36 |
| 114 | The Pathophysiology of Preeclampsia Involves Altered Levels of Angiogenic Factors Promoted by Hypoxia and Autoantibody-Mediated Mechanisms1. Biology of Reproduction, 2012, 87, 36. | 1.2 | 37 |
| 115 | Single-nucleotide polymorphisms in genes involved in placental function and unexplained stillbirth. American Journal of Obstetrics and Gynecology, 2012, 207, 316.e1-316.e7. | 0.7 | 5 |
| 116 | Maternal tumor necrosis factor receptor 2 gene variants associated with pre-eclampsia in Tunisian women. Journal of Obstetrics and Gynaecology Research, 2013, 39, 1301-1307. | 0.6 | 6 |
| 117 | Elucidating Immune Mechanisms Causing Hypertension During Pregnancy. Physiology, 2013, 28, 225-233. | 1.6 | 78 |
| 118 | Serum levels of adrenomedullin and inflammatory cytokines in women with term idiopathic intrauterine growth restriction. Journal of Obstetrics and Gynaecology, 2013, 33, 135-139. | 0.4 | 13 |
| 119 | Placental trophoblast cell differentiation: Physiological regulation and pathological relevance to preeclampsia. Molecular Aspects of Medicine, 2013, 34, 981-1023. | 2.7 | 306 |
| 120 | Preeclampsia-associated stresses activate Gadd45a signaling and sFlt-1 in placental explants. Journal of Cellular Physiology, 2013, 228, 362-370. | 2.0 | 39 |
| 121 | Expression of inflammatory cytokines in placentas from pregnancies complicated with preeclampsia and HELLP syndrome. Journal of Maternal-Fetal and Neonatal Medicine, 2013, 26, 680-685. | 0.7 | 13 |
| 123 | Cytokines as Key Players in the Pathophysiology of Preeclampsia. Medical Principles and Practice, 2013, 22, 8-19. | 1.1 | 154 |
| 124 | Tumor Necrosis Factor-Alpha Polymorphism at Position -238 in Preeclampsia. Iranian Red Crescent Medical Journal, 2014, 16, e11195. | 0.5 | 18 |
| 125 | Monocytes and Macrophages in Pregnancy and Pre-Eclampsia. Frontiers in Immunology, 2014, 5, 298. | 2.2 | 172 |
| 126 | IL-27 Activates Human Trophoblasts to Express IP-10 and IL-6: Implications in the Immunopathophysiology of Preeclampsia. Mediators of Inflammation, 2014, 2014, 1-10. | 1.4 | 25 |
| 127 | Down-Regulation of TIMP-3 Leads to Increase in TACE Expression and TNF- α Production by Placental Trophoblast Cells. American Journal of Reproductive Immunology, 2014, 71, 427-433. | 1.2 | 10 |
| 128 | Nitric oxide (NO) reversed TNF- α inhibition of trophoblast interaction with endothelial cellular networks. Placenta, 2014, 35, 417-421. | 0.7 | 26 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 129 | Progranulin shows cytoprotective effects on trophoblast cells in vitro but does not antagonize TNF- α -induced apoptosis. Archives of Gynecology and Obstetrics, 2014, 290, 867-873. | 0.8 | 8 |
| 130 | An integrative view on the physiology of human early placental villi. Progress in Biophysics and Molecular Biology, 2014, 114, 33-48. | 1.4 | 32 |
| 131 | Antihypertensive drugs methyldopa, labetalol, hydralazine, and clonidine improve trophoblast interaction with endothelial cellular networks in vitro. Journal of Hypertension, 2014, 32, 1075-1083. | 0.3 | 27 |
| 132 | Placental Regulation of Inflammation and Hypoxia after TNF- α Infusion in Mice. American Journal of Reproductive Immunology, 2015, 74, 407-418. | 1.2 | 42 |
| 133 | Vitamin D and Inflammatory Cytokines in Healthy and Preeclamptic Pregnancies. Nutrients, 2015, 7, 6465-6490. | 1.7 | 66 |
| 134 | Increased risk for the development of preeclampsia in obese pregnancies: weighing in on the mechanisms. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2015, 309, R1326-R1343. | 0.9 | 99 |
| 135 | Oxidative Stress in Preeclampsia. , 2015, , 283-290. | | 1 |
| 136 | Aberrant maternal inflammation as a cause of pregnancy complications: A potential therapeutic target?. Placenta, 2015, 36, 960-966. | 0.7 | 91 |
| 137 | Placenta-derived angiogenic proteins and their contribution to the pathogenesis of preeclampsia. Angiogenesis, 2015, 18, 115-123. | 3.7 | 43 |
| 138 | Placental tumor necrosis factor- α protein expression during normal human gestation. Journal of Maternal-Fetal and Neonatal Medicine, 2016, 29, 3934-3938. | 0.7 | 14 |
| 139 | The paternal polymorphism rs5370 in the EDN1 gene decreases the risk of preeclampsia. Pregnancy Hypertension, 2016, 6, 327-332. | 0.6 | 14 |
| 140 | Inventory of Novel Animal Models Addressing Etiology of Preeclampsia in the Development of New Therapeutic/Intervention Opportunities. American Journal of Reproductive Immunology, 2016, 75, 402-410. | 1.2 | 30 |
| 141 | Effect of Factors Secreted by the Placenta on Phenotype of THP-1 Cells Cultured on a 3D Scaffold. Bulletin of Experimental Biology and Medicine, 2016, 161, 162-167. | 0.3 | 2 |
| 142 | Evaluation of 5-HT ₇ receptor expression in the placentae of normal and pre-eclamptic women. Clinical and Experimental Hypertension, 2016, 38, 189-193. | 0.5 | 8 |
| 143 | Effect of THP-1 Cells on the Formation of Vascular Tubes by Endothelial EA.hy926 Cells in the Presence of Placenta Secretory Products. Bulletin of Experimental Biology and Medicine, 2017, 162, 545-551. | 0.3 | 2 |
| 144 | Precision Medicine in Gynecology and Obstetrics. Comprehensive Gynecology and Obstetrics, 2017, , . | 0.0 | 1 |
| 145 | The protective effect of apolipoprotein in models of trophoblast invasion and preeclampsia. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2017, 312, R40-R48. | 0.9 | 18 |
| 146 | Loss of placental thrombomodulin in oocyte donation pregnancies. Fertility and Sterility, 2017, 107, 119-129.e5. | 0.5 | 11 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 147 | Antihypertensive methyldopa, labetalol, hydralazine, and clonidine reversed tumour necrosis factor- α -induced endothelial nitric oxide synthase expression in endothelial-trophoblast cellular networks. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2017, 44, 421-427. | 0.9 | 14 |
| 148 | Uncovering novel roles for matrix metalloproteinases in preeclampsia. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2017, 313, H687-H689. | 1.5 | 1 |
| 149 | A pronounced uterine pro-inflammatory response at parturition is an ancient feature in mammals. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2017, 284, 20171694. | 1.2 | 32 |
| 150 | Maternal monocytes in pregnancy and preeclampsia in humans and in rats. <i>Journal of Reproductive Immunology</i> , 2017, 119, 91-97. | 0.8 | 46 |
| 151 | Innate immune cells in the placental bed in healthy pregnancy and preeclampsia. <i>Placenta</i> , 2018, 69, 125-133. | 0.7 | 94 |
| 152 | Gene expression analysis of MMPs in women with preeclampsia using cell-free fetal RNA in maternal plasma. <i>Pregnancy Hypertension</i> , 2019, 17, 261-268. | 0.6 | 4 |
| 153 | The role of TNF- α and TLR4 polymorphisms in the placenta of pregnant women complicated by preeclampsia and in silico analysis. <i>International Journal of Biological Macromolecules</i> , 2019, 134, 1205-1215. | 3.6 | 16 |
| 154 | PREECLAMPSIA: Parte 2 ^o : fisiopatologia e trattamento. <i>Giornale De Tecniche Nefrologiche & Dialitiche</i> , 2019, 31, 79-88. | 0.1 | 0 |
| 155 | Rosiglitazone blocks first trimester in-vitro placental injury caused by NF- κ B-mediated inflammation. <i>Scientific Reports</i> , 2019, 9, 2018. | 1.6 | 18 |
| 156 | Preeclampsia: Disease biology and burden, its management strategies with reference to India. <i>Pregnancy Hypertension</i> , 2019, 15, 23-31. | 0.6 | 51 |
| 157 | Regulation of CX3CL1 Expression in Human First-Trimester Decidual Cells: Implications for Preeclampsia. <i>Reproductive Sciences</i> , 2019, 26, 1256-1265. | 1.1 | 17 |
| 158 | Placental fractalkine immunoreactivity in preeclampsia and its correlation with histopathological changes in the placenta and adverse pregnancy outcomes*. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2020, 33, 806-815. | 0.7 | 14 |
| 159 | Regulation of Uterine Spiral Artery Remodeling: a Review. <i>Reproductive Sciences</i> , 2020, 27, 1932-1942. | 1.1 | 47 |
| 160 | Hypoxia and Mitochondrial Dysfunction in Pregnancy Complications. <i>Antioxidants</i> , 2021, 10, 405. | 2.2 | 33 |
| 161 | Uteroplacental Circulation in Normal Pregnancy and Preeclampsia: Functional Adaptation and Maladaptation. <i>International Journal of Molecular Sciences</i> , 2021, 22, 8622. | 1.8 | 16 |
| 162 | Immunological comparison of pregnant Dahl salt-sensitive and Sprague-Dawley rats commonly used to model characteristics of preeclampsia. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2021, 321, R125-R138. | 0.9 | 3 |
| 163 | Immunomodulatory Analogies Between Trophoblastic and Cancer Cells and Their Hosts. , 2001, , 190-208. | | 1 |
| 164 | Gadd45 Stress Sensors in Preeclampsia. <i>Advances in Experimental Medicine and Biology</i> , 2013, 793, 121-129. | 0.8 | 12 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 165 | Inhibition of TGF- β 23 restores the invasive capability of extravillous trophoblasts in preeclamptic pregnancies. <i>Journal of Clinical Investigation</i> , 1999, 103, 1641-1650. | 3.9 | 341 |
| 166 | Pro-Inflammatory Profile of Preeclamptic Placental Mesenchymal Stromal Cells: New Insights into the Etiopathogenesis of Preeclampsia. <i>PLoS ONE</i> , 2013, 8, e59403. | 1.1 | 59 |
| 167 | Placental cytokines and preeclampsia. <i>Frontiers in Bioscience - Landmark</i> , 2007, 12, 2706. | 3.0 | 50 |
| 168 | Microcirculation in Pregnancy. <i>Physiological Research</i> , 2014, 63, 395-408. | 0.4 | 10 |
| 169 | Correlation of Tumor Necrosis Factor Alpha, Interleukin 6 and Interleukin 10 With Blood Pressure, Risk of Preeclampsia and Low Birth Weight in Gestational Diabetes. <i>Physiological Research</i> , 2019, 68, 395-408. | 0.4 | 22 |
| 170 | The Role of Heme Oxygenase in Pregnancy. , 2002, , 67-82. | | 0 |
| 171 | Hypertensive Disorders in Pregnancy. , 2004, , 43-67. | | 0 |
| 172 | Magnesium Role in Cytokine Regulation of Hypoxic Placentas Related to Certain Placental Pathology. , 2013, , 51-63. | | 0 |
| 173 | Pathogenesis of Preeclampsia. <i>Comprehensive Gynecology and Obstetrics</i> , 2017, , 211-224. | 0.0 | 0 |
| 174 | The role of immune activation in contributing to vascular dysfunction and the pathophysiology of hypertension during preeclampsia. <i>Minerva Ginecologica</i> , 2010, 62, 105-20. | 0.8 | 77 |
| 175 | Endothelial dysfunction. An important mediator in the pathophysiology of hypertension during pre-eclampsia. <i>Minerva Ginecologica</i> , 2012, 64, 309-20. | 0.8 | 70 |
| 176 | Effects of tumor necrosis factor on undifferentiated and syncytialised placental choriocarcinoma BeWo cells. <i>Toxicology in Vitro</i> , 2022, 80, 105327. | 1.1 | 0 |
| 177 | Gadd45 in Preeclampsia. <i>Advances in Experimental Medicine and Biology</i> , 2022, 1360, 101-108. | 0.8 | 1 |
| 178 | LPS-mediated activation of TLR4 controls <i>Toxoplasma gondii</i> growth in human trophoblast cell (BeWo) and human villous explants in a dependent-manner of TRIF, MyD88, NF- κ B and cytokines. <i>Tissue and Cell</i> , 2022, 78, 101907. | 1.0 | 3 |
| 179 | Role of ADAM and ADAMTS disintegrin and metalloproteinases in normal pregnancy and preeclampsia. <i>Biochemical Pharmacology</i> , 2022, 206, 115266. | 2.0 | 6 |
| 180 | A Deficiency of Placental IL-10 in Preeclampsia. <i>Journal of Immunology</i> , 1999, 163, 3491-3495. | 0.4 | 199 |
| 181 | Placental Mitochondrial Function and Dysfunction in Preeclampsia. <i>International Journal of Molecular Sciences</i> , 2023, 24, 4177. | 1.8 | 11 |