

CITATION REPORT

List of articles citing

**Acute renal failure after coronary intervention:
incidence, risk factors, and relationship to mortality**

DOI: 10.1016/s0002-9343(97)00150-2

American Journal of Medicine, 1997, 103, 368-75.

Source: <https://exaly.com/paper-pdf/28286061/citation-report.pdf>

Version: 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| # | Paper | IF | Citations |
|------|--|-----|-----------|
| 1449 | Mode of Coronary Revascularization and Short term Clinical Outcomes in Patients with Chronic Kidney Disease. 2014 , 30, 1180-5 | | 2 |
| 1448 | Clarithromycin and digoxin toxicity. 1998 , 128, 604 | | 12 |
| 1447 | Positive-pressure ventilation in amyotrophic lateral sclerosis. 1998 , 128, 601-2 | | |
| 1446 | Ultrasonographic Screening for Deep Venous Thrombosis after Arthroplasty. 1998 , 128, 601 | | 2 |
| 1445 | Encouraging physical activity. 1998 , 128, 605-6 | | |
| 1444 | Hepatitis C virus genotype distribution in B-cell non-Hodgkin lymphoma. 1998 , 128, 602 | | 15 |
| 1443 | Patient care after percutaneous coronary artery interventions. 1998 , 128, 598; author reply 599-600 | | |
| 1442 | Catheter-related bacteremia in patients undergoing hemodialysis. 1998 , 128, 600 | | 6 |
| 1441 | Catheter-Related Bacteremia in Patients Undergoing Hemodialysis. 1998 , 128, 600 | | |
| 1440 | Anemia secondary to vitamin D intoxication. 1998 , 128, 602-3 | | 4 |
| 1439 | Patient Care after Percutaneous Coronary Artery Interventions. 1998 , 128, 599 | | |
| 1438 | The toxic shock syndrome and Staphylococcus lugdunensis bacteremia. 1998 , 128, 603-4 | | 28 |
| 1437 | Encouraging Physical Activity. 1998 , 128, 606 | | |
| 1436 | AL amyloidosis combined with acquired factor V deficiency. 1998 , 128, 604-5 | | 15 |
| 1435 | Aspirin and ticlopidine after routine coronary stenting: the gold standard as of 1999. 1999 , 7, 233-9 | | 3 |
| 1434 | Changes in renal function with percutaneous transluminal coronary angioplasty. <i>International Journal of Cardiology</i> , 1999 , 70, 127-31 | 3.2 | 7 |
| 1433 | Prophylaxis against contrast-induced nephropathy. 1999 , 353, 1638-9 | | 19 |

| | | |
|------|---|-----|
| 1432 | Public disclosure of performance data: does the public get what the public wants?. 1999 , 353, 1639-40 | 34 |
| 1431 | A prospective randomized trial of prevention measures in patients at high risk for contrast nephropathy: results of the P.R.I.N.C.E. Study. Prevention of Radiocontrast Induced Nephropathy Clinical Evaluation. 1999 , 33, 403-11 | 290 |
| 1430 | Early use of coronary angiography and intervention. 1999 , 17, 373-86 | 4 |
| 1429 | Radiological contrast agents. 1999 , 22, 498-506 | |
| 1428 | Influence of dialysis procedure, membrane surface and membrane material on iopromide elimination in patients with reduced kidney function. 2000 , 20, 300-4 | 24 |
| 1427 | New perspectives for prevention/treatment of acute renal failure. 2000 , 13, 105-12 | 7 |
| 1426 | Five French (5 Fr) guiding catheters for percutaneous coronary angioplasty and stent placement: An initial feasibility study. 2000 , 51, 352-7 | 12 |
| 1425 | Renal artery angioplasty and stent placement: predictors of a favorable outcome. 2000 , 139, 64-71 | 106 |
| 1424 | Nephrotoxicity related to contrast media. 2000 , 34, 317-22 | 81 |
| 1423 | An evaluation of pharmacological strategies for the prevention and treatment of acute renal failure. 2000 , 59, 79-91 | 50 |
| 1422 | Risks associated with renal dysfunction in patients in the coronary care unit. 2000 , 36, 679-84 | 196 |
| 1421 | The prognostic implications of further renal function deterioration within 48 h of interventional coronary procedures in patients with pre-existent chronic renal insufficiency. 2000 , 36, 1542-8 | 572 |
| 1420 | Postoperative renal insufficiency. 2001 , 85, 1241-54 | 3 |
| 1419 | Contrast media-induced nephrotoxicity: identification of patients at risk and algorithms for prevention. 2001 , 12, 3-9 | 162 |
| 1418 | Determinants of mortality after myocardial infarction in patients with advanced renal dysfunction. 2001 , 37, 1191-200 | 195 |
| 1417 | Severity scores and outcomes with acute renal failure in the ICU setting. 2001 , 132, 181-95 | 13 |
| 1416 | Acute renal failure requiring dialysis after percutaneous coronary interventions. 2001 , 52, 409-16 | 175 |
| 1415 | Prediction and prevention of contrast nephropathy. 2001 , 14, 547-58 | 18 |

| | | | |
|------|---|------|-----|
| 1414 | Use of fenoldopam for prevention of radiocontrast nephropathy in the cardiac catheterization laboratory: a case series. 2001 , 14, 179-85 | | 9 |
| 1413 | Prevention of contrast nephropathy after cardiac catheterisation. 2001 , 85, 361-2 | | 31 |
| 1412 | [Treatment for acute renal failure--concepts and controversies. 1. Pharmacologic treatment and experimental strategies]. 2001 , 36, 134-42 | | 3 |
| 1411 | Prevention of contrast media-associated nephropathy: randomized comparison of 2 hydration regimens in 1620 patients undergoing coronary angioplasty. 2002 , 162, 329-36 | | 557 |
| 1410 | The independent association of renal dysfunction and arrhythmias in critically ill patients. 2002 , 122, 669-77 | | 38 |
| 1409 | Coronary anatomy predicts presence or absence of renal artery stenosis. A prospective study in patients undergoing cardiac catheterization for suspected coronary artery disease. 2002 , 23, 1684-91 | | 100 |
| 1408 | Emergency evaluation of chest pain in patients with advanced kidney disease. 2002 , 162, 2464-8 | | 26 |
| 1407 | Interventional therapy for coronary artery disease. 2002 , 166, 791-6 | | 12 |
| 1406 | Role of the kidney in perioperative inflammatory responses. 2002 , 88, 330-4 | | 21 |
| 1405 | Effect of diabetes on renal medullary oxygenation during water diuresis. 2002 , 25, 575-8 | | 88 |
| 1404 | Outcomes of patients with chronic renal insufficiency in the bypass angioplasty revascularization investigation. <i>Circulation</i> , 2002 , 105, 2253-8 | 16.7 | 184 |
| 1403 | Prognostic implications of abnormalities in renal function in patients with acute coronary syndromes. <i>Circulation</i> , 2002 , 106, 974-80 | 16.7 | 379 |
| 1402 | Incidence, risk factors, and prognosis of a moderate increase in plasma creatinine early after cardiac surgery. 2002 , 30, 1495-8 | | 124 |
| 1401 | Contrast-associated nephropathy. 2002 , 4, 372-9 | | 13 |
| 1400 | Cardiovascular disease and the kidney. Tracking a killer in chronic kidney disease. 2002 , 111, 53-60 | | 31 |
| 1399 | Haemodialysis for the Prevention of Contrast-Induced Nephropathy. 2002 , 37, 471-481 | | 28 |
| 1398 | Cardiology Grand Rounds from The University of North Carolina at Chapel Hill. Contrast induced nephropathy after angiography. 2002 , 323, 252-8 | | 20 |
| 1397 | Prognosis of patients who develop acute renal failure during the first 24 hours of cardiogenic shock after myocardial infarction. <i>American Journal of Medicine</i> , 2002 , 112, 115-9 | 2.4 | 88 |

| | | |
|------|--|-----|
| 1396 | Acetylcysteine and contrast agent-associated nephrotoxicity. 2002 , 40, 298-303 | 269 |
| 1395 | Acetylcysteine protects against acute renal damage in patients with abnormal renal function undergoing a coronary procedure. 2002 , 40, 1383-8 | 235 |
| 1394 | Comparative tolerability of contrast media used for coronary interventions. 2002 , 25, 1079-98 | 13 |
| 1393 | Fenoldopam mesylate blocks reductions in renal plasma flow after radiocontrast dye infusion: a pilot trial in the prevention of contrast nephropathy. 2002 , 143, 894-903 | 176 |
| 1392 | Two vessels or not two vessels? That is the question. 2002 , 143, 948-9 | 1 |
| 1391 | Benefits of aspirin and beta-blockade after myocardial infarction in patients with chronic kidney disease. 2002 , 144, 226-32 | 176 |
| 1390 | Pseudoaccountability. 2002 , 136, 631 | |
| 1389 | Appropriate antibiotic use for acute pharyngitis. 2002 , 136, 632-3; author reply 632-3 | |
| 1388 | Long-Term Prediction of Coronary Heart Disease in Young Men. 2002 , 136, 631 | |
| 1387 | William of Occam and Occam's razor. 2002 , 136, 634-5 | 18 |
| 1386 | Correction: cost-effectiveness of statins in older patients with myocardial infarction. 2002 , 136, 635 | 3 |
| 1385 | Cardiovascular Outcomes and Renal Disease. 2002 , 136, 634 | 4 |
| 1384 | Changing trends in incidence and predictors of radiographic contrast nephropathy after percutaneous coronary intervention with use of fenoldopam. 2002 , 89, 999-1002 | 65 |
| 1383 | Aortic distensibility is increasing in elite athletes. 2002 , 89, 1002-4 | 19 |
| 1382 | Nephropathy requiring dialysis after percutaneous coronary intervention and the critical role of an adjusted contrast dose. 2002 , 90, 1068-73 | 255 |
| 1381 | A protocol for prevention of radiographic contrast nephropathy during percutaneous coronary intervention: effect of selective dopamine receptor agonist fenoldopam. 2002 , 55, 169-73 | 59 |
| 1380 | Continuous versus intermittent renal replacement therapy: a meta-analysis. 2002 , 28, 29-37 | 284 |
| 1379 | The dialytic management of acute renal failure in the elderly. 2002 , 15, 127-32 | 8 |

| | | |
|------|---|-----|
| 1378 | Determinants of serum creatinine trajectory in acute contrast nephropathy. 2002 , 15, 349-54 | 104 |
| 1377 | Contrast nephropathy or contrast creatininopathy. 2003 , 26, 53-4 | 5 |
| 1376 | Acute coronary syndromes in patients with renal failure. 2003 , 5, 266-70 | 13 |
| 1375 | Incidence and circumstances of serum creatinine increase after abdominal aortic surgery. 2003 , 29, 1821-4 | 18 |
| 1374 | Interface between heart disease and renal dysfunction: from association to action. 2003 , 12, 20-24 | |
| 1373 | Contrast agent--associated nephrotoxicity. 2003 , 45, 493-503 | 82 |
| 1372 | Meta-analysis of randomized clinical trials on the usefulness of acetylcysteine for prevention of contrast nephropathy. 2003 , 92, 1454-8 | 114 |
| 1371 | Effectiveness of theophylline prophylaxis of renal impairment after coronary angiography in patients with chronic renal insufficiency. 2003 , 91, 1157-62 | 84 |
| 1370 | Prevention of radiocontrast-induced nephropathy. 2003 , 58, 532-8 | 7 |
| 1369 | Continuous veno-venous hemofiltration for the treatment of contrast-induced acute renal failure after percutaneous coronary interventions. 2003 , 58, 59-64 | 34 |
| 1368 | Oral acetylcysteine does not protect renal function from moderate to high doses of intravenous radiographic contrast. 2003 , 58, 336-41 | 118 |
| 1367 | Percutaneous coronary intervention-associated nephropathy foreshadows increased risk of late adverse events in patients with normal baseline serum creatinine. 2003 , 59, 338-43 | 121 |
| 1366 | More contrast, no more brightness. 2003 , 59, 344-5 | |
| 1365 | Prophylaxis of contrast-induced nephropathy in patients undergoing coronary angiography. 2003 , 60, 458-61 | 72 |
| 1364 | The effect of N-acetylcysteine on renal function, nitric oxide, and oxidative stress after angiography. 2003 , 64, 2182-7 | 72 |
| 1363 | Gadolinium as a nonnephrotoxic contrast agent for catheter-based arteriographic evaluation of renal arteries in patients with azotemia. 2003 , 37, 346-52 | 36 |
| 1362 | Utilisation des produits de contraste au cours des examens radiologiques réalisés en urgence : risques et précautions à prendre / Use of contrast media in emergency investigations: risks and guidelines for their prevention. 2003 , 12, 510-518 | 1 |
| 1361 | Radiocontrast media-induced renal injury--saline is effective in prevention. 2003 , 93, C5-6 | 7 |

| | | | |
|------|--|------|------|
| 1360 | A randomized prospective trial to assess the role of saline hydration on the development of contrast nephrotoxicity. 2003 , 93, C29-34 | | 285 |
| 1359 | Did contrast nephropathy in rappid really occur? reply. 2003 , 42, 2167-2169 | | 0 |
| 1358 | Influence of concurrent renal dysfunction on outcomes of patients with acute coronary syndromes and implications of the use of glycoprotein IIb/IIIa inhibitors. 2003 , 41, 718-24 | | 140 |
| 1357 | A rapid protocol for the prevention of contrast-induced renal dysfunction: the RAPPID study. 2003 , 41, 2114-8 | | 262 |
| 1356 | Cholesterol emboli after invasive cardiac procedures. 2003 , 42, 217-8 | | 37 |
| 1355 | Kidney disease as a risk factor for development of cardiovascular disease: a statement from the American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. <i>Circulation</i> , 2003 , 108, 2154-69 | 16.7 | 2515 |
| 1354 | Acetylcysteine in the prevention of contrast-induced nephropathy after coronary angiography. 2003 , 146, E23 | | 74 |
| 1353 | Acetylcysteine for prevention of contrast nephropathy: meta-analysis. 2003 , 362, 598-603 | | 385 |
| 1352 | A drug to prevent renal failure?. 2003 , 362, 589-90 | | 14 |
| 1351 | Contrast nephropathy : an evidence-based approach to prevention. 2003 , 3, 395-405 | | 19 |
| 1350 | Kidney disease as a risk factor for development of cardiovascular disease: a statement from the American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. 2003 , 42, 1050-65 | | 802 |
| 1349 | Radiocontrast-induced nephropathy and percutaneous coronary intervention: a review of preventive measures. 2003 , 4, 639-52 | | 5 |
| 1348 | Impact of renal insufficiency in patients undergoing primary angioplasty for acute myocardial infarction. <i>Circulation</i> , 2003 , 108, 2769-75 | 16.7 | 319 |
| 1347 | Acetylcysteine for prevention of acute deterioration of renal function following elective coronary angiography and intervention: a randomized controlled trial. 2003 , 289, 553-8 | | 300 |
| 1346 | Nephrotoxic effects in high-risk patients undergoing angiography. 2003 , 348, 491-9 | | 857 |
| 1345 | Outcome in a post-cardiac surgery population with acute renal failure requiring dialysis: does age make a difference?. 2003 , 18, 732-6 | | 47 |
| 1344 | Fenoldopam mesylate for the prevention of contrast-induced nephropathy: a randomized controlled trial. 2003 , 290, 2284-91 | | 317 |
| 1343 | How to manage the renal patient with coronary heart disease: the agony and the ecstasy of opinion-based medicine. 2003 , 14, 2556-72 | | 65 |

| | | | |
|------|---|-----|-----|
| 1342 | The prevention of radiocontrast-agent-induced nephropathy by hemofiltration. 2003 , 349, 1333-40 | | 382 |
| 1341 | Radiocontrast-induced nephropathy. 2003 , 10, 137-47 | | 40 |
| 1340 | Acetylcysteine and Nephropathy. 2003 , 103, 64AA | | 1 |
| 1339 | Interventions for preventing contrast-induced nephropathy. 2003 , | | |
| 1338 | Acetylcysteine, coronary procedure and prevention of contrast-induced worsening of renal function: which benefit for which patient?. <i>Acta Cardiologica</i> , 2003 , 58, 555-60 | 0.9 | 34 |
| 1337 | N-acetylcysteine for radiocontrast-induced nephropathy: potential role in the emergency department?. 2004 , 6, 253-8 | | 2 |
| 1336 | Addendum. 2004 , 27, 1061-1062 | | |
| 1335 | Contrast nephropathy, pathophysiology and prevention. 2004 , 27, 1054-65 | | 7 |
| 1334 | . 2004 , | | 4 |
| 1333 | Complications of Cardiopulmonary Bypass and Cardioplegia. 280-348 | | |
| 1332 | Frequently asked questions: iodinated contrast agents. 2004 , 24 Suppl 1, S3-10 | | 66 |
| 1331 | N-acetylcysteine for the prevention of radiocontrast induced nephropathy: a meta-analysis of prospective controlled trials. 2004 , 15, 761-9 | | 202 |
| 1330 | Contrast-induced nephropathy. 2004 , 183, 1673-89 | | 306 |
| 1329 | Oral N-Acetylcysteine (NAC) to Prevent Contrast-Induced Nephrotoxicity: Is NAC the Answer?. 2004 , 137, 31-36 | | 1 |
| 1328 | How does minor renal dysfunction influence cardiovascular risk and the management of cardiovascular disease?. 2004 , 15, 517-23 | | 44 |
| 1327 | Prevention of radiocontrast nephropathy: back to basics. 2004 , 291, 2376-7 | | 18 |
| 1326 | Renal function and long term mortality after unstable angina/non-ST segment elevation myocardial infarction treated very early and predominantly with percutaneous coronary intervention. 2004 , 90, 902-7 | | 21 |
| 1325 | Provider use of preventive strategies for radiocontrast nephropathy in high-risk patients. 2004 , 96, c56-62 | | 13 |

| | | |
|------|---|----------|
| 1324 | N-acetylcysteine in the prevention of radiocontrast-induced nephropathy. 2004 , 15, 251-60 | 151 |
| 1323 | Prevention of contrast-induced nephropathy with sodium bicarbonate: a randomized controlled trial. 2004 , 291, 2328-34 | 781 |
| 1322 | Survival after coronary revascularization among patients with kidney disease. <i>Circulation</i> , 2004 , 110, 1890-5 | 16.7 196 |
| 1321 | Theophylline for the prevention of radiocontrast nephropathy: a meta-analysis. 2004 , 19, 2747-53 | 82 |
| 1320 | Nephrotoxic effects of X-ray contrast media. 2004 , 42, 325-32 | 21 |
| 1319 | Standard vs double dose of N-acetylcysteine to prevent contrast agent associated nephrotoxicity. 2004 , 25, 206-11 | 185 |
| 1318 | Free floating thrombus in left atrium. 2004 , 90, 907 | 1 |
| 1317 | Systematic review of the impact of N-acetylcysteine on contrast nephropathy. 2004 , 65, 1366-74 | 189 |
| 1316 | Abbreviated dosing of N-acetylcysteine prevents contrast-induced nephropathy after elective and urgent coronary angiography and intervention. 2004 , 17, 159-65 | 46 |
| 1315 | Residual renal function: considerations on its importance and preservation in dialysis patients. 2004 , 17, 196-201 | 69 |
| 1314 | Effect of N-acetylcysteine for prevention of contrast nephropathy in patients with moderate to severe renal insufficiency: a randomized trial. 2004 , 43, 801-8 | 74 |
| 1313 | Prevention of radiocontrast-induced nephropathy. 2004 , 44, 12-24 | 75 |
| 1312 | Impact of nephropathy after percutaneous coronary intervention and a method for risk stratification. 2004 , 93, 1515-9 | 360 |
| 1311 | Impact of chronic kidney disease on prognosis of patients with diabetes mellitus treated with percutaneous coronary intervention. 2004 , 94, 300-5 | 141 |
| 1310 | Prognostic significance of renal insufficiency in patients presenting with acute coronary syndrome (the Prospective Multicenter SYCOMORE study). 2004 , 94, 1543-7 | 44 |
| 1309 | Clinical and laboratory diagnosis of acute renal failure. 2004 , 18, 1-20 | 18 |
| 1308 | Percutaneous coronary intervention in diabetics. 2004 , 5, 277-85 | 6 |
| 1307 | Role of N-acetylcysteine in prevention of contrast-induced nephropathy after cardiovascular procedures: a meta-analysis. 2004 , 27, 607-10 | 46 |

| | | | |
|------|--|------|------|
| 1306 | Rotational X-ray coronary angiography. 2004 , 63, 201-7 | | 29 |
| 1305 | SCAI statement on ad hoc versus the separate performance of diagnostic cardiac catheterization and coronary intervention. 2004 , 63, 444-51 | | 16 |
| 1304 | Contrast nephropathy after coronary angiography. 2004 , 79, 211-9 | | 81 |
| 1303 | Ascorbic acid prevents contrast-mediated nephropathy in patients with renal dysfunction undergoing coronary angiography or intervention. <i>Circulation</i> , 2004 , 110, 2837-42 | 16.7 | 230 |
| 1302 | N-Acetylcysteine versus fenoldopam mesylate to prevent contrast agent-associated nephrotoxicity. 2004 , 44, 762-5 | | 75 |
| 1301 | Contrast nephropathy: review focusing on prevention. 2004 , 44, 1763-71 | | 177 |
| 1300 | Contrast-induced nephropathy in patients undergoing primary angioplasty for acute myocardial infarction. 2004 , 44, 1780-5 | | 530 |
| 1299 | Relation between renal dysfunction and cardiovascular outcomes after myocardial infarction. 2004 , 351, 1285-95 | | 1464 |
| 1298 | [Value of N-acetylcysteine to prevent nephrotoxicity from iodinated contrast agents]. 2004 , 85, 725-7 | | 3 |
| 1297 | Cardiovascular risk in stage 4 and 5 nephropathy. 2004 , 11, 116-33 | | 23 |
| 1296 | Prevention of contrast-induced nephropathy in vascular patients undergoing angiography: a randomized controlled trial of intravenous N-acetylcysteine. 2004 , 40, 1136-41 | | 75 |
| 1295 | Prevention of contrast media nephrotoxicity--the story so far. 2004 , 59, 381-9 | | 63 |
| 1294 | Renal insufficiency and mortality from acute coronary syndromes. 2004 , 147, 623-9 | | 88 |
| 1293 | Slowing the progression of diabetic nephropathy and its cardiovascular consequences. 2004 , 148, 243-51 | | 25 |
| 1292 | Is acetylcysteine effective in preventing contrast-related nephropathy? A meta-analysis. <i>American Journal of Medicine</i> , 2004 , 117, 938-47 | 2.4 | 148 |
| 1291 | A simple risk score for prediction of contrast-induced nephropathy after percutaneous coronary intervention: development and initial validation. 2004 , 44, 1393-9 | | 676 |
| 1290 | The incidence and risk of acute renal failure after cardiac surgery. 2004 , 18, 442-5 | | 197 |
| 1289 | Preventing contrast nephropathy: what is the best strategy? A review of the literature. 2004 , 44, 327-37 | | 38 |

| | | |
|------|--|---------|
| 1288 | Evidence-based practice for the use of N-acetylcysteine. 2004 , 23, 270-3 | 3 |
| 1287 | Renal disease: value of functional magnetic resonance imaging with flow and perfusion measurements. 2004 , 39, 698-705 | 64 |
| 1286 | Reducing radiocontrast use in the cardiac catheterization laboratory--a behavioral modification approach. 2004 , 13, 188-92 | 4 |
| 1285 | A simple risk score for prediction of contrast-induced nephropathy after percutaneous coronary interventionDevelopment and initial validation. 2004 , 44, 1393-1399 | 1392 |
| 1284 | Pharmacological interventions for preventing contrast-induced nephropathy. 2004 , | 1 |
| 1283 | Prevention of radiocontrast nephropathy. 2005 , 4, 274-279 | |
| 1282 | Contrast dose-to-creatinine clearance ratio as a potential indicator of risk for radiocontrast-induced nephropathy: correlation of D/CrCL with area under the contrast concentration-time curve using iodixanol. 2005 , 40, 598-603 | 37 |
| 1281 | Recent advances in the prevention of radiocontrast-induced nephropathy. 2004 , 10, 505-9 | 6 |
| 1280 | Use of iso-osmolar nonionic dimeric contrast media in multidetector row computed tomography angiography for patients with renal impairment. 2005 , 40, 672-5 | 38 |
| 1279 | Prophylactic acetylcysteine usage for prevention of contrast nephropathy after coronary angiography. 2005 , 46, 464-7 | 16 |
| 1278 | Renal replacement therapy for preventing contrast-induced nephropathy. 2005 , | 1 |
| 1277 | Cardiovascular interventional magnetic resonance imaging. <i>Circulation</i> , 2005 , 112, 3009-17 | 16.7 79 |
| 1276 | Prevention of radiocontrast nephropathy. 2005 , 14, 105-10 | 32 |
| 1275 | In which patients should serum creatinine be measured before iodinated contrast medium administration?. 2005 , 15, 749-54 | 66 |
| 1274 | The current state of nonuremic applications for extracorporeal blood purification. 2005 , 18, 380-90 | 11 |
| 1273 | Venous mapping using venography and the risk of radiocontrast-induced nephropathy. 2005 , 18, 239-42 | 45 |
| 1272 | N-acetylcysteine for the prevention of contrast-induced nephropathy. A systematic review and meta-analysis. 2005 , 20, 193-200 | 86 |
| 1271 | Nephrotoxicity of low-osmolality versus iso-osmolality contrast agents: impact of N-acetylcysteine. 2005 , 68, 2250-5 | 47 |

| | | |
|------|--|-----|
| 1270 | The role of osmolality in the incidence of contrast-induced nephropathy: a systematic review of angiographic contrast media in high risk patients. 2005 , 68, 2256-63 | 150 |
| 1269 | Low hematocrit predicts contrast-induced nephropathy after percutaneous coronary interventions. 2005 , 67, 706-13 | 124 |
| 1268 | Postprocedural drop in hematocrit versus contrast-induced nephropathy: eggs or chickens?. 2005 , 68, 1371-2; author reply 1372 | 1 |
| 1267 | Reply from the Authors. 2005 , 68, 1372 | |
| 1266 | PAI-1 in diabetic nephropathy. 2005 , 68, 1372-3; author reply 1373 | 0 |
| 1265 | Gadodiamide-based coronary angiography in a patient with severe renal insufficiency. 2005 , 18, 379-83 | 4 |
| 1264 | Contrast material-induced renal failure: an overview. 2005 , 18, 417-23 | 36 |
| 1263 | Contrast-induced nephropathy after percutaneous coronary interventions in relation to chronic kidney disease and hemodynamic variables. 2005 , 95, 13-9 | 431 |
| 1262 | The diagnosis of deep venous thrombosis and pulmonary embolism in medical-surgical intensive care unit patients. 2005 , 20, 314-9 | 15 |
| 1261 | Renal failure after percutaneous coronary intervention is associated with high mortality. 2005 , 64, 442-8 | 122 |
| 1260 | Effect of nonionic radiocontrast agents on the occurrence of contrast-induced nephropathy in patients with mild-moderate chronic renal insufficiency: pooled analysis of the randomized trials. 2005 , 65, 386-93 | 62 |
| 1259 | The clinical epidemiology of contrast-induced nephropathy. 2005 , 28 Suppl 2, S3-11 | 77 |
| 1258 | Reducing the risks for contrast-induced nephropathy. 2005 , 28 Suppl 2, S12-8 | 10 |
| 1257 | Contrast media: safety, viscosity, and volume. 2005 , 15 Suppl 4, D62-4 | 8 |
| 1256 | Safety of contrast venography prior to caval interruption in patients with renal insufficiency. 2005 , 19, 347-51 | 2 |
| 1255 | Economic analysis of the use of contrast media during percutaneous coronary interventions in France and Spain. 2005 , 8, 111-130 | |
| 1254 | Uroangiographic Contrast Media-Induced Nephropathy: Correlations between Their Physicochemical Properties and Renal Damage. 2005 , 72, 446-456 | |
| 1253 | Potential mechanisms behind contrast medium-induced nephropathy. 2005 , 9, 17 | |

| | | | |
|------|--|------|-----|
| 1252 | Late Adverse Events Following Administration of Iodinated Contrast Media: An Update. 2005 , 121-131 | | |
| 1251 | Contrast nephropathy post cardiac resynchronization therapy: an under-recognized complication with important morbidity. 2005 , 7, 899-903 | | 31 |
| 1250 | Contrast-medium-Induced nephropathy correlated to the ratio between dose in gram iodine and estimated GFR in ml/min. 2005 , 46, 830-42 | | 88 |
| 1249 | Lights and shadows on the pathogenesis of contrast-induced nephropathy: state of the art. 2005 , 20, 1542-50 | | 118 |
| 1248 | N-acetylcysteine does not prevent contrast induced nephropathy after cardiac catheterisation with an ionic low osmolality contrast medium: a multicentre clinical trial. 2005 , 91, 774-8 | | 51 |
| 1247 | Dobutamine stress echocardiography and cardiac troponin T for the detection of significant coronary artery disease and predicting outcome in renal transplant candidates. 2005 , 6, 327-35 | | 35 |
| 1246 | Contrast-induced nephropathy: what we know, what we think we know, and what we don't know. 2005 , 22, 108-13 | | 2 |
| 1245 | Contrast medium-induced nephrotoxicity: which pathway?. 2005 , 235, 752-5 | | 57 |
| 1244 | Radiocontrast-induced acute renal failure. 2005 , 20, 63-75 | | 71 |
| 1243 | Transoesophageal echocardiography for the identification of a giant aortic thrombus. 2005 , 91, 778 | | 1 |
| 1242 | Complications of Percutaneous Coronary Interventions. 2005 , | | 1 |
| 1241 | The ACIST power injection system reduces the amount of contrast media delivered to the patient, as well as fluoroscopy time, during diagnostic and interventional cardiac procedures. 2005 , 7, 183-7 | | 19 |
| 1240 | Effect of pravastatin on rate of kidney function loss in people with or at risk for coronary disease. <i>Circulation</i> , 2005 , 112, 171-8 | 16.7 | 161 |
| 1239 | Preventing in-hospital cardiac and renal complications in high-risk PCI patients. 2005 , 7, G13-G24 | | 2 |
| 1238 | Prehydration alone is sufficient to prevent contrast-induced nephropathy after day-only angiography procedures--a randomised controlled trial. 2005 , 14, 245-51 | | 21 |
| 1237 | [Is dialysis efficient in the prevention of ionic contrast agent-induced nephropathy?]. 2005 , 34, 803-8 | | 1 |
| 1236 | Contrast-induced nephropathy. 2005 , 21, 261-80 | | 139 |
| 1235 | Renal effects of CO2 and iodinated contrast media in patients undergoing renovascular intervention: a prospective, randomized study. 2005 , 16, 57-65 | | 37 |

| | | | |
|------|--|------|-----|
| 1234 | Bench-to-bedside review: preventive measures for contrast-induced nephropathy in critically ill patients. 2005 , 9, 361-70 | | 12 |
| 1233 | Short- and long-term renal outcomes of immediate prophylactic hemodialysis after cardiovascular catheterizations in patients with severe renal insufficiency. <i>International Journal of Cardiology</i> , 2005 , 101, 407-13 | 3.2 | 16 |
| 1232 | Contrast-induced nephropathy: a review. 2005 , 6, 82-8 | | 31 |
| 1231 | Diagnostic clinique et biologique de l'insuffisance rénale aiguë 2005 , 2, 31-45 | | |
| 1230 | Association of chronic kidney disease with clinical outcomes after coronary revascularization: the Arterial Revascularization Therapies Study (ARTS). 2005 , 149, 512-9 | | 146 |
| 1229 | Outcomes in patients admitted for chest pain with renal failure and troponin I elevations. 2005 , 150, 674-80 | | 13 |
| 1228 | Statin therapy reduces contrast-induced nephropathy: an analysis of contemporary percutaneous interventions. <i>American Journal of Medicine</i> , 2005 , 118, 843-9 | 2.4 | 165 |
| 1227 | [Which are the characteristics of the surgical situations at risk of acute renal failure?]. 2005 , 24, 177-81 | | 1 |
| 1226 | La protection rénale préopératoire. Perioperative acute renal failure. Recommandations du jury □ Texte long. 2005 , 24, 87-119 | | 5 |
| 1225 | Importancia de los factores clínicos y anatómicos en el intervencionismo coronario. 2005 , 58, 430-441 | | 0 |
| 1224 | Influence of Clinical and Anatomical Factors on the Outcome of Percutaneous Coronary Interventions. 2005 , 58, 430-441 | | |
| 1223 | Percutaneous coronary interventions in the high-risk renal patient: strategies for renal protection and vascular protection. 2005 , 23, 299-310 | | 5 |
| 1222 | Nephropathy induced by contrast media: pathogenesis, risk factors and preventive strategies. 2005 , 172, 1461-71 | | 243 |
| 1221 | Prevention of contrast media-induced nephrotoxicity after angiographic procedures. 2005 , 16, 13-23 | | 74 |
| 1220 | Iodinated contrast induced renal vasoconstriction is due in part to the downregulation of renal cortical and medullary nitric oxide synthesis. 2006 , 44, 383-91 | | 25 |
| 1219 | Renal safety of iodixanol. 2006 , 4, 655-61 | | 9 |
| 1218 | Contrast-induced nephropathy: a clinical and evidence-based approach. <i>Circulation</i> , 2006 , 113, 1799-806 | 16.7 | 326 |
| 1217 | Risk factors and therapy strategies for contrast-induced nephropathy. 2006 , 28, 365-81 | | 13 |

| | | | |
|------|--|-----|-----|
| 1216 | Prevention and treatment of contrast-induced nephropathy. 2006 , 9, 42-9 | | 20 |
| 1215 | Failure of ICD-9-CM codes to identify patients with comorbid chronic kidney disease in diabetes. 2006 , 41, 564-80 | | 138 |
| 1214 | Concomitant unilateral internal iliac artery embolization and endovascular infrarenal aortic aneurysm repair. 2006 , 43, 903-7 | | 46 |
| 1213 | Drug-associated renal dysfunction. 2006 , 22, 357-74, viii | | 95 |
| 1212 | Prognostic Value of Serum Creatinine in Non-ST-Elevation Acute Coronary Syndrome. 2006 , 59, 209-216 | | 1 |
| 1211 | Valor pronóstico de la creatinina sérica en el síndrome coronario agudo sin elevación del segmento ST. 2006 , 59, 209-216 | | 11 |
| 1210 | Clinical practice. Preventing nephropathy induced by contrast medium. 2006 , 354, 379-86 | | 455 |
| 1209 | A novel contrast removal system from the coronary sinus using an adsorbing column during coronary angiography in a porcine model. 2006 , 47, 1866-70 | | 16 |
| 1208 | A meta-analysis of the renal safety of isosmolar iodixanol compared with low-osmolar contrast media. 2006 , 48, 692-9 | | 254 |
| 1207 | Renal toxicity evaluation and comparison between visipaque (iodixanol) and hexabrix (ioxaglate) in patients with renal insufficiency undergoing coronary angiography: the RECOVER study: a randomized controlled trial. 2006 , 48, 924-30 | | 220 |
| 1206 | Contrast-induced nephropathy--prevention and risk reduction. 2006 , 21, i11-23 | | 62 |
| 1205 | Comparison of two hemofiltration protocols for prevention of contrast-induced nephropathy in high-risk patients. <i>American Journal of Medicine</i> , 2006 , 119, 155-62 | 2.4 | 143 |
| 1204 | Reply. <i>American Journal of Medicine</i> , 2006 , 119, e17-e18 | 2.4 | 35 |
| 1203 | Impact of renal insufficiency on outcome after contemporary percutaneous coronary intervention. 2006 , 151, 146-52 | | 76 |
| 1202 | Acetylcysteine In Diabetes (AID): a randomized study of acetylcysteine for the prevention of contrast nephropathy in diabetics. 2006 , 151, 1032.e9-12 | | 43 |
| 1201 | European Society of Urogenital Radiology (ESUR) guidelines on the safe use of iodinated contrast media. 2006 , 60, 307-13 | | 63 |
| 1200 | Hypertension is an independent risk factor for contrast nephropathy after percutaneous coronary intervention. <i>International Journal of Cardiology</i> , 2006 , 110, 237-41 | 3.2 | 16 |
| 1199 | Tratamento endovascular do aneurisma da aorta abdominal em paciente com insuficiência renal crônica. 2006 , 5, 325-330 | | 1 |

| | | |
|------|--|-----|
| 1198 | Insuffisance et aggression rñales aiguë pñiopñatoires. 2006 , 1, 1-11 | |
| 1197 | Toxicitñ rñale des produits de contraste iodé. 2006 , 1, 1-6 | |
| 1196 | Cost-effectiveness of hemofiltration to prevent contrast nephropathy in patients with chronic kidney disease. 2006 , 34, 1044-51 | 25 |
| 1195 | Facts and fallacies concerning the prevention of contrast medium-induced nephropathy. 2006 , 34, 2060-8 | 41 |
| 1194 | Contrast-induced nephropathy in urgent coronary interventions. 2006 , 7, 737-41 | 14 |
| 1193 | [Carboplatin plus gemcitabine therapy versus carboplatin plus weekly paclitaxel therapy for advanced non-small cell lung cancer]. 2006 , 33, 773-7 | 7 |
| 1192 | Contrast media and nephropathy: findings from systematic analysis and Food and Drug Administration reports of adverse effects. 2006 , 41, 651-60 | 94 |
| 1191 | Renal replacement therapy in the treatment of acute renal failure-intermittent and continuous. 2006 , 19, 455-64 | 12 |
| 1190 | Coronary venous capture of contrast during angiography. 2006 , 19, 401-4 | 2 |
| 1189 | Magnetic navigation in percutaneous coronary intervention. 2006 , 19, 558-65 | 29 |
| 1188 | Contrast-induced nephropathy: definition, epidemiology, and patients at risk. 2006 , S11-5 | 517 |
| 1187 | Selection of contrast media. 2006 , S39-45 | 20 |
| 1186 | Follow-up of patients with contrast-induced nephropathy. 2006 , S46-50 | 45 |
| 1185 | BOLD-MRI for the assessment of renal oxygenation in humans: acute effect of nephrotoxic xenobiotics. 2006 , 70, 144-50 | 91 |
| 1184 | Renal failure in 57 925 patients undergoing coronary procedures using iso-osmolar or low-osmolar contrast media. 2006 , 70, 1811-7 | 114 |
| 1183 | Eighty-peak kilovoltage 16-channel multidetector computed tomography and reduced contrast-medium doses tailored to body weight to diagnose pulmonary embolism in azotaemic patients. 2006 , 16, 1165-76 | 50 |
| 1182 | Is radiocontrast-induced nephropathy avoidable? The scientific evidence in 2006. 2006 , 1, 103-118 | |
| 1181 | Concomitant drugs with exposure to contrast media. 2006 , S20-4 | 16 |

| | | |
|------|--|-----|
| 1180 | Renal complications associated with contrast media. 2006 , 16, D11-D16 | 1 |
| 1179 | Should we screen MDCT patients for risk of CIN?. 2006 , 16, D17-D21 | |
| 1178 | Safety of gadodiamide mixed with a small quantity of iohexol in patients with impaired renal function undergoing coronary angiography. 2006 , 21, 141-5 | 6 |
| 1177 | Iloprost for prevention of contrast-mediated nephropathy in high-risk patients undergoing a coronary procedure. Results of a randomized pilot study. 2006 , 62, 589-95 | 19 |
| 1176 | Acetazolamide for prevention of contrast-induced nephropathy: a new use for an old drug. 2006 , 27, 238-42 | 37 |
| 1175 | ESUR guidelines on contrast media. 2006 , 31, 131-40 | 50 |
| 1174 | Empfehlungen zur Diagnostik und Behandlung von Patienten mit koronarer Herzkrankheit und Niereninsuffizienz. 2006 , 1, 8-30 | 3 |
| 1173 | Empfehlungen zur Diagnostik und Behandlung von Patienten mit koronarer Herzkrankheit und Niereninsuffizienz. 2006 , 1, 103-117 | 1 |
| 1172 | Effect of a rapid intra-arterial infusion of dextrose 5% prior to coronary angiography on frequency of contrast-induced nephropathy in high-risk patients. 2006 , 97, 981-3 | 16 |
| 1171 | Epidemiology and prognostic implications of contrast-induced nephropathy. 2006 , 98, 5K-13K | 286 |
| 1170 | Risk prediction of contrast-induced nephropathy. 2006 , 98, 27K-36K | 280 |
| 1169 | Contrast medium use. 2006 , 98, 42K-58K | 119 |
| 1168 | Gadolinium-based contrast agents and nephrotoxicity in patients undergoing coronary artery procedures. 2006 , 67, 175-80 | 79 |
| 1167 | Hyperuricemia as a risk factor for contrast-induced nephropathy in patients with chronic kidney disease. 2006 , 67, 227-35 | 42 |
| 1166 | Outcomes of contrast-induced nephropathy: experience in patients undergoing cardiovascular intervention. 2006 , 67, 335-43 | 75 |
| 1165 | Effect of radio contrast media on residual renal function in peritoneal dialysis patients--a prospective study. 2006 , 21, 1334-9 | 48 |
| 1164 | Prophylaxis of contrast material-induced nephropathy in patients in intensive care: acetylcysteine, theophylline, or both? A randomized study. 2006 , 239, 793-804 | 73 |
| 1163 | Gadolinium-based contrast media for multi-detector row spiral CT pulmonary angiography in patients with renal insufficiency. 2006 , 238, 1077; author reply 1077-8 | 3 |

| | | |
|------|---|----------|
| 1162 | Pediatric radiology. 2006 , 238, 1072-3; author reply 1073-4 | 1 |
| 1161 | IV contrast administration for CT: a survey of practices for the screening and prevention of contrast nephropathy. 2006 , 186, 1651-8 | 60 |
| 1160 | Computer-aided detection with screening mammography: improving performance or simply shifting the operating point?. 2006 , 239, 916-7; author reply 917-8 | 6 |
| 1159 | In vitro study of various iodinated contrast media. 2006 , 239, 918-9; author reply 919-21 | |
| 1158 | The clinical and renal consequences of contrast-induced nephropathy. 2006 , 21, i2-10 | 115 |
| 1157 | Preventing nephropathy induced by contrast medium. 2006 , 354, 1853-5; author reply 1853-5 | 14 |
| 1156 | N-acetylcysteine and contrast-induced nephropathy. 2006 , 355, 1497-8; author reply 1499-500 | 1 |
| 1155 | Associations of increases in serum creatinine with mortality and length of hospital stay after coronary angiography. 2006 , 17, 2871-7 | 194 |
| 1154 | Radiocontrast-Induced Acute Renal Failure—Impact beyond the Acute Phase. 2006 , 17, 595-599 | 2 |
| 1153 | Prophylaxis strategies for contrast-induced nephropathy. 2006 , 295, 2765-79 | 283 |
| 1152 | Risk factors for contrast-induced nephropathy. 2006 , 29, 84-93 | 42 |
| 1151 | Propionyl-L-carnitine therapy: effects on endothelin-1 and homocysteine levels in patients with peripheral arterial disease and end-stage renal disease. 2006 , 29, 100-7 | 17 |
| 1150 | N-acetylcysteine and contrast-induced nephropathy in primary angioplasty. 2006 , 354, 2773-82 | 474 |
| 1149 | Renal Insufficiency Following Contrast Media Administration Trial (REMEDIAL): a randomized comparison of 3 preventive strategies. <i>Circulation</i> , 2007 , 115, 1211-7 | 16.7 364 |
| 1148 | N-acetylcysteine and sodium bicarbonate versus N-acetylcysteine and standard hydration for the prevention of radiocontrast-induced nephropathy following coronary angiography. 2007 , 41, 46-50 | 30 |
| 1147 | N-acetylcysteine reduces urinary albumin excretion following contrast administration: evidence of biological effect. 2007 , 22, 2520-4 | 20 |
| 1146 | Comparison between gadolinium and iodine contrast for percutaneous intervention in atherosclerotic renal artery stenosis: clinical outcomes. 2008 , 23, 1233-40 | 32 |
| 1145 | N-acetylcysteine and/or theophylline as a preventive measure for contrast agent induced nephropathy: we should use better markers for detection of renal function. 2007 , 245, 918-9; author reply 918-9 | 0 |

| | | | |
|------|--|------|----|
| 1144 | [Nephrogenic systemic fibrosis--a new interdisciplinary challenge]. 2007 , 132, 2706-10 | | 3 |
| 1143 | Prevention of Contrast Induced Nephropathy. 2007 , 2, 3-9 | | |
| 1142 | Mortality, kidney disease and cardiac procedures following acute coronary syndrome. 2008 , 23, 934-40 | | 28 |
| 1141 | Aortic root catheter-directed coronary CT angiography in swine: coronary enhancement with minimum volume of iodinated contrast material. 2007 , 188, W415-22 | | 5 |
| 1140 | Contrast-induced nephropathy. 2007 , 99, 474-83 | | 61 |
| 1139 | Targeted renal therapy and contrast-induced nephropathy during endovascular abdominal aortic aneurysm repair: results of a feasibility pilot trial. 2007 , 14, 520-7 | | 20 |
| 1138 | Contrast-induced nephropathy. 2007 , 14, 92-100 | | 31 |
| 1137 | ACCF/AHA 2007 clinical competence statement on vascular imaging with computed tomography and magnetic resonance: a report of the American College of Cardiology Foundation/American Heart Association/American College of Physicians Task Force on Clinical Competence and Training: developed in collaboration with the Society of Atherosclerosis Imaging and Prevention, the Society for Cardiovascular Angiography and Interventions, the Society of Cardiovascular Computed Tomography, the Society for Cardiovascular Magn. 2007 , 116, 1318-35 | 16.7 | 18 |
| 1136 | Preventing contrast nephropathy in catheter laboratory. 2007 , 93, 654-5 | | 2 |
| 1135 | ACCF/AHA 2007 Clinical Competence Statement on vascular imaging with computed tomography and magnetic resonance. 2007 , 12, 359-78 | | 7 |
| 1134 | Risk markers for contrast-induced nephropathy. 2007 , 334, 283-90 | | 45 |
| 1133 | Iodixanol: risk of subsequent contrast nephropathy in cancer patients with underlying renal insufficiency undergoing diagnostic computed tomography examinations. 2007 , 31, 493-8 | | 24 |
| 1132 | European Society of Urogenital Radiology guidelines on contrast media application. 2007 , 17, 70-6 | | 50 |
| 1131 | Determinants of amount of contrast utilized in patients undergoing percutaneous coronary procedures. 2007 , 18, 275-82 | | 4 |
| 1130 | The incidence and prognostic significance of acute kidney injury. 2007 , 16, 227-36 | | 90 |
| 1129 | Prevention of contrast-induced nephropathy: a critical review. 2007 , 16, 336-47 | | 32 |
| 1128 | Radiocontrast nephropathy. 2007 , 13, 26-31 | | |
| 1127 | Renal replacement strategies in the ICU. 2007 , 132, 1379-88 | | 51 |

| | | | |
|------|---|-----|-----|
| 1126 | Prognostic value of dobutamine stress echocardiography in patients with chronic kidney disease. 2007 , 153, 385-91 | | 52 |
| 1125 | Sodium bicarbonate, N-acetylcysteine, and saline for prevention of radiocontrast-induced nephropathy. A comparison of 3 regimens for protecting contrast-induced nephropathy in patients undergoing coronary procedures. A single-center prospective controlled trial. 2007 , 154, 539-44 | | 149 |
| 1124 | Discordance between serum creatinine and creatinine clearance for identification of ED patients with abdominal pain at risk for contrast-induced nephropathy. 2007 , 25, 268-72 | | 24 |
| 1123 | Patients at high risk of adverse events from intravenous contrast media after computed tomography examination. 2007 , 62, 26-32 | | 4 |
| 1122 | Physiological effects of contrast media for use in multidetector row computed tomography. 2007 , 62, 14-25 | | 6 |
| 1121 | Efficacy of N-acetylcysteine and hydration versus placebo and hydration in decreasing contrast-induced renal dysfunction in patients undergoing coronary angiography with or without concomitant percutaneous coronary intervention. 2007 , 36, 195-204 | | 20 |
| 1120 | Intravenous N-acetylcysteine for preventing contrast-induced nephropathy: a randomised trial. <i>International Journal of Cardiology</i> , 2007 , 115, 57-62 | 3-2 | 47 |
| 1119 | Trimetazidine in the prevention of contrast-induced nephropathy after coronary procedures. 2007 , 93, 698-702 | | 47 |
| 1118 | Economic burden of contrast-induced nephropathy: implications for prevention strategies. 2007 , 10, 119-34 | | 82 |
| 1117 | Imaging diagnosis of acute pulmonary embolism. 2007 , 5, 519-29 | | 7 |
| 1116 | [Contrast-induced nephropathy and acute renal failure following emergent cardiac catheterization: incidence, risk factors and prognosis]. 2007 , 60, 1026-34 | | 43 |
| 1115 | Volume-to-creatinine clearance ratio: a pharmacokinetically based risk factor for prediction of early creatinine increase after percutaneous coronary intervention. 2007 , 50, 584-90 | | 275 |
| 1114 | Renal protection for coronary angiography in advanced renal failure patients by prophylactic hemodialysis. A randomized controlled trial. 2007 , 50, 1015-20 | | 74 |
| 1113 | ACCF/AHA 2007 clinical competence statement on vascular imaging with computed tomography and magnetic resonance. A report of the American College of Cardiology Foundation/American Heart Association/American College of Physicians Task Force on Clinical Competence and Training. 2007 , 50, 1097-114 | | 25 |
| 1112 | Percutaneous Coronary Revascularization in Patients with End-Stage Renal Disease. 2007 , 19-50 | | |
| 1111 | . 2007 , | | 1 |
| 1110 | Prevention of contrast induced nephropathy: recommendations for the high risk patient undergoing cardiovascular procedures. 2007 , 69, 135-40 | | 122 |
| 1109 | The impact of advanced chronic kidney disease on in-hospital mortality following percutaneous coronary intervention for acute myocardial infarction. 2007 , 70, 701-5 | | 16 |

| | | |
|------|---|-----|
| 1108 | Acute myocardial infarction percutaneous coronary intervention in the patient with chronic renal disease--two strikes trying to avoid the third. 2007 , 70, 706-7 | 0 |
| 1107 | Evidence-based emergency medicine review. Prevention of contrast-induced nephropathy in the emergency department. 2007 , 50, 335-45, 345.e1-2 | 20 |
| 1106 | Nephrogenic systemic fibrosis: is any contrast safe in renal failure?. 2007 , 37, 429-31 | 9 |
| 1105 | Can continuous venovenous hemofiltration prevent contrast-induced nephropathy: is the dye already cast?. 2007 , 20, 93-5 | 2 |
| 1104 | Contrast nephropathy following computed tomography angiography of the chest for pulmonary embolism in the emergency department. 2007 , 5, 50-4 | 106 |
| 1103 | Contrast-induced nephropathy remains a serious complication of PCI. 2007 , 20, 236-40 | 26 |
| 1102 | A meta-analysis of N-acetylcysteine in contrast-induced nephrotoxicity: unsupervised clustering to resolve heterogeneity. 2007 , 5, 32 | 108 |
| 1101 | Comparison of usefulness of sodium bicarbonate versus sodium chloride to prevent contrast-induced nephropathy in patients undergoing an emergent coronary procedure. 2007 , 100, 781-6 | 131 |
| 1100 | References. 2007 , 49, S160-S179 | |
| 1099 | The role of CT and MRI in the assessment of peripheral vascular disease. 2007 , 9, 412-9 | 4 |
| 1098 | [Iodinated contrast agent-induced nephropathy]. 2007 , 47, 761-7 | 6 |
| 1097 | Conservative outpatient renoprotective protocol in patients with low GFR undergoing contrast angiography: a case series. 2007 , 11, 209-213 | 14 |
| 1096 | Conventional or Gadolinium containing contrast media: the choice between acute renal failure or Nephrogenic Systemic Fibrosis?. 2007 , 119, 271-5 | 9 |
| 1095 | A randomized controlled trial comparing hydration therapy to additional hemodialysis or N-acetylcysteine for the prevention of contrast medium-induced nephropathy: the Dialysis-versus-Diuresis (DVD) Trial. 2007 , 96, 130-9 | 67 |
| 1094 | Low rate of contrast-induced Nephropathy after CT perfusion and CT angiography in acute stroke patients. 2007 , 254, 1491-7 | 60 |
| 1093 | Kardiovaskuläre Erkrankungen bei Patienten mit Nierenerkrankungen. 2007 , 2, S9-S14 | 0 |
| 1092 | The effect of withdrawal of ACE inhibitors or angiotensin receptor blockers prior to coronary angiography on the incidence of contrast-induced nephropathy. 2008 , 40, 749-55 | 75 |
| 1091 | Using proteomics to identify preprocedural risk factors for contrast induced nephropathy. 2008 , 2, 1058-1064 | 17 |

| | | |
|------|---|--------|
| 1090 | Contrast-induced nephropathy. 2008 , 71, 62-72 | 78 |
| 1089 | The incidence of clinically significant contrast-induced nephropathy following non-emergent coronary angiography. 2008 , 71, 879-85 | 13 |
| 1088 | Transient and persistent renal dysfunction are predictors of survival after percutaneous coronary intervention: insights from the Dartmouth Dynamic Registry. 2008 , 72, 347-354 | 83 |
| 1087 | A prospective, double-blind, randomized, controlled trial on the efficacy and cardiorenal safety of iodixanol vs. iopromide in patients with chronic kidney disease undergoing coronary angiography with or without percutaneous coronary intervention. 2008 , 72, 958-65 | 42 |
| 1086 | B-Natriuretic Peptide for the Prevention of Radiocontrast-Induced Nephropathy. 2008 , 37, 168-173 | 1 |
| 1085 | Development and evaluation of a liquid chromatography-mass spectrometry assay and its application for the assessment of renal function. 2008 , 1189, 410-6 | 13 |
| 1084 | Contrast-Induced Nephropathy: A Review Focusing on X-ray and CT Applications and the Iso-Osmolar Controversy. 2008 , 11, 1-7 | |
| 1083 | Contrast-induced nephropathy. 2008 , 21, 74-85 | 75 |
| 1082 | Major adverse noncardiac events after PCI as predictors of long-term mortality. 2008 , 21, 395-402 | 3 |
| 1081 | The effect of mannitol on renal function after cardiopulmonary bypass in patients with established renal dysfunction. 2008 , 63, 701-4 | 42 |
| 1080 | A comparison of contemporary definitions of contrast nephropathy in patients undergoing percutaneous coronary intervention and a proposal for a novel nephropathy grading system. 2008 , 101, 812-9 | 140 |
| 1079 | Incidence, predictors, and outcomes of post-percutaneous coronary intervention nephropathy in patients with diabetes mellitus and normal baseline serum creatinine levels. 2008 , 101, 1544-9 | 23 |
| 1078 | Removal of contrast medium from the coronary sinus during coronary angiography: feasibility of a simple and available technique for the prevention of nephropathy. 2008 , 9, 9-13 | 14 |
| 1077 | Renal effects of contrast media in diabetic patients undergoing diagnostic or interventional coronary angiography. 2008 , 22, 171-7 | 34 |
| 1076 | Contrast-induced acute kidney injury. 2008 , 51, 1419-28 | 703 |
| 1075 | Clinical relevance. 2008 , 3-28 | 1 |
| 1074 | Drug-associated acute kidney injury in the intensive care unit. 2008 , 29-41 | 2 |
| 1073 | Clinical outcomes of contrast-induced nephropathy in patients undergoing percutaneous coronary intervention: a prospective, multicenter, randomized study to analyze the effect of hydration and acetylcysteine. <i>International Journal of Cardiology</i> , 2008 , 126, 407-13 | 3.2 72 |

| | | | |
|------|--|-----|-----|
| 1072 | Impact of saphenous vein graft radiographic markers on clinical events and angiographic parameters. 2008 , 85, 520-4 | | 5 |
| 1071 | Prevention of radiocontrast medium-induced nephropathy using short-term high-dose simvastatin in patients with renal insufficiency undergoing coronary angiography (PROMISS) trial--a randomized controlled study. 2008 , 155, 499.e1-8 | | 126 |
| 1070 | Current role of sodium bicarbonate-based preprocedural hydration for the prevention of contrast-induced acute kidney injury: a meta-analysis. 2008 , 156, 414-21 | | 87 |
| 1069 | Reducing the risk of iodine-based and MRI contrast media administration: recommendation for a questionnaire at the time of booking. 2008 , 66, 225-9 | | 14 |
| 1068 | Radiologists' knowledge and perceptions of the impact of contrast-induced nephropathy and its risk factors when performing computed tomography examinations: a survey of European radiologists. 2008 , 66, 235-45 | | 17 |
| 1067 | Predictors for contrast media-induced nephropathy and long-term survival: prospectively assessed data from the randomized controlled Dialysis-Versus-Diuresis (DVD) trial. 2008 , 24, 845-50 | | 41 |
| 1066 | Radiographic contrast-induced nephropathy and patient mortality. 2008 , 83, 1412; author reply 1413 | | 1 |
| 1065 | Protection rñale et administration de produits de contraste. 2008 , 12, 328-334 | | |
| 1064 | Mortality associated with nephropathy after radiographic contrast exposure. 2008 , 83, 1095-100 | | 126 |
| 1063 | Does renin-angiotensin aldosterone system blockade exacerbate contrast-induced nephropathy in patients with chronic kidney disease? A prospective 50-month Mayo Clinic study. 2008 , 30, 67-72 | | 21 |
| 1062 | Contrast medium dose-to-GFR ratio: a measure of systemic exposure to predict contrast-induced nephropathy after percutaneous coronary intervention. 2008 , 49, 658-67 | | 102 |
| 1061 | Incidence and outcomes of contrast-induced AKI following computed tomography. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008 , 3, 1274-81 | 6.9 | 150 |
| 1060 | Sodium bicarbonate is associated with an increased incidence of contrast nephropathy: a retrospective cohort study of 7977 patients at mayo clinic. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008 , 3, 10-8 | 6.9 | 94 |
| 1059 | Incidence of contrast-induced nephropathy with volume supplementation--insights from a large cohort. 2008 , 17, 409-14 | | 5 |
| 1058 | Prevention, incidence, and outcomes of contrast-induced acute kidney injury. 2008 , 168, 1325-32 | | 77 |
| 1057 | Contrast-induced nephropathy: what are the true clinical consequences?. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008 , 3, 263-72 | 6.9 | 94 |
| 1056 | Acute renal haemodynamic effects of radiocontrast media in patients undergoing left ventricular and coronary angiography. 2008 , 23, 1588-94 | | 19 |
| 1055 | Protective effect of erythropoietin against radiocontrast-induced renal tubular epithelial cell injury. 2008 , 28, 203-9 | | 21 |

| | | | |
|------|--|-----|-----|
| 1054 | Acute kidney injury in the diabetic rat: studies in the isolated perfused and intact kidney. 2008 , 28, 831-9 | | 21 |
| 1053 | Diagnosis, epidemiology and outcomes of acute kidney injury. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008 , 3, 844-61 | 6.9 | 345 |
| 1052 | Contrast agents and renal cell apoptosis. 2008 , 29, 2569-76 | | 154 |
| 1051 | Prevention of contrast-induced nephropathy with volume expansion. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008 , 3, 273-80 | 6.9 | 128 |
| 1050 | Contrast-induced acute kidney injury: is there a risk after intravenous contrast?. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008 , 3, 1242-3 | 6.9 | 48 |
| 1049 | Contrast-induced nephropathy: the wheel has turned 360 degrees. 2008 , 49, 646-57 | | 89 |
| 1048 | Activation of hypoxia-inducible factors ameliorates hypoxic distal tubular injury in the isolated perfused rat kidney. 2008 , 23, 3472-8 | | 76 |
| 1047 | What are radiologists doing to prevent contrast-induced nephropathy (CIN) compared with measures supported by current evidence? A survey of European radiologists on CIN associated with computed tomography. 2008 , 49, 310-20 | | 15 |
| 1046 | Renal safety of CT angiography and perfusion imaging in the emergency evaluation of acute stroke. 2008 , 29, 1826-30 | | 99 |
| 1045 | Adaptation to hypoxia in the diabetic rat kidney. 2008 , 73, 34-42 | | 142 |
| 1044 | Iso-osmolality versus low-osmolality iodinated contrast medium at intravenous contrast-enhanced CT: effect on kidney function. 2008 , 248, 97-105 | | 94 |
| 1043 | Contrast volume reduction with superior vena cava catheter-directed coronary CT angiography: comparison with peripheral i.v. contrast enhancement in a swine model. 2008 , 190, W247-54 | | 4 |
| 1042 | Which study when? Iodinated contrast-enhanced CT versus gadolinium-enhanced MR imaging. 2008 , 249, 9-15 | | 34 |
| 1041 | The PREDICT study: a randomized double-blind comparison of contrast-induced nephropathy after low- or isoosmolar contrast agent exposure. 2008 , 191, 151-7 | | 121 |
| 1040 | Diabetes and radiocontrast media increase endothelin converting enzyme-1 in the kidney. 2008 , 74, 91-100 | | 30 |
| 1039 | Contrast-induced nephropathy: contrast material not required?. 2008 , 191, 383-6 | | 45 |
| 1038 | The epidemiology of drug-induced disorders: the kidney. 2008 , 7, 679-90 | | 56 |
| 1037 | Valuable markers for contrast-induced nephropathy in patients undergoing cardiac catheterization. 2008 , 72, 1499-505 | | 57 |

| | | |
|------|--|-----|
| 1036 | Sodium bicarbonate improves long-term clinical outcomes compared with sodium chloride in patients with chronic kidney disease undergoing an emergent coronary procedure. 2008 , 72, 1610-4 | 13 |
| 1035 | Adverse reactions to Iodinated Contrast Media. 2008 , 97-106 | 1 |
| 1034 | Renal Stem Cells. 2008 , 419-443 | 1 |
| 1033 | Meta-analysis: effectiveness of drugs for preventing contrast-induced nephropathy. 2008 , 148, 284-94 | 291 |
| 1032 | CT in the ED: Special Considerations. 399-403 | |
| 1031 | Prevention of intravenous contrast-induced nephropathy in hospital inpatients. 2008 , 7, 1-4 | 4 |
| 1030 | The ACTIVE Trial: comparison of the effects on renal function of iomeprol-400 and iodixanol-320 in patients with chronic kidney disease undergoing abdominal computed tomography. 2008 , 43, 170-8 | 114 |
| 1029 | Acute kidney injury with iodinated contrast. 2008 , 36, S204-11 | 59 |
| 1028 | The incidence of contrast-induced nephropathy in trauma patients. 2008 , 15, 134-9 | 48 |
| 1027 | Clinical significance and preventive strategies for contrast-induced nephropathy. 2008 , 17, 616-23 | 34 |
| 1026 | Renal Insufficiency Following Radiocontrast Exposure Trial (REINFORCE): a randomized comparison of sodium bicarbonate versus sodium chloride hydration for the prevention of contrast-induced nephropathy. 2008 , 19, 413-9 | 75 |
| 1025 | Contrast safety in the cancer patient: preventing contrast-induced nephropathy. 2008 , 8 Spec No A, S124-7 | 11 |
| 1024 | Introduction to cardiac catheterization part 1: Diagnostic coronary angiography. 2008 , 3, 398-406 | 2 |
| 1023 | The role of extracorporeal blood purification therapies in the prevention of radiocontrast-induced nephropathy. 2008 , 31, 515-24 | 14 |
| 1022 | DNA-free recombinant SV40 capsids protect mice from acute renal failure by inducing stress response, survival pathway and apoptotic arrest. 2008 , 3, e2998 | 5 |
| 1021 | Nefroproteção relacionada ao uso de meio de contraste iodado: atenção de enfermagem. 2008 , 21, 187-191 | 2 |
| 1020 | Incidence of contrast-induced nephropathy in intensive care patients undergoing computerised tomography and prevalence of risk factors. 2009 , 37, 968-75 | 39 |
| 1019 | Provider knowledge of contrast-induced acute kidney injury. 2009 , 338, 280-6 | 6 |

| | | | |
|------|--|------|-----|
| 1018 | Nephrotoxicity of iso-osmolar iodixanol compared with nonionic low-osmolar contrast media: meta-analysis of randomized controlled trials. 2009 , 250, 68-86 | | 225 |
| 1017 | Contrast-induced nephropathy and long-term adverse events: cause and effect?. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2009 , 4, 1162-9 | 6.9 | 195 |
| 1016 | Is carbon dioxide a safe and good alternative for diatrizoate meglumine as a contrast in digital subtraction angiography?. 2009 , 19, 15-9 | | 2 |
| 1015 | Systematic review: sodium bicarbonate treatment regimens for the prevention of contrast-induced nephropathy. 2009 , 151, 631-8 | | 133 |
| 1014 | Safety and performance of targeted renal therapy: the Be-RITe! Registry. 2009 , 16, 1-12 | | 40 |
| 1013 | Determinant factors of renal failure after coronary artery bypass grafting with on-pump technique. 2009 , 18, 300-4 | | 6 |
| 1012 | Iloprost prevents contrast-induced nephropathy in patients with renal dysfunction undergoing coronary angiography or intervention. <i>Circulation</i> , 2009 , 120, 1793-9 | 16.7 | 60 |
| 1011 | Assessment of renal vasoconstriction in vivo after intra-arterial administration of the isosmotic contrast medium iodixanol compared to the low-osmotic contrast medium iopamidol. 2009 , 24, 1478-85 | | 18 |
| 1010 | Probability of reduced renal function after contrast-enhanced CT: a model based on serum creatinine level, patient age, and estimated glomerular filtration rate. 2009 , 193, 494-500 | | 19 |
| 1009 | Background fluctuation of kidney function versus contrast-induced nephrotoxicity. 2009 , 192, 711-8 | | 163 |
| 1008 | Cystatin C for enhancement of risk stratification in non-ST elevation acute coronary syndrome patients with an increased troponin T. 2009 , 55, 1118-25 | | 35 |
| 1007 | Prevention of contrast-induced nephropathy: A randomized controlled trial of sodium bicarbonate and N-acetylcysteine. 2009 , 18, 193-7 | | 12 |
| 1006 | Reversible acute kidney injury following contrast exposure and the risk of long-term mortality. 2009 , 29, 136-44 | | 74 |
| 1005 | [Therapeutic approach in patients undergoing percutaneous coronary interventions]. 2009 , 62, 331-6 | | 3 |
| 1004 | Choice of contrast medium in patients with impaired renal function undergoing percutaneous coronary intervention. 2009 , 2, 430-7 | | 39 |
| 1003 | The cardiorenal syndrome. 2009 , 27, 114-26 | | 57 |
| 1002 | Intraoperative continuous venovenous hemofiltration during coronary surgery. 2009 , 17, 462-6 | | 8 |
| 1001 | Influence of renal function on the effects of early revascularization in non-ST-elevation myocardial infarction: data from the Swedish Web-System for Enhancement and Development of Evidence-Based Care in Heart Disease Evaluated According to Recommended Therapies (SWEDEHEART). <i>Circulation</i> , 2009 , 120, 851-8 | 16.7 | 175 |

| | | | |
|------|---|-----|-----|
| 1000 | Defining contrast-induced nephropathy. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2009 , 4, 1151-3 | 6.9 | 9 |
| 999 | Early angiography in patients with chronic kidney disease: a collaborative systematic review. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2009 , 4, 1032-43 | 6.9 | 73 |
| 998 | Effect of elective coronary angiography on glomerular filtration rate in patients with advanced chronic kidney disease. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2009 , 4, 1907-13 | 6.9 | 18 |
| 997 | Contrast medium volume to estimated glomerular filtration rate ratio as a predictor of contrast-induced nephropathy developing after elective percutaneous coronary intervention. 2009 , 54, 214-20 | | 26 |
| 996 | Sodium bicarbonate-based hydration prevents contrast-induced nephropathy: a meta-analysis. 2009 , 7, 23 | | 99 |
| 995 | Impact of the definition utilized on the rate of contrast-induced nephropathy in percutaneous coronary intervention. 2009 , 103, 1657-62 | | 51 |
| 994 | Efficacy of single-bolus administration of sodium bicarbonate to prevent contrast-induced nephropathy in patients with mild renal insufficiency undergoing an elective coronary procedure. 2009 , 104, 921-5 | | 50 |
| 993 | [Transradial approach and kidney disease]. 2009 , 58, 348-54 | | 1 |
| 992 | Sodium bicarbonate therapy for prevention of contrast-induced nephropathy: a systematic review and meta-analysis. 2009 , 53, 617-27 | | 161 |
| 991 | Forced euvolemic diuresis with mannitol and furosemide for prevention of contrast-induced nephropathy in patients with CKD undergoing coronary angiography: a randomized controlled trial. 2009 , 54, 602-9 | | 84 |
| 990 | Serum creatinine ratio: a novel predictor of mortality after percutaneous coronary intervention in patients with normal and abnormal renal function. 2009 , 74, 49-55 | | 4 |
| 989 | The use of radiographic contrast media during PCI: a focused review: a position statement of the Society of Cardiovascular Angiography and Interventions. 2009 , 74, 728-46 | | 35 |
| 988 | Evidence-based approach for prevention of radiocontrast-induced nephropathy. 2009 , 4, 500-6 | | 10 |
| 987 | Evidence-based practice guideline for the treatment of CKD. 2009 , 13, 537-66 | | 95 |
| 986 | [Contrast induced nephropathy]. 2009 , 121, 15-32 | | 4 |
| 985 | Preventing contrast-induced nephropathy in patients with baseline renal dysfunction undergoing coronary angiography. 2009 , 11, 71-8 | | 1 |
| 984 | Prevention and treatment of contrast-associated nephropathy in interventional cardiology. 2009 , 11, 377-83 | | 16 |
| 983 | High prevalence of stage 3 chronic kidney disease in older adults despite normal serum creatinine. 2009 , 24, 86-92 | | 28 |

| | | | |
|-----|---|-----|----|
| 982 | Factors associated with the use of preventive care for contrast-induced acute kidney injury. 2009 , 24, 289-98 | | 6 |
| 981 | Rationale, design, and baseline characteristics of the Acetylcystein for Contrast-Induced nephropaThy (ACT) Trial: a pragmatic randomized controlled trial to evaluate the efficacy of acetylcysteine for the prevention of contrast-induced nephropathy. 2009 , 10, 38 | | 18 |
| 980 | N-acetylcysteine does not prevent contrast-induced nephropathy after cardiac catheterization in patients with diabetes mellitus and chronic kidney disease: a randomized clinical trial. 2009 , 10, 45 | | 39 |
| 979 | N-acetylcysteine versus N-acetylcysteine + theophylline for the prevention of contrast nephropathy. 2009 , 39, 793-9 | | 25 |
| 978 | N-Acetylcysteine added to volume expansion with sodium bicarbonate does not further prevent contrast-induced nephropathy: results from the cardiac angiography in renally impaired patients study. 2009 , 22, 261-5 | | 13 |
| 977 | Nephrotoxic effects of iodixanol and iopromide in patients with abnormal renal function receiving N-acetylcysteine and hydration before coronary angiography and intervention: a randomized trial. 2009 , 39, 25-31 | | 26 |
| 976 | Ionic low-osmolar versus nonionic iso-osmolar contrast media to obviate worsening nephropathy after angioplasty in chronic renal failure patients: the ICON (Ionic versus non-ionic Contrast to Obviate worsening Nephropathy after angioplasty in chronic renal failure patients) study. <i>JACC: Cardiovascular Interventions</i> , 2009 , 2, 415-21 | 5 | 53 |
| 975 | Sodium bicarbonate plus N-acetylcysteine prophylaxis: a meta-analysis. <i>JACC: Cardiovascular Interventions</i> , 2009 , 2, 1116-24 | 5 | 83 |
| 974 | The critically ill kidney. 2009 , 15, 21-26 | | |
| 973 | Pediatric Nephrology in the ICU. 2009 , | | 4 |
| 972 | The value of hydration and acetylcysteine in the prevention of contrast-induced nephropathy: a potentially catastrophic complication of the percutaneous coronary interventions. <i>International Journal of Cardiology</i> , 2009 , 134, 431-3; author reply 434 | 3.2 | 3 |
| 971 | N-acetylcysteine versus AScorbic acid for preventing contrast-Induced nephropathy in patients with renal insufficiency undergoing coronary angiography NASPI study-a prospective randomized controlled trial. 2009 , 157, 576-83 | | 56 |
| 970 | Nephrotoxicity of iodixanol versus iopamidol in patients with chronic kidney disease and diabetes mellitus undergoing coronary angiographic procedures. 2009 , 158, 822-828.e3 | | 62 |
| 969 | High-dose N-acetylcysteine for the prevention of contrast-induced nephropathy. <i>American Journal of Medicine</i> , 2009 , 122, 874.e9-15 | 2.4 | 71 |
| 968 | D-dimer testing in patients with suspected pulmonary embolism and impaired renal function. <i>American Journal of Medicine</i> , 2009 , 122, 1050-3 | 2.4 | 45 |
| 967 | Whole blood reflectance for assessment of hematologic condition and detection of angiographic contrast media. 2009 , 48, 2435-43 | | 6 |
| 966 | Preventing acute decrease in renal function induced by coronary angiography (PRECORD): a prospective randomized trial. 2009 , 102, 761-7 | | 5 |
| 965 | Renal protective effects and the prevention of contrast-induced nephropathy by atrial natriuretic peptide. 2009 , 53, 1040-6 | | 81 |

| | | |
|-----|---|-----|
| 964 | Contrast-Induced Acute Kidney Injury: Specialty-Specific Protocols for Interventional Radiology, Diagnostic Computed Tomography Radiology, and Interventional Cardiology. 2009 , 84, 170-179 | 105 |
| 963 | Urinary fatty acid-binding protein 1: an early predictive biomarker of kidney injury. 2009 , 296, F669-79 | 103 |
| 962 | Sodium bicarbonate versus normal saline for protection against contrast nephropathy. 2009 , 31, 118-23 | 9 |
| 961 | [Myocardial revascularization in patients with chronic renal failure]. 2009 , 132 Suppl 1, 55-60 | 0 |
| 960 | Contrast-induced nephropathy in patients with type 2 diabetes during coronary angiography: risk-factors and prognostic value. 2009 , 86 Suppl 1, S63-9 | 18 |
| 959 | Use of gadolinium for carotid artery angiography and stenting in patients with renal insufficiency. 2009 , 35, 211-8 | 7 |
| 958 | Comparison of iodixanol and iohexol in patients undergoing intravenous pyelography: a prospective controlled study. 2009 , 31, 181-8 | 19 |
| 957 | Contrast Medium-Induced Nephropathy. 2009 , 63-80 | |
| 956 | Contrast-induced acute kidney injury (CIAKI). 2009 , 47, 783-8, v | 25 |
| 955 | In emergency situations, should serum creatinine be checked in all patients before performing second contrast CT examinations within 24 hours?. 2009 , 6, 268-73 | 42 |
| 954 | Mitigating contrast-induced acute kidney injury associated with cardiac catheterization. 2009 , 17, 263-9 | 16 |
| 953 | Measurement of change in estimated glomerular filtration rate in patients with renal insufficiency after contrast-enhanced computed tomography: a case-control study. 2009 , 33, 455-9 | 22 |
| 952 | The diagnostic and prognostic value of tissue Doppler imaging during dobutamine stress echocardiography in end-stage renal disease. 2009 , 20, 230-7 | 12 |
| 951 | Contrast volume during primary percutaneous coronary intervention and subsequent contrast-induced nephropathy and mortality. 2009 , 150, 170-7 | 234 |
| 950 | Incidence of and risk factors for contrast-induced nephropathy after cardiac catheterization in Japanese patients. 2009 , 73, 1518-22 | 25 |
| 949 | Renal Artery Stenting: Efficacy and Complications. 2009 , 5, 7-12 | 1 |
| 948 | Differentiation of Bosniak categories IIF and III cystic masses: what radiologists should know. 2010 , 34, 847-54 | 9 |
| 947 | Functional contrast-enhanced CT for evaluation of acute ischemic stroke does not increase the risk of contrast-induced nephropathy. 2010 , 31, 817-21 | 69 |

| | | |
|-----|--|-----|
| 946 | Renal ultrasonography in the evaluation of acute kidney injury: developing a risk stratification framework. 2010 , 170, 1900-7 | 38 |
| 945 | Acute kidney injury following transcatheter aortic valve implantation: predictive factors, prognostic value, and comparison with surgical aortic valve replacement. 2010 , 31, 865-74 | 355 |
| 944 | Creatinine increases after intravenous contrast administration: incidence and impact. 2010 , 45, 471-6 | 12 |
| 943 | Preprocedural score for risk of contrast-induced nephropathy in elective coronary angiography and intervention. 2010 , 11, 444-9 | 53 |
| 942 | Strategies for the prevention of contrast-induced acute kidney injury. 2010 , 19, 539-49 | 43 |
| 941 | The contrast media and nephrotoxicity following coronary revascularization by primary angioplasty for acute myocardial infarction study: design and rationale of the CONTRAST-AMI study. 2010 , 11, 199-206 | 3 |
| 940 | Contrast-induced nephropathy in postmenopausal women undergoing percutaneous coronary intervention for acute myocardial infarction. 2010 , 221, 211-9 | 8 |
| 939 | 1. Radiocontrast-induced Acute Kidney Injury.. 2010 , 99, 938-942 | |
| 938 | Contrast-induced acute kidney injury in patients with renal dysfunction undergoing a coronary procedure and receiving non-ionic low-osmolar versus iso-osmolar contrast media. 2010 , 339, 25-30 | 21 |
| 937 | Contrast-induced nephropathy: a contemporary and simplified review. 2010 , 2, 233-242 | 4 |
| 936 | Prognostic value of different laboratory measures of renal function for long-term mortality after contrast media-associated renal impairment. 2010 , 33, E51-9 | 11 |
| 935 | Contrast-induced nephropathy: pathogenesis and prevention. 2010 , 25, 191-204 | 40 |
| 934 | Impact of catheter down-sizing and power injector use on the amount of contrast medium delivered. 2010 , 25, 24-8 | 5 |
| 933 | Predictive value of serum cystatin C, β_2 -microglobulin, and urinary liver-type fatty acid-binding protein on the development of contrast-induced nephropathy. 2010 , 25, 85-90 | 20 |
| 932 | Incidence and outcomes of contrast-induced nephropathy after computed tomography in patients with CKD: a quality improvement report. 2010 , 55, 1018-25 | 75 |
| 931 | Efficacy of N-acetylcysteine and aminophylline in preventing contrast-induced nephropathy. 2010 , 55, 174-9 | 19 |
| 930 | Effect of contrast-induced nephropathy on cardiac outcomes after use of nonionic isosmolar contrast media during coronary procedure. 2010 , 56, 300-6 | 18 |
| 929 | The effect of cardiac angiography timing, contrast media dose, and preoperative renal function on acute renal failure after coronary artery bypass grafting. 2010 , 139, 1539-44 | 69 |

| | | |
|-----|--|--------|
| 928 | Novel approach for the prevention of contrast nephropathy. 2010 , 62, 81-9 | 26 |
| 927 | Usefulness of atorvastatin (80 mg) in prevention of contrast-induced nephropathy in patients with chronic renal disease. 2010 , 105, 288-92 | 124 |
| 926 | Contrast-induced nephropathy in patients undergoing emergency percutaneous coronary intervention for acute coronary syndrome. 2010 , 105, 624-8 | 101 |
| 925 | Endovascular detection and removal of radiographic contrast. 2010 , 11, 114-5 | 1 |
| 924 | Background incidence of serum creatinine threshold rises in a predominantly female clinical trial population without underlying renal disease. 2010 , 58, 297-300 | |
| 923 | Evidence-based management of patients undergoing PCI: contrast-induced acute kidney injury. 2010 , 75 Suppl 1, S15-20 | 33 |
| 922 | ACCF/ACR/AHA/NASCI/SAIP/SCAI/SCCT 2010 expert consensus document on coronary computed tomographic angiography: a report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents. 2010 , 76, E1-42 | 37 |
| 921 | The prevention of contrast induced nephropathy by sarpogrelate in patients with chronic kidney disease: a study protocol for a prospective randomized controlled clinical trial. 2010 , 11, 122 | 6 |
| 920 | Safety of low volume iodinated contrast administration for arteriovenous fistula intervention in chronic kidney disease stage 4 or 5 utilizing a bicarbonate prophylaxis strategy. 2010 , 23, 638-42 | 7 |
| 919 | Low Incidence of Contrast Induced Nephropathy after Coronary Angiography in Patients with Advanced Chronic Kidney Disease Treated with Prophylactic Continuous Veno-venous Hemofiltration. 2010 , 2, | |
| 918 | Small dose contrast venography as venous mapping in predialysis patients. 2010 , 11, 122-7 | 12 |
| 917 | Is there an association between non-steroidal anti-inflammatory drugs and contrast nephropathy?. 2010 , 95, 726-31 | 7 |
| 916 | Pediatric Interventional Cardiovascular Magnetic Resonance. 2010 , 593-609 | 1 |
| 915 | Intravenous contrast medium administration and scan timing at CT: considerations and approaches. 2010 , 256, 32-61 | 633 |
| 914 | Iodixanol versus low-osmolar contrast media for prevention of contrast induced nephropathy: meta-analysis of randomized, controlled trials. 2010 , 3, 351-8 | 71 |
| 913 | The risk of acute radiocontrast-mediated kidney injury following endovascular therapy for acute ischemic stroke is low. 2010 , 31, 1584-7 | 16 |
| 912 | Chronic kidney disease: an inherent risk factor for acute kidney injury?. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2010 , 5, 1690-5 | 6.9 68 |
| 911 | Safety of performing CT angiography in stroke patients treated with intravenous thrombolysis. 2010 , 81, 783-7 | 33 |

| | | | |
|-----|--|------|-----|
| 910 | Contrast-induced nephropathy in patients undergoing percutaneous coronary intervention. 2010 , 2010, | | 12 |
| 909 | Mehran contrast-induced nephropathy risk score predicts short- and long-term clinical outcomes in patients with ST-elevation-myocardial infarction. 2010 , 3, 491-8 | | 86 |
| 908 | Curbing the use of ultrasonography in the diagnosis of acute kidney injury: Penny wise or pound foolish?: Comment on "Renal ultrasonography in the evaluation of acute kidney injury". 2010 , 170, 1907-8 | | 6 |
| 907 | Impact of female gender on frequency of contrast medium-induced nephropathy: post hoc analysis of dialysis versus diuresis trial. 2010 , 19, 1363-8 | | 11 |
| 906 | Prevention of cardio-renal syndromes: workgroup statements from the 7th ADQI Consensus Conference. 2010 , 25, 1777-84 | | 29 |
| 905 | Epidemiology of cardio-renal syndromes: workgroup statements from the 7th ADQI Consensus Conference. 2010 , 25, 1406-16 | | 152 |
| 904 | Acute kidney injury: kidney injury after contrast media: marker or mediator?. 2010 , 6, 634-6 | | 6 |
| 903 | Chronic kidney injury in patients after cardiac catheterisation or percutaneous coronary intervention: a comparison of radial and femoral approaches (from the British Columbia Cardiac and Renal Registries). 2010 , 96, 1538-42 | | 61 |
| 902 | Contrast-induced nephropathy: current practices among cardiologists. 2010 , 32, 928-34 | | 1 |
| 901 | Contrast-induced nephropathy in patients undergoing computed tomography (CONNECT) - a clinical problem in daily practice? A multicenter observational study. 2010 , 51, 741-50 | | 32 |
| 900 | ACCF/ACR/AHA/NASCI/SAIP/SCAI/SCCT 2010 expert consensus document on coronary computed tomographic angiography: a report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents. <i>Circulation</i> , 2010 , 121, 2509-43 | 16.7 | 139 |
| 899 | Contrast medium nephrotoxicity after renal artery and coronary angioplasty. 2010 , 51, 462-6 | | 3 |
| 898 | Impact of renin-angiotensin-aldosterone blockade by angiotensin-converting enzyme inhibitors or AT-1 blockers on frequency of contrast medium-induced nephropathy: a post-hoc analysis from the Dialysis-versus-Diuresis (DVD) trial. 2010 , 25, 759-64 | | 49 |
| 897 | Acute kidney injury following coronary angiography is associated with a long-term decline in kidney function. 2010 , 78, 803-9 | | 157 |
| 896 | Prevention of Contrast-Induced Nephropathy (CIN) in Interventional Radiology Practice. 2010 , 27, 348-59 | | 37 |
| 895 | Efficacy of short-term high-dose atorvastatin for prevention of contrast-induced nephropathy in patients undergoing coronary angiography. <i>Angiology</i> , 2010 , 61, 711-4 | 2.1 | 80 |
| 894 | Contrast Media-Induced Nephropathy and Nephrogenic Systemic Fibrosis. 2010 , 831-844 | | 1 |
| 893 | Pre-procedural glucose levels and the risk for contrast-induced acute kidney injury in patients undergoing coronary angiography. 2010 , 55, 1433-40 | | 71 |

| | | | | |
|-----|--|-----|--|-----|
| 892 | ACCF/ACR/AHA/NASCI/SAIP/SCAI/SCCT 2010 expert consensus document on coronary computed tomographic angiography: a report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents. 2010 , 55, 2663-99 | | | 219 |
| 891 | Removal of contrast media from the coronary sinus attenuates renal injury after coronary angiography and intervention. 2010 , 56, 525-6 | | | 19 |
| 890 | Distinct patterns of autoantibodies against G-protein-coupled receptors in Chagas' cardiomyopathy and megacolon: their potential impact for early risk assessment in asymptomatic Chagas' patients. 2010 , 56, 526-7 | | | 9 |
| 889 | Contrast-induced nephropathy after a second contrast exposure. 2010 , 32, 796-801 | | | 39 |
| 888 | N-acetylcysteine: short-term clinical benefits after coronary angiography in high-risk renal patients. 2010 , 63, 12-9 | | | 8 |
| 887 | Sodium bicarbonate for prevention of contrast-induced acute kidney injury: a systematic review and meta-analysis. 2010 , 25, 747-58 | | | 86 |
| 886 | Il paziente a rischio di nefropatia da mezzi di contrasto iodati. 2010 , 4, 223-229 | | | |
| 885 | Effect of N-acetylcysteine on cystatin C-based renal function after elective coronary angiography (ENABLE Study): a prospective, randomized trial. <i>International Journal of Cardiology</i> , 2010 , 138, 239-45 | 3.2 | | 25 |
| 884 | Frequency and predictors of contrast-induced nephropathy after angioplasty for chronic total occlusions. <i>International Journal of Cardiology</i> , 2010 , 139, 68-74 | 3.2 | | 58 |
| 883 | [Contrast-induced nephropathy]. 2010 , 39, 807-14 | | | 1 |
| 882 | [Contrast-induced nephropathy in patients at risk of renal failure undergoing computed tomography: systematic review and meta-analysis of randomized controlled trials]. 2010 , 135, 343-50 | | | 7 |
| 881 | Can urine dipstick predict an elevated serum creatinine?. 2010 , 28, 613-6 | | | 2 |
| 880 | Coronary revascularisation in chronic kidney disease. Part 1: stable coronary artery disease. 2010 , 36 Suppl 1, 106-17 | | | 5 |
| 879 | Oral hydration and alkalization is noninferior to intravenous therapy for prevention of contrast-induced nephropathy in patients with chronic kidney disease. 2010 , 23, 460-6 | | | 27 |
| 878 | N-acetilcisteína: beneficio clínico a corto plazo tras coronariografía en pacientes renales de alto riesgo. 2010 , 63, 12-19 | | | 21 |
| 877 | Radiation dosimetry and safety issues in the investigation of pulmonary embolism. 2010 , 40, 442-54 | | | 77 |
| 876 | Cardio-renal syndromes: report from the consensus conference of the acute dialysis quality initiative. 2010 , 31, 703-11 | | | 581 |
| 875 | Contrast nephropathy: can we see the whole picture. 2010 , 16, 35-41 | | | |

| | | | |
|-----|--|------|------|
| 874 | Development of inpatient risk stratification models of acute kidney injury for use in electronic health records. 2010 , 30, 639-50 | | 65 |
| 873 | Prevention of contrast-induced nephropathy with haemofiltration in high-risk patients after percutaneous coronary intervention. 2011 , 13, 164-9 | | |
| 872 | Contrast-induced acute kidney injury. 2011 , 17, 193-195 | | 2 |
| 871 | Acetylcysteine for prevention of renal outcomes in patients undergoing coronary and peripheral vascular angiography: main results from the randomized Acetylcysteine for Contrast-induced nephropathy Trial (ACT). <i>Circulation</i> , 2011 , 124, 1250-9 | 16.7 | 263 |
| 870 | WITHDRAWN: Acute kidney injury following cardiac surgery, diagnostic value of plasma neutrophil gelatinase-associated lipocalin. 2011 , | | |
| 869 | The Canadian Association of Radiologists guidelines for the prevention of contrast-induced nephropathy: a critical appraisal. 2011 , 62, 238-42 | | 6 |
| 868 | L'utilisation du Gadolinium pour les interventions artérielles. 2011 , 25, 393-403 | | |
| 867 | Contrast-induced acute kidney injury: short- and long-term implications. 2011 , 31, 300-9 | | 48 |
| 866 | Standardized endpoint definitions for Transcatheter Aortic Valve Implantation clinical trials: a consensus report from the Valve Academic Research Consortium. 2011 , 57, 253-69 | | 662 |
| 865 | 2011 ACCF/AHA/SCAI Guideline for Percutaneous Coronary Intervention: Executive Summary: A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. 2011 , 58, 2550-2583 | | 99 |
| 864 | 2011 ACCF/AHA/SCAI Guideline for Percutaneous Coronary Intervention. A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. 2011 , 58, e44-122 | | 1703 |
| 863 | Short-term, high-dose statins in the prevention of contrast-induced nephropathy: a systematic review and meta-analysis. 2011 , 76, 475-83 | | 19 |
| 862 | N-Acetylcysteine in Kidney Disease. 2011 , 367-388 | | 1 |
| 861 | Current use of CT in the evaluation and management of injured patients. 2011 , 91, 233-48 | | 20 |
| 860 | Statins for the prevention of contrast-induced nephropathy: a systematic review and meta-analysis. 2011 , 33, 344-51 | | 70 |
| 859 | The use of gadolinium for arterial interventions. 2011 , 25, 366-76 | | 8 |
| 858 | Incidence and predictors of acute kidney injury in patients undergoing transcatheter aortic valve implantation. 2011 , 161, 735-9 | | 121 |
| 857 | Impact of contrast-induced acute kidney injury definition on clinical outcomes. 2011 , 161, 963-71 | | 51 |

| | | | |
|-----|--|-----|-----|
| 856 | Contrast-induced nephropathy risk assessment in real world practice. <i>American Journal of Medicine</i> , 2011 , 124, 1127-8 | 2.4 | 5 |
| 855 | Potentials and limitations of low-concentration contrast medium (150 mg iodine/ml) in CT pulmonary angiography. 2011 , 66, 43-9 | | 5 |
| 854 | Importance of oral fluid intake after coronary computed tomography angiography: an observational study. 2011 , 77, 118-22 | | 22 |
| 853 | Sodium bicarbonate for the prevention of contrast induced nephropathy: a meta-analysis of published clinical trials. 2011 , 79, 48-55 | | 36 |
| 852 | Predictive value of myocardial and coronary imaging in the long-term outcome of potential renal transplant recipients. <i>International Journal of Cardiology</i> , 2011 , 146, 191-6 | 3.2 | 16 |
| 851 | [Contrast-induced nephropathy]. 2011 , 137, 84-90 | | 1 |
| 850 | High incidence of nephropathy in neurosurgical patients after intra-arterial administration of low-osmolar and iso-osmolar contrast media. 2011 , 52, 422-9 | | 8 |
| 849 | Prevention of Radiocontrast-Induced Nephropathy after Coronary Angiography: N-Acetylcysteine plus Saline Hydration versus Saline Hydration. 2011 , 10, | | |
| 848 | Defining Incidence and Outcome of Contrast-Induced Nephropathy Among Trauma: Is It Overhyped?. 2011 , 77, 686-689 | | 11 |
| 847 | Proteinuria as a Predictor of Renal Dysfunction in Trauma Patients Receiving Intravenous Contrast. 2011 , 77, 1194-1200 | | 6 |
| 846 | Contrast nephropathy in patients undergoing percutaneous coronary intervention. 2011 , 2, 165-172 | | 1 |
| 845 | Imaging patients with kidney disease: how do we approach contrast-related toxicity?. 2011 , 341, 215-21 | | 18 |
| 844 | Risks and complications of coronary angiography: a comprehensive review. 2012 , 4, 65-93 | | 118 |
| 843 | Determination of safe contrast media dosage to estimated glomerular filtration rate ratios to avoid contrast-induced nephropathy after elective percutaneous coronary intervention. 2011 , 41, 265-71 | | 11 |
| 842 | Cystatin C: a possible sensitive marker for detecting potential kidney injury after computed tomography coronary angiography. 2011 , 35, 240-5 | | 10 |
| 841 | Contrast-enhanced magnetic resonance angiography for preoperative imaging of deep inferior epigastric artery perforator flaps: advantages and disadvantages compared with computed tomography angiography: a United Kingdom perspective. 2011 , 67, 671-4 | | 23 |
| 840 | Current approaches to prevention of contrast induced acute kidney injury. 2011 , 9, 247-52 | | 1 |
| 839 | Preprocedural evaluation and endovascular treatment of iliofemoral artery disease without contrast media for patients with pre-existing renal insufficiency. 2011 , 75, 179-84 | | 9 |

| | | |
|-----|--|-----|
| 838 | Prediction of chronic renal insufficiency after coronary angiography by an early increase in oxidative stress and decrease in glomerular filtration rate. 2011 , 75, 437-42 | 8 |
| 837 | The impact of hyperuricemia on in-hospital mortality and incidence of acute kidney injury in patients undergoing percutaneous coronary intervention. 2011 , 75, 692-7 | 42 |
| 836 | Does N-acetylcysteine reduce the incidence of contrast-induced nephropathy and clinical events in patients undergoing primary angioplasty for acute myocardial infarction?. 2011 , 50, 673-7 | 21 |
| 835 | [Noninvasive assessment of coronary artery with substantial contrast volume reduction using 320-detector row computed tomography: a feasibility study]. 2011 , 67, 145-51 | |
| 834 | Elective percutaneous coronary intervention in the elderly patient. 2011 , 7, 271-281 | 3 |
| 833 | The effect of coronary angiography on renal function in preemptive renal transplant candidates. 2011 , 25, 594-9 | 4 |
| 832 | An invasive management strategy is associated with improved outcomes in high-risk acute coronary syndromes in patients with chronic kidney disease. 2011 , 41, 743-50 | 7 |
| 831 | Efficacy of statin pretreatment for the prevention of contrast-induced nephropathy: a meta-analysis of randomised controlled trials. 2011 , 65, 624-30 | 20 |
| 830 | Endovascular procedures with CTA and MRA roadmapping. 2011 , 21, 259-62 | 11 |
| 829 | Visualizing the future: enhancing neuroimaging with nanotechnology. 2011 , 75, 626-37; discussion 618-9 | 30 |
| 828 | Preventive strategies of renal insufficiency in patients with diabetes undergoing intervention or arteriography (the PREVENT Trial). 2011 , 107, 1447-52 | 35 |
| 827 | Management of non-ST-segment elevation acute myocardial infarction in patients with chronic kidney disease (from the Korea Acute Myocardial Infarction Registry). 2011 , 108, 206-13 | 21 |
| 826 | Systemic hypothermia to prevent radiocontrast nephropathy (from the COOL-RCN Randomized Trial). 2011 , 108, 741-6 | 7 |
| 825 | Angiotensin-converting enzyme inhibitors and their effects on contrast-induced nephropathy after cardiac catheterization or percutaneous coronary intervention. 2011 , 12, 90-3 | 10 |
| 824 | 2011 ACCF/AHA/SCAI guideline for percutaneous coronary intervention: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. 2013 , 82, E266-355 | 81 |
| 823 | Effect on renal function of an iso-osmolar contrast agent in patients with monoclonal gammopathies. 2011 , 21, 63-9 | 10 |
| 822 | Contrast induced nephropathy: updated ESUR Contrast Media Safety Committee guidelines. 2011 , 21, 2527-41 | 618 |
| 821 | Epidemiology and outcome of the cardio-renal syndrome. 2011 , 16, 531-42 | 34 |

| | | |
|-----|--|-----|
| 820 | Evaluation of brain and kidney energy metabolism in an animal model of contrast-induced nephropathy. 2011 , 26, 115-22 | 6 |
| 819 | Intravenous glutathione prevents renal oxidative stress after coronary angiography more effectively than oral N-acetylcysteine. 2011 , 26, 465-72 | 18 |
| 818 | La nefropatia da mezzi di contrasto: i fattori di rischio e l'ènomalia italiana. 2011 , 7, 36-41 | |
| 817 | Prèvention der Kontrastmittelneuropathie. 2011 , 6, 128-134 | 1 |
| 816 | Are we still wondering or wandering in the dark?. 2011 , 6, 485-6 | |
| 815 | Renal function at the time of a myocardial infarction maintains prognostic value for more than 10 years. 2011 , 11, 37 | 7 |
| 814 | Early detection and intervention using neutrophil gelatinase-associated lipocalin (NGAL) may improve renal outcome of acute contrast media induced nephropathy: a randomized controlled trial in patients undergoing intra-arterial angiography (ANTI-CIN Study). 2011 , 12, 39 | 19 |
| 813 | Safety and efficacy of dual-axis rotational coronary angiography vs. standard coronary angiography. 2011 , 77, 820-7 | 32 |
| 812 | Use of performance curves in estimating number of procedures required to achieve proficiency in coronary angiography. 2011 , 78, 387-93 | 7 |
| 811 | The ratio of contrast volume to glomerular filtration rate predicts outcomes after percutaneous coronary intervention for ST-segment elevation acute myocardial infarction. 2011 , 78, 198-201 | 46 |
| 810 | Frequency, determinants, and prognostic effects of acute kidney injury and red blood cell transfusion in patients undergoing transcatheter aortic valve implantation. 2011 , 77, 881-9 | 102 |
| 809 | The synthesis and evaluation of unsymmetrical dimeric X-ray contrast agents. 2011 , 52, 3068-3071 | 6 |
| 808 | The synthesis and evaluation of trimeric X-ray contrast agents. 2011 , 52, 3065-3067 | 3 |
| 807 | Standardized endpoint definitions for transcatheter aortic valve implantation clinical trials: a consensus report from the Valve Academic Research Consortium. 2011 , 32, 205-17 | 510 |
| 806 | Oral prostaglandin e1 in combination with sodium bicarbonate and normal saline in the prevention of contrast-induced nephropathy: a pilot study. 2011 , 20, 229-34 | 6 |
| 805 | Reducing the dose of contrast medium in angiography by use of a highly sensitive receiver and synchrotron radiation system. 2011 , 197, W508-13 | 4 |
| 804 | Bedeutung der Niereninsuffizienz in der Behandlung der Herzinsuffizienz - kardiorenales und renokardiales Syndrom. 2011 , 7, 25-37 | |
| 803 | Feasibility of applying low contrast media volume in cervicocranial artery angiography with 256-slice CT. 2011 , | |

| | | | |
|-----|---|------|------|
| 802 | Prolonged carotid sinus reflex is a risk factor for contrast-induced nephropathy following carotid artery stenting. 2011 , 32, 441-5 | | 11 |
| 801 | Computed Tomography Angiography in the Assessment of Patients With Stroke/TIA. 2011 , 1, 187-99 | | 41 |
| 800 | Baseline incidence and severity of renal insufficiency evaluated by estimated glomerular filtration rates in patients scheduled for contrast-enhanced CT. 2011 , 52, 581-6 | | 12 |
| 799 | Comparison of the effect of low- and iso-osmolar contrast agents on heart rate during chest CT angiography: results of a prospective randomized multicenter study. 2011 , 258, 930-7 | | 11 |
| 798 | Associations between acute kidney injury and cardiovascular and renal outcomes after coronary angiography. <i>Circulation</i> , 2011 , 123, 409-16 | 16.7 | 250 |
| 797 | 2011 ACCF/AHA/SCAI Guideline for Percutaneous Coronary Intervention: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. <i>Circulation</i> , 2011 , 124, 2574-603 | 16.7 | 389 |
| 796 | Cardiac resynchronization therapy in the cardiorenal syndrome. 2011 , 2011, 168461 | | 3 |
| 795 | Micronutrients and cardiorenal disease: insights into novel assessments and treatment. 2011 , 31, 177-85 | | 5 |
| 794 | Cardiorenal syndromes. 2011 , 3, 1-9 | | 21 |
| 793 | Abdominal dynamic CT in patients with renal dysfunction: contrast agent dose reduction with low tube voltage and high tube current-time product settings at 256-detector row CT. 2011 , 261, 467-76 | | 89 |
| 792 | Acute kidney injury in older adults. 2011 , 22, 28-38 | | 136 |
| 791 | Studies on Renal Disorders. 2011 , | | 3 |
| 790 | Risk of acute kidney injury after minimally invasive transapical aortic valve implantation in 270 patients. 2011 , 39, 835-42; discussion 842-3 | | 87 |
| 789 | Reference renal artery diameter is a stronger predictor of contrast-induced nephropathy than chronic kidney disease in patients with high cardiovascular risk. 2011 , 1, 38-44 | | 3 |
| 788 | Timing of coronary angiography and outcome in patients with non-ST elevation acute coronary syndromes and kidney disease: real-world data from the acute coronary syndromes Israeli survey. 2011 , 119, 224-34 | | 5 |
| 787 | Contrast-induced nephropathy in primary percutaneous coronary intervention. 2011 , 97, 1723-5 | | 10 |
| 786 | Impact of contrast-induced acute kidney injury with transient or persistent renal dysfunction on long-term outcomes of patients with acute myocardial infarction undergoing percutaneous coronary intervention. 2011 , 97, 1753-7 | | 124 |
| 785 | 2011 ACCF/AHA/SCAI Guideline for Percutaneous Coronary Intervention: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. <i>Circulation</i> , 2011 , 124, e574-651 | 16.7 | 1039 |

| | | |
|-----|---|-----|
| 784 | The role of nebivolol in the prevention of contrast-induced nephropathy in patients with renal dysfunction. 2011 , 11, 613-7 | 16 |
| 783 | Cardiorenal syndromes: pathophysiology to prevention. 2010 , 2011, 762590 | 21 |
| 782 | Heart-kidney interaction: epidemiology of cardiorenal syndromes. 2010 , 2011, 351291 | 38 |
| 781 | Design of clinical trials in AKI: a report from an NIDDK workshop. Trials of patients with sepsis and in selected hospital settings. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012 , 7, 856-60 ⁶⁹ | 33 |
| 780 | An x-space magnetic particle imaging scanner. 2012 , 83, 033708 | 88 |
| 779 | Antioxidant and renoprotective effects of paricalcitol on experimental contrast-induced nephropathy model. 2012 , 85, 1038-43 | 30 |
| 778 | The effects of chronic usage of angiotensin-converting enzyme inhibitors and angiotensin receptor blockers on contrast-induced nephropathy in low-risk patients [corrected]. 2013 , 13, 245-50 | 11 |
| 777 | Efficacy of short-term high-dose statin in preventing contrast-induced nephropathy: a meta-analysis of seven randomized controlled trials. 2012 , 7, e34450 | 57 |
| 776 | Contrast-induced nephropathy risk due to emergency contrast-enhanced computed tomography. 2012 , 19, 20-3 | 2 |
| 775 | ACE inhibitor or angiotensin II receptor blocker use is a risk factor for contrast-induced nephropathy. 2012 , 25, 776-81 | 18 |
| 774 | Contrast-induced nephropathy in critical care. 2012 , 32, 15-23; quiz 1p following page 23 | 5 |
| 773 | Prevention of contrast-induced AKI--the ongoing saga. 2012 , 8, 260-1 | 2 |
| 772 | Evaluation of potential renal transplant recipients with computed tomography angiography. 2012 , 147, 1114-22 | 8 |
| 771 | References. 2012 , 2, 124-138 | 33 |
| 770 | Epidemiology of contrast material-induced nephropathy in the era of hydration. 2012 , 263, 706-13 | 71 |
| 769 | Contrast-induced acute kidney injury: the continuous quest for pharmacological prevention. 2012 , 5, 741-3 | 2 |
| 768 | Low contrast agent and radiation dose protocol for hepatic dynamic CT of thin adults at 256-detector row CT: effect of low tube voltage and hybrid iterative reconstruction algorithm on image quality. 2012 , 264, 445-54 | 112 |
| 767 | Chronic kidney disease is not associated with a higher risk for mortality or acute kidney injury in transcatheter aortic valve implantation. 2012 , 27, 3502-8 | 34 |

| | | |
|-----|---|----------|
| 766 | Intrarenal application of N-acetylcysteine for the prevention of contrast medium-induced nephropathy in primary angioplasty. 2012 , 23, 265-70 | 24 |
| 765 | Preventing radiocontrast-induced nephropathy in chronic kidney disease patients undergoing coronary angiography. 2012 , 4, 157-72 | 14 |
| 764 | Risk assessment and prevention of contrast-induced nephropathy in patients undergoing coronary angiography. 2012 , 51, 519-21 | 2 |
| 763 | N-acetylcysteine for the prevention of contrast-induced nephropathy in the emergency department. 2012 , 51, 2709-14 | 19 |
| 762 | Pre-procedural glucose levels and the risk for contrast-induced acute kidney injury in patients undergoing emergency coronary intervention. 2012 , 76, 1848-55 | 16 |
| 761 | Sodium bicarbonate therapy for the prevention of contrast-induced acute kidney injury [a systematic review and meta-analysis] 2012 , 76, 2255-65 | 73 |
| 760 | Comparison of Dual-Axis Rotational Coronary Angiography (XPERSWING) Versus Conventional Technique in Routine Practice. 2012 , 65, 434-439 | 1 |
| 759 | Contrast-induced nephropathy: attributable incidence and potential harm. 2012 , 16, 127 | 6 |
| 758 | Section 4: Contrast-induced AKI. 2012 , 2, 69-88 | 46 |
| 757 | Contrast-induced nephropathy in patients with renal insufficiency undergoing contrast-enhanced MDCT. 2012 , 22, 2147-52 | 31 |
| 756 | Imaging of the carotid artery. 2012 , 220, 294-309 | 49 |
| 755 | Contrast-induced nephropathy and risk of acute kidney injury and mortality after cardiac operations. 2012 , 94, 772-6 | 34 |
| 754 | Renal impairment and transapical aortic valve implantation: impact of contrast medium dose on kidney function and survival. 2012 , 41, 1225-32 | 22 |
| 753 | Renal function in patients treated with abdominal aortic stentgraft implantation with an intentional occlusion of accessory renal artery. 2012 , 26, 299-305 | 13 |
| 752 | The effect of renin-angiotensin-aldosterone system blockade on contrast-induced acute kidney injury: a propensity-matched study. 2012 , 60, 576-82 | 51 |
| 751 | Contrast-induced kidney injury: mechanisms, risk factors, and prevention. 2012 , 33, 2007-15 | 303 |
| 750 | How do centres begin the process to prevent contrast-induced acute kidney injury: a report from a new regional collaborative. 2012 , 21, 54-62 | 25 |
| 749 | Ischemic preconditioning for prevention of contrast medium-induced nephropathy: randomized pilot RenPro Trial (Renal Protection Trial). <i>Circulation</i> , 2012 , 126, 296-303 | 16.7 198 |

| | | | |
|-----|---|-----|-----|
| 748 | Comparison of dual-axis rotational coronary angiography (XPERSWING) versus conventional technique in routine practice. 2012 , 65, 434-9 | | 15 |
| 747 | Use and overuse of left ventriculography. 2012 , 163, 617-23.e1 | | 14 |
| 746 | Incidence and predictors of acute kidney injury after transcatheter aortic valve replacement. 2012 , 163, 1031-6 | | 112 |
| 745 | Safety of iodinated intravenous contrast medium administration in sickle cell disease. <i>American Journal of Medicine</i> , 2012 , 125, 100.e11-6 | 2.4 | 12 |
| 744 | N-acetylcysteine for the prevention of contrast-induced nephropathy in rats. 2012 , 36, 210-215 | | |
| 743 | Risk factors for contrast induced nephropathy: a study among Italian patients. 2012 , 64, 484-91 | | 18 |
| 742 | Prevention of contrast nephropathy by furosemide with matched hydration: the MYTHOS (Induced Diuresis With Matched Hydration Compared to Standard Hydration for Contrast Induced Nephropathy Prevention) trial. <i>JACC: Cardiovascular Interventions</i> , 2012 , 5, 90-7 | 5 | 156 |
| 741 | [Glycaemic management in type 1 and 2 diabetes patients undergoing interventional cardiology procedures. Heart and Diabetes Working Group. Sociedad Española de Cardiología. Sociedad Española de Diabetes]. 2012 , 138, 207.e1-5 | | 0 |
| 740 | The economics of renal failure and kidney disease in critically ill patients. 2012 , 28, 99-111, vii | | 26 |
| 739 | Cardiological Society of India practice guidelines for angiography in patients with renal dysfunction. 2012 , 64 Suppl 2, S18-43 | | 2 |
| 738 | Cardio-renal syndrome type 1: epidemiology, pathophysiology, and treatment. 2012 , 32, 18-25 | | 31 |
| 737 | Determination of lower limb microvasculature by intrafemoral arterial injection using computed tomography-assisted angiography. 2012 , 36, 1376-81 | | 2 |
| 736 | Imaging in Clinical Trials. 2012 , 597-617 | | |
| 735 | Desmoid Tumors. 2012 , | | 3 |
| 734 | Acute kidney injury biomarkers for patients in a coronary care unit: a prospective cohort study. 2012 , 7, e32328 | | 37 |
| 733 | Association of serum bilirubin with contrast-induced nephropathy and future cardiovascular events in patients undergoing coronary intervention. 2012 , 7, e42594 | | 23 |
| 732 | Comparison of oral and intravenous hydration strategies for the prevention of contrast-induced nephropathy in patients undergoing coronary angiography or angioplasty: a randomized clinical trial. <i>Acta Cardiologica</i> , 2012 , 67, 565-9 | 0.9 | 31 |
| 731 | Contrast Nephropathy: A Paradigm for Cardiorenal Interactions in Clinical Practice. 2012 , | | |

| | | |
|-----|---|-----|
| 730 | n-Acetylcysteine for prevention of iodinated contrast-induced nephropathy in computed tomography angiography procedures in patients with chronic kidney disease. 2012 , 6, 181-189 | |
| 729 | Hydration with sodium bicarbonate does not prevent contrast nephropathy: a multicenter clinical trial. 2012 , 99, 1129-34 | 19 |
| 728 | Prevention of contrast-induced acute kidney injury in patients with stable chronic renal disease undergoing elective percutaneous coronary and peripheral interventions: randomized comparison of two preventive strategies. 2012 , 79, 929-37 | 28 |
| 727 | Double-vessel coronary stenting via 5 French diagnostic catheters. 2012 , 80, 630-3 | 5 |
| 726 | Novel use of the GuideLiner catheter to minimize contrast use during PCI in a patient with chronic kidney disease. 2012 , 80, 453-5 | 18 |
| 725 | Staging of multivessel percutaneous coronary interventions: an expert consensus statement from the Society for Cardiovascular Angiography and Interventions. 2012 , 79, 1138-52 | 19 |
| 724 | Impact of severity of renal dysfunction on determinants of in-hospital mortality among patients undergoing percutaneous coronary intervention. 2012 , 80, 352-7 | 43 |
| 723 | 2011 ACCF/AHA/SCAI Guideline for Percutaneous Coronary Intervention: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. 2012 , 79, 453-95 | 121 |
| 722 | Does nebivolol prevent contrast-induced nephropathy in humans?. 2012 , 35, 250-4 | 28 |
| 721 | Age-related differences in treatment strategies and clinical outcomes in unselected cohort of patients with ST-segment elevation myocardial infarction transferred for primary angioplasty. 2012 , 34, 214-21 | 23 |
| 720 | [Use of N-acetylcysteine for the prevention of contrast-induced nephropathy in rats]. 2012 , 36, 210-5 | 3 |
| 719 | Effect of an electronic alert on risk of contrast-induced acute kidney injury in hospitalized patients undergoing computed tomography. 2012 , 60, 74-81 | 35 |
| 718 | Effectiveness of theophylline in preventing contrast-induced nephropathy after coronary angiographic procedures. 2012 , 25, 404-10 | 21 |
| 717 | Incidence, morbidity, and mortality of contrast-induced acute kidney injury in a surgical intensive care unit: a prospective cohort study. 2012 , 27, 322.e1-5 | 32 |
| 716 | Urinary liver-type fatty acid-binding protein level as a predictive biomarker of contrast-induced acute kidney injury. 2012 , 42, 557-63 | 48 |
| 715 | Contrast-induced nephropathy: protective role of fenoldopam. 2012 , 39, 497-505 | 5 |
| 714 | Prevention and management of contrast-induced acute kidney injury. 2012 , 14, 1-7 | 4 |
| 713 | Guidelines on the use of iodinated contrast media in patients with kidney disease 2012: digest version. JSN, JRS, and JCS Joint Working Group. 2013 , 31, 546-84 | 9 |

| | | | | |
|-----|---|-----|--|----|
| 712 | Development of contrast-induced nephropathy in subarachnoid hemorrhage: a single center perspective. 2013 , 19, 150-6 | | | 13 |
| 711 | N-acetyl-cysteine reduces the occurrence of contrast-induced acute kidney injury in patients with renal dysfunction: a single-center randomized controlled trial. 2013 , 17, 396-404 | | | 8 |
| 710 | Guidelines on the use of iodinated contrast media in patients with kidney disease 2012: digest version : JSN, JRS, and JCS Joint Working Group. 2013 , 17, 441-79 | | | 19 |
| 709 | Invasive Coronary Angiography. 2013 , 77-89 | | | 0 |
| 708 | A correlation between acute kidney injury and myonecrosis after scheduled percutaneous coronary intervention. 2013 , 14, 713-20 | | | 2 |
| 707 | 20-Hour preprocedural hydration is not superior to 5-hour preprocedural hydration in the prevention of contrast-induced increases in serum creatinine and cystatin C. <i>International Journal of Cardiology</i> , 2013 , 167, 2200-3 | 3.2 | | 10 |
| 706 | Prevention of contrast induced nephropathy during coronary angiography with a coronary sinus contrast removal system sited from the femoral vein. <i>International Journal of Cardiology</i> , 2013 , 165, e9-10 ² | | | 5 |
| 705 | Ascorbic Acid for the prevention of contrast-induced nephropathy after coronary angiography in patients with chronic renal impairment: a randomized controlled trial. 2013 , 17, 384-90 | | | 16 |
| 704 | Contrast-induced acute kidney injury following coronary angiography: a cohort study of hospitalized patients with or without chronic kidney disease. 2013 , 28, 1463-71 | | | 40 |
| 703 | Relation of the severity of contrast induced nephropathy to SYNTAX score and long term prognosis in patients treated with primary percutaneous coronary intervention. <i>International Journal of Cardiology</i> , 2013 , 168, 3480-5 | 3.2 | | 21 |
| 702 | Gene polymorphisms of interleukin-10 and tumor necrosis factor- α are associated with contrast-induced nephropathy. 2013 , 37, 110-7 | | | 23 |
| 701 | Atherosclerosis: Clinical Perspectives Through Imaging. 2013 , | | | 2 |
| 700 | Current concepts of contrast-induced nephropathy: a brief review. 2013 , 76, 673-81 | | | 28 |
| 699 | Age, glomerular filtration rate, ejection fraction, and the AGEF score predict contrast-induced nephropathy in patients with acute myocardial infarction undergoing primary percutaneous coronary intervention. 2013 , 82, 878-85 | | | 55 |
| 698 | Development of an easily applicable risk score model for contrast-induced nephropathy prediction after percutaneous coronary intervention: a novel approach tailored to current practice. <i>International Journal of Cardiology</i> , 2013 , 163, 46-55 | 3.2 | | 52 |
| 697 | Low contrast- and low radiation dose protocol for cardiac CT of thin adults at 256-row CT: usefulness of low tube voltage scans and the hybrid iterative reconstruction algorithm. 2013 , 29, 913-23 | | | 32 |
| 696 | Chapter 5: Referral to specialists and models of care. 2013 , 3, 112-119 | | | 30 |
| 695 | Prevention of contrast-induced nephropathy with Na/K citrate. 2013 , 34, 2362-7 | | | 25 |

| | | | |
|-----|--|-----|-----|
| 694 | Contrast-induced acute kidney injury: how much contrast is safe?. 2013 , 28, 1376-83 | | 45 |
| 693 | Renal function after percutaneous coronary interventions depending on the type of hydration. 2013 , 58, 369-75 | | 1 |
| 692 | Efficacy of allopurinol pretreatment for prevention of contrast-induced nephropathy: a randomized controlled trial. <i>International Journal of Cardiology</i> , 2013 , 167, 1396-9 | 3.2 | 40 |
| 691 | Acute kidney injury in patients undergoing cardiac surgery and coronary angiography on the same day. 2013 , 95, 513-9 | | 42 |
| 690 | Evaluation of renal function tests by age and sex to determine emergency department patients' eligibility for cardiac computed tomography. 2013 , 45, 220-7 | | 1 |
| 689 | Contrast-induced acute kidney injury and clinical outcomes after intra-arterial and intravenous contrast administration: risk comparison adjusted for patient characteristics by design. 2013 , 165, 793-99, 799.e1 | | 42 |
| 688 | Risk factors for acute kidney injury in the elderly undergoing contrast-enhanced computed tomography in the emergency department. 2013 , 76, 271-6 | | 14 |
| 687 | Contrast volume use in manual vs automated contrast injection systems for diagnostic coronary angiography and percutaneous coronary interventions. 2013 , 29, 372-6 | | 10 |
| 686 | Effect of short-term different statins loading dose on renal function and CI-AKI incidence in patients undergoing invasive coronary procedures. <i>International Journal of Cardiology</i> , 2013 , 168, 5101-3 ^{3.2} | | 6 |
| 685 | Renal function-based contrast dosing predicts acute kidney injury following transcatheter aortic valve implantation. <i>JACC: Cardiovascular Interventions</i> , 2013 , 6, 479-86 | 5 | 90 |
| 684 | N-acetylcysteine plus intravenous fluids versus intravenous fluids alone to prevent contrast-induced nephropathy in emergency computed tomography. 2013 , 62, 511-520.e25 | | 24 |
| 683 | High dose statins should be used in all patients undergoing percutaneous coronary intervention. <i>International Journal of Cardiology</i> , 2013 , 168, 5103-4 | 3.2 | |
| 682 | Automated 3D analysis of pre-procedural MDCT to predict annulus plane angulation and C-arm positioning: benefit on procedural outcome in patients referred for TAVR. 2013 , 6, 238-48 | | 49 |
| 681 | Feasibility study of the RenalGuard [®] balanced hydration system: a novel strategy for the prevention of contrast-induced nephropathy in high risk patients. <i>International Journal of Cardiology</i> , 2013 , 166, 482-6 | 3.2 | 34 |
| 680 | Prevention of contrast-induced nephropathy in STEMI patients undergoing primary percutaneous coronary intervention: a systematic review. 2013 , 26, 97-105 | | 12 |
| 679 | Risk factors for radiocontrast nephropathy after emergency department contrast-enhanced computerized tomography. 2013 , 20, 40-5 | | 20 |
| 678 | [Acute renal failure after cardiac surgery: evaluation of the RIFLE criteria]. 2013 , 9, 108-14 | | 11 |
| 677 | Contrast material-induced nephrotoxicity and intravenous low-osmolality iodinated contrast material. 2013 , 267, 94-105 | | 140 |

| | | |
|-----|---|---------|
| 676 | Chapter 3: Management of progression and complications of CKD. 2013 , 3, 73-90 | 86 |
| 675 | References. 2013 , 3, 136-150 | 66 |
| 674 | Specific Etiologies. 2013 , 199-236 | |
| 673 | Intravenous Contrast in CT and MR Imaging: Risks. 2013 , 81-94 | |
| 672 | A decision analysis to determine a testing threshold for computed tomographic angiography and D-dimer in the evaluation of aortic dissection. 2013 , 31, 1047-55 | 16 |
| 671 | A reduced contrast volume acquisition regimen based on cardiovascular dynamics improves visualisation of head and neck vasculature with carotid MDCT angiography. 2013 , 82, e64-9 | 15 |
| 670 | Renal protection: preconditioning for the prevention of contrast-induced nephropathy. 2013 , 26, 144-9 | 3 |
| 669 | Incidence and Main Determinants of Contrast-Induced Nephropathy following Coronary Angiography or Subsequent Balloon Angioplasty. 2013 , 3, 128-135 | 13 |
| 668 | Sichere Anwendung von Kontrastmittel in der elektiven CT-Diagnostik. 2013 , 36, 163-168 | |
| 667 | Why is diabetes mellitus a risk factor for contrast-induced nephropathy?. 2013 , 2013, 123589 | 67 |
| 666 | Risk factors for hemodialysis and mortality in patients with contrast-induced nephropathy. 2013 , 20, 607-12 | 5 |
| 665 | Association of central pulse pressure with contrast-induced nephropathy and clinical outcomes in patients undergoing coronary intervention. 2013 , 31, 2187-94 | 9 |
| 664 | Acute renal dysfunction: time from coronary angiography to cardiac surgery. 2013 , 21, 649-54 | 5 |
| 663 | Disparities in revascularization rates after acute myocardial infarction between aboriginal and non-aboriginal people in Australia. <i>Circulation</i> , 2013 , 127, 811-9 | 16.7 23 |
| 662 | Contrast-induced acute kidney injury and risk of adverse clinical outcomes after coronary angiography: a systematic review and meta-analysis. 2013 , 6, 37-43 | 180 |
| 661 | Variation in use of left ventriculography in the Veterans Affairs Health Care System. 2013 , 6, 687-93 | 4 |
| 660 | Frequency of acute kidney injury following intravenous contrast medium administration: a systematic review and meta-analysis. 2013 , 267, 119-28 | 252 |
| 659 | Contrast-induced Nephropathy. 2013 , 14, 106-16 | 106 |

| | | | |
|-----|--|-----|----|
| 658 | Outcome of acute renal injury in diabetic mice with experimental endotoxemia: role of hypoxia-inducible factor-1 β . 2013 , 2013, 254529 | | 12 |
| 657 | Risk score for the prediction of contrast-induced nephropathy in elderly patients undergoing percutaneous coronary intervention. <i>Angiology</i> , 2013 , 64, 188-94 | 2.1 | 47 |
| 656 | Dosing of iodinated contrast volume: a new simple algorithm to stratify the risk of contrast-induced nephropathy in patients with acute coronary syndrome. 2013 , 82, 888-97 | | 12 |
| 655 | An uninvestigated risk factor for contrast-induced nephropathy in chronic kidney disease: proteinuria. 2013 , 35, 62-5 | | 11 |
| 654 | Contrast-induced acute kidney injury following PCI. 2013 , 43, 483-90 | | 47 |
| 653 | Renal function following fistulography in patients with advanced chronic kidney disease. 2013 , 35, 791-5 | | 3 |
| 652 | Acute kidney injury after endovascular repair of abdominal aortic aneurysm. 2013 , 20, 315-30 | | 58 |
| 651 | Risk of contrast-induced nephropathy in patients undergoing endovascular treatment of acute ischemic stroke. 2013 , 5, 543-5 | | 25 |
| 650 | Acetylcysteine for the prevention of renal outcomes in patients with diabetes mellitus undergoing coronary and peripheral vascular angiography: a substudy of the acetylcysteine for contrast-induced nephropathy trial. 2013 , 6, 139-45 | | 19 |
| 649 | Guidelines on the use of iodinated contrast media in patients with kidney disease 2012: digest version. 2013 , 77, 1883-914 | | 10 |
| 648 | Rosuvastatin versus atorvastatin to prevent contrast induced nephropathy in patients undergoing primary percutaneous coronary intervention (ROSA-CIN trial). <i>Acta Cardiologica</i> , 2013 , 68, 489-94 | 0.9 | 26 |
| 647 | Low-contrast dose protection protocol for diagnostic computed tomography in patients at high-risk for contrast-induced nephropathy. 2013 , 37, 289-96 | | 6 |
| 646 | Kidney enhancement on X-rays following emergency percutaneous coronary procedure predicts poor short- and long-term clinical outcomes. 2013 , 77, 982-7 | | 0 |
| 645 | Preventive effect of pretreatment with intravenous nicorandil on contrast-induced nephropathy in patients with renal dysfunction undergoing coronary angiography (PRINCIPLE Study). 2013 , 54, 957-64 | | 20 |
| 644 | Epidemiology, Clinical Features and Diagnosis of Contrast Induced Nephropathy: A Brief Review. 2013 , 01, | | |
| 643 | Radiographic contrast-media-induced acute kidney injury: pathophysiology and prophylactic strategies. 2013 , 2013, 496438 | | 23 |
| 642 | The efficacy of the cystatin C based glomerular filtration rate in the estimation of safe contrast media volume. 2013 , 43, 622-7 | | 3 |
| 641 | Renal Cortical Lactate Dehydrogenase: A Useful, Accurate, Quantitative Marker of In Vivo Tubular Injury and Acute Renal Failure. 2013 , 8, e66776 | | 45 |

| | | |
|-----|--|-----|
| 640 | Contrast-Induced Nephropathy: Risk Factors, Clinical Implication, Diagnostics Approach, Prevention. 2013 , | |
| 639 | Patients at risk for contrast-induced acute kidney injury. 2013 , 223-229 | |
| 638 | Development of contrast-induced acute kidney injury after elective contrast media exposure in patients with type 2 diabetes mellitus: effect of albuminuria. 2014 , 9, e106454 | 12 |
| 637 | Side effects of radiographic contrast media: pathogenesis, risk factors, and prevention. 2014 , 2014, 741018 | 139 |
| 636 | Prevention of contrast-induced nephropathy through a knowledge of its pathogenesis and risk factors. 2014 , 2014, 823169 | 68 |
| 635 | The Choice of the Iodinated Radiographic Contrast Media to Prevent Contrast-Induced Nephropathy. 2014 , 2014, 1-11 | 8 |
| 634 | Follow up for emergency department patients after intravenous contrast and risk of nephropathy. 2014 , 15, 276-81 | 8 |
| 633 | Urinary metabolomic analysis to detect changes after intravenous, non-ionic, low osmolar iodinated radiocontrast for computerized tomographic imaging. 2014 , 15, 152-7 | 3 |
| 632 | Acetylcysteine for preventing contrast-induced nephropathy. 2014 , | |
| 631 | Incidence and Predictors of Contrast Induced Nephropathy after Coronary Intervention at College of Medical Sciences Teaching Hospital, Bharatpur. 2014 , 11, 3-11 | |
| 630 | Do tubular changes in the diabetic kidney affect the susceptibility to acute kidney injury?. 2014 , 127, 133-8 | 16 |
| 629 | Accuracy of point-of-care serum creatinine devices for detecting patients at risk of contrast-induced nephropathy: a critical overview. 2014 , 51, 332-43 | 15 |
| 628 | Benefit of statin pretreatment in prevention of contrast-induced nephropathy in different adult patient population: systematic review and meta-analysis. 2014 , 1, e000127 | 20 |
| 627 | Practical Manual of Interventional Cardiology. 2014 , | 2 |
| 626 | Pre-diabetes and the risk of contrast induced nephropathy in patients undergoing coronary angiography or percutaneous intervention. 2014 , 106, 458-64 | 15 |
| 625 | Cystatin C as biomarker of contrast-induced nephropathy in pediatric cardiac angiography. 2014 , 44, 178-85 | 3 |
| 624 | Comparative study of renal protective effects of allopurinol and N-acetyl-cysteine on contrast induced nephropathy in patients undergoing cardiac catheterization. 2014 , 8, HC03-7 | 16 |
| 623 | Extended renal outcomes with use of iodixanol versus iohexol after coronary angiography. 2014 , 2014, 506479 | 5 |

| | | |
|-----|--|--------|
| 622 | Incidence, risk factors, and prognosis of acute kidney injury following transarterial chemoembolization in patients with hepatocellular carcinoma: a prospective cohort study. 2015 , 51 Suppl 2, e3-8 | 8 |
| 621 | Acute kidney injury by radiographic contrast media: pathogenesis and prevention. 2014 , 2014, 362725 | 77 |
| 620 | Pharmacological strategies to prevent contrast-induced acute kidney injury. 2014 , 2014, 236930 | 25 |
| 619 | The role of antiplatelet therapy in aneurysm coiling. 2014 , 36, 383-8 | 19 |
| 618 | Contrast Medium-Induced Nephropathy. 2014 , 81-104 | 4 |
| 617 | Neutrophil gelatinase-associated lipocalin as an early marker of contrast-induced nephropathy after coronary angiography. <i>Angiology</i> , 2014 , 65, 216-23 | 2.1 24 |
| 616 | Intravenous contrast material exposure is not an independent risk factor for dialysis or mortality. 2014 , 273, 714-25 | 163 |
| 615 | Incidence of contrast material-induced nephropathy after neuroendovascular procedures. 2014 , 273, 853-8 | 19 |
| 614 | Quality improvement guidelines for diagnostic arteriography. 2014 , 25, 1873-81 | 39 |
| 613 | A simple preprocedural score for risk of contrast-induced acute kidney injury after percutaneous coronary intervention. 2014 , 83, E8-16 | 38 |
| 612 | A randomized trial comparing dual axis rotational versus conventional coronary angiography in a population with a high prevalence of coronary artery disease. 2014 , 27, 456-64 | 7 |
| 611 | Association between N-terminal pro-brain natriuretic peptide levels and contrast-induced nephropathy in patients undergoing percutaneous coronary intervention for acute coronary syndrome. 2014 , 37, 485-92 | 22 |
| 610 | Reducing iodinated contrast volume by manipulating injection pressure during coronary angiography. 2014 , 83, 741-5 | 9 |
| 609 | Pre-procedural fasting for coronary interventions: is it time to change practice?. 2014 , 100, 658-61 | 12 |
| 608 | Renal function-adjusted contrast volume redefines the baseline estimation of contrast-induced acute kidney injury risk in patients undergoing primary percutaneous coronary intervention. 2014 , 7, 465-72 | 49 |
| 607 | Acute kidney injury after primary angioplasty: effect of different hydration treatments. 2014 , 15, 60-7 | 11 |
| 606 | Does intravenous or oral high-dose N-acetylcysteine in addition to saline prevent contrast-induced nephropathy assessed by cystatin C?. 2014 , 25, 111-7 | 9 |
| 605 | Effect of statins in preventing contrast-induced nephropathy: an updated meta-analysis. 2014 , 25, 565-74 | 8 |

| | | | |
|-----|--|-----|-----|
| 604 | The use of nitrates in the prevention of contrast-induced nephropathy in patients hospitalized after undergoing percutaneous coronary intervention. 2014 , 19, 310-4 | | 6 |
| 603 | Frequency and risk factors of contrast-induced nephropathy after cardiac catheterization in type II diabetic patients: a study among Egyptian patients. 2014 , 36, 191-7 | | 8 |
| 602 | Validated contemporary risk model of acute kidney injury in patients undergoing percutaneous coronary interventions: insights from the National Cardiovascular Data Registry Cath-PCI Registry. 2014 , 3, e001380 | | 114 |
| 601 | Low-contrast-dose protocol in cardiac CT: 20% contrast dose reduction using 100 kVp and high-tube-current-time setting in 256-slice CT. 2014 , 55, 545-53 | | 11 |
| 600 | Oral hydration for prevention of contrast-induced acute kidney injury in elective radiological procedures: a systematic review and meta-analysis of randomized controlled trials. 2014 , 6, 618-24 | | 30 |
| 599 | Contrast-induced acute kidney injury: definition, epidemiology, and outcome. 2014 , 2014, 859328 | | 34 |
| 598 | Contrast media: are there differences in nephrotoxicity among contrast media?. 2014 , 2014, 934947 | | 34 |
| 597 | Parkinsonian tremor-induced electrocardiographic artifacts mimicking atrial flutter/fibrillation or ventricular tachycardia. <i>International Journal of Cardiology</i> , 2014 , 173, 597-600 | 3.2 | 10 |
| 596 | Impact of continuous deterioration of kidney function 6 to 8 months after percutaneous coronary intervention for acute coronary syndrome. 2014 , 113, 1647-51 | | 32 |
| 595 | High BNP level as risk factor for acute kidney injury and predictor of all-cause mortality in STEMI patients. 2014 , 39, 507-14 | | 20 |
| 594 | Incidence of contrast-induced nephropathy in hospitalised patients with cancer. 2014 , 24, 184-90 | | 37 |
| 593 | Lower blood vitamin D levels are associated with an increased incidence of contrast-induced nephropathy in patients undergoing coronary angiography. 2014 , 30, 428-33 | | 11 |
| 592 | [Perioperative acute kidney injury and failure]. 2014 , 10, 121-31 | | 3 |
| 591 | Prevention of contrast-induced nephropathy with N-acetylcysteine or sodium bicarbonate in patients with ST-segment-myocardial infarction: a prospective, randomized, open-labeled trial. 2014 , 7, 216-24 | | 39 |
| 590 | Canadian Association of Radiologists consensus guidelines for the prevention of contrast-induced nephropathy: update 2012. 2014 , 65, 96-105 | | 105 |
| 589 | Neutrophil-to-lymphocyte ratio predicts contrast-induced nephropathy in patients undergoing primary percutaneous coronary intervention. <i>Angiology</i> , 2014 , 65, 51-6 | 2.1 | 46 |
| 588 | Contemporary incidence, predictors, and outcomes of acute kidney injury in patients undergoing percutaneous coronary interventions: insights from the NCDR Cath-PCI registry. <i>JACC: Cardiovascular Interventions</i> , 2014 , 7, 1-9 | 5 | 315 |
| 587 | Remote ischemic preconditioning and renoprotection: from myth to a novel therapeutic option?. 2014 , 25, 216-24 | | 99 |

| | | | |
|-----|---|-----|----|
| 586 | Peter Andrew McCullough, MD, MPH: an interview with the editor. 2014 , 114, 1772-85 | | 0 |
| 585 | Ultra low-dose of gadobenate dimeglumine for late gadolinium enhancement (LGE) imaging in acute myocardial infarction: a feasibility study. 2014 , 83, 2151-2158 | | 9 |
| 584 | Neuroimaging in intracerebral hemorrhage. 2014 , 45, 903-8 | | 75 |
| 583 | Efficacy of oral hydration in the prevention of contrast-induced acute kidney injury in patients undergoing coronary angiography or intervention. 2014 , 128, 95-100 | | 19 |
| 582 | Predictive factors of contrast-induced nephropathy in patients undergoing primary coronary angioplasty. 2014 , 107, 424-32 | | 15 |
| 581 | Prediction of contrast-induced nephropathy in patients with serum creatinine levels in the upper normal range by cystatin C: a prospective study in 374 patients. 2014 , 202, 452-8 | | 14 |
| 580 | Perioperative Management of the Patient with Diabetes. 2014 , 32, 171-190 | | |
| 579 | A Practical Approach to Preventing Renal Complications in the Catheterization Laboratory. 2014 , 3, 429-439 | | |
| 578 | Predicting Contrast-induced Renal Complications in the Catheterization Laboratory. 2014 , 3, 369-377 | | 4 |
| 577 | The influence of contrast media on kidney function in patients with stable coronary artery disease. 2014 , 48, 234-40 | | |
| 576 | The role of inflammation in contrast-induced nephropathy. 2014 , 87, 20130738 | | 38 |
| 575 | Acute kidney injury after CABG versus PCI: an observational study using 2 cohorts. 2014 , 64, 985-94 | | 28 |
| 574 | The spectrum of risk factors for contrast induced nephropathy in patients undergoing coronary angiography or intervention. 2014 , 4, 157-161 | | 2 |
| 573 | Contrast-induced nephropathy in patients with ST elevation myocardial infarction treated with primary percutaneous coronary intervention. <i>Angiology</i> , 2014 , 65, 37-42 | 2.1 | 27 |
| 572 | Contrast between innovator drug- and generic drug-induced renal dysfunction on coronary angiography (CONTRAST study). 2014 , 29, 603-10 | | 2 |
| 571 | Remote ischemic preconditioning to reduce contrast-induced nephropathy: study protocol for a randomized controlled trial. 2014 , 15, 119 | | 8 |
| 570 | Acute kidney injury after transcatheter aortic valve implantation: incidence, predictors and impact on mortality. 2014 , 107, 133-9 | | 81 |
| 569 | The incidence of acute kidney injury after cardiac catheterization or PCI: a comparison of radial vs. femoral approach. <i>International Journal of Cardiology</i> , 2014 , 173, 595-7 | 3.2 | 22 |

| | | | |
|-----|---|-----|----|
| 568 | Contrast-induced nephropathy: the sin of primary percutaneous coronary intervention?. 2014 , 35, 1504-6 | | 16 |
| 567 | Statins and contrast-induced acute kidney injury with coronary angiography. <i>American Journal of Medicine</i> , 2014 , 127, 987-1000 | 2.4 | 43 |
| 566 | Risk scoring system to predict contrast induced nephropathy following percutaneous coronary intervention. 2014 , 66, 517-24 | | 21 |
| 565 | Relation of contrast-induced nephropathy to long-term mortality after percutaneous coronary intervention. 2014 , 114, 362-8 | | 69 |
| 564 | Pitfalls and complications management in the endovascular treatment of aneurysms. 2014 , 25, 405-13 | | 1 |
| 563 | CT angiography of the neck: value of contrast medium dose reduction with low tube voltage and high tube current in a 64-detector row CT. 2014 , 69, e183-9 | | 15 |
| 562 | Elevation of urinary liver-type fatty acid-binding protein as predicting factor for occurrence of contrast-induced acute kidney injury and its reduction by hemodiafiltration with blood suction from right atrium. 2014 , 29, 191-7 | | 12 |
| 561 | Acute Kidney Injury in Intensive Care. 2014 , 647-686 | | |
| 560 | Requests for Imaging Using Contrast Media: What Information Must be Provided. 2014 , 13-16 | | |
| 559 | Kriterien für die Notwendigkeit und Dauer von Krankenhausbehandlung bei Koronarangiografien und -interventionen. 2015 , 9, 295-302 | | 4 |
| 558 | Impact of high dose versus low dose atorvastatin on contrast induced nephropathy in diabetic patients with acute coronary syndrome undergoing early percutaneous coronary intervention. 2015 , 67, 329-336 | | 3 |
| 557 | Acute Kidney Injury After Carotid Artery Stenting. <i>JACC: Cardiovascular Interventions</i> , 2015 , 8, 1515-1517 | | 6 |
| 556 | Pharmacological interventions for preventing contrast-induced nephropathy. 2015 , | | |
| 555 | Systematic oral hydration with water is similar to parenteral hydration for prevention of contrast-induced nephropathy: an updated meta-analysis of randomised clinical data. 2015 , 2, e000317 | | 21 |
| 554 | Renal replacement therapy for preventing contrast-induced nephropathy. 2015 , | | |
| 553 | Real-Time MRI-Guided Catheter Tracking Using Hyperpolarized Silicon Particles. <i>Scientific Reports</i> , 2015 , 5, 12842 | 4.9 | 20 |
| 552 | Remote ischemic preconditioning for prevention of contrast-medium-induced nephropathy: (RenoProtection-trial). 2015 , 5, 297-304 | | |
| 551 | Kontrastmittelnephropathie. 2015 , 10, 399-400 | | |

| | | |
|-----|---|----|
| 550 | Noninvasive Qureshi Grading Scheme Predicts 90-Day mRS in Patients with Acute Ischemic Stroke. 2015 , 25, 761-5 | 1 |
| 549 | Endovascular therapy by CO ₂ angiography to prevent contrast-induced nephropathy in patients with chronic kidney disease: a prospective multicenter trial of CO ₂ angiography registry. 2015 , 85, 870-7 | 53 |
| 548 | Temporal trends of fluoroscopy time and contrast utilization in coronary chronic total occlusion revascularization: insights from a multicenter United States registry. 2015 , 85, 393-9 | 43 |
| 547 | Renal safety of intravenous gadolinium-enhanced magnetic resonance imaging in patients awaiting liver transplantation. 2015 , 21, 1340-6 | 8 |
| 546 | 3-Hydroxy-3-methylglutaryl-CoA reductase enzyme inhibitors for prevention of contrast-induced nephropathy: a meta-analysis of prospective randomized controlled studies. 2015 , 22, e158-66 | 7 |
| 545 | Novel Biomarkers of Acute Kidney Injury After Contrast Coronary Angiography. 2015 , 23, 240-6 | 20 |
| 544 | Short-term rosuvastatin treatment for the prevention of contrast-induced acute kidney injury in patients receiving moderate or high volumes of contrast media: a sub-analysis of the TRACK-D study. 2015 , 128, 784-9 | 4 |
| 543 | Cardiorenal Syndromes: Advances in Determining Diagnosis, Prognosis and Therapy. 2015 , 03, | 3 |
| 542 | Korean Guidelines for Interventional Recanalization of Lower Extremity Arteries. 2015 , 16, 696-722 | 6 |
| 541 | Current status of sodium bicarbonate in coronary angiography: an updated comprehensive meta-analysis and systematic review. 2015 , 2015, 690308 | 5 |
| 540 | The Effect of Renin-Angiotensin-Aldosterone System Blockade Medications on Contrast-Induced Nephropathy in Patients Undergoing Coronary Angiography: A Meta-Analysis. 2015 , 10, e0129747 | 16 |
| 539 | Acetylcysteine for preventing contrast-induced nephropathy. 2015 , | |
| 538 | Contrast-Induced Nephropathy: Identifying the Risks, Choosing the Right Agent, and Reviewing Effective Prevention and Management Methods. 2015 , 44, 501-4 | 26 |
| 537 | Contrast-induced nephropathy following chronic total occlusion percutaneous coronary intervention in patients with chronic kidney disease. 2015 , 25, 2274-81 | 10 |
| 536 | The EUROpean and Chinese cardiac and renal Remote Ischemic Preconditioning Study (EURO-CRIPS): study design and methods. 2015 , 16, 246-52 | 8 |
| 535 | High-dose atorvastatin for preventing contrast-induced nephropathy in primary percutaneous coronary intervention. 2015 , 16, 213-9 | 12 |
| 534 | Remote Ischemic Preconditioning To Reduce Contrast-Induced Nephropathy: A Randomized Controlled Trial. 2015 , 50, 527-32 | 42 |
| 533 | Pathogenesis and treatment of the cardiorenal syndrome: Implications of L-arginine-nitric oxide pathway impairment. 2015 , 154, 1-12 | 27 |

| | | | |
|-----|---|-----|----|
| 532 | Application of low tube voltage coronary CT angiography with low-dose iodine contrast agent in patients with a BMI of 26-30 kg/m ² . 2015 , 70, 138-45 | | 24 |
| 531 | [Atherosclerotic ischemic renal disease: clinical challenges]. 2015 , 144, 163-5 | | |
| 530 | The usefulness of carvedilol and nebivolol in preventing contrast nephropathy in rats. 2015 , 37, 511-7 | | 7 |
| 529 | Trimetazidine Decreases Risk of Contrast-Induced Nephropathy in Patients With Chronic Kidney Disease: A Meta-Analysis of Randomized Controlled Trials. 2015 , 20, 539-46 | | 16 |
| 528 | Continuous intravenous infusion of nicorandil for 4 hours before and 24 hours after percutaneous coronary intervention protects against contrast-induced nephropathy in patients with poor renal function. <i>International Journal of Cardiology</i> , 2015 , 195, 228-34 | 3.2 | 25 |
| 527 | Novel risk score of contrast-induced nephropathy after percutaneous coronary intervention. 2015 , 20, 544-51 | | 18 |
| 526 | Incidence of Acute Kidney Injury (AKI) after Endovascular Abdominal Aortic Aneurysm Repair (EVAR) and Impact on Outcome. 2015 , 49, 534-40 | | 77 |
| 525 | Balloon pulmonary angioplasty using contrast agents improves impaired renal function in patients with chronic thromboembolic pulmonary hypertension. <i>International Journal of Cardiology</i> , 2015 , 188, 41-2 | 3.2 | 15 |
| 524 | Prevalencia y pronóstico de la nefropatía tras intervencionismo coronario percutáneo de pacientes con síndrome coronario agudo y función renal normal. 2015 , 68, 310-316 | | 2 |
| 523 | Association between Platelet-to-Lymphocyte Ratio and Contrast-Induced Nephropathy in Patients Undergoing Percutaneous Coronary Intervention for Acute Coronary Syndrome. 2015 , 5, 96-104 | | 18 |
| 522 | Effect of Contrast-Induced Nephropathy on the Long-Term Outcome of Patients with Non-ST Segment Elevation Myocardial Infarction. 2015 , 5, 116-24 | | 13 |
| 521 | Contrast media controversies in 2015: imaging patients with renal impairment or risk of contrast reaction. 2015 , 204, 1174-81 | | 67 |
| 520 | Prevalence and prognosis of percutaneous coronary intervention-associated nephropathy in patients with acute coronary syndrome and normal kidney function. 2015 , 68, 310-6 | | 2 |
| 519 | Acute kidney injury after aortic valve replacement: incidence, risk factors and outcomes. 2015 , 13, 301-16 | | 75 |
| 518 | Do Cardiovascular Risk Factors and Coronary SYNTAX Score Predict Contrast Volume Use During Cardiac Catheterization?. <i>Angiology</i> , 2015 , 66, 933-40 | 2.1 | 6 |
| 517 | Time to revisit the problem of CIN? The low incidence of acute kidney injury with and without contrast in hospitalized patients: an observational cohort study. 2015 , 2, 38 | | 11 |
| 516 | Relative and absolute changes in urinary neutrophil gelatinase-associated lipocalin and correlation with small increases in serum creatinine levels after coronary angiography: an observational study. 2015 , 129, 84-90 | | 4 |
| 515 | Renoprevention revisited--The impact of preemptive withdrawal of RAAS blockade prior to iodinated contrast exposure in older CKD patients: results of a new meta-analysis. 2015 , 130, 25-6 | | 4 |

| | | | |
|-----|---|-----|-----|
| 514 | Contrast-associated Acute Kidney Injury. 2015 , 31, 725-35 | | 3 |
| 513 | Risk prediction models for contrast induced nephropathy: systematic review. 2015 , 351, h4395 | | 108 |
| 512 | Carfilzomib-related acute kidney injury may be prevented by N-acetyl-L-cysteine. 2015 , 21, 313-6 | | 21 |
| 511 | The Preinterventional Cystatin-Creatinine-Ratio: A Prognostic Marker for Contrast Medium-Induced Acute Kidney Injury and Long-Term All-Cause Mortality. 2015 , 131, 59-65 | | 10 |
| 510 | Impact of Contrast-Induced Acute Kidney Injury After Percutaneous Coronary Intervention on Short- and Long-Term Outcomes: Pooled Analysis From the HORIZONS-AMI and ACUITY Trials. 2015 , 8, e002475 | | 97 |
| 509 | Comparison of combination therapy of high-dose oral N-acetylcysteine and intravenous sodium bicarbonate hydration with individual therapies in the reduction of Contrast-induced Nephropathy during Cardiac Catheterisation and Percutaneous Coronary Intervention (CONTRAST): A multi-centre, randomised, controlled trial. <i>International Journal of Cardiology</i> , 2015 , 201, 237-42 | 3.2 | 29 |
| 508 | Atherosclerotic ischemic renal disease: Clinical challenges. 2015 , 144, 163-165 | | |
| 507 | Society of Thoracic Surgeons score predicts kidney injury in patients not undergoing bypass surgery. 2015 , 99, 123-9 | | 8 |
| 506 | Comprehensive Management of High Risk Cardiovascular Patients. 2016 , | | 1 |
| 505 | Urine metabolomic analysis to detect metabolites associated with the development of contrast induced nephropathy. 2016 , 3, 204-212 | | 3 |
| 504 | Angiotensin-Converting Enzyme Inhibitor Prevents the Worsening of Renal Function in the Late Phase after Percutaneous Coronary Intervention. <i>Journal of Atherosclerosis and Thrombosis</i> , 2016 , 23, 233-40 | 4 | 4 |
| 503 | [Not Available]. 2016 , 106, 699-703 | | 11 |
| 502 | Contrast Induced Nephropathy with Intravenous Iodinated Contrast Media in Routine Diagnostic Imaging: An Initial Experience in a Tertiary Care Hospital. 2016 , 2016, 8792984 | | 6 |
| 501 | Long-Term Follow-Up Evaluation of Renal Function in Patients with Chronic Kidney Disease Undergoing Cardiac Surgery. 2016 , 2016, 9680718 | | 2 |
| 500 | 7 Endovascular Surgery: General Technique. 2016 , | | |
| 499 | Iodinated contrast media. 2016 , 239-297 | | |
| 498 | Remote Ischemic Preconditioning for the Prevention of Contrast-Induced Acute Kidney Injury in Diabetics Receiving Elective Percutaneous Coronary Intervention. 2016 , 11, e0164256 | | 12 |
| 497 | Contrast Medium Exposure During Computed Tomography and Risk of Development of End-Stage Renal Disease in Patients With Chronic Kidney Disease: A Nationwide Population-Based, Propensity Score-Matched, Longitudinal Follow-Up Study. 2016 , 95, e3388 | | 17 |

| | | |
|-----|---|--------|
| 496 | Role of N-Acetylcysteine to Prevent Contrast-Induced Nephropathy: A Meta-analysis. 2016 , 23, e172-83 | 12 |
| 495 | AKIGUARD (Acute Kidney Injury GUARding Device) trial: in-hospital and one-year outcomes. 2016 , 17, 530-7 | 21 |
| 494 | Elevated serum fibrinogen levels and risk of contrast-induced acute kidney injury in patients undergoing a percutaneous coronary intervention for the treatment of acute coronary syndrome. 2016 , 27, 13-8 | 18 |
| 493 | Sodium Bicarbonate Prevents Contrast-Induced Nephropathy in Addition to Theophylline: A Randomized Controlled Trial. 2016 , 95, e3720 | 10 |
| 492 | Clinical Significance of Persistent Global and Focal Computed Tomography Nephrograms After Cardiac Catheterization and Their Relationships to Urinary Biomarkers of Kidney Damage and Procedural Factors: Pilot Study. 2016 , 51, 797-803 | 3 |
| 491 | Acute Kidney Injury After Transcatheter Aortic Valve Replacement. 2016 , 31, 416-22 | 23 |
| 490 | A Comparison of Nephrotoxicity of Contrast Medium in Elderly Patients who Underwent Renal or Peripheral Arterial Vascular Intervention. 2016 , 55, 9-14 | 9 |
| 489 | Relation of contrast nephropathy to adverse events in pulmonary emboli patients diagnosed with contrast CT. 2016 , 34, 1247-50 | 11 |
| 488 | Predicting Acute Kidney Injury in the Catheterization Laboratory. 2016 , 67, 1723-4 | 2 |
| 487 | Renal Function is the Main Predictor of Acute Kidney Injury after Endovascular Abdominal Aortic Aneurysm Repair. 2016 , 31, 52-9 | 29 |
| 486 | Efficacy of Short-Term Statin Treatment for the Prevention of Contrast-Induced Acute Kidney Injury in Patients Undergoing Coronary Angiography/ Percutaneous Coronary Intervention: A Meta-Analysis of 21 Randomized Controlled Trials. 2016 , 16, 201-19 | 32 |
| 485 | The effect of N-acetylcysteine on the incidence of contrast-induced kidney injury: A systematic review and trial sequential analysis. <i>International Journal of Cardiology</i> , 2016 , 209, 319-27 | 3.2 25 |
| 484 | Incidence and In-Hospital Mortality of Acute Kidney Injury (AKI) and Dialysis-Requiring AKI (AKI-D) After Cardiac Catheterization in the National Inpatient Sample. 2016 , 5, e002739 | 21 |
| 483 | Real-time magnetic resonance imaging-guided transcatheter aortic valve replacement. 2016 , 151, 1269-77 | 7 |
| 482 | Robot-assisted real-time magnetic resonance image-guided transcatheter aortic valve replacement. 2016 , 151, 1407-12 | 7 |
| 481 | Effects of combination therapy of statin and N-acetylcysteine for the prevention of contrast-induced nephropathy in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. <i>International Journal of Cardiology</i> , 2016 , 212, 100-6 | 3.2 15 |
| 480 | Long-Term (3 Years) Prognosis of Contrast-Induced Acute Kidney Injury After Coronary Angiography. 2016 , 117, 1741-6 | 9 |
| 479 | Short-Term High-Dose Vitamin E to Prevent Contrast Medium-Induced Acute Kidney Injury in Patients With Chronic Kidney Disease Undergoing Elective Coronary Angiography: A Randomized Placebo-Controlled Trial. 2016 , 5, e002919 | 21 |

| | | | |
|-----|---|-----|----|
| 478 | Effectiveness of N-Acetylcysteine for the Prevention of Contrast-Induced Nephropathy: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. 2016 , 5, | | 39 |
| 477 | Should nurses use mechanical chest compression devices during CPR?. 2016 , 34, 2044-2045 | | 6 |
| 476 | Contrast-induced nephropathy. 2016 , 480-482 | | |
| 475 | Asymmetric dimethylarginine predicts the risk of contrast-induced acute kidney injury in patients undergoing cardiac catheterization. 2016 , 254, 161-166 | | 8 |
| 474 | Comparison of Two Risk Models in Predicting the Incidence of Contrast-Induced Nephropathy after Percutaneous Coronary Intervention. 2016 , 29, 447-453 | | 6 |
| 473 | Association of radial versus femoral access with contrast-induced acute kidney injury in patients undergoing primary percutaneous coronary intervention for ST-elevation myocardial infarction. 2016 , 17, 546-551 | | 13 |
| 472 | Prediction of fluoroscopic angulations for transcatheter aortic valve implantation by CT angiography: influence on procedural parameters. 2017 , 18, 906-914 | | 20 |
| 471 | Contrast-induced nephropathy related to adverse events in pulmonary embolism patients: causation or conflation? Reply. 2016 , 34, 2043-2044 | | 1 |
| 470 | Contrast induced nephropathy in vascular surgery. 2016 , 117 Suppl 2, ii63-ii73 | | 14 |
| 469 | Hyperuricemia and contrast-induced acute kidney injury: A systematic review and meta-analysis. <i>International Journal of Cardiology</i> , 2016 , 224, 286-294 | 3.2 | 24 |
| 468 | The efficacy of N-acetylcysteine plus sodium bicarbonate in the prevention of contrast-induced nephropathy after cardiac catheterization and percutaneous coronary intervention: A meta-analysis of randomized controlled trials. <i>International Journal of Cardiology</i> , 2016 , 221, 251-9 | 3.2 | 13 |
| 467 | Carotid and cerebral CT angiography using low volume of iodinated contrast material and low tube voltage. 2016 , 97, 1173-1179 | | 14 |
| 466 | Determination of a Testing Threshold for Lumbar Puncture in the Diagnosis of Subarachnoid Hemorrhage after a Negative Head Computed Tomography: A Decision Analysis. 2016 , 23, 1119-1127 | | 7 |
| 465 | Outcomes, risk factors and health burden of contrast-induced acute kidney injury: an observational study of one million hospitalizations with image-guided cardiovascular procedures. 2016 , 17, 167 | | 34 |
| 464 | Renal Protection Using Remote Ischemic Peri-Conditioning During Inter-Facility Helicopter Transport of Patients With ST-Segment Elevation Myocardial Infarction: A Retrospective Study. 2016 , 29, 603-611 | | 7 |
| 463 | Serum hyperchloremia as a risk factor for acute kidney injury in patients with ST-segment elevation myocardial infarction undergoing percutaneous coronary intervention. 2016 , 29, 7-11 | | 15 |
| 462 | Outcome of 1051 Octogenarian Patients With ST-Segment Elevation Myocardial Infarction Treated With Primary Percutaneous Coronary Intervention: Observational Cohort From the London Heart Attack Group. 2016 , 5, | | 19 |
| 461 | Prevention of Contrast-Induced Renal Failure for the Interventional Cardiologist. 2016 , 9, | | 2 |

| | | |
|-----|--|-------|
| 460 | Feasibility of 320-row area detector CT coronary angiography using 40 mL of contrast material: assessment of image quality and diagnostic accuracy. 2016 , 26, 3802-3810 | 9 |
| 459 | Safety enhancement in adult body computed tomography scanning: comparison of iodixanol versus iohexol. 2016 , 5, 148 | 7 |
| 458 | Contrast-Induced Acute Kidney Injury: An Update. 2016 , 30, 215-28 | 50 |
| 457 | Is the Presence of a Solitary Kidney an Independent Risk Factor for Acute Kidney Injury after Contrast-enhanced CT?. 2016 , 278, 74-81 | 14 |
| 456 | Contrast volume reduction using third generation dual source computed tomography for the evaluation of patients prior to transcatheter aortic valve implantation. 2016 , 26, 4497-4504 | 35 |
| 455 | The prevalence and prognosis of contrast-induced acute kidney injury according to the definition in patients with acute myocardial infarction who underwent primary percutaneous coronary intervention. 2016 , 13, 29-33 | 3 |
| 454 | Imaging patients with renal impairment. 2016 , 41, 1108-21 | 2 |
| 453 | Contrast-induced nephropathy following angiography and cardiac interventions. 2016 , 102, 638-48 | 108 |
| 452 | Incidence and Risk Factors for Postcontrast Acute Kidney Injury in Survivors of Sudden Cardiac Arrest. 2016 , 67, 469-476.e1 | 5 |
| 451 | [Contrast-induced nephropathy: An update]. 2016 , 65, 87-94 | 2 |
| 450 | Contrast Induced Nephropathy and Long-term Renal Decline After Percutaneous Transluminal Angioplasty for Symptomatic Peripheral Arterial Disease. 2016 , 51, 386-93 | 29 |
| 449 | Incidence and impact of acute kidney injury in patients with acute coronary syndromes treated with coronary artery bypass grafting: Insights from the Harmonizing Outcomes With Revascularization and Stents in Acute Myocardial Infarction (HORIZONS-AMI) and Acute Catheterization and Urgent Intervention Triage Strategy (ACUTE) trials. 2016 , 171, 10-7 | 24 |
| 448 | Contrast-Induced Nephropathy: From Pathophysiology to Preventive Strategies. 2016 , 32, 247-55 | 144 |
| 447 | Automatic spectral imaging protocol selection and iterative reconstruction in abdominal CT with reduced contrast agent dose: initial experience. 2017 , 27, 374-383 | 31 |
| 446 | Changing Paradigms in Contrast Nephropathy. 2017 , 28, 397-399 | 12 |
| 445 | Risk of contrast-induced acute kidney injury in ST-elevation myocardial infarction patients undergoing multi-vessel intervention-meta-analysis of randomized trials and risk prediction modeling study using observational data. 2017 , 90, 205-212 | 13 |
| 444 | Efficacy and Safety of Coronary Sinus Aspiration During Coronary Angiography to Attenuate the Risk of Contrast-Induced Acute Kidney Injury in Predisposed Patients. 2017 , 10, e004348 | 6 |
| 443 | Race and contrast-induced nephropathy in patients undergoing coronary angiography and cardiac catheterization. <i>International Journal of Cardiology</i> , 2017 , 230, 610-613 | 3-2 4 |

| | | | |
|-----|---|-----|----|
| 442 | A New Preprocedure Risk Score for Predicting Contrast-Induced Acute Kidney Injury. 2017 , 33, 714-723 | | 22 |
| 441 | Prevention of Contrast-Induced Acute Kidney Injury by Furosemide With Matched Hydration in Patients Undergoing Interventional Procedures: A Systematic Review and Meta-Analysis of Randomized Trials. <i>JACC: Cardiovascular Interventions</i> , 2017 , 10, 355-363 | 5 | 60 |
| 440 | Trimetazidine in the prevention of contrast-induced nephropathy in chronic kidney disease. 2017 , 18, 315-319 | | 13 |
| 439 | Effect of radiographic contrast media on renal perfusion - First results. 2016 , 64, 287-295 | | 9 |
| 438 | Mono-Energy Coronary Angiography with a Compact Synchrotron Source. <i>Scientific Reports</i> , 2017 , 7, 42211 | 4-9 | 20 |
| 437 | In situ lactate dehydrogenase activity: a novel renal cortical imaging biomarker of tubular injury?. 2017 , 312, F465-F473 | | 26 |
| 436 | Contrast-induced nephropathy in patients undergoing endovascular peripheral vascular intervention: Incidence, risk factors, and outcomes as observed in the Blue Cross Blue Shield of Michigan Cardiovascular Consortium. 2017 , 30, 274-280 | | 34 |
| 435 | Longitudinal Risk of Adverse Events in Patients With Acute Kidney Injury After Percutaneous Coronary Intervention: Insights From the National Cardiovascular Data Registry. 2017 , 10, | | 39 |
| 434 | Serum cystatin c is not superior to serum creatinine for early diagnosis of contrast-induced nephropathy in patients who underwent angiography. 2017 , 31, | | 7 |
| 433 | Contrast-induced acute kidney injury in interventional cardiology: Emerging evidence and unifying mechanisms of protection by remote ischemic conditioning. 2017 , 18, 549-553 | | 13 |
| 432 | Proteinuria as an independent risk factor for contrast-induced acute kidney injury and mortality in patients with stroke undergoing cerebral angiography. 2017 , 9, 445-448 | | 9 |
| 431 | Insight of forced diuresis with matched controlled hydration strategy to prevent contrast-induced acute kidney injury in patients undergoing cardiovascular intervention. <i>International Journal of Cardiology</i> , 2017 , 242, 18 | 3-2 | |
| 430 | A retrospective observational cohort study investigating the association between acute kidney injury and all-cause mortality among patients undergoing endovascular repair of abdominal aortic aneurysms. 2017 , 31, 686-691 | | 2 |
| 429 | Successful Repair of Acute Type B and Retrograde Type A Aortic Dissection With Kidney Ischemia. 2017 , 51, 342-345 | | 1 |
| 428 | Predictive value of inflammatory factors on contrast-induced acute kidney injury in patients who underwent an emergency percutaneous coronary intervention. 2017 , 40, 719-725 | | 51 |
| 427 | Cerebral bone subtraction CT angiography using 80 kVp and sinogram-affirmed iterative reconstruction: contrast medium and radiation dose reduction with improvement of image quality. 2017 , 59, 127-134 | | 4 |
| 426 | The quest for equilibrium: exploring the thin red line between bleeding and ischaemic risks in the management of acute coronary syndromes in chronic kidney disease patients. 2017 , 32, 1967-1976 | | 11 |
| 425 | Sodium Bicarbonate Versus Sodium Chloride for Preventing Contrast-Associated Acute Kidney Injury in Critically Ill Patients: A Randomized Controlled Trial. 2017 , 45, 637-644 | | 17 |

| | | | |
|-----|--|-----|----|
| 424 | Effects of staged versus ad hoc percutaneous coronary interventions on renal function-Is there a benefit to staging?. 2017 , 18, 344-348 | | 9 |
| 423 | On Contrast-Induced Acute Kidney Injury, Risk Prediction, and the Future of Predictive Model Development. 2017 , 33, 711-713 | | 3 |
| 422 | Ascorbic acid ameliorates renal injury in a murine model of contrast-induced nephropathy. 2017 , 18, 101 | | 5 |
| 421 | Endovascular aneurysm repair (EVAR)- and transcatheter aortic valve replacement (TAVR)-associated acute kidney injury. 2017 , 91, 1312-1323 | | 28 |
| 420 | Effect of Atrial Fibrillation on Contrast-Induced Nephropathy Development in Patients With Non-ST-Segment Elevation Myocardial Infarction. <i>Angiology</i> , 2017 , 68, 871-876 | 2.1 | 4 |
| 419 | Impact of Transient or Persistent Contrast-induced Nephropathy on Long-term Mortality After Elective Percutaneous Coronary Intervention. 2017 , 120, 2146-2153 | | 17 |
| 418 | Effect of Access Site Choice on Acute Kidney Injury After Percutaneous Coronary Intervention. 2017 , 120, 2141-2145 | | 11 |
| 417 | Incidence of contrast induced acute kidney injury in patients undergoing percutaneous coronary intervention in North Indian population. 2017 , 7, 143-148 | | |
| 416 | Effect of iodinated contrast media on renal perfusion: A randomized comparison study in pigs using quantitative contrast-enhanced ultrasound (CEUS). <i>Scientific Reports</i> , 2017 , 7, 13125 | 4.9 | 13 |
| 415 | Contrast Induced Nephropathy. 2017 , 347-353 | | |
| 414 | Oral N-acetylcysteine for prophylaxis of contrast-induced nephropathy in patients following coronary angioplasty: A meta-analysis. 2017 , 14, 1568-1576 | | 12 |
| 413 | Contrast-Induced Acute Kidney Injury: Pathophysiology, Manifestations, Prevention, and Management. 2017 , 25, 743-753 | | 16 |
| 412 | Comparison of the Frequency of Acute Kidney Injury in Patients With Renal Transplant Who Underwent Percutaneous Coronary Intervention Versus Coronary Artery Bypass Graft Surgery. 2017 , 120, 1104-1109 | | 2 |
| 411 | Image quality and contrast agent exposure in cardiac computed tomography angiography prior to transcatheter aortic valve implantation procedures using different acquisition protocols. 2017 , 4, 75-83 | | 3 |
| 410 | Pre-procedural peripheral endothelial function is associated with increased serum creatinine following percutaneous coronary procedure in stable patients with a preserved estimated glomerular filtration rate. 2017 , 70, 461-469 | | 3 |
| 409 | Decreased mortality in patients prescribed vancomycin after implementation of antimicrobial stewardship program. 2017 , 45, 1194-1197 | | 9 |
| 408 | Current Risk of Contrast-Induced Acute Kidney Injury After Coronary Angiography and Intervention: A Reappraisal of the Literature. 2017 , 33, 1225-1228 | | 46 |
| 407 | Maximum Arterial Contrast Concentrations With Computed Tomography and Left Ventriculography: Implications for Contrast Nephrotoxicity Risk. 2017 , 41, 976-982 | | 8 |

| | | | |
|-----|---|-----|-----|
| 406 | Contrast-Induced Nephropathy Is Less Common in Patients with Good Coronary Collateral Circulation. 2017 , 7, 316-323 | | 0 |
| 405 | Kontrastmittel. 2017 , 21, 454-457 | | |
| 404 | Contrast-induced nephropathy: Basic concepts, pathophysiological implications and prevention strategies. 2017 , 180, 99-112 | | 101 |
| 403 | Contrast-induced kidney injury: how does it affect long-term cardiac mortality?. 2017 , 18, 908-915 | | 8 |
| 402 | Risk of contrast induced nephropathy in diabetic patients affected by critical limb ischemia and diabetic foot ulcers treated by percutaneous transluminal angioplasty of lower limbs. 2017 , 33, e2866 | | 7 |
| 401 | Irreversible deterioration of estimated glomerular filtration rate in patients with acute myocardial infarction after primary percutaneous coronary intervention. 2017 , 32, 318-324 | | |
| 400 | CT in the ED: Special Considerations. 401-514 | | |
| 399 | Risk Factors of Contrast-induced Acute Kidney Injury in Patients Undergoing Emergency Percutaneous Coronary Intervention. 2017 , 130, 45-50 | | 16 |
| 398 | Analysis of Urine Composition in Type II Diabetic Mice after Intervention Therapy Using Holothurian Polypeptides. 2017 , 5, 54 | | 3 |
| 397 | Contrast-Induced Nephropathy in Renal Transplant Recipients: A Single Center Experience. <i>Frontiers in Medicine</i> , 2017 , 4, 64 | 4-9 | 7 |
| 396 | The impact of N-acetylcysteine and ascorbic acid in contrast-induced nephropathy in critical care patients: an open-label randomized controlled study. 2017 , 21, 269 | | 20 |
| 395 | Efficacy and safety of a balanced salt solution versus a 0.9% saline infusion for the prevention of contrast-induced acute kidney injury (BASIC trial): a study protocol for a randomized controlled trial. 2017 , 18, 461 | | 2 |
| 394 | Contrast-induced acute kidney injury: A review of practical points. 2017 , 6, 86-99 | | 73 |
| 393 | Serum Creatinine Level doesn't Change much at 2nd Day after Percutaneous Coronary Intervention (PCI): Amount of Contrast Injected may be the Key Factor. 2017 , 9, 155-158 | | |
| 392 | The Pathogenesis, Outcomes, and Prevention of Contrast-Associated Acute Kidney Injury. 2018 , 274-303 | | |
| 391 | CKD-EPI versus Cockcroft-Gault formula for predicting contrast-induced nephropathy following percutaneous coronary intervention in patients without significant renal impairment. 2018 , 37, 25-33 | | |
| 390 | The EUROpean and Chinese cardiac and renal Remote Ischemic Preconditioning Study (EURO-CRIPS CardioGroup I): A randomized controlled trial. <i>International Journal of Cardiology</i> , 2018 , 257, 1-6 | 3-2 | 35 |
| 389 | Safety of cerebral angiography and neuroendovascular therapy in patients with chronic kidney disease. 2018 , 60, 529-533 | | 9 |

| | | | |
|-----|---|-----|----|
| 388 | Comparison of the Effect of Dual-Axis Rotational Coronary Angiography Versus Conventional Coronary Angiography on Frequency of Acute Kidney Injury, X-Ray Exposure Time, and Quantity of Contrast Medium Injected. 2018 , 121, 1046-1050 | | 5 |
| 387 | Prospective evaluation of the development of contrast-induced nephropathy in patients with acute coronary syndrome undergoing rotational coronary angiography vs. conventional coronary angiography: CINERAMA study. 2018 , 38, 169-178 | | 2 |
| 386 | Contrast-Induced Nephropathy in STEMI Patients With and Without Chronic Kidney Disease. 2018 , 17, 25-31 | | 7 |
| 385 | CKD-EPI versus Cockcroft-Gault formula for predicting contrast-induced nephropathy following percutaneous coronary intervention in patients without significant renal impairment. 2018 , 37, 25-33 | | 5 |
| 384 | Coronary artery disease in decompensated patients undergoing liver transplantation evaluation. 2018 , 24, 333-342 | | 46 |
| 383 | Contrast-induced nephropathy - an entity to bear in mind and to prevent: A nephrological perspective. 2018 , 37, 35-36 | | 1 |
| 382 | Prediction of contrast induced acute kidney injury using novel biomarkers following contrast coronary angiography. 2018 , 111, 103-110 | | 17 |
| 381 | Iso-osmolar contrast media and adverse renal and cardiac events after percutaneous cardiovascular intervention. 2018 , 7, 331-341 | | 6 |
| 380 | Magnetic iodixanol - a novel contrast agent and its early characterization. 2018 , 43, 10-19 | | 0 |
| 379 | Contrast-induced nephropathy - an entity to bear in mind and to prevent: A nephrological perspective. 2018 , 37, 35-36 | | |
| 378 | Dynamic CTA in Native Kidneys Using a Multiphase CT Protocol-Potential of Significant Reduction of Contrast Medium. 2018 , 25, 842-849 | | 2 |
| 377 | Platelet to Lymphocyte Ratio Predicts Contrast-Induced Nephropathy in Patients With ST-Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. <i>Angiology</i> , 2018 , 69, 71-78 | 2.1 | 30 |
| 376 | Predictors of acute kidney injury in patients admitted with ST-elevation myocardial infarction - results from the Bremen STEMI-Registry. 2018 , 7, 710-722 | | 28 |
| 375 | Prevention of contrast-induced nephropathy by limb ischemic preconditioning: underlying mechanisms and clinical effects. 2018 , 314, F319-F328 | | 8 |
| 374 | Prospective evaluation of the development of contrast-induced nephropathy in patients with acute coronary syndrome undergoing rotational coronary angiography vs. conventional coronary angiography: CINERAMA study. 2018 , 38, 169-178 | | 8 |
| 373 | The Assessment of the Risk of Acute Kidney Injury in Patients Undergoing an Urgent Endovascular Treatment Due to Severe Renal Bleeding. 2018 , 41, 398-405 | | 4 |
| 372 | A New Risk Factor Profile for Contrast-Induced Acute Kidney Injury in Patients Who Underwent an Emergency Percutaneous Coronary Intervention. <i>Angiology</i> , 2018 , 69, 523-531 | 2.1 | 9 |
| 371 | Can SGLT2 Inhibitors Cause Acute Renal Failure? Plausible Role for Altered Glomerular Hemodynamics and Medullary Hypoxia. 2018 , 41, 239-252 | | 40 |

| | | |
|-----|--|-----|
| 370 | Contrast-Induced Nephropathy and Oxygen Pretreatment in Patients With Impaired Renal Function. 2018 , 3, 65-72 | 2 |
| 369 | Cardiorenal Syndrome Type 1: Definition, Etiopathogenesis, Diagnostics and Treatment. 2018 , 19, 73-80 | |
| 368 | Outcomes after Angiography with Sodium Bicarbonate and Acetylcysteine. 2018 , 378, 603-614 | 258 |
| 367 | Measurement of Vascular Diameter in Computed Tomography Angiography With Reduced Iodine Load: Comparison of Virtual Monochromatic Imaging in Dual-Energy Computed Tomography and Conventional Polychromatic Scan In Vitro. 2018 , 42, 919-924 | 1 |
| 366 | The Effect of the First Computed Tomography Examination on Renal Function During the Initial Cancer Work-up in Newly Diagnosed Cancer Patients without Chronic/Kidney Disease: A Retrospective, Single-institution, Observational Study. 2018 , 10, | |
| 365 | Relationship between High Level of Estimated Glomerular Filtration Rate and Contrast-Induced Acute Kidney Injury in Patients who Underwent an Emergency Percutaneous Coronary Intervention. 2018 , 131, 2041-2048 | 2 |
| 364 | Meta-analysis of rosuvastatin efficacy in prevention of contrast-induced acute kidney injury. 2018 , 12, 3685-3690 | 9 |
| 363 | Prevention of postcontrast acute kidney injury after percutaneous transluminal angioplasty by inducing RenalGuard controlled furosemide forced diuresis with matched hydration: study protocol for a randomised controlled trial. 2018 , 8, e021842 | 4 |
| 362 | Contrast-Associated Acute Kidney Injury. 2018 , 95-112 | |
| 361 | A New and Simple Risk Predictor of Contrast-Induced Nephropathy in Patients Undergoing Primary Percutaneous Coronary Intervention: TIMI Risk Index. 2018 , 2018, 5908215 | 6 |
| 360 | Urinary semaphorin 3A as an early biomarker to predict contrast-induced acute kidney injury in patients undergoing percutaneous coronary intervention. 2018 , 51, e6487 | 9 |
| 359 | Baseline atrial fibrillation is associated with contrast-induced nephropathy after cardiac catheterization in coronary artery disease: Systemic review and meta-analysis. 2018 , 41, 1555-1562 | 6 |
| 358 | Contrast volume to estimated glomerular filtration rate ratio for prediction of contrast-induced acute kidney injury after cardiac catheterization in adults with congenital heart disease. 2018 , 92, 1301-1308 | 2 |
| 357 | Evaluation of Safety Guidelines on the Use of Iodinated Contrast Material: Conundrum Continued. 2018 , 53, 616-622 | 17 |
| 356 | Complications of Coronary Artery Interventions: Overview. 2018 , 951-974 | |
| 355 | Contrast medium induced acute kidney injury: a narrative review. 2018 , 31, 797-812 | 40 |
| 354 | Diagnostic use of contrast medium and the kidney: A challenge or a ghost?. 2018 , 2, 6-15 | |
| 353 | Low-voltage (80-kVp) abdominopelvic computed tomography allows 60% contrast dose reduction in patients at risk of contrast-induced nephropathy. 2018 , 51, 352-355 | 10 |

| | | | |
|-----|--|-----|----|
| 352 | Anemia (Hemoglobin \leq 3 g/dL) as a Risk Factor for Contrast-Induced Acute Kidney Injury Following Coronary Angiography. 2018 , 122, 961-965 | | 20 |
| 351 | Association of Syntax Score II with Contrast-induced Nephropathy and Hemodialysis Requirement in Patients with ST Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. 2018 , 48, 59-70 | | 6 |
| 350 | Effects of hyperglycaemia and elevated glycosylated haemoglobin on contrast-induced nephropathy after coronary angiography. 2018 , 16, 377-383 | | 7 |
| 349 | Emergency coronary angioplasty with stenting using Cordis diagnostic coronary catheters when there is difficulty in engaging guide catheters and bench evaluation of diagnostic and guide catheters. 2018 , 37, 117-125 | | 4 |
| 348 | Iodixanol versus iopromide in patients with renal insufficiency undergoing coronary angiography with or without PCI. 2018 , 97, e0617 | | 12 |
| 347 | Circulating Fibroblast Growth Factor 21 is Associated with Subsequent Renal Injury Events in Patients Undergoing Coronary Angiography. <i>Scientific Reports</i> , 2018 , 8, 12425 | 4-9 | 9 |
| 346 | MR-guided Cardiac Interventions. 2018 , 27, 115-128 | | 14 |
| 345 | Emergency coronary angioplasty with stenting using Cordis diagnostic coronary catheters when there is difficulty in engaging guide catheters and bench evaluation of diagnostic and guide catheters. 2018 , 37, 117-125 | | 1 |
| 344 | Left ventricular end diastolic pressure and contrast-induced acute kidney injury in patients with acute coronary syndrome undergoing percutaneous coronary intervention. 2018 , 19, 16-20 | | 5 |
| 343 | Prevention of contrast-induced nephropathy by adequate hydration combined with isosorbide dinitrate for patients with renal insufficiency and congestive heart failure. 2019 , 42, 21-25 | | 7 |
| 342 | Comparison of Outcomes After Percutaneous Coronary Interventions in Patients of Eighty Years and Above Compared With Those Less Than 80 Years. 2019 , 124, 1372-1379 | | 2 |
| 341 | Predicting contrast induced nephropathy in patients undergoing percutaneous coronary intervention. 2019 , 11, 2672-2674 | | 1 |
| 340 | A Nephrology Consult in the Neurocritical Care Unit. 2019 , 359-369 | | |
| 339 | Effect of iodinated contrast media on the oxygen tension in the renal cortico-medullary region of pigs. 2019 , 73, 261-270 | | 5 |
| 338 | Left ventricular end-diastolic pressure-guided hydration for the prevention of contrast-induced acute kidney injury in patients with stable ischemic heart disease: the LAKESIDE trial. 2019 , 51, 1815-1822 | | 5 |
| 337 | Sometimes neither water nor fire are more useful than friendship-a new risk score for prediction of contrast-induced nephropathy (CIN) and long-term adverse outcomes in patients undergoing coronary angiography. 2019 , 11, 2675-2679 | | 2 |
| 336 | Comparative Performance of Prediction Models for Contrast-Associated Acute Kidney Injury After Percutaneous Coronary Intervention. 2019 , 12, e005854 | | 5 |
| 335 | Cardiac Evaluation in Liver Transplantation. 2019 , 6, 328-337 | | |

| | | | |
|-----|---|-----|-----|
| 334 | Contrast Volume and Decline in Kidney Function in Optical Coherence Tomography-Guided Percutaneous Coronary Intervention. 2019 , 60, 1022-1029 | | 1 |
| 333 | Procedural Strategies to Reduce the Incidence of Contrast-Induced Acute Kidney Injury During Percutaneous Coronary Intervention. <i>JACC: Cardiovascular Interventions</i> , 2019 , 12, 1877-1888 | 5 | 42 |
| 332 | A prospective study of contrast preservation using ultra-low contrast delivery technique versus standard automated contrast injector system in coronary procedures. 2019 , 71, 297-302 | | 1 |
| 331 | Metabolomics assessment reveals oxidative stress and altered energy production in the heart after ischemic acute kidney injury in mice. 2019 , 95, 590-610 | | 31 |
| 330 | Comparison of Abdominal Computed Tomographic Enhancement and Organ Lesion Depiction Between Weight-Based Scanner Software Contrast Dosing and a Fixed-Dose Protocol in a Tertiary Care Oncologic Center. 2019 , 43, 155-162 | | 11 |
| 329 | Bimodal intravascular volumetric imaging combining OCT and MPI. 2019 , 46, 1371-1383 | | 3 |
| 328 | Use of Intracardiac Compared With Transesophageal Echocardiography for Left Atrial Appendage Occlusion in the Amulet Observational Study. <i>JACC: Cardiovascular Interventions</i> , 2019 , 12, 1030-1039 | 5 | 19 |
| 327 | Contrast-Associated Acute Kidney Injury. 2019 , 380, 2146-2155 | | 191 |
| 326 | Efficacy of post-procedural oral hydration volume on risk of contrast-induced acute kidney injury following primary percutaneous coronary intervention: study protocol for a randomized controlled trial. 2019 , 20, 290 | | 1 |
| 325 | Risk Prediction for Contrast-Induced Nephropathy in Cancer Patients Undergoing Computed Tomography under Preventive Measures. 2019 , 2019, 8736163 | | 6 |
| 324 | Effect of renal insufficiency and diabetes mellitus on in-hospital mortality after acute coronary syndromes treated with primary PCI. Results from the ALKK PCI Registry. <i>International Journal of Cardiology</i> , 2019 , 292, 43-49 | 3.2 | 3 |
| 323 | Safety and Efficacy of Tolvaptan for the Prevention of Contrast-Induced Acute Kidney Injury in Patients with Heart Failure and Chronic Kidney Disease. 2019 , 5, 100-106 | | 1 |
| 322 | The Role of CT Angiography in Evaluating Lower Extremity Trauma: 157 Patient Case Series at a Military Treatment Facility. 2019 , 184, e490-e493 | | 5 |
| 321 | Nephrotoxicity of iodinated contrast media: From pathophysiology to prevention strategies. 2019 , 116, 231-241 | | 51 |
| 320 | Negligible Risk of Acute Renal Failure Among Hospitalized Patients After Contrast-Enhanced Imaging With Iodinated Versus Gadolinium-Based Agents. 2019 , 54, 312-318 | | 12 |
| 319 | Immediate and delayed hypersensitivity after intra-arterial injection of iodinated contrast media: a prospective study in patients with coronary angiography. 2019 , 29, 5314-5321 | | 7 |
| 318 | Evaluation of contrast nephropathy in percutaneous treatment of chronic total occlusions. 2019 , 11, 95-100 | | 0 |
| 317 | Contrast Induced Acute Kidney Injury and its Impact on Mid-Term Kidney Function, Cardiovascular Events and Mortality. <i>Scientific Reports</i> , 2019 , 9, 16896 | 4.9 | 8 |

| | | |
|-----|---|----|
| 316 | Guideline on the Use of Iodinated Contrast Media in Patients With Kidney Disease 2018. 2019 , 83, 2572-2607 | 4 |
| 315 | Impact of pretreatment with carnitine and tadalafil on contrast-induced nephropathy in CKD patients. 2019 , 41, 976-986 | 6 |
| 314 | Persistent Renal Dysfunction in Patients Undergoing Primary Percutaneous Coronary Intervention for Acute Myocardial Infarction. 2019 , 8, e014096 | 8 |
| 313 | Comparison of full-iodine conventional CT and half-iodine virtual monochromatic imaging: advantages and disadvantages. 2019 , 29, 1400-1407 | 10 |
| 312 | Association of the blood urea nitrogen-to-left ventricular ejection fraction ratio with contrast-induced nephropathy in patients with acute coronary syndrome who underwent percutaneous coronary intervention. 2019 , 51, 475-481 | 1 |
| 311 | Contrast-induced nephropathy is associated with new-onset atrial fibrillation in acute coronary syndrome after cardiac catheterization: Systemic review and meta-analysis. 2019 , 24, e12625 | 1 |
| 310 | The Epidemiology of Acute Kidney Injury. 2019 , 684-697.e4 | |
| 309 | [Epidemiology of acute renal failure]. 2019 , 15, 63-69 | 4 |
| 308 | Clinical impact of baseline chronic kidney disease in patients undergoing transcatheter or surgical aortic valve replacement. 2019 , 93, 740-748 | 11 |
| 307 | Acute effects of Iodixanol on renal function after intra-arterial administration in patients with end-stage kidney disease. 2018 , 70, 391-398 | 2 |
| 306 | Contrast-induced nephropathy after peripheral vascular intervention: Long-term renal outcome and risk factors for progressive renal dysfunction. 2019 , 69, 913-920 | 9 |
| 305 | Duplex-guided versus Conventional Percutaneous Transluminal Angioplasty of Iliac TASC II A and B Lesion: A Randomized Controlled Trial. 2019 , 55, 138-147 | 5 |
| 304 | The C-Reactive Protein to Albumin Ratio Predicts Acute Kidney Injury in Patients With ST-Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. 2019 , 28, 1638-1645 ²² | |
| 303 | Examining antecedents of reconciliation following service failure and recovery. 2020 , 28, 417-433 | 4 |
| 302 | Kidney Biomarkers of Injury and Repair as Predictors of Contrast-Associated AKI: A Substudy of the PRESERVE Trial. 2020 , 75, 187-194 | 18 |
| 301 | Incidence, predictors, and relevance of acute kidney injury in patients undergoing left atrial appendage closure with Amplatzer occluders: a multicentre observational study. 2020 , 109, 444-453 | 7 |
| 300 | Guideline on the use of iodinated contrast media in patients with kidney disease 2018. 2020 , 24, 1-44 | 13 |
| 299 | Guideline on the use of iodinated contrast media in patients with kidney disease 2018. 2020 , 38, 3-46 | 2 |

| | | |
|-----|---|-------|
| 298 | Acute kidney injury after platinum-based transcatheter arterial chemoembolization and transarterial infusion chemotherapy in patients with hepatocellular carcinoma. 2020 , 50, 36-43 | 3 |
| 297 | Remote ischemic preconditioning reduces the incidence of contrast-induced nephropathy in patients undergoing coronary angiography/intervention: Systematic review and meta-analysis of randomized controlled trials. 2020 , 96, 1200-1212 | 4 |
| 296 | Editor's Choice - Acute Kidney Injury (AKI) in Aortic Intervention: Findings From the Midlands Aortic Renal Injury (MARI) Cohort Study. 2020 , 59, 899-909 | 16 |
| 295 | New Fast kVp Switching Dual-Energy CT: Reduced Severity of Beam Hardening Artifacts and Improved Image Quality in Reduced-Iodine Virtual Monochromatic Imaging. 2020 , 27, 1586-1593 | 8 |
| 294 | Acute Kidney Injury After Radiocontrast-Enhanced Computerized Tomography in Hospitalized Patients With Advanced Renal Failure: A Propensity-Score-Matching Analysis. 2020 , 55, 677-687 | 9 |
| 293 | Effect of N-acetylcysteine on prevention of contrast-associated acute kidney injury in patients with STEMI undergoing primary percutaneous coronary intervention: a systematic review and meta-analysis of randomised controlled trials. 2020 , 10, e039009 | 2 |
| 292 | Nicorandil for CIN Prevention - A Hope or Hype?. 2020 , 21, 1128-1130 | |
| 291 | Contrast-Induced Nephropathy. 2020 , | |
| 290 | Association between contrast-induced nephrotoxicity and contrast enhanced computed tomography followed by endoscopic retrograde cholangiopancreatography. 2020 , 129, 109074 | 0 |
| 289 | CO-based C-arm computed tomography (CACT) of the pelvic arteries: feasibility and diagnostic performance in comparison to CO-angiography in patients with peripheral arterial disease. 2021 , 62, 1707-1715 | |
| 288 | Kidney Disease in the Cardiac Catheterization Laboratory. 2020 , | |
| 287 | Clinical risk factors for the prediction of acute kidney injury post cardiac resynchronization therapy in an elderly population. 2020 , 30, 100594 | |
| 286 | Predictors of Acute Kidney Injury Following Surgical Valve Replacement. 2021 , 69, 396-404 | 0 |
| 285 | Impact of creatinine screening on contrast-induced nephropathy following computerized tomography for stroke. 2021 , 45, 420-425 | 1 |
| 284 | Prevalence and outcome of contrast-induced nephropathy in major trauma patients. 2020 , 1 | 2 |
| 283 | Postangiography Increases in Serum Creatinine and Biomarkers of Injury and Repair. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2020 , 15, 1240-1250 | 6.9 4 |
| 282 | Kontrastmittelinduzierte akute Nierenschädigung [Konsensuspapier der Arbeitsgemeinschaft Herz & Niere der Deutschen Gesellschaft für Kardiologie & Herz- und Kreislaufforschung e. V. und der Deutschen Gesellschaft für Nephrologie e. V.. 2020 , 14, 494-504 | 5 |
| 281 | Trials and tribulations of diagnosing and preventing contrast-induced acute kidney injury. 2021 , 162, 1581-1586 | 2 |

| | | | |
|-----|---|-----|---|
| 280 | Risk factors and incidence of contrast-induced acute kidney injury associated with diagnostic or interventional coronary angiography. 2020 , 9, 21-24 | | |
| 279 | Optical Coherence Tomography-Guided Percutaneous Coronary Intervention With Low-Molecular-Weight Dextran - Effect on Renal Function. 2020 , 84, 917-925 | | 8 |
| 278 | Pre-Procedural Hyperglycemia Increases the Risk of Contrast-Induced Nephropathy in Patients Undergoing Coronary Angiography: A Systematic Review and Meta-Analysis. 2020 , 21, 1377-1385 | | 1 |
| 277 | Acute Kidney Injury after Endovascular Treatment in Patients with Acute Ischemic Stroke. <i>Journal of Clinical Medicine</i> , 2020 , 9, | 5.1 | 1 |
| 276 | The conundrum of contrast-induced acute kidney injury. 2020 , 12, 1721-1727 | | 9 |
| 275 | Contrast-Induced Acute Kidney Injury-Definitions, Epidemiology, and Implications. 2020 , 9, 299-309 | | 8 |
| 274 | Comparison of the preventive efficacy of rosuvastatin versus atorvastatin in post-contrast acute kidney injury in patients with ST-segment elevation myocardial infarction undergoing percutaneous coronary intervention. 2020 , 128, 110336 | | 1 |
| 273 | Predicting Contrast-Induced Renal Complications. 2020 , 9, 321-333 | | |
| 272 | Biomarkers of Contrast-Induced Nephropathy:: Which Ones are Clinically Important?. 2020 , 9, 335-344 | | 6 |
| 271 | Contrast Media-Different Types of Contrast Media, Their History, Chemical Properties, and Relative Nephrotoxicity. 2020 , 9, 279-292 | | 2 |
| 270 | A Practical Approach to Preventing Renal Complications in the Catheterization Laboratory. 2020 , 9, 403-407 | | |
| 269 | Radiocontrast Toxicity. 2020 , 27, 50-55 | | 3 |
| 268 | Dietary restriction for prevention of contrast-induced acute kidney injury in patients undergoing percutaneous coronary angiography: a randomized controlled trial. <i>Scientific Reports</i> , 2020 , 10, 5202 | 4.9 | 8 |
| 267 | Experimental studies on computed tomographic peritoneography. Peritoneal and residual renal function tolerance to iodinated contrast media injected into the peritoneal cavity. 2020 , 33, 163-169 | | 2 |
| 266 | The Relationship Between Contrast Associated Nephropathy and Coronary Collateral Circulation in very Old Patients. 2020 , 56, | | |
| 265 | High White Blood Cell Count Is a Risk Factor for Contrast-Induced Nephropathy following Mechanical Thrombectomy for Acute Ischemic Stroke. 2020 , 10, 59-65 | | 2 |
| 264 | Hydration Strategies for Preventing Contrast-Induced Acute Kidney Injury: A Systematic Review and Bayesian Network Meta-Analysis. 2020 , 2020, 7292675 | | 5 |
| 263 | Subclinical Contrast-Induced Acute Kidney Injury in Patients Undergoing Cerebral Computed Tomography. 2020 , 10, 125-136 | | 6 |

| | | | |
|-----|---|-----|----|
| 262 | DyeVertPLUS EZ System for Preventing Contrast-Induced Acute Kidney Injury in Patients Undergoing Diagnostic Coronary Angiography and/or Percutaneous Coronary Intervention: A UK-Based Cost-Utility Analysis. 2020 , 4, 459-472 | | 2 |
| 261 | Elevated left ventricular end diastolic pressure is associated with increased risk of contrast-induced acute kidney injury in patients undergoing percutaneous coronary intervention. <i>International Journal of Cardiology</i> , 2020 , 306, 196-202 | 3.2 | 2 |
| 260 | Increased circulating galectin-1 levels are associated with the progression of kidney function decline in patients undergoing coronary angiography. <i>Scientific Reports</i> , 2020 , 10, 1435 | 4.9 | 5 |
| 259 | Nicorandil Reduces the Incidence of Contrast-Induced Nephropathy in Patients Undergoing Coronary Angiography/Intervention - Systematic Review and Meta-Analysis of Randomized Controlled Trials Including GRADE Qualification. 2020 , 21, 1121-1127 | | 2 |
| 258 | Risk of Acute Kidney Injury Following Contrast-enhanced CT in Hospitalized Pediatric Patients: A Propensity Score Analysis. 2020 , 294, 548-556 | | 12 |
| 257 | The Association of Fractional Pulse Pressure with Acute Kidney Injury in Patients Undergoing Coronary Intervention due to ST-Segment Elevated Myocardial Infarction. 2020 , 29, 572-579 | | |
| 256 | Variation in contrast-associated acute kidney injury prophylaxis for percutaneous coronary intervention: insights from the Veterans Affairs Clinical Assessment, Reporting, and Tracking (CART) program. 2020 , 21, 150 | | 2 |
| 255 | Contrast-induced acute kidney injury. 2020 , 35, 209-217 | | 24 |
| 254 | Application of 17 Contrast-Induced Acute Kidney Injury Risk Prediction Models. 2020 , 10, 162-174 | | 2 |
| 253 | Incidence and Risk Factors of Postcontrast Acute Kidney Injury in Patients with Acute Ischemic Stroke. 2020 , 2020, 7182826 | | 4 |
| 252 | Serum levels of neutrophil Gelatinase associated Lipocalin (NGAL) predicts hemodialysis after coronary angiography in high risk patients with acute coronary syndrome. 2020 , 21, 143 | | 2 |
| 251 | In-Vitro MPI-guided IVOCT catheter tracking in real time for motion artifact compensation. 2020 , 15, e0230821 | | 4 |
| 250 | Current comments on contrast media administration in patients with renal insufficiency. 2021 , 69, 37-44 | | 2 |
| 249 | Cumulative iodinated contrast exposure for computed tomography during acute kidney injury and major adverse kidney events. 2021 , 31, 3258-3266 | | 1 |
| 248 | Obesity as a Risk Factor for Radiographic Contrast-Induced Nephropathy. <i>Angiology</i> , 2021 , 72, 274-278 | 2.1 | 6 |
| 247 | Refining Safe Contrast Limits for Preventing Acute Kidney Injury After Percutaneous Coronary Intervention. 2021 , 10, e018890 | | 3 |
| 246 | AKI and CIN: Real or Hype?. 2021 , 53-58 | | |
| 245 | Acute kidney injury following endovascular intervention for peripheral artery disease. 2021 , 108, 152-159 | | 1 |

| | | | |
|-----|--|-----|---|
| 244 | Contrast-Induced Nephropathy Post Percutaneous Interventional Procedures. 2021 , 321-326 | | |
| 243 | Remote Ischemic Preconditioning for Prevention of Contrast-Induced Acute Kidney Injury in Patients of CKD Stage III and IV Undergoing Elective Coronary Angiography: A Randomized Controlled Trial. 2021 , 31, 116-123 | | |
| 242 | Impact of Angiotensin-Converting Enzyme Inhibitors/Angiotensin Receptor Blockers on Renal Function in Chronic Kidney Disease Patients Undergoing Coronary Angiography. <i>Cureus</i> , 2021 , 13, e12808 ^{1,2} | | 1 |
| 241 | Lokale und systemische Komplikationen nach perkutanen Interventionen in der Therapie der peripheren arteriellen Verschlusskrankheit. 2021 , 26, 219-224 | | 0 |
| 240 | Alpha-tocopherol and contrast-induced nephropathy: A meta-analysis of randomized controlled trials. 2021 , 91, 188-196 | | 5 |
| 239 | Poor Prognosis of Contrast-Induced Nephropathy during Long Term Follow Up. 2021 , 57, 197-203 | | 0 |
| 238 | Acute Kidney Injury After Repeated Exposure to Contrast Material for Coronary Angiography. 2021 , 5, 46-54 | | 1 |
| 237 | High-Dose Atorvastatin Raises Threshold of Contrast-Induced Nephropathy in Diabetic Patients Undergoing Elective Coronary Intervention: A Randomized Controlled Study. 2021 , 2021, 1-8 | | 0 |
| 236 | Diagnostic accuracy of cardiac testing for coronary artery disease in potential liver transplant recipients: A systematic review and meta-analysis. 2021 , 32, 100714 | | 3 |
| 235 | Post-contrast Acute Kidney Injury Following Emergency Transcatheter Arterial Embolization for Uncontrollable Postpartum Hemorrhage. <i>Angiology</i> , 2021 , 72, 533-538 | 2.1 | |
| 234 | NITRATE-CIN Study: Protocol of a Randomized (1:1) Single-Center, UK, Double-Blind Placebo-Controlled Trial Testing the Effect of Inorganic Nitrate on Contrast-Induced Nephropathy in Patients Undergoing Coronary Angiography for Acute Coronary Syndromes. 2021 , 26, 303-309 | | 2 |
| 233 | Trimetazidine reduces contrast-induced nephropathy in patients with renal insufficiency undergoing coronary angiography and angioplasty: A systematic review and meta-analysis (PRISMA). 2021 , 100, e24603 | | 0 |
| 232 | A Novel Risk Assessment Model Using Urinary System Contrast Blush Grading to Predict Contrast-Induced Acute Kidney Injury in Low-Risk Profile Patients. <i>Angiology</i> , 2021 , 72, 524-532 | 2.1 | 3 |
| 231 | Impact of radial access on contrast-induced acute kidney injury in patients with coronary artery bypass grafts. 2021 , | | |
| 230 | Contrast-Induced Nephropathy: A Review of Mechanisms and Risks. <i>Cureus</i> , 2021 , 13, e14842 | 1.2 | 5 |
| 229 | Should Percutaneous Coronary Intervention be the Standard Treatment Strategy for Significant Coronary Artery Disease in all Octogenarians?. 2021 , 17, 244-259 | | 1 |
| 228 | Therapy with lopinavir/ritonavir and hydroxychloroquine is associated with acute kidney injury in COVID-19 patients. 2021 , 16, e0249760 | | 4 |
| 227 | The Relationship Between HbA1c and Contrast-Induced Nephropathy in Patients with Non-ST Elevation Myocardial Infarction and Non- Established Diabetes Mellitus. | | |

| | | | |
|-----|--|-----|---|
| 226 | RenalGuard system and conventional hydration for preventing contrast-associated acute kidney injury in patients undergoing cardiac interventional procedures: A systematic review and meta-analysis. <i>International Journal of Cardiology</i> , 2021 , 333, 83-89 | 3.2 | 3 |
| 225 | Disease a month: Update and review of contrast-associated acute kidney injury. 2021 , 101272 | | |
| 224 | Significance of serum FGF-23 for risk assessment of contrast-associated acute kidney injury and clinical outcomes in patients undergoing coronary angiography. 2021 , 16, e0254835 | | 2 |
| 223 | Tailored Versus Standard Hydration to Prevent Acute Kidney Injury After Percutaneous Coronary Intervention: Network Meta-Analysis. 2021 , 10, e021342 | | 2 |
| 222 | Systemic Immune-Inflammation Index May Predict the Development of Contrast-Induced Nephropathy in Patients With ST-Segment Elevation Myocardial Infarction. <i>Angiology</i> , 2021 , 33197211030053 | 2.1 | 7 |
| 221 | Factors Contributing to an Efficacious Endovascular Treatment for Acute Ischemic Stroke in Asian Population. 2021 , 16, 91-110 | | 2 |
| 220 | Acute Kidney Injury Following Percutaneous Coronary Intervention for Acute Coronary Syndrome: Incidence, Aetiology, Risk Factors and Outcomes. <i>Angiology</i> , 2021 , 33197211040375 | 2.1 | 2 |
| 219 | Biomarkers for the early prediction of contrast-induced nephropathy after percutaneous coronary intervention in adults: A systematic review and meta-analysis. <i>Angiology</i> , 2021 , 33197211039921 | 2.1 | 1 |
| 218 | One-year Outcome of Different Unprotected Left Main Percutaneous Coronary Interventions Techniques in Acute Coronary Syndromes. 2020 , 9, 878-884 | | |
| 217 | Risk of Worsening Renal Function Following Repeated Exposures to Contrast Media During Percutaneous Coronary Interventions. 2021 , 10, e021473 | | 0 |
| 216 | Does intravenous contrast medium administration result in altered renal biomarkers? A study in clinically stable cats with and without azotemia. 2021 , 1098612X211038535 | | |
| 215 | A Comparison Between Two Definitions of Contrast-Associated Acute Kidney Injury for Long-Term Mortality in Elderly and Non-elderly Patients After Elective Percutaneous Coronary Intervention. 2021 , 8, 720857 | | 1 |
| 214 | Magnitude and associated factors of contrast induced nephropathy among patients undergoing coronary angiography and interventions at a cardiac referral hospital in Tanzania - a cross-sectional study. 2021 , 38, 311 | | 1 |
| 213 | Reducing the Risk of Contrast Medium Induced Nephropathy. 2006 , 35-45 | | 2 |
| 212 | Iatrogenic Injury. 2005 , 351-439 | | 7 |
| 211 | Hypoxia, Oxidative Stress, and the Pathophysiology of Contrast-Media-Induced Nephropathy. 2011 , 229-256 | | 3 |
| 210 | Renal Function Monitoring. 2010 , 1443-1475 | | 1 |
| 209 | What Is the Best Means of Preventing Perioperative Renal Injury?. 2009 , 226-239 | | 1 |

| | | |
|-----|--|-----|
| 208 | Contrast Nephropathy. 2008 , 41-46 | 1 |
| 207 | DDAH-2 alleviates contrast medium iopromide-induced acute kidney injury through nitric oxide synthase. 2019 , 133, 2361-2378 | 10 |
| 206 | Coronary anatomy, anatomic variations and anomalies: a retrospective coronary angiography study. 2015 , 56, 339-45 | 14 |
| 205 | Renal Safety of Iodinated Contrast Media Depending on Their Osmolarity - Current Outlooks. 2016 , 81, 157-65 | 20 |
| 204 | Prevention of contrast-induced acute kidney injury: is simple oral hydration similar to intravenous? A systematic review of the evidence. 2013 , 8, e60009 | 54 |
| 203 | Sildenafil citrate for prophylaxis of nephropathy in an animal model of contrast-induced acute kidney injury. 2014 , 9, e113598 | 36 |
| 202 | Randomly and Non-Randomly Missing Renal Function Data in the Strong Heart Study: A Comparison of Imputation Methods. 2015 , 10, e0138923 | 6 |
| 201 | A web-based tool to predict acute kidney injury in patients with ST-elevation myocardial infarction: Development, internal validation and comparison. 2017 , 12, e0181658 | 8 |
| 200 | Contrast-induced nephropathy in patients with diabetes mellitus between iso- and low-osmolar contrast media: A meta-analysis of full-text prospective, randomized controlled trials. 2018 , 13, e0194330 | 14 |
| 199 | Incidence of contrast-induced acute kidney injury (CI-AKI) in high-risk oncology patients undergoing contrast-enhanced CT with a reduced dose of the iso-osmolar iodinated contrast medium iodixanol. 2020 , 15, e0233433 | 6 |
| 198 | Practical Risk Stratification Score for Prediction of Contrast-Induced Nephropathy After Primary Percutaneous Coronary Intervention in Patients With Acute ST-Segment Elevation Myocardial Infarction. 2019 , 10, 350-357 | 8 |
| 197 | Contrast induced nephropathy among patients with normal renal function undergoing coronary angiography. 2016 , 5, 21-4 | 10 |
| 196 | Incidence of contrast-induced nephropathy a prospective study. 2017 , 6, 192-198 | 4 |
| 195 | Targeted Renal Therapy and Contrast-Induced Nephropathy During Endovascular Abdominal Aortic Aneurysm Repair:Results of a Feasibility Pilot Trial. 2007 , 14, 520-527 | 7 |
| 194 | N-acetylcysteine in the prevention of contrast agent-induced nephrotoxicity in patients undergoing computed tomography studies. 2006 , 3, 773-777 | 1 |
| 193 | Contrast nephropathy. 2000 , 11, 177-182 | 286 |
| 192 | Toward proteomics in uroscopy: urinary protein profiles after radiocontrast medium administration. 2001 , 12, 1026-1035 | 75 |
| 191 | Akut koroner sendromda kontrast nefropati geliimi ile Syntax Skoru arasındaki ilişi. 2019 , 44, 1181-1188 | 1 |

| | | |
|-----|---|----|
| 190 | The association between post-procedural oral hydration and risk of contrast-induced acute kidney injury among ST-elevation myocardial infarction patients undergoing primary percutaneous coronary intervention. 2019 , 7, 321 | 3 |
| 189 | Contrast-induced nephropathy in interventional cardiology. 2011 , 4, 85-99 | 21 |
| 188 | Pathophysiology of Contrast-Induced Acute Kidney Injury (CIAKI). 2019 , 25, 4642-4647 | 12 |
| 187 | Acute Kidney Injury Post Cardiac Catheterization: Does Vascular Access Route Matter?. 2019 , 15, 96-101 | 1 |
| 186 | Preventing Cardio-renal Syndrome Rather than Treating It: Could Statins Play A Role?. 2011 , 5, 226-30 | 3 |
| 185 | Direct coronary stenting in reducing radiation and radiocontrast consumption. 2010 , 44, 153-7 | 7 |
| 184 | COMPARISON OF DIFFERENT AGENTS ON RADIOGRAPHIC CONTRAST AGENT INDUCED NEPHROPATHY. 2008 , 5, | 3 |
| 183 | Contrast medium-induced nephropathy. Aspects on incidence, consequences, risk factors and prevention. 2007 , 2, 118-124 | 2 |
| 182 | Acetylcysteine and Fenoldopam. 2003 , 23, 39-46 | 7 |
| 181 | N-acetylcysteine for Prevention of Contrast-Induced Nephropathy: A Narrative Review. 2011 , 41, 695-702 | 9 |
| 180 | The role of neutrophil-gelatinase-associated lipocalin in early diagnosis of contrast nephropathy. 2015 , 25, 292-6 | 11 |
| 179 | N-acetyl cysteine versus allopurinol in the prevention of contrast nephropathy in patients with chronic kidney disease: A randomized controlled trial. 2017 , 27, 93-98 | 4 |
| 178 | Effect of N-acetyl cysteine in prevention of contrast nephropathy on patients under intravenous pyelography and contrast CT. 2012 , 1, 28 | 2 |
| 177 | Contrast-induced nephropathy in urological imaging: A comparison with cardiology interventions. 2017 , 33, 241-245 | 1 |
| 176 | Impact of platelet glycoprotein IIb/IIIa receptor inhibitors on renal function in patients with ST-segment elevation myocardial infarction treated with primary or rescue percutaneous coronary intervention. 2009 , 5, 604-9 | 2 |
| 175 | Predictors of contrast-induced nephropathy in chronic total occlusion percutaneous coronary intervention. 2014 , 9, 1173-80 | 24 |
| 174 | Carbon dioxide-angiography for patients with peripheral arterial disease at risk of contrast-induced nephropathy. 2020 , 12, 76-90 | 5 |
| 173 | The risk of renal function deterioration in abdominal aortic stent graft patients with and without previous kidney function failure - an analysis of risk factors. 2020 , 85, e643-e649 | 2 |

| | | |
|-----|--|---|
| 172 | Predictive factors for contrast-induced acute kidney injury in high-risk patients given N-acetylcysteine prophylaxis. 2018 , 38, 269-276 | 2 |
| 171 | "Contrast nephropathy" in renal transplantation: Is it real?. 2016 , 6, 682-688 | 4 |
| 170 | Preventive Effect of Atorvastatin (80 mg) on Contrast-Induced Nephropathy After Angiography in High-Risk Patients: Double-Blind Randomized Clinical Trial. 2016 , 8, e29574 | 8 |
| 169 | Risk of acute kidney injury requiring hemodialysis after contrast-enhanced imaging after traumatic injury: A National Trauma Databank analysis. 2021 , | 0 |
| 168 | Computed Tomography Angiography (CTA) before Reconstructive Jaw Surgery Using Fibula Free Flap: Retrospective Analysis of Vascular Architecture. 2021 , 11, | 1 |
| 167 | Urinary proteomics investigations into contrast-induced acute kidney injury. 2021 , 16, e0258736 | 0 |
| 166 | Impact of contrast-induced acute kidney injury on long-term major adverse cardiovascular events and kidney function after percutaneous coronary intervention: insights from a territory-wide cohort study in Hong Kong.. 2022 , 15, 338-346 | 0 |
| 165 | The Kidney in Sepsis. 2000 , 91-101 | |
| 164 | X-Ray Coronary Angiography. 2002 , 602-635 | |
| 163 | COMPLICATIONS OF PERCUTANEOUS CORONARY INTERVENTIONS. 2004 , 162-193 | |
| 162 | Who should not go to the cathlab?. 2004 , 1-7 | |
| 161 | Contrast agents for preventing contrast-induced nephropathy. | |
| 160 | Early versus Late Complications. 2005 , 92-112 | |
| 159 | How can we reduce the incidence of contrast-induced acute renal failure?. 2005 , 9, 58-60 | |
| 158 | Systemic Complications of Endovascular Therapy. 2005 , 55-69 | |
| 157 | Epidemiology and predictors of contrast-induced nephropathy. 2005 , 19-33 | |
| 156 | Pharmacologic prophylaxis. 2005 , 121-132 | |
| 155 | Renal replacement therapies. 2005 , 133-148 | |

- 154 Clinical features and prognostic implications of contrast-induced nephropathy. **2005**, 35-45
- 153 Chronic Kidney Disease. **2006**, 463-500
- 152 Chronic total occlusion. **2006**, 199-209
- 151 Renal complications after percutaneous intervention. **2006**, 255-265
- 150 Optimal patient preparation and selection to avoid complications. **2006**, 17-37
- 149 Continuous renal replacement modalities for acute kidney failure in critically ill patients.
- 148 Contrast medium-induced nephropathy. Aspects on incidence, consequences, risk factors and prevention. **2007**, 2, 118-24 2
- 147 The kidney and percutaneous coronary interventions. **2007**, 9-26
- 146 Contrast-induced nephropathy after percutaneous coronary interventions. **2007**, 493-501
- 145 Akutes Nierenversagen. **2008**, 543-562
- 144 Contrast Media Safety and Managing At-Risk Patients: Update 2008. **2008**, 25-30
- 143 The Role of Extracorporeal Blood Purification Therapies in the Prevention of Radiocontrast Agent-Induced Nephropathy. **2009**, 1000-1006
- 142 Contrast Nephropathy and Its Prevention. **2009**, 1807-1812
- 141 Nephrotoxicity of Contrast Media. **2009**, 280-284
- 140 Risk Factors for Nosocomial Renal Failure. **2009**, 91-95
- 139 Drug-Induced Acute Renal Failure. **2009**, 317-324 1
- 138 Intermittent Hemofiltration for Management of Fluid Overload and Administration of Contrast Media. **2009**, 1276-1281
- 137 Interventional Cardiology. **2010**, 847-866

136 Systemic Complications. **2010**, 627-642

135 Renovascular Disease. **2010**, 2216-2225

134 The Epidemiology of Acute Kidney Injury. **2010**, 643-653

133 Contrast agents. **2010**, 28-37

132 Contrast agents for preventing contrast-induced nephropathy.

131 Understanding Endovascular Aortic Surgery: Endovascular Repair of Aortic Aneurysmal Disease. **2011**, 257-279

130 Nierenerkrankungen und Hypertonie. **2011**, 1423-1436

129 Dialyse auf der Intensivstation. **2011**, 215-226

128 The cardiac catheterization patient. **2011**, 570-574

127 Contrast-Induced Nephropathy. **2011**, 909-912

126 Dietary Approaches to Kidney Diseases. **2011**, 2170-2204

2

125 Venous Thromboembolism in Medical-Surgical Critically Ill Patients. **2011**, 1145-1149

124 Continuous renal replacement modalities for acute kidney failure in critically ill patients.

123 Use of trimetazidine in preventing contrast-induced nephropathy: a nephrologist's perspective. **2011**, 53, 264-266

122 Renal Insufficiency and the Impact of Contrast Agents. 464-475

121 Imaging Techniques in Desmoid Tumors. **2012**, 47-75

120 Contrast-induced nephropathy. **2012**, 101-105

119 Contrast-Induced Acute Kidney Injury and the Role of Chronic Kidney Disease in PCI. **2012**, 66-74

1

- 118 Magnetic Particle Imaging for Angiography, Stem Cell Tracking, Cancer Imaging and Inflammation Imaging. **2012**, 523-540 1
- 117 Evaluation of a Dedicated Everolimus Coated Side Branch Access (SBA) System. 136-144
- 116 Contrast-induced nephropathy in cardiology practice: Risk factors and approaches to prevention. **2012**, 18, 222-227
- 115 CTO. 279-287
- 114 MR Enhancing Implants. **2014**, 725-737
- 113 Computed Tomography Angiography before Intravenous Thrombolysis Does Not Increase the Risk of Renal Dysfunction. **2013**, 2013, 1-6
- 112 Guideline for Interventional Recanalization of Lower Extremity Artery. **2014**, 71, 205 2
- 111 Dialyse auf der Intensivstation. **2014**, 225-236
- 110 Posttraumatic acute kidney injury and effect of the use of contrast media in elderly patients. **2014**, 25, 201-207
- 109 Contrast Media-Induced Nephropathy and Nephrogenic Systemic Fibrosis. **2014**, 825-837
- 108 Contrast-Induced Nephropathy Post Percutaneous Interventional Procedures. **2014**, 299-303
- 107 Determination of Risk Factors Regarding the Administration of Contrast Media. **1999**, 96-117
- 106 Contrast Agents and Contrast-Induced Nephropathy. **2015**, 06, 451-457
- 105 Contrast-Induced Nephropathy. **2015**, 88, 375 2
- 104 RISK OF CONTRAST INDUCED NEPHROPATHY WITH INTRAVENOUS ADMINISTRATION. **2015**, 4, 12846-12851
- 103 Contrast-Enhanced Magnetic Resonance Angiography for Preoperative Imaging in DIEP Flap Breast Reconstruction. **2016**, 163-170 1
- 102 Prevention Strategies of Contrast Medium Induced Nephropathy (CIN): A Review of the Current Literature. **2016**, 06, 98-110 1
- 101 INCIDENCE AND MAIN DETERMINANTS OF CONTRAST-INDUCED NEPHROPATHY FOLLOWING CORONARY ANGIOGRAPHY OR SUBSEQUENT BALLOON ANGIOPLASTY. **2016**, 3, 4459-4462

| | | | |
|-----|---|-----|---|
| 100 | The Risk Factors for Developing Contrast-induced Nephropathy after the Evaluation of Trauma Patients at a Regional Trauma Center in Korea. 2016 , 29, 124-128 | | |
| 99 | Contrast Induced Nephropathy - A Review. 001-003 | | |
| 98 | Risk Factors and Pathophysiology of Contrast- Induced Nephropathy after Contrast Enhanced Multidetector Computed Tomography. 2017 , 103-115 | | 1 |
| 97 | CONTRAST INDUCED NEPHROPATHY IN PATIENTS UNDERGOING ELECTIVE PCI IN A TERTIARY CARE CENTRE IN COASTAL KERALA. 2017 , 6, 5128-5131 | | |
| 96 | Nierenerkrankungen und Hypertonie in der Neurologie. 2018 , 1-20 | | |
| 95 | Contrast-induced Nephropathy Following Percutaneous Coronary Intervention at a Tertiary Cardiac Center in Nepal. <i>Cureus</i> , 2018 , 10, e3331 | 1.2 | 1 |
| 94 | [Impact of rotational coronary angiography in the amount of iodinated contrast and the exposure to ionising radiations in patients undergoing invasive coronary procedures: A systematic review and meta-analysis]. 2018 , 88, 277-286 | | |
| 93 | Pediatric Interventional Cardiovascular Magnetic Resonance. 2019 , 554-567.e5 | | |
| 92 | A Pilot Study of Prediction of Creatinine Clearance by Ellipsoid Volumetry of Kidney Using Noncontrast Computed Tomography. 2019 , 2, 60-66 | | 2 |
| 91 | Perkaban Koroner Girişim Yapılan Koroner Yavaş Akım Olgularında, Serum Urotensin II Düzeyi ve Kontrast Kaynaklı Nefropati Gelişimi ile Bir Yıllık Klinik Takip Bulgular-Arasındaki İlişki. 2019 , 9, 442-454 | | |
| 90 | Aqueous Extract of Sorindeia Juglandifolia Leaves Protects Methotrexate-Induced Liver and Kidney Damage in Rat. 2019 , 4, 127-140 | | 1 |
| 89 | DyeVert PLUS EZ system for Preventing Contrast-Induced Acute Kidney Injury in Patients Undergoing Diagnostic Coronary Angiography and/or Percutaneous Coronary Intervention: A UK-Based Cost-Utility Analysis. | | |
| 88 | Intravascular Iodinated Contrast Is an Independent Cause of Acute Kidney Injury Following Coronary Angiography. 2019 , 112, 541-546 | | 1 |
| 87 | The relationship between coronary collateral circulation and contrast induced nephropathy in patients with non-ST elevation myocardial infarction. 2019 , 178, | | |
| 86 | Biomarkers of Acute Kidney Injury and Scope of Utilization in the Cardiac Catheterization Laboratory. 2020 , 429-448 | | |
| 85 | Predictive Value of Lymphocyte-to-monocyte Ratio in Patients with Contrast-induced Nephropathy After Percutaneous Coronary Intervention for Acute Coronary Syndrome. 2021 , 9, 123-130 | | 2 |
| 84 | Predictive value of plasma volume status for contrast-induced nephropathy in patients with heart failure undergoing PCI. 2021 , | | 1 |
| 83 | Albuminuria Pre-emptively Identifies Cardiac Patients at Risk of Contrast-Induced Nephropathy. <i>Journal of Clinical Medicine</i> , 2021 , 10, | 5.1 | 0 |

| | | |
|----|---|----|
| 82 | Contrast-Induced Acute Kidney Injury: Epidemiology, Risk Stratification, and Prognosis. 2020 , 183-207 | |
| 81 | Angiotensin-Converting Enzyme Inhibitors and Contrast-Associated Acute Kidney Injury After Coronary Angiography and Intervention. 2021 , 21, 487-497 | 0 |
| 80 | [Contrast medium-induced acute kidney injury-Consensus paper of the working group "Heart and Kidney" of the German Cardiac Society and the German Society of Nephrology]. 2021 , 62, 111-120 | 1 |
| 79 | Impact of Geriatric Nutritional Index in Contrast-Induced Nephropathy Developed in Patients with Non-ST Segment Elevation Myocardial Infarction who Underwent Percutaneous Coronary Intervention. 2020 , 35, 47-54 | |
| 78 | The Economic Impact of Kidney Disease in the Cardiac Catheterization Laboratory. 2020 , 409-427 | |
| 77 | Contrast-Induced Nephropathy After Peripheral Vascular Intervention. 2020 , 313-333 | |
| 76 | Nierenerkrankungen und Hypertonie in der Neurologie. 2020 , 1763-1781 | |
| 75 | Problematiche relative all'impiego dei mezzi di contrasto nell'anziano. 2006 , 21-28 | |
| 74 | Ischémie/reperfusion, stress oxydant, pręconditionnement et insuffisance rñale aiguę 2007 , 85-105 | |
| 73 | Requests for Imaging Using Contrast Agents: What Information Must be Provided. 2009 , 11-14 | |
| 72 | Contrast-Induced Nephropathy. 2009 , 275-280 | |
| 71 | Contrast-induced acute kidney injury: specialty-specific protocols for interventional radiology, diagnostic computed tomography radiology, and interventional cardiology. 2009 , 84, 170-9 | 36 |
| 70 | Acute renal failure. 2008 , 2008, | 1 |
| 69 | Influence of the timing of cardiac catheterization and amount of contrast media on acute renal failure after cardiac surgery. 2011 , 16, 502-8 | 6 |
| 68 | Acute kidney injury. 2011 , 2011, | 6 |
| 67 | Effect of renin-angiotensin-aldosterone system blockade therapy on incidence of contrast-induced nephropathy in patients with chronic kidney disease. 2012 , 6, 432-6 | 7 |
| 66 | Decreased GFR estimated by MDRD or Cockcroft-Gault equation predicts incident CVD: the strong heart study. 2009 , 22, 373-80 | 19 |
| 65 | Atorvastatin and prevention of contrast induced nephropathy following coronary angiography. 2015 , 20, 1-6 | 9 |

| | | |
|----|--|-------|
| 64 | Additional furosemide treatment beyond saline hydration for the prevention of contrast-induced nephropathy: a meta-analysis of randomized controlled trials. 2015 , 8, 387-94 | 6 |
| 63 | Comparison of the efficacy of recombinant human brain natriuretic peptide with saline hydration in preventing contrast-induced nephropathy in patients undergoing coronary angiography with or without concomitant percutaneous coronary intervention. 2015 , 8, 14166-72 | 4 |
| 62 | Cause of Emergency Department Mortality; a Case-control Study. 2014 , 2, 30-5 | 5 |
| 61 | The Current Status of Performing Left Ventriculography in Taiwan. 2016 , 32, 49-54 | 1 |
| 60 | Prevention of contrast induced nephropathy; a cardiology point of view. 2015 , 4, 23-25 | 1 |
| 59 | Pentoxifylline and prevention of contrast-induced nephropathy: Is it efficient in patients with myocardial infarction undergoing coronary angioplasty?. 2016 , 12, 238-242 | 4 |
| 58 | Nephroprotective Effects of L-Carnitine against Contrast-Induced Nephropathy in Patients Undergoing Percutaneous Coronary Intervention: A Randomized Open-Labeled Clinical Trial. 2017 , 12, 57-64 | |
| 57 | Contrast Utilization During Chronic Total Occlusion Percutaneous Coronary Intervention: Insights From a Contemporary Multicenter Registry. 2016 , 28, 288-94 | 12 |
| 56 | A first attempt of inferior vena cava filter successfully guided by a mixed-reality system: a case report. 2019 , 16, 575-577 | 3 |
| 55 | Impact of One-Catheter Strategy with TIG I Catheter on Coronary Catheterization Performance and Economic Costs. 2019 , 113, 960-968 | 0 |
| 54 | Comparing Trimetazidine with Allopurinol in Prevention of Contrast Induced Nephropathy After Coronary Angiography. 2020 , 32, 451-455 | |
| 53 | Pericoronary fat attenuation index in computed tomography angiography is associated with mortality in end-stage renal disease. 2021 , | 0 |
| 52 | 2021 ACC/AHA/SCAI Guideline for Coronary Artery Revascularization: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines.. 2021 , | 42 |
| 51 | Comparing Trimetazidine with Allopurinol in Prevention of Contrast Induced Nephropathy After Coronary Angiography. 2020 , 32, 451-455 | 0 |
| 50 | Effect of Oral Hydration on Contrast-Induced Acute Kidney Injury among Patients after Primary Percutaneous Coronary Intervention. 2021 , 1-9 | 0 |
| 49 | PREDICTIVE VALUE OF C-REACTIVE PROTEIN/ALBUMIN RATIO IN THE DEVELOPMENT OF CONTRAST-INDUCED NEPHROPATHY IN PATIENTS WITH ACUTE ISCHEMIC STROKE TREATED PERCUTANEOUSLY. <i>ÖzÜniversitesi Sađık Hizmetleri Meslek Yüksek Okulu Dergisi</i> , | 0.2 |
| 48 | Statin Use Influence on the Occurrence of Acute Kidney Injury in Patients with Peripheral Arterial Disease.. <i>Journal of Atherosclerosis and Thrombosis</i> , 2022 , | 4 0 |
| 47 | Association of the Novel Inflammatory Marker Systemic Immune-Inflammation index and Contrast-Induced Nephropathy in Patients Undergoing Transcatheter Aortic Valve Replacement for Severe Aortic Stenosis.. <i>Angiology</i> , 2022 , 33197211045031 | 2.1 3 |

| | | | |
|----|--|------|----|
| 46 | 2021 ACC/AHA/SCAI Guideline for Coronary Artery Revascularization: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. <i>Circulation</i> , 2021 , CIR0000000000001038 | 16.7 | 23 |
| 45 | Recommendations in pre-procedural imaging assessment for TAVI intervention: SIC-SIRM position paper part 2 (CT and MR angiography, standard medical reporting, future perspectives).. <i>Radiologia Medica</i> , 2022 , 1 | 6.5 | 2 |
| 44 | The influence of timing of coronary angiography on acute kidney injury in out-of-hospital cardiac arrest patients: a retrospective cohort study.. <i>Annals of Intensive Care</i> , 2022 , 12, 12 | 8.9 | |
| 43 | Renal impairment and the prognosis of endovascular thrombectomy: a meta-analysis and systematic review. <i>Therapeutic Advances in Neurological Disorders</i> , 2022 , 15, 175628642210836 | 6.6 | |
| 42 | Acute Kidney Injury Following Transcatheter Aortic Valve Implantation: Association with Contrast Media Dosage and Contrast Media Based Risk Predication Models.. <i>Journal of Clinical Medicine</i> , 2022 , 11, | 5.1 | 0 |
| 41 | The predictive value of the prognostic nutritional index for contrast-induced nephropathy. <i>Journal of Clinical Medicine of Kazakhstan</i> , 2022 , 19, 46-49 | 0.2 | 0 |
| 40 | Mehran risk score model for predicting contrast-induced nephropathy after cardiac resynchronization therapy in patients with heart failure. <i>Gulhane Medical Journal</i> , 2022 , 64, 40-46 | | |
| 39 | Admission electrolyte and osmotic pressure levels are associated with the incidence of contrast-associated acute kidney injury.. <i>Scientific Reports</i> , 2022 , 12, 4714 | 4.9 | |
| 38 | Long-Term Clinical Impact of Contrast-Associated Acute Kidney Injury Following PCI: An ADAPT-DES Substudy.. <i>JACC: Cardiovascular Interventions</i> , 2022 , | 5 | 3 |
| 37 | Hyperglycemia on Admission Predicts Acute Kidney Failure and Renal Functional Recovery among Inpatients.. <i>Journal of Clinical Medicine</i> , 2021 , 11, | 5.1 | 3 |
| 36 | Contrast-Associated Acute Kidney Injury After Endovascular Therapy for Acute Ischemic Stroke: A Meta-Analysis. | | 0 |
| 35 | Contrast-Associated Acute Kidney Injury.. <i>Journal of Clinical Medicine</i> , 2022 , 11, | 5.1 | 1 |
| 34 | Periprocedural planning for neuroendovascular procedures. 163-179 | | |
| 33 | A Novel Predictor of Contrast-Induced Nephropathy in Patients With Carotid Artery Disease; the Systemic Immune Inflammation Index.. <i>Angiology</i> , 2022 , 33197211061919 | 2.1 | 3 |
| 32 | Clinical impact of noncontrast percutaneous coronary intervention in patients with acute coronary syndrome.. <i>Journal of Medical Investigation</i> , 2022 , 69, 57-64 | 1.2 | |
| 31 | Low-Osmolar vs. Iso-Osmolar Contrast Media on the Risk of Contrast-Induced Acute Kidney Injury: A Propensity Score Matched Study.. <i>Frontiers in Medicine</i> , 2022 , 9, 862023 | 4.9 | |
| 30 | Optimizing the Outcomes of Percutaneous Coronary Intervention in Patients with Chronic Kidney Disease.. <i>Journal of Clinical Medicine</i> , 2022 , 11, | 5.1 | 1 |
| 29 | Evaluating the Effectiveness of Pretreatment With Intravenous Fluid in Reducing the Risk of Developing Contrast-Induced Nephropathy: A Systematic Review and Meta-Analysis. <i>Cureus</i> , 2022 , | 1.2 | 0 |

28 Contrast-medium-induced nephrotoxicity. 18-24

27 Impact of chronic total occlusion in a non-infarct-related coronary artery on contrast-associated nephropathy in acute ST-elevation myocardial infarction. *Acta Cardiologica*, 1-6 0.9 1

26 Percutaneous transluminal renal artery stenting using digital subtraction angiography with diluted contrast medium in a patient with severe chronic kidney disease. *Journal of Cardiology Cases*, 2022, 0.6

25 Reduction of contrast agent volume utilization for cardiac catheterization in current clinical practice. *International Journal of Cardiology*, 2022, 3.2

24 Radiographic Contrast Media and the Kidney. *Clinical Journal of the American Society of Nephrology: CJASN*, CJN.16311221 6.9 1

23 Ultra-low CONTRAST PCI vs conventional PCI in patients of ACS with increased risk of CI-AKI (CONSAVE-AKI). 2022, 1

22 When Prevention is Truly Better than Cure: Contrast-Associated Acute Kidney Injury in Percutaneous Coronary Intervention. 2022, 18, 73-85 0

21 The target vessel SYNTAX score: A novel pre-procedural predictor for contrast-induced acute kidney injury after rotational atherectomy. 2022, 0

20 Progress on the Timing of Early Initiation of Continuous Renal Replacement Therapy for Acute Kidney Injury after Cardiac Surgery. 2022, 12, 9125-9131 0

19 Incidence, predictors and clinical implications of new renal impairment following percutaneous coronary intervention. 2022, 9, e001876 0

18 C-Reactive Protein to Albumin Ratio as a Predictor of Contrast-Induced Nephropathy After Carotid Angiography. 000331972211359 1

17 Iodinated Contrast and Acute Kidney Injury. 2022, 145-162 0

16 Evaluating the Image Quality of Neck Structures Scanned on Chest CT with Low-Concentration-Iodine Contrast Media. 2022, 8, 2854-2863 0

15 Implementation of an Electronic Health Records-Based Safe Contrast Limit for Preventing Contrast-Associated Acute Kidney Injury After Percutaneous Coronary Intervention. 0

14 Contrast-induced acute kidney injury and its contemporary prevention. 9, 0

13 Risk Factors and Clinical Outcomes for Contrast-induced Nephropathy After Percutaneous Coronary Intervention in Patients with Normal Serum Creatinine. 2010, 39, 374-380 3

12 Assessment of Vascular and Valvular Calcification Improves Screening for Coronary Artery Disease Before Liver Transplantation. 2023, 191, 23-31 0

11 Quality and safety in the literature: January 2023. 2023, 32, 56-60 0

- 10 Coronary Angiography. **2013**, 37-59 ○
- 9 Kidney Function, Male Gender, and Aneurysm Diameter Are Predictors of Acute Kidney Injury in Patients with Abdominal Aortic Aneurysms Treated Endovascularly. **2023**, 15, 130 ○
- 8 Differential renal proteomics analysis in a novel rat model of iodinated contrast-induced acute kidney injury. **2023**, 45, ○
- 7 A Novel Preoperative Risk Assessment Tool to Identify Patients at Risk of Contrast-Associated Acute Kidney Injury After Endovascular Abdominal Aortic Aneurysm Repair. **2023**, ○
- 6 Interleukin-18 and Gelsolin Are Associated with Acute Kidney Disease after Cardiac Catheterization. **2023**, 13, 487 ○
- 5 The Combined Treatment of Glutathione Sodium Salt and Ascorbic Acid for Preventing Contrast-Associated Acute Kidney Injury in ST-Elevation Myocardial Infarction Patients Undergoing Primary PCI: A Hypothesis to Be Validated. **2023**, 12, 773 ○
- 4 What Is the Best Means of Preventing Perioperative Renal Injury?. **2013**, 233-248 ○
- 3 Procedural and Technological Innovations Facilitating Ultra-low Contrast Percutaneous Coronary Interventions. 18, ○
- 2 Cardiac Catheterization and Coronary Arteriography. **2023**, 237-266 ○
- 1 Association between dose reduction of renin-angiotensin-aldosterone system inhibitors before coronary artery angiography and acute kidney injury: a propensity score-matched study. 1-12 ○