

# Additional modules for versatile and economical PCR-b modification in *Saccharomyces cerevisiae*

DOI: 10.1002/(sici)1097-0061(199807)14:10<953::aid-yea293>3.0.co;2-u

Citation Report

| #  | ARTICLE   | IF   | CITATIONS |
|----|---|------|-----------|
| 1  | Heterologous modules for efficient and versatile PCR-based gene targeting in <i>Schizosaccharomyces pombe</i> . <i>Yeast</i> , 1998, 14, 943-951.   | 0.8  | 2,105     |
| 2  | Role of the Yeast Gin4p Protein Kinase in Septin Assembly and the Relationship between Septin Assembly and Septin Function. <i>Journal of Cell Biology</i> , 1998, 143, 719-736.  | 2.3  | 207       |
| 3  | Involvement of an Actomyosin Contractile Ring in <i>Saccharomyces cerevisiae</i> Cytokinesis. <i>Journal of Cell Biology</i> , 1998, 142, 1301-1312.  | 2.3  | 372       |
| 4  | The Src Homology Domain 3 (SH3) of a Yeast Type I Myosin, Myo5p, Binds to Verprolin and Is Required for Targeting to Sites of Actin Polarization. <i>Journal of Cell Biology</i> , 1998, 141, 1357-1370.                            | 2.3  | 129       |
| 5  | Chapter 19 Using Green Fluorescent Protein Fusion Proteins to Quantitate Microtubule and Spindle Dynamics in Budding Yeast. <i>Methods in Cell Biology</i> , 1998, 61, 369-383.   | 0.5  | 18        |
| 6  | Distinct Regions of the Swi5 and Ace2 Transcription Factors Are Required for Specific Gene Activation. <i>Journal of Biological Chemistry</i> , 1999, 274, 21029-21036.   | 1.6  | 79        |
| 7  | Ubiquitin-dependent degradation of multiple F-box proteins by an autocatalytic mechanism. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 9124-9129.                             | 3.3  | 248       |
| 8  | Mammalian Cdk5 is a functional homologue of the budding yeast Pho85 cyclin-dependent protein kinase. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 14445-14450.                | 3.3  | 67        |
| 9  | Formation of AP-3 transport intermediates requires Vps41 function. <i>Nature Cell Biology</i> , 1999, 1, 346-353.   | 4.6  | 122       |
| 10 | The TOR signalling pathway controls nuclear localization of nutrient-regulated transcription factors. <i>Nature</i> , 1999, 402, 689-692.   | 13.7 | 883       |
| 11 | Chromosomal promoter replacement in <i>Saccharomyces cerevisiae</i> : construction of conditional lethal strains for the cloning of glycosyltransferases from various organisms. <i>Glycoconjugate Journal</i> , 1999, 16, 673-679. | 1.4  | 5         |
| 12 | Heterologous URA3MX cassettes for gene replacement in <i>Saccharomyces cerevisiae</i> . , 1999, 15, 507-511.  |      | 116       |
| 13 | Epitope tagging of yeast genes using a PCR-based strategy: more tags and improved practical routines. , 1999, 15, 963-972.  |      | 946       |
| 14 | High-efficiency gene targeting in <i>Schizosaccharomyces pombe</i> using a modular, PCR-based approach with long tracts of flanking homology. <i>Yeast</i> , 1999, 15, 1419-1427.   | 0.8  | 167       |
| 15 | The trp1- $\hat{r}$ FA designer deletion for PCR-based gene functional analysis in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1999, 15, 1769-1774.  | 0.8  | 5         |
| 16 | Control of the Terminal Step of Intracellular Membrane Fusion by Protein Phosphatase 1 &nbsp;. <i>Science</i> , 1999, 285, 1084-1087.   | 6.0  | 152       |
| 17 | ROC1, a Homolog of APC11, Represents a Family of Cullin Partners with an Associated Ubiquitin Ligase Activity. <i>Molecular Cell</i> , 1999, 3, 535-541.  | 4.5  | 429       |
| 18 | Relocalization of Telomeric Ku and SIR Proteins in Response to DNA Strand Breaks in Yeast. <i>Cell</i> , 1999, 97, 621-633.   | 13.5 | 438       |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | The RNA export factor Gle1p is located on the cytoplasmic fibrils of the NPC and physically interacts with the FG-nucleoporin Rip1p, the DEAD-box protein Rat8p/Dbp5p and a new protein Ymr255p. <i>EMBO Journal</i> , 1999, 18, 5761-5777.         | 3.5 | 140       |
| 20 | The Morphogenesis Checkpoint in <i>Saccharomyces cerevisiae</i> : Cell Cycle Control of Swe1p Degradation by Hsl1p and Hsl7p. <i>Molecular and Cellular Biology</i> , 1999, 19, 6929-6939.  | 1.1 | 156       |
| 21 | Oxidative Stress-Induced Destruction of the Yeast C-Type Cyclin Ume3p Requires Phosphatidylinositol-Specific Phospholipase C and the 26S Proteasome. <i>Molecular and Cellular Biology</i> , 1999, 19, 3338-3348.                                   | 1.1 | 70        |
| 22 | Visualizing protein dynamics in yeast with green fluorescent protein. <i>Methods in Enzymology</i> , 2000, 327, 61-69.  | 0.4 | 2         |
| 23 | The beauty of the yeast: Live cell microscopy at the limits of optical resolution. <i>Microscopy Research and Technique</i> , 2000, 51, 511-529.  | 1.2 | 48        |
| 24 | FAP1, a homologue of human transcription factor NF-X1, competes with rapamycin for binding to FKBP12 in yeast. <i>Molecular Microbiology</i> , 2000, 37, 1480-1493.   | 1.2 | 20        |
| 25 | A yeast DNAJ protein required for uncoating of clathrin-coated vesicles in vivo. <i>Nature Cell Biology</i> , 2000, 2, 958-963.   | 4.6 | 101       |
| 26 | Nuclear export of yeast signal recognition particle lacking Srp54p by the Xpo1p/Crm1p NES-dependent pathway. <i>Current Biology</i> , 2000, 10, 1256-1264.  | 1.8 | 60        |
| 27 | The Bfa1/Bub2 GAP complex comprises a universal checkpoint required to prevent mitotic exit. <i>Current Biology</i> , 2000, 10, 1379-1382.  | 1.8 | 93        |
| 28 | The forkhead protein Fkh2 is a component of the yeast cell cycle transcription factor SFF. <i>EMBO Journal</i> , 2000, 19, 3750-3761.   | 3.5 | 125       |
| 29 | A Role for Myosin-I in Actin Assembly through Interactions with Vrp1p, Bee1p, and the Arp2/3 Complex. <i>Journal of Cell Biology</i> , 2000, 148, 353-362.  | 2.3 | 227       |
| 30 | The yeast antiviral proteins Ski2p, Ski3p, and Ski8p exist as a complex in vivo. <i>Rna</i> , 2000, 6, 449-457.   | 1.6 | 149       |
| 31 | REF, an evolutionarily conserved family of hnRNP-like proteins, interacts with TAP/Mex67p and participates in mRNA nuclear export. <i>Rna</i> , 2000, 6, 638-650.   | 1.6 | 331       |
| 32 | Nascent 60S ribosomal subunits enter the free pool bound by Nmd3p. <i>Rna</i> , 2000, 6, 1625-1634.   | 1.6 | 48        |
| 33 | Inhibition of TATA-Binding Protein Function by SAGA Subunits Spt3 and Spt8 at Gcn4-Activated Promoters. <i>Molecular and Cellular Biology</i> , 2000, 20, 634-647.  | 1.1 | 133       |
| 34 | YLL031c Belongs to a Novel Family of Membrane Proteins Involved in the Transfer of Ethanolaminephosphate onto the Core Structure of Glycosylphosphatidylinositol Anchors in Yeast. <i>Journal of Biological Chemistry</i> , 2000, 275, 24458-24465. | 1.6 | 62        |
| 35 | Increased Rates of Genomic Deletions Generated by Mutations in the Yeast Gene Encoding DNA Polymerase $\epsilon$ or by Decreases in the Cellular Levels of DNA Polymerase $\epsilon$ . <i>Molecular and Cellular Biology</i> , 2000, 20, 7490-7504. | 1.1 | 87        |
| 36 | A Homolog of Voltage-Gated Ca <sup>2+</sup> Channels Stimulated by Depletion of Secretory Ca <sup>2+</sup> in Yeast. <i>Molecular and Cellular Biology</i> , 2000, 20, 6686-6694.   | 1.1 | 207       |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | PNG1, a Yeast Gene Encoding a Highly Conserved Peptide:N-Glycanase. <i>Journal of Cell Biology</i> , 2000, 149, 1039-1052.  | 2.3 | 211       |
| 38 | Repair of Intermediate Structures Produced at DNA Interstrand Cross-Links in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2000, 20, 3425-3433.   | 1.1 | 145       |
| 39 | Distinct Protein Domains of the Yeast Golgi GDP-mannose Transporter Mediate Oligomer Assembly and Export from the Endoplasmic Reticulum. <i>Journal of Biological Chemistry</i> , 2000, 275, 17718-17727.   | 1.6 | 75        |
| 40 | Roles of Hof1p, Bni1p, Bnr1p, and Myo1p in Cytokinesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2000, 11, 593-611.   | 0.9 | 210       |
| 41 | Mdv1p Is a Wd Repeat Protein That Interacts with the Dynamin-Related Gtpase, Dnm1p, to Trigger Mitochondrial Division. <i>Journal of Cell Biology</i> , 2000, 151, 353-366.   | 2.3 | 347       |
| 42 | The myosin motor, Myo4p, binds Ash1 mRNA via the adapter protein, She3p. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 5273-5278.  | 3.3 | 187       |
| 43 | Studies on the role of the hydrophobic domain of Ost4p in interactions with other subunits of yeast oligosaccharyl transferase. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 1516-1520.   | 3.3 | 29        |
| 44 | Purification of GimC from <i>Saccharomyces cerevisiae</i> . , 2000, 140, 185-193.   |     | 1         |
| 45 | The Yeast Inositol Polyphosphate 5-Phosphatases Inp52p and Inp53p Translocate to Actin Patches following Hyperosmotic Stress: Mechanism for Regulating Phosphatidylinositol 4,5-Bisphosphate at Plasma Membrane Invaginations. <i>Molecular and Cellular Biology</i> , 2000, 20, 9376-9390. | 1.1 | 36        |
| 46 | Septin-Dependent Assembly of a Cell Cycle-Regulatory Module in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2000, 20, 4049-4061.   | 1.1 | 250       |
| 47 | Multiple Signals from Dysfunctional Mitochondria Activate the Pleiotropic Drug Resistance Pathway in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2000, 275, 37347-37356.   | 1.6 | 175       |
| 48 | Telomerase-dependent repeat divergence at the 3' ends of yeast telomeres. <i>Nucleic Acids Research</i> , 2000, 28, 2690-2694.  | 6.5 | 110       |
| 49 | Prm1p, a Pheromone-Regulated Multispanning Membrane Protein, Facilitates Plasma Membrane Fusion during Yeast Mating. <i>Journal of Cell Biology</i> , 2000, 151, 719-730.   | 2.3 | 166       |
| 50 | Phosphorylation by Cdc28 Activates the Cdc20-Dependent Activity of the Anaphase-Promoting Complex. <i>Journal of Cell Biology</i> , 2000, 149, 1377-1390.   | 2.3 | 268       |
| 51 | The APC11 RING-H2 Finger Mediates E2-Dependent Ubiquitination. <i>Molecular Biology of the Cell</i> , 2000, 11, 2315-2325.  | 0.9 | 167       |
| 52 | Nmd3p Is a Crm1p-Dependent Adapter Protein for Nuclear Export of the Large Ribosomal Subunit. <i>Journal of Cell Biology</i> , 2000, 151, 1057-1066.  | 2.3 | 258       |
| 53 | Essential Function of the Polo Box of Cdc5 in Subcellular Localization and Induction of Cytokinetic Structures. <i>Molecular and Cellular Biology</i> , 2000, 20, 286-298.  | 1.1 | 151       |
| 54 | Functional Cus1p Is Found with Hsh155p in a Multiprotein Splicing Factor Associated with U2 snRNA. <i>Molecular and Cellular Biology</i> , 2000, 20, 2176-2185.   | 1.1 | 28        |

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 55 | Cdc28 Activates Exit from Mitosis in Budding Yeast. <i>Journal of Cell Biology</i> , 2000, 149, 1361-1376.   | 2.3  | 83        |
| 56 | The Secretory Pathway Mediates Localization of the Cell Polarity Regulator Aip3p/Bud6p. <i>Molecular Biology of the Cell</i> , 2000, 11, 647-661.  | 0.9  | 63        |
| 57 | Multiple Signals Regulate GAL Transcription in Yeast. <i>Molecular and Cellular Biology</i> , 2000, 20, 3880-3886.   | 1.1  | 69        |
| 58 | Ama1p is a meiosis-specific regulator of the anaphase promoting complex/cyclosome in yeast. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 14548-14553.  | 3.3  | 135       |
| 59 | A <i>Saccharomyces</i> gene family involved in invasive growth, cell-cell adhesion, and mating. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 12158-12163.  | 3.3  | 357       |
| 60 | Characterization of Yeast Yea4p, a Uridine Diphosphate-N-acetylglucosamine Transporter Localized in the Endoplasmic Reticulum and Required for Chitin Synthesis. <i>Journal of Biological Chemistry</i> , 2000, 275, 13580-13587.  | 1.6  | 51        |
| 61 | Identification of Two Novel TAF Subunits of the Yeast <i>Saccharomyces cerevisiae</i> TFIID Complex. <i>Journal of Biological Chemistry</i> , 2000, 275, 13895-13900.  | 1.6  | 69        |
| 62 | Molecular Characterization of FMN1, the Structural Gene for the Monofunctional Flavokinase of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2000, 275, 28618-28624.   | 1.6  | 61        |
| 63 | Mechanism of Metabolic Control. <i>Journal of Cell Biology</i> , 2000, 151, 863-878.   | 2.3  | 196       |
| 64 | Disruption of Cytoplasmic and Mitochondrial Polyglutamate Synthetase Activity in <i>Saccharomyces cerevisiae</i> . <i>Archives of Biochemistry and Biophysics</i> , 2000, 376, 299-312.  | 1.4  | 29        |
| 65 | The <i>Saccharomyces cerevisiae</i> homologue YPA1 of the mammalian phosphotyrosyl phosphatase activator of protein phosphatase 2A controls progression through the G1 phase of the yeast cell cycle. Edited by J. Karn. <i>Journal of Molecular Biology</i> , 2000, 302, 103-119. | 2.0  | 28        |
| 66 | A novel multi-purpose cassette for repeated integrative epitope tagging of genes in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 2000, 246, 179-185.  | 1.0  | 99        |
| 67 | The Bub2p Spindle Checkpoint Links Nuclear Migration with Mitotic Exit. <i>Molecular Cell</i> , 2000, 6, 1-10.   | 4.5  | 299       |
| 68 | A Yeast Intronic Splicing Enhancer and Nam8p Are Required for Mer1p-Activated Splicing. <i>Molecular Cell</i> , 2000, 6, 329-338.  | 4.5  | 73        |
| 69 | Class C Vps Protein Complex Regulates Vacuolar SNARE Pairing and Is Required for Vesicle Docking/Fusion. <i>Molecular Cell</i> , 2000, 6, 661-671.   | 4.5  | 250       |
| 70 | DNA Polymerase $\eta$ Introduces Multiple Mutations when Bypassing Spontaneous DNA Damage in <i>Saccharomyces cerevisiae</i> . <i>Molecular Cell</i> , 2000, 6, 1491-1499.   | 4.5  | 114       |
| 71 | The Isw2 Chromatin Remodeling Complex Represses Early Meiotic Genes upon Recruitment by Ume6p. <i>Cell</i> , 2000, 103, 423-433.   | 13.5 | 280       |
| 72 | Functional and Genomic Analyses Reveal an Essential Coordination between the Unfolded Protein Response and ER-Associated Degradation. <i>Cell</i> , 2000, 101, 249-258.  | 13.5 | 1,777     |

| #  | ARTICLE   | IF   | CITATIONS |
|----|---|------|-----------|
| 73 | A novel strategy for constructing N-terminal chromosomal fusions to green fluorescent protein in the yeast <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 2000, 485, 29-34.  | 1.3  | 41        |
| 74 | Molecular Linkage Underlying Microtubule Orientation Toward Cortical Sites in Yeast. <i>Science</i> , 2000, 287, 2257-2259.   | 6.0  | 200       |
| 75 | Plasma Membrane Compartmentalization in Yeast by Messenger RNA Transport and a Septin Diffusion Barrier. <i>Science</i> , 2000, 290, 341-344.   | 6.0  | 400       |
| 76 | The Yeast Nuclear Pore Complex. <i>Journal of Cell Biology</i> , 2000, 148, 635-652.  | 2.3  | 1,329     |
| 77 | The yeast mitotic cyclin Clb2 cannot substitute for S phase cyclins in replication origin firing. <i>EMBO Reports</i> , 2000, 1, 507-512.   | 2.0  | 33        |
| 78 | <i>Saccharomyces cerevisiae</i> Ras/cAMP pathway controls post-diauxic shift element-dependent transcription through the zinc finger protein Gis1. <i>EMBO Journal</i> , 2000, 19, 2569-2579.                                 | 3.5  | 187       |
| 79 | Protein Phosphatase Methyltransferase 1 (Ppm1p) Is the Sole Activity Responsible for Modification of the Major Forms of Protein Phosphatase 2A in Yeast. <i>Archives of Biochemistry and Biophysics</i> , 2001, 395, 239-245. | 1.4  | 29        |
| 80 | Alkaline Response Genes of <i>Saccharomyces cerevisiae</i> and Their Relationship to the RIM101 Pathway. <i>Journal of Biological Chemistry</i> , 2001, 276, 1850-1856.   | 1.6  | 205       |
| 81 | SAGA is an essential in vivo target of the yeast acidic activator Gal4p. <i>Genes and Development</i> , 2001, 15, 1935-1945.  | 2.7  | 258       |
| 82 | Identification of RFC(Ctf18p, Ctf8p, Dcc1p). <i>Molecular Cell</i> , 2001, 7, 959-970.  | 4.5  | 308       |
| 83 | Composition and Functional Characterization of Yeast 66S Ribosome Assembly Intermediates. <i>Molecular Cell</i> , 2001, 8, 505-515.   | 4.5  | 280       |
| 84 | TIP41 Interacts with TAP42 and Negatively Regulates the TOR Signaling Pathway. <i>Molecular Cell</i> , 2001, 8, 1017-1026.  | 4.5  | 208       |
| 85 | Autophagy and the cytoplasm to vacuole targeting pathway both require Aut10p. <i>FEBS Letters</i> , 2001, 508, 23-28.   | 1.3  | 102       |
| 86 | The Transcription Factor Associated Ccr4 and Caf1 Proteins Are Components of the Major Cytoplasmic mRNA Deadenylation in <i>Saccharomyces cerevisiae</i> . <i>Cell</i> , 2001, 104, 377-386.                                  | 13.5 | 523       |
| 87 | Maturation and Intranuclear Transport of Pre-Ribosomes Requires Noc Proteins. <i>Cell</i> , 2001, 105, 499-509.   | 13.5 | 206       |
| 88 | Ubiquitin-Dependent Sorting into the Multivesicular Body Pathway Requires the Function of a Conserved Endosomal Protein Sorting Complex, ESCRT-I. <i>Cell</i> , 2001, 106, 145-155.   | 13.5 | 1,228     |
| 89 | Multiple Gln/Asn-Rich Prion Domains Confer Susceptibility to Induction of the Yeast [PSI <sup>+</sup> ] Prion. <i>Cell</i> , 2001, 106, 183-194.  | 13.5 | 303       |
| 90 | Regulation of the Bub2/Bfa1 GAP Complex by Cdc5 and Cell Cycle Checkpoints. <i>Cell</i> , 2001, 107, 655-665.   | 13.5 | 235       |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 91  | Bms1p, a novel GTP-binding protein, and the related Tsr1p are required for distinct steps of 40S ribosome biogenesis in yeast. <i>Rna</i> , 2001, 7, 1268-1283.  | 1.6 | 117       |
| 92  | Phosphorylation by Sky1p promotes Npl3p shuttling and mRNA dissociation. <i>Rna</i> , 2001, 7, 302-313.  | 1.6 | 123       |
| 93  | A Streamlined Process to Phenotypically Profile Heterologous cDNAs in Parallel Using Yeast Cell-Based Assays. <i>Genome Research</i> , 2001, 11, 1899-1912.  | 2.4 | 30        |
| 94  | Effect of chromosomal locus, GC content and length of homology on PCR-mediated targeted gene replacement in <i>Saccharomyces</i> . <i>Nucleic Acids Research</i> , 2001, 29, 5156-5162.  | 6.5 | 36        |
| 95  | Molecular dissection of yeast spindle pole bodies by two hybrid, in vitro binding, and co-purification. <i>Methods in Cell Biology</i> , 2001, 67, 71-94.  | 0.5 | 10        |
| 96  | Chapter 15 Epitope tagging and visualization of nuclear-encoded mitochondrial proteins in yeast. <i>Methods in Cell Biology</i> , 2001, 65, 257-276.   | 0.5 | 4         |
| 97  | The use of transgenic plants in the bioremediation of soils contaminated with trace elements. <i>Applied Microbiology and Biotechnology</i> , 2001, 55, 661-672.   | 1.7 | 216       |
| 98  | Cgr1p, a Novel Nucleolar Protein Encoded by <i>Saccharomyces cerevisiae</i> Orf YGL0292w. <i>Current Microbiology</i> , 2001, 42, 65-69.   | 1.0 | 11        |
| 99  | Overlapping and distinct roles of the duplicated yeast transcription factors Ace2p and Swi5p. <i>Molecular Microbiology</i> , 2001, 40, 422-432.   | 1.2 | 122       |
| 100 | Genetic analysis of the signalling pathway activated by external amino acids in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2001, 41, 489-502.   | 1.2 | 54        |
| 101 | Vrp1p Functions in Both Actomyosin Ring-Dependent and Hof1p-Dependent Pathways of Cytokinesis. <i>Traffic</i> , 2001, 2, 189-201.  | 1.3 | 29        |
| 102 | Hut1 proteins identified in <i>Saccharomyces cerevisiae</i> and <i>Schizosaccharomyces pombe</i> are functional homologues involved in the protein-folding process at the endoplasmic reticulum. <i>Yeast</i> , 2001, 18, 543-554. | 0.8 | 51        |
| 103 | Overexpression of HUT1 gene stimulates in vivo galactosylation by enhancing UDP-galactose transport activity in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2001, 18, 533-541.  | 0.8 | 33        |
| 104 | Constitutive promoter modules for PCR-based gene modification in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2001, 18, 723-728.   | 0.8 | 8         |
| 105 | Yeast Artificial Chromosome Targeting Technology: An Approach for the Deletion of Genes in the C57BL/6 Mouse. <i>Analytical Biochemistry</i> , 2001, 296, 270-278.   | 1.1 | 6         |
| 106 | Cortex, a <i>Drosophila</i> gene required to complete oocyte meiosis, is a member of the Cdc20/fizzy protein family. <i>Genesis</i> , 2001, 29, 141-152.   | 0.8 | 63        |
| 107 | A specific DNA-dependent DNA polymerase is associated with <i>Saccharomyces cerevisiae</i> telomerase. <i>Biochemistry (Moscow)</i> , 2001, 66, 1361-1367.   | 0.7 | 3         |
| 108 | Sec63p and Kar2p are required for the translocation of SRP-dependent precursors into the yeast endoplasmic reticulum in vivo. <i>EMBO Journal</i> , 2001, 20, 262-271.   | 3.5 | 145       |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 109 | Cic1, an adaptor protein specifically linking the 26S proteasome to its substrate, the SCF component Cdc4. <i>EMBO Journal</i> , 2001, 20, 4423-4431.   | 3.5 | 44        |
| 110 | A cytosolic NAD-dependent deacetylase, Hst2p, can modulate nucleolar and telomeric silencing in yeast. <i>EMBO Journal</i> , 2001, 20, 197-209.   | 3.5 | 147       |
| 111 | Spt16-Pob3 and the HMG protein Nhp6 combine to form the nucleosome-binding factor SPN. <i>EMBO Journal</i> , 2001, 20, 3506-3517.   | 3.5 | 237       |
| 112 | A nuclear AAA-type ATPase (Rix7p) is required for biogenesis and nuclear export of 60S ribosomal subunits. <i>EMBO Journal</i> , 2001, 20, 3695-3704.   | 3.5 | 81        |
| 113 | Nog2p, a putative GTPase associated with pre-60S subunits and required for late 60S maturation steps. <i>EMBO Journal</i> , 2001, 20, 6475-6484.  | 3.5 | 171       |
| 114 | The intracellular location of two aminoacyl-tRNA synthetases depends on complex formation with Arc1p. <i>EMBO Journal</i> , 2001, 20, 6889-6898.  | 3.5 | 68        |
| 115 | Functions of Vrp1p in cytokinesis and actin patches are distinct and neither requires a WH2/V domain. <i>EMBO Journal</i> , 2001, 20, 6979-6989.  | 3.5 | 19        |
| 116 | Post-transcriptional regulation through the HO 3'-UTR by Mpt5, a yeast homolog of Pumilio and FBF. <i>EMBO Journal</i> , 2001, 20, 552-561.   | 3.5 | 126       |
| 117 | Mrc1 transduces signals of DNA replication stress to activate Rad53. <i>Nature Cell Biology</i> , 2001, 3, 958-965.   | 4.6 | 474       |
| 118 | Yeast Eps15-like endocytic protein, Pan1p, activates the Arp2/3 complex. <i>Nature Cell Biology</i> , 2001, 3, 687-690.   | 4.6 | 158       |
| 119 | Exonuclease activity is required for sequence addition and Cdc13p loading at a de novo telomere. <i>Current Biology</i> , 2001, 11, 1336-1340.  | 1.8 | 136       |
| 120 | Arp2/3 complex and actin dynamics are required for actin-based mitochondrial motility in yeast. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2001, 98, 3162-3167. | 3.3 | 180       |
| 121 | The Sda1 Protein Is Required for Passage through Start. <i>Molecular Biology of the Cell</i> , 2001, 12, 201-219.   | 0.9 | 42        |
| 122 | A Novel Function of <i>Saccharomyces cerevisiae</i> CDC5 in Cytokinesis. <i>Journal of Cell Biology</i> , 2001, 152, 451-470.   | 2.3 | 114       |
| 123 | Yeast Gga Coat Proteins Function with Clathrin in Golgi to Endosome Transport. <i>Molecular Biology of the Cell</i> , 2001, 12, 1885-1896.  | 0.9 | 136       |
| 124 | Yng2p-dependent NuA4 Histone H4 Acetylation Activity Is Required for Mitotic and Meiotic Progression. <i>Journal of Biological Chemistry</i> , 2001, 276, 43653-43662.  | 1.6 | 55        |
| 125 | Bud8p and Bud9p, Proteins That May Mark the Sites for Bipolar Budding in Yeast. <i>Molecular Biology of the Cell</i> , 2001, 12, 2497-2518.   | 0.9 | 90        |
| 126 | Erv41p and Erv46p. <i>Journal of Cell Biology</i> , 2001, 152, 503-518.   | 2.3 | 135       |



| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 127 | <i>SPO21</i> Is Required for Meiosis-specific Modification of the Spindle Pole Body in Yeast. <i>Molecular Biology of the Cell</i> , 2001, 12, 1611-1621.   | 0.9 | 53        |
| 128 | <i>Candida albicans</i> Int1p Interacts with the Septin Ring in Yeast and Hyphal Cells. <i>Molecular Biology of the Cell</i> , 2001, 12, 3538-3549.   | 0.9 | 76        |
| 129 | The Yeast hnRNP-Like Proteins Yra1p and Yra2p Participate in mRNA Export through Interaction with Mex67p. <i>Molecular and Cellular Biology</i> , 2001, 21, 4219-4232.  | 1.1 | 134       |
| 130 | Dynamic Localization and Function of Bni1p at the Sites of Directed Growth in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2001, 21, 827-839.  | 1.1 | 136       |
| 131 | A Family of Small Coiled-Coil-forming Proteins Functioning at the Late Endosome in Yeast. <i>Molecular Biology of the Cell</i> , 2001, 12, 711-723.   | 0.9 | 73        |
| 132 | Sac1 Lipid Phosphatase and Stt4 Phosphatidylinositol 4-Kinase Regulate a Pool of Phosphatidylinositol 4-Phosphate That Functions in the Control of the Actin Cytoskeleton and Vacuole Morphology. <i>Molecular Biology of the Cell</i> , 2001, 12, 2396-2411. | 0.9 | 216       |
| 133 | CAC3 ( MS11 ) Suppression of RAS2 G19V Is Independent of Chromatin Assembly Factor I and Mediated by NPR1. <i>Molecular and Cellular Biology</i> , 2001, 21, 1784-1794.   | 1.1 | 28        |
| 134 | Degradation of Endoplasmic Reticulum (ER) Quality Control Substrates Requires Transport between the ER and Golgi. <i>Journal of Biological Chemistry</i> , 2001, 276, 23296-23303.  | 1.6 | 159       |
| 135 | Identification of 12 New Yeast Mitochondrial Ribosomal Proteins Including 6 That Have No Prokaryotic Homologues. <i>Journal of Biological Chemistry</i> , 2001, 276, 15861-15867.   | 1.6 | 67        |
| 136 | Xbp1-Mediated Repression of CLB Gene Expression Contributes to the Modifications of Yeast Cell Morphology and Cell Cycle Seen during Nitrogen-Limited Growth. <i>Molecular and Cellular Biology</i> , 2001, 21, 3714-3724.                                    | 1.1 | 28        |
| 137 | Role of positively charged transmembrane segments in the insertion and assembly of mitochondrial inner-membrane proteins. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2001, 98, 13814-13819.                     | 3.3 | 22        |
| 138 | <i>Saccharomyces cerevisiae</i> Mob1p Is Required for Cytokinesis and Mitotic Exit. <i>Molecular and Cellular Biology</i> , 2001, 21, 6972-6983.  | 1.1 | 135       |
| 139 | Maf1p, a Negative Effector of RNA Polymerase III in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2001, 21, 5031-5040.  | 1.1 | 163       |
| 140 | A protein interaction map for cell polarity development. <i>Journal of Cell Biology</i> , 2001, 154, 549-576.   | 2.3 | 294       |
| 141 | New Function of CDC13 in Positive Telomere Length Regulation. <i>Molecular and Cellular Biology</i> , 2001, 21, 4233-4245.  | 1.1 | 17        |
| 142 | Cortical Num1p Interacts with the Dynein Intermediate Chain Pac11p and Cytoplasmic Microtubules in Budding Yeast. <i>Journal of Cell Biology</i> , 2001, 152, 251-262.  | 2.3 | 108       |
| 143 | The nucleoporin Nup60p functions as a Gsp1p-GTP-sensitive tether for Nup2p at the nuclear pore complex. <i>Journal of Cell Biology</i> , 2001, 154, 937-950.  | 2.3 | 84        |
| 144 | The yeast Xrs2 complex functions in S phase checkpoint regulation. <i>Genes and Development</i> , 2001, 15, 2238-2249.  | 2.7 | 194       |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 145 | Asf1 links Rad53 to control of chromatin assembly. <i>Genes and Development</i> , 2001, 15, 1061-1066.   | 2.7 | 146       |
| 146 | Dad1p, Third Component of the Duo1p/Dam1p Complex Involved in Kinetochores Function and Mitotic Spindle Integrity. <i>Molecular Biology of the Cell</i> , 2001, 12, 2601-2613.                                   | 0.9 | 60        |
| 147 | Vps41p Function in the Alkaline Phosphatase Pathway Requires Homo-oligomerization and Interaction with AP-3 through Two Distinct Domains. <i>Molecular Biology of the Cell</i> , 2001, 12, 37-51.                | 0.9 | 80        |
| 148 | Polyploids require Bik1 for kinetochores microtubule attachment. <i>Journal of Cell Biology</i> , 2001, 155, 1173-1184.  | 2.3 | 98        |
| 149 | The Npr1 Kinase Controls Biosynthetic and Endocytic Sorting of the Yeast Gap1 Permease. <i>Journal of Biological Chemistry</i> , 2001, 276, 43939-43948.   | 1.6 | 147       |
| 150 | A conserved ubiquitin ligase of the nuclear envelope/endoplasmic reticulum that functions in both ER-associated and Matalpha 2 repressor degradation. <i>Genes and Development</i> , 2001, 15, 2660-2674.        | 2.7 | 421       |
| 151 | Skp1p and the F-Box Protein Rcy1p Form a Non-SCF Complex Involved in Recycling of the SNARE Snc1p in Yeast. <i>Molecular and Cellular Biology</i> , 2001, 21, 3105-3117.   | 1.1 | 157       |
| 152 | A Cdc28 Mutant Uncouples G1 Cyclin Phosphorylation and Ubiquitination from G1 Cyclin Proteolysis. <i>Journal of Biological Chemistry</i> , 2001, 276, 41725-41732.   | 1.6 | 4         |
| 153 | The Swi5 activator recruits the Mediator complex to the HO promoter without RNA polymerase II. <i>Genes and Development</i> , 2001, 15, 2457-2469.   | 2.7 | 116       |
| 154 | The yeast SAS (something about silencing) protein complex contains a MYST-type putative acetyltransferase and functions with chromatin assembly factor ASF1. <i>Genes and Development</i> , 2001, 15, 3155-3168. | 2.7 | 127       |
| 155 | In vivo chromatin remodeling by yeast ISWI homologs Isw1p and Isw2p. <i>Genes and Development</i> , 2001, 15, 619-626.   | 2.7 | 108       |
| 156 | NHEJ regulation by mating type is exercised through a novel protein, Lif2p, essential to the Ligase IV pathway. <i>Genes and Development</i> , 2001, 15, 3005-3012.  | 2.7 | 156       |
| 157 | Yeast PalA/AIP1/Alix Homolog Rim20p Associates with a PEST-Like Region and Is Required for Its Proteolytic Cleavage. <i>Journal of Bacteriology</i> , 2001, 183, 6917-6923.                                      | 1.0 | 113       |
| 158 | Transcriptional Coregulation by the Cell Integrity Mitogen-Activated Protein Kinase Slt2 and the Cell Cycle Regulator Swi4. <i>Molecular and Cellular Biology</i> , 2001, 21, 6515-6528.                         | 1.1 | 109       |
| 159 | The brefeldin A resistance protein Bfr1p is a component of polyribosome-associated mRNP complexes in yeast. <i>Nucleic Acids Research</i> , 2001, 29, 2567-2574.   | 6.5 | 45        |
| 160 | A Gip1p-Glc7p phosphatase complex regulates septin organization and spore wall formation. <i>Journal of Cell Biology</i> , 2001, 155, 797-808.   | 2.3 | 88        |
| 161 | Implication of a novel multiprotein Dam1p complex in outer kinetochores function. <i>Journal of Cell Biology</i> , 2001, 155, 1137-1146.   | 2.3 | 167       |
| 162 | A Novel Form of Transcriptional Silencing by Sum1-1 Requires Hst1 and the Origin Recognition Complex. <i>Molecular and Cellular Biology</i> , 2001, 21, 3514-3522.   | 1.1 | 83        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 163 | Multiple roles for the yeast SUB2/yUAP56 gene in splicing. <i>Genes and Development</i> , 2001, 15, 36-41.   | 2.7 | 111       |
| 164 | Aux1p/Swa2p Is Required for Cortical Endoplasmic Reticulum Inheritance in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2001, 12, 2614-2628.  | 0.9 | 52        |
| 165 | Cooperative Binding of the Cytoplasm to Vacuole Targeting Pathway Proteins, Cvt13 and Cvt20, to Phosphatidylinositol 3-Phosphate at the Pre-autophagosomal Structure Is Required for Selective Autophagy. <i>Journal of Biological Chemistry</i> , 2002, 277, 30198-30207. | 1.6 | 176       |
| 166 | Yeast Oxidosqualene Cyclase (Erg7p) Is a Major Component of Lipid Particles. <i>Journal of Biological Chemistry</i> , 2002, 277, 2406-2412.  | 1.6 | 80        |
| 167 | The SANT Domain of Ada2 Is Required for Normal Acetylation of Histones by the Yeast SAGA Complex. <i>Journal of Biological Chemistry</i> , 2002, 277, 8178-8186.   | 1.6 | 106       |
| 168 | A novel mitochondrial protein, Tar1p, is encoded on the antisense strand of the nuclear 25S rDNA. <i>Genes and Development</i> , 2002, 16, 2755-2760.  | 2.7 | 67        |
| 169 | A Lack of SUMO Conjugation Affects cNLS-dependent Nuclear Protein Import in Yeast. <i>Journal of Biological Chemistry</i> , 2002, 277, 49554-49561.  | 1.6 | 63        |
| 170 | Sgt1p Contributes to Cyclic AMP Pathway Activity and Physically Interacts with the Adenylyl Cyclase Cyr1p/Cdc35p in Budding Yeast. <i>Eukaryotic Cell</i> , 2002, 1, 568-582.  | 3.4 | 92        |
| 171 | Protein Kinase A Operates a Molecular Switch That Governs Yeast Pseudohyphal Differentiation. <i>Molecular and Cellular Biology</i> , 2002, 22, 3981-3993.   | 1.1 | 172       |
| 172 | An Essential Function of Yeast Cyclin-dependent Kinase Cdc28 Maintains Chromosome Stability. <i>Journal of Biological Chemistry</i> , 2002, 277, 48627-48634.  | 1.6 | 16        |
| 173 | Genetic analysis of iron citrate toxicity in yeast: Implications for mammalian iron homeostasis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 16922-16927.   | 3.3 | 42        |
| 174 | Identification of Interdependent Signals Required for Anterograde Traffic of the ATP-binding Cassette Transporter Protein Yor1p. <i>Journal of Biological Chemistry</i> , 2002, 277, 34860-34869.  | 1.6 | 23        |
| 175 | The RAVE Complex Is Essential for Stable Assembly of the Yeast V-ATPase. <i>Journal of Biological Chemistry</i> , 2002, 277, 13831-13839.  | 1.6 | 160       |
| 176 | A second set of loxP marker cassettes for Cre-mediated multiple gene knockouts in budding yeast. <i>Nucleic Acids Research</i> , 2002, 30, 23e-23.   | 6.5 | 855       |
| 177 | Uptake of the anticancer drug cisplatin mediated by the copper transporter Ctr1 in yeast and mammals. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 14298-14302.  | 3.3 | 770       |
| 178 | Bcl-x Complements <i>Saccharomyces cerevisiae</i> Genes That Facilitate the Switch from Glycolytic to Oxidative Metabolism. <i>Journal of Biological Chemistry</i> , 2002, 277, 44870-44876.   | 1.6 | 59        |
| 179 | The role of cell cycle-regulated expression in the localization of spatial landmark proteins in yeast. <i>Journal of Cell Biology</i> , 2002, 156, 829-841.  | 2.3 | 60        |
| 180 | The Rpb9 Subunit of RNA Polymerase II Binds Transcription Factor TFIIIE and Interferes with the SAGA and Elongator Histone Acetyltransferases. <i>Journal of Biological Chemistry</i> , 2002, 277, 10220-10225.  | 1.6 | 52        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 181 | Actin cable dynamics in budding yeast. Proceedings of the National Academy of Sciences of the United States of America, 2002, 99, 751-756.   | 3.3 | 180       |
| 182 | PCR-based engineering of yeast genome. Methods in Enzymology, 2002, 350, 445-469.  | 0.4 | 85        |
| 183 | Regulation of the <i>Saccharomyces cerevisiae</i> Slt2 Kinase Pathway by the Stress-inducible Sdp1 Dual Specificity Phosphatase. Journal of Biological Chemistry, 2002, 277, 21278-21284.  | 1.6 | 104       |
| 184 | Lysine methylation within the globular domain of histone H3 by Dot1 is important for telomeric silencing and Sir protein association. Genes and Development, 2002, 16, 1518-1527.  | 2.7 | 471       |
| 185 | SALSA, a variant of yeast SAGA, contains truncated Spt7, which correlates with activated transcription. Proceedings of the National Academy of Sciences of the United States of America, 2002, 99, 11622-11627.                        | 3.3 | 111       |
| 186 | The TOR-controlled transcription activators GLN3, RTG1, and RTG3 are regulated in response to intracellular levels of glutamine. Proceedings of the National Academy of Sciences of the United States of America, 2002, 99, 6784-6789. | 3.3 | 287       |
| 187 | The Yeast Homolog of Human PinX1 Is Involved in rRNA and Small Nucleolar RNA Maturation, Not in Telomere Elongation Inhibition. Journal of Biological Chemistry, 2002, 277, 35712-35719.   | 1.6 | 74        |
| 188 | Nucleotide Exchange Factor for the Yeast Hsp70 Molecular Chaperone Ssa1p. Molecular and Cellular Biology, 2002, 22, 4677-4689.   | 1.1 | 133       |
| 189 | Spo13 regulates cohesin cleavage. Genes and Development, 2002, 16, 1672-1681.  | 2.7 | 46        |
| 190 | The Glc7p-Interacting Protein Bud14p Attenuates Polarized Growth, Pheromone Response, and Filamentous Growth in <i>Saccharomyces cerevisiae</i> . Eukaryotic Cell, 2002, 1, 884-894.   | 3.4 | 28        |
| 191 | The NC2 alpha and beta subunits play different roles in vivo. Genes and Development, 2002, 16, 3265-3276.  | 2.7 | 27        |
| 192 | The Retinoblastoma Family of Proteins Directly Represses Transcription in <i>Saccharomyces cerevisiae</i> . Journal of Biological Chemistry, 2002, 277, 8797-8801.   | 1.6 | 5         |
| 193 | Synthesis of Triacylglycerols by the Acyl-Coenzyme A:Diacyl-Glycerol Acyltransferase Dga1p in Lipid Particles of the Yeast <i>Saccharomyces cerevisiae</i> . Journal of Bacteriology, 2002, 184, 519-524.                              | 1.0 | 200       |
| 194 | Ctf3p, the Mis6 budding yeast homolog, interacts with Mcm22p and Mcm16p at the yeast outer kinetochore. Genes and Development, 2002, 16, 101-113.  | 2.7 | 111       |
| 195 | Mps3p is a novel component of the yeast spindle pole body that interacts with the yeast centrin homologue Cdc31p. Journal of Cell Biology, 2002, 159, 945-956.   | 2.3 | 146       |
| 196 | Mutation of YCS4, a Budding Yeast Condensin Subunit, Affects Mitotic and Nonmitotic Chromosome Behavior. Molecular Biology of the Cell, 2002, 13, 632-645.   | 0.9 | 167       |
| 197 | A Ypt32p Exchange Factor Is a Putative Effector of Ypt1p. Molecular Biology of the Cell, 2002, 13, 3336-3343.  | 0.9 | 56        |
| 198 | The Roles of Bud-Site-Selection Proteins during Haploid Invasive Growth in Yeast. Molecular Biology of the Cell, 2002, 13, 2990-3004.  | 0.9 | 75        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 199 | Methylation of histone H3 Lys 4 in coding regions of active genes. Proceedings of the National Academy of Sciences of the United States of America, 2002, 99, 8695-8700.   | 3.3 | 673       |
| 200 | Ubiquitination of Histone H2B by Rad6 Is Required for Efficient Dot1-mediated Methylation of Histone H3 Lysine 79. Journal of Biological Chemistry, 2002, 277, 34655-34657.  | 1.6 | 365       |
| 201 | Interaction between Not1p, a Component of the Ccr4-Not Complex, a Global Regulator of Transcription, and Dhh1p, a Putative RNA Helicase. Journal of Biological Chemistry, 2002, 277, 2835-2842.                                    | 1.6 | 86        |
| 202 | Yeast Cells Lacking the ARV1 Gene Harbor Defects in Sphingolipid Metabolism. Journal of Biological Chemistry, 2002, 277, 36152-36160.  | 1.6 | 59        |
| 203 | The Shq1p-Naf1p Complex Is Required for Box H/ACA Small Nucleolar Ribonucleoprotein Particle Biogenesis. Journal of Biological Chemistry, 2002, 277, 45235-45242.  | 1.6 | 61        |
| 204 | Regulation of Fab1 Phosphatidylinositol 3-Phosphate 5-Kinase Pathway by Vac7 Protein and Fig4, a Polyphosphoinositide Phosphatase Family Member. Molecular Biology of the Cell, 2002, 13, 1238-1251.                               | 0.9 | 159       |
| 205 | Role of Ndt80, Sum1, and Swe1 as Targets of the Meiotic Recombination Checkpoint That Control Exit from Pachytene and Spore Formation in Saccharomyces cerevisiae. Molecular and Cellular Biology, 2002, 22, 6430-6440.            | 1.1 | 55        |
| 206 | Conserved homeodomain proteins interact with MADS box protein Mcm1 to restrict ECB-dependent transcription to the M/G1 phase of the cell cycle. Genes and Development, 2002, 16, 3034-3045.  | 2.7 | 144       |
| 207 | Mitochondrial Import of the ADP/ATP Carrier: the Essential TIM Complex of the Intermembrane Space Is Required for Precursor Release from the TOM Complex. Molecular and Cellular Biology, 2002, 22, 7780-7789.                     | 1.1 | 97        |
| 208 | ER-Golgi Traffic Is a Prerequisite for Efficient ER Degradation. Molecular Biology of the Cell, 2002, 13, 1806-1818.   | 0.9 | 105       |
| 209 | The Mnd1 Protein Forms a Complex with Hop2 To Promote Homologous Chromosome Pairing and Meiotic Double-Strand Break Repair. Molecular and Cellular Biology, 2002, 22, 3078-3088.   | 1.1 | 138       |
| 210 | Molecular Characterization of Saccharomyces cerevisiae TFIID. Molecular and Cellular Biology, 2002, 22, 6000-6013.   | 1.1 | 98        |
| 211 | Bfa1 can regulate Tem1 function independently of Bub2 in the mitotic exit network of Saccharomyces cerevisiae. Proceedings of the National Academy of Sciences of the United States of America, 2002, 99, 5436-5441.               | 3.3 | 46        |
| 212 | Dissection of a Complex Phenotype by Functional Genomics Reveals Roles for the Yeast Cyclin-Dependent Protein Kinase Pho85 in Stress Adaptation and Cell Integrity. Molecular and Cellular Biology, 2002, 22, 5076-5088.           | 1.1 | 85        |
| 213 | Identification of a Multifunctional Domain in Autonomously Replicating Sequence-Binding Factor 1 Required for Transcriptional Activation, DNA Replication, and Gene Silencing. Molecular and Cellular Biology, 2002, 22, 505-516.  | 1.1 | 38        |
| 214 | Proteomics Analysis Reveals Stable Multiprotein Complexes in Both Fission and Budding Yeasts Containing Myb-Related Cdc5p/Cef1p, Novel Pre-mRNA Splicing Factors, and snRNAs. Molecular and Cellular Biology, 2002, 22, 2011-2024. | 1.1 | 193       |
| 215 | Microtubule capture by the cleavage apparatus is required for proper spindle positioning in yeast. Genes and Development, 2002, 16, 1627-1639.   | 2.7 | 132       |
| 216 | The Saccharomyces cerevisiae LSB6 Gene Encodes Phosphatidylinositol 4-Kinase Activity. Journal of Biological Chemistry, 2002, 277, 47709-47718.  | 1.6 | 75        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 217 | Cloning-free genome alterations in <i>saccharomyces cerevisiae</i> using adaptamer-mediated PCR. <i>Methods in Enzymology</i> , 2002, 350, 258-277.   | 0.4 | 91        |
| 218 | Genome-wide location and regulated recruitment of the RSC nucleosome-remodeling complex. <i>Genes and Development</i> , 2002, 16, 806-819.  | 2.7 | 239       |
| 219 | Glycopeptide export from the endoplasmic reticulum into cytosol is mediated by a mechanism distinct from that for export of misfolded glycoprotein. <i>Glycobiology</i> , 2002, 12, 803-811.  | 1.3 | 12        |
| 220 | Ntg2p, a <i>Saccharomyces cerevisiae</i> DNAN-Glycosylase/Apurinic or Apyrimidinic Lyase Involved in Base Excision Repair of Oxidative DNA Damage, Interacts with the DNA Mismatch Repair Protein Mlh1p. <i>Journal of Biological Chemistry</i> , 2002, 277, 29963-29972. | 1.6 | 32        |
| 221 | ROX1 and ERG Regulation in <i>Saccharomyces cerevisiae</i> : Implications for Antifungal Susceptibility. <i>Eukaryotic Cell</i> , 2002, 1, 1041-1044.   | 3.4 | 42        |
| 222 | Spo13 protects meiotic cohesin at centromeres in meiosis I. <i>Genes and Development</i> , 2002, 16, 1659-1671.   | 2.7 | 85        |
| 223 | Yeast Vps55p, a Functional Homolog of Human Obesity Receptor Gene-related Protein, Is Involved in Late Endosome to Vacuole Trafficking. <i>Molecular Biology of the Cell</i> , 2002, 13, 1694-1708.   | 0.9 | 40        |
| 224 | Sterol-dependent Regulation of Sphingolipid Metabolism in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2002, 277, 26177-26184.  | 1.6 | 53        |
| 225 | Exosome-Mediated Recognition and Degradation of mRNAs Lacking a Termination Codon. <i>Science</i> , 2002, 295, 2262-2264.   | 6.0 | 493       |
| 226 | Stable mRNP Formation and Export Require Cotranscriptional Recruitment of the mRNA Export Factors Yra1p and Sub2p by Hpr1p. <i>Molecular and Cellular Biology</i> , 2002, 22, 8241-8253.  | 1.1 | 274       |
| 227 | The <i>Saccharomyces cerevisiae</i> Mob2p-Cbk1p kinase complex promotes polarized growth and acts with the mitotic exit network to facilitate daughter cell-specific localization of Ace2p transcription factor. <i>Journal of Cell Biology</i> , 2002, 158, 885-900.     | 2.3 | 194       |
| 228 | A Third Osmosensing Branch in <i>Saccharomyces cerevisiae</i> Requires the Msb2 Protein and Functions in Parallel with the Sho1 Branch. <i>Molecular and Cellular Biology</i> , 2002, 22, 4739-4749.  | 1.1 | 113       |
| 229 | A Specific Structural Requirement for Ergosterol in Long-chain Fatty Acid Synthesis Mutants Important for Maintaining Raft Domains in Yeast. <i>Molecular Biology of the Cell</i> , 2002, 13, 4414-4428.  | 0.9 | 112       |
| 230 | Snf1 Protein Kinase and the Repressors Nrg1 and Nrg2 Regulate FLO11 , Haploid Invasive Growth, and Diploid Pseudohyphal Differentiation. <i>Molecular and Cellular Biology</i> , 2002, 22, 3994-4000.   | 1.1 | 192       |
| 231 | Proteomics of the Eukaryotic Transcription Machinery: Identification of Proteins Associated with Components of Yeast TFIID by Multidimensional Mass Spectrometry. <i>Molecular and Cellular Biology</i> , 2002, 22, 4723-4738.  | 1.1 | 285       |
| 232 | Yeast Protein Kinases and the RHO1 Exchange Factor TUS1 Are Novel Components of the Cell Integrity Pathway in Yeast. <i>Molecular and Cellular Biology</i> , 2002, 22, 1329-1339.   | 1.1 | 127       |
| 233 | Cyclophilin A Peptidyl-Prolyl Isomerase Activity Promotes Zpr1 Nuclear Export. <i>Molecular and Cellular Biology</i> , 2002, 22, 6993-7003.   | 1.1 | 58        |
| 234 | Novel Proteins Linking the Actin Cytoskeleton to the Endocytic Machinery in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2002, 13, 3646-3661.   | 0.9 | 52        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 235 | Saccharomyces cerevisiae C-Type Cyclin Ume3p/Srb11p Is Required for Efficient Induction and Execution of Meiotic Development. Eukaryotic Cell, 2002, 1, 66-74.                                       | 3.4 | 27        |
| 236 | Analysis of the size and shape of protein complexes from yeast. Methods in Enzymology, 2002, 351, 150-168.   | 0.4 | 14        |
| 237 | Erv14p Directs a Transmembrane Secretory Protein into COPII-coated Transport Vesicles. Molecular Biology of the Cell, 2002, 13, 880-891.   | 0.9 | 109       |
| 238 | Intron status and 3'-end formation control cotranscriptional export of mRNA. Genes and Development, 2002, 16, 2761-2766.   | 2.7 | 98        |
| 239 | CAK1 Promotes Meiosis and Spore Formation in Saccharomyces cerevisiae in a CDC28 -Independent Fashion. Molecular and Cellular Biology, 2002, 22, 57-68.  | 1.1 | 43        |
| 240 | Bni5p, a Septin-Interacting Protein, Is Required for Normal Septin Function and Cytokinesis in Saccharomyces cerevisiae. Molecular and Cellular Biology, 2002, 22, 6906-6920.                        | 1.1 | 65        |
| 241 | Manipulation of a Nuclear NAD <sup>+</sup> Salvage Pathway Delays Aging without Altering Steady-state NAD <sup>+</sup> Levels. Journal of Biological Chemistry, 2002, 277, 18881-18890.              | 1.6 | 269       |
| 242 | Regulation of the Premiddle and Middle Phases of Expression of the NDT80 Gene during Sporulation of Saccharomyces cerevisiae. Molecular and Cellular Biology, 2002, 22, 6417-6429.                   | 1.1 | 83        |
| 243 | Mutations in genes of Saccharomyces cerevisiae encoding pre-mRNA splicing factors cause cell cycle arrest through activation of the spindle checkpoint. Nucleic Acids Research, 2002, 30, 4361-4370. | 6.5 | 44        |
| 244 | Direct regulation of Arp2/3 complex activity and function by the actin binding protein coronin. Journal of Cell Biology, 2002, 159, 993-1004.  | 2.3 | 179       |
| 245 | Esc1, a Nuclear Periphery Protein Required for Sir4-Based Plasmid Anchoring and Partitioning. Molecular and Cellular Biology, 2002, 22, 8292-8301.   | 1.1 | 131       |
| 246 | Yeast hnRNP K-Like Genes Are Involved in Regulation of the Telomeric Position Effect and Telomere Length. Molecular and Cellular Biology, 2002, 22, 286-297.   | 1.1 | 39        |
| 247 | ORC and the intra-S-phase checkpoint: a threshold regulates Rad53p activation in S phase. Genes and Development, 2002, 16, 3236-3252.  | 2.7 | 188       |
| 248 | NuA4 Subunit Yng2 Function in Intra-S-Phase DNA Damage Response. Molecular and Cellular Biology, 2002, 22, 8215-8225.  | 1.1 | 117       |
| 249 | The MEK Kinase Ssk2p Promotes Actin Cytoskeleton Recovery After Osmotic Stress. Molecular Biology of the Cell, 2002, 13, 2869-2880.  | 0.9 | 46        |
| 250 | Stu1p Is Physically Associated with $\hat{\gamma}$ -Tubulin and Is Required for Structural Integrity of the Mitotic Spindle. Molecular Biology of the Cell, 2002, 13, 1881-1892.                     | 0.9 | 44        |
| 251 | An active precursor in assembly of yeast nuclear ribonuclease P. Rna, 2002, 8, 1348-1360.  | 1.6 | 30        |
| 252 | Studying yeast vacuoles. Methods in Enzymology, 2002, 351, 408-432.  | 0.4 | 48        |

| #   | ARTICLE   | IF   | CITATIONS |
|-----|---|------|-----------|
| 253 | Functional Assays for Mammalian G-Protein-Coupled Receptors in Yeast. <i>Methods in Enzymology</i> , 2002, 344, 92-111.   | 0.4  | 16        |
| 254 | [14] Glutaredoxins and oxidative stress defense in yeast. <i>Methods in Enzymology</i> , 2002, 348, 136-146.  | 0.4  | 19        |
| 255 | Nuclear protein transport. <i>Methods in Enzymology</i> , 2002, 351, 587-607.   | 0.4  | 37        |
| 256 | Expression of the essential mRNA export factor Yra1p is autoregulated by a splicing-dependent mechanism. <i>Rna</i> , 2002, 8, 969-980.   | 1.6  | 46        |
| 257 | Requirement of Hos2 Histone Deacetylase for Gene Activity in Yeast. <i>Science</i> , 2002, 298, 1412-1414.  | 6.0  | 245       |
| 258 | A role for cell polarity proteins in mitotic exit. <i>EMBO Journal</i> , 2002, 21, 4851-4862.   | 3.5  | 104       |
| 259 | Hmo1, an HMG-box protein, belongs to the yeast ribosomal DNA transcription system. <i>EMBO Journal</i> , 2002, 21, 5498-5507.   | 3.5  | 98        |
| 260 | Separase, Polo Kinase, the Kinetochores Protein Slk19, and Spo12 Function in a Network that Controls Cdc14 Localization during Early Anaphase. <i>Cell</i> , 2002, 108, 207-220.                                | 13.5 | 414       |
| 261 | Vacuole Fusion at a Ring of Vertex Docking Sites Leaves Membrane Fragments within the Organelle. <i>Cell</i> , 2002, 108, 357-369.  | 13.5 | 211       |
| 262 | Chromatin Boundaries in Budding Yeast. <i>Cell</i> , 2002, 109, 551-562.  | 13.5 | 336       |
| 263 | Yph1p, an ORC-Interacting Protein. <i>Cell</i> , 2002, 109, 835-848.  | 13.5 | 172       |
| 264 | Noc3p, a bHLH Protein, Plays an Integral Role in the Initiation of DNA Replication in Budding Yeast. <i>Cell</i> , 2002, 109, 849-860.  | 13.5 | 112       |
| 265 | Phospho-Regulation of Kinetochores-Microtubule Attachments by the Aurora Kinase Ipl1p. <i>Cell</i> , 2002, 111, 163-172.  | 13.5 | 575       |
| 266 | Function and Selectivity of Bromodomains in Anchoring Chromatin-Modifying Complexes to Promoter Nucleosomes. <i>Cell</i> , 2002, 111, 369-379.  | 13.5 | 483       |
| 267 | Supercomplex formation between Mlh1-Mlh3 and Sgs1-Top3 heterocomplexes in meiotic yeast cells. <i>Biochemical and Biophysical Research Communications</i> , 2002, 296, 949-953.                                 | 1.0  | 40        |
| 268 | The ATP-binding cassette (ABC) transporter Bpt1p mediates vacuolar sequestration of glutathione conjugates in yeast. <i>FEBS Letters</i> , 2002, 520, 63-67.  | 1.3  | 78        |
| 269 | Genome-wide Nuclear Morphology Screen Identifies Novel Genes Involved in Nuclear Architecture and Gene-silencing in <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 2002, 321, 551-561. | 2.0  | 38        |
| 270 | Crp1p, A New Cruciform DNA-binding Protein in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 2002, 323, 685-700.   | 2.0  | 8         |



| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 271 | Lsm Proteins Are Required for Normal Processing of Pre-tRNAs and Their Efficient Association with La-Homologous Protein Lhp1p. <i>Molecular and Cellular Biology</i> , 2002, 22, 5248-5256.  | 1.1 | 61        |
| 272 | Myosin-like proteins 1 and 2 are not required for silencing or telomere anchoring, but act in the Tel1 pathway of telomere length control. <i>Journal of Structural Biology</i> , 2002, 140, 79-91.  | 1.3 | 76        |
| 273 | Rad9 Phosphorylation Sites Couple Rad53 to the <i>Saccharomyces cerevisiae</i> DNA Damage Checkpoint. <i>Molecular Cell</i> , 2002, 9, 1055-1065.  | 4.5 | 205       |
| 274 | Two TOR Complexes, Only One of which Is Rapamycin Sensitive, Have Distinct Roles in Cell Growth Control. <i>Molecular Cell</i> , 2002, 10, 457-468.  | 4.5 | 1,685     |
| 275 | The FAD- and O <sub>2</sub> -Dependent Reaction Cycle of Ero1-Mediated Oxidative Protein Folding in the Endoplasmic Reticulum. <i>Molecular Cell</i> , 2002, 10, 983-994.  | 4.5 | 374       |
| 276 | Transient Stability of DNA Ends Allows Nonhomologous End Joining to Precede Homologous Recombination. <i>Molecular Cell</i> , 2002, 10, 1189-1199.   | 4.5 | 176       |
| 277 | Stt4 PI 4-Kinase Localizes to the Plasma Membrane and Functions in the Pkc1-Mediated MAP Kinase Cascade. <i>Developmental Cell</i> , 2002, 2, 593-605.   | 3.1 | 236       |
| 278 | Endosome-Associated Complex, ESCRT-II, Recruits Transport Machinery for Protein Sorting at the Multivesicular Body. <i>Developmental Cell</i> , 2002, 3, 283-289.  | 3.1 | 589       |
| 279 | Studies of Cargo Delivery to the Vacuole Mediated by Autophagosomes in <i>Saccharomyces cerevisiae</i> . <i>Developmental Cell</i> , 2002, 3, 815-824.   | 3.1 | 96        |
| 280 | Rlp7p is associated with 60S preribosomes, restricted to the granular component of the nucleolus, and required for pre-rRNA processing. <i>Journal of Cell Biology</i> , 2002, 157, 941-952.   | 2.3 | 73        |
| 281 | Yeast Ysl2p, Homologous to Sec7 Domain Guanine Nucleotide Exchange Factors, Functions in Endocytosis and Maintenance of Vacuole Integrity and Interacts with the Arf-Like Small GTPase Arl1p. <i>Molecular and Cellular Biology</i> , 2002, 22, 4914-4928. | 1.1 | 42        |
| 282 | Expression and genomic analysis of midasin, a novel and highly conserved AAA protein distantly related to dynein. <i>BMC Genomics</i> , 2002, 3, 18.   | 1.2 | 88        |
| 283 | Cancer therapeutics in yeast. <i>Cancer Cell</i> , 2002, 2, 267-273.   | 7.7 | 39        |
| 284 | Mks1 in Concert with TOR Signaling Negatively Regulates RTC Target Gene Expression in <i>S. cerevisiae</i> . <i>Current Biology</i> , 2002, 12, 389-395.   | 1.8 | 108       |
| 285 | Control of Replication Timing by a Transcriptional Silencer. <i>Current Biology</i> , 2002, 12, 869-875.   | 1.8 | 60        |
| 286 | Live Imaging of Telomeres. <i>Current Biology</i> , 2002, 12, 2076-2089.   | 1.8 | 276       |
| 287 | Control of Lte1 Localization by Cell Polarity Determinants and Cdc14. <i>Current Biology</i> , 2002, 12, 2098-2110.  | 1.8 | 76        |
| 288 | Transcriptional and posttranslational regulation of a membrane nutrient transporter. <i>Biochemistry and Molecular Biology Education</i> , 2002, 30, 388-393.  | 0.5 | 5         |

| #   | ARTICLE   | IF   | CITATIONS |
|-----|---|------|-----------|
| 289 | Mutation of TRS130 , which encodes a component of the TRAPP II complex, activates transcription of OCH1 in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2002, 42, 85-93. | 0.8  | 23        |
| 290 | The vesicular transport protein Cgp1p/Vps54p/Tcs3p/Luv1p is required for the integrity of the actin cytoskeleton. <i>Molecular Genetics and Genomics</i> , 2002, 268, 190-205.          | 1.0  | 10        |
| 291 | Mouse GSPT2, but not GSPT1, can substitute for yeast eRF3in vivo. <i>Genes To Cells</i> , 2002, 7, 1043-1057.   | 0.5  | 28        |
| 292 | Rpa12p, a conserved RNA polymerase I subunit with two functional domains. <i>Molecular Microbiology</i> , 2002, 43, 1105-1113.  | 1.2  | 45        |
| 293 | Depression of <i>Saccharomyces cerevisiae</i> invasive growth on non-glucose carbon sources requires the Snf1 kinase. <i>Molecular Microbiology</i> , 2002, 45, 453-469.                | 1.2  | 46        |
| 294 | Antifungal activity in <i>Saccharomyces cerevisiae</i> is modulated by calcium signalling. <i>Molecular Microbiology</i> , 2002, 46, 257-268.   | 1.2  | 109       |
| 295 | Cdc4 is involved in the transcriptional control of OCH1, a gene encoding $\beta$ -1,6-mannosyltransferase in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2002, 19, 69-77.          | 0.8  | 9         |
| 296 | SRC1: an intron-containing yeast gene involved in sister chromatid segregation. <i>Yeast</i> , 2002, 19, 43-54.   | 0.8  | 45        |
| 297 | Marker-fusion PCR for one-step mutagenesis of essential genes in yeast. <i>Yeast</i> , 2002, 19, 141-149.   | 0.8  | 23        |
| 298 | Genetic interactions link ARF1, YPT31/32 and TRS130. <i>Yeast</i> , 2002, 19, 1075-1086.  | 0.8  | 28        |
| 299 | A large nucleolar U3 ribonucleoprotein required for 18S ribosomal RNA biogenesis. <i>Nature</i> , 2002, 417, 967-970.   | 13.7 | 618       |
| 300 | Protein dislocation from the ER requires polyubiquitination and the AAA-ATPase Cdc48. <i>Nature Cell Biology</i> , 2002, 4, 134-139.  | 4.6  | 489       |
| 301 | Nuclear architecture and spatial positioning help establish transcriptional states of telomeres in yeast. <i>Nature Cell Biology</i> , 2002, 4, 214-221.                                | 4.6  | 253       |
| 302 | Interdependent nuclear accumulation of budding yeast Cdt1 and Mcm2â€™7 during G1 phase. <i>Nature Cell Biology</i> , 2002, 4, 198-207.  | 4.6  | 245       |
| 303 | The DEAD box protein Dhh1 stimulates the decapping enzyme Dcp1. <i>EMBO Journal</i> , 2002, 21, 2788-2797.  | 3.5  | 153       |
| 304 | MRM2 encodes a novel yeast mitochondrial 21S rRNA methyltransferase. <i>EMBO Journal</i> , 2002, 21, 1139-1147.   | 3.5  | 73        |
| 305 | The Khd1 protein, which has three KH RNA-binding motifs, is required for proper localization of ASH1 mRNA in yeast. <i>EMBO Journal</i> , 2002, 21, 1158-1167.                          | 3.5  | 96        |
| 306 | Ccr4p is the catalytic subunit of a Ccr4p/Pop2p/Notp mRNA deadenylase complex in <i>Saccharomyces cerevisiae</i> . <i>EMBO Journal</i> , 2002, 21, 1427-1436.                           | 3.5  | 291       |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 307 | The Paf1 complex physically and functionally associates with transcription elongation factors in vivo. <i>EMBO Journal</i> , 2002, 21, 1764-1774.  | 3.5 | 280       |
| 308 | 60S pre-ribosome formation viewed from assembly in the nucleolus until export to the cytoplasm. <i>EMBO Journal</i> , 2002, 21, 5539-5547.   | 3.5 | 307       |
| 309 | The mRNA export machinery requires the novel Sac3p-Thp1p complex to dock at the nucleoplasmic entrance of the nuclear pores. <i>EMBO Journal</i> , 2002, 21, 5843-5852.  | 3.5 | 238       |
| 310 | Cell cycle-dependent nuclear export of Cdh1p may contribute to the inactivation of APC/CCdh1. <i>EMBO Journal</i> , 2002, 21, 6515-6526.   | 3.5 | 89        |
| 311 | In <i>Saccharomyces cerevisiae</i> , ATP2 mRNA sorting to the vicinity of mitochondria is essential for respiratory function. <i>EMBO Journal</i> , 2002, 21, 6893-6904.   | 3.5 | 98        |
| 312 | The path from nucleolar 90S to cytoplasmic 40S pre-ribosomes. <i>EMBO Journal</i> , 2003, 22, 1370-1380.   | 3.5 | 264       |
| 313 | Systematic analysis of essential yeast TAFs in genome-wide transcription and preinitiation complex assembly. <i>EMBO Journal</i> , 2003, 22, 3395-3402.  | 3.5 | 93        |
| 314 | Elg1 forms an alternative RFC complex important for DNA replication and genome integrity. <i>EMBO Journal</i> , 2003, 22, 4304-4313.   | 3.5 | 148       |
| 315 | Genome-wide location of yeast RNA polymerase III transcription machinery. <i>EMBO Journal</i> , 2003, 22, 4738-4747.   | 3.5 | 145       |
| 316 | PCR-based methods facilitate targeted gene manipulations and cloning procedures. <i>Current Genetics</i> , 2003, 44, 115-123.  | 0.8 | 68        |
| 317 | Mutations in the yeast cyclin-dependent kinase Cdc28 reveal a role in the spindle assembly checkpoint. <i>Molecular Genetics and Genomics</i> , 2003, 269, 672-684.  | 1.0 | 16        |
| 318 | Autocrine activation of the pheromone response pathway in <i>mat<math>\Delta</math>2</i> - cells is attenuated by SST2- and ASG7-dependent mechanisms. <i>Molecular Genetics and Genomics</i> , 2003, 270, 225-233.          | 1.0 | 14        |
| 319 | Activity of phosphoforms and truncated versions of Ndt80, a checkpoint-regulated sporulation-specific transcription factor of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 2003, 270, 324-336. | 1.0 | 18        |
| 320 | A WASp Homolog Powers Actin Polymerization-Dependent Motility of Endosomes In Vivo. <i>Current Biology</i> , 2003, 13, 455-463.  | 1.8 | 53        |
| 321 | Negative Regulation of Yeast WASp by Two SH3 Domain-Containing Proteins. <i>Current Biology</i> , 2003, 13, 1000-1008.   | 1.8 | 138       |
| 322 | Where do all the proteins go?. <i>Targets</i> , 2003, 2, 237-244.  | 0.3 | 3         |
| 323 | Yeast Functional Analysis: Identification of Two Essential Genes Involved in ER to Golgi Trafficking. <i>Traffic</i> , 2003, 4, 607-617.   | 1.3 | 35        |
| 324 | Cell cycle- and age-dependent activation of Sod1p drives the formation of stress resistant cell subpopulations within clonal yeast cultures. <i>Molecular Microbiology</i> , 2003, 50, 857-870.                              | 1.2 | 44        |

| #   | ARTICLE   | IF   | CITATIONS |
|-----|---|------|-----------|
| 325 | The nuclear actin-related protein Act3p/Arp4p of <i>Saccharomyces cerevisiae</i> is involved in transcription regulation of stress genes. <i>Molecular Microbiology</i> , 2003, 50, 1155-1171.                                  | 1.2  | 37        |
| 326 | The upstream regulator, Rsr1p, and downstream effectors, Gic1p and Gic2p, of the Cdc42p small GTPase coordinately regulate initiation of budding in <i>Saccharomyces cerevisiae</i> . <i>Genes To Cells</i> , 2003, 8, 235-250. | 0.5  | 29        |
| 327 | Mos10 (Vps60) is required for normal filament maturation in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2003, 49, 1267-1285.  | 1.2  | 13        |
| 328 | New ?marker swap? plasmids for converting selectable markers on budding yeast gene disruptions and plasmids. <i>Yeast</i> , 2003, 20, 985-993.  | 0.8  | 93        |
| 329 | Four ARF GAPs in <i>Saccharomyces cerevisiae</i> have both overlapping and distinct functions. <i>Yeast</i> , 2003, 20, 315-330.  | 0.8  | 40        |
| 330 | A ?marker switch? approach for targeted mutagenesis of genes in <i>Schizosaccharomyces pombe</i> . <i>Yeast</i> , 2003, 20, 587-594.  | 0.8  | 32        |
| 331 | KIROM2 encodes an essential GEF homologue in <i>Kluyveromyces lactis</i> . <i>Yeast</i> , 2003, 20, 611-624.  | 0.8  | 16        |
| 332 | Construction of a set of <i>Saccharomyces cerevisiae</i> vectors designed for recombinational cloning. <i>Yeast</i> , 2003, 20, 739-746.  | 0.8  | 51        |
| 333 | Global analysis of protein localization in budding yeast. <i>Nature</i> , 2003, 425, 686-691.   | 13.7 | 3,884     |
| 334 | Global analysis of protein expression in yeast. <i>Nature</i> , 2003, 425, 737-741.   | 13.7 | 3,407     |
| 335 | Targets of the cyclin-dependent kinase Cdk1. <i>Nature</i> , 2003, 425, 859-864.  | 13.7 | 835       |
| 336 | Ubp3 requires a cofactor, Bre5, to specifically de-ubiquitinate the COPII protein, Sec23. <i>Nature Cell Biology</i> , 2003, 5, 661-667.  | 4.6  | 146       |
| 337 | Yeast epsin-related proteins required for Golgi endosome traffic define a $\beta$ -adaptn ear-binding motif. <i>Nature Cell Biology</i> , 2003, 5, 77-81.   | 4.6  | 95        |
| 338 | Lem3p Is Essential for the Uptake and Potency of Alkylphosphocholine Drugs, Edelfosine and Miltefosine. <i>Journal of Biological Chemistry</i> , 2003, 278, 36041-36050.  | 1.6  | 115       |
| 339 | Sequential Protein Association with Nascent 60S Ribosomal Particles. <i>Molecular and Cellular Biology</i> , 2003, 23, 4449-4460.   | 1.1  | 180       |
| 340 | Regulation of PI4,5P2 synthesis by nuclear-cytoplasmic shuttling of the Mss4 lipid kinase. <i>EMBO Journal</i> , 2003, 22, 4223-4236.   | 3.5  | 103       |
| 341 | Dual Roles for Spt5 in Pre-mRNA Processing and Transcription Elongation Revealed by Identification of Spt5-Associated Proteins. <i>Molecular and Cellular Biology</i> , 2003, 23, 1368-1378.                                    | 1.1  | 242       |
| 342 | The Importance of Genetic Recombination for Fidelity of Chromosome Pairing in Meiosis. <i>Developmental Cell</i> , 2003, 5, 915-925.  | 3.1  | 134       |

| #   | ARTICLE   | IF   | CITATIONS |
|-----|---|------|-----------|
| 343 | Bsp1p/Ypr171p is an adapter that directly links some synaptojanin family members to the cortical actin cytoskeleton in yeast. <i>FEBS Letters</i> , 2003, 537, 35-41.   | 1.3  | 19        |
| 344 | The yeast deubiquitinating enzyme Ubp16 is anchored to the outer mitochondrial membrane. <i>FEBS Letters</i> , 2003, 549, 135-140.  | 1.3  | 40        |
| 345 | Exit from Exit. <i>Cell</i> , 2003, 112, 697-709.   | 13.5 | 73        |
| 346 | Transcriptional regulation of phosphate-responsive genes in low-affinity phosphate-transporter-defective mutants in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2003, 306, 843-850.        | 1.0  | 55        |
| 347 | Deubiquitination, a New Player in Golgi to Endoplasmic Reticulum Retrograde Transport. <i>Journal of Biological Chemistry</i> , 2003, 278, 51989-51992.   | 1.6  | 68        |
| 348 | Coupling tandem affinity purification and quantitative tyrosine iodination to determine subunit stoichiometry of protein complexes. <i>Methods</i> , 2003, 31, 104-109.   | 1.9  | 8         |
| 349 | The Q Motif. <i>Molecular Cell</i> , 2003, 11, 127-138.   | 4.5  | 303       |
| 350 | RNAse III-Mediated Degradation of Unspliced Pre-mRNAs and Lariat Introns. <i>Molecular Cell</i> , 2003, 11, 1279-1289.  | 4.5  | 133       |
| 351 | TOR and PKA Signaling Pathways Converge on the Protein Kinase Rim15 to Control Entry into G0. <i>Molecular Cell</i> , 2003, 12, 1607-1613.  | 4.5  | 277       |
| 352 | Histone H2A/H2B Dimer Exchange by ATP-Dependent Chromatin Remodeling Activities. <i>Molecular Cell</i> , 2003, 12, 1599-1606.   | 4.5  | 175       |
| 353 | The COP9 signalosome-like complex in <i>S. cerevisiae</i> and links to other PCI complexes. <i>International Journal of Biochemistry and Cell Biology</i> , 2003, 35, 706-715.  | 1.2  | 54        |
| 354 | The Cdc14 Phosphatase and the FEAR Network Control Meiotic Spindle Disassembly and Chromosome Segregation. <i>Developmental Cell</i> , 2003, 4, 711-726.  | 3.1  | 118       |
| 355 | An Mtw1 Complex Promotes Kinetochore Biorientation that Is Monitored by the Ipl1/Aurora Protein Kinase. <i>Developmental Cell</i> , 2003, 5, 735-745.   | 3.1  | 94        |
| 356 | Amino acid changes in Xrs2p, Dun1p, and Rfa2p that remove the preferred targets of the ATM family of protein kinases do not affect DNA repair or telomere length in <i>Saccharomyces cerevisiae</i> . <i>DNA Repair</i> , 2003, 2, 1041-1064. | 1.3  | 45        |
| 357 | Decapping and Decay of Messenger RNA Occur in Cytoplasmic Processing Bodies. <i>Science</i> , 2003, 300, 805-808.   | 6.0  | 1,168     |
| 358 | Nob1p Is Required for Cleavage of the 5' End of 18S rRNA. <i>Molecular and Cellular Biology</i> , 2003, 23, 1798-1807.  | 1.1  | 144       |
| 359 | The Role of Cdc42p GTPase-activating Proteins in Assembly of the Septin Ring in Yeast. <i>Molecular Biology of the Cell</i> , 2003, 14, 4051-4066.  | 0.9  | 221       |
| 360 | Urmlylation: A Ubiquitin-like Pathway that Functions during Invasive Growth and Budding in Yeast. <i>Molecular Biology of the Cell</i> , 2003, 14, 4329-4341.   | 0.9  | 102       |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 361 | A Protein Complex Containing Mdm10p, Mdm12p, and Mmm1p Links Mitochondrial Membranes and DNA to the Cytoskeleton-based Segregation Machinery. <i>Molecular Biology of the Cell</i> , 2003, 14, 4618-4627.  | 0.9 | 242       |
| 362 | Vps20p and Vta1p interact with Vps4p and function in multivesicular body sorting and endosomal transport in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 2003, 116, 3957-3970.   | 1.2 | 101       |
| 363 | The Role of the Polo Kinase Cdc5 in Controlling Cdc14 Localization. <i>Molecular Biology of the Cell</i> , 2003, 14, 4486-4498.  | 0.9 | 84        |
| 364 | Telomeric Protein Distributions and Remodeling Through the Cell Cycle in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2003, 14, 556-570.   | 0.9 | 83        |
| 365 | The budding yeast Ipl1/Aurora protein kinase regulates mitotic spindle disassembly. <i>Journal of Cell Biology</i> , 2003, 160, 329-339.   | 2.3 | 133       |
| 366 | Spindle orientation in <i>Saccharomyces cerevisiae</i> depends on the transport of microtubule ends along polarized actin cables. <i>Journal of Cell Biology</i> , 2003, 161, 483-488.   | 2.3 | 170       |
| 367 | Mutations in the histone fold domain of the TAF12 gene show synthetic lethality with the TAF1 gene lacking the TAF N-terminal domain (TAND) by different mechanisms from those in the SPT15 gene encoding the TATA box-binding protein (TBP). <i>Nucleic Acids Research</i> , 2003, 31, 1261-1274. | 6.5 | 10        |
| 368 | The Transcription Factor Rim101p Governs Ion Tolerance and Cell Differentiation by Direct Repression of the Regulatory Genes NRG1 and SMP1 in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2003, 23, 677-686.   | 1.1 | 239       |
| 369 | Signal-dependent N-CoR Requirement for Repression by the Ski Oncoprotein. <i>Journal of Biological Chemistry</i> , 2003, 278, 24858-24864.   | 1.6 | 41        |
| 370 | Regulation of the Yeast Amphiphysin Homologue Rvs167p by Phosphorylation. <i>Molecular Biology of the Cell</i> , 2003, 14, 3027-3040.  | 0.9 | 33        |
| 371 | Chl4p and Iml3p Are Two New Members of the Budding Yeast Outer Kinetochores. <i>Molecular Biology of the Cell</i> , 2003, 14, 460-476.   | 0.9 | 57        |
| 372 | EAP1, a <i>Candida albicans</i> Gene Involved in Binding Human Epithelial Cells. <i>Eukaryotic Cell</i> , 2003, 2, 1266-1273.  | 3.4 | 168       |
| 373 | A Gradient of Affinity for the Karyopherin Kap95p along the Yeast Nuclear Pore Complex. <i>Journal of Biological Chemistry</i> , 2003, 278, 42699-42709.   | 1.6 | 104       |
| 374 | War1p, a Novel Transcription Factor Controlling Weak Acid Stress Response in Yeast. <i>Molecular and Cellular Biology</i> , 2003, 23, 1775-1785.   | 1.1 | 129       |
| 375 | A Multiplicity of Coactivators Is Required by Gcn4p at Individual Promoters In Vivo. <i>Molecular and Cellular Biology</i> , 2003, 23, 2800-2820.  | 1.1 | 131       |
| 376 | Direct Evidence for a Critical Role of Myosin II in Budding Yeast Cytokinesis and the Evolvability of New Cytokinetic Mechanisms in the Absence of Myosin II. <i>Molecular Biology of the Cell</i> , 2003, 14, 798-809.  | 0.9 | 75        |
| 377 | Recruitment of SWI/SNF by Gcn4p Does Not Require Snf2p or Gcn5p but Depends Strongly on SWI/SNF Integrity, SRB Mediator, and SAGA. <i>Molecular and Cellular Biology</i> , 2003, 23, 8829-8845.  | 1.1 | 52        |
| 378 | Gts1p Activates SNF1-dependent Derepression of HSP104 and TPS1 in the Stationary Phase of Yeast Growth. <i>Journal of Biological Chemistry</i> , 2003, 278, 29760-29768.   | 1.6 | 12        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 379 | The roles of endonucleolytic cleavage and exonucleolytic digestion in the 5'-end processing of <i>S. cerevisiae</i> box C/D snoRNAs. <i>Rna</i> , 2003, 9, 1362-1370.  | 1.6 | 47        |
| 380 | Grr1-dependent Inactivation of Mth1 Mediates Glucose-induced Dissociation of Rgt1 from HXT Gene Promoters. <i>Molecular Biology of the Cell</i> , 2003, 14, 3230-3241.   | 0.9 | 138       |
| 381 | Chemical Genetic Analysis of Apg1 Reveals A Non-kinase Role in the Induction of Autophagy. <i>Molecular Biology of the Cell</i> , 2003, 14, 477-490.   | 0.9 | 152       |
| 382 | She4p/Dim1p Interacts with the Motor Domain of Unconventional Myosins in the Budding Yeast, <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2003, 14, 2237-2249.                                    | 0.9 | 65        |
| 383 | Molecular Dissection of a Yeast Septin: Distinct Domains Are Required for Septin Interaction, Localization, and Function. <i>Molecular and Cellular Biology</i> , 2003, 23, 2762-2777.                                       | 1.1 | 170       |
| 384 | Far3 and Five Interacting Proteins Prevent Premature Recovery from Pheromone Arrest in the Budding Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2003, 23, 1750-1763.                      | 1.1 | 101       |
| 385 | Loss of Sin3/Rpd3 Histone Deacetylase Restores the DNA Damage Response in Checkpoint-Deficient Strains of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2003, 23, 4522-4531.                     | 1.1 | 51        |
| 386 | Synthetic Lethal Interaction of the Mitochondrial Phosphatidylethanolamine Biosynthetic Machinery with the Prohibitin Complex of <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2003, 14, 370-383. | 0.9 | 87        |
| 387 | Identification and Characterization of a Novel RanGTP-binding Protein in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2003, 278, 15397-15405.  | 1.6 | 10        |
| 388 | RNA-protein interactions promote asymmetric sorting of the ASH1 mRNA ribonucleoprotein complex. <i>Rna</i> , 2003, 9, 1383-1399.   | 1.6 | 34        |
| 389 | Sac3 Is an mRNA Export Factor That Localizes to Cytoplasmic Fibrils of Nuclear Pore Complex. <i>Molecular Biology of the Cell</i> , 2003, 14, 836-847.   | 0.9 | 65        |
| 390 | Vps51 Is Part of the Yeast Vps Fifty-three Tethering Complex Essential for Retrograde Traffic from the Early Endosome and Cvt Vesicle Completion. <i>Journal of Biological Chemistry</i> , 2003, 278, 5009-5020.             | 1.6 | 91        |
| 391 | Use of a Genetically Introduced Cross-linker to Identify Interaction Sites of Acidic Activators within Native Transcription Factor IID and SAGA. <i>Journal of Biological Chemistry</i> , 2003, 278, 6779-6786.              | 1.6 | 35        |
| 392 | Ykt6p Is a Multifunctional Yeast R-SNARE That Is Required for Multiple Membrane Transport Pathways to the Vacuole. <i>Molecular Biology of the Cell</i> , 2003, 14, 1868-1881.   | 0.9 | 75        |
| 393 | Hierarchy of protein assembly at the vertex ring domain for yeast vacuole docking and fusion. <i>Journal of Cell Biology</i> , 2003, 160, 365-374.   | 2.3 | 126       |
| 394 | An Rpb4/Rpb7-Like Complex in Yeast RNA Polymerase III Contains the Orthologue of Mammalian CGRP-RCP. <i>Molecular and Cellular Biology</i> , 2003, 23, 195-205.  | 1.1 | 46        |
| 395 | Mitotic Exit Regulation through Distinct Domains within the Protein Kinase Cdc15. <i>Molecular and Cellular Biology</i> , 2003, 23, 5018-5030.   | 1.1 | 42        |
| 396 | Functional interactions between the transcription and mRNA 3' end processing machineries mediated by Ssu72 and Sub1. <i>Genes and Development</i> , 2003, 17, 1030-1042.   | 2.7 | 127       |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 397 | Histone H1 Is Required for Proper Regulation of Pyruvate Decarboxylase Gene Expression in <i>Neurospora crassa</i> . <i>Eukaryotic Cell</i> , 2003, 2, 341-350.   | 3.4 | 37        |
| 398 | Topological analysis of a plant vacuolar Na <sup>+</sup> /H <sup>+</sup> antiporter reveals a luminal C terminus that regulates antiporter cation selectivity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 12510-12515. | 3.3 | 161       |
| 399 | Pleiotropic Effects of Ubp6 Loss on Drug Sensitivities and Yeast Prion Are Due to Depletion of the Free Ubiquitin Pool. <i>Journal of Biological Chemistry</i> , 2003, 278, 52102-52115.  | 1.6 | 102       |
| 400 | An Early Function during Transcription for the Yeast mRNA Export Factor Dbp5p/Rat8p Suggested by Its Genetic and Physical Interactions with Transcription Factor IIH Components. <i>Molecular Biology of the Cell</i> , 2003, 14, 1664-1676.                                    | 0.9 | 86        |
| 401 | Yeast recombination pathways triggered by topoisomerase II-mediated DNA breaks. <i>Nucleic Acids Research</i> , 2003, 31, 4373-4384.  | 6.5 | 36        |
| 402 | Association of the RENT complex with nontranscribed and coding regions of rDNA and a regional requirement for the replication fork block protein Fob1 in rDNA silencing. <i>Genes and Development</i> , 2003, 17, 2162-2176.  | 2.7 | 203       |
| 403 | Spatial Organization of Protein-RNA Interactions in the Branch Site-3' Splice Site Region during pre-mRNA Splicing in Yeast. <i>Molecular and Cellular Biology</i> , 2003, 23, 4174-4186.   | 1.1 | 64        |
| 404 | Cdc50p, a Conserved Endosomal Membrane Protein, Controls Polarized Growth in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2003, 14, 730-747.  | 0.9 | 46        |
| 405 | Sec3p Is Needed for the Spatial Regulation of Secretion and for the Inheritance of the Cortical Endoplasmic Reticulum. <i>Molecular Biology of the Cell</i> , 2003, 14, 4770-4782.  | 0.9 | 104       |
| 406 | Widespread cytoplasmic mRNA transport in yeast: Identification of 22 bud-localized transcripts using DNA microarray analysis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 11429-11434.                                  | 3.3 | 257       |
| 407 | A Role for <i>Saccharomyces cerevisiae</i> Cul8 Ubiquitin Ligase in Proper Anaphase Progression. <i>Journal of Biological Chemistry</i> , 2003, 278, 22828-22837.   | 1.6 | 45        |
| 408 | Yeast Kinetochores Do Not Stabilize Stu2p-dependent Spindle Microtubule Dynamics. <i>Molecular Biology of the Cell</i> , 2003, 14, 4181-4195.   | 0.9 | 75        |
| 409 | The GAP activity of Msb3p and Msb4p for the Rab GTPase Sec4p is required for efficient exocytosis and actin organization. <i>Journal of Cell Biology</i> , 2003, 162, 635-646.  | 2.3 | 77        |
| 410 | Set2-Catalyzed Methylation of Histone H3 Represses Basal Expression of GAL4 in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2003, 23, 5972-5978.   | 1.1 | 59        |
| 411 | Tor Kinases Are in Distinct Membrane-associated Protein Complexes in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2003, 14, 1204-1220.  | 0.9 | 222       |
| 412 | A novel and essential mechanism determining specificity and activity of protein phosphatase 2A (PP2A) in vivo. <i>Genes and Development</i> , 2003, 17, 2138-2150.  | 2.7 | 89        |
| 413 | RAM: A Conserved Signaling Network That Regulates Ace2p Transcriptional Activity and Polarized Morphogenesis. <i>Molecular Biology of the Cell</i> , 2003, 14, 3782-3803.   | 0.9 | 203       |
| 414 | New Findings on Interactions among the Yeast Oligosaccharyl Transferase Subunits Using a Chemical Cross-linker. <i>Journal of Biological Chemistry</i> , 2003, 278, 33078-33087.  | 1.6 | 29        |



| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 415 | Transcriptional activation via sequential histone H2B ubiquitylation and deubiquitylation, mediated by SAGA-associated Ubp8. <i>Genes and Development</i> , 2003, 17, 2648-2663.  | 2.7 | 598       |
| 416 | Ask10p Mediates the Oxidative Stress-Induced Destruction of the <i>Saccharomyces cerevisiae</i> C-Type Cyclin Ume3p/Srb11p. <i>Eukaryotic Cell</i> , 2003, 2, 962-970.  | 3.4 | 31        |
| 417 | A Noc Complex Specifically Involved in the Formation and Nuclear Export of Ribosomal 40 S Subunits. <i>Journal of Biological Chemistry</i> , 2003, 278, 4072-4081.  | 1.6 | 110       |
| 418 | YMR313c/TGL3 Encodes a Novel Triacylglycerol Lipase Located in Lipid Particles of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2003, 278, 23317-23323.                                      | 1.6 | 177       |
| 419 | Synthetic Lethal Analysis Implicates Ste20p, a p21-activated Protein Kinase, in Polarisome Activation. <i>Molecular Biology of the Cell</i> , 2003, 14, 1501-1516.  | 0.9 | 88        |
| 420 | G1 Transcription Factors Are Differentially Regulated in <i>Saccharomyces cerevisiae</i> by the Swi6-Binding Protein Stb1. <i>Molecular and Cellular Biology</i> , 2003, 23, 5064-5077.                                   | 1.1 | 60        |
| 421 | Vps51p Mediates the Association of the GARP (Vps52/53/54) Complex with the Late Golgi t-SNARE Tlg1p. <i>Molecular Biology of the Cell</i> , 2003, 14, 1610-1623.  | 0.9 | 162       |
| 422 | Control of landmark events in meiosis by the CDK Cdc28 and the meiosis-specific kinase Ime2. <i>Genes and Development</i> , 2003, 17, 1524-1539.  | 2.7 | 260       |
| 423 | Biochemical Characterization of the Ran-RanBP1-RanGAP System: Are RanBP Proteins and the Acidic Tail of RanGAP Required for the Ran-RanGAP GTPase Reaction?. <i>Molecular and Cellular Biology</i> , 2003, 23, 8124-8136. | 1.1 | 48        |
| 424 | Use of Modular Substrates Demonstrates Mechanistic Diversity and Reveals Differences in Chaperone Requirement of ERAD. <i>Journal of Biological Chemistry</i> , 2003, 278, 35903-35913.                                   | 1.6 | 169       |
| 425 | Oma1, a Novel Membrane-bound Metallopeptidase in Mitochondria with Activities Overlapping with the m-AAA Protease. <i>Journal of Biological Chemistry</i> , 2003, 278, 46414-46423.                                       | 1.6 | 135       |
| 426 | Phosphorelay-Regulated Degradation of the Yeast Ssk1p Response Regulator by the Ubiquitin-Proteasome System. <i>Molecular and Cellular Biology</i> , 2003, 23, 6662-6671.   | 1.1 | 51        |
| 427 | Interaction with Tap42 Is Required for the Essential Function of Sit4 and Type 2A Phosphatases. <i>Molecular Biology of the Cell</i> , 2003, 14, 4342-4351.   | 0.9 | 74        |
| 428 | The Putative GTPases Nog1p and Lsg1p Are Required for 60S Ribosomal Subunit Biogenesis and Are Localized to the Nucleus and Cytoplasm, Respectively. <i>Molecular and Cellular Biology</i> , 2003, 23, 4344-4355.         | 1.1 | 126       |
| 429 | SWI/SNF-dependent chromatin remodeling of RNR3 requires TAFII and the general transcription machinery. <i>Genes and Development</i> , 2003, 17, 502-515.  | 2.7 | 74        |
| 430 | Intravacuolar Membrane Lysis in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2003, 278, 7810-7821.  | 1.6 | 59        |
| 431 | The Rtf1 Component of the Paf1 Transcriptional Elongation Complex Is Required for Ubiquitination of Histone H2B. <i>Journal of Biological Chemistry</i> , 2003, 278, 33625-33628.   | 1.6 | 250       |
| 432 | Identification of p21-Activated Kinase Specificity Determinants in Budding Yeast: a Single Amino Acid Substitution Imparts Ste20 Specificity to Cla4. <i>Molecular and Cellular Biology</i> , 2003, 23, 1569-1580.        | 1.1 | 13        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 433 | An N <sup>ε</sup> -Acetyltransferase Responsible for Acetylation of the N-terminal Residues of Histones H4 and H2A. <i>Journal of Biological Chemistry</i> , 2003, 278, 38109-38112.   | 1.6 | 115       |
| 434 | In yeast, the pseudohyphal phenotype induced by isoamyl alcohol results from the operation of the morphogenesis checkpoint. <i>Journal of Cell Science</i> , 2003, 116, 3423-3431.   | 1.2 | 43        |
| 435 | Attachment of the Ubiquitin-Related Protein Urm1p to the Antioxidant Protein Ahp1p. <i>Eukaryotic Cell</i> , 2003, 2, 930-936.   | 3.4 | 98        |
| 436 | Atg23 Is Essential for the Cytoplasm to Vacuole Targeting Pathway and Efficient Autophagy but Not Pexophagy. <i>Journal of Biological Chemistry</i> , 2003, 278, 48445-48452.  | 1.6 | 74        |
| 437 | Yaf9, a Novel NuA4 Histone Acetyltransferase Subunit, Is Required for the Cellular Response to Spindle Stress in Yeast. <i>Molecular and Cellular Biology</i> , 2003, 23, 6086-6102.   | 1.1 | 92        |
| 438 | mRNAs Encoding Telomerase Components and Regulators Are Controlled by UPF Genes in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2003, 2, 134-142.  | 3.4 | 62        |
| 439 | Lysine-79 of histone H3 is hypomethylated at silenced loci in yeast and mammalian cells: A potential mechanism for position-effect variegation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 1820-1825. | 3.3 | 278       |
| 440 | Bro1 is an endosome-associated protein that functions in the MVB pathway in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 2003, 116, 1893-1903.   | 1.2 | 189       |
| 441 | The intramitochondrial dynamin-related GTPase, Mgm1p, is a component of a protein complex that mediates mitochondrial fusion. <i>Journal of Cell Biology</i> , 2003, 160, 303-311.   | 2.3 | 221       |
| 442 | Ady4p and Spo74p Are Components of the Meiotic Spindle Pole Body That Promote Growth of the Prospore Membrane in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2003, 2, 431-445.  | 3.4 | 48        |
| 443 | Myo4p and She3p are required for cortical ER inheritance in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 2003, 163, 1255-1266.   | 2.3 | 168       |
| 444 | Cdc48p is required for the cell cycle commitment point at Start via degradation of the G1-CDK inhibitor Far1p. <i>Journal of Cell Biology</i> , 2003, 163, 21-26.  | 2.3 | 69        |
| 445 | Mrc1 is a replication fork component whose phosphorylation in response to DNA replication stress activates Rad53. <i>Genes and Development</i> , 2003, 17, 1755-1767.  | 2.7 | 332       |
| 446 | FHA Domain-Mediated DNA Checkpoint Regulation of Rad53. <i>Cell Cycle</i> , 2003, 2, 381-394.  | 1.3 | 48        |
| 447 | Interaction of Gal4p with Components of Transcription Machinery In Vivo. <i>Methods in Enzymology</i> , 2003, 370, 445-454.  | 0.4 | 39        |
| 448 | Functional Analysis of TFIID Components Using Conditional Mutants. <i>Methods in Enzymology</i> , 2003, 370, 415-430.  | 0.4 | 12        |
| 449 | LST8 negatively regulates amino acid biosynthesis as a component of the TOR pathway. <i>Journal of Cell Biology</i> , 2003, 161, 333-347.  | 2.3 | 126       |
| 450 | Monitoring Changes in the Subcellular Location of Proteins in <i>S. cerevisiae</i> . , 2004, 241, 299-312.   |     | 1         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 451 | Gcn4p and Novel Upstream Activating Sequences Regulate Targets of the Unfolded Protein Response. <i>PLoS Biology</i> , 2004, 2, e246.  | 2.6 | 143       |
| 452 | Heterodimeric complexes of Hop2 and Mnd1 function with Dmc1 to promote meiotic homolog juxtaposition and strand assimilation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004, 101, 10572-10577.                     | 3.3 | 110       |
| 453 | Mmm2p, a mitochondrial outer membrane protein required for yeast mitochondrial shape and maintenance of mtDNA nucleoids. <i>Journal of Cell Biology</i> , 2004, 164, 677-688.  | 2.3 | 136       |
| 454 | Requirement for Bbp1p in the Proper Mitotic Functions of Cdc5p in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2004, 15, 1711-1723.  | 0.9 | 15        |
| 455 | Identification of a Distinctive Mutation Spectrum Associated with High Levels of Transcription in Yeast. <i>Molecular and Cellular Biology</i> , 2004, 24, 4801-4809.  | 1.1 | 21        |
| 456 | GTP Hydrolysis by eRF3 Facilitates Stop Codon Decoding during Eukaryotic Translation Termination. <i>Molecular and Cellular Biology</i> , 2004, 24, 7769-7778.   | 1.1 | 160       |
| 457 | Interspore bridges: a new feature of the <i>Saccharomyces cerevisiae</i> spore wall. <i>Microbiology (United Kingdom)</i> , 2004, 154, 107-115.  | 0.7 | 54        |
| 458 | Sister-chromatid cohesion mediated by the alternative RF-Ctf18/Dcc1/Ctf8, the helicase Chl1 and the polymerase- $\alpha$ -associated protein Ctf4 is essential for chromatid disjunction during meiosis II. <i>Journal of Cell Science</i> , 2004, 117, 3547-3559. | 1.2 | 130       |
| 459 | The ctf13-30/CTF13 Genomic Haploinsufficiency Modifier Screen Identifies the Yeast Chromatin Remodeling Complex RSC, Which Is Required for the Establishment of Sister Chromatid Cohesion. <i>Molecular and Cellular Biology</i> , 2004, 24, 1232-1244.            | 1.1 | 132       |
| 460 | Positive and Negative Regulation of a SNARE Protein by Control of Intracellular Localization. <i>Molecular Biology of the Cell</i> , 2004, 15, 1802-1815.  | 0.9 | 168       |
| 461 | Fus1p Interacts With Components of the Hog1p Mitogen-Activated Protein Kinase and Cdc42p Morphogenesis Signaling Pathways to Control Cell Fusion During Yeast Mating. <i>Genetics</i> , 2004, 166, 67-77.  | 1.2 | 60        |
| 462 | Cbf1p Is Required for Chromatin Remodeling at Promoter-proximal CACGTG Motifs in Yeast. <i>Journal of Biological Chemistry</i> , 2004, 279, 27116-27123.   | 1.6 | 53        |
| 463 | A Type V Myosin (Myo2p) and a Rab-like G-Protein (Ypt11p) Are Required for Retention of Newly Inherited Mitochondria in Yeast Cells during Cell Division. <i>Molecular Biology of the Cell</i> , 2004, 15, 3994-4002.  | 0.9 | 101       |
| 464 | Changes in the Localization of the <i>Saccharomyces cerevisiae</i> Anaphase-Promoting Complex Upon Microtubule Depolymerization and Spindle Checkpoint Activation. <i>Genetics</i> , 2004, 167, 1079-1094.   | 1.2 | 18        |
| 465 | Regulation of Spindle Pole Function by an Intermediary Metabolite. <i>Molecular Biology of the Cell</i> , 2004, 15, 2606-2616.   | 0.9 | 23        |
| 466 | GPI7 Involved in Glycosylphosphatidylinositol Biosynthesis Is Essential for Yeast Cell Separation. <i>Journal of Biological Chemistry</i> , 2004, 279, 51869-51879.  | 1.6 | 40        |
| 467 | The Arf activator Gea2p and the P-type ATPase Drs2p interact at the Golgi in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 2004, 117, 711-722.  | 1.2 | 97        |
| 468 | The External Amino Acid Signaling Pathway Promotes Activation of Stp1 and Uga35/Dal81 Transcription Factors for Induction of the AGP1 Gene in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2004, 166, 1727-1739.  | 1.2 | 45        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 469 | Diminished S-Phase Cyclin-Dependent Kinase Function Elicits Vital Rad53-Dependent Checkpoint Responses in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2004, 24, 10208-10222.  | 1.1 | 41        |
| 470 | The Soh1/MED31 Protein Is an Ancient Component of <i>Schizosaccharomyces pombe</i> and <i>Saccharomyces cerevisiae</i> Mediator. <i>Journal of Biological Chemistry</i> , 2004, 279, 49455-49459.   | 1.6 | 39        |
| 471 | TOR Complex 1 Includes a Novel Component, Tco89p (YPL180w), and Cooperates with Ssd1p to Maintain Cellular Integrity in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2004, 279, 14752-14762.                              | 1.6 | 218       |
| 472 | The Protein Kinase Snf1 Is Required for Tolerance to the Ribonucleotide Reductase Inhibitor Hydroxyurea. <i>Molecular and Cellular Biology</i> , 2004, 24, 2560-2572.   | 1.1 | 46        |
| 473 | Physical interactions between the Alg1, Alg2, and Alg11 mannosyltransferases of the endoplasmic reticulum. <i>Glycobiology</i> , 2004, 14, 559-570.   | 1.3 | 70        |
| 474 | Activation of the DNA Damage Checkpoint in Yeast Lacking the Histone Chaperone Anti-Silencing Function 1. <i>Molecular and Cellular Biology</i> , 2004, 24, 10313-10327.  | 1.1 | 90        |
| 475 | The NEF4 Complex Regulates Rad4 Levels and Utilizes Snf2/Swi2-Related ATPase Activity for Nucleotide Excision Repair. <i>Molecular and Cellular Biology</i> , 2004, 24, 6362-6378.  | 1.1 | 56        |
| 476 | Deficiencies in the Endoplasmic Reticulum (ER)-Membrane Protein Gab1p Perturb Transfer of Glycosylphosphatidylinositol to Proteins and Cause Perinuclear ER-associated Actin Bar Formation. <i>Molecular Biology of the Cell</i> , 2004, 15, 2758-2770. | 0.9 | 29        |
| 477 | Sgt1p and Skp1p Modulate the Assembly and Turnover of CBF3 Complexes Required for Proper Kinetochores Function. <i>Molecular Biology of the Cell</i> , 2004, 15, 3366-3378.   | 0.9 | 43        |
| 478 | Arginine methyltransferase affects interactions and recruitment of mRNA processing and export factors. <i>Genes and Development</i> , 2004, 18, 2024-2035.  | 2.7 | 119       |
| 479 | Novel regulation of mitotic exit by the Cdc42 effectors Gic1 and Gic2. <i>Journal of Cell Biology</i> , 2004, 164, 219-231.   | 2.3 | 41        |
| 480 | The post-replication repair RAD18 and RAD6 genes are involved in the prevention of spontaneous mutations caused by 7,8-dihydro-8-oxoguanine in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2004, 32, 5003-5010.                   | 6.5 | 48        |
| 481 | Glycerophosphoinositol, a Novel Phosphate Source Whose Transport Is Regulated by Multiple Factors in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2004, 279, 31937-31942.   | 1.6 | 33        |
| 482 | The Small Subunit Processome Is Required for Cell Cycle Progression at G1. <i>Molecular Biology of the Cell</i> , 2004, 15, 5038-5046.  | 0.9 | 68        |
| 483 | In <i>Candida albicans</i> , the Nim1 kinases Gin4 and Hsl1 negatively regulate pseudohypha formation and Gin4 also controls septin organization. <i>Journal of Cell Biology</i> , 2004, 164, 581-591.  | 2.3 | 96        |
| 484 | Analysis of Snu13p mutations reveals differential interactions with the U4 snRNA and U3 snoRNA. <i>Rna</i> , 2004, 10, 308-320.   | 1.6 | 25        |
| 485 | The Differential Roles of Budding Yeast Tem1p, Cdc15p, and Bub2p Protein Dynamics in Mitotic Exit. <i>Molecular Biology of the Cell</i> , 2004, 15, 1519-1532.  | 0.9 | 89        |
| 486 | Distinct Roles for the Hsp40 and Hsp90 Molecular Chaperones during Cystic Fibrosis Transmembrane Conductance Regulator Degradation in Yeast. <i>Molecular Biology of the Cell</i> , 2004, 15, 4787-4797.  | 0.9 | 149       |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 487 | Neuropathy Target Esterase and Its Yeast Homologue Degrade Phosphatidylcholine to Glycerophosphocholine in Living Cells. <i>Journal of Biological Chemistry</i> , 2004, 279, 24024-24033.                           | 1.6 | 192       |
| 488 | Rea1, a Dynein-related Nuclear AAA-ATPase, Is Involved in Late rRNA Processing and Nuclear Export of 60 S Subunits. <i>Journal of Biological Chemistry</i> , 2004, 279, 55411-55418.                                | 1.6 | 63        |
| 489 | Identification and functional analysis of the essential and regulatory light chains of the only type II myosin Myo1p in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 2004, 165, 843-855.      | 2.3 | 58        |
| 490 | An Array of Coactivators Is Required for Optimal Recruitment of TATA Binding Protein and RNA Polymerase II by Promoter-Bound Gcn4p. <i>Molecular and Cellular Biology</i> , 2004, 24, 4104-4117.                    | 1.1 | 83        |
| 491 | Five Genes Involved in Biosynthesis of the Pyruvylated Gal $\beta$ 1,3-Epitope in <i>Schizosaccharomyces pombe</i> N-Linked Glycans. <i>Journal of Biological Chemistry</i> , 2004, 279, 35644-35655.               | 1.6 | 37        |
| 492 | The Essential ATP-binding Cassette Protein RLI1 Functions in Translation by Promoting Preinitiation Complex Assembly. <i>Journal of Biological Chemistry</i> , 2004, 279, 42157-42168.                              | 1.6 | 155       |
| 493 | Genetic and Biochemical Interactions Among Yar1, Ltv1 and RpS3 Define Novel Links Between Environmental Stress and Ribosome Biogenesis in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2004, 168, 1877-1889. | 1.2 | 54        |
| 494 | Nrg1 and Nrg2 Transcriptional Repressors Are Differently Regulated in Response to Carbon Source. <i>Eukaryotic Cell</i> , 2004, 3, 311-317.   | 3.4 | 34        |
| 495 | Mhr1p-dependent Concatemeric Mitochondrial DNA Formation for Generating Yeast Mitochondrial Homoplasmic Cells. <i>Molecular Biology of the Cell</i> , 2004, 15, 310-322.  | 0.9 | 63        |
| 496 | Binding of the Replication Terminator Protein Fob1p to the Ter Sites of Yeast Causes Polar Fork Arrest. <i>Journal of Biological Chemistry</i> , 2004, 279, 1932-1941.  | 1.6 | 53        |
| 497 | Counting of Rif1p and Rif2p on <i>Saccharomyces cerevisiae</i> Telomeres Regulates Telomere Length. <i>Molecular and Cellular Biology</i> , 2004, 24, 10857-10867.  | 1.1 | 118       |
| 498 | Cyclin B-Cdk Activity Stimulates Meiotic Rereplication in Budding Yeast. <i>Genetics</i> , 2004, 167, 1621-1628.  | 1.2 | 11        |
| 499 | In vivo target of a transcriptional activator revealed by fluorescence resonance energy transfer. <i>Genes and Development</i> , 2004, 18, 333-343.   | 2.7 | 165       |
| 500 | Interactions among Rax1p, Rax2p, Bud8p, and Bud9p in Marking Cortical Sites for Bipolar Bud-site Selection in Yeast. <i>Molecular Biology of the Cell</i> , 2004, 15, 5145-5157.                                    | 0.9 | 48        |
| 501 | Enhanced Membrane Fusion in Sterol-enriched Vacuoles Bypasses the Vrp1p Requirement. <i>Molecular Biology of the Cell</i> , 2004, 15, 4609-4621.  | 0.9 | 24        |
| 502 | The Amino Terminus of the <i>Saccharomyces cerevisiae</i> DNA Helicase Rrm3p Modulates Protein Function Altering Replication and Checkpoint Activity. <i>Genetics</i> , 2004, 168, 1205-1218.                       | 1.2 | 21        |
| 503 | Integration of Stress Responses: Modulation of Calcineurin Signaling in <i>Saccharomyces cerevisiae</i> by Protein Kinase A. <i>Eukaryotic Cell</i> , 2004, 3, 1147-1153.   | 3.4 | 79        |
| 504 | Gyp5p and Gyl1p are involved in the control of polarized exocytosis in budding yeast. <i>Journal of Cell Science</i> , 2004, 117, 4757-4767.  | 1.2 | 22        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 505 | Chl1p, a DNA Helicase-Like Protein in Budding Yeast, Functions in Sister-Chromatid Cohesion. <i>Genetics</i> , 2004, 166, 33-42.   | 1.2 | 139       |
| 506 | Npa1p, a Component of Very Early Pre-60S Ribosomal Particles, Associates with a Subset of Small Nucleolar RNPs Required for Peptidyl Transferase Center Modification. <i>Molecular and Cellular Biology</i> , 2004, 24, 6324-6337. | 1.1 | 71        |
| 507 | TOR Controls Transcriptional and Translational Programs via Sap-Sit4 Protein Phosphatase Signaling Effectors. <i>Molecular and Cellular Biology</i> , 2004, 24, 8332-8341.   | 1.1 | 74        |
| 508 | The Interaction between Sgt1p and Skp1p Is Regulated by HSP90 Chaperones and Is Required for Proper CBF3 Assembly. <i>Molecular and Cellular Biology</i> , 2004, 24, 8938-8950.  | 1.1 | 67        |
| 509 | Novel Functional Dissection of the Localization-Specific Roles of Budding Yeast Polo Kinase Cdc5p. <i>Molecular and Cellular Biology</i> , 2004, 24, 9873-9886.  | 1.1 | 34        |
| 510 | Yeast Chromatin Assembly Complex 1 Protein Excludes Nonacetylatable Forms of Histone H4 from Chromatin and the Nucleus. <i>Molecular and Cellular Biology</i> , 2004, 24, 10180-10192.   | 1.1 | 38        |
| 511 | Posttranscriptional Regulation of HO Expression by the Mkt1-Pbp1 Complex. <i>Molecular and Cellular Biology</i> , 2004, 24, 3670-3681.   | 1.1 | 52        |
| 512 | The GTPase-Activating Enzyme Gyp1p Is Required for Recycling of Internalized Membrane Material by Inactivation of the Rab/Ypt GTPase Ypt1p. <i>Molecular and Cellular Biology</i> , 2004, 24, 3815-3826.                           | 1.1 | 30        |
| 513 | Identification of Protein Complexes Required for Efficient Sister Chromatid Cohesion. <i>Molecular Biology of the Cell</i> , 2004, 15, 1736-1745.  | 0.9 | 221       |
| 514 | Cystic Fibrosis Transmembrane Conductance Regulator Degradation Depends on the Lectins Htm1p/EDEM and the Cdc48 Protein Complex in Yeast. <i>Molecular Biology of the Cell</i> , 2004, 15, 4125-4135.                              | 0.9 | 90        |
| 515 | Pxl1p, a Paxillin-like Protein in <i>Saccharomyces cerevisiae</i> , May Coordinate Cdc42p and Rho1p Functions during Polarized Growth. <i>Molecular Biology of the Cell</i> , 2004, 15, 3977-3985.                                 | 0.9 | 30        |
| 516 | <i>Saccharomyces cerevisiae</i> JEN1 Promoter Activity Is Inversely Related to Concentration of Repressing Sugar. <i>Applied and Environmental Microbiology</i> , 2004, 70, 8-17.  | 1.4 | 32        |
| 517 | Control of Yeast Filamentous-Form Growth by Modules in an Integrated Molecular Network. <i>Genome Research</i> , 2004, 14, 380-390.  | 2.4 | 78        |
| 518 | The Yeast Pdr15p ATP-binding Cassette (ABC) Protein Is a General Stress Response Factor Implicated in Cellular Detoxification. <i>Journal of Biological Chemistry</i> , 2004, 279, 11593-11599.                                    | 1.6 | 53        |
| 519 | The Yeast Vacuolar Proton-translocating ATPase Contains a Subunit Homologous to the <i>Manduca sexta</i> and Bovine e Subunits That Is Essential for Function. <i>Journal of Biological Chemistry</i> , 2004, 279, 17361-17365.    | 1.6 | 66        |
| 520 | Akr1p-dependent Palmitoylation of Yck2p Yeast Casein Kinase 1 Is Necessary and Sufficient for Plasma Membrane Targeting. <i>Journal of Biological Chemistry</i> , 2004, 279, 27138-27147.  | 1.6 | 59        |
| 521 | Functional Characterization of Pwp2, a WD Family Protein Essential for the Assembly of the 90 S Pre-ribosomal Particle. <i>Journal of Biological Chemistry</i> , 2004, 279, 37385-37397.   | 1.6 | 76        |
| 522 | Live cell imaging of the assembly, disassembly, and actin cable-dependent movement of endosomes and actin patches in the budding yeast, <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 2004, 167, 519-530.     | 2.3 | 166       |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 523 | Sgt1 Associates with Hsp90: an Initial Step of Assembly of the Core Kinetochore Complex. <i>Molecular and Cellular Biology</i> , 2004, 24, 8069-8079.  | 1.1 | 77        |
| 524 | Multivesicular Body-ESCRT Components Function in pH Response Regulation in <i>Saccharomyces cerevisiae</i> and <i>Candida albicans</i> . <i>Molecular Biology of the Cell</i> , 2004, 15, 5528-5537.   | 0.9 | 155       |
| 525 | Stress Induction and Mitochondrial Localization of Oxr1 Proteins in Yeast and Humans. <i>Molecular and Cellular Biology</i> , 2004, 24, 3180-3187.   | 1.1 | 97        |
| 526 | Essential Role for the Myotubularin-related Phosphatase Ymr1p and the Synaptojanin-like Phosphatases Sjl2p and Sjl3p in Regulation of Phosphatidylinositol 3-Phosphate in Yeast. <i>Molecular Biology of the Cell</i> , 2004, 15, 3567-3579. | 0.9 | 79        |
| 527 | The Omp85 family of proteins is essential for outer membrane biogenesis in mitochondria and bacteria. <i>Journal of Cell Biology</i> , 2004, 164, 19-24.   | 2.3 | 335       |
| 528 | Direct Sorting of the Yeast Uracil Permease to the Endosomal System Is Controlled by Uracil Binding and Rsp5p-dependent Ubiquitylation. <i>Molecular Biology of the Cell</i> , 2004, 15, 883-895.  | 0.9 | 108       |
| 529 | Septin Ring Assembly Requires Concerted Action of Polarisome Components, a PAK Kinase Cla4p, and the Actin Cytoskeleton in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2004, 15, 5329-5345.                     | 0.9 | 94        |
| 530 | Spt10-Dependent Transcriptional Activation in <i>Saccharomyces cerevisiae</i> Requires both the Spt10 Acetyltransferase Domain and Spt21. <i>Molecular and Cellular Biology</i> , 2004, 24, 135-143.   | 1.1 | 34        |
| 531 | The Novel Yeast PAS Kinase Rim15 Orchestrates G0-Associated Antioxidant Defense Mechanisms. <i>Cell Cycle</i> , 2004, 3, 460-466.  | 1.3 | 154       |
| 532 | A synthetic A tail rescues yeast nuclear accumulation of a ribozyme-terminated transcript. <i>Rna</i> , 2004, 10, 1888-1899.   | 1.6 | 107       |
| 533 | The Type 1 Phosphatase Reg1p-Glc7p Is Required for the Glucose-induced Degradation of Fructose-1,6-bisphosphatase in the Vacuole. <i>Journal of Biological Chemistry</i> , 2004, 279, 9713-9724.   | 1.6 | 22        |
| 534 | Telomere Cap Components Influence the Rate of Senescence in Telomerase-Deficient Yeast Cells. <i>Molecular and Cellular Biology</i> , 2004, 24, 837-845.   | 1.1 | 33        |
| 535 | The Small-Subunit Processome Is a Ribosome Assembly Intermediate. <i>Eukaryotic Cell</i> , 2004, 3, 1619-1626.   | 3.4 | 152       |
| 536 | Role for Nhp6, Gcn5, and the Swi/Snf Complex in Stimulating Formation of the TATA-Binding Protein-TFIID-DNA Complex. <i>Molecular and Cellular Biology</i> , 2004, 24, 8312-8321.  | 1.1 | 34        |
| 537 | The yeast Apq12 protein affects nucleocytoplasmic mRNA transport. <i>Rna</i> , 2004, 10, 1352-1358.  | 1.6 | 18        |
| 538 | The Deubiquitinating Enzyme Doa4p Protects Cells from DNA Topoisomerase I Poisons. <i>Journal of Biological Chemistry</i> , 2004, 279, 21271-21281.  | 1.6 | 23        |
| 539 | Clb6/Cdc28 and Cdc14 Regulate Phosphorylation Status and Cellular Localization of Swi6. <i>Molecular and Cellular Biology</i> , 2004, 24, 2277-2285.   | 1.1 | 62        |
| 540 | Activation of the RAS/Cyclic AMP Pathway Suppresses a TOR Deficiency in Yeast. <i>Molecular and Cellular Biology</i> , 2004, 24, 338-351.  | 1.1 | 239       |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 541 | Applicability of Tandem Affinity Purification MudPIT to Pathway Proteomics in Yeast. <i>Molecular and Cellular Proteomics</i> , 2004, 3, 226-237.  | 2.5 | 130       |
| 542 | The Budding Yeast Mei5 and Sae3 Proteins Act Together With Dmc1 During Meiotic Recombination. <i>Genetics</i> , 2004, 168, 1219-1230.  | 1.2 | 83        |
| 543 | Stm1p, a G4 Quadruplex and Purine Motif Triplex Nucleic Acid-binding Protein, Interacts with Ribosomes and Subtelomeric Yâ€² DNA in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2004, 279, 24323-24333.                     | 1.6 | 37        |
| 544 | Degradation of the Gluconeogenic Enzymes Fructose-1,6-bisphosphatase and Malate Dehydrogenase Is Mediated by Distinct Proteolytic Pathways and Signaling Events. <i>Journal of Biological Chemistry</i> , 2004, 279, 49138-49150.                          | 1.6 | 100       |
| 545 | The Role of Hot13p and Redox Chemistry in the Mitochondrial TIM22 Import Pathway. <i>Journal of Biological Chemistry</i> , 2004, 279, 43744-43751.   | 1.6 | 78        |
| 546 | ASH1 mRNA Anchoring Requires Reorganization of the Myo4p-She3p-She2p Transport Complex. <i>Journal of Biological Chemistry</i> , 2004, 279, 46286-46294.   | 1.6 | 21        |
| 547 | The Origin Recognition Complex and Sir4 Protein Recruit Sir1p to Yeast Silent Chromatin through Independent Interactions Requiring a Common Sir1p Domain. <i>Molecular and Cellular Biology</i> , 2004, 24, 774-786.                                       | 1.1 | 64        |
| 548 | Identification of Edc3p as an Enhancer of mRNA Decapping in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2004, 166, 729-739.  | 1.2 | 135       |
| 549 | The Presequence Translocase-associated Protein Import Motor of Mitochondria. <i>Journal of Biological Chemistry</i> , 2004, 279, 38047-38054.  | 1.6 | 109       |
| 550 | Histone Fold Protein Dls1p Is Required for Isw2-Dependent Chromatin Remodeling In Vivo. <i>Molecular and Cellular Biology</i> , 2004, 24, 2605-2613.   | 1.1 | 44        |
| 551 | Two-hybrid Mpp10p interaction-defective Imp4 proteins are not interaction defective in vivo but do confer specific pre-rRNA processing defects in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2004, 32, 1404-1413.                   | 6.5 | 19        |
| 552 | Budding Yeast Silencing Complexes and Regulation of Sir2 Activity by Protein-Protein Interactions. <i>Molecular and Cellular Biology</i> , 2004, 24, 6931-6946.  | 1.1 | 73        |
| 553 | Yeast ARL1 encodes a regulator of K <sup>+</sup> influx. <i>Journal of Cell Science</i> , 2004, 117, 2309-2320.  | 1.2 | 37        |
| 554 | Distinct Machinery Is Required in <i>Saccharomyces cerevisiae</i> for the Endoplasmic Reticulum-associated Degradation of a Multispanning Membrane Protein and a Soluble Luminal Protein. <i>Journal of Biological Chemistry</i> , 2004, 279, 38369-38378. | 1.6 | 232       |
| 555 | Tor Signaling and Nutrient-based Signals Converge on Mks1p Phosphorylation to Regulate Expression of Rtg1p- and Rtg3p-dependent Target Genes. <i>Journal of Biological Chemistry</i> , 2004, 279, 46527-46535.   | 1.6 | 80        |
| 556 | On the Mechanism of Constitutive Pdr1 Activator-mediated PDR5 Transcription in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2004, 279, 42677-42686.  | 1.6 | 34        |
| 557 | Pil1p and Lsp1p Negatively Regulate the 3-Phosphoinositide-dependent Protein Kinase-like Kinase Pkh1p and Downstream Signaling Pathways Pkc1p and Ypk1p. <i>Journal of Biological Chemistry</i> , 2004, 279, 22030-22038.                                  | 1.6 | 99        |
| 558 | YBP1 and Its Homologue YBP2/YBH1 Influence Oxidative-Stress Tolerance by Nonidentical Mechanisms in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2004, 3, 318-330.   | 3.4 | 29        |



| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 559 | Regulation and Recognition of SCF Grr1 Targets in the Glucose and Amino Acid Signaling Pathways. <i>Molecular and Cellular Biology</i> , 2004, 24, 8994-9005.  | 1.1 | 77        |
| 560 | Anatomy and Dynamics of DNA Replication Fork Movement in Yeast Telomeric Regions. <i>Molecular and Cellular Biology</i> , 2004, 24, 4019-4031.   | 1.1 | 133       |
| 561 | Importance of the Sir3 N Terminus and Its Acetylation for Yeast Transcriptional Silencing. <i>Genetics</i> , 2004, 168, 547-551.   | 1.2 | 68        |
| 562 | A Yeast Mitochondrial Membrane Methyltransferase-like Protein Can Compensate for oxa1 Mutations. <i>Journal of Biological Chemistry</i> , 2004, 279, 47464-47472.  | 1.6 | 38        |
| 563 | Rad6 plays a role in transcriptional activation through ubiquitylation of histone H2B. <i>Genes and Development</i> , 2004, 18, 184-195.   | 2.7 | 186       |
| 564 | Negative Regulation of Phosphatidylinositol 4,5-Bisphosphate Levels by the INP51-associated Proteins TAX4 and IRS4. <i>Journal of Biological Chemistry</i> , 2004, 279, 39604-39610.   | 1.6 | 20        |
| 565 | The <i>Caulobacter crescentus</i> CgtA C Protein Cosediments with the Free 50S Ribosomal Subunit. <i>Journal of Bacteriology</i> , 2004, 186, 481-489.   | 1.0 | 66        |
| 566 | Autonomous Function of the Amino-Terminal Inhibitory Domain of TAF1 in Transcriptional Regulation. <i>Molecular and Cellular Biology</i> , 2004, 24, 3089-3099.  | 1.1 | 10        |
| 567 | A high resolution protein interaction map of the yeast Mediator complex. <i>Nucleic Acids Research</i> , 2004, 32, 5379-5391.  | 6.5 | 192       |
| 568 | Mutant Telomere Sequences Lead to Impaired Chromosome Separation and a Unique Checkpoint Response. <i>Molecular Biology of the Cell</i> , 2004, 15, 1623-1634.   | 0.9 | 39        |
| 569 | Pak1 Protein Kinase Regulates Activation and Nuclear Localization of Snf1-Gal83 Protein Kinase. <i>Molecular and Cellular Biology</i> , 2004, 24, 8255-8263.   | 1.1 | 82        |
| 570 | Atg21 Is a Phosphoinositide Binding Protein Required for Efficient Lipidation and Localization of Atg8 during Uptake of Amino-peptidase I by Selective Autophagy. <i>Molecular Biology of the Cell</i> , 2004, 15, 3553-3566.      | 0.9 | 203       |
| 571 | Vesicles carry most exocyst subunits to exocytic sites marked by the remaining two subunits, Sec3p and Exo70p. <i>Journal of Cell Biology</i> , 2004, 167, 889-901.  | 2.3 | 270       |
| 572 | Cdc50p, a Protein Required for Polarized Growth, Associates with the Drs2p P-Type ATPase Implicated in Phospholipid Translocation in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2004, 15, 3418-3432. | 0.9 | 238       |
| 573 | A Ddc2-Rad53 Fusion Protein Can Bypass the Requirements for RAD9 and MRC1 in Rad53 Activation. <i>Molecular Biology of the Cell</i> , 2004, 15, 5443-5455.   | 0.9 | 27        |
| 574 | Gid8p (Dcr1p) and Dcr2p Function in a Common Pathway To Promote START Completion in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2004, 3, 1627-1638.   | 3.4 | 15        |
| 575 | The Sec1/Munc18 Protein, Vps33p, Functions at the Endosome and the Vacuole of <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2004, 15, 2593-2605.  | 0.9 | 52        |
| 576 | CTD kinase I is involved in RNA polymerase I transcription. <i>Nucleic Acids Research</i> , 2004, 32, 5851-5860.   | 6.5 | 19        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 577 | The Yeast Elongator Histone Acetylase Requires Sit4-dependent Dephosphorylation for Toxin-Target Capacity. <i>Molecular Biology of the Cell</i> , 2004, 15, 1459-1469.                              | 0.9 | 57        |
| 578 | Gene Recruitment of the Activated INO1 Locus to the Nuclear Membrane. <i>PLoS Biology</i> , 2004, 2, e342.  | 2.6 | 357       |
| 579 | Stable and Dynamic Axes of Polarity Use Distinct Formin Isoforms in Budding Yeast. <i>Molecular Biology of the Cell</i> , 2004, 15, 4971-4989.  | 0.9 | 142       |
| 580 | Antagonistic Roles of ESCRT and Vps Class C/HOPS Complexes in the Recycling of Yeast Membrane Proteins. <i>Molecular Biology of the Cell</i> , 2004, 15, 4203-4214.                                 | 0.9 | 45        |
| 581 | The Ubiquitin Ligase Rsp5p is Required for Modification and Sorting of Membrane Proteins into Multivesicular Bodies. <i>Traffic</i> , 2004, 5, 383-392.   | 1.3 | 61        |
| 582 | Two genes encoding <i>Arabidopsis halleri</i> MTP1 metal transport proteins co-segregate with zinc tolerance and account for high MTP1 transcript levels. <i>Plant Journal</i> , 2004, 39, 425-439. | 2.8 | 274       |
| 583 | Dissection of septin actin interactions using actin overexpression in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2004, 53, 469-483.  | 1.2 | 22        |
| 584 | Translation termination factors function outside of translation: yeast eRF1 interacts with myosin light chain, Mlc1p, to effect cytokinesis. <i>Molecular Microbiology</i> , 2004, 53, 687-696.     | 1.2 | 31        |
| 585 | Rck1 and Rck2 MAPKAP kinases and the HOG pathway are required for oxidative stress resistance. <i>Molecular Microbiology</i> , 2004, 53, 1743-1756.   | 1.2 | 162       |
| 586 | The Function of the Endocytic Scaffold Protein Pan1p Depends on Multiple Domains. <i>Traffic</i> , 2004, 5, 963-978.  | 1.3 | 23        |
| 587 | Late-G1 cyclin-CDK activity is essential for control of cell morphogenesis in budding yeast. <i>Nature Cell Biology</i> , 2004, 6, 59-66.   | 4.6 | 81        |
| 588 | GGA proteins bind ubiquitin to facilitate sorting at the trans-Golgi network. <i>Nature Cell Biology</i> , 2004, 6, 252-259.  | 4.6 | 155       |
| 589 | Golgi targeting of ARF-like GTPase Arl3p requires its N <sup>1</sup> -acetylation and the integral membrane protein Sys1p. <i>Nature Cell Biology</i> , 2004, 6, 414-419.                           | 4.6 | 168       |
| 590 | The SNARE Ykt6 mediates protein palmitoylation during an early stage of homotypic vacuole fusion. <i>EMBO Journal</i> , 2004, 23, 45-53.  | 3.5 | 72        |
| 591 | Physical and functional interactions between nucleotide excision repair and DNA damage checkpoint. <i>EMBO Journal</i> , 2004, 23, 429-438.   | 3.5 | 112       |
| 592 | Ubiquitin interactions of NZF zinc fingers. <i>EMBO Journal</i> , 2004, 23, 1411-1421.  | 3.5 | 238       |
| 593 | Separation of silencing from perinuclear anchoring functions in yeast Ku80, Sir4 and Esc1 proteins. <i>EMBO Journal</i> , 2004, 23, 1301-1312.  | 3.5 | 237       |
| 594 | Genome-wide lethality screen identifies new PI4,5P2 effectors that regulate the actin cytoskeleton. <i>EMBO Journal</i> , 2004, 23, 3747-3757.  | 3.5 | 124       |

| #   | ARTICLE  | IF   | CITATIONS |
|-----|--|------|-----------|
| 595 | Inactivation of the RRB1-Pescadillo pathway involved in ribosome biogenesis induces chromosomal instability. <i>Oncogene</i> , 2004, 23, 8597-8602.  | 2.6  | 61        |
| 596 | The conserved kinetochore protein shugoshin protects centromeric cohesion during meiosis. <i>Nature</i> , 2004, 427, 510-517.  | 13.7 | 523       |
| 597 | PMT family of <i>Candida albicans</i> : five protein mannosyltransferase isoforms affect growth, morphogenesis and antifungal resistance. <i>Molecular Microbiology</i> , 2004, 55, 546-560. | 1.2  | 149       |
| 598 | Genetic rescue of a <i>Toxoplasma gondii</i> conditional cell cycle mutant. <i>Molecular Microbiology</i> , 2004, 55, 1060-1071.   | 1.2  | 28        |
| 599 | The histones H2A/H2B and H3/H4 are imported into the yeast nucleus by different mechanisms. <i>European Journal of Cell Biology</i> , 2004, 83, 511-520.                                     | 1.6  | 30        |
| 600 | The Replication Fork Block Protein Fob1 Functions as a Negative Regulator of the FEAR Network. <i>Current Biology</i> , 2004, 14, 467-480.   | 1.8  | 56        |
| 601 | Spo13 Maintains Centromeric Cohesion and Kinetochore Coorientation during Meiosis I. <i>Current Biology</i> , 2004, 14, 2168-2182.   | 1.8  | 80        |
| 602 | Heat shock-induced degradation of Msn2p, a <i>Saccharomyces cerevisiae</i> transcription factor, occurs in the nucleus. <i>Molecular Genetics and Genomics</i> , 2004, 272, 353-362.         | 1.0  | 26        |
| 603 | Genetic aspects of targeted insertion mutagenesis in yeasts. <i>FEMS Microbiology Reviews</i> , 2004, 28, 201-223.   | 3.9  | 36        |
| 604 | Der1p, a protein required for degradation of malformed soluble proteins of the endoplasmic reticulum: topology and Der1-like proteins. <i>FEMS Yeast Research</i> , 2004, 4, 721-729.        | 1.1  | 74        |
| 605 | A new enrichment approach identifies genes that alter cell cycle progression in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2004, 45, 350-359.                               | 0.8  | 17        |
| 606 | Regulation by phosphorylation of Pho81p, a cyclin-dependent kinase inhibitor in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2004, 46, 10-9.                                  | 0.8  | 11        |
| 607 | Hym1p affects cell cycle progression in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2004, 46, 183-192.   | 0.8  | 8         |
| 608 | Optimized cassettes for fluorescent protein tagging in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2004, 21, 661-670.   | 0.8  | 639       |
| 609 | A versatile toolbox for PCR-based tagging of yeast genes: new fluorescent proteins, more markers and promoter substitution cassettes. <i>Yeast</i> , 2004, 21, 947-962.                      | 0.8  | 1,837     |
| 610 | pDUAL, a multipurpose, multicopy vector capable of chromosomal integration in fission yeast. <i>Yeast</i> , 2004, 21, 1289-1305.   | 0.8  | 119       |
| 611 | Redundancy in the function of mitochondrial phosphate transport in <i>Saccharomyces cerevisiae</i> and <i>Arabidopsis thaliana</i> . <i>Molecular Microbiology</i> , 2004, 51, 307-317.      | 1.2  | 75        |
| 612 | Hif1 Is a Component of Yeast Histone Acetyltransferase B, a Complex Mainly Localized in the Nucleus. <i>Journal of Biological Chemistry</i> , 2004, 279, 16033-16043.                        | 1.6  | 88        |

| #   | ARTICLE   | IF   | CITATIONS |
|-----|---|------|-----------|
| 613 | Role of the yeast Rrp1 protein in the dynamics of pre-ribosome maturation. <i>Rna</i> , 2004, 10, 813-827.  | 1.6  | 89        |
| 614 | Global Analyses of Sumoylated Proteins in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2004, 279, 32262-32268.  | 1.6  | 284       |
| 615 | The <i>S. cerevisiae</i> HtrA-like protein Nma111p is a nuclear serine protease that mediates yeast apoptosis. <i>Journal of Cell Science</i> , 2004, 117, 115-126.                   | 1.2  | 191       |
| 616 | Functional specialization within a vesicle tethering complex. <i>Journal of Cell Biology</i> , 2004, 167, 875-887.  | 2.3  | 75        |
| 617 | Mapping Global Histone Acetylation Patterns to Gene Expression. <i>Cell</i> , 2004, 117, 721-733.   | 13.5 | 561       |
| 618 | Cln3 Activates G1-Specific Transcription via Phosphorylation of the SBF Bound Repressor Whi5. <i>Cell</i> , 2004, 117, 887-898.   | 13.5 | 373       |
| 619 | TOR Regulates Ribosomal Protein Gene Expression via PKA and the Forkhead Transcription Factor FHL1. <i>Cell</i> , 2004, 119, 969-979.   | 13.5 | 418       |
| 620 | Fus3-Regulated Tec1 Degradation through SCFCdc4 Determines MAPK Signaling Specificity during Mating in Yeast. <i>Cell</i> , 2004, 119, 981-990.                                       | 13.5 | 133       |
| 621 | Identification and characterization of the rlp1+, the novel Rad51 paralog in the fission yeast <i>Schizosaccharomyces pombe</i> . <i>DNA Repair</i> , 2004, 3, 1363-1374.             | 1.3  | 22        |
| 622 | DNA decay and limited Rad53 activation after liquid holding of UV-treated nucleotide excision repair deficient <i>S. cerevisiae</i> cells. <i>DNA Repair</i> , 2004, 3, 1591-1599.    | 1.3  | 14        |
| 623 | Regulatory mechanisms controlling biogenesis of ubiquitin and the proteasome. <i>FEBS Letters</i> , 2004, 567, 259-264.   | 1.3  | 79        |
| 624 | A genome-wide screen identifies Yos9p as essential for ER-associated degradation of glycoproteins. <i>FEBS Letters</i> , 2004, 577, 422-426.  | 1.3  | 117       |
| 625 | GFP as a tool to analyze the organization, dynamics and function of nuclei and microtubules in <i>Neurospora crassa</i> . <i>Fungal Genetics and Biology</i> , 2004, 41, 897-910.     | 0.9  | 306       |
| 626 | Chromatin Disassembly Mediated by the Histone Chaperone Asf1 Is Essential for Transcriptional Activation of the Yeast PHO5 and PHO8 Genes. <i>Molecular Cell</i> , 2004, 14, 657-666. | 4.5  | 275       |
| 627 | Targeted mRNA Degradation by Deadenylation-Independent Decapping. <i>Molecular Cell</i> , 2004, 15, 5-15.   | 4.5  | 140       |
| 628 | A Pre-Ribosome with a Tadpole-like Structure Functions in ATP-Dependent Maturation of 60S Subunits. <i>Molecular Cell</i> , 2004, 15, 295-301.  | 4.5  | 62        |
| 629 | Expanded CAG Repeats Activate the DNA Damage Checkpoint Pathway. <i>Molecular Cell</i> , 2004, 15, 287-293.   | 4.5  | 69        |
| 630 | Binding of Chromatin-Modifying Activities to Phosphorylated Histone H2A at DNA Damage Sites. <i>Molecular Cell</i> , 2004, 16, 979-990.   | 4.5  | 513       |

| #   | ARTICLE  | IF   | CITATIONS |
|-----|--|------|-----------|
| 631 | Mcm10 and Cdc45 Cooperate in Origin Activation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 2004, 340, 195-202.  | 2.0  | 78        |
| 632 | The Atg1-Atg13 Complex Regulates Atg9 and Atg23 Retrieval Transport from the Pre-Autophagosomal Structure. <i>Developmental Cell</i> , 2004, 6, 79-90.   | 3.1  | 429       |
| 633 | The Glc7p Nuclear Phosphatase Promotes mRNA Export by Facilitating Association of Mex67p with mRNA. <i>Molecular Cell</i> , 2004, 13, 201-212.   | 4.5  | 162       |
| 634 | The Nuclear Hat1p/Hat2p Complex. <i>Molecular Cell</i> , 2004, 14, 195-205.  | 4.5  | 121       |
| 635 | The Carboxy-Terminal Extension of Yeast Ribosomal Protein S14 Is Necessary for Maturation of 43S Preribosomes. <i>Molecular Cell</i> , 2004, 14, 331-342.  | 4.5  | 83        |
| 636 | Nuclear Retention of Unspliced mRNAs in Yeast Is Mediated by Perinuclear Mlp1. <i>Cell</i> , 2004, 116, 63-73.   | 13.5 | 310       |
| 637 | Functional roles for evolutionarily conserved Spt4p at centromeres and heterochromatin in <i>Saccharomyces cerevisiae</i> . <i>EMBO Journal</i> , 2004, 23, 1804-1814.   | 3.5  | 49        |
| 638 | The C-terminus of Bfa1p in budding yeast is essential to induce mitotic arrest in response to diverse checkpoint-activating signals. <i>Genes To Cells</i> , 2004, 9, 399-418.   | 0.5  | 10        |
| 639 | An Lsm2-Lsm7 Complex in <i>Saccharomyces cerevisiae</i> Associates with the Small Nucleolar RNA snR5. <i>Molecular Biology of the Cell</i> , 2004, 15, 2842-2852.  | 0.9  | 41        |
| 640 | Nuclear Pre-mRNA Decapping and 5' Degradation in Yeast Require the Lsm2-8p Complex. <i>Molecular and Cellular Biology</i> , 2004, 24, 9646-9657.   | 1.1  | 58        |
| 641 | Living recombinant <i>Saccharomyces cerevisiae</i> secreting proteins or peptides as a new drug delivery system in the gut. <i>Journal of Biotechnology</i> , 2004, 110, 37-49.  | 1.9  | 31        |
| 642 | Acyl-CoA-binding protein, Acb1p, is required for normal vacuole function and ceramide synthesis in <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 2004, 380, 907-918.                                    | 1.7  | 73        |
| 643 | A novel role of Mgm1p, a dynamin-related GTPase, in ATP synthase assembly and cristae formation/maintenance. <i>Biochemical Journal</i> , 2004, 381, 19-23.  | 1.7  | 71        |
| 644 | Gene CATCHR--Gene Cloning And Tagging for <i>Caenorhabditis elegans</i> using yeast Homologous Recombination: a novel approach for the analysis of gene expression. <i>Nucleic Acids Research</i> , 2005, 33, e163-e163. | 6.5  | 18        |
| 645 | Subcellular localization and functional expression of the glycerol uptake protein 1 (GUP1) of <i>Saccharomyces cerevisiae</i> tagged with green fluorescent protein. <i>Biochemical Journal</i> , 2005, 390, 145-155.    | 1.7  | 21        |
| 646 | Slx4 becomes phosphorylated after DNA damage in a Mec1/Tel1-dependent manner and is required for repair of DNA alkylation damage. <i>Biochemical Journal</i> , 2005, 391, 325-333.                                       | 1.7  | 45        |
| 647 | Two-step method for constructing unmarked insertions, deletions and allele substitutions in the yeast genome. <i>FEMS Microbiology Letters</i> , 2005, 248, 31-36.   | 0.7  | 28        |
| 648 | Identification of <i>Candida albicans</i> Genes that Induce <i>Saccharomyces cerevisiae</i> Cell Adhesion and Morphogenesis. <i>Biotechnology Progress</i> , 2005, 21, 1601-1609.  | 1.3  | 26        |

| #   | ARTICLE  | IF   | CITATIONS |
|-----|--|------|-----------|
| 649 | <i>Aspergillus nidulans</i> HOG pathway is activated only by two-component signalling pathway in response to osmotic stress. <i>Molecular Microbiology</i> , 2005, 56, 1246-1261.  | 1.2  | 174       |
| 650 | Swi/SNF-GCN5-dependent chromatin remodelling determines induced expression of GDH3, one of the paralogous genes responsible for ammonium assimilation and glutamate biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2005, 57, 291-305. | 1.2  | 33        |
| 651 | The HEAT repeat protein Blm10 regulates the yeast proteasome by capping the core particle. <i>Nature Structural and Molecular Biology</i> , 2005, 12, 294-303.   | 3.6  | 176       |
| 652 | Ubx2 links the Cdc48 complex to ER-associated protein degradation. <i>Nature Cell Biology</i> , 2005, 7, 993-998.  | 4.6  | 257       |
| 653 | The roles of MAD1, MAD2 and MAD3 in meiotic progression and the segregation of nonexchange chromosomes. <i>Nature Genetics</i> , 2005, 37, 756-760.  | 9.4  | 72        |
| 654 | Mechanistically distinct roles for Sgs1p in checkpoint activation and replication fork maintenance. <i>EMBO Journal</i> , 2005, 24, 405-417.   | 3.5  | 132       |
| 655 | Central role of Lfh1p-Fhl1p interaction in the synthesis of yeast ribosomal proteins. <i>EMBO Journal</i> , 2005, 24, 533-542.   | 3.5  | 161       |
| 656 | The yeast EDC1 mRNA undergoes deadenylation-independent decapping stimulated by Not2p, Not4p, and Not5p. <i>EMBO Journal</i> , 2005, 24, 1033-1045.  | 3.5  | 80        |
| 657 | Histone H3 phosphorylation can promote TBP recruitment through distinct promoter-specific mechanisms. <i>EMBO Journal</i> , 2005, 24, 997-1008.  | 3.5  | 89        |
| 658 | Sec17p and HOPS, in distinct SNARE complexes, mediate SNARE complex disruption or assembly for fusion. <i>EMBO Journal</i> , 2005, 24, 1775-1786.  | 3.5  | 104       |
| 659 | The Rsp5 ubiquitin ligase is coupled to and antagonized by the Ubp2 deubiquitinating enzyme. <i>EMBO Journal</i> , 2005, 24, 2414-2424.  | 3.5  | 138       |
| 660 | Rap1 prevents telomere fusions by nonhomologous end joining. <i>EMBO Journal</i> , 2005, 24, 3117-3127.  | 3.5  | 149       |
| 661 | The Bud14p-Glc7p complex functions as a cortical regulator of dynein in budding yeast. <i>EMBO Journal</i> , 2005, 24, 3000-3011.  | 3.5  | 36        |
| 662 | Regulation of G0 entry by the Pho80-Pho85 cyclin-CDK complex. <i>EMBO Journal</i> , 2005, 24, 4271-4278.   | 3.5  | 135       |
| 663 | Telomerase limits the extent of base pairing between template RNA and telomeric DNA. <i>EMBO Reports</i> , 2005, 6, 361-366.   | 2.0  | 61        |
| 664 | Cyclin specificity in the phosphorylation of cyclin-dependent kinase substrates. <i>Nature</i> , 2005, 434, 104-108.   | 13.7 | 343       |
| 665 | Endonucleolytic processing of covalent protein-linked DNA double-strand breaks. <i>Nature</i> , 2005, 436, 1053-1057.  | 13.7 | 536       |
| 666 | ERS1 encodes a functional homologue of the human lysosomal cystine transporter. <i>FEBS Journal</i> , 2005, 272, 2497-2511.  | 2.2  | 45        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 667 | Identification of ATP-NADH kinase isozymes and their contribution to supply of NADP(H) in <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 2005, 272, 3337-3349.   | 2.2 | 69        |
| 668 | Replication-Independent Histone Deposition by the HIR Complex and Asf1. <i>Current Biology</i> , 2005, 15, 2044-2049.   | 1.8 | 189       |
| 669 | Biochemical and genetic analysis of the distinct proliferating cell nuclear antigens of <i>Toxoplasma gondii</i> . <i>Molecular and Biochemical Parasitology</i> , 2005, 142, 56-65.  | 0.5 | 22        |
| 670 | Construction, Verification and Experimental Use of Two Epitope-Tagged Collections of Budding Yeast Strains. <i>Comparative and Functional Genomics</i> , 2005, 6, 2-16.   | 2.0 | 80        |
| 671 | A role for KEM1 at the START of the cell cycle in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2005, 48, 300-309.  | 0.8 | 2         |
| 672 | A set of epitope-tagging integration vectors for functional analysis in. <i>FEMS Yeast Research</i> , 2005, 5, 943-950.   | 1.1 | 9         |
| 673 | Biochemical and genetic characterization of Yra1p in budding yeast. <i>Yeast</i> , 2005, 22, 43-56.   | 0.8 | 23        |
| 674 | New modules for the repeated internal and N-terminal epitope tagging of genes in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2005, 22, 1-12.   | 0.8 | 108       |
| 675 | Svf1 inhibits reactive oxygen species generation and promotes survival under conditions of oxidative stress in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2005, 22, 641-652.  | 0.8 | 27        |
| 676 | Three novel antibiotic marker cassettes for gene disruption and marker switching in <i>Schizosaccharomyces pombe</i> . <i>Yeast</i> , 2005, 22, 1013-1019.  | 0.8 | 244       |
| 677 | Additional vectors for PCR-based gene tagging in <i>Saccharomyces cerevisiae</i> and <i>Schizosaccharomyces pombe</i> using nourseothricin resistance. <i>Yeast</i> , 2005, 22, 1061-1068.  | 0.8 | 92        |
| 678 | Disruption of LRG1 inhibits mother-daughter separation in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2005, 22, 1117-1132.   | 0.8 | 10        |
| 680 | Yos1p Is a Novel Subunit of the Yip1p/Yif1p Complex and Is Required for Transport between the Endoplasmic Reticulum and the Golgi Complex. <i>Molecular Biology of the Cell</i> , 2005, 16, 1673-1683.                                | 0.9 | 51        |
| 681 | The Mitotic Cyclins Clb2p and Clb4p Affect Morphogenesis in <i>Candida albicans</i> . <i>Molecular Biology of the Cell</i> , 2005, 16, 3387-3400.   | 0.9 | 90        |
| 682 | The Pleckstrin Homology Domain Proteins Slm1 and Slm2 Are Required for Actin Cytoskeleton Organization in Yeast and Bind Phosphatidylinositol-4,5-Bisphosphate and TORC2. <i>Molecular Biology of the Cell</i> , 2005, 16, 1883-1900. | 0.9 | 120       |
| 683 | Yeast Recombination Enhancer Is Stimulated by Transcription Activation. <i>Molecular and Cellular Biology</i> , 2005, 25, 7976-7987.  | 1.1 | 18        |
| 684 | Cyc2p, a Membrane-bound Flavoprotein Involved in the Maturation of Mitochondrial c-Type Cytochromes. <i>Journal of Biological Chemistry</i> , 2005, 280, 39852-39859.   | 1.6 | 42        |
| 685 | De Novo Kinetochores Assembly Requires the Centromeric Histone H3 Variant. <i>Molecular Biology of the Cell</i> , 2005, 16, 5649-5660.  | 0.9 | 67        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 686 | A Role for Jsn1p in Recruiting the Arp2/3 Complex to Mitochondria in Budding Yeast. <i>Molecular Biology of the Cell</i> , 2005, 16, 5094-5102.   | 0.9 | 50        |
| 687 | The Ccr4-Not Complex Independently Controls both Msn2-Dependent Transcriptional Activation via a Newly Identified Glc7/Bud14 Type I Protein Phosphatase Module and TFIIID Promoter Distribution. <i>Molecular and Cellular Biology</i> , 2005, 25, 488-498.         | 1.1 | 61        |
| 688 | The Deubiquitinating Enzyme Ubp1 Affects Sorting of the ATP-binding Cassette-Transporter Ste6 in the Endocytic Pathway. <i>Molecular Biology of the Cell</i> , 2005, 16, 1319-1329.   | 0.9 | 33        |
| 689 | Yeast phosphatidylinositol 4-kinase, Ptk1, has essential roles at the Golgi and in the nucleus. <i>Journal of Cell Biology</i> , 2005, 171, 967-979.  | 2.3 | 119       |
| 690 | Defining the Order in Which Nmd3p and Rpl10p Load onto Nascent 60S Ribosomal Subunits. <i>Molecular and Cellular Biology</i> , 2005, 25, 3802-3813.   | 1.1 | 107       |
| 691 | Activator Gcn4p and Cyc8p/Tup1p Are Interdependent for Promoter Occupancy at ARG1 In Vivo. <i>Molecular and Cellular Biology</i> , 2005, 25, 11171-11183.   | 1.1 | 30        |
| 692 | The Actin Cytoskeleton Is Required for Selective Types of Autophagy, but Not Nonspecific Autophagy, in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2005, 16, 5843-5856.  | 0.9 | 139       |
| 693 | The Putative NTPase Fap7 Mediates Cytoplasmic 20S Pre-rRNA Processing through a Direct Interaction with Rps14. <i>Molecular and Cellular Biology</i> , 2005, 25, 10352-10364.   | 1.1 | 85        |
| 694 | Septin-dependent compartmentalization of the endoplasmic reticulum during yeast polarized growth. <i>Journal of Cell Biology</i> , 2005, 169, 897-908.  | 2.3 | 145       |
| 695 | Fzf1p Regulates an Inducible Response to Nitrosative Stress in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2005, 16, 4781-4791.  | 0.9 | 60        |
| 696 | PtdIns(3)P accumulation in triple lipid-phosphatase-deletion mutants triggers lethal hyperactivation of the Rho1p/Plc1p cell-integrity MAP kinase pathway. <i>Journal of Cell Science</i> , 2005, 118, 5589-5601.   | 1.2 | 17        |
| 697 | Components of the ESCRT Pathway, DFG16, and YGR122w Are Required for Rim101 To Act as a Corepressor with Nrg1 at the Negative Regulatory Element of the DIT1 Gene of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2005, 25, 6772-6788. | 1.1 | 70        |
| 698 | A Yeast Polyamine Acetyltransferase. <i>Journal of Biological Chemistry</i> , 2005, 280, 16659-16664.   | 1.6 | 38        |
| 699 | Defects in Structural Integrity of Ergosterol and the Cdc50p-Drs2p Putative Phospholipid Translocase Cause Accumulation of Endocytic Membranes, onto Which Actin Patches Are Assembled in Yeast. <i>Molecular Biology of the Cell</i> , 2005, 16, 5592-5609.        | 0.9 | 45        |
| 700 | Role for the SCFCDC4 Ubiquitin Ligase in <i>Candida albicans</i> Morphogenesis. <i>Molecular Biology of the Cell</i> , 2005, 16, 2772-2785.   | 0.9 | 87        |
| 701 | Role for Hsp70 Chaperone in <i>Saccharomyces cerevisiae</i> Prion Seed Replication. <i>Eukaryotic Cell</i> , 2005, 4, 289-297.  | 3.4 | 91        |
| 702 | Cyclophilin A Is Localized to the Nucleus and Controls Meiosis in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2005, 4, 17-29.  | 3.4 | 63        |
| 703 | Sensitivity to Phosphonoacetic Acid. <i>Genetics</i> , 2005, 170, 569-580.  | 1.2 | 56        |



| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 704 | Dynamic Changes in Protein-Protein Interaction and Protein Phosphorylation Probed with Amine-reactive Isotope Tag. <i>Molecular and Cellular Proteomics</i> , 2005, 4, 1358-1369.   | 2.5 | 71        |
| 705 | End3p-Mediated Endocytosis Is Required for Spore Wall Formation in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2005, 170, 1561-1574.  | 1.2 | 22        |
| 706 | The 9-1-1 Checkpoint Clamp Physically Interacts with Pol $\eta$ and Is Partially Required for Spontaneous Pol $\eta$ -dependent Mutagenesis in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2005, 280, 38657-38665. | 1.6 | 104       |
| 707 | Cell Cycle-regulated Trafficking of Chs2 Controls Actomyosin Ring Stability during Cytokinesis. <i>Molecular Biology of the Cell</i> , 2005, 16, 2529-2543.   | 0.9 | 110       |
| 708 | Dissection of Arp2/3 Complex Actin Nucleation Mechanism and Distinct Roles for Its Nucleation-Promoting Factors in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2005, 171, 35-47.  | 1.2 | 38        |
| 709 | <i>Saccharomyces cerevisiae</i> Sps1p Regulates Trafficking of Enzymes Required for Spore Wall Synthesis. <i>Eukaryotic Cell</i> , 2005, 4, 536-544.  | 3.4 | 31        |
| 710 | Defects in SUMO (Small Ubiquitin-related Modifier) Conjugation and Deconjugation Alter Cell Sensitivity to DNA Topoisomerase I-induced DNA Damage. <i>Journal of Biological Chemistry</i> , 2005, 280, 23566-23575.                               | 1.6 | 45        |
| 711 | Control of Golgi Morphology and Function by Sed5 $\tau$ -SNARE Phosphorylation. <i>Molecular Biology of the Cell</i> , 2005, 16, 4918-4930.   | 0.9 | 36        |
| 712 | Cell Polarity Protein Spa2P Associates With Proteins Involved In Actin Function In <i>Saccharomyces Cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2005, 16, 4595-4608.  | 0.9 | 33        |
| 713 | Nop53p is required for late 60S ribosome subunit maturation and nuclear export in yeast. <i>Rna</i> , 2005, 11, 1215-1224.  | 1.6 | 38        |
| 714 | Molecular Genetic Analysis of the Yeast Repressor Rfx1/Crt1 Reveals a Novel Two-Step Regulatory Mechanism. <i>Molecular and Cellular Biology</i> , 2005, 25, 7399-7411.   | 1.1 | 35        |
| 715 | The Phosphoinositide Phosphatase Sjl2 Is Recruited to Cortical Actin Patches in the Control of Vesicle Formation and Fission during Endocytosis. <i>Molecular and Cellular Biology</i> , 2005, 25, 2910-2923.                                     | 1.1 | 72        |
| 716 | The DNA Damage Checkpoint Response Requires Histone H2B Ubiquitination by Rad6-Bre1 and H3 Methylation by Dot1. <i>Journal of Biological Chemistry</i> , 2005, 280, 9879-9886.  | 1.6 | 249       |
| 717 | Novel Response to Microtubule Perturbation in Meiosis. <i>Molecular and Cellular Biology</i> , 2005, 25, 4767-4781.   | 1.1 | 49        |
| 718 | A telomeric repeat sequence adjacent to a DNA double-stranded break produces an antieckpoint. <i>Genes and Development</i> , 2005, 19, 2546-2559.   | 2.7 | 66        |
| 719 | The Yeast GTPase Mtg2p Is Required for Mitochondrial Translation and Partially Suppresses an rRNA Methyltransferase Mutant, mrm2. <i>Molecular Biology of the Cell</i> , 2005, 16, 954-963.   | 0.9 | 63        |
| 720 | APC and EB1 Function Together in Mitosis to Regulate Spindle Dynamics and Chromosome Alignment. <i>Molecular Biology of the Cell</i> , 2005, 16, 4609-4622.   | 0.9 | 205       |
| 721 | Yeast poly(A)-binding protein Pab1 shuttles between the nucleus and the cytoplasm and functions in mRNA export. <i>Rna</i> , 2005, 11, 517-531.   | 1.6 | 115       |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 722 | DNA Interstrand Cross-Link Repair in the <i>Saccharomyces cerevisiae</i> Cell Cycle: Overlapping Roles for PSO2 ( SNM1 ) with MutS Factors and EXO1 during S Phase. <i>Molecular and Cellular Biology</i> , 2005, 25, 2297-2309.            | 1.1 | 68        |
| 723 | The nuclear pore complex-associated protein, Mlp2p, binds to the yeast spindle pole body and promotes its efficient assembly. <i>Journal of Cell Biology</i> , 2005, 170, 225-235.  | 2.3 | 81        |
| 724 | Identification of Sumoylated Proteins by Systematic Immunoprecipitation of the Budding Yeast Proteome. <i>Molecular and Cellular Proteomics</i> , 2005, 4, 73-83.   | 2.5 | 103       |
| 725 | Yeast Mon2p is a highly conserved protein that functions in the cytoplasm-to-vacuole transport pathway and is required for Golgi homeostasis. <i>Journal of Cell Science</i> , 2005, 118, 4751-4764.  | 1.2 | 35        |
| 726 | Genetic Analysis of the Kinetochore DASH Complex Reveals an Antagonistic Relationship with the Ras/Protein Kinase A Pathway and a Novel Subunit Required for AskI Association. <i>Molecular and Cellular Biology</i> , 2005, 25, 767-778.   | 1.1 | 29        |
| 727 | Control of Ste6 Recycling by Ubiquitination in the Early Endocytic Pathway in Yeast. <i>Molecular Biology of the Cell</i> , 2005, 16, 2809-2821.  | 0.9 | 17        |
| 728 | Atg9 Cycles Between Mitochondria and the Pre-Autophagosomal Structure in Yeasts. <i>Autophagy</i> , 2005, 1, 101-109.   | 4.3 | 234       |
| 729 | Protein Kinase A Regulates Constitutive Expression of Small Heat-Shock Genes in an Msn2/4p-Independent and Hsf1p-Dependent Manner in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2005, 169, 1203-1214.                              | 1.2 | 59        |
| 730 | Two Complexes of Spindle Checkpoint Proteins Containing Cdc20 and Mad2 Assemble during Mitosis Independently of the Kinetochore in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2005, 4, 867-878.                             | 3.4 | 45        |
| 731 | Recombination protein Tid1p controls resolution of cohesin-dependent linkages in meiosis in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 2005, 171, 241-253.  | 2.3 | 30        |
| 732 | Atg17 Regulates the Magnitude of the Autophagic Response. <i>Molecular Biology of the Cell</i> , 2005, 16, 3438-3453.   | 0.9 | 207       |
| 733 | Glycerophosphocholine-dependent Growth Requires Gde1p (YPL110c) and Git1p in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2005, 280, 36110-36117.   | 1.6 | 64        |
| 734 | A Yeast Recombination-Based Cloning System to Produce Chimeric HIV-1 Viruses and Express HIV-1 Genes. , 2005, 304, 369-386.   |     | 8         |
| 735 | The <i>Saccharomyces cerevisiae</i> YLL012/YEH1, YLR020/YEH2, and TGL1 Genes Encode a Novel Family of Membrane-Anchored Lipases That Are Required for Steryl Ester Hydrolysis. <i>Molecular and Cellular Biology</i> , 2005, 25, 1655-1668. | 1.1 | 124       |
| 736 | Tgl4p and Tgl5p, Two Triacylglycerol Lipases of the Yeast <i>Saccharomyces cerevisiae</i> Are Localized to Lipid Particles. <i>Journal of Biological Chemistry</i> , 2005, 280, 37301-37309.  | 1.6 | 184       |
| 737 | Functional specificity of shuttling hnRNPs revealed by genome-wide analysis of their RNA binding profiles. <i>Rna</i> , 2005, 11, 383-393.  | 1.6 | 88        |
| 738 | Processing bodies require RNA for assembly and contain nontranslating mRNAs. <i>Rna</i> , 2005, 11, 371-382.  | 1.6 | 583       |
| 739 | HST2 Mediates SIR2-Independent Life-Span Extension by Calorie Restriction. <i>Science</i> , 2005, 309, 1861-1864.   | 6.0 | 213       |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 740 | Systematic yeast synthetic lethal and synthetic dosage lethal screens identify genes required for chromosome segregation. Proceedings of the National Academy of Sciences of the United States of America, 2005, 102, 13956-13961.                 | 3.3 | 126       |
| 741 | Instability of the Mitofusin Fzo1 Regulates Mitochondrial Morphology during the Mating Response of the Yeast <i>Saccharomyces cerevisiae</i> . Journal of Biological Chemistry, 2005, 280, 18598-18603.  | 1.6 | 81        |
| 742 | A contact-activated kinase signals <i>Candida albicans</i> invasive growth and biofilm development. Proceedings of the National Academy of Sciences of the United States of America, 2005, 102, 5576-5581.   | 3.3 | 189       |
| 743 | Characterization of the Yeast Trimeric-SAS Acetyltransferase Complex. Journal of Biological Chemistry, 2005, 280, 11987-11994.   | 1.6 | 76        |
| 744 | Unbiased selection of localization elements reveals cis-acting determinants of mRNA bud localization in <i>Saccharomyces cerevisiae</i> . Proceedings of the National Academy of Sciences of the United States of America, 2005, 102, 18005-18010. | 3.3 | 62        |
| 745 | Silencing Near tRNA Genes Requires Nucleolar Localization. Journal of Biological Chemistry, 2005, 280, 8637-8639.  | 1.6 | 75        |
| 746 | An early step in wobble uridine tRNA modification requires the Elongator complex. Rna, 2005, 11, 424-436.  | 1.6 | 382       |
| 747 | DNA-Bound Bas1 Recruits Pho2 To Activate ADE Genes in <i>Saccharomyces cerevisiae</i> . Eukaryotic Cell, 2005, 4, 1725-1735.   | 3.4 | 16        |
| 748 | Constitutive Activation of the pH-Responsive Rim101 Pathway in Yeast Mutants Defective in Late Steps of the MVB/ESCRT Pathway. Molecular and Cellular Biology, 2005, 25, 9478-9490.  | 1.1 | 89        |
| 749 | Characterizing the Sphingolipid Signaling Pathway That Remediate Defects Associated with Loss of the Yeast Amphiphysin-like Orthologs, Rvs161p and Rvs167p. Journal of Biological Chemistry, 2005, 280, 4270-4278.                                 | 1.6 | 24        |
| 750 | Molecular Organization of Target of Rapamycin Complex 2. Journal of Biological Chemistry, 2005, 280, 30697-30704.  | 1.6 | 197       |
| 751 | Transcriptional Regulation by Lge1p Requires a Function Independent of Its Role in Histone H2B Ubiquitination*. Journal of Biological Chemistry, 2005, 280, 2759-2770.   | 1.6 | 30        |
| 752 | The <i>Saccharomyces cerevisiae</i> Srb8-Srb11 Complex Functions with the SAGA Complex during Gal4-Activated Transcription. Molecular and Cellular Biology, 2005, 25, 114-123.   | 1.1 | 68        |
| 753 | Identification and Characterization of Elf1, a Conserved Transcription Elongation Factor in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 2005, 25, 10122-10135.   | 1.1 | 72        |
| 754 | G $\beta$ Subunit Gpa2 Recruits Kelch Repeat Subunits That Inhibit Receptor-G Protein Coupling during cAMP-induced Dimorphic Transitions in <i>Saccharomyces cerevisiae</i> . Molecular Biology of the Cell, 2005, 16, 4557-4571.                  | 0.9 | 67        |
| 755 | A GTP:AMP Phosphotransferase, Adk2p, in <i>Saccharomyces cerevisiae</i> . Journal of Biological Chemistry, 2005, 280, 18604-18609.   | 1.6 | 8         |
| 756 | Hsp110 Cooperates with Different Cytosolic HSP70 Systems in a Pathway for de Novo Folding. Journal of Biological Chemistry, 2005, 280, 41252-41261.  | 1.6 | 82        |
| 757 | The mRNA Nuclear Export Factor Hpr1 Is Regulated by Rsp5-mediated Ubiquitylation. Journal of Biological Chemistry, 2005, 280, 13401-13405.   | 1.6 | 61        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 758 | The core centromere and Sgo1 establish a 50-kb cohesin-protected domain around centromeres during meiosis I. <i>Genes and Development</i> , 2005, 19, 3017-3030.  | 2.7 | 87        |
| 759 | The DNA Repair Protein yKu80 Regulates the Function of Recombination Enhancer during Yeast Mating Type Switching. <i>Molecular and Cellular Biology</i> , 2005, 25, 8476-8485.  | 1.1 | 9         |
| 760 | Regulation of Cell Polarity by Interactions of Msb3 and Msb4 with Cdc42 and Polarisome Components. <i>Molecular and Cellular Biology</i> , 2005, 25, 8567-8580.   | 1.1 | 64        |
| 761 | The Yeast Phosphotyrosyl Phosphatase Activator Is Part of the Tap42-Phosphatase Complexes. <i>Molecular Biology of the Cell</i> , 2005, 16, 2119-2127.  | 0.9 | 38        |
| 762 | Genetic Interactions Between Mediator and the Late G1-Specific Transcription Factor Swi6 in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2005, 171, 477-488.   | 1.2 | 13        |
| 763 | Yeast prions [URE3] and [PSI <sup>+</sup> ] are diseases. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 10575-10580.  | 3.3 | 243       |
| 764 | YEH2/YLR020c Encodes a Novel Steryl Ester Hydrolase of the Yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2005, 280, 13321-13328.   | 1.6 | 60        |
| 765 | Mannosyl-Diinositolphospho-Ceramide, the Major Yeast Plasma Membrane Sphingolipid, Governs Toxicity of <i>Kluyveromyces lactis</i> Zymocin. <i>Eukaryotic Cell</i> , 2005, 4, 879-889.  | 3.4 | 28        |
| 766 | Oxidative DNA Damage Causes Mitochondrial Genomic Instability in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2005, 25, 5196-5204.   | 1.1 | 134       |
| 767 | An intracellular phosphate buffer filters transient fluctuations in extracellular phosphate levels. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 9565-9570.  | 3.3 | 85        |
| 768 | Swe1 regulation and transcriptional control restrict the activity of mitotic cyclins toward replication proteins in <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 8910-8915.      | 3.3 | 38        |
| 769 | The <i>Kluyveromyces lactis</i> $\hat{A}$ -toxin targets tRNA anticodons. <i>Rna</i> , 2005, 11, 1648-1654.   | 1.6 | 187       |
| 770 | Pex3p Initiates the Formation of a Preperoxisomal Compartment from a Subdomain of the Endoplasmic Reticulum in <i>Saccharomyces cerevisiae</i> *. <i>Journal of Biological Chemistry</i> , 2005, 280, 34933-34939.  | 1.6 | 149       |
| 771 | The Smk1p MAP kinase negatively regulates Gsc2p, a 1,3- $\hat{A}$ -glucan synthase, during spore wall morphogenesis in <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 12431-12436. | 3.3 | 35        |
| 772 | Role of Dot1-Dependent Histone H3 Methylation in G1 and S Phase DNA Damage Checkpoint Functions of Rad9. <i>Molecular and Cellular Biology</i> , 2005, 25, 8430-8443.   | 1.1 | 268       |
| 773 | UVA radiation is highly mutagenic in cells that are unable to repair 7,8-dihydro-8-oxoguanine in <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 13538-13543.                       | 3.3 | 119       |
| 774 | A data integration methodology for systems biology: Experimental verification. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 17302-17307.   | 3.3 | 124       |
| 775 | Ice2p is important for the distribution and structure of the cortical ER network in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 2005, 118, 65-77.  | 1.2 | 48        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 776 | The complete set of H/ACA snoRNAs that guide rRNA pseudouridylations in <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 2005, 11, 928-938.  | 1.6 | 51        |
| 777 | Rrp15p, a novel component of pre-ribosomal particles required for 60S ribosome subunit maturation. <i>Rna</i> , 2005, 11, 495-502.   | 1.6 | 26        |
| 778 | The 2 $\mu$ m Plasmid Causes Cell Death in <i>Saccharomyces cerevisiae</i> with a Mutation in Ulp1 Protease. <i>Molecular and Cellular Biology</i> , 2005, 25, 4299-4310.  | 1.1 | 39        |
| 779 | Direct Activation of Genes Involved in Intracellular Iron Use by the Yeast Iron-Responsive Transcription Factor Aft2 without Its Paralog Aft1. <i>Molecular and Cellular Biology</i> , 2005, 25, 6760-6771.                  | 1.1 | 100       |
| 780 | Modulation of Yeast Genome Expression in Response to Defective RNA Polymerase III-Dependent Transcription. <i>Molecular and Cellular Biology</i> , 2005, 25, 8631-8642.  | 1.1 | 36        |
| 781 | The Histone Chaperone Anti-Silencing Function 1 Is a Global Regulator of Transcription Independent of Passage through S Phase. <i>Molecular and Cellular Biology</i> , 2005, 25, 652-660.                                    | 1.1 | 51        |
| 782 | Mapping and Functional Characterization of the TAF11 Interaction with TFIIA. <i>Molecular and Cellular Biology</i> , 2005, 25, 945-957.  | 1.1 | 19        |
| 783 | A Role for the <i>Saccharomyces cerevisiae</i> Regulation of Ace2 and Polarized Morphogenesis Signaling Network in Cell Integrity. <i>Genetics</i> , 2005, 171, 443-455.   | 1.2 | 44        |
| 784 | A Genomic Screen for Yeast Vacuolar Membrane ATPase Mutants. <i>Genetics</i> , 2005, 170, 1539-1551.   | 1.2 | 86        |
| 785 | The Yeast Histone Chaperone Chromatin Assembly Factor 1 Protects Against Double-Strand DNA-Damaging Agents. <i>Genetics</i> , 2005, 171, 1513-1522.  | 1.2 | 64        |
| 786 | The Ama1-Directed Anaphase-Promoting Complex Regulates the Smk1 Mitogen-Activated Protein Kinase During Meiosis in Yeast. <i>Genetics</i> , 2005, 171, 901-911.  | 1.2 | 25        |
| 787 | TFIID and Spt-Ada-Gcn5-Acetyltransferase Functions Probed by Genome-wide Synthetic Genetic Array Analysis Using a <i>Saccharomyces cerevisiae</i> taf9-ts Allele. <i>Genetics</i> , 2005, 171, 959-973.                      | 1.2 | 18        |
| 788 | Genome-Wide Synthetic Lethal Screens Identify an Interaction Between the Nuclear Envelope Protein, Apq12p, and the Kinetochore in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2005, 171, 489-501.                    | 1.2 | 16        |
| 789 | ATP-independent Control of Vac8 Palmitoylation by a SNARE Subcomplex on Yeast Vacuoles. <i>Journal of Biological Chemistry</i> , 2005, 280, 15348-15355.   | 1.6 | 17        |
| 790 | Pml39, a Novel Protein of the Nuclear Periphery Required for Nuclear Retention of Improper Messenger Ribonucleoparticles. <i>Molecular Biology of the Cell</i> , 2005, 16, 5258-5268.  | 0.9 | 76        |
| 791 | The kinetochore protein Ndc10p is required for spindle stability and cytokinesis in yeast. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 5408-5413.                    | 3.3 | 50        |
| 792 | Genome-wide Analysis Reveals Inositol, Not Choline, as the Major Effector of Ino2p-Ino4p and Unfolded Protein Response Target Gene Expression in Yeast. <i>Journal of Biological Chemistry</i> , 2005, 280, 9106-9118.       | 1.6 | 112       |
| 793 | Role of the Novel Metallopeptidase MoP112 and Saccharolysin for the Complete Degradation of Proteins Residing in Different Subcompartments of Mitochondria. <i>Journal of Biological Chemistry</i> , 2005, 280, 20132-20139. | 1.6 | 77        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 794 | Forcing interactions as a genetic screen to identify proteins that exert a defined activity. <i>Genome Research</i> , 2005, 15, 560-565.   | 2.4 | 8         |
| 795 | Loss of Meiotic Rereplication Block in <i>Saccharomyces cerevisiae</i> Cells Defective in Cdc28p Regulation. <i>Eukaryotic Cell</i> , 2005, 4, 55-62.  | 3.4 | 15        |
| 796 | H2B Ubiquitin Protease Ubp8 and Sgf11 Constitute a Discrete Functional Module within the <i>Saccharomyces cerevisiae</i> SAGA Complex. <i>Molecular and Cellular Biology</i> , 2005, 25, 1162-1172.  | 1.1 | 126       |
| 797 | Gene function prediction from congruent synthetic lethal interactions in yeast. <i>Molecular Systems Biology</i> , 2005, 1, 2005.0026.   | 3.2 | 118       |
| 798 | Differential Activities and Regulation of <i>Saccharomyces cerevisiae</i> Formin Proteins Bni1 and Bnr1 by Bud6. <i>Journal of Biological Chemistry</i> , 2005, 280, 28023-28033.  | 1.6 | 134       |
| 799 | Mso1 Is a Novel Component of the Yeast Exocytic SNARE Complex. <i>Journal of Biological Chemistry</i> , 2005, 280, 34033-34041.  | 1.6 | 13        |
| 800 | Alg14 Recruits Alg13 to the Cytoplasmic Face of the Endoplasmic Reticulum to Form a Novel Bipartite UDP-N-acetylglucosamine Transferase Required for the Second Step of N-Linked Glycosylation. <i>Journal of Biological Chemistry</i> , 2005, 280, 36254-36262. | 1.6 | 102       |
| 801 | Mutants in trs120 disrupt traffic from the early endosome to the late Golgi. <i>Journal of Cell Biology</i> , 2005, 171, 823-833.  | 2.3 | 109       |
| 802 | The mobile nucleoporin Nup2p and chromatin-bound Prp20p function in endogenous NPC-mediated transcriptional control. <i>Journal of Cell Biology</i> , 2005, 171, 955-965.  | 2.3 | 114       |
| 803 | Synthetic Genetic Array Analysis of the PtdIns 4-kinase Pik1p Identifies Components in a Golgi-specific Ypt31/rab-GTPase Signaling Pathway. <i>Molecular Biology of the Cell</i> , 2005, 16, 776-793.  | 0.9 | 112       |
| 804 | Tor and Cyclic AMP-Protein Kinase A: Two Parallel Pathways Regulating Expression of Genes Required for Cell Growth. <i>Eukaryotic Cell</i> , 2005, 4, 63-71.   | 3.4 | 106       |
| 805 | Comparison of Cell Wall Localization among Pir Family Proteins and Functional Dissection of the Region Required for Cell Wall Binding and Bud Scar Recruitment of Pir1p. <i>Eukaryotic Cell</i> , 2005, 4, 1872-1881.  | 3.4 | 40        |
| 806 | A role for the CPF 3'-end processing machinery in RNAP II-dependent gene looping. <i>Genes and Development</i> , 2005, 19, 2969-2978.  | 2.7 | 219       |
| 807 | The Novel ATP-Binding Cassette Protein ARB1 Is a Shuttling Factor That Stimulates 40S and 60S Ribosome Biogenesis. <i>Molecular and Cellular Biology</i> , 2005, 25, 9859-9873.  | 1.1 | 60        |
| 808 | A Nonhistone Protein-Protein Interaction Required for Assembly of the SIR Complex and Silent Chromatin. <i>Molecular and Cellular Biology</i> , 2005, 25, 4514-4528.   | 1.1 | 85        |
| 809 | Cotranscriptional Recruitment of the Pseudouridylsynthetase Cbf5p and of the RNA Binding Protein Naf1p during H/ACA snoRNP Assembly. <i>Molecular and Cellular Biology</i> , 2005, 25, 3295-3304.  | 1.1 | 65        |
| 810 | Interaction of the <i>Saccharomyces cerevisiae</i> Cortical Actin Patch Protein Rvs167p With Proteins Involved in ER to Golgi Vesicle Trafficking. <i>Genetics</i> , 2005, 170, 555-568.   | 1.2 | 20        |
| 811 | Physical and Genetic Interactions Link the Yeast Protein Zds1p with mRNA Nuclear Export. <i>Journal of Biological Chemistry</i> , 2005, 280, 9691-9697.  | 1.6 | 13        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 812 | Chromatin assembly factor Asf1p-dependent occupancy of the SAS histone acetyltransferase complex at the silent mating-type locus HML $\alpha$ . <i>Nucleic Acids Research</i> , 2005, 33, 2742-2750.                   | 6.5 | 5         |
| 813 | Regulation of Histone Deposition Proteins Asf1/Hir1 by Multiple DNA Damage Checkpoint Kinases in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2005, 171, 885-899.   | 1.2 | 30        |
| 814 | Mutations in Mre11 Phosphoesterase Motif I That Impair <i>Saccharomyces cerevisiae</i> Mre11-Rad50-Xrs2 Complex Stability in Addition to Nuclease Activity. <i>Genetics</i> , 2005, 171, 1561-1570.                    | 1.2 | 76        |
| 815 | Spindle Checkpoint Maintenance Requires Ame1 and Okp1. <i>Cell Cycle</i> , 2005, 4, 1448-1456.   | 1.3 | 29        |
| 816 | FEAR but not MEN Genes are Required for Exit from Meiosis I. <i>Cell Cycle</i> , 2005, 4, 4093-4098.   | 1.3 | 30        |
| 817 | Evidence that the Tfg1/Tfg2 dimer interface of TFIIF lies near the active center of the RNA polymerase II initiation complex. <i>Nucleic Acids Research</i> , 2005, 33, 5045-5052.                                     | 6.5 | 41        |
| 818 | The Active Form of the <i>Saccharomyces cerevisiae</i> Ribonucleotide Reductase Small Subunit Is a Heterodimer <i>In Vitro</i> and <i>In Vivo</i> . <i>Biochemistry</i> , 2005, 44, 15366-15377.                       | 1.2 | 33        |
| 819 | Utilization of Target-Specific, Hypersensitive Strains of <i>Saccharomyces cerevisiae</i> To Determine the Mode of Action of Antifungal Compounds. <i>Antimicrobial Agents and Chemotherapy</i> , 2005, 49, 2558-2560. | 1.4 | 6         |
| 820 | Chromosome looping in yeast. <i>Journal of Cell Biology</i> , 2005, 168, 375-387.  | 2.3 | 186       |
| 821 | Discrimination Between Defects in Elongation Fidelity and Termination Efficiency Provides Mechanistic Insights into Translational Readthrough. <i>Journal of Molecular Biology</i> , 2005, 348, 801-815.               | 2.0 | 102       |
| 822 | The Structure of Sif2p, a WD Repeat Protein Functioning in the SET3 Corepressor Complex. <i>Journal of Molecular Biology</i> , 2005, 351, 923-935.   | 2.0 | 45        |
| 823 | Conserved Mechanism of Oxa1 Insertion into the Mitochondrial Inner Membrane. <i>Journal of Molecular Biology</i> , 2005, 354, 520-528.   | 2.0 | 21        |
| 824 | The TOR and EGO Protein Complexes Orchestrate Microautophagy in Yeast. <i>Molecular Cell</i> , 2005, 19, 15-26.  | 4.5 | 305       |
| 825 | The Protein Kinase Kin4 Inhibits Exit from Mitosis in Response to Spindle Position Defects. <i>Molecular Cell</i> , 2005, 19, 223-234.   | 4.5 | 131       |
| 826 | Histone H2B Ubiquitylation Controls Processive Methylation but Not Monomethylation by Dot1 and Set1. <i>Molecular Cell</i> , 2005, 19, 271-277.  | 4.5 | 224       |
| 827 | Mediator Expression Profiling Epistasis Reveals a Signal Transduction Pathway with Antagonistic Submodules and Highly Specific Downstream Targets. <i>Molecular Cell</i> , 2005, 19, 511-522.                          | 4.5 | 192       |
| 828 | Exploration of the Topological Requirements of ERAD Identifies Yos9p as a Lectin Sensor of Misfolded Glycoproteins in the ER Lumen. <i>Molecular Cell</i> , 2005, 19, 741-751.   | 4.5 | 204       |
| 829 | The Putative RNA Helicase Dbp4p Is Required for Release of the U14 snoRNA from Preribosomes in <i>Saccharomyces cerevisiae</i> . <i>Molecular Cell</i> , 2005, 20, 53-64.  | 4.5 | 83        |

| #   | ARTICLE   | IF   | CITATIONS |
|-----|---|------|-----------|
| 830 | An Essential GTPase Promotes Assembly of Preribosomal RNA Processing Complexes. <i>Molecular Cell</i> , 2005, 20, 633-643.  | 4.5  | 65        |
| 831 | Eaf3 Chromodomain Interaction with Methylated H3-K36 Links Histone Deacetylation to Pol II Elongation. <i>Molecular Cell</i> , 2005, 20, 971-978.   | 4.5  | 480       |
| 832 | Gpi17p does not stably interact with other subunits of glycosylphosphatidylinositol transamidase in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2005, 1735, 79-88.  | 1.2  | 20        |
| 833 | Flux of sterol intermediates in a yeast strain deleted of the lanosterol C-14 demethylase Erg11p. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2005, 1735, 111-118.  | 1.2  | 23        |
| 834 | Anc1 interacts with the catalytic subunits of the general transcription factors TFIID and TFIIF, the chromatin remodeling complexes RSC and INO80, and the histone acetyltransferase complex NuA3. <i>Biochemical and Biophysical Research Communications</i> , 2005, 332, 398-403. | 1.0  | 43        |
| 835 | Sen34p depletion blocks tRNA splicing in vivo and delays rRNA processing. <i>Biochemical and Biophysical Research Communications</i> , 2005, 337, 89-94.  | 1.0  | 25        |
| 836 | The budding yeast protein Chl1p has a role in transcriptional silencing, rDNA recombination, and aging. <i>Biochemical and Biophysical Research Communications</i> , 2005, 337, 167-172.  | 1.0  | 16        |
| 837 | Characterization of a novel autophagy-specific gene, ATG29. <i>Biochemical and Biophysical Research Communications</i> , 2005, 338, 1884-1889.  | 1.0  | 92        |
| 838 | The transcriptional activator HAP4 is a high copy suppressor of an oxa1 yeast mutation. <i>Gene</i> , 2005, 354, 53-57.   | 1.0  | 8         |
| 839 | Sterile 20 Kinase Phosphorylates Histone H2B at Serine 10 during Hydrogen Peroxide-Induced Apoptosis in <i>S. cerevisiae</i> . <i>Cell</i> , 2005, 120, 25-36.  | 13.5 | 226       |
| 840 | Acetylation in Histone H3 Globular Domain Regulates Gene Expression in Yeast. <i>Cell</i> , 2005, 121, 375-385.   | 13.5 | 362       |
| 841 | Contribution of the Endoplasmic Reticulum to Peroxisome Formation. <i>Cell</i> , 2005, 122, 85-95.  | 13.5 | 416       |
| 842 | The FK506 Binding Protein Fpr3 Counteracts Protein Phosphatase 1 to Maintain Meiotic Recombination Checkpoint Activity. <i>Cell</i> , 2005, 122, 861-873.   | 13.5 | 137       |
| 843 | In Vivo Dynamics of Clathrin and Its Adaptor-Dependent Recruitment to the Actin-Based Endocytic Machinery in Yeast. <i>Developmental Cell</i> , 2005, 9, 87-98.   | 3.1  | 165       |
| 844 | Glucose deprivation mediates interaction between CTDK-I and Snf1 in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 2005, 579, 5318-5324.   | 1.3  | 7         |
| 845 | Cell integrity signaling activation in response to hyperosmotic shock in yeast. <i>FEBS Letters</i> , 2005, 579, 6186-6190.   | 1.3  | 52        |
| 846 | <i>Candida albicans</i> CHT3 encodes the functional homolog of the Cts1 chitinase of <i>Saccharomyces cerevisiae</i> . <i>Fungal Genetics and Biology</i> , 2005, 42, 935-947.  | 0.9  | 88        |
| 847 | Metabolic Engineering of the Purine Pathway for Riboflavin Production in <i>Ashbya gossypii</i> . <i>Applied and Environmental Microbiology</i> , 2005, 71, 5743-5751.  | 1.4  | 106       |



| #   | ARTICLE  | IF   | CITATIONS |
|-----|--|------|-----------|
| 848 | Modulation of the transcription regulatory program in yeast cells committed to sporulation. <i>Genome Biology</i> , 2006, 7, R20.  | 13.9 | 63        |
| 849 | Increased Recombination Between Active tRNA Genes. <i>DNA and Cell Biology</i> , 2006, 25, 359-364.  | 0.9  | 22        |
| 850 | Nuclear Localization Signal and Protein Context both Mediate Importin $\beta$ Specificity of Nuclear Import Substrates. <i>Molecular and Cellular Biology</i> , 2006, 26, 8697-8709.   | 1.1  | 80        |
| 851 | Screening for Genetic Modifiers of Amyloid Toxicity in Yeast. <i>Methods in Enzymology</i> , 2006, 412, 201-222.   | 0.4  | 15        |
| 852 | Yeast as a Model for Human Disease. <i>Current Protocols in Human Genetics</i> , 2006, 48, Unit 15.6.  | 3.5  | 36        |
| 853 | Yeast ( <i>Saccharomyces cerevisiae</i> ). , 2006, 344, 465-473.   |      | 6         |
| 854 | Modulation of Prion Formation, Aggregation, and Toxicity by the Actin Cytoskeleton in Yeast. <i>Molecular and Cellular Biology</i> , 2006, 26, 617-629.  | 1.1  | 133       |
| 855 | <i>Saccharomyces cerevisiae</i> -Based Molecular Tool Kit for Manipulation of Genes from Gram-Negative Bacteria. <i>Applied and Environmental Microbiology</i> , 2006, 72, 5027-5036.  | 1.4  | 384       |
| 856 | Cotranscriptional Recruitment to the mRNA Export Receptor Mex67p Contributes to Nuclear Pore Anchoring of Activated Genes. <i>Molecular and Cellular Biology</i> , 2006, 26, 7858-7870.  | 1.1  | 173       |
| 857 | Sjl2p is specifically involved in early steps of endocytosis intimately linked to actin dynamics via the Ark1p/Prk1p kinases. <i>FEBS Letters</i> , 2006, 580, 633-641.  | 1.3  | 7         |
| 858 | The NoCut Pathway Links Completion of Cytokinesis to Spindle Midzone Function to Prevent Chromosome Breakage. <i>Cell</i> , 2006, 125, 85-98.  | 13.5 | 267       |
| 859 | Single-Molecule Analysis of Dynein Processivity and Stepping Behavior. <i>Cell</i> , 2006, 126, 335-348.   | 13.5 | 571       |
| 860 | Mutational analysis of the Lem3p-Dnf1p putative phospholipid-translocating P-type ATPase reveals novel regulatory roles for Lem3p and a carboxyl-terminal region of Dnf1p independent of the phospholipid-translocating activity of Dnf1p in yeast. <i>Biochemical and Biophysical Research Communications</i> , 2006, 344, 323-331. | 1.0  | 29        |
| 861 | Ubiquinone accumulates in the mitochondria of yeast mutated in the ubiquinone binding protein, Qcr8p. <i>Biochemical and Biophysical Research Communications</i> , 2006, 344, 241-245.   | 1.0  | 5         |
| 862 | Gcn5p contributes to the bidirectional character of the UGA3â€œGLT1 yeast promoter. <i>Biochemical and Biophysical Research Communications</i> , 2006, 348, 989-996.   | 1.0  | 2         |
| 863 | Esc4/Rtt107 and the control of recombination during replication. <i>DNA Repair</i> , 2006, 5, 618-628.   | 1.3  | 41        |
| 864 | Histone H2A phosphorylation and H3 methylation are required for a novel Rad9 DSB repair function following checkpoint activation. <i>DNA Repair</i> , 2006, 5, 693-703.  | 1.3  | 114       |
| 865 | Trapping of DNA topoisomerase I on nick-containing DNA in cell free extracts of <i>Saccharomyces cerevisiae</i> . <i>DNA Repair</i> , 2006, 5, 799-809.  | 1.3  | 21        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 866 | Interplay between Ino80 and Swr1 chromatin remodeling enzymes regulates cell cycle checkpoint adaptation in response to DNA damage. <i>Genes and Development</i> , 2006, 20, 2437-2449.                       | 2.7 | 162       |
| 867 | The C-terminal TPR Domain of Tom70 Defines a Family of Mitochondrial Protein Import Receptors Found only in Animals and Fungi. <i>Journal of Molecular Biology</i> , 2006, 358, 1010-1022.                    | 2.0 | 97        |
| 868 | Transcriptional Activators Are Dispensable for Transcription in the Absence of Spt6-Mediated Chromatin Reassembly of Promoter Regions. <i>Molecular Cell</i> , 2006, 21, 405-416.                             | 4.5 | 150       |
| 869 | Constraining G1-Specific Transcription to Late G1 Phase: The MBF-Associated Corepressor Nrm1 Acts via Negative Feedback. <i>Molecular Cell</i> , 2006, 23, 483-496.   | 4.5 | 121       |
| 870 | The DNA Repair Helicases XPD and FancJ Have Essential Iron-Sulfur Domains. <i>Molecular Cell</i> , 2006, 23, 801-808.   | 4.5 | 298       |
| 871 | Transcription Termination and Nuclear Degradation of Cryptic Unstable Transcripts: A Role for the Nrd1-Nab3 Pathway in Genome Surveillance. <i>Molecular Cell</i> , 2006, 23, 853-864.                        | 4.5 | 209       |
| 872 | DNA Degradation at Unprotected Telomeres in Yeast Is Regulated by the CDK1 (Cdc28/Clb) Cell-Cycle Kinase. <i>Molecular Cell</i> , 2006, 24, 127-137.  | 4.5 | 117       |
| 873 | Differential Recruitment of the Splicing Machinery during Transcription Predicts Genome-Wide Patterns of mRNA Splicing. <i>Molecular Cell</i> , 2006, 24, 903-915.  | 4.5 | 60        |
| 874 | Analysis of in vivo targets of transcriptional activators by fluorescence resonance energy transfer. <i>Methods</i> , 2006, 40, 353-359.  | 1.9 | 6         |
| 875 | Quantitative genetic analysis in <i>Saccharomyces cerevisiae</i> using epistatic miniarray profiles (E-MAPs) and its application to chromatin functions. <i>Methods</i> , 2006, 40, 344-352.                  | 1.9 | 76        |
| 876 | System of centromeric, episomal, and integrative vectors based on drug resistance markers for <i>Saccharomyces cerevisiae</i> . <i>BioTechniques</i> , 2006, 40, 73-78.                                       | 0.8 | 174       |
| 877 | A novel magnetic resonance-based method to measure gene expression in living cells. <i>Nucleic Acids Research</i> , 2006, 34, e51-e51.  | 6.5 | 14        |
| 878 | Rtt107/Esc4 binds silent chromatin and DNA repair proteins using different BRCT motifs. <i>BMC Molecular Biology</i> , 2006, 7, 40.   | 3.0 | 27        |
| 879 | The yeast potassium transporter TRK2 is able to substitute for TRK1 in its biological function under low K and low pH conditions. <i>Yeast</i> , 2006, 23, 581-589.   | 0.8 | 18        |
| 880 | HB tag modules for PCR-based gene tagging and tandem affinity purification in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2006, 23, 623-632.   | 0.8 | 53        |
| 881 | A two-step method for the introduction of single or multiple defined point mutations into the genome of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2006, 23, 825-831.                                   | 0.8 | 46        |
| 882 | The <i>Saccharomyces cerevisiae</i> orthologue of the human protein phosphatase 4 core regulatory subunit R2 confers resistance to the anticancer drug cisplatin. <i>FEBS Journal</i> , 2006, 273, 3322-3334. | 2.2 | 28        |
| 883 | Oligomerization of the Mg <sup>2+</sup> -transport proteins Alr1p and Alr2p in yeast plasma membrane. <i>FEBS Journal</i> , 2006, 273, 4236-4249.   | 2.2 | 33        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 884 | Accounting for strain-specific differences during RTG target gene regulation in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2006, 6, 112-119.   | 1.1 | 15        |
| 885 | Differing responses of Gat1 and Gln3 phosphorylation and localization to rapamycin and methionine sulfoximine treatment in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2006, 6, 218-229.          | 1.1 | 51        |
| 886 | Control by sugar of <i>Saccharomyces cerevisiae</i> flocculation for industrial ethanol production. <i>FEMS Yeast Research</i> , 2006, 6, 280-287.   | 1.1 | 28        |
| 887 | Double-strand breaks trigger MRX- and Mec1-dependent, but Tel1-independent, checkpoint activation. <i>FEMS Yeast Research</i> , 2006, 6, 836-847.  | 1.1 | 17        |
| 888 | The mechanism by which overexpression of Gts1p induces flocculation in a FLO8-inactive strain of the yeast <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2006, 6, 914-923.                          | 1.1 | 13        |
| 889 | Yeast-based assay for the measurement of positive and negative influences on microsatellite stability. <i>FEMS Yeast Research</i> , 2006, 6, 716-725.  | 1.1 | 10        |
| 890 | Genome-wide analysis of the response to protein glycosylation deficiency in yeast. <i>FEMS Yeast Research</i> , 2006, 6, 1264-1273.  | 1.1 | 16        |
| 891 | Partial Rescue of pos5 Mutants by YEF1 and UTR1 Genes in <i>Saccharomyces cerevisiae</i> . <i>Acta Biochimica Et Biophysica Sinica</i> , 2006, 38, 293-298.  | 0.9 | 6         |
| 892 | Coordinated Nuclear Import of RNA Polymerase III Subunits. <i>Traffic</i> , 2006, 7, 465-473.  | 1.3 | 20        |
| 893 | Clathrin is Important for Normal Actin Dynamics and Progression of Sla2p-Containing Patches During Endocytosis in Yeast. <i>Traffic</i> , 2006, 7, 574-588.  | 1.3 | 56        |
| 894 | Vac8p, an Armadillo Repeat Protein, Coordinates Vacuole Inheritance With Multiple Vacuolar Processes. <i>Traffic</i> , 2006, 7, 1368-1377.   | 1.3 | 44        |
| 895 | Sel1p/Ubx2p Participates in a Distinct Cdc48p-Dependent Endoplasmic Reticulum-Associated Degradation Pathway. <i>Traffic</i> , 2006, 7, 1213-1223.   | 1.3 | 20        |
| 896 | Formation and Nuclear Export of Preribosomes Are Functionally Linked to the Small-Ubiquitin-Related Modifier Pathway. <i>Traffic</i> , 2006, 7, 1311-1321.   | 1.3 | 87        |
| 897 | tRNA <sup>Glu</sup> wobble uridine methylation by Trm9 identifies Elongator's key role for zymocin-induced cell death in yeast. <i>Molecular Microbiology</i> , 2006, 59, 677-688.                                   | 1.2 | 98        |
| 898 | The UGA3-GLT1 intergenic region constitutes a promoter whose bidirectional nature is determined by chromatin organization in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2006, 59, 1790-1806.  | 1.2 | 18        |
| 899 | Proteasome- and SCF-dependent degradation of yeast adenine deaminase upon transition from proliferation to quiescence requires a new F-box protein named Saf1p. <i>Molecular Microbiology</i> , 2006, 60, 1014-1025. | 1.2 | 31        |
| 900 | The Ipl1-Aurora protein kinase activates the spindle checkpoint by creating unattached kinetochores. <i>Nature Cell Biology</i> , 2006, 8, 78-83.  | 4.6 | 272       |
| 901 | Molecular architecture of a kinetochore microtubule attachment site. <i>Nature Cell Biology</i> , 2006, 8, 581-585.  | 4.6 | 263       |

| #   | ARTICLE   | IF   | CITATIONS |
|-----|---|------|-----------|
| 902 | A conserved GTPase-containing complex is required for intracellular sorting of the general amino-acid permease in yeast. <i>Nature Cell Biology</i> , 2006, 8, 657-667.                                       | 4.6  | 169       |
| 903 | Linear chromosome maintenance in the absence of essential telomere-capping proteins. <i>Nature Cell Biology</i> , 2006, 8, 734-740.   | 4.6  | 39        |
| 904 | A complex of Yos9p and the HRD ligase integrates endoplasmic reticulum quality control into the degradation machinery. <i>Nature Cell Biology</i> , 2006, 8, 849-854.   | 4.6  | 204       |
| 905 | Proteasome-dependent degradation of Est1p regulates the cell cycle-restricted assembly of telomerase in <i>Saccharomyces cerevisiae</i> . <i>Nature Structural and Molecular Biology</i> , 2006, 13, 720-728. | 3.6  | 81        |
| 906 | Eisosomes mark static sites of endocytosis. <i>Nature</i> , 2006, 439, 998-1003.  | 13.7 | 304       |
| 907 | Hrr25-dependent phosphorylation state regulates organization of the pre-40S subunit. <i>Nature</i> , 2006, 441, 651-655.  | 13.7 | 191       |
| 908 | A subcomplex of RNA polymerase III subunits involved in transcription termination and reinitiation. <i>EMBO Journal</i> , 2006, 25, 118-128.  | 3.5  | 119       |
| 909 | DNA interstrand crosslink repair during G1 involves nucleotide excision repair and DNA polymerase $\eta$ . <i>EMBO Journal</i> , 2006, 25, 1285-1294.   | 3.5  | 149       |
| 910 | Purification of active HOPS complex reveals its affinities for phosphoinositides and the SNARE Vam7p. <i>EMBO Journal</i> , 2006, 25, 1579-1589.  | 3.5  | 224       |
| 911 | The Hrd1p ligase complex forms a linchpin between ER-luminal substrate selection and Cdc48p recruitment. <i>EMBO Journal</i> , 2006, 25, 1827-1835.   | 3.5  | 129       |
| 912 | TOR regulates late steps of ribosome maturation in the nucleoplasm via Nog1 in response to nutrients. <i>EMBO Journal</i> , 2006, 25, 3832-3842.  | 3.5  | 54        |
| 913 | Ptc1p regulates cortical ER inheritance via Slr2p. <i>EMBO Journal</i> , 2006, 25, 4413-4422.   | 3.5  | 66        |
| 914 | An essential role for Orc6 in DNA replication through maintenance of pre-replicative complexes. <i>EMBO Journal</i> , 2006, 25, 5150-5158.  | 3.5  | 55        |
| 915 | CDK Pho85 targets CDK inhibitor Sic1 to relieve yeast G1 checkpoint arrest after DNA damage. <i>Nature Structural and Molecular Biology</i> , 2006, 13, 908-914.  | 3.6  | 36        |
| 916 | Kinase Cak1 functionally interacts with the PAF1 complex and phosphatase Ssu72 via kinases Ctk1 and Bur1. <i>Molecular Genetics and Genomics</i> , 2006, 275, 136-147.  | 1.0  | 8         |
| 917 | Residues of the yeast ALR1 protein that are critical for Magnesium uptake. <i>Current Genetics</i> , 2006, 49, 7-20.  | 0.8  | 26        |
| 918 | Coordinate regulation of multiple and distinct biosynthetic pathways by TOR and PKA kinases in <i>S. cerevisiae</i> . <i>Current Genetics</i> , 2006, 49, 281-293.  | 0.8  | 46        |
| 919 | <i>Agrobacterium</i> T-DNA-mediated integration and gene replacement in the brown rot pathogen <i>Monilinia fructicola</i> . <i>Current Genetics</i> , 2006, 49, 309-322.                                     | 0.8  | 43        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 920 | Structural involvement in substrate recognition of an essential aspartate residue conserved in Mep/Amt and Rh-type ammonium transporters. <i>Current Genetics</i> , 2006, 49, 364-374.  | 0.8 | 62        |
| 921 | Roles of the RAM signaling network in cell cycle progression in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2006, 49, 384-392.  | 0.8 | 20        |
| 922 | Sequences in the N-terminal cytoplasmic domain of <i>Saccharomyces cerevisiae</i> maltose permease are required for vacuolar degradation but not glucose-induced internalization. <i>Current Genetics</i> , 2006, 50, 101-114.                  | 0.8 | 17        |
| 923 | Ypt32p regulates the translocation of Chs3p from an internal pool to the plasma membrane. <i>European Journal of Cell Biology</i> , 2006, 85, 107-116.  | 1.6 | 21        |
| 924 | IQGAP and mitotic exit network (MEN) proteins are required for cytokinesis and re-polarization of the actin cytoskeleton in the budding yeast, <i>Saccharomyces cerevisiae</i> . <i>European Journal of Cell Biology</i> , 2006, 85, 1201-1215. | 1.6 | 27        |
| 925 | CTD kinase I is required for the integrity of the rDNA tandem array. <i>Nucleic Acids Research</i> , 2006, 34, 4996-5006.   | 6.5 | 14        |
| 926 | Limiting amounts of budding yeast Rad53 S-phase checkpoint activity results in increased resistance to DNA alkylation damage. <i>Nucleic Acids Research</i> , 2006, 34, 5852-5862.  | 6.5 | 22        |
| 927 | SAGA-associated Sgf73p facilitates formation of the preinitiation complex assembly at the promoters either in a HAT-dependent or independent manner in vivo. <i>Nucleic Acids Research</i> , 2006, 34, 6225-6232.                               | 6.5 | 34        |
| 928 | Low abundance of telomerase in yeast: Implications for telomerase haploinsufficiency. <i>Rna</i> , 2006, 12, 1721-1737.   | 1.6 | 108       |
| 929 | The nucleolar protein Esf2 interacts directly with the DExD/H box RNA helicase, Dbp8, to stimulate ATP hydrolysis. <i>Nucleic Acids Research</i> , 2006, 34, 3189-3199.   | 6.5 | 54        |
| 930 | The WD40-repeat protein Pwp1p associates in vivo with 25S ribosomal chromatin in a histone H4 tail-dependent manner. <i>Nucleic Acids Research</i> , 2006, 34, 3555-3567.   | 6.5 | 11        |
| 931 | SIR2 modifies histone H4-K16 acetylation and affects superhelicity in the ARS region of plasmid chromatin in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2006, 34, 5426-5437.   | 6.5 | 15        |
| 932 | Predicting essential genes in fungal genomes. <i>Genome Research</i> , 2006, 16, 1126-1135.   | 2.4 | 109       |
| 933 | The Yeast Ccr4-Not Complex Controls Ubiquitination of the Nascent-associated Polypeptide (NAC-EGD) Complex. <i>Journal of Biological Chemistry</i> , 2006, 281, 31389-31398.  | 1.6 | 100       |
| 934 | The Flo8 Transcription Factor Is Essential for Hyphal Development and Virulence in <i>Candida albicans</i> . <i>Molecular Biology of the Cell</i> , 2006, 17, 295-307.  | 0.9 | 193       |
| 935 | The Arf-GTPase-Activating Protein Gcs1p Is Essential for Sporulation and Regulates the Phospholipase D Spo14p. <i>Eukaryotic Cell</i> , 2006, 5, 112-124.   | 3.4 | 18        |
| 936 | Roles of type II myosin and a tropomyosin isoform in retrograde actin flow in budding yeast. <i>Journal of Cell Biology</i> , 2006, 175, 957-969.   | 2.3 | 80        |
| 937 | Connection Between the Rag4 Glucose Sensor and the KIRgt1 Repressor in <i>Kluyveromyces lactis</i> . <i>Genetics</i> , 2006, 174, 617-626.  | 1.2 | 28        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 938 | Laa1p, a Conserved AP-1 Accessory Protein Important for AP-1 Localization in Yeast. <i>Molecular Biology of the Cell</i> , 2006, 17, 3304-3317.   | 0.9 | 27        |
| 939 | Expression of the <i>Candida albicans</i> Morphogenesis Regulator Gene CZF1 and Its Regulation by Efg1p and Czf1p. <i>Eukaryotic Cell</i> , 2006, 5, 825-835.   | 3.4 | 59        |
| 940 | The RSC Chromatin Remodeling Complex Bears an Essential Fungal-Specific Protein Module With Broad Functional Roles. <i>Genetics</i> , 2006, 172, 795-809.   | 1.2 | 61        |
| 941 | Alternative 3' Pre-mRNA Processing in <i>Saccharomyces cerevisiae</i> Is Modulated by Nab4/Hrp1 In Vivo. <i>PLoS Biology</i> , 2006, 5, e6.   | 2.6 | 34        |
| 942 | Analysis of kinesin motor function at budding yeast kinetochores. <i>Journal of Cell Biology</i> , 2006, 172, 861-874.  | 2.3 | 127       |
| 943 | Phospholipase D and the SNARE Sso1p are necessary for vesicle fusion during sporulation in yeast. <i>Journal of Cell Science</i> , 2006, 119, 1406-1415.  | 1.2 | 110       |
| 944 | The polarity-establishment component Bem1p interacts with the exocyst complex through the Sec15p subunit. <i>Journal of Cell Science</i> , 2006, 119, 876-888.  | 1.2 | 42        |
| 945 | Actin-Induced Hyperactivation of the Ras Signaling Pathway Leads to Apoptosis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2006, 26, 6487-6501.                           | 1.1 | 135       |
| 946 | Ammonium Toxicity and Potassium Limitation in Yeast. <i>PLoS Biology</i> , 2006, 4, e351.   | 2.6 | 123       |
| 947 | The transcriptional activity of RNA polymerase I is a key determinant for the level of all ribosome components. <i>Genes and Development</i> , 2006, 20, 2030-2040.                                       | 2.7 | 175       |
| 948 | Nuclear congression is driven by cytoplasmic microtubule plus end interactions in <i>S. cerevisiae</i> . <i>Journal of Cell Biology</i> , 2006, 172, 27-39.   | 2.3 | 45        |
| 949 | Acetylation of H2AZ Lys 14 is associated with genome-wide gene activity in yeast. <i>Genes and Development</i> , 2006, 20, 711-722.   | 2.7 | 241       |
| 950 | Rsc4 Connects the Chromatin Remodeler RSC to RNA Polymerases. <i>Molecular and Cellular Biology</i> , 2006, 26, 4920-4933.  | 1.1 | 98        |
| 951 | Swt1, a Novel Yeast Protein, Functions in Transcription. <i>Journal of Biological Chemistry</i> , 2006, 281, 36518-36525.   | 1.6 | 16        |
| 952 | KIRHO1 and KIPKC1 are essential for cell integrity signalling in <i>Kluyveromyces lactis</i> . <i>Microbiology (United Kingdom)</i> , 2006, 152, 2635-2649.   | 0.7 | 25        |
| 953 | Phosphatase 2A Negatively Regulates Mitotic Exit in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2006, 17, 80-89.   | 0.9 | 65        |
| 954 | Sumoylation of the budding yeast kinetochore protein Ndc10 is required for Ndc10 spindle localization and regulation of anaphase spindle elongation. <i>Journal of Cell Biology</i> , 2006, 174, 653-663. | 2.3 | 74        |
| 955 | Genetic and molecular interactions of the Erv41p-Erv46p complex involved in transport between the endoplasmic reticulum and Golgi complex. <i>Journal of Cell Science</i> , 2006, 119, 4730-4740.         | 1.2 | 19        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 956 | Yeh1 Constitutes the Major Steryl Ester Hydrolase under Heme-Deficient Conditions in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2006, 5, 1018-1025.   | 3.4 | 31        |
| 957 | The Regulation of Microtubule Dynamics in <i>Saccharomyces cerevisiae</i> by Three Interacting Plus-End Tracking Proteins. <i>Molecular Biology of the Cell</i> , 2006, 17, 2789-2798.  | 0.9 | 71        |
| 958 | Rtn1p Is Involved in Structuring the Cortical Endoplasmic Reticulum. <i>Molecular Biology of the Cell</i> , 2006, 17, 3009-3020.  | 0.9 | 118       |
| 959 | Evidence that Spt2/Sin1, an HMG-Like Factor, Plays Roles in Transcription Elongation, Chromatin Structure, and Genome Stability in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2006, 26, 1496-1509. | 1.1 | 82        |
| 960 | Distinct Roles for TGN/Endosome Epsin-like Adaptors Ent3p and Ent5p. <i>Molecular Biology of the Cell</i> , 2006, 17, 3907-3920.  | 0.9 | 69        |
| 961 | Eukaryotic Translation Initiation Factor 3 (eIF3) and eIF2 Can Promote mRNA Binding to 40S Subunits Independently of eIF4G in Yeast. <i>Molecular and Cellular Biology</i> , 2006, 26, 1355-1372.                                 | 1.1 | 111       |
| 962 | Target hub proteins serve as master regulators of development in yeast. <i>Genes and Development</i> , 2006, 20, 435-448.   | 2.7 | 153       |
| 963 | Role of Cell Cycle-regulated Expression in the Localized Incorporation of Cell Wall Proteins in Yeast. <i>Molecular Biology of the Cell</i> , 2006, 17, 3267-3280.  | 0.9 | 31        |
| 964 | Yeast Gcn4p Stabilization Is Initiated by the Dissociation of the Nuclear Pho85p/Pcl5p Complex. <i>Molecular Biology of the Cell</i> , 2006, 17, 2952-2962.   | 0.9 | 22        |
| 965 | Suppressed Disassembly of Autolyzing p94/CAPN3 by N2A Connectin/Titin in a Genetic Reporter System. <i>Journal of Biological Chemistry</i> , 2006, 281, 18519-18531.  | 1.6 | 59        |
| 966 | Inositol Deacylation by Bst1p Is Required for the Quality Control of Glycosylphosphatidylinositol-anchored Proteins. <i>Molecular Biology of the Cell</i> , 2006, 17, 834-850.  | 0.9 | 86        |
| 967 | Role of a Cdc42p Effector Pathway in Recruitment of the Yeast Septins to the Presumptive Bud Site. <i>Molecular Biology of the Cell</i> , 2006, 17, 1110-1125.  | 0.9 | 127       |
| 968 | The Multi-KH Domain Protein of <i>Saccharomyces cerevisiae</i> Scp160p Contributes to the Regulation of Telomeric Silencing. <i>Journal of Biological Chemistry</i> , 2006, 281, 18227-18235.                                     | 1.6 | 15        |
| 969 | Diverse Functions of Spindle Assembly Checkpoint Genes in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2006, 172, 53-65.   | 1.2 | 53        |
| 970 | Function of the MAPK scaffold protein, Ste5, requires a cryptic PH domain. <i>Genes and Development</i> , 2006, 20, 1946-1958.  | 2.7 | 54        |
| 971 | Amino Acids Regulate Retrieval of the Yeast General Amino Acid Permease from the Vacuolar Targeting Pathway. <i>Molecular Biology of the Cell</i> , 2006, 17, 3031-3050.  | 0.9 | 63        |
| 972 | Nitrogen Availability and TOR Regulate the Snf1 Protein Kinase in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2006, 5, 1831-1837.  | 3.4 | 77        |
| 973 | Cell Cycle-Dependent Regulation of <i>Saccharomyces cerevisiae</i> Donor Preference during Mating-Type Switching by SBF (Swi4/Swi6) and Fkh1. <i>Molecular and Cellular Biology</i> , 2006, 26, 5470-5480.                        | 1.1 | 21        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 974 | Yeast GTB1 Encodes a Subunit of Glucosidase II Required for Glycoprotein Processing in the Endoplasmic Reticulum. <i>Journal of Biological Chemistry</i> , 2006, 281, 6325-6333.  | 1.6 | 50        |
| 975 | Glc7/Protein Phosphatase 1 Regulatory Subunits Can Oppose the Ipl1/Aurora Protein Kinase by Redistributing Glc7. <i>Molecular and Cellular Biology</i> , 2006, 26, 2648-2660.   | 1.1 | 102       |
| 976 | The role of the Brr5/Ysh1 C-terminal domain and its homolog Syc1 in mRNA 3'-end processing in <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 2006, 12, 435-445.   | 1.6 | 26        |
| 977 | The Yng1p Plant Homeodomain Finger Is a Methyl-Histone Binding Module That Recognizes Lysine 4-Methylated Histone H3. <i>Molecular and Cellular Biology</i> , 2006, 26, 7871-7879.  | 1.1 | 142       |
| 978 | Yeast Nuclear Envelope Subdomains with Distinct Abilities to Resist Membrane Expansion. <i>Molecular Biology of the Cell</i> , 2006, 17, 1768-1778.   | 0.9 | 88        |
| 979 | The yeast Igl family member Sro7p is an effector of the secretory Rab GTPase Sec4p. <i>Journal of Cell Biology</i> , 2006, 172, 55-66.  | 2.3 | 81        |
| 980 | SIZ1/SIZ2 Control of Chromosome Transmission Fidelity Is Mediated by the Sumoylation of Topoisomerase II. <i>Genetics</i> , 2006, 172, 783-794.   | 1.2 | 81        |
| 981 | The Role of Cdc55 in the Spindle Checkpoint Is through Regulation of Mitotic Exit in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2006, 17, 658-666.  | 0.9 | 59        |
| 982 | The Phosphatidylinositol 4,5-Biphosphate and TORC2 Binding Proteins Slm1 and Slm2 Function in Sphingolipid Regulation. <i>Molecular and Cellular Biology</i> , 2006, 26, 5861-5875.   | 1.1 | 125       |
| 983 | Nuclear Recycling of the Pre-60S Ribosomal Subunit-Associated Factor Arx1 Depends on Rei1 in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2006, 26, 3718-3727.                                   | 1.1 | 66        |
| 984 | Glycosylphosphatidylinositol-anchored Proteins Are Required for the Transport of Detergent-resistant Microdomain-associated Membrane Proteins Tat2p and Fur4p. <i>Journal of Biological Chemistry</i> , 2006, 281, 4013-4023. | 1.6 | 45        |
| 985 | Analysis of Mitogen-Activated Protein Kinase Signaling Specificity in Response to Hyperosmotic Stress: Use of an Analog-Sensitive HOG1 Allele. <i>Eukaryotic Cell</i> , 2006, 5, 1215-1228.                                   | 3.4 | 70        |
| 986 | A subset of Mer1p-dependent introns requires Bud13p for splicing activation and nuclear retention. <i>Rna</i> , 2006, 12, 1361-1372.  | 1.6 | 31        |
| 987 | Cortical Recruitment and Nuclear Cytoplasmic Shuttling of Scd5p, a Protein Phosphatase-1-targeting Protein Involved in Actin Organization and Endocytosis. <i>Molecular Biology of the Cell</i> , 2006, 17, 251-262.          | 0.9 | 15        |
| 988 | Role of the Transcription Activator Ste12p as a Repressor of PRY3 Expression. <i>Molecular and Cellular Biology</i> , 2006, 26, 7901-7912.  | 1.1 | 11        |
| 989 | Four Novel Suppressors of <i>gic1 gic2</i> and Their Roles in Cytokinesis and Polarized Cell Growth in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2006, 174, 665-678.  | 1.2 | 19        |
| 990 | Actin-based Motility during Endocytosis in Budding Yeast. <i>Molecular Biology of the Cell</i> , 2006, 17, 1354-1363.   | 0.9 | 59        |
| 991 | Methylation of Histone H3 Mediates the Association of the NuA3 Histone Acetyltransferase with Chromatin. <i>Molecular and Cellular Biology</i> , 2006, 26, 3018-3028.   | 1.1 | 65        |



| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 992  | Ubp8p, a Histone Deubiquitinase Whose Association with SAGA Is Mediated by Sgf11p, Differentially Regulates Lysine 4 Methylation of Histone H3 In Vivo. <i>Molecular and Cellular Biology</i> , 2006, 26, 3339-3352.                            | 1.1 | 83        |
| 993  | PER1 Is Required for GPI-Phospholipase A2 Activity and Involved in Lipid Remodeling of GPI-anchored Proteins. <i>Molecular Biology of the Cell</i> , 2006, 17, 5253-5264.   | 0.9 | 103       |
| 994  | Skp1p Regulates Soi3p/Rav1p Association with Endosomal Membranes but Is Not Required for Vacuolar ATPase Assembly. <i>Eukaryotic Cell</i> , 2006, 5, 2104-2113.   | 3.4 | 15        |
| 995  | Contributions of Trf4p- and Trf5p-dependent polyadenylation to the processing and degradative functions of the yeast nuclear exosome. <i>Rna</i> , 2006, 12, 26-32.   | 1.6 | 100       |
| 996  | Glc7p Regulates Reg1 Phosphatase Signals to Yck1,2 Casein Kinase 1 to Regulate Transport Activity and Glucose-Induced Inactivation of <i>Saccharomyces Maltose Permease</i> . <i>Genetics</i> , 2006, 172, 1427-1439.                           | 1.2 | 38        |
| 997  | Chs5/6 Complex: A Multiprotein Complex That Interacts with and Conveys Chitin Synthase III from the Trans-Golgi Network to the Cell Surface. <i>Molecular Biology of the Cell</i> , 2006, 17, 4157-4166.  | 0.9 | 68        |
| 998  | Purine Biosynthesis, Riboflavin Production, and Trophic-Phase Span Are Controlled by a Myb-Related Transcription Factor in the Fungus <i>Ashbya gossypii</i> . <i>Applied and Environmental Microbiology</i> , 2006, 72, 5052-5060.             | 1.4 | 60        |
| 999  | The role of the integral membrane nucleoporins Ndc1p and Pom152p in nuclear pore complex assembly and function. <i>Journal of Cell Biology</i> , 2006, 173, 361-371.  | 2.3 | 86        |
| 1000 | Identification of Novel Mutations in ACT1 and SLA2 That Suppress the Actin-Cable-Overproducing Phenotype Caused by Overexpression of a Dominant Active Form of Bni1p in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2006, 173, 527-539. | 1.2 | 8         |
| 1001 | The E and G Subunits of the Yeast V-ATPase Interact Tightly and Are Both Present at More Than One Copy per V1 Complex. <i>Journal of Biological Chemistry</i> , 2006, 281, 22752-22760.   | 1.6 | 65        |
| 1002 | Adaptive evolution by mutations in the FLO11 gene. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 11228-11233.   | 3.3 | 197       |
| 1003 | Global Replication-Independent Histone H4 Exchange in Budding Yeast. <i>Eukaryotic Cell</i> , 2006, 5, 1780-1787.   | 3.4 | 22        |
| 1004 | Doa1 Is a Cdc48 Adapter That Possesses a Novel Ubiquitin Binding Domain. <i>Molecular and Cellular Biology</i> , 2006, 26, 822-830.   | 1.1 | 78        |
| 1005 | Functional genomics of genes with small open reading frames (sORFs) in <i>S. cerevisiae</i> . <i>Genome Research</i> , 2006, 16, 365-373.   | 2.4 | 193       |
| 1006 | Characterization of the Yeast Amphiphysins Rvs161p and Rvs167p Reveals Roles for the Rvs Heterodimer In Vivo. <i>Molecular Biology of the Cell</i> , 2006, 17, 1306-1321.   | 0.9 | 54        |
| 1007 | Erv26p Directs Pro-Alkaline Phosphatase into Endoplasmic Reticulum-derived Coat Protein Complex II Transport Vesicles. <i>Molecular Biology of the Cell</i> , 2006, 17, 4780-4789.  | 0.9 | 34        |
| 1008 | SUMO modifications control assembly of synaptonemal complex and polycomplex in meiosis of <i>Saccharomyces cerevisiae</i> . <i>Genes and Development</i> , 2006, 20, 2067-2081.   | 2.7 | 236       |
| 1009 | Phosphorylation of the Chromosomal Passenger Protein Bir1 Is Required for Localization of Ndc10 to the Spindle during Anaphase and Full Spindle Elongation. <i>Molecular Biology of the Cell</i> , 2006, 17, 1065-1074.                         | 0.9 | 39        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1010 | Regulation of the Oxidative Stress Response Through Slt2p-Dependent Destruction of Cyclin C in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2006, 172, 1477-1486.   | 1.2 | 55        |
| 1011 | An FHA domain-mediated protein interaction network of Rad53 reveals its role in polarized cell growth. <i>Journal of Cell Biology</i> , 2006, 175, 743-753.  | 2.3 | 85        |
| 1012 | The mRNA Export Factor Sus1 Is Involved in Spt/Ada/Gcn5 Acetyltransferase-mediated H2B Deubiquitinylation through Its Interaction with Ubp8 and Sgf11. <i>Molecular Biology of the Cell</i> , 2006, 17, 4228-4236. | 0.9 | 115       |
| 1013 | Recruitment of Atg9 to the preautophagosomal structure by Atg11 is essential for selective autophagy in budding yeast. <i>Journal of Cell Biology</i> , 2006, 175, 925-935.  | 2.3 | 185       |
| 1014 | Multiple Endoplasmic Reticulum-to-Nucleus Signaling Pathways Coordinate Phospholipid Metabolism with Gene Expression by Distinct Mechanisms. <i>Journal of Biological Chemistry</i> , 2006, 281, 24070-24083.      | 1.6 | 51        |
| 1015 | A Genomewide Screen for Petite-negative Yeast Strains Yields a New Subunit of the i-AAA Protease Complex. <i>Molecular Biology of the Cell</i> , 2006, 17, 213-226.  | 0.9 | 73        |
| 1016 | Enolase Activates Homotypic Vacuole Fusion and Protein Transport to the Vacuole in Yeast. <i>Journal of Biological Chemistry</i> , 2006, 281, 14523-14528.   | 1.6 | 56        |
| 1017 | Regulation of Mating and Filamentation Genes by Two Distinct Ste12 Complexes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2006, 26, 4794-4805.                                     | 1.1 | 126       |
| 1018 | Independent Recruitment of Mediator and SAGA by the Activator Met4. <i>Molecular and Cellular Biology</i> , 2006, 26, 3149-3163.   | 1.1 | 36        |
| 1019 | The Stress-activated Mitogen-activated Protein Kinase Signaling Cascade Promotes Exit from Mitosis. <i>Molecular Biology of the Cell</i> , 2006, 17, 3136-3146.  | 0.9 | 31        |
| 1020 | Phosphorylation of histone H4 Ser1 regulates sporulation in yeast and is conserved in fly and mouse spermatogenesis. <i>Genes and Development</i> , 2006, 20, 2580-2592.   | 2.7 | 94        |
| 1021 | Ndj1, a Telomere-Associated Protein, Promotes Meiotic Recombination in Budding Yeast. <i>Molecular and Cellular Biology</i> , 2006, 26, 3683-3694.   | 1.1 | 50        |
| 1022 | Autophagy Counterbalances Endoplasmic Reticulum Expansion during the Unfolded Protein Response. <i>PLoS Biology</i> , 2006, 4, e423.   | 2.6 | 860       |
| 1023 | The Brl Domain in Sec63p Is Required for Assembly of Functional Endoplasmic Reticulum Translocons. <i>Journal of Biological Chemistry</i> , 2006, 281, 7899-7906.  | 1.6 | 44        |
| 1024 | Consequences of Defective Tubulin Folding on Heterodimer Levels, Mitosis and Spindle Morphology in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2006, 173, 635-646.   | 1.2 | 20        |
| 1025 | Comprehensive Mutational Analysis of Yeast DEXD/H Box RNA Helicases Required for Small Ribosomal Subunit Synthesis. <i>Molecular and Cellular Biology</i> , 2006, 26, 1183-1194.                                   | 1.1 | 62        |
| 1026 | $\beta$ -Tubulin Is Required for Proper Recruitment and Assembly of Kar9-Bim1 Complexes in Budding Yeast. <i>Molecular Biology of the Cell</i> , 2006, 17, 4420-4434.  | 0.9 | 44        |
| 1027 | Multiple Approaches to Study <i>S. cerevisiae</i> Rad9, a Prototypical Checkpoint Protein. <i>Methods in Enzymology</i> , 2006, 409, 131-150.  | 0.4 | 9         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1028 | Yeast Ntr1/Spp382 Mediates Prp43 Function in Postspliosomes. <i>Molecular and Cellular Biology</i> , 2006, 26, 6016-6023.  | 1.1 | 56        |
| 1029 | Ccr4 contributes to tolerance of replication stress through control of CRT1 mRNA poly(A) tail length. <i>Journal of Cell Science</i> , 2006, 119, 5178-5192.   | 1.2 | 57        |
| 1030 | Budding yeast Hed1 down-regulates the mitotic recombination machinery when meiotic recombination is impaired. <i>Genes and Development</i> , 2006, 20, 1766-1775.  | 2.7 | 124       |
| 1031 | Chromatin-Modifying Enzymes Are Essential When the <i>Saccharomyces cerevisiae</i> Morphogenesis Checkpoint Is Constitutively Activated. <i>Genetics</i> , 2006, 174, 1135-1149.   | 1.2 | 16        |
| 1032 | Suppressor Analysis of a Histone Defect Identifies a New Function for the Hda1 Complex in Chromosome Segregation. <i>Genetics</i> , 2006, 173, 435-450.  | 1.2 | 7         |
| 1033 | ELM1 Is Required for Multidrug Resistance in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2006, 173, 1919-1937.   | 1.2 | 17        |
| 1034 | <i>Saccharomyces cerevisiae</i> Mer2, Mei4 and Rec114 Form a Complex Required for Meiotic Double-Strand Break Formation. <i>Genetics</i> , 2006, 173, 1969-1981.   | 1.2 | 110       |
| 1035 | Chemical Inactivation of Cdc7 Kinase in Budding Yeast Results in a Reversible Arrest That Allows Efficient Cell Synchronization Prior to Meiotic Recombination. <i>Genetics</i> , 2006, 174, 1767-1774.  | 1.2 | 56        |
| 1036 | Comprehensive Mutational Analysis of Yeast DEX/H Box RNA Helicases Involved in Large Ribosomal Subunit Biogenesis. <i>Molecular and Cellular Biology</i> , 2006, 26, 1195-1208.  | 1.1 | 63        |
| 1037 | Evidence Suggesting that Pif1 Helicase Functions in DNA Replication with the Dna2 Helicase/Nuclease and DNA Polymerase $\epsilon$ . <i>Molecular and Cellular Biology</i> , 2006, 26, 2490-2500.   | 1.1 | 184       |
| 1038 | A conserved late endosome targeting signal required for Doa4 deubiquitylating enzyme function. <i>Journal of Cell Biology</i> , 2006, 175, 825-835.  | 2.3 | 41        |
| 1039 | Assortment of Phosphatidylinositol 3-Kinase Complexes Atg14p Directs Association of Complex I to the Pre-autophagosomal Structure in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2006, 17, 1527-1539.                     | 0.9 | 202       |
| 1040 | Domains within the GARP Subunit Vps54 Confer Separate Functions in Complex Assembly and Early Endosome Recognition. <i>Molecular Biology of the Cell</i> , 2006, 17, 1859-1870.  | 0.9 | 43        |
| 1041 | Yeast-to-Hyphal Transition Triggers Formin-dependent Golgi Localization to the Growing Tip in <i>Candida albicans</i> . <i>Molecular Biology of the Cell</i> , 2006, 17, 4364-4378.  | 0.9 | 52        |
| 1042 | Novel Function of Clathrin Light Chain in Promoting Endocytic Vesicle Formation. <i>Molecular Biology of the Cell</i> , 2006, 17, 4343-4352.   | 0.9 | 41        |
| 1043 | Yeast G1 DNA damage checkpoint regulation by H2A phosphorylation is independent of chromatin remodeling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 13771-13776.                              | 3.3 | 77        |
| 1044 | Histone chaperone Asf1 is required for histone H3 lysine 56 acetylation, a modification associated with S phase in mitosis and meiosis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 6988-6993. | 3.3 | 232       |
| 1045 | Genome-wide Transcriptional Dependence on TAF1 Functional Domains. <i>Journal of Biological Chemistry</i> , 2006, 281, 6404-6412.  | 1.6 | 13        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1046 | Nuclear Import of Ho Endonuclease Utilizes Two Nuclear Localization Signals and Four Importins of the Ribosomal Import System. <i>Journal of Biological Chemistry</i> , 2006, 281, 12218-12226.   | 1.6 | 17        |
| 1047 | Determinants of Rbp1p Localization in Specific Cytoplasmic mRNA-processing Foci, P-bodies. <i>Journal of Biological Chemistry</i> , 2006, 281, 29379-29390.   | 1.6 | 12        |
| 1048 | Mutual Antagonism of Target of Rapamycin and Calcineurin Signaling. <i>Journal of Biological Chemistry</i> , 2006, 281, 33000-33007.  | 1.6 | 64        |
| 1049 | Nutrient-dependent Multimerization of the Mammalian Target of Rapamycin through the N-terminal HEAT Repeat Region. <i>Journal of Biological Chemistry</i> , 2006, 281, 28605-28614.   | 1.6 | 67        |
| 1050 | Biofilm formation by <i>Pseudomonas fluorescens</i> WCS365: a role for LapD. <i>Microbiology (United Kingdom)</i> , 2006, 150, 1111-1120.   | 0.7 | 111       |
| 1051 | The Forkhead transcription factor Hcm1 regulates chromosome segregation genes and fills the S-phase gap in the transcriptional circuitry of the cell cycle. <i>Genes and Development</i> , 2006, 20, 2266-2278.                                   | 2.7 | 250       |
| 1052 | Pheromone-Induced Degradation of Ste12 Contributes to Signal Attenuation and the Specificity of Developmental Fate. <i>Eukaryotic Cell</i> , 2006, 5, 2147-2160.  | 3.4 | 17        |
| 1053 | Sir3 C-Terminal Domain Involvement in the Initiation and Spreading of Heterochromatin. <i>Molecular and Cellular Biology</i> , 2006, 26, 7616-7631.   | 1.1 | 23        |
| 1054 | Role of Doa1 in the <i>Saccharomyces cerevisiae</i> DNA Damage Response. <i>Molecular and Cellular Biology</i> , 2006, 26, 4122-4133.   | 1.1 | 35        |
| 1055 | Palmitoylation by the DHHC protein Pfa4 regulates the ER exit of Chs3. <i>Journal of Cell Biology</i> , 2006, 174, 19-25.   | 2.3 | 104       |
| 1056 | Phosphoregulation of Cbk1 is critical for RAM network control of transcription and morphogenesis. <i>Journal of Cell Biology</i> , 2006, 175, 755-766.  | 2.3 | 57        |
| 1057 | Distinct Roles for the Essential MYST Family HAT Esa1p in Transcriptional Silencing. <i>Molecular Biology of the Cell</i> , 2006, 17, 1744-1757.  | 0.9 | 48        |
| 1058 | The <i>Saccharomyces cerevisiae</i> Spindle Pole Body (SPB) Component Nbp1p Is Required for SPB Membrane Insertion and Interacts with the Integral Membrane Proteins Ndc1p and Mps2p. <i>Molecular Biology of the Cell</i> , 2006, 17, 1959-1970. | 0.9 | 42        |
| 1059 | In Vivo Targeting of the Yeast Pop2 Deadenylase Subunit to Reporter Transcripts Induces Their Rapid Degradation and Generates New Decay Intermediates. <i>Journal of Biological Chemistry</i> , 2006, 281, 25940-25947.                           | 1.6 | 31        |
| 1060 | Arp10p Is a Pointed-End-associated Component of Yeast Dynactin. <i>Molecular Biology of the Cell</i> , 2006, 17, 738-748.   | 0.9 | 17        |
| 1061 | SAS-mediated acetylation of histone H4 Lys 16 is required for H2A.Z incorporation at subtelomeric regions in <i>Saccharomyces cerevisiae</i> . <i>Genes and Development</i> , 2006, 20, 2507-2512.  | 2.7 | 90        |
| 1062 | Exomer: a coat complex for transport of select membrane proteins from the trans-Golgi network to the plasma membrane in yeast. <i>Journal of Cell Biology</i> , 2006, 174, 973-983.   | 2.3 | 110       |
| 1063 | A Tandem Affinity Tag for Two-step Purification under Fully Denaturing Conditions. <i>Molecular and Cellular Proteomics</i> , 2006, 5, 737-748.   | 2.5 | 315       |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1064 | Telomeric heterochromatin boundaries require NuA4-dependent acetylation of histone variant H2A.Z in <i>Saccharomyces cerevisiae</i> . <i>Genes and Development</i> , 2006, 20, 700-710.  | 2.7 | 176       |
| 1065 | Kap120 Functions as a Nuclear Import Receptor for Ribosome Assembly Factor Rpf1 in Yeast. <i>Molecular and Cellular Biology</i> , 2006, 26, 3170-3180.   | 1.1 | 17        |
| 1066 | Caffeine Targets TOR Complex I and Provides Evidence for a Regulatory Link between the FRB and Kinase Domains of Tor1p. <i>Journal of Biological Chemistry</i> , 2006, 281, 31616-31626.   | 1.6 | 164       |
| 1067 | The Regulatory Roles of the Galactose Permease and Kinase in the Induction Response of the GAL Network in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2006, 281, 13485-13492.   | 1.6 | 54        |
| 1068 | Pbn1p: An essential endoplasmic reticulum membrane protein required for protein processing in the endoplasmic reticulum of budding yeast. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 939-944. | 3.3 | 24        |
| 1069 | Dimeric organization of the yeast oligosaccharyl transferase complex. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 8947-8952.   | 3.3 | 37        |
| 1070 | Nucleosome Depletion Activates Poised RNA Polymerase III at Unconventional Transcription Sites in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2006, 281, 29155-29164.   | 1.6 | 34        |
| 1071 | Multiple Degradation Pathways for Misfolded Mutants of the Yeast Plasma Membrane ATPase, PMA1. <i>Journal of Biological Chemistry</i> , 2006, 281, 31457-31466.  | 1.6 | 23        |
| 1072 | Phosphatidylinositol 4-Phosphate Is Required for Translation Initiation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2006, 281, 38139-38149.  | 1.6 | 13        |
| 1073 | The Transcriptional Response of the Yeast Na <sup>+</sup> -ATPase ENA1 Gene to Alkaline Stress Involves Three Main Signaling Pathways*. <i>Journal of Biological Chemistry</i> , 2006, 281, 36632-36642.   | 1.6 | 80        |
| 1074 | Mapping the Functional Domains of Yeast NMD3, the Nuclear Export Adapter for the 60 S Ribosomal Subunit. <i>Journal of Biological Chemistry</i> , 2006, 281, 36579-36587.  | 1.6 | 23        |
| 1075 | Transduction of the Nitrogen Signal Activating Gln3-mediated Transcription Is Independent of Npr1 Kinase and Rsp5-Bul1/2 Ubiquitin Ligase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2006, 281, 28546-28554.        | 1.6 | 30        |
| 1076 | Ups1p, a conserved intermembrane space protein, regulates mitochondrial shape and alternative topogenesis of Mgm1p. <i>Journal of Cell Biology</i> , 2006, 173, 651-658.   | 2.3 | 92        |
| 1077 | The CPSF30 Binding Site on the NS1A Protein of Influenza A Virus Is a Potential Antiviral Target. <i>Journal of Virology</i> , 2006, 80, 3957-3965.  | 1.5 | 157       |
| 1078 | The Histone Deubiquitinating Enzyme Ubp10 Is Involved in rDNA Locus Control in <i>Saccharomyces cerevisiae</i> by Affecting Sir2p Association. <i>Genetics</i> , 2006, 174, 2249-2254.   | 1.2 | 13        |
| 1079 | The <i>S. cerevisiae</i> Rrm3p DNA helicase moves with the replication fork and affects replication of all yeast chromosomes. <i>Genes and Development</i> , 2006, 20, 3104-3116.  | 2.7 | 154       |
| 1080 | Structure and Function of the <i>Saccharomyces cerevisiae</i> Sir3 BAH Domain. <i>Molecular and Cellular Biology</i> , 2006, 26, 3256-3265.  | 1.1 | 56        |
| 1081 | Nuclear RNA surveillance in <i>Saccharomyces cerevisiae</i> : Trf4p-dependent polyadenylation of nascent hypomethylated tRNA and an aberrant form of 5S rRNA. <i>Rna</i> , 2006, 12, 508-521.  | 1.6 | 181       |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1082 | A Conserved Motif Is Prerequisite for the Interaction of NAC with Ribosomal Protein L23 and Nascent Chains. <i>Journal of Biological Chemistry</i> , 2006, 281, 2847-2857.  | 1.6 | 75        |
| 1083 | A conserved G protein (Drg1p) plays a role in regulation of invasive filamentation in <i>Candida albicans</i> . <i>Microbiology (United Kingdom)</i> , 2006, 152, 3691-3700.  | 0.7 | 13        |
| 1084 | Ribosomal DNA Transcription-Dependent Processes Interfere with Chromosome Segregation. <i>Molecular and Cellular Biology</i> , 2006, 26, 6239-6247.   | 1.1 | 38        |
| 1085 | Inhibition of homologous recombination by a cohesin-associated clamp complex recruited to the rDNA recombination enhancer. <i>Genes and Development</i> , 2006, 20, 2887-2901.  | 2.7 | 144       |
| 1086 | Linking DNA-binding proteins to their recognition sequences by using protein microarrays. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 9940-9945.  | 3.3 | 63        |
| 1087 | Novel Role for the C Terminus of <i>Saccharomyces cerevisiae</i> Rev1 in Mediating Protein-Protein Interactions. <i>Molecular and Cellular Biology</i> , 2006, 26, 8173-8182.   | 1.1 | 51        |
| 1088 | The Function and Regulation of Budding Yeast Swe1 in Response to Interrupted DNA Synthesis. <i>Molecular Biology of the Cell</i> , 2006, 17, 2746-2756.   | 0.9 | 45        |
| 1089 | RNA polymerase II elongation factors Spt4p and Spt5p play roles in transcription elongation by RNA polymerase I and rRNA processing. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 12707-12712. | 3.3 | 102       |
| 1090 | Long Chain Base Tolerance in <i>Saccharomyces cerevisiae</i> Is Induced by Retrograde Signals from the Mitochondria. <i>Journal of Biological Chemistry</i> , 2006, 281, 6376-6384.   | 1.6 | 47        |
| 1091 | Bioinformatic and Comparative Localization of Rab Proteins Reveals Functional Insights into the Uncharacterized GTPases Ypt10p and Ypt11p. <i>Molecular and Cellular Biology</i> , 2006, 26, 7299-7317.   | 1.1 | 69        |
| 1092 | Autoregulation of the mRNA export factor Yra1p requires inefficient splicing of its pre-mRNA. <i>Rna</i> , 2006, 12, 994-1006.  | 1.6 | 43        |
| 1093 | Did2 coordinates Vps4-mediated dissociation of ESCRT-III from endosomes. <i>Journal of Cell Biology</i> , 2006, 175, 715-720.   | 2.3 | 149       |
| 1094 | Characterization of Mmp37p, a <i>Saccharomyces cerevisiae</i> Mitochondrial Matrix Protein with a Role in Mitochondrial Protein Import. <i>Molecular Biology of the Cell</i> , 2006, 17, 4051-4062.   | 0.9 | 56        |
| 1095 | Fine-Structure Analysis of Ribosomal Protein Gene Transcription. <i>Molecular and Cellular Biology</i> , 2006, 26, 4853-4862.   | 1.1 | 89        |
| 1096 | The La protein functions redundantly with tRNA modification enzymes to ensure tRNA structural stability. <i>Rna</i> , 2006, 12, 644-654.  | 1.6 | 55        |
| 1097 | Dynamic microtubules are essential for efficient chromosome capture and biorientation in <i>S. cerevisiae</i> . <i>Journal of Cell Biology</i> , 2006, 175, 17-23.  | 2.3 | 19        |
| 1098 | Yeast transcripts cleaved by an internal ribozyme provide new insight into the role of the cap and poly(A) tail in translation and mRNA decay. <i>Rna</i> , 2006, 12, 1323-1337.  | 1.6 | 87        |
| 1099 | A novel role for the yeast protein kinase Dbf2p in vacuolar H <sup>+</sup> -ATPase function and sorbic acid stress tolerance. <i>Microbiology (United Kingdom)</i> , 2007, 153, 4016-4026.  | 0.7 | 25        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1100 | Cdc6 Stability Is Regulated by the Huwe1 Ubiquitin Ligase after DNA Damage. <i>Molecular Biology of the Cell</i> , 2007, 18, 3340-3350.  | 0.9 | 124       |
| 1101 | Potential Interface between Ribosomal Protein Production and Pre-rRNA Processing. <i>Molecular and Cellular Biology</i> , 2007, 27, 4815-4824.   | 1.1 | 99        |
| 1102 | Hmo1 Is Required for TOR-Dependent Regulation of Ribosomal Protein Gene Transcription. <i>Molecular and Cellular Biology</i> , 2007, 27, 8015-8026.  | 1.1 | 74        |
| 1103 | The Isw2 Chromatin-Remodeling ATPase Cooperates with the Fkh2 Transcription Factor To Repress Transcription of the B-Type Cyclin Gene CLB2. <i>Molecular and Cellular Biology</i> , 2007, 27, 2848-2860.         | 1.1 | 31        |
| 1104 | The Biologically Relevant Targets and Binding Affinity Requirements for the Function of the Yeast Actin-Binding Protein 1 Src-Homology 3 Domain Vary With Genetic Context. <i>Genetics</i> , 2007, 176, 193-208. | 1.2 | 35        |
| 1105 | Yeast P4-ATPases Drs2p and Dnf1p Are Essential Cargos of the NPFXD/Sla1p Endocytic Pathway. <i>Molecular Biology of the Cell</i> , 2007, 18, 487-500.  | 0.9 | 63        |
| 1106 | Effects of Ubiquitin System Alterations on the Formation and Loss of a Yeast Prion. <i>Journal of Biological Chemistry</i> , 2007, 282, 3004-3013.   | 1.6 | 74        |
| 1107 | The Ccr4-Not Complex Regulates Skn7 through Srb10 Kinase. <i>Eukaryotic Cell</i> , 2007, 6, 2251-2259.   | 3.4 | 25        |
| 1108 | Temporal and Spatial Control of HGC1 Expression Results in Hgc1 Localization to the Apical Cells of Hyphae in <i>Candida albicans</i> . <i>Eukaryotic Cell</i> , 2007, 6, 253-261.                               | 3.4 | 22        |
| 1109 | Regulation of Iron Homeostasis Mediated by the Heme-binding Protein Dap1 (Damage Resistance Protein) Tj ETQq1.1 0.784314 rgBT 1.6 55   | 1.6 | 55        |
| 1110 | Mutational Analysis of the Ribosomal Protein Rpl10 from Yeast. <i>Journal of Biological Chemistry</i> , 2007, 282, 32630-32639.  | 1.6 | 35        |
| 1111 | Aup1p, a Yeast Mitochondrial Protein Phosphatase Homolog, Is Required for Efficient Stationary Phase Mitophagy and Cell Survival. <i>Journal of Biological Chemistry</i> , 2007, 282, 5617-5624.                 | 1.6 | 232       |
| 1112 | Hyperphosphorylation of the C-terminal Repeat Domain of RNA Polymerase II Facilitates Dissociation of Its Complex with Mediator. <i>Journal of Biological Chemistry</i> , 2007, 282, 14113-14120.                | 1.6 | 99        |
| 1113 | Assembly factors Rpf2 and Rrs1 recruit 5S rRNA and ribosomal proteins rpL5 and rpL11 into nascent ribosomes. <i>Genes and Development</i> , 2007, 21, 2580-2592.   | 2.7 | 175       |
| 1114 | Hxk2 Regulates the Phosphorylation State of Mig1 and Therefore Its Nucleocytoplasmic Distribution. <i>Journal of Biological Chemistry</i> , 2007, 282, 4485-4493.  | 1.6 | 146       |
| 1115 | Analysis of Transcriptional Activation at a Distance in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2007, 27, 5575-5586.   | 1.1 | 98        |
| 1116 | Chromatin Disassembly from the <i>PHO5</i> Promoter Is Essential for the Recruitment of the General Transcription Machinery and Coactivators. <i>Molecular and Cellular Biology</i> , 2007, 27, 6372-6382.       | 1.1 | 83        |
| 1117 | Puf3p, a Pumilio family RNA binding protein, localizes to mitochondria and regulates mitochondrial biogenesis and motility in budding yeast. <i>Journal of Cell Biology</i> , 2007, 176, 197-207.                | 2.3 | 161       |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1118 | Identification of Yeast IQGAP (Iqg1p) as an Anaphase-Promoting-Complex Substrate and Its Role in Actomyosin-Ring-Independent Cytokinesis. <i>Molecular Biology of the Cell</i> , 2007, 18, 5139-5153.  | 0.9 | 59        |
| 1119 | The Aurora kinase Ipl1 maintains the centromeric localization of PP2A to protect cohesin during meiosis. <i>Journal of Cell Biology</i> , 2007, 176, 911-918.  | 2.3 | 53        |
| 1120 | Succinate inhibition of $\alpha$ -ketoglutarate-dependent enzymes in a yeast model of paraganglioma. <i>Human Molecular Genetics</i> , 2007, 16, 3136-3148.  | 1.4 | 155       |
| 1121 | Constitutively high dNTP concentration inhibits cell cycle progression and the DNA damage checkpoint in yeast <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 1183-1188. | 3.3 | 118       |
| 1122 | Protein characterization of <i>Saccharomyces cerevisiae</i> RNA polymerase II after <i>in vivo</i> cross-linking. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 19948-19953.                             | 3.3 | 54        |
| 1123 | Microtubule disruption stimulates P-body formation. <i>Rna</i> , 2007, 13, 493-502.  | 1.6 | 68        |
| 1124 | MSN2 and MSN4 Link Calorie Restriction and TOR to Sirtuin-Mediated Lifespan Extension in <i>Saccharomyces cerevisiae</i> . <i>PLoS Biology</i> , 2007, 5, e261.  | 2.6 | 273       |
| 1125 | Limited Functional Redundancy and Oscillation of Cyclins in Multinucleated <i>Ashbya gossypii</i> Fungal Cells. <i>Eukaryotic Cell</i> , 2007, 6, 473-486.   | 3.4 | 25        |
| 1126 | Negative Regulation of Yeast Eps15-like Arp2/3 Complex Activator, Pan1p, by the Hip1R-related Protein, Sla2p, during Endocytosis. <i>Molecular Biology of the Cell</i> , 2007, 18, 658-668.  | 0.9 | 39        |
| 1127 | Rmd9p Controls the Processing/Stability of Mitochondrial mRNAs and Its Overexpression Compensates for a Partial Deficiency of Oxa1p in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2007, 175, 1105-1115.   | 1.2 | 40        |
| 1128 | O-Mannosylation is Required for Degradation of the Endoplasmic Reticulum-associated Degradation Substrate Gas1* <sub>p</sub> via the Ubiquitin/Proteasome Pathway in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biochemistry</i> , 2007, 143, 555-567.    | 0.9 | 47        |
| 1129 | Bub1 Kinase Targets Sgo1 to Ensure Efficient Chromosome Biorientation in Budding Yeast Mitosis. <i>PLoS Genetics</i> , 2007, 3, e213.  | 1.5 | 103       |
| 1130 | 4 A Guided Tour to PCR-based Genomic Manipulations of (PCR-targeting). <i>Methods in Microbiology</i> , 2007, , 55-78.   | 0.4 | 6         |
| 1131 | Analysis of P-Body Assembly in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2007, 18, 2274-2287.   | 0.9 | 210       |
| 1132 | Requirements for nuclear localization of the Lsm2-8p complex and competition between nuclear and cytoplasmic Lsm complexes. <i>Journal of Cell Science</i> , 2007, 120, 4310-4320.   | 1.2 | 25        |
| 1133 | Direct Binding to Rsp5 Mediates Ubiquitin-independent Sorting of Sna3 via the Multivesicular Body Pathway. <i>Molecular Biology of the Cell</i> , 2007, 18, 697-706.   | 0.9 | 79        |
| 1134 | Edc3p and a glutamine/asparagine-rich domain of Lsm4p function in processing body assembly in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 2007, 179, 437-449.   | 2.3 | 411       |
| 1135 | Analysis of Origin Recognition Complex in <i>Saccharomyces cerevisiae</i> by Use of Degron Mutants. <i>Journal of Biochemistry</i> , 2007, 143, 455-465.   | 0.9 | 7         |



| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1136 | Identification and Characterization of Vnx1p, a Novel Type of Vacuolar Monovalent Cation/H <sup>+</sup> Antiporter of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2007, 282, 24284-24293.  | 1.6 | 74        |
| 1137 | Yeast Rrp14p is required for ribosomal subunit synthesis and for correct positioning of the mitotic spindle during mitosis. <i>Nucleic Acids Research</i> , 2007, 35, 1354-1366.  | 6.5 | 39        |
| 1138 | Histone H3 K4 Demethylation during Activation and Attenuation of <i>GAL1</i> Transcription in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2007, 27, 7856-7864.  | 1.1 | 40        |
| 1139 | Roles of the HEAT repeat proteins Utp10 and Utp20 in 40S ribosome maturation. <i>Rna</i> , 2007, 13, 1516-1527.   | 1.6 | 43        |
| 1140 | 5 Studying Essential Genes: Generating and Using Promoter Fusions and Conditional Alleles. <i>Methods in Microbiology</i> , 2007, 36, 79-102.   | 0.4 | 1         |
| 1141 | In Vivo Deletion Analysis of the Architecture of a Multiprotein Complex of Translation Initiation Factors. <i>Methods in Enzymology</i> , 2007, 431, 15-32.   | 0.4 | 16        |
| 1142 | Visualization of Mitochondria in Budding Yeast. <i>Methods in Cell Biology</i> , 2007, 80, 591-626.   | 0.5 | 10        |
| 1143 | Heme Levels Switch the Function of Hap1 of <i>Saccharomyces cerevisiae</i> between Transcriptional Activator and Transcriptional Repressor. <i>Molecular and Cellular Biology</i> , 2007, 27, 7414-7424.  | 1.1 | 115       |
| 1144 | A novel role in cytokinesis reveals a housekeeping function for the unfolded protein response. <i>Journal of Cell Biology</i> , 2007, 177, 1017-1027.   | 2.3 | 53        |
| 1145 | The Golgi-resident protease Kex2 acts in conjunction with Prm1 to facilitate cell fusion during yeast mating. <i>Journal of Cell Biology</i> , 2007, 176, 209-222.  | 2.3 | 34        |
| 1146 | mRNAs Encoding Polarity and Exocytosis Factors Are Cotransported with the Cortical Endoplasmic Reticulum to the Incipient Bud in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2007, 27, 3441-3455.   | 1.1 | 120       |
| 1147 | trans-SNARE complex assembly and yeast vacuole membrane fusion. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 8755-8760.  | 3.3 | 76        |
| 1148 | Ribosome Biogenesis Is Sensed at the Start Cell Cycle Checkpoint. <i>Molecular Biology of the Cell</i> , 2007, 18, 953-964.   | 0.9 | 116       |
| 1149 | Split-Ubiquitin Two-Hybrid Assay To Analyze Protein-Protein Interactions at the Endosome: Application to <i>Saccharomyces cerevisiae</i> Bro1 Interacting with ESCRT Complexes, the Doa4 Ubiquitin Hydrolase, and the Rsp5 Ubiquitin Ligase. <i>Eukaryotic Cell</i> , 2007, 6, 1266-1277. | 3.4 | 31        |
| 1150 | Endocytic Recycling in Yeast Is Regulated by Putative Phospholipid Translocases and the Ypt31p/32p/Rcy1p Pathway. <i>Molecular Biology of the Cell</i> , 2007, 18, 295-312.   | 0.9 | 145       |
| 1151 | Increased association of telomerase with short telomeres in yeast. <i>Genes and Development</i> , 2007, 21, 1726-1730.  | 2.7 | 117       |
| 1152 | Interaction of the Endocytic Scaffold Protein Pan1 with the Type I Myosins Contributes to the Late Stages of Endocytosis. <i>Molecular Biology of the Cell</i> , 2007, 18, 2893-2903.   | 0.9 | 36        |
| 1153 | Yeast Formins Bni1 and Bnr1 Utilize Different Modes of Cortical Interaction during the Assembly of Actin Cables. <i>Molecular Biology of the Cell</i> , 2007, 18, 1826-1838.  | 0.9 | 109       |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1154 | Membrane-active Compounds Activate the Transcription Factors Pdr1 and Pdr3 Connecting Pleiotropic Drug Resistance and Membrane Lipid Homeostasis in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2007, 18, 4932-4944. | 0.9 | 47        |
| 1155 | Erv14 family cargo receptors are necessary for ER exit during sporulation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 2007, 120, 908-916.   | 1.2 | 50        |
| 1156 | Misfolded Proteins Traffic from the Endoplasmic Reticulum (ER) Due to ER Export Signals. <i>Molecular Biology of the Cell</i> , 2007, 18, 455-463.  | 0.9 | 96        |
| 1157 | Accumulation of Polyadenylated mRNA, Pab1p, eIF4E, and eIF4G with P-Bodies in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2007, 18, 2592-2602.   | 0.9 | 138       |
| 1158 | Efficient Tor Signaling Requires a Functional Class C Vps Protein Complex in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2007, 176, 2139-2150.  | 1.2 | 81        |
| 1159 | The Plasma Membrane Proteins Prm1 and Fig1 Ascertain Fidelity of Membrane Fusion during Yeast Mating. <i>Molecular Biology of the Cell</i> , 2007, 18, 547-556.   | 0.9 | 87        |
| 1160 | Cdc37p Is Required for Stress-Induced High-Osmolarity Glycerol and Protein Kinase C Mitogen-Activated Protein Kinase Pathway Functionality by Interaction with Hog1p and Slt2p (Mpk1p). <i>Eukaryotic Cell</i> , 2007, 6, 521-532.                | 3.4 | 58        |
| 1161 | Specific Defects in Different Transcription Complexes Compensate for the Requirement of the Negative Cofactor 2 Repressor in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2007, 176, 125-138.  | 1.2 | 11        |
| 1162 | Alternative Modes of Organellar Segregation during Sporulation in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2007, 6, 2009-2017.  | 3.4 | 52        |
| 1163 | Nucleotide Exchange Factors for Hsp70s Are Required for [URE3] Prion Propagation in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2007, 18, 2149-2154.   | 0.9 | 98        |
| 1164 | Ent3p and Ent5p Exhibit Cargo-specific Functions in Trafficking Proteins between the Trans-Golgi Network and the Endosomes in Yeast. <i>Molecular Biology of the Cell</i> , 2007, 18, 1803-1815.  | 0.9 | 74        |
| 1165 | An Interrelationship Between Autophagy and Filamentous Growth in Budding Yeast. <i>Genetics</i> , 2007, 177, 205-214.   | 1.2 | 36        |
| 1166 | Mitotic Cdc6 Stabilizes Anaphase-Promoting Complex Substrates by a Partially Cdc28-Independent Mechanism, and This Stabilization Is Suppressed by Deletion of Cdc55. <i>Molecular and Cellular Biology</i> , 2007, 27, 1158-1171.                 | 1.1 | 26        |
| 1167 | Adjacent positioning of cellular structures enabled by a Cdc42 GTPase-activating protein-mediated zone of inhibition. <i>Journal of Cell Biology</i> , 2007, 179, 1375-1384.  | 2.3 | 106       |
| 1168 | Variable and Tissue-Specific Subunit Composition of Mitochondrial m-AAA Protease Complexes Linked to Hereditary Spastic Paraplegia. <i>Molecular and Cellular Biology</i> , 2007, 27, 758-767.  | 1.1 | 172       |
| 1169 | Probing the Membrane Environment of the TOR Kinases Reveals Functional Interactions between TORC1, Actin, and Membrane Trafficking in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2007, 18, 2779-2794.               | 0.9 | 91        |
| 1170 | The specialized cytosolic J-protein, Jjj1, functions in 60S ribosomal subunit biogenesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 1558-1563.  | 3.3 | 80        |
| 1171 | A SAGA-Independent Function of SPT3 Mediates Transcriptional Deregulation in a Mutant of the Ccr4-Not Complex in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2007, 177, 123-135.  | 1.2 | 19        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1172 | Atg18 Regulates Organelle Morphology and Fab1 Kinase Activity Independent of Its Membrane Recruitment by Phosphatidylinositol 3,5-Bisphosphate. <i>Molecular Biology of the Cell</i> , 2007, 18, 4232-4244.   | 0.9 | 112       |
| 1173 | SCFCdc4-mediated Degradation of the Hac1p Transcription Factor Regulates the Unfolded Protein Response in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2007, 18, 426-440.         | 0.9 | 30        |
| 1174 | Involvement of Specific COPI Subunits in Protein Sorting from the Late Endosome to the Vacuole in Yeast. <i>Molecular and Cellular Biology</i> , 2007, 27, 526-540.   | 1.1 | 61        |
| 1175 | Spc24 and Stu2 Promote Spindle Integrity When DNA Replication Is Stalled. <i>Molecular Biology of the Cell</i> , 2007, 18, 2805-2816.   | 0.9 | 22        |
| 1176 | Contribution of Trf4/5 and the Nuclear Exosome to Genome Stability Through Regulation of Histone mRNA Levels in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2007, 175, 993-1010.                      | 1.2 | 59        |
| 1177 | Sequential and Distinct Roles of the Cadherin Domain-containing Protein Axl2p in Cell Polarization in Yeast Cell Cycle. <i>Molecular Biology of the Cell</i> , 2007, 18, 2542-2560.                           | 0.9 | 58        |
| 1178 | Nuclear fusion during yeast mating occurs by a three-step pathway. <i>Journal of Cell Biology</i> , 2007, 179, 659-670.   | 2.3 | 42        |
| 1179 | Flo11p-Independent Control of $\alpha$ -Mat $\alpha$ -Formation by Hsp70 Molecular Chaperones and Nucleotide Exchange Factors in Yeast. <i>Genetics</i> , 2007, 177, 1679-1689.                               | 1.2 | 25        |
| 1180 | The p23 molecular chaperone promotes functional telomerase complexes through DNA dissociation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 5765-5770. | 3.3 | 52        |
| 1181 | Stable Preanaphase Spindle Positioning Requires Bud6p and an Apparent Interaction between the Spindle Pole Bodies and the Neck. <i>Eukaryotic Cell</i> , 2007, 6, 797-807.                                    | 3.4 | 8         |
| 1182 | DNA breaks are masked by multiple Rap1 binding in yeast: implications for telomere capping and telomerase regulation. <i>Genes and Development</i> , 2007, 21, 292-302.                                       | 2.7 | 81        |
| 1183 | The decapping activator Lsm1p-7p-Pat1p complex has the intrinsic ability to distinguish between oligoadenylated and polyadenylated RNAs. <i>Rna</i> , 2007, 13, 998-1016.                                     | 1.6 | 130       |
| 1184 | Identification of a Vacuole-associated Metalloreductase and Its Role in Ctr2-mediated Intracellular Copper Mobilization. <i>Journal of Biological Chemistry</i> , 2007, 282, 21629-21638.                     | 1.6 | 95        |
| 1185 | Transcriptional responses to fatty acid are coordinated by combinatorial control. <i>Molecular Systems Biology</i> , 2007, 3, 115.  | 3.2 | 58        |
| 1186 | A Cadmium-transporting P1B-type ATPase in Yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2007, 282, 947-955.  | 1.6 | 73        |
| 1187 | Atg27 Is Required for Autophagy-dependent Cycling of Atg9. <i>Molecular Biology of the Cell</i> , 2007, 18, 581-593.  | 0.9 | 162       |
| 1188 | Ure2p Function Is Enhanced by Its Prion Domain in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2007, 176, 1557-1565.   | 1.2 | 72        |
| 1189 | Novelsfi1 Alleles Uncover Additional Functions for Sfi1p in Bipolar Spindle Assembly and Function. <i>Molecular Biology of the Cell</i> , 2007, 18, 2047-2056.  | 0.9 | 27        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1190 | An Approach to Studying the Localization and Dynamics of Eukaryotic Translation Factors in Live Yeast Cells. <i>Methods in Enzymology</i> , 2007, 431, 33-45.   | 0.4 | 4         |
| 1191 | The Overexpression of a <i>Saccharomyces cerevisiae</i> Centromeric Histone H3 Variant Mutant Protein Leads to a Defect in Kinetochore Biorientation. <i>Genetics</i> , 2007, 175, 513-525.                                 | 1.2 | 29        |
| 1192 | Prohibitins Interact Genetically with Atp23, a Novel Processing Peptidase and Chaperone for the F1FO-ATP Synthase. <i>Molecular Biology of the Cell</i> , 2007, 18, 627-635.  | 0.9 | 124       |
| 1193 | Grd19/Snx3p functions as a cargo-specific adapter for retromer-dependent endocytic recycling. <i>Journal of Cell Biology</i> , 2007, 177, 115-125.  | 2.3 | 153       |
| 1194 | Regulation of rRNA Synthesis by TATA-Binding Protein-Associated Factor Mot1. <i>Molecular and Cellular Biology</i> , 2007, 27, 2886-2896.   | 1.1 | 13        |
| 1195 | Role for the Ssu72 C-Terminal Domain Phosphatase in RNA Polymerase II Transcription Elongation. <i>Molecular and Cellular Biology</i> , 2007, 27, 926-936.  | 1.1 | 50        |
| 1196 | TFIIS elongation factor and Mediator act in conjunction during transcription initiation <i>in vivo</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 16062-16067.   | 3.3 | 61        |
| 1197 | The Transmembrane Domain of Acid Trehalase Mediates Ubiquitin-independent Multivesicular Body Pathway Sorting. <i>Molecular Biology of the Cell</i> , 2007, 18, 2511-2524.  | 0.9 | 24        |
| 1198 | A Protein Interaction Map of the Mitotic Spindle. <i>Molecular Biology of the Cell</i> , 2007, 18, 3800-3809.   | 0.9 | 88        |
| 1199 | <i>Saccharomyces cerevisiae</i> Ebs1p is a putative ortholog of human Smg7 and promotes nonsense-mediated mRNA decay. <i>Nucleic Acids Research</i> , 2007, 35, 7688-7697.  | 6.5 | 63        |
| 1200 | Btn2, a Hook1 Ortholog and Potential Batten Disease-Related Protein, Mediates Late Endosome-Golgi Protein Sorting in Yeast. <i>Molecular and Cellular Biology</i> , 2007, 27, 605-621.                                      | 1.1 | 68        |
| 1201 | DNA Recombination-Initiation Plays a Role in the Extremely Biased Inheritance of Yeast [ $\rho$ ] Mitochondrial DNA That Contains the Replication Origin ori5. <i>Molecular and Cellular Biology</i> , 2007, 27, 1133-1145. | 1.1 | 49        |
| 1202 | Assembly of Regulatory Factors on rRNA and Ribosomal Protein Genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2007, 27, 6686-6705.  | 1.1 | 69        |
| 1203 | Effect of 21 Different Nitrogen Sources on Global Gene Expression in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2007, 27, 3065-3086.   | 1.1 | 237       |
| 1204 | Composite synthetic lethal identification of membrane traffic inhibitors. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 6235-6240.                                    | 3.3 | 18        |
| 1205 | Mechanism of Dun1 Activation by Rad53 Phosphorylation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2007, 282, 986-995.   | 1.6 | 68        |
| 1206 | The Central Role of PDR1 in the Foundation of Yeast Drug Resistance. <i>Journal of Biological Chemistry</i> , 2007, 282, 5063-5074.   | 1.6 | 64        |
| 1207 | Phosphatidylethanolamine, a Limiting Factor of Autophagy in Yeast Strains Bearing a Defect in the Carboxypeptidase Y Pathway of Vacuolar Targeting. <i>Journal of Biological Chemistry</i> , 2007, 282, 16736-16743.        | 1.6 | 35        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 1208 | RAVE Is Essential for the Efficient Assembly of the C Subunit with the Vacuolar H <sup>+</sup> -ATPase. <i>Journal of Biological Chemistry</i> , 2007, 282, 26185-26194.   | 1.6  | 73        |
| 1209 | Membrane Topology of the Alg14 Endoplasmic Reticulum UDP-GlcNAc Transferase Subunit. <i>Journal of Biological Chemistry</i> , 2007, 282, 29081-29088.  | 1.6  | 20        |
| 1210 | Hse1, a Component of the Yeast Hrs-STAM Ubiquitin-sorting Complex, Associates with Ubiquitin Peptidases and a Ligase to Control Sorting Efficiency into Multivesicular Bodies. <i>Molecular Biology of the Cell</i> , 2007, 18, 324-335. | 0.9  | 63        |
| 1211 | NPFXD-mediated Endocytosis Is Required for Polarity and Function of a Yeast Cell Wall Stress Sensor. <i>Molecular Biology of the Cell</i> , 2007, 18, 57-65.   | 0.9  | 85        |
| 1212 | Mechanism for the degradation of origin recognition complex containing Orc5p with a defective Walker A motif and its suppression by over-production of Orc4p in yeast cells. <i>Biochemical Journal</i> , 2007, 402, 397-403.            | 1.7  | 8         |
| 1213 | Transcriptional regulation of phospholipid biosynthesis is linked to fatty acid metabolism by an acyl-CoA-binding-protein-dependent mechanism in <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 2007, 407, 219-230.      | 1.7  | 45        |
| 1214 | The ergosterol biosynthesis inhibitor zaragozic acid promotes vacuolar degradation of the tryptophan permease Tat2p in yeast. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2007, 1768, 1681-1690.                               | 1.4  | 18        |
| 1215 | Cis1/Atg31 is required for autophagosome formation in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2007, 356, 405-410.   | 1.0  | 91        |
| 1216 | Activation of a novel pathway involving Mms1 and Rad59 in sgs1 cells. <i>Biochemical and Biophysical Research Communications</i> , 2007, 356, 1031-1037.   | 1.0  | 5         |
| 1217 | Kinetochores Orientation during Meiosis Is Controlled by Aurora B and the Monopolin Complex. <i>Cell</i> , 2007, 128, 477-490.   | 13.5 | 131       |
| 1218 | Sequential E2s Drive Polyubiquitin Chain Assembly on APC Targets. <i>Cell</i> , 2007, 130, 127-139.  | 13.5 | 227       |
| 1219 | Functional Architecture of RNA Polymerase I. <i>Cell</i> , 2007, 131, 1260-1272.   | 13.5 | 192       |
| 1220 | Yeast Mitochondrial Division and Distribution Require the Cortical Num1 Protein. <i>Developmental Cell</i> , 2007, 12, 363-375.  | 3.1  | 95        |
| 1221 | Mutual Interdependence of MS11 (CAC3) and YAK1 in <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 2007, 368, 30-43.  | 2.0  | 10        |
| 1222 | Crystal Structure, Biochemical and Genetic Characterization of Yeast and <i>E. cuniculi</i> TAF15 N-terminal Domain: Implications for TFIID Assembly. <i>Journal of Molecular Biology</i> , 2007, 368, 1292-1306.                        | 2.0  | 21        |
| 1223 | The role of Pif1p, a DNA helicase in <i>Saccharomyces cerevisiae</i> , in maintaining mitochondrial DNA. <i>Mitochondrion</i> , 2007, 7, 211-222.  | 1.6  | 43        |
| 1224 | H2B Ubiquitylation Acts as a Barrier to Ctk1 Nucleosomal Recruitment Prior to Removal by Ubp8 within a SAGA-Related Complex. <i>Molecular Cell</i> , 2007, 27, 275-288.  | 4.5  | 196       |
| 1225 | Evolution of Ime2 Phosphorylation Sites on Cdk1 Substrates Provides a Mechanism to Limit the Effects of the Phosphatase Cdc14 in Meiosis. <i>Molecular Cell</i> , 2007, 25, 689-702.   | 4.5  | 70        |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 1226 | Meiosis-Specific Destruction of the Ume6p Repressor by the Cdc20-Directed APC/C. <i>Molecular Cell</i> , 2007, 27, 951-961.   | 4.5  | 63        |
| 1227 | Structural Organization of the Anaphase-Promoting Complex Bound to the Mitotic Activator Slp1. <i>Molecular Cell</i> , 2007, 28, 871-885.   | 4.5  | 54        |
| 1228 | Interplay of Chromatin Modifiers on a Short Basic Patch of Histone H4 Tail Defines the Boundary of Telomeric Heterochromatin. <i>Molecular Cell</i> , 2007, 28, 1002-1014.  | 4.5  | 188       |
| 1229 | Role of the Conserved Sir3-BAH Domain in Nucleosome Binding and Silent Chromatin Assembly. <i>Molecular Cell</i> , 2007, 28, 1015-1028.   | 4.5  | 145       |
| 1230 | The utilization of a <i>Saccharomyces cerevisiae</i> HUG1P-GFP promoter reporter construct for the selective detection of DNA damage. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2007, 633, 21-34. | 0.9  | 40        |
| 1231 | Urmylation controls Nil1p and Gln3p-dependent expression of nitrogen-catabolite repressed genes in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 2007, 581, 541-550.  | 1.3  | 25        |
| 1232 | Functional analyses of the extra- and intracellular domains of the yeast cell wall integrity sensors Mid2 and Wsc1. <i>FEBS Letters</i> , 2007, 581, 4495-4500.   | 1.3  | 56        |
| 1233 | Spore germination in <i>Saccharomyces cerevisiae</i> : global gene expression patterns and cell cycle landmarks. <i>Genome Biology</i> , 2007, 8, R241.   | 13.9 | 54        |
| 1234 | The Sas3p and Gcn5p histone acetyltransferases are recruited to similar genes. <i>Genome Biology</i> , 2007, 8, R119.   | 13.9 | 43        |
| 1235 | Fluorescence Imaging of Mitochondria in Yeast. <i>Methods in Molecular Biology</i> , 2007, 372, 433-459.  | 0.4  | 4         |
| 1236 | Cell-Free Production of Membrane Proteins in the Presence of Detergents. , 0, , 165-178.  |      | 1         |
| 1237 | Disrupting Mediator with a Short Peptide Ligand. <i>ChemBioChem</i> , 2007, 8, 1233-1236.   | 1.3  | 4         |
| 1238 | Immobility, inheritance and plasticity of shape of the yeast nucleus. <i>BMC Cell Biology</i> , 2007, 8, 47.  | 3.0  | 30        |
| 1239 | Inactivation of NMD increases viability of sup45 nonsense mutants in <i>Saccharomyces cerevisiae</i> . <i>BMC Molecular Biology</i> , 2007, 8, 71.  | 3.0  | 15        |
| 1240 | Disintegrator vectors for single-copy yeast chromosomal integration. <i>Yeast</i> , 2007, 24, 447-455.  | 0.8  | 47        |
| 1241 | Delayed accumulation of the yeast G1 cyclins Cln1 and Cln2 and the F-box protein Grr1 in response to glucose. <i>Yeast</i> , 2007, 24, 419-429.   | 0.8  | 5         |
| 1242 | The effect of tea tree oil and antifungal agents on a reporter for yeast cell integrity signalling. <i>Yeast</i> , 2007, 24, 321-334.   | 0.8  | 34        |
| 1243 | Dosage rescue by UBC4 restores cell wall integrity in <i>Saccharomyces cerevisiae</i> lacking the myosin type II gene MYO1. <i>Yeast</i> , 2007, 24, 343-355.   | 0.8  | 8         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1244 | Bimolecular fluorescence complementation analysis system for in vivo detection of protein-protein interaction in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2007, 24, 767-775. | 0.8 | 173       |
| 1245 | Cdk and APC activities limit the spindle-stabilizing function of Fin1 to anaphase. <i>Nature Cell Biology</i> , 2007, 9, 106-112.  | 4.6 | 110       |
| 1246 | Cdk1 coordinates cell-surface growth with the cell cycle. <i>Nature Cell Biology</i> , 2007, 9, 506-515.   | 4.6 | 132       |
| 1247 | TOR signalling regulates mitotic commitment through the stress MAP kinase pathway and the Polo and Cdc2 kinases. <i>Nature Cell Biology</i> , 2007, 9, 1263-1272.                    | 4.6 | 161       |
| 1248 | Genetic basis of proteome variation in yeast. <i>Nature Genetics</i> , 2007, 39, 1369-1375.  | 9.4 | 767       |
| 1249 | Regulation of histone modification and cryptic transcription by the Bur1 and Paf1 complexes. <i>EMBO Journal</i> , 2007, 26, 4646-4656.  | 3.5 | 118       |
| 1250 | Pkh-kinases control eisosome assembly and organization. <i>EMBO Journal</i> , 2007, 26, 4946-4955.   | 3.5 | 117       |
| 1251 | prp8 mutations that cause human retinitis pigmentosa lead to a U5 snRNP maturation defect in yeast. <i>Nature Structural and Molecular Biology</i> , 2007, 14, 1077-1083.            | 3.6 | 93        |
| 1252 | Hierarchy of Atg proteins in pre-autophagosomal structure organization. <i>Genes To Cells</i> , 2007, 12, 209-218.   | 0.5 | 602       |
| 1253 | The yeast penta-EF protein Pef1p is involved in cation-dependent budding and cell polarization. <i>Molecular Microbiology</i> , 2007, 65, 1122-1138.                                 | 1.2 | 8         |
| 1254 | Nucleocytoplasmic Trafficking Is Required for Functioning of the Adaptor Protein Sla1p in Endocytosis. <i>Traffic</i> , 2007, 8, 347-358.  | 1.3 | 16        |
| 1255 | Evidence for a Direct Role of the Doa4 Deubiquitinating Enzyme in Protein Sorting into the MVB Pathway. <i>Traffic</i> , 2007, 8, 566-581.   | 1.3 | 46        |
| 1256 | Snc1p v-εSNARE Transport to the Prospore Membrane During Yeast Sporulation is Dependent on Endosomal Retrieval Pathways. <i>Traffic</i> , 2007, 8, 1231-1245.                        | 1.3 | 19        |
| 1257 | Regulation of Copper-Dependent Endocytosis and Vacuolar Degradation of the Yeast Copper Transporter, Ctr1p, by the Rsp5 Ubiquitin Ligase. <i>Traffic</i> , 2007, 8, 1375-1384.       | 1.3 | 99        |
| 1258 | Trafficking of Siderophore Transporters in <i>Saccharomyces cerevisiae</i> and Intracellular Fate of Ferrioxamine B Conjugates. <i>Traffic</i> , 2007, 8, 1601-1616.                 | 1.3 | 36        |
| 1259 | Dominant-Negative Behavior of Mammalian Vps35 in Yeast Requires a Conserved PRLYL Motif Involved in Retromer Assembly. <i>Traffic</i> , 2007, 8, 1829-1840.                          | 1.3 | 31        |
| 1260 | Cdc37p is involved in osmoadaptation and controls high osmolarity-induced cross-talk via the MAP kinase Kss1p. <i>FEMS Yeast Research</i> , 2007, 7, 796-807.                        | 1.1 | 11        |
| 1261 | Lot6p from <i>Saccharomyces cerevisiae</i> is a FMN-dependent reductase with a potential role in quinone detoxification. <i>FEBS Journal</i> , 2007, 274, 1328-1339.                 | 2.2 | 45        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1262 | A genetic screen identifies mutations in the yeast WAR1 gene, linking transcription factor phosphorylation to weak-acid stress adaptation. <i>FEBS Journal</i> , 2007, 274, 3094-3107.  | 2.2 | 13        |
| 1263 | The phosphatidylethanolamine level of yeast mitochondria is affected by the mitochondrial components Oxa1p and Yme1p. <i>FEBS Journal</i> , 2007, 274, 6180-6190.   | 2.2 | 43        |
| 1264 | Pericentric Chromatin Is an Elastic Component of the Mitotic Spindle. <i>Current Biology</i> , 2007, 17, 741-748.   | 1.8 | 78        |
| 1265 | Yeast Rev1 is cell cycle regulated, phosphorylated in response to DNA damage and its binding to chromosomes is dependent upon MEC1. <i>DNA Repair</i> , 2007, 6, 121-127.   | 1.3 | 53        |
| 1266 | Transcription-associated mutagenesis in yeast is directly proportional to the level of gene expression and influenced by the direction of DNA replication. <i>DNA Repair</i> , 2007, 6, 1285-1296.                            | 1.3 | 100       |
| 1267 | Mrc1 protects uncapped budding yeast telomeres from exonuclease EXO1. <i>DNA Repair</i> , 2007, 6, 1607-1617.   | 1.3 | 20        |
| 1268 | Localization of Gts1p in cortical actin patches of yeast and its possible role in endocytosis. <i>European Journal of Cell Biology</i> , 2007, 86, 275-285.   | 1.6 | 4         |
| 1269 | The arginine methyltransferase Rmt2 is enriched in the nucleus and co-purifies with the nuclear porins Nup49, Nup57 and Nup100. <i>Experimental Cell Research</i> , 2007, 313, 1778-1789.                                     | 1.2 | 7         |
| 1270 | MPK1 gene is required for filamentous growth induced by isoamyl alcohol in <i>Saccharomyces cerevisiae</i> strains from the alcoholic fermentation. <i>Applied Microbiology and Biotechnology</i> , 2007, 75, 111-115.        | 1.7 | 4         |
| 1271 | Mutations in the Atp1p and Atp3p subunits of yeast ATP synthase differentially affect respiration and fermentation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioenergetics and Biomembranes</i> , 2007, 39, 127-144. | 1.0 | 27        |
| 1272 | Characterisation and expression of monosaccharide transporters in lupins, <i>Lupinus polyphyllus</i> and <i>L. albus</i> . <i>Journal of Plant Research</i> , 2007, 120, 697-705.   | 1.2 | 9         |
| 1273 | In vivo functional characterization of the <i>Saccharomyces cerevisiae</i> 60S biogenesis GTPase Nog1. <i>Molecular Genetics and Genomics</i> , 2007, 278, 105-123.   | 1.0 | 31        |
| 1274 | Characterization of MYR1, a dosage suppressor of YPT6 and RIC1 deficient mutants. <i>Current Genetics</i> , 2008, 53, 235-247.  | 0.8 | 5         |
| 1275 | Regulation of <i>Saccharomyces cerevisiae</i> DNA polymerase $\delta$ transcript and protein. <i>Radiation and Environmental Biophysics</i> , 2008, 47, 157-168.  | 0.6 | 20        |
| 1276 | Rapid regulation of protein activity in fission yeast. <i>BMC Cell Biology</i> , 2008, 9, 23.   | 3.0 | 11        |
| 1277 | Membrane topology and post-translational modification of the essential protein Rot1. <i>Yeast</i> , 2008, 25, 93-106.   | 0.8 | 9         |
| 1278 | Control of 5-FOA and 5-FU resistance by <i>Saccharomyces cerevisiae</i> YJL055W. <i>Yeast</i> , 2008, 25, 155-160.  | 0.8 | 16        |
| 1279 | Modules for cloning-free chromatin tagging in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2008, 25, 235-239.   | 0.8 | 63        |



| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1280 | A vector system for efficient and economical switching of C-terminal epitope tags in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2008, 25, 301-311.  | 0.8 | 40        |
| 1281 | Dominant marker vectors for selecting yeast mating products. <i>Yeast</i> , 2008, 25, 595-599.  | 0.8 | 12        |
| 1282 | A small molecule-directed approach to control protein localization and function. <i>Yeast</i> , 2008, 25, 577-594.  | 0.8 | 31        |
| 1283 | Photoactivatable GFP tagging cassettes for protein tracking studies in the budding yeast <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2008, 25, 651-659.  | 0.8 | 25        |
| 1284 | A PCR-based strategy to generate yeast strains expressing endogenous levels of amino-terminal epitope-tagged proteins. <i>Biotechnology Journal</i> , 2008, 3, 524-529.                                     | 1.8 | 15        |
| 1285 | Interdependence of the Ypt/RabGAP Gyp5p and Gyl1p for Recruitment to the Sites of Polarized Growth. <i>Traffic</i> , 2008, 9, 608-622.  | 1.3 | 3         |
| 1286 | Yeast Arf3p Modulates Plasma Membrane PtdIns(4,5)P2 Levels to Facilitate Endocytosis. <i>Traffic</i> , 2008, 9, 559-573.  | 1.3 | 31        |
| 1287 | Irs4p and Tax4p: Two Redundant EH Domain Proteins Involved in Autophagy. <i>Traffic</i> , 2008, 9, 755-769.   | 1.3 | 13        |
| 1288 | Multiple Pathways Regulate Endocytic Coat Disassembly in <i>Saccharomyces cerevisiae</i> for Optimal Downstream Trafficking. <i>Traffic</i> , 2008, 9, 848-859.   | 1.3 | 60        |
| 1289 | A Cryosectioning Procedure for the Ultrastructural Analysis and the Immunogold Labelling of Yeast <i>Saccharomyces cerevisiae</i> . <i>Traffic</i> , 2008, 9, 1060-1072.                                    | 1.3 | 83        |
| 1290 | Components of the Vid30c are needed for the rapamycin-induced degradation of the high-affinity hexose transporter Hxt7p in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2008, 8, 204-216. | 1.1 | 18        |
| 1291 | Hyperpolarized growth of <i>Saccharomyces cerevisiae</i> cak1Δ and cla4 mutants weakens cell walls and renders cells dependent on chitin synthase 3. <i>FEMS Yeast Research</i> , 2008, 8, 362-373.         | 1.1 | 9         |
| 1292 | Site specificity of yeast histone acetyltransferase B complex <i>in vivo</i> . <i>FEBS Journal</i> , 2008, 275, 2122-2136.  | 2.2 | 26        |
| 1293 | Phenyl hydroquinone, an Ames test-negative carcinogen, induces Hog1-dependent stress response signaling. <i>FEBS Journal</i> , 2008, 275, 5733-5744.  | 2.2 | 10        |
| 1294 | Structure of the archaeal Kae1/Bud32 fusion protein MJ1130: a model for the eukaryotic EKC/KEOPS subcomplex. <i>EMBO Journal</i> , 2008, 27, 2340-2351.   | 3.5 | 62        |
| 1295 | Distinct roles for yeast Stn1 in telomere capping and telomerase inhibition. <i>EMBO Journal</i> , 2008, 27, 2328-2339.   | 3.5 | 94        |
| 1296 | Checkpoint-dependent phosphorylation of Exo1 modulates the DNA damage response. <i>EMBO Journal</i> , 2008, 27, 2400-2410.  | 3.5 | 151       |
| 1297 | Curing of the [URE3] prion by Btn2p, a Batten disease-related protein. <i>EMBO Journal</i> , 2008, 27, 2725-2735.   | 3.5 | 94        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 1298 | Histone methyltransferase Dot1 and Rad9 inhibit single-stranded DNA accumulation at DSBs and uncapped telomeres. <i>EMBO Journal</i> , 2008, 27, 1502-12.  | 3.5  | 159       |
| 1299 | Yeast Rap1 contributes to genomic integrity by activating DNA damage repair genes. <i>EMBO Journal</i> , 2008, 27, 1575-1584.  | 3.5  | 30        |
| 1300 | Quantitative analysis of snoRNA association with pre-ribosomes and release of snR30 by Rok1 helicase. <i>EMBO Reports</i> , 2008, 9, 1230-1236.  | 2.0  | 72        |
| 1301 | A mechanism for asymmetric segregation of age during yeast budding. <i>Nature</i> , 2008, 454, 728-734.  | 13.7 | 298       |
| 1302 | Sae2, Exo1 and Sgs1 collaborate in DNA double-strand break processing. <i>Nature</i> , 2008, 455, 770-774.   | 13.7 | 876       |
| 1303 | The Ino80 chromatin-remodeling enzyme regulates replisome function and stability. <i>Nature Structural and Molecular Biology</i> , 2008, 15, 338-345.  | 3.6  | 143       |
| 1304 | Cascading transcriptional effects of a naturally occurring frameshift mutation in <i>Saccharomyces cerevisiae</i> . <i>Molecular Ecology</i> , 2008, 17, 2985-2997.  | 2.0  | 37        |
| 1305 | Protein N-glycosylation determines functionality of the <i>Saccharomyces cerevisiae</i> cell wall integrity sensor Mid2p. <i>Molecular Microbiology</i> , 2008, 68, 1438-1449.   | 1.2  | 39        |
| 1306 | Yeast $\beta$ -tubulin suppressor Ats1/Kti13 relates to the Elongator complex and interacts with Elongator partner protein Kti11. <i>Molecular Microbiology</i> , 2008, 69, 175-187.                                       | 1.2  | 33        |
| 1307 | A versatile partner of eukaryotic protein complexes that is involved in multiple biological processes: Kti11/Dph3. <i>Molecular Microbiology</i> , 2008, 69, 1221-1233.  | 1.2  | 48        |
| 1308 | tRNA and protein methylase complexes mediate zymocin toxicity in yeast. <i>Molecular Microbiology</i> , 2008, 69, 1266-1277.   | 1.2  | 56        |
| 1309 | The periplasmic regulator ExoR inhibits ExoS/ChvI two-component signalling in <i>Sinorhizobium meliloti</i> . <i>Molecular Microbiology</i> , 2008, 69, 1290-1303.   | 1.2  | 59        |
| 1310 | Genetic interaction network of the <i>Saccharomyces cerevisiae</i> type 1 phosphatase Glc7. <i>BMC Genomics</i> , 2008, 9, 336.  | 1.2  | 14        |
| 1311 | Correlation between the secondary structure of pre-mRNA introns and the efficiency of splicing in <i>Saccharomyces cerevisiae</i> . <i>BMC Genomics</i> , 2008, 9, 355.  | 1.2  | 43        |
| 1312 | Oxidative stress and chronological aging in glycogen-phosphorylase-deleted yeast. <i>Free Radical Biology and Medicine</i> , 2008, 45, 1446-1456.  | 1.3  | 46        |
| 1313 | The F658G substitution in <i>Saccharomyces cerevisiae</i> cohesin Irr1/Scs3 is semi-dominant in the diploid and disturbs mitosis, meiosis and the cell cycle. <i>European Journal of Cell Biology</i> , 2008, 87, 831-844. | 1.6  | 4         |
| 1314 | Intrinsic 5'-deoxyribose-5-phosphate lyase activity in <i>Saccharomyces cerevisiae</i> Trf4 protein with a possible role in base excision DNA repair. <i>DNA Repair</i> , 2008, 7, 187-198.                                | 1.3  | 21        |
| 1315 | Identification of pathways controlling DNA damage induced mutation in <i>Saccharomyces cerevisiae</i> . <i>DNA Repair</i> , 2008, 7, 801-810.  | 1.3  | 25        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 1316 | Structure of the Oligosaccharyl Transferase Complex at 12 Å.. Resolution. <i>Structure</i> , 2008, 16, 432-440.  | 1.6  | 45        |
| 1317 | Chapter 14 Methods to Determine mRNA Half-Life in <i>Saccharomyces cerevisiae</i> . <i>Methods in Enzymology</i> , 2008, 448, 267-284.   | 0.4  | 28        |
| 1318 | Genome adaptation to chemical stress: clues from comparative transcriptomics in <i>Saccharomyces cerevisiae</i> and <i>Candida glabrata</i> . <i>Genome Biology</i> , 2008, 9, R164. | 13.9 | 48        |
| 1319 | Antioxidant Small Molecules Confer Variable Protection against Oxidative Damage in Yeast Mutants. <i>Journal of Agricultural and Food Chemistry</i> , 2008, 56, 11740-11751.         | 2.4  | 32        |
| 1320 | Efficient termination of transcription by RNA polymerase I requires the 5' exonuclease Rat1 in yeast. <i>Genes and Development</i> , 2008, 22, 1069-1081.                            | 2.7  | 102       |
| 1321 | Cell cycle-dependent kinetochore localization of condensin complex in <i>Saccharomyces cerevisiae</i> . <i>Journal of Structural Biology</i> , 2008, 162, 248-259.                   | 1.3  | 25        |
| 1322 | The Transmembrane Segment of Tom20 Is Recognized by Mim1 for Docking to the Mitochondrial TOM Complex. <i>Journal of Molecular Biology</i> , 2008, 376, 694-704.                     | 2.0  | 88        |
| 1323 | Role of Î³-Subunit N- and C-Termini in Assembly of the Mitochondrial ATP Synthase in Yeast. <i>Journal of Molecular Biology</i> , 2008, 377, 1314-1323.                              | 2.0  | 6         |
| 1324 | Nonsense-Mediated mRNA Decay in Yeast Does Not Require PAB1 or a Poly(A) Tail. <i>Molecular Cell</i> , 2008, 29, 134-140.  | 4.5  | 59        |
| 1325 | Dcn1 Functions as a Scaffold-Type E3 Ligase for Cullin Neddylation. <i>Molecular Cell</i> , 2008, 29, 23-35.   | 4.5  | 163       |
| 1326 | Mus81/Mms4 Endonuclease and Sgs1 Helicase Collaborate to Ensure Proper Recombination Intermediate Metabolism during Meiosis. <i>Molecular Cell</i> , 2008, 31, 313-323.              | 4.5  | 134       |
| 1327 | Widespread Impact of Nonsense-Mediated mRNA Decay on the Yeast Intronome. <i>Molecular Cell</i> , 2008, 31, 360-370.   | 4.5  | 105       |
| 1328 | Futile Cycle of Transcription Initiation and Termination Modulates the Response to Nucleotide Shortage in <i>S. cerevisiae</i> . <i>Molecular Cell</i> , 2008, 31, 671-682.          | 4.5  | 93        |
| 1329 | Dissecting sensor functions in cell wall integrity signaling in <i>Kluyveromyces lactis</i> . <i>Fungal Genetics and Biology</i> , 2008, 45, 422-435.                                | 0.9  | 48        |
| 1330 | Detection and localisation of protein-protein interactions in <i>Saccharomyces cerevisiae</i> using a split-GFP method. <i>Fungal Genetics and Biology</i> , 2008, 45, 597-604.      | 0.9  | 18        |
| 1331 | Identification of Tup1 and Cyc8 mutations defective in the responses to osmotic stress. <i>Biochemical and Biophysical Research Communications</i> , 2008, 368, 50-55.               | 1.0  | 14        |
| 1332 | A novel role for thioredoxin reductase in the iron metabolism of <i>S. cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2008, 371, 63-68.               | 1.0  | 9         |
| 1333 | In vitro reconstitution of Î³-secretase activity using yeast microsomes. <i>Biochemical and Biophysical Research Communications</i> , 2008, 377, 141-145.                            | 1.0  | 37        |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 1334 | The role of the <i>Saccharomyces cerevisiae</i> Cdc7-Dbf4 complex in the replication checkpoint. <i>Gene</i> , 2008, 414, 32-40.  | 1.0  | 35        |
| 1335 | Meiosis I Is Established through Division-Specific Translational Control of a Cyclin. <i>Cell</i> , 2008, 133, 280-291.   | 13.5 | 202       |
| 1336 | The Structural Basis for Activation of the Rab Ypt1p by the TRAPP Membrane-Tethering Complexes. <i>Cell</i> , 2008, 133, 1202-1213.   | 13.5 | 166       |
| 1337 | A Network of Nuclear Envelope Membrane Proteins Linking Centromeres to Microtubules. <i>Cell</i> , 2008, 134, 427-438.  | 13.5 | 165       |
| 1338 | The GET Complex Mediates Insertion of Tail-Anchored Proteins into the ER Membrane. <i>Cell</i> , 2008, 134, 634-645.  | 13.5 | 460       |
| 1339 | Regulation of Ceramide Biosynthesis by TOR Complex 2. <i>Cell Metabolism</i> , 2008, 7, 148-158.  | 7.2  | 174       |
| 1340 | Golgi Localization of Glycosyltransferases Requires a Vps74p Oligomer. <i>Developmental Cell</i> , 2008, 14, 523-534.   | 3.1  | 164       |
| 1341 | Global Analysis of the Meiotic Crossover Landscape. <i>Developmental Cell</i> , 2008, 15, 401-415.  | 3.1  | 197       |
| 1342 | P bodies promote stress granule assembly in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 2008, 183, 441-455.  | 2.3  | 455       |
| 1343 | Chapter 9 Counting Proteins in Living Cells by Quantitative Fluorescence Microscopy with Internal Standards. <i>Methods in Cell Biology</i> , 2008, 89, 253-273.  | 0.5  | 59        |
| 1344 | Tor Pathway Control of the Nitrogen-responsive DAL5 Gene Bifurcates at the Level of Gln3 and Gat1 Regulation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2008, 283, 8919-8929.            | 1.6  | 52        |
| 1345 | APC/C-Cdh1-mediated degradation of the Polo kinase Cdc5 promotes the return of Cdc14 into the nucleolus. <i>Genes and Development</i> , 2008, 22, 79-90.  | 2.7  | 80        |
| 1346 | A cryptic unstable transcript mediates transcriptional silencing of the Ty1 retrotransposon in <i>S. cerevisiae</i> . <i>Genes and Development</i> , 2008, 22, 615-626.   | 2.7  | 173       |
| 1347 | 60S ribosomal subunit assembly dynamics defined by semi-quantitative mass spectrometry of purified complexes. <i>Nucleic Acids Research</i> , 2008, 36, 4988-4999.  | 6.5  | 47        |
| 1348 | A functional proteomics approach links the ubiquitin-related modifier Urm1 to a tRNA modification pathway. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 18255-18260. | 3.3  | 97        |
| 1349 | The SBF- and MBF-associated Protein Msa1 Is Required for Proper Timing of G1-specific Transcription in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2008, 283, 6040-6049.                     | 1.6  | 27        |
| 1350 | The Vacuolar Import and Degradation Pathway Merges with the Endocytic Pathway to Deliver Fructose-1,6-bisphosphatase to the Vacuole for Degradation. <i>Journal of Biological Chemistry</i> , 2008, 283, 26116-26127.       | 1.6  | 40        |
| 1351 | Mitophagy in Yeast Occurs through a Selective Mechanism. <i>Journal of Biological Chemistry</i> , 2008, 283, 32386-32393.   | 1.6  | 296       |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1352 | Cell Wall Integrity MAPK Pathway Is Essential for Lipid Homeostasis. <i>Journal of Biological Chemistry</i> , 2008, 283, 34204-34217.   | 1.6 | 38        |
| 1353 | The AAA ATPase Rix7 powers progression of ribosome biogenesis by stripping Nsa1 from pre-60S particles. <i>Journal of Cell Biology</i> , 2008, 181, 935-944.  | 2.3 | 78        |
| 1354 | Binding interactions control SNARE specificity in vivo. <i>Journal of Cell Biology</i> , 2008, 183, 1089-1100.  | 2.3 | 15        |
| 1355 | Novel Ist1-Did2 Complex Functions at a Late Step in Multivesicular Body Sorting. <i>Molecular Biology of the Cell</i> , 2008, 19, 475-484.  | 0.9 | 118       |
| 1356 | Self-Interaction Is Critical for Atg9 Transport and Function at the Phagophore Assembly Site during Autophagy. <i>Molecular Biology of the Cell</i> , 2008, 19, 5506-5516.  | 0.9 | 101       |
| 1357 | Mitogen-Activated Protein Kinase Hog1 Mediates Adaptation to G <sub>1</sub> Checkpoint Arrest during Arsenite and Hyperosmotic Stress. <i>Eukaryotic Cell</i> , 2008, 7, 1309-1317.   | 3.4 | 27        |
| 1358 | Mitochondrial nucleoids undergo remodeling in response to metabolic cues. <i>Journal of Cell Science</i> , 2008, 121, 1861-1868.  | 1.2 | 96        |
| 1359 | Stb1 Collaborates with Other Regulators To Modulate the G <sub>1</sub> -Specific Transcriptional Circuit. <i>Molecular and Cellular Biology</i> , 2008, 28, 6919-6928.  | 1.1 | 37        |
| 1360 | Coa2 Is an Assembly Factor for Yeast Cytochrome <i>c</i> Oxidase Biogenesis That Facilitates the Maturation of Cox1. <i>Molecular and Cellular Biology</i> , 2008, 28, 4927-4939.   | 1.1 | 55        |
| 1361 | Nucleocytoplasmic Shuttling of the Golgi Phosphatidylinositol 4-Kinase Pik1 Is Regulated by 14-3-3 Proteins and Coordinates Golgi Function with Cell Growth. <i>Molecular Biology of the Cell</i> , 2008, 19, 1046-1061.    | 0.9 | 64        |
| 1362 | Catalytic-Site Mutations in the MYST Family Histone Acetyltransferase Esa1. <i>Genetics</i> , 2008, 178, 1209-1220.   | 1.2 | 29        |
| 1363 | Interspecies variation reveals a conserved repressor of $\hat{\pm}$ -specific genes in <i>Saccharomyces</i> yeasts. <i>Genes and Development</i> , 2008, 22, 1704-1716.   | 2.7 | 26        |
| 1364 | A Proteomics Analysis of Yeast Mot1p Protein-Protein Associations. <i>Molecular and Cellular Proteomics</i> , 2008, 7, 2090-2106.   | 2.5 | 16        |
| 1365 | Acetylation of Rsc4p by Gcn5p Is Essential in the Absence of Histone H3 Acetylation. <i>Molecular and Cellular Biology</i> , 2008, 28, 6967-6972.   | 1.1 | 21        |
| 1366 | The <i>Saccharomyces cerevisiae</i> Actin Cytoskeletal Component Bsp1p Has an Auxiliary Role in Actomyosin Ring Function and in the Maintenance of Bud-Neck Structure. <i>Genetics</i> , 2008, 178, 1903-1914.              | 1.2 | 7         |
| 1367 | Identification of Clb2 Residues Required for Swe1 Regulation of Clb2-Cdc28 in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2008, 179, 863-874.   | 1.2 | 17        |
| 1368 | The Molecular Basis for Relative Physiological Functionality of the ADP/ATP Carrier Isoforms in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2008, 179, 1285-1299.   | 1.2 | 18        |
| 1369 | Synthetic Lethal Screens Identify Gene Silencing Processes in Yeast and Implicate the Acetylated Amino Terminus of Sir3 in Recognition of the Nucleosome Core. <i>Molecular and Cellular Biology</i> , 2008, 28, 3861-3872. | 1.1 | 53        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1370 | A Yeast Sir2 Mutant Temperature Sensitive for Silencing. <i>Genetics</i> , 2008, 180, 1955-1962.   | 1.2 | 11        |
| 1371 | Involvement of <i>Saccharomyces cerevisiae</i> Avo3p/Tsc11p in Maintaining TOR Complex 2 Integrity and Coupling to Downstream Signaling. <i>Eukaryotic Cell</i> , 2008, 7, 1328-1343.                        | 3.4 | 15        |
| 1372 | Functional Dissection of the NuA4 Histone Acetyltransferase Reveals Its Role as a Genetic Hub and that Eaf1 Is Essential for Complex Integrity. <i>Molecular and Cellular Biology</i> , 2008, 28, 2244-2256. | 1.1 | 101       |
| 1373 | Nap1 Links Transcription Elongation, Chromatin Assembly, and Messenger RNP Complex Biogenesis. <i>Molecular and Cellular Biology</i> , 2008, 28, 2113-2124.  | 1.1 | 29        |
| 1374 | Dissecting the localization and function of Atg18, Atg21 and Ygr223c. <i>Autophagy</i> , 2008, 4, 896-910.   | 4.3 | 90        |
| 1375 | Tel2 mediates activation and localization of ATM/Tel1 kinase to a double-strand break. <i>Genes and Development</i> , 2008, 22, 854-859.   | 2.7 | 44        |
| 1376 | Unconventional Genomic Architecture in the Budding Yeast <i>Saccharomyces cerevisiae</i> Masks the Nested Antisense Gene <i>NAG1</i> . <i>Eukaryotic Cell</i> , 2008, 7, 1289-1298.                          | 3.4 | 17        |
| 1377 | Localization of autophagy-related proteins in yeast using a versatile plasmid-based resource of fluorescent protein fusions. <i>Autophagy</i> , 2008, 4, 792-800.  | 4.3 | 11        |
| 1378 | Opposing Activities of the Snx3-Retromer Complex and ESCRT Proteins Mediate Regulated Cargo Sorting at a Common Endosome. <i>Molecular Biology of the Cell</i> , 2008, 19, 4694-4706.                        | 0.9 | 55        |
| 1379 | Mat Formation in <i>Saccharomyces cerevisiae</i> Requires Nutrient and pH Gradients. <i>Eukaryotic Cell</i> , 2008, 7, 122-130.  | 3.4 | 48        |
| 1380 | Superoxide anions regulate TORC1 and its ability to bind Fpr1:rapamycin complex. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 15166-15171.            | 3.3 | 48        |
| 1381 | Role of Nucleotide Binding in Septin-Septin Interactions and Septin Localization in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2008, 28, 5120-5137.                           | 1.1 | 47        |
| 1382 | Characterization of Kl <i>GRR1</i> and <i>SMS1</i> Genes, Two New Elements of the Glucose Signaling Pathway of <i>Kluyveromyces lactis</i> . <i>Eukaryotic Cell</i> , 2008, 7, 1299-1308.                    | 3.4 | 17        |
| 1383 | Dre2, a Conserved Eukaryotic Fe/S Cluster Protein, Functions in Cytosolic Fe/S Protein Biogenesis. <i>Molecular and Cellular Biology</i> , 2008, 28, 5569-5582.  | 1.1 | 145       |
| 1384 | Structure-system correlation identifies a gene regulatory Mediator submodule. <i>Genes and Development</i> , 2008, 22, 872-877.  | 2.7 | 31        |
| 1385 | Autophosphorylation-Induced Degradation of the Pho85 Cyclin Pcl5 Is Essential for Response to Amino Acid Limitation. <i>Molecular and Cellular Biology</i> , 2008, 28, 6858-6869.                            | 1.1 | 14        |
| 1386 | Rtr1 Is the <i>Saccharomyces cerevisiae</i> Homolog of a Novel Family of RNA Polymerase II-Binding Proteins. <i>Eukaryotic Cell</i> , 2008, 7, 938-948.  | 3.4 | 45        |
| 1387 | Nuclear Export Receptor Xpo1/Crm1 Is Physically and Functionally Linked to the Spindle Pole Body in Budding Yeast. <i>Molecular and Cellular Biology</i> , 2008, 28, 5348-5358.                              | 1.1 | 15        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1388 | Oxygen-regulated isoforms of cytochrome <i>c</i> oxidase have differential effects on its nitric oxide production and on hypoxic signaling. Proceedings of the National Academy of Sciences of the United States of America, 2008, 105, 8203-8208. | 3.3 | 113       |
| 1389 | Yeast UCS proteins promote actomyosin interactions and limit myosin turnover in cells. Proceedings of the National Academy of Sciences of the United States of America, 2008, 105, 8014-8019.  | 3.3 | 33        |
| 1390 | Yeast Pgc1p (YPL206c) Controls the Amount of Phosphatidylglycerol via a Phospholipase C-type Degradation Mechanism. Journal of Biological Chemistry, 2008, 283, 17107-17115.   | 1.6 | 46        |
| 1391 | Regulation of the Yeast Ace2 Transcription Factor during the Cell Cycle*. Journal of Biological Chemistry, 2008, 283, 11135-11145.   | 1.6 | 58        |
| 1392 | Functional Surfaces on the p35/ARPC2 Subunit of Arp2/3 Complex Required for Cell Growth, Actin Nucleation, and Endocytosis. Journal of Biological Chemistry, 2008, 283, 16950-16959.   | 1.6 | 30        |
| 1393 | Elongating RNA Polymerase II Is Disassembled through Specific Degradation of Its Largest but Not Other Subunits in Response to DNA Damage in Vivo. Journal of Biological Chemistry, 2008, 283, 6897-6905.  | 1.6 | 31        |
| 1394 | The <i>Saccharomyces cerevisiae</i> 60 S Ribosome Biogenesis Factor Tif6p Is Regulated by Hrr25p-mediated Phosphorylation. Journal of Biological Chemistry, 2008, 283, 9681-9691.  | 1.6 | 33        |
| 1395 | Sus1 is recruited to coding regions and functions during transcription elongation in association with SAGA and TREX2. Genes and Development, 2008, 22, 2811-2822.  | 2.7 | 90        |
| 1396 | Large-Scale Analysis of Yeast Filamentous Growth by Systematic Gene Disruption and Overexpression. Molecular Biology of the Cell, 2008, 19, 284-296.   | 0.9 | 118       |
| 1397 | Role of Dot1 in the Response to Alkylating DNA Damage in <i>Saccharomyces cerevisiae</i> : Regulation of DNA Damage Tolerance by the Error-Prone Polymerases Pol $\eta$ /Rev1. Genetics, 2008, 179, 1197-1210.                                     | 1.2 | 47        |
| 1398 | Visual Analysis of the Yeast 5S rRNA Gene Transcriptome: Regulation and Role of La Protein. Molecular and Cellular Biology, 2008, 28, 4576-4587.   | 1.1 | 42        |
| 1399 | Protein Kinases Fpk1p and Fpk2p are Novel Regulators of Phospholipid Asymmetry. Molecular Biology of the Cell, 2008, 19, 1783-1797.  | 0.9 | 82        |
| 1400 | In vivo reconstitution of autophagy in <i>Saccharomyces cerevisiae</i> . Journal of Cell Biology, 2008, 182, 703-713.  | 2.3 | 61        |
| 1401 | The Sur7 Protein Regulates Plasma Membrane Organization and Prevents Intracellular Cell Wall Growth in <i>Candida albicans</i> . Molecular Biology of the Cell, 2008, 19, 5214-5225.   | 0.9 | 77        |
| 1402 | Counting Kinetochore Protein Numbers in Budding Yeast Using Genetically Encoded Fluorescent Proteins. Methods in Cell Biology, 2008, 85, 127-151.  | 0.5 | 54        |
| 1403 | Dissection of the Carboxyl-Terminal Domain of the Proteasomal Subunit Rpn11 in Maintenance of Mitochondrial Structure and Function. Molecular Biology of the Cell, 2008, 19, 1022-1031.  | 0.9 | 36        |
| 1404 | Cleavage of the signaling mucin Msb2 by the aspartyl protease Yps1 is required for MAPK activation in yeast. Journal of Cell Biology, 2008, 181, 1073-1081.  | 2.3 | 107       |
| 1405 | Assembly of the PtdIns 4-kinase Stt4 complex at the plasma membrane requires Ypp1 and Efr3. Journal of Cell Biology, 2008, 183, 1061-1074.   | 2.3 | 150       |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1406 | Hos2p/Set3p Deacetylase Complex Signals Secretory Stress through the Mpk1p Cell Integrity Pathway. <i>Eukaryotic Cell</i> , 2008, 7, 1191-1199.  | 3.4 | 28        |
| 1407 | A Block in Endoplasmic Reticulum-to-Golgi Trafficking Inhibits Phospholipid Synthesis and Induces Neutral Lipid Accumulation. <i>Journal of Biological Chemistry</i> , 2008, 283, 25735-25751.   | 1.6 | 63        |
| 1408 | The Rsp5 E3 Ligase Mediates Turnover of Low Affinity Phosphate Transporters in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2008, 283, 5327-5334.  | 1.6 | 16        |
| 1409 | Interactions between the Yeast SM22 Homologue Scp1 and Actin Demonstrate the Importance of Actin Bundling in Endocytosis. <i>Journal of Biological Chemistry</i> , 2008, 283, 15037-15046.   | 1.6 | 41        |
| 1410 | A genome-wide screen identifies genes required for formation of the wobble nucleoside 5-methoxycarbonylmethyl-2-thiouridine in <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 2008, 14, 2183-2194.   | 1.6 | 170       |
| 1411 | <i>Saccharomyces cerevisiae</i> HMO1 interacts with TFIID and participates in start site selection by RNA polymerase II. <i>Nucleic Acids Research</i> , 2008, 36, 1343-1357.  | 6.5 | 30        |
| 1412 | Nuclear translocation of Gln3 in response to nutrient signals requires Golgi-to-endosome trafficking in <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 7194-7199. | 3.3 | 49        |
| 1413 | Requirement for the Budding Yeast Polo Kinase Cdc5 in Proper Microtubule Growth and Dynamics. <i>Eukaryotic Cell</i> , 2008, 7, 444-453.   | 3.4 | 35        |
| 1414 | The function of two closely related Rho proteins is determined by an atypical switch I region. <i>Journal of Cell Science</i> , 2008, 121, 1065-1075.  | 1.2 | 21        |
| 1415 | The Cytoplasmic Region of $\alpha$ -1,6-Mannosyltransferase Mnn9p Is Crucial for Retrograde Transport from the Golgi Apparatus to the Endoplasmic Reticulum in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2008, 7, 310-318.              | 3.4 | 25        |
| 1416 | <i>Saccharomyces cerevisiae</i> Afr1 Protein Is a Protein Phosphatase 1/Glc7-Targeting Subunit That Regulates the Septin Cytoskeleton during Mating. <i>Eukaryotic Cell</i> , 2008, 7, 1246-1255.  | 3.4 | 16        |
| 1417 | Interaction between Epsin/Yap180 Adaptors and the Scaffolds Ede1/Pan1 Is Required for Endocytosis. <i>Molecular Biology of the Cell</i> , 2008, 19, 2936-2948.   | 0.9 | 54        |
| 1418 | The malate-aspartate NADH shuttle components are novel metabolic longevity regulators required for calorie restriction-mediated life span extension in yeast. <i>Genes and Development</i> , 2008, 22, 931-944.  | 2.7 | 130       |
| 1419 | An Increase in Mitochondrial DNA Promotes Nuclear DNA Replication in Yeast. <i>PLoS Genetics</i> , 2008, 4, e1000047.  | 1.5 | 31        |
| 1420 | Novel Low Abundance and Transient RNAs in Yeast Revealed by Tiling Microarrays and Ultra High-Throughput Sequencing Are Not Conserved Across Closely Related Yeast Species. <i>PLoS Genetics</i> , 2008, 4, e1000299.                                    | 1.5 | 31        |
| 1421 | The NDR/LATS Family Kinase Cbk1 Directly Controls Transcriptional Asymmetry. <i>PLoS Biology</i> , 2008, 6, e203.  | 2.6 | 92        |
| 1422 | Role of PUG1 in Inducible Porphyrin and Heme Transport in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2008, 7, 859-871.   | 3.4 | 53        |
| 1423 | Levels of SCS7/FA2H-Mediated Fatty Acid 2-Hydroxylation Determine the Sensitivity of Cells to Antitumor PM02734. <i>Cancer Research</i> , 2008, 68, 9779-9787.   | 0.4 | 57        |



| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1424 | Nsf1/Ypl230w participates in transcriptional activation during non-fermentative growth and in response to salt stress in <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> , 2008, 154, 2482-2491.                          | 0.7 | 22        |
| 1425 | Dynamics and Design Principles of a Basic Regulatory Architecture Controlling Metabolic Pathways. <i>PLoS Biology</i> , 2008, 6, e146.   | 2.6 | 43        |
| 1426 | DNA Damage Activates the SAC in an ATM/ATR-Dependent Manner, Independently of the Kinetochores. <i>PLoS Genetics</i> , 2008, 4, e1000015.  | 1.5 | 68        |
| 1427 | The Coordination of Centromere Replication, Spindle Formation, and Kinetochores-Microtubule Interaction in Budding Yeast. <i>PLoS Genetics</i> , 2008, 4, e1000262.  | 1.5 | 27        |
| 1428 | A Novel Tel1/ATM N-Terminal Motif, TAN, Is Essential for Telomere Length Maintenance and a DNA Damage Response. <i>Molecular and Cellular Biology</i> , 2008, 28, 5736-5746.   | 1.1 | 21        |
| 1429 | A Chemical Genomics Study Identifies Snf1 as a Repressor of GCN4 Translation. <i>Journal of Biological Chemistry</i> , 2008, 283, 35889-35898.   | 1.6 | 47        |
| 1430 | Phosphorylation of the Budding Yeast 9-1-1 Complex Is Required for Dpb11 Function in the Full Activation of the UV-Induced DNA Damage Checkpoint. <i>Molecular and Cellular Biology</i> , 2008, 28, 4782-4793.                                   | 1.1 | 109       |
| 1431 | Cdc7-Dbf4 Regulates <i>NDT80</i> Transcription as Well as Reductional Segregation during Budding Yeast Meiosis. <i>Molecular Biology of the Cell</i> , 2008, 19, 4956-4967.  | 0.9 | 45        |
| 1432 | Stress-Dependent Dynamics of Global Chromatin Remodeling in Yeast: Dual Role for SWI/SNF in the Heat Shock Stress Response. <i>Molecular and Cellular Biology</i> , 2008, 28, 2221-2234.   | 1.1 | 131       |
| 1433 | HOPS Proofreads the <i>trans</i> -SNARE Complex for Yeast Vacuole Fusion. <i>Molecular Biology of the Cell</i> , 2008, 19, 2500-2508.  | 0.9 | 109       |
| 1434 | The Checkpoint Kinase Hsl1p Is Activated by Elm1p-dependent Phosphorylation. <i>Molecular Biology of the Cell</i> , 2008, 19, 4675-4686.   | 0.9 | 35        |
| 1435 | The dual control of TFIIIB recruitment by NC2 is gene specific. <i>Nucleic Acids Research</i> , 2008, 36, 539-549.   | 6.5 | 81        |
| 1436 | The Atg1 Kinase Complex Is Involved in the Regulation of Protein Recruitment to Initiate Sequestering Vesicle Formation for Nonspecific Autophagy in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2008, 19, 668-681. | 0.9 | 233       |
| 1437 | Alg13p, the Catalytic Subunit of the Endoplasmic Reticulum UDP-GlcNAc Glycosyltransferase, Is a Target for Proteasomal Degradation. <i>Molecular Biology of the Cell</i> , 2008, 19, 2169-2178.  | 0.9 | 18        |
| 1438 | Pseudosubstrate Inhibition of the Anaphase-Promoting Complex by Acm1: Regulation by Proteolysis and Cdc28 Phosphorylation. <i>Molecular and Cellular Biology</i> , 2008, 28, 4653-4664.  | 1.1 | 42        |
| 1439 | ESCRT-III Protein Snf7 Mediates High-Level Expression of the <i>SUC2</i> Gene via the Rim101 Pathway. <i>Eukaryotic Cell</i> , 2008, 7, 1888-1894.   | 3.4 | 16        |
| 1440 | Translation of <i>ASH1</i> mRNA is repressed by Puf6p-Fun12p/elf5B interaction and released by CK2 phosphorylation. <i>Genes and Development</i> , 2008, 22, 1037-1050.  | 2.7 | 92        |
| 1441 | Different Domains of the UBL-UBA Ubiquitin Receptor, Ddi1/Vsm1, Are Involved in Its Multiple Cellular Roles. <i>Molecular Biology of the Cell</i> , 2008, 19, 3625-3637.   | 0.9 | 57        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1442 | Afi1p Functions as an Arf3p Polarization-specific Docking Factor for Development of Polarity. <i>Journal of Biological Chemistry</i> , 2008, 283, 16915-16927.   | 1.6 | 12        |
| 1443 | Assembly of a Fab1 Phosphoinositide Kinase Signaling Complex Requires the Fig4 Phosphoinositide Phosphatase. <i>Molecular Biology of the Cell</i> , 2008, 19, 4273-4286.   | 0.9 | 120       |
| 1444 | 14-3-3 Interaction with Histone H3 Involves a Dual Modification Pattern of Phosphoacetylation. <i>Molecular and Cellular Biology</i> , 2008, 28, 2840-2849.  | 1.1 | 77        |
| 1445 | Chapter Four Fluorescence Microscopy-Based Assays for Monitoring Yeast Atg Protein Trafficking. <i>Methods in Enzymology</i> , 2008, 451, 43-56.   | 0.4 | 14        |
| 1446 | Regulation of Mih1/Cdc25 by protein phosphatase 2A and casein kinase 1. <i>Journal of Cell Biology</i> , 2008, 180, 931-945.   | 2.3 | 57        |
| 1447 | Eaf1 Is the Platform for NuA4 Molecular Assembly That Evolutionarily Links Chromatin Acetylation to ATP-Dependent Exchange of Histone H2A Variants. <i>Molecular and Cellular Biology</i> , 2008, 28, 2257-2270.   | 1.1 | 155       |
| 1448 | Interactions among Ytm1, Erb1, and Nop7 Required for Assembly of the Nop7-Subcomplex in Yeast Preribosomes. <i>Molecular Biology of the Cell</i> , 2008, 19, 2844-2856.  | 0.9 | 52        |
| 1449 | Myo2p, a class V myosin in budding yeast, associates with a large ribonucleic acid-protein complex that contains mRNAs and subunits of the RNA-processing body. <i>Rna</i> , 2008, 14, 491-502.  | 1.6 | 29        |
| 1450 | Yeast Golgi-localized, $\beta$ -Ear-containing, ADP-Ribosylation Factor-binding Proteins Are but Adaptor Protein-1 Is Not Required for Cell-free Transport of Membrane Proteins from the Trans-Golgi Network to the Prevacuolar Compartment. <i>Molecular Biology of the Cell</i> , 2008, 19, 4826-4836. | 0.9 | 22        |
| 1451 | Quantitative analysis of autophagy-related protein stoichiometry by fluorescence microscopy. <i>Journal of Cell Biology</i> , 2008, 182, 129-140.  | 2.3 | 108       |
| 1452 | Protein Phosphatase Type 1 Directs Chitin Synthesis at the Bud Neck in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2008, 19, 3040-3051.   | 0.9 | 30        |
| 1453 | Clustering of yeast tRNA genes is mediated by specific association of condensin with tRNA gene transcription complexes. <i>Genes and Development</i> , 2008, 22, 2204-2214.  | 2.7 | 200       |
| 1454 | The Yeast GID Complex, a Novel Ubiquitin Ligase (E3) Involved in the Regulation of Carbohydrate Metabolism. <i>Molecular Biology of the Cell</i> , 2008, 19, 3323-3333.  | 0.9 | 132       |
| 1455 | Analysis of the Yeast Kinome Reveals a Network of Regulated Protein Localization during Filamentous Growth. <i>Molecular Biology of the Cell</i> , 2008, 19, 2708-2717.  | 0.9 | 50        |
| 1456 | The Rho GDI Rdi1 Regulates Rho GTPases by Distinct Mechanisms. <i>Molecular Biology of the Cell</i> , 2008, 19, 2885-2896.   | 0.9 | 68        |
| 1457 | Recruitment and Dissociation of Nonhomologous End Joining Proteins at a DNA Double-Strand Break in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2008, 178, 1237-1249.   | 1.2 | 115       |
| 1458 | The Temporal Program of Chromosome Replication: Genomewide Replication in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2008, 180, 1833-1847.  | 1.2 | 96        |
| 1459 | Autoregulation of Npl3, a Yeast SR Protein, Requires a Novel Downstream Region and Serine Phosphorylation. <i>Molecular and Cellular Biology</i> , 2008, 28, 3873-3881.  | 1.1 | 19        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1460 | The Clathrin Adaptor Gga2p Is a Phosphatidylinositol 4-phosphate Effector at the Golgi Exit. <i>Molecular Biology of the Cell</i> , 2008, 19, 1991-2002.  | 0.9 | 66        |
| 1461 | Roles for Ctk1 and Spt6 in Regulating the Different Methylation States of Histone H3 Lysine 36. <i>Molecular and Cellular Biology</i> , 2008, 28, 4915-4926.  | 1.1 | 140       |
| 1462 | Different Levels of Bfa1/Bub2 GAP Activity Are Required to Prevent Mitotic Exit of Budding Yeast Depending on the Type of Perturbations. <i>Molecular Biology of the Cell</i> , 2008, 19, 4328-4340.                      | 0.9 | 16        |
| 1463 | Shugoshin Promotes Sister Kinetochore Biorientation in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2008, 19, 1199-1209.  | 0.9 | 43        |
| 1464 | Competition between the Rex1 exonuclease and the La protein affects both Trf4p-mediated RNA quality control and pre-tRNA maturation. <i>Rna</i> , 2008, 14, 1214-1227.  | 1.6 | 79        |
| 1465 | A Random Mutagenesis Approach to Isolate Dominant-Negative Yeast <i>sec1</i> Mutants Reveals a Functional Role for Domain 3a in Yeast and Mammalian Sec1/Munc18 Proteins. <i>Genetics</i> , 2008, 180, 165-178.           | 1.2 | 34        |
| 1466 | Arp2 Links Autophagic Machinery with the Actin Cytoskeleton. <i>Molecular Biology of the Cell</i> , 2008, 19, 1962-1975.  | 0.9 | 111       |
| 1467 | Nonapoptotic Death of <i>Saccharomyces cerevisiae</i> Cells That Is Stimulated by Hsp90 and Inhibited by Calcineurin and Cmk2 in Response to Endoplasmic Reticulum Stresses. <i>Eukaryotic Cell</i> , 2008, 7, 2037-2051. | 3.4 | 49        |
| 1468 | Swf1p, a Member of the DHHC-CRD Family of Palmitoyltransferases, Regulates the Actin Cytoskeleton and Polarized Secretion Independently of Its DHHC Motif. <i>Molecular Biology of the Cell</i> , 2008, 19, 4454-4468.    | 0.9 | 20        |
| 1469 | Molecular chaperone Hsp90 stabilizes Pih1/Nop17 to maintain R2TP complex activity that regulates snoRNA accumulation. <i>Journal of Cell Biology</i> , 2008, 180, 563-578.  | 2.3 | 159       |
| 1470 | G1/S Cyclin-dependent Kinase Regulates Small GTPase Rho1p through Phosphorylation of RhoGEF Tus1p in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2008, 19, 1763-1771.                        | 0.9 | 47        |
| 1471 | Yeast Est2p Affects Telomere Length by Influencing Association of Rap1p with Telomeric Chromatin. <i>Molecular and Cellular Biology</i> , 2008, 28, 2380-2390.  | 1.1 | 15        |
| 1472 | Absolute Gene Occupancies by RNA Polymerase III, TFIIB, and TFIIC in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2008, 283, 26568-26576.   | 1.6 | 23        |
| 1473 | Ubiquitin Ligase Hul5 Is Required for Fragment-specific Substrate Degradation in Endoplasmic Reticulum-associated Degradation. <i>Journal of Biological Chemistry</i> , 2008, 283, 16374-16383.                           | 1.6 | 46        |
| 1474 | The Peripheral Membrane Subunits of the SAM Complex Function Codependently in Mitochondrial Outer Membrane Biogenesis. <i>Molecular Biology of the Cell</i> , 2008, 19, 126-136.  | 0.9 | 115       |
| 1475 | Divergent functions of three <i>Candida albicans</i> zinc-cluster transcription factors (CTA4, ASG1 and) Tj ETQq1 1 0.784314 rgBT /Overl  | 0.7 | 37        |
| 1476 | Role of the <i>Saccharomyces cerevisiae</i> Rad51 Paralogs in Sister Chromatid Recombination. <i>Genetics</i> , 2008, 178, 113-126.   | 1.2 | 63        |
| 1477 | Distinct roles for Khd1p in the localization and expression of bud-localized mRNAs in yeast. <i>Rna</i> , 2008, 14, 2333-2347.  | 1.6 | 65        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1478 | Evidence for lesion bypass by yeast replicative DNA polymerases during DNA damage. <i>Nucleic Acids Research</i> , 2008, 36, 5660-5667.   | 6.5 | 80        |
| 1479 | Organization of the Pre-autophagosomal Structure Responsible for Autophagosome Formation. <i>Molecular Biology of the Cell</i> , 2008, 19, 2039-2050.   | 0.9 | 233       |
| 1480 | Rad53 regulates replication fork restart after DNA damage in <i>Saccharomyces cerevisiae</i> . <i>Genes and Development</i> , 2008, 22, 1906-1920.  | 2.7 | 81        |
| 1481 | Ubiquitinâ€“Proteasome-dependent Degradation of a Mitofusin, a Critical Regulator of Mitochondrial Fusion. <i>Molecular Biology of the Cell</i> , 2008, 19, 2457-2464.  | 0.9 | 146       |
| 1482 | Disrupting Vesicular Trafficking at the Endosome Attenuates Transcriptional Activation by Gcn4. <i>Molecular and Cellular Biology</i> , 2008, 28, 6796-6818.  | 1.1 | 23        |
| 1483 | Mrd1p binds to pre-rRNA early during transcription independent of U3 snoRNA and is required for compaction of the pre-rRNA into small subunit processomes. <i>Nucleic Acids Research</i> , 2008, 36, 4364-4380. | 6.5 | 18        |
| 1484 | The Telotype Defines the Telomere State in <i>Saccharomyces cerevisiae</i> and Is Inherited as a Dominant Non-Mendelian Characteristic in Cells Lacking Telomerase. <i>Genetics</i> , 2008, 178, 245-257.       | 1.2 | 18        |
| 1485 | Ypi1, a Positive Regulator of Nuclear Protein Phosphatase Type 1 Activity in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2008, 19, 1032-1045.                                      | 0.9 | 41        |
| 1486 | Mgr3p and Mgr1p Are Adaptors for the Mitochondrial i-AAA Protease Complex. <i>Molecular Biology of the Cell</i> , 2008, 19, 5387-5397.  | 0.9 | 49        |
| 1487 | Functional Dissection of a HECT Ubiquitin E3 Ligase. <i>Molecular and Cellular Proteomics</i> , 2008, 7, 35-45.   | 2.5 | 70        |
| 1488 | Gcn5p Plays an Important Role in Centromere Kinetochores Function in Budding Yeast. <i>Molecular and Cellular Biology</i> , 2008, 28, 988-996.  | 1.1 | 39        |
| 1489 | Human Peroxiredoxin Prxl Is an Orthologue of Yeast Tsa1, Capable of Suppressing Genome Instability in <i>Saccharomyces cerevisiae</i> . <i>Cancer Research</i> , 2008, 68, 1055-1063.                           | 0.4 | 24        |
| 1490 | Recruitment of <i>Saccharomyces cerevisiae</i> Dnl4â€“Lif1 Complex to a Double-Strand Break Requires Interactions With Yku80 and the Xrs2 FHA Domain. <i>Genetics</i> , 2008, 180, 1809-1819.                   | 1.2 | 74        |
| 1491 | Identification and Verification of Sro7p as an Effector of the Sec4p Rab GTPase. <i>Methods in Enzymology</i> , 2008, 438, 95-108.  | 0.4 | 0         |
| 1492 | The Polo-like kinase Cdc5 interacts with FEAR network components and Cdc14. <i>Cell Cycle</i> , 2008, 7, 3262-3272.   | 1.3 | 24        |
| 1493 | Toxicity of Nickel Compounds Mediated by HTZ1, Histone Variant H2A.Z, in <i>Saccharomyces cerevisiae</i> . <i>Biological and Pharmaceutical Bulletin</i> , 2008, 31, 2007-2011.                                 | 0.6 | 1         |
| 1494 | Analysis of mutant origin recognition complex with reduced ATPase activity <i>in vivo</i> and <i>in vitro</i> . <i>Biochemical Journal</i> , 2008, 413, 535-543.  | 1.7 | 8         |
| 1495 | Chromatin- and Transcription-Related Factors Repress Transcription from within Coding Regions throughout the <i>Saccharomyces cerevisiae</i> Genome. <i>PLoS Biology</i> , 2008, 6, e277.                       | 2.6 | 268       |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1496 | The Septins Function in G1 Pathways that Influence the Pattern of Cell Growth in Budding Yeast. <i>PLoS ONE</i> , 2008, 3, e2022.  | 1.1 | 41        |
| 1497 | Whi5 Regulation by Site Specific CDK-Phosphorylation in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2009, 4, e4300.  | 1.1 | 61        |
| 1498 | Cdc7p-Dbf4p Regulates Mitotic Exit by Inhibiting Polo Kinase. <i>PLoS Genetics</i> , 2009, 5, e1000498.  | 1.5 | 38        |
| 1499 | The Rts1 Regulatory Subunit of Protein Phosphatase 2A Is Required for Control of G1 Cyclin Transcription and Nutrient Modulation of Cell Size. <i>PLoS Genetics</i> , 2009, 5, e1000727.   | 1.5 | 31        |
| 1500 | A Newly Identified Essential Complex, Dre2-Tah18, Controls Mitochondria Integrity and Cell Death after Oxidative Stress in Yeast. <i>PLoS ONE</i> , 2009, 4, e4376.  | 1.1 | 64        |
| 1501 | Multisite Phosphorylation of the Guanine Nucleotide Exchange Factor Cdc24 during Yeast Cell Polarization. <i>PLoS ONE</i> , 2009, 4, e6563.  | 1.1 | 33        |
| 1502 | The CCR4-NOT Complex Physically and Functionally Interacts with TRAMP and the Nuclear Exosome. <i>PLoS ONE</i> , 2009, 4, e6760.   | 1.1 | 56        |
| 1503 | A Screen for Spore Wall Permeability Mutants Identifies a Secreted Protease Required for Proper Spore Wall Assembly. <i>PLoS ONE</i> , 2009, 4, e7184.   | 1.1 | 36        |
| 1504 | Sulfur Metabolism Actively Promotes Initiation of Cell Division in Yeast. <i>PLoS ONE</i> , 2009, 4, e8018.  | 1.1 | 17        |
| 1506 | Coassembly of Mgm1 isoforms requires cardiolipin and mediates mitochondrial inner membrane fusion. <i>Journal of Cell Biology</i> , 2009, 186, 793-803.  | 2.3 | 243       |
| 1507 | Nbl1p: A Borealin/Dasra/CSC-1-like Protein Essential for Aurora/Ipl1 Complex Function and Integrity in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2009, 20, 1772-1784.   | 0.9 | 32        |
| 1508 | The Anaphase Promoting Complex Targeting Subunit Ama1 Links Meiotic Exit to Cytokinesis during Sporulation in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2009, 20, 134-145.  | 0.9 | 65        |
| 1509 | Genome-wide replication profiles indicate an expansive role for Rpd3L in regulating replication initiation timing or efficiency, and reveal genomic loci of Rpd3 function in <i>Saccharomyces cerevisiae</i> . <i>Genes and Development</i> , 2009, 23, 1077-1090. | 2.7 | 116       |
| 1510 | Histone deacetylase Rpd3 antagonizes Sir2-dependent silent chromatin propagation. <i>Nucleic Acids Research</i> , 2009, 37, 3699-3713.   | 6.5 | 56        |
| 1511 | The nucleoporins Nup170p and Nup157p are essential for nuclear pore complex assembly. <i>Journal of Cell Biology</i> , 2009, 185, 459-473.   | 2.3 | 78        |
| 1512 | Gene Looping Is Conferred by Activator-dependent Interaction of Transcription Initiation and Termination Machineries. <i>Journal of Biological Chemistry</i> , 2009, 284, 25015-25025.   | 1.6 | 75        |
| 1513 | The CORVET Subunit Vps8 Cooperates with the Rab5 Homolog Vps21 to Induce Clustering of Late Endosomal Compartments. <i>Molecular Biology of the Cell</i> , 2009, 20, 5276-5289.  | 0.9 | 83        |
| 1514 | Kinase-Independent Functions of <i>TEL1</i> in Telomere Maintenance. <i>Molecular and Cellular Biology</i> , 2009, 29, 5193-5202.  | 1.1 | 27        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1515 | A nonproteolytic proteasome activity controls organelle fission in yeast. <i>Journal of Cell Science</i> , 2009, 122, 3673-3683.  | 1.2 | 17        |
| 1516 | Phosphorylation of the Transcription Elongation Factor Spt5 by Yeast Bur1 Kinase Stimulates Recruitment of the PAF Complex. <i>Molecular and Cellular Biology</i> , 2009, 29, 4852-4863.  | 1.1 | 155       |
| 1517 | Nej1 recruits the Srs2 helicase to DNA double-strand breaks and supports repair by a single-strand annealing-like mechanism. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 12037-12042.           | 3.3 | 19        |
| 1518 | Stress-induced Ceramide-activated Protein Phosphatase Can Compensate for Loss of Amphiphysin-like Activity In <i>Saccharomyces cerevisiae</i> and Functions to Reinitiate Endocytosis. <i>Journal of Biological Chemistry</i> , 2009, 284, 11930-11941. | 1.6 | 14        |
| 1519 | Contrasting Roles of Checkpoint Proteins as Recombination Modulators at Fob1- <i>Ter</i> Complexes with or without Fork Arrest. <i>Eukaryotic Cell</i> , 2009, 8, 487-495.  | 3.4 | 23        |
| 1520 | Gga2 Mediates Sequential Ubiquitin-independent and Ubiquitin-dependent Steps in the Trafficking of ARN1 from the trans-Golgi Network to the Vacuole. <i>Journal of Biological Chemistry</i> , 2009, 284, 23830-23841.                                   | 1.6 | 37        |
| 1521 | Fungal Functional Genomics: Tunable Knockout-Knock-in Expression and Tagging Strategies. <i>Eukaryotic Cell</i> , 2009, 8, 800-804.   | 3.4 | 31        |
| 1522 | Mutational Uncoupling of the Role of Sus1 in Nuclear Pore Complex Targeting of an mRNA Export Complex and Histone H2B Deubiquitination. <i>Journal of Biological Chemistry</i> , 2009, 284, 12049-12056.  | 1.6 | 21        |
| 1523 | NTE1-encoded Phosphatidylcholine Phospholipase B Regulates Transcription of Phospholipid Biosynthetic Genes. <i>Journal of Biological Chemistry</i> , 2009, 284, 36034-36046.   | 1.6 | 35        |
| 1524 | The Major Role of the Rab Ypt7p in Vacuole Fusion Is Supporting HOPS Membrane Association. <i>Journal of Biological Chemistry</i> , 2009, 284, 16118-16125.   | 1.6 | 68        |
| 1525 | The Gcn5 Bromodomain of the SAGA Complex Facilitates Cooperative and Cross-tail Acetylation of Nucleosomes. <i>Journal of Biological Chemistry</i> , 2009, 284, 9411-9417.  | 1.6 | 68        |
| 1526 | Lipoic Acid Synthesis and Attachment in Yeast Mitochondria. <i>Journal of Biological Chemistry</i> , 2009, 284, 23234-23242.  | 1.6 | 110       |
| 1527 | The Yeast RNA Polymerase II-associated Factor Iwr1p Is Involved in the Basal and Regulated Transcription of Specific Genes. <i>Journal of Biological Chemistry</i> , 2009, 284, 28958-28967.  | 1.6 | 15        |
| 1528 | The 19 S Proteasome Subcomplex Establishes a Specific Protein Interaction Network at the Promoter for Stimulated Transcriptional Initiation in Vivo. <i>Journal of Biological Chemistry</i> , 2009, 284, 35714-35724.                                   | 1.6 | 37        |
| 1529 | Reconstitution of Rad53 Activation by Mec1 through Adaptor Protein Mrc1. <i>Journal of Biological Chemistry</i> , 2009, 284, 18593-18604.   | 1.6 | 42        |
| 1530 | Control of Protein and Sterol Trafficking by Antagonistic Activities of a Type IV P-type ATPase and Oxysterol Binding Protein Homologue. <i>Molecular Biology of the Cell</i> , 2009, 20, 2920-2931.  | 0.9 | 41        |
| 1531 | Genome-wide Analysis of AP-3-dependent Protein Transport in Yeast. <i>Molecular Biology of the Cell</i> , 2009, 20, 1592-1604.  | 0.9 | 43        |
| 1532 | A network biology approach to aging in yeast. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 1145-1150.  | 3.3 | 69        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1533 | Distinct Subsets of Sit4 Holophosphatases Are Required for Inhibition of <i>Saccharomyces cerevisiae</i> Growth by Rapamycin and Zymocin. <i>Eukaryotic Cell</i> , 2009, 8, 1637-1647.   | 3.4 | 21        |
| 1534 | <i>Saccharomyces cerevisiae</i> Rbg1 Protein and Its Binding Partner Gir2 Interact on Polyribosomes with Gcn1. <i>Eukaryotic Cell</i> , 2009, 8, 1061-1071.  | 3.4 | 29        |
| 1535 | The protein phosphatase 2A functions in the spindle position checkpoint by regulating the checkpoint kinase Kin4. <i>Genes and Development</i> , 2009, 23, 1639-1649.  | 2.7 | 53        |
| 1536 | Role of the Ndc1 interaction network in yeast nuclear pore complex assembly and maintenance. <i>Journal of Cell Biology</i> , 2009, 185, 475-491.  | 2.3 | 139       |
| 1537 | An Internal Domain of Exo70p Is Required for Actin-independent Localization and Mediates Assembly of Specific Exocyst Components. <i>Molecular Biology of the Cell</i> , 2009, 20, 153-163.                                      | 0.9 | 30        |
| 1538 | The Exomer Coat Complex Transports Fus1p to the Plasma Membrane via a Novel Plasma Membrane Sorting Signal in Yeast. <i>Molecular Biology of the Cell</i> , 2009, 20, 4985-4996.   | 0.9 | 55        |
| 1539 | Mediator complex association with constitutively transcribed genes in yeast. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 16734-16739.                                    | 3.3 | 59        |
| 1540 | RNA Helicase Prp43 and Its Co-factor Pfa1 Promote 20 to 18 S rRNA Processing Catalyzed by the Endonuclease Nob1. <i>Journal of Biological Chemistry</i> , 2009, 284, 35079-35091.  | 1.6 | 166       |
| 1541 | Aup1-mediated Regulation of Rtg3 during Mitophagy. <i>Journal of Biological Chemistry</i> , 2009, 284, 35885-35895.  | 1.6 | 79        |
| 1542 | Nascent Peptide-dependent Translation Arrest Leads to Not4p-mediated Protein Degradation by the Proteasome. <i>Journal of Biological Chemistry</i> , 2009, 284, 10343-10352.   | 1.6 | 220       |
| 1543 | Role of Inn1 and its interactions with Hof1 and Cyk3 in promoting cleavage furrow and septum formation in <i>S. cerevisiae</i> . <i>Journal of Cell Biology</i> , 2009, 185, 995-1012.   | 2.3 | 87        |
| 1544 | The Cdc42 Effectors Ste20, Cla4, and Skm1 Down-Regulate the Expression of Genes Involved in Sterol Uptake by a Mitogen-activated Protein Kinase-independent Pathway. <i>Molecular Biology of the Cell</i> , 2009, 20, 4826-4837. | 0.9 | 22        |
| 1545 | Characterization of a Highly Conserved Binding Site of Mlh1 Required for Exonuclease I-Dependent Mismatch Repair. <i>Molecular and Cellular Biology</i> , 2009, 29, 907-918.   | 1.1 | 56        |
| 1546 | Telomere Maintenance and Survival in <i>Saccharomyces cerevisiae</i> in the Absence of Telomerase and RAD52. <i>Genetics</i> , 2009, 182, 671-684.   | 1.2 | 25        |
| 1547 | The N-terminal PIN domain of the exosome subunit Rrp44 harbors endonuclease activity and tethers Rrp44 to the yeast core exosome. <i>Nucleic Acids Research</i> , 2009, 37, 1127-1140.   | 6.5 | 202       |
| 1548 | The WASP Homologue Las17 Activates the Novel Actin-regulatory Activity of Ysc84 to Promote Endocytosis in Yeast. <i>Molecular Biology of the Cell</i> , 2009, 20, 1618-1628.   | 0.9 | 35        |
| 1549 | Mitochondrial outer and inner membrane fusion requires a modified carrier protein. <i>Journal of Cell Biology</i> , 2009, 184, 569-581.  | 2.3 | 69        |
| 1550 | The Ime2 Protein Kinase Enhances the Disassociation of the Sum1 Repressor from Middle Meiotic Promoters. <i>Molecular and Cellular Biology</i> , 2009, 29, 4352-4362.  | 1.1 | 32        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1551 | <i>Candida glabrata</i> PHO4 Is Necessary and Sufficient for Pho2-Independent Transcription of Phosphate Starvation Genes. <i>Genetics</i> , 2009, 182, 471-479.   | 1.2 | 28        |
| 1552 | In <i>Saccharomyces cerevisiae</i> , yKu and Subtelomeric Core X Sequences Repress Homologous Recombination Near Telomeres as Part of the Same Pathway. <i>Genetics</i> , 2009, 183, 441-451.                          | 1.2 | 22        |
| 1553 | Physical and genetic interactions of yeast Cwc21p, an ortholog of human SRm300/SRRM2, suggest a role at the catalytic center of the spliceosome. <i>Rna</i> , 2009, 15, 2161-2173.                                     | 1.6 | 51        |
| 1554 | p21-Activated Kinases Cla4 and Ste20 Regulate Vacuole Inheritance in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2009, 8, 560-572.  | 3.4 | 27        |
| 1555 | A CAAX motif can compensate for the PH domain of Num1 for cortical dynein attachment. <i>Cell Cycle</i> , 2009, 8, 3182-3190.  | 1.3 | 47        |
| 1556 | Bck2 is a phase-independent activator of cell cycle-regulated genes in yeast. <i>Cell Cycle</i> , 2009, 8, 239-252.  | 1.3 | 28        |
| 1557 | The SUMO-targeted ubiquitin ligase subunit Slx5 resides in nuclear foci and at sites of DNA breaks. <i>Cell Cycle</i> , 2009, 8, 1080-1089.  | 1.3 | 40        |
| 1558 | Ultrastructural Analysis of Nanogold-labeled Endocytic Compartments of Yeast <i>Saccharomyces cerevisiae</i> Using a Cryosectioning Procedure. <i>Journal of Histochemistry and Cytochemistry</i> , 2009, 57, 801-809. | 1.3 | 13        |
| 1559 | KIAft, the <i>Kluyveromyces lactis</i> Ortholog of Aft1 and Aft2, Mediates Activation of Iron-Responsive Transcription Through the PuCACCC Aft-Type Sequence. <i>Genetics</i> , 2009, 183, 93-106.                     | 1.2 | 32        |
| 1560 | Mutations in the U5 snRNA result in altered splicing of subsets of pre-mRNAs and reduced stability of Prp8. <i>Rna</i> , 2009, 15, 1292-1304.  | 1.6 | 13        |
| 1561 | The RNA binding protein Cwc2 interacts directly with the U6 snRNA to link the nineteen complex to the spliceosome during pre-mRNA splicing. <i>Nucleic Acids Research</i> , 2009, 37, 4205-4217.                       | 6.5 | 35        |
| 1562 | Shugoshin prevents cohesin cleavage by PP2A <sup>Cdc55</sup> -dependent inhibition of separase. <i>Genes and Development</i> , 2009, 23, 766-780.  | 2.7 | 59        |
| 1563 | Establishment of Cohesion at the Pericentromere by the Ctf19 Kinetochore Subcomplex and the Replication Fork-Associated Factor, Csm3. <i>PLoS Genetics</i> , 2009, 5, e1000629.  | 1.5 | 87        |
| 1564 | Dual Regulation by Pairs of Cyclin-Dependent Protein Kinases and Histone Deacetylases Controls G1 Transcription in Budding Yeast. <i>PLoS Biology</i> , 2009, 7, e1000188.   | 2.6 | 42        |
| 1565 | The cap binding complex influences H2B ubiquitination by facilitating splicing of the <i>SUS1</i> pre-mRNA. <i>Rna</i> , 2009, 15, 1515-1527.  | 1.6 | 31        |
| 1566 | Bayesian Modeling of the Yeast SH3 Domain Interactome Predicts Spatiotemporal Dynamics of Endocytosis Proteins. <i>PLoS Biology</i> , 2009, 7, e1000218.   | 2.6 | 172       |
| 1567 | Peroxiredoxin Tsa1 Is the Key Peroxidase Suppressing Genome Instability and Protecting against Cell Death in <i>Saccharomyces cerevisiae</i> . <i>PLoS Genetics</i> , 2009, 5, e1000524.                               | 1.5 | 58        |
| 1568 | The Origin Recognition Complex Interacts with a Subset of Metabolic Genes Tightly Linked to Origins of Replication. <i>PLoS Genetics</i> , 2009, 5, e1000755.  | 1.5 | 25        |



| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1569 | Yeast telomerase and the SUN domain protein Mps3 anchor telomeres and repress subtelomeric recombination. <i>Genes and Development</i> , 2009, 23, 928-938.  | 2.7 | 164       |
| 1570 | A Genome-Wide Screen for Regulators of TORC1 in Response to Amino Acid Starvation Reveals a Conserved Npr2/3 Complex. <i>PLoS Genetics</i> , 2009, 5, e1000515.  | 1.5 | 134       |
| 1571 | Loss of Yeast Peroxiredoxin Tsa1p Induces Genome Instability through Activation of the DNA Damage Checkpoint and Elevation of dNTP Levels. <i>PLoS Genetics</i> , 2009, 5, e1000697.   | 1.5 | 38        |
| 1572 | Sem1 is a functional component of the nuclear pore complex-associated messenger RNA export machinery. <i>Journal of Cell Biology</i> , 2009, 184, 833-846.   | 2.3 | 96        |
| 1573 | Sgt1 Dimerization Is Negatively Regulated by Protein Kinase CK2-mediated Phosphorylation at Ser361. <i>Journal of Biological Chemistry</i> , 2009, 284, 18692-18698.   | 1.6 | 27        |
| 1574 | Budding Yeast <i>SSD1-V</i> Regulates Transcript Levels of Many Longevity Genes and Extends Chronological Life Span in Purified Quiescent Cells. <i>Molecular Biology of the Cell</i> , 2009, 20, 3851-3864.                   | 0.9 | 46        |
| 1575 | PCNA monoubiquitylation and DNA polymerase $\beta$ ubiquitin-binding domain are required to prevent 8-oxoguanine-induced mutagenesis in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2009, 37, 2549-2559. | 6.5 | 34        |
| 1576 | Actin-Mediated Endocytosis Limits Intracellular Cr Accumulation and Cr Toxicity during Chromate Stress. <i>Toxicological Sciences</i> , 2009, 111, 437-446.  | 1.4 | 28        |
| 1577 | Role of the Cell Wall Integrity and Filamentous Growth Mitogen-Activated Protein Kinase Pathways in Cell Wall Remodeling during Filamentous Growth. <i>Eukaryotic Cell</i> , 2009, 8, 1118-1133.                               | 3.4 | 45        |
| 1578 | The functional importance of telomere clustering: Global changes in gene expression result from SIR factor dispersion. <i>Genome Research</i> , 2009, 19, 611-625.   | 2.4 | 110       |
| 1579 | Biofilm Matrix Regulation by <i>Candida albicans</i> Zap1. <i>PLoS Biology</i> , 2009, 7, e1000133.  | 2.6 | 286       |
| 1580 | The Synaptonemal Complex Protein Zip1 Promotes Bi-Orientation of Centromeres at Meiosis I. <i>PLoS Genetics</i> , 2009, 5, e1000771.   | 1.5 | 68        |
| 1581 | Nucleosome Remodeling and Transcriptional Repression Are Distinct Functions of Isw1 in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2009, 29, 2419-2430.  | 1.1 | 29        |
| 1582 | Two Dot1 isoforms in <i>Saccharomyces cerevisiae</i> as a result of leaky scanning by the ribosome. <i>Nucleic Acids Research</i> , 2009, 37, 7047-7058.   | 6.5 | 10        |
| 1583 | Two structurally distinct domains of the nucleoporin Nup170 cooperate to tether a subset of nucleoporins to nuclear pores. <i>Journal of Cell Biology</i> , 2009, 185, 387-395.  | 2.3 | 35        |
| 1584 | Alternative Splicing of PTC7 in <i>Saccharomyces cerevisiae</i> Determines Protein Localization. <i>Genetics</i> , 2009, 183, 185-194.   | 1.2 | 71        |
| 1585 | Molecular Dissection of Erv26p Identifies Separable Cargo Binding and Coat Protein Sorting Activities. <i>Journal of Biological Chemistry</i> , 2009, 284, 24049-24060.  | 1.6 | 25        |
| 1586 | A Genomic Screen for Yeast Mutants Defective in Selective Mitochondria Autophagy. <i>Molecular Biology of the Cell</i> , 2009, 20, 4730-4738.  | 0.9 | 229       |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1587 | Deleting the 14-3-3 Protein Bmh1 Extends Life Span in <i>Saccharomyces cerevisiae</i> by Increasing Stress Response. <i>Genetics</i> , 2009, 183, 1373-1384.   | 1.2 | 33        |
| 1588 | Tap42-associated protein phosphatase type 2A negatively regulates induction of autophagy. <i>Autophagy</i> , 2009, 5, 616-624.   | 4.3 | 78        |
| 1589 | Mrd1p Is Required for Release of Base-Paired U3 snoRNA within the Preribosomal Complex. <i>Molecular and Cellular Biology</i> , 2009, 29, 5763-5774.   | 1.1 | 9         |
| 1590 | Sgt1 Dimerization Is Required for Yeast Kinetochores Assembly. <i>Journal of Biological Chemistry</i> , 2009, 284, 3586-3592.  | 1.6 | 26        |
| 1591 | The Dot1 Histone Methyltransferase and the Rad9 Checkpoint Adaptor Contribute to Cohesin-Dependent Double-Strand Break Repair by Sister Chromatid Recombination in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2009, 182, 437-446. | 1.2 | 57        |
| 1592 | Ipl1-dependent phosphorylation of Dam1 is reduced by tension applied on kinetochores. <i>Journal of Cell Science</i> , 2009, 122, 4375-4382.   | 1.2 | 61        |
| 1593 | Pds1p Is Required for Meiotic Recombination and Prophase I Progression in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2009, 181, 65-79.  | 1.2 | 11        |
| 1594 | Dynamic and complex transcription factor binding during an inducible response in yeast. <i>Genes and Development</i> , 2009, 23, 1351-1363.  | 2.7 | 89        |
| 1595 | Altering Sphingolipid Metabolism in <i>Saccharomyces cerevisiae</i> Cells Lacking the Amphiphysin Ortholog Rvs161 Reinitiates Sugar Transporter Endocytosis. <i>Eukaryotic Cell</i> , 2009, 8, 779-789.                                    | 3.4 | 6         |
| 1596 | A Novel Proteomics Approach for the Discovery of Chromatin-associated Protein Networks. <i>Molecular and Cellular Proteomics</i> , 2009, 8, 870-882.   | 2.5 | 95        |
| 1597 | Functional Redundancy of Yeast Proteins Reh1 and Rei1 in Cytoplasmic 60S Subunit Maturation. <i>Molecular and Cellular Biology</i> , 2009, 29, 4014-4023.  | 1.1 | 27        |
| 1598 | Membrane expansion alleviates endoplasmic reticulum stress independently of the unfolded protein response. <i>Journal of Cell Biology</i> , 2009, 187, 525-536.  | 2.3 | 451       |
| 1599 | Dynein-Driven Mitotic Spindle Positioning Restricted to Anaphase by She1p Inhibition of Dynactin Recruitment. <i>Molecular Biology of the Cell</i> , 2009, 20, 3003-3011.  | 0.9 | 38        |
| 1600 | PtdIns4P recognition by Vps74/GOLPH3 links PtdIns 4-kinase signaling to retrograde Golgi trafficking. <i>Journal of Cell Biology</i> , 2009, 187, 967-975.   | 2.3 | 147       |
| 1601 | Direct Link between RACK1 Function and Localization at the Ribosome In Vivo. <i>Molecular and Cellular Biology</i> , 2009, 29, 1626-1634.  | 1.1 | 136       |
| 1602 | Hyphal Chain Formation in <i>Candida albicans</i> : Cdc28-Hgc1 Phosphorylation of Efg1 Represses Cell Separation Genes. <i>Molecular and Cellular Biology</i> , 2009, 29, 4406-4416.   | 1.1 | 71        |
| 1603 | Stringent Mating-Type-Regulated Auxotrophy Increases the Accuracy of Systematic Genetic Interaction Screens with <i>Saccharomyces cerevisiae</i> Mutant Arrays. <i>Genetics</i> , 2009, 181, 289-300.                                      | 1.2 | 20        |
| 1604 | Protein Phosphatase Type 1-Interacting Protein Ysw1 Is Involved in Proper Septin Organization and Prospore Membrane Formation during Sporulation. <i>Eukaryotic Cell</i> , 2009, 8, 1027-1037.   | 3.4 | 11        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1605 | Sem1p and Ubp6p orchestrate telomeric silencing by modulating histone H2B ubiquitination and H3 acetylation. <i>Nucleic Acids Research</i> , 2009, 37, 1843-1853.   | 6.5 | 22        |
| 1606 | The NatA Acetyltransferase Couples Sup35 Prion Complexes to the [PSI <sup>+</sup> ] Phenotype. <i>Molecular Biology of the Cell</i> , 2009, 20, 1068-1080.  | 0.9 | 20        |
| 1607 | Control of transcriptional elongation and cotranscriptional histone modification by the yeast BUR kinase substrate Spt5. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 6956-6961. | 3.3 | 127       |
| 1608 | Cooperative Regulation of ADE3 Transcription by Gcn4p and Bas1p in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2009, 8, 1268-1277.   | 3.4 | 13        |
| 1609 | Regulation of Nuclear Positioning and Dynamics of the Silent Mating Type Loci by the Yeast Ku70/Ku80 Complex. <i>Molecular and Cellular Biology</i> , 2009, 29, 835-848.  | 1.1 | 71        |
| 1610 | Early-Arriving Syp1p and Ede1p Function in Endocytic Site Placement and Formation in Budding Yeast. <i>Molecular Biology of the Cell</i> , 2009, 20, 4640-4651.   | 0.9 | 110       |
| 1611 | Dynamic Compartmentalization of Base Excision Repair Proteins in Response to Nuclear and Mitochondrial Oxidative Stress. <i>Molecular and Cellular Biology</i> , 2009, 29, 794-807.   | 1.1 | 40        |
| 1612 | Ribosome Association and Stability of the Nascent Polypeptide-Associated Complex Is Dependent Upon Its Own Ubiquitination. <i>Genetics</i> , 2009, 181, 447-460.  | 1.2 | 37        |
| 1613 | Role of the Histone Variant H2A.Z/Htz1p in TBP Recruitment, Chromatin Dynamics, and Regulated Expression of Oleate-Responsive Genes. <i>Molecular and Cellular Biology</i> , 2009, 29, 2346-2358.                                       | 1.1 | 77        |
| 1614 | Trans-acting antisense RNAs mediate transcriptional gene cosuppression in <i>S. cerevisiae</i> . <i>Genes and Development</i> , 2009, 23, 1534-1545.  | 2.7 | 138       |
| 1615 | Genetic Identification of Factors That Modulate Ribosomal DNA Transcription in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2009, 182, 105-119.  | 1.2 | 35        |
| 1616 | Regulation of Cell Polarity through Phosphorylation of Bni4 by Pho85 G1 Cyclin-dependent Kinases in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2009, 20, 3239-3250.                                       | 0.9 | 31        |
| 1617 | The Signaling Mucins Msb2 and Hkr1 Differentially Regulate the Filamentation Mitogen-activated Protein Kinase Pathway and Contribute to a Multimodal Response. <i>Molecular Biology of the Cell</i> , 2009, 20, 3101-3114.              | 0.9 | 87        |
| 1618 | Methionine sulphoxide reductases protect iron-sulphur clusters from oxidative inactivation in yeast. <i>Microbiology (United Kingdom)</i> , 2009, 155, 612-623.   | 0.7 | 24        |
| 1619 | Distinct Roles for Key Karyogamy Proteins during Yeast Nuclear Fusion. <i>Molecular Biology of the Cell</i> , 2009, 20, 3773-3782.  | 0.9 | 24        |
| 1620 | The Yeast GATA Factor Gat1 Occupies a Central Position in Nitrogen Catabolite Repression-Sensitive Gene Activation. <i>Molecular and Cellular Biology</i> , 2009, 29, 3803-3815.  | 1.1 | 70        |
| 1621 | Multicopy suppressor analysis of thermosensitive YIP1 alleles implicates GOT1 in transport from the ER. <i>Journal of Cell Science</i> , 2009, 122, 1540-1550.  | 1.2 | 38        |
| 1622 | A Phosphatidylinositol-Transfer Protein and Phosphatidylinositol-4-phosphate 5-Kinase Control Cdc42 to Regulate the Actin Cytoskeleton and Secretory Pathway in Yeast. <i>Molecular Biology of the Cell</i> , 2009, 20, 3583-3597.      | 0.9 | 38        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1623 | The tRNA Modification Complex Elongator Regulates the Cdc42-Dependent Mitogen-Activated Protein Kinase Pathway That Controls Filamentous Growth in Yeast. <i>Eukaryotic Cell</i> , 2009, 8, 1362-1372.                                       | 3.4 | 24        |
| 1624 | New High-Throughput Screening Assay To Reveal Similarities and Differences in Inhibitory Sensitivities of Multidrug ATP-Binding Cassette Transporters. <i>Antimicrobial Agents and Chemotherapy</i> , 2009, 53, 1516-1527.                   | 1.4 | 21        |
| 1625 | The Direct Binding of Mrc1, a Checkpoint Mediator, to Mcm6, a Replication Helicase, Is Essential for the Replication Checkpoint against Methyl Methanesulfonate-Induced Stress. <i>Molecular and Cellular Biology</i> , 2009, 29, 5008-5019. | 1.1 | 60        |
| 1626 | Yvh1 is required for a late maturation step in the 60S biogenesis pathway. <i>Journal of Cell Biology</i> , 2009, 186, 863-880.  | 2.3 | 106       |
| 1627 | Antagonistic regulation of Fus2p nuclear localization by pheromone signaling and the cell cycle. <i>Journal of Cell Biology</i> , 2009, 184, 409-422.  | 2.3 | 22        |
| 1628 | The Association of yKu With Subtelomeric Core X Sequences Prevents Recombination Involving Telomeric Sequences. <i>Genetics</i> , 2009, 183, 453-467.  | 1.2 | 28        |
| 1629 | The Multiple Roles of Cohesin in Meiotic Chromosome Morphogenesis and Pairing. <i>Molecular Biology of the Cell</i> , 2009, 20, 1030-1047.   | 0.9 | 85        |
| 1630 | <i>PTC1</i> Is Required for Vacuole Inheritance and Promotes the Association of the Myosin-V Vacuole-specific Receptor Complex. <i>Molecular Biology of the Cell</i> , 2009, 20, 1312-1323.  | 0.9 | 40        |
| 1631 | Evidence of carbon monoxide-mediated phase advancement of the yeast metabolic cycle. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 14293-14296.  | 3.3 | 20        |
| 1632 | Nonsense-mediated mRNA decay mutes the splicing defects of spliceosome component mutations. <i>Rna</i> , 2009, 15, 2236-2247.  | 1.6 | 30        |
| 1633 | Phylogenetic Conservation and Homology Modeling Help Reveal a Novel Domain within the Budding Yeast Heterochromatin Protein Sir1. <i>Molecular and Cellular Biology</i> , 2009, 29, 687-702.   | 1.1 | 9         |
| 1634 | The Structure of the Scaffold Nucleoporin Nup120 Reveals a New and Unexpected Domain Architecture. <i>Structure</i> , 2009, 17, 1082-1091.   | 1.6 | 56        |
| 1635 | Checkpoint kinase phosphorylation in response to endogenous oxidative DNA damage in repair-deficient stationary-phase <i>Saccharomyces cerevisiae</i> . <i>Mechanisms of Ageing and Development</i> , 2009, 130, 501-508.                    | 2.2 | 14        |
| 1636 | The human actin-related protein hArap5: Nucleo-cytoplasmic shuttling and involvement in DNA repair. <i>Experimental Cell Research</i> , 2009, 315, 206-217.  | 1.2 | 37        |
| 1637 | Motor- and Tail-Dependent Targeting of Dynein to Microtubule Plus Ends and the Cell Cortex. <i>Current Biology</i> , 2009, 19, 196-205.  | 1.8 | 102       |
| 1638 | Regulation of Spo12 Phosphorylation and Its Essential Role in the FEAR Network. <i>Current Biology</i> , 2009, 19, 449-460.  | 1.8 | 39        |
| 1639 | Protein Phosphatase 1 Regulates Exit from the Spindle Checkpoint in Budding Yeast. <i>Current Biology</i> , 2009, 19, 1182-1187.   | 1.8 | 138       |
| 1640 | Mitochondrial Inheritance Is Required for MEN-Regulated Cytokinesis in Budding Yeast. <i>Current Biology</i> , 2009, 19, 1730-1735.  | 1.8 | 31        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1641 | Reciprocal regulation of nuclear import of the yeast MutS-like DNA mismatch repair proteins Msh2 and Msh6. <i>DNA Repair</i> , 2009, 8, 739-751.  | 1.3 | 17        |
| 1642 | Jid1 is a protein functioning in the mitochondrial matrix, unable to directly participate in endoplasmic reticulum associated protein degradation. <i>FEBS Letters</i> , 2009, 583, 2954-2958.  | 1.3 | 7         |
| 1643 | Isolation and partial purification of the <i>Saccharomyces cerevisiae</i> cytokinetic apparatus. <i>Cytoskeleton</i> , 2010, 67, 13-22.   | 4.4 | 19        |
| 1644 | Prp45 affects Prp22 partition in spliceosomal complexes and splicing efficiency of non-consensus substrates. <i>Journal of Cellular Biochemistry</i> , 2009, 106, 139-151.  | 1.2 | 31        |
| 1645 | A genetic screen in <i>Saccharomyces cerevisiae</i> identifies new genes that interact with mex67-5, a temperature-sensitive allele of the gene encoding the mRNA export receptor. <i>Molecular Genetics and Genomics</i> , 2009, 281, 125-134. | 1.0 | 15        |
| 1646 | Loss of mitochondrial DNA under genotoxic stress conditions in the absence of the yeast DNA helicase Pif1p occurs independently of the DNA helicase Rrm3p. <i>Molecular Genetics and Genomics</i> , 2009, 281, 635-645.                         | 1.0 | 18        |
| 1647 | Cyk3 acts in actomyosin ring independent cytokinesis by recruiting Inn1 to the yeast bud neck. <i>Molecular Genetics and Genomics</i> , 2009, 282, 437-451.   | 1.0 | 44        |
| 1648 | Ubiquitin ligase Rsp5p is involved in the gene expression changes during nutrient limitation in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2009, 26, 1-15.  | 0.8 | 12        |
| 1649 | Plasmids with E2 epitope tags: tagging modules for N-terminal and C-terminal PCR-based gene targeting in both budding and fission yeast, and inducible expression vectors for fission yeast. <i>Yeast</i> , 2009, 26, 55-66.                    | 0.8 | 4         |
| 1650 | Additional cassettes for epitope and fluorescent fusion proteins in <i>Candida albicans</i> . <i>Yeast</i> , 2009, 26, 399-406.   | 0.8 | 65        |
| 1651 | Atg17 recruits Atg9 to organize the pre-autophagosomal structure. <i>Genes To Cells</i> , 2009, 14, 525-538.  | 0.5 | 137       |
| 1652 | Ctf4 coordinates the progression of helicase and DNA polymerase $\delta$ . <i>Genes To Cells</i> , 2009, 14, 807-820.   | 0.5 | 82        |
| 1653 | Linear ubiquitin fusion to Rps31 and its subsequent cleavage are required for the efficient production and functional integrity of 40S ribosomal subunits. <i>Molecular Microbiology</i> , 2009, 72, 69-84.                                     | 1.2 | 61        |
| 1654 | Elongator function depends on antagonistic regulation by casein kinase Hrr25 and protein phosphatase Sit4. <i>Molecular Microbiology</i> , 2009, 73, 869-881.   | 1.2 | 47        |
| 1655 | Lte1, Cdc14 and MEN-controlled Cdk inactivation in yeast coordinate rDNA decompaction with late telophase progression. <i>EMBO Journal</i> , 2009, 28, 1562-1575.   | 3.5 | 19        |
| 1656 | Cotranslational assembly of the yeast SET1C histone methyltransferase complex. <i>EMBO Journal</i> , 2009, 28, 2959-2970.   | 3.5 | 73        |
| 1657 | Syp1 is a conserved endocytic adaptor that contains domains involved in cargo selection and membrane tubulation. <i>EMBO Journal</i> , 2009, 28, 3103-3116.   | 3.5 | 138       |
| 1658 | The MRX complex stabilizes the replisome independently of the S phase checkpoint during replication stress. <i>EMBO Journal</i> , 2009, 28, 1142-1156.  | 3.5 | 79        |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 1659 | RAD6 and RAD18 are RAD5-pathway-dependent tolerance to chronic low-dose ultraviolet light. <i>Nature</i> , 2009, 457, 612-615.  | 13.7 | 73        |
| 1660 | Messenger RNA targeting to endoplasmic reticulum stress signalling sites. <i>Nature</i> , 2009, 457, 736-740.   | 13.7 | 297       |
| 1661 | Histone levels are regulated by phosphorylation and ubiquitylation-dependent proteolysis. <i>Nature Cell Biology</i> , 2009, 11, 925-933.   | 4.6  | 187       |
| 1662 | A Dam1-based artificial kinetochore is sufficient to promote chromosome segregation in budding yeast. <i>Nature Cell Biology</i> , 2009, 11, 1109-1115.   | 4.6  | 48        |
| 1663 | Recruiting a microtubule-binding complex to DNA directs chromosome segregation in budding yeast. <i>Nature Cell Biology</i> , 2009, 11, 1116-1120.  | 4.6  | 54        |
| 1664 | The establishment of gene silencing at single-cell resolution. <i>Nature Genetics</i> , 2009, 41, 800-806.  | 9.4  | 71        |
| 1665 | A Flp-nick system to study repair of a single protein-bound nick in vivo. <i>Nature Methods</i> , 2009, 6, 753-757.   | 9.0  | 52        |
| 1666 | Telomerase from yeast <i>Saccharomyces cerevisiae</i> is active in vitro as a monomer. <i>Biochemistry (Moscow)</i> , 2009, 74, 749-755.  | 0.7  | 5         |
| 1667 | Search for genes influencing the maintenance of the [ISP +] prion-like antisuppressor determinant in yeast with the use of an insertion gene library. <i>Molecular Biology</i> , 2009, 43, 360-366. | 0.4  | 4         |
| 1668 | Distinct promoter dynamics of the basal transcription factor TBP across the yeast genome. <i>Nature Structural and Molecular Biology</i> , 2009, 16, 1043-1048.                                     | 3.6  | 79        |
| 1669 | Negative Regulation of the Actin-Regulating Kinase Prk1p by Patch Localization-Induced Autophosphorylation. <i>Traffic</i> , 2009, 10, 35-41.   | 1.3  | 6         |
| 1670 | Identification of the Novel TRAPP Associated Protein Tca17. <i>Traffic</i> , 2009, 10, 713-723.   | 1.3  | 44        |
| 1671 | Modulation of sterol homeostasis by the Cdc42p effectors Cla4p and Ste20p in the yeast <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 2009, 276, 7253-7264.                                | 2.2  | 11        |
| 1672 | <i>ETP1/YHL010C</i> is a novel gene needed for the adaptation of <i>Saccharomyces cerevisiae</i> to ethanol. <i>FEMS Yeast Research</i> , 2009, 9, 372-380.   | 1.1  | 26        |
| 1673 | Effects of amiodarone on $K^{+}$ , internal pH and $Ca^{2+}$ homeostasis in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2009, 9, 832-848.  | 1.1  | 21        |
| 1674 | NQR1 controls lifespan by regulating the promotion of respiratory metabolism in yeast. <i>Aging Cell</i> , 2009, 8, 140-151.  | 3.0  | 37        |
| 1675 | Calorie restriction reduces rDNA recombination independently of rDNA silencing. <i>Aging Cell</i> , 2009, 8, 624-632.   | 3.0  | 43        |
| 1676 | Tools for Fungal Proteomics: Multifunctional Neurospora Vectors for Gene Replacement, Protein Expression and Protein Purification. <i>Genetics</i> , 2009, 182, 11-23.                              | 1.2  | 114       |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 1677 | Specific roles for the Ccr4-Not complex subunits in expression of the genome. <i>Rna</i> , 2009, 15, 377-383.   | 1.6  | 56        |
| 1678 | Set2-Dependent K36 Methylation Is Regulated by Novel Intratail Interactions within H3. <i>Molecular and Cellular Biology</i> , 2009, 29, 6413-6426.   | 1.1  | 16        |
| 1679 | Structure and Functional Studies of the CS Domain of the Essential H/ACA Ribonucleoparticle Assembly Protein SHQ1. <i>Journal of Biological Chemistry</i> , 2009, 284, 1906-1916.                               | 1.6  | 26        |
| 1680 | Expression of Bacterial Rho Factor in Yeast Identifies New Factors Involved in the Functional Interplay between Transcription and mRNP Biogenesis. <i>Molecular and Cellular Biology</i> , 2009, 29, 4033-4044. | 1.1  | 17        |
| 1681 | Functional Expression of a Bacterial Xylose Isomerase in <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 2009, 75, 2304-2311.   | 1.4  | 241       |
| 1682 | Effects of Age on Meiosis in Budding Yeast. <i>Developmental Cell</i> , 2009, 16, 844-855.  | 3.1  | 22        |
| 1683 | Atg32 Is a Mitochondrial Protein that Confers Selectivity during Mitophagy. <i>Developmental Cell</i> , 2009, 17, 98-109.   | 3.1  | 709       |
| 1684 | A Yeast Killer Toxin Screen Provides Insights into A/B Toxin Entry, Trafficking, and Killing Mechanisms. <i>Developmental Cell</i> , 2009, 17, 552-560.   | 3.1  | 68        |
| 1685 | A plasmid collection for PCR-based gene targeting in the filamentous ascomycete <i>Ashbya gossypii</i> . <i>Fungal Genetics and Biology</i> , 2009, 46, 595-603.  | 0.9  | 27        |
| 1686 | Mechanochemical Removal of Ribosome Biogenesis Factors from Nascent 60S Ribosomal Subunits. <i>Cell</i> , 2009, 138, 911-922.   | 13.5 | 141       |
| 1687 | Recombinational Repair within Heterochromatin Requires ATP-Dependent Chromatin Remodeling. <i>Cell</i> , 2009, 138, 1109-1121.  | 13.5 | 73        |
| 1688 | Determination of the core promoter regions of the <i>Saccharomyces cerevisiae</i> RPS3 gene. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2009, 1789, 741-750.                           | 0.9  | 6         |
| 1689 | Mobilization of steryl esters from lipid particles of the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2009, 1791, 118-124.            | 1.2  | 32        |
| 1690 | Lap3 is a selective target of autophagy in yeast, <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2009, 378, 551-557.  | 1.0  | 37        |
| 1691 | SBF transcription factor complex positively regulates UV mutagenesis in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2009, 379, 1009-1014.                    | 1.0  | 4         |
| 1692 | A phosphorylation-independent role for the yeast cyclin-dependent kinase activating kinase Cak1. <i>Gene</i> , 2009, 447, 97-105.   | 1.0  | 1         |
| 1693 | Multiple Defects in the Respiratory Chain Lead to the Repression of Genes Encoding Components of the Respiratory Chain and TCA Cycle Enzymes. <i>Journal of Molecular Biology</i> , 2009, 387, 1081-1091.       | 2.0  | 6         |
| 1694 | Functional Analysis of Yeast <i>bcs1</i> Mutants Highlights the Role of Bcs1p-Specific Amino Acids in the AAA Domain. <i>Journal of Molecular Biology</i> , 2009, 388, 252-261.                                 | 2.0  | 25        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1695 | Stimulation of mRNA Export by an F-box Protein, Mdm30p, in Vivo. <i>Journal of Molecular Biology</i> , 2009, 389, 238-247.  | 2.0 | 16        |
| 1696 | The Shuttling Protein Npl3 Promotes Translation Termination Accuracy in <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 2009, 394, 410-422.   | 2.0 | 23        |
| 1697 | Sus1, Cdc31, and the Sac3 CID Region Form a Conserved Interaction Platform that Promotes Nuclear Pore Association and mRNA Export. <i>Molecular Cell</i> , 2009, 33, 727-737.   | 4.5 | 128       |
| 1698 | BAX Inhibitor-1 Is a Negative Regulator of the ER Stress Sensor IRE1 $\pm$ . <i>Molecular Cell</i> , 2009, 33, 679-691.   | 4.5 | 281       |
| 1699 | Prp43 Bound at Different Sites on the Pre-rRNA Performs Distinct Functions in Ribosome Synthesis. <i>Molecular Cell</i> , 2009, 36, 583-592.  | 4.5 | 152       |
| 1700 | Usa1 Functions as a Scaffold of the HRD-Ubiquitin Ligase. <i>Molecular Cell</i> , 2009, 36, 782-793.  | 4.5 | 114       |
| 1701 | Rrp17p Is a Eukaryotic Exonuclease Required for 5' End Processing of Pre-60S Ribosomal RNA. <i>Molecular Cell</i> , 2009, 36, 768-781.  | 4.5 | 83        |
| 1702 | Monitoring mitophagy in yeast: The Om45-GFP processing assay. <i>Autophagy</i> , 2009, 5, 1186-1189.  | 4.3 | 81        |
| 1703 | Analysis of the dual function of the ESCRT-III protein Snf7 in endocytic trafficking and in gene expression. <i>Biochemical Journal</i> , 2009, 424, 89-97.   | 1.7 | 24        |
| 1704 | Identification of a conserved F-box protein 6 interactor essential for endocytosis and cytokinesis in fission yeast. <i>Biochemical Journal</i> , 2009, 420, 169-180.   | 1.7 | 12        |
| 1705 | Functional domains of yeast hexokinase 2. <i>Biochemical Journal</i> , 2010, 432, 181-190.  | 1.7 | 51        |
| 1706 | The role of the Rdh54 protein in regulation of DNA repair in yeast <i>Saccharomyces cerevisiae</i> . <i>Russian Journal of Genetics</i> , 2010, 46, 170-177.  | 0.2 | 2         |
| 1707 | The yeast dynamin-like protein Vps1:vps1 mutations perturb the internalization and the motility of endocytic vesicles and endosomes via disorganization of the actin cytoskeleton. <i>European Journal of Cell Biology</i> , 2010, 89, 499-508. | 1.6 | 40        |
| 1708 | Interaction between Sec7p and Pik1p: The first clue for the regulation of a coincidence detection signal. <i>European Journal of Cell Biology</i> , 2010, 89, 575-583.  | 1.6 | 16        |
| 1709 | Association of the yeast DNA helicase Pif1p with mitochondrial membranes and mitochondrial DNA. <i>European Journal of Cell Biology</i> , 2010, 89, 742-747.  | 1.6 | 9         |
| 1710 | Proteolytic Instability and the Action of Nonclassical Transcriptional Activators. <i>Current Biology</i> , 2010, 20, 868-871.  | 1.8 | 22        |
| 1711 | Reconstitution and Protein Composition Analysis of Endocytic Actin Patches. <i>Current Biology</i> , 2010, 20, 1890-1899.   | 1.8 | 59        |
| 1712 | The Rtt109 histone acetyltransferase facilitates error-free replication to prevent CAG/CTG repeat contractions. <i>DNA Repair</i> , 2010, 9, 414-420.   | 1.3 | 22        |



| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1713 | Mec1/Tel1-dependent phosphorylation of Slx4 stimulates Rad1-dependent cleavage of non-homologous DNA tails. <i>DNA Repair</i> , 2010, 9, 718-726.   | 1.3 | 54        |
| 1714 | Regulation of tolerance to DNA alkylating damage by Dot1 and Rad53 in <i>Saccharomyces cerevisiae</i> . <i>DNA Repair</i> , 2010, 9, 1038-1049.   | 1.3 | 35        |
| 1715 | Swa2-dependent clathrin dynamics is critical for Flo11p processing and Mat formation in the yeast <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 2010, 584, 1149-1155.                     | 1.3 | 8         |
| 1716 | Cln3 blocks IME1 transcription and the Ime1-Ume6 interaction to cause the sporulation incompetence in a sake yeast, Kyokai no. 7. <i>Journal of Bioscience and Bioengineering</i> , 2010, 110, 1-7. | 1.1 | 12        |
| 1717 | The Rsp5 ubiquitin ligase and the AAA-ATPase Cdc48 control the ubiquitin-mediated degradation of the COPII component Sec23. <i>Experimental Cell Research</i> , 2010, 316, 3351-3357.               | 1.2 | 17        |
| 1718 | Involvement of Mitochondrial Ferredoxin and Para-Aminobenzoic Acid in Yeast Coenzyme Q Biosynthesis. <i>Chemistry and Biology</i> , 2010, 17, 449-459.  | 6.2 | 100       |
| 1719 | Traffic of a Viral Movement Protein Complex to the Highly Curved Tubules of the Cortical Endoplasmic Reticulum. <i>Traffic</i> , 2010, 11, 912-930.   | 1.3 | 39        |
| 1720 | Quantitative Analysis of Endocytosis with Cytoplasmic pHluorin Chimeras. <i>Traffic</i> , 2010, 11, 1141-1150.  | 1.3 | 53        |
| 1721 | Targeting and membrane insertion into the endoplasmic reticulum membrane of <i>Saccharomyces cerevisiae</i> essential protein Rot1. <i>FEMS Yeast Research</i> , 2010, 10, 639-647.                 | 1.1 | 5         |
| 1722 | The budding yeast protein Sum1 functions independently of its binding partners Hst1 and Sir2 histone deacetylases to regulate microtubule assembly. <i>FEMS Yeast Research</i> , 2010, 10, 660-673. | 1.1 | 5         |
| 1723 | Suppression of Ycf1p function by Cka1p-dependent phosphorylation is attenuated in response to salt stress. <i>FEMS Yeast Research</i> , 2010, 10, 839-857.  | 1.1 | 15        |
| 1724 | A mutational analysis reveals new functional interactions between domains of the Oxa1 protein in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2010, 75, 474-488.               | 1.2 | 6         |
| 1725 | A <i>Candida albicans</i> cell wall-linked protein promotes invasive filamentation into semi-solid medium. <i>Molecular Microbiology</i> , 2010, 76, 733-748.                                       | 1.2 | 36        |
| 1726 | Molecular architecture of the TRAPP-II complex and implications for vesicle tethering. <i>Nature Structural and Molecular Biology</i> , 2010, 17, 1298-1304.  | 3.6 | 70        |
| 1727 | Molecular organization of the COG vesicle tethering complex. <i>Nature Structural and Molecular Biology</i> , 2010, 17, 1292-1297.  | 3.6 | 79        |
| 1728 | Reduced Rif2 and lack of Mec1 target short telomeres for elongation rather than double-strand break repair. <i>Nature Structural and Molecular Biology</i> , 2010, 17, 1438-1445.                   | 3.6 | 65        |
| 1729 | Inactivation of the Sas2 histone acetyltransferase delays senescence driven by telomere dysfunction. <i>EMBO Journal</i> , 2010, 29, 158-170.   | 3.5 | 45        |
| 1730 | Mdm35p imports Ups proteins into the mitochondrial intermembrane space by functional complex formation. <i>EMBO Journal</i> , 2010, 29, 2875-2887.  | 3.5 | 68        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 1731 | Intercalation of a new tier of transcription regulation into an ancient circuit. <i>Nature</i> , 2010, 468, 959-963.   | 13.7 | 74        |
| 1732 | Tension directly stabilizes reconstituted kinetochore-microtubule attachments. <i>Nature</i> , 2010, 468, 576-579.   | 13.7 | 408       |
| 1733 | Chemical genetics screen for enhancers of rapamycin identifies a specific inhibitor of an SCF family E3 ubiquitin ligase. <i>Nature Biotechnology</i> , 2010, 28, 738-742.                                     | 9.4  | 132       |
| 1734 | DNA zip codes control an ancient mechanism for gene targeting to the nuclear periphery. <i>Nature Cell Biology</i> , 2010, 12, 111-118.  | 4.6  | 170       |
| 1735 | The N-end rule pathway is mediated by a complex of the RING-type Ubr1 and HECT-type Ufd4 ubiquitin ligases. <i>Nature Cell Biology</i> , 2010, 12, 1177-1185.  | 4.6  | 121       |
| 1736 | Boric Acid Disturbs Cell Wall Synthesis in <i>Saccharomyces cerevisiae</i> . <i>International Journal of Microbiology</i> , 2010, 2010, 1-10.  | 0.9  | 25        |
| 1737 | Functional Overlap between eIF4G Isoforms in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2010, 5, e9114.   | 1.1  | 50        |
| 1738 | A Local Role for the Small Ribosomal Subunit Primary Binder rpS5 in Final 18S rRNA Processing in Yeast. <i>PLoS ONE</i> , 2010, 5, e10194.   | 1.1  | 22        |
| 1739 | Temporal Characterization of Homology-Independent Centromere Coupling in Meiotic Prophase. <i>PLoS ONE</i> , 2010, 5, e10336.  | 1.1  | 38        |
| 1740 | Single-Molecule Atomic Force Microscopy Reveals Clustering of the Yeast Plasma-Membrane Sensor Wsc1. <i>PLoS ONE</i> , 2010, 5, e11104.  | 1.1  | 91        |
| 1741 | The Roles of the <i>Saccharomyces cerevisiae</i> RecQ Helicase SGS1 in Meiotic Genome Surveillance. <i>PLoS ONE</i> , 2010, 5, e15380.   | 1.1  | 10        |
| 1742 | Expanded Roles of the Origin Recognition Complex in the Architecture and Function of Silenced Chromatin in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2010, 30, 626-639.        | 1.1  | 24        |
| 1743 | The AAA-ATPase Cdc48 and cofactor Shp1 promote chromosome bi-orientation by balancing Aurora B activity. <i>Journal of Cell Science</i> , 2010, 123, 2025-2034.  | 1.2  | 39        |
| 1744 | Double-Strand Break Repair Pathways Protect against CAG/CTG Repeat Expansions, Contractions and Repeat-Mediated Chromosomal Fragility in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2010, 184, 65-77. | 1.2  | 59        |
| 1745 | Exit from the Golgi Is Required for the Expansion of the Autophagosomal Phagophore in Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2010, 21, 2270-2284.                      | 0.9  | 160       |
| 1746 | Post-Golgi Sec Proteins Are Required for Autophagy in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2010, 21, 2257-2269.  | 0.9  | 159       |
| 1747 | Release of SF3 from the intron branchpoint activates the first step of pre-mRNA splicing. <i>Rna</i> , 2010, 16, 516-528.  | 1.6  | 142       |
| 1748 | Cdc14p resets the competency of replication licensing by dephosphorylating multiple initiation proteins during mitotic exit in budding yeast. <i>Journal of Cell Science</i> , 2010, 123, 3933-3943.           | 1.2  | 32        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1749 | Comprehensive Screening of Human Genes with Inhibitory Effects on Yeast Growth and Validation of a Yeast Cell-Based System for Screening Chemicals. <i>Journal of Biomolecular Screening</i> , 2010, 15, 368-378.                     | 2.6 | 15        |
| 1750 | Abasic Sites in the Transcribed Strand of Yeast DNA Are Removed by Transcription-Coupled Nucleotide Excision Repair. <i>Molecular and Cellular Biology</i> , 2010, 30, 3206-3215.   | 1.1 | 58        |
| 1751 | A Guanine Nucleotide Exchange Factor Is a Component of the Meiotic Spindle Pole Body in <i>Schizosaccharomyces pombe</i> . <i>Molecular Biology of the Cell</i> , 2010, 21, 1272-1281.  | 0.9 | 12        |
| 1752 | The $\hat{1}^2$ /Gcd7 Subunit of Eukaryotic Translation Initiation Factor 2B (eIF2B), a Guanine Nucleotide Exchange Factor, Is Crucial for Binding eIF2 <i>in Vivo</i> . <i>Molecular and Cellular Biology</i> , 2010, 30, 5218-5233. | 1.1 | 35        |
| 1753 | Structure and Function of the PLAA/Ufd3-p97/Cdc48 Complex. <i>Journal of Biological Chemistry</i> , 2010, 285, 365-372.   | 1.6 | 30        |
| 1754 | Rad26p, a transcription-coupled repair factor, is recruited to the site of DNA lesion in an elongating RNA polymerase II-dependent manner <i>in vivo</i> . <i>Nucleic Acids Research</i> , 2010, 38, 1461-1477.                       | 6.5 | 50        |
| 1755 | Identification and Analysis of the Interaction between Edc3 and Dcp2 in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2010, 30, 1446-1456.  | 1.1 | 57        |
| 1756 | Yeast arginine methyltransferase Hmt1p regulates transcription elongation and termination by methylating Npl3p. <i>Nucleic Acids Research</i> , 2010, 38, 2217-2228.  | 6.5 | 31        |
| 1757 | Npr2, Yeast Homolog of the Human Tumor Suppressor <i>NPRL2</i> , Is a Target of Grr1 Required for Adaptation to Growth on Diverse Nitrogen Sources. <i>Eukaryotic Cell</i> , 2010, 9, 592-601.  | 3.4 | 23        |
| 1758 | Translational repression by PUF proteins <i>in vitro</i> . <i>Rna</i> , 2010, 16, 1217-1225.  | 1.6 | 38        |
| 1759 | The Na <sup>+</sup> /H <sup>+</sup> exchanger Nhx1p regulates the initiation of <i>Saccharomyces cerevisiae</i> vacuole fusion. <i>Journal of Cell Science</i> , 2010, 123, 3266-3275.  | 1.2 | 44        |
| 1760 | Mutation of High-Affinity Methionine Permease Contributes to Selenomethionyl Protein Production in <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 2010, 76, 6351-6359.                             | 1.4 | 12        |
| 1761 | Dicentric breakage at telomere fusions. <i>Genes and Development</i> , 2010, 24, 720-733.   | 2.7 | 39        |
| 1762 | Asf1 Can Promote Trimethylation of H3 K36 by Set2. <i>Molecular and Cellular Biology</i> , 2010, 30, 1116-1129.   | 1.1 | 32        |
| 1763 | Vesicle Docking to the Spindle Pole Body Is Necessary to Recruit the Exocyst During Membrane Formation in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2010, 21, 3693-3707.                               | 0.9 | 19        |
| 1764 | Ecm1 is a new pre-ribosomal factor involved in pre-60S particle export. <i>Rna</i> , 2010, 16, 1007-1017.   | 1.6 | 38        |
| 1765 | Novel <i>trans</i> -Tail Regulation of H2B Ubiquitylation and H3K4 Methylation by the N Terminus of Histone H2A. <i>Molecular and Cellular Biology</i> , 2010, 30, 3635-3645.   | 1.1 | 43        |
| 1766 | Formation of the Redox Cofactor Centers during Cox1 Maturation in Yeast Cytochrome Oxidase. <i>Molecular and Cellular Biology</i> , 2010, 30, 1004-1017.  | 1.1 | 88        |



| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1785 | The RNA Polymerase-associated Factor 1 Complex (Paf1C) Directly Increases the Elongation Rate of RNA Polymerase I and Is Required for Efficient Regulation of rRNA Synthesis. <i>Journal of Biological Chemistry</i> , 2010, 285, 14152-14159. | 1.6 | 50        |
| 1786 | Vesicle trafficking maintains nuclear shape in <i>Saccharomyces cerevisiae</i> during membrane proliferation. <i>Journal of Cell Biology</i> , 2010, 191, 1079-1088.   | 2.3 | 41        |
| 1787 | Coa3 and Cox14 are essential for negative feedback regulation of <i>COX1</i> translation in mitochondria. <i>Journal of Cell Biology</i> , 2010, 191, 141-154.   | 2.3 | 99        |
| 1788 | Regulators of Vps4 ATPase Activity at Endosomes Differentially Influence the Size and Rate of Formation of Intraluminal Vesicles. <i>Molecular Biology of the Cell</i> , 2010, 21, 1023-1032.  | 0.9 | 77        |
| 1789 | Dissecting BAR Domain Function in the Yeast Amphiphysins Rvs161 and Rvs167 during Endocytosis. <i>Molecular Biology of the Cell</i> , 2010, 21, 3054-3069.   | 0.9 | 73        |
| 1790 | A role for the dynamin-like protein Vps1 during endocytosis in yeast. <i>Journal of Cell Science</i> , 2010, 123, 3496-3506.   | 1.2 | 96        |
| 1791 | Nonsense-mediated mRNA decay maintains translational fidelity by limiting magnesium uptake. <i>Genes and Development</i> , 2010, 24, 1491-1495.  | 2.7 | 47        |
| 1792 | Peroxisomal Membrane Proteins Insert into the Endoplasmic Reticulum. <i>Molecular Biology of the Cell</i> , 2010, 21, 2057-2065.   | 0.9 | 154       |
| 1793 | Requirements for Transitional Endoplasmic Reticulum Site Structure and Function in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2010, 21, 1530-1545.   | 0.9 | 54        |
| 1794 | Multiple Signals Converge on a Differentiation MAPK Pathway. <i>PLoS Genetics</i> , 2010, 6, e1000883.   | 1.5 | 54        |
| 1795 | Transcriptional plasticity through differential assembly of a multiprotein activation complex. <i>Nucleic Acids Research</i> , 2010, 38, 4998-5014.  | 6.5 | 19        |
| 1796 | Meiotic Cohesin Promotes Pairing of Nonhomologous Centromeres in Early Meiotic Prophase. <i>Molecular Biology of the Cell</i> , 2010, 21, 1799-1809.   | 0.9 | 34        |
| 1797 | Stepwise Assembly of Dimeric F <sub>1</sub> F <sub>o</sub> -ATP Synthase in Mitochondria Involves the Small F <sub>o</sub> -Subunits k and i. <i>Molecular Biology of the Cell</i> , 2010, 21, 1494-1504.                                      | 0.9 | 40        |
| 1798 | Uga3 and Uga35/Dal81 Transcription Factors Regulate UGA4 Transcription in Response to <sup>13</sup> C-Aminobutyric Acid and Leucine. <i>Eukaryotic Cell</i> , 2010, 9, 1262-1271.  | 3.4 | 18        |
| 1799 | The Sua5 Protein Is Essential for Normal Translational Regulation in Yeast. <i>Molecular and Cellular Biology</i> , 2010, 30, 354-363.   | 1.1 | 65        |
| 1800 | Cdk Phosphorylation of a Nucleoporin Controls Localization of Active Genes through the Cell Cycle. <i>Molecular Biology of the Cell</i> , 2010, 21, 3421-3432.   | 0.9 | 38        |
| 1801 | Slk19p of <i>Saccharomyces cerevisiae</i> Regulates Anaphase Spindle Dynamics Through Two Independent Mechanisms. <i>Genetics</i> , 2010, 186, 1247-1260.  | 1.2 | 10        |
| 1802 | Activation of a Poised RNAPII-Dependent Promoter Requires Both SAGA and Mediator. <i>Genetics</i> , 2010, 184, 659-672.  | 1.2 | 21        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1803 | Functional Genomics Analysis of the <i>Saccharomyces cerevisiae</i> Iron Responsive Transcription Factor Aft1 Reveals Iron-Independent Functions. <i>Genetics</i> , 2010, 185, 1111-1128.  | 1.2 | 34        |
| 1804 | The Transcription Activity of Gis1 Is Negatively Modulated by Proteasome-mediated Limited Proteolysis. <i>Journal of Biological Chemistry</i> , 2010, 285, 6465-6476.  | 1.6 | 22        |
| 1805 | Phosphorylation of Lte1 by Cdk prevents polarized growth during mitotic arrest in <i>S. cerevisiae</i> . <i>Journal of Cell Biology</i> , 2010, 191, 1097-1112.  | 2.3 | 24        |
| 1806 | Mitotic Expression of Spo13 Alters M-Phase Progression and Nucleolar Localization of Cdc14 in Budding Yeast. <i>Genetics</i> , 2010, 185, 841-854.   | 1.2 | 7         |
| 1807 | Biphasic targeting and cleavage furrow ingression directed by the tail of a myosin II. <i>Journal of Cell Biology</i> , 2010, 191, 1333-1350.  | 2.3 | 99        |
| 1808 | Protein Arginine Methylation Facilitates Cotranscriptional Recruitment of Pre-mRNA Splicing Factors. <i>Molecular and Cellular Biology</i> , 2010, 30, 5245-5256.  | 1.1 | 43        |
| 1809 | Multiple Myo4 motors enhance ASH1 mRNA transport in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 2010, 189, 755-767.   | 2.3 | 45        |
| 1810 | Genome-wide localization analysis of a complete set of Tafs reveals a specific effect of the taf1 mutation on Taf2 occupancy and provides indirect evidence for different TFIID conformations at different promoters. <i>Nucleic Acids Research</i> , 2010, 38, 1805-1820. | 6.5 | 14        |
| 1811 | Interruption of Inositol Sphingolipid Synthesis Triggers Stt4p-dependent Protein Kinase C Signaling*. <i>Journal of Biological Chemistry</i> , 2010, 285, 41947-41960.   | 1.6 | 39        |
| 1812 | Selective Transport of $\alpha$ -Mannosidase by Autophagic Pathways. <i>Journal of Biological Chemistry</i> , 2010, 285, 30019-30025.  | 1.6 | 103       |
| 1813 | Yet1p and Yet3p, the Yeast Homologs of BAP29 and BAP31, Interact with the Endoplasmic Reticulum Translocation Apparatus and Are Required for Inositol Prototrophy. <i>Journal of Biological Chemistry</i> , 2010, 285, 18252-18261.  | 1.6 | 28        |
| 1814 | Directed Evolution Reveals Hidden Properties of VMAT, a Neurotransmitter Transporter. <i>Journal of Biological Chemistry</i> , 2010, 285, 5076-5084.   | 1.6 | 24        |
| 1815 | Late Phase of the Endoplasmic Reticulum Stress Response Pathway Is Regulated by Hog1 MAP Kinase. <i>Journal of Biological Chemistry</i> , 2010, 285, 17545-17555.  | 1.6 | 54        |
| 1816 | Replication Fork Arrest and rDNA Silencing Are Two Independent and Separable Functions of the Replication Terminator Protein Fob1 of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2010, 285, 12612-12619.                                    | 1.6 | 23        |
| 1817 | Vacuolar Cation/H <sup>+</sup> Antiporters of <i>Saccharomyces cerevisiae</i> *. <i>Journal of Biological Chemistry</i> , 2010, 285, 33914-33922.  | 1.6 | 39        |
| 1818 | Phosphatidylserine Is Involved in the Ferrichrome-induced Plasma Membrane Trafficking of Arn1 in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2010, 285, 39564-39573.  | 1.6 | 13        |
| 1819 | Dbf4 Regulates the Cdc5 Polo-like Kinase through a Distinct Non-canonical Binding Interaction. <i>Journal of Biological Chemistry</i> , 2010, 285, 41244-41254.  | 1.6 | 33        |
| 1820 | The p40/ARPC1 Subunit of Arp2/3 Complex Performs Multiple Essential Roles in WASp-regulated Actin Nucleation. <i>Journal of Biological Chemistry</i> , 2010, 285, 8481-8491.   | 1.6 | 24        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1821 | A Profile of Differentially Abundant Proteins at the Yeast Cell Periphery during Pseudohyphal Growth. <i>Journal of Biological Chemistry</i> , 2010, 285, 15476-15488.   | 1.6 | 32        |
| 1822 | Activator Gcn4 Employs Multiple Segments of Med15/Gal11, Including the KIX Domain, to Recruit Mediator to Target Genes in Vivo. <i>Journal of Biological Chemistry</i> , 2010, 285, 2438-2455.   | 1.6 | 68        |
| 1823 | The DEAD-box RNA helicase-like Utp25 is an SSU processome component. <i>Rna</i> , 2010, 16, 2156-2169.   | 1.6 | 39        |
| 1824 | A dual function for chaperones SSB and RAC and the NAC nascent polypeptide-associated complex on ribosomes. <i>Journal of Cell Biology</i> , 2010, 189, 57-68.   | 2.3 | 189       |
| 1825 | The RasGAP Proteins Ira2 and Neurofibromin Are Negatively Regulated by Gpb1 in Yeast and ETEA in Humans. <i>Molecular and Cellular Biology</i> , 2010, 30, 2264-2279.  | 1.1 | 56        |
| 1826 | Compartment-specific Synthesis of Phosphatidylethanolamine Is Required for Normal Heavy Metal Resistance. <i>Molecular Biology of the Cell</i> , 2010, 21, 443-455.  | 0.9 | 71        |
| 1827 | Identification of Ypk1 as a Novel Selective Substrate for Nitrogen Starvation-triggered Proteolysis Requiring Autophagy System and Endosomal Sorting Complex Required for Transport (ESCRT) Machinery Components. <i>Journal of Biological Chemistry</i> , 2010, 285, 36984-36994. | 1.6 | 14        |
| 1828 | Splicing of Yeast $\alpha 5^{\text{th}}$ Group I Intron Requires SUV3 to Recycle MRS1 via Mitochondrial Degradosome-promoted Decay of Excised Intron Ribonucleoprotein (RNP). <i>Journal of Biological Chemistry</i> , 2010, 285, 8585-8594.                                       | 1.6 | 22        |
| 1829 | Sss1p Is Required to Complete Protein Translocon Activation. <i>Journal of Biological Chemistry</i> , 2010, 285, 32671-32677.  | 1.6 | 7         |
| 1830 | Evolutionarily Conserved Function of <i>RRP36</i> in Early Cleavages of the Pre-rRNA and Production of the 40S Ribosomal Subunit. <i>Molecular and Cellular Biology</i> , 2010, 30, 1130-1144.   | 1.1 | 25        |
| 1831 | The conserved oligomeric Golgi complex is involved in double-membrane vesicle formation during autophagy. <i>Journal of Cell Biology</i> , 2010, 188, 101-114.   | 2.3 | 179       |
| 1832 | Degradation of the <i>Saccharomyces cerevisiae</i> Mating-Type Regulator $\hat{1}$ : Genetic Dissection of <i>Cis</i> -determinants and <i>Trans</i> -acting Pathways. <i>Genetics</i> , 2010, 185, 497-511.   | 1.2 | 17        |
| 1833 | Membrane assembly modulates the stability of the meiotic spindle-pole body. <i>Journal of Cell Science</i> , 2010, 123, 2481-2490.   | 1.2 | 9         |
| 1834 | Differential Roles of Transcriptional Mediator Subunits in Regulation of Multidrug Resistance Gene Expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2010, 21, 2469-2482.   | 0.9 | 49        |
| 1835 | Changes in the Nuclear Envelope Environment Affect Spindle Pole Body Duplication in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2010, 186, 867-883.  | 1.2 | 40        |
| 1836 | Actin-Related Protein Arp6 Influences H2A.Z-Dependent and -Independent Gene Expression and Links Ribosomal Protein Genes to Nuclear Pores. <i>PLoS Genetics</i> , 2010, 6, e1000910.   | 1.5 | 52        |
| 1837 | Trehalose Is a Key Determinant of the Quiescent Metabolic State That Fuels Cell Cycle Progression upon Return to Growth. <i>Molecular Biology of the Cell</i> , 2010, 21, 1982-1990.   | 0.9 | 112       |
| 1838 | Co-Evolution of Transcriptional Silencing Proteins and the DNA Elements Specifying Their Assembly. <i>PLoS Biology</i> , 2010, 8, e1000550.  | 2.6 | 36        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1839 | Elevated Levels of the Polo Kinase Cdc5 Override the Mec1/ATR Checkpoint in Budding Yeast by Acting at Different Steps of the Signaling Pathway. <i>PLoS Genetics</i> , 2010, 6, e1000763.                        | 1.5 | 49        |
| 1840 | Free Oligosaccharides to Monitor Glycoprotein Endoplasmic Reticulum-associated Degradation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2010, 285, 12390-12404.                  | 1.6 | 84        |
| 1841 | Roles for H2A.Z and Its Acetylation in GAL1 Transcription and Gene Induction, but Not GAL1-Transcriptional Memory. <i>PLoS Biology</i> , 2010, 8, e1000401.   | 2.6 | 75        |
| 1842 | Dynamics of Rad9 Chromatin Binding and Checkpoint Function Are Mediated by Its Dimerization and Are Cell Cycle-Regulated by CDK1 Activity. <i>PLoS Genetics</i> , 2010, 6, e1001047.                              | 1.5 | 59        |
| 1843 | Mek1 Suppression of Meiotic Double-Strand Break Repair Is Specific to Sister Chromatids, Chromosome Autonomous and Independent of Rec8 Cohesin Complexes. <i>Genetics</i> , 2010, 185, 771-782.                   | 1.2 | 41        |
| 1844 | Mapping of Vps21 and HOPS Binding Sites in Vps8 and Effect of Binding Site Mutants on Endocytic Trafficking. <i>Eukaryotic Cell</i> , 2010, 9, 602-610.   | 3.4 | 18        |
| 1845 | Hsp90 Nuclear Accumulation in Quiescence Is Linked to Chaperone Function and Spore Development in Yeast. <i>Molecular Biology of the Cell</i> , 2010, 21, 63-72.  | 0.9 | 41        |
| 1846 | Initial Polarized Bud Growth by Endocytic Recycling in the Absence of Actin Cable-dependent Vesicle Transport in Yeast. <i>Molecular Biology of the Cell</i> , 2010, 21, 1237-1252.                               | 0.9 | 21        |
| 1847 | Cadmium regulates copper homeostasis by inhibiting the activity of Mac1, a transcriptional activator of the copper regulon, in <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 2010, 431, 257-265. | 1.7 | 17        |
| 1848 | The Lrs4-Csm1 monopolin complex associates with kinetochores during anaphase and is required for accurate chromosome segregation. <i>Cell Cycle</i> , 2010, 9, 3611-3618.   | 1.3 | 19        |
| 1849 | Determining Atg protein stoichiometry at the phagophore assembly site by fluorescence microscopy. <i>Autophagy</i> , 2010, 6, 144-147.  | 4.3 | 13        |
| 1850 | Atg8 regulates vacuolar membrane dynamics in a lipidation-independent manner in <i>Pichia pastoris</i> . <i>Journal of Cell Science</i> , 2010, 123, 4107-4116.   | 1.2 | 52        |
| 1851 | Î±-Arrestins Aly1 and Aly2 Regulate Intracellular Trafficking in Response to Nutrient Signaling. <i>Molecular Biology of the Cell</i> , 2010, 21, 3552-3566.  | 0.9 | 97        |
| 1852 | Dynamic Localization of Fus3 Mitogen-Activated Protein Kinase Is Necessary To Evoke Appropriate Responses and Avoid Cytotoxic Effects. <i>Molecular and Cellular Biology</i> , 2010, 30, 4293-4307.               | 1.1 | 18        |
| 1853 | Coevolution within a transcriptional network by compensatory <i>trans</i> and <i>cis</i> mutations. <i>Genome Research</i> , 2010, 20, 1672-1678.   | 2.4 | 62        |
| 1854 | Strategies for Isolating Constitutively Active and Dominant-Negative Pheromone Receptor Mutants in Yeast. <i>Methods in Enzymology</i> , 2010, 485, 329-348.  | 0.4 | 3         |
| 1855 | Bulk Segregant Analysis by High-Throughput Sequencing Reveals a Novel Xylose Utilization Gene from <i>Saccharomyces cerevisiae</i> . <i>PLoS Genetics</i> , 2010, 6, e1000942.                                    | 1.5 | 177       |
| 1856 | The Final Step in 5.8S rRNA Processing Is Cytoplasmic in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2010, 30, 976-984.   | 1.1 | 64        |



| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 1857 | Spinning-Disk Confocal Microscopy of Yeast. <i>Methods in Enzymology</i> , 2010, 470, 581-602.   | 0.4  | 11        |
| 1858 | ChIP-Seq. <i>Methods in Enzymology</i> , 2010, 470, 77-104.  | 0.4  | 23        |
| 1859 | Sensitive and Convenient Yeast Reporter Assay for High-Throughput Analysis by Using a Secretory Luciferase from <i>Cypridina noctiluca</i> . <i>Analytical Chemistry</i> , 2010, 82, 5768-5776.  | 3.2  | 30        |
| 1860 | Zebrafish ( <i>Danio rerio</i> ) Endomembrane Antiporter Similar to a Yeast Cation/H <sup>+</sup> Transporter Is Required for Neural Crest Development. <i>Biochemistry</i> , 2010, 49, 6557-6566.   | 1.2  | 19        |
| 1861 | Analyzing P-Bodies and Stress Granules in <i>Saccharomyces cerevisiae</i> . <i>Methods in Enzymology</i> , 2010, 470, 619-640.   | 0.4  | 69        |
| 1862 | The AAA-ATPase Rea1 Drives Removal of Biogenesis Factors during Multiple Stages of 60S Ribosome Assembly. <i>Molecular Cell</i> , 2010, 38, 712-721.   | 4.5  | 114       |
| 1863 | Spindle Position Is Coordinated with Cell-Cycle Progression through Establishment of Mitotic Exit-Activating and -Inhibitory Zones. <i>Molecular Cell</i> , 2010, 39, 444-454.   | 4.5  | 44        |
| 1864 | The transcription activation domain of Rpn4 is separate from its degrons. <i>International Journal of Biochemistry and Cell Biology</i> , 2010, 42, 282-286.   | 1.2  | 7         |
| 1865 | Phosphatidylethanolamine synthesized by four different pathways is supplied to the plasma membrane of the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2010, 1801, 480-486. | 1.2  | 33        |
| 1866 | The Cdc48-Ufd1-Npl4 complex is central in ubiquitin-proteasome triggered catabolite degradation of fructose-1,6-bisphosphatase. <i>Biochemical and Biophysical Research Communications</i> , 2010, 394, 335-341.                                     | 1.0  | 16        |
| 1867 | Dissection of the assembly pathway of the proteasome lid in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2010, 396, 1048-1053.   | 1.0  | 63        |
| 1868 | Subcellular localization of the interaction of bipolar landmarks Bud8p and Bud9p with Rax2p in <i>Saccharomyces cerevisiae</i> diploid cells. <i>Biochemical and Biophysical Research Communications</i> , 2010, 399, 525-530.                       | 1.0  | 1         |
| 1869 | A Surveillance Pathway Monitors the Fitness of the Endoplasmic Reticulum to Control Its Inheritance. <i>Cell</i> , 2010, 142, 256-269.   | 13.5 | 95        |
| 1870 | The Monopolin Complex Crosslinks Kinetochores Components to Regulate Chromosome-Microtubule Attachments. <i>Cell</i> , 2010, 142, 556-567.   | 13.5 | 119       |
| 1871 | Identification of Aneuploidy-Tolerating Mutations. <i>Cell</i> , 2010, 143, 71-83.   | 13.5 | 352       |
| 1872 | Retrotranslocation of a Misfolded Luminal ER Protein by the Ubiquitin-Ligase Hrd1p. <i>Cell</i> , 2010, 143, 579-591.  | 13.5 | 262       |
| 1873 | Functional Overlap and Regulatory Links Shape Genetic Interactions between Signaling Pathways. <i>Cell</i> , 2010, 143, 991-1004.  | 13.5 | 123       |
| 1874 | Chromate-induced sulfur starvation and mRNA mistranslation in yeast are linked in a common mechanism of Cr toxicity. <i>Toxicology in Vitro</i> , 2010, 24, 1764-1767.   | 1.1  | 22        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 1875 | Unconventional secretion of Acb1 is mediated by autophagosomes. <i>Journal of Cell Biology</i> , 2010, 188, 527-536.   | 2.3  | 360       |
| 1876 | The Eukaryotic Translation Elongation Factor 1B <sup>3</sup> Has a Non-guanine Nucleotide Exchange Factor Role in Protein Metabolism. <i>Journal of Biological Chemistry</i> , 2010, 285, 37995-38004. | 1.6  | 19        |
| 1877 | An Atg9-containing compartment that functions in the early steps of autophagosome biogenesis. <i>Journal of Cell Biology</i> , 2010, 190, 1005-1022.   | 2.3  | 412       |
| 1878 | The Awesome Power of Yeast Evolutionary Genetics: New Genome Sequences and Strain Resources for the <i>Saccharomyces sensu stricto</i> Genus. <i>G3: Genes, Genomes, Genetics</i> , 2011, 1, 11-25.    | 0.8  | 348       |
| 1879 | Control of Chromatin Structure by Spt6: Different Consequences in Coding and Regulatory Regions. <i>Molecular and Cellular Biology</i> , 2011, 31, 531-541.  | 1.1  | 68        |
| 1880 | Histone H4 Lysine 20 of <i>Saccharomyces cerevisiae</i> Is Monomethylated and Functions in Subtelomeric Silencing. <i>Biochemistry</i> , 2011, 50, 10473-10483.  | 1.2  | 24        |
| 1881 | Advanced Methods for High-Throughput Microscopy Screening of Genetically Modified Yeast Libraries. <i>Methods in Molecular Biology</i> , 2011, 781, 127-159.   | 0.4  | 101       |
| 1882 | Conditional Depletion of Nuclear Proteins by the Anchor Away System. <i>Current Protocols in Molecular Biology</i> , 2011, 93, Unit13.10B.   | 2.9  | 9         |
| 1884 | Global Regulation of H2A.Z Localization by the INO80 Chromatin-Remodeling Enzyme Is Essential for Genome Integrity. <i>Cell</i> , 2011, 144, 200-213.  | 13.5 | 345       |
| 1885 | Osh Proteins Regulate Phosphoinositide Metabolism at ER-Plasma Membrane Contact Sites. <i>Cell</i> , 2011, 144, 389-401.   | 13.5 | 442       |
| 1886 | DNA Replication through G-Quadruplex Motifs Is Promoted by the <i>Saccharomyces cerevisiae</i> Pif1 DNA Helicase. <i>Cell</i> , 2011, 145, 678-691.  | 13.5 | 492       |
| 1887 | Eukaryotic Origin-Dependent DNA Replication In Vitro Reveals Sequential Action of DDK and S-CDK Kinases. <i>Cell</i> , 2011, 146, 80-91.   | 13.5 | 276       |
| 1888 | Motility and Segregation of Hsp104-Associated Protein Aggregates in Budding Yeast. <i>Cell</i> , 2011, 147, 1186-1196.   | 13.5 | 114       |
| 1889 | Regulated Offloading of Cytoplasmic Dynein from Microtubule Plus Ends to the Cortex. <i>Developmental Cell</i> , 2011, 20, 639-651.  | 3.1  | 95        |
| 1890 | The mating projections of <i>Saccharomyces cerevisiae</i> and <i>Candida albicans</i> show key characteristics of hyphal growth. <i>Fungal Biology</i> , 2011, 115, 547-556.                           | 1.1  | 18        |
| 1891 | Optimized protocols and plasmids for in vivo cloning in yeast. <i>Gene</i> , 2011, 484, 86-89.   | 1.0  | 9         |
| 1892 | Functions of the mitotic B-type cyclins CLB1, CLB2, and CLB3 at mitotic exit antagonized by the CDC14 phosphatase. <i>Fungal Genetics and Biology</i> , 2011, 48, 966-978.                             | 0.9  | 5         |
| 1893 | Expressing the <i>Erwinia amylovora</i> type III effector DspA/E in the yeast <i>Saccharomyces cerevisiae</i> strongly alters cellular trafficking. <i>FEBS Open Bio</i> , 2011, 1, 23-28.             | 1.0  | 10        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 1894 | Metabolic link between phosphatidylethanolamine and triacylglycerol metabolism in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2011, 1811, 1030-1037.   | 1.2  | 49        |
| 1895 | Ketoconazole and miconazole alter potassium homeostasis in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2011, 1808, 433-445.  | 1.4  | 13        |
| 1896 | Gln3 and Gcn4 hybrid transcriptional activator determines catabolic and biosynthetic gene expression in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2011, 404, 859-864. | 1.0  | 11        |
| 1897 | N-terminal short fragment of TUP1 confers resistance to 5-bromodeoxyuridine in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2011, 411, 25-31.                            | 1.0  | 3         |
| 1898 | Cdk1-Dependent Destruction of Eco1 Prevents Cohesion Establishment after S Phase. <i>Molecular Cell</i> , 2011, 42, 378-389.   | 4.5  | 72        |
| 1899 | Acetyl-CoA Induces Cell Growth and Proliferation by Promoting the Acetylation of Histones at Growth Genes. <i>Molecular Cell</i> , 2011, 42, 426-437.  | 4.5  | 583       |
| 1900 | Sgs1 Truncations Induce Genome Rearrangements but Suppress Detrimental Effects of BLM Overexpression in <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 2011, 405, 877-891.                                  | 2.0  | 15        |
| 1901 | Destabilization and Recovery of a Yeast Prion after Mild Heat Shock. <i>Journal of Molecular Biology</i> , 2011, 408, 432-448.   | 2.0  | 73        |
| 1902 | Structure of H/ACA RNP Protein Nhp2p Reveals Cis/Trans Isomerization of a Conserved Proline at the RNA and Nop10 Binding Interface. <i>Journal of Molecular Biology</i> , 2011, 411, 927-942.  | 2.0  | 30        |
| 1903 | Genomic Promoter Replacement Cassettes to Alter Gene Expression in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Methods in Molecular Biology</i> , 2011, 765, 275-294.   | 0.4  | 10        |
| 1904 | XUTs are a class of Xrn1-sensitive antisense regulatory non-coding RNA in yeast. <i>Nature</i> , 2011, 475, 114-117.   | 13.7 | 329       |
| 1905 | Sensing of Replication Stress and Mec1 Activation Act through Two Independent Pathways Involving the 9-1-1 Complex and DNA Polymerase $\delta$ . <i>PLoS Genetics</i> , 2011, 7, e1002022.   | 1.5  | 40        |
| 1906 | Identification of an Amphipathic Helix Important for the Formation of Ectopic Septin Spirals and Axial Budding in Yeast Axial Landmark Protein Bud3p. <i>PLoS ONE</i> , 2011, 6, e16744.   | 1.1  | 14        |
| 1907 | A Viable Hypomorphic Allele of the Essential IMP3 Gene Reveals Novel Protein Functions in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2011, 6, e19500.   | 1.1  | 5         |
| 1908 | Unexpected Accumulation of mcm5U and mcm5s2U in a trm9 Mutant Suggests an Additional Step in the Synthesis of mcm5U and mcm5s2U. <i>PLoS ONE</i> , 2011, 6, e20783.  | 1.1  | 66        |
| 1909 | Dia2 Controls Transcription by Mediating Assembly of the RSC Complex. <i>PLoS ONE</i> , 2011, 6, e21172.   | 1.1  | 6         |
| 1910 | Insights into the Role of Yeast eIF2A in IRES-Mediated Translation. <i>PLoS ONE</i> , 2011, 6, e24492.   | 1.1  | 25        |
| 1911 | Regulation of Septin Dynamics by the <i>Saccharomyces cerevisiae</i> Lysine Acetyltransferase NuA4. <i>PLoS ONE</i> , 2011, 6, e25336.   | 1.1  | 33        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1912 | Multiple TORC1-Associated Proteins Regulate Nitrogen Starvation-Dependent Cellular Differentiation in <i>Saccharomyces cerevisiae</i> . PLoS ONE, 2011, 6, e26081.                                 | 1.1 | 18        |
| 1913 | The Inheritance of Histone Modifications Depends upon the Location in the Chromosome in <i>Saccharomyces cerevisiae</i> . PLoS ONE, 2011, 6, e28980.   | 1.1 | 2         |
| 1914 | <i>In vivo</i> and <i>in vitro</i> Studies of Adaptor-clathrin Interaction. Journal of Visualized Experiments, 2011, , .   | 0.2 | 8         |
| 1915 | Functional analysis of RhoGDI inhibitory activity on vacuole membrane fusion. Biochemical Journal, 2011, 434, 445-457.   | 1.7 | 4         |
| 1916 | Rim2, a pyrimidine nucleotide exchanger, is needed for iron utilization in mitochondria. Biochemical Journal, 2011, 440, 137-146.  | 1.7 | 42        |
| 1917 | A Bnr-like formin links actin to the spindle pole body during sporulation in the filamentous fungus <i>Ashbya gossypii</i> . Molecular Microbiology, 2011, 80, 1276-1295.                          | 1.2 | 8         |
| 1918 | Nucleocytoplasmic shuttling of Ssd1 defines the destiny of its bound mRNAs. Molecular Microbiology, 2011, 81, 831-849.   | 1.2 | 33        |
| 1919 | Organization and Assembly of the TRAPP II Complex. Traffic, 2011, 12, 715-725.   | 1.3 | 60        |
| 1920 | Vacuolar Import of Phosphatidylcholine Requires the ATP-binding Cassette Transporter Ybt1. Traffic, 2011, 12, 1257-1268.   | 1.3 | 34        |
| 1921 | The Sterol-binding Protein Kes1/Osh4p Is a Regulator of Polarized Exocytosis. Traffic, 2011, 12, 1521-1536.  | 1.3 | 65        |
| 1922 | Ume6p is required for germination and early colony development of yeast ascospores. FEMS Yeast Research, 2011, 11, 104-113.  | 1.1 | 11        |
| 1923 | An efficient method to optimize <i>Kluyveromyces lactis</i> gene targeting. FEMS Yeast Research, 2011, 11, 509-513.  | 1.1 | 13        |
| 1924 | Gcn4p-mediated transcriptional repression of ribosomal protein genes under amino-acid starvation. EMBO Journal, 2011, 30, 859-872.   | 3.5 | 43        |
| 1925 | The highly conserved KEOPS/EKC complex is essential for a universal tRNA modification, t6A. EMBO Journal, 2011, 30, 873-881.   | 3.5 | 149       |
| 1926 | Mitochondria regulate autophagy by conserved signalling pathways. EMBO Journal, 2011, 30, 2101-2114.   | 3.5 | 156       |
| 1927 | The SWI/SNF complex acts to constrain distribution of the centromeric histone variant Cse4. EMBO Journal, 2011, 30, 1919-1927.   | 3.5 | 47        |
| 1928 | A stable yeast strain efficiently producing cholesterol instead of ergosterol is functional for tryptophan uptake, but not weak organic acid resistance. Metabolic Engineering, 2011, 13, 555-569. | 3.6 | 95        |
| 1929 | Modeling Vesicle Traffic Reveals Unexpected Consequences for Cdc42p-Mediated Polarity Establishment. Current Biology, 2011, 21, 184-194.   | 1.8 | 111       |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1930 | RAD51-independent inverted-repeat recombination by a strand-annealing mechanism. <i>DNA Repair</i> , 2011, 10, 408-415.  | 1.3 | 33        |
| 1931 | Ddc1 checkpoint protein and DNA polymerase $\epsilon$ interact with nick-containing DNA repair intermediate in cell free extracts of <i>Saccharomyces cerevisiae</i> . <i>DNA Repair</i> , 2011, 10, 815-825.                | 1.3 | 10        |
| 1932 | Srs2 enables checkpoint recovery by promoting disassembly of DNA damage foci from chromatin. <i>DNA Repair</i> , 2011, 10, 1213-1222.  | 1.3 | 26        |
| 1933 | The dCMP transferase activity of yeast Rev1 is biologically relevant during the bypass of endogenously generated AP sites. <i>DNA Repair</i> , 2011, 10, 1262-1271.  | 1.3 | 36        |
| 1934 | Pil1, an eisosome organizer, plays an important role in the recruitment of synaptojanins and amphiphysins to facilitate receptor-mediated endocytosis in yeast. <i>European Journal of Cell Biology</i> , 2011, 90, 825-833. | 1.6 | 25        |
| 1935 | Recognition of an ERAD-L substrate analyzed by site-specific in vivo photocrosslinking. <i>FEBS Letters</i> , 2011, 585, 1281-1286.  | 1.3 | 25        |
| 1936 | Molecular response of <i>Saccharomyces cerevisiae</i> wine and laboratory strains to high sugar stress conditions. <i>International Journal of Food Microbiology</i> , 2011, 145, 211-220.                                   | 2.1 | 44        |
| 1937 | Coenzyme Q Biosynthesis: Coq6 Is Required for the C5-Hydroxylation Reaction and Substrate Analogs Rescue Coq6 Deficiency. <i>Chemistry and Biology</i> , 2011, 18, 1134-1142.  | 6.2 | 92        |
| 1938 | Assembly of <i>Saccharomyces cerevisiae</i> 60S ribosomal subunits: role of factors required for 27S pre-rRNA processing. <i>EMBO Journal</i> , 2011, 30, 4020-4032.   | 3.5 | 84        |
| 1939 | The PIAS homologue Siz2 regulates perinuclear telomere position and telomerase activity in budding yeast. <i>Nature Cell Biology</i> , 2011, 13, 867-874.  | 4.6 | 88        |
| 1940 | Probing the role of point mutations in the cyp51A gene from <i>Aspergillus fumigatus</i> in the model yeast <i>Saccharomyces cerevisiae</i> . <i>Medical Mycology</i> , 2011, 49, 276-284.                                   | 0.3 | 27        |
| 1941 | Blm10 Protein Promotes Proteasomal Substrate Turnover by an Active Gating Mechanism. <i>Journal of Biological Chemistry</i> , 2011, 286, 42830-42839.  | 1.6 | 74        |
| 1942 | Ribosome Assembly Factors Prevent Premature Translation Initiation by 40S Assembly Intermediates. <i>Science</i> , 2011, 333, 1449-1453.   | 6.0 | 199       |
| 1943 | A Mutation in the Golgi Qb-SNARE Gene GOSR2 Causes Progressive Myoclonus Epilepsy with Early Ataxia. <i>American Journal of Human Genetics</i> , 2011, 88, 657-663.  | 2.6 | 166       |
| 1944 | Tracking chromosome dynamics in live yeast cells: coordinated movement of rDNA homologs and anaphase disassembly of the nucleolus during meiosis. <i>Chromosome Research</i> , 2011, 19, 1013-1026.                          | 1.0 | 11        |
| 1945 | Genome-wide screen for inositol auxotrophy in <i>Saccharomyces cerevisiae</i> implicates lipid metabolism in stress response signaling. <i>Molecular Genetics and Genomics</i> , 2011, 285, 125-149.                         | 1.0 | 53        |
| 1946 | The <i>Saccharomyces cerevisiae</i> flavodoxin-like proteins Ycp4 and Rfs1 play a role in stress response and in the regulation of genes related to metabolism. <i>Archives of Microbiology</i> , 2011, 193, 515-525.        | 1.0 | 15        |
| 1947 | Requirements of Slm proteins for proper eisosome organization, endocytic trafficking and recycling in the yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Biosciences</i> , 2011, 36, 79-96.                           | 0.5 | 28        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 1948 | Identification of a functional docking site in the Rpn1 LRR domain for the UBA-UBL domain protein Ddi1. <i>BMC Biology</i> , 2011, 9, 33.   | 1.7 | 62        |
| 1949 | Rec10 and Rec12 independent recombination in meiosis of <i>Schizosaccharomyces pombe</i> . <i>Yeast</i> , 2011, 28, 405-421.  | 0.8 | 2         |
| 1950 | Selection of STOP-free sequences from random mutagenesis for loss of interaction™ two hybrid studies. <i>Yeast</i> , 2011, 28, 535-545.   | 0.8 | 4         |
| 1951 | Quantitative analysis of Pac1/LIS1 mediated dynein targeting: Implications for regulation of dynein activity in budding yeast. <i>Cytoskeleton</i> , 2011, 68, 157-174.                           | 1.0 | 63        |
| 1952 | A transcriptome screen in yeast identifies a novel assembly factor for the mitochondrial complex III. <i>Mitochondrion</i> , 2011, 11, 391-396.   | 1.6 | 20        |
| 1953 | Diverse protein kinase interactions identified by protein microarrays reveal novel connections between cellular processes. <i>Genes and Development</i> , 2011, 25, 767-778.                      | 2.7 | 60        |
| 1954 | <sup>3</sup> H2A is a component of yeast heterochromatin required for telomere elongation. <i>Cell Cycle</i> , 2011, 10, 293-300.   | 1.3 | 24        |
| 1955 | Rapamycin increases rDNA stability by enhancing association of Sir2 with rDNA in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2011, 39, 1336-1350.                           | 6.5 | 70        |
| 1956 | Polymerase Epsilon Is Required To Maintain Replicative Senescence. <i>Molecular and Cellular Biology</i> , 2011, 31, 1637-1645.   | 1.1 | 11        |
| 1957 | Mitotic Exit Control of the <i>Saccharomyces cerevisiae</i> Ndr/LATS Kinase Cbk1 Regulates Daughter Cell Separation after Cytokinesis. <i>Molecular and Cellular Biology</i> , 2011, 31, 721-735. | 1.1 | 49        |
| 1958 | Intergenic transcription causes repression by directing nucleosome assembly. <i>Genes and Development</i> , 2011, 25, 29-40.  | 2.7 | 154       |
| 1959 | Bro1 binding to Snf7 regulates ESCRT-III membrane scission activity in yeast. <i>Journal of Cell Biology</i> , 2011, 192, 295-306.  | 2.3 | 78        |
| 1960 | Mechanism of activation of methyltransferases involved in translation by the Trm112 hub™ protein. <i>Nucleic Acids Research</i> , 2011, 39, 6249-6259.  | 6.5 | 65        |
| 1961 | Clustering heterochromatin: Sir3 promotes telomere clustering independently of silencing in yeast. <i>Journal of Cell Biology</i> , 2011, 192, 417-431.   | 2.3 | 64        |
| 1962 | Identification of novel proteins associated with yeast snR30 small nucleolar RNA. <i>Nucleic Acids Research</i> , 2011, 39, 9659-9670.  | 6.5 | 16        |
| 1963 | Clathrin light chain directs endocytosis by influencing the binding of the yeast Hip1R homologue, Sla2, to F-actin. <i>Molecular Biology of the Cell</i> , 2011, 22, 3699-3714.                   | 0.9 | 37        |
| 1964 | Existence of a novel clathrin-independent endocytic pathway in yeast that depends on Rho1 and formin. <i>Journal of Cell Biology</i> , 2011, 195, 657-671.  | 2.3 | 62        |
| 1965 | Viral protein targeting to the cortical endoplasmic reticulum is required for cell-cell spreading in plants. <i>Journal of Cell Biology</i> , 2011, 193, 521-535.                                 | 2.3 | 81        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1966 | Intranuclear Function for Protein Phosphatase 2A: Pph21 and Pph22 Are Required for Rapamycin-Induced GATA Factor Binding to the <i>DAL5</i> Promoter in Yeast. <i>Molecular and Cellular Biology</i> , 2011, 31, 92-104. | 1.1 | 22        |
| 1967 | Systematic Screen of <i>Schizosaccharomyces pombe</i> Deletion Collection Uncovers Parallel Evolution of the Phosphate Signal Transduction Pathway in Yeasts. <i>Eukaryotic Cell</i> , 2011, 10, 198-206.                | 3.4 | 46        |
| 1968 | A mitochondrial-focused genetic interaction map reveals a scaffold-like complex required for inner membrane organization in mitochondria. <i>Journal of Cell Biology</i> , 2011, 195, 323-340.                           | 2.3 | 402       |
| 1969 | CENP-A exceeds microtubule attachment sites in centromere clusters of both budding and fission yeast. <i>Journal of Cell Biology</i> , 2011, 195, 563-572.   | 2.3 | 126       |
| 1970 | Cohesin Plays a Dual Role in Gene Regulation and Sister-Chromatid Cohesion During Meiosis in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2011, 187, 1041-1051.   | 1.2 | 15        |
| 1971 | Cytosolic Localization of Acetohydroxyacid Synthase Ilv2 and Its Impact on Diacetyl Formation during Beer Fermentation. <i>Applied and Environmental Microbiology</i> , 2011, 77, 727-731.                               | 1.4 | 31        |
| 1972 | Spt10 and Spt21 Are Required for Transcriptional Silencing in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2011, 10, 118-129.  | 3.4 | 15        |
| 1973 | The Swi2-like protein Uls1 is involved in replication stress response. <i>Nucleic Acids Research</i> , 2011, 39, 8765-8777.  | 6.5 | 26        |
| 1974 | Gex1 is a yeast glutathione exchanger that interferes with pH and redox homeostasis. <i>Molecular Biology of the Cell</i> , 2011, 22, 2054-2067.   | 0.9 | 57        |
| 1975 | The mRNA cap-binding complex stimulates the formation of pre-initiation complex at the promoter via its interaction with Mot1p in vivo. <i>Nucleic Acids Research</i> , 2011, 39, 2188-2209.                             | 6.5 | 29        |
| 1976 | A genomic integration method for the simultaneous visualization of endogenous mRNAs and their translation products in living yeast. <i>Rna</i> , 2011, 17, 2249-2255.  | 1.6 | 22        |
| 1977 | The eisosome core is composed of BAR domain proteins. <i>Molecular Biology of the Cell</i> , 2011, 22, 2360-2372.  | 0.9 | 91        |
| 1978 | Dual function of Rpn5 in two PCI complexes, the 26S proteasome and COP9 signalosome. <i>Molecular Biology of the Cell</i> , 2011, 22, 911-920.   | 0.9 | 40        |
| 1979 | Dynamic regulation of mitochondrial respiratory chain efficiency in <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> , 2011, 157, 3500-3511.   | 0.7 | 19        |
| 1980 | The yeast Batten disease orthologue Btn1 controls endosome-Golgi retrograde transport via SNARE assembly. <i>Journal of Cell Biology</i> , 2011, 195, 203-215.   | 2.3 | 44        |
| 1981 | Cdc55 coordinates spindle assembly and chromosome disjunction during meiosis. <i>Journal of Cell Biology</i> , 2011, 193, 1213-1228.   | 2.3 | 28        |
| 1982 | Scaffold number in yeast signaling system sets tradeoff between system output and dynamic range. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 20265-20270.        | 3.3 | 57        |
| 1983 | Cell Polarity in <i>Saccharomyces cerevisiae</i> Depends on Proper Localization of the Bud9 Landmark Protein by the EKC/KEOPS Complex. <i>Genetics</i> , 2011, 188, 871-882.   | 1.2 | 7         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 1984 | Hap2-3-5-Gln3 determine transcriptional activation of GDH1 and ASN1 under repressive nitrogen conditions in the yeast <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> , 2011, 157, 879-889. | 0.7 | 14        |
| 1985 | Glucose regulates clathrin adaptors at the trans-Golgi network and endosomes. <i>Molecular Biology of the Cell</i> , 2011, 22, 3671-3683.  | 0.9 | 25        |
| 1986 | Cellular Morphogenesis Under Stress Is Influenced by the Sphingolipid Pathway Gene <i>ISC1</i> and DNA Integrity Checkpoint Genes in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2011, 189, 533-547.       | 1.2 | 20        |
| 1987 | Coordinated regulation of sulfur and phospholipid metabolism reflects the importance of methylation in the growth of yeast. <i>Molecular Biology of the Cell</i> , 2011, 22, 4192-4204.                            | 0.9 | 43        |
| 1988 | Cbk1 kinase and Bck2 control MAP kinase activation and inactivation during heat shock. <i>Molecular Biology of the Cell</i> , 2011, 22, 4892-4907.   | 0.9 | 19        |
| 1989 | Direct regulation of nucleosome density by the conserved AAA-ATPase Yta7. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, E1302-11.                            | 3.3 | 39        |
| 1990 | The spindle checkpoint protein Mad2 regulates APC/C activity during prometaphase and metaphase of meiosis I in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2011, 22, 2848-2861.       | 0.9 | 33        |
| 1991 | Transcription Regulation by the Noncoding RNA <i>SRG1</i> Requires Spt2-Dependent Chromatin Deposition in the Wake of RNA Polymerase II. <i>Molecular and Cellular Biology</i> , 2011, 31, 1288-1300.              | 1.1 | 56        |
| 1992 | Pathways for Holliday Junction Processing during Homologous Recombination in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2011, 31, 1921-1933.  | 1.1 | 58        |
| 1993 | The Zds proteins control entry into mitosis and target protein phosphatase 2A to the Cdc25 phosphatase. <i>Molecular Biology of the Cell</i> , 2011, 22, 20-32.  | 0.9 | 34        |
| 1994 | The yeast Cbk1 kinase regulates mRNA localization via the mRNA-binding protein Ssd1. <i>Journal of Cell Biology</i> , 2011, 192, 583-598.  | 2.3 | 54        |
| 1995 | Trs65p, a subunit of the Ypt1p GEF TRAPP <sup>II</sup> , interacts with the Arf1p exchange factor Gea2p to facilitate COPI-mediated vesicle traffic. <i>Molecular Biology of the Cell</i> , 2011, 22, 3634-3644.   | 0.9 | 26        |
| 1996 | Functional Elements in Initiation Factors 1, 1A, and 2 <sup>̂</sup> Discriminate against Poor AUG Context and Non-AUG Start Codons. <i>Molecular and Cellular Biology</i> , 2011, 31, 4814-4831.                   | 1.1 | 71        |
| 1997 | Luminal interactions in nuclear pore complex assembly and stability. <i>Molecular Biology of the Cell</i> , 2011, 22, 1375-1388.   | 0.9 | 56        |
| 1998 | The yeast p24 complex regulates GPI-anchored protein transport and quality control by monitoring anchor remodeling. <i>Molecular Biology of the Cell</i> , 2011, 22, 2924-2936.                                    | 0.9 | 113       |
| 1999 | Functional Interplay of the Mre11 Nuclease and Ku in the Response to Replication-Associated DNA Damage. <i>Molecular and Cellular Biology</i> , 2011, 31, 4379-4389.   | 1.1 | 94        |
| 2000 | Spatial regulation of Cdc55 <sup>PP2A</sup> by Zds1/Zds2 controls mitotic entry and mitotic exit in budding yeast. <i>Journal of Cell Biology</i> , 2011, 193, 445-454.  | 2.3 | 53        |
| 2001 | New Regulators of a High Affinity Ca <sup>2+</sup> Influx System Revealed through a Genome-wide Screen in Yeast. <i>Journal of Biological Chemistry</i> , 2011, 286, 10744-10754.                                  | 1.6 | 92        |



| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2002 | Development and validation of a yeast high-throughput screen for inhibitors of A $\beta$ 242 oligomerization. <i>DMM Disease Models and Mechanisms</i> , 2011, 4, 822-831.                                    | 1.2 | 43        |
| 2003 | RNA-Binding Protein Khd1 and Ccr4 Deadenyase Play Overlapping Roles in the Cell Wall Integrity Pathway in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2011, 10, 1340-1347.                     | 3.4 | 21        |
| 2004 | Chemical and Synthetic Genetic Array Analysis Identifies Genes that Suppress Xylose Utilization and Fermentation in <i>Saccharomyces cerevisiae</i> . <i>G3: Genes, Genomes, Genetics</i> , 2011, 1, 247-258. | 0.8 | 22        |
| 2005 | The Cik1/Kar3 Motor Complex Is Required for the Proper Kinetochore-Microtubule Interaction After Stressful DNA Replication. <i>Genetics</i> , 2011, 187, 397-407.   | 1.2 | 14        |
| 2006 | The Ddc2/ATRIP checkpoint protein monitors meiotic recombination intermediates. <i>Journal of Cell Science</i> , 2011, 124, 2488-2500.  | 1.2 | 41        |
| 2007 | Sequential requirements for the GTPase domain of the mitofusin Fzo1 and the ubiquitin ligase SCFMdm30 in mitochondrial outer membrane fusion. <i>Journal of Cell Science</i> , 2011, 124, 1403-1410.          | 1.2 | 65        |
| 2008 | Genome-Wide Transcriptional Dependence on Conserved Regions of Mot1. <i>Molecular and Cellular Biology</i> , 2011, 31, 2253-2261.   | 1.1 | 5         |
| 2009 | Tem1 localization to the spindle pole bodies is essential for mitotic exit and impairs spindle checkpoint function. <i>Journal of Cell Biology</i> , 2011, 192, 599-614.                                      | 2.3 | 58        |
| 2010 | Yet1p interacts with Scs2p-Opi1p to regulate ER localization of the Opi1p repressor. <i>Molecular Biology of the Cell</i> , 2011, 22, 1430-1439.  | 0.9 | 15        |
| 2011 | Histone H2B ubiquitylation and H3 lysine 4 methylation prevent ectopic silencing of euchromatic loci important for the cellular response to heat. <i>Molecular Biology of the Cell</i> , 2011, 22, 2741-2753. | 0.9 | 13        |
| 2012 | Two novel WD40 domain-containing proteins, Ere1 and Ere2, function in the retromer-mediated endosomal recycling pathway. <i>Molecular Biology of the Cell</i> , 2011, 22, 4093-4107.                          | 0.9 | 41        |
| 2013 | Exposure of bipartite hydrophobic signal triggers nuclear quality control of Ndc10 at the endoplasmic reticulum/nuclear envelope. <i>Molecular Biology of the Cell</i> , 2011, 22, 4726-4739.                 | 0.9 | 55        |
| 2014 | The budding yeast polo-like kinase Cdc5 regulates the Ndt80 branch of the meiotic recombination checkpoint pathway. <i>Molecular Biology of the Cell</i> , 2011, 22, 3478-3490.                               | 0.9 | 34        |
| 2015 | Selective regulation of autophagy by the Iml1-Npr2-Npr3 complex in the absence of nitrogen starvation. <i>Molecular Biology of the Cell</i> , 2011, 22, 4124-4133.  | 0.9 | 81        |
| 2016 | Two Proteins with Different Functions Are Derived from the <i>KIHEM13</i> Gene. <i>Eukaryotic Cell</i> , 2011, 10, 1331-1339.   | 3.4 | 1         |
| 2017 | Highly redundant function of multiple AT-rich sequences as core promoter elements in the TATA-less <i>RPS5</i> promoter of <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2011, 39, 59-75. | 6.5 | 34        |
| 2018 | Roles of Dim2 in Ribosome Assembly. <i>Journal of Biological Chemistry</i> , 2011, 286, 2578-2586.  | 1.6 | 43        |
| 2019 | Transcriptional Regulation by Asf1. <i>Journal of Biological Chemistry</i> , 2011, 286, 7082-7092.  | 1.6 | 26        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2020 | UBC9 Autosumoylation Negatively Regulates Sumoylation of Septins in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2011, 286, 21826-21834.  | 1.6 | 20        |
| 2021 | Investigation of in Vivo Diferric Tyrosyl Radical Formation in <i>Saccharomyces cerevisiae</i> Rnr2 Protein. <i>Journal of Biological Chemistry</i> , 2011, 286, 41499-41509.   | 1.6 | 46        |
| 2022 | The Hog1 Mitogen-Activated Protein Kinase Mediates a Hypoxic Response in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2011, 188, 325-338.  | 1.2 | 57        |
| 2023 | Ama1p-activated anaphase-promoting complex regulates the destruction of Cdc20p during meiosis II. <i>Molecular Biology of the Cell</i> , 2011, 22, 315-326.   | 0.9 | 17        |
| 2024 | Topoisomerase 1 provokes the formation of short deletions in repeated sequences upon high transcription in <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 692-697. | 3.3 | 72        |
| 2025 | Lipid droplets are functionally connected to the endoplasmic reticulum in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 2011, 124, 2424-2437.  | 1.2 | 356       |
| 2026 | Key Functional Regions in the Histone Variant H2A.Z C-Terminal Docking Domain. <i>Molecular and Cellular Biology</i> , 2011, 31, 3871-3884.   | 1.1 | 25        |
| 2027 | Evidence that a septin diffusion barrier is dispensable for cytokinesis in budding yeast. <i>Biological Chemistry</i> , 2011, 392, 813-829.   | 1.2 | 71        |
| 2028 | Quantitative Proteomic Analysis of Chromatin Reveals that Ctf18 Acts in the DNA Replication Checkpoint. <i>Molecular and Cellular Proteomics</i> , 2011, 10, M110.005561.   | 2.5 | 60        |
| 2029 | A generalized model for multi-marker analysis of cell cycle progression in synchrony experiments. <i>Bioinformatics</i> , 2011, 27, i295-i303.  | 1.8 | 8         |
| 2030 | Yeast Transcription Elongation Factor Spt5 Associates with RNA Polymerase I and RNA Polymerase II Directly. <i>Journal of Biological Chemistry</i> , 2011, 286, 18825-18833.  | 1.6 | 45        |
| 2031 | Phosphate-responsive Signaling Pathway Is a Novel Component of NAD <sup>+</sup> Metabolism in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2011, 286, 14271-14281.  | 1.6 | 50        |
| 2032 | Distinct Functions of Evolutionary Conserved MSF1 and Late Embryogenesis Abundant (LEA)-like Domains in Mitochondria. <i>Journal of Biological Chemistry</i> , 2011, 286, 39141-39152.  | 1.6 | 16        |
| 2033 | A Conserved Residue in the Yeast Bem1p SH3 Domain Maintains the High Level of Binding Specificity Required for Function. <i>Journal of Biological Chemistry</i> , 2011, 286, 19470-19477.   | 1.6 | 9         |
| 2034 | Mne1 Is a Novel Component of the Mitochondrial Splicing Apparatus Responsible for Processing of a COX1 Group I Intron in Yeast. <i>Journal of Biological Chemistry</i> , 2011, 286, 10137-10146.  | 1.6 | 17        |
| 2035 | Rtt107 Is Required for Recruitment of the SMC5/6 Complex to DNA Double Strand Breaks. <i>Journal of Biological Chemistry</i> , 2011, 286, 26250-26257.  | 1.6 | 43        |
| 2036 | An Unusual Transmembrane Helix in the Endoplasmic Reticulum Ubiquitin Ligase Doa10 Modulates Degradation of Its Cognate E2 Enzyme. <i>Journal of Biological Chemistry</i> , 2011, 286, 20163-20174.   | 1.6 | 35        |
| 2037 | Functional Adaptation between Yeast Actin and Its Cognate Myosin Motors. <i>Journal of Biological Chemistry</i> , 2011, 286, 30384-30392.   | 1.6 | 9         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2038 | Î±-Synuclein and ALPS motifs are membrane curvature sensors whose contrasting chemistry mediates selective vesicle binding. <i>Journal of Cell Biology</i> , 2011, 194, 89-103.  | 2.3 | 177       |
| 2039 | lpl1/Aurora-dependent phosphorylation of Sli15/INCENP regulates CPC's spindle interaction to ensure proper microtubule dynamics. <i>Journal of Cell Biology</i> , 2011, 194, 137-153.  | 2.3 | 39        |
| 2040 | Chromosome passenger complexes control anaphase duration and spindle elongation via a kinesin-5 brake. <i>Journal of Cell Biology</i> , 2011, 193, 285-294.  | 2.3 | 30        |
| 2041 | The Conserved Foot Domain of RNA Pol II Associates with Proteins Involved in Transcriptional Initiation and/or Early Elongation. <i>Genetics</i> , 2011, 189, 1235-1248.   | 1.2 | 17        |
| 2042 | Maf1 Protein, Repressor of RNA Polymerase III, Indirectly Affects tRNA Processing. <i>Journal of Biological Chemistry</i> , 2011, 286, 39478-39488.  | 1.6 | 44        |
| 2043 | Lte1 promotes mitotic exit by controlling the localization of the spindle position checkpoint kinase Kin4. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 12584-12590.                            | 3.3 | 36        |
| 2044 | The Cyc8-Tup1 complex inhibits transcription primarily by masking the activation domain of the recruiting protein. <i>Genes and Development</i> , 2011, 25, 2525-2539.   | 2.7 | 118       |
| 2045 | Determinants of endocytic membrane geometry, stability, and scission. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, E979-88.   | 3.3 | 75        |
| 2046 | Role for topoisomerase 1 in transcription-associated mutagenesis in yeast. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 698-703.  | 3.3 | 80        |
| 2047 | Structural Studies and the Assembly of the Heptameric Post-translational Translocon Complex. <i>Journal of Biological Chemistry</i> , 2011, 286, 2956-2965.  | 1.6 | 35        |
| 2048 | Discovery of a Small Molecule Targeting <i>IRA2</i> Deletion in Budding Yeast and Neurofibromin Loss in Malignant Peripheral Nerve Sheath Tumor Cells. <i>Molecular Cancer Therapeutics</i> , 2011, 10, 1740-1750.                                     | 1.9 | 10        |
| 2049 | The DExD/H box ATPase Dhh1 functions in translational repression, mRNA decay, and processing body dynamics. <i>Journal of Cell Biology</i> , 2011, 194, 527-537.   | 2.3 | 77        |
| 2050 | Symmetry, asymmetry, and kinetics of silencing establishment in <i>Saccharomyces cerevisiae</i> revealed by single-cell optical assays. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 1209-1216. | 3.3 | 26        |
| 2051 | A Cdc48p-associated Factor Modulates Endoplasmic Reticulum-associated Degradation, Cell Stress, and Ubiquitinated Protein Homeostasis. <i>Journal of Biological Chemistry</i> , 2011, 286, 5744-5755.  | 1.6 | 41        |
| 2052 | Genome-wide analysis of yeast stress survival and tolerance acquisition to analyze the central trade-off between growth rate and cellular robustness. <i>Molecular Biology of the Cell</i> , 2011, 22, 4435-4446.                                      | 0.9 | 138       |
| 2053 | The multifunctional Ccr4-Not complex directly promotes transcription elongation. <i>Genes and Development</i> , 2011, 25, 581-593.   | 2.7 | 144       |
| 2054 | The conserved GTPase Gem1 regulates endoplasmic reticulum-mitochondria connections. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 14151-14156.   | 3.3 | 312       |
| 2055 | Isolation and characterization of novel mutations in CDC50, the non-catalytic subunit of the Drs2p phospholipid flippase. <i>Journal of Biochemistry</i> , 2011, 149, 423-432.   | 0.9 | 22        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2056 | Proteasomal degradation of Sfp1 contributes to the repression of ribosome biogenesis during starvation and is mediated by the proteasome activator Blm10. <i>Molecular Biology of the Cell</i> , 2011, 22, 528-540.        | 0.9 | 40        |
| 2057 | Bub1-Mediated Adaptation of the Spindle Checkpoint. <i>PLoS Genetics</i> , 2011, 7, e1001282.  | 1.5 | 10        |
| 2058 | Vacuolar Protein Sorting Genes Regulate Mat Formation in <i>Saccharomyces cerevisiae</i> by Flo11p-Dependent and -Independent Mechanisms. <i>Eukaryotic Cell</i> , 2011, 10, 1516-1526.                                    | 3.4 | 13        |
| 2059 | N-Terminal Acetylation Inhibits Protein Targeting to the Endoplasmic Reticulum. <i>PLoS Biology</i> , 2011, 9, e1001073.   | 2.6 | 167       |
| 2060 | Quantitative Epistasis Analysis and Pathway Inference from Genetic Interaction Data. <i>PLoS Computational Biology</i> , 2011, 7, e1002048.  | 1.5 | 15        |
| 2061 | Srf1 Is a Novel Regulator of Phospholipase D Activity and Is Essential to Buffer the Toxic Effects of C16:0 Platelet Activating Factor. <i>PLoS Genetics</i> , 2011, 7, e1001299.  | 1.5 | 12        |
| 2062 | Pch2 Acts through Xrs2 and Tel1/ATM to Modulate Interhomolog Bias and Checkpoint Function during Meiosis. <i>PLoS Genetics</i> , 2011, 7, e1002351.  | 1.5 | 66        |
| 2063 | The mitochondrial contact site complex, a determinant of mitochondrial architecture. <i>EMBO Journal</i> , 2011, 30, 4356-4370.  | 3.5 | 395       |
| 2064 | Cryptic Transcription Mediates Repression of Subtelomeric Metal Homeostasis Genes. <i>PLoS Genetics</i> , 2011, 7, e1002163.   | 1.5 | 23        |
| 2065 | The SUN Protein Mps3 Is Required for Spindle Pole Body Insertion into the Nuclear Membrane and Nuclear Envelope Homeostasis. <i>PLoS Genetics</i> , 2011, 7, e1002365.   | 1.5 | 89        |
| 2066 | Regulation of the CgPdr1 Transcription Factor from the Pathogen <i>Candida glabrata</i> . <i>Eukaryotic Cell</i> , 2011, 10, 187-197.  | 3.4 | 65        |
| 2067 | BAX inhibitor-1 regulates autophagy by controlling the IRE1 $\alpha$ branch of the unfolded protein response. <i>EMBO Journal</i> , 2011, 30, 4465-4478.   | 3.5 | 105       |
| 2068 | Elevated Proteasome Capacity Extends Replicative Lifespan in <i>Saccharomyces cerevisiae</i> . <i>PLoS Genetics</i> , 2011, 7, e1002253.   | 1.5 | 202       |
| 2069 | A Novel Checkpoint and RPA Inhibitory Pathway Regulated by Rif1. <i>PLoS Genetics</i> , 2011, 7, e1002417.   | 1.5 | 46        |
| 2070 | The nucleolar protein Nop19p interacts preferentially with Utp25p and Dhr2p and is essential for the production of the 40S ribosomal subunit in <i>Saccharomyces cerevisiae</i> . <i>RNA Biology</i> , 2011, 8, 1158-1172. | 1.5 | 5         |
| 2071 | Mechanism and cellular function of Bud6 as an actin nucleation-promoting factor. <i>Molecular Biology of the Cell</i> , 2011, 22, 4016-4028.   | 0.9 | 58        |
| 2072 | DNA-end capping by the budding yeast transcription factor and subtelomeric binding protein Tbf1. <i>EMBO Journal</i> , 2012, 31, 138-149.  | 3.5 | 28        |
| 2073 | Separation of DNA Replication from the Assembly of Break-Competent Meiotic Chromosomes. <i>PLoS Genetics</i> , 2012, 8, e1002643.  | 1.5 | 81        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2074 | TERRA Promotes Telomere Shortening through Exonuclease 1-Mediated Resection of Chromosome Ends. <i>PLoS Genetics</i> , 2012, 8, e1002747.  | 1.5 | 132       |
| 2075 | Reversal of PCNA Ubiquitylation by Ubp10 in <i>Saccharomyces cerevisiae</i> . <i>PLoS Genetics</i> , 2012, 8, e1002826.  | 1.5 | 46        |
| 2076 | Role of Mex67-Mtr2 in the Nuclear Export of 40S Pre-Ribosomes. <i>PLoS Genetics</i> , 2012, 8, e1002915.   | 1.5 | 62        |
| 2077 | The Enigmatic Conservation of a Rap1 Binding Site in the <i>Saccharomyces cerevisiae</i> HMRE Silencer. <i>G3: Genes, Genomes, Genetics</i> , 2012, 2, 1555-1562.                                      | 0.8 | 5         |
| 2078 | Rad53 is essential for a mitochondrial DNA inheritance checkpoint regulating G1 to S progression. <i>Journal of Cell Biology</i> , 2012, 198, 793-798.   | 2.3 | 28        |
| 2079 | Construction of histidine-tagged yeast mitochondrial cytochrome <i>c</i> oxidase for facile purification of mutant forms. <i>Biochemical Journal</i> , 2012, 444, 199-204.                             | 1.7 | 29        |
| 2080 | Regulation of clathrin coat assembly by Eps15 homology domain-mediated interactions during endocytosis. <i>Molecular Biology of the Cell</i> , 2012, 23, 687-700.                                      | 0.9 | 13        |
| 2081 | Cell cycle-dependent regulation of the nuclease activity of Mus81-Eme1/Mms4. <i>Nucleic Acids Research</i> , 2012, 40, 8325-8335.  | 6.5 | 109       |
| 2082 | Polyglutamine Toxicity Is Controlled by Prion Composition and Gene Dosage in Yeast. <i>PLoS Genetics</i> , 2012, 8, e1002634.  | 1.5 | 45        |
| 2083 | The Rix1 (Ipi1p-2p-3p) complex is a critical determinant of DNA replication licensing independent of their roles in ribosome biogenesis. <i>Cell Cycle</i> , 2012, 11, 1325-1339.                      | 1.3 | 14        |
| 2084 | Posttranscriptional Regulation of Cell-Cell Interaction Protein-Encoding Transcripts by Zfs1p in <i>Schizosaccharomyces pombe</i> . <i>Molecular and Cellular Biology</i> , 2012, 32, 4206-4214.       | 1.1 | 23        |
| 2085 | The Linker Histone Plays a Dual Role during Gametogenesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2012, 32, 2771-2783.  | 1.1 | 28        |
| 2086 | Cdc5-Dependent Asymmetric Localization of Bfa1 Fine-Tunes Timely Mitotic Exit. <i>PLoS Genetics</i> , 2012, 8, e1002450.   | 1.5 | 24        |
| 2087 | A Yeast GSK-3 Kinase Mck1 Promotes Cdc6 Degradation to Inhibit DNA Re-Replication. <i>PLoS Genetics</i> , 2012, 8, e1003099.   | 1.5 | 29        |
| 2088 | Persulfide formation on mitochondrial cysteine desulfurase: enzyme activation by a eukaryote-specific interacting protein and Fe-S cluster synthesis. <i>Biochemical Journal</i> , 2012, 448, 171-187. | 1.7 | 58        |
| 2089 | Distinct role of Mediator tail module in regulation of SAGA-dependent, TATA-containing genes in yeast. <i>EMBO Journal</i> , 2012, 31, 44-57.  | 3.5 | 67        |
| 2090 | The yeast vacuolar Rab GTPase Ypt7p has an activity beyond membrane recruitment of the homotypic fusion and protein sorting Class C Vps complex. <i>Biochemical Journal</i> , 2012, 443, 205-211.      | 1.7 | 6         |
| 2091 | Characterizing the roles of Met31 and Met32 in coordinating Met4-activated transcription in the absence of Met30. <i>Molecular Biology of the Cell</i> , 2012, 23, 1928-1942.                          | 0.9 | 25        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2092 | MYST protein acetyltransferase activity requires active site lysine autoacetylation. <i>EMBO Journal</i> , 2012, 31, 58-70.   | 3.5 | 101       |
| 2093 | The dual PH domain protein Opy1 functions as a sensor and modulator of PtdIns(4,5)P <sub>2</sub> synthesis. <i>EMBO Journal</i> , 2012, 31, 2882-2894.  | 3.5 | 20        |
| 2094 | The Yeast Ubr1 Ubiquitin Ligase Participates in a Prominent Pathway That Targets Cytosolic Thermosensitive Mutants for Degradation. <i>G3: Genes, Genomes, Genetics</i> , 2012, 2, 619-628.       | 0.8 | 35        |
| 2095 | Sumoylation regulates Kap114-mediated nuclear transport. <i>EMBO Journal</i> , 2012, 31, 2461-2472.   | 3.5 | 33        |
| 2096 | DNA replication stress differentially regulates G1/S genes via Rad53-dependent inactivation of Nrm1. <i>EMBO Journal</i> , 2012, 31, 1811-1822.   | 3.5 | 65        |
| 2097 | The yeast vacuolar ABC transporter Ybt1p regulates membrane fusion through Ca <sup>2+</sup> transport modulation. <i>Biochemical Journal</i> , 2012, 448, 365-372.                                | 1.7 | 26        |
| 2098 | A Conserved Role of IQGAP1 in Regulating TOR Complex 1. <i>Journal of Cell Science</i> , 2012, 125, 2041-52.  | 1.2 | 41        |
| 2099 | <i>SPO71</i> Mediates Prospore Membrane Size and Maturation in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2012, 11, 1191-1200.  | 3.4 | 14        |
| 2100 | Functional Antagonism between Sas3 and Gcn5 Acetyltransferases and ISWI Chromatin Remodelers. <i>PLoS Genetics</i> , 2012, 8, e1002994.   | 1.5 | 26        |
| 2101 | Adaptor Autoregulation Promotes Coordinated Binding within Clathrin Coats. <i>Journal of Biological Chemistry</i> , 2012, 287, 17398-17407.   | 1.6 | 11        |
| 2102 | A Systematic Approach to Pair Secretory Cargo Receptors with Their Cargo Suggests a Mechanism for Cargo Selection by Erv14. <i>PLoS Biology</i> , 2012, 10, e1001329.                             | 2.6 | 87        |
| 2103 | Sequestration of Highly Expressed mRNAs in Cytoplasmic Granules, P-Bodies, and Stress Granules Enhances Cell Viability. <i>PLoS Genetics</i> , 2012, 8, e1002527.                                 | 1.5 | 75        |
| 2104 | Cdc7-Dbf4 Is a Gene-Specific Regulator of Meiotic Transcription in Yeast. <i>Molecular and Cellular Biology</i> , 2012, 32, 541-557.  | 1.1 | 21        |
| 2105 | Nsi1 plays a significant role in the silencing of ribosomal DNA in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2012, 40, 4892-4903.   | 6.5 | 21        |
| 2106 | Yeast mRNA cap-binding protein Cbc1/Sto1 is necessary for the rapid reprogramming of translation after hyperosmotic shock. <i>Molecular Biology of the Cell</i> , 2012, 23, 137-150.              | 0.9 | 45        |
| 2107 | The SWI/SNF Klsnf2 Subunit Controls the Glucose Signaling Pathway To Coordinate Glycolysis and Glucose Transport in <i>Kluyveromyces lactis</i> . <i>Eukaryotic Cell</i> , 2012, 11, 1382-1390.   | 3.4 | 8         |
| 2108 | Vacuolar H <sup>+</sup> -ATPase (V-ATPase) Promotes Vacuolar Membrane Permeabilization and Nonapoptotic Death in Stressed Yeast. <i>Journal of Biological Chemistry</i> , 2012, 287, 19029-19039. | 1.6 | 46        |
| 2109 | The RNA helicase Dhh1p cooperates with Rbp1p to promote porin mRNA decay via its non-conserved C-terminal domain. <i>Nucleic Acids Research</i> , 2012, 40, 1331-1344.                            | 6.5 | 15        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2110 | Natural Variation in the Yeast Glucose-Signaling Network Reveals a New Role for the Mig3p Transcription Factor. <i>C3: Genes, Genomes, Genetics</i> , 2012, 2, 1607-1612.                           | 0.8 | 21        |
| 2111 | Impaired Manganese Metabolism Causes Mitotic Misregulation. <i>Journal of Biological Chemistry</i> , 2012, 287, 18717-18729.  | 1.6 | 14        |
| 2112 | Mammalian HCA66 protein is required for both ribosome synthesis and centriole duplication. <i>Nucleic Acids Research</i> , 2012, 40, 6270-6289.   | 6.5 | 10        |
| 2113 | Alg14 organizes the formation of a multiglycosyltransferase complex involved in initiation of lipid-linked oligosaccharide biosynthesis. <i>Glycobiology</i> , 2012, 22, 504-516.                   | 1.3 | 44        |
| 2114 | <i>VPS13</i> Regulates Membrane Morphogenesis During Sporulation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 2012, 125, 3004-11.  | 1.2 | 90        |
| 2115 | Regulation of the formin Bnr1 by septins and a MARK/Par1-family septin-associated kinase. <i>Molecular Biology of the Cell</i> , 2012, 23, 4041-4053.   | 0.9 | 34        |
| 2116 | Ebp2 and Brx1 function cooperatively in 60S ribosomal subunit assembly in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2012, 40, 4574-4588.                                    | 6.5 | 31        |
| 2117 | The tail-module of yeast Mediator complex is required for telomere heterochromatin maintenance. <i>Nucleic Acids Research</i> , 2012, 40, 581-593.  | 6.5 | 29        |
| 2118 | Hierarchical recruitment into nascent ribosomes of assembly factors required for 27SB pre-rRNA processing in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2012, 40, 8646-8661. | 6.5 | 67        |
| 2119 | Role of mitochondrial inner membrane organizing system in protein biogenesis of the mitochondrial outer membrane. <i>Molecular Biology of the Cell</i> , 2012, 23, 3948-3956.                       | 0.9 | 108       |
| 2120 | The C-terminus of Dpb2 is required for interaction with Pol2 and for cell viability. <i>Nucleic Acids Research</i> , 2012, 40, 11545-11553.   | 6.5 | 26        |
| 2121 | Comparative dynamic transcriptome analysis (cDTA) reveals mutual feedback between mRNA synthesis and degradation. <i>Genome Research</i> , 2012, 22, 1350-1359.                                     | 2.4 | 255       |
| 2122 | Phosphatidylinositol 4-Kinases Are Required for Autophagic Membrane Trafficking. <i>Journal of Biological Chemistry</i> , 2012, 287, 37964-37972.   | 1.6 | 62        |
| 2123 | Seg1 controls eisosome assembly and shape. <i>Journal of Cell Biology</i> , 2012, 198, 405-420.   | 2.3 | 54        |
| 2124 | A novel patch assembly domain in Num1 mediates dynein anchoring at the cortex during spindle positioning. <i>Journal of Cell Biology</i> , 2012, 196, 743-756.                                      | 2.3 | 53        |
| 2125 | High-curvature domains of the ER are important for the organization of ER exit sites in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 2012, 125, 3412-20.                      | 1.2 | 73        |
| 2126 | Regulation of Vacuolar H <sup>+</sup> -ATPase Activity by the Cdc42 Effector Ste20 in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2012, 11, 442-451.                                 | 3.4 | 11        |
| 2127 | <i>Saccharomyces cerevisiae</i> Ribosomal Protein L26 Is Not Essential for Ribosome Assembly and Function. <i>Molecular and Cellular Biology</i> , 2012, 32, 3228-3241.                             | 1.1 | 48        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2128 | Rab GTPase regulation of retromer-mediated cargo export during endosome maturation. <i>Molecular Biology of the Cell</i> , 2012, 23, 2505-2515.  | 0.9 | 99        |
| 2129 | Cdc42p regulation of the yeast formin Bni1p mediated by the effector Gic2p. <i>Molecular Biology of the Cell</i> , 2012, 23, 3814-3826.  | 0.9 | 38        |
| 2130 | Sequential RNA degradation pathways provide a fail-safe mechanism to limit the accumulation of unspliced transcripts in <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 2012, 18, 1563-1572.                        | 1.6 | 30        |
| 2131 | Histone Chaperone Rtt106 Promotes Nucleosome Formation Using (H3-H4) <sub>2</sub> Tetramers. <i>Journal of Biological Chemistry</i> , 2012, 287, 10753-10760.  | 1.6 | 45        |
| 2132 | Direct Interplay among Histones, Histone Chaperones, and a Chromatin Boundary Protein in the Control of Histone Gene Expression. <i>Molecular and Cellular Biology</i> , 2012, 32, 4337-4349.                      | 1.1 | 24        |
| 2133 | Dual roles of Atg8 <sup>~</sup> PE deconjugation by Atg4 in autophagy. <i>Autophagy</i> , 2012, 8, 883-892.  | 4.3 | 196       |
| 2134 | Suppression Analysis of <i>esa1</i> Mutants in <i>Saccharomyces cerevisiae</i> Links NAB3 to Transcriptional Silencing and Nucleolar Functions. <i>G3: Genes, Genomes, Genetics</i> , 2012, 2, 1223-1232.          | 0.8 | 9         |
| 2135 | Sphingolipids Regulate the Yeast High-Osmolarity Glycerol Response Pathway. <i>Molecular and Cellular Biology</i> , 2012, 32, 2861-2870.   | 1.1 | 56        |
| 2136 | A pre-initiation complex at the 3' end of genes drives antisense transcription independent of divergent sense transcription. <i>Nucleic Acids Research</i> , 2012, 40, 2432-2444.                                  | 6.5 | 61        |
| 2137 | The Importance of Conserved Features of Yeast Actin-Binding Protein 1 (Abp1p): The Conditional Nature of Essentiality. <i>Genetics</i> , 2012, 191, 1199-1211.   | 1.2 | 10        |
| 2138 | The Mck1 GSK-3 kinase inhibits the activity of Clb2-Cdk1 post-nuclear division. <i>Cell Cycle</i> , 2012, 11, 3421-3432.   | 1.3 | 9         |
| 2139 | Desumoylation of the Endoplasmic Reticulum Membrane VAP Family Protein Scs2 by Ulp1 and SUMO Regulation of the Inositol Synthesis Pathway. <i>Molecular and Cellular Biology</i> , 2012, 32, 64-75.                | 1.1 | 24        |
| 2140 | Ribosomal proteins L7 and L8 function in concert with six A <sub>3</sub> assembly factors to propagate assembly of domains I and II of 25S rRNA in yeast 60S ribosomal subunits. <i>Rna</i> , 2012, 18, 1805-1822. | 1.6 | 59        |
| 2141 | Tye7 regulates yeast Ty1 retrotransposon sense and antisense transcription in response to adenyllic nucleotides stress. <i>Nucleic Acids Research</i> , 2012, 40, 5271-5282.                                       | 6.5 | 33        |
| 2142 | Multiple phosphorylation of Rad9 by CDK is required for DNA damage checkpoint activation. <i>Cell Cycle</i> , 2012, 11, 3792-3800.   | 1.3 | 27        |
| 2143 | The SMS domain of Trs23p is responsible for the in vitro appearance of the TRAPP I complex in <i>Saccharomyces cerevisiae</i> . <i>Cellular Logistics</i> , 2012, 2, 28-42.  | 0.9 | 16        |
| 2144 | Sister chromatid cohesion establishment occurs in concert with lagging strand synthesis. <i>Cell Cycle</i> , 2012, 11, 2114-2121.  | 1.3 | 42        |
| 2145 | ATPase-dependent role of the atypical kinase Rio2 on the evolving pre-40S ribosomal subunit. <i>Nature Structural and Molecular Biology</i> , 2012, 19, 1316-1323.   | 3.6 | 137       |



| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2146 | The exomer cargo adaptor structure reveals a novel GTPase-binding domain. <i>EMBO Journal</i> , 2012, 31, 4191-4203.   | 3.5 | 34        |
| 2147 | The Stability of the Small Nucleolar Ribonucleoprotein (snoRNP) Assembly Protein Pih1 in <i>Saccharomyces cerevisiae</i> Is Modulated by Its C Terminus. <i>Journal of Biological Chemistry</i> , 2012, 287, 43205-43214.  | 1.6 | 21        |
| 2148 | Apn1 AP-endonuclease is essential for the repair of oxidatively damaged DNA bases in yeast frataxin-deficient cells. <i>Human Molecular Genetics</i> , 2012, 21, 4060-4072.  | 1.4 | 9         |
| 2149 | NEDD8 links cullin-RING ubiquitin ligase function to the p97 pathway. <i>Nature Structural and Molecular Biology</i> , 2012, 19, 511-516.  | 3.6 | 74        |
| 2150 | Anticheckpoint pathways at telomeres in yeast. <i>Nature Structural and Molecular Biology</i> , 2012, 19, 307-313.   | 3.6 | 78        |
| 2151 | Phosphoinositide-mediated clathrin adaptor progression at the trans-Golgi network. <i>Nature Cell Biology</i> , 2012, 14, 239-248.   | 4.6 | 140       |
| 2152 | Yeast Adaptor Protein, Nbp2p, Is Conserved Regulator of Fungal Ptc1p Phosphatases and Is Involved in Multiple Signaling Pathways. <i>Journal of Biological Chemistry</i> , 2012, 287, 22133-22141.   | 1.6 | 14        |
| 2153 | Nuclear Envelope Phosphatase 1-Regulatory Subunit 1 (Formerly TMEM188) Is the Metazoan Spo7p Ortholog and Functions in the Lipin Activation Pathway. <i>Journal of Biological Chemistry</i> , 2012, 287, 3123-3137.  | 1.6 | 86        |
| 2154 | Yeast Lipin 1 Orthologue Pah1p Regulates Vacuole Homeostasis and Membrane Fusion. <i>Journal of Biological Chemistry</i> , 2012, 287, 2221-2236.   | 1.6 | 88        |
| 2155 | The Hect Domain E3 Ligase Tom1 and the F-box Protein Dia2 Control Cdc6 Degradation in G1 Phase. <i>Journal of Biological Chemistry</i> , 2012, 287, 44212-44220.   | 1.6 | 21        |
| 2156 | The SUMO E3 Ligase Siz2 Exerts a Locus-Dependent Effect on Gene Silencing in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2012, 11, 452-462.   | 3.4 | 14        |
| 2157 | <i>YPR139c/LOA1</i> encodes a novel lysophosphatidic acid acyltransferase associated with lipid droplets and involved in TAG homeostasis. <i>Molecular Biology of the Cell</i> , 2012, 23, 233-246.  | 0.9 | 39        |
| 2158 | Essential Role of the NH <sub>2</sub> -Terminal Region of Cdc24 Guanine Nucleotide Exchange Factor in Its Initial Polarized Localization in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2012, 11, 2-15.   | 3.4 | 3         |
| 2159 | Exploring the Topology of the Gid Complex, the E3 Ubiquitin Ligase Involved in Catabolite-induced Degradation of Gluconeogenic Enzymes. <i>Journal of Biological Chemistry</i> , 2012, 287, 25602-25614.   | 1.6 | 84        |
| 2160 | The Conserved Bud20 Zinc Finger Protein Is a New Component of the Ribosomal 60S Subunit Export Machinery. <i>Molecular and Cellular Biology</i> , 2012, 32, 4898-4912.   | 1.1 | 42        |
| 2161 | Oxidative-stress-induced nuclear to cytoplasmic relocalization is required for Not4-dependent cyclin C destruction. <i>Journal of Cell Science</i> , 2012, 125, 1015-1026.   | 1.2 | 53        |
| 2162 | Novel Transcript Truncating Function of Rap1p Revealed by Synthetic Codon-Optimized Ty1 Retrotransposon. <i>Genetics</i> , 2012, 190, 523-535.   | 1.2 | 13        |
| 2163 | Target of Rapamycin Complex 2 Signals to Downstream Effector Yeast Protein Kinase 2 (Ypk2) through Adheres-Voraciously-to-Target-of-Rapamycin-2 Protein 1 (Avo1) in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2012, 287, 6089-6099. | 1.6 | 23        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2164 | Class E compartments form in response to ESCRT dysfunction in yeast due to hyperactivity of the Vps21 Rab GTPase. <i>Journal of Cell Science</i> , 2012, 125, 5208-20.   | 1.2 | 46        |
| 2165 | In Vivo Phosphorylation of Ser21 and Ser83 during Nutrient-induced Activation of the Yeast Protein Kinase A (PKA) Target Trehalase. <i>Journal of Biological Chemistry</i> , 2012, 287, 44130-44142.           | 1.6 | 64        |
| 2166 | Yeast Irc6p is a novel type of conserved clathrin coat accessory factor related to small G proteins. <i>Molecular Biology of the Cell</i> , 2012, 23, 4416-4429.   | 0.9 | 9         |
| 2167 | Mechanism of Start Site Selection by RNA Polymerase II. <i>Journal of Biological Chemistry</i> , 2012, 287, 557-567.   | 1.6 | 24        |
| 2168 | Rlp24 activates the AAA-ATPase Drg1 to initiate cytoplasmic pre-60S maturation. <i>Journal of Cell Biology</i> , 2012, 199, 771-782.   | 2.3 | 67        |
| 2169 | Mitotic exit kinase Dbf2 directly phosphorylates chitin synthase Chs2 to regulate cytokinesis in budding yeast. <i>Molecular Biology of the Cell</i> , 2012, 23, 2445-2456.                                    | 0.9 | 56        |
| 2170 | Structural basis for the assembly and nucleic acid binding of the TREX-2 transcription-export complex. <i>Nature Structural and Molecular Biology</i> , 2012, 19, 328-336.                                     | 3.6 | 90        |
| 2171 | Modular architecture of eukaryotic RNase P and RNase MRP revealed by electron microscopy. <i>Nucleic Acids Research</i> , 2012, 40, 3275-3288.   | 6.5 | 23        |
| 2172 | Hog1 Controls Global Reallocation of RNA Pol II upon Osmotic Shock in <i>Saccharomyces cerevisiae</i> . <i>G3: Genes, Genomes, Genetics</i> , 2012, 2, 1129-1136.  | 0.8 | 33        |
| 2173 | Genetic Analysis of Mps3 SUN Domain Mutants in <i>Saccharomyces cerevisiae</i> Reveals an Interaction with the SUN-Like Protein Slp1. <i>G3: Genes, Genomes, Genetics</i> , 2012, 2, 1703-1718.                | 0.8 | 32        |
| 2174 | Frameshift Mutagenesis: The Roles of Primerâ€“Template Misalignment and the Nonhomologous End-Joining Pathway in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2012, 190, 501-510.                       | 1.2 | 18        |
| 2175 | A Conserved Transcriptional Regulator Governs Fungal Morphology in Widely Diverged Species. <i>Genetics</i> , 2012, 190, 511-521.  | 1.2 | 77        |
| 2176 | Suppressors of <i>ipl1-2</i> in Components of a Glc7 Phosphatase Complex, Cdc48 AAA ATPase, TORC1, and the Kinetochore. <i>G3: Genes, Genomes, Genetics</i> , 2012, 2, 1687-1701.                              | 0.8 | 10        |
| 2177 | Two surfaces on the histone chaperone Rtt106 mediate histone binding, replication, and silencing. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, E144-53. | 3.3 | 43        |
| 2178 | Novel Attributes of Hed1 Affect Dynamics and Activity of the Rad51 Presynaptic Filament during Meiotic Recombination. <i>Journal of Biological Chemistry</i> , 2012, 287, 1566-1575.                           | 1.6 | 31        |
| 2179 | A Conserved Function for the H2A.Z C Terminus. <i>Journal of Biological Chemistry</i> , 2012, 287, 19148-19157.  | 1.6 | 32        |
| 2180 | Iron-responsive Transcription Factor Aft1 Interacts with Kinetochore Protein Iml3 and Promotes Pericentromeric Cohesin. <i>Journal of Biological Chemistry</i> , 2012, 287, 4139-4147.                         | 1.6 | 14        |
| 2181 | Structure-based Analyses Reveal Distinct Binding Sites for Atg2 and Phosphoinositides in Atg18. <i>Journal of Biological Chemistry</i> , 2012, 287, 31681-31690.   | 1.6 | 112       |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 2182 | Structure–function mapping of a heptameric module in the nuclear pore complex. <i>Journal of Cell Biology</i> , 2012, 196, 419-434.  | 2.3  | 110       |
| 2183 | Control of the mitotic exit network during meiosis. <i>Molecular Biology of the Cell</i> , 2012, 23, 3122-3132.  | 0.9  | 21        |
| 2184 | Retrotransposon profiling of RNA polymerase III initiation sites. <i>Genome Research</i> , 2012, 22, 681-692.  | 2.4  | 33        |
| 2185 | Rad26p regulates the occupancy of histone H2A–H2B dimer at the active genes in vivo. <i>Nucleic Acids Research</i> , 2012, 40, 3348-3363.  | 6.5  | 21        |
| 2186 | Overcoming natural replication barriers: differential helicase requirements. <i>Nucleic Acids Research</i> , 2012, 40, 1091-1105.  | 6.5  | 76        |
| 2187 | Quality control of MAT a1 splicing and exon skipping by nuclear RNA degradation. <i>Nucleic Acids Research</i> , 2012, 40, 1787-1796.  | 6.5  | 24        |
| 2188 | The 19S proteasome subcomplex promotes the targeting of NuA4 HAT to the promoters of ribosomal protein genes to facilitate the recruitment of TFIID for transcriptional initiation in vivo. <i>Nucleic Acids Research</i> , 2012, 40, 1969-1983. | 6.5  | 38        |
| 2189 | Flavinylation and Assembly of Succinate Dehydrogenase Are Dependent on the C-terminal Tail of the Flavoprotein Subunit. <i>Journal of Biological Chemistry</i> , 2012, 287, 40670-40679.   | 1.6  | 37        |
| 2190 | Interplay between the transcription factors acting on the GATA- and GABA-responsive elements of <i>Saccharomyces cerevisiae</i> UGA promoters. <i>Microbiology (United Kingdom)</i> , 2012, 158, 925-935.  | 0.7  | 10        |
| 2191 | Histone dosage regulates DNA damage sensitivity in a checkpoint-independent manner by the homologous recombination pathway. <i>Nucleic Acids Research</i> , 2012, 40, 9604-9620.   | 6.5  | 49        |
| 2192 | Analysis of yeast endocytic site formation and maturation through a regulatory transition point. <i>Molecular Biology of the Cell</i> , 2012, 23, 657-668.   | 0.9  | 75        |
| 2193 | PAS Kinase Promotes Cell Survival and Growth Through Activation of Rho1. <i>Science Signaling</i> , 2012, 5, ra9.  | 1.6  | 12        |
| 2194 | Mutation in the Fe–S scaffold protein Isu bypasses frataxin deletion. <i>Biochemical Journal</i> , 2012, 441, 473-480.   | 1.7  | 43        |
| 2195 | Loss of Function of the Cik1/Kar3 Motor Complex Results in Chromosomes with Syntelic Attachment That Are Sensed by the Tension Checkpoint. <i>PLoS Genetics</i> , 2012, 8, e1002492.   | 1.5  | 22        |
| 2196 | Confinement to Organelle-Associated Inclusion Structures Mediates Asymmetric Inheritance of Aggregated Protein in Budding Yeast. <i>Cell Reports</i> , 2012, 2, 738-747.   | 2.9  | 173       |
| 2197 | A Ribosome-Bound Quality Control Complex Triggers Degradation of Nascent Peptides and Signals Translation Stress. <i>Cell</i> , 2012, 151, 1042-1054.  | 13.5 | 536       |
| 2198 | The impact of protein glycosylation on Flo11-dependent adherence in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2012, 12, 809-818.  | 1.1  | 11        |
| 2199 | The Filamentous Growth MAPK Pathway Responds to Glucose Starvation Through the Mig1/2 Transcriptional Repressors in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2012, 192, 869-887.  | 1.2  | 51        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 2200 | Oxidative Stress Responses Involve Oxidation of a Conserved Ubiquitin Pathway Enzyme. <i>Molecular and Cellular Biology</i> , 2012, 32, 4472-4481.                                       | 1.1  | 25        |
| 2201 | Identification of RNA Helicase Target Sites by UV Cross-Linking and Analysis of cDNA. <i>Methods in Enzymology</i> , 2012, 511, 275-288.   | 0.4  | 56        |
| 2202 | A negative feedback loop at the nuclear periphery regulates <i>GAL</i> gene expression. <i>Molecular Biology of the Cell</i> , 2012, 23, 1367-1375.                                      | 0.9  | 50        |
| 2203 | Myosin-V Is Activated by Binding Secretory Cargo and Released in Coordination with Rab/Exocyst Function. <i>Developmental Cell</i> , 2012, 23, 769-781.                                  | 3.1  | 63        |
| 2204 | In Vivo Approaches to Dissecting the Function of RNA Helicases in Eukaryotic Ribosome Assembly. <i>Methods in Enzymology</i> , 2012, 511, 289-321.                                       | 0.4  | 1         |
| 2205 | Construction of a cytosolic firefly luciferase reporter cassette for use in PCR-mediated gene deletion and fusion in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2012, 29, 505-517. | 0.8  | 3         |
| 2206 | An improved short-lived fluorescent protein transcriptional reporter for <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2012, 29, 519-530.   | 0.8  | 48        |
| 2207 | Sic1 plays a role in timing and oscillatory behaviour of B-type cyclins. <i>Biotechnology Advances</i> , 2012, 30, 108-130.  | 6.0  | 29        |
| 2208 | Transcription of Two Long Noncoding RNAs Mediates Mating-Type Control of Gametogenesis in Budding Yeast. <i>Cell</i> , 2012, 150, 1170-1181.   | 13.5 | 235       |
| 2209 | Lis1 Acts as a "Clutch" between the ATPase and Microtubule-Binding Domains of the Dynein Motor. <i>Cell</i> , 2012, 150, 975-986.  | 13.5 | 209       |
| 2210 | Protein Modularity, Cooperative Binding, and Hybrid Regulatory States Underlie Transcriptional Network Diversification. <i>Cell</i> , 2012, 151, 80-95.                                  | 13.5 | 89        |
| 2211 | Balanced Ero1 activation and inactivation establishes ER redox homeostasis. <i>Journal of Cell Biology</i> , 2012, 196, 713-725.   | 2.3  | 65        |
| 2212 | Cargo ubiquitination is essential for multivesicular body intraluminal vesicle formation. <i>EMBO Reports</i> , 2012, 13, 331-338.   | 2.0  | 76        |
| 2213 | The structure of purified kinetochores reveals multiple microtubule-attachment sites. <i>Nature Structural and Molecular Biology</i> , 2012, 19, 925-929.                                | 3.6  | 77        |
| 2214 | Sgf29p Facilitates the Recruitment of TATA Box Binding Protein but Does Not Alter SAGA's Global Structural Integrity in Vivo. <i>Biochemistry</i> , 2012, 51, 706-714.                   | 1.2  | 19        |
| 2215 | Yar1 Protects the Ribosomal Protein Rps3 from Aggregation. <i>Journal of Biological Chemistry</i> , 2012, 287, 21806-21815.  | 1.6  | 58        |
| 2216 | Molecular basis of the functional distinction between Cln1 and Cln2 cyclins. <i>Cell Cycle</i> , 2012, 11, 3117-3131.  | 1.3  | 11        |
| 2217 | Role of Hog1, Tps1 and Sod1 in boric acid tolerance of <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> 10.1093/mic/dks071   | 0.7  | 16        |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 2218 | High-Resolution View of the Yeast Meiotic Program Revealed by Ribosome Profiling. <i>Science</i> , 2012, 335, 552-557.  | 6.0  | 496       |
| 2219 | cis - and trans -Acting Localization Determinants of pH Response Regulator Rim13 in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2012, 11, 1201-1209.   | 3.4  | 10        |
| 2220 | Ume6 transcription factor is part of a signaling cascade that regulates autophagy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 11206-11210.   | 3.3  | 100       |
| 2221 | Glucose levels regulate the nucleo-mitochondrial distribution of Mig2. <i>Mitochondrion</i> , 2012, 12, 370-380.  | 1.6  | 13        |
| 2222 | Atl1 Regulates Choice between Global Genome and Transcription-Coupled Repair of O6-Alkylguanines. <i>Molecular Cell</i> , 2012, 47, 50-60.  | 4.5  | 31        |
| 2223 | <i>Saccharomyces cerevisiae</i> Tel2 plays roles in TORC signaling and telomere maintenance that can be mutationally separated. <i>Biochemical and Biophysical Research Communications</i> , 2012, 417, 1182-1187.  | 1.0  | 3         |
| 2224 | Oxidative stress induces mitochondrial fragmentation in frataxin-deficient cells. <i>Biochemical and Biophysical Research Communications</i> , 2012, 418, 336-341.  | 1.0  | 34        |
| 2225 | GABA induction of the <i>Saccharomyces cerevisiae</i> UGA4 gene depends on the quality of the carbon source: Role of the key transcription factors acting in this process. <i>Biochemical and Biophysical Research Communications</i> , 2012, 421, 572-577. | 1.0  | 6         |
| 2226 | A versatile PCR-based tandem epitope tagging system for <i>Streptomyces coelicolor</i> genome. <i>Biochemical and Biophysical Research Communications</i> , 2012, 424, 22-27.   | 1.0  | 4         |
| 2227 | Autophagosome formation can be achieved in the absence of Atg18 by expressing engineered PAS $\alpha$ -targeted Atg2. <i>FEBS Letters</i> , 2012, 586, 2473-2478.   | 1.3  | 29        |
| 2228 | The Sec7 Arf-GEF Is Recruited to the trans-Golgi Network by Positive Feedback. <i>Developmental Cell</i> , 2012, 22, 799-810.   | 3.1  | 89        |
| 2229 | Forkhead Transcription Factors Establish Origin Timing and Long-Range Clustering in <i>S. cerevisiae</i> . <i>Cell</i> , 2012, 148, 99-111.   | 13.5 | 174       |
| 2230 | Biochemically Distinct Vesicles from the Endoplasmic Reticulum Fuse to Form Peroxisomes. <i>Cell</i> , 2012, 149, 397-409.  | 13.5 | 183       |
| 2231 | Exploring the Yeast Acetylome Using Functional Genomics. <i>Cell</i> , 2012, 149, 936-948.  | 13.5 | 63        |
| 2232 | Negative Feedback Enhances Robustness in the Yeast Polarity Establishment Circuit. <i>Cell</i> , 2012, 149, 322-333.  | 13.5 | 192       |
| 2233 | Regulation of Chaperone Effects on a Yeast Prion by Cochaperone Sgt2. <i>Molecular and Cellular Biology</i> , 2012, 32, 4960-4970.  | 1.1  | 56        |
| 2234 | Yeast importin- $\beta$ is required for nuclear import of the Mig2 repressor. <i>BMC Cell Biology</i> , 2012, 13, 31.   | 3.0  | 8         |
| 2235 | Yeast glucose pathways converge on the transcriptional regulation of trehalose biosynthesis. <i>BMC Genomics</i> , 2012, 13, 239.   | 1.2  | 30        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2236 | The <i>Saccharomyces cerevisiae</i> Hot1p regulated gene YHR087W (HG11) has a role in translation upon high glucose concentration stress. <i>BMC Molecular Biology</i> , 2012, 13, 19. | 3.0 | 20        |
| 2237 | A quantitative model of the initiation of DNA replication in <i>Saccharomyces cerevisiae</i> predicts the effects of system perturbations. <i>BMC Systems Biology</i> , 2012, 6, 78.   | 3.0 | 6         |
| 2238 | <i>De novo</i> generation of a phosphate starvation-regulated promoter in <i>Candida glabrata</i> . <i>FEMS Yeast Research</i> , 2012, 12, 980-989.                                    | 1.1 | 9         |
| 2239 | A role for Atg8â€PE deconjugation in autophagosome biogenesis. <i>Autophagy</i> , 2012, 8, 780-793.  | 4.3 | 184       |
| 2240 | Preferential Repair of DNA Double-strand Break at the Active Gene in Vivo. <i>Journal of Biological Chemistry</i> , 2012, 287, 36414-36422.  | 1.6 | 21        |
| 2241 | The histone acetyltransferase <i>Hat1</i> facilitates DNA damage repair and morphogenesis in <i>Candida albicans</i> . <i>Molecular Microbiology</i> , 2012, 86, 1197-1214.            | 1.2 | 42        |
| 2242 | Membrane Protein Rim21 Plays a Central Role in Sensing Ambient pH in <i>Saccharomyces cerevisiae</i> *. <i>Journal of Biological Chemistry</i> , 2012, 287, 38473-38481.               | 1.6 | 58        |
| 2243 | Lack of the Glc7 phosphatase regulatory subunit Ypi1 activates the morphogenetic checkpoint. <i>International Journal of Biochemistry and Cell Biology</i> , 2012, 44, 1862-1871.      | 1.2 | 6         |
| 2244 | Accumulation of linear mitochondrial DNA fragments in the nucleus shortens the chronological life span of yeast. <i>European Journal of Cell Biology</i> , 2012, 91, 782-788.          | 1.6 | 19        |
| 2245 | Studying Repair of a Single Protein-Bound Nick In Vivo Using the Flp-Nick System. <i>Methods in Molecular Biology</i> , 2012, 920, 393-415.  | 0.4 | 2         |
| 2246 | FAT10 and NUB1L bind to the VWA domain of Rpn10 and Rpn1 to enable proteasome-mediated proteolysis. <i>Nature Communications</i> , 2012, 3, 749.                                       | 5.8 | 65        |
| 2247 | Synchronizing Nuclear Import of Ribosomal Proteins with Ribosome Assembly. <i>Science</i> , 2012, 338, 666-671.  | 6.0 | 95        |
| 2248 | TULIPs: tunable, light-controlled interacting protein tags for cell biology. <i>Nature Methods</i> , 2012, 9, 379-384.   | 9.0 | 433       |
| 2249 | N-Myristoylation of the Rpt2 Subunit Regulates Intracellular Localization of the Yeast 26S Proteasome. <i>Biochemistry</i> , 2012, 51, 8856-8866.                                      | 1.2 | 46        |
| 2250 | Structure of the pre-60S ribosomal subunit with nuclear export factor Arx1 bound at the exit tunnel. <i>Nature Structural and Molecular Biology</i> , 2012, 19, 1234-1241.             | 3.6 | 103       |
| 2251 | Components of a Fanconi-Like Pathway Control Pso2-Independent DNA Interstrand Crosslink Repair in Yeast. <i>PLoS Genetics</i> , 2012, 8, e1002884.                                     | 1.5 | 41        |
| 2252 | Analysis of the Key Elements of FFAT-Like Motifs Identifies New Proteins That Potentially Bind VAP on the ER, Including Two AKAPs and FAPP2. <i>PLoS ONE</i> , 2012, 7, e30455.        | 1.1 | 77        |
| 2253 | Regulation of Mat Responses by a Differentiation MAPK Pathway in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2012, 7, e32294.  | 1.1 | 21        |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 2254 | A Deubiquitylating Complex Required for Neosynthesis of a Yeast Mitochondrial ATP Synthase Subunit. PLoS ONE, 2012, 7, e38071.  | 1.1  | 8         |
| 2255 | FRUIT, a Scar-Free System for Targeted Chromosomal Mutagenesis, Epitope Tagging, and Promoter Replacement in Escherichia coli and Salmonella enterica. PLoS ONE, 2012, 7, e44841.   | 1.1  | 73        |
| 2256 | Paralogous ALT1 and ALT2 Retention and Diversification Have Generated Catalytically Active and Inactive Aminotransferases in Saccharomyces cerevisiae. PLoS ONE, 2012, 7, e45702.   | 1.1  | 28        |
| 2257 | Apparent Defect in Yeast Bud-Site Selection Due to a Specific Failure to Splice the Pre-mRNA of a Regulator of Cell-Type-Specific Transcription. PLoS ONE, 2012, 7, e47621.   | 1.1  | 13        |
| 2258 | Identification and Functional Analysis of the erh1+ Gene Encoding Enhancer of Rudimentary Homolog from the Fission Yeast Schizosaccharomyces pombe. PLoS ONE, 2012, 7, e49059.  | 1.1  | 17        |
| 2259 | Regulation of Hxt3 and Hxt7 Turnover Converges on the Vid30 Complex and Requires Inactivation of the Ras/cAMP/PKA Pathway in Saccharomyces cerevisiae. PLoS ONE, 2012, 7, e50458.   | 1.1  | 34        |
| 2260 | The defects in cell wall integrity and G <sub>2</sub> →M transition of the <i>htl1</i> mutant are interconnected. Yeast, 2012, 29, 45-57.   | 0.8  | 5         |
| 2261 | Cassette series designed for live-cell imaging of proteins and high-resolution techniques in yeast. Yeast, 2012, 29, 119-136.   | 0.8  | 38        |
| 2262 | Interactions of Sen1, Nrd1, and Nab3 with Multiple Phosphorylated Forms of the Rpb1 C-Terminal Domain in Saccharomyces cerevisiae. Eukaryotic Cell, 2012, 11, 417-429.  | 3.4  | 46        |
| 2263 | ER Cargo Properties Specify a Requirement for COPII Coat Rigidity Mediated by Sec13p. Science, 2012, 335, 1359-1362.  | 6.0  | 124       |
| 2264 | Role of Scd5, a protein phosphatase-1 targeting protein, in phosphoregulation of Sla1 during endocytosis. Journal of Cell Science, 2012, 125, 4728-39.  | 1.2  | 16        |
| 2265 | Prion formation by a yeast GLFG nucleoporin. Prion, 2012, 6, 391-399.   | 0.9  | 74        |
| 2266 | Structure of yeast Argonaute with guide RNA. Nature, 2012, 486, 368-374.  | 18.7 | 314       |
| 2267 | Identification of RCN1 and RSA3 as ethanol-tolerant genes in Saccharomyces cerevisiae using a high copy barcoded library. FEMS Yeast Research, 2012, 12, 48-60.   | 1.1  | 17        |
| 2268 | Abasic sites linked to dUTP incorporation in DNA are a major cause of spontaneous mutations in absence of base excision repair and Rad17-Mec3-Ddc1 (9 <sup>11</sup> ) DNA damage checkpoint clamp in Saccharomyces cerevisiae. DNA Repair, 2012, 11, 294-303. | 1.3  | 12        |
| 2269 | Cyclin-dependent kinase-dependent phosphorylation of Lif1 and Sae2 controls imprecise nonhomologous end joining accompanied by double-strand break resection. Genes To Cells, 2012, 17, 473-493.  | 0.5  | 26        |
| 2270 | Peroxisome Formation Requires the Endoplasmic Reticulum Channel Protein <i>scpS</i> . Traffic, 2012, 13, 599-609.   | 1.3  | 42        |
| 2271 | Oligomerization studies show that the kinase domain of the tomato pollen receptor kinase LePRK2 is necessary for interaction with LePRK1. Plant Physiology and Biochemistry, 2012, 53, 40-45.   | 2.8  | 5         |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 2272 | Close Encounters of the Collaborative Kind. <i>Computer</i> , 2012, 45, 24-30.  | 1.2  | 6         |
| 2273 | Inositol phosphate kinase Vip1p interacts with histone chaperone Asf1p in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology Reports</i> , 2012, 39, 4989-4996.                                     | 1.0  | 4         |
| 2274 | Sulfur Amino Acids Regulate Translational Capacity and Metabolic Homeostasis through Modulation of tRNA Thiolation. <i>Cell</i> , 2013, 154, 416-429.   | 13.5 | 189       |
| 2275 | Methionine Inhibits Autophagy and Promotes Growth by Inducing the SAM-Responsive Methylation of PP2A. <i>Cell</i> , 2013, 154, 403-415.   | 13.5 | 203       |
| 2276 | Competition between Pre-mRNAs for the Splicing Machinery Drives Global Regulation of Splicing. <i>Molecular Cell</i> , 2013, 51, 338-348.   | 4.5  | 99        |
| 2277 | Mapping the functional yeast ABC transporter interactome. <i>Nature Chemical Biology</i> , 2013, 9, 565-572.  | 3.9  | 93        |
| 2278 | <i>Drosophila</i> GAGA factor polyglutamine domains exhibit prion-like behavior. <i>BMC Genomics</i> , 2013, 14, 374.   | 1.2  | 26        |
| 2279 | In vivo and in silico analysis of PCNA ubiquitylation in the activation of the Post Replication Repair pathway in <i>S. cerevisiae</i> . <i>BMC Systems Biology</i> , 2013, 7, 24.                      | 3.0  | 7         |
| 2280 | Genetic manipulation of longevity-related genes as a tool to regulate yeast life span and metabolite production during winemaking. <i>Microbial Cell Factories</i> , 2013, 12, 1.                       | 1.9  | 135       |
| 2281 | Activation of Aro80 transcription factor by heat-induced aromatic amino acid influx in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2013, 438, 43-47. | 1.0  | 11        |
| 2282 | Inactivation of Tor proteins affects the dynamics of endocytic proteins in early stage of endocytosis. <i>Journal of Biosciences</i> , 2013, 38, 351-361.   | 0.5  | 6         |
| 2283 | Mutually dependent degradation of Ama1p and Cdc20p terminates APC/C ubiquitin ligase activity at the completion of meiotic development in yeast. <i>Cell Division</i> , 2013, 8, 9.                     | 1.1  | 8         |
| 2284 | Quality Control and Substrate-Dependent Downregulation of the Nutrient Transporter Fur4. <i>Traffic</i> , 2013, 14, 412-427.  | 1.3  | 88        |
| 2285 | An Iml3-Chl4 Heterodimer Links the Core Centromere to Factors Required for Accurate Chromosome Segregation. <i>Cell Reports</i> , 2013, 5, 29-36.   | 2.9  | 56        |
| 2286 | Divergent Contributions of Conserved Active Site Residues to Transcription by Eukaryotic RNA Polymerases I and II. <i>Cell Reports</i> , 2013, 4, 974-984.  | 2.9  | 34        |
| 2287 | Mutations in SNF1 complex genes affect yeast cell wall strength. <i>European Journal of Cell Biology</i> , 2013, 92, 383-395.   | 1.6  | 31        |
| 2288 | Mutations in LYRM4, encoding iron-sulfur cluster biogenesis factor ISD11, cause deficiency of multiple respiratory chain complexes. <i>Human Molecular Genetics</i> , 2013, 22, 4460-4473.              | 1.4  | 97        |
| 2289 | Integration of Multiple Nutrient Cues and Regulation of Lifespan by Ribosomal Transcription Factor Ifh1. <i>Cell Reports</i> , 2013, 4, 1063-1071.  | 2.9  | 36        |



| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 2290 | The THO complex component Thp2 counteracts telomeric R-loops and telomere shortening. <i>EMBO Journal</i> , 2013, 32, 2861-2871.  | 3.5  | 125       |
| 2291 | Dissection of the elements of osmotic stress response transcription factor Hot1 involved in the interaction with MAPK Hog1 and in the activation of transcription. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2013, 1829, 1111-1125. | 0.9  | 15        |
| 2292 | Phosphorylation of Chs2p regulates interaction with COPII. <i>Journal of Cell Science</i> , 2013, 126, 2151-6.  | 1.2  | 14        |
| 2293 | Synthesis and study of antiproliferative, antitopoisomerase II, DNA-intercalating and DNA-damaging activities of aryl naphthalimides. <i>Bioorganic and Medicinal Chemistry</i> , 2013, 21, 6484-6495.  | 1.4  | 21        |
| 2294 | Construction of Mutant Alleles in <i>Saccharomyces cerevisiae</i> without Cloning: Overview and the Delitto Perfetto Method. <i>Current Protocols in Molecular Biology</i> , 2013, 104, 13.10C.1-13.10C.17.   | 2.9  | 9         |
| 2295 | Inositol Pyrophosphates Regulate Cell Growth and the Environmental Stress Response by Activating the HDAC Rpd3L. <i>Cell Reports</i> , 2013, 3, 1476-1482.  | 2.9  | 67        |
| 2296 | Involvement of mitochondrial dynamics in the segregation of mitochondrial matrix proteins during stationary phase mitophagy. <i>Nature Communications</i> , 2013, 4, 2789.  | 5.8  | 95        |
| 2297 | Casein Kinase 2 Associates with the Yeast Chromatin Reassembly Factor Spt2/Sin1 To Regulate Its Function in the Repression of Spurious Transcription. <i>Molecular and Cellular Biology</i> , 2013, 33, 4198-4211.  | 1.1  | 15        |
| 2298 | Key events during the transition from rapid growth to quiescence in budding yeast require posttranscriptional regulators. <i>Molecular Biology of the Cell</i> , 2013, 24, 3697-3709.   | 0.9  | 56        |
| 2299 | Placing a Disrupted Degradation Motif at the C Terminus of Proteasome Substrates Attenuates Degradation without Impairing Ubiquitylation. <i>Journal of Biological Chemistry</i> , 2013, 288, 12645-12653.  | 1.6  | 21        |
| 2300 | Translation Initiation Requires Cell Division Cycle 123 (Cdc123) to Facilitate Biogenesis of the Eukaryotic Initiation Factor 2 (eIF2). <i>Journal of Biological Chemistry</i> , 2013, 288, 21537-21546.  | 1.6  | 30        |
| 2301 | Yeast Upf1 CH domain interacts with Rps26 of the 40S ribosomal subunit. <i>Rna</i> , 2013, 19, 1105-1115.   | 1.6  | 33        |
| 2302 | Spatial sequestration of misfolded proteins by a dynamic chaperone pathway enhances cellular fitness during stress. <i>Nature Cell Biology</i> , 2013, 15, 1231-1243.   | 4.6  | 281       |
| 2303 | Cytoplasmic proteasomes are not indispensable for cell growth in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2013, 436, 372-376.   | 1.0  | 19        |
| 2304 | A Transcriptome-wide Atlas of RNP Composition Reveals Diverse Classes of mRNAs and lncRNAs. <i>Cell</i> , 2013, 154, 996-1009.  | 13.5 | 212       |
| 2305 | The Energetic State of Mitochondria Modulates Complex III Biogenesis through the ATP-Dependent Activity of Bcs1. <i>Cell Metabolism</i> , 2013, 18, 567-577.  | 7.2  | 18        |
| 2306 | Changes in Cell Morphology Are Coordinated with Cell Growth through the TORC1 Pathway. <i>Current Biology</i> , 2013, 23, 1269-1279.  | 1.8  | 38        |
| 2307 | A Histone Acetylation Switch Regulates H2A.Z Deposition by the SWR-C Remodeling Enzyme. <i>Science</i> , 2013, 340, 195-199.  | 6.0  | 168       |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 2308 | A Wee1 checkpoint inhibits anaphase onset. <i>Journal of Cell Biology</i> , 2013, 201, 843-862.   | 2.3  | 60        |
| 2309 | Characterization of fungal <i>RTG2</i> genes in retrograde signaling of <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2013, 13, 495-503.   | 1.1  | 5         |
| 2310 | Cdc48 Chaperone and Adaptor Ubx4 Distribute the Proteasome in the Nucleus for Anaphase Proteolysis. <i>Journal of Biological Chemistry</i> , 2013, 288, 37180-37191.  | 1.6  | 10        |
| 2311 | Cell-cycle and DNA damage regulation of the DNA mismatch repair protein Msh2 occurs at the transcriptional and post-transcriptional level. <i>DNA Repair</i> , 2013, 12, 97-109.  | 1.3  | 13        |
| 2312 | Effects of the Paf1 Complex and Histone Modifications on snoRNA 3' End Formation Reveal Broad and Locus-Specific Regulation. <i>Molecular and Cellular Biology</i> , 2013, 33, 170-182.   | 1.1  | 22        |
| 2313 | High levels of transcription stimulate transversions at GC base pairs in yeast. <i>Environmental and Molecular Mutagenesis</i> , 2013, 54, 44-53.   | 0.9  | 9         |
| 2314 | Mechanism-Oriented Controllability of Intracellular Quantum Dots Formation: The Role of Glutathione Metabolic Pathway. <i>ACS Nano</i> , 2013, 7, 2240-2248.  | 7.3  | 96        |
| 2315 | Regulation of mating in the budding yeast <i>Saccharomyces cerevisiae</i> by the zinc cluster proteins Sut1 and Sut2. <i>Biochemical and Biophysical Research Communications</i> , 2013, 438, 66-70.                                    | 1.0  | 7         |
| 2316 | Elevated Rad53 kinase activity influences formation and interhomolog repair of meiotic DNA double-strand breaks in budding yeast. <i>Biochemical and Biophysical Research Communications</i> , 2013, 441, 593-599.                      | 1.0  | 2         |
| 2317 | Maf1, repressor of tRNA transcription, is involved in the control of gluconeogenic genes in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 2013, 526, 16-22.   | 1.0  | 16        |
| 2318 | Tid1/Rdh54 translocase is phosphorylated through a Mec1- and Rad53-dependent manner in the presence of DSB lesions in budding yeast. <i>DNA Repair</i> , 2013, 12, 347-355.   | 1.3  | 18        |
| 2319 | Chaperone Nap1 Shields Histone Surfaces Used in a Nucleosome and Can Put H2A-H2B in an Unconventional Tetrameric Form. <i>Molecular Cell</i> , 2013, 51, 662-677.   | 4.5  | 69        |
| 2320 | A short C-terminal tail prevents mis-targeting of hydrophobic mitochondrial membrane proteins to the ER. <i>FEBS Letters</i> , 2013, 587, 3480-3486.  | 1.3  | 3         |
| 2321 | Global Analysis of Eukaryotic mRNA Degradation Reveals Xrn1-Dependent Buffering of Transcript Levels. <i>Molecular Cell</i> , 2013, 52, 52-62.  | 4.5  | 218       |
| 2322 | DNA resection proteins Sgs1 and Exo1 are required for G1 checkpoint activation in budding yeast. <i>DNA Repair</i> , 2013, 12, 751-760.   | 1.3  | 13        |
| 2323 | Role of polymerase $\gamma$ in mitochondrial mutagenesis of <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2013, 431, 270-273.  | 1.0  | 9         |
| 2324 | Endoplasmic reticulum-associated mitochondria-cortex tether functions in the distribution and inheritance of mitochondria. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, E458-67. | 3.3  | 162       |
| 2325 | A Role for the Nucleoporin Nup170p in Chromatin Structure and Gene Silencing. <i>Cell</i> , 2013, 152, 969-983.   | 13.5 | 141       |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 2326 | Mps1 and Ipl1/Aurora B Act Sequentially to Correctly Orient Chromosomes on the Meiotic Spindle of Budding Yeast. <i>Science</i> , 2013, 339, 1071-1074.  | 6.0  | 76        |
| 2327 | Genetic Networks Inducing Invasive Growth in <i>Saccharomyces cerevisiae</i> Identified Through Systematic Genome-Wide Overexpression. <i>Genetics</i> , 2013, 193, 1297-1310.   | 1.2  | 44        |
| 2328 | Tension sensing by Aurora B kinase is independent of survivin-based centromere localization. <i>Nature</i> , 2013, 497, 118-121.   | 13.7 | 129       |
| 2329 | Carotenoid-based phenotypic screen of the yeast deletion collection reveals new genes with roles in isoprenoid production. <i>Metabolic Engineering</i> , 2013, 15, 174-183.   | 3.6  | 157       |
| 2330 | Determination and physiological roles of the glycosylphosphatidylinositol lipid remodelling pathway in yeast. <i>Molecular Microbiology</i> , 2013, 88, 140-155.   | 1.2  | 27        |
| 2331 | Suppression of Chromosome Healing and Anticheckpoint Pathways in Yeast Postsenescence Survivors. <i>Genetics</i> , 2013, 194, 403-408.   | 1.2  | 0         |
| 2332 | Vps1 in the late endosome-to-vacuole traffic. <i>Journal of Biosciences</i> , 2013, 38, 73-83.   | 0.5  | 18        |
| 2333 | DNA Replication Checkpoint Signaling Depends on a Rad53-Dbf4 N-Terminal Interaction in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2013, 194, 389-401.   | 1.2  | 24        |
| 2334 | Using the cre-lox system to randomize target gene expression states and generate diverse phenotypes. <i>Biotechnology and Bioengineering</i> , 2013, 110, 2677-2686.   | 1.7  | 0         |
| 2335 | Epigenetic Silencing Mediates Mitochondria Stress-Induced Longevity. <i>Cell Metabolism</i> , 2013, 17, 954-964.   | 7.2  | 171       |
| 2336 | Ubiquitin Binding by a CUE Domain Regulates Ubiquitin Chain Formation by ERAD E3 Ligases. <i>Molecular Cell</i> , 2013, 50, 528-539.   | 4.5  | 54        |
| 2337 | Pif1 family helicases suppress genome instability at G-quadruplex motifs. <i>Nature</i> , 2013, 497, 458-462.  | 13.7 | 403       |
| 2338 | Doa4 function in ILV budding is restricted through its interaction with the Vps20 subunit of ESCRT-III. <i>Journal of Cell Science</i> , 2013, 126, 1881-90.   | 1.2  | 19        |
| 2339 | Investigation of in Vivo Roles of the C-terminal Tails of the Small Subunit ( $\beta^2$ ) of <i>Saccharomyces cerevisiae</i> Ribonucleotide Reductase. <i>Journal of Biological Chemistry</i> , 2013, 288, 13951-13959.    | 1.6  | 9         |
| 2340 | Screening for Hydrolytic Enzymes Reveals Ayr1p as a Novel Triacylglycerol Lipase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2013, 288, 36061-36072.                                     | 1.6  | 39        |
| 2341 | Heterochromatin protein Sir3 induces contacts between the amino terminus of histone H4 and nucleosomal DNA. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, 8495-8500. | 3.3  | 57        |
| 2342 | Uth1 is a mitochondrial inner membrane protein dispensable for postlog phase and rapamycin-induced mitophagy. <i>FEBS Journal</i> , 2013, 280, 4970-4982.  | 2.2  | 37        |
| 2343 | Eukaryotic Oligosaccharyltransferase Generates Free Oligosaccharides during N-Glycosylation. <i>Journal of Biological Chemistry</i> , 2013, 288, 32673-32684.  | 1.6  | 45        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2344 | Highly expressed loci are vulnerable to misleading ChIP localization of multiple unrelated proteins. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, 18602-18607.                           | 3.3 | 373       |
| 2345 | Trans-Lesion DNA Polymerases May Be Involved in Yeast Meiosis. <i>G3: Genes, Genomes, Genetics</i> , 2013, 3, 633-644.  | 0.8 | 13        |
| 2346 | Temporal regulation of the Mus81-Mms4 endonuclease ensures cell survival under conditions of DNA damage. <i>Nucleic Acids Research</i> , 2013, 41, 8943-8958.   | 6.5 | 30        |
| 2347 | Interaction Maps of the <i>Saccharomyces cerevisiae</i> ESCRT-III Protein Snf7. <i>Eukaryotic Cell</i> , 2013, 12, 1538-1546.   | 3.4 | 4         |
| 2348 | Non-FG mediated transport of the large pre-ribosomal subunit through the nuclear pore complex by the mRNA export factor Gle2. <i>Nucleic Acids Research</i> , 2013, 41, 8266-8279.  | 6.5 | 18        |
| 2349 | Has1 regulates consecutive maturation and processing steps for assembly of 60S ribosomal subunits. <i>Nucleic Acids Research</i> , 2013, 41, 7889-7904.   | 6.5 | 52        |
| 2350 | Yeast-Gene Replacement Using PCR Products. <i>Methods in Enzymology</i> , 2013, 533, 43-55.   | 0.4 | 4         |
| 2351 | The Yeast Copper Response Is Regulated by DNA Damage. <i>Molecular and Cellular Biology</i> , 2013, 33, 4041-4050.  | 1.1 | 29        |
| 2352 | Distinct roles of Rho1, Cdc42, and Cyk3 in septum formation and abscission during yeast cytokinesis. <i>Journal of Cell Biology</i> , 2013, 202, 311-329.   | 2.3 | 71        |
| 2353 | Activation of the Smk1 Mitogen-Activated Protein Kinase by Developmentally Regulated Autophosphorylation. <i>Molecular and Cellular Biology</i> , 2013, 33, 688-700.  | 1.1 | 22        |
| 2354 | Inheritance of yeast nuclear pore complexes requires the Nsp1p subcomplex. <i>Journal of Cell Biology</i> , 2013, 203, 187-196.   | 2.3 | 43        |
| 2355 | High-Throughput Genetic and Gene Expression Analysis of the RNAPII-CTD Reveals Unexpected Connections to SRB10/CDK8. <i>PLoS Genetics</i> , 2013, 9, e1003758.  | 1.5 | 21        |
| 2356 | Application of Metabolic <sup>13</sup> C Labeling in Conjunction with High-Field Nuclear Magnetic Resonance Spectroscopy for Comparative Conformational Analysis of High Mannose-Type Oligosaccharides. <i>Biomolecules</i> , 2013, 3, 108-123. | 1.8 | 37        |
| 2357 | Inhibition of the Smc5/6 Complex during Meiosis Perturbs Joint Molecule Formation and Resolution without Significantly Changing Crossover or Non-crossover Levels. <i>PLoS Genetics</i> , 2013, 9, e1003898.                                    | 1.5 | 45        |
| 2358 | The Ecm11-Gmc2 Complex Promotes Synaptonemal Complex Formation through Assembly of Transverse Filaments in Budding Yeast. <i>PLoS Genetics</i> , 2013, 9, e1003194.   | 1.5 | 93        |
| 2359 | Loss of a Conserved tRNA Anticodon Modification Perturbs Cellular Signaling. <i>PLoS Genetics</i> , 2013, 9, e1003675.  | 1.5 | 181       |
| 2360 | A Highly Redundant Gene Network Controls Assembly of the Outer Spore Wall in <i>S. cerevisiae</i> . <i>PLoS Genetics</i> , 2013, 9, e1003700.   | 1.5 | 52        |
| 2361 | Xbp1 Directs Global Repression of Budding Yeast Transcription during the Transition to Quiescence and Is Important for the Longevity and Reversibility of the Quiescent State. <i>PLoS Genetics</i> , 2013, 9, e1003854.                        | 1.5 | 64        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2362 | Insights into Diphthamide, Key Diphtheria Toxin Effector. <i>Toxins</i> , 2013, 5, 958-968.   | 1.5 | 21        |
| 2363 | Dot1-Dependent Histone H3K79 Methylation Promotes Activation of the Mek1 Meiotic Checkpoint Effector Kinase by Regulating the Hop1 Adaptor. <i>PLoS Genetics</i> , 2013, 9, e1003262.   | 1.5 | 67        |
| 2364 | An RNA-Binding Complex Involved in Ribosome Biogenesis Contains a Protein with Homology to tRNA CCA-Adding Enzyme. <i>PLoS Biology</i> , 2013, 11, e1001669.  | 2.6 | 32        |
| 2365 | Oxidative Stress and Replication-Independent DNA Breakage Induced by Arsenic in <i>Saccharomyces cerevisiae</i> . <i>PLoS Genetics</i> , 2013, 9, e1003640.   | 1.5 | 34        |
| 2366 | Rad52 Sumoylation Prevents the Toxicity of Unproductive Rad51 Filaments Independently of the Anti-Recombinase Srs2. <i>PLoS Genetics</i> , 2013, 9, e1003833.   | 1.5 | 20        |
| 2367 | Fkh1 and Fkh2 associate with Sir2 to control CLB2 transcription under normal and oxidative stress conditions. <i>Frontiers in Physiology</i> , 2013, 4, 173.  | 1.3 | 27        |
| 2368 | Reverse PCA, a Systematic Approach for Identifying Genes Important for the Physical Interaction between Protein Pairs. <i>PLoS Genetics</i> , 2013, 9, e1003838.  | 1.5 | 16        |
| 2369 | The Mub1/Ubr2 Ubiquitin Ligase Complex Regulates the Conserved Dsn1 Kinetochore Protein. <i>PLoS Genetics</i> , 2013, 9, e1003216.  | 1.5 | 29        |
| 2370 | Rpd3- and Spt16-Mediated Nucleosome Assembly and Transcriptional Regulation on Yeast Ribosomal DNA Genes. <i>Molecular and Cellular Biology</i> , 2013, 33, 2748-2759.  | 1.1 | 22        |
| 2371 | Repression of G<sub>1</sub>/S Transcription Is Mediated via Interaction of the GTB Motifs of Nrm1 and Whi5 with Swi6. <i>Molecular and Cellular Biology</i> , 2013, 33, 1476-1486.  | 1.1 | 30        |
| 2372 | Yeast hEST1A/B (SMG5/6)-Like Proteins Contribute to Environment-Sensing Adaptive Gene Expression Responses. <i>G3: Genes, Genomes, Genetics</i> , 2013, 3, 1649-1659.   | 0.8 | 4         |
| 2373 | Yeast polypeptide exit tunnel ribosomal proteins L17, L35 and L37 are necessary to recruit late-assembling factors required for 27SB pre-rRNA processing. <i>Nucleic Acids Research</i> , 2013, 41, 1965-1983.                                | 6.5 | 51        |
| 2374 | The rRNA methyltransferase Bud23 shows functional interaction with components of the SSU processome and RNase MRP. <i>Rna</i> , 2013, 19, 828-840.  | 1.6 | 31        |
| 2375 | Arf3p GTPase is a key regulator of Bud2p activation for invasive growth in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2013, 24, 2328-2339.  | 0.9 | 10        |
| 2376 | The Yeast Snt2 Protein Coordinates the Transcriptional Response to Hydrogen Peroxide-Mediated Oxidative Stress. <i>Molecular and Cellular Biology</i> , 2013, 33, 3735-3748.  | 1.1 | 53        |
| 2377 | TORC1-regulated protein kinase Npr1 phosphorylates Orm to stimulate complex sphingolipid synthesis. <i>Molecular Biology of the Cell</i> , 2013, 24, 870-881.   | 0.9 | 88        |
| 2378 | The head module of Mediator directs activation of preloaded RNAPII in vivo. <i>Nucleic Acids Research</i> , 2013, 41, 10124-10134.  | 6.5 | 8         |
| 2379 | Genome-Wide Investigation of the Role of the tRNA Nuclear-Cytoplasmic Trafficking Pathway in Regulation of the Yeast <i>Saccharomyces cerevisiae</i> Transcriptome and Proteome. <i>Molecular and Cellular Biology</i> , 2013, 33, 4241-4254. | 1.1 | 25        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2380 | Pbp1 Is Involved in Ccr4- and Khd1-Mediated Regulation of Cell Growth through Association with Ribosomal Proteins Rpl12a and Rpl12b. <i>Eukaryotic Cell</i> , 2013, 12, 864-874.   | 3.4 | 16        |
| 2381 | A developmentally regulated translational control pathway establishes the meiotic chromosome segregation pattern. <i>Genes and Development</i> , 2013, 27, 2147-2163.  | 2.7 | 90        |
| 2382 | Dissecting the roles of the histone chaperones reveals the evolutionary conserved mechanism of transcription-coupled deposition of H3.3. <i>Nucleic Acids Research</i> , 2013, 41, 5199-5209.  | 6.5 | 19        |
| 2383 | Energy metabolism regulates clathrin adaptors at the <i>trans</i> -Golgi network and endosomes. <i>Molecular Biology of the Cell</i> , 2013, 24, 832-847.  | 0.9 | 26        |
| 2384 | Comprehensive in vivo RNA-binding site analyses reveal a role of Prp8 in spliceosomal assembly. <i>Nucleic Acids Research</i> , 2013, 41, 3805-3818.   | 6.5 | 22        |
| 2385 | Targeting and functional mechanisms of the cytokinesis-related Fâ€BAR protein Hof1 during the cell cycle. <i>Molecular Biology of the Cell</i> , 2013, 24, 1305-1320.  | 0.9 | 47        |
| 2386 | A transient Î±-helical molecular recognition element in the disordered N-terminus of the Sgs1 helicase is critical for chromosome stability and binding of Top3/Rmi1. <i>Nucleic Acids Research</i> , 2013, 41, 10215-10227.         | 6.5 | 21        |
| 2387 | Loss of Vacuolar H <sup>+</sup> -ATPase (V-ATPase) Activity in Yeast Generates an Iron Deprivation Signal That Is Moderated by Induction of the Peroxiredoxin TSA2. <i>Journal of Biological Chemistry</i> , 2013, 288, 11366-11377. | 1.6 | 41        |
| 2388 | Ligand-induced activation of a forminâ€NPF pair leads to collaborative actin nucleation. <i>Journal of Cell Biology</i> , 2013, 201, 595-611.  | 2.3 | 35        |
| 2389 | The level of origin firing inversely affects the rate of replication fork progression. <i>Journal of Cell Biology</i> , 2013, 201, 373-383.  | 2.3 | 91        |
| 2390 | TEAK: Topology Enrichment Analysis framework for detecting activated biological subpathways. <i>Nucleic Acids Research</i> , 2013, 41, 1425-1437.  | 6.5 | 59        |
| 2391 | Different polarisome components play distinct roles in Slt2p-regulated cortical ER inheritance in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2013, 24, 3145-3154.                                      | 0.9 | 12        |
| 2392 | The BAH domain of Rsc2 is a histone H3 binding domain. <i>Nucleic Acids Research</i> , 2013, 41, 9168-9182.  | 6.5 | 31        |
| 2393 | Sit4p/PP6 regulates ER-to-Golgi traffic by controlling the dephosphorylation of COPII coat subunits. <i>Molecular Biology of the Cell</i> , 2013, 24, 2727-2738.   | 0.9 | 43        |
| 2394 | Biogenesis of RNA Polymerases II and III Requires the Conserved GPN Small GTPases in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2013, 193, 853-864.   | 1.2 | 45        |
| 2395 | The Zinc Cluster Protein Sut1 Contributes to Filamentation in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2013, 12, 244-253.  | 3.4 | 19        |
| 2396 | Multiple RNA interactions position Mrd1 at the site of the small subunit pseudoknot within the 90S pre-ribosome. <i>Nucleic Acids Research</i> , 2013, 41, 1178-1190.  | 6.5 | 18        |
| 2397 | The Cyclomodulin Cycle Inhibiting Factor (CIF) Alters Cullin Neddylation Dynamics. <i>Journal of Biological Chemistry</i> , 2013, 288, 14716-14726.  | 1.6 | 16        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2398 | Genome-wide haploinsufficiency screen reveals a novel role for $\hat{1}^3$ -TuSC in spindle organization and genome stability. <i>Molecular Biology of the Cell</i> , 2013, 24, 2753-2763.   | 0.9 | 11        |
| 2399 | Mechanisms of Antisense Transcription Initiation from the 3' End of the <i>GAL10</i> Coding Sequence <i>In Vivo</i> . <i>Molecular and Cellular Biology</i> , 2013, 33, 3549-3567.   | 1.1 | 20        |
| 2400 | Dual mechanisms prevent premature chromosome segregation during meiosis. <i>Genes and Development</i> , 2013, 27, 2139-2146.   | 2.7 | 34        |
| 2401 | Mediator is an intrinsic component of the basal RNA polymerase II machinery in vivo. <i>Nucleic Acids Research</i> , 2013, 41, 9651-9662.  | 6.5 | 24        |
| 2402 | Alterations in the Ure2 $\hat{1}$ Cap Domain Elicit Different GATA Factor Responses to Rapamycin Treatment and Nitrogen Limitation. <i>Journal of Biological Chemistry</i> , 2013, 288, 1841-1855.                                       | 1.6 | 23        |
| 2403 | The C Terminus of the Histone Chaperone Asf1 Cross-Links to Histone H3 in Yeast and Promotes Interaction with Histones H3 and H4. <i>Molecular and Cellular Biology</i> , 2013, 33, 605-621.   | 1.1 | 20        |
| 2404 | Rtp1p Is a Karyopherin-Like Protein Required for RNA Polymerase II Biogenesis. <i>Molecular and Cellular Biology</i> , 2013, 33, 1756-1767.  | 1.1 | 13        |
| 2405 | Las1 interacts with Grc3 polynucleotide kinase and is required for ribosome synthesis in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2013, 41, 1135-1150.  | 6.5 | 40        |
| 2406 | A novel role for Sem1 and TREX-2 in transcription involves their impact on recruitment and H2B deubiquitylation activity of SAGA. <i>Nucleic Acids Research</i> , 2013, 41, 5655-5668.   | 6.5 | 11        |
| 2407 | Immobile myosin-II plays a scaffolding role during cytokinesis in budding yeast. <i>Journal of Cell Biology</i> , 2013, 200, 271-286.  | 2.3 | 47        |
| 2408 | Functional overlap among distinct G1/S inhibitory pathways allows robust G1 arrest by yeast mating pheromones. <i>Molecular Biology of the Cell</i> , 2013, 24, 3675-3688.   | 0.9 | 9         |
| 2409 | Rab GAP cascade regulates dynamics of Ypt6 in the Golgi traffic. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, 18976-18981.  | 3.3 | 62        |
| 2410 | Insulin-induced Gene Protein (INSIG)-dependent Sterol Regulation of Hmg2 Endoplasmic Reticulum-associated Degradation (ERAD) in Yeast. <i>Journal of Biological Chemistry</i> , 2013, 288, 8519-8530.                                    | 1.6 | 22        |
| 2411 | The Yeast ATP-binding Cassette (ABC) Transporter Ycf1p Enhances the Recruitment of the Soluble SNARE Vam7p to Vacuoles for Efficient Membrane Fusion. <i>Journal of Biological Chemistry</i> , 2013, 288, 18300-18310.                   | 1.6 | 20        |
| 2412 | Activation of an Essential Calcium Signaling Pathway in <i>Saccharomyces cerevisiae</i> by Kch1 and Kch2, Putative Low-Affinity Potassium Transporters. <i>Eukaryotic Cell</i> , 2013, 12, 204-214.                                      | 3.4 | 31        |
| 2413 | The Preference for Error-Free or Error-Prone Postreplication Repair in <i>Saccharomyces cerevisiae</i> Exposed to Low-Dose Methyl Methanesulfonate Is Cell Cycle Dependent. <i>Molecular and Cellular Biology</i> , 2013, 33, 1515-1527. | 1.1 | 42        |
| 2414 | Cks1 Enhances Transcription Efficiency at the <i>GAL1</i> Locus by Linking the Paf1 Complex to the 19S Proteasome. <i>Eukaryotic Cell</i> , 2013, 12, 1192-1201.   | 3.4 | 6         |
| 2415 | Phospholipid Flippases Lem3p-Dnf1p and Lem3p-Dnf2p Are Involved in the Sorting of the Tryptophan Permease Tat2p in Yeast. <i>Journal of Biological Chemistry</i> , 2013, 288, 3594-3608.   | 1.6 | 33        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2416 | Feedback control of Swe1p degradation in the yeast morphogenesis checkpoint. <i>Molecular Biology of the Cell</i> , 2013, 24, 914-922.  | 0.9 | 19        |
| 2417 | Characterization of an M28 metalloprotease family member residing in the yeast vacuole. <i>FEMS Yeast Research</i> , 2013, 13, 471-484.   | 1.1 | 9         |
| 2418 | An expanded tool kit for the auxin-inducible degron system in budding yeast. <i>Yeast</i> , 2013, 30, 341-351.  | 0.8 | 342       |
| 2419 | Iml3p, a component of the Ctf19 complex of the budding yeast kinetochore is required to maintain kinetochore integrity under conditions of spindle stress. <i>FEMS Yeast Research</i> , 2013, 13, 375-385.                            | 1.1 | 7         |
| 2420 | Pyruvate decarboxylase activity is regulated by the Ser/Thr protein phosphatase Sit4p in the yeast <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2013, 13, 518-528.  | 1.1 | 19        |
| 2421 | Interplay of Aro80 and GATA activators in regulation of genes for catabolism of aromatic amino acids in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2013, 88, 1120-1134.  | 1.2 | 52        |
| 2422 | Mitotic phosphorylation of Exo84 disrupts exocyst assembly and arrests cell growth. <i>Journal of Cell Biology</i> , 2013, 202, 97-111.   | 2.3 | 27        |
| 2423 | Mod5 protein binds to tRNA gene complexes and affects local transcriptional silencing. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, E3081-9.                                   | 3.3 | 33        |
| 2424 | mChIP-KAT-MS, a method to map protein interactions and acetylation sites for lysine acetyltransferases. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, E1641-50.                 | 3.3 | 38        |
| 2425 | DNA damage checkpoint triggers autophagy to regulate the initiation of anaphase. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, E41-9.   | 3.3 | 59        |
| 2426 | Regulation of the Yeast Triacylglycerol Lipase Tgl3p by Formation of Nonpolar Lipids. <i>Journal of Biological Chemistry</i> , 2013, 288, 19939-19948.  | 1.6 | 38        |
| 2427 | Functional Analysis of Bre1p, an E3 Ligase for Histone H2B Ubiquitylation, in Regulation of RNA Polymerase II Association with Active Genes and Transcription in Vivo. <i>Journal of Biological Chemistry</i> , 2013, 288, 9619-9633. | 1.6 | 18        |
| 2428 | Kch1 Family Proteins Mediate Essential Responses to Endoplasmic Reticulum Stresses in the Yeasts <i>Saccharomyces cerevisiae</i> and <i>Candida albicans</i> . <i>Journal of Biological Chemistry</i> , 2013, 288, 34861-34870.       | 1.6 | 14        |
| 2429 | An E2 enzyme Ubc11 is required for ubiquitination of Slp1/Cdc20 and spindle checkpoint silencing in fission yeast. <i>Cell Cycle</i> , 2013, 12, 961-971.   | 1.3 | 7         |
| 2430 | Sequential primed kinases create a damage-responsive phosphodegron on Eco1. <i>Nature Structural and Molecular Biology</i> , 2013, 20, 194-201.   | 3.6 | 70        |
| 2431 | Formation and dissociation of proteasome storage granules are regulated by cytosolic pH. <i>Journal of Cell Biology</i> , 2013, 201, 663-671.   | 2.3 | 76        |
| 2432 | Identification of the binding site of Rlp7 on assembling 60S ribosomal subunits in <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 2013, 19, 1639-1647.  | 1.6 | 23        |
| 2433 | Identification of a second amphipathic lipid-packing sensor-like motif that contributes to Gcs1p function in the early endosome-to-TGN pathway. <i>Journal of Biochemistry</i> , 2013, 153, 573-587.                                  | 0.9 | 4         |



| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2434 | Break-induced replication occurs by conservative DNA synthesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, 13475-13480.  | 3.3 | 156       |
| 2435 | Functional Analysis of Rad14p, a DNA Damage Recognition Factor in Nucleotide Excision Repair, in Regulation of Transcription in Vivo. <i>Journal of Biological Chemistry</i> , 2013, 288, 793-806.                        | 1.6 | 9         |
| 2436 | Ras/cAMP-dependent Protein Kinase (PKA) Regulates Multiple Aspects of Cellular Events by Phosphorylating the Whi3 Cell Cycle Regulator in Budding Yeast. <i>Journal of Biological Chemistry</i> , 2013, 288, 10558-10566. | 1.6 | 23        |
| 2437 | The pre-mRNA retention and splicing complex controls tRNA maturation by promoting TAN1 expression. <i>Nucleic Acids Research</i> , 2013, 41, 5669-5678.   | 6.5 | 26        |
| 2438 | Synergistic effects of TOR and proteasome pathways on the yeast transcriptome and cell growth. <i>Open Biology</i> , 2013, 3, 120137.   | 1.5 | 21        |
| 2439 | New vectors for epitope tagging and gene disruption in <i>Schizosaccharomyces pombe</i> . <i>BioTechniques</i> , 2013, 55, 257-263.   | 0.8 | 16        |
| 2440 | The Prefoldin Bud27 Mediates the Assembly of the Eukaryotic RNA Polymerases in an Rpb5-Dependent Manner. <i>PLoS Genetics</i> , 2013, 9, e1003297.  | 1.5 | 69        |
| 2441 | The Anaphase Promoting Complex Contributes to the Degradation of the <i>S. cerevisiae</i> Telomerase Recruitment Subunit Est1p. <i>PLoS ONE</i> , 2013, 8, e55055.  | 1.1 | 14        |
| 2442 | The SWI/SNF Chromatin Remodeling Complex Influences Transcription by RNA Polymerase I in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2013, 8, e56793.   | 1.1 | 17        |
| 2443 | Centromere Binding and a Conserved Role in Chromosome Stability for SUMO-Dependent Ubiquitin Ligases. <i>PLoS ONE</i> , 2013, 8, e65628.  | 1.1 | 29        |
| 2444 | Msb1 Interacts with Cdc42, Boi1, and Boi2 and May Coordinate Cdc42 and Rho1 Functions during Early Stage of Bud Development in Budding Yeast. <i>PLoS ONE</i> , 2013, 8, e66321.  | 1.1 | 9         |
| 2445 | Integrity of the <i>Saccharomyces cerevisiae</i> Rpn11 Protein Is Critical for Formation of Proteasome Storage Granules (PSC) and Survival in Stationary Phase. <i>PLoS ONE</i> , 2013, 8, e70357.                        | 1.1 | 28        |
| 2446 | Role of Endocytosis in Localization and Maintenance of the Spatial Markers for Bud-Site Selection in Yeast. <i>PLoS ONE</i> , 2013, 8, e72123.  | 1.1 | 8         |
| 2447 | The FEAR Protein Slk19 Restricts Cdc14 Phosphatase to the Nucleus until the End of Anaphase, Regulating Its Participation in Mitotic Exit in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2013, 8, e73194.         | 1.1 | 5         |
| 2448 | Mutations in Non-Acid Patch Residues Disrupt H2A.Z's Association with Chromatin through Multiple Mechanisms. <i>PLoS ONE</i> , 2013, 8, e76394.   | 1.1 | 6         |
| 2449 | Cell Cycle Regulated Interaction of a Yeast Hippo Kinase and Its Activator MO25/Hym1. <i>PLoS ONE</i> , 2013, 8, e78334.  | 1.1 | 11        |
| 2450 | The Role of Ctk1 Kinase in Termination of Small Non-Coding RNAs. <i>PLoS ONE</i> , 2013, 8, e80495.   | 1.1 | 15        |
| 2451 | A Rad53 Independent Function of Rad9 Becomes Crucial for Genome Maintenance in the Absence of the RecQ Helicase Sgs1. <i>PLoS ONE</i> , 2013, 8, e81015.  | 1.1 | 13        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2452 | Analysis of the Tolerance to DNA Alkylating Damage in MEC1 and RAD53 Checkpoint Mutants of <i>Saccharomyces cerevisiae</i> . PLoS ONE, 2013, 8, e81108.   | 1.1 | 5         |
| 2453 | Mak5 and Ebp2 Act Together on Early Pre-60S Particles and Their Reduced Functionality Bypasses the Requirement for the Essential Pre-60S Factor Nsa1. PLoS ONE, 2013, 8, e82741.  | 1.1 | 29        |
| 2454 | Whi3, an <i>S. cerevisiae</i> RNA-Binding Protein, Is a Component of Stress Granules That Regulates Levels of Its Target mRNAs. PLoS ONE, 2013, 8, e84060.  | 1.1 | 17        |
| 2455 | Trx2p-dependent Regulation of <i>Saccharomyces cerevisiae</i> Oxidative Stress Response by the Skn7p Transcription Factor under Respiring Conditions. PLoS ONE, 2013, 8, e85404.  | 1.1 | 3         |
| 2456 | The Cell Wall Sensors Mtl1, Wsc1, and Mid2 Are Required for Stress-Induced Nuclear to Cytoplasmic Translocation of Cyclin C and Programmed Cell Death in Yeast. <i>Oxidative Medicine and Cellular Longevity</i> , 2013, 2013, 1-15.  | 1.9 | 23        |
| 2457 | Extending the <i>Schizosaccharomyces pombe</i> Molecular Genetic Toolbox. PLoS ONE, 2014, 9, e97683.  | 1.1 | 51        |
| 2458 | Cohesin without Cohesion: A Novel Role for Pds5 in <i>Saccharomyces cerevisiae</i> . PLoS ONE, 2014, 9, e100470.  | 1.1 | 27        |
| 2459 | SPO24 Is a Transcriptionally Dynamic, Small ORF-Encoding Locus Required for Efficient Sporulation in <i>Saccharomyces cerevisiae</i> . PLoS ONE, 2014, 9, e105058.  | 1.1 | 6         |
| 2460 | The GCKIII Kinase Sps1 and the 14-3-3 Isoforms, Bmh1 and Bmh2, Cooperate to Ensure Proper Sporulation in <i>Saccharomyces cerevisiae</i> . PLoS ONE, 2014, 9, e113528.  | 1.1 | 22        |
| 2461 | Ribosome Profiling Reveals Pervasive Translation Outside of Annotated Protein-Coding Genes. <i>Cell Reports</i> , 2014, 8, 1365-1379.   | 2.9 | 591       |
| 2462 | Reciprocal conversion of Gtr1 and Gtr2 nucleotide-binding states by Npr2-Npr3 inactivates TORC1 and induces autophagy. <i>Autophagy</i> , 2014, 10, 1565-1578.  | 4.3 | 58        |
| 2463 | Conserved electron donor complex Dre2-Tah18 is required for ribonucleotide reductase metallocofactor assembly and DNA synthesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, E1695-704.                               | 3.3 | 40        |
| 2464 | Rad9 interacts with Aft1 to facilitate genome surveillance in fragile genomic sites under non-DNA damage-inducing conditions in <i>S. cerevisiae</i> . <i>Nucleic Acids Research</i> , 2014, 42, 12650-12667.   | 6.5 | 19        |
| 2465 | Molecular genetic techniques for gene manipulation in <i>Candida albicans</i> . <i>Virulence</i> , 2014, 5, 507-520.  | 1.8 | 16        |
| 2466 | Intramolecular Disulfide Bond of Tim22 Protein Maintains Integrity of the TIM22 Complex in the Mitochondrial Inner Membrane. <i>Journal of Biological Chemistry</i> , 2014, 289, 4827-4838.   | 1.6 | 27        |
| 2467 | Glucose Activates TORC2-Gad8 Protein via Positive Regulation of the cAMP/cAMP-dependent Protein Kinase A (PKA) Pathway and Negative Regulation of the Pmk1 Protein-Mitogen-activated Protein Kinase Pathway. <i>Journal of Biological Chemistry</i> , 2014, 289, 21727-21737. | 1.6 | 35        |
| 2468 | Integrity of SRP RNA is ensured by La and the nuclear RNA quality control machinery. <i>Nucleic Acids Research</i> , 2014, 42, 10698-10710.   | 6.5 | 7         |
| 2469 | Genetic Architecture of Ethanol-Responsive Transcriptome Variation in <i>Saccharomyces cerevisiae</i> Strains. <i>Genetics</i> , 2014, 198, 369-382.  | 1.2 | 58        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2470 | Signaling Events of the Rim101 Pathway Occur at the Plasma Membrane in a Ubiquitination-Dependent Manner. <i>Molecular and Cellular Biology</i> , 2014, 34, 3525-3534.                              | 1.1 | 42        |
| 2471 | Yeast histone H3 lysine 4 demethylase Jhd2 regulates mitotic ribosomal DNA condensation. <i>BMC Biology</i> , 2014, 12, 75.   | 1.7 | 29        |
| 2472 | Chd1 co-localizes with early transcription elongation factors independently of H3K36 methylation and releases stalled RNA polymerase II at introns. <i>Epigenetics and Chromatin</i> , 2014, 7, 32. | 1.8 | 25        |
| 2473 | <i>N</i> -Acetylglucosamine (GlcNAc)-Inducible Gene <i>GIG2</i> Is a Novel Component of GlcNAc Metabolism in <i>Candida albicans</i> . <i>Eukaryotic Cell</i> , 2014, 13, 66-76.                    | 3.4 | 9         |
| 2474 | The C-terminal domains of human neurofibromin and its budding yeast homologs Ira1 and Ira2 regulate the metaphase to anaphase transition. <i>Cell Cycle</i> , 2014, 13, 2780-2789.                  | 1.3 | 23        |
| 2475 | YCL047C/POF1 Is a Novel Nicotinamide Mononucleotide Adenylyltransferase (NMNAT) in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2014, 289, 15577-15587.               | 1.6 | 23        |
| 2476 | Sgo1 recruits PP2A to chromosomes to ensure sister chromatid bi-orientation in mitosis. <i>Journal of Cell Science</i> , 2014, 127, 4974-83.  | 1.2 | 39        |
| 2477 | A cluster of methylations in the domain IV of 25S rRNA is required for ribosome stability. <i>Rna</i> , 2014, 20, 1632-1644.  | 1.6 | 64        |
| 2478 | Autoregulation of the <i>Kluyveromyces lactis</i> pyruvate decarboxylase gene KIPDC1 involves the regulatory gene RAG3. <i>Microbiology (United Kingdom)</i> , 2014, 160, 1369-1378.                | 0.7 | 3         |
| 2479 | The crystal structure of <i>S. cerevisiae</i> Sad1, a catalytically inactive deubiquitinase that is broadly required for pre-mRNA splicing. <i>Rna</i> , 2014, 20, 656-669.                         | 1.6 | 30        |
| 2480 | The Yeast Sks1p Kinase Signaling Network Regulates Pseudohyphal Growth and Glucose Response. <i>PLoS Genetics</i> , 2014, 10, e1004183.   | 1.5 | 28        |
| 2481 | Down-Regulation of Rad51 Activity during Meiosis in Yeast Prevents Competition with Dmc1 for Repair of Double-Strand Breaks. <i>PLoS Genetics</i> , 2014, 10, e1004005.                             | 1.5 | 53        |
| 2482 | Rrp12 and the Exportin Crm1 Participate in Late Assembly Events in the Nucleolus during 40S Ribosomal Subunit Biogenesis. <i>PLoS Genetics</i> , 2014, 10, e1004836.                                | 1.5 | 17        |
| 2483 | Eaf5/7/3 form a functionally independent NuA4 submodule linked to <i>scp</i> RNA polymerase <i>scp</i> II-coupled nucleosome recycling. <i>EMBO Journal</i> , 2014, 33, 1397-1415.                  | 3.5 | 55        |
| 2484 | Pooled Segregant Sequencing Reveals Genetic Determinants of Yeast Pseudohyphal Growth. <i>PLoS Genetics</i> , 2014, 10, e1004570.   | 1.5 | 24        |
| 2485 | Stress-Induced Nuclear RNA Degradation Pathways Regulate Yeast Bromodomain Factor 2 to Promote Cell Survival. <i>PLoS Genetics</i> , 2014, 10, e1004661.  | 1.5 | 18        |
| 2486 | Controlling Meiotic Recombinational Repair – Specifying the Roles of ZMMs, Sgs1 and Mus81/Mms4 in Crossover Formation. <i>PLoS Genetics</i> , 2014, 10, e1004690.                                   | 1.5 | 55        |
| 2487 | <i>Msp1</i> maintains mitochondrial function by facilitating the degradation of mislocalized tail-anchored proteins. <i>EMBO Journal</i> , 2014, 33, 1548-1564.                                     | 3.5 | 172       |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2488 | Accumulation of a Threonine Biosynthetic Intermediate Attenuates General Amino Acid Control by Accelerating Degradation of Gcn4 via Pho85 and Cdk8. <i>PLoS Genetics</i> , 2014, 10, e1004534.                                    | 1.5 | 11        |
| 2489 | A Neurotoxic Glycerophosphocholine Impacts PtdIns-4, 5-Bisphosphate and TORC2 Signaling by Altering Ceramide Biosynthesis in Yeast. <i>PLoS Genetics</i> , 2014, 10, e1004010.  | 1.5 | 4         |
| 2490 | Positive Feedback of NDT80 Expression Ensures Irreversible Meiotic Commitment in Budding Yeast. <i>PLoS Genetics</i> , 2014, 10, e1004398.  | 1.5 | 39        |
| 2491 | The telomerase inhibitor Gno1p/PINX1 activates the helicase Prp43p during ribosome biogenesis. <i>Nucleic Acids Research</i> , 2014, 42, 7330-7345.   | 6.5 | 41        |
| 2492 | A Novel Role for Cet1p mRNA 5â€²-Triphosphatase in Promoter Proximal Accumulation of RNA Polymerase II in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2014, 196, 161-176.   | 1.2 | 13        |
| 2493 | Licensing of Yeast Centrosome Duplication Requires Phosphoregulation of Sfi1. <i>PLoS Genetics</i> , 2014, 10, e1004666.  | 1.5 | 37        |
| 2494 | Comparative genetic analysis of PP2A-Cdc55 regulators in budding yeast. <i>Cell Cycle</i> , 2014, 13, 2073-2083.  | 1.3 | 9         |
| 2495 | A Method for Sporulating Budding Yeast Cells That Allows for Unbiased Identification of Kinase Substrates Using Stable Isotope Labeling by Amino Acids in Cell Culture. <i>G3: Genes, Genomes, Genetics</i> , 2014, 4, 2125-2135. | 0.8 | 12        |
| 2496 | Counting Molecules Within Cells. <i>Colloquium Series on Quantitative Cell Biology</i> , 2014, 1, 1-74.   | 0.5 | 4         |
| 2497 | Divergence in a master variator generates distinct phenotypes and transcriptional responses. <i>Genes and Development</i> , 2014, 28, 409-421.  | 2.7 | 19        |
| 2498 | Tracking of the dynamic localization of the Rab-specific HOPS subunits reveal their distinct interaction with Ypt7 and vacuoles. <i>Cellular Logistics</i> , 2014, 4, e29191.   | 0.9 | 15        |
| 2499 | Assessing Regulated Nuclear Transport in <i>Saccharomyces cerevisiae</i> . <i>Methods in Cell Biology</i> , 2014, 122, 311-330.   | 0.5 | 3         |
| 2500 | Localization and Function of Budding Yeast CENP-A Depends upon Kinetochore Protein Interactions and Is Independent of Canonical Centromere Sequence. <i>Cell Reports</i> , 2014, 9, 2027-2033.                                    | 2.9 | 20        |
| 2501 | Identification of Candidate Substrates for the Golgi Tul1 E3 Ligase Using Quantitative diGly Proteomics in Yeast. <i>Molecular and Cellular Proteomics</i> , 2014, 13, 2871-2882.   | 2.5 | 32        |
| 2502 | Osh4p is needed to reduce the level of phosphatidylinositol-4-phosphate on secretory vesicles as they mature. <i>Molecular Biology of the Cell</i> , 2014, 25, 3389-3400.   | 0.9 | 47        |
| 2503 | A selfish DNA element engages a meiosis-specific motor and telomeres for germ-line propagation. <i>Journal of Cell Biology</i> , 2014, 205, 643-661.  | 2.3 | 15        |
| 2504 | Proline Biosynthesis Is Required for Endoplasmic Reticulum Stress Tolerance in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2014, 289, 27794-27806.   | 1.6 | 37        |
| 2505 | ER-phagy mediates selective degradation of endoplasmic reticulum independently of the core autophagy machinery. <i>Journal of Cell Science</i> , 2014, 127, 4078-88.  | 1.2 | 221       |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2506 | Bifurcation of the endocytic pathway into Rab5-dependent and -independent transport to the vacuole. <i>Nature Communications</i> , 2014, 5, 3498.  | 5.8 | 27        |
| 2507 | Primers4Yeast: a comprehensive web tool for planning primers for <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2014, 31, 77-80.   | 0.8 | 41        |
| 2508 | Improper protein trafficking contributes to artemisinin sensitivity in cells lacking the KDAC Rpd3p. <i>FEBS Letters</i> , 2014, 588, 4018-4025.   | 1.3 | 10        |
| 2509 | Asymmetric distribution of phosphatidylserine is generated in the absence of phospholipid flippases in <i>Saccharomyces cerevisiae</i> . <i>MicrobiologyOpen</i> , 2014, 3, 803-821.   | 1.2 | 14        |
| 2510 | Loss of Ypk1, the Yeast Homolog to the Human Serum- and Glucocorticoid-induced Protein Kinase, Accelerates Phospholipase B1-mediated Phosphatidylcholine Deacylation. <i>Journal of Biological Chemistry</i> , 2014, 289, 31591-31604. | 1.6 | 15        |
| 2511 | Tor $\epsilon$ 3 deficiency activates catabolism of the ketone body-like acetic acid to promote trehalose accumulation and longevity. <i>Aging Cell</i> , 2014, 13, 457-467.   | 3.0 | 48        |
| 2512 | Fpk1/2 kinases regulate cellular sphingoid long-chain base abundance and alter cellular resistance to LCB elevation or depletion. <i>MicrobiologyOpen</i> , 2014, 3, 196-212.  | 1.2 | 12        |
| 2513 | Loss of Vacuolar H <sup>+</sup> -ATPase Activity in Organelles Signals Ubiquitination and Endocytosis of the Yeast Plasma Membrane Proton pump Pma1p. <i>Journal of Biological Chemistry</i> , 2014, 289, 32316-32326.                 | 1.6 | 39        |
| 2514 | NuA4 Links Methylation of Histone H3 Lysines 4 and 36 to Acetylation of Histones H4 and H3. <i>Journal of Biological Chemistry</i> , 2014, 289, 32656-32670.   | 1.6 | 30        |
| 2515 | Yeast Endocytic Adaptor AP $\epsilon$ 2 Binds the Stress Sensor Mid2 and Functions in Polarized Cell Responses. <i>Traffic</i> , 2014, 15, 546-557.  | 1.3 | 31        |
| 2516 | Regulation of Mitotic Spindle Disassembly by an Environmental Stress-Sensing Pathway in Budding Yeast. <i>Genetics</i> , 2014, 198, 1043-1057.   | 1.2 | 11        |
| 2517 | Co-transcriptional recruitment of Puf6 by She2 couples translational repression to mRNA localization. <i>Nucleic Acids Research</i> , 2014, 42, 8692-8704.   | 6.5 | 22        |
| 2518 | A conserved and essential basic region mediates tRNA binding to the Elp1 subunit of the Elongator complex. <i>Molecular Microbiology</i> , 2014, 92, 1227-1242.  | 1.2 | 18        |
| 2519 | Glutathione is essential to preserve nuclear function and cell survival under oxidative stress. <i>Free Radical Biology and Medicine</i> , 2014, 75, S25-S26.  | 1.3 | 17        |
| 2520 | Shared Genetic Pathways Contribute to the Tolerance of Endogenous and Low-Dose Exogenous DNA Damage in Yeast. <i>Genetics</i> , 2014, 198, 519-530.  | 1.2 | 6         |
| 2521 | Efficient chromosomal gene modification with CRISPR/cas9 and PCR-based homologous recombination donors in cultured <i>Drosophila</i> cells. <i>Nucleic Acids Research</i> , 2014, 42, e89-e89.   | 6.5 | 110       |
| 2522 | The budding yeast Polo-like kinase Cdc5 is released from the nucleus during anaphase for timely mitotic exit. <i>Cell Cycle</i> , 2014, 13, 3260-3270.   | 1.3 | 15        |
| 2523 | H2B ubiquitylation modulates spliceosome assembly and function in budding yeast. <i>Biology of the Cell</i> , 2014, 106, 126-138.  | 0.7 | 19        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2524 | Binding to Any <sc>ESCRT</sc> Can Mediate Ubiquitinâ€independent Cargo Sorting. <i>Traffic</i> , 2014, 15, 212-229.   | 1.3 | 17        |
| 2525 | Eukaryotic translation initiation factor eIF5 promotes the accuracy of start codon recognition by regulating Pi release and conformational transitions of the preinitiation complex. <i>Nucleic Acids Research</i> , 2014, 42, 9623-9640. | 6.5 | 30        |
| 2526 | Rrd1p, an RNA polymerase II-specific prolyl isomerase and activator of phosphoprotein phosphatase, promotes transcription independently of rapamycin response. <i>Nucleic Acids Research</i> , 2014, 42, 9892-9907.                       | 6.5 | 11        |
| 2527 | Compositional and structural analysis of selected chromosomal domains from <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2014, 42, e2-e2.   | 6.5 | 25        |
| 2528 | Med13p prevents mitochondrial fission and programmed cell death in yeast through nuclear retention of cyclin C. <i>Molecular Biology of the Cell</i> , 2014, 25, 2807-2816.   | 0.9 | 39        |
| 2529 | Secretion of a foreign protein from budding yeasts is enhanced by cotranslational translocation and by suppression of vacuolar targeting. <i>Microbial Cell Factories</i> , 2014, 13, 125.  | 1.9 | 93        |
| 2530 | Dissecting Ribosome Assembly and Transport in Budding Yeast. <i>Methods in Cell Biology</i> , 2014, 122, 437-461.   | 0.5 | 16        |
| 2531 | Deficiency of the exportomer components Pex1, Pex6, and Pex15 causes enhanced pexophagy in <i>Saccharomyces cerevisiae</i> . <i>Autophagy</i> , 2014, 10, 835-845.  | 4.3 | 58        |
| 2532 | A Highly Conserved Region within H2B Is Important for FACT To Act on Nucleosomes. <i>Molecular and Cellular Biology</i> , 2014, 34, 303-314.  | 1.1 | 32        |
| 2533 | OM14 is a mitochondrial receptor for cytosolic ribosomes that supports co-translational import into mitochondria. <i>Nature Communications</i> , 2014, 5, 5711.   | 5.8 | 106       |
| 2534 | Cotranscriptional recruitment of yeast TRAMP complex to intronic sequences promotes optimal pre-mRNA splicing. <i>Nucleic Acids Research</i> , 2014, 42, 643-660.   | 6.5 | 403       |
| 2535 | Brr2p carboxy-terminal Sec63 domain modulates Prp16 splicing RNA helicase. <i>Nucleic Acids Research</i> , 2014, 42, 13897-13910.   | 6.5 | 13        |
| 2536 | Defective histone supply causes condensin-dependent chromatin alterations, SAC activation and chromosome decatenation impairment. <i>Nucleic Acids Research</i> , 2014, 42, 12469-12482.  | 6.5 | 13        |
| 2537 | Separate responses of karyopherins to glucose and amino acid availability regulate nucleocytoplasmic transport. <i>Molecular Biology of the Cell</i> , 2014, 25, 2840-2852.   | 0.9 | 24        |
| 2538 | Ribosome Assembly Factors Pwp1 and Nop12 Are Important for Folding of 5.8S rRNA during Ribosome Biogenesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2014, 34, 1863-1877.                              | 1.1 | 33        |
| 2539 | Global Regulation of a Differentiation MAPK Pathway in Yeast. <i>Genetics</i> , 2014, 198, 1309-1328.   | 1.2 | 33        |
| 2540 | The Î²-1,3-glucanosyltransferase Gas1 regulates Sir2-mediated rDNA stability in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2014, 42, 8486-8499.  | 6.5 | 17        |
| 2541 | Length-dependent processing of telomeres in the absence of telomerase. <i>Nucleic Acids Research</i> , 2014, 42, 3648-3665.   | 6.5 | 63        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2542 | The signaling lipid PI(3,5)P <sub>2</sub> stabilizes V <sub>1</sub> V <sub>o</sub> sector interactions and activates the V-ATPase. <i>Molecular Biology of the Cell</i> , 2014, 25, 1251-1262.  | 0.9 | 117       |
| 2543 | Acetylation of the Transcriptional Repressor Ume6p Allows Efficient Promoter Release and Timely Induction of the Meiotic Transient Transcription Program in Yeast. <i>Molecular and Cellular Biology</i> , 2014, 34, 631-642.   | 1.1 | 22        |
| 2544 | The analysis of <i>S. cerevisiae</i> cells deleted for mitotic cyclin Clb2 reveals a novel requirement of Sgs1 DNA helicase and Exonuclease 1 when replication forks break in the presence of alkylation damage. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2014, 769, 80-92. | 0.4 | 5         |
| 2545 | Suppression of Expression Between Adjacent Genes Within Heterologous Modules in Yeast. <i>G3: Genes, Genomes, Genetics</i> , 2014, 4, 109-116.  | 0.8 | 12        |
| 2546 | Npr2 inhibits TORC1 to prevent inappropriate utilization of glutamine for biosynthesis of nitrogen-containing metabolites. <i>Science Signaling</i> , 2014, 7, ra120.   | 1.6 | 42        |
| 2547 | Assays for Autophagy II: Mitochondrial Autophagy. <i>Methods in Molecular Biology</i> , 2014, 1163, 165-173.  | 0.4 | 3         |
| 2548 | Nucleolar proteins Bfr2 and Enp2 interact with DEAD-box RNA helicase Dbp4 in two different complexes. <i>Nucleic Acids Research</i> , 2014, 42, 3194-3206.  | 6.5 | 26        |
| 2549 | Conditional Genetic Interactions of <i>RTT107</i> , <i>SLX4</i> , and <i>HRQ1</i> Reveal Dynamic Networks upon DNA Damage in <i>S. cerevisiae</i> . <i>G3: Genes, Genomes, Genetics</i> , 2014, 4, 1059-1069.   | 0.8 | 13        |
| 2550 | Immuno- and Correlative Light Microscopy- Electron Tomography Methods for 3D Protein Localization in Yeast. <i>Traffic</i> , 2014, 15, 1164-1178.   | 1.3 | 17        |
| 2551 | tRNA <sup>2</sup> processing in yeast involves tRNase Z, Rex1, and Rrp6. <i>Rna</i> , 2014, 20, 115-130.  | 1.6 | 37        |
| 2552 | A SUMO-targeted ubiquitin ligase is involved in the degradation of the nuclear pool of the SUMO E3 ligase Siz1. <i>Molecular Biology of the Cell</i> , 2014, 25, 1-16.  | 0.9 | 37        |
| 2553 | Slr2p phosphorylation induces cyclin C nuclear-to-cytoplasmic translocation in response to oxidative stress. <i>Molecular Biology of the Cell</i> , 2014, 25, 1396-1407.  | 0.9 | 37        |
| 2554 | A comprehensive protein-protein interactome for yeast PAS kinase 1 reveals direct inhibition of respiration through the phosphorylation of Cbf1. <i>Molecular Biology of the Cell</i> , 2014, 25, 2199-2215.  | 0.9 | 13        |
| 2555 | Role of Phosphatidylserine in Phospholipid Flippase-Mediated Vesicle Transport in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2014, 13, 363-375.   | 3.4 | 28        |
| 2556 | The function of ORAOV1/LTO1, a gene that is overexpressed frequently in cancer: essential roles in the function and biogenesis of the ribosome. <i>Oncogene</i> , 2014, 33, 484-494.  | 2.6 | 36        |
| 2557 | Semi-quantitative colony immunoassay for determining and optimizing protein expression in <i>Saccharomyces cerevisiae</i> and <i>Escherichia coli</i> . <i>Analytical Biochemistry</i> , 2014, 447, 82-89.  | 1.1 | 2         |
| 2558 | An ER Protein Functionally Couples Neutral Lipid Metabolism on Lipid Droplets to Membrane Lipid Synthesis in the ER. <i>Cell Reports</i> , 2014, 6, 44-55.  | 2.9 | 99        |
| 2559 | Nuclear FKBP, Fpr3 and Fpr4 affect genome-wide genes transcription. <i>Molecular Genetics and Genomics</i> , 2014, 289, 125-136.  | 1.0 | 8         |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 2560 | A role for the proteasome in the turnover of <i>Scp5p</i> and in [ <i>PSI</i> <sup>+</sup> ] prion propagation. <i>Molecular Microbiology</i> , 2014, 92, 507-528.   | 1.2  | 17        |
| 2561 | Yeast Nkp2 is required for accurate chromosome segregation and interacts with several components of the central kinetochore. <i>Molecular Biology Reports</i> , 2014, 41, 787-797.   | 1.0  | 6         |
| 2562 | Cdc37 engages in stable, S14A mutation-reinforced association with the most atypical member of the yeast kinome, Cdk-activating kinase (Cak1). <i>Cell Stress and Chaperones</i> , 2014, 19, 695-703.  | 1.2  | 2         |
| 2563 | Nuclear Localization of Haa1, Which Is Linked to Its Phosphorylation Status, Mediates Lactic Acid Tolerance in <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 2014, 80, 3488-3495.  | 1.4  | 31        |
| 2564 | The Drug Diazaborine Blocks Ribosome Biogenesis by Inhibiting the AAA-ATPase Drg1. <i>Journal of Biological Chemistry</i> , 2014, 289, 3913-3922.  | 1.6  | 48        |
| 2565 | Functional characterization of kinetochore protein, <i>Ctf19</i> in meiosis <i>I</i> : an implication of differential impact of <i>Ctf19</i> on the assembly of mitotic and meiotic kinetochores in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2014, 91, 1179-1199. | 1.2  | 27        |
| 2566 | IpO: plasmids and methods for simplified, PCR-based DNA transplant in yeast. <i>Yeast</i> , 2014, 31, 185-193.   | 0.8  | 1         |
| 2567 | DNA replication and spindle checkpoints cooperate during S phase to delay mitosis and preserve genome integrity. <i>Journal of Cell Biology</i> , 2014, 204, 165-175.  | 2.3  | 35        |
| 2568 | Constitutive and Nitrogen Catabolite Repression-sensitive Production of Gat1 Isoforms. <i>Journal of Biological Chemistry</i> , 2014, 289, 2918-2933.  | 1.6  | 9         |
| 2569 | YCF1-Mediated Cadmium Resistance in Yeast Is Dependent on Copper Metabolism and Antioxidant Enzymes. <i>Antioxidants and Redox Signaling</i> , 2014, 21, 1475-1489.  | 2.5  | 23        |
| 2570 | Identification and characterization of a drug-sensitive strain enables puromycin-based translational assays in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2014, 31, 167-178.   | 0.8  | 10        |
| 2571 | Dynamic association of the PI3P-interacting Mon1-Ccz1 GEF with vacuoles is controlled through its phosphorylation by the type 1 casein kinase Yck3. <i>Molecular Biology of the Cell</i> , 2014, 25, 1608-1619.  | 0.9  | 54        |
| 2572 | Enhanced eIF1 binding to the 40S ribosome impedes conformational rearrangements of the preinitiation complex and elevates initiation accuracy. <i>Rna</i> , 2014, 20, 150-167.   | 1.6  | 36        |
| 2573 | Glutamine methylation in histone H2A is an RNA-polymerase-I-dedicated modification. <i>Nature</i> , 2014, 505, 564-568.  | 13.7 | 186       |
| 2574 | A hierarchical model for assembly of eukaryotic 60S ribosomal subunit domains. <i>Genes and Development</i> , 2014, 28, 198-210.   | 2.7  | 115       |
| 2575 | The N-Terminal Methionine of Cellular Proteins as a Degradation Signal. <i>Cell</i> , 2014, 156, 158-169.  | 13.5 | 147       |
| 2576 | Coupled GTPase and remodelling ATPase activities form a checkpoint for ribosome export. <i>Nature</i> , 2014, 505, 112-116.  | 13.7 | 132       |
| 2577 | Isp7 Is a Novel Regulator of Amino Acid Uptake in the TOR Signaling Pathway. <i>Molecular and Cellular Biology</i> , 2014, 34, 794-806.  | 1.1  | 22        |



| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2578 | Glutathione is essential to preserve nuclear function and cell survival under oxidative stress. <i>Free Radical Biology and Medicine</i> , 2014, 67, 103-114.  | 1.3 | 40        |
| 2579 | Principles of ER cotranslational translocation revealed by proximity-specific ribosome profiling. <i>Science</i> , 2014, 346, 1257-521.  | 6.0 | 350       |
| 2580 | Mediator, TATA-binding Protein, and RNA Polymerase II Contribute to Low Histone Occupancy at Active Gene Promoters in Yeast. <i>Journal of Biological Chemistry</i> , 2014, 289, 14981-14995.  | 1.6 | 25        |
| 2581 | Rsp5/Nedd4 is the main ubiquitin ligase that targets cytosolic misfolded proteins following heat stress. <i>Nature Cell Biology</i> , 2014, 16, 1227-1237.   | 4.6 | 161       |
| 2582 | Cell cycle-dependent regulation of mitochondrial preprotein translocase. <i>Science</i> , 2014, 346, 1109-1113.  | 6.0 | 128       |
| 2583 | Targeted Disruption of an EHD domain Protein Endocytic Complex, Pan1. <i>Traffic</i> , 2014, 15, 43-59.  | 1.3 | 10        |
| 2584 | Der1 promotes movement of misfolded proteins through the endoplasmic reticulum membrane. <i>Nature Cell Biology</i> , 2014, 16, 77-86.   | 4.6 | 132       |
| 2585 | Multiple mechanisms determine the order of APC/C substrate degradation in mitosis. <i>Journal of Cell Biology</i> , 2014, 207, 23-39.  | 2.3 | 68        |
| 2586 | A Visual Screen of Protein Localization during Sporulation Identifies New Components of Prospore Membrane-Associated Complexes in Budding Yeast. <i>Eukaryotic Cell</i> , 2014, 13, 383-391.   | 3.4 | 26        |
| 2587 | On the molecular relationships between high-zinc tolerance and aconitase (Aco1) in <i>Saccharomyces cerevisiae</i> . <i>Metallomics</i> , 2014, 6, 634-645.  | 1.0 | 4         |
| 2588 | Interaction between Ribosome Assembly Factors Krr1 and Faf1 Is Essential for Formation of Small Ribosomal Subunit in Yeast. <i>Journal of Biological Chemistry</i> , 2014, 289, 22692-22703.   | 1.6 | 26        |
| 2589 | Identification of the phytosphingosine metabolic pathway leading to odd-numbered fatty acids. <i>Nature Communications</i> , 2014, 5, 5338.  | 5.8 | 81        |
| 2590 | <i>Saccharomyces cerevisiae</i> Kelch Proteins and Bud14 Protein Form a Stable 520-kDa Formin Regulatory Complex That Controls Actin Cable Assembly and Cell Morphogenesis. <i>Journal of Biological Chemistry</i> , 2014, 289, 18290-18301. | 1.6 | 23        |
| 2591 | Kinetochores require oligomerization of Dam1 complex to maintain microtubule attachments against tension and promote biorientation. <i>Nature Communications</i> , 2014, 5, 4951.  | 5.8 | 51        |
| 2592 | A unique DNA entry gate serves for regulated loading of the eukaryotic replicative helicase MCM2-7 onto DNA. <i>Genes and Development</i> , 2014, 28, 1653-1666.   | 2.7 | 105       |
| 2593 | Sec23 Homolog Nel1 Is a Novel GTPase-activating Protein for Sar1 but Does Not Function as a Subunit of the Coat Protein Complex II (COPII) Coat. <i>Journal of Biological Chemistry</i> , 2014, 289, 21423-21432.                            | 1.6 | 3         |
| 2594 | Rpb4 Subunit Functions Mainly in mRNA Synthesis by RNA Polymerase II. <i>Journal of Biological Chemistry</i> , 2014, 289, 17446-17452.   | 1.6 | 41        |
| 2595 | TOR complex 2-Ypk1 signaling is an essential positive regulator of the general amino acid control response and autophagy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 10586-10591.   | 3.3 | 43        |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 2596 | The yeast oligopeptide transporter Opt2 is localized to peroxisomes and affects glutathione redox homeostasis. <i>FEMS Yeast Research</i> , 2014, 14, n/a-n/a.  | 1.1  | 29        |
| 2597 | The FACT complex interacts with the E3 ubiquitin ligase Psh1 to prevent ectopic localization of CENP-A. <i>Genes and Development</i> , 2014, 28, 1815-1826.   | 2.7  | 66        |
| 2598 | Presequence Recognition by the Tom40 Channel Contributes to Precursor Translocation into the Mitochondrial Matrix. <i>Molecular and Cellular Biology</i> , 2014, 34, 3473-3485.   | 1.1  | 32        |
| 2599 | Comparative proteomics of mitosis and meiosis in <i>Saccharomyces cerevisiae</i> . <i>Journal of Proteomics</i> , 2014, 109, 1-15.  | 1.2  | 12        |
| 2600 | Translation of Small Open Reading Frames within Unannotated RNA Transcripts in <i>Saccharomyces cerevisiae</i> . <i>Cell Reports</i> , 2014, 7, 1858-1866.  | 2.9  | 150       |
| 2601 | Genome-wide Translational Changes Induced by the Prion [PSI <sup>+</sup> ]. <i>Cell Reports</i> , 2014, 8, 439-448.   | 2.9  | 57        |
| 2602 | <i>In Vivo</i> Metabolic Fingerprinting of Neutral Lipids with Hyperspectral Stimulated Raman Scattering Microscopy. <i>Journal of the American Chemical Society</i> , 2014, 136, 8820-8828.  | 6.6  | 169       |
| 2603 | Regulation of cytokinesis in the milk yeast <i>Kluyveromyces lactis</i> . <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2014, 1843, 2685-2697.   | 1.9  | 4         |
| 2604 | Promoter sequences direct cytoplasmic localization and translation of mRNAs during starvation in yeast. <i>Nature</i> , 2014, 514, 117-121.   | 13.7 | 191       |
| 2605 | Interaction of Ddc1 and RPA with single-stranded/double-stranded DNA junctions in yeast whole cell extracts: Proteolytic degradation of the large subunit of replication protein A in <i>ddc1<sup>Δ</sup></i> strains. <i>DNA Repair</i> , 2014, 22, 30-40. | 1.3  | 4         |
| 2606 | Defects in triacylglycerol lipolysis affect synthesis of triacylglycerols and steryl esters in the yeast. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2014, 1841, 1393-1402.  | 1.2  | 8         |
| 2607 | The SUN protein Mps3 controls Ndc1 distribution and function on the nuclear membrane. <i>Journal of Cell Biology</i> , 2014, 204, 523-539.  | 2.3  | 50        |
| 2608 | Enzymatic and Regulatory Attributes of Trehalose 6-Phosphate Phosphatase from <i>Candida utilis</i> and its Role During Thermal Stress. <i>Journal of Cellular Physiology</i> , 2014, 229, 1245-1255.   | 2.0  | 6         |
| 2609 | Surveillance of Nuclear Pore Complex Assembly by ESCRT-III/Vps4. <i>Cell</i> , 2014, 159, 388-401.  | 13.5 | 211       |
| 2610 | Condensin suppresses recombination and regulates double-strand break processing at the repetitive ribosomal DNA array to ensure proper chromosome segregation during meiosis in budding yeast. <i>Molecular Biology of the Cell</i> , 2014, 25, 2934-2947.  | 0.9  | 20        |
| 2611 | Amyloid-associated activity contributes to the severity and toxicity of a prion phenotype. <i>Nature Communications</i> , 2014, 5, 4384.  | 5.8  | 39        |
| 2612 | Nutritional status modulates box C/D snoRNP biogenesis by regulated subcellular relocalization of the R2TP complex. <i>Genome Biology</i> , 2014, 15, 404.  | 3.8  | 40        |
| 2613 | Coregulated Expression of the Na <sup>+</sup> /Phosphate Pho89 Transporter and Ena1 Na <sup>+</sup> -ATPase Allows Their Functional Coupling under High-pH Stress. <i>Molecular and Cellular Biology</i> , 2014, 34, 4420-4435.                             | 1.1  | 39        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 2614 | Peroxisomes are juxtaposed to strategic sites on mitochondria. <i>Molecular BioSystems</i> , 2014, 10, 1742-1748.  | 2.9  | 95        |
| 2615 | The F-BAR protein Hof1 tunes formin activity to sculpt actin cables during polarized growth. <i>Molecular Biology of the Cell</i> , 2014, 25, 1730-1743.   | 0.9  | 30        |
| 2616 | Regulation of Clathrin-Mediated Endocytosis by Dynamic Ubiquitination and Deubiquitination. <i>Current Biology</i> , 2014, 24, 951-959.  | 1.8  | 40        |
| 2617 | Analyzing G1â€™S Transcriptional Control. <i>Methods in Molecular Biology</i> , 2014, 1170, 463-476.   | 0.4  | 1         |
| 2618 | The molecular topography of silenced chromatin in <i>Saccharomyces cerevisiae</i> . <i>Genes and Development</i> , 2014, 28, 245-258.  | 2.7  | 66        |
| 2619 | Large-Scale Genetic Perturbations Reveal Regulatory Networks and an Abundance of Gene-Specific Repressors. <i>Cell</i> , 2014, 157, 740-752.   | 13.5 | 248       |
| 2620 | Fission of SNX-BARâ€™coated endosomal retrograde transport carriers is promoted by the dynamin-related protein Vps1. <i>Journal of Cell Biology</i> , 2014, 204, 793-806.                                | 2.3  | 75        |
| 2621 | A Dynamic Interface between Vacuoles and Mitochondria in Yeast. <i>Developmental Cell</i> , 2014, 30, 95-102.  | 3.1  | 321       |
| 2622 | Tension-dependent removal of pericentromeric shugoshin is an indicator of sister chromosome biorientation. <i>Genes and Development</i> , 2014, 28, 1291-1309.   | 2.7  | 65        |
| 2623 | Glucose Starvation Inhibits Autophagy via Vacuolar Hydrolysis and Induces Plasma Membrane Internalization by Down-regulating Recycling. <i>Journal of Biological Chemistry</i> , 2014, 289, 16736-16747. | 1.6  | 87        |
| 2624 | Stress-dependent Proteolytic Processing of the Actin Assembly Protein Lsb1 Modulates a Yeast Prion. <i>Journal of Biological Chemistry</i> , 2014, 289, 27625-27639.                                     | 1.6  | 29        |
| 2625 | Quality control of inner nuclear membrane proteins by the Asi complex. <i>Science</i> , 2014, 346, 751-755.  | 6.0  | 171       |
| 2626 | Swi2/Snf2-like protein Uls1 functions in the Sgs1-dependent pathway of maintenance of rDNA stability and alleviation of replication stress. <i>DNA Repair</i> , 2014, 21, 24-35.                         | 1.3  | 8         |
| 2627 | Pat1 contributes to the RNA binding activity of the Lsm1-7â€™Pat1 complex. <i>Rna</i> , 2014, 20, 1465-1475.   | 1.6  | 39        |
| 2628 | Sister kinetochores are mechanically fused during meiosis I in yeast. <i>Science</i> , 2014, 346, 248-251.   | 6.0  | 68        |
| 2629 | Pseudouridine profiling reveals regulated mRNA pseudouridylation in yeast and human cells. <i>Nature</i> , 2014, 515, 143-146.   | 13.7 | 800       |
| 2630 | Use of Yeast Spores for Microencapsulation of Enzymes. <i>Applied and Environmental Microbiology</i> , 2014, 80, 4502-4510.  | 1.4  | 19        |
| 2631 | Frataxin-bypassing Isu1: characterization of the bypass activity in cells and mitochondria. <i>Biochemical Journal</i> , 2014, 459, 71-81.   | 1.7  | 34        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2632 | Mitochondrially-targeted bacterial phosphatidylethanolamine methyltransferase sustained phosphatidylcholine synthesis of a <i>Saccharomyces cerevisiae</i> $\Delta$ pem1 $\Delta$ pem2 double mutant without exogenous choline supply. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2014, 1841, 1264-1271. | 1.2 | 8         |
| 2633 | Applied Usage of Yeast Spores as Chitosan Beads. <i>Applied and Environmental Microbiology</i> , 2014, 80, 5098-5105.   | 1.4 | 20        |
| 2634 | Budding Yeast for Budding Geneticists: A Primer on the <i>Saccharomyces cerevisiae</i> Model System. <i>Genetics</i> , 2014, 197, 33-48.  | 1.2 | 168       |
| 2635 | Transcriptional Regulation by Pho23 Modulates the Frequency of Autophagosome Formation. <i>Current Biology</i> , 2014, 24, 1314-1322.   | 1.8 | 87        |
| 2636 | Sus1p Facilitates Pre-Initiation Complex Formation at the SAGA-Regulated Genes Independently of Histone H2B De-Ubiquitylation. <i>Journal of Molecular Biology</i> , 2014, 426, 2928-2941.  | 2.0 | 16        |
| 2637 | Analysis of subcellular localization and function of the yeast Rab6 homologue, Ypt6p, using a novel amino-terminal tagging strategy. <i>Biochemical and Biophysical Research Communications</i> , 2014, 450, 519-525.   | 1.0 | 8         |
| 2638 | The Wsc1p Cell Wall Signaling Protein Controls Biofilm (Mat) Formation Independently of Flo11p in <i>Saccharomyces cerevisiae</i> . <i>G3: Genes, Genomes, Genetics</i> , 2014, 4, 199-207.   | 0.8 | 11        |
| 2639 | The C-terminal Cytosolic Region of Rim21 Senses Alterations in Plasma Membrane Lipid Composition. <i>Journal of Biological Chemistry</i> , 2015, 290, 30797-30805.  | 1.6 | 25        |
| 2640 | Atg23 and Atg27 Act at the Early Stages of Atg9 Trafficking in <i>S. cerevisiae</i> . <i>Traffic</i> , 2015, 16, 172-190.   | 1.3 | 44        |
| 2641 | Improved Plasmids for Fluorescent Protein Tagging of Microtubules in <i>Saccharomyces cerevisiae</i> . <i>Traffic</i> , 2015, 16, 773-786.  | 1.3 | 57        |
| 2643 | Loss of APD1 in Yeast Confers Hydroxyurea Sensitivity Suppressed by Yap1p Transcription Factor. <i>Scientific Reports</i> , 2015, 5, 7897.  | 1.6 | 9         |
| 2644 | Functional roles of a predicted branched chain aminotransferase encoded by the LkBAT1 gene of the yeast <i>Lachancea kluyveri</i> . <i>Fungal Genetics and Biology</i> , 2015, 85, 71-82.   | 0.9 | 9         |
| 2645 | Srv2/CAP is required for polarized actin cable assembly and patch internalization during clathrin-mediated endocytosis. <i>Journal of Cell Science</i> , 2016, 129, 367-79.   | 1.2 | 8         |
| 2646 | Yeast Cyc8p and Tup1p proteins function as coactivators for transcription of Stp1/2p-dependent amino acid transporter genes. <i>Biochemical and Biophysical Research Communications</i> , 2015, 468, 32-38.   | 1.0 | 23        |
| 2647 | Dissecting seipin function: the localized accumulation of phosphatidic acid at ER/LD junctions in the absence of seipin is suppressed by Sei1p <sup>N</sup> term only in combination with Ldb16p. <i>BMC Cell Biology</i> , 2015, 16, 29.   | 3.0 | 57        |
| 2648 | Ubiquitin binding by the CUE domain promotes endosomal localization of the Rab5 GEF Vps9. <i>Molecular Biology of the Cell</i> , 2015, 26, 1345-1356.   | 0.9 | 27        |
| 2649 | A Pan1/End3/Sla1 complex links Arp2/3-mediated actin assembly to sites of clathrin-mediated endocytosis. <i>Molecular Biology of the Cell</i> , 2015, 26, 3841-3856.  | 0.9 | 58        |
| 2650 | Quantitation of yeast cell-cell fusion using multicolor flow cytometry. <i>Cytometry Part A: the Journal of the International Society for Analytical Cytology</i> , 2015, 87, 843-854.  | 1.1 | 3         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2651 | Depletion of yeast PDK1 orthologs triggers a stress-like transcriptional response. <i>BMC Genomics</i> , 2015, 16, 719.   | 1.2 | 3         |
| 2652 | Quantitative framework for ordered degradation of APC/C substrates. <i>BMC Biology</i> , 2015, 13, 96.  | 1.7 | 23        |
| 2653 | iAID: an improved auxin-inducible degron system for the construction of a "tight" conditional mutant in the budding yeast <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2015, 32, 567-581.               | 0.8 | 40        |
| 2654 | Metallothionein as a clonable tag for protein localization by electron microscopy of cells. <i>Journal of Microscopy</i> , 2015, 260, 20-29.  | 0.8 | 28        |
| 2655 | TORC1 Regulates Developmental Responses to Nitrogen Stress via Regulation of the GATA Transcription Factor Gaf1. <i>MBio</i> , 2015, 6, e00959.   | 1.8 | 32        |
| 2656 | A novel protein, <i>Rsf1</i> , is critical for the single-strand annealing pathway of double-strand break repair in <i>Schizosaccharomyces pombe</i> . <i>Molecular Microbiology</i> , 2015, 96, 1211-1225. | 1.2 | 2         |
| 2657 | Development of a Tightly Controlled Off Switch for <i>Saccharomyces cerevisiae</i> Regulated by Camphor, a Low-Cost Natural Product. <i>G3: Genes, Genomes, Genetics</i> , 2015, 5, 1983-1990.              | 0.8 | 27        |
| 2658 | Sec66-Dependent Regulation of Yeast Spindle-Pole Body Duplication Through Pom152. <i>Genetics</i> , 2015, 201, 1479-1495.   | 1.2 | 11        |
| 2659 | Reliable cell cycle commitment in budding yeast is ensured by signal integration. <i>ELife</i> , 2015, 4, .   | 2.8 | 67        |
| 2660 | Expression of Subtelomeric lncRNAs Links Telomeres Dynamics to RNA Decay in <i>S. cerevisiae</i> . <i>Non-coding RNA</i> , 2015, 1, 94-126.   | 1.3 | 9         |
| 2661 | Electron microscopy for ultrastructural analysis and protein localization in <i>Saccharomyces cerevisiae</i> . <i>Microbial Cell</i> , 2015, 2, 412-428.  | 1.4 | 23        |
| 2662 | PRIMED: PRIMER Database for Deleting and Tagging All Fission and Budding Yeast Genes Developed Using the Open-Source Genome Retrieval Script (GRS). <i>PLoS ONE</i> , 2015, 10, e0116657.                   | 1.1 | 6         |
| 2663 | Inositol Depletion Restores Vesicle Transport in Yeast Phospholipid Flippase Mutants. <i>PLoS ONE</i> , 2015, 10, e0120108.   | 1.1 | 9         |
| 2664 | A Genomic Safe Haven for Mutant Complementation in <i>Cryptococcus neoformans</i> . <i>PLoS ONE</i> , 2015, 10, e0122916.   | 1.1 | 83        |
| 2665 | The CAF-1 and Hir Histone Chaperones Associate with Sites of Meiotic Double-Strand Breaks in Budding Yeast. <i>PLoS ONE</i> , 2015, 10, e0125965.   | 1.1 | 13        |
| 2666 | Differential Chromosome Conformations as Hallmarks of Cellular Identity Revealed by Mathematical Polymer Modeling. <i>PLoS Computational Biology</i> , 2015, 11, e1004306.                                  | 1.5 | 28        |
| 2667 | Competition between Heterochromatic Loci Allows the Abundance of the Silencing Protein, Sir4, to Regulate de novo Assembly of Heterochromatin. <i>PLoS Genetics</i> , 2015, 11, e1005425.                   | 1.5 | 9         |
| 2668 | The <i>Saccharomyces cerevisiae</i> AMPK, Snf1, Negatively Regulates the Hog1 MAPK Pathway in ER Stress Response. <i>PLoS Genetics</i> , 2015, 11, e1005491.  | 1.5 | 45        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2669 | The RNAPII-CTD Maintains Genome Integrity through Inhibition of Retrotransposon Gene Expression and Transposition. <i>PLoS Genetics</i> , 2015, 11, e1005608.  | 1.5 | 4         |
| 2670 | The Eukaryotic Mismatch Recognition Complexes Track with the Replisome during DNA Synthesis. <i>PLoS Genetics</i> , 2015, 11, e1005719.  | 1.5 | 17        |
| 2671 | Cdc14 Early Anaphase Release, FEAR, Is Limited to the Nucleus and Dispensable for Efficient Mitotic Exit. <i>PLoS ONE</i> , 2015, 10, e0128604.  | 1.1 | 18        |
| 2672 | A Random Screen Using a Novel Reporter Assay System Reveals a Set of Sequences That Are Preferred as the TATA or TATA-Like Elements in the <i>CYC1</i> Promoter of <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2015, 10, e0129357. | 1.1 | 4         |
| 2673 | A Yeast Mutant Deleted of <i>GPH1</i> Bears Defects in Lipid Metabolism. <i>PLoS ONE</i> , 2015, 10, e0136957.   | 1.1 | 7         |
| 2674 | Structural Motifs Critical for In Vivo Function and Stability of the RecQ-Mediated Genome Instability Protein Rmi1. <i>PLoS ONE</i> , 2015, 10, e0145466.  | 1.1 | 3         |
| 2675 | Genes of Different Catabolic Pathways Are Coordinately Regulated by Dal81 in <i>Saccharomyces cerevisiae</i> . <i>Journal of Amino Acids</i> , 2015, 2015, 1-8.  | 5.8 | 7         |
| 2676 | Maturation of the proteasome core particle induces an affinity switch that controls regulatory particle association. <i>Nature Communications</i> , 2015, 6, 6384.   | 5.8 | 39        |
| 2677 | Identification of <i>Dck1</i> and <i>Lmo1</i> as upstream regulators of the small <i>GTPase Rho5</i> in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2015, 96, 306-324.   | 1.2 | 23        |
| 2678 | A functional Rim101 complex is required for proper accumulation of the Ena1 Na <sup>+</sup> -ATPase protein in response to salt stress in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2015, 15, fov017.                 | 1.1 | 18        |
| 2679 | Ndj1, a telomere-associated protein, regulates centrosome separation in budding yeast meiosis. <i>Journal of Cell Biology</i> , 2015, 209, 247-259.  | 2.3 | 26        |
| 2680 | A defect of the vacuolar putative lipase <i>Atg15</i> accelerates degradation of lipid droplets through lipolysis. <i>Autophagy</i> , 2015, 11, 1247-1258.   | 4.3 | 32        |
| 2681 | TORC2 and eisosomes are spatially interdependent, requiring optimal level of phosphatidylinositol 4, 5-bisphosphate for their integrity. <i>Journal of Biosciences</i> , 2015, 40, 299-311.  | 0.5 | 13        |
| 2683 | The longevity of <i>tor1<sup>Δ</sup></i> , <i>sch9<sup>Δ</sup></i> , and <i>ras2<sup>Δ</sup></i> mutants depends on actin dynamics in <i>Saccharomyces cerevisiae</i> . <i>Cell and Bioscience</i> , 2015, 5, 18.                          | 2.1 | 8         |
| 2684 | A second essential function of the Est1-binding arm of yeast telomerase RNA. <i>Rna</i> , 2015, 21, 862-876.   | 1.6 | 20        |
| 2685 | A protein interaction map of the LSU processome. <i>Genes and Development</i> , 2015, 29, 862-875.   | 2.7 | 41        |
| 2686 | A fluorescent tool set for yeast Atg proteins. <i>Autophagy</i> , 2015, 11, 954-960.   | 4.3 | 22        |
| 2687 | Bub3 promotes Cdc20-dependent activation of the APC/C in <i>S. cerevisiae</i> . <i>Journal of Cell Biology</i> , 2015, 209, 519-527.   | 2.3 | 21        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2688 | Insights into molecular plasticity in protein complexes from Trm9-Trm112 tRNA modifying enzyme crystal structure. <i>Nucleic Acids Research</i> , 2015, 43, 10989-11002.                                       | 6.5 | 38        |
| 2689 | Combinatorial genetic analysis of a network of actin disassembly-promoting factors. <i>Cytoskeleton</i> , 2015, 72, 349-361.   | 1.0 | 18        |
| 2690 | NlpA-mediated modulation of outer membrane vesicle production through peptidoglycan dynamics in <i>Escherichia coli</i> . <i>MicrobiologyOpen</i> , 2015, 4, 375-389.  | 1.2 | 85        |
| 2691 | Regulation of Telomere Length Requires a Conserved N-Terminal Domain of Rif2 in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2015, 201, 573-586.  | 1.2 | 21        |
| 2692 | Atg41/Icy2 regulates autophagosome formation. <i>Autophagy</i> , 2015, 11, 2288-2299.  | 4.3 | 88        |
| 2693 | The seipin complex Fld1/Ldb16 stabilizes ER-lipid droplet contact sites. <i>Journal of Cell Biology</i> , 2015, 211, 829-844.  | 2.3 | 171       |
| 2694 | Experimental testing of a new integrated model of the budding yeast S<sc> tart</sc> transition. <i>Molecular Biology of the Cell</i> , 2015, 26, 3966-3984.  | 0.9 | 25        |
| 2695 | Yeast cell-based analysis of human lactate dehydrogenase isoforms. <i>Journal of Biochemistry</i> , 2015, 158, mvv061.   | 0.9 | 10        |
| 2696 | Influence of Ogg1 repair on the genetic stability of <i>ccc2</i> mutant of <i>Saccharomyces cerevisiae</i> chemically challenged with 4-nitroquinoline-1-oxide (4-NQO). <i>Mutagenesis</i> , 2016, 31, gev062. | 1.0 | 5         |
| 2697 | Cwc21p promotes the second step conformation of the spliceosome and modulates 3' splice site selection. <i>Nucleic Acids Research</i> , 2015, 43, 3309-3317.   | 6.5 | 17        |
| 2698 | Enzymatic attributes of an l-isoaspartyl methyltransferase from <i>Candida utilis</i> and its role in cell survival. <i>Biochemistry and Biophysics Reports</i> , 2015, 4, 59-75.                              | 0.7 | 2         |
| 2699 | A large-scale analysis of autophagy-related gene expression identifies new regulators of autophagy. <i>Autophagy</i> , 2015, 11, 2114-2122.  | 4.3 | 57        |
| 2700 | Evidence for a Nonendosomal Function of the <i>Saccharomyces cerevisiae</i> ESCRT-III-Like Protein Chm7. <i>Genetics</i> , 2015, 201, 1439-1452.   | 1.2 | 40        |
| 2701 | INO80 Chromatin Remodeler Facilitates Release of RNA Polymerase II from Chromatin for Ubiquitin-Mediated Proteasomal Degradation. <i>Molecular Cell</i> , 2015, 60, 784-796.                                   | 4.5 | 64        |
| 2702 | Coordinated Ribosomal ITS2 RNA Processing by the Las1 Complex Integrating Endonuclease, Polynucleotide Kinase, and Exonuclease Activities. <i>Molecular Cell</i> , 2015, 60, 808-815.                          | 4.5 | 83        |
| 2703 | Genome-Wide Association of Mediator and RNA Polymerase II in Wild-Type and Mediator Mutant Yeast. <i>Molecular and Cellular Biology</i> , 2015, 35, 331-342.   | 1.1 | 48        |
| 2704 | An Engineered Minimal WASP-Myosin Fusion Protein Reveals Essential Functions for Endocytosis. <i>Developmental Cell</i> , 2015, 35, 281-294.   | 3.1 | 35        |
| 2705 | Control of lipid organization and actin assembly during clathrin-mediated endocytosis by the cytoplasmic tail of the rhomboid protein Rbd2. <i>Molecular Biology of the Cell</i> , 2015, 26, 1509-1522.        | 0.9 | 13        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2706 | The dynein cortical anchor Num1 activates dynein motility by relieving Pac1/LIS1-mediated inhibition. <i>Journal of Cell Biology</i> , 2015, 211, 309-322.   | 2.3 | 64        |
| 2707 | Glucose-Regulated Phosphorylation of the PUF Protein Puf3 Regulates the Translational Fate of Its Bound mRNAs and Association with RNA Granules. <i>Cell Reports</i> , 2015, 11, 1638-1650.                                    | 2.9 | 83        |
| 2708 | Replication-Coupled PCNA Unloading by the Elg1 Complex Occurs Genome-wide and Requires Okazaki Fragment Ligation. <i>Cell Reports</i> , 2015, 12, 774-787.   | 2.9 | 100       |
| 2709 | Deletion of Mitochondrial Porin Alleviates Stress Sensitivity in the Yeast Model of Shwachman-Diamond Syndrome. <i>Journal of Genetics and Genomics</i> , 2015, 42, 671-684.   | 1.7 | 8         |
| 2710 | A vacuolar membrane protein Avt7p is involved in transport of amino acid and spore formation in <i>Saccharomyces cerevisiae</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 2015, 79, 190-195.                       | 0.6 | 15        |
| 2711 | Rph1/KDM4 Mediates Nutrient-Limitation Signaling that Leads to the Transcriptional Induction of Autophagy. <i>Current Biology</i> , 2015, 25, 546-555.   | 1.8 | 96        |
| 2712 | Feedback control of prion formation and propagation by the ribosome-associated chaperone complex. <i>Molecular Microbiology</i> , 2015, 96, 621-632.   | 1.2 | 36        |
| 2713 | Pan1 regulates transitions between stages of clathrin-mediated endocytosis. <i>Molecular Biology of the Cell</i> , 2015, 26, 1371-1385.  | 0.9 | 20        |
| 2714 | Proteasome assembly from 15S precursors involves major conformational changes and recycling of the Pba1-Pba2 chaperone. <i>Nature Communications</i> , 2015, 6, 6123.  | 5.8 | 42        |
| 2715 | Rbs1, a New Protein Implicated in RNA Polymerase III Biogenesis in Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2015, 35, 1169-1181.  | 1.1 | 30        |
| 2716 | Cytokinesis breaks dicentric chromosomes preferentially at pericentromeric regions and telomere fusions. <i>Genes and Development</i> , 2015, 29, 322-336.   | 2.7 | 48        |
| 2717 | Maintenance of Nucleosomal Balance in <i>Cis</i> by Conserved AAA-ATPase Yta7. <i>Genetics</i> , 2015, 199, 105-116.   | 1.2 | 18        |
| 2718 | Storage lipid synthesis is necessary for autophagy induced by nitrogen starvation. <i>FEBS Letters</i> , 2015, 589, 269-276.   | 1.3 | 52        |
| 2719 | The Erv41-Erv46 complex serves as a retrograde receptor to retrieve escaped ER proteins. <i>Journal of Cell Biology</i> , 2015, 208, 197-209.  | 2.3 | 40        |
| 2720 | The Anillin-Related Region of Bud4 Is the Major Functional Determinant for Bud4's Function in Septin Organization during Bud Growth and Axial Bud Site Selection in Budding Yeast. <i>Eukaryotic Cell</i> , 2015, 14, 241-251. | 3.4 | 19        |
| 2721 | Constitutively active ESCRT-II suppresses the MVB-sorting phenotype of ESCRT-0 and ESCRT-I mutants. <i>Molecular Biology of the Cell</i> , 2015, 26, 554-568.  | 0.9 | 21        |
| 2722 | Yeast as a model system to screen purine derivatives against human CDK1 and CDK2 kinases. <i>Journal of Biotechnology</i> , 2015, 195, 30-36.  | 1.9 | 4         |
| 2723 | The interplay of Hrd3 and the molecular chaperone system ensures efficient degradation of malformed secretory proteins. <i>Molecular Biology of the Cell</i> , 2015, 26, 185-194.  | 0.9 | 32        |



| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2724 | <sc>Y</sc>ap7 is a transcriptional repressor of nitric oxide oxidase in yeasts, which arose from neofunctionalization after whole genome duplication. <i>Molecular Microbiology</i> , 2015, 96, 951-972.        | 1.2 | 43        |
| 2725 | Glycolysis Controls Plasma Membrane Glucose Sensors To Promote Glucose Signaling in Yeasts. <i>Molecular and Cellular Biology</i> , 2015, 35, 747-757.  | 1.1 | 18        |
| 2726 | Yeast Translation Elongation Factor-1A Binds Vacuole-localized Rho1p to Facilitate Membrane Integrity through F-actin Remodeling. <i>Journal of Biological Chemistry</i> , 2015, 290, 4705-4716.                | 1.6 | 15        |
| 2727 | tRNA thiolation links translation to stress responses in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2015, 26, 270-282.  | 0.9 | 61        |
| 2728 | Essential role of the iron-sulfur cluster binding domain of the primase regulatory subunit Pri2 in DNA replication initiation. <i>Protein and Cell</i> , 2015, 6, 194-210.                                      | 4.8 | 30        |
| 2729 | Regulatory link between steryl ester formation and hydrolysis in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2015, 1851, 977-986. | 1.2 | 15        |
| 2730 | Hrr25/CK1 $\hat{c}$ -directed release of Ltv1 from pre-40S ribosomes is necessary for ribosome assembly and cell growth. <i>Journal of Cell Biology</i> , 2015, 208, 745-759.                                   | 2.3 | 71        |
| 2731 | Rab5-family guanine nucleotide exchange factors bind retromer and promote its recruitment to endosomes. <i>Molecular Biology of the Cell</i> , 2015, 26, 1119-1128.   | 0.9 | 36        |
| 2732 | The Chromatin and Transcriptional Landscape of Native <i>Saccharomyces cerevisiae</i> Telomeres and Subtelomeric Domains. <i>Genetics</i> , 2015, 200, 505-521.   | 1.2 | 92        |
| 2733 | The translation termination factor eRF1 (Sup45p) of <i>Saccharomyces cerevisiae</i> is required for pseudohyphal growth and invasion. <i>FEMS Yeast Research</i> , 2015, 15, fov033.                            | 1.1 | 4         |
| 2734 | Lam6 Regulates the Extent of Contacts between Organelles. <i>Cell Reports</i> , 2015, 12, 7-14.   | 2.9 | 173       |
| 2735 | Distribution of Sec24 isoforms to each ER exit site is dynamically regulated in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 2015, 589, 1234-1239.   | 1.3 | 11        |
| 2736 | Significantly Diverged Did2/Vps46 Orthologues from the Protozoan Parasite <i>Giardia lamblia</i> . <i>Current Microbiology</i> , 2015, 71, 333-340.   | 1.0 | 5         |
| 2737 | Phosphorylation of Sae2 Mediates Forkhead-associated (FHA) Domain-specific Interaction and Regulates Its DNA Repair Function. <i>Journal of Biological Chemistry</i> , 2015, 290, 10751-10763.                  | 1.6 | 44        |
| 2738 | Fungal Mediator Tail Subunits Contain Classical Transcriptional Activation Domains. <i>Molecular and Cellular Biology</i> , 2015, 35, 1363-1375.  | 1.1 | 17        |
| 2739 | Functional Interplay between the 53BP1-Ortholog Rad9 and the Mre11 Complex Regulates Resection, End-Tethering and Repair of a Double-Strand Break. <i>PLoS Genetics</i> , 2015, 11, e1004928.                   | 1.5 | 103       |
| 2740 | Phosphorylation of Elp1 by Hrr25 Is Required for Elongator-Dependent tRNA Modification in Yeast. <i>PLoS Genetics</i> , 2015, 11, e1004931.   | 1.5 | 38        |
| 2741 | Examination of Sec22 Homodimer Formation and Role in SNARE-dependent Membrane Fusion. <i>Journal of Biological Chemistry</i> , 2015, 290, 10657-10666.  | 1.6 | 16        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 2742 | Conformational Flexibility and Subunit Arrangement of the Modular Yeast Spt-Ada-Gcn5 Acetyltransferase Complex. <i>Journal of Biological Chemistry</i> , 2015, 290, 10057-10070.   | 1.6  | 59        |
| 2743 | Chaperoning 5S RNA assembly. <i>Genes and Development</i> , 2015, 29, 1432-1446.   | 2.7  | 54        |
| 2744 | Yeast <i>Saccharomyces cerevisiae</i> adiponectin receptor homolog Izh2 is involved in the regulation of zinc, phospholipid and pH homeostasis. <i>Metallomics</i> , 2015, 7, 1338-1351.   | 1.0  | 8         |
| 2745 | Tracking individual secretory vesicles during exocytosis reveals an ordered and regulated process. <i>Journal of Cell Biology</i> , 2015, 210, 181-189.  | 2.3  | 75        |
| 2746 | A conserved mechanism of TOR-dependent RCK-mediated mRNA degradation regulates autophagy. <i>Nature Cell Biology</i> , 2015, 17, 930-942.  | 4.6  | 91        |
| 2747 | The DEAH-box Helicase Dhr1 Dissociates U3 from the Pre-rRNA to Promote Formation of the Central Pseudoknot. <i>PLoS Biology</i> , 2015, 13, e1002083.  | 2.6  | 70        |
| 2748 | Eaf1p Is Required for Recruitment of NuA4 in Targeting TFIID to the Promoters of the Ribosomal Protein Genes for Transcriptional Initiation <i>In Vivo</i> . <i>Molecular and Cellular Biology</i> , 2015, 35, 2947-2964.                | 1.1  | 23        |
| 2749 | The Protein Quality Control Machinery Regulates Its Misassembled Proteasome Subunits. <i>PLoS Genetics</i> , 2015, 11, e1005178.   | 1.5  | 52        |
| 2750 | Role for RNA:DNA hybrids in origin-independent replication priming in a eukaryotic system. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, 5779-5784.                                | 3.3  | 90        |
| 2751 | Diversification of Paralogous $\pm$ -Isopropylmalate Synthases by Modulation of Feedback Control and Hetero-Oligomerization in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2015, 14, 564-577.                             | 3.4  | 29        |
| 2752 | Phosphorylation of Atg31 is required for autophagy. <i>Protein and Cell</i> , 2015, 6, 288-296.  | 4.8  | 13        |
| 2753 | Fungal cytochrome P450 monooxygenases of <i>Fusarium oxysporum</i> for the synthesis of $\omega$ -hydroxy fatty acids in engineered <i>Saccharomyces cerevisiae</i> . <i>Microbial Cell Factories</i> , 2015, 14, 45.                    | 1.9  | 56        |
| 2754 | A reduced VWA domain-containing proteasomal ubiquitin receptor of <i>Giardia lamblia</i> localizes to the flagellar pore regions in microtubule-dependent manner. <i>Parasites and Vectors</i> , 2015, 8, 120.                           | 1.0  | 11        |
| 2755 | Compartmentalization of a Bistable Switch Enables Memory to Cross a Feedback-Driven Transition. <i>Cell</i> , 2015, 160, 1182-1195.  | 13.5 | 45        |
| 2756 | Polyploidy can drive rapid adaptation in yeast. <i>Nature</i> , 2015, 519, 349-352.  | 13.7 | 376       |
| 2757 | Both the autophagy and proteasomal pathways facilitate the Ubp3p-dependent depletion of a subset of translation and RNA turnover factors during nitrogen starvation in <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 2015, 21, 898-910. | 1.6  | 27        |
| 2758 | Head-to-tail regulation is critical for the in vivo function of myosin V. <i>Journal of Cell Biology</i> , 2015, 209, 359-365.   | 2.3  | 29        |
| 2759 | Traffic of p24 Proteins and COPII Coat Composition Mutually Influence Membrane Scaffolding. <i>Current Biology</i> , 2015, 25, 1296-1305.  | 1.8  | 29        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2760 | Coordinated Ribosomal L4 Protein Assembly into the Pre-Ribosome Is Regulated by Its Eukaryote-Specific Extension. <i>Molecular Cell</i> , 2015, 58, 854-862.  | 4.5 | 69        |
| 2761 | Hsp70 clears misfolded kinases that partitioned into distinct quality-control compartments. <i>Molecular Biology of the Cell</i> , 2015, 26, 1583-1600.   | 0.9 | 7         |
| 2762 | Bimolecular Fluorescence Complementation (BiFC) Analysis: Advances and Recent Applications for Genome-Wide Interaction Studies. <i>Journal of Molecular Biology</i> , 2015, 427, 2039-2055.                           | 2.0 | 207       |
| 2763 | Control of Plasma Membrane Permeability by ABC Transporters. <i>Eukaryotic Cell</i> , 2015, 14, 442-453.  | 3.4 | 39        |
| 2764 | Identifying a Rab Effector on the Macroautophagy Pathway. <i>Methods in Molecular Biology</i> , 2015, 1298, 117-125.  | 0.4 | 1         |
| 2765 | Role of Phosphatidylinositol Phosphate Signaling in the Regulation of the Filamentous-Growth Mitogen-Activated Protein Kinase Pathway. <i>Eukaryotic Cell</i> , 2015, 14, 427-440.                                    | 3.4 | 13        |
| 2766 | The N-terminal domains determine cellular localization and functions of the Doa4 and Ubp5 deubiquitinating enzymes. <i>Biochemical and Biophysical Research Communications</i> , 2015, 467, 570-576.                  | 1.0 | 7         |
| 2767 | Starvation-Dependent Regulation of Golgi Quality Control Links the TOR Signaling and Vacuolar Protein Sorting Pathways. <i>Cell Reports</i> , 2015, 12, 1876-1886.  | 2.9 | 46        |
| 2768 | Dynamic single cell measurements of kinase activity by synthetic kinase activity relocation sensors. <i>BMC Biology</i> , 2015, 13, 55.   | 1.7 | 34        |
| 2769 | PKA, PHO and stress response pathways regulate the expression of UDP-glucose pyrophosphorylase through Msn2/4 in budding yeast. <i>FEBS Letters</i> , 2015, 589, 2409-2416.   | 1.3 | 8         |
| 2770 | Complex regulation of Hsf1-Skn7 activities by the catalytic subunits of PKA in <i>Saccharomyces cerevisiae</i> : experimental and computational evidences. <i>BMC Systems Biology</i> , 2015, 9, 42.                  | 3.0 | 10        |
| 2771 | Different pathways for the nuclear import of yeast RNA polymerase II. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2015, 1849, 1354-1362.  | 0.9 | 12        |
| 2772 | Premature termination of <i>GAT1</i> transcription explains paradoxical negative correlation between nitrogen-responsive mRNA, but constitutive low-level protein production. <i>RNA Biology</i> , 2015, 12, 824-837. | 1.5 | 9         |
| 2773 | ChEC-seq kinetics discriminates transcription factor binding sites by DNA sequence and shape in vivo. <i>Nature Communications</i> , 2015, 6, 8733.   | 5.8 | 153       |
| 2774 | The Fab1/PIKfyve Phosphoinositide Phosphate Kinase Is Not Necessary to Maintain the pH of Lysosomes and of the Yeast Vacuole. <i>Journal of Biological Chemistry</i> , 2015, 290, 9919-9928.                          | 1.6 | 46        |
| 2775 | Response of yeast cells to high glucose involves molecular and physiological differences when compared to other osmopressure conditions. <i>FEMS Yeast Research</i> , 2015, 15, fov039.                               | 1.1 | 36        |
| 2776 | Atg27 Tyrosine Sorting Motif is Important for Its Trafficking and Atg9 Localization. <i>Traffic</i> , 2015, 16, 365-378.  | 1.3 | 36        |
| 2777 | Plasmids for C-terminal tagging in <i>Saccharomyces cerevisiae</i> that contain improved GFP proteins, Envy and Ivy. <i>Yeast</i> , 2015, 32, 379-387.  | 0.8 | 60        |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 2778 | Termination of Replication Stress Signaling via Concerted Action of the Slx4 Scaffold and the PP4 Phosphatase. <i>Genetics</i> , 2015, 201, 937-949.  | 1.2  | 21        |
| 2779 | Analysis of phosphorylation sites on autophagy proteins. <i>Protein and Cell</i> , 2015, 6, 698-701.  | 4.8  | 5         |
| 2780 | STUBL-mediated degradation of the transcription factor MAT1±2 requires degradation elements that coincide with corepressor binding sites. <i>Molecular Biology of the Cell</i> , 2015, 26, 3401-3412.           | 0.9  | 20        |
| 2781 | Snf1/AMP-activated protein kinase activates Arf3p to promote invasive yeast growth via a non-canonical GEF domain. <i>Nature Communications</i> , 2015, 6, 7840.  | 5.8  | 12        |
| 2782 | Polarity establishment requires localized activation of Cdc42. <i>Journal of Cell Biology</i> , 2015, 211, 19-26.   | 2.3  | 50        |
| 2783 | Spatial reorganization of telomeres in long-lived quiescent cells. <i>Genome Biology</i> , 2015, 16, 206.   | 3.8  | 75        |
| 2784 | Spt6 Is Essential for rRNA Synthesis by RNA Polymerase I. <i>Molecular and Cellular Biology</i> , 2015, 35, 2321-2331.  | 1.1  | 13        |
| 2785 | Lipid partitioning at the nuclear envelope controls membrane biogenesis. <i>Molecular Biology of the Cell</i> , 2015, 26, 3641-3657.  | 0.9  | 113       |
| 2786 | The Exosome Is Recruited to RNA Substrates through Specific Adaptor Proteins. <i>Cell</i> , 2015, 162, 1029-1038.   | 13.5 | 170       |
| 2787 | Comparative Analysis of Transmembrane Regulators of the Filamentous Growth Mitogen-Activated Protein Kinase Pathway Uncovers Functional and Regulatory Differences. <i>Eukaryotic Cell</i> , 2015, 14, 868-883. | 3.4  | 13        |
| 2788 | ER-mitochondrial junctions can be bypassed by dominant mutations in the endosomal protein Vps13. <i>Journal of Cell Biology</i> , 2015, 210, 883-890.   | 2.3  | 203       |
| 2789 | Sequestration of mRNAs Modulates the Timing of Translation during Meiosis in Budding Yeast. <i>Molecular and Cellular Biology</i> , 2015, 35, 3448-3458.  | 1.1  | 28        |
| 2790 | The fusogenic lipid phosphatidic acid promotes the biogenesis of mitochondrial outer membrane protein Ugo1. <i>Journal of Cell Biology</i> , 2015, 210, 951-960.  | 2.3  | 36        |
| 2791 | Ipl1/Aurora-B is necessary for kinetochore restructuring in meiosis I in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2015, 26, 2986-3000.  | 0.9  | 30        |
| 2792 | Control of the spindle checkpoint by lateral kinetochore attachment and limited Mad1 recruitment. <i>Molecular Biology of the Cell</i> , 2015, 26, 2620-2639.   | 0.9  | 8         |
| 2793 | A LAPF/phafin1-like protein regulates TORC1 and lysosomal membrane permeabilization in response to endoplasmic reticulum membrane stress. <i>Molecular Biology of the Cell</i> , 2015, 26, 4631-4645.           | 0.9  | 68        |
| 2794 | The synaptonemal complex is assembled by a polySUMOylation-driven feedback mechanism in yeast. <i>Journal of Cell Biology</i> , 2015, 211, 785-793.   | 2.3  | 40        |
| 2795 | Molecular Interactions and Cellular Itinerary of the Yeast RAVE (Regulator of the H <sup>+</sup> -ATPase of) Tj ETQq1 1 0.784314 rgBT /Overlock 10  | 1.6  | 28        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2796 | Chromatin Dynamics and the RNA Exosome Function in Concert to Regulate Transcriptional Homeostasis. <i>Cell Reports</i> , 2015, 13, 1610-1622.   | 2.9 | 34        |
| 2797 | Overexpression of alpha-synuclein at non-toxic levels increases dopaminergic cell death induced by copper exposure via modulation of protein degradation pathways. <i>Neurobiology of Disease</i> , 2015, 81, 76-92. | 2.1 | 57        |
| 2798 | DEAD-Box RNA Helicase Dbp4 Is Required for Small-Subunit Processome Formation and Function. <i>Molecular and Cellular Biology</i> , 2015, 35, 816-830.   | 1.1 | 17        |
| 2799 | Spatiotemporal Cascade of Transcription Factor Binding Required for Promoter Activation. <i>Molecular and Cellular Biology</i> , 2015, 35, 688-698.  | 1.1 | 14        |
| 2800 | Both the Charged Linker Region and ATPase Domain of Hsp90 Are Essential for Rad51-Dependent DNA Repair. <i>Eukaryotic Cell</i> , 2015, 14, 64-77.  | 3.4 | 20        |
| 2801 | Identification of proteins associated with RNA polymerase III using a modified tandem chromatin affinity purification. <i>Gene</i> , 2015, 556, 51-60.   | 1.0 | 8         |
| 2802 | Assays for the biochemical and ultrastructural measurement of selective and nonselective types of autophagy in the yeast <i>Saccharomyces cerevisiae</i> . <i>Methods</i> , 2015, 75, 141-150.                       | 1.9 | 38        |
| 2803 | Cdc42p-Interacting Protein Bem4p Regulates the Filamentous-Growth Mitogen-Activated Protein Kinase Pathway. <i>Molecular and Cellular Biology</i> , 2015, 35, 417-436.   | 1.1 | 25        |
| 2804 | Heat Stress-Induced Cup9-Dependent Transcriptional Regulation of <i>SIR2</i> . <i>Molecular and Cellular Biology</i> , 2015, 35, 437-450.  | 1.1 | 12        |
| 2805 | Functional Profiling of Human Fungal Pathogen Genomes. <i>Cold Spring Harbor Perspectives in Medicine</i> , 2015, 5, a019596-a019596.  | 2.9 | 9         |
| 2806 | The yeast Arf-GAP Glo3p is required for the endocytic recycling of cell surface proteins. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2015, 1853, 144-156.                                      | 1.9 | 14        |
| 2807 | The glyoxylate shunt is essential for desiccation tolerance in <i>C. elegans</i> and budding yeast. <i>ELife</i> , 2016, 5, .  | 2.8 | 64        |
| 2808 | The Phospholipid:Diacylglycerol Acyltransferase Lro1 Is Responsible for Hepatitis C Virus Core-Induced Lipid Droplet Formation in a Yeast Model System. <i>PLoS ONE</i> , 2016, 11, e0159324.                        | 1.1 | 8         |
| 2809 | Screening the yeast genome for energetic metabolism pathways involved in a phenotypic response to the anti-cancer agent 3-bromopyruvate. <i>Oncotarget</i> , 2016, 7, 10153-10173.                                   | 0.8 | 18        |
| 2810 | An evidence based hypothesis on the existence of two pathways of mitochondrial crista formation. <i>ELife</i> , 2016, 5, .   | 2.8 | 81        |
| 2811 | Dynamic control of Hsf1 during heat shock by a chaperone switch and phosphorylation. <i>ELife</i> , 2016, 5, .   | 2.8 | 185       |
| 2812 | Role of Nfu1 and Bol3 in iron-sulfur cluster transfer to mitochondrial clients. <i>ELife</i> , 2016, 5, .  | 2.8 | 107       |
| 2813 | The TRAPP Subunit Trs130p Interacts with the GAP Gyp6p to Mediate Ypt6p Dynamics at the Late Golgi. <i>Frontiers in Cell and Developmental Biology</i> , 2016, 4, 48.  | 1.8 | 11        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2814 | A Network of Paralogous Stress Response Transcription Factors in the Human Pathogen <i>Candida glabrata</i> . <i>Frontiers in Microbiology</i> , 2016, 7, 645.  | 1.5 | 31        |
| 2815 | Exploiting Single-Cell Quantitative Data to Map Genetic Variants Having Probabilistic Effects. <i>PLoS Genetics</i> , 2016, 12, e1006213.   | 1.5 | 11        |
| 2816 | Structural Analysis of Free N-Glycans in Î±-Glucosidase Mutants of <i>Saccharomyces cerevisiae</i> : Lack of the Evidence for the Occurrence of Catabolic Î±-Glucosidase Acting on the N-Glycans. <i>PLoS ONE</i> , 2016, 11, e0151891. | 1.1 | 5         |
| 2817 | Msa1 and Msa2 Modulate G1-Specific Transcription to Promote G1 Arrest and the Transition to Quiescence in Budding Yeast. <i>PLoS Genetics</i> , 2016, 12, e1006088.   | 1.5 | 18        |
| 2818 | Scarless Gene Tagging with One-Step Transformation and Two-Step Selection in <i>Saccharomyces cerevisiae</i> and <i>Schizosaccharomyces pombe</i> . <i>PLoS ONE</i> , 2016, 11, e0163950.   | 1.1 | 15        |
| 2819 | Analysis of the Physiological Activities of Scd6 through Its Interaction with Hmt1. <i>PLoS ONE</i> , 2016, 11, e0164773.   | 1.1 | 12        |
| 2820 | Yeast cell wall integrity sensors form specific plasma membrane microdomains important for signalling. <i>Cellular Microbiology</i> , 2016, 18, 1251-1267.  | 1.1 | 52        |
| 2821 | Role of mitochondrial processing peptidase and AAA proteases in processing of the yeast acetoacetyl synthase precursor. <i>FEBS Open Bio</i> , 2016, 6, 765-773.  | 1.0 | 2         |
| 2822 | Ubiquilin/Dsk2 promotes inclusion body formation and vacuole (lysosome)-mediated disposal of mutated huntingtin. <i>Molecular Biology of the Cell</i> , 2016, 27, 2025-2036.  | 0.9 | 27        |
| 2823 | Creating a chimeric clathrin heavy chain that functions independently of yeast clathrin light chain. <i>Traffic</i> , 2016, 17, 754-768.  | 1.3 | 2         |
| 2824 | Yeast nitrogen catabolite repression is sustained by signals distinct from glutamine and glutamate reservoirs. <i>Molecular Microbiology</i> , 2016, 99, 360-379.   | 1.2 | 23        |
| 2825 | Endoplasmic Reticulum-associated Degradation of Pca1p, a Polytopic Protein, via Interaction with the Proteasome at the Membrane. <i>Journal of Biological Chemistry</i> , 2016, 291, 15082-15092.                                       | 1.6 | 9         |
| 2826 | Ribosomal proteins produced in excess are degraded by the ubiquitin-proteasome system. <i>Molecular Biology of the Cell</i> , 2016, 27, 2642-2652.  | 0.9 | 105       |
| 2827 | Altered RNA processing and export lead to retention of mRNAs near transcription sites and nuclear pore complexes or within the nucleolus. <i>Molecular Biology of the Cell</i> , 2016, 27, 2742-2756.                                   | 0.9 | 24        |
| 2828 | Analysis of <scp>COPII</scp> Vesicles Indicates a Role for the Emp47-Ssp120 Complex in Transport of Cell Surface Glycoproteins. <i>Traffic</i> , 2016, 17, 191-210.   | 1.3 | 9         |
| 2829 | Autophagy-independent function of Atg8 in lipid droplet dynamics in yeast. <i>Journal of Biochemistry</i> , 2016, 161, mvw078.  | 0.9 | 10        |
| 2830 | Deubiquitinase activity is required for the proteasomal degradation of misfolded cytosolic proteins upon heat-stress. <i>Nature Communications</i> , 2016, 7, 12907.  | 5.8 | 45        |
| 2831 | A Signaling Lipid Associated with Alzheimer's Disease Promotes Mitochondrial Dysfunction. <i>Scientific Reports</i> , 2016, 6, 19332.   | 1.6 | 25        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 2832 | A role for Mfb1p in region-specific anchorage of high-functioning mitochondria and lifespan in <i>Saccharomyces cerevisiae</i> . <i>Nature Communications</i> , 2016, 7, 10595.  | 5.8  | 46        |
| 2833 | The SND proteins constitute an alternative targeting route to the endoplasmic reticulum. <i>Nature</i> , 2016, 540, 134-138.   | 13.7 | 168       |
| 2834 | Calcium channel regulator Mid1 links TORC2-mediated changes in mitochondrial respiration to autophagy. <i>Journal of Cell Biology</i> , 2016, 215, 779-788.  | 2.3  | 18        |
| 2835 | The Rtr1p CTD phosphatase autoregulates its mRNA through a degradation pathway involving the REX exonucleases. <i>Rna</i> , 2016, 22, 559-570.   | 1.6  | 6         |
| 2836 | Chm7 and Heh1 collaborate to link nuclear pore complex quality control with nuclear envelope sealing. <i>EMBO Journal</i> , 2016, 35, 2447-2467.   | 3.5  | 116       |
| 2837 | High resolution microscopy reveals the nuclear shape of budding yeast during cell cycle and in various biological states. <i>Journal of Cell Science</i> , 2016, 129, 4480-4495.   | 1.2  | 33        |
| 2838 | Orderly progression through S-phase requires dynamic ubiquitylation and deubiquitylation of PCNA. <i>Scientific Reports</i> , 2016, 6, 25513.  | 1.6  | 19        |
| 2839 | ATP hydrolysis by UPF1 is required for efficient translation termination at premature stop codons. <i>Nature Communications</i> , 2016, 7, 14021.  | 5.8  | 49        |
| 2840 | Temporal Expression of a Master Regulator Drives Synchronous Sporulation in Budding Yeast. <i>G3: Genes, Genomes, Genetics</i> , 2016, 6, 3553-3560.   | 0.8  | 40        |
| 2841 | An acridine derivative, [4,5-bis{(N-carboxy methyl imidazolium)methyl}acridine] dibromide, shows anti-TDP-43 aggregation effect in ALS disease models. <i>Scientific Reports</i> , 2016, 6, 39490.                                       | 1.6  | 35        |
| 2842 | Sch 9p kinase and the Gcn4p transcription factor regulate glycerol production during winemaking. <i>FEMS Yeast Research</i> , 2016, 17, fow106.  | 1.1  | 14        |
| 2843 | Ole1, fatty acid desaturase, is required for Atg9 delivery and isolation membrane expansion during autophagy in <i>Saccharomyces cerevisiae</i> . <i>Biology Open</i> , 2017, 6, 35-40.  | 0.6  | 16        |
| 2844 | Structure of the <i>Saccharomyces cerevisiae</i> Hrr25:Mam1 monopolin subcomplex reveals a novel kinase regulator. <i>EMBO Journal</i> , 2016, 35, 2139-2151.  | 3.5  | 22        |
| 2845 | Prevention of DNA Rereplication Through a Meiotic Recombination Checkpoint Response. <i>G3: Genes, Genomes, Genetics</i> , 2016, 6, 3869-3881.   | 0.8  | 2         |
| 2846 | Disruption of ribosome assembly in yeast blocks cotranscriptional pre-rRNA processing and affects the global hierarchy of ribosome biogenesis. <i>Rna</i> , 2016, 22, 852-866.   | 1.6  | 37        |
| 2847 | The zinc cluster proteins U <sub>pc2</sub> and E <sub>cm22</sub> promote filamentation in <i>Saccharomyces cerevisiae</i> by sterol biosynthesis-dependent and -independent pathways. <i>Molecular Microbiology</i> , 2016, 99, 512-527. | 1.2  | 15        |
| 2848 | Doa1 targets ubiquitinated substrates for mitochondria-associated degradation. <i>Journal of Cell Biology</i> , 2016, 213, 49-63.  | 2.3  | 68        |
| 2849 | The transcription factor Ace2 and its paralog Swi5 regulate ethanol production during static fermentation through their targets Cts1 and Rps4a in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2016, 16, fow022.       | 1.1  | 9         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 2850 | DNA Replication Stress Phosphoproteome Profiles Reveal Novel Functional Phosphorylation Sites on Xrs2 in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2016, 203, 353-368.  | 1.2 | 2         |
| 2851 | Rlm1 mediates a positive autoregulatory transcriptional feedback essential for Slt2 MAPK dependent gene expression. <i>Journal of Cell Science</i> , 2016, 129, 1649-60.  | 1.2 | 33        |
| 2852 | Direct interaction of Ste11 and Mkk1/2 through Nst1 integrates high osmolarity glycerol and pheromone pathways to the cell wall integrity MAPK pathway. <i>FEBS Letters</i> , 2016, 590, 148-160.   | 1.3 | 18        |
| 2853 | Chromatin dynamics during DNA replication. <i>Genome Research</i> , 2016, 26, 1245-1256.  | 2.4 | 56        |
| 2854 | The Proteasome Subunit Rpn8 Interacts with the Small Nucleolar RNA Protein (snRNP) Assembly Protein Pih1 and Mediates Its Ubiquitin-independent Degradation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2016, 291, 11761-11775. | 1.6 | 5         |
| 2855 | RNA binding protein Pub1p regulates glycerol production and stress tolerance by controlling Gpd1p activity during winemaking. <i>Applied Microbiology and Biotechnology</i> , 2016, 100, 5017-5027.   | 1.7 | 8         |
| 2856 | Cyclin-Dependent Kinase Co-Ordinates Carbohydrate Metabolism and Cell Cycle in <i>S. cerevisiae</i> . <i>Molecular Cell</i> , 2016, 62, 546-557.  | 4.5 | 71        |
| 2857 | Distinct temporal requirements for autophagy and the proteasome in yeast meiosis. <i>Autophagy</i> , 2016, 12, 671-688.   | 4.3 | 42        |
| 2858 | Regulation of a Spindle Positioning Factor at Kinetochores by SUMO-Targeted Ubiquitin Ligases. <i>Developmental Cell</i> , 2016, 36, 415-427.   | 3.1 | 24        |
| 2859 | Enhancing cytochrome P450-mediated conversions in <i>P. pastoris</i> through RAD52 over-expression and optimizing the cultivation conditions. <i>Fungal Genetics and Biology</i> , 2016, 89, 114-125.   | 0.9 | 22        |
| 2860 | Functions for diverse metabolic activities in heterochromatin. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, E1526-35.  | 3.3 | 14        |
| 2861 | Overexpression of Sly41 suppresses COPII vesicle tethering deficiencies by elevating intracellular calcium levels. <i>Molecular Biology of the Cell</i> , 2016, 27, 1635-1649.  | 0.9 | 4         |
| 2862 | Rtt107 BRCT domains act as a targeting module in the DNA damage response. <i>DNA Repair</i> , 2016, 37, 22-32.  | 1.3 | 15        |
| 2863 | The plasma membrane protein Rch1 is a negative regulator of cytosolic calcium homeostasis and positively regulated by the calcium/calineurin signaling pathway in budding yeast. <i>European Journal of Cell Biology</i> , 2016, 95, 164-174.                     | 1.6 | 34        |
| 2864 | Promoter architecture and transcriptional regulation of Abf1-dependent ribosomal protein genes in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2016, 44, 6113-6126.  | 6.5 | 28        |
| 2865 | A Complex of Htm1 and the Oxidoreductase Pdi1 Accelerates Degradation of Misfolded Glycoproteins. <i>Journal of Biological Chemistry</i> , 2016, 291, 12195-12207.  | 1.6 | 30        |
| 2866 | Mechanism of Regulation of Intrachromatid Recombination and Long-Range Chromosome Interactions in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2016, 36, 1451-1463.  | 1.1 | 7         |
| 2867 | Cox26 is a novel stoichiometric subunit of the yeast cytochrome c oxidase. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2016, 1863, 1624-1632.  | 1.9 | 21        |



| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2868 | Biochemical Analysis of Yeast Suppressor of Ty 4/5 (Spt4/5) Reveals the Importance of Nucleic Acid Interactions in the Prevention of RNA Polymerase II Arrest. <i>Journal of Biological Chemistry</i> , 2016, 291, 9853-9870.                                | 1.6 | 49        |
| 2869 | A Conserved C-terminal Element in the Yeast Doa10 and Human MARCH6 Ubiquitin Ligases Required for Selective Substrate Degradation. <i>Journal of Biological Chemistry</i> , 2016, 291, 12105-12118.  | 1.6 | 36        |
| 2870 | Early Loss of Telomerase Action in Yeast Creates a Dependence on the DNA Damage Response Adaptor Proteins. <i>Molecular and Cellular Biology</i> , 2016, 36, 1908-1919.  | 1.1 | 13        |
| 2871 | The Activity-Dependent Regulation of Protein Kinase Stability by the Localization to P-Bodies. <i>Genetics</i> , 2016, 203, 1191-1202.   | 1.2 | 19        |
| 2872 | Vtc5, a Novel Subunit of the Vacuolar Transporter Chaperone Complex, Regulates Polyphosphate Synthesis and Phosphate Homeostasis in Yeast. <i>Journal of Biological Chemistry</i> , 2016, 291, 22262-22275.  | 1.6 | 67        |
| 2873 | Structure of Bor1 supports an elevator transport mechanism for SLC4 anion exchangers. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 10542-10546.   | 3.3 | 83        |
| 2874 | Transcription factor Reb1 is required for proper transcriptional start site usage at the divergently transcribed TFC6-ESC2 locus in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 2016, 594, 108-116.  | 1.0 | 9         |
| 2875 | Attenuation of transcriptional and signaling responses limits viability of <i>Saccharomyces cerevisiae</i> during periods of glucose deprivation. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2016, 1860, 2563-2575.                           | 1.1 | 2         |
| 2876 | Loss of the SUMO protease Ulp2 triggers a specific multichromosome aneuploidy. <i>Genes and Development</i> , 2016, 30, 1881-1894.   | 2.7 | 37        |
| 2877 | Role of CK2-dependent phosphorylation of Ifh1 and Crf1 in transcriptional regulation of ribosomal protein genes in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2016, 1859, 1004-1013.              | 0.9 | 16        |
| 2878 | Endoplasmic reticulum stress affects the transport of phosphatidylethanolamine from mitochondria to the endoplasmic reticulum in <i>S. cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2016, 1861, 1959-1967. | 1.2 | 7         |
| 2879 | Physical Association of <i>Saccharomyces cerevisiae</i> Polo-like Kinase Cdc5 with Chromosomal Cohesin Facilitates DNA Damage Response. <i>Journal of Biological Chemistry</i> , 2016, 291, 17228-17246.   | 1.6 | 9         |
| 2880 | New families of single integration vectors and gene tagging plasmids for genetic manipulations in budding yeast. <i>Molecular Genetics and Genomics</i> , 2016, 291, 2231-2240.  | 1.0 | 28        |
| 2881 | PCNA Retention on DNA into G2/M Phase Causes Genome Instability in Cells Lacking Elg1. <i>Cell Reports</i> , 2016, 16, 684-695.  | 2.9 | 65        |
| 2882 | Cbr1 is a Dph3 reductase required for the tRNA wobble uridine modification. <i>Nature Chemical Biology</i> , 2016, 12, 995-997.  | 3.9 | 14        |
| 2883 | Simple rules for passive diffusion through the nuclear pore complex. <i>Journal of Cell Biology</i> , 2016, 215, 57-76.  | 2.3 | 337       |
| 2884 | Stimulating <i>S</i> -adenosyl- <i>methionine</i> synthesis extends lifespan via activation of AMPK. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 11913-11918.  | 3.3 | 35        |
| 2885 | Different Regulations of <i>ROM2</i> and <i>LRG1</i> Expression by Ccr4, Pop2, and Dhh1 in the <i>Saccharomyces cerevisiae</i> Cell Wall Integrity Pathway. <i>MSphere</i> , 2016, 1, .  | 1.3 | 12        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 2886 | Multicellularity makes somatic differentiation evolutionarily stable. Proceedings of the National Academy of Sciences of the United States of America, 2016, 113, 8362-8367.   | 3.3  | 18        |
| 2887 | A Targeted Mutation Identified through pK Measurements Indicates a Postrecruitment Role for Fis1 in Yeast Mitochondrial Fission. Journal of Biological Chemistry, 2016, 291, 20329-20344.  | 1.6  | 14        |
| 2888 | Autophagy in <i>Saccharomyces cerevisiae</i> requires the monomeric GTP-binding proteins, Arl1 and Ypt6. Autophagy, 2016, 12, 1721-1737.   | 4.3  | 35        |
| 2889 | Reduced kinase activity of polo kinase Cdc5 affects chromosome stability and DNA damage response in <i>S. cerevisiae</i> . Cell Cycle, 2016, 15, 2906-2919.  | 1.3  | 16        |
| 2890 | Split-ubiquitin yeast two-hybrid interaction reveals a novel interaction between a natural resistance associated macrophage protein and a membrane bound thioredoxin in Brassica juncea. Plant Molecular Biology, 2016, 92, 519-537. | 2.0  | 4         |
| 2891 | Identification of multi-copy suppressors for endoplasmic reticulum-mitochondria tethering proteins in <i>Saccharomyces cerevisiae</i> . FEBS Letters, 2016, 590, 3061-3070.  | 1.3  | 11        |
| 2892 | Kar9 controls the nucleocytoplasmic distribution of yeast EB1. Cell Cycle, 2016, 15, 2860-2866.  | 1.3  | 4         |
| 2893 | Interaction of the cotranslational Hsp70 Ssb with ribosomal proteins and rRNA depends on its lid domain. Nature Communications, 2016, 7, 13563.  | 5.8  | 49        |
| 2894 | Histone Deacetylases with Antagonistic Roles in <i>Saccharomyces cerevisiae</i> Heterochromatin Formation. Genetics, 2016, 204, 177-190.   | 1.2  | 18        |
| 2895 | The DEAD-Box Protein Dhh1p Couples mRNA Decay and Translation by Monitoring Codon Optimality. Cell, 2016, 167, 122-132.e9.   | 13.5 | 232       |
| 2896 | Subnuclear positioning and interchromosomal clustering of the <i>GAL1-10</i> locus are controlled by separable, interdependent mechanisms. Molecular Biology of the Cell, 2016, 27, 2980-2993.                                       | 0.9  | 42        |
| 2897 | Dosage Mutator Genes in <i>Saccharomyces cerevisiae</i> : A Novel Mutator Mode-of-Action of the Mph1 DNA Helicase. Genetics, 2016, 204, 975-986.   | 1.2  | 14        |
| 2898 | Lycopene overproduction in <i>Saccharomyces cerevisiae</i> through combining pathway engineering with host engineering. Microbial Cell Factories, 2016, 15, 113.   | 1.9  | 158       |
| 2899 | Evolving specificity of tRNA 3-methyl-cytidine-32 ( $m^3C32$ ) modification: a subset of tRNAs <sup>Ser</sup> requires $N^6$ -isopentenylation of A37. Rna, 2016, 22, 1400-1410.   | 1.6  | 64        |
| 2900 | Architecture of the 90S Pre-ribosome: A Structural View on the Birth of the Eukaryotic Ribosome. Cell, 2016, 166, 380-393.   | 13.5 | 184       |
| 2901 | Role of the yeast ribosomal protein L16 in ribosome biogenesis. FEBS Journal, 2016, 283, 2968-2985.  | 2.2  | 9         |
| 2902 | The Chromatin Remodeler ISW1 Is a Quality Control Factor that Surveys Nuclear mRNP Biogenesis. Cell, 2016, 167, 1201-1214.e15.   | 13.5 | 34        |
| 2903 | <i>LTE1</i> promotes exit from mitosis by multiple mechanisms. Molecular Biology of the Cell, 2016, 27, 3991-4001.   | 0.9  | 22        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2904 | Ribosome-stalk biogenesis is coupled with recruitment of nuclear-export factor to the nascent 60S subunit. <i>Nature Structural and Molecular Biology</i> , 2016, 23, 1074-1082.   | 3.6 | 36        |
| 2905 | Protein Folding Activity of the Ribosome is Involved in Yeast Prion Propagation. <i>Scientific Reports</i> , 2016, 6, 32117.   | 1.6 | 19        |
| 2906 | Overlapping Role of Respiratory Supercomplex Factor Rcf2 and Its N-terminal Homolog Rcf3 in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2016, 291, 23769-23778.                               | 1.6 | 22        |
| 2907 | The pleiotropic deubiquitinase Ubp3 confers aneuploidy tolerance. <i>Genes and Development</i> , 2016, 30, 2259-2271.  | 2.7 | 22        |
| 2908 | Combined Action of Histone Reader Modules Regulates NuA4 Local Acetyltransferase Function but Not Its Recruitment on the Genome. <i>Molecular and Cellular Biology</i> , 2016, 36, 2768-2781.                                | 1.1 | 32        |
| 2909 | Hsp90 and p23 Molecular Chaperones Control Chromatin Architecture by Maintaining the Functional Pool of the RSC Chromatin Remodeler. <i>Molecular Cell</i> , 2016, 64, 888-899.  | 4.5 | 42        |
| 2910 | A Molecular Titration System Coordinates Ribosomal Protein Gene Transcription with Ribosomal RNA Synthesis. <i>Molecular Cell</i> , 2016, 64, 720-733.   | 4.5 | 47        |
| 2911 | A family of membrane-shaping proteins at ER subdomains regulates pre-peroxisomal vesicle biogenesis. <i>Journal of Cell Biology</i> , 2016, 215, 515-529.  | 2.3 | 74        |
| 2912 | Identification and Characterisation of a pH-stable GFP. <i>Scientific Reports</i> , 2016, 6, 28166.  | 1.6 | 50        |
| 2913 | Phosphorylation of the C-terminal tail of proteasome subunit $\beta 7$ is required for binding of the proteasome quality control factor Ecm29. <i>Scientific Reports</i> , 2016, 6, 27873.                                   | 1.6 | 23        |
| 2914 | The yeast Aft2 transcription factor determines selenite toxicity by controlling the low affinity phosphate transport system. <i>Scientific Reports</i> , 2016, 6, 32836.   | 1.6 | 22        |
| 2915 | Ribosome recycling defects modify the balance between the synthesis and assembly of specific subunits of the oxidative phosphorylation complexes in yeast mitochondria. <i>Nucleic Acids Research</i> , 2016, 44, 5785-5797. | 6.5 | 14        |
| 2916 | Spatial control of lipid droplet proteins by the ERAD ubiquitin ligase Doa10. <i>EMBO Journal</i> , 2016, 35, 1644-1655.   | 3.5 | 65        |
| 2917 | The <i>Saccharomyces cerevisiae</i> poly(A) binding protein Pab1 as a target for eliciting stress tolerant phenotypes. <i>Scientific Reports</i> , 2016, 5, 18318.   | 1.6 | 15        |
| 2918 | Loop 5 region is important for the activity of the long-chain base transporter Rsb1. <i>Journal of Biochemistry</i> , 2016, 161, mvw059.   | 0.9 | 2         |
| 2919 | Regulation of the yeast triacylglycerol lipases Tgl4p and Tgl5p by the presence/absence of nonpolar lipids. <i>Molecular Biology of the Cell</i> , 2016, 27, 2014-2024.  | 0.9 | 14        |
| 2920 | Phosphopeptide binding by Sld3 links Dbf4-dependent kinase to MCM replicative helicase activation. <i>EMBO Journal</i> , 2016, 35, 961-973.  | 3.5 | 122       |
| 2921 | An improved reversibly dimerizing mutant of the FK506-binding protein FKBP. <i>Cellular Logistics</i> , 2016, 6, e1204848.   | 0.9 | 17        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2922 | Applications of pHluorin for Quantitative, Kinetic and High-throughput Analysis of Endocytosis in Budding Yeast. <i>Journal of Visualized Experiments</i> , 2016, , .  | 0.2 | 12        |
| 2923 | Timely Closure of the Prospore Membrane Requires <i>SPS1</i> and <i>SPO77</i> in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2016, 203, 1203-1216.   | 1.2 | 9         |
| 2924 | Recurrent Mutations in the MTOR Regulator RRAGC in Follicular Lymphoma. <i>Clinical Cancer Research</i> , 2016, 22, 5383-5393.   | 3.2 | 36        |
| 2925 | Gad8 Protein Is Found in the Nucleus Where It Interacts with the Mlul Cell Cycle Box-binding Factor (MBF) Transcriptional Complex to Regulate the Response to DNA Replication Stress. <i>Journal of Biological Chemistry</i> , 2016, 291, 9371-9381. | 1.6 | 23        |
| 2926 | Concerted removal of the Erb1-Ytm1 complex in ribosome biogenesis relies on an elaborate interface. <i>Nucleic Acids Research</i> , 2016, 44, 926-939.   | 6.5 | 27        |
| 2927 | Cooperation between SAGA and SWI/SNF complexes is required for efficient transcriptional responses regulated by the yeast MAPK Slt2. <i>Nucleic Acids Research</i> , 2016, 44, gkw324.   | 6.5 | 35        |
| 2928 | The Plasma Membrane Protein Nce102 Implicated in Eisosome Formation Rescues a Heme Defect in Mitochondria. <i>Journal of Biological Chemistry</i> , 2016, 291, 17417-17426.  | 1.6 | 11        |
| 2929 | Lipid droplets maintain lipid homeostasis during anaphase for efficient cell separation in budding yeast. <i>Molecular Biology of the Cell</i> , 2016, 27, 2368-2380.  | 0.9 | 31        |
| 2930 | Yeast Vps13 promotes mitochondrial function and is localized at membrane contact sites. <i>Molecular Biology of the Cell</i> , 2016, 27, 2435-2449.  | 0.9 | 143       |
| 2931 | Kinesin-related Smy1 enhances the Rab-dependent association of myosin-V with secretory cargo. <i>Molecular Biology of the Cell</i> , 2016, 27, 2450-2462.  | 0.9 | 13        |
| 2932 | Telomere Dysfunction Triggers Palindrome Formation Independently of Double-Strand Break Repair Mechanisms. <i>Genetics</i> , 2016, 203, 1659-1668.   | 1.2 | 8         |
| 2933 | Num1 anchors mitochondria to the plasma membrane via two domains with different lipid binding specificities. <i>Journal of Cell Biology</i> , 2016, 213, 513-524.  | 2.3 | 68        |
| 2934 | Fin1-PP1 Helps Clear Spindle Assembly Checkpoint Protein Bub1 from Kinetochores in Anaphase. <i>Cell Reports</i> , 2016, 14, 1074-1085.  | 2.9 | 18        |
| 2935 | Sensing a bud in the yeast morphogenesis checkpoint: a role for Elm1. <i>Molecular Biology of the Cell</i> , 2016, 27, 1764-1775.  | 0.9 | 26        |
| 2936 | Defects in the NC2 repressor affect both canonical and non-coding RNA polymerase II transcription initiation in yeast. <i>BMC Genomics</i> , 2016, 17, 183.  | 1.2 | 9         |
| 2937 | Regulation of the Yeast Hxt6 Hexose Transporter by the Rod1 Arrestin, the Snf1 Protein Kinase, and the Bmh2 14-3-3 Protein. <i>Journal of Biological Chemistry</i> , 2016, 291, 14973-14985.   | 1.6 | 41        |
| 2938 | Radicol-Mediated Inhibition of Topoisomerase VIB-VIA Activity of the Human Malaria Parasite <i>Plasmodium falciparum</i> . <i>MSphere</i> , 2016, 1, .   | 1.3 | 16        |
| 2939 | The K <sup>+</sup> -dependent GTPase Nug1 is implicated in the association of the helicase Dbp10 to the immature peptidyl transferase centre during ribosome maturation. <i>Nucleic Acids Research</i> , 2016, 44, 1800-1812.                        | 6.5 | 36        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2940 | The Dysferlin Domain-Only Protein, Spo73, Is Required for Prospore Membrane Extension in <i>Saccharomyces cerevisiae</i> . <i>MSphere</i> , 2016, 1, .   | 1.3 | 10        |
| 2941 | Distinct types of translation termination generate substrates for ribosome-associated quality control. <i>Nucleic Acids Research</i> , 2016, 44, 6840-6852.  | 6.5 | 17        |
| 2942 | Hsp90 induces increased genomic instability toward DNA-damaging agents by tuning down <i>RAD53</i> transcription. <i>Molecular Biology of the Cell</i> , 2016, 27, 2463-2478.  | 0.9 | 14        |
| 2943 | Protein cofactor competition regulates the action of a multifunctional RNA helicase in different pathways. <i>RNA Biology</i> , 2016, 13, 320-330.   | 1.5 | 39        |
| 2944 | Vcx1-D1 (M383I), the Vcx1 mutant with a calcineurin-independent vacuolar Ca <sup>2+</sup> /H <sup>+</sup> exchanger activity, confers calcineurin-independent Mn <sup>2+</sup> tolerance in <i>Saccharomyces cerevisiae</i> . <i>Canadian Journal of Microbiology</i> , 2016, 62, 475-484. | 0.8 | 3         |
| 2945 | Posttranslational marks control architectural and functional plasticity of the nuclear pore complex basket. <i>Journal of Cell Biology</i> , 2016, 212, 167-180.   | 2.3 | 39        |
| 2946 | Proteasome Impairment Induces Recovery of Mitochondrial Membrane Potential and an Alternative Pathway of Mitochondrial Fusion. <i>Molecular and Cellular Biology</i> , 2016, 36, 347-362.  | 1.1 | 6         |
| 2947 | Ty1 Integrase Interacts with RNA Polymerase III-specific Subcomplexes to Promote Insertion of Ty1 Elements Upstream of Polymerase (Pol) III-transcribed Genes. <i>Journal of Biological Chemistry</i> , 2016, 291, 6396-6411.  | 1.6 | 24        |
| 2948 | Zds1/Zds2-PP2A <sup>Cdc55</sup> complex specifies signaling output from Rho1 GTPase. <i>Journal of Cell Biology</i> , 2016, 212, 51-61.  | 2.3 | 15        |
| 2949 | Efficient engineering of marker-free synthetic allotetraploids of <i>Saccharomyces</i> . <i>Fungal Genetics and Biology</i> , 2016, 89, 10-17.   | 0.9 | 50        |
| 2950 | Contributions of the Prion Protein Sequence, Strain, and Environment to the Species Barrier. <i>Journal of Biological Chemistry</i> , 2016, 291, 1277-1288.  | 1.6 | 23        |
| 2951 | Polymerases $\mu$ and $\alpha$ , repair dysfunctional telomeres facilitated by salt. <i>Nucleic Acids Research</i> , 2016, 44, 3728-3738.  | 6.5 | 1         |
| 2952 | Rpb1 foot mutations demonstrate a major role of Rpb4 in mRNA stability during stress situations in yeast. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2016, 1859, 731-743.   | 0.9 | 23        |
| 2953 | Dynamics of CDKN1A in Single Cells Defined by an Endogenous Fluorescent Tagging Toolkit. <i>Cell Reports</i> , 2016, 14, 1800-1811.  | 2.9 | 85        |
| 2954 | Molecular detection of intrauterine microbial colonization in women with endometriosis. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2016, 199, 69-75.   | 0.5 | 120       |
| 2955 | New integrative modules for multicolor-protein labeling and live-cell imaging in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2016, 16, fow027.  | 1.1 | 22        |
| 2956 | Expression homeostasis during DNA replication. <i>Science</i> , 2016, 351, 1087-1090.  | 6.0 | 101       |
| 2957 | Histone Chaperone Nap1 Is a Major Regulator of Histone H2A-H2B Dynamics at the Inducible GAL Locus. <i>Molecular and Cellular Biology</i> , 2016, 36, 1287-1296.   | 1.1 | 24        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2958 | Hexokinase 2 Is an Intracellular Glucose Sensor of Yeast Cells That Maintains the Structure and Activity of Mig1 Protein Repressor Complex. <i>Journal of Biological Chemistry</i> , 2016, 291, 7267-7285.   | 1.6 | 77        |
| 2959 | Clathrin binding by the adaptor Ent5 promotes late stages of clathrin coat maturation. <i>Molecular Biology of the Cell</i> , 2016, 27, 1143-1153.   | 0.9 | 11        |
| 2960 | Chromosome-Specific and Global Effects of Aneuploidy in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2016, 202, 1395-1409.  | 1.2 | 37        |
| 2961 | Proteasome storage granules are transiently associated with the insoluble protein deposit (IPOD). <i>Journal of Cell Science</i> , 2016, 129, 1190-7.  | 1.2 | 30        |
| 2962 | Combined Interactions of Plant Homeodomain and Chromodomain Regulate NuA4 Activity at DNA Double-Strand Breaks. <i>Genetics</i> , 2016, 202, 77-92.  | 1.2 | 6         |
| 2963 | Adjustable under-expression of yeast mating pathway proteins in <i>Saccharomyces cerevisiae</i> using a programmed ribosomal frameshift. <i>Applied Microbiology and Biotechnology</i> , 2016, 100, 4997-5005.   | 1.7 | 0         |
| 2964 | Regulation of Antisense Transcription by NuA4 Histone Acetyltransferase and Other Chromatin Regulatory Factors. <i>Molecular and Cellular Biology</i> , 2016, 36, 992-1006.  | 1.1 | 12        |
| 2965 | The alternate AP-1 adaptor subunit Apm2 interacts with the Mil1 regulatory protein and confers differential cargo sorting. <i>Molecular Biology of the Cell</i> , 2016, 27, 588-598.   | 0.9 | 16        |
| 2966 | Evolution and Functional Trajectory of Sir1 in Gene Silencing. <i>Molecular and Cellular Biology</i> , 2016, 36, 1164-1179.  | 1.1 | 11        |
| 2967 | Potassium and the K <sup>+</sup> /H <sup>+</sup> Exchanger Kha1p Promote Binding of Copper to ApoFet3p Multi-copper Ferroxidase. <i>Journal of Biological Chemistry</i> , 2016, 291, 9796-9806.  | 1.6 | 20        |
| 2968 | Lipid droplet-mediated ER homeostasis regulates autophagy and cell survival during starvation. <i>Journal of Cell Biology</i> , 2016, 212, 621-631.  | 2.3 | 158       |
| 2969 | The signaling pathways underlying starvation-induced upregulation of $\alpha$ -mannosidase Ams1 in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2016, 1860, 1192-1201.  | 1.1 | 12        |
| 2970 | Characterization of the <i>S. cerevisiae</i> inp51 mutant links phosphatidylinositol 4,5-bisphosphate levels with lipid content, membrane fluidity and cold growth. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2016, 1861, 213-226. | 1.2 | 23        |
| 2971 | The casein kinases Yck1p and Yck2p act in the secretory pathway, in part, by regulating the Rab exchange factor Sec2p. <i>Molecular Biology of the Cell</i> , 2016, 27, 686-701.   | 0.9 | 12        |
| 2972 | The Late S-Phase Transcription Factor Hcm1 Is Regulated through Phosphorylation by the Cell Wall Integrity Checkpoint. <i>Molecular and Cellular Biology</i> , 2016, 36, 941-953.  | 1.1 | 11        |
| 2973 | Synthetic biology and molecular genetics in non-conventional yeasts: Current tools and future advances. <i>Fungal Genetics and Biology</i> , 2016, 89, 126-136.  | 0.9 | 166       |
| 2974 | Ceramide signals for initiation of yeast mating-specific cell cycle arrest. <i>Cell Cycle</i> , 2016, 15, 441-454.   | 1.3 | 12        |
| 2975 | Architecture of the Rix1- <i>Rea1</i> checkpoint machinery during pre-60S-ribosome remodeling. <i>Nature Structural and Molecular Biology</i> , 2016, 23, 37-44.   | 3.6 | 104       |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 2976 | Dynamic relocation of the TORC1-Gtr1/2-Ego1/2/3 complex is regulated by Gtr1 and Gtr2. <i>Molecular Biology of the Cell</i> , 2016, 27, 382-396.   | 0.9 | 59        |
| 2977 | Elucidation of novel budding yeast separase mutants. <i>Bioscience, Biotechnology and Biochemistry</i> , 2016, 80, 473-478.  | 0.6 | 1         |
| 2978 | Slx4 and Rtt107 control checkpoint signalling and DNA resection at double-strand breaks. <i>Nucleic Acids Research</i> , 2016, 44, 669-682.  | 6.5 | 59        |
| 2979 | Structural Heterogeneity in Pre-40S Ribosomes. <i>Structure</i> , 2017, 25, 329-340.   | 1.6 | 60        |
| 2980 | Translational control of lipogenic enzymes in the cell cycle of synchronous, growing yeast cells. <i>EMBO Journal</i> , 2017, 36, 487-502.   | 3.5 | 59        |
| 2981 | Cell wall synthesis and central carbohydrate metabolism are interconnected by the SNF1/Mig1 pathway in <i>Kluyveromyces lactis</i> . <i>European Journal of Cell Biology</i> , 2017, 96, 70-81.        | 1.6 | 10        |
| 2982 | Roles of an N-terminal coiled-coil-containing domain in the localization and function of Bem3, a Rho GTPase-activating protein in budding yeast. <i>Fungal Genetics and Biology</i> , 2017, 99, 40-51. | 0.9 | 5         |
| 2983 | Chromatin potentiates transcription. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, 1536-1541.  | 3.3 | 52        |
| 2984 | Selection and stabilization of endocytic sites by Ede1, a yeast functional homologue of human Eps15. <i>Molecular Biology of the Cell</i> , 2017, 28, 567-575.   | 0.9 | 28        |
| 2985 | Retromer-driven membrane tubulation separates endosomal recycling from Rab7/Ypt7-dependent fusion. <i>Molecular Biology of the Cell</i> , 2017, 28, 783-791.   | 0.9 | 32        |
| 2986 | Adaptation of a Genetic Screen Reveals an Inhibitor for Mitochondrial Protein Import Component Tim44. <i>Journal of Biological Chemistry</i> , 2017, 292, 5429-5442.                                   | 1.6 | 18        |
| 2987 | The budding yeast Polo-like kinase localizes to distinct populations at centrosomes during mitosis. <i>Molecular Biology of the Cell</i> , 2017, 28, 1011-1020.  | 0.9 | 15        |
| 2988 | The novel mitochondrial matrix protease Ste23 is required for efficient presequence degradation and processing. <i>Molecular Biology of the Cell</i> , 2017, 28, 997-1002.                             | 0.9 | 19        |
| 2989 | Predicted RNA Binding Proteins Pes4 and Mip6 Regulate mRNA Levels, Translation, and Localization during Sporulation in Budding Yeast. <i>Molecular and Cellular Biology</i> , 2017, 37, .              | 1.1 | 17        |
| 2990 | Yeast dynamin interaction with ESCRT proteins at the endosome. <i>Cell Biology International</i> , 2017, 41, 484-494.  | 1.4 | 1         |
| 2991 | Ypk1/Ypk2 kinases maintain Rho1 at the plasma membrane by flippase-dependent lipid remodelling after membrane stresses. <i>Journal of Cell Science</i> , 2017, 130, 1169-1178.                         | 1.2 | 26        |
| 2992 | When sensing is gambling: An experimental system reveals how plasticity can generate tunable betting strategies. <i>Evolution; International Journal of Organic Evolution</i> , 2017, 71, 859-871.     | 1.1 | 20        |
| 2993 | Pnc1 piggy-back import into peroxisomes relies on Gpd1 homodimerisation. <i>Scientific Reports</i> , 2017, 7, 42579.   | 1.6 | 16        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 2994 | Synthesis, debugging, and effects of synthetic chromosome consolidation: synVI and beyond. <i>Science</i> , 2017, 355, .   | 6.0  | 184       |
| 2995 | The pre-mRNA retention and splicing complex controls expression of the Mediator subunit Med20. <i>RNA Biology</i> , 2017, 14, 1411-1417.   | 1.5  | 4         |
| 2996 | Regulation of yeast ESCRT-III membrane scission activity by the Doa4 ubiquitin hydrolase. <i>Molecular Biology of the Cell</i> , 2017, 28, 661-672.  | 0.9  | 15        |
| 2997 | Cytosolic proteostasis through importing of misfolded proteins into mitochondria. <i>Nature</i> , 2017, 543, 443-446.  | 13.7 | 363       |
| 2998 | A Ribosomopathy Reveals Decoding Defective Ribosomes Driving Human Dysmorphism. <i>American Journal of Human Genetics</i> , 2017, 100, 506-522.  | 2.6  | 69        |
| 2999 | Coordination of Double Strand Break Repair and Meiotic Progression in Yeast by a Mek1-Ndt80 Negative Feedback Loop. <i>Genetics</i> , 2017, 206, 497-512.  | 1.2  | 49        |
| 3000 | Nmd3 is a structural mimic of $\text{eIF}5\text{A}$ , and activates the $\text{cpGTPase Lsg1}$ during 60S ribosome biogenesis. <i>EMBO Journal</i> , 2017, 36, 854-868.  | 3.5  | 67        |
| 3001 | Nuclear RNA Decay Pathways Aid Rapid Remodeling of Gene Expression in Yeast. <i>Molecular Cell</i> , 2017, 65, 787-800.e5.   | 4.5  | 57        |
| 3002 | Paf1 Has Distinct Roles in Transcription Elongation and Differential Transcript Fate. <i>Molecular Cell</i> , 2017, 65, 685-698.e8.  | 4.5  | 55        |
| 3003 | Overexpression of mitochondrial oxodicarboxylate carrier (ODC1) preserves oxidative phosphorylation in a yeast model of the Barth syndrome. <i>DMM Disease Models and Mechanisms</i> , 2017, 10, 439-450.                    | 1.2  | 10        |
| 3004 | Mitochondrial depolarization in yeast zygotes inhibits clonal expansion of selfish mtDNA. <i>Journal of Cell Science</i> , 2017, 130, 1274-1284.   | 1.2  | 25        |
| 3005 | An mRNA Capping Enzyme Targets FACT to the Active Gene To Enhance the Engagement of RNA Polymerase II into Transcriptional Elongation. <i>Molecular and Cellular Biology</i> , 2017, 37, .                                   | 1.1  | 19        |
| 3006 | Chromatin determinants impart camptothecin sensitivity. <i>EMBO Reports</i> , 2017, 18, 1000-1012.   | 2.0  | 18        |
| 3007 | Protein arginine methylation of Npl3 promotes splicing of the SUS1 intron harboring non-consensus 5â€² splice site and branch site. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2017, 1860, 730-739. | 0.9  | 7         |
| 3008 | MX Cassettes for Knocking Out Genes in Yeast. <i>Cold Spring Harbor Protocols</i> , 2017, 2017, pdb.top080689.   | 0.2  | 3         |
| 3009 | Quantitative high-content imaging identifies novel regulators of Neo1 trafficking at endosomes. <i>Molecular Biology of the Cell</i> , 2017, 28, 1539-1550.  | 0.9  | 31        |
| 3010 | Temperature-dependent regulation of rDNA condensation in <i>Saccharomyces cerevisiae</i> . <i>Cell Cycle</i> , 2017, 16, 1118-1127.  | 1.3  | 16        |
| 3011 | To CURE or not to CURE? Differential effects of the chaperone sorting factor Cur1 on yeast prions are mediated by the chaperone Sis1. <i>Molecular Microbiology</i> , 2017, 105, 242-257.                                    | 1.2  | 26        |



| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3012 | Triosephosphates as intermediates of the Crabtree effect. <i>Biochemistry (Moscow)</i> , 2017, 82, 458-464.   | 0.7 | 3         |
| 3013 | Cfs1p, a Novel Membrane Protein in the PQ-Loop Family, Is Involved in Phospholipid Flippase Functions in Yeast. <i>G3: Genes, Genomes, Genetics</i> , 2017, 7, 179-192.                       | 0.8 | 12        |
| 3014 | Rad6â€œBre1-mediated H2B ubiquitination regulates telomere replication by promoting telomere-end resection. <i>Nucleic Acids Research</i> , 2017, 45, 3308-3322.                              | 6.5 | 28        |
| 3015 | The LSH/HELLS homolog Irc5 contributes to cohesin association with chromatin in yeast. <i>Nucleic Acids Research</i> , 2017, 45, 6404-6416.   | 6.5 | 16        |
| 3016 | The Nucleoporin Nup2 Contains a Meiotic-Autonomous Region that Promotes the Dynamic Chromosome Events of Meiosis. <i>Genetics</i> , 2017, 206, 1319-1337.                                     | 1.2 | 18        |
| 3017 | Phosphoinositide binding profiles of the PX domains of <i>Giardia lamblia</i> . <i>Parasitology International</i> , 2017, 66, 606-614.  | 0.6 | 7         |
| 3018 | NuA4 Lysine Acetyltransferase Complex Contributes to Phospholipid Homeostasis in <i>Saccharomyces cerevisiae</i> . <i>G3: Genes, Genomes, Genetics</i> , 2017, 7, 1799-1809.                  | 0.8 | 7         |
| 3019 | Abf1 and other general regulatory factors control ribosome biogenesis gene expression in budding yeast. <i>Nucleic Acids Research</i> , 2017, 45, 4493-4506.                                  | 6.5 | 41        |
| 3020 | The histone variant H2A.Z promotes efficient cotranscriptional splicing in <i>S. cerevisiae</i> . <i>Genes and Development</i> , 2017, 31, 702-717.   | 2.7 | 35        |
| 3021 | The essential cell division protein FtsN contains a critical disulfide bond in a nonâ€œessential domain. <i>Molecular Microbiology</i> , 2017, 103, 413-422.                                  | 1.2 | 17        |
| 3022 | Histone H4 Facilitates the Proteolysis of the Budding Yeast CENP-ACse4 Centromeric Histone Variant. <i>Genetics</i> , 2017, 205, 113-124.   | 1.2 | 13        |
| 3023 | The histone methyltransferases Set5 and Set1 have overlapping functions in gene silencing and telomere maintenance. <i>Epigenetics</i> , 2017, 12, 93-104.                                    | 1.3 | 22        |
| 3024 | Meiotic Centromere Coupling and Pairing Function by Two Separate Mechanisms in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2017, 205, 657-671.  | 1.2 | 30        |
| 3025 | Expression control of the AMPK regulatory subunit and its functional significance in yeast ER stress response. <i>Scientific Reports</i> , 2017, 7, 46713.                                    | 1.6 | 8         |
| 3026 | Inâ€œVivo Ubiquitin Linkage-type Analysis Reveals that the Cdc48-Rad23/Dsk2 Axis Contributes to K48-Linked Chain Specificity of the Proteasome. <i>Molecular Cell</i> , 2017, 66, 488-502.e7. | 4.5 | 111       |
| 3027 | A New Method, â€œReverse Yeast Two-Hybrid Arrayâ€œ (RYTHA), Identifies Mutants that Dissociate the Physical Interaction Between Elg1 and Slx5. <i>Genetics</i> , 2017, 206, 1683-1697.        | 1.2 | 0         |
| 3028 | Yeast dynamin associates with the GARP tethering complex for endosome-to-Golgi traffic. <i>European Journal of Cell Biology</i> , 2017, 96, 612-621.  | 1.6 | 6         |
| 3029 | Methods to Assess Autophagy and Chronological Aging in Yeast. <i>Methods in Enzymology</i> , 2017, 588, 367-394.  | 0.4 | 20        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 3030 | The CCAAT-Binding Complex Controls Respiratory Gene Expression and Iron Homeostasis in <i>Candida Glabrata</i> . <i>Scientific Reports</i> , 2017, 7, 3531.  | 1.6  | 21        |
| 3031 | The Arl3 and Arl1 GTPases cooperate with Cog8 to regulate selective autophagy via Atg9 trafficking. <i>Traffic</i> , 2017, 18, 580-589.  | 1.3  | 15        |
| 3032 | Self-protective responses to norvaline-induced stress in a leucyl-tRNA synthetase editing-deficient yeast strain. <i>Nucleic Acids Research</i> , 2017, 45, 7367-7381.                                   | 6.5  | 14        |
| 3033 | Transcriptomic analysis of <i>Saccharomyces cerevisiae</i> upon honokiol treatment. <i>Research in Microbiology</i> , 2017, 168, 626-635.  | 1.0  | 12        |
| 3034 | Integrative proteomics and biochemical analyses define Ptc6p as the <i>Saccharomyces cerevisiae</i> pyruvate dehydrogenase phosphatase. <i>Journal of Biological Chemistry</i> , 2017, 292, 11751-11759. | 1.6  | 25        |
| 3035 | Absence of Gim proteins, but not GimC complex, alters stress-induced transcription. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2017, 1860, 773-781.                             | 0.9  | 9         |
| 3036 | Cellular conditions that modulate the fungicidal activity of occidiofungin. <i>Journal of Applied Microbiology</i> , 2017, 123, 380-391.   | 1.4  | 8         |
| 3037 | Mutations in KDSR Cause Recessive Progressive Symmetric Erythrokeratoderma. <i>American Journal of Human Genetics</i> , 2017, 100, 978-984.  | 2.6  | 67        |
| 3038 | The replicative lifespan-extending deletion of <i>SGF73</i> results in altered ribosomal gene expression in yeast. <i>Aging Cell</i> , 2017, 16, 785-796.  | 3.0  | 14        |
| 3039 | Rif1 maintains telomeres and mediates DNA repair by encasing DNA ends. <i>Nature Structural and Molecular Biology</i> , 2017, 24, 588-595.   | 3.6  | 51        |
| 3040 | An ubiquitin-dependent balance between mitofusin turnover and fatty acids desaturation regulates mitochondrial fusion. <i>Nature Communications</i> , 2017, 8, 15832.                                    | 5.8  | 17        |
| 3041 | Systematic analysis of $Ca^{2+}$ homeostasis in <i>Saccharomyces cerevisiae</i> based on chemical-genetic interaction profiles. <i>Molecular Biology of the Cell</i> , 2017, 28, 3415-3427.              | 0.9  | 10        |
| 3042 | Epigenetic Transcriptional Memory of GAL Genes Depends on Growth in Glucose and the Tup1 Transcription Factor in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2017, 206, 1895-1907.               | 1.2  | 35        |
| 3043 | Analysis of Meiotic Chromosome-Associated Protein Dynamics Using Conditional Expression in Budding Yeast. <i>Methods in Molecular Biology</i> , 2017, 1471, 157-174.                                     | 0.4  | 2         |
| 3044 | Fungicide Xylariasp. BCC 1067 extract induces reactive oxygen species and activates multidrug resistance system in <i>Saccharomyces cerevisiae</i> . <i>Future Microbiology</i> , 2017, 12, 417-440.     | 1.0  | 7         |
| 3045 | Mitophagy in Yeast: A Screen of Mitophagy-Deficient Mutants. <i>Methods in Molecular Biology</i> , 2017, 1759, 95-104.   | 0.4  | 1         |
| 3046 | Introducing MX Cassettes into <i>Saccharomyces cerevisiae</i> . <i>Cold Spring Harbor Protocols</i> , 2017, 2017, pdb.prot088104.  | 0.2  | 5         |
| 3047 | Aneuploidy Causes Non-genetic Individuality. <i>Cell</i> , 2017, 169, 229-242.e21.   | 13.5 | 81        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3048 | Conserved Atg8 recognition sites mediate Atg4 association with autophagosomal membranes and Atg8 deconjugation. <i>EMBO Reports</i> , 2017, 18, 765-780.  | 2.0 | 59        |
| 3049 | Protein Kinase C Controls Binding of Igo/ENSA Proteins to Protein Phosphatase 2A in Budding Yeast. <i>Journal of Biological Chemistry</i> , 2017, 292, 4925-4941.   | 1.6 | 13        |
| 3050 | A Metabolic Function for Phospholipid and Histone Methylation. <i>Molecular Cell</i> , 2017, 66, 180-193.e8.  | 4.5 | 189       |
| 3051 | Yeast cells as an assay system for in vivo O -GlcNAc modification. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2017, 1861, 1159-1167.   | 1.1 | 10        |
| 3052 | Deleting the <sc>DAG</sc> kinase Dgk1 augments yeast vacuole fusion through increased Ypt7 activity and altered membrane fluidity. <i>Traffic</i> , 2017, 18, 315-329.  | 1.3 | 43        |
| 3053 | Conserved role for Gga proteins in phosphatidylinositol 4-kinase localization to the <i>trans</i>-Golgi network. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, 3433-3438. | 3.3 | 52        |
| 3054 | A conserved inter-domain communication mechanism regulates the ATPase activity of the AAA-protein Drg1. <i>Scientific Reports</i> , 2017, 7, 44751.   | 1.6 | 4         |
| 3055 | Yeast dynamin Vps1 associates with clathrin to facilitate vesicular trafficking and controls Golgi homeostasis. <i>European Journal of Cell Biology</i> , 2017, 96, 182-197.  | 1.6 | 6         |
| 3056 | An engineered minimal chromosomal passenger complex reveals a role for INCENP/Sli15 spindle association in chromosome biorientation. <i>Journal of Cell Biology</i> , 2017, 216, 911-923.                                       | 2.3 | 25        |
| 3057 | Deubiquitinating enzymes Ubp2 and Ubp15 regulate endocytosis by limiting ubiquitination and degradation of ARTs. <i>Molecular Biology of the Cell</i> , 2017, 28, 1271-1283.  | 0.9 | 32        |
| 3058 | Congenital valvular defects associated with deleterious mutations in the PLD1 gene. <i>Journal of Medical Genetics</i> , 2017, 54, 278-286.   | 1.5 | 36        |
| 3059 | The Rim101 pathway contributes to ER stress adaptation through sensing the state of plasma membrane. <i>Biochemical Journal</i> , 2017, 474, 51-63.   | 1.7 | 14        |
| 3060 | Distinct complexes of yeast Snx4 family <sc>SNXâ€BARs</sc> mediate retrograde trafficking of Snc1 and Atg27. <i>Traffic</i> , 2017, 18, 134-144.  | 1.3 | 56        |
| 3061 | Interactions between intersubunit transmembrane domains regulate the chaperone-dependent degradation of an oligomeric membrane protein. <i>Biochemical Journal</i> , 2017, 474, 357-376.  | 1.7 | 23        |
| 3062 | Production of encapsulated creatinase using yeast spores. <i>Bioengineered</i> , 2017, 8, 411-419.  | 1.4 | 5         |
| 3063 | Lipid droplets are central organelles for meiosis II progression during yeast sporulation. <i>Molecular Biology of the Cell</i> , 2017, 28, 440-451.  | 0.9 | 26        |
| 3064 | Pex35 is a regulator of peroxisome abundance. <i>Journal of Cell Science</i> , 2017, 130, 791-804.  | 1.2 | 34        |
| 3065 | Premature Silencing of the Spindle Assembly Checkpoint Is Prevented by the Bub1-H2A-Sgo1-PP2A Axis in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2017, 205, 1169-1178.   | 1.2 | 12        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 3066 | <i>ICE</i> plasmids for generating terminal 3xFLAG tagged genes that allow inducible, constitutive or endogenous expression in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2017, 34, 223-235.                                   | 0.8  | 13        |
| 3067 | Dynamic localization of a yeast development-specific PP1 complex during prospore membrane formation is dependent on multiple localization signals and complex formation. <i>Molecular Biology of the Cell</i> , 2017, 28, 3881-3895. | 0.9  | 9         |
| 3068 | A tethering complex drives the terminal stage of SNARE-dependent membrane fusion. <i>Nature</i> , 2017, 551, 634-638.  | 13.7 | 92        |
| 3069 | Rewiring a Rab regulatory network reveals a possible inhibitory role for the vesicle tether, <i>Uso1</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E8637-E8645.          | 3.3  | 17        |
| 3070 | <i>Ecm22</i> and <i>Upc2</i> regulate yeast mating through control of expression of the mating genes <i>PRM1</i> and <i>PRM4</i> . <i>Biochemical and Biophysical Research Communications</i> , 2017, 493, 1485-1490.                | 1.0  | 3         |
| 3071 | <i>Pib2</i> and EGO Complex are both required for activation of TORC1. <i>Journal of Cell Science</i> , 2017, 130, 3878-3890.  | 1.2  | 41        |
| 3072 | The duration of mitosis and daughter cell size are modulated by nutrients in budding yeast. <i>Journal of Cell Biology</i> , 2017, 216, 3463-3470.   | 2.3  | 57        |
| 3073 | Structure and RNA recognition of ribosome assembly factor <i>Utp30</i> . <i>Rna</i> , 2017, 23, 1936-1945.   | 1.6  | 1         |
| 3074 | A conserved signaling network monitors delivery of sphingolipids to the plasma membrane in budding yeast. <i>Molecular Biology of the Cell</i> , 2017, 28, 2589-2599.  | 0.9  | 28        |
| 3075 | The Kinetochore Receptor for the Cohesin Loading Complex. <i>Cell</i> , 2017, 171, 72-84.e13.  | 13.5 | 88        |
| 3076 | Natively Unfolded FG Repeats Stabilize the Structure of the Nuclear Pore Complex. <i>Cell</i> , 2017, 171, 904-917.e19.  | 13.5 | 99        |
| 3077 | Molecular insights into the m-AAA protease-mediated dislocation of transmembrane helices in the mitochondrial inner membrane. <i>Journal of Biological Chemistry</i> , 2017, 292, 20058-20066.                                       | 1.6  | 9         |
| 3078 | Molecular basis for inner kinetochore configuration through <i>RWD</i> domain-peptide interactions. <i>EMBO Journal</i> , 2017, 36, 3458-3482.   | 3.5  | 34        |
| 3079 | Late rDNA Condensation Ensures Timely <i>Cdc14</i> Release and Coordination of Mitotic Exit Signaling with Nucleolar Segregation. <i>Current Biology</i> , 2017, 27, 3248-3263.e5.   | 1.8  | 26        |
| 3080 | Proteomic identification of histone post-translational modifications and proteins enriched at a DNA double-strand break. <i>Nucleic Acids Research</i> , 2017, 45, 10923-10940.  | 6.5  | 12        |
| 3081 | Evidence for ESCRT- and clathrin-dependent microautophagy. <i>Journal of Cell Biology</i> , 2017, 216, 3263-3274.  | 2.3  | 127       |
| 3082 | Mitochondria-driven assembly of a cortical anchor for mitochondria and dynein. <i>Journal of Cell Biology</i> , 2017, 216, 3061-3071.  | 2.3  | 41        |
| 3083 | Microtubule cross-linking activity of <i>She1</i> ensures spindle stability for spindle positioning. <i>Journal of Cell Biology</i> , 2017, 216, 2759-2775.  | 2.3  | 9         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3084 | The impact of nonpolar lipids on the regulation of the steryl ester hydrolases Tgl1p and Yeh1p in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2017, 1862, 1491-1501.                            | 1.2 | 2         |
| 3085 | Hof1 and Chs4 Interact via F-BAR Domain and Sel1-like Repeats to Control Extracellular Matrix Deposition during Cytokinesis. <i>Current Biology</i> , 2017, 27, 2878-2886.e5.   | 1.8 | 19        |
| 3086 | Diversification of Transcriptional Regulation Determines Subfunctionalization of Paralogous Branched Chain Aminotransferases in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2017, 207, 975-991.   | 1.2 | 27        |
| 3087 | Firing of Replication Origins Frees Dbf4-Cdc7 to Target Eco1 for Destruction. <i>Current Biology</i> , 2017, 27, 2849-2855.e2.  | 1.8 | 18        |
| 3088 | GATOR1 regulates nitrogenic cataplerotic reactions of the mitochondrial TCA cycle. <i>Nature Chemical Biology</i> , 2017, 13, 1179-1186.  | 3.9 | 16        |
| 3089 | Cytosine deamination and base excision repair cause R-loop-induced CAG repeat fragility and instability in <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E8392-E8401.                 | 3.3 | 72        |
| 3090 | Interaction between repressor Opi1p and ER membrane protein Scs2p facilitates transit of phosphatidic acid from the ER to mitochondria and is essential for INO1 gene expression in the presence of choline. <i>Journal of Biological Chemistry</i> , 2017, 292, 18713-18728. | 1.6 | 19        |
| 3091 | Phosphorylation and nuclear transit modulate the balance between normal function and terminal aggregation of the yeast RNA-binding protein Ssd1. <i>Molecular Biology of the Cell</i> , 2017, 28, 3057-3069.  | 0.9 | 11        |
| 3092 | Yeast silencing factor Sir4 and a subset of nucleoporins form a complex distinct from nuclear pore complexes. <i>Journal of Cell Biology</i> , 2017, 216, 3145-3159.  | 2.3 | 40        |
| 3093 | The chromatin remodeling Isw1a complex is regulated by SUMOylation. <i>Biochemical Journal</i> , 2017, 474, 3455-3469.  | 1.7 | 3         |
| 3094 | The phosphorylation of a kinetochore protein Dam1 by Aurora B/Ipl1 kinase promotes chromosome bipolar attachment in yeast. <i>Scientific Reports</i> , 2017, 7, 11880.  | 1.6 | 11        |
| 3095 | Endonucleolytic cleavage in the expansion segment 7 of 25S rRNA is an early marker of low-level oxidative stress in yeast. <i>Journal of Biological Chemistry</i> , 2017, 292, 18469-18485.   | 1.6 | 27        |
| 3096 | A membrane-inserted structural model of the yeast mitofusin Fzo1. <i>Scientific Reports</i> , 2017, 7, 10217.   | 1.6 | 25        |
| 3097 | The ATPase Fap7 Tests the Ability to Carry Out Translocation-like Conformational Changes and Releases Dim1 during 40S Ribosome Maturation. <i>Molecular Cell</i> , 2017, 67, 990-1000.e3.   | 4.5 | 48        |
| 3098 | The diferric-tyrosyl radical cluster of ribonucleotide reductase and cytosolic iron-sulfur clusters have distinct and similar biogenesis requirements. <i>Journal of Biological Chemistry</i> , 2017, 292, 11445-11451.   | 1.6 | 19        |
| 3099 | <i>1,6</i> -glucan synthesis-associated genes are required for proper spore wall formation in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2017, 34, 431-446.   | 0.8 | 12        |
| 3100 | Nineteen complex-related factor Prp45 is required for the early stages of cotranscriptional spliceosome assembly. <i>Rna</i> , 2017, 23, 1512-1524.   | 1.6 | 7         |
| 3101 | Glycolytic Enzymes Coalesce in G Bodies under Hypoxic Stress. <i>Cell Reports</i> , 2017, 20, 895-908.  | 2.9 | 139       |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 3102 | The forkhead-like transcription factor (Fhl1p) maintains yeast replicative lifespan by regulating ribonucleotide reductase 1 ( RNR1 ) gene transcription. <i>Biochemical and Biophysical Research Communications</i> , 2017, 488, 218-223.  | 1.0  | 7         |
| 3103 | Novel function of a dynein light chain in actin assembly during clathrin-mediated endocytosis. <i>Journal of Cell Biology</i> , 2017, 216, 2565-2580.   | 2.3  | 14        |
| 3104 | Syp1 regulates the clathrin-mediated and clathrin-independent endocytosis of multiple cargo proteins through a novel sorting motif. <i>Molecular Biology of the Cell</i> , 2017, 28, 2434-2448.   | 0.9  | 13        |
| 3105 | Phosphatidylinositol 3,5-bisphosphate is involved in methylglyoxal-induced activation of the Mpk1 mitogen-activated protein kinase cascade in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2017, 292, 15039-15048.                                      | 1.6  | 8         |
| 3106 | Recombination at subtelomeres is regulated by physical distance, double-strand break resection and chromatin status. <i>EMBO Journal</i> , 2017, 36, 2609-2625.   | 3.5  | 46        |
| 3107 | Ubiquitination of stalled ribosome triggers ribosome-associated quality control. <i>Nature Communications</i> , 2017, 8, 159.   | 5.8  | 249       |
| 3108 | Identification and characterization of roles for Puf1 and Puf2 proteins in the yeast response to high calcium. <i>Scientific Reports</i> , 2017, 7, 3037.   | 1.6  | 15        |
| 3109 | Pwp2 mediates UTP-B assembly via two structurally independent domains. <i>Scientific Reports</i> , 2017, 7, 3169.   | 1.6  | 9         |
| 3110 | Rps26 directs mRNA-specific translation by recognition of Kozak sequence elements. <i>Nature Structural and Molecular Biology</i> , 2017, 24, 700-707.  | 3.6  | 118       |
| 3111 | Direct interaction of the Golgi V-ATPase a-subunit isoform with PI(4)P drives localization of Golgi V-ATPases in yeast. <i>Molecular Biology of the Cell</i> , 2017, 28, 2518-2530.   | 0.9  | 44        |
| 3112 | In vitro reconstitution of the yeast spore wall dityrosine layer discloses the mechanism of its assembly. <i>Journal of Biological Chemistry</i> , 2017, 292, 15880-15891.  | 1.6  | 8         |
| 3113 | The SAGA complex, together with transcription factors and the endocytic protein Rvs167p, coordinates the reprofiling of gene expression in response to changes in sterol composition in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2017, 28, 2637-2649. | 0.9  | 11        |
| 3114 | CAT-tailing as a fail-safe mechanism for efficient degradation of stalled nascent polypeptides. <i>Science</i> , 2017, 357, 414-417.  | 6.0  | 113       |
| 3115 | Regulation of hetDNA Length during Mitotic Double-Strand Break Repair in Yeast. <i>Molecular Cell</i> , 2017, 67, 539-549.e4.   | 4.5  | 36        |
| 3116 | Ubiquitin orchestrates proteasome dynamics between proliferation and quiescence in yeast. <i>Molecular Biology of the Cell</i> , 2017, 28, 2479-2491.   | 0.9  | 41        |
| 3117 | The golgin protein Coy1 functions in intra-Golgi retrograde transport and interacts with the COG complex and Golgi SNAREs. <i>Molecular Biology of the Cell</i> , 2017, 28, 2686-2700.  | 0.9  | 12        |
| 3118 | A Functional Link Between Bir1 and the <i>Saccharomyces cerevisiae</i> Ctf19 Kinetochore Complex Revealed Through Quantitative Fitness Analysis. <i>G3: Genes, Genomes, Genetics</i> , 2017, 7, 3203-3215.  | 0.8  | 5         |
| 3119 | Visualizing the Assembly Pathway of Nucleolar Pre-60S Ribosomes. <i>Cell</i> , 2017, 171, 1599-1610.e14.  | 13.5 | 162       |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3120 | Distinct roles for plasma membrane PtdIns(4)P and PtdIns(4,5)P <sub>2</sub> during yeast receptor-mediated endocytosis. <i>Journal of Cell Science</i> , 2018, 131, .   | 1.2 | 14        |
| 3121 | Repression of Middle Sporulation Genes in <i>Saccharomyces cerevisiae</i> by the Sum1-Rfm1-Hst1 Complex Is Maintained by Set1 and H3K4 Methylation. <i>G3: Genes, Genomes, Genetics</i> , 2017, 7, 3971-3982.       | 0.8 | 13        |
| 3122 | Stress-response transcription factors Msn2 and Msn4 couple TORC2-Ypk1 signaling and mitochondrial respiration to <i>ATG8</i> gene expression and autophagy. <i>Autophagy</i> , 2017, 13, 1804-1812.                 | 4.3 | 24        |
| 3123 | Preribosomes escaping from the nucleus are caught during translation by cytoplasmic quality control. <i>Nature Structural and Molecular Biology</i> , 2017, 24, 1107-1115.  | 3.6 | 35        |
| 3124 | Genetic and Epigenetic Strategies Potentiate Gal4 Activation to Enhance Fitness in Recently Diverged Yeast Species. <i>Current Biology</i> , 2017, 27, 3591-3602.e3.  | 1.8 | 18        |
| 3125 | Long-Chain Polyprenols Promote Spore Wall Formation in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2017, 207, 1371-1386.  | 1.2 | 18        |
| 3126 | GSK-3 $\beta$ Homolog Rim11 and the Histone Deacetylase Complex Ume6-Sin3-Rpd3 Are Involved in Replication Stress Response Caused by Defects in Dna2. <i>Genetics</i> , 2017, 206, 829-842.                         | 1.2 | 4         |
| 3127 | Identification and Mode of Action of a Plant Natural Product Targeting Human Fungal Pathogens. <i>Antimicrobial Agents and Chemotherapy</i> , 2017, 61, .   | 1.4 | 35        |
| 3128 | Nucleolin directly mediates Epstein-Barr virus immune evasion through binding to G-quadruplexes of EBNA1 mRNA. <i>Nature Communications</i> , 2017, 8, 16043.   | 5.8 | 94        |
| 3129 | Sequential phosphorylation of CST subunits by different cyclin-Cdk1 complexes orchestrate telomere replication. <i>Cell Cycle</i> , 2017, 16, 1271-1287.  | 1.3 | 18        |
| 3130 | Determining if an mRNA is a Substrate of Nonsense-Mediated mRNA Decay in <i>Saccharomyces cerevisiae</i> . <i>Methods in Molecular Biology</i> , 2017, 1507, 169-177.   | 0.4 | 8         |
| 3131 | Ubiquitination of ERMES components by the E3 ligase Rsp5 is involved in mitophagy. <i>Autophagy</i> , 2017, 13, 114-132.  | 4.3 | 39        |
| 3132 | Mitochondrial Superoxide Dismutase and Yap1p Act as a Signaling Module Contributing to Ethanol Tolerance of the Yeast <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 2017, 83, . | 1.4 | 22        |
| 3133 | Consecutive hydrolysis of creatinine using creatininase and creatinase encapsulated in <i>Saccharomyces cerevisiae</i> spores. <i>Biotechnology Letters</i> , 2017, 39, 261-267.                                    | 1.1 | 2         |
| 3134 | Structural studies of protein arginine methyltransferase 2 reveal its interactions with potential substrates and inhibitors. <i>FEBS Journal</i> , 2017, 284, 77-96.  | 2.2 | 25        |
| 3135 | Transcription of the mating-type-regulated lncRNA IRT1 is governed by TORC1 and PKA. <i>Current Genetics</i> , 2017, 63, 325-329.   | 0.8 | 11        |
| 3136 | Cargo selectivity of yeast sorting nexins. <i>Traffic</i> , 2017, 18, 110-122.  | 1.3 | 43        |
| 3137 | Interdependent action of KH domain proteins Krr1 and Dim2 drive the 40S platform assembly. <i>Nature Communications</i> , 2017, 8, 2213.  | 5.8 | 38        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3138 | Bypass of Activation Loop Phosphorylation by Aspartate 836 in Activation of the Endoribonuclease Activity of Ire1. <i>Molecular and Cellular Biology</i> , 2017, 37, .  | 1.1 | 12        |
| 3139 | She1 affects dynein through direct interactions with the microtubule and the dynein microtubule-binding domain. <i>Nature Communications</i> , 2017, 8, 2151.   | 5.8 | 25        |
| 3140 | The assembly factor Erb1 functions in multiple remodeling events during 60S ribosomal subunit assembly in <i>S. cerevisiae</i> . <i>Nucleic Acids Research</i> , 2017, 45, gkw1361.   | 6.5 | 16        |
| 3141 | The RES complex is required for efficient transformation of the precatalytic B spliceosome into an activated B <sup>act</sup> complex. <i>Genes and Development</i> , 2017, 31, 2416-2429.  | 2.7 | 18        |
| 3142 | A new method for post-translationally labeling proteins in live cells for fluorescence imaging and tracking. <i>Protein Engineering, Design and Selection</i> , 2017, 30, 771-780.  | 1.0 | 18        |
| 3143 | Cdc14 phosphatase directs centrosome re-duplication at the meiosis I to meiosis II transition in budding yeast. <i>Wellcome Open Research</i> , 2017, 2, 2.   | 0.9 | 20        |
| 3144 | New plasmids for the disruption and repeated use of selection markers in <i>Saccharomyces cerevisiae</i> . <i>Journal of General and Applied Microbiology</i> , 2017, 63, 199-202.  | 0.4 | 1         |
| 3145 | AtOMA1 Affects the OXPHOS System and Plant Growth in Contrast to Other Newly Identified ATP-Independent Proteases in Arabidopsis Mitochondria. <i>Frontiers in Plant Science</i> , 2017, 8, 1543.                                   | 1.7 | 17        |
| 3146 | Assays to Monitor Autophagy in <i>Saccharomyces cerevisiae</i> . <i>Cells</i> , 2017, 6, 23.  | 1.8 | 53        |
| 3147 | A Novel Assay Reveals a Maturation Process during Ascospore Wall Formation. <i>Journal of Fungi (Basel, Switzerland)</i> , 2017, 3, 54.   | 1.5 | 5         |
| 3148 | Investigating symmetry breaking in yeast. <i>Methods in Cell Biology</i> , 2017, 139, 23-50.  | 0.5 | 0         |
| 3149 | CRISPR-UnLOCK: Multipurpose Cas9-Based Strategies for Conversion of Yeast Libraries and Strains. <i>Frontiers in Microbiology</i> , 2017, 8, 1773.  | 1.5 | 21        |
| 3150 | Utilizing Biotinylated Proteins Expressed in Yeast to Visualize DNA-Protein Interactions at the Single-Molecule Level. <i>Frontiers in Microbiology</i> , 2017, 8, 2062.  | 1.5 | 6         |
| 3151 | E3 ubiquitin ligase Bre1 couples sister chromatid cohesion establishment to DNA replication in <i>Saccharomyces cerevisiae</i> . <i>ELife</i> , 2017, 6, .  | 2.8 | 21        |
| 3152 | The AAA protein Msp1 mediates clearance of excess tail-anchored proteins from the peroxisomal membrane. <i>ELife</i> , 2017, 6, .   | 2.8 | 65        |
| 3153 | A CDC25 family protein phosphatase gates cargo recognition by the Vps26 retromer subunit. <i>ELife</i> , 2017, 6, .   | 2.8 | 14        |
| 3154 | ATP synthesis in the energy metabolism pathway: a new perspective for manipulating CdSe quantum dots biosynthesized in <i>Saccharomyces cerevisiae</i> . <i>International Journal of Nanomedicine</i> , 2017, Volume 12, 3865-3879. | 3.3 | 10        |
| 3155 | A method for labeling proteins with tags at the native genomic loci in budding yeast. <i>PLoS ONE</i> , 2017, 12, e0176184.   | 1.1 | 10        |



| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3156 | Small molecule inhibitors uncover synthetic genetic interactions of human flap endonuclease 1 (FEN1) with DNA damage response genes. <i>PLoS ONE</i> , 2017, 12, e0179278.                                     | 1.1 | 36        |
| 3157 | Activation of Dun1 in response to nuclear DNA instability accounts for the increase in mitochondrial point mutations in Rad27/FEN1 deficient <i>S. cerevisiae</i> . <i>PLoS ONE</i> , 2017, 12, e0180153.      | 1.1 | 2         |
| 3158 | Mpp10 represents a platform for the interaction of multiple factors within the 90S pre-ribosome. <i>PLoS ONE</i> , 2017, 12, e0183272.   | 1.1 | 15        |
| 3159 | Developmentally regulated internal transcription initiation during meiosis in budding yeast. <i>PLoS ONE</i> , 2017, 12, e0188001.   | 1.1 | 16        |
| 3160 | Chimeric proteins tagged with specific 3xHA cassettes may present instability and functional problems. <i>PLoS ONE</i> , 2017, 12, e0183067.   | 1.1 | 19        |
| 3161 | Lysine acetyltransferase NuA4 and acetyl-CoA regulate glucose-deprived stress granule formation in <i>Saccharomyces cerevisiae</i> . <i>PLoS Genetics</i> , 2017, 13, e1006626.                                | 1.5 | 20        |
| 3162 | The T-cell leukemia related rpl10-R98S mutant traps the 60S export adapter Nmd3 in the ribosomal P site in yeast. <i>PLoS Genetics</i> , 2017, 13, e1006894.   | 1.5 | 19        |
| 3163 | Reduced dosage of the chromosome axis factor Red1 selectively disrupts the meiotic recombination checkpoint in <i>Saccharomyces cerevisiae</i> . <i>PLoS Genetics</i> , 2017, 13, e1006928.                    | 1.5 | 11        |
| 3164 | The translesion DNA polymerases Pol $\eta$ and Rev1 are activated independently of PCNA ubiquitination upon UV radiation in mutants of DNA polymerase $\epsilon$ . <i>PLoS Genetics</i> , 2017, 13, e1007119.  | 1.5 | 22        |
| 3165 | Development of an in silico method for the identification of subcomplexes involved in the biogenesis of multiprotein complexes in <i>Saccharomyces cerevisiae</i> . <i>BMC Systems Biology</i> , 2017, 11, 67. | 3.0 | 1         |
| 3166 | Atg4 proteolytic activity can be inhibited by Atg1 phosphorylation. <i>Nature Communications</i> , 2017, 8, 295.   | 5.8 | 70        |
| 3167 | Not5-dependent co-translational assembly of Ada2 and Spt20 is essential for functional integrity of SAGA. <i>Nucleic Acids Research</i> , 2017, 45, 1186-1199.   | 6.5 | 36        |
| 3168 | Monitoring the Formation of Autophagosomal Precursor Structures in Yeast <i>Saccharomyces cerevisiae</i> . <i>Methods in Enzymology</i> , 2017, 588, 323-365.  | 0.4 | 2         |
| 3169 | Defects in methylation of arginine 37 on CENP-A/Cse4 are compensated by the ubiquitin ligase complex Ubr2/Mub1. <i>FEMS Yeast Research</i> , 2017, 17, .   | 1.1 | 4         |
| 3170 | Biochemical characterization of the helicase Sen1 provides new insights into the mechanisms of non-coding transcription termination. <i>Nucleic Acids Research</i> , 2017, 45, 1355-1370.                      | 6.5 | 52        |
| 3171 | Role of the Hof1-Cyk3 interaction in cleavage-furrow ingression and primary-septum formation during yeast cytokinesis. <i>Molecular Biology of the Cell</i> , 2018, 29, 597-609.                               | 0.9 | 13        |
| 3172 | The exoribonuclease Xrn1 is a post-transcriptional negative regulator of autophagy. <i>Autophagy</i> , 2018, 14, 898-912.  | 4.3 | 30        |
| 3173 | Stu2 acts as a microtubule destabilizer in metaphase budding yeast spindles. <i>Molecular Biology of the Cell</i> , 2018, 29, 247-255.   | 0.9 | 15        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3174 | Autophagy balances mtDNA synthesis and degradation by DNA polymerase POLG during starvation. <i>Journal of Cell Biology</i> , 2018, 217, 1601-1611.   | 2.3 | 47        |
| 3175 | A regulatory circuit of two lncRNAs and a master regulator directs cell fate in yeast. <i>Nature Communications</i> , 2018, 9, 780.   | 5.8 | 38        |
| 3176 | Maturation of the 90S pre-ribosome requires Mrd1 dependent U3 snoRNA and 35S pre-rRNA structural rearrangements. <i>Nucleic Acids Research</i> , 2018, 46, 3692-3706.   | 6.5 | 8         |
| 3177 | Analysis of the thresholds for transcriptional activation by the yeast MAP kinases Fus3 and Kss1. <i>Molecular Biology of the Cell</i> , 2018, 29, 669-682.   | 0.9 | 13        |
| 3178 | Turnover of aberrant pre-40S pre-ribosomal particles is initiated by a novel endonucleolytic decay pathway. <i>Nucleic Acids Research</i> , 2018, 46, 4699-4714.  | 6.5 | 15        |
| 3179 | Yeast Aim21/Tda2 both regulates free actin by reducing barbed end assembly and forms a complex with Cap1/Cap2 to balance actin assembly between patches and cables. <i>Molecular Biology of the Cell</i> , 2018, 29, 923-936. | 0.9 | 19        |
| 3180 | Control of vacuole membrane homeostasis by a resident PI-3,5-kinase inhibitor. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018, 115, 4684-4689.                                 | 3.3 | 19        |
| 3181 | Lipid droplet consumption is functionally coupled to vacuole homeostasis independent of lipophagy. <i>Journal of Cell Science</i> , 2018, 131, .  | 1.2 | 26        |
| 3182 | Hierarchical recruitment of ribosomal proteins and assembly factors remodels nucleolar pre-60S ribosomes. <i>Journal of Cell Biology</i> , 2018, 217, 2503-2518.  | 2.3 | 33        |
| 3183 | An interplay between Shugoshin and Spo13 for centromeric cohesin protection and sister kinetochore mono-orientation during meiosis I in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2018, 64, 1141-1152.      | 0.8 | 9         |
| 3184 | Visualizing multiple inter-organelle contact sites using the organelle-targeted split-GFP system. <i>Scientific Reports</i> , 2018, 8, 6175.  | 1.6 | 85        |
| 3185 | The Functional Specialization of Exomer as a Cargo Adaptor During the Evolution of Fungi. <i>Genetics</i> , 2018, 208, 1483-1498.   | 1.2 | 8         |
| 3186 | Osw2 is required for proper assembly of glucan and/or mannan layers of the yeast spore wall. <i>Journal of Biochemistry</i> , 2018, 163, 293-304.   | 0.9 | 4         |
| 3187 | A Flow Cytometry-Based Phenotypic Screen To Identify Novel Endocytic Factors in <i>Saccharomyces cerevisiae</i> . <i>G3: Genes, Genomes, Genetics</i> , 2018, 8, 1497-1512.   | 0.8 | 5         |
| 3188 | Regulation of yeast fatty acid desaturase in response to iron deficiency. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2018, 1863, 657-668.  | 1.2 | 29        |
| 3190 | Development of a yeast internal-subunit eGFP labeling strategy and its application in subunit identification in eukaryotic group II chaperonin TRiC/CCT. <i>Scientific Reports</i> , 2018, 8, 2374.                           | 1.6 | 17        |
| 3191 | Cdc48 and ubiquilins confer selective anterograde protein sorting and entry into the multivesicular body in yeast. <i>Molecular Biology of the Cell</i> , 2018, 29, 948-963.  | 0.9 | 9         |
| 3192 | Antisense transcription-dependent chromatin signature modulates sense transcript dynamics. <i>Molecular Systems Biology</i> , 2018, 14, e8007.  | 3.2 | 42        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3193 | Cdc48/VCP Promotes Chromosome Morphogenesis by Releasing Condensin from Self-Entrapment in Chromatin. <i>Molecular Cell</i> , 2018, 69, 664-676.e5.   | 4.5 | 53        |
| 3194 | Differential abundance and transcription of 14-3-3 proteins during vegetative growth and sexual reproduction in budding yeast. <i>Scientific Reports</i> , 2018, 8, 2145.   | 1.6 | 5         |
| 3195 | Stu2 uses a 15-nm parallel coiled coil for kinetochore localization and concomitant regulation of the mitotic spindle. <i>Molecular Biology of the Cell</i> , 2018, 29, 285-294.  | 0.9 | 5         |
| 3196 | Membrane protein insertion through a mitochondrial $\beta$ -barrel gate. <i>Science</i> , 2018, 359, .  | 6.0 | 111       |
| 3197 | Heterologous Expression of a Novel Drug Transporter from the Malaria Parasite Alters Resistance to Quinoline Antimalarials. <i>Scientific Reports</i> , 2018, 8, 2464.  | 1.6 | 26        |
| 3198 | Two subunits of the exocyst, Sec3p and Exo70p, can function exclusively on the plasma membrane. <i>Molecular Biology of the Cell</i> , 2018, 29, 736-750.   | 0.9 | 17        |
| 3199 | Tuning CRISPR-Cas9 Gene Drives in <i>Saccharomyces cerevisiae</i> . <i>G3: Genes, Genomes, Genetics</i> , 2018, 8, 999-1018.  | 0.8 | 40        |
| 3200 | Transcription rate strongly affects splicing fidelity and cotranscriptionality in budding yeast. <i>Genome Research</i> , 2018, 28, 203-213.  | 2.4 | 88        |
| 3201 | The Na <sup>+</sup> /K <sup>+</sup> /H <sup>+</sup> exchanger Nhx1 controls multivesicular body- $\rightarrow$ vacuolar lysosome fusion. <i>Molecular Biology of the Cell</i> , 2018, 29, 317-325.  | 0.9 | 17        |
| 3202 | Alternative routes for synthesis of N-linked glycans by Alg2 mannosyltransferase. <i>FASEB Journal</i> , 2018, 32, 2492-2506.   | 0.2 | 15        |
| 3203 | Lack of the NAD <sup>+</sup> -dependent glycerol 3-phosphate dehydrogenase impairs the function of transcription factors Sip4 and Cat8 required for ethanol utilization in <i>Kluyveromyces lactis</i> . <i>Fungal Genetics and Biology</i> , 2018, 111, 16-29. | 0.9 | 9         |
| 3204 | Site-Specific Installation of Succinyl Lysine Analog into Histones Reveals the Effect of H2BK34 Succinylation on Nucleosome Dynamics. <i>Cell Chemical Biology</i> , 2018, 25, 166-174.e7.  | 2.5 | 42        |
| 3205 | Cells Escape an Operational Mitotic Checkpoint through a Stochastic Process. <i>Current Biology</i> , 2018, 28, 28-37.e7.   | 1.8 | 17        |
| 3206 | Compensatory Internalization of Pma1 in V-ATPase Mutants in <i>Saccharomyces cerevisiae</i> Requires Calcium- and Glucose-Sensitive Phosphatases. <i>Genetics</i> , 2018, 208, 655-672.   | 1.2 | 15        |
| 3207 | A functional link between NAD <sup>+</sup> homeostasis and N-terminal protein acetylation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2018, 293, 2927-2938.   | 1.6 | 18        |
| 3208 | A Role for the Respiratory Chain in Regulating Meiosis Initiation in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2018, 208, 1181-1194.  | 1.2 | 19        |
| 3209 | DNA binding by the MAT $\alpha$ 2 transcription factor controls its access to alternative ubiquitin-modification pathways. <i>Molecular Biology of the Cell</i> , 2018, 29, 542-556.  | 0.9 | 12        |
| 3210 | A role for CENP-A/Cse4 phosphorylation on serine 33 in deposition at the centromere. <i>FEMS Yeast Research</i> , 2018, 18, .   | 1.1 | 20        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3211 | Viewing pre-60S maturation at a minuteâ€™s timescale. <i>Nucleic Acids Research</i> , 2018, 46, 3140-3151.  | 6.5 | 18        |
| 3212 | Cell Size and Growth Rate Are Modulated by TORC2-Dependent Signals. <i>Current Biology</i> , 2018, 28, 196-210.e4.  | 1.8 | 44        |
| 3213 | Mps1 promotes chromosome meiotic chromosome biorientation through Dam1. <i>Molecular Biology of the Cell</i> , 2018, 29, 479-489.   | 0.9 | 15        |
| 3214 | Activation of Rab GTPase Sec4 by its GEF Sec2 is required for prospore membrane formation during sporulation in yeast <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2018, 18, .  | 1.1 | 6         |
| 3215 | Nuclear fate of yeast snoRNA is determined by co-transcriptional Rnt1 cleavage. <i>Nature Communications</i> , 2018, 9, 1783.   | 5.8 | 24        |
| 3216 | The SAGA/TREX-2 subunit Sus1 binds widely to transcribed genes and affects mRNA turnover globally. <i>Epigenetics and Chromatin</i> , 2018, 11, 13.   | 1.8 | 19        |
| 3217 | Mitochondria Export Sulfur Species Required for Cytosolic tRNA Thiolation. <i>Cell Chemical Biology</i> , 2018, 25, 738-748.e3.   | 2.5 | 28        |
| 3218 | A High-Throughput Mutational Scan of an Intrinsically Disordered Acidic Transcriptional Activation Domain. <i>Cell Systems</i> , 2018, 6, 444-455.e6.   | 2.9 | 135       |
| 3219 | CDK and MAPK Synergistically Regulate Signaling Dynamics via a Shared Multi-site Phosphorylation Region on the Scaffold Protein Ste5. <i>Molecular Cell</i> , 2018, 69, 938-952.e6.               | 4.5 | 39        |
| 3220 | Phospholipid flippases and Sfk1p, a novel regulator of phospholipid asymmetry, contribute to low permeability of the plasma membrane. <i>Molecular Biology of the Cell</i> , 2018, 29, 1203-1218. | 0.9 | 30        |
| 3221 | A Screen for Candidate Targets of Lysine Polyphosphorylation Uncovers a Conserved Network Implicated in Ribosome Biogenesis. <i>Cell Reports</i> , 2018, 22, 3427-3439.                           | 2.9 | 61        |
| 3222 | Kinesins relocate the chromosomal passenger complex to the midzone for spindle disassembly. <i>Journal of Cell Biology</i> , 2018, 217, 1687-1700.  | 2.3 | 13        |
| 3223 | Rif1 acts through Protein Phosphatase 1 but independent of replication timing to suppress telomere extension in budding yeast. <i>Nucleic Acids Research</i> , 2018, 46, 3993-4003.               | 6.5 | 24        |
| 3224 | Characterization of a yeast sporulation-specific P450 family protein, Dit2, using an in vitro assay to crosslink formyl tyrosine. <i>Journal of Biochemistry</i> , 2018, 163, 123-131.            | 0.9 | 2         |
| 3225 | Living cell synthesis of CdSe quantum dots: Manipulation based on the transformation mechanism of intracellular Se-precursors. <i>Nano Research</i> , 2018, 11, 2498-2511.                        | 5.8 | 23        |
| 3226 | A complex molecular switch directs stress-induced cyclin C nuclear release through SCF<sup>Grr1</sup>-mediated degradation of Med13. <i>Molecular Biology of the Cell</i> , 2018, 29, 363-375.    | 0.9 | 29        |
| 3227 | Regulation of lipid droplets by metabolically controlled Ldo isoforms. <i>Journal of Cell Biology</i> , 2018, 217, 127-138.   | 2.3 | 86        |
| 3228 | Identification of seipin-linked factors that act as determinants of a lipid droplet subpopulation. <i>Journal of Cell Biology</i> , 2018, 217, 269-282.   | 2.3 | 99        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3229 | The minimal ESCRT machinery of <i>Giardia lamblia</i> has altered inter-subunit interactions within the ESCRT-II and ESCRT-III complexes. <i>European Journal of Cell Biology</i> , 2018, 97, 44-62. | 1.6 | 24        |
| 3230 | The evolutionarily conserved factor Sus1/ENY2 plays a role in telomere length maintenance. <i>Current Genetics</i> , 2018, 64, 635-644.  | 0.8 | 6         |
| 3231 | Distinct features of multivesicular body-lysosome fusion revealed by a new cell-free content-mixing assay. <i>Traffic</i> , 2018, 19, 138-149.   | 1.3 | 18        |
| 3232 | Genome-Wide Mapping of Decay Factor-mRNA Interactions in Yeast Identifies Nutrient-Responsive Transcripts as Targets of the Deadenyase Ccr4. <i>G3: Genes, Genomes, Genetics</i> , 2018, 8, 315-330. | 0.8 | 27        |
| 3233 | Specialized ribosomes and specific ribosomal protein paralogs control translation of mitochondrial proteins. <i>Journal of Cell Biology</i> , 2018, 217, 117-126.                                    | 2.3 | 82        |
| 3234 | Mitochondrial inner-membrane protease Yme1 degrades outer-membrane proteins Tom22 and Om45. <i>Journal of Cell Biology</i> , 2018, 217, 139-149.   | 2.3 | 33        |
| 3235 | The Sla1 adaptor-clathrin interaction regulates coat formation and progression of endocytosis. <i>Traffic</i> , 2018, 19, 446-462.   | 1.3 | 9         |
| 3236 | Ssd1 and Gcn2 suppress global translation efficiency in replicatively aged yeast while their activation extends lifespan. <i>ELife</i> , 2018, 7, .  | 2.8 | 61        |
| 3237 | The chaperone Chs7 forms a stable complex with Chs3 and promotes its activity at the cell surface. <i>Traffic</i> , 2018, 19, 285-295.   | 1.3 | 8         |
| 3238 | Combinatorial Genetic Control of Rpd3S Through Histone H3K4 and H3K36 Methylation in Budding Yeast. <i>G3: Genes, Genomes, Genetics</i> , 2018, 8, 3411-3420.  | 0.8 | 15        |
| 3239 | Evidence of <i>Zip1</i> Promoting Sister Kinetochore Mono-orientation During Meiosis in Budding Yeast. <i>G3: Genes, Genomes, Genetics</i> , 2018, 8, 3691-3701.                                     | 0.8 | 13        |
| 3240 | Formyl-methionine as an N-degron of a eukaryotic N-end rule pathway. <i>Science</i> , 2018, 362, .   | 6.0 | 65        |
| 3241 | The Npa1p complex chaperones the assembly of the earliest eukaryotic large ribosomal subunit precursor. <i>PLoS Genetics</i> , 2018, 14, e1007597.   | 1.5 | 23        |
| 3242 | Yeast thioredoxin reductase Trr1p controls TORC1-regulated processes. <i>Scientific Reports</i> , 2018, 8, 16500.  | 1.6 | 14        |
| 3243 | RNA Recognition-like Motifs Activate a Mitogen-Activated Protein Kinase. <i>Biochemistry</i> , 2018, 57, 6878-6887.  | 1.2 | 3         |
| 3244 | Comparative Transcriptomics Highlights New Features of the Iron Starvation Response in the Human Pathogen <i>Candida glabrata</i> . <i>Frontiers in Microbiology</i> , 2018, 9, 2689.                | 1.5 | 7         |
| 3245 | Genetic interactions between specific chromosome copy number alterations dictate complex aneuploidy patterns. <i>Genes and Development</i> , 2018, 32, 1485-1498.                                    | 2.7 | 37        |
| 3246 | Identification of Elg1 interaction partners and effects on post-replication chromatin re-formation. <i>PLoS Genetics</i> , 2018, 14, e1007783.   | 1.5 | 15        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 3247 | Engineering allosteric regulation in protein kinases. <i>Science Signaling</i> , 2018, 11, .   | 1.6  | 18        |
| 3248 | Investigation of Ldb19/Art1 localization and function at the late Golgi. <i>PLoS ONE</i> , 2018, 13, e0206944.   | 1.1  | 8         |
| 3249 | Hsp90 Is Essential for Chl1-Mediated Chromosome Segregation and Sister Chromatid Cohesion. <i>MSphere</i> , 2018, 3, .   | 1.3  | 4         |
| 3250 | The intraluminal fragment pathway mediates ESCRT-independent surface transporter down-regulation. <i>Nature Communications</i> , 2018, 9, 5358.  | 5.8  | 19        |
| 3251 | A role for Gic1 and Gic2 in Cdc42 polarization at elevated temperature. <i>PLoS ONE</i> , 2018, 13, e0200863.  | 1.1  | 8         |
| 3252 | Repression of Divergent Noncoding Transcription by a Sequence-Specific Transcription Factor. <i>Molecular Cell</i> , 2018, 72, 942-954.e7.   | 4.5  | 34        |
| 3253 | RNA helicases mediate structural transitions and compositional changes in pre-ribosomal complexes. <i>Nature Communications</i> , 2018, 9, 5383.   | 5.8  | 30        |
| 3254 | A suppressor of a wtf poison-antidote meiotic driver acts via mimicry of the driver's antidote. <i>PLoS Genetics</i> , 2018, 14, e1007836.   | 1.5  | 29        |
| 3255 | Mek1 coordinates meiotic progression with DNA break repair by directly phosphorylating and inhibiting the yeast pachytene exit regulator Ndt80. <i>PLoS Genetics</i> , 2018, 14, e1007832. | 1.5  | 33        |
| 3256 | The APT complex is involved in non-coding RNA transcription and is distinct from CPF. <i>Nucleic Acids Research</i> , 2018, 46, 11528-11538.   | 6.5  | 17        |
| 3257 | Budding Yeast <i>BFA1</i> Has Multiple Positive Roles in Directing Late Mitotic Events. <i>G3: Genes, Genomes, Genetics</i> , 2018, 8, 3397-3410.  | 0.8  | 2         |
| 3258 | Suppressor mutations in <i>Rps26A</i> or <i>Rpl5</i> bypass the <i>Cgr1</i> function for pre-ribosomal 5S RNP-rotation. <i>Nature Communications</i> , 2018, 9, 4094.                      | 5.8  | 22        |
| 3259 | Protein AMPylation by an Evolutionarily Conserved Pseudokinase. <i>Cell</i> , 2018, 175, 809-821.e19.  | 13.5 | 149       |
| 3260 | Direct membrane binding and self-interaction contribute to Mmr1 function in mitochondrial inheritance. <i>Molecular Biology of the Cell</i> , 2018, 29, 2346-2357.                         | 0.9  | 9         |
| 3261 | Tel1 and Rif2 oppositely regulate telomere protection at uncapped telomeres in <i>Saccharomyces cerevisiae</i> . <i>Journal of Genetics and Genomics</i> , 2018, 45, 467-476.              | 1.7  | 2         |
| 3262 | Rab-Effector-Kinase Interplay Modulates Intraluminal Fragment Formation during Vacuole Fusion. <i>Developmental Cell</i> , 2018, 47, 80-97.e6.   | 3.1  | 7         |
| 3263 | A simple and inexpensive quantitative technique for determining chemical sensitivity in <i>Saccharomyces cerevisiae</i> . <i>Scientific Reports</i> , 2018, 8, 11919.                      | 1.6  | 25        |
| 3264 | Centromeres License the Mitotic Condensation of Yeast Chromosome Arms. <i>Cell</i> , 2018, 175, 780-795.e15.   | 13.5 | 37        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3265 | Hierarchical Organization Endows the Kinase Domain with Regulatory Plasticity. <i>Cell Systems</i> , 2018, 7, 371-383.e4.   | 2.9 | 20        |
| 3266 | Filamentation of asparagine synthetase in <i>Saccharomyces cerevisiae</i> . <i>PLoS Genetics</i> , 2018, 14, e1007737.  | 1.5 | 34        |
| 3267 | Slr2 MAPK association with chromatin is required for transcriptional activation of Rlm1 dependent genes upon cell wall stress. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2018, 1861, 1029-1039. | 0.9 | 16        |
| 3268 | Ribosome biogenesis factor Ltv1 chaperones the assembly of the small subunit head. <i>Journal of Cell Biology</i> , 2018, 217, 4141-4154.   | 2.3 | 40        |
| 3269 | Identification of factors that promote biogenesis of tRNA <sup>CGA</sup> . <i>RNA Biology</i> , 2018, 15, 1286-1294.  | 1.5 | 6         |
| 3270 | Cox15 interacts with the cytochrome bc1 dimer within respiratory supercomplexes as well as in the absence of cytochrome c oxidase. <i>Journal of Biological Chemistry</i> , 2018, 293, 16426-16439.                       | 1.6 | 8         |
| 3271 | AAA ATPase Afg1 preserves organellar fidelity and cellular healthspan by maintaining mitochondrial matrix proteostasis. <i>Journal of Cell Science</i> , 2018, 131, .   | 1.2 | 7         |
| 3272 | Single-Molecule Analysis Reveals Linked Cycles of RSC Chromatin Remodeling and Ace1p Transcription Factor Binding in Yeast. <i>Molecular Cell</i> , 2018, 72, 875-887.e9.   | 4.5 | 50        |
| 3273 | Induction of Ptp2 and Cmp2 protein phosphatases is crucial for the adaptive response to ER stress in <i>Saccharomyces cerevisiae</i> . <i>Scientific Reports</i> , 2018, 8, 13078.  | 1.6 | 8         |
| 3274 | Heat-stress triggers MAPK crosstalk to turn on the hyperosmotic response pathway. <i>Scientific Reports</i> , 2018, 8, 15168.   | 1.6 | 46        |
| 3275 | A ZIP1 separation-of-function allele reveals that centromere pairing drives meiotic segregation of achiasmate chromosomes in budding yeast. <i>PLoS Genetics</i> , 2018, 14, e1007513.                                    | 1.5 | 8         |
| 3276 | Gene drive inhibition by the anti-CRISPR proteins AcrIIA2 and AcrIIA4 in <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> , 2018, 164, 464-474.   | 0.7 | 85        |
| 3277 | Rad6-Bre1 mediated histone H2Bub1 protects uncapped telomeres from exonuclease Exo1 in <i>Saccharomyces cerevisiae</i> . <i>DNA Repair</i> , 2018, 72, 64-76.   | 1.3 | 6         |
| 3278 | Pulsatile inputs achieve tunable attenuation of gene expression variability and graded multi-gene regulation. <i>Nature Communications</i> , 2018, 9, 3521.   | 5.8 | 98        |
| 3279 | Cargo induces retromer-mediated membrane remodeling on membranes. <i>Molecular Biology of the Cell</i> , 2018, 29, 2709-2719.   | 0.9 | 19        |
| 3280 | Key phosphorylation events in Spc29 and Spc42 guide multiple steps of yeast centrosome duplication. <i>Molecular Biology of the Cell</i> , 2018, 29, 2280-2291.   | 0.9 | 3         |
| 3281 | Phosphorylation and Proteasome Recognition of the mRNA-Binding Protein Cth2 Facilitates Yeast Adaptation to Iron Deficiency. <i>MBio</i> , 2018, 9, .   | 1.8 | 11        |
| 3282 | Molecular basis for disassembly of an importin:ribosomal protein complex by the escortin Tsr2. <i>Nature Communications</i> , 2018, 9, 3669.  | 5.8 | 17        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3283 | Lso2 is a conserved ribosome-bound protein required for translational recovery in yeast. <i>PLoS Biology</i> , 2018, 16, e2005903.   | 2.6 | 31        |
| 3284 | The half-bridge component Kar1 promotes centrosome separation and duplication during budding yeast meiosis. <i>Molecular Biology of the Cell</i> , 2018, 29, 1798-1810.  | 0.9 | 4         |
| 3285 | Tsc3 regulates SPT amino acid choice in <i>Saccharomyces cerevisiae</i> by promoting alanine in the sphingolipid pathway. <i>Journal of Lipid Research</i> , 2018, 59, 2126-2139.  | 2.0 | 11        |
| 3286 | Feedback regulation of TORC1 by its downstream effectors Npr1 and Par32. <i>Molecular Biology of the Cell</i> , 2018, 29, 2751-2765.   | 0.9 | 13        |
| 3287 | Translational Reprogramming Provides a Blueprint for Cellular Adaptation. <i>Cell Chemical Biology</i> , 2018, 25, 1372-1379.e3.   | 2.5 | 14        |
| 3288 | Integration of Multiple Metabolic Signals Determines Cell Fate Prior to Commitment. <i>Molecular Cell</i> , 2018, 71, 733-744.e11.   | 4.5 | 41        |
| 3289 | Suppression of respiratory growth defect of mutant deficient in mitochondrial phospholipase A1 by overexpression of genes involved in coenzyme Q synthesis in <i>Saccharomyces cerevisiae</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 2018, 82, 1633-1639. | 0.6 | 1         |
| 3290 | Marker-free insertion of a series of C-terminal epitopes based on the 50:50 method in <i>Saccharomyces cerevisiae</i> . <i>Journal of General and Applied Microbiology</i> , 2018, 64, 99-102.   | 0.4 | 1         |
| 3291 | Interrogation of ethnomedicinal plants for synthetic lethality effects in combination with deficiency in the DNA repair endonuclease RAD1 using a yeast cell-based assay. <i>Journal of Ethnopharmacology</i> , 2018, 223, 10-21.  | 2.0 | 1         |
| 3292 | Select $\beta$ -arrestins control cell-surface abundance of the mammalian Kir2.1 potassium channel in a yeast model. <i>Journal of Biological Chemistry</i> , 2018, 293, 11006-11021.  | 1.6 | 17        |
| 3293 | Ribosomal protein L14 contributes to the early assembly of 60S ribosomal subunits in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2018, 46, 4715-4732.  | 6.5 | 12        |
| 3294 | Atg9 establishes Atg2-dependent contact sites between the endoplasmic reticulum and phagophores. <i>Journal of Cell Biology</i> , 2018, 217, 2743-2763.  | 2.3 | 194       |
| 3295 | Biofilm formation and toxin production provide a fitness advantage in mixed colonies of environmental yeast isolates. <i>Ecology and Evolution</i> , 2018, 8, 5541-5550.   | 0.8 | 22        |
| 3296 | Methods for Controlled Protein Depletion to Study Protein Function during Meiosis. <i>Methods in Enzymology</i> , 2018, 601, 331-357.  | 0.4 | 16        |
| 3297 | Multiple kinases inhibit origin licensing and helicase activation to ensure reductive cell division during meiosis. <i>ELife</i> , 2018, 7, .  | 2.8 | 22        |
| 3298 | Intracellular mechanism by which arsenite activates the yeast stress MAPK Hog1. <i>Molecular Biology of the Cell</i> , 2018, 29, 1904-1915.  | 0.9 | 28        |
| 3299 | New ubiquitin-dependent mechanisms regulating the Aurora B-Protein Phosphatase 1 balance in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 2018, 131, .  | 1.2 | 2         |
| 3300 | Combined Enrichment/Enzymatic Approach To Study Tightly Clustered Multisite Phosphorylation on Ser-Rich Domains. <i>Journal of Proteome Research</i> , 2018, 17, 3050-3060.  | 1.8 | 3         |



| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3301 | Endoplasmic reticulum-plasma membrane contact sites integrate sterol and phospholipid regulation. <i>PLoS Biology</i> , 2018, 16, e2003864.   | 2.6 | 132       |
| 3302 | Fungal Light-Oxygen-Voltage Domains for Optogenetic Control of Gene Expression and Flocculation in Yeast. <i>MBio</i> , 2018, 9, .  | 1.8 | 34        |
| 3303 | A newly characterized vacuolar serine carboxypeptidase, Atg42/Ybr139w, is required for normal vacuole function and the terminal steps of autophagy in the yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2018, 29, 1089-1099. | 0.9 | 60        |
| 3304 | Using the Flp Recombinase to Induce Site-Specific Protein-DNA Nicks. <i>Methods in Enzymology</i> , 2018, 601, 1-25.  | 0.4 | 1         |
| 3305 | Live-cell imaging of early coat protein dynamics during clathrin-mediated endocytosis. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2018, 1865, 1566-1578.  | 1.9 | 1         |
| 3306 | Seipin and the membrane-shaping protein Pex30 cooperate in organelle budding from the endoplasmic reticulum. <i>Nature Communications</i> , 2018, 9, 2939.  | 5.8 | 107       |
| 3307 | Toxicity of dihydroxyacetone is exerted through the formation of methylglyoxal in <i>Saccharomyces cerevisiae</i> : effects on actin polarity and nuclear division. <i>Biochemical Journal</i> , 2018, 475, 2637-2652.  | 1.7 | 6         |
| 3308 | Quantitative imaging of chromatin decompaction in living cells. <i>Molecular Biology of the Cell</i> , 2018, 29, 1763-1777.   | 0.9 | 20        |
| 3309 | PI(3,5)P <sub>2</sub> controls vacuole potassium transport to support cellular osmoregulation. <i>Molecular Biology of the Cell</i> , 2018, 29, 1718-1731.  | 0.9 | 19        |
| 3310 | Deregulation of the G1/S-phase transition is the proximal cause of mortality in old yeast mother cells. <i>Genes and Development</i> , 2018, 32, 1075-1084.   | 2.7 | 46        |
| 3311 | Integrated control of formin-mediated actin assembly by a stationary inhibitor and a mobile activator. <i>Journal of Cell Biology</i> , 2018, 217, 3512-3530.   | 2.3 | 20        |
| 3312 | An isolated CLASP TOG domain suppresses microtubule catastrophe and promotes rescue. <i>Molecular Biology of the Cell</i> , 2018, 29, 1359-1375.  | 0.9 | 36        |
| 3313 | Phosphatidylserine and GTPase activation control Cdc42 nanoclustering to counter dissipative diffusion. <i>Molecular Biology of the Cell</i> , 2018, 29, 1299-1310.   | 0.9 | 40        |
| 3314 | Inositol polyphosphates regulate and predict yeast pseudohyphal growth phenotypes. <i>PLoS Genetics</i> , 2018, 14, e1007493.   | 1.5 | 15        |
| 3315 | Genome-wide C-SWAT library for high-throughput yeast genome tagging. <i>Nature Methods</i> , 2018, 15, 598-600.   | 9.0 | 57        |
| 3316 | Global Proteome Remodeling during ER Stress Involves Hac1-Driven Expression of Long Undecoded Transcript Isoforms. <i>Developmental Cell</i> , 2018, 46, 219-235.e8.  | 3.1 | 45        |
| 3317 | Competitive organelle-specific adaptors recruit Vps13 to membrane contact sites. <i>Journal of Cell Biology</i> , 2018, 217, 3593-3607.   | 2.3 | 122       |
| 3318 | A Genome-Wide Screen Reveals a Role for the HIR Histone Chaperone Complex in Preventing Mislocalization of Budding Yeast CENP-A. <i>Genetics</i> , 2018, 210, 203-218.  | 1.2 | 20        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3319 | Modulation of TORC2 Signaling by a Conserved Lkb1 Signaling Axis in Budding Yeast. <i>Genetics</i> , 2018, 210, 155-170.  | 1.2 | 17        |
| 3320 | Phosphorylation-Mediated Clearance of Amyloid-like Assemblies in Meiosis. <i>Developmental Cell</i> , 2018, 45, 392-405.e6.   | 3.1 | 66        |
| 3321 | PCYT1A Regulates Phosphatidylcholine Homeostasis from the Inner Nuclear Membrane in Response to Membrane Stored Curvature Elastic Stress. <i>Developmental Cell</i> , 2018, 45, 481-495.e8.                   | 3.1 | 99        |
| 3322 | Control of mitotic chromosome condensation by the fission yeast transcription factor Zasl. <i>Journal of Cell Biology</i> , 2018, 217, 2383-2401.   | 2.3 | 3         |
| 3323 | TOR Facilitates the Targeting of the 19S Proteasome Subcomplex To Enhance Transcription Complex Assembly at the Promoters of the Ribosomal Protein Genes. <i>Molecular and Cellular Biology</i> , 2018, 38, . | 1.1 | 9         |
| 3324 | Prolonged cyclin-dependent kinase inhibition results in septin perturbations during return to growth and mitosis. <i>Journal of Cell Biology</i> , 2018, 217, 2429-2443.                                      | 2.3 | 7         |
| 3325 | <scp>SNARE</scp> -mediated membrane fusion arrests at pore expansion to regulate the volume of an organelle. <i>EMBO Journal</i> , 2018, 37, .  | 3.5 | 39        |
| 3326 | Genome-wide screening of budding yeast with honokiol to associate mitochondrial function with lipid metabolism. <i>Traffic</i> , 2018, 19, 867-878.   | 1.3 | 8         |
| 3327 | Lipid trafficking by yeast Snx4 family SNX-BAR proteins promotes autophagy and vacuole membrane fusion. <i>Molecular Biology of the Cell</i> , 2018, 29, 2190-2200.   | 0.9 | 43        |
| 3328 | Set4 is a chromatin-associated protein, promotes survival during oxidative stress, and regulates stress response genes in yeast. <i>Journal of Biological Chemistry</i> , 2018, 293, 14429-14443.             | 1.6 | 23        |
| 3329 | Reconstitution reveals Ykt6 as the autophagosomal SNARE in autophagosome-vacuole fusion. <i>Journal of Cell Biology</i> , 2018, 217, 3656-3669.   | 2.3 | 88        |
| 3330 | Ivy1 is a negative regulator of Gtr-dependent TORC1 activation. <i>Journal of Cell Science</i> , 2018, 131, .   | 1.2 | 10        |
| 3331 | The Small Yeast GTPase Rho5 and Its Dimeric GEF Dck1/Lmo1 Respond to Glucose Starvation. <i>International Journal of Molecular Sciences</i> , 2018, 19, 2186.   | 1.8 | 16        |
| 3332 | Tma64/eIF2D, Tma20/MCT-1, and Tma22/DENR Recycle Post-termination 40S Subunits In Vivo. <i>Molecular Cell</i> , 2018, 71, 761-774.e5.   | 4.5 | 62        |
| 3333 | Lipid moiety of glycosylphosphatidylinositol-anchored proteins contributes to the determination of their final destination in yeast. <i>Genes To Cells</i> , 2018, 23, 880-892.                               | 0.5 | 19        |
| 3334 | Negative regulation of <i>Candida glabrata</i> Pdr1 by the deubiquitinase subunit Bre5 occurs in a ubiquitin independent manner. <i>Molecular Microbiology</i> , 2018, 110, 309-323.                          | 1.2 | 9         |
| 3335 | Investigating the long-term stability of protein immunogen(s) for whole recombinant yeast-based vaccines. <i>FEMS Yeast Research</i> , 2018, 18, .  | 1.1 | 17        |
| 3336 | Functional Analysis of the Yeast LINC Complex Using Fluctuation Spectroscopy and Super-Resolution Imaging. <i>Methods in Molecular Biology</i> , 2018, 1840, 137-161.   | 0.4 | 10        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3337 | Error-free DNA damage tolerance pathway is facilitated by the Irc5 translocase through cohesin. <i>EMBO Journal</i> , 2018, 37, .  | 3.5 | 14        |
| 3338 | Structures of the fungal dynamin-related protein Vps1 reveal a unique, open helical architecture. <i>Journal of Cell Biology</i> , 2018, 217, 3608-3624.   | 2.3 | 12        |
| 3339 | Eisosomes are metabolically regulated storage compartments for APC-type nutrient transporters. <i>Molecular Biology of the Cell</i> , 2018, 29, 2113-2127.   | 0.9 | 40        |
| 3340 | Functional Impact of the H2A.Z Histone Variant During Meiosis in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2018, 209, 997-1015.  | 1.2 | 19        |
| 3341 | Evidence for interaction between Hsp90 and the ER membrane complex. <i>Cell Stress and Chaperones</i> , 2018, 23, 1101-1115.   | 1.2 | 11        |
| 3342 | Target of rapamycin complex 2-dependent phosphorylation of the coat protein Pan1 by Akl1 controls endocytosis dynamics in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2018, 293, 12043-12053. | 1.6 | 23        |
| 3343 | The combination of NAD <sup>+</sup> -dependent deacetylase gene deletion and the interruption of gluconeogenesis causes increased glucose metabolism in budding yeast. <i>PLoS ONE</i> , 2018, 13, e0194942.                 | 1.1 | 11        |
| 3344 | PiggyBac-based screening identified BEM4 as a suppressor to rescue growth defects in och1-disrupted yeast cells. <i>Bioscience, Biotechnology and Biochemistry</i> , 2018, 82, 1497-1507.                                    | 0.6 | 2         |
| 3345 | P-Body Localization of the Hrr25/Casein Kinase 1 Protein Kinase Is Required for the Completion of Meiosis. <i>Molecular and Cellular Biology</i> , 2018, 38, .   | 1.1 | 10        |
| 3346 | Cdk1 phosphorylation of Esp1/Separase functions with PP2A and Slk19 to regulate pericentric Cohesin and anaphase onset. <i>PLoS Genetics</i> , 2018, 14, e1007029.   | 1.5 | 15        |
| 3347 | DNA strand-exchange patterns associated with double-strand break-induced and spontaneous mitotic crossovers in <i>Saccharomyces cerevisiae</i> . <i>PLoS Genetics</i> , 2018, 14, e1007302.                                  | 1.5 | 14        |
| 3348 | Histone depletion prevents telomere fusions in pre-senescent cells. <i>PLoS Genetics</i> , 2018, 14, e1007407.   | 1.5 | 6         |
| 3349 | Casein kinase 2 mediated phosphorylation of Spt6 modulates histone dynamics and regulates spurious transcription. <i>Nucleic Acids Research</i> , 2018, 46, 7612-7630.   | 6.5 | 22        |
| 3350 | Integrated TORC1 and PKA signaling control the temporal activation of glucose-induced gene expression in yeast. <i>Nature Communications</i> , 2019, 10, 3558.   | 5.8 | 49        |
| 3351 | The prion-like protein kinase Sky1 is required for efficient stress granule disassembly. <i>Nature Communications</i> , 2019, 10, 3614.  | 5.8 | 36        |
| 3352 | Systematic analysis of F-box proteins reveals a new branch of the yeast mating pathway. <i>Journal of Biological Chemistry</i> , 2019, 294, 14717-14731.   | 1.6 | 8         |
| 3353 | Mutations in topoisomerase III <sup>2</sup> result in a B cell immunodeficiency. <i>Nature Communications</i> , 2019, 10, 3644.  | 5.8 | 37        |
| 3354 | Compartmentalized Synthesis of Triacylglycerol at the Inner Nuclear Membrane Regulates Nuclear Organization. <i>Developmental Cell</i> , 2019, 50, 755-766.e6.   | 3.1 | 52        |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 3355 | Thermophile 90S Pre-ribosome Structures Reveal the Reverse Order of Co-transcriptional 18S rRNA Subdomain Integration. <i>Molecular Cell</i> , 2019, 75, 1256-1269.e7.  | 4.5  | 48        |
| 3356 | Asymmetric inheritance of spindle microtubule-organizing centres preserves replicative lifespan. <i>Nature Cell Biology</i> , 2019, 21, 952-965.  | 4.6  | 25        |
| 3357 | Construction of functional chimeras of syntaxin-1A and its yeast orthologue, and their application to the yeast cell-based assay for botulinum neurotoxin serotype C. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2019, 1863, 129396. | 1.1  | 2         |
| 3358 | Functional interplay between DEAD-box RNA helicases Ded1 and Dbp1 in preinitiation complex attachment and scanning on structured mRNAs in vivo. <i>Nucleic Acids Research</i> , 2019, 47, 8785-8806.  | 6.5  | 32        |
| 3359 | Free glycans derived from O-mannosylated glycoproteins suggest the presence of an O-glycoprotein degradation pathway in yeast. <i>Journal of Biological Chemistry</i> , 2019, 294, 15900-15911.   | 1.6  | 4         |
| 3360 | Structure and assembly of the mitochondrial membrane remodelling GTPase Mgm1. <i>Nature</i> , 2019, 571, 429-433.   | 13.7 | 86        |
| 3361 | Crosstalk between chromatin structure, cohesin activity and transcription. <i>Epigenetics and Chromatin</i> , 2019, 12, 47.   | 1.8  | 17        |
| 3362 | Mec1ATR Autophosphorylation and Ddc2ATRIP Phosphorylation Regulates DNA Damage Checkpoint Signaling. <i>Cell Reports</i> , 2019, 28, 1090-1102.e3.  | 2.9  | 19        |
| 3363 | Mitochondrial cysteinyl-tRNA synthetase is expressed via alternative transcriptional initiation regulated by energy metabolism in yeast cells. <i>Journal of Biological Chemistry</i> , 2019, 294, 13781-13788.                                     | 1.6  | 16        |
| 3364 | Single-molecule localization microscopy and tracking with red-shifted states of conventional BODIPY conjugates in living cells. <i>Nature Communications</i> , 2019, 10, 3400.  | 5.8  | 48        |
| 3365 | Multisite phosphorylation code of CDK. <i>Nature Structural and Molecular Biology</i> , 2019, 26, 649-658.  | 3.6  | 64        |
| 3366 | The Bub1-TPR Domain Interacts Directly with Mad3 to Generate Robust Spindle Checkpoint Arrest. <i>Current Biology</i> , 2019, 29, 2407-2414.e7.   | 1.8  | 19        |
| 3367 | Comprehensive and quantitative analysis of G1 cyclins. A tool for studying the cell cycle. <i>PLoS ONE</i> , 2019, 14, e0218531.  | 1.1  | 4         |
| 3368 | A COPII subunit acts with an autophagy receptor to target endoplasmic reticulum for degradation. <i>Science</i> , 2019, 365, 53-60.   | 6.0  | 114       |
| 3369 | The Role of Transcription Factors and Nuclear Pore Proteins in Controlling the Spatial Organization of the Yeast Genome. <i>Developmental Cell</i> , 2019, 49, 936-947.e4.  | 3.1  | 44        |
| 3370 | Ratiometric GPCR signaling enables directional sensing in yeast. <i>PLoS Biology</i> , 2019, 17, e3000484.  | 2.6  | 27        |
| 3371 | The Hsp90 Molecular Chaperone Regulates the Transcription Factor Network Controlling Chromatin Accessibility. <i>Journal of Molecular Biology</i> , 2019, 431, 4993-5003.   | 2.0  | 8         |
| 3372 | Seipin negatively regulates sphingolipid production at the ER-LD contact site. <i>Journal of Cell Biology</i> , 2019, 218, 3663-3680.   | 2.3  | 19        |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 3373 | Cell Cycle Inhibitor Whi5 Records Environmental Information to Coordinate Growth and Division in Yeast. <i>Cell Reports</i> , 2019, 29, 987-994.e5.  | 2.9  | 38        |
| 3374 | Activity of a ubiquitin ligase adaptor is regulated by disordered insertions in its arrestin domain. <i>Molecular Biology of the Cell</i> , 2019, 30, 3057-3072.   | 0.9  | 15        |
| 3375 | Reciprocal interactions between mtDNA and lifespan control in budding yeast. <i>Molecular Biology of the Cell</i> , 2019, 30, 2943-2952.   | 0.9  | 11        |
| 3376 | Ubiquitin receptors are required for substrate-mediated activation of the proteasome's unfolding ability. <i>Scientific Reports</i> , 2019, 9, 14506.  | 1.6  | 20        |
| 3377 | The Ulp2 <sc>SUMO</sc> protease promotes transcription elongation through regulation of histone sumoylation. <i>EMBO Journal</i> , 2019, 38, e102003.  | 3.5  | 28        |
| 3378 | Auxin-Inducible Degron System for Depletion of Proteins in <i>Saccharomyces cerevisiae</i> . <i>Current Protocols in Molecular Biology</i> , 2019, 128, e104.  | 2.9  | 18        |
| 3380 | Gcn5 and Esa1 function as histone crotonyltransferases to regulate crotonylation-dependent transcription. <i>Journal of Biological Chemistry</i> , 2019, 294, 20122-20134.   | 1.6  | 66        |
| 3381 | Kinetochore-associated Stu2 promotes chromosome biorientation in vivo. <i>PLoS Genetics</i> , 2019, 15, e1008423.  | 1.5  | 26        |
| 3382 | PCNA Deubiquitylases Control DNA Damage Bypass at Replication Forks. <i>Cell Reports</i> , 2019, 29, 1323-1335.e5.   | 2.9  | 14        |
| 3383 | Psp2, a novel regulator of autophagy that promotes autophagy-related protein translation. <i>Cell Research</i> , 2019, 29, 994-1008.   | 5.7  | 23        |
| 3384 | Analysis of Functional Domains in Rho5, the Yeast Homolog of Human Rac1 GTPase, in Oxidative Stress Response. <i>International Journal of Molecular Sciences</i> , 2019, 20, 5550.   | 1.8  | 8         |
| 3385 | Effects of Sterols on the Interaction of SDS, Benzalkonium Chloride, and A Novel Compound, Kor105, with Membranes. <i>Biomolecules</i> , 2019, 9, 627.   | 1.8  | 10        |
| 3386 | The E3 ubiquitin ligase Pib1 regulates effective gluconeogenic shutdown upon glucose availability. <i>Journal of Biological Chemistry</i> , 2019, 294, 17209-17223.  | 1.6  | 5         |
| 3387 | Mechanical stiffness of reconstituted actin patches correlates tightly with endocytosis efficiency. <i>PLoS Biology</i> , 2019, 17, e3000500.  | 2.6  | 19        |
| 3388 | Live-Cell Imaging of mRNP-NPC Interactions in Budding Yeast. <i>Methods in Molecular Biology</i> , 2019, 2038, 131-150.  | 0.4  | 3         |
| 3389 | Phosphorylation of the lipid droplet localized glycerol-3-phosphate acyltransferase Gpt2 prevents a futile triacylglycerol cycle in yeast. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2019, 1864, 158509. | 1.2  | 17        |
| 3390 | DEAD-box ATPases are global regulators of phase-separated organelles. <i>Nature</i> , 2019, 573, 144-148.  | 13.7 | 274       |
| 3391 | Kinetic assay of starvation sensitivity in yeast autophagy mutants allows for the identification of intermediary phenotypes. <i>BMC Research Notes</i> , 2019, 12, 505.  | 0.6  | 1         |

| #    | ARTICLE  | IF   | CITATIONS |
|------|--|------|-----------|
| 3392 | Fine-tuning the expression of target genes using a DDI2 promoter gene switch in budding yeast. <i>Scientific Reports</i> , 2019, 9, 12538.   | 1.6  | 9         |
| 3393 | A Bacterial Effector Mimics a Host HSP90 Client to Undermine Immunity. <i>Cell</i> , 2019, 179, 205-218.e21.   | 13.5 | 53        |
| 3394 | Regulation of Skn7-dependent, oxidative stress-induced genes by the RNA polymerase II-CTD phosphatase, Fcp1, and Mediator kinase subunit, Cdk8, in yeast. <i>Journal of Biological Chemistry</i> , 2019, 294, 16080-16094. | 1.6  | 9         |
| 3395 | Proteomics-Based Monitoring of Pathway Activity Reveals that Blocking Diacylglycerol Biosynthesis Rescues from Alpha-Synuclein Toxicity. <i>Cell Systems</i> , 2019, 9, 309-320.e8.  | 2.9  | 12        |
| 3396 | Blocking late stages of splicing quickly limits pre-spliceosome assembly in vivo. <i>RNA Biology</i> , 2019, 16, 1775-1784.  | 1.5  | 4         |
| 3397 | Phosphorylation of mitochondrial matrix proteins regulates their selective mitophagic degradation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 20517-20527.        | 3.3  | 26        |
| 3398 | Alternative Experimental Models for Studying Influenza Proteins, Host-Virus Interactions and Anti-Influenza Drugs. <i>Pharmaceuticals</i> , 2019, 12, 147.   | 1.7  | 13        |
| 3399 | Glutarylation of Histone H4 Lysine 91 Regulates Chromatin Dynamics. <i>Molecular Cell</i> , 2019, 76, 660-675.e9.  | 4.5  | 112       |
| 3400 | PP2C phosphatases promote autophagy by dephosphorylation of the Atg1 complex. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 1613-1620.                               | 3.3  | 48        |
| 3401 | Identification of Suppressor of Clathrin Deficiency-1 ( <i>SCD1</i> ) and Its Connection to Clathrin-Mediated Endocytosis in <i>Saccharomyces cerevisiae</i> . <i>G3: Genes, Genomes, Genetics</i> , 2019, 9, 867-877.     | 0.8  | 7         |
| 3402 | Adaptor protein complex-1 (AP-1) is recruited by the HEATR5 protein Laa1 and its co-factor Laa2 in yeast. <i>Journal of Biological Chemistry</i> , 2019, 294, 1410-1419.   | 1.6  | 13        |
| 3403 | Expression of TorsinA in a heterologous yeast system reveals interactions with luminal domains of LINC and nuclear pore complex components. <i>Molecular Biology of the Cell</i> , 2019, 30, 530-541.                      | 0.9  | 14        |
| 3404 | Gyp1 has a dual function as Ypt1 GAP and interaction partner of Atg8 in selective autophagy. <i>Autophagy</i> , 2019, 15, 1031-1050.   | 4.3  | 8         |
| 3405 | Coenzyme Q biosynthetic proteins assemble in a substrate-dependent manner into domains at ER-mitochondria contacts. <i>Journal of Cell Biology</i> , 2019, 218, 1353-1369.   | 2.3  | 69        |
| 3406 | The cell-cycle transcriptional network generates and transmits a pulse of transcription once each cell cycle. <i>Cell Cycle</i> , 2019, 18, 363-378.   | 1.3  | 15        |
| 3407 | <i>Saccharomyces cerevisiae</i> Cytosolic Thioredoxins Control Glycolysis, Lipid Metabolism, and Protein Biosynthesis under Wine-Making Conditions. <i>Applied and Environmental Microbiology</i> , 2019, 85, .            | 1.4  | 9         |
| 3408 | Methods for studying the regulation of membrane traffic by ubiquitin and the ESCRT pathway. <i>Methods in Enzymology</i> , 2019, 619, 269-291.   | 0.4  | 1         |
| 3409 | Type I myosins anchor actin assembly to the plasma membrane during clathrin-mediated endocytosis. <i>Journal of Cell Biology</i> , 2019, 218, 1138-1147.   | 2.3  | 31        |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 3410 | Tsr4 Is a Cytoplasmic Chaperone for the Ribosomal Protein Rps2 in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2019, 39, .   | 1.1  | 19        |
| 3411 | The nucleosome acidic patch directly interacts with subunits of the Paf1 and FACT complexes and controls chromatin architecture in vivo. <i>Nucleic Acids Research</i> , 2019, 47, 8410-8423.                             | 6.5  | 27        |
| 3412 | H3K36 Methylation and the Chromodomain Protein Eaf3 Are Required for Proper Cotranscriptional Spliceosome Assembly. <i>Cell Reports</i> , 2019, 27, 3760-3769.e4.   | 2.9  | 37        |
| 3413 | Cyclin-dependent kinase-mediated phosphorylation of the exocyst subunit Exo84 in late G1 phase suppresses exocytic secretion and cell growth in yeast. <i>Journal of Biological Chemistry</i> , 2019, 294, 11323-11332.   | 1.6  | 5         |
| 3414 | Regulation of the anaphase promoting complex/cyclosome by the degradation of its unassembled catalytic subunit, Apc11. <i>FASEB Journal</i> , 2019, 33, 9752-9761.  | 0.2  | 5         |
| 3415 | Conserved juxtamembrane domains in the yeast golgin Coy1 drive assembly of a megadalton-sized complex and mediate binding to tethering and SNARE proteins. <i>Journal of Biological Chemistry</i> , 2019, 294, 9690-9705. | 1.6  | 2         |
| 3416 | Rsp5 and Mdm30 reshape the mitochondrial network in response to age-induced vacuole stress. <i>Molecular Biology of the Cell</i> , 2019, 30, 2141-2154.   | 0.9  | 15        |
| 3417 | <i>Pichia pastoris</i> protease-deficient and auxotrophic strains generated by a novel, user-friendly vector toolbox for gene deletion. <i>Yeast</i> , 2019, 36, 557-570.   | 0.8  | 17        |
| 3418 | Reversible DNA damage checkpoint activation at the presenescent stage in telomerase-deficient cells of <i>Saccharomyces cerevisiae</i> . <i>Genes To Cells</i> , 2019, 24, 546-558.                                       | 0.5  | 1         |
| 3419 | The identification of small molecule inhibitors of the plant inositol phosphorylceramide synthase which demonstrate herbicidal activity. <i>Scientific Reports</i> , 2019, 9, 8083.                                       | 1.6  | 7         |
| 3420 | Ssd1 and the cell wall integrity pathway promote entry, maintenance, and recovery from quiescence in budding yeast. <i>Molecular Biology of the Cell</i> , 2019, 30, 2205-2217.   | 0.9  | 20        |
| 3421 | SAGA DUBm-mediated surveillance regulates prompt export of stress-inducible transcripts for proteostasis. <i>Nature Communications</i> , 2019, 10, 2458.  | 5.8  | 11        |
| 3422 | Evidence that Ergosterol Biosynthesis Modulates Activity of the Pdr1 Transcription Factor in <i>Candida glabrata</i> . <i>MBio</i> , 2019, 10, .  | 1.8  | 42        |
| 3423 | Inhibiting eukaryotic ribosome biogenesis. <i>BMC Biology</i> , 2019, 17, 46.   | 1.7  | 39        |
| 3424 | Multiple Negative Regulators Restrict Recruitment of the SWI/SNF Chromatin Remodeler to the <i>HO</i> Promoter in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2019, 212, 1181-1204.                               | 1.2  | 9         |
| 3425 | Rif1 S-acylation mediates DNA double-strand break repair at the inner nuclear membrane. <i>Nature Communications</i> , 2019, 10, 2535.  | 5.8  | 34        |
| 3426 | Structure and function of Vms1 and Arb1 in RQC and mitochondrial proteome homeostasis. <i>Nature</i> , 2019, 570, 538-542.  | 13.7 | 63        |
| 3427 | Condensin-Mediated Chromosome Folding and Internal Telomeres Drive Dicentric Severing by Cytokinesis. <i>Molecular Cell</i> , 2019, 75, 131-144.e3.   | 4.5  | 24        |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 3428 | Protein aggregation mediates stoichiometry of protein complexes in aneuploid cells. <i>Genes and Development</i> , 2019, 33, 1031-1047.   | 2.7  | 83        |
| 3429 | Proper insertion and topogenesis of membrane proteins in the ER depend on Sec63. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2019, 1863, 1371-1380.   | 1.1  | 11        |
| 3430 | Mitochondrial protein translocation-associated degradation. <i>Nature</i> , 2019, 569, 679-683.   | 13.7 | 203       |
| 3431 | Effective mismatch repair depends on timely control of PCNA retention on DNA by the Elg1 complex. <i>Nucleic Acids Research</i> , 2019, 47, 6826-6841.  | 6.5  | 20        |
| 3432 | Proximity-dependent biotinylation by TurboID to identify protein-protein interaction networks in yeast. <i>Journal of Cell Science</i> , 2019, 132, .   | 1.2  | 51        |
| 3433 | Tsr4 and Nap1, two novel members of the ribosomal protein chaperOME. <i>Nucleic Acids Research</i> , 2019, 47, 6984-7002.   | 6.5  | 28        |
| 3434 | Retrograde trafficking and quality control of yeast synaptobrevin, Snc1, are conferred by its transmembrane domain. <i>Molecular Biology of the Cell</i> , 2019, 30, 1729-1742.                             | 0.9  | 20        |
| 3435 | Cyclin-Specific Docking Mechanisms Reveal the Complexity of M-CDK Function in the Cell Cycle. <i>Molecular Cell</i> , 2019, 75, 76-89.e3.   | 4.5  | 50        |
| 3436 | A Novel G-Quadruplex Binding Protein in Yeastâ€”Slx9. <i>Molecules</i> , 2019, 24, 1774.  | 1.7  | 16        |
| 3437 | Chromosome detachment from the nuclear envelope is required for genomic stability in closed mitosis. <i>Molecular Biology of the Cell</i> , 2019, 30, 1578-1586.  | 0.9  | 2         |
| 3438 | Pop2 phosphorylation at S39 contributes to the glucose repression of stress response genes, HSP12 and HSP26. <i>PLoS ONE</i> , 2019, 14, e0215064.  | 1.1  | 5         |
| 3439 | Hsp104 facilitates the endoplasmicâ€”reticulumâ€”associated degradation of diseaseâ€”associated and aggregationâ€”prone substrates. <i>Protein Science</i> , 2019, 28, 1290-1306.                           | 3.1  | 16        |
| 3440 | Mitochondria export ironâ€”sulfur and sulfur intermediates to the cytoplasm for ironâ€”sulfur cluster assembly and tRNA thiolation in yeast. <i>Journal of Biological Chemistry</i> , 2019, 294, 9489-9502. | 1.6  | 54        |
| 3441 | Yeast centrosome components form a noncanonical LINC complex at the nuclear envelope insertion site. <i>Journal of Cell Biology</i> , 2019, 218, 1478-1490.   | 2.3  | 33        |
| 3442 | AP-2-Dependent Endocytic Recycling of the Chitin Synthase Chs3 Regulates Polarized Growth in <i>Candida albicans</i> . <i>MBio</i> , 2019, 10, .  | 1.8  | 19        |
| 3443 | Ribosomal mutation in helix 32 of 18S rRNA alters fidelity of eukaryotic translation start site selection. <i>FEBS Letters</i> , 2019, 593, 852-867.  | 1.3  | 3         |
| 3444 | Clathrin Adaptor Complex-interacting Protein Irc6 Functions through the Conserved C-Terminal Domain. <i>Scientific Reports</i> , 2019, 9, 4436.   | 1.6  | 4         |
| 3445 | Glycolytic flux in <i>Saccharomyces cerevisiae</i> is dependent on RNA polymerase III and its negative regulator Maf1. <i>Biochemical Journal</i> , 2019, 476, 1053-1082.                                   | 1.7  | 9         |



| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 3446 | Sequential Ubiquitination of Ribosomal Protein uS3 Triggers the Degradation of Non-functional 18S rRNA. <i>Cell Reports</i> , 2019, 26, 3400-3415.e7.   | 2.9  | 67        |
| 3447 | Transcription-dependent spreading of the Dal80 yeast GATA factor across the body of highly expressed genes. <i>PLoS Genetics</i> , 2019, 15, e1007999.  | 1.5  | 8         |
| 3448 | Live cell monitoring of double strand breaks in <i>S. cerevisiae</i> . <i>PLoS Genetics</i> , 2019, 15, e1008001.   | 1.5  | 28        |
| 3449 | Promethin Is a Conserved Seipin Partner Protein. <i>Cells</i> , 2019, 8, 268.   | 1.8  | 52        |
| 3450 | Ribosome profiling analysis of eEF3-depleted <i>Saccharomyces cerevisiae</i> . <i>Scientific Reports</i> , 2019, 9, 3037.   | 1.6  | 18        |
| 3451 | Interphase Microtubules Safeguard Mitotic Progression by Suppressing an Aurora B-Dependent Arrest Induced by DNA Replication Stress. <i>Cell Reports</i> , 2019, 26, 2875-2889.e3.                      | 2.9  | 8         |
| 3452 | Mismatch recognition and subsequent processing have distinct effects on mitotic recombination intermediates and outcomes in yeast. <i>Nucleic Acids Research</i> , 2019, 47, 4554-4568.                 | 6.5  | 23        |
| 3453 | RNase H activities counteract a toxic effect of Polymerase $\beta$ in cells replicating with depleted dNTP pools. <i>Nucleic Acids Research</i> , 2019, 47, 4612-4623.                                  | 6.5  | 21        |
| 3454 | SUMOylation of the nuclear pore complex basket is involved in sensing cellular stresses. <i>Journal of Cell Science</i> , 2019, 132, .  | 1.2  | 15        |
| 3455 | Mitochondrial AAA-ATPase Msp1 detects mislocalized tail-anchored proteins through a dual-recognition mechanism. <i>EMBO Reports</i> , 2019, 20, .   | 2.0  | 41        |
| 3456 | Aggregate Filamentous Growth Responses in Yeast. <i>MSphere</i> , 2019, 4, .  | 1.3  | 20        |
| 3457 | Functional complementation reveals that 9 of the 13 human V-ATPase subunits can functionally substitute for their yeast orthologs. <i>Journal of Biological Chemistry</i> , 2019, 294, 8273-8285.       | 1.6  | 4         |
| 3458 | The Endoplasmic Reticulum-Mitochondria Encounter Structure Complex Coordinates Coenzyme Q Biosynthesis. <i>Contact (Thousand Oaks (Ventura County, Calif))</i> , 2019, 2, 251525641882540.              | 0.4  | 35        |
| 3459 | Suppression of respiratory growth defect of mitochondrial phosphatidylserine decarboxylase deficient mutant by overproduction of Sfh1, a Sec14 homolog, in yeast. <i>PLoS ONE</i> , 2019, 14, e0215009. | 1.1  | 6         |
| 3460 | Yeast Dop1 is required for glycosyltransferase retrieval from the trans-Golgi network. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2019, 1863, 1147-1157.                                 | 1.1  | 4         |
| 3461 | Yeast Ataxin-2 Forms an Intracellular Condensate Required for the Inhibition of TORC1 Signaling during Respiratory Growth. <i>Cell</i> , 2019, 177, 697-710.e17.  | 13.5 | 73        |
| 3462 | Redox State Controls Phase Separation of the Yeast Ataxin-2 Protein via Reversible Oxidation of Its Methionine-Rich Low-Complexity Domain. <i>Cell</i> , 2019, 177, 711-721.e8.                         | 13.5 | 121       |
| 3463 | KEOPS complex promotes homologous recombination via DNA resection. <i>Nucleic Acids Research</i> , 2019, 47, 5684-5697.   | 6.5  | 20        |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 3464 | Ccr4â€œNot maintains genomic integrity by controlling the ubiquitylation and degradation of arrested RNAPII. <i>Genes and Development</i> , 2019, 33, 705-717.  | 2.7  | 23        |
| 3465 | Following Anterograde Transport of Phosphatidylserine in Yeast in Real Time. <i>Methods in Molecular Biology</i> , 2019, 1949, 35-46.   | 0.4  | 5         |
| 3466 | A conserved genetic interaction between Spt6 and Set2 regulates H3K36 methylation. <i>Nucleic Acids Research</i> , 2019, 47, 3888-3903.   | 6.5  | 33        |
| 3467 | Excessive Cell Growth Causes Cytoplasm Dilution And Contributes to Senescence. <i>Cell</i> , 2019, 176, 1083-1097.e18.  | 13.5 | 347       |
| 3468 | Dual Role of Mitochondrial Porin in Metabolite Transport across the Outer Membrane and Protein Transfer to the Inner Membrane. <i>Molecular Cell</i> , 2019, 73, 1056-1065.e7.  | 4.5  | 51        |
| 3469 | The anaphase-promoting complex regulates the degradation of the inner nuclear membrane protein Mps3. <i>Journal of Cell Biology</i> , 2019, 218, 839-854.   | 2.3  | 27        |
| 3470 | A synthetic non-histone substrate to study substrate targeting by the Gcn5 HAT and sirtuin HDACs. <i>Journal of Biological Chemistry</i> , 2019, 294, 6227-6239.  | 1.6  | 9         |
| 3471 | The C-terminal region of Net1 is an activator of RNA polymerase I transcription with conserved features from yeast to human. <i>PLoS Genetics</i> , 2019, 15, e1008006.   | 1.5  | 7         |
| 3472 | Signalling through the yeast MAPK Cell Wall Integrity pathway controls P-body assembly upon cell wall stress. <i>Scientific Reports</i> , 2019, 9, 3186.  | 1.6  | 16        |
| 3473 | Demethylation of the Protein Phosphatase PP2A Promotes Demethylation of Histones to Enable Their Function as a Methyl Group Sink. <i>Molecular Cell</i> , 2019, 73, 1115-1126.e6.   | 4.5  | 28        |
| 3474 | A subtelomeric region affects telomerase-negative replicative senescence in <i>Saccharomyces cerevisiae</i> . <i>Scientific Reports</i> , 2019, 9, 1845.  | 1.6  | 9         |
| 3475 | The copper-sensing transcription factor Mac1, the histone deacetylase Hst1, and nicotinic acid regulate de novo NAD+ biosynthesis in budding yeast. <i>Journal of Biological Chemistry</i> , 2019, 294, 5562-5575.                                    | 1.6  | 15        |
| 3476 | Distinct Functions of the Cap-Binding Complex in Stimulation of Nuclear mRNA Export. <i>Molecular and Cellular Biology</i> , 2019, 39, .  | 1.1  | 15        |
| 3477 | A role for the <i>Saccharomyces cerevisiae</i> ABCF protein New1 in translation termination/recycling. <i>Nucleic Acids Research</i> , 2019, 47, 8807-8820.   | 6.5  | 26        |
| 3478 | A Stress-Responsive Signaling Network Regulating Pseudohyphal Growth and Ribonucleoprotein Granule Abundance in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2019, 213, 705-720.   | 1.2  | 6         |
| 3479 | Spatiotemporal dissection of the <i>trans</i> -Golgi network. <i>Journal of Cell Science</i> , 2019, 132, .   | 1.2  | 42        |
| 3480 | Deletions associated with stabilization of the Top1 cleavage complex in yeast are products of the nonhomologous end-joining pathway. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 22683-22691. | 3.3  | 15        |
| 3481 | Analysis of novel hyperosmotic shock response suggests â€œbeads in liquidâ€•cytosol structure. <i>Biology Open</i> , 2019, 8, .   | 0.6  | 18        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3482 | Actin and Nuclear Envelope Components Influence Ectopic Recombination in the Absence of Swr1. <i>Genetics</i> , 2019, 213, 819-834.   | 1.2 | 11        |
| 3483 | A Conserved PP2A Regulatory Subunit Enforces Proportional Relationships Between Cell Size and Growth Rate. <i>Genetics</i> , 2019, 213, 517-528.  | 1.2 | 8         |
| 3484 | Vac8 spatially confines autophagosome formation at the vacuole. <i>Journal of Cell Science</i> , 2019, 132, .   | 1.2 | 48        |
| 3485 | Tricalbins Contribute to Cellular Lipid Flux and Form Curved ER-PM Contacts that Are Bridged by Rod-Shaped Structures. <i>Developmental Cell</i> , 2019, 51, 488-502.e8.                                      | 3.1 | 72        |
| 3486 | Tricalbin-Mediated Contact Sites Control ER Curvature to Maintain Plasma Membrane Integrity. <i>Developmental Cell</i> , 2019, 51, 476-487.e7.  | 3.1 | 87        |
| 3487 | Role of the Cell Asymmetry Apparatus and Ribosome-Associated Chaperones in the Destabilization of a <i>Saccharomyces cerevisiae</i> Prion by Heat Shock. <i>Genetics</i> , 2019, 212, 757-771.                | 1.2 | 11        |
| 3488 | Proteins That Interact with the Mucin-Type Glycoprotein Msb2p Include a Regulator of the Actin Cytoskeleton. <i>Biochemistry</i> , 2019, 58, 4842-4856.   | 1.2 | 8         |
| 3489 | Protein Methylation and Translation: Role of Lysine Modification on the Function of Yeast Elongation Factor 1A. <i>Biochemistry</i> , 2019, 58, 4997-5010.  | 1.2 | 12        |
| 3490 | DNA variants affecting the expression of numerous genes in trans have diverse mechanisms of action and evolutionary histories. <i>PLoS Genetics</i> , 2019, 15, e1008375.                                     | 1.5 | 34        |
| 3491 | Overexpression of the peroxin Pex34p suppresses impaired acetate utilization in yeast lacking the mitochondrial aspartate/glutamate carrier Agc1p. <i>FEMS Yeast Research</i> , 2019, 19, .                   | 1.1 | 4         |
| 3492 | Yeast R2TP Interacts with Extended Termini of Client Protein Nop58p. <i>Scientific Reports</i> , 2019, 9, 20228.  | 1.6 | 5         |
| 3493 | Pol5 is required for recycling of small subunit biogenesis factors and for formation of the peptide exit tunnel of the large ribosomal subunit. <i>Nucleic Acids Research</i> , 2019, 48, 405-420.            | 6.5 | 7         |
| 3494 | A novel ER membrane protein Ehg1/May24 plays a critical role in maintaining multiple nutrient permeases in yeast under high-pressure perturbation. <i>Scientific Reports</i> , 2019, 9, 18341.                | 1.6 | 9         |
| 3495 | Pol5 is an essential ribosome biogenesis factor required for 60S ribosomal subunit maturation in <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 2019, 25, 1561-1575.  | 1.6 | 9         |
| 3496 | A kinase-dependent checkpoint prevents escape of immature ribosomes into the translating pool. <i>PLoS Biology</i> , 2019, 17, e3000329.  | 2.6 | 30        |
| 3497 | Yeast dynamin and Ypt6 function in parallel for the endosome-to-Golgi retrieval of Snc1. <i>Cell Biology International</i> , 2019, 43, 1137-1151.   | 1.4 | 3         |
| 3498 | Rpd3L Contributes to the DNA Damage Sensitivity of <i>Saccharomyces cerevisiae</i> Checkpoint Mutants. <i>Genetics</i> , 2019, 211, 503-513.  | 1.2 | 9         |
| 3499 | Glucose Signaling Is Connected to Chromosome Segregation Through Protein Kinase A Phosphorylation of the Dam1 Kinetochores Subunit in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2019, 211, 531-547. | 1.2 | 8         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3500 | Global analysis of protein homomerization in <i>Saccharomyces cerevisiae</i> . <i>Genome Research</i> , 2019, 29, 135-145.  | 2.4 | 12        |
| 3501 | Avidity-driven polarity establishment via multivalent lipid-GTPase module interactions. <i>EMBO Journal</i> , 2019, 38, .   | 3.5 | 27        |
| 3502 | Determinants of the assembly, integrity and maintenance of the endoplasmic reticulum-peroxisome tether. <i>Traffic</i> , 2019, 20, 213-225.   | 1.3 | 7         |
| 3503 | Collided ribosomes form a unique structural interface to induce Hel2-driven quality control pathways. <i>EMBO Journal</i> , 2019, 38, .   | 3.5 | 232       |
| 3504 | A cell separation checkpoint that enforces the proper order of late cytokinetic events. <i>Journal of Cell Biology</i> , 2019, 218, 150-170.  | 2.3 | 10        |
| 3505 | Proteins and RNA sequences required for the transition of the t-Utp complex into the SSU processome. <i>FEMS Yeast Research</i> , 2019, 19, .   | 1.1 | 6         |
| 3506 | Mrc1 and Tof1 prevent fragility and instability at long CAG repeats by their fork stabilizing function. <i>Nucleic Acids Research</i> , 2019, 47, 794-805.  | 6.5 | 29        |
| 3507 | Yeast chemogenomic screen identifies distinct metabolic pathways required to tolerate exposure to phenolic fermentation inhibitors ferulic acid, 4-hydroxybenzoic acid and coniferyl aldehyde. <i>Metabolic Engineering</i> , 2019, 52, 98-109. | 3.6 | 39        |
| 3508 | Cell-cycle control of cell polarity in yeast. <i>Journal of Cell Biology</i> , 2019, 218, 171-189.  | 2.3 | 41        |
| 3509 | Budding yeast Rtt107 prevents checkpoint hyperactivation after replicative stress by limiting DNA damage. <i>DNA Repair</i> , 2019, 74, 1-16.   | 1.3 | 4         |
| 3510 | A conserved mechanism for mitochondria-dependent dynein anchoring. <i>Molecular Biology of the Cell</i> , 2019, 30, 691-702.  | 0.9 | 19        |
| 3511 | Universal plasmids to facilitate gene deletion and gene tagging in filamentous fungi. <i>Fungal Genetics and Biology</i> , 2019, 125, 28-35.  | 0.9 | 8         |
| 3512 | Compositional reorganization of the nucleolus in budding yeast mitosis. <i>Molecular Biology of the Cell</i> , 2019, 30, 591-606.   | 0.9 | 10        |
| 3513 | Maintenance of Cardiolipin and Crista Structure Requires Cooperative Functions of Mitochondrial Dynamics and Phospholipid Transport. <i>Cell Reports</i> , 2019, 26, 518-528.e6.  | 2.9 | 48        |
| 3514 | The kinetochore module Okp1 <sup>CENP-C</sup> /Ame1 <sup>CENP-U</sup> is a reader for N-terminal modifications on the centromeric histone Cse4 <sup>CENP-A</sup> . <i>EMBO Journal</i> , 2019, 38, .  | 3.5 | 34        |
| 3515 | A fast and tuneable auxin-inducible degron for depletion of target proteins in budding yeast. <i>Yeast</i> , 2019, 36, 75-81.   | 0.8 | 27        |
| 3516 | Nuclear envelope expansion in budding yeast is independent of cell growth and does not determine nuclear volume. <i>Molecular Biology of the Cell</i> , 2019, 30, 131-145.  | 0.9 | 38        |
| 3517 | A Steric Gating Mechanism Dictates the Substrate Specificity of a Rab-GEF. <i>Developmental Cell</i> , 2019, 48, 100-114.e9.  | 3.1 | 49        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3518 | Developmental regulation of an organelle tether coordinates mitochondrial remodeling in meiosis. <i>Journal of Cell Biology</i> , 2019, 218, 559-579.   | 2.3 | 57        |
| 3519 | The Pat1-Lsm Complex Stabilizes ATG mRNA during Nitrogen Starvation-Induced Autophagy. <i>Molecular Cell</i> , 2019, 73, 314-324.e4.  | 4.5 | 28        |
| 3520 | The glycerophosphocholine acyltransferase Gpc1 is part of a phosphatidylcholine (PC)-remodeling pathway that alters PC species in yeast. <i>Journal of Biological Chemistry</i> , 2019, 294, 1189-1201.                   | 1.6 | 14        |
| 3521 | Interactions in the ESCRT-III network of the yeast <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2019, 65, 607-619.   | 0.8 | 13        |
| 3522 | Yeast genetic interaction screens in the age of CRISPR/Cas. <i>Current Genetics</i> , 2019, 65, 307-327.  | 0.8 | 29        |
| 3523 | An Early mtUPR: Redistribution of the Nuclear Transcription Factor Rox1 to Mitochondria Protects against Intramitochondrial Proteotoxic Aggregates. <i>Molecular Cell</i> , 2020, 77, 180-188.e9.                         | 4.5 | 53        |
| 3524 | The carboxy terminus of yeast Atg13 binds phospholipid membrane via motifs that overlap with the Vac8-interacting domain. <i>Autophagy</i> , 2020, 16, 1007-1020.   | 4.3 | 17        |
| 3525 | The transcription factor Spt4-Spt5 complex regulates the expression of <i>ATG8</i> and <i>ATG41</i> . <i>Autophagy</i> , 2020, 16, 1172-1185.   | 4.3 | 9         |
| 3526 | Single amino acid mutations in the <i>Saccharomyces cerevisiae</i> rhomboid peptidase, Pcp1p, alter mitochondrial morphology. <i>Cell Biology International</i> , 2020, 44, 200-215.                                      | 1.4 | 2         |
| 3527 | Crosstalk between <i>Saccharomyces cerevisiae</i> SAPKs Hog1 and Mpk1 is mediated by glycerol accumulation. <i>Fungal Biology</i> , 2020, 124, 361-367.   | 1.1 | 9         |
| 3528 | Quality Control of Protein Complex Assembly by a Transmembrane Recognition Factor. <i>Molecular Cell</i> , 2020, 77, 108-119.e9.  | 4.5 | 67        |
| 3529 | Pho85 and PI(4,5)P2 regulate different lipid metabolic pathways in response to cold. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2020, 1865, 158557.                                    | 1.2 | 10        |
| 3530 | Thermosensitive Nucleosome Editing Reveals the Role of DNA Sequence in Targeted Histone Variant Deposition. <i>Cell Reports</i> , 2020, 30, 257-268.e5.   | 2.9 | 16        |
| 3531 | Aurora B-dependent Ndc80 degradation regulates kinetochore composition in meiosis. <i>Genes and Development</i> , 2020, 34, 209-225.  | 2.7 | 16        |
| 3532 | SWI/SNF chromatin remodelling complex contributes to clearance of cytoplasmic protein aggregates and regulates unfolded protein response in <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 2020, 287, 3024-3041. | 2.2 | 7         |
| 3533 | MRX Increases Chromatin Accessibility at Stalled Replication Forks to Promote Nascent DNA Resection and Cohesin Loading. <i>Molecular Cell</i> , 2020, 77, 395-410.e3.  | 4.5 | 49        |
| 3534 | Cyclin-dependent kinase Pho85p and its cyclins are involved in replicative lifespan through multiple pathways in yeast. <i>FEBS Letters</i> , 2020, 594, 1166-1175.   | 1.3 | 7         |
| 3535 | SYCP2 Translocation-Mediated Dysregulation and Frameshift Variants Cause Human Male Infertility. <i>American Journal of Human Genetics</i> , 2020, 106, 41-57.  | 2.6 | 66        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3536 | Activity of the yeast cytoplasmic Hsp70 nucleotide-exchange factor Fes1 is regulated by reversible methionine oxidation. <i>Journal of Biological Chemistry</i> , 2020, 295, 552-569.                         | 1.6 | 19        |
| 3537 | A Regulatory Cysteine Residue Mediates Reversible Inactivation of NAD <sup>+</sup> -Dependent Aldehyde Dehydrogenases to Promote Oxidative Stress Response. <i>ACS Chemical Biology</i> , 2020, 15, 28-32.    | 1.6 | 4         |
| 3538 | ESCRT machinery mediates selective microautophagy of endoplasmic reticulum in yeast. <i>EMBO Journal</i> , 2020, 39, e102586.   | 3.5 | 77        |
| 3539 | Mechanism of Long-Range Chromosome Motion Triggered by Gene Activation. <i>Developmental Cell</i> , 2020, 52, 309-320.e5.   | 3.1 | 32        |
| 3540 | FACT and Ash1 promote long-range and bidirectional nucleosome eviction at the HO promoter. <i>Nucleic Acids Research</i> , 2020, 48, 10877-10889.   | 6.5 | 7         |
| 3541 | Yeast Rgd3 is a phospho-regulated F-BAR-containing RhoGAP involved in the regulation of Rho3 distribution and cell morphology. <i>Molecular Biology of the Cell</i> , 2020, 31, 2570-2582.                    | 0.9 | 4         |
| 3542 | Snf1 AMPK positively regulates ER-phagy via expression control of Atg39 autophagy receptor in yeast ER stress response. <i>PLoS Genetics</i> , 2020, 16, e1009053.  | 1.5 | 14        |
| 3543 | The yeast cell wall protein Pry3 inhibits mating through highly conserved residues within the CAP domain. <i>Biology Open</i> , 2020, 9, .  | 0.6 | 5         |
| 3544 | Homeostatic Control of Meiotic Prophase Checkpoint Function by Pch2 and Hop1. <i>Current Biology</i> , 2020, 30, 4413-4424.e5.  | 1.8 | 32        |
| 3545 | The pears and lemons protein UvPal1 regulates development and virulence of <i>Ustilago violacea</i> . <i>Environmental Microbiology</i> , 2020, 22, 5414-5432.  | 1.8 | 19        |
| 3546 | Cdk1 Phosphorylation of the Dam1 Complex Strengthens Kinetochores-Microtubule Attachments. <i>Current Biology</i> , 2020, 30, 4491-4499.e5.   | 1.8 | 17        |
| 3547 | Cooperativity between the Ribosome-Associated Chaperone Ssb/RAC and the Ubiquitin Ligase Ltn1 in Ubiquitination of Nascent Polypeptides. <i>International Journal of Molecular Sciences</i> , 2020, 21, 6815. | 1.8 | 4         |
| 3548 | Degradation of Non-coding RNAs Promotes Recycling of Termination Factors at Sites of Transcription. <i>Cell Reports</i> , 2020, 32, 107942.   | 2.9 | 19        |
| 3549 | The Sur7/Pal family transmembrane protein Tos7 (Yol019w) plays a role in secretion in budding yeast. <i>Fungal Genetics and Biology</i> , 2020, 144, 103467.  | 0.9 | 0         |
| 3550 | Membrane dynamics and protein targets of lipid droplet microautophagy during ER stress-induced proteostasis in the budding yeast, <i>Saccharomyces cerevisiae</i> . <i>Autophagy</i> , 2021, 17, 2363-2383.   | 4.3 | 27        |
| 3551 | Vps13 is required for the packaging of the ER into autophagosomes during ER-phagy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 18530-18539.           | 3.3 | 42        |
| 3552 | Structural insights into telomere protection and homeostasis regulation by yeast CST complex. <i>Nature Structural and Molecular Biology</i> , 2020, 27, 752-762.   | 3.6 | 23        |
| 3553 | Chromatin remodeler Ino80C acts independently of H2A.Z to evict promoter nucleosomes and stimulate transcription of highly expressed genes in yeast. <i>Nucleic Acids Research</i> , 2020, 48, 8408-8430.     | 6.5 | 15        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3554 | Construction of the Central Protuberance and L1 Stalk during 60S Subunit Biogenesis. <i>Molecular Cell</i> , 2020, 79, 615-628.e5.  | 4.5 | 48        |
| 3555 | Free convective PCR: From principle study to commercial applicationsâ€”A critical review. <i>Analytica Chimica Acta</i> , 2020, 1108, 177-197.  | 2.6 | 27        |
| 3556 | Roles of the PH, coiled-coil and SAM domains of the yeast polarity protein Boi2 in polarity-site localization and function in polarized growth. <i>Current Genetics</i> , 2020, 66, 1101-1115.                | 0.8 | 4         |
| 3557 | Poly(A)-Binding Protein Regulates the Efficiency of Translation Termination. <i>Cell Reports</i> , 2020, 33, 108399.  | 2.9 | 25        |
| 3558 | Irc20 Regulates the Yeast Endogenous 2-Î¼m Plasmid Levels by Controlling Flp1. <i>Frontiers in Molecular Biosciences</i> , 2020, 7, 221.  | 1.6 | 0         |
| 3559 | Histone sumoylation promotes Set3 histone-deacetylase complex-mediated transcriptional regulation. <i>Nucleic Acids Research</i> , 2020, 48, 12151-12168.   | 6.5 | 28        |
| 3560 | Ribosomal protein S7 ubiquitination during ER stress in yeast is associated with selective mRNA translation and stress outcome. <i>Scientific Reports</i> , 2020, 10, 19669.                                  | 1.6 | 21        |
| 3561 | The Proteasome Activators Blm10/PA200 Enhance the Proteasomal Degradation of N-Terminal Huntingtin. <i>Biomolecules</i> , 2020, 10, 1581.   | 1.8 | 9         |
| 3562 | Nutrient Signaling, Stress Response, and Inter-organelle Communication Are Non-canonical Determinants of Cell Fate. <i>Cell Reports</i> , 2020, 33, 108446.   | 2.9 | 20        |
| 3563 | Sphingolipids and Inositol Phosphates Regulate the Tau Protein Phosphorylation Status in Humanized Yeast. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 592159.                               | 1.8 | 7         |
| 3564 | The catalytic activity of the translation termination factor methyltransferase Mtq2-Trm112 complex is required for large ribosomal subunit biogenesis. <i>Nucleic Acids Research</i> , 2020, 48, 12310-12325. | 6.5 | 9         |
| 3565 | SAGAâ€™CORE subunit Spt7 is required for correct Ubp8 localization, chromatin association and deubiquitinase activity. <i>Epigenetics and Chromatin</i> , 2020, 13, 46.                                       | 1.8 | 3         |
| 3566 | Semisynthesis of site-specifically succinylated histone reveals that succinylation regulates nucleosome unwrapping rate and DNA accessibility. <i>Nucleic Acids Research</i> , 2020, 48, 9538-9549.           | 6.5 | 34        |
| 3567 | Plasma membrane to vacuole traffic induced by glucose starvation requires Gga2â€™dependent sorting at the trans-Golgi network. <i>Biology of the Cell</i> , 2020, 112, 349-367.                               | 0.7 | 12        |
| 3568 | Aggregation and Prion-Inducing Properties of the G-Protein Gamma Subunit Ste18 are Regulated by Membrane Association. <i>International Journal of Molecular Sciences</i> , 2020, 21, 5038.                    | 1.8 | 4         |
| 3569 | Coupling of 5S RNP rotation with maturation of functional centers during large ribosomal subunit assembly. <i>Nature Communications</i> , 2020, 11, 3751.   | 5.8 | 24        |
| 3570 | Rrp6 Moonlights in an RNA Exosome-Independent Manner to Promote Cell Survival and Gene Expression during Stress. <i>Cell Reports</i> , 2020, 31, 107754.  | 2.9 | 16        |
| 3571 | Yeast pericentrin/Spc110 contains multiple domains required for tethering the Î³-tubulin complex to the centrosome. <i>Molecular Biology of the Cell</i> , 2020, 31, 1437-1452.                               | 0.9 | 2         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3572 | Translation Initiation Site Profiling Reveals Widespread Synthesis of Non-AUG-Initiated Protein Isoforms in Yeast. <i>Cell Systems</i> , 2020, 11, 145-160.e5.   | 2.9 | 41        |
| 3573 | Senataxin Ortholog Sen1 Limits DNA:RNA Hybrid Accumulation at DNA Double-Strand Breaks to Control End Resection and Repair Fidelity. <i>Cell Reports</i> , 2020, 31, 107603.   | 2.9 | 35        |
| 3574 | The Cdc48 Complex Alleviates the Cytotoxicity of Misfolded Proteins by Regulating Ubiquitin Homeostasis. <i>Cell Reports</i> , 2020, 32, 107898.   | 2.9 | 22        |
| 3575 | The LKB1-like Kinase Elm1 Controls Septin Hourglass Assembly and Stability by Regulating Filament Pairing. <i>Current Biology</i> , 2020, 30, 2386-2394.e4.  | 1.8 | 12        |
| 3576 | A Noncanonical Hippo Pathway Regulates Spindle Disassembly and Cytokinesis During Meiosis in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2020, 216, 447-462.   | 1.2 | 4         |
| 3577 | GPCR receptor phosphorylation and endocytosis are not necessary to switch polarized growth between internal cues during pheromone response in <i>S. cerevisiae</i> . <i>Communicative and Integrative Biology</i> , 2020, 13, 128-139. | 0.6 | 1         |
| 3578 | SWR1-Independent Association of H2A.Z to the LINC Complex Promotes Meiotic Chromosome Motion. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 594092.  | 1.8 | 10        |
| 3579 | Structural insights into assembly of the ribosomal nascent polypeptide exit tunnel. <i>Nature Communications</i> , 2020, 11, 5111.   | 5.8 | 21        |
| 3580 | The ESCRT-III complex is required for nuclear pore complex sequestration and regulates gamete replicative lifespan in budding yeast meiosis. <i>Nucleus</i> , 2020, 11, 219-236.   | 0.6 | 11        |
| 3581 | Mechanisms coordinating ribosomal protein gene transcription in response to stress. <i>Nucleic Acids Research</i> , 2020, 48, 11408-11420.   | 6.5 | 27        |
| 3582 | The conserved elongation factor Spn1 is required for normal transcription, histone modifications, and splicing in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2020, 48, 10241-10258.                             | 6.5 | 21        |
| 3583 | Conserved Ark1-related kinases function in a TORC2 signaling network. <i>Molecular Biology of the Cell</i> , 2020, 31, 2057-2069.  | 0.9 | 4         |
| 3584 | Functional information from clinically-derived drug resistant forms of the <i>Candida glabrata</i> Pdr1 transcription factor. <i>PLoS Genetics</i> , 2020, 16, e1009005.   | 1.5 | 26        |
| 3585 | An optogenetic system to control membrane phospholipid asymmetry through flippase activation in budding yeast. <i>Scientific Reports</i> , 2020, 10, 12474.  | 1.6 | 6         |
| 3586 | Excess diacylglycerol at the endoplasmic reticulum disrupts endomembrane homeostasis and autophagy. <i>BMC Biology</i> , 2020, 18, 107.  | 1.7 | 12        |
| 3587 | PCNA promotes context-specific sister chromatid cohesion establishment separate from that of chromatin condensation. <i>Cell Cycle</i> , 2020, 19, 2436-2450.  | 1.3 | 7         |
| 3588 | Overexpression of Mdm36 reveals Num1 foci that mediate dynein-dependent microtubule sliding in budding yeast. <i>Journal of Cell Science</i> , 2020, 133, .  | 1.2 | 7         |
| 3589 | Methylglyoxal inhibits nuclear division through alterations in vacuolar morphology and accumulation of Atg18 on the vacuolar membrane in <i>Saccharomyces cerevisiae</i> . <i>Scientific Reports</i> , 2020, 10, 13887.                | 1.6 | 6         |



| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 3590 | Cdc48 cofactor Shp1 regulates signal-induced SCF <sup>Met30</sup> disassembly. Proceedings of the National Academy of Sciences of the United States of America, 2020, 117, 21319-21327.   | 3.3  | 7         |
| 3591 | Loss of the seipin gene perturbs eggshell formation in <i>C. elegans</i> . Development (Cambridge), 2020, 147, .  | 1.2  | 19        |
| 3592 | A complex genetic interaction implicates that phospholipid asymmetry and phosphate homeostasis regulate Golgi functions. PLoS ONE, 2020, 15, e0236520.  | 1.1  | 5         |
| 3593 | Maturation Kinetics of a Multiprotein Complex Revealed by Metabolic Labeling. Cell, 2020, 183, 1785-1800.e26.   | 13.5 | 34        |
| 3594 | Proline-Rich Motifs Control G2-CDK Target Phosphorylation and Priming an Anchoring Protein for Polo Kinase Localization. Cell Reports, 2020, 31, 107757.  | 2.9  | 16        |
| 3595 | Deletion of the GAPDH gene contributes to genome stability in <i>Saccharomyces cerevisiae</i> . Scientific Reports, 2020, 10, 21146.  | 1.6  | 4         |
| 3596 | Swc4 positively regulates telomere length independently of its roles in NuA4 and SWR1 complexes. Nucleic Acids Research, 2020, 48, 12792-12803.   | 6.5  | 8         |
| 3597 | A nanobody suite for yeast scaffold nucleoporins provides details of the nuclear pore complex structure. Nature Communications, 2020, 11, 6179.   | 5.8  | 12        |
| 3598 | An F-Box Protein, Mdm30, Interacts with TREX Subunit Sub2 To Regulate Cellular Abundance Cotranscriptionally in Orchestrating mRNA Export Independently of Splicing and Mitochondrial Function. Molecular and Cellular Biology, 2020, 40, . | 1.1  | 6         |
| 3599 | Stability and nuclear localization of yeast telomerase depend on protein components of RNase P/MRP. Nature Communications, 2020, 11, 2173.  | 5.8  | 23        |
| 3600 | Relocation of Collapsed Forks to the Nuclear Pore Complex Depends on Sumoylation of DNA Repair Proteins and Permits Rad51 Association. Cell Reports, 2020, 31, 107635.  | 2.9  | 43        |
| 3601 | The degradation-promoting roles of deubiquitinases Ubp6 and Ubp3 in cytosolic and ER protein quality control. PLoS ONE, 2020, 15, e0232755.   | 1.1  | 3         |
| 3602 | Regulated repression governs the cell fate promoter controlling yeast meiosis. Nature Communications, 2020, 11, 2271.   | 5.8  | 22        |
| 3603 | Cdc24 interacts with the septins to create a positive feedback during bud site assembly in yeast. Journal of Cell Science, 2020, 133, .   | 1.2  | 12        |
| 3604 | Growth-Dependent Activation of Protein Kinases Suggests a Mechanism for Measuring Cell Growth. Genetics, 2020, 215, 729-746.  | 1.2  | 10        |
| 3605 | Phosphorylation of the RecQ Helicase Sgs1/BLM Controls Its DNA Unwinding Activity during Meiosis and Mitosis. Developmental Cell, 2020, 53, 706-723.e5.   | 3.1  | 26        |
| 3606 | Functions for Cdc42p BEM adaptors in regulating a differentiation-type MAP kinase pathway. Molecular Biology of the Cell, 2020, 31, 491-510.  | 0.9  | 16        |
| 3607 | <i>Saccharomyces cerevisiae</i> nutrient signaling pathways show an unexpected early activation pattern during winemaking. Microbial Cell Factories, 2020, 19, 124.   | 1.9  | 10        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3608 | Identification of an extracytoplasmic function sigma factor that facilitates <i>σ</i> <sup>70</sup> type cytochrome maturation and current generation under electrolyte flow conditions in <i>Shewanella oneidensis</i> . <i>Environmental Microbiology</i> , 2020, 22, 3671-3684. | 1.8 | 9         |
| 3609 | Nucleolar localization of the yeast RNA exosome subunit Rrp44 hints at early pre-rRNA processing as its main function. <i>Journal of Biological Chemistry</i> , 2020, 295, 11195-11213.  | 1.6 | 4         |
| 3610 | Vac8 determines phagophore assembly site vacuolar localization during nitrogen starvation-induced autophagy. <i>Autophagy</i> , 2021, 17, 1636-1648.   | 4.3 | 22        |
| 3611 | Doa4 directly binds Snf7 to inhibit the recruitment of ESCRT-III remodeling factors. <i>Journal of Cell Science</i> , 2020, 133, .   | 1.2 | 10        |
| 3612 | High density of unrepaired genomic ribonucleotides leads to Topoisomerase 1-mediated severe growth defects in absence of ribonucleotide reductase. <i>Nucleic Acids Research</i> , 2020, 48, 4274-4297.  | 6.5 | 8         |
| 3613 | Mec1 Is Activated at the Onset of Normal S Phase by Low-dNTP Pools Impeding DNA Replication. <i>Molecular Cell</i> , 2020, 78, 396-410.e4.   | 4.5 | 48        |
| 3614 | An Essential and Cell-Cycle-Dependent ORC Dimerization Cycle Regulates Eukaryotic Chromosomal DNA Replication. <i>Cell Reports</i> , 2020, 30, 3323-3338.e6.   | 2.9 | 8         |
| 3615 | Labile heme critically regulates mitochondrial biogenesis through the transcriptional co-activator Hap4p in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2020, 295, 5095-5109.   | 1.6 | 10        |
| 3616 | Critical Roles of a RhoGEF-Anillin Module in Septin Architectural Remodeling during Cytokinesis. <i>Current Biology</i> , 2020, 30, 1477-1490.e3.  | 1.8 | 25        |
| 3617 | RQT complex dissociates ribosomes collided on endogenous RQC substrate SDD1. <i>Nature Structural and Molecular Biology</i> , 2020, 27, 323-332.   | 3.6 | 97        |
| 3618 | Recruitment of an Activated Gene to the Yeast Nuclear Pore Complex Requires Sumoylation. <i>Frontiers in Genetics</i> , 2020, 11, 174.   | 1.1 | 7         |
| 3619 | Specific proteomic adaptation to distinct environments in <i>Vibrio parahaemolyticus</i> includes significant fluctuations in expression of essential proteins. <i>Environmental Microbiology</i> , 2020, 22, 4279-4294.   | 1.8 | 3         |
| 3620 | Plasma membrane tension regulates eisosome structure and function. <i>Molecular Biology of the Cell</i> , 2020, 31, 287-303.   | 0.9 | 38        |
| 3621 | Multi-kinase control of environmental stress responsive transcription. <i>PLoS ONE</i> , 2020, 15, e0230246.   | 1.1 | 12        |
| 3622 | An intergenic safe haven region in <i>Aspergillus fumigatus</i> . <i>Medical Mycology</i> , 2020, 58, 1178-1186.   | 0.3 | 12        |
| 3623 | A heat-sensitive Osh protein controls PI4P polarity. <i>BMC Biology</i> , 2020, 18, 28.  | 1.7 | 11        |
| 3624 | Role of conserved regions of Tim22 in the structural organization of the carrier translocase. <i>Journal of Cell Science</i> , 2020, 133, .  | 1.2 | 6         |
| 3625 | Inhibitory effect of arginine on proline utilization in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2020, 37, 531-540.  | 0.8 | 16        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3626 | The histone H3-H4 tetramer is a copper reductase enzyme. <i>Science</i> , 2020, 369, 59-64.   | 6.0 | 60        |
| 3627 | PP2ACdc55 dephosphorylates Pds1 and inhibits spindle elongation. <i>Journal of Cell Science</i> , 2020, 133, .  | 1.2 | 4         |
| 3628 | A method for efficient quantitative analysis of genomic subtelomere Yâ€² element abundance in yeasts. <i>Yeast</i> , 2020, 37, 373-388.   | 0.8 | 3         |
| 3629 | Nascent Transcript Folding Plays a Major Role in Determining RNA Polymerase Elongation Rates. <i>Molecular Cell</i> , 2020, 79, 488-503.e11.  | 4.5 | 52        |
| 3630 | Rad9/53BP1 promotes DNA repair via crossover recombination by limiting the Sgs1 and Mph1 helicases. <i>Nature Communications</i> , 2020, 11, 3181.  | 5.8 | 18        |
| 3631 | Regulation of the Ysh1 endonuclease of the mRNA cleavage/polyadenylation complex by ubiquitin-mediated degradation. <i>RNA Biology</i> , 2020, 17, 689-702.   | 1.5 | 7         |
| 3632 | A VPS13D spastic ataxia mutation disrupts the conserved adaptor-binding site in yeast Vps13. <i>Human Molecular Genetics</i> , 2020, 29, 635-648.   | 1.4 | 35        |
| 3633 | Skp, Cullin, F-box (SCF)-Met30 and SCF-Cdc4-Mediated Proteolysis of CENP-A Prevents Mislocalization of CENP-A for Chromosomal Stability in Budding Yeast. <i>PLoS Genetics</i> , 2020, 16, e1008597.  | 1.5 | 28        |
| 3634 | mRNA with Mammalian Codon Bias Accumulates in Yeast Mutants with Constitutive Stress Granules. <i>International Journal of Molecular Sciences</i> , 2020, 21, 1234.   | 1.8 | 3         |
| 3635 | The ATAD2/ANCCA homolog Yta7 cooperates with Scm3 <sup>HJURP</sup> to deposit Cse4 <sup>CENP-A</sup> at the centromere in yeast. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 5386-5393. | 3.3 | 16        |
| 3636 | Interaction between the yeast RAVE complex and Vph1-containing Vo sectors is a central glucose-sensitive interaction required for V-ATPase reassembly. <i>Journal of Biological Chemistry</i> , 2020, 295, 2259-2269.                           | 1.6 | 15        |
| 3637 | Extranuclear Structural Components that Mediate Dynamic Chromosome Movements in Yeast Meiosis. <i>Current Biology</i> , 2020, 30, 1207-1216.e4.   | 1.8 | 24        |
| 3638 | Yeast transformation stress, together with loss of Pah1, phosphatidic acid phosphatase, leads to Ty1 retrotransposon insertion into theINO4gene. <i>FASEB Journal</i> , 2020, 34, 4749-4763.  | 0.2 | 3         |
| 3639 | Pre-emptive Quality Control of a Misfolded Membrane Protein by Ribosome-Driven Effects. <i>Current Biology</i> , 2020, 30, 854-864.e5.  | 1.8 | 36        |
| 3640 | Functional interrelationships between carbohydrate and lipid storage, and mitochondrial activity during sporulation in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2020, 37, 269-279.  | 0.8 | 3         |
| 3641 | Autophagy of an Amyloid-like Translational Repressor Regulates Meiotic Exit. <i>Developmental Cell</i> , 2020, 52, 141-151.e5.  | 3.1 | 32        |
| 3642 | Structural and functional characterization of the dominant negative Pâ€loop lysine mutation in the dynamin superfamily protein Vps1. <i>Protein Science</i> , 2020, 29, 1416-1428.  | 3.1 | 6         |
| 3643 | Sec16 function in ER export and autophagy is independent of its phosphorylation in<i>Saccharomyces cerevisiae</i>. <i>Molecular Biology of the Cell</i> , 2020, 31, 149-156.  | 0.9 | 5         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3644 | LAM Genes Contribute to Environmental Stress Tolerance but Sensibilize Yeast Cells to Azoles. <i>Frontiers in Microbiology</i> , 2020, 11, 38.   | 1.5 | 14        |
| 3645 | Genetic Analysis of Peroxisomal Genes Required for Longevity in a Yeast Model of Citrin Deficiency. <i>Diseases (Basel, Switzerland)</i> , 2020, 8, 2.   | 1.0 | 2         |
| 3646 | Overexpression of budding yeast protein phosphatase Ppz1 impairs translation. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2020, 1867, 118727.   | 1.9 | 13        |
| 3647 | The coordinate actions of calcineurin and Hog1 mediate the stress response through multiple nodes of the cell cycle network. <i>PLoS Genetics</i> , 2020, 16, e1008600.  | 1.5 | 12        |
| 3648 | Mitochondrial Fission and Fusion Dynamics Generate Efficient, Robust, and Evenly Distributed Network Topologies in Budding Yeast Cells. <i>Cell Systems</i> , 2020, 10, 287-297.e5.                                  | 2.9 | 37        |
| 3649 | Yeast synaptobrevin, Snc1, engages distinct routes of postendocytic recycling mediated by a sorting nexin, Rcy1-COPI, and retromer. <i>Molecular Biology of the Cell</i> , 2020, 31, 944-962.                        | 0.9 | 22        |
| 3650 | Investigation of Heterologously Expressed Glucose-6-Phosphate Dehydrogenase Genes in a Yeast <i>zwf1</i> Deletion. <i>Microorganisms</i> , 2020, 8, 546.   | 1.6 | 19        |
| 3651 | Regulation of intrinsic polarity establishment by a differentiation-type MAPK pathway. <i>Journal of Cell Science</i> , 2020, 133, .   | 1.2 | 11        |
| 3652 | Live-cell imaging of mitochondrial motility and interactions in <i>Drosophila</i> neurons and yeast. <i>Methods in Cell Biology</i> , 2020, 155, 519-544.  | 0.5 | 3         |
| 3653 | A processive phosphorylation circuit with multiple kinase inputs and mutually diversional routes controls G1/S decision. <i>Nature Communications</i> , 2020, 11, 1836.  | 5.8 | 12        |
| 3654 | N-terminal protein acetylation by NatB modulates the levels of Nmnats, the NAD <sup>+</sup> biosynthetic enzymes in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2020, 295, 7362-7375. | 1.6 | 6         |
| 3655 | <scp>ESCRT</scp>-dependent protein sorting is required for the viability of yeast clathrin-mediated endocytosis mutants. <i>Traffic</i> , 2020, 21, 430-450.   | 1.3 | 4         |
| 3656 | The Mgs1/WRNIP1 ATPase is required to prevent a recombination salvage pathway at damaged replication forks. <i>Science Advances</i> , 2020, 6, eaaz3327.   | 4.7 | 11        |
| 3657 | The Ccr4-Not complex monitors the translating ribosome for codon optimality. <i>Science</i> , 2020, 368, .   | 6.0 | 180       |
| 3658 | A genetic interaction map centered on cohesin reveals auxiliary factors in sister chromatid cohesion. <i>Journal of Cell Science</i> , 2020, 133, .  | 1.2 | 5         |
| 3659 | Atg21 organizes Atg8 lipidation at the contact of the vacuole with the phagophore. <i>Autophagy</i> , 2021, 17, 1458-1478.   | 4.3 | 23        |
| 3660 | TORC1 Determines Fab1 Lipid Kinase Function at Signaling Endosomes and Vacuoles. <i>Current Biology</i> , 2021, 31, 297-309.e8.  | 1.8 | 31        |
| 3661 | Meiotic Cells Counteract Programmed Retrotransposon Activation via RNA-Binding Translational Repressor Assemblies. <i>Developmental Cell</i> , 2021, 56, 22-35.e7.   | 3.1 | 8         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3662 | CG32803 is the fly homolog of LDAF1 and influences lipid storage in vivo. <i>Insect Biochemistry and Molecular Biology</i> , 2021, 133, 103512.  | 1.2 | 6         |
| 3663 | MÃ¶ssbauer and LC-ICP-MS investigation of iron trafficking between vacuoles and mitochondria in vma21 <sup>Δ</sup> <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2021, 296, 100141. | 1.6 | 8         |
| 3664 | Hypoosmolality impedes cytoophidium integrity during nitrogen starvation. <i>Yeast</i> , 2021, 38, 276-289.  | 0.8 | 10        |
| 3665 | Phosphatidylserine flux into mitochondria unveiled by organelle-targeted <i>Escherichia coli</i> phosphatidylserine synthase PssA. <i>FEBS Journal</i> , 2021, 288, 3285-3299.                                   | 2.2 | 5         |
| 3666 | The ATPase Irc20 facilitates Rad51 chromatin enrichment during homologous recombination in yeast <i>Saccharomyces cerevisiae</i> . <i>DNA Repair</i> , 2021, 97, 103019.   | 1.3 | 0         |
| 3667 | The G-patch activators Pfa1 and PINX1 exhibit different modes of interaction with the Prp43 RNA helicase. <i>RNA Biology</i> , 2021, 18, 510-522.  | 1.5 | 7         |
| 3668 | Mitochondrial Origin of Cytosolic Protein Aggregation. <i>SSRN Electronic Journal</i> , 0, , .   | 0.4 | 1         |
| 3673 | Autophagy sustains glutamate and aspartate synthesis in <i>Saccharomyces cerevisiae</i> during nitrogen starvation. <i>Nature Communications</i> , 2021, 12, 57.   | 5.8 | 24        |
| 3674 | The Smk1 MAPK and Its Activator, Ssp2, Are Required for Late Prospore Membrane Development in Sporulating <i>Saccharomyces cerevisiae</i> . <i>Journal of Fungi (Basel, Switzerland)</i> , 2021, 7, 53.          | 1.5 | 2         |
| 3675 | Metabolic regulation of telomere silencing by SESAME complex-catalyzed H3T11 phosphorylation. <i>Nature Communications</i> , 2021, 12, 594.  | 5.8 | 18        |
| 3676 | The vacuolar amino acid transport system is a novel, direct target of GATA transcription factors. <i>Bioscience, Biotechnology and Biochemistry</i> , 2021, 85, 587-599.   | 0.6 | 2         |
| 3679 | Proteasome activator Blm10 levels and autophagic degradation directly impact the proteasome landscape. <i>Journal of Biological Chemistry</i> , 2021, 296, 100468.   | 1.6 | 6         |
| 3680 | High-resolution analysis of cell-state transitions in yeast suggests widespread transcriptional tuning by alternative starts. <i>Genome Biology</i> , 2021, 22, 34.  | 3.8 | 30        |
| 3681 | Direct binding of ESCRT protein Chm7 to phosphatidic acid-rich membranes at nuclear envelope herniations. <i>Journal of Cell Biology</i> , 2021, 220, .  | 2.3 | 40        |
| 3683 | Snd3 controls nucleus-vacuole junctions in response to glucose signaling. <i>Cell Reports</i> , 2021, 34, 108637.  | 2.9 | 22        |
| 3684 | Bil2 Is a Novel Inhibitor of the Yeast Formin Bnr1 Required for Proper Actin Cable Organization and Polarized Secretion. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 634587.                   | 1.8 | 1         |
| 3688 | Identification of novel lipid droplet factors that regulate lipophagy and cholesterol efflux in macrophage foam cells. <i>Autophagy</i> , 2021, 17, 3671-3689.   | 4.3 | 90        |
| 3690 | Structural basis of Stu2 recruitment to yeast kinetochores. <i>ELife</i> , 2021, 10, .   | 2.8 | 11        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3691 | The UBA domain of conjugating enzyme Ubc1/Ube2K facilitates assembly of K48/K63-linked branched ubiquitin chains. <i>EMBO Journal</i> , 2021, 40, e106094.  | 3.5 | 25        |
| 3692 | The yeast ISW1b ATP-dependent chromatin remodeler is critical for nucleosome spacing and dinucleosome resolution. <i>Scientific Reports</i> , 2021, 11, 4195.   | 1.6 | 12        |
| 3693 | Chemical-Genetic Interactions of <i>Bacopa monnieri</i> Constituents in Cells Deficient for the DNA Repair Endonuclease RAD1 Appear Linked to Vacuolar Disruption. <i>Molecules</i> , 2021, 26, 1207.   | 1.7 | 4         |
| 3694 | Pre-existing bilayer stresses modulate triglyceride accumulation in the ER versus lipid droplets. <i>ELife</i> , 2021, 10, .  | 2.8 | 55        |
| 3695 | Arf1 orchestrates Rab GTPase conversion at the trans-Golgi network. <i>Molecular Biology of the Cell</i> , 2021, 32, mbc.E20-10-0664.   | 0.9 | 15        |
| 3697 | A Conserved Histone H3-H4 Interface Regulates DNA Damage Tolerance and Homologous Recombination during the Recovery from Replication Stress. <i>Molecular and Cellular Biology</i> , 2021, 41, .  | 1.1 | 3         |
| 3698 | OXPHOS deficiency activates global adaptation pathways to maintain mitochondrial membrane potential. <i>EMBO Reports</i> , 2021, 22, e51606.  | 2.0 | 34        |
| 3699 | The eIF4E-binding protein Eap1 has similar but independent roles in cell growth and gene expression with the cytoplasmic deadenylase Ccr4. <i>Bioscience, Biotechnology and Biochemistry</i> , 2021, 85, 1452-1459.                                   | 0.6 | 1         |
| 3700 | Rad9, a 53BP1 Ortholog of Budding Yeast, Is Insensitive to Spo11-Induced Double-Strand Breaks During Meiosis. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 635383.   | 1.8 | 8         |
| 3701 | The ribosome collision sensor Hel2 functions as preventive quality control in the secretory pathway. <i>Cell Reports</i> , 2021, 34, 108877.  | 2.9 | 14        |
| 3704 | Reprogramming of translation in yeast cells impaired for ribosome recycling favors short, efficiently translated mRNAs. <i>ELife</i> , 2021, 10, .  | 2.8 | 22        |
| 3706 | Phosphoproteomics reveals a distinctive Mec1/ATR signaling response upon DNA end hyper-resection. <i>EMBO Journal</i> , 2021, 40, e104566.  | 3.5 | 17        |
| 3709 | A novel allele of <i>SIR2</i> reveals a heritable intermediate state of gene silencing. <i>Genetics</i> , 2021, 218, .  | 1.2 | 1         |
| 3711 | Regulation of translation by methylation multiplicity of 18S rRNA. <i>Cell Reports</i> , 2021, 34, 108825.  | 2.9 | 16        |
| 3712 | A nuclear-based quality control pathway for non-imported mitochondrial proteins. <i>ELife</i> , 2021, 10, .   | 2.8 | 46        |
| 3714 | Complex Mechanisms of Antimony Genotoxicity in Budding Yeast Involves Replication and Topoisomerase I-Associated DNA Lesions, Telomere Dysfunction and Inhibition of DNA Repair. <i>International Journal of Molecular Sciences</i> , 2021, 22, 4510. | 1.8 | 4         |
| 3718 | Pbp1 mediates the aberrant expression of genes involved in growth defect of <i>ccr4</i> and <i>pop2</i> mutants in yeast <i>Saccharomyces cerevisiae</i> . <i>Genes To Cells</i> , 2021, 26, 381-398.   | 0.5 | 5         |
| 3719 | Mechanisms that ensure monogamous mating in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2021, 32, 638-644.   | 0.9 | 6         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3720 | Genetic interaction between glyoxylate pathway regulator UCC1 and Laâ€motifâ€encoding SRO9 regulates stress response and growth rate improvement in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biochemical and Molecular Toxicology</i> , 2021, 35, e22781. | 1.4 | 2         |
| 3723 | Sir4 Deficiency Reverses Cell Senescence by Sub-Telomere Recombination. <i>Cells</i> , 2021, 10, 778.  | 1.8 | 4         |
| 3725 | Quorum sensing regulates rRNA synthesis in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 2021, 776, 145442.  | 1.0 | 2         |
| 3726 | How cells determine the number of polarity sites. <i>ELife</i> , 2021, 10, .   | 2.8 | 15        |
| 3727 | Coevolution of ribosomal RNA expansion segment 7L and assembly factor Noc2p specializes the ribosome biogenesis pathway between <i>Saccharomyces cerevisiae</i> and <i>Candida albicans</i> . <i>Nucleic Acids Research</i> , 2021, 49, 4655-4667.               | 6.5 | 4         |
| 3728 | The Pif1 helicase is actively inhibited during meiotic recombination which restrains gene conversion tract length. <i>Nucleic Acids Research</i> , 2021, 49, 4522-4533.  | 6.5 | 16        |
| 3731 | Multiplexed mRNA assembly into ribonucleoprotein particles plays an operon-like role in the control of yeast cell physiology. <i>ELife</i> , 2021, 10, .   | 2.8 | 16        |
| 3732 | Exomer complex regulates protein traffic at the TGN through differential interactions with cargos and clathrin adaptor complexes. <i>FASEB Journal</i> , 2021, 35, e21615.   | 0.2 | 9         |
| 3733 | Yeast Fin1-PP1 dephosphorylates an Ipl1 substrate, Ndc80, to remove Bub1-Bub3 checkpoint proteins from the kinetochore during anaphase. <i>PLoS Genetics</i> , 2021, 17, e1009592.   | 1.5 | 5         |
| 3734 | Arf1 directly recruits the Pik1-Frq1 PI4K complex to regulate the final stages of Golgi maturation. <i>Molecular Biology of the Cell</i> , 2021, 32, 1064-1080.  | 0.9 | 32        |
| 3735 | Mps1 promotes poleward chromosome movements in meiotic prometaphase. <i>Molecular Biology of the Cell</i> , 2021, 32, 1020-1032.   | 0.9 | 2         |
| 3736 | Dynamics of SAS-I mediated H4 K16 acetylation during DNA replication in yeast. <i>PLoS ONE</i> , 2021, 16, e0251660.   | 1.1 | 1         |
| 3738 | The N-terminal domain of the A12.2 subunit stimulates RNA polymerase I transcription elongation. <i>Biophysical Journal</i> , 2021, 120, 1883-1893.  | 0.2 | 17        |
| 3742 | Essential role of the endocytic site-associated protein Ecm25 in stress-induced cell elongation. <i>Cell Reports</i> , 2021, 35, 109122.   | 2.9 | 3         |
| 3743 | ABC transporter Pdr5 is required for cantharidin resistance in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2021, 553, 141-147.  | 1.0 | 3         |
| 3745 | Natural variants suppress mutations in hundreds of essential genes. <i>Molecular Systems Biology</i> , 2021, 17, e10138.   | 3.2 | 13        |
| 3746 | Pbp1, the yeast ortholog of human Ataxin-2, functions in the cell growth on non-fermentable carbon sources. <i>PLoS ONE</i> , 2021, 16, e0251456.  | 1.1 | 4         |
| 3748 | Atg32 dependent mitophagy sustains spermidine and nitric oxide required for heat stress tolerance in <i>S. cerevisiae</i> . <i>Journal of Cell Science</i> , 2021, 134, .  | 1.2 | 7         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3751 | Exploratory polarization facilitates mating partner selection in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2021, 32, 1048-1063.   | 0.9 | 12        |
| 3754 | Functional control of Eco1 through the MCM complex in sister chromatid cohesion. <i>Gene</i> , 2021, 784, 145584.  | 1.0 | 7         |
| 3755 | Roles of High Osmolarity Glycerol and Cell Wall Integrity Pathways in Cadmium Toxicity in <i>Saccharomyces cerevisiae</i> . <i>International Journal of Molecular Sciences</i> , 2021, 22, 6169.   | 1.8 | 10        |
| 3756 | Co-translational assembly and localized translation of nucleoporins in nuclear pore complex biogenesis. <i>Molecular Cell</i> , 2021, 81, 2417-2427.e5.  | 4.5 | 45        |
| 3759 | Msn2/4 transcription factors positively regulate expression of Atg39 ER-phagy receptor. <i>Scientific Reports</i> , 2021, 11, 11919.   | 1.6 | 11        |
| 3761 | Genetic Dissection of Vps13 Regulation in Yeast Using Disease Mutations from Human Orthologs. <i>International Journal of Molecular Sciences</i> , 2021, 22, 6200.   | 1.8 | 8         |
| 3763 | Ume6 Acts as a Stable Platform To Coordinate Repression and Activation of Early Meiosis-Specific Genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2021, 41, e0037820.  | 1.1 | 3         |
| 3764 | The SESAME complex regulates cell senescence through the generation of acetyl-CoA. <i>Nature Metabolism</i> , 2021, 3, 983-1000.   | 5.1 | 23        |
| 3765 | Bro1 stimulates Vps4 to promote intraluminal vesicle formation during multivesicular body biogenesis. <i>Journal of Cell Biology</i> , 2021, 220, .  | 2.3 | 10        |
| 3766 | Assembly factors chaperone ribosomal RNA folding by isolating helical junctions that are prone to misfolding. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .  | 3.3 | 20        |
| 3767 | Scaling of subcellular actin structures with cell length through decelerated growth. <i>ELife</i> , 2021, 10, .  | 2.8 | 10        |
| 3768 | Ctf3/CENP-I provides a docking site for the desumoylase Ulp2 at the kinetochore. <i>Journal of Cell Biology</i> , 2021, 220, .   | 2.3 | 4         |
| 3769 | Genetic interaction between F-box motif encoding YDR131C and retrograde signaling-related RTG1 regulates the stress response and apoptosis in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biochemical and Molecular Toxicology</i> , 2021, 35, e22864. | 1.4 | 2         |
| 3770 | The RNA-binding protein Puf5 contributes to buffering of mRNA upon chromatin-mediated changes in nascent transcription. <i>Journal of Cell Science</i> , 2021, 134, .  | 1.2 | 0         |
| 3771 | Increased mitochondrial protein import and cardiolipin remodelling upon early mtUPR. <i>PLoS Genetics</i> , 2021, 17, e1009664.  | 1.5 | 19        |
| 3773 | Cloning and characterization of glycogen branching and debranching enzymes from the parasitic protist <i>Trichomonas vaginalis</i> . <i>Biochimie</i> , 2021, 186, 59-72.  | 1.3 | 2         |
| 3774 | Phospho-regulated Bim1/EB1 interactions trigger Dam1c ring assembly at the budding yeast outer kinetochore. <i>EMBO Journal</i> , 2021, 40, e108004.   | 3.5 | 12        |
| 3776 | Heh2/Man1 may be an evolutionarily conserved sensor of NPC assembly state. <i>Molecular Biology of the Cell</i> , 2021, 32, 1359-1373.   | 0.9 | 10        |



| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3777 | Genetic analysis argues for a coactivator function for the <i>Saccharomyces cerevisiae</i> Tup1 corepressor. <i>Genetics</i> , 2021, 219, .   | 1.2 | 3         |
| 3778 | Phospholipid flippases and Sfk1 are essential for the retention of ergosterol in the plasma membrane. <i>Molecular Biology of the Cell</i> , 2021, 32, 1374-1392.   | 0.9 | 19        |
| 3779 | Physical interactions between MCM and Rad51 facilitate replication fork lesion bypass and ssDNA gap filling by non-recombinogenic functions. <i>Cell Reports</i> , 2021, 36, 109440.                                | 2.9 | 12        |
| 3780 | MTM1 plays an important role in the regulation of zinc tolerance in <i>Saccharomyces cerevisiae</i> . <i>Journal of Trace Elements in Medicine and Biology</i> , 2021, 66, 126759.                                  | 1.5 | 10        |
| 3781 | The histone chaperone HIR maintains chromatin states to control nitrogen assimilation and fungal virulence. <i>Cell Reports</i> , 2021, 36, 109406.   | 2.9 | 10        |
| 3782 | GFP-fused yeast cells as whole-cell biosensors for genotoxicity evaluation of nitrosamines. <i>Applied Microbiology and Biotechnology</i> , 2021, 105, 5607-5616.   | 1.7 | 6         |
| 3785 | Glycerol-3-phosphate biosynthesis regenerates cytosolic NAD <sup>+</sup> to alleviate mitochondrial disease. <i>Cell Metabolism</i> , 2021, 33, 1974-1987.e9.   | 7.2 | 55        |
| 3787 | Pex30-like proteins function as adaptors at distinct ER membrane contact sites. <i>Journal of Cell Biology</i> , 2021, 220, .   | 2.3 | 18        |
| 3788 | Cell adaptation to aneuploidy by the environmental stress response dampens induction of the cytosolic unfolded-protein response. <i>Molecular Biology of the Cell</i> , 2021, 32, 1557-1564.                        | 0.9 | 9         |
| 3789 | Modular and Molecular Optimization of a LOV (Light-Oxygen-Voltage)-Based Optogenetic Switch in Yeast. <i>International Journal of Molecular Sciences</i> , 2021, 22, 8538.  | 1.8 | 8         |
| 3790 | Simple-to-use CRISPR-SpCas9/SaCas9/AsCas12a vector series for genome editing in <i>Saccharomyces cerevisiae</i> . <i>G3: Genes, Genomes, Genetics</i> , 2021, 11, .   | 0.8 | 6         |
| 3792 | Target of Rapamycin Complex 1 (TORC1), Protein Kinase A (PKA) and Cytosolic pH Regulate a Transcriptional Circuit for Lipid Droplet Formation. <i>International Journal of Molecular Sciences</i> , 2021, 22, 9017. | 1.8 | 9         |
| 3793 | Role of Calcium/Calcineurin Signalling in Regulating Intracellular Reactive Oxygen Species Homeostasis in <i>Saccharomyces cerevisiae</i> . <i>Genes</i> , 2021, 12, 1311.  | 1.0 | 4         |
| 3795 | Puf6 primes 60S pre-ribosome nuclear export at low temperature. <i>Nature Communications</i> , 2021, 12, 4696.  | 5.8 | 8         |
| 3796 | Vacuolar H <sup>+</sup> -ATPase dysfunction rescues intraluminal vesicle cargo sorting in yeast lacking PI(3,5)P2 or Doa4. <i>Journal of Cell Science</i> , 2021, 134, .  | 1.2 | 7         |
| 3797 | Translational autoregulation of the <i>S. cerevisiae</i> high-affinity polyamine transporter Hol1. <i>Molecular Cell</i> , 2021, 81, 3904-3918.e6.  | 4.5 | 6         |
| 3798 | Spatiotemporal control of pathway sensors and cross-pathway feedback regulate a differentiation MAPK pathway in yeast. <i>Journal of Cell Science</i> , 2021, 134, .  | 1.2 | 6         |
| 3799 | Pumilio2 Promotes Growth of Mature Neurons. <i>International Journal of Molecular Sciences</i> , 2021, 22, 8998.  | 1.8 | 8         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3800 | Not4 and Not5 modulate translation elongation by Rps7A ubiquitination, Rli1 moonlighting, and condensates that exclude eIF5A. <i>Cell Reports</i> , 2021, 36, 109633.                                   | 2.9 | 20        |
| 3801 | Proteome plasticity in response to persistent environmental change. <i>Molecular Cell</i> , 2021, 81, 3294-3309.e12.  | 4.5 | 12        |
| 3802 | Nhp2 is a reader of H2A <sup>Q105</sup> me and part of a network integrating metabolism with rRNA synthesis. <i>EMBO Reports</i> , 2021, 22, e52435.  | 2.0 | 5         |
| 3803 | A conserved ATG2 binding site in WIPI4 and yeast Hsv2 is disrupted by mutations causing $\hat{2}$ -propeller protein-associated neurodegeneration. <i>Human Molecular Genetics</i> , 2021, 31, 111-121. | 1.4 | 11        |
| 3804 | The Hob proteins are novel and conserved lipid-binding proteins at ER-PM contact sites. <i>Journal of Cell Science</i> , 2022, 135, .   | 1.2 | 19        |
| 3805 | <i>IRC3</i> Regulates Mitochondrial Translation in Response to Metabolic Cues in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2021, 41, e0023321.                          | 1.1 | 2         |
| 3806 | Connection of core and tail Mediator modules restrains transcription from TFIIID-dependent promoters. <i>PLoS Genetics</i> , 2021, 17, e1009529.  | 1.5 | 9         |
| 3808 | Global analysis of <i>Saccharomyces cerevisiae</i> growth in mucin. <i>G3: Genes, Genomes, Genetics</i> , 2021, 11, .   | 0.8 | 1         |
| 3810 | Repeated strand invasion and extensive branch migration are hallmarks of meiotic recombination. <i>Molecular Cell</i> , 2021, 81, 4258-4270.e4.   | 4.5 | 32        |
| 3811 | Investigation of RNA metabolism through large-scale genetic interaction profiling in yeast. <i>Nucleic Acids Research</i> , 2021, 49, 8535-8555.  | 6.5 | 4         |
| 3812 | Regulation of sphingolipid synthesis by the G1/S transcription factor Swi4. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2021, 1866, 158983.                           | 1.2 | 3         |
| 3813 | The budding yeast protein Chl1p is required for delaying progression through G1/S phase after DNA damage. <i>Cell Division</i> , 2021, 16, 4.   | 1.1 | 2         |
| 3814 | The Cdc14 Phosphatase Controls Resolution of Recombination Intermediates and Crossover Formation during Meiosis. <i>International Journal of Molecular Sciences</i> , 2021, 22, 9811.                   | 1.8 | 7         |
| 3815 | Spt4 facilitates the movement of RNA polymerase II through the +2 nucleosomal barrier. <i>Cell Reports</i> , 2021, 36, 109755.  | 2.9 | 11        |
| 3816 | Disruption of origin chromatin structure by helicase activation in the absence of DNA replication. <i>Genes and Development</i> , 2021, 35, 1339-1355.  | 2.7 | 4         |
| 3818 | Pan2-Pan3 complex, together with Ccr4-Not complex, has a role in the cell growth on non-fermentable carbon sources. <i>Biochemical and Biophysical Research Communications</i> , 2021, 570, 125-130.    | 1.0 | 1         |
| 3819 | Docking to a Basic Helix Promotes Specific Phosphorylation by G1-Cdk1. <i>International Journal of Molecular Sciences</i> , 2021, 22, 9514.   | 1.8 | 2         |
| 3820 | Cooperative regulation of endocytic vesicle transport by yeast Eps15-like protein Pan1p and epsins. <i>Journal of Biological Chemistry</i> , 2021, 297, 101254.   | 1.6 | 2         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3821 | Regulation of sphingolipid biosynthesis in the endoplasmic reticulum via signals from the plasma membrane in budding yeast. <i>FEBS Journal</i> , 2022, 289, 457-472.   | 2.2 | 8         |
| 3822 | Multiplexed profiling facilitates robust m6A quantification at site, gene and sample resolution. <i>Nature Methods</i> , 2021, 18, 1060-1067.   | 9.0 | 57        |
| 3823 | A zygote-based assay to evaluate intranuclear shuttling in <i>S. cerevisiae</i> . <i>STAR Protocols</i> , 2021, 2, 100736.  | 0.5 | 0         |
| 3824 | Measuring in vivo protein turnover and exchange in yeast macromolecular assemblies. <i>STAR Protocols</i> , 2021, 2, 100800.  | 0.5 | 3         |
| 3825 | Imaging the Actin Cytoskeleton in Live Budding Yeast Cells. <i>Methods in Molecular Biology</i> , 2022, 2364, 53-80.  | 0.4 | 0         |
| 3827 | A CRISPR/Cas9-based method for seamless N-terminal protein tagging in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2021, 38, 592-600.   | 0.8 | 2         |
| 3828 | The histone chaperone Spt6 is required for normal recruitment of the capping enzyme Abd1 to transcribed regions. <i>Journal of Biological Chemistry</i> , 2021, 297, 101205.  | 1.6 | 2         |
| 3829 | Cotranslational Targeting and Posttranslational Translocation can Cooperate in Spc3 Topogenesis. <i>Journal of Molecular Biology</i> , 2021, 433, 167109.   | 2.0 | 3         |
| 3831 | Substrate ubiquitination retains misfolded membrane proteins in the endoplasmic reticulum for degradation. <i>Cell Reports</i> , 2021, 36, 109717.  | 2.9 | 9         |
| 3832 | Analysis of local protein accumulation kinetics by live-cell imaging in yeast systems. <i>STAR Protocols</i> , 2021, 2, 100733.   | 0.5 | 8         |
| 3833 | Defining Functions of Mannoproteins in <i>Saccharomyces cerevisiae</i> by High-Dimensional Morphological Phenotyping. <i>Journal of Fungi (Basel, Switzerland)</i> , 2021, 7, 769.  | 1.5 | 6         |
| 3834 | Compound heterozygous variants in <i>SHQ1</i> are associated with a spectrum of neurological features, including early-onset dystonia. <i>Human Molecular Genetics</i> , 2022, 31, 614-624.                               | 1.4 | 12        |
| 3835 | Shortening of membrane lipid acyl chains compensates for phosphatidylcholine deficiency in choline auxotroph yeast. <i>EMBO Journal</i> , 2021, 40, e107966.  | 3.5 | 12        |
| 3837 | Quantitative proteomics identifies the universally conserved ATPase Ola1p as a positive regulator of heat shock response in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2021, 297, 101050. | 1.6 | 6         |
| 3838 | The N-Terminal Tail of Histone H3 Regulates Copper Homeostasis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2021, 41, .   | 1.1 | 5         |
| 3839 | Haspin Modulates the G2/M Transition Delay in Response to Polarization Failures in Budding Yeast. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 625717.   | 1.8 | 5         |
| 3840 | Defining steps in RAVE-catalyzed V-ATPase assembly using purified RAVE and V-ATPase subcomplexes. <i>Journal of Biological Chemistry</i> , 2021, 296, 100703.   | 1.6 | 14        |
| 3841 | SUMO is a pervasive regulator of meiosis. <i>ELife</i> , 2021, 10, .  | 2.8 | 50        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3842 | Tos4 mediates gene expression homeostasis through interaction with HDAC complexes independently of H3K56 acetylation. <i>Journal of Biological Chemistry</i> , 2021, 296, 100533.  | 1.6 | 4         |
| 3843 | Transcription levels of a noncoding RNA orchestrate opposing regulatory and cell fate outcomes in yeast. <i>Cell Reports</i> , 2021, 34, 108643.   | 2.9 | 11        |
| 3844 | Loss of Smi1, a protein involved in cell wall synthesis, extends replicative life span by enhancing rDNA stability in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2021, 296, 100258.          | 1.6 | 4         |
| 3845 | Construction of Tight Conditional Mutants Using the Improved Auxin-Inducible Degron (iAID) Method in the Budding Yeast <i>Saccharomyces cerevisiae</i> . <i>Methods in Molecular Biology</i> , 2021, 2196, 15-26.            | 0.4 | 2         |
| 3846 | Fluorescence Staining of Mitochondria for Morphology Analysis in <i>Saccharomyces cerevisiae</i> . <i>Methods in Molecular Biology</i> , 2014, 1163, 131-152.  | 0.4 | 4         |
| 3847 | PCR-Mediated Epitope Tagging of Genes in Yeast. <i>Methods in Molecular Biology</i> , 2014, 1205, 37-44.   | 0.4 | 2         |
| 3848 | Manipulating the Yeast Genome: Deletion, Mutation, and Tagging by PCR. <i>Methods in Molecular Biology</i> , 2014, 1205, 45-78.  | 0.4 | 69        |
| 3849 | Live-Cell Imaging of Mitochondria and the Actin Cytoskeleton in Budding Yeast. <i>Methods in Molecular Biology</i> , 2016, 1365, 25-62.  | 0.4 | 7         |
| 3850 | Real-Time Visualization and Quantification of Contractile Ring Proteins in Single Living Cells. <i>Methods in Molecular Biology</i> , 2016, 1369, 9-23.  | 0.4 | 8         |
| 3851 | A Review of Fluorescent Proteins for Use in Yeast. <i>Methods in Molecular Biology</i> , 2016, 1369, 309-346.  | 0.4 | 11        |
| 3852 | Analysis of Autophagosome Membrane Cycling by Fluorescence Microscopy. <i>Methods in Molecular Biology</i> , 2008, 445, 135-145.   | 0.4 | 3         |
| 3853 | In Vivo Detection and Characterization of Sumoylation Targets in <i>Saccharomyces cerevisiae</i> . <i>Methods in Molecular Biology</i> , 2009, 497, 81-103.  | 0.4 | 47        |
| 3854 | Live-Cell Imaging of the Cytoskeleton and Mitochondrial Cytoskeletal Interactions in Budding Yeast. <i>Methods in Molecular Biology</i> , 2009, 586, 41-68.  | 0.4 | 4         |
| 3855 | Casein kinase II phosphorylates the C-terminal region of Lif1 to promote the Lif1-Xrs2 interaction needed for non-homologous end joining. <i>Biochemical and Biophysical Research Communications</i> , 2018, 501, 1080-1084. | 1.0 | 2         |
| 3856 | The Yeast Ccr4-Not Complex Controls Ubiquitination of the Nascent-associated Polypeptide (NAC-EGD) Complex. <i>Journal of Biological Chemistry</i> , 2006, 281, 31389-31398.   | 1.6 | 36        |
| 3857 | Caffeine Targets TOR Complex I and Provides Evidence for a Regulatory Link between the FRB and Kinase Domains of Tor1p. <i>Journal of Biological Chemistry</i> , 2006, 281, 31616-31626.                                     | 1.6 | 57        |
| 3858 | Downregulation of autophagy by Met30-mediated Atg9 ubiquitination. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .   | 3.3 | 10        |
| 3859 | HIV-1 Gag release from yeast reveals ESCRT interaction with the Gag N-terminal protein region. <i>Journal of Biological Chemistry</i> , 2020, 295, 17950-17972.  | 1.6 | 5         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3860 | Uncoupling the hydrolysis of lipid-linked oligosaccharide from the oligosaccharyl transfer reaction by point mutations in yeast oligosaccharyltransferase. <i>Journal of Biological Chemistry</i> , 2020, 295, 16072-16085. | 1.6 | 6         |
| 3861 | Differential requirement for Bub1 and Bub3 in regulation of meiotic versus mitotic chromosome segregation. <i>Journal of Cell Biology</i> , 2020, 219, .  | 2.3 | 14        |
| 3862 | Calcineurin-dependent regulation of endocytosis by a plasma membrane ubiquitin ligase adaptor, Rcr1. <i>Journal of Cell Biology</i> , 2020, 219, .  | 2.3 | 9         |
| 3863 | Cdc42 GTPase regulates ESCRTs in nuclear envelope sealing and ER remodeling. <i>Journal of Cell Biology</i> , 2020, 219, .  | 2.3 | 17        |
| 3864 | Seipin and Nem1 establish discrete ER subdomains to initiate yeast lipid droplet biogenesis. <i>Journal of Cell Biology</i> , 2020, 219, .  | 2.3 | 68        |
| 3865 | The mitotic exit network regulates the spatiotemporal activity of Cdc42 to maintain cell size. <i>Journal of Cell Biology</i> , 2021, 220, .  | 2.3 | 6         |
| 3866 | Spatial regulation of clathrin-mediated endocytosis through position-dependent site maturation. <i>Journal of Cell Biology</i> , 2020, 219, .   | 2.3 | 18        |
| 3867 | MICOS subcomplexes assemble independently on the mitochondrial inner membrane in proximity to ER contact sites. <i>Journal of Cell Biology</i> , 2020, 219, .   | 2.3 | 27        |
| 3868 | The EB1â€™Kinesin-14 complex is required for efficient metaphase spindle assembly and kinetochore bi-orientation. <i>Journal of Cell Biology</i> , 2020, 219, .   | 2.3 | 12        |
| 3869 | Precisely calibrated and spatially informed illumination for conventional fluorescence and improved PALM imaging applications. <i>Methods and Applications in Fluorescence</i> , 2020, 8, 025004.                           | 1.1 | 6         |
| 3870 | Roles for the <i>Saccharomyces cerevisiae</i> SDS3, CBK1 and HYM1 Genes in Transcriptional Repression by SIN3. <i>Genetics</i> , 2000, 154, 573-586.  | 1.2 | 41        |
| 3871 | Bipartite Structure of the SGS1 DNA Helicase in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2000, 154, 1101-1114.   | 1.2 | 135       |
| 3872 | Genetic Analysis Reveals That <i>FLO11</i> Upregulation and Cell Polarization Independently Regulate Invasive Growth in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2000, 156, 1005-1023.                           | 1.2 | 88        |
| 3873 | Requirement for Three Novel Protein Complexes in the Absence of the Sgs1 DNA Helicase in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2001, 157, 103-118.  | 1.2 | 377       |
| 3874 | The <i>Saccharomyces cerevisiae</i> Suppressor of Choline Sensitivity ( <i>SCS2</i> ) Gene Is a Multicopy Suppressor of <i>mec1</i> Telomeric Silencing Defects. <i>Genetics</i> , 2001, 158, 145-154.                      | 1.2 | 20        |
| 3875 | Interaction of the Repressors Nrg1 and Nrg2 With the Snf1 Protein Kinase in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2001, 158, 563-572.   | 1.2 | 60        |
| 3876 | Genes Involved in Sister Chromatid Separation and Segregation in the Budding Yeast <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2001, 159, 453-470.  | 1.2 | 133       |
| 3877 | The Kar3-Interacting Protein Cik1p Plays a Critical Role in Passage Through Meiosis I in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2001, 159, 939-951.  | 1.2 | 18        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 3878 | The Spo12 Protein of <i>Saccharomyces cerevisiae</i> : A Regulator of Mitotic Exit Whose Cell Cycle-Dependent Degradation Is Mediated by the Anaphase-Promoting Complex. <i>Genetics</i> , 2001, 159, 965-980.                                | 1.2 | 28        |
| 3879 | Phosphate Transport and Sensing in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2001, 159, 1491-1499.  | 1.2 | 205       |
| 3880 | A Role for <i>MMS4</i> in the Processing of Recombination Intermediates During Meiosis in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2001, 159, 1511-1525.   | 1.2 | 101       |
| 3881 | The Novel Adaptor Protein, Mti1p, and Vrp1p, a Homolog of Wiskott-Aldrich Syndrome Protein-Interacting Protein (WIP), May Antagonistically Regulate Type I Myosins in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2002, 160, 923-934. | 1.2 | 49        |
| 3882 | Ady3p Links Spindle Pole Body Function to Spore Wall Synthesis in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2002, 160, 1439-1450.   | 1.2 | 39        |
| 3883 | IBD2 Encodes a Novel Component of the Bub2p-Dependent Spindle Checkpoint in the Budding Yeast <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2002, 161, 595-609.   | 1.2 | 7         |
| 3884 | The <i>Saccharomyces cerevisiae</i> RNase Mitochondrial RNA Processing Is Critical for Cell Cycle Progression at the End of Mitosis. <i>Genetics</i> , 2002, 161, 1029-1042.  | 1.2 | 66        |
| 3885 | Loss of Ypk1 Function Causes Rapamycin Sensitivity, Inhibition of Translation Initiation and Synthetic Lethality in 14-3-3-Deficient Yeast. <i>Genetics</i> , 2002, 161, 1453-1464.   | 1.2 | 38        |
| 3886 | Alleles of the Yeast PMS1 Mismatch-Repair Gene That Differentially Affect Recombination- and Replication-Related Processes. <i>Genetics</i> , 2002, 162, 1131-1145.   | 1.2 | 28        |
| 3887 | Maintenance of Mitochondrial Morphology Is Linked to Maintenance of the Mitochondrial Genome in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2002, 162, 1147-1156.   | 1.2 | 50        |
| 3888 | Genetic and Biochemical Basis for Viability of Yeast Lacking Mitochondrial Genomes. <i>Genetics</i> , 2002, 162, 1595-1604.   | 1.2 | 44        |
| 3889 | Loss of CDC5 Function in <i>Saccharomyces cerevisiae</i> Leads to Defects in Swe1p Regulation and Bfa1p/Bub2p-Independent Cytokinesis. <i>Genetics</i> , 2003, 163, 21-33.  | 1.2 | 38        |
| 3890 | The Mus81/Mms4 Endonuclease Acts Independently of Double-Holliday Junction Resolution to Promote a Distinct Subset of Crossovers During Meiosis in Budding Yeast. <i>Genetics</i> , 2003, 164, 81-94.   | 1.2 | 346       |
| 3891 | Opposite Roles of the F-Box Protein Rcy1p and the GTPase-Activating Protein Gyp2p During Recycling of Internalized Proteins in Yeast. <i>Genetics</i> , 2003, 164, 469-477.   | 1.2 | 16        |
| 3892 | The Role of Cdh1p in Maintaining Genomic Stability in Budding Yeast. <i>Genetics</i> , 2003, 165, 489-503.  | 1.2 | 28        |
| 3893 | A Novel Step in $\beta$ -Tubulin Folding Is Important for Heterodimer Formation in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2003, 165, 531-541.  | 1.2 | 39        |
| 3894 | Identification of Edc3p as an Enhancer of mRNA Decapping in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2004, 166, 729-739.   | 1.2 | 35        |
| 3895 | The External Amino Acid Signaling Pathway Promotes Activation of Stp1 and Uga35/Dal81 Transcription Factors for Induction of the <i>AGP1</i> Gene in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2004, 166, 1727-1739.                | 1.2 | 20        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3896 | Protein quality control degron-containing substrates are differentially targeted in the cytoplasm and nucleus by ubiquitin ligases. <i>Genetics</i> , 2021, 217, 1-19.   | 1.2 | 14        |
| 3897 | Characterization of long G4-rich enhancer-associated genomic regions engaging in a novel loop:loop $\hat{a}$ C4 Kissing $\hat{a}$ ™ interaction. <i>Nucleic Acids Research</i> , 2020, 48, 5907-5925.  | 6.5 | 17        |
| 3942 | Nuclear export of Far1p in response to pheromones requires the export receptor Msn5p/Ste21p. <i>Genes and Development</i> , 1999, 13, 2284-2300.   | 2.7 | 90        |
| 3943 | The yeast Sgs1p helicase acts upstream of Rad53p in the DNA replication checkpoint and colocalizes with Rad53p in S-phase-specific foci. <i>Genes and Development</i> , 2000, 14, 81-96.   | 2.7 | 229       |
| 3944 | The Something About Silencing protein, Sas3, is the catalytic subunit of NuA3, a yTAF<sub>II</sub>30-containing HAT complex that interacts with the Spt16 subunit of the yeast CP (Cdc68/Pob3) $\hat{a}$ FACT complex. <i>Genes and Development</i> , 2000, 14, 1196-1208. | 2.7 | 194       |
| 3945 | Deletion of MUD2, the yeast homolog of U2AF65, can bypass the requirement for Sub2, an essential spliceosomal ATPase. <i>Genes and Development</i> , 2001, 15, 42-49.  | 2.7 | 171       |
| 3946 | Two checkpoint complexes are independently recruited to sites of DNA damage in vivo. <i>Genes and Development</i> , 2001, 15, 2809-2821.   | 2.7 | 333       |
| 3947 | PCR-Mediated Gene Disruption: One-Step Method. <i>Cold Spring Harbor Protocols</i> , 2006, 2006, pdb.prot4169.   | 0.2 | 4         |
| 3948 | Follicular lymphoma $\hat{a}$ associated mutations in vacuolar ATPase ATP6V1B2 activate autophagic flux and mTOR. <i>Journal of Clinical Investigation</i> , 2019, 129, 1626-1640.   | 3.9 | 23        |
| 3949 | Dynamic, Rho1p-dependent localization of Pkc1p to sites of polarized growth. <i>Journal of Cell Science</i> , 2000, 113, 2685-2693.  | 1.2 | 63        |
| 3950 | A novel EH domain protein of <i>Saccharomyces cerevisiae</i> , Ede1p, involved in endocytosis. <i>Journal of Cell Science</i> , 2000, 113, 3309-3319.  | 1.2 | 77        |
| 3951 | The budding yeast Dbf2 protein kinase localises to the centrosome and moves to the bud neck in late mitosis. <i>Journal of Cell Science</i> , 2000, 113, 3399-3408.  | 1.2 | 99        |
| 3952 | Regulation of cytokinesis by the Elm1 protein kinase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 2000, 113, 1435-1445.   | 1.2 | 97        |
| 3953 | The Bub2-dependent mitotic pathway in yeast acts every cell cycle and regulates cytokinesis. <i>Journal of Cell Science</i> , 2001, 114, 2345-2354.  | 1.2 | 55        |
| 3954 | The life cycle of actin patches in mating yeast. <i>Journal of Cell Science</i> , 2001, 114, 1505-1513.  | 1.2 | 67        |
| 3955 | Rho5p downregulates the yeast cell integrity pathway. <i>Journal of Cell Science</i> , 2002, 115, 3139-3148.   | 1.2 | 43        |
| 3956 | In budding yeast, contraction of the actomyosin ring and formation of the primary septum at cytokinesis depend on each other. <i>Journal of Cell Science</i> , 2002, 115, 293-302.   | 1.2 | 166       |
| 3957 | Sla1p couples the yeast endocytic machinery to proteins regulating actin dynamics. <i>Journal of Cell Science</i> , 2002, 115, 1703-1715.  | 1.2 | 104       |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3958 | Robust G1 checkpoint arrest in budding yeast: dependence on DNA damage signaling and repair. <i>Journal of Cell Science</i> , 2002, 115, 1749-1757.                            | 1.2 | 47        |
| 3959 | An open interface in the pre-80S ribosome coordinated by ribosome assembly factors Tsr1 and Dim1 enables temporal regulation of Fap7. <i>Rna</i> , 2021, 27, 221-233.          | 1.6 | 20        |
| 3960 | Cdc14 phosphatase directs centrosome re-duplication at the meiosis I to meiosis II transition in budding yeast. <i>Wellcome Open Research</i> , 0, 2, 2.                       | 0.9 | 9         |
| 3961 | Mapping targets for small nucleolar RNAs in yeast. <i>Wellcome Open Research</i> , 2018, 3, 120.   | 0.9 | 18        |
| 3962 | Spo13 prevents premature cohesin cleavage during meiosis. <i>Wellcome Open Research</i> , 2019, 4, 29.   | 0.9 | 9         |
| 3963 | Spo13 prevents premature cohesin cleavage during meiosis. <i>Wellcome Open Research</i> , 2019, 4, 29.   | 0.9 | 14        |
| 3964 | A Protein Complex Containing the Conserved Swi2/Snf2-Related ATPase Swr1p Deposits Histone Variant H2A.Z into Euchromatin. <i>PLoS Biology</i> , 2004, 2, e131.                | 2.6 | 512       |
| 3965 | Phosphorylation of the Synaptonemal Complex Protein Zip1 Regulates the Crossover/Noncrossover Decision during Yeast Meiosis. <i>PLoS Biology</i> , 2015, 13, e1002329.         | 2.6 | 43        |
| 3966 | The Yeast GSK-3 Homologue Mck1 Is a Key Controller of Quiescence Entry and Chronological Lifespan. <i>PLoS Genetics</i> , 2015, 11, e1005282.                                  | 1.5 | 18        |
| 3967 | Identification of a Novel Regulatory Mechanism of Nutrient Transport Controlled by TORC1-Npr1-Amu1/Par32. <i>PLoS Genetics</i> , 2015, 11, e1005382.                           | 1.5 | 30        |
| 3968 | Large-Scale Analysis of Kinase Signaling in Yeast Pseudohyphal Development Identifies Regulation of Ribonucleoprotein Granules. <i>PLoS Genetics</i> , 2015, 11, e1005564.     | 1.5 | 24        |
| 3969 | The Dedicated Chaperone Acl4 Escorts Ribosomal Protein Rpl4 to Its Nuclear Pre-60S Assembly Site. <i>PLoS Genetics</i> , 2015, 11, e1005565.                                   | 1.5 | 59        |
| 3970 | Stable Pseudohyphal Growth in Budding Yeast Induced by Synergism between Septin Defects and Altered MAP-kinase Signaling. <i>PLoS Genetics</i> , 2015, 11, e1005684.           | 1.5 | 19        |
| 3971 | Regulation of Budding Yeast CENP-A levels Prevents Misincorporation at Promoter Nucleosomes and Transcriptional Defects. <i>PLoS Genetics</i> , 2016, 12, e1005930.            | 1.5 | 39        |
| 3972 | Nutrient Control of Yeast Gametogenesis Is Mediated by TORC1, PKA and Energy Availability. <i>PLoS Genetics</i> , 2016, 12, e1006075.  | 1.5 | 36        |
| 3973 | Mek1 Down Regulates Rad51 Activity during Yeast Meiosis by Phosphorylation of Hed1. <i>PLoS Genetics</i> , 2016, 12, e1006226.   | 1.5 | 76        |
| 3974 | Chronological Lifespan in Yeast Is Dependent on the Accumulation of Storage Carbohydrates Mediated by Yak1, Mck1 and Rim15 Kinases. <i>PLoS Genetics</i> , 2016, 12, e1006458. | 1.5 | 51        |
| 3975 | Cleavage of the SUN-domain protein Mps3 at its N-terminus regulates centrosome disjunction in budding yeast meiosis. <i>PLoS Genetics</i> , 2017, 13, e1006830.                | 1.5 | 22        |



| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3976 | The PHD finger protein Spp1 has distinct functions in the Set1 and the meiotic DSB formation complexes. <i>PLoS Genetics</i> , 2018, 14, e1007223.   | 1.5 | 41        |
| 3977 | Phenomic screen identifies a role for the yeast lysine acetyltransferase NuA4 in the control of Bcy1 subcellular localization, glycogen biosynthesis, and mitochondrial morphology. <i>PLoS Genetics</i> , 2020, 16, e1009220. | 1.5 | 5         |
| 3978 | Ybp2 Associates with the Central Kinetochore of <i>Saccharomyces cerevisiae</i> and Mediates Proper Mitotic Progression. <i>PLoS ONE</i> , 2008, 3, e1617.   | 1.1 | 13        |
| 3979 | Interaction of the Deubiquitinating Enzyme Ubp2 and the E3 Ligase Rsp5 Is Required for Transporter/Receptor Sorting in the Multivesicular Body Pathway. <i>PLoS ONE</i> , 2009, 4, e4259.                                      | 1.1 | 24        |
| 3980 | Analysis of SEC9 Suppression Reveals a Relationship of SNARE Function to Cell Physiology. <i>PLoS ONE</i> , 2009, 4, e5449.  | 1.1 | 7         |
| 3981 | rRNA Maturation in Yeast Cells Depleted of Large Ribosomal Subunit Proteins. <i>PLoS ONE</i> , 2009, 4, e8249.   | 1.1 | 104       |
| 3982 | Extra N-Terminal Residues Have a Profound Effect on the Aggregation Properties of the Potential Yeast Prion Protein Mca1. <i>PLoS ONE</i> , 2010, 5, e9929.  | 1.1 | 15        |
| 3983 | Localization to, and Effects of Pbp1, Pbp4, Lsm12, Dhh1, and Pab1 on Stress Granules in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2010, 5, e10006.   | 1.1 | 107       |
| 3984 | The Yeast Cell Fusion Protein Prm1p Requires Covalent Dimerization to Promote Membrane Fusion. <i>PLoS ONE</i> , 2010, 5, e10593.  | 1.1 | 9         |
| 3985 | The SWR1 Histone Replacement Complex Causes Genetic Instability and Genome-Wide Transcription Misregulation in the Absence of H2A.Z. <i>PLoS ONE</i> , 2010, 5, e12143.  | 1.1 | 103       |
| 3986 | The Unfolded Protein Response Is Not Necessary for the G1/S Transition, but It Is Required for Chromosome Maintenance in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2010, 5, e12732.                                  | 1.1 | 25        |
| 3987 | <i>Saccharomyces cerevisiae</i> Bat1 and Bat2 Aminotransferases Have Functionally Diverged from the Ancestral-Like <i>Kluyveromyces lactis</i> Orthologous Enzyme. <i>PLoS ONE</i> , 2011, 6, e16099.                          | 1.1 | 62        |
| 3988 | Cdc48 and Cofactors Npl4-Ufd1 Are Important for G1 Progression during Heat Stress by Maintaining Cell Wall Integrity in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2011, 6, e18988.                                   | 1.1 | 13        |
| 3989 | SWI/SNF and Asf1 Independently Promote Derepression of the DNA Damage Response Genes under Conditions of Replication Stress. <i>PLoS ONE</i> , 2011, 6, e21633.  | 1.1 | 7         |
| 3990 | tRNA Gene Identity Affects Nuclear Positioning. <i>PLoS ONE</i> , 2011, 6, e29267.   | 1.1 | 11        |
| 3991 | Yeast Transcription Termination Factor Rtt103 Functions in DNA Damage Response. <i>PLoS ONE</i> , 2012, 7, e31288.   | 1.1 | 17        |
| 3992 | Yeast IME2 Functions Early in Meiosis Upstream of Cell Cycle-Regulated SBF and MBF Targets. <i>PLoS ONE</i> , 2012, 7, e31575.   | 1.1 | 9         |
| 3993 | Novel E3 Ubiquitin Ligases That Regulate Histone Protein Levels in the Budding Yeast <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2012, 7, e36295.  | 1.1 | 52        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 3994 | Med5(Nut1) and Med17(Srb4) Are Direct Targets of Mediator Histone H4 Tail Interactions. <i>PLoS ONE</i> , 2012, 7, e38416.   | 1.1 | 27        |
| 3995 | Rsp5 Ubiquitin Ligase Is Required for Protein Trafficking in <i>Saccharomyces cerevisiae</i> COPI Mutants. <i>PLoS ONE</i> , 2012, 7, e39582.  | 1.1 | 18        |
| 3996 | Characterization of Rad51 from Apicomplexan Parasite <i>Toxoplasma gondii</i> : An Implication for Inefficient Gene Targeting. <i>PLoS ONE</i> , 2012, 7, e41925.  | 1.1 | 11        |
| 3997 | Evolutionary Conservation of the Ribosomal Biogenesis Factor Rbm19/Mrd1: Implications for Function. <i>PLoS ONE</i> , 2012, 7, e43786.   | 1.1 | 10        |
| 3998 | Sorting Signals That Mediate Traffic of Chitin Synthase III between the TGN/Endosomes and to the Plasma Membrane in Yeast. <i>PLoS ONE</i> , 2012, 7, e46386.  | 1.1 | 34        |
| 3999 | The Fate of Linear DNA in <i>Saccharomyces cerevisiae</i> and <i>Candida glabrata</i> : The Role of Homologous and Non-Homologous End Joining. <i>PLoS ONE</i> , 2013, 8, e69628.  | 1.1 | 20        |
| 4000 | The N-Terminus of Vps74p Is Essential for the Retention of Glycosyltransferases in the Golgi but Not for the Modulation of Apical Polarized Growth in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2013, 8, e74715. | 1.1 | 5         |
| 4001 | Effects of the Yeast RNA-Binding Protein Whi3 on the Half-Life and Abundance of CLN3 mRNA and Other Targets. <i>PLoS ONE</i> , 2013, 8, e84630.  | 1.1 | 43        |
| 4002 | The Budding Yeast Amphiphysin Complex Is Required for Contractile Actin Ring (CAR) Assembly and Post-Contraction GEF-Independent Accumulation of Rho1-GTP. <i>PLoS ONE</i> , 2014, 9, e97663.                              | 1.1 | 3         |
| 4003 | Unidirectional P-Body Transport during the Yeast Cell Cycle. <i>PLoS ONE</i> , 2014, 9, e99428.  | 1.1 | 17        |
| 4004 | PHD and TFIIIS-Like Domains of the Bye1 Transcription Factor Determine Its Multivalent Genomic Distribution. <i>PLoS ONE</i> , 2014, 9, e102464.   | 1.1 | 14        |
| 4005 | PCR Duplication: A One-Step Cloning-Free Method to Generate Duplicated Chromosomal Loci and Interference-Free Expression Reporters in Yeast. <i>PLoS ONE</i> , 2014, 9, e114590.   | 1.1 | 16        |
| 4006 | SPO73 and SPO71 Function Cooperatively in Prospore Membrane Elongation During Sporulation in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2015, 10, e0143571.   | 1.1 | 8         |
| 4007 | Characterization of Fluorescent Proteins for Three- and Four-Color Live-Cell Imaging in <i>S. cerevisiae</i> . <i>PLoS ONE</i> , 2016, 11, e0146120.   | 1.1 | 25        |
| 4008 | Recruitment of <i>Saccharomyces cerevisiae</i> Cmr1/Ydl156w to Coding Regions Promotes Transcription Genome Wide. <i>PLoS ONE</i> , 2016, 11, e0148897.  | 1.1 | 8         |
| 4009 | Kin2, the Budding Yeast Ortholog of Animal MARK/PAR-1 Kinases, Localizes to the Sites of Polarized Growth and May Regulate Septin Organization and the Cell Wall. <i>PLoS ONE</i> , 2016, 11, e0153992.                    | 1.1 | 11        |
| 4010 | Identification of Putative Mek1 Substrates during Meiosis in <i>Saccharomyces cerevisiae</i> Using Quantitative Phosphoproteomics. <i>PLoS ONE</i> , 2016, 11, e0155931.   | 1.1 | 13        |
| 4011 | Regulation of the Na <sup>+</sup> /K <sup>+</sup> -ATPase Ena1 Expression by Calcineurin/Crz1 under High pH Stress: A Quantitative Study. <i>PLoS ONE</i> , 2016, 11, e0158424.  | 1.1 | 19        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 4012 | Cytoplasmic deadenylase Ccr4 is required for translational repression of LRG1 mRNA in the stationary phase. PLoS ONE, 2017, 12, e0172476.   | 1.1 | 15        |
| 4013 | Low RNA Polymerase III activity results in up regulation of HXT2 glucose transporter independently of glucose signaling and despite changing environment. PLoS ONE, 2017, 12, e0185516. | 1.1 | 4         |
| 4014 | Chl1 DNA helicase and Scc2 function in chromosome condensation through cohesin deposition. PLoS ONE, 2017, 12, e0188739.  | 1.1 | 9         |
| 4015 | The absence of specific yeast heat-shock proteins leads to abnormal aggregation and compromised autophagic clearance of mutant Huntingtin proteins. PLoS ONE, 2018, 13, e0191490.       | 1.1 | 16        |
| 4016 | Gene cloning of a neutral ceramidase from the sphingolipid metabolic pathway based on transcriptome analysis of <i>Amorphophallus muelleri</i> . PLoS ONE, 2018, 13, e0194863.          | 1.1 | 2         |
| 4017 | Mitochondrial dysfunction reduces yeast replicative lifespan by elevating RAS-dependent ROS production by the ER-localized NADPH oxidase Yno1. PLoS ONE, 2018, 13, e0198619.            | 1.1 | 28        |
| 4018 | PCNA antagonizes cohesin-dependent roles in genomic stability. PLoS ONE, 2020, 15, e0235103.  | 1.1 | 4         |
| 4019 | Tomato heat stress protein Hsp16.1-CIII represents a member of a new class of nucleocytoplasmic small heat stress proteins in plants. Cell Stress and Chaperones, 2003, 8, 381.         | 1.2 | 33        |
| 4020 | Termination of non-coding transcription in yeast relies on both an RNA Pol II CTD interaction domain and a CTD-mimicking region in Sen1. EMBO Journal, 2020, 39, e101548.               | 3.5 | 23        |
| 4021 | A new linear cyclin docking motif that mediates exclusively S-phase CDK-specific signaling. EMBO Journal, 2021, 40, e105839.  | 3.5 | 23        |
| 4022 | Non-recombinogenic roles for Rad52 in translesion synthesis during DNA damage tolerance. EMBO Reports, 2021, 22, e50410.  | 2.0 | 15        |
| 4023 | Atg1 kinase regulates autophagosome-vacuole fusion by controlling SNARE bundling. EMBO Reports, 2020, 21, e51869.   | 2.0 | 26        |
| 4024 | The MAPKKs Ste11 and Bck1 jointly transduce the high oxidative stress signal through the cell wall integrity MAP kinase pathway. Microbial Cell, 2015, 2, 329-342.                      | 1.4 | 14        |
| 4025 | INO1 transcriptional memory leads to DNA zip code-dependent interchromosomal clustering. Microbial Cell, 2015, 2, 481-490.  | 1.4 | 30        |
| 4026 | Decreasing cytosolic translation is beneficial to yeast and human Tafazzin-deficient cells. Microbial Cell, 2018, 5, 220-232.   | 1.4 | 13        |
| 4027 | Snf1 cooperates with the CWI MAPK pathway to mediate the degradation of Med13 following oxidative stress. Microbial Cell, 2018, 5, 357-370.   | 1.4 | 19        |
| 4028 | Systematic analysis of nuclear gene function in respiratory growth and expression of the mitochondrial genome in <i>S. cerevisiae</i> . Microbial Cell, 2020, 7, 234-249.               | 1.4 | 20        |
| 4029 | Slm35 links mitochondrial stress response and longevity through TOR signaling pathway. Aging, 2016, 8, 3255-3271.   | 1.4 | 13        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 4030 | Cell cycle regulation of condensin Smc4. <i>Oncotarget</i> , 2019, 10, 263-276.   | 0.8 | 12        |
| 4031 | Unbiased yeast screens identify cellular pathways affected in Niemann-Pick disease type C. <i>Life Science Alliance</i> , 2020, 3, e201800253.  | 1.3 | 10        |
| 4032 | Ncl1-mediated metabolic rewiring critical during metabolic stress. <i>Life Science Alliance</i> , 2019, 2, e201900360.  | 1.3 | 2         |
| 4033 | Endogenous RNAi pathway evolutionarily shapes the destiny of the antisense lncRNAs transcriptome. <i>Life Science Alliance</i> , 2019, 2, e201900407.   | 1.3 | 9         |
| 4034 | A time-resolved interaction analysis of Bem1 reconstructs the flow of Cdc42 during polar growth. <i>Life Science Alliance</i> , 2020, 3, e202000813.  | 1.3 | 8         |
| 4035 | Mps2 links Csm4 and Mps3 to form a telomere-associated LINC complex in budding yeast. <i>Life Science Alliance</i> , 2020, 3, e202000824.   | 1.3 | 8         |
| 4036 | Yeast Translation Elongation Factor eIF5A Expression Is Regulated by Nutrient Availability through Different Signalling Pathways. <i>International Journal of Molecular Sciences</i> , 2021, 22, 219. | 1.8 | 10        |
| 4037 | Targeting Copper Homeostasis Improves Functioning of vps13 <sup>Δ</sup> Yeast Mutant Cells, a Model of VPS13-Related Diseases. <i>International Journal of Molecular Sciences</i> , 2021, 22, 2248.   | 1.8 | 6         |
| 4038 | Meiosis I chromosome segregation is established through regulation of microtubule-kinetochore interactions. <i>ELife</i> , 2012, 1, e00117.   | 2.8 | 85        |
| 4039 | ATR/Mec1 prevents lethal meiotic recombination initiation on partially replicated chromosomes in budding yeast. <i>ELife</i> , 2013, 2, e00844.   | 2.8 | 44        |
| 4040 | Sterol homeostasis requires regulated degradation of squalene monooxygenase by the ubiquitin ligase Doa10/Teb4. <i>ELife</i> , 2013, 2, e00953.   | 2.8 | 167       |
| 4041 | Shugoshin biases chromosomes for biorientation through condensin recruitment to the pericentromere. <i>ELife</i> , 2014, 3, e01374.   | 2.8 | 74        |
| 4042 | Aim24 and MICOS modulate respiratory function, tafazzin-related cardiolipin modification and mitochondrial architecture. <i>ELife</i> , 2014, 3, e01684.  | 2.8 | 64        |
| 4043 | Evolution of histone 2A for chromatin compaction in eukaryotes. <i>ELife</i> , 2014, 3, .   | 2.8 | 19        |
| 4044 | Lis1 regulates dynein by sterically blocking its mechanochemical cycle. <i>ELife</i> , 2014, 3, .   | 2.8 | 89        |
| 4045 | A RanGTP-independent mechanism allows ribosomal protein nuclear import for ribosome assembly. <i>ELife</i> , 2014, 3, e03473.   | 2.8 | 63        |
| 4046 | Redox signaling via the molecular chaperone BiP protects cells against endoplasmic reticulum-derived oxidative stress. <i>ELife</i> , 2014, 3, e03496.  | 2.8 | 93        |
| 4047 | Transcription mediated insulation and interference direct gene cluster expression switches. <i>ELife</i> , 2014, 3, e03635.   | 2.8 | 35        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 4048 | Heritable capture of heterochromatin dynamics in <i>Saccharomyces cerevisiae</i> . <i>ELife</i> , 2015, 4, e05007.  | 2.8 | 76        |
| 4049 | Specificity in endoplasmic reticulum-stress signaling in yeast entails a step-wise engagement of HAC1 mRNA to clusters of the stress sensor Ire1. <i>ELife</i> , 2014, 3, e05031. | 2.8 | 44        |
| 4050 | A non-canonical mechanism for Crm1-export cargo complex assembly. <i>ELife</i> , 2015, 4, .   | 2.8 | 34        |
| 4051 | Structural evidence for Scc4-dependent localization of cohesin loading. <i>ELife</i> , 2015, 4, e06057.   | 2.8 | 69        |
| 4052 | The chromatin remodelers RSC and ISW1 display functional and chromatin-based promoter antagonism. <i>ELife</i> , 2015, 4, e06073.   | 2.8 | 68        |
| 4053 | The ER Stress Surveillance (ERSU) pathway regulates daughter cell ER protein aggregate inheritance. <i>ELife</i> , 2015, 4, .   | 2.8 | 22        |
| 4054 | MICOS coordinates with respiratory complexes and lipids to establish mitochondrial inner membrane architecture. <i>ELife</i> , 2015, 4, .   | 2.8 | 212       |
| 4055 | The $\hat{I}^2$ -hairpin of 40S exit channel protein Rps5/uS7 promotes efficient and accurate translation initiation in vivo. <i>ELife</i> , 2015, 4, e07939.                     | 2.8 | 24        |
| 4056 | The vacuole/lysosome is required for cell-cycle progression. <i>ELife</i> , 2015, 4, .  | 2.8 | 49        |
| 4057 | Yeast Eps15-like endocytic protein Pan1p regulates the interaction between endocytic vesicles, endosomes and the actin cytoskeleton. <i>ELife</i> , 2016, 5, .                    | 2.8 | 16        |
| 4058 | The ribosomal protein Asc1/RACK1 is required for efficient translation of short mRNAs. <i>ELife</i> , 2016, 5, .  | 2.8 | 107       |
| 4059 | Role of competition between polarity sites in establishing a unique front. <i>ELife</i> , 2015, 4, .  | 2.8 | 56        |
| 4060 | Spatial signals link exit from mitosis to spindle position. <i>ELife</i> , 2016, 5, .   | 2.8 | 26        |
| 4061 | Constitutive turnover of histone H2A.Z at yeast promoters requires the preinitiation complex. <i>ELife</i> , 2016, 5, .   | 2.8 | 73        |
| 4062 | Heterochromatin assembly by interrupted Sir3 bridges across neighboring nucleosomes. <i>ELife</i> , 2016, 5, .  | 2.8 | 30        |
| 4063 | ATPase activity of the DEAD-box protein Dhh1 controls processing body formation. <i>ELife</i> , 2016, 5, .  | 2.8 | 115       |
| 4064 | Transcriptional rewiring over evolutionary timescales changes quantitative and qualitative properties of gene expression. <i>ELife</i> , 2016, 5, .                               | 2.8 | 54        |
| 4065 | Prefabrication of a ribosomal protein subcomplex essential for eukaryotic ribosome formation. <i>ELife</i> , 2016, 5, .   | 2.8 | 25        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 4066 | Concerted action of the MutL <sup>2</sup> heterodimer and Mer3 helicase regulates the global extent of meiotic gene conversion. <i>ELife</i> , 2017, 6, .    | 2.8 | 67        |
| 4067 | Nucleosomes influence multiple steps during replication initiation. <i>ELife</i> , 2017, 6, .  | 2.8 | 58        |
| 4068 | Gene regulatory network plasticity predates a switch in function of a conserved transcription regulator. <i>ELife</i> , 2017, 6, .                           | 2.8 | 46        |
| 4069 | Functional mapping of yeast genomes by saturated transposition. <i>ELife</i> , 2017, 6, .  | 2.8 | 126       |
| 4070 | The dynamic three-dimensional organization of the diploid yeast genome. <i>ELife</i> , 2017, 6, .  | 2.8 | 57        |
| 4071 | Cell cycle entry triggers a switch between two modes of Cdc42 activation during yeast polarization. <i>ELife</i> , 2017, 6, .                                | 2.8 | 59        |
| 4072 | Kinetochore inactivation by expression of a repressive mRNA. <i>ELife</i> , 2017, 6, .   | 2.8 | 66        |
| 4073 | Transcription of a 5' extended mRNA isoform directs dynamic chromatin changes and interference of a downstream promoter. <i>ELife</i> , 2017, 6, .           | 2.8 | 68        |
| 4074 | Aneuploidy as a cause of impaired chromatin silencing and mating-type specification in budding yeast. <i>ELife</i> , 2017, 6, .                              | 2.8 | 18        |
| 4075 | Switch-like Arp2/3 activation upon WASP and WIP recruitment to an apparent threshold level by multivalent linker proteins in vivo. <i>ELife</i> , 2017, 6, . | 2.8 | 47        |
| 4076 | Cryo-EM structure of a late pre-40S ribosomal subunit from <i>Saccharomyces cerevisiae</i> . <i>ELife</i> , 2017, 6, .                                       | 2.8 | 77        |
| 4077 | Non-invasive measurement of mRNA decay reveals translation initiation as the major determinant of mRNA stability. <i>ELife</i> , 2018, 7, .                  | 2.8 | 174       |
| 4078 | Cortical dynein pulling mechanism is regulated by differentially targeted attachment molecule Num1. <i>ELife</i> , 2018, 7, .                                | 2.8 | 30        |
| 4079 | Temporal profiling of redox-dependent heterogeneity in single cells. <i>ELife</i> , 2018, 7, .   | 2.8 | 27        |
| 4080 | An assay for de novo kinetochore assembly reveals a key role for the CENP-T pathway in budding yeast. <i>ELife</i> , 2018, 7, .                              | 2.8 | 46        |
| 4081 | Role of the pre-initiation complex in Mediator recruitment and dynamics. <i>ELife</i> , 2018, 7, .   | 2.8 | 30        |
| 4082 | Pat1 promotes processing body assembly by enhancing the phase separation of the DEAD-box ATPase Dhh1 and RNA. <i>ELife</i> , 2019, 8, .                      | 2.8 | 50        |
| 4083 | Dissociation rate compensation mechanism for budding yeast pioneer transcription factors. <i>ELife</i> , 2019, 8, .  | 2.8 | 68        |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 4084 | The proteasome biogenesis regulator Rpn4 cooperates with the unfolded protein response to promote ER stress resistance. <i>ELife</i> , 2019, 8, .   | 2.8 | 42        |
| 4086 | The structure of the Ctf19c/CCAN from budding yeast. <i>ELife</i> , 2019, 8, .  | 2.8 | 71        |
| 4087 | An ESCRT-LEM protein surveillance system is poised to directly monitor the nuclear envelope and nuclear transport system. <i>ELife</i> , 2019, 8, .   | 2.8 | 92        |
| 4088 | Dynamic relocalization of replication origins by Fkh1 requires execution of DDK function and Cdc45 loading at origins. <i>ELife</i> , 2019, 8, .  | 2.8 | 21        |
| 4089 | Metabolic constraints drive self-organization of specialized cell groups. <i>ELife</i> , 2019, 8, .   | 2.8 | 42        |
| 4090 | Meiotic cellular rejuvenation is coupled to nuclear remodeling in budding yeast. <i>ELife</i> , 2019, 8, .  | 2.8 | 51        |
| 4091 | Molecular basis for dyneinopathies reveals insight into dynein regulation and dysfunction. <i>ELife</i> , 2019, 8, .  | 2.8 | 39        |
| 4092 | The structure of the yeast Ctf3 complex. <i>ELife</i> , 2019, 8, .  | 2.8 | 15        |
| 4093 | The endonuclease Cue2 cleaves mRNAs at stalled ribosomes during No Go Decay. <i>ELife</i> , 2019, 8, .  | 2.8 | 139       |
| 4094 | Specific lid-base contacts in the 26s proteasome control the conformational switching required for substrate degradation. <i>ELife</i> , 2019, 8, .   | 2.8 | 27        |
| 4095 | A protein quality control pathway at the mitochondrial outer membrane. <i>ELife</i> , 2020, 9, .  | 2.8 | 38        |
| 4096 | The GTPase Nog1 co-ordinates the assembly, maturation and quality control of distant ribosomal functional centers. <i>ELife</i> , 2020, 9, .  | 2.8 | 36        |
| 4097 | Inhibition of post-termination ribosome recycling at premature termination codons in UPF1 ATPase mutants. <i>ELife</i> , 2020, 9, .   | 2.8 | 14        |
| 4098 | Peroxiredoxin promotes longevity and H <sub>2</sub> O <sub>2</sub> -resistance in yeast through redox-modulation of protein kinase A. <i>ELife</i> , 2020, 9, .                             | 2.8 | 20        |
| 4099 | chTOG is a conserved mitotic error correction factor. <i>ELife</i> , 2020, 9, .   | 2.8 | 27        |
| 4100 | Association and dissociation between the mitochondrial Far complex and Atg32 regulate mitophagy. <i>ELife</i> , 2020, 9, .  | 2.8 | 14        |
| 4101 | The GAR domain integrates functions that are necessary for the proper localization of fibrillarlin (FBL) inside eukaryotic cells. <i>PeerJ</i> , 2020, 8, e9029.                            | 0.9 | 9         |
| 4102 | An Inducible System for Silencing Establishment Reveals a Stepwise Mechanism in Which Anchoring at the Nuclear Periphery Precedes Heterochromatin Formation. <i>Cells</i> , 2021, 10, 2810. | 1.8 | 3         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 4104 | Cell cycle-independent integration of stress signals by Xbp1 promotes Non-G1/G0 quiescence entry. <i>Journal of Cell Biology</i> , 2022, 221, .  | 2.3 | 7         |
| 4105 | Genome-Wide Regulations of the Preinitiation Complex Formation and Elongating RNA Polymerase II by an E3 Ubiquitin Ligase, San1. <i>Molecular and Cellular Biology</i> , 2022, 42, MCB0036821. | 1.1 | 3         |
| 4106 | TORC1 signaling modulates Cdk8-dependent <i>GAL</i> gene expression in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2021, 219, .  | 1.2 | 1         |
| 4107 | Association of snR190 snoRNA chaperone with early pre-60S particles is regulated by the RNA helicase Dbp7 in yeast. <i>Nature Communications</i> , 2021, 12, 6153.                             | 5.8 | 19        |
| 4108 | Decreasing mitochondrial RNA polymerase activity reverses biased inheritance of hypersuppressive mtDNA. <i>PLoS Genetics</i> , 2021, 17, e1009808.   | 1.5 | 2         |
| 4109 | Set4 regulates stress response genes and coordinates histone deacetylases within yeast subtelomeres. <i>Life Science Alliance</i> , 2021, 4, e202101126.                                       | 1.3 | 7         |
| 4110 | Unexplored Cdc42 functions at the budding yeast nucleus suggested by subcellular localization. <i>Small GTPases</i> , 2021, , 1-12.  | 0.7 | 0         |
| 4111 | Conserved heterodimeric GTPase Rbg1/Tma46 promotes efficient translation in eukaryotic cells. <i>Cell Reports</i> , 2021, 37, 109877.  | 2.9 | 10        |
| 4114 | Epistasis, aneuploidy, and functional mutations underlie evolution of resistance to induced microtubule depolymerization. <i>EMBO Journal</i> , 2021, 40, e108225.                             | 3.5 | 11        |
| 4115 | Mechanism of lipid droplet formation by the yeast Sei1/Ldb16 Seipin complex. <i>Nature Communications</i> , 2021, 12, 5892.  | 5.8 | 40        |
| 4116 | Programmed cortical ER collapse drives selective ER degradation and inheritance in yeast meiosis. <i>Journal of Cell Biology</i> , 2021, 220, .  | 2.3 | 14        |
| 4117 | Ice2 promotes ER membrane biogenesis in yeast by inhibiting the conserved lipin phosphatase complex. <i>EMBO Journal</i> , 2021, 40, e107958.  | 3.5 | 26        |
| 4118 | Translational control of lipogenesis links protein synthesis and phosphoinositide signaling with nuclear division in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2022, 220, .          | 1.2 | 7         |
| 4119 | Sterol Composition Modulates the Response of <i>Saccharomyces cerevisiae</i> to Iron Deficiency. <i>Journal of Fungi (Basel, Switzerland)</i> , 2021, 7, 901.                                  | 1.5 | 8         |
| 4121 | Protein phosphatase 2A (PP2A) promotes anaphase entry after DNA replication stress in budding yeast. <i>Molecular Biology of the Cell</i> , 2021, 32, ar36.                                    | 0.9 | 1         |
| 4122 | Cnm1 mediates nucleus-mitochondria contact site formation in response to phospholipid levels. <i>Journal of Cell Biology</i> , 2021, 220, .  | 2.3 | 29        |
| 4123 | Roles for L <sub>o</sub> microdomains and ESCRT in ER stress-induced lipid droplet microautophagy in budding yeast. <i>Molecular Biology of the Cell</i> , 2021, 32, br12.                     | 0.9 | 20        |
| 4124 | A preferred sequence for organelle inheritance during polarized cell growth. <i>Journal of Cell Science</i> , 2021, 134, .   | 1.2 | 8         |



| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 4125 | Yeast as a model system in the analysis of DNA topoisomerase I poisons. , 2003, , 89-107.   |     | 0         |
| 4127 | Large-Scale Protein Localization in Yeast. , 2006, , 179-183.   |     | 0         |
| 4128 | Multiple Degradation Pathways for Misfolded Mutants of the Yeast Plasma Membrane ATPase, PMA1. Journal of Biological Chemistry, 2006, 281, 31457-31466.                       | 1.6 | 6         |
| 4129 | The Spindle Checkpoint: Assays for the Analysis of Spindle Checkpoint Arrest and Recovery. Methods in Molecular Biology, 2009, 545, 243-258.                                  | 0.4 | 0         |
| 4131 | Multisite Phosphorylation of the Guanine Nucleotide Exchange Factor Cdc24 during Yeast Cell Polarization. , 2014, , 79-106.   |     | 0         |
| 4134 | Paf1C mediates regulation of gene expression by the MAP Kinase Slt2. Postdoc Journal, 0, , .  | 0.4 | 0         |
| 4138 | Mapping paths: new approaches to dissect eukaryotic signaling circuitry. F1000Research, 2016, 5, 1853.  | 0.8 | 0         |
| 4159 | Hof1 and Chs4 interact via F-BAR domain and Sel1-like repeats to control extracellular matrix deposition during cytokinesis in budding yeast. SSRN Electronic Journal, 0, , . | 0.4 | 0         |
| 4160 | Quantitative Analysis of Clathrin-Mediated Endocytosis in Yeast by Live Cell Fluorescence Microscopy. Methods in Molecular Biology, 2018, 1847, 225-237.                      | 0.4 | 1         |
| 4185 | Mapping targets for small nucleolar RNAs in yeast. Wellcome Open Research, 2018, 3, 120.  | 0.9 | 7         |
| 4249 | Fluorescence Microscopy for Analysis of Relocalization of Structure-Specific Endonucleases. Methods in Molecular Biology, 2021, 2153, 521-534.                                | 0.4 | 0         |
| 4252 | Resource plasticity-driven carbon-nitrogen budgeting enables specialization and division of labor in a clonal community. ELife, 2020, 9, .                                    | 2.8 | 8         |
| 4255 | Atg39 selectively captures inner nuclear membrane into luminal vesicles for delivery to the autophagosome. Journal of Cell Biology, 2021, 220, .                              | 2.3 | 17        |
| 4256 | Growth-dependent signals drive an increase in early G1 cyclin concentration to link cell cycle entry with cell growth. ELife, 2021, 10, .                                     | 2.8 | 21        |
| 4257 | The MAP She1 coordinates directional spindle positioning by spatially restricting dynein activity. Journal of Cell Science, 2021, 134, .                                      | 1.2 | 3         |
| 4258 | Methylated PP2A stabilizes Gcn4 to enable a methionine-induced anabolic program. Journal of Biological Chemistry, 2020, 295, 18390-18405.                                     | 1.6 | 4         |
| 4270 | Cohesin regulates homology search during recombinational DNA repair. Nature Cell Biology, 2021, 23, 1176-1186.  | 4.6 | 43        |
| 4271 | The Shu complex prevents mutagenesis and cytotoxicity of single-strand specific alkylation lesions. ELife, 2021, 10, .  | 2.8 | 3         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 4285 | Polo-like kinase acts as a molecular timer that safeguards the asymmetric fate of spindle microtubule-organizing centers. <i>ELife</i> , 2020, 9, .  | 2.8 | 3         |
| 4286 | The yeast Sgs1p helicase acts upstream of Rad53p in the DNA replication checkpoint and colocalizes with Rad53p in S-phase-specific foci. <i>Genes and Development</i> , 2000, 14, 81-96.   | 2.7 | 221       |
| 4287 | The something about silencing protein, Sas3, is the catalytic subunit of NuA3, a $\gamma$ TAF(II)30-containing HAT complex that interacts with the Spt16 subunit of the yeast CP (Cdc68/Pob3)-FACT complex. <i>Genes and Development</i> , 2000, 14, 1196-208. | 2.7 | 186       |
| 4311 | Atg27p co-fractionates with clathrin-coated vesicles in budding yeast. <i>MicroPublication Biology</i> , 2021, 2021, .   | 0.1 | 0         |
| 4312 | Atg27p localization is clathrin- and Ent3p/5p-dependent. <i>MicroPublication Biology</i> , 2021, 2021, .   | 0.1 | 0         |
| 4313 | YML018C protein localizes to the vacuolar membrane independently of Atg27p. <i>MicroPublication Biology</i> , 2021, 2021, .  | 0.1 | 0         |
| 4314 | A mutation in budding yeast affecting nuclear envelope insertion of the spindle pole body. <i>MicroPublication Biology</i> , 2021, 2021, .   | 0.1 | 1         |
| 4315 | <i>VID22</i> counteracts G-quadruplex-induced genome instability. <i>Nucleic Acids Research</i> , 2021, 49, 12785-12804.   | 6.5 | 5         |
| 4316 | ER-localized phosphatidylethanolamine synthase plays a conserved role in lipid droplet formation. <i>Molecular Biology of the Cell</i> , 2022, 33, mbcE21110558T.  | 0.9 | 7         |
| 4317 | Yap5 Competes With Hap4 for the Regulation of Iron Homeostasis Genes in the Human Pathogen <i>Candida glabrata</i> . <i>Frontiers in Cellular and Infection Microbiology</i> , 2021, 11, 731988.   | 1.8 | 0         |
| 4318 | Multiple epistatic DNA variants in a single gene affect gene expression in <i>trans</i> . <i>Genetics</i> , 2022, 220, .   | 1.2 | 6         |
| 4319 | Widespread use of unconventional targeting signals in mitochondrial ribosome proteins. <i>EMBO Journal</i> , 2022, 41, e109519.  | 3.5 | 12        |
| 4321 | Telomerase subunit Est2 marks internal sites that are prone to accumulate DNA damage. <i>BMC Biology</i> , 2021, 19, 247.  | 1.7 | 4         |
| 4322 | Sir3 heterochromatin protein promotes non-homologous end joining by direct inhibition of Sae2. <i>EMBO Journal</i> , 2021, , e108813.  | 3.5 | 5         |
| 4323 | Phosphorylation-dependent mitotic SUMOylation drives nuclear envelope-chromatin interactions. <i>Journal of Cell Biology</i> , 2021, 220, .  | 2.3 | 13        |
| 4326 | Golgi membrane protein Erd1 Is essential for recycling a subset of Golgi glycosyltransferases. <i>ELife</i> , 2021, 10, .  | 2.8 | 6         |
| 4327 | An improved auxin-inducible degron system for fission yeast. <i>G3: Genes, Genomes, Genetics</i> , 2022, 12, .   | 0.8 | 16        |
| 4329 | Adaptive Role of Cell Death in Yeast Communities Stressed with Macrolide Antifungals. <i>MSphere</i> , 2021, 6, e0074521.  | 1.3 | 3         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 4330 | Immunoaffinity purification of endogenous proteins from <i>S.Âcerevisiae</i> for post-translational modification and protein interaction analysis. STAR Protocols, 2021, 2, 100945.                   | 0.5 | 0         |
| 4331 | Characterization and optimization of the <i>LAC4</i> upstream region for lowâ€leakage expression in <i>Kluyveromyces marxianus</i>. Yeast, 2022, 39, 283-296.   | 0.8 | 4         |
| 4332 | Deletion of <i>MUD2,</i> the yeast homolog of U2AF65, can bypass the requirement for Sub2, an essential spliceosomal ATPase. Genes and Development, 2001, 15, 42-49.                                  | 2.7 | 63        |
| 4333 | Modulated termination of non-coding transcription partakes in the regulation of gene expression. Nucleic Acids Research, 2022, 50, 1430-1448.   | 6.5 | 4         |
| 4334 | Induction of autophagy and endoplasmic reticulum autophagy caused by cadmium telluride quantum dots are protective mechanisms of yeast cell. Journal of Applied Toxicology, 2022, 42, 1146-1158.      | 1.4 | 6         |
| 4335 | Structural Probing with MNase Tethered to Ribosome Assembly Factors Resolves Flexible RNA Regions within the Nascent Pre-Ribosomal RNA. Non-coding RNA, 2022, 8, 1.                                   | 1.3 | 3         |
| 4336 | Systems Biology Feedback (of the Collaborative Kind). Computer, 2019, , 1-1.  | 1.2 | 0         |
| 4338 | What makes a histone variant a variant: Changing H2A to become H2A.Z. PLoS Genetics, 2021, 17, e1009950.  | 1.5 | 7         |
| 4339 | Blocking Mitophagy Does Not Significantly Improve Fuel Ethanol Production in Bioethanol Yeast <i>Saccharomyces cerevisiae</i> . Applied and Environmental Microbiology, 2022, 88, aem0206821.         | 1.4 | 5         |
| 4340 | Stepwise assembly of the eukaryotic translation initiation factor 2 complex. Journal of Biological Chemistry, 2022, 298, 101583.  | 1.6 | 6         |
| 4342 | ER-phagy requires the assembly of actin at sites of contact between the cortical ER and endocytic pits. Proceedings of the National Academy of Sciences of the United States of America, 2022, 119, . | 3.3 | 16        |
| 4343 | Structural basis for assembly of TRAPPII complex and specific activation of GTPase Ypt31/32. Science Advances, 2022, 8, eabi5603.   | 4.7 | 10        |
| 4344 | The modifying enzyme Tsr3 establishes the hierarchy of Rio kinase binding in 40S ribosome assembly. Rna, 2022, 28, 568-582.   | 1.6 | 7         |
| 4345 | Mapping Critical Residues in ATG11â€™s Coiled-Coil 2 Domain that Block Multiple Interactions and Disrupt Selective Autophagy. Frontiers in Cell and Developmental Biology, 2021, 9, 775364.           | 1.8 | 1         |
| 4346 | Glucose starvation induces a switch in the histone acetylome for activation of gluconeogenic and fat metabolism genes. Molecular Cell, 2022, 82, 60-74.e5.  | 4.5 | 43        |
| 4347 | Application of yeast spores as Î²-glucan particles. Particuology, 2022, 71, 34-40.  | 2.0 | 2         |
| 4348 | Chl1 helicase controls replication fork progression by regulating dNTP pools. Life Science Alliance, 2022, 5, e202101153.   | 1.3 | 1         |
| 4349 | Bro1 binds the Vps20 subunit of ESCRTâ€™II and promotes ESCRTâ€™II regulation by Doa4. Traffic, 2022, 23, 109-119.  | 1.3 | 1         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 4350 | Non-linear elastic properties of actin patches to partially rescue yeast endocytosis efficiency in the absence of the cross-linker Sac6. <i>Soft Matter</i> , 2022, , .   | 1.2 | 0         |
| 4351 | Structural basis for cytoplasmic dynein-1 regulation by Lis1. <i>ELife</i> , 2022, 11, .  | 2.8 | 29        |
| 4353 | Contribution of domain structure to the function of the yeast DEDD family exoribonuclease and RNase T functional homolog, Rex1. <i>Rna</i> , 2022, 28, 493-507.   | 1.6 | 1         |
| 4354 | Vps501, a novel vacuolar <scp>SNX&#x2013;BAR</scp> protein cooperates with the <scp>SEA</scp> complex to regulate <scp>TORC1</scp> signaling. <i>Traffic</i> , 2022, , .  | 1.3 | 3         |
| 4356 | Substrate-induced differential degradation and partitioning of the two tryptophan permeases Tat1 and Tat2 into eisosomes in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2022, 1864, 183858. | 1.4 | 3         |
| 4357 | A transcriptional roadblock protects yeast centromeres. <i>Nucleic Acids Research</i> , 2022, 50, 7801-7815.  | 6.5 | 11        |
| 4358 | Investigating cell autonomy in microorganisms. <i>Current Genetics</i> , 2022, 68, 305.   | 0.8 | 0         |
| 4360 | SWI/SNF senses carbon starvation with a pH-sensitive low-complexity sequence. <i>ELife</i> , 2022, 11, .  | 2.8 | 23        |
| 4363 | RNA modifications detection by comparative Nanopore direct RNA sequencing. <i>Nature Communications</i> , 2021, 12, 7198.   | 5.8 | 163       |
| 4364 | Elongating RNA polymerase II and RNA:DNA hybrids hinder fork progression and gene expression at sites of head-on replication-transcription collisions. <i>Nucleic Acids Research</i> , 2021, 49, 12769-12784.                           | 6.5 | 28        |
| 4365 | A New Role for the Mitochondrial Processing Peptidase in Protein Translocation into Mitochondria. <i>SSRN Electronic Journal</i> , 0, , .   | 0.4 | 0         |
| 4367 | Bur1 functions with TORC1 for vacuole&#x2013;mediated cell cycle progression. <i>EMBO Reports</i> , 2022, 23, e53477.   | 2.0 | 8         |
| 4368 | Mitochondrial energy metabolism regulates the nutrient import activity and endocytosis of APC transporters. <i>FEBS Letters</i> , 2022, , .   | 1.3 | 0         |
| 4369 | The chaperone Tsr2 regulates Rps26 release and reincorporation from mature ribosomes to enable a reversible, ribosome-mediated response to stress. <i>Science Advances</i> , 2022, 8, eabl4386.   | 4.7 | 17        |
| 4370 | Uip4p modulates nuclear pore complex function in <i>Saccharomyces cerevisiae</i>. <i>Nucleus</i> , 2022, 13, 79-94.   | 0.6 | 2         |
| 4371 | Rewiring phospholipid biosynthesis reveals resilience to membrane perturbations and uncovers regulators of lipid homeostasis. <i>EMBO Journal</i> , 2022, 41, e109998.  | 3.5 | 21        |
| 4372 | Iron Supplementation Delays Aging and Extends Cellular Lifespan through Potentiation of Mitochondrial Function. <i>Cells</i> , 2022, 11, 862.   | 1.8 | 10        |
| 4374 | Cdc48Ufd1/Npl4 segregase removes mislocalized centromeric histone H3 variant CENP-A from non-centromeric chromatin. <i>Nucleic Acids Research</i> , 2022, 50, 3276-3291.  | 6.5 | 6         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 4375 | Linkage reprogramming by tailor-made E3s reveals polyubiquitin chain requirements in DNA-damage bypass. <i>Molecular Cell</i> , 2022, 82, 1589-1602.e5.  | 4.5 | 4         |
| 4376 | An adaptable live-cell imaging protocol to analyze organelle morphology in <i>Saccharomyces cerevisiae</i> . <i>STAR Protocols</i> , 2022, 3, 101124.  | 0.5 | 1         |
| 4377 | Collision of germline POLE and PMS2 variants in a young patient treated with immune checkpoint inhibitors. <i>Npj Precision Oncology</i> , 2022, 6, 15.  | 2.3 | 11        |
| 4379 | Pheromone Guidance of Polarity Site Movement in Yeast. <i>Biomolecules</i> , 2022, 12, 502.  | 1.8 | 1         |
| 4381 | The microtubule plus-end tracking protein Bik1 is required for chromosome congression. <i>Molecular Biology of the Cell</i> , 2022, 33, mbcE21100500.  | 0.9 | 1         |
| 4382 | Identification of 14-3-3 proteins, Polo kinase, and RNA-binding protein Pes4 as key regulators of meiotic commitment in budding yeast. <i>Current Biology</i> , 2022, 32, 1534-1547.e9.  | 1.8 | 12        |
| 4383 | Tom70-based transcriptional regulation of mitochondrial biogenesis and aging. <i>ELife</i> , 2022, 11, .   | 2.8 | 16        |
| 4384 | The Nup2 meiotic-autonomous region relieves inhibition of Nup60 to promote progression of meiosis and sporulation in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2022, 221, .  | 1.2 | 4         |
| 4385 | Cdc14 spatiotemporally dephosphorylates Atg13 to activate autophagy during meiotic divisions. <i>Journal of Cell Biology</i> , 2022, 221, .  | 2.3 | 9         |
| 4386 | Dedicated chaperones coordinate co-translational regulation of ribosomal protein production with ribosome assembly to preserve proteostasis. <i>ELife</i> , 2022, 11, .  | 2.8 | 11        |
| 4387 | Follicular lymphoma-associated mutations in the V-ATPase chaperone VMA21 activate autophagy creating a targetable dependency. <i>Autophagy</i> , 2022, 18, 1982-2000.  | 4.3 | 9         |
| 4389 | The flavonoid corylin exhibits lifespan extension properties in mouse. <i>Nature Communications</i> , 2022, 13, 1238.  | 5.8 | 10        |
| 4390 | The Mnn10/Anp1-dependent N-linked outer chain glycan is dispensable for <i>Candida albicans</i> cell wall integrity. <i>Genetics</i> , 2022, 221, .  | 1.2 | 4         |
| 4392 | Emergence of the primordial pre-60S from the 90S pre-ribosome. <i>Cell Reports</i> , 2022, 39, 110640.   | 2.9 | 17        |
| 4393 | Deletion of autophagy related, ATG1 and F-box motif encoding YDR131C, together, lead to synthetic growth defects and flocculation behavior in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biochemical and Molecular Toxicology</i> , 2022, , e23064. | 1.4 | 0         |
| 4395 | Coordinated regulation of TORC2 signaling by MCC/eisosome-associated proteins, Pil1 and tetraspan membrane proteins during the stress response. <i>Molecular Microbiology</i> , 2022, 117, 1227-1244.  | 1.2 | 9         |
| 4396 | Clathrin-mediated trafficking of phospholipid flippases is required for local plasma membrane/cell wall damage repair in budding yeast. <i>Biochemical and Biophysical Research Communications</i> , 2022, 606, 156-162.                                 | 1.0 | 0         |
| 4397 | An optimized protocol to analyze membrane protein degradation in yeast using quantitative western blot and flow cytometry. <i>STAR Protocols</i> , 2022, 3, 101274.  | 0.5 | 3         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 4398 | Antagonism between killer yeast strains as an experimental model for biological nucleation dynamics. <i>ELife</i> , 2021, 10, .   | 2.8 | 8         |
| 4401 | A Set of Plasmid-Based Modules for Easy Switching of C-Terminal Epitope Tags in <i>Saccharomyces cerevisiae</i> . <i>Microorganisms</i> , 2021, 9, 2505.  | 1.6 | 1         |
| 4402 | A pathogenic role for histone H3 copper reductase activity in a yeast model of Friedreich's ataxia. <i>Science Advances</i> , 2021, 7, eabj9889.  | 4.7 | 6         |
| 4403 | Spatial control of avidity regulates initiation and progression of selective autophagy. <i>Nature Communications</i> , 2021, 12, 7194.  | 5.8 | 14        |
| 4404 | Analysis of Ubiquitylation and SUMOylation of Yeast Nuclear Pore Complex Proteins. <i>Methods in Molecular Biology</i> , 2022, 2502, 259-269.   | 0.4 | 0         |
| 4406 | Dimerization-dependent membrane tethering by Atg23 is essential for yeast autophagy. <i>Cell Reports</i> , 2022, 39, 110702.  | 2.9 | 8         |
| 4407 | Proteolysis of adaptor protein Mmr1 during budding is necessary for mitochondrial homeostasis in <i>Saccharomyces cerevisiae</i> . <i>Nature Communications</i> , 2022, 13, 2005.                                       | 5.8 | 6         |
| 4408 | Cryo-ET detects bundled triple helices but not ladders in meiotic budding yeast. <i>PLoS ONE</i> , 2022, 17, e0266035.  | 1.1 | 2         |
| 4409 | Tricalbin proteins regulate plasma membrane phospholipid homeostasis. <i>Life Science Alliance</i> , 2022, 5, e202201430.   | 1.3 | 9         |
| 4411 | Reciprocal regulation of phosphatidylcholine synthesis and H3K36 methylation programs metabolic adaptation. <i>Cell Reports</i> , 2022, 39, 110672.   | 2.9 | 3         |
| 4412 | Vps68 cooperates with ESCRT-III in intraluminal vesicle formation. <i>Journal of Cell Science</i> , 2022, 135, .  | 1.2 | 3         |
| 4413 | Cla4p Kinase Activity Is Down-Regulated by Fus3p during Yeast Mating. <i>Biomolecules</i> , 2022, 12, 598.  | 1.8 | 4         |
| 4414 | SWI/SNF and the histone chaperone Rtt106 drive expression of the Pleiotropic Drug Resistance network genes. <i>Nature Communications</i> , 2022, 13, 1968.  | 5.8 | 3         |
| 4415 | Hsp90 and phosphorylation of the Slr2(Mpk1) MAP kinase activation loop are essential for catalytic, but not non-catalytic, Slr2-mediated transcription in yeast. <i>Cell Stress and Chaperones</i> , 2022, 27, 295-304. | 1.2 | 2         |
| 4416 | Utilizing chemically induced dimerization of FKBP to analyze endocytosis by live-cell imaging in budding yeast. <i>STAR Protocols</i> , 2022, 3, 101323.  | 0.5 | 1         |
| 4438 | Meiotic cDNA libraries reveal gene truncations and mitochondrial proteins important for competitive fitness in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2022, , .  | 1.2 | 0         |
| 4439 | Multiple nutrient transporters enable cells to mitigate a rate-affinity tradeoff. <i>PLoS Computational Biology</i> , 2022, 18, e1010060.   | 1.5 | 4         |
| 4444 | SATurated Transposon Analysis in Yeast (SATAY) for Deep Functional Mapping of Yeast Genomes. <i>Methods in Molecular Biology</i> , 2022, 2477, 349-379.   | 0.4 | 1         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 4447 | Pls1 Is a Peroxisomal Matrix Protein with a Role in Regulating Lysine Biosynthesis. <i>Cells</i> , 2022, 11, 1426.  | 1.8 | 3         |
| 4448 | Regulation of CLB6 expression by the cytoplasmic deadenylase Ccr4 through its coding and 3' UTR regions. <i>PLoS ONE</i> , 2022, 17, e0268283.  | 1.1 | 2         |
| 4450 | Clearance of an amyloid-like translational repressor is governed by 14-3-3 proteins. <i>Cell Reports</i> , 2022, 39, 110753.  | 2.9 | 9         |
| 4451 | The activator/repressor Hap1 binds to the yeast <i>eIF5A</i> -encoding gene <i>TIF51A</i> to adapt its expression to the mitochondrial functional status. <i>FEBS Letters</i> , 2022, 596, 1809-1826.   | 1.3 | 5         |
| 4452 | Vacuole fragmentation depends on a novel Atg18-containing retromer-complex. <i>Autophagy</i> , 2023, 19, 278-295.   | 4.3 | 13        |
| 4456 | Mechanism of ER stress-mediated ER-phagy by CdTe-QDs in yeast cells. <i>Toxicology Letters</i> , 2022, 365, 36-45.  | 0.4 | 2         |
| 4457 | Fine-tuning acetyl-CoA carboxylase 1 activity through localization: functional genomics reveals a role for the lysine acetyltransferase NuA4 and sphingolipid metabolism in regulating Acc1 activity and localization. <i>Genetics</i> , 2022, 221, . | 1.2 | 7         |
| 4458 | Follicular lymphoma-associated mutations in the V-ATPase chaperone Vma21 activate autophagy by dysfunctional V-ATPase assembly. , 2022, 1, 226-233.   |     | 0         |
| 4459 | Ribosomal protein eL39 is important for maturation of the nascent polypeptide exit tunnel and proper protein folding during translation. <i>Nucleic Acids Research</i> , 2022, 50, 6453-6473.   | 6.5 | 8         |
| 4462 | METALIC reveals interorganelle lipid flux in live cells by enzymatic mass tagging. <i>Nature Cell Biology</i> , 2022, 24, 996-1004.   | 4.6 | 26        |
| 4463 | Cip1 tunes cell cycle arrest duration upon calcineurin activation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022, 119, .  | 3.3 | 1         |
| 4465 | The MyLO CRISPR-Cas9 toolkit: a markerless yeast localization and overexpression CRISPR-Cas9 toolkit. <i>G3: Genes, Genomes, Genetics</i> , 2022, 12, .   | 0.8 | 7         |
| 4467 | Vacuole-Specific Lipid Release for Tracking Intracellular Lipid Metabolism and Transport in <i>Saccharomyces cerevisiae</i> . <i>ACS Chemical Biology</i> , 2022, 17, 1485-1494.  | 1.6 | 4         |
| 4468 | Phosphatidic acid species 34:1 mediates expression of the myo-inositol 3-phosphate synthase gene <i>INO1</i> for lipid synthesis in yeast. <i>Journal of Biological Chemistry</i> , 2022, 298, 102148.  | 1.6 | 5         |
| 4469 | G1-Cyclin2 (Cln2) promotes chromosome hypercondensation in <i>eco1/ctf7 rad61</i> null cells during hyperthermic stress in <i>Saccharomyces cerevisiae</i> . <i>G3: Genes, Genomes, Genetics</i> , 2022, 12, .  | 0.8 | 0         |
| 4470 | Copper inhibits protein maturation in the secretory pathway by targeting the Sec61 translocon in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2022, 298, 102170.  | 1.6 | 3         |
| 4472 | Establishment of a Cre-loxP System Based on a Leaky LAC4 Promoter and an Unstable panARS Element in <i>Kluyveromyces marxianus</i> . <i>Microorganisms</i> , 2022, 10, 1240.  | 1.6 | 2         |
| 4473 | Architecture of the linker-scaffold in the nuclear pore. <i>Science</i> , 2022, 376, .  | 6.0 | 51        |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 4474 | Identification of 8-Hydroxyquinoline Derivatives That Decrease Cystathionine Beta Synthase (CBS) Activity. <i>International Journal of Molecular Sciences</i> , 2022, 23, 6769.  | 1.8 | 2         |
| 4475 | Nuclear pore complex acetylation regulates <scp>mRNA</scp> export and cell cycle commitment in budding yeast. <i>EMBO Journal</i> , 2022, 41, .  | 3.5 | 4         |
| 4476 | The histone H2B Arg95 residue links the pheromone response pathway to rapamycin-induced G1 arrest in yeast. <i>Scientific Reports</i> , 2022, 12, .  | 1.6 | 3         |
| 4477 | Cytostatic effects of structurally different ginsenosides on yeast cells with altered sterol biosynthesis and transport. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2022, 1864, 183993.   | 1.4 | 4         |
| 4478 | DIP2 is a unique regulator of diacylglycerol lipid homeostasis in eukaryotes. <i>ELife</i> , 0, 11, .  | 2.8 | 5         |
| 4481 | Adaptor linked K63 di-ubiquitin activates Nedd4/Rsp5 E3 ligase. <i>ELife</i> , 0, 11, .  | 2.8 | 3         |
| 4482 | Sus1 maintains a normal lifespan through regulation of TREX-2 complex-mediated mRNA export. <i>Aging</i> , 2022, , 4990-5012.  | 1.4 | 3         |
| 4484 | The Intracellular Distribution of the Small GTPase Rho5 and Its Dimeric Guanidine Nucleotide Exchange Factor Dck1/Lmo1 Determine Their Function in Oxidative Stress Response. <i>International Journal of Molecular Sciences</i> , 2022, 23, 7896. | 1.8 | 3         |
| 4485 | An integrated model for termination of RNA polymerase III transcription. <i>Science Advances</i> , 2022, 8, .  | 4.7 | 14        |
| 4486 | Distinct functions of three chromatin remodelers in activator binding and preinitiation complex assembly. <i>PLoS Genetics</i> , 2022, 18, e1010277.   | 1.5 | 3         |
| 4487 | Sen1 is a key regulator of transcription-driven conflicts. <i>Molecular Cell</i> , 2022, 82, 2952-2966.e6.   | 4.5 | 14        |
| 4488 | Reconstitution of kinetochore motility and microtubule dynamics reveals a role for a kinesin-8 in establishing end-on attachments. <i>ELife</i> , 0, 11, .   | 2.8 | 5         |
| 4489 | Haploinsufficiency of the sex-determining genes at MAT $\hat{1}$ ± restricts genome expansion in <i>Saccharomyces cerevisiae</i> . <i>IScience</i> , 2022, 25, 104783.   | 1.9 | 2         |
| 4490 | The N-Terminal Domain of Bfa1 Coordinates Mitotic Exit Independent of GAP Activity in <i>Saccharomyces cerevisiae</i> . <i>Cells</i> , 2022, 11, 2179.   | 1.8 | 0         |
| 4491 | Ddp1 Cooperates with Ppx1 to Counter a Stress Response Initiated by Nonvacuolar Polyphosphate. <i>MBio</i> , 2022, 13, .   | 1.8 | 10        |
| 4493 | Calmodulin kinase 2 genetically interacts with Rch1p to negatively regulate calcium import into <i>Saccharomyces cerevisiae</i> after extracellular calcium pulse. <i>Archives of Microbiology</i> , 2022, 204, .                                  | 1.0 | 2         |
| 4495 | The RNA-binding protein Puf5 and the HMGB protein Ixr1 contribute to cell cycle progression through the regulation of cell cycle-specific expression of CLB1 in <i>Saccharomyces cerevisiae</i> . <i>PLoS Genetics</i> , 2022, 18, e1010340.       | 1.5 | 1         |
| 4496 | Ubiquitination drives COPI priming and Golgi SNARE localization. <i>ELife</i> , 0, 11, .   | 2.8 | 5         |



| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 4499 | Glutamine transport as a possible regulator of nitrogen catabolite repression in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2022, 39, 493-507.   | 0.8 | 1         |
| 4500 | Reversal of histone H2B mono-ubiquitination is required for replication stress recovery. <i>DNA Repair</i> , 2022, 119, 103387.  | 1.3 | 1         |
| 4501 | Phosphoregulation of the ATP synthase beta subunit stimulates mitochondrial activity for G2/M progression. <i>Advances in Biological Regulation</i> , 2022, 85, 100905.  | 1.4 | 2         |
| 4502 | Aurora B activity is promoted by cooperation between discrete localization sites in budding yeast. <i>Molecular Biology of the Cell</i> , 2022, 33, .  | 0.9 | 4         |
| 4504 | Receptor for advanced glycation end-products (RAGE) mediates phagocytosis in nonprofessional phagocytes. <i>Communications Biology</i> , 2022, 5, .  | 2.0 | 3         |
| 4506 | Heterologous (Over) Expression of Human SoLute Carrier (SLC) in Yeast: A Well-Recognized Tool for Human Transporter Function/Structure Studies. <i>Life</i> , 2022, 12, 1206.  | 1.1 | 1         |
| 4508 | The lipid flippase Drs2 regulates anterograde transport of Atg9 during autophagy. , 2022, 1, 345-367.  |     | 1         |
| 4509 | The histone deacetylases Rpd3 and Hst1 antagonistically regulate de novo NAD <sup>+</sup> metabolism in the budding yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2022, , 102410.   | 1.6 | 3         |
| 4511 | Diploid-associated adaptation to chronic low-dose UV irradiation requires homologous recombination in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 0, , .   | 1.2 | 0         |
| 4515 | Rvb1/Rvb2 proteins couple transcription and translation during glucose starvation. <i>ELife</i> , 0, 11, .   | 2.8 | 2         |
| 4516 | Mechanism of commitment to a mating partner in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2022, 33, .  | 0.9 | 2         |
| 4517 | Expression level of Sec62 modulates membrane insertion of marginally hydrophobic segments. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2022, 1864, 184051.   | 1.4 | 0         |
| 4518 | Acquired resistance to severe ethanol stress-induced inhibition of proteasomal proteolysis in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2022, 1866, 130241.          | 1.1 | 4         |
| 4520 | Repression of essential cell cycle genes increases cellular fitness. <i>PLoS Genetics</i> , 2022, 18, e1010349.  | 1.5 | 2         |
| 4522 | A balancing act: interactions within NuA4/TIP60 regulate picNuA4 function in <i>Saccharomyces cerevisiae</i> and humans. <i>Genetics</i> , 0, , .  | 1.2 | 0         |
| 4523 | Rpd3 regulates single-copy origins independently of the rDNA array by opposing Fkh1-mediated origin stimulation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022, 119, . | 3.3 | 9         |
| 4527 | Sensing of individual stalled 80S ribosomes by Fap1 for nonfunctional rRNA turnover. <i>Molecular Cell</i> , 2022, 82, 3424-3437.e8.   | 4.5 | 15        |
| 4529 | Eisosome protein Pil1 regulates mitochondrial morphology, mitophagy, and cell death in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2022, 298, 102533.                                   | 1.6 | 2         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 4531 | Ferric reductase-related proteins mediate fungal heme acquisition. <i>ELife</i> , 0, 11, .  | 2.8 | 3         |
| 4532 | Loss of Num1-mediated cortical dynein anchoring negatively impacts respiratory growth. <i>Journal of Cell Science</i> , 2022, 135, .  | 1.2 | 2         |
| 4534 | Causative links between ER stress and oxidative damage in a yeast model of human N88S seipinopathy. <i>Free Radical Biology and Medicine</i> , 2022, 192, 165-181.                                      | 1.3 | 3         |
| 4535 | Kap- $\beta$ 2/Transportin mediates $\beta$ -catenin nuclear transport in Wnt signaling. <i>ELife</i> , 0, 11, .  | 2.8 | 5         |
| 4536 | The role of bivalent ions in the regulation of D-loop extension mediated by DMC1 during meiotic recombination. <i>IScience</i> , 2022, 25, 105439.  | 1.9 | 2         |
| 4537 | Termination of the unfolded protein response is guided by ER stress-induced HAC1 mRNA nuclear retention. <i>Nature Communications</i> , 2022, 13, .   | 5.8 | 3         |
| 4538 | Structure of the pre-mRNA leakage 39-kDa protein reveals a single domain of integrated zf-C3HC and Rsm1 modules. <i>Scientific Reports</i> , 2022, 12, .  | 1.6 | 1         |
| 4539 | Generation and characterization of conditional yeast mutants affecting each of the 2 essential functions of the scaffolding proteins Boi1/2 and Bem1. <i>G3: Genes, Genomes, Genetics</i> , 2022, 12, . | 0.8 | 1         |
| 4540 | A Simple, Improved Method for Scarless Genome Editing of Budding Yeast Using CRISPR-Cas9. <i>Methods and Protocols</i> , 2022, 5, 79.   | 0.9 | 0         |
| 4541 | A method for the isolation and characterization of autophagic bodies from yeast provides a key tool to investigate cargos of autophagy. <i>Journal of Biological Chemistry</i> , 2022, 298, 102641.     | 1.6 | 5         |
| 4542 | DNA damage-induced nuclear import of HSP90 $\alpha$ is promoted by Aha1. <i>Molecular Biology of the Cell</i> , 2022, 33, .   | 0.9 | 1         |
| 4544 | Hypusinated eIF5A Promotes Ribosomal Frameshifting during Decoding of ODC Antizyme mRNA in <i>Saccharomyces cerevisiae</i> . <i>International Journal of Molecular Sciences</i> , 2022, 23, 12972.      | 1.8 | 2         |
| 4546 | A two-tiered system for selective receptor and transporter protein degradation. <i>PLoS Genetics</i> , 2022, 18, e1010446.  | 1.5 | 1         |
| 4547 | Dissociation of ERMES clusters plays a key role in attenuating the endoplasmic reticulum stress. <i>IScience</i> , 2022, 25, 105362.  | 1.9 | 4         |
| 4548 | Inducible transcriptional condensates drive 3D genome reorganization in the heat shock response. <i>Molecular Cell</i> , 2022, 82, 4386-4399.e7.  | 4.5 | 20        |
| 4552 | The Polo kinase Cdc5 is regulated at multiple levels in the adaptation response to telomere dysfunction. <i>Genetics</i> , 0, , .   | 1.2 | 1         |
| 4553 | High-resolution secretory timeline from vesicle formation at the Golgi to fusion at the plasma membrane in <i>S. cerevisiae</i> . <i>ELife</i> , 0, 11, .   | 2.8 | 4         |
| 4555 | Regulation of Cdc42 protein turnover modulates the filamentous growth MAPK pathway. <i>Journal of Cell Biology</i> , 2022, 221, .   | 2.3 | 4         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 4556 | Atg9 interactions via its transmembrane domains are required for phagophore expansion during autophagy. <i>Autophagy</i> , 2023, 19, 1459-1478.  | 4.3 | 16        |
| 4557 | SETD2 transcriptional control of ATG14L/S isoforms regulates autophagosome-lysosome fusion. <i>Cell Death and Disease</i> , 2022, 13, .  | 2.7 | 7         |
| 4560 | Lipid saturation induces degradation of squalene epoxidase for sterol homeostasis and cell survival. <i>Life Science Alliance</i> , 2023, 6, e202201612.   | 1.3 | 5         |
| 4561 | Set1 regulates telomere function via H3K4 methylation-dependent and -independent pathways and calibrates the abundance of telomere maintenance factors. <i>Molecular Biology of the Cell</i> , 2023, 34, . | 0.9 | 2         |
| 4562 | Cms1 coordinates stepwise local 90S pre-ribosome assembly with timely snR83 release. <i>Cell Reports</i> , 2022, 41, 111684.   | 2.9 | 1         |
| 4563 | Dbf4 Zn-Finger Motif Is Specifically Required for Stimulation of Ctf19-Activated Origins in <i>Saccharomyces cerevisiae</i> . <i>Genes</i> , 2022, 13, 2202.   | 1.0 | 0         |
| 4564 | Mitotic phosphorylation inhibits the Golgi mannosidase MAN1A1. <i>Cell Reports</i> , 2022, 41, 111679.   | 2.9 | 3         |
| 4565 | Autophagy promotes cell survival by maintaining NAD levels. <i>Developmental Cell</i> , 2022, 57, 2584-2598.e11.   | 3.1 | 31        |
| 4566 | N6-methyladenosine (m6A) reader Pho92 is recruited co-transcriptionally and couples translation to mRNA decay to promote meiotic fitness in yeast. <i>ELife</i> , 0, 11, .                                 | 2.8 | 17        |
| 4568 | Rate of transcription elongation and sequence-specific pausing by RNA polymerase I directly influence rRNA processing. <i>Journal of Biological Chemistry</i> , 2022, 298, 102730.                         | 1.6 | 4         |
| 4570 | Tup1 is critical for transcriptional repression in Quiescence in <i>S. cerevisiae</i> . <i>PLoS Genetics</i> , 2022, 18, e1010559.   | 1.5 | 5         |
| 4571 | SESAME-catalyzed H3T11 phosphorylation inhibits Dot1-catalyzed H3K79me3 to regulate autophagy and telomere silencing. <i>Nature Communications</i> , 2022, 13, .   | 5.8 | 7         |
| 4572 | A single 2'-O-methylation of ribosomal RNA gates assembly of a functional ribosome. <i>Nature Structural and Molecular Biology</i> , 2023, 30, 91-98.  | 3.6 | 14        |
| 4573 | Adaptation to high rates of chromosomal instability and aneuploidy through multiple pathways in budding yeast. <i>EMBO Journal</i> , 2023, 42, .   | 3.5 | 10        |
| 4574 | Rad51-mediated interhomolog recombination during budding yeast meiosis is promoted by the meiotic recombination checkpoint and the conserved Pif1 helicase. <i>PLoS Genetics</i> , 2022, 18, e1010407.     | 1.5 | 2         |
| 4575 | Ultrastructure expansion microscopy reveals the cellular architecture of budding and fission yeast. <i>Journal of Cell Science</i> , 2022, 135, .  | 1.2 | 12        |
| 4576 | Meiotic nuclear pore complex remodeling provides key insights into nuclear basket organization. <i>Journal of Cell Biology</i> , 2023, 222, .  | 2.3 | 8         |
| 4577 | Bidirectional promoter activity from expression cassettes can drive off-target repression of neighboring gene translation. <i>ELife</i> , 0, 11, .   | 2.8 | 4         |

| #    | ARTICLE   | IF   | CITATIONS |
|------|---|------|-----------|
| 4578 | Coupling Cell Communication and Optogenetics: Implementation of a Light-Inducible Intercellular System in Yeast. <i>ACS Synthetic Biology</i> , 2023, 12, 71-82.  | 1.9  | 4         |
| 4579 | Live Cell Imaging of Yeast Golgi Dynamics. <i>Methods in Molecular Biology</i> , 2023, , 3-15.  | 0.4  | 0         |
| 4580 | Hsp70 and Hsp110 Chaperones Promote Early Steps of Proteasome Assembly. <i>Biomolecules</i> , 2023, 13, 11.   | 1.8  | 1         |
| 4581 | Pab1 acetylation at K131 decreases stress granule formation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2023, 299, 102834.  | 1.6  | 2         |
| 4585 | New insights into the regulation of Cystathionine beta synthase (CBS), an enzyme involved in intellectual deficiency in Down syndrome. <i>Frontiers in Neuroscience</i> , 0, 16, .  | 1.4  | 3         |
| 4587 | A toolbox for systematic discovery of stable and transient protein interactors in baker's yeast. <i>Molecular Systems Biology</i> , 2023, 19, .   | 3.2  | 9         |
| 4589 | Using Budding Yeast as a Model to Understand Dynein-Mediated Cargo Transport. <i>Methods in Molecular Biology</i> , 2023, , 25-42.  | 0.4  | 2         |
| 4590 | Decoding of the ubiquitin code for clearance of colliding ribosomes by the RQT complex. <i>Nature Communications</i> , 2023, 14, .  | 5.8  | 11        |
| 4591 | An essential Noc3p dimerization cycle mediates ORC double-hexamer formation in replication licensing. <i>Life Science Alliance</i> , 2023, 6, e202201594.   | 1.3  | 0         |
| 4593 | Two modes of Cue2-mediated mRNA cleavage with distinct substrate recognition initiate no-go decay. <i>Nucleic Acids Research</i> , 2023, 51, 253-270.   | 6.5  | 9         |
| 4594 | The RNA export and RNA decay complexes THO and TRAMP prevent transcription-replication conflicts, DNA breaks, and CAG repeat contractions. <i>PLoS Biology</i> , 2022, 20, e3001940.  | 2.6  | 2         |
| 4596 | Phosphosite Scanning reveals a complex phosphorylation code underlying CDK-dependent activation of Hcm1. <i>Nature Communications</i> , 2023, 14, .   | 5.8  | 7         |
| 4597 | Lysate-to-grid: Rapid Isolation of Native Complexes from Budding Yeast for Cryo-EM Imaging. <i>Bio-protocol</i> , 2023, 13, .   | 0.2  | 1         |
| 4598 | Hotspot of de novo telomere addition stabilizes linear amplicons in yeast grown in sulfate-limiting conditions. <i>Genetics</i> , 2023, 224, .  | 1.2  | 2         |
| 4599 | Complementation Assay Using Fusion of Split-GFP and TurboID (CsFiND) Enables Simultaneous Visualization and Proximity Labeling of Organelle Contact Sites in Yeast. <i>Contact (Thousand Oaks)</i> Tj ETQq0 0 0 rgBT4/Overlook 10 Tf 50 |      |           |
| 4600 | Mitochondrial complexome reveals quality-control pathways of protein import. <i>Nature</i> , 2023, 614, 153-159.  | 13.7 | 47        |
| 4601 | Multi-site phosphorylation of yeast Mif2/CENP-C promotes inner kinetochore assembly. <i>Current Biology</i> , 2023, 33, 688-696.e6.   | 1.8  | 3         |
| 4602 | Concurrent remodelling of nucleolar 60S subunit precursors by the Rea1 ATPase and Spb4 RNA helicase. <i>ELife</i> , 0, 12, .  | 2.8  | 3         |

| #    | ARTICLE  | IF  | CITATIONS |
|------|--|-----|-----------|
| 4603 | Meiotic cells escape prolonged spindle checkpoint activity through kinetochore silencing and slippage. <i>PLoS Genetics</i> , 2023, 19, e1010707.  | 1.5 | 1         |
| 4604 | Exploring the consequences of redirecting an exocytic Rab onto endocytic vesicles. <i>Molecular Biology of the Cell</i> , 2023, 34, .  | 0.9 | 0         |
| 4605 | Systematic analysis of microtubule plus-end networks defines EB-cargo complexes critical for mitosis in budding yeast. <i>Molecular Biology of the Cell</i> , 2023, 34, .  | 0.9 | 1         |
| 4606 | Helicase activities of Rad5 and Rrm3 genetically interact in the prevention of recombinogenic DNA lesions in <i>Saccharomyces cerevisiae</i> . <i>DNA Repair</i> , 2023, 126, 103488.                                    | 1.3 | 2         |
| 4607 | Protocol for monitoring and analyzing single nucleotide incorporation by <i>S. cerevisiae</i> RNA polymerases. <i>STAR Protocols</i> , 2023, 4, 102191.  | 0.5 | 2         |
| 4608 | Spontaneous deamination of cytosine to uracil is biased to the non-transcribed DNA strand in yeast. <i>DNA Repair</i> , 2023, 126, 103489.   | 1.3 | 1         |
| 4609 | Differential requirements for Gcn5 and NuA4 HAT activities in the starvation-induced versus basal transcriptomes. <i>Nucleic Acids Research</i> , 0, , .   | 6.5 | 1         |
| 4610 | Evolutionary tuning of barbed end competition allows simultaneous construction of architecturally distinct actin structures. <i>Journal of Cell Biology</i> , 2023, 222, .   | 2.3 | 10        |
| 4611 | Rad51 filaments assembled in the absence of the complex formed by the Rad51 paralogs Rad55 and Rad57 are outcompeted by translesion DNA polymerases on UV-induced ssDNA gaps. <i>PLoS Genetics</i> , 2023, 19, e1010639. | 1.5 | 2         |
| 4612 | Essential mitochondrial role in iron-sulfur cluster assembly of the cytoplasmic isopropylmalate isomerase Leu1 in <i>Saccharomyces cerevisiae</i> . <i>Mitochondrion</i> , 2023, 69, 104-115.                            | 1.6 | 2         |
| 4613 | TFIID dependency of steady-state mRNA transcription altered epigenetically by simultaneous functional loss of Taf1 and Spt3 is Hsp104-dependent. <i>PLoS ONE</i> , 2023, 18, e0281233.                                   | 1.1 | 0         |
| 4617 | Molecular basis of eIF5A-dependent CAT tailing in eukaryotic ribosome-associated quality control. <i>Molecular Cell</i> , 2023, 83, 607-621.e4.  | 4.5 | 9         |
| 4618 | Discovery of a rapidly evolving yeast defense factor, <i>KTD1</i> , against the secreted killer toxin K28. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2023, 120, .         | 3.3 | 1         |
| 4620 | Parental histone distribution and location of the replication obstacle at nascent strands control homologous recombination. <i>Cell Reports</i> , 2023, 42, 112174.  | 2.9 | 1         |
| 4622 | Recruitment of the lipid kinase Mss4 to the meiotic spindle pole promotes prospore membrane formation in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2023, 34, .                            | 0.9 | 1         |
| 4624 | Phosphorylation controls the oligomeric state of She2 and mRNA localization in yeast. <i>Rna</i> , 2023, 29, 745-755.  | 1.6 | 0         |
| 4626 | Maize Antifungal Protein AFP1 Elevates Fungal Chitin Levels by Targeting Chitin Deacetylases and Other Glycoproteins. <i>MBio</i> , 2023, 14, .  | 1.8 | 4         |
| 4627 | Transcription termination complex, Rtt103-Rai1-Rat1, regulates sub-telomeric transcripts in <i>Saccharomyces cerevisiae</i> . <i>RNA Biology</i> , 2023, 20, 95-108.   | 1.5 | 0         |

| #    | ARTICLE   | IF  | CITATIONS |
|------|---|-----|-----------|
| 4631 | SIR telomere silencing depends on nuclear envelope lipids and modulates sensitivity to a lysolipid. <i>Journal of Cell Biology</i> , 2023, 222, .                                 | 2.3 | 1         |
| 4632 | The dynamin Vps1 mediates Atg9 transport to the sites of autophagosome formation. <i>Journal of Biological Chemistry</i> , 2023, 299, 104712.                                     | 1.6 | 1         |
| 4633 | A systematic proximity ligation approach to studying proteinâ€s substrate specificity identifies the substrate spectrum of the Ssh1 translocon. <i>EMBO Journal</i> , 2023, 42, . | 3.5 | 3         |
| 4634 | The UbiB family member Cqd1 forms a novel membrane contact site in mitochondria. <i>Journal of Cell Science</i> , 0, , .  | 1.2 | 1         |
| 4635 | Exo1 protects DNA nicks from ligation to promote crossover formation during meiosis. <i>PLoS Biology</i> , 2023, 21, e3002085.  | 2.6 | 5         |
| 4636 | Chaperone-directed ribosome repair after oxidative damage. <i>Molecular Cell</i> , 2023, 83, 1527-1537.e5.  | 4.5 | 8         |