

CITATION REPORT

List of articles citing

Angiographic assessment of myocardial reperfusion in patients treated with primary angioplasty for acute myocardial infarction: myocardial blush grade. Zwolle Myocardial Infarction Study Group

DOI: 10.1161/01.cir.97.23.2302
Circulation, 1998, 97, 2302-6.

Source: <https://exaly.com/paper-pdf/29644538/citation-report.pdf>

Version: 2024-04-23

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
1108	Determinants and prognostic implications of persistent ST-segment elevation after primary angioplasty for acute myocardial infarction: importance of microvascular reperfusion injury on clinical outcome. <i>Circulation</i> , 1999 , 99, 1972-7	16.7	313
1107	[Patency, perfusion and prognosis in acute myocardial infarct]. 1999 , 24, 421-9		8
1106	The role of myocardial viability in deriving benefit from reestablishing infarct-related artery flow after acute myocardial infarction. 2000 , 42, 455-470		10
1105	The use of the electrocardiogram to identify epicardial coronary and tissue reperfusion in acute myocardial infarction. 2000 , 10, 137-47		13
1104	The use of the electrocardiogram to identify epicardial coronary and tissue reperfusion in acute myocardial infarction. 2000 , 10, 5-14		3
1103	Treatment strategies for microvascular dysfunction following acute myocardial infarction. 2000 , 2, 405-10		1
1102	Reperfusion therapy and strategies in acute myocardial infarction. 2000 , 34, 355-6		
1101	Combination reperfusion therapy with abciximab and reduced dose reteplase: results from TIMI 14. The Thrombolysis in Myocardial Infarction (TIMI) 14 Investigators. 2000 , 21, 1944-53		68
1100	Primary angioplasty versus thrombolysis for acute myocardial infarction. 2000 , 342, 890-1; author reply 891-2		7
1099	Relationship of TIMI myocardial perfusion grade to mortality after administration of thrombolytic drugs. <i>Circulation</i> , 2000 , 101, 125-30	16.7	707
1098	Absence of preinfarction angina is associated with a risk of no-reflow phenomenon after primary coronary angioplasty for a first anterior wall acute myocardial infarction. 2000 , 75, 253-60		19
1097	Angiographic no-reflow phenomenon as a predictor of adverse long-term outcome in patients treated with percutaneous transluminal coronary angioplasty for first acute myocardial infarction. 2000 , 36, 1202-9		438
1096	Shifting the open-artery hypothesis downstream: the quest for optimal reperfusion. 2001 , 37, 9-18		201
1095	Postinfarctional remodeling: increased dye intensity in the myocardial risk area after angioplasty of infarct-related coronary artery is associated with reduction of ventricular volumes. 2001 , 37, 1239-45		12
1094	The reperfusion wars: where will the next battles be fought?. 2001 , 110, 65-6		2
1093	Direct infarct artery stenting without predilation and no-reflow in patients with acute myocardial infarction. 2001 , 142, 684-90		66
1092	ST segment resolution as a tool for assessing the efficacy of reperfusion therapy. 2001 , 38, 1283-94		212

1091	A randomized double-blind trial of intravenous trimetazidine as adjunctive therapy to primary angioplasty for acute myocardial infarction. 2001 , 77, 263-73		41
1090	[Biochemical markers of ischemic and non-ischemic myocardial damage]. 2001 , 96, 144-56		21
1089	Determinants and prognostic implications of terminal QRS complex distortion in patients treated with primary angioplasty for acute myocardial infarction. 2001 , 88, 210-3		34
1088	Association of noninvasive markers of coronary artery reperfusion to assess microvascular obstruction in patients with acute myocardial infarction treated with primary angioplasty. 2001 , 88, 342-6		23
1087	Predictive value of markers of myocardial reperfusion in acute myocardial infarction for follow-up left ventricular function. 2001 , 88, 1358-63		40
1086	Interventions in acute myocardial infarction. 2001 , 26, 619-72		3
1085	Integrated analysis of myocardial blush and ST-segment elevation recovery after successful primary angioplasty: Real-time grading of microvascular reperfusion and prediction of early and late recovery of left ventricular function. <i>Circulation</i> , 2002 , 106, 313-8	16.7	165
1084	Relationship of the TIMI myocardial perfusion grades, flow grades, frame count, and percutaneous coronary intervention to long-term outcomes after thrombolytic administration in acute myocardial infarction. <i>Circulation</i> , 2002 , 105, 1909-13	16.7	305
1083	Incidence and clinical significance of distal embolization during primary angioplasty for acute myocardial infarction. 2002 , 23, 1112-7		463
1082	Coronary flow velocity pattern immediately after percutaneous coronary intervention as a predictor of complications and in-hospital survival after acute myocardial infarction. <i>Circulation</i> , 2002 , 106, 3051-6	16.7	78
1081	Intracoronary thrombectomy with the X-sizer catheter system improves epicardial flow and accelerates ST-segment resolution in patients with acute coronary syndrome: a prospective, randomized, controlled study. <i>Circulation</i> , 2002 , 105, 2355-60	16.7	143
1080	Lessons learned from recent cardiovascular clinical trials: Part I. <i>Circulation</i> , 2002 , 106, 746-51	16.7	101
1079	Continuous vectorcardiography is superior to standard electrocardiography in the prediction of long-term outcome after thrombolysis in patients with acute myocardial infarction. 2002 , 13, 169-75		5
1078	How to best counteract the enemies? By ensuring adequate oxygen delivery. 2002 , 4, G35-G42		
1077	Impact of normalized myocardial perfusion after successful angioplasty in acute myocardial infarction. 2002 , 39, 591-7		322
1076	Role of coronary angiography in acute coronary artery syndromes. 2002 , 27, 411-45		
1075	Reliability of resolution of ST-segment elevation after coronary reperfusion in predicting myocardial salvage in anterior wall acute myocardial infarction. 2002 , 90, 227-32		7
1074	Methodologic and clinical validation of the TIMI myocardial perfusion grade in acute myocardial infarction. 2002 , 14, 233-7		21

1073	ST-segment monitoring in patients with acute coronary syndromes. 2003 , 5, 278-83	7
1072	Usefulness of myocardial blush grade early and late after primary coronary angioplasty for acute myocardial infarction in predicting left ventricular function. 2003 , 92, 1015-9	46
1071	Effect of the PercuSurge GuardWire device on the integrity of microvasculature and clinical outcomes during primary transradial coronary intervention in acute myocardial infarction. 2003 , 92, 1331-5	94
1070	Choc cardiogénique primaire à la phase aiguë de l'infarctus traité par angioplastie: expérience de trois centres alsaciens publics non universitaires. 2003 , 52, 344-348	
1069	Outcome in patients treated with primary angioplasty for acute myocardial infarction due to left main coronary artery occlusion. 2003 , 91, 235-8	54
1068	Frequency and sequelae of ST elevation acute myocardial infarction caused by spontaneous distal embolization from unstable coronary lesions. 2003 , 91, 708-11	18
1067	Angiographic assessment of myocardial perfusion in patients with isolated coronary artery ectasia. 2003 , 91, 996-9, A7	34
1066	Volumetric intravascular ultrasound evidence that distal embolization during acute infarct intervention contributes to inadequate myocardial perfusion grade. 2003 , 92, 728-32	39
1065	Modern management of acute myocardial infarction. 2003 , 28, 7-127	
1064	Evaluation of the PercuSurge Guardwire Plus Temporary Occlusion and Aspiration System during primary angioplasty in acute myocardial infarction. 2003 , 60, 443-51	21
1063	Prediction of clinical outcome after mechanical revascularization in acute myocardial infarction by markers of myocardial reperfusion. 2003 , 41, 532-8	83
1062	Thrombolysis in myocardial infarction myocardial perfusion grade in angiography correlates with myocardial salvage in patients with acute myocardial infarction treated with stenting or thrombolysis. 2003 , 41, 925-9	52
1061	Glucose-insulin-potassium infusion in patients treated with primary angioplasty for acute myocardial infarction: the glucose-insulin-potassium study: a randomized trial. 2003 , 42, 784-91	242
1060	Symptom-onset-to-balloon time and mortality in patients with acute myocardial infarction treated by primary angioplasty. 2003 , 42, 991-7	328
1059	Intracoronary thrombectomy improves myocardial reperfusion in patients undergoing direct angioplasty for acute myocardial infarction. 2003 , 42, 1395-402	158
1058	Diabetes mellitus and acute myocardial infarction: more data supporting a poorer microvasculature reperfusion. 2003 , 146, E6	2
1057	Use of the electrocardiogram in acute myocardial infarction. 2003 , 348, 933-40	242
1056	[Primary cardiogenic shock in the acute phase of myocardial infarction treated by angioplasty: experience of three non-university public centers in Alsace]. 2003 , 52, 344-8	

1055	Glycoprotein IIb/IIIa receptor antagonists: a comparative review of their use in percutaneous coronary intervention. 2003 , 3, 423-36		18
1054	Angiographic assessment of reperfusion in acute myocardial infarction by myocardial blush grade. <i>Circulation</i> , 2003 , 107, 2115-9	16.7	298
1053	Relation of coronary flow pattern to myocardial blush grade in patients with first acute myocardial infarction. 2003 , 89, 1147-51		30
1052	Primary stenting produces earlier and more efficient myocardial reperfusion than primary PTCA alone in patients with acute ST segment elevation MI. 2003 , 54, 195-203		
1051	Myocardial contrast echocardiography is superior to other known modalities for assessing myocardial reperfusion after acute myocardial infarction. 2003 , 89, 139-44		52
1050	Prognostic value of ST-segment resolution-when and what to measure. 2003 , 24, 337-45		34
1049	Mechanical prevention of distal embolization during primary angioplasty: safety, feasibility, and impact on myocardial reperfusion. <i>Circulation</i> , 2003 , 108, 171-6	16.7	128
1048	Effects of abciximab on microvascular integrity and left ventricular functional recovery in patients with acute infarction treated by primary coronary angioplasty. 2003 , 24, 67-76		64
1047	Incomplete resolution of ST-segment elevation is a marker of transient microcirculatory dysfunction after stenting for acute myocardial infarction. <i>Circulation</i> , 2003 , 107, 2684-9	16.7	53
1046	Resolution of ST-segment elevation after thrombolytic therapy in elderly patients with acute myocardial infarction. 2003 , 10, 83-7		1
1045	Effectiveness of thrombectomy before stent implantation in acute myocardial infarction. 2003 , 67, 951-4		10
1044	Relation between the timing of the last preinfarction angina and microvascular reperfusion in patients with recanalized acute myocardial infarction. 2003 , 44, 845-54		3
1043	Comparison of TIMI Frame Count and TIMI Myocardial Perfusion Grade in Assessment of Myocardial Reperfusion after Primary Angioplasty for Acute Myocardial Infarction. 2003 , 33, 864		2
1042	Efficacy of Embolic Protection using PercuSurge GuardWire System During Primary Percutaneous Coronary Intervention with the Lesions Suggesting Large Thrombus Burden. 2003 , 33, 1103		
1041	Relationship between collateral blood flow and microvascular perfusion after reperfused acute myocardial infarction. 2003 , 44, 855-63		8
1040	Glycoprotein IIb/IIIa Receptor Inhibitors. 2004 , 97-108		
1039	The Clinical Significance of Serum-Hydroxyproline in Acute Coronary Syndrome. 2004 , 34, 1148		
1038	Comparison of TIMI Myocardial Perfusion Grade with Coronary Flow Reserve for Prediction of Recovery of LV Function and LV Remodeling in Acute Myocardial Infarction. 2004 , 34, 247		6

1037	Influence of isoprostone F2alpha-III on reflow after myocardial infarction. 2004 , 25, 847-53	8
1036	Time-to-treatment significantly affects the extent of ST-segment resolution and myocardial blush in patients with acute myocardial infarction treated by primary angioplasty. 2004 , 25, 1009-13	154
1035	Assessing myocardial perfusion with the transthoracic Doppler technique in patients with reperfused anterior myocardial infarction: comparison with angiographic, enzymatic and electrocardiographic indices. 2004 , 25, 1526-33	33
1034	Impaired myocardial perfusion is a major explanation of the poor outcome observed in patients undergoing primary angioplasty for ST-segment-elevation myocardial infarction and signs of heart failure. <i>Circulation</i> , 2004 , 109, 958-61	16.7 43
1033	Role of collateral circulation in the acute phase of ST-segment-elevation myocardial infarction treated with primary coronary intervention. 2004 , 25, 854-8	88
1032	Effects and safety of intracoronary thrombectomy using transradial application of the PercuSurge distal balloon protection system in patients with early or recent myocardial infarction. 2004 , 102, 206-14	8
1031	The effect of ITF-1697 on reperfusion in patients undergoing primary angioplasty. Safety and efficacy of a novel tetrapeptide, ITF-1697. 2004 , 25, 392-400	8
1030	Facilitation of primary coronary angioplasty by early start of a glycoprotein 2b/3a inhibitor: results of the ongoing tirofiban in myocardial infarction evaluation (On-TIME) trial. 2004 , 25, 837-46	124
1029	Excimer laser coronary angioplasty in acute myocardial infarction. 2004 , 12, 291-5	1
1028	Relation between coronary pressure derived collateral flow, myocardial perfusion grade, and outcome in left ventricular function after rescue percutaneous coronary intervention. 2004 , 90, 1450-4	15
1027	New tools for assessing microvascular obstruction in patients with ST elevation myocardial infarction. 2004 , 90, 119-20	5
1026	Reperfusion treatment of ST-elevation acute myocardial infarction. 2004 , 47, 131-57	15
1025	Myocardial perfusion grade and survival after percutaneous transluminal coronary angioplasty in patients with cardiogenic shock. 2004 , 93, 1081-5	21
1024	Effect of intracoronary adenosine infusion during coronary intervention on myocardial reperfusion injury in patients with acute myocardial infarction. 2004 , 94, 9-13	65
1023	Comparison in patients having primary coronary angioplasty of abciximab versus tirofiban on recovery of left ventricular function. 2004 , 94, 35-9	55
1022	[How to assess coronary lesion, reperfusion quality, and PCI result?]. 2004 , 53, 222-8	
1021	Relation of lateral ST-segment elevation pattern to myocardial salvage in patients with recanalized anterolateral acute myocardial infarction. 2004 , 27, 106-11	2
1020	The effect of ST resolution on QT dispersion after interventional treatment in acute myocardial infarction. 2004 , 27, 159-62	4

1019	Thrombus aspiration prior to primary angioplasty in acute myocardial infarction: estimation of rescued myocardial tissue by return of ST-segment elevation. 2004 , 27, 451-4	2
1018	Reperfusion of myocardium does not necessarily equate with epicardial coronary artery blood flow. 2004 , 27, 589-90	
1017	Markers of myocardial reperfusion as predictors of left ventricular function recovery in acute myocardial infarction treated with primary angioplasty. 2004 , 27, 683-8	15
1016	[Thrombolysis in ST-elevation myocardial infarction. Current role in the light of recent studies]. 2004 , 53, 445-54	1
1015	Ubiquitin-free routes into the proteasome. 2004 , 61, 1596-600	82
1014	Transradial application of PercuSurge GuardWire device during primary percutaneous intervention of infarct-related artery with high-burden thrombus formation. 2004 , 61, 503-11	9
1013	Relationship between pattern of slow-reflow and tissue level of perfusion after coronary stenting: A single-patient report. 2004 , 61, 222-6	1
1012	Angiographic predictors of left ventricular ejection fraction after successful angioplasty in acute myocardial infarction: an angiographic risk score for use in the catheterization laboratory. 2004 , 61, 338-43	3
1011	Myonecrosis following stent placement: association between impaired TIMI myocardial perfusion grade and MRI visualization of microinfarction. 2004 , 61, 472-6	41
1010	Increase of myocardial salvage and left ventricular function recovery with intracoronary abciximab downstream of the coronary occlusion in patients with acute myocardial infarction treated with primary coronary intervention. 2004 , 62, 186-92	76
1009	Assessment of the safety and efficacy of the novel tetrapeptide ITF-1697 on infarct size after primary PTCA in acute myocardial infarction: a randomised, placebo-controlled pilot trial. 2004 , 5, 141-51	2
1008	Comment ?valuer le retentissement fonctionnel d'une lésion coronaire, la qualité de la reperfusion coronaire et le résultat d'une angioplastie?. 2004 ,	
1007	Preprocedural TIMI flow and mortality in patients with acute myocardial infarction treated by primary angioplasty. 2004 , 43, 1363-7	116
1006	Early assessment of coronary flow reserve by transthoracic Doppler echocardiography predicts late remodeling in reperfused anterior myocardial infarction. 2004 , 17, 750-5	39
1005	Short- and long-term recovery of left ventricular function predicted at the time of primary percutaneous coronary intervention in anterior myocardial infarction. 2004 , 43, 534-41	74
1004	Frequency, correlates, and clinical implications of myocardial perfusion after primary angioplasty and stenting, with and without glycoprotein IIb/IIIa inhibition, in acute myocardial infarction. 2004 , 44, 305-12	150
1003	Achieved platelet aggregation inhibition after different antiplatelet regimens during percutaneous coronary intervention for ST-segment elevation myocardial infarction. 2004 , 44, 1187-93	80
1002	Prognostic utility of comparative methods for assessment of ST-segment resolution after primary angioplasty for acute myocardial infarction: the Controlled Abciximab and Device Investigation to Lower Late Angioplasty Complications (CADILLAC) trial. 2004 , 44, 1215-23	43

1001	Pressure response to vasopressors and mortality after direct angioplasty for cardiogenic shock. 2004 , 94, 197-202	6
1000	Single lead ST-segment recovery: a simple, reliable measure of successful fibrinolysis after acute myocardial infarction. 2004 , 147, 275-80	12
999	Sex-related differences in outcome after ST-segment elevation myocardial infarction treated by primary angioplasty: data from the Zwolle Myocardial Infarction study. 2004 , 148, 852-6	59
998	Relation between white blood cell counts and myocardial reperfusion in patients with recanalized anterior acute myocardial infarction. 2004 , 68, 526-31	13
997	Effects of a distal protection device during primary stenting in patients with acute anterior myocardial infarction. 2004 , 68, 763-8	39
996	NT-proBNP: a marker for successful myocardial reperfusion in AMI patients treated with primary percutaneous coronary intervention. 2004 , 6, 749-52	2
995	Prognostic utility of comparative methods for assessment of ST-segment resolution after primary angioplasty for acute myocardial infarction: The controlled abciximab and device investigation to lower late angioplasty complications (CADILLAC) trial. 2004 , 44, 1215-1223	139
994	Efficacy of primary PCI: the microvessel perspective. 2005 , 7, 14-19	3
993	Distal microcirculatory protection during percutaneous coronary intervention in acute ST-segment elevation myocardial infarction: a randomized controlled trial. 2005 , 293, 1063-72	438
992	Persistent hyperglycemia is associated with left ventricular dysfunction in patients with acute myocardial infarction. 2005 , 69, 23-8	20
991	Comparison of the effects of a distal embolic protection device and an aspiration catheter during percutaneous coronary intervention in patients with acute myocardial infarction. 2005 , 69, 49-54	16
990	Effects of primary percutaneous coronary intervention on P wave dispersion. 2005 , 10, 342-7	14
989	Impact of ST-segment depression resolution on mortality after successful mechanical reperfusion in patients with ST-segment elevation acute myocardial infarction. 2005 , 95, 234-6	11
988	Relation of myocardial perfusion to mortality after primary percutaneous coronary intervention. 2005 , 95, 980-2	5
987	Comparison of myocardial perfusion after successful primary percutaneous coronary intervention in patients with ST-elevation myocardial infarction with versus without diabetes mellitus. 2005 , 95, 1375-7	54
986	Absence of seasonal variation in myocardial perfusion, enzymatic infarct size, and mortality in patients with ST-segment elevation myocardial infarction treated with primary angioplasty. 2005 , 95, 1459-61	10
985	Relation of aspirin resistance to coronary flow reserve in patients undergoing elective percutaneous coronary intervention. 2005 , 96, 760-3	34
984	Angiographic evaluation of myocardial perfusion in patients with syndrome X. 2005 , 96, 803-5	20

983	Impact of beta-blocker therapy at discharge on long-term mortality after primary angioplasty for ST-segment elevation myocardial infarction. 2005 , 96, 806-9	16
982	Prediction of decrease in myocardial perfusion defect size and severity during a 3-month follow-up by the degree of acute resolution of electrocardiographic changes. 2005 , 38, 100-5	4
981	Six-month angiographic results of primary angioplasty with adjunctive PercuSurge GuardWire device support: evaluation of the restenotic rate of the target lesion and the fate of the distal balloon occlusion site. 2005 , 64, 35-42	12
980	Intracoronary ST segment evolution during primary coronary stenting predicts infarct zone recovery. 2005 , 64, 53-60	12
979	Facilitated primary coronary intervention with abciximab and very low dose of alteplase during off-hours compared with direct primary intervention during regular hours. 2005 , 65, 484-91	4
978	Use of abciximab prevents microcirculatory impairment in patients treated with coronary angioplasty for unstable angina: results of a prospective randomized study. 2005 , 66, 165-9	10
977	[Coronary microcirculation. Pathophysiology, clinical relevance, and importance for regenerative therapy after myocardial infarction]. 2005 , 30, 641-50	3
976	Coronary circulation and interventional cardiology. 2005 , 33, 1735-42	9
975	Current Assessments of the Adequacy of Myocardial Perfusion During Acute MI. 2005 , 7, 25-34	
974	Coronary clearance frame count: a new index of microvascular perfusion. 2005 , 19, 97-100	12
973	Preprocedural hs-CRP Level Serves as a Marker for Procedure-Related Myocardial Injury During Coronary Stenting. 2005 , 35, 140	4
972	Efficacy of Thrombosuction using the Export Aspiration Catheter before Primary Percutaneous Coronary Intervention in Acute Myocardial Infarction. 2005 , 35, 172	2
971	Systemic inflammation and reperfusion injury in patients with acute myocardial infarction. 2005 , 2005, 385-9	40
970	Relation between the TIMI frame count and the degree of microvascular injury after primary coronary angioplasty in patients with acute anterior myocardial infarction. 2005 , 91, 64-7	14
969	Randomized evaluation of the effects of filter-based distal protection on myocardial perfusion and infarct size after primary percutaneous catheter intervention in myocardial infarction with and without ST-segment elevation. <i>Circulation</i> , 2005 , 112, 1462-9	16.7 190
968	The ECG in acute coronary syndromes: new tricks from an old dog. 2005 , 91, 851-3	7
967	Effects of percutaneous coronary thrombectomy with the X-sizer catheter on epicardial flow and microvascular function in acute coronary syndromes. 2005 , 11, 461-6	
966	"Sea anemones" on a degenerated bicuspid aortic valve. 2005 , 91, 853	

965	Combined prognostic utility of ST-segment recovery and myocardial blush after primary percutaneous coronary intervention in acute myocardial infarction. 2005 , 26, 667-74		145
964	Distal protection improved reperfusion and reduced left ventricular dysfunction in patients with acute myocardial infarction who had angioscopically defined ruptured plaque. <i>Circulation</i> , 2005 , 112, 1001-7	16.7	86
963	Is routine stenting for acute myocardial infarction superior to balloon angioplasty? A randomised comparison in a large cohort of unselected patients. 2005 , 91, 641-5		52
962	Postconditioning the human heart. <i>Circulation</i> , 2005 , 112, 2143-8	16.7	831
961	Clinical implications of persistent ST segment depression after admission in patients with non-ST segment elevation acute coronary syndrome. 2005 , 91, 95-6		4
960	The effect of interventional treatment in acute myocardial infarction on ST resolution: a comparison of coronary angioplasty with excimer laser angioplasty. 2005 , 56, 377-84		10
959	Prognostic value of circulating levels of endothelin-1 in patients after acute myocardial infarction undergoing primary coronary angioplasty. 2005 , 127, 1491-7		62
958	ST-segment depression in lead aVR: a useful predictor of impaired myocardial reperfusion in patients with inferior acute myocardial infarction. 2005 , 128, 780-6		25
957	Ageing, impaired myocardial perfusion, and mortality in patients with ST-segment elevation myocardial infarction treated by primary angioplasty. 2005 , 26, 662-6		76
956	A quantitative analysis of the benefits of pre-hospital infarct angioplasty triage on outcome in patients undergoing primary angioplasty for acute myocardial infarction. 2005 , 7, K36-K40		2
955	A dynamic model forecasting myocardial infarct size before, during, and after reperfusion therapy: an ASSENT-2 ECG/VCG substudy. 2005 , 26, 1726-33		10
954	How should we evaluate an open artery in STEMI patients?. 2005 , 26, 634-6		5
953	Prediction of mortality after primary percutaneous coronary intervention for acute myocardial infarction: the CADILLAC risk score. 2005 , 45, 1397-405		367
952	Oxidative stress and inflammatory response during and following coronary interventions for acute myocardial infarction. 2005 , 39, 629-36		41
951	Angiographic evaluation of the effect of intracoronary abciximab administration in patients undergoing urgent PCI. 2005 , 105, 250-5		38
950	Thrombectomy prior to PCI may improve ST-segment resolution in acute myocardial infarction. Commentary. 2005 , 9, 314-7		
949	Glucose-insulin-potassium and reperfusion in acute myocardial infarction: rationale and design of the Glucose-Insulin-Potassium Study-2 (GIPS-2). 2005 , 149, 585-91		24
948	Levosimendan improves hemodynamics and coronary flow reserve after percutaneous coronary intervention in patients with acute myocardial infarction and left ventricular dysfunction. 2005 , 150, 563-8		51

947	Unsuccessful reperfusion in patients with ST-segment elevation myocardial infarction treated by primary angioplasty. 2005 , 150, 557-62	84
946	Preprocedural Thrombolysis in Myocardial Infarction (TIMI) flow significantly affects the extent of ST-segment resolution and myocardial blush in patients with acute anterior myocardial infarction treated by primary angioplasty. 2005 , 150, 827-31	14
945	Distal embolization during primary angioplasty: histopathologic features and predictability. 2005 , 150, 102-8	80
944	Prognostic significance of residual cumulative ST-segment deviation after mechanical reperfusion in patients with ST-segment elevation myocardial infarction. 2005 , 150, 1248-54	39
943	Circadian variation in myocardial perfusion and mortality in patients with ST-segment elevation myocardial infarction treated by primary angioplasty. 2005 , 150, 1185-9	45
942	ST-segment elevation resolution and outcome in patients treated with primary angioplasty and glucose-insulin-potassium infusion. 2005 , 149, 1135	11
941	Intervencionismo en el contexto del infarto de miocardio. Conceptos actuales. 2005 , 58, 567-584	7
940	Percutaneous Coronary Intervention in Myocardial Infarction. Current Concepts. 2005 , 58, 567-584	1
939	Impact of diabetes mellitus on myocardial perfusion after primary angioplasty in patients with acute myocardial infarction. 2005 , 45, 508-14	99
938	X-sizer for thrombectomy in acute myocardial infarction improves ST-segment resolution: results of the X-sizer in AMI for negligible embolization and optimal ST resolution (X AMINE ST) trial. 2005 , 46, 246-52	158
937	Manual thrombus-aspiration improves myocardial reperfusion: the randomized evaluation of the effect of mechanical reduction of distal embolization by thrombus-aspiration in primary and rescue angioplasty (REMEDIA) trial. 2005 , 46, 371-6	286
936	Myocardial perfusion in ST-elevation myocardial infarction treated successfully with primary angioplasty. 2006 , 40, 96-104	10
935	Association of C-reactive protein and myocardial perfusion in patients with ST-elevation acute myocardial infarction. 2006 , 186, 177-83	26
934	Impact of statin therapy at discharge on 1-year mortality in patients with ST-segment elevation myocardial infarction treated with primary angioplasty. 2006 , 189, 186-92	9
933	Electrocardiographic markers of reperfusion in ST-elevation myocardial infarction. 2006 , 24, 367-76, viii	12
932	Value of Coronary Blood Flow Pattern as a Predictor of Functional Recovery and Short-Term Left Ventricular Remodeling After Primary Coronary Angioplasty. A Transthoracic Doppler Study. 2006 , 59, 352-359	
931	Limitations of Myocardial Blush Grade in the Evaluation of Myocardial Perfusion in Patients With Acute Myocardial Infarction and TIMI Grade 3 Flow. 2006 , 59, 575-581	1
930	Coronary Angiography: Beyond Coronary Anatomy. 2006 , 59, 596-608	1

929	Valor del patrón de flujo coronario tras angioplastia primaria como predictor de recuperación funcional y remodelado ventricular a corto plazo. Estudio mediante ecocardiografía Doppler transtorácica. 2006 , 59, 352-359	0
928	Limitaciones del blush miocárdico para el análisis de la perfusión miocárdica en pacientes con infarto agudo de miocardio y flujo TIMI 3. 2006 , 59, 575-581	10
927	Coronariografía: más allá de la anatomía coronaria. 2006 , 59, 596-608	3
926	Value of platelet reactivity in predicting response to treatment and clinical outcome in patients undergoing primary coronary intervention: insights into the STRATEGY Study. 2006 , 48, 2178-85	125
925	Rheolytic thrombectomy with percutaneous coronary intervention for infarct size reduction in acute myocardial infarction: 30-day results from a multicenter randomized study. 2006 , 48, 244-52	222
924	Thrombus aspiration before primary angioplasty improves myocardial reperfusion in acute myocardial infarction: the DEAR-MI (Dethrombosis to Enhance Acute Reperfusion in Myocardial Infarction) study. 2006 , 48, 1552-9	199
923	Final myocardial blush grade predicts troponin I elevation in unstable angina patients undergoing percutaneous intervention. 2006 , 15, 291-6	
922	Successful reperfusion for acute ST elevation myocardial infarction is associated with a decrease in WBC count. 2006 , 147, 321-6	4
921	User-defined ROI tracking for estimation of the myocardial blush grade.	
920	Platelet microaggregation inhibition in patients with acute myocardial infarction pretreated with tirofiban and relationship with angiographic and clinical outcome. 2006 , 151, 1102-7	8
919	Sestamibi single photon emission computed tomography immediately after primary percutaneous coronary intervention identifies patients at risk for large infarcts. 2006 , 151, 1108-14	13
918	Relationship between infarct artery location, epicardial flow, and myocardial perfusion after primary percutaneous revascularization in acute myocardial infarction. 2006 , 151, 1288-95	42
917	Comparison of very early treatment with either fibrinolysis or percutaneous coronary intervention facilitated with abciximab with respect to ST recovery and infarct-related artery epicardial flow in patients with acute ST-segment elevation myocardial infarction: the Swedish Early Decision (SWEDES) reperfusion trial. 2006 , 151, 798.e1-7	27
916	Thrombus Aspiration during Percutaneous coronary intervention in Acute myocardial infarction Study (TAPAS)--study design. 2006 , 151, 597.e1-597.e7	32
915	Impact of routine stenting on myocardial perfusion and the extent of myocardial necrosis in patients undergoing primary angioplasty for ST-segment elevation myocardial infarction. 2006 , 151, 1296.e1-6	4
914	Feasibility and benefit of prehospital diagnosis, triage, and therapy by paramedics only in patients who are candidates for primary angioplasty for acute myocardial infarction. 2006 , 151, 1255.e1-5	40
913	Association between thrombolysis in myocardial infarction myocardial perfusion grade, biomarkers, and clinical outcomes among patients with moderate- to high-risk acute coronary syndromes: observations from the randomized trial to evaluate the relative PROTECTION against post-PCI microvascular dysfunction and post-PCI ischemia among antiplatelet and antithrombotic agents in myocardial infarction 30 (PROTECTION-30) (Percutaneous Coronary Intervention, 2006, 151, 1255.e1-5)	27
912	Intracoronary nitroprusside for the prevention of the no-reflow phenomenon after primary percutaneous coronary intervention in acute myocardial infarction. A randomized, double-blind, placebo-controlled clinical trial. 2006 , 152, 887.e9-14	87

911	Early abciximab administration in acute myocardial infarction treated with primary coronary intervention. 2006 , 108, 36-42	36
910	Balloon-Based Distal Protection Device Improves the Recovery of Left Ventricular Function after Primary Coronary Angioplasty. 2006 , 36, 84	
909	The influence of admission glucose on epicardial and microvascular flow after primary angioplasty. 2006 , 119, 95-102	7
908	Effect of preinfarction angina pectoris on myocardial blush grade after reperfusion in first anterior wall acute myocardial infarction. 2006 , 70, 698-702	3
907	Plain computed tomography for assessment of early coronary microcirculatory damage after revascularization therapy in acute myocardial infarction. 2006 , 70, 1475-80	3
906	Impact of PercuSurge device conjugative with intracoronary administration of nitroprusside on no-reflow phenomenon following primary percutaneous coronary intervention. 2006 , 70, 1538-42	18
905	The distal protection during primary percutaneous coronary intervention alleviates the adverse effects of large thrombus burden on myocardial reperfusion. 2006 , 70, 232-8	12
904	Full Length Article. 2006 , 48, 1552-1559	164
903	Impact of metabolic syndrome on myocardial perfusion grade after primary percutaneous coronary intervention in patients with acute ST elevation myocardial infarction. 2006 , 17, 339-43	14
902	The effect of verapamil on the restoration of myocardial perfusion and functional recovery in patients with angiographic no-reflow after primary percutaneous coronary intervention. 2006 , 27, 247-54	12
901	Impact of myocardial perfusion on left atrial remodeling following primary angioplasty for acute myocardial infarction. 2006 , 17, 597-603	9
900	Imaging microvascular obstruction and its clinical significance following acute myocardial infarction. 2006 , 11, 305-12	19
899	ST-segment analysis to predict infarct size and functional outcome in acute myocardial infarction treated with primary coronary intervention and adjunctive abciximab therapy. 2006 , 97, 48-54	25
898	Effect of reactive hyperemia after coronary recanalization on myocardial tissue reperfusion by thrombolysis in myocardial infarction flow grade in acute myocardial infarction. 2006 , 97, 617-23	7
897	Association between level of platelet inhibition after early use of abciximab and myocardial reperfusion in ST-elevation acute myocardial infarction treated by primary percutaneous coronary intervention. 2006 , 97, 798-803	8
896	Time course of microvascular resistance of the infarct and noninfarct coronary artery following an anterior wall acute myocardial infarction. 2006 , 97, 1131-6	47
895	Effect of coronary occlusion site on angiographic and clinical outcome in acute myocardial infarction patients treated with early coronary intervention. 2006 , 97, 1137-41	49
894	Effect of impaired myocardial reperfusion on left ventricular remodeling in patients with anterior wall acute myocardial infarction treated with primary coronary intervention. 2006 , 98, 725-8	37

893	Tratamiento coadyuvante en el intervencionismo coronario percutáneo del infarto agudo de miocardio. 2006 , 6, 39H-48H		
892	Electrocardiogram dynamics for risk stratification in ST-segment elevation myocardial infarction--immediate and serially updated information on outcome. 2006 , 39, S75-8		1
891	Management of acute myocardial infarction. 2006 , 34, 188-194		
890	A pilot study with a new, rapid-exchange, thrombus-aspirating device in patients with thrombus-containing lesions: the Diver C.E. study. 2006 , 67, 887-93		23
889	Successful thrombus extraction with the Rescue thrombus management system during acute percutaneous coronary intervention improves flow but does not necessarily restore optimal myocardial tissue perfusion. 2006 , 67, 879-86		4
888	Early administration of abciximab bolus in the emergency department improves angiographic outcome after primary PCI as assessed by TIMI frame count: results of the early ReoPro administration in myocardial infarction (ERAMI) trial. 2006 , 68, 218-24		37
887	Potential significance of spontaneous and interventional ST-changes in patients transferred for primary percutaneous coronary intervention: observations from the ST-MONitoring in Acute Myocardial Infarction study (The MONAMI study). 2006 , 27, 267-75		57
886	Significance of total and differential leucocyte count in patients with acute myocardial infarction treated with primary coronary angioplasty. 2006 , 27, 2511-5		57
885	Endothelin-1 and acute myocardial infarction: a no-reflow mediator after successful percutaneous myocardial revascularization. 2006 , 27, 1793-8		90
884	No-reflow: a heterogeneous clinical phenomenon with multiple therapeutic strategies. 2006 , 12, 3807-15		10
883	Predictors of elevated cardiac troponin T on admission in ST-segment elevation myocardial infarction. 2006 , 43, 281-6		8
882	Routine thrombectomy in percutaneous coronary intervention for acute ST-segment-elevation myocardial infarction: a randomized, controlled trial. <i>Circulation</i> , 2006 , 114, 40-7	16.7	216
881	Impact of intracoronary aspiration thrombectomy during primary angioplasty on left ventricular remodelling in patients with anterior ST elevation myocardial infarction. 2006 , 92, 951-7		121
880	Prospective, randomised, controlled trial to study the effect of intracoronary injection of verapamil and adenosine on coronary blood flow during percutaneous coronary intervention in patients with acute coronary syndromes. 2006 , 92, 1278-84		59
879	Impact of routine stenting on clinical outcome in diabetic patients undergoing primary angioplasty for ST-segment elevation myocardial infarction. 2006 , 29, 920-3		9
878	An Essential Guide to Cardiac Cell Therapy. 2006 ,		2
877	Distal embolic protection during percutaneous coronary intervention in patients with acute coronary syndromes: the RUBY study. 2006 , 8, 148-54		8
876	Hyperaemic microvascular resistance is not increased in viable myocardium after chronic myocardial infarction. 2007 , 28, 2320-5		30

875	Apical and midventricular transient left ventricular dysfunction syndrome (tako-tsubo cardiomyopathy): frequency, mechanisms, and prognosis. 2007 , 132, 809-16	437
874	Intracoronary streptokinase after primary percutaneous coronary intervention. 2007 , 356, 1823-34	132
873	Primary angioplasty vs. early routine post-fibrinolysis angioplasty for acute myocardial infarction with ST-segment elevation: the GRACIA-2 non-inferiority, randomized, controlled trial. 2007 , 28, 949-60	106
872	Impact of catheter sizes and intracoronary glyceryl trinitrate on the TIMI frame count when digital angiograms are acquired at lower frame rates during elective angiography and PCI. 2007 , 9, 231-8	6
871	Myocardial blush grade: to evaluate myocardial viability in patients with acute myocardial infarction. 2007 , 58, 556-60	1
870	Coronary blood flow in patients with cardiac syndrome X. 2007 , 18, 45-8	5
869	Predictors and clinical significance of angiographically detected distal embolization after primary percutaneous coronary interventions. 2007 , 18, 443-9	16
868	Temporal changes in TIMI myocardial perfusion grade in relation to epicardial flow, ST-resolution and left ventricular function after primary percutaneous coronary intervention. 2007 , 18, 513-8	6
867	Primary angioplasty and routine utilization of thrombus aspiration devices: feasibility and results in a consecutive series of 486 patients. 2007 , 8, 258-64	11
866	Plasma asymmetric dimethylarginine and L-arginine levels in patients with cardiac syndrome X. 2007 , 18, 539-44	17
865	Adjunctive mechanical devices to prevent distal embolization in patients undergoing mechanical revascularization for acute myocardial infarction: a meta-analysis of randomized trials. 2007 , 153, 343-53	117
864	The Outcome of PCI for stent-Thrombosis Multicentre Study (OPTIMIST): rationale and design of a multicenter registry. 2007 , 153, 377.e1-5	7
863	Quantitative evaluation of myocardial blush to assess tissue level reperfusion in patients with acute ST-elevation myocardial infarction: incremental prognostic value compared with visual assessment. 2007 , 153, 612-20	28
862	Effects of balloon-based distal protection during primary percutaneous coronary intervention on early and late infarct size and left ventricular remodeling: a pilot study using serial contrast-enhanced magnetic resonance imaging. 2007 , 153, 665.e1-8	13
861	Baseline platelet reactivity in acute myocardial infarction treated with primary angioplasty--influence on myocardial reperfusion, left ventricular performance, and clinical events. 2007 , 154, 62-70	23
860	Impact of collateral flow on myocardial reperfusion and infarct size in patients undergoing primary angioplasty for acute myocardial infarction. 2007 , 154, 379-84	28
859	Coronary angiography for follow-up of heart transplant recipients: insights from TIMI frame count and TIMI myocardial perfusion grade. 2007 , 26, 593-7	10
858	Effect of microvascular reperfusion on prognosis and left ventricular function in anterior wall myocardial infarction treated with primary angioplasty. 2007 , 114, 183-7	13

857	Excimer laser thrombus elimination for prevention of distal embolization and no-reflow in patients with acute ST elevation myocardial infarction: results from the randomized LaserAMI study. 2007 , 116, 20-6	40
856	Combination of electrocardiographic and angiographic markers of reperfusion in the prediction of infarct size in patients with ST-segment elevation myocardial infarction undergoing successful primary angioplasty. 2007 , 117, 232-7	16
855	Comparison between stenting and balloon in elderly patients undergoing primary angioplasty for ST-segment elevation myocardial infarction. 2007 , 119, 306-9	12
854	Nitric oxide inhalation improves microvascular flow and decreases infarction size after myocardial ischemia and reperfusion. 2007 , 50, 808-17	97
853	Export aspiration system in patients with acute coronary syndrome and visible thrombus provides no substantial benefit. 2007 , 70, 35-42	9
852	Significance of additional ST segment elevation in patients with no reflow after angioplasty for acute myocardial infarction. 2007 , 20, 262-9	1
851	Baseline systemic inflammatory status and no-reflow phenomenon after percutaneous coronary angioplasty for acute myocardial infarction. 2007 , 117, 306-11	39
850	Randomized early versus late abciximab in acute myocardial infarction treated with primary coronary intervention (RELAX-AMI Trial). 2007 , 49, 1517-24	90
849	Adjunctive thrombectomy in acute myocardial infarction: for some but not for all. 2007 , 49, 1586; author reply 1586-7	2
848	Reply. 2007 , 49, 1586-1587	
847	Angiographic stent thrombosis after routine use of drug-eluting stents in ST-segment elevation myocardial infarction: the importance of thrombus burden. 2007 , 50, 573-83	360
846	Impact of plaque rupture on infarct size in ST-segment elevation anterior acute myocardial infarction. 2007 , 50, 1230-7	64
845	Sex Differences of the Clinical Characteristics and Early Management in the Korea Acute Myocardial Infarction Registry. 2007 , 37, 64	16
844	Myocardial blush grade: a predictor for major adverse cardiac events after primary PTCA with stent implantation for acute myocardial infarction. 2007 , 62, 445-51	13
843	Thrombosuction utilizing an export aspiration catheter during primary percutaneous coronary intervention in acute myocardial infarction. 2007 , 48, 261-9	1
842	Selective intracoronary administration of nitroprusside before balloon dilatation prevents slow reflow during percutaneous coronary intervention in patients with acute myocardial infarction. 2007 , 48, 423-33	28
841	Percutaneous coronary arterial thrombectomy for acute myocardial infarction reduces no-reflow phenomenon and protects against left ventricular remodeling related to the proximal left anterior descending and right coronary artery. 2007 , 48, 287-302	10
840	High dose adenosine for suboptimal myocardial reperfusion after primary PCI: A randomized placebo-controlled pilot study. 2008 , 71, 283-9	61

839	Time-dependent benefit of initial thrombosuction on myocardial reperfusion in primary percutaneous coronary intervention. 2008 , 62, 555-61	42
838	Thrombus aspiration with export catheter in ST elevation myocardial infarction. 2007 , 20, 38-43	10
837	Prospective, multicenter study of thrombectomy in patients with acute myocardial infarction: the X-Tract AMI registry. 2007 , 20, 44-50	14
836	Intracoronary thrombectomy with the export aspiration catheter before angioplasty in patients with ST-segment elevation myocardial infarction. 2007 , 20, 136-42	5
835	Rheolytic thrombectomy during percutaneous coronary intervention improves long-term outcome in high-risk patients with acute myocardial infarction. 2007 , 20, 292-8	21
834	Relationship between microvascular resistance and perfusion in patients with reperfused acute myocardial infarction. 2007 , 20, 340-50	10
833	Angiographic no-reflow phenomenon and plaque characteristics by virtual histology intravascular ultrasound in patients with acute myocardial infarction. 2007 , 20, 335-9	34
832	Angiographic and clinical characteristics associated with the removable plaque components by means of thrombectomy catheters in patients with myocardial infarction. 2007 , 8, 236-42	2
831	Safety and efficacy of the FilterWire EZ in acute ST-segment elevation myocardial infarction. 2007 , 99, 911-5	35
830	Relation between leucocyte count, myonecrosis, myocardial perfusion, and outcomes following primary angioplasty. 2007 , 99, 1067-71	28
829	Relation of myocardial blush grade to microvascular perfusion and myocardial infarct size after primary or rescue percutaneous coronary intervention. 2007 , 99, 1671-3	44
828	[Intracoronary thrombectomy with the Export Aspiration Catheter before angioplasty in patients with ST-segment elevation myocardial infarction]. 2007 , 56, 48-53	3
827	The value of both ST-segment and QRS complex changes during acute coronary occlusion for prediction of reperfusion-induced myocardial salvage in a canine model. 2007 , 40, 18-25	29
826	Effect of a distal protection device on epicardial blood flow and myocardial perfusion in primary percutaneous coronary intervention. 2007 , 8, 575-9	5
825	Renal frame count and renal blush grade: quantitative measures that predict the success of renal stenting in hypertensive patients with renal artery stenosis. 2008 , 1, 286-92	34
824	A comparison of 2 thrombus aspiration devices with histopathological analysis of retrieved material in patients presenting with ST-segment elevation myocardial infarction. 2008 , 1, 258-64	31
823	Upfront thrombus aspiration in primary coronary intervention for patients with ST-segment elevation acute myocardial infarction: report of the VAMPIRE (VAcuum asPIration thrombus REmoval) trial. 2008 , 1, 424-31	112
822	The effects of aspirin and clopidogrel response on myonecrosis after percutaneous coronary intervention: a BRIEF-PCI (Brief Infusion of Intravenous Eptifibatide Following Successful Percutaneous Coronary Intervention) trial substudy. 2008 , 1, 654-9	10

821	Angiographic core laboratory reproducibility analyses: implications for planning clinical trials using coronary angiography and left ventriculography end-points. 2008 , 24, 453-62	10
820	Estimate of myocardial salvage in late presentation acute myocardial infarction by comparing functional and perfusion abnormalities in predischARGE gated SPECT. 2008 , 35, 906-11	8
819	Early intervention: which patients and how early?. 2008 , 10, 303-11	2
818	Ultrasound enhanced prehospital thrombolysis using microbubbles infusion in patients with acute ST elevation myocardial infarction: rationale and design of the Sonolysis study. 2008 , 9, 72	8
817	The effect of thrombectomy on myocardial blush in primary angioplasty: the Randomized Evaluation of Thrombus Aspiration by two thrombectomy devices in acute Myocardial Infarction (RETAMI) trial. 2008 , 71, 84-91	34
816	Feasibility of primary clot extraction prior to percutaneous coronary intervention in acute myocardial infarction. 2008 , 71, 870-6	2
815	Concordant improvements in coronary flow reserve and ST-segment resolution during percutaneous coronary intervention for acute myocardial infarction: a benefit of postconditioning. 2008 , 72, 212-20	95
814	Adjunctive use of the Rinspiration system for fluidic thrombectomy during primary angioplasty: the Rinspiration international registry. 2008 , 72, 196-203	10
813	The feasibility and safety of routine thrombus aspiration in patients with non-ST-elevation myocardial infarction. 2008 , 72, 937-42	25
812	Distribution and determinants of myocardial perfusion grade following late mechanical recanalization of occluded infarct-related arteries postmyocardial infarction: a report from the occluded artery trial. 2008 , 72, 783-9	5
811	Clinical implications of distal embolization during coronary interventional procedures in patients with acute myocardial infarction: quantitative study with Doppler guidewire. 2008 , 1, 268-76	38
810	'No-reflow' after acute myocardial infarction: direct visualisation of microvascular obstruction by gadolinium-enhanced CMR. 2008 , 16, 179-81	27
809	Direct stenting after thrombus removal before primary angioplasty in acute myocardial infarction. 2008 , 21, 300-6	12
808	Myocardial perfusion characterization from contrast angiography spectral distribution. 2008 , 27, 641-9	12
807	Effect of statin therapy before Q-wave myocardial infarction on myocardial perfusion. 2008 , 101, 139-43	17
806	Comparison of usefulness of C-reactive protein versus white blood cell count to predict outcome after primary percutaneous coronary intervention for ST elevation myocardial infarction. 2008 , 101, 446-51	32
805	Relation of left ventricular infarct transmuralitY and infarct size after primary percutaneous coronary angioplasty to time from symptom onset to balloon inflation. 2008 , 102, 1163-9	18
804	Reperfusion strategies in acute ST-elevation myocardial infarction: an overview of current status. 2008 , 50, 352-82	69

803 Assessment and management of acute coronary syndrome. **2008**, 4, 99-105

802 Antithrombotic therapies in primary angioplasty: rationale, results and future directions. **2008**, 68, 2325-44 6

801 [Analysis of mortality in myocardial infarction patients treated with primary angioplasty]. **2008**, 131, 481-6

800 Effect of Persistent Microvascular Obstruction on Post-infarction Ventricular Remodeling Following Intracoronary Bone-Marrow Cell Transplantation: A Contrast-Enhanced Cardiac Magnetic Resonance Study. **2008**, 61, 602-610

799 The extent of microvascular damage during myocardial contrast echocardiography is superior to other known indexes of post-infarct reperfusion in predicting left ventricular remodeling: results of the multicenter AMICI study. **2008**, 51, 552-9 167

798 Reply. **2008**, 51, 1324-1325

797 ST-segment elevation myocardial infarction due to early and late stent thrombosis a new group of high-risk patients. **2008**, 51, 2396-402 87

796 Functional recovery after acute myocardial infarction: comparison between angiography, electrocardiography, and cardiovascular magnetic resonance measures of microvascular injury. **2008**, 52, 181-9 279

795 Reperfusion injury in acute myocardial infarction: from bench to cath lab. Part II: Clinical issues and therapeutic options. **2008**, 101, 565-75 28

794 Angiographic and physiologic assessment of coronary flow and myocardial perfusion in the cardiac catheterization laboratory. **2008**, 10, 69-78 3

793 Thrombus aspiration during primary percutaneous coronary intervention. **2008**, 358, 557-67 796

792 Adjunctive devices in primary or rescue PCI: a meta-analysis of randomized trials. **2008**, 123, 313-21 61

791 Distal protection device protects microvascular integrity during primary percutaneous intervention in acute myocardial infarction: a prospective, randomized, multicenter trial. **2008**, 123, 162-8 19

790 Is it possible to predict which patients need distal protection during primary angioplasty?. **2008**, 127, 179-85 9

789 Gender differences of success rate of percutaneous coronary intervention and short term cardiac events in Korea Acute Myocardial Infarction Registry. **2008**, 130, 227-34 47

788 Excimer laser in acute myocardial infarction: single centre experience on 66 patients. **2008**, 127, 98-102 33

787 Contemporary outcomes of rescue percutaneous coronary intervention for acute myocardial infarction: comparison with primary angioplasty and the role of distal protection devices (EMERALD trial). **2008**, 155, 1090-6 13

786 Change of white blood cell count more prognostic important than baseline values after primary percutaneous coronary intervention for ST elevation myocardial infarction. **2008**, 122, 185-9 5

785	von Willebrand factor in cardiovascular disease: focus on acute coronary syndromes. <i>Circulation</i> , 2008 , 117, 1449-59	16.7	294
784	Impacto de la obstrucción microvascular persistente en el remodelado ventricular postinfarto tras el implante intracoronario de células mononucleadas de médula ósea: un estudio de cardi resonancia con contraste. 2008 , 61, 602-610		3
783	Cardiac death and reinfarction after 1 year in the Thrombus Aspiration during Percutaneous coronary intervention in Acute myocardial infarction Study (TAPAS): a 1-year follow-up study. 2008 , 371, 1915-20		638
782	Prehospital initiation of tirofiban in patients with ST-elevation myocardial infarction undergoing primary angioplasty (On-TIME 2): a multicentre, double-blind, randomised controlled trial. 2008 , 372, 537-46		345
781	ST-segment recovery and outcome after primary percutaneous coronary intervention for ST-elevation myocardial infarction: insights from the Assessment of Pexelizumab in Acute Myocardial Infarction (APEX-AMI) trial. <i>Circulation</i> , 2008 , 118, 1335-46	16.7	121
780	Myocardial perfusion assessed by contrast echocardiography correlates with angiographic perfusion parameters in patients with a first acute myocardial infarction successfully treated with angioplasty. 2008 , 24, 633-9		16
779	Relationship between collateral circulation and successful myocardial reperfusion in acute myocardial infarction: a subanalysis of the PREMIAR trial. 2008 , 59, 587-92		9
778	Primary Percutaneous Coronary Intervention. 2008 , 91-117		
777	Myocyte Protection by Device Therapy. 2008 , 248-264		
776	Adjunctive antithrombotic therapy during primary percutaneous coronary intervention. 2008 , 10, J2-J14		
775	Adverse outcomes in fibrinolytic-based facilitated percutaneous coronary intervention: insights from the ASSENT-4 PCI electrocardiographic substudy. 2008 , 29, 871-9		25
774	Reperfusion after primary angioplasty for ST-elevation myocardial infarction: predictors of success and relationship to clinical outcomes in the APEX-AMI angiographic study. 2008 , 29, 1127-35		42
773	Plasma levels of thromboxane A2 on admission are associated with no-reflow after primary percutaneous coronary intervention. 2008 , 29, 1843-50		58
772	Randomized comparison between tirofiban and abciximab to promote complete ST-resolution in primary angioplasty: results of the facilitated angioplasty with tirofiban or abciximab (FATA) in ST-elevation myocardial infarction trial. 2008 , 29, 2972-80		35
771	Reperfusion ventricular arrhythmia 'bursts' predict larger infarct size despite TIMI 3 flow restoration with primary angioplasty for anterior ST-elevation myocardial infarction. 2009 , 30, 757-64		33
770	Early glycoprotein IIb/IIIa inhibitors in primary angioplasty (EGYPT) cooperation: an individual patient data meta-analysis. 2008 , 94, 1548-58		107
769	Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation: the Task Force on the Management of ST-Segment Elevation Acute Myocardial Infarction of the European Society of Cardiology. 2008 , 29, 2909-45		1774
768	Postprocedural single-lead ST-segment deviation and long-term mortality in patients with ST-segment elevation myocardial infarction treated by primary angioplasty. 2008 , 94, 44-7		15

767	Beneficial effect of post-procedural abciximab in patients undergoing primary coronary angioplasty and presenting with the no-reflow phenomenon. 2008 , 10, 100-3	2
766	Role of adjunctive thrombectomy and embolic protection devices in acute myocardial infarction: a comprehensive meta-analysis of randomized trials. 2008 , 29, 2989-3001	193
765	Adjunctive manual thrombectomy improves myocardial perfusion and mortality in patients undergoing primary percutaneous coronary intervention for ST-elevation myocardial infarction: a meta-analysis of randomized trials. 2008 , 29, 3002-10	194
764	Predictors and time-related impact of distal embolization during primary angioplasty. 2009 , 30, 305-13	71
763	Incidence and clinical consequences of distal embolization on the coronary angiogram after percutaneous coronary intervention for ST-elevation myocardial infarction. 2009 , 30, 908-15	108
762	Angiographic and clinical outcome of invasively managed patients with thrombosed coronary bare metal or drug-eluting stents: the OPTIMIST study. 2008 , 29, 3011-21	37
761	No-reflow: the next challenge in treatment of ST-elevation acute myocardial infarction. 2008 , 29, 1795-7	34
760	Early validation study of 64-slice multidetector computed tomography for the assessment of myocardial viability and the prediction of left ventricular remodelling after acute myocardial infarction. 2008 , 29, 490-8	51
759	Relationship of admission QRS duration and changes in QRS duration with myocardial reperfusion in patients with acute ST segment elevation myocardial infarction (STEMI) treated with fibrinolytic therapy. 2008 , 72, 873-9	7
758	Correlation of serial cardiac magnetic resonance imaging parameters with early resolution of ST-segment elevation after primary percutaneous coronary intervention. 2008 , 72, 1621-6	24
757	Evaluation of myocardial perfusion and deformation in patients with acute myocardial infarction treated with primary angioplasty and stent placement. 2008 , 19, 497-506	9
756	Circulating CD34-positive cell number is related to effective myocardial reperfusion in acute myocardial infarction treated with primary coronary angioplasty. 2008 , 9, 677-82	2
755	The no-reflow phenomenon in acute myocardial infarction after primary angioplasty: incidence, predictive factors, and long-term outcomes. 2008 , 9, 59-63	16
754	Total blush score: a new index for the assessment of microvascular perfusion in idiopathic dilated cardiomyopathy. 2008 , 19, 181-5	7
753	PCI in the presence of significant intraluminal thrombus. 2008 , 177-186	
752	Acute Coronary Syndromes and Acute Myocardial Infarction. 2008 , 589-646	
751	Individual patient-data meta-analysis comparing clinical outcome in patients with ST-elevation myocardial infarction treated with percutaneous coronary intervention with or without prior thrombectomy. ATTEMPT study: a pooled Analysis of Trials on ThrombEctomy in acute Myocardial infarction based on individual Patient data. 2009 , 5, 243-7	15
750	The success of primary angioplasty: beyond TIMI flow summary of a presentation held at acute cardiac care meeting in Paris, October 2008. 2009 , 11, 66-8	1

749	Early stent thrombosis in patients with acute coronary syndromes treated with drug-eluting and bare metal stents: the Acute Catheterization and Urgent Intervention Triage Strategy trial. <i>Circulation</i> , 2009 , 119, 687-98	16.7	154
748	Computer-assisted myocardial blush quantification after percutaneous coronary angioplasty for acute myocardial infarction: a substudy from the TAPAS trial. 2009 , 30, 594-9		49
747	Obstructive sleep apnea in patients admitted for acute myocardial infarction. Prevalence, predictors, and effect on microvascular perfusion. 2009 , 135, 1488-1495		114
746	Comparison of two visual angiographic perfusion grades in acute myocardial infarction. 2009 , 114, 149-53		3
745	Assessment of microvascular obstruction and prediction of short-term remodeling after acute myocardial infarction: cardiac MR imaging study. 2009 , 250, 363-70		98
744	Pathophysiology of the no-reflow phenomenon. 2009 , 11, 69-76		26
743	Role of thrombectomy in acute myocardial infarction. 2009 , 7, 289-97		
742	Total absence of ST-segment resolution after failed thrombolysis is correlated with unfavorable short- and long-term outcomes despite successful rescue angioplasty. 2009 , 42, 73-8		5
741	Predictive value of Q waves on the 12-lead electrocardiogram after reperfusion therapy for ST elevation myocardial infarction. 2009 , 42, 310-8		4
740	Impact of acute hyperglycemia during primary stent implantation in patients with ST-elevation myocardial infarction. 2009 , 53, 272-7		30
739	Intravenous administration of adenosine triphosphate disodium during primary percutaneous coronary intervention attenuates the transient rapid improvement of myocardial wall motion, not myocardial stunning, shortly after recanalization in acute anterior myocardial infarction. 2009 , 54, 289-96		6
738	Posterior wall involvement attenuates predictive value of ST-segment elevation in lead V4R for right ventricular involvement in inferior acute myocardial infarction. 2009 , 54, 386-93		10
737	Temporal course of pregnancy-associated plasma protein-A in angioplasty-treated ST-elevation myocardial infarction patients and potential significance of concomitant heparin administration. 2009 , 103, 29-35		40
736	Prophylactic use of manual thrombectomy in ST-segment elevation myocardial infarction. 2009 , 10, 224-8		
735	Myocardium selective densitometric perfusion assessment after acute myocardial infarction. 2009 , 10, 49-54		5
734	Incidence, predictors, and outcome of reinfarction and stent thrombosis within one year after primary percutaneous coronary intervention for ST-elevation myocardial infarction. 2009 , 73, 627-34		10
733	Comparison of the angiographic myocardial blush grade with delayed-enhanced cardiac magnetic resonance for the assessment of microvascular obstruction in acute myocardial infarctions. 2009 , 74, 1000-7		29
732	Microvascular obstruction after percutaneous coronary intervention. 2010 , 75, 369-77		10

731	Vessel masking improves densitometric myocardial perfusion assessment. 2009 , 25, 229-36	8
730	Impact of vessel size on distal embolization, myocardial perfusion and clinical outcome in patients undergoing primary angioplasty for ST-segment elevation myocardial infarction. 2009 , 27, 198-203	11
729	Benefits of pharmacological facilitation with glycoprotein IIb-IIIa inhibitors in diabetic patients undergoing primary angioplasty for STEMI. A subanalysis of the EGYPT cooperation. 2009 , 28, 288-98	11
728	Scintigraphic evaluation of routine filterwire distal protection in percutaneous coronary intervention for acute ST-segment elevation myocardial infarction: a randomized controlled trial. 2009 , 16, 784-91	10
727	Appearance of microvascular obstruction on high resolution first-pass perfusion, early and late gadolinium enhancement CMR in patients with acute myocardial infarction. 2009 , 11, 33	71
726	Intracoronary versus intravenous abciximab in ST-segment elevation myocardial infarction: rationale and design of the CICERO trial in patients undergoing primary percutaneous coronary intervention with thrombus aspiration. 2009 , 10, 90	9
725	Angiographic quantification of thrombus in ST-elevation acute myocardial infarction presenting with an occluded infarct-related artery and its relationship with results of percutaneous intervention. 2009 , 22, 207-15	17
724	Relation between door-to-balloon time and microvascular perfusion as evaluated by myocardial blush grade, corrected TIMI frame count, and ST-segment resolution in treatment of acute myocardial infarction. 2009 , 22, 437-43	8
723	Randomized comparison of primary percutaneous coronary intervention with combined proximal embolic protection and thrombus aspiration versus primary percutaneous coronary intervention alone in ST-segment elevation myocardial infarction: the PREPARE (PROximal Embolic Protection in Acute myocardial infarction and Resolution of ST-Elevation) study. 2009 , 2, 934-43	42
722	Thrombus Aspiration during Percutaneous coronary intervention in Acute non-ST-elevation myocardial infarction Study (TAPAS II)-Study design. 2009 , 17, 409-13	11
721	Upstream high-dose tirofiban does not reduce myocardial infarct size in patients undergoing primary percutaneous coronary intervention: a magnetic resonance imaging pilot study. 2009 , 32, 321-6	14
720	Therapeutic effect of urapidil on myocardial perfusion in patients with ST-elevation acute coronary syndrome. 2009 , 20, 152-7	11
719	Effect of thrombus aspiration on infarct size and left ventricular function in high-risk patients with acute myocardial infarction treated by percutaneous coronary intervention. Results of a prospective controlled pilot study. 2009 , 157, 583.e1-7	34
718	Management of ST-segment elevation myocardial infarction: Comparison of the updated guidelines from North America and Europe. 2009 , 158, 695-705	12
717	Contribution of angiographic and electrocardiographic parameters of reperfusion to prediction of mortality and morbidity after acute ST-elevation myocardial infarction: Insights from the Assessment of Pexelizumab in Acute Myocardial Infarction trial. 2009 , 158, 755-60	5
716	Thrombus aspiration during primary percutaneous coronary intervention improves myocardial reperfusion and reduces infarct size: the EXPIRA (thrombectomy with export catheter in infarct-related artery during primary percutaneous coronary intervention) prospective, randomized trial. 2009 , 53, 309-15	287
715	Myocardial no-reflow in humans. 2009 , 54, 281-92	569
714	Intracoronary injection of in situ forming alginate hydrogel reverses left ventricular remodeling after myocardial infarction in Swine. 2009 , 54, 1014-23	263

713	Early assessment of myocardial viability by the use of delayed enhancement computed tomography after primary percutaneous coronary intervention. 2009 , 2, 1072-81	33
712	Early electrocardiographic findings and MR imaging-verified microvascular injury and myocardial infarct size. 2009 , 2, 1187-94	39
711	Toward Robust Myocardial Blush Grade Estimation in Contrast Angiography. 2009 , 249-256	1
710	Role of ischemia modified albumin to ST-segment resolution after mechanical reperfusion in patients with ST-segment elevation myocardial infarction. 2009 , 203, 576-80	6
709	Intracoronary versus intravenous abciximab administration in patients with ST-elevation myocardial infarction undergoing thrombus aspiration during primary percutaneous coronary intervention--effects on soluble CD40 ligand concentrations. 2009 , 206, 523-7	33
708	Diabetes mellitus is associated with distal embolization, impaired myocardial perfusion, and higher mortality in patients with ST-segment elevation myocardial infarction treated with primary angioplasty and glycoprotein IIb-IIIa inhibitors. 2009 , 207, 181-5	77
707	[Not Available]. 2009 , 62, 293.e1-293.e47	5
706	Microvascular damage prevention with thromb aspiration during primary percutaneous intervention in acute myocardial infarction. 2009 , 20, 51-7	2
705	Glycoprotein IIB/IIIA inhibitor to reduce postpercutaneous coronary intervention myonecrosis and improve coronary flow in diabetics: the 'OPTIMIZE-IT' pilot randomized study. 2009 , 10, 245-51	16
704	Randomized evaluation of intracoronary nitroprusside vs. adenosine after thrombus aspiration during primary percutaneous coronary intervention for the prevention of no-reflow in acute myocardial infarction: the REOPEN-AMI study protocol. 2009 , 10, 585-92	8
703	Myocardial abnormalities underlying persistent ST-segment elevation after anterior myocardial infarction. 2009 , 10, 44-50	9
702	Significant association between neutrophil aggregation in aspirated thrombus and myocardial damage in patients with ST-segment elevation acute myocardial infarction. 2009 , 73, 139-44	34
701	Clinical impact of thrombectomy in acute ST-elevation myocardial infarction: an individual patient-data pooled analysis of 11 trials. 2009 , 30, 2193-203	201
700	Acute myocardial infarction: the enduring challenge for cardiac protection and survival. 2009 , 73, 2000-8	53
699	Cathétisme cardiaque, angiocardigraphie et coronarographie. Technique et résultats. 2009 , 4, 1-28	0
698	NT pro-B-type natriuretic peptide levels are related to microvascular reperfusion in patients undergoing direct percutaneous transluminal coronary angioplasty for anterior ST-segment elevation myocardial infarction. 2010 , 11, 359-64	2
697	Predictive value of tissue Doppler imaging for left ventricular ejection fraction, remodelling, and infarct size after percutaneous coronary intervention for acute myocardial infarction. 2010 , 11, 596-601	11
696	Early and aggressive treatment of patients with ST-segment elevation myocardial infarction: deciphering recent clinical trials and the timing of optimal platelet inhibition. 2010 , 12, D24-D35	

695	The first clinical pilot study of intravenous adrenomedullin administration in patients with acute myocardial infarction. 2010 , 56, 413-9	54
694	Assessment of coronary artery flow velocity pattern as a long-term predictor of left ventricular function and cardiac events after percutaneous coronary intervention in anterior acute myocardial infarction. 2010 , 49, 1693-701	3
693	Rheolytic thrombectomy: any role left?. 2010 , 2, 57-65	2
692	Predictors of ventricular tachyarrhythmia in high-risk myocardial infarction patients treated with primary coronary intervention. 2010 , 18, 122-8	7
691	Thrombus aspiration in a series of patients with stable or unstable angina pectoris and lesion-site thrombus formation. 2010 , 18, 423-9	6
690	Proximal embolic protection in patients undergoing primary angioplasty for acute myocardial infarction (PREPARE): core lab adjudicated angiographic outcomes of a randomised controlled trial. 2010 , 18, 531-6	3
689	Primary percutaneous coronary interventions in nonagenarians. 2010 , 33, 157-61	11
688	Comparison of immediate vs early invasive strategy in patients with first acute non-ST-elevation myocardial infarction. 2010 , 33, 650-655	16
687	Manual versus nonmanual thrombectomy in primary and rescue percutaneous coronary angioplasty. 2010 , 25, 275-81	10
686	Noninvasive assessment of myocardial damage after acute anterior myocardial infarction: myocardial blush grade in conjunction with analysis of coronary flow pattern. 2010 , 25, 299-305	3
685	Feasibility of an accurate assessment of myocardial salvage by comparing functional and perfusion abnormalities in post-reperfusion gated SPECT. 2010 , 17, 825-30	3
684	Thrombectomy during primary angioplasty: methods, devices, and clinical trial data. 2010 , 12, 422-8	6
683	Distal protection beneficial?. 2010 , 26, 135-8	1
682	Impact of distal embolization on myocardial perfusion and survival among patients undergoing primary angioplasty with glycoprotein IIb-IIIa inhibitors: insights from the EGYPT cooperation. 2010 , 30, 23-8	11
681	Gender-related differences in outcome after ST-segment elevation myocardial infarction treated by primary angioplasty and glycoprotein IIb-IIIa inhibitors: insights from the EGYPT cooperation. 2010 , 30, 342-6	31
680	Impact of early abciximab administration on myocardial reperfusion in patients with ST-segment elevation myocardial infarction pretreated with 600 mg of clopidogrel before percutaneous coronary intervention. 2010 , 30, 347-53	6
679	Efficacy of thrombectomy for acute myocardial infarction-Special focus on its efficacy according to different infarct-related arteries. 2010 , 55, 189-95	6
678	Predictive value of ST resolution analysis performed immediately versus at ninety minutes after primary percutaneous coronary intervention. 2010 , 105, 467-74	7

677	Relation of elevated levels of plasma myeloperoxidase to impaired myocardial microcirculation after reperfusion in patients with acute myocardial infarction. 2010 , 105, 922-9	20
676	Intracoronary compared to intravenous Abciximab and high-dose bolus compared to standard dose in patients with ST-segment elevation myocardial infarction undergoing transradial primary percutaneous coronary intervention: a two-by-two factorial placebo-controlled randomized study. 2010 , 105, 1520-7	43
675	Angiographic predictors of recurrent stent thrombosis (from the Outcome of PCI for stent-Thrombosis Multicentre Study [OPTIMIST]). 2010 , 105, 1710-5	9
674	Impact of thrombectomy with EXPort Catheter in Infarct-Related Artery during Primary Percutaneous Coronary Intervention (EXPIRA Trial) on cardiac death. 2010 , 106, 624-9	45
673	Comparison of 600 versus 300-mg Clopidogrel loading dose in patients with ST-segment elevation myocardial infarction undergoing primary coronary angioplasty. 2010 , 106, 1208-11	47
672	Circulating cytochrome c as potential biomarker of impaired reperfusion in ST-segment elevation acute myocardial infarction. 2010 , 106, 1443-9	12
671	Usefulness of intraplatelet melatonin levels to predict angiographic no-reflow after primary percutaneous coronary intervention in patients with ST-segment elevation myocardial infarction. 2010 , 106, 1540-4	13
670	Impact of thrombus aspiration use and direct stenting on final myocardial blush score in patients presenting with ST-elevation myocardial infarction. 2010 , 11, 149-54	3
669	Multicentre experience with MGuard net protective stent in ST-elevation myocardial infarction: safety, feasibility, and impact on myocardial reperfusion. 2010 , 75, 715-21	25
668	TIMI myocardial perfusion frame count: a new method to assess myocardial perfusion and its predictive value for short-term prognosis. 2010 , 75, 722-32	28
667	Feasibility and applicability of computer-assisted myocardial blush quantification after primary percutaneous coronary intervention for ST-segment elevation myocardial infarction. 2010 , 75, 701-6	13
666	Impact of abciximab on mortality and reinfarction in patients with acute ST-segment elevation myocardial infarction treated with primary stenting. 2010 , 75, 895-902	9
665	Thrombus aspiration in primary percutaneous coronary intervention in high-risk patients with ST-elevation myocardial infarction: a real-world registry. 2010 , 76, 70-6	27
664	Early assessment of infarct size and prediction of functional recovery by quantitative myocardial blush grade in patients with acute coronary syndromes treated according to current guidelines. 2010 , 76, 502-10	9
663	Rebuttal: TIMI myocardial perfusion frame count: an angiographic method to assess flow in the myocardium but not in the epicardial artery. 2010 , 76, 1073-5	1
662	Efficacy of high-dose atorvastatin loading before primary percutaneous coronary intervention in ST-segment elevation myocardial infarction: the STATIN STEMI trial. 2010 , 3, 332-9	119
661	Local delivery versus intracoronary infusion of abciximab in patients with acute coronary syndromes. 2010 , 3, 928-34	65
660	Safety and efficacy of thrombectomy in patients undergoing primary percutaneous coronary intervention for acute ST elevation MI: a meta-analysis of randomized controlled trials. 2010 , 10, 10	54

659	The contribution of intramyocardial hemorrhage to the "no-reflow phenomenon": a study performed by cardiac magnetic resonance. 2010 , 27, 1120-9		8
658	Clinical experience with a novel intracoronary perfusion catheter to treat no-reflow phenomenon in acute coronary syndromes. 2010 , 23, 109-13		11
657	Perfusion assessed by real-time contrast echocardiography correlates with clinical and echocardiographic parameters in patients with first STEMI treated with PCI - 6-month follow-up. 2010 , 6, 176-82		6
656	Stent direto na intervençāo coronāria primāria: avaliaçāo de seu uso na prātica clīnica contemporānea. 2010 , 18, 412-418		2
655	Thrombectomy during percutaneous coronary intervention in patients with acute myocardial infarction. 2010 , 1, 30-37		
654	Intracoronary versus intravenous administration of abciximab in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention with thrombus aspiration: the comparison of intracoronary versus intravenous abciximab administration during percutaneous reperfusion of ST-segment elevation myocardial infarction (CICERO) trial	16.7	106
653	The role of PET with ¹³ N-ammonia and ¹⁸ F-FDG in the assessment of myocardial perfusion and metabolism in patients with recent AMI and intracoronary stem cell injection. 2010 , 51, 1908-16		35
652	Is the myocardial blush grade scored by the operator during primary percutaneous coronary intervention of prognostic value in patients with ST-elevation myocardial infarction in routine clinical practice?. 2010 , 3, 216-23		15
651	Outcome of primary percutaneous intervention in patients with infarct-related coronary artery ectasia. 2010 , 61, 574-9		21
650	Thrombus aspiration in ST elevation myocardial infarction: comparative efficacy in patients treated early and late after onset of symptoms. 2010 , 96, 1287-90		21
649	Early administration of small-molecule glycoprotein IIb/IIIa inhibitors before primary percutaneous coronary intervention for ST-elevation myocardial infarction: insights from randomized clinical trials. 2010 , 15, 135-44		3
648	Predictors of outcomes in medically treated patients with acute coronary syndromes after angiographic triage: an Acute Catheterization And Urgent Intervention Triage Strategy (ACUITY) substudy. <i>Circulation</i> , 2010 , 121, 853-62	16.7	28
647	Myocardial perfusion grade after late infarct artery recanalization is associated with global and regional left ventricular function at one year: analysis from the Total Occlusion Study of Canada-2. 2010 , 3, 549-55		7
646	Influence of bone marrow stem cells on left ventricle perfusion and ejection fraction in patients with acute myocardial infarction of anterior wall: randomized clinical trial: Impact of bone marrow stem cell intracoronary infusion on improvement of microcirculation. 2010 , 31, 691-702		81
645	Predicting the no-reflow phenomenon following successful percutaneous coronary intervention. 2010 , 4, 403-20		5
644	Design and methodologies of the POSTconditioning during coronary angioplasty in acute myocardial infarction (POST-AMI) trial. 2010 , 116, 110-6		20
643	Microvascular obstruction: underlying pathophysiology and clinical diagnosis. 2010 , 55, 1649-60		191
642	Clinical predictors of successful thrombectomy with the Export [®] aspiration catheter in the acute phase of myocardial infarction. Data from the RICO survey working group. 2010 , 103, 522-9		2

641	Successful reperfusion therapy: from epicardial to myocardial salvage. 2010 , 63, 757-9	1
640	Myocardial perfusion grade, myocardial salvage indices and long-term mortality in patients with acute myocardial infarction and full restoration of epicardial blood flow after primary percutaneous coronary intervention. 2010 , 63, 770-8	4
639	The sum of ST-segment elevation is the best predictor of microvascular obstruction in patients treated successfully by primary percutaneous coronary intervention. Cardiovascular magnetic resonance study. 2010 , 63, 1145-54	8
638	Clinical correlates of arterial lactate levels in patients with ST-segment elevation myocardial infarction at admission: a descriptive study. 2010 , 14, R164	37
637	Myeloperoxidase may contribute to the no-reflow phenomenon in patients with acute myocardial infarction. 2010 , 139, 187-92	9
636	Serum B-type natriuretic peptide on admission can predict the 'no-reflow' phenomenon after primary drug-eluting stent implantation for ST-segment elevation myocardial infarction. 2010 , 141, 175-81	20
635	Baseline von Willebrand factor plasma levels and no-reflow phenomenon after primary percutaneous coronary intervention for ST segment elevation myocardial infarction. 2010 , 145, 230-232	6
634	Extent of ST-segment resolution after fibrinolysis adds improved risk stratification to clinical risk score for ST-segment elevation myocardial infarction. 2010 , 159, 55-62	6
633	Quantitative myocardial blush grade for the detection of cardiac allograft vasculopathy. 2010 , 159, 643-651.e216	
632	Endothelin-1 release in acute myocardial infarction as a predictor of long-term prognosis and no-reflow assessed by contrast-enhanced magnetic resonance imaging. 2010 , 159, 882-90	53
631	Relationship between myocardial blush grades, staining, and severe microvascular damage after primary percutaneous coronary intervention a study performed with contrast-enhanced magnetic resonance in a large consecutive series of patients. 2010 , 159, 1124-32	21
630	Differential effects of post-dilation after stent deployment in patients presenting with and without acute myocardial infarction. 2010 , 160, 979-986.e1	34
629	Thrombus aspiration followed by direct stenting: a novel strategy of primary percutaneous coronary intervention in ST-segment elevation myocardial infarction. Results of the Polish-Italian-Hungarian RANdomized ThrombEctomy Trial (PIHRATE Trial). 2010 , 160, 966-72	73
628	Angiographic assessment of microvascular perfusion--myocardial blush in clinical practice. 2010 , 160, 1015-22	30
627	Impact of diabetes on survival in patients with ST-segment elevation myocardial infarction treated by primary angioplasty: insights from the POLISH STEMI registry. 2010 , 210, 516-20	32
626	Association of CD40 ligand levels in the culprit coronary arteries with subsequent prognosis of acute myocardial infarction. 2010 , 213, 268-72	5
625	Impact of statin pretreatment on the incidence of plaque rupture in ST-elevation acute myocardial infarction. 2010 , 213, 505-11	21
624	[Not Available]. 2010 , 63, 757-9	1

623	Grado de perfusión miocárdica, índices de rescate miocárdico y mortalidad a largo plazo en pacientes con infarto agudo de miocardio y restablecimiento completo del flujo sanguíneo epicárdico tras intervención coronaria percutánea primaria. 2010 , 63, 770-778	8
622	La suma de la elevación del segmento ST predice mejor la obstrucción microvascular en pacientes tratados con éxito con una intervención coronaria percutánea primaria. Un estudio de resonancia magnética cardiovascular. 2010 , 63, 1145-1154	20
621	A no-reflow prediction model in patients with ST-elevation acute myocardial infarction and primary drug-eluting stenting. 2011 , 45, 98-104	6
620	Composition of coronary thrombus in acute myocardial infarction. 2011 , 57, 1359-67	266
619	Optimal therapy for ST-segment elevation myocardial infarction: the role of residual thrombus. 2011 , 57, 1874-6	4
618	Rationale and design of the INFUSE-AMI study: A 2x2 factorial, randomized, multicenter, single-blind evaluation of intracoronary abciximab infusion and aspiration thrombectomy in patients undergoing percutaneous coronary intervention for anterior ST-segment elevation myocardial infarction. 2011 , 161, 178-181.e7	34
617	Value of the SYNTAX score in patients treated by primary percutaneous coronary intervention for acute ST-elevation myocardial infarction: The MI SYNTAXscore study. 2011 , 161, 771-81	89
616	Association of initial thrombolysis in myocardial infarction flow grade with mortality among patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention: a National Registry of Myocardial Infarction-5 (NRM-5) analysis. 2011 , 162, 178-83	9
615	Quantitative Blush Evaluator accurately quantifies microvascular dysfunction in patients with ST-elevation myocardial infarction: comparison with cardiovascular magnetic resonance. 2011 , 162, 372-381.e2 ¹⁶	
614	Effects of atorvastatin pretreatment on infarct size in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2011 , 162, 1026-33	37
613	Relationship between angiographic dynamic and densitometric assessment of myocardial reperfusion and survival in patients with acute myocardial infarction treated with primary percutaneous coronary intervention: the harmonizing outcomes with revascularization and stents in AMI (HORIZONS-AMI) trial. 2011 , 162, 1044-51	19
612	Effect of delayed vs immediate stent implantation on myocardial perfusion and cardiac function in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous intervention with thrombus aspiration. 2011 , 27, 541-7	27
611	Endothelin-1 levels predict endothelial progenitor cell mobilization after acute myocardial infarction. 2011 , 82, 177-81	16
610	An uncommon cause of myocardial infarction. 2011 , 20, 202-4	1
609	Dynamic Changes in ST Segment Resolution After Myocardial Infarction and the Association with Microvascular Injury on Cardiac Magnetic Resonance Imaging. 2011 , 20, 111-8	20
608	Evolution of myocardial perfusion during primary angioplasty in spontaneously reperfused infarct-related artery: impact on long-term clinical outcomes and left ventricular function recovery. 2011 , 147, 25-31	10
607	Laser for complex coronary lesions: impact of excimer lasers and technical advancements. 2011 , 146, 296-9	15
606	Do currently recommended risk prediction tools help primary prevention of cardiovascular disease in Chinese population?. 2011 , 146, 299-301	2

605	Predictors of myocardial microvascular obstruction in patients treated by primary percutaneous coronary intervention and a short ischemic time. 2011 , 153, 113-5	3
604	Usefulness of endothelin-1 assessment in acute myocardial infarction. 2011 , 64, 105-10	19
603	Usefulness of Endothelin-1 Assessment in Acute Myocardial Infarction. 2011 , 64, 105-110	3
602	Effect of Aspiration Thrombectomy on Microvascular Dysfunction in ST-Segment Elevation Myocardial Infarction With an Elevated Neutrophil Count. 2011 , 41, 68-75	5
601	Intervenç�o coron�ria pelas vias radial ou femoral no infarto agudo do mioc�rdio com supradesnivelamento do segmento ST: uma vis�o da pr�tica cl�nica contempor�nea. 2011 , 19, 266-271	
600	MRI discriminates thrombus composition and ST resolution after percutaneous coronary intervention in patients with ST-elevation myocardial infarction. 2011 , 6, e18459	7
599	Effect of primary percutaneous coronary intervention on myocardial repolarization. 2011 , 12, 795-9	2
598	Association of matrix metalloproteinase-1, matrix metalloproteinase-9, tissue inhibitor of matrix metalloproteinase-1, and interleukin-6 with epicardial and myocardial perfusion. 2011 , 22, 253-8	7
597	Myocardial blood flow and infarct size after CD133+ cell injection in large myocardial infarction with good recanalization and poor reperfusion: results from a randomized controlled trial. 2011 , 12, 239-48	39
596	Ischemic preconditioning effect of prodromal angina is attenuated in acute myocardial infarction patients with hypertensive left ventricular hypertrophy. 2011 , 75, 1192-9	18
595	High QRS score on admission strongly predicts impaired myocardial reperfusion in patients with a first anterior acute myocardial infarction. 2011 , 75, 626-32	13
594	Residual platelet reactivity after clopidogrel loading in patients with ST-elevation myocardial infarction undergoing an unexpectedly delayed primary percutaneous coronary intervention. - Impact on intracoronary thrombus burden and myocardial perfusion-. 2011 , 75, 2105-12	15
593	Effect of pretreatment with pioglitazone on reperfusion injury in diabetic patients with acute myocardial infarction. 2011 , 75, 1968-74	18
592	Angiographic patterns of myocardial reperfusion after primary angioplasty and ventricular remodeling. 2011 , 22, 507-14	11
591	Endogenous serum erythropoietin and no-reflow in patients with ST-elevation myocardial infarction. 2011 , 41, 1210-9	10
590	Early glycoprotein IIb/IIIa inhibitors in primary angioplasty-abciximab long-term results (EGYPT-ALT) cooperation: individual patient's data meta-analysis. 2011 , 9, 2361-70	100
589	Facilitated reperfusion with prehospital glycoprotein IIb/IIIa inhibition: predictors of complete ST-segment resolution before primary percutaneous coronary intervention in the On-TIME 2 trial: correlates of reperfusion before primary PCI. 2011 , 44, 42-8	5
588	Grade 3 ischemia on the admission electrocardiogram is associated with severe microvascular injury on cardiac magnetic resonance imaging after ST elevation myocardial infarction. 2011 , 44, 49-57	25

587	Predictors and outcome of grade 3 ischemia in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2011 , 44, 516-22	18
586	Management of the no-reflow phenomenon. 2011 , 132, 72-85	21
585	Comparison between the effect of intracoronary bolus of tirofiban versus eptifibatide as adjunctive antiplatelet therapy on the outcome of primary coronary intervention in patients with acute anterior ST segment elevation myocardial infarction. 2011 , 63, 109-115	
584	Impact of chronic pre-treatment of statins on the level of systemic inflammation and myocardial perfusion in patients undergoing primary angioplasty. 2011 , 107, 179-85	24
583	Relation of circulating osteoprotegerin levels on admission to microvascular obstruction after primary percutaneous coronary intervention. 2011 , 107, 857-62	8
582	New method of intracoronary adenosine injection to prevent microvascular reperfusion injury in patients with acute myocardial infarction undergoing percutaneous coronary intervention. 2011 , 107, 1131-5	39
581	Quantitative analysis of the impact of total ischemic time on myocardial perfusion and clinical outcome in patients with ST-elevation myocardial infarction. 2011 , 108, 1536-41	24
580	Usefulness of thrombus aspiration for the treatment of coronary stent thrombosis. 2011 , 108, 1721-7	15
579	TIMI myocardial perfusion frame count: turning blush into numbers. 2011 , 77, 459	
578	Impact of chronic total occlusions on markers of reperfusion, infarct size, and long-term mortality: a substudy from the TAPAS-trial. 2011 , 77, 484-91	50
577	Computer-assisted quantification of myocardial reperfusion after primary percutaneous coronary intervention predicts functional and contrast-enhanced cardiovascular magnetic resonance outcomes in patients with ST-segment elevation myocardial infarction. 2011 , 77, 174-81	6
576	Aspiration coronary thrombectomy for acute myocardial infarction increases myocardial salvage: single center randomized study. 2011 , 78, 523-31	33
575	Prognostic significance of coronary thrombus in patients undergoing percutaneous coronary intervention for acute coronary syndromes: a subanalysis of the ACUITY (Acute Catheterization and Urgent Intervention Triage strategY) trial. 2011 , 4, 769-77	27
574	Acute Myocardial Infarction with Severe ST Segment Elevation Treated with Percutaneous Coronary Intervention More than Two Days after Onset: A Case with Remarkable Recovery. 2011 , 20, 103-6	1
573	High-dose intracoronary adenosine for myocardial salvage in patients with acute ST-segment elevation myocardial infarction. 2011 , 32, 867-77	111
572	Ischaemic postconditioning revisited: lack of effects on infarct size following primary percutaneous coronary intervention. 2012 , 33, 103-12	181
571	Operator versus core laboratory assessment of angiographic reperfusion markers in patients undergoing primary percutaneous coronary intervention for ST-segment-elevation myocardial infarction. 2012 , 5, 563-9	8
570	Delayed versus immediate stenting for the treatment of ST-elevation acute myocardial infarction with a high thrombus burden. 2012 , 23, 497-506	31

569	Clinical impact of intracoronary abciximab in patients undergoing primary percutaneous coronary intervention: an individual patient data pooled analysis of randomised studies. 2012 ,	4
568	Prevalence, predictors and clinical impact of unique and multiple chronic total occlusion in non-infarct-related artery in patients presenting with ST-elevation myocardial infarction. 2012 , 98, 1732-7	20
567	Recent Advances in the Treatment of ST-Segment Elevation Myocardial Infarction. 2012 , 2012, 683683	8
566	Prognostic impact of a chronic total occlusion in a non-infarct-related artery in patients with ST-segment elevation myocardial infarction: 3-year results from the HORIZONS-AMI trial. 2012 , 33, 768-75	157
565	Measuring myocardial salvage. 2012 , 94, 266-75	47
564	Coronary microvascular dysfunction in the clinical setting: from mystery to reality. 2012 , 33, 2771-2782b	150
563	Remote ischemic conditioning: the cardiologist's perspective. 2012 , 13, 667-74	15
562	Effect of serum gamma-glutamyl transferase levels on myocardial perfusion and long-term prognosis after primary angioplasty in patients with acute ST-elevation myocardial infarction. 2012 , 60, 1186-93	16
561	Efficacy of clotinab in acute myocardial infarction trial-ST elevation myocardial infarction (ECLAT-STEMI). 2012 , 76, 405-13	5
560	Clinical implications of electrocardiograms for patients with anterior wall ST-segment elevation acute myocardial infarction in the interventional era. 2012 , 76, 32-40	18
559	Intracoronary hemodynamic effects of pressure-controlled intermittent coronary sinus occlusion (PICSO): results from the First-In-Man Prepare PICSO Study. 2012 , 25, 549-56	13
558	Glycoprotein IIb-IIIa inhibitors. 2012 , 30, e242-54	20
557	Intracoronary EKG assessment: getting the right data in the nick of time. 2012 , 80, 754-5	
556	Comparison of magnetic resonance imaging findings in non-ST-segment elevation versus ST-segment elevation myocardial infarction patients undergoing early invasive intervention. 2012 , 28, 1487-97	18
555	Cardiac magnetic resonance derived late microvascular obstruction assessment post ST-segment elevation myocardial infarction is the best predictor of left ventricular function: a comparison of angiographic and cardiac magnetic resonance derived measurements. 2012 , 28, 1971-81	37
554	Plasma osteoprotegerin level on admission is associated with no-reflow phenomenon after primary angioplasty and subsequent left ventricular remodeling in patients with acute ST-segment elevation myocardial infarction. 2012 , 221, 254-9	11
553	Serum uric acid: a marker of metabolic syndrome and subclinical atherosclerosis in Korean men. 2012 , 63, 420-8	25
552	Characterizing myocardial edema and hemorrhage using quantitative T2 and T2* mapping at multiple time intervals post ST-segment elevation myocardial infarction. 2012 , 5, 566-72	79

551	Intracoronary microparticles and microvascular obstruction in patients with ST elevation myocardial infarction undergoing primary percutaneous intervention. 2012 , 33, 2928-38	78
550	A high loading dose of clopidogrel reduces myocardial infarct size in patients undergoing primary percutaneous coronary intervention: a magnetic resonance imaging study. 2012 , 163, 500-7	22
549	Multiple biomarkers at admission are associated with angiographic, electrocardiographic, and imaging cardiovascular mechanistic markers of outcomes in patients undergoing primary percutaneous coronary intervention for acute ST-elevation myocardial infarction. 2012 , 163, 783-9	4
548	Impact of early abciximab administration on infarct size in patients with ST-elevation myocardial infarction. 2012 , 155, 230-5	14
547	Additive benefit of glycoprotein IIb/IIIa inhibition and adjunctive thrombus aspiration during primary coronary intervention: results of the Initial Thrombosuction and Tirofiban Infusion (ITTI) trial. 2012 , 156, 174-9	18
546	Relationship between microcirculatory dysfunction and resolution of ST-segment elevation in the early phase after primary angioplasty in patients with ST-segment elevation myocardial infarction. 2012 , 159, 144-9	5
545	Angiographically-derived SYNTAX score and its prognostic value in dialysis patients: comparison with the Khan index. 2012 , 28, 450-7	6
544	Prognostic value at 5 years of microvascular obstruction after acute myocardial infarction assessed by cardiovascular magnetic resonance. 2012 , 14, 46	69
543	Study design for the "effect of METOp ^{ro} lol in CARDioproteCtioN during an acute myocardial InfarCtion" (METOCARD-CNIC): a randomized, controlled parallel-group, observer-blinded clinical trial of early pre-reperfusion metoprolol administration in ST-segment elevation myocardial infarction. 2012 , 164, 473-480.e5	35
542	Postconditioning during coronary angioplasty in acute myocardial infarction: the POST-AMI trial. 2012 , 162, 33-8	104
541	Prognostic value of myocardial contrast delayed enhancement with 64-slice multidetector computed tomography after acute myocardial infarction. 2012 , 59, 730-8	50
540	Time-dependent detrimental effects of distal embolization on myocardium and microvasculature during primary percutaneous coronary intervention. 2012 , 5, 1170-7	22
539	A prospective randomized trial of thrombectomy versus no thrombectomy in patients with ST-segment elevation myocardial infarction and thrombus-rich lesions: MUSTELA (MULTIdevice Thrombectomy in Acute ST-Segment Elevation Acute Myocardial Infarction) trial. 2012 , 5, 1223-30	60
538	Superior outcome with direct catheter laboratory access vs ED-activated primary percutaneous coronary intervention. 2012 , 30, 1118-24	6
537	Imaging the myocardial microcirculation post-myocardial infarction. 2012 , 9, 282-92	13
536	Linear quantification correction for myocardial perfusion imaging from x-ray coronary angiography. 2012 ,	3
535	Safety and efficacy of intracoronary adenosine administration in patients with acute myocardial infarction undergoing primary percutaneous coronary intervention: a meta-analysis of randomized controlled trials. 2012 , 6, 101-14	22
534	High-throughput 13-parameter immunophenotyping identifies shifts in the circulating T-cell compartment following reperfusion in patients with acute myocardial infarction. 2012 , 7, e47155	24

533	[Aortic valve stenosis is associated with reduced myocardial perfusion as assessed by videodensitometry in coronary angiograms]. 2012 , 153, 1256-62	0
532	Acute effect of sodium nitroprusside on microvascular dysfunction in patients who underwent percutaneous coronary intervention for acute ST-segment elevation myocardial infarction. 2012 , 53, 337-40	7
531	The effect of pre-hospital glycoprotein IIb/IIIa inhibitors on angiographic outcome in STEMI patients who are candidates for primary PCI. 2012 , 79, 956-64	13
530	Collateral blood flow can predict myocardial blush grade in primary coronary intervention. 2012 , 80, 67-70	5
529	MGUard versus bare-metal stents plus manual thrombectomy in ST-elevation myocardial infarction patients-(GUARDIAN) trial: study design and rationale. 2012 , 79, 1118-26	15
528	Early statin treatment prior to primary PCI for acute myocardial infarction: REPERATOR, a randomized placebo-controlled pilot trial. 2012 , 80, 756-65	33
527	Inverse relation of total serum bilirubin to coronary artery calcification score detected by multidetector computed tomography in males. 2012 , 35, 301-6	33
526	Intracoronary versus intravenous high-dose bolus plus maintenance administration of tirofiban in patients undergoing primary percutaneous coronary intervention for acute ST elevation myocardial infarction. 2012 , 34, 65-72	7
525	Toward a comprehensive approach to pharmacoinvasive therapy for patients with ST segment elevation acute myocardial infarction. 2012 , 34, 180-6	7
524	Independent no-reflow predictors in female patients with ST-elevation acute myocardial infarction treated with primary percutaneous coronary intervention. 2012 , 27, 243-9	24
523	A comparison of intracoronary with intravenous glycoprotein IIb/IIIa inhibitors during percutaneous coronary intervention in patients with acute coronary syndrome: a meta-analysis of randomized controlled trials. 2012 , 25, 223-34	6
522	Impact of vascular approach (transradial vs. transfemoral) on the efficacy of thrombus aspiration in acute myocardial infarction patients. 2012 , 13, 79-83	5
521	The impact of intracoronary thrombus aspiration on STEMI outcomes. 2012 , 13, 167-71	6
520	Implications of ST-segment elevation in leads V5 and V6 in patients with reperfused inferior wall acute myocardial infarction. 2012 , 109, 314-9	4
519	Usefulness of the SYNTAX score to predict "no reflow" in patients treated with primary percutaneous coronary intervention for ST-segment elevation myocardial infarction. 2012 , 109, 601-6	64
518	Thrombus-aspiration through 5 Fr guiding catheter with transradial approach in acute coronary syndromes: feasibility of a mini-invasive strategy. 2012 , 25, 323-9	1
517	Clinical impact of thrombus aspiration during primary percutaneous coronary intervention: results from Korea Acute Myocardial Infarction Registry. 2012 , 59, 249-57	19
516	Thrombus aspiration during primary percutaneous coronary intervention is associated with reduced myocardial edema, hemorrhage, microvascular obstruction and left ventricular remodeling. 2012 , 14, 19	17

515	Effect of postconditioning in patients with ST-elevation acute myocardial infarction. 2012 , 27, 14-8	8
514	Pharmacoinvasive therapy for ST elevation myocardial infarction in China: a pilot study. 2012 , 33, 101-8	2
513	Thrombus Aspiration in Thrombus containing culprit lesions in Non-ST-Elevation Myocardial Infarction (TATORT-NSTEMI): study protocol for a randomized controlled trial. 2013 , 14, 110	9
512	Impact of multivessel disease on myocardial perfusion and survival among patients undergoing primary percutaneous coronary intervention with glycoprotein IIb/IIIa inhibitors. 2013 , 106, 155-61	6
511	Validation of videodensitometric myocardial perfusion assessment. 2013 , 8, 600-607	
510	Optical coherence tomography guided in-stent thrombus removal in patients with acute coronary syndromes. 2013 , 29, 989-96	13
509	What is the optimum adjunctive reperfusion strategy for primary percutaneous coronary intervention?. 2013 , 382, 633-43	15
508	Left ventricular apical aneurysm following primary percutaneous coronary intervention. 2013 , 28, 677-83	9
507	Usefulness of local delivery of thrombolytics before thrombectomy in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention (the delivery of thrombolytics before thrombectomy in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention [DISSOLUTION] randomized	25
506	Effects of total bilirubin on the prevalence of osteoporosis in postmenopausal women without potential liver disease. 2013 , 31, 637-43	10
505	Association between proximal stent edge positioning on atherosclerotic plaques containing lipid pools and postprocedural myocardial infarction (from the CLI-POOL Study). 2013 , 111, 526-31	30
504	Complementary prognostic utility of myocardial blush grade and ST-segment resolution after primary percutaneous coronary intervention: analysis from the HORIZONS-AMI trial. 2013 , 166, 676-83	20
503	Comparison of the reperfusion efficacy of thrombus aspiration with and without distal protection during primary percutaneous coronary intervention in patients with acute ST-segment elevation myocardial infarction. 2013 , 112, 1725-9	6
502	Remote ischemic post-conditioning of the lower limb during primary percutaneous coronary intervention safely reduces enzymatic infarct size in anterior myocardial infarction: a randomized controlled trial. 2013 , 6, 1055-63	163
501	Interaction of chronic total occlusion and chronic kidney disease in patients undergoing primary percutaneous coronary intervention for acute ST-elevation myocardial infarction. 2013 , 112, 194-9	11
500	Relationship between myocardial reperfusion, infarct size, and mortality: the INFUSE-AMI (Intracoronary Abciximab and Aspiration Thrombectomy in Patients With Large Anterior Myocardial Infarction) trial. 2013 , 6, 718-24	34
499	Prognostic value of microvascular resistance index immediately after primary percutaneous coronary intervention on left ventricular remodeling in patients with reperfused anterior acute ST-segment elevation myocardial infarction. 2013 , 6, 1046-54	25
498	Meta-analysis of randomized controlled trials on the efficacy and safety of intracoronary administration of tirofiban for no-reflow phenomenon. 2013 , 13, 68	10

497	Chronic pretreatment of metformin is associated with the reduction of the no-reflow phenomenon in patients with diabetes mellitus after primary angioplasty for acute myocardial infarction. 2013 , 31, 60-4	16
496	A meta-analytic overview of thrombectomy during primary angioplasty. 2013 , 166, 606-12	123
495	Case-control registry of excimer laser coronary angioplasty versus distal protection devices in patients with acute coronary syndromes due to saphenous vein graft disease. 2013 , 112, 1586-91	24
494	Microvascular coronary flow comparison in acute myocardial infarction angioplasty treated with a mesh covered stent (MGUARD stent) versus bare metal stent: MICAMI-MGUARD. 2013 , 14, 4-8	6
493	Predictors of cardiovascular magnetic resonance-derived microvascular obstruction on patient admission in STEMI. 2013 , 166, 77-84	18
492	Closed versus open cell stent for high-risk percutaneous coronary interventions in ST-elevation acute myocardial infarction: the Closed versus Open Cells stent for High risk percutaneous coronary Interventions in ST-Elevation acute myocardial infarction (COHISE) pilot study. 2013 , 165, 115-20	4
491	Intracoronary ECG during primary percutaneous coronary intervention for ST-segment elevation myocardial infarction predicts microvascular obstruction and infarct size. 2013 , 165, 61-6	12
490	Effect of manual thrombus aspiration during primary percutaneous coronary intervention on infarct size: evaluation with cardiac computed tomography. 2013 , 168, 4328-30	13
489	Usefulness of monocyte chemoattractant protein-1 to predict no-reflow and three-year mortality in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2013 , 112, 187-93	14
488	Effect of macroscopic-positive thrombus retrieval during primary percutaneous coronary intervention with thrombus aspiration on myocardial infarct size and microvascular obstruction. 2013 , 111, 159-65	5
487	Contrast-induced nephropathy in patients undergoing primary percutaneous coronary intervention without acute left ventricular ejection fraction impairment. 2013 , 111, 684-8	31
486	Angiographic assessment of myocardial perfusion in Tako-Tsubo syndrome. 2013 , 168, 4717-22	12
485	The never ending quest for an ideal angiographic surrogate of coronary reperfusion. 2013 , 65, 7-11	
484	Association of fragmented QRS complex with myocardial reperfusion in acute ST-elevated myocardial infarction. 2013 , 18, 69-74	18
483	Use of thrombectomy devices in primary percutaneous coronary intervention: a systematic review and meta-analysis. 2013 , 163, 229-241	46
482	The neutrophil to lymphocyte ratio was associated with impaired myocardial perfusion and long term adverse outcome in patients with ST-elevated myocardial infarction undergoing primary coronary intervention. 2013 , 228, 203-10	91
481	Development and validation of a clinical risk score predicting the no-reflow phenomenon in patients treated with primary percutaneous coronary intervention for ST-segment elevation myocardial infarction. 2013 , 124, 153-60	15
480	Effect of new method of intracoronary adenosine injection during primary percutaneous coronary intervention on microvascular reperfusion injury - clinical outcome and 1-year follow-up. 2013 , 124, 199-206	4

479	Myocardial 'no-reflow'--diagnosis, pathophysiology and treatment. 2013 , 167, 1798-806		34
478	Utility of Thrombectomy in Primary Percutaneous Coronary Intervention. 2013 , 2, 361-374		
477	Predictors of post-stress LVEF drop 6 months after reperfused myocardial infarction: a gated myocardial perfusion SPECT study. 2013 , 27, 112-22		3
476	Implications of myocardial reperfusion on survival in women versus men with acute myocardial infarction undergoing primary coronary intervention. 2013 , 112, 1087-92		6
475	No-reflow reversibility: a study based on serial assessment of multiple biomarkers. 2013 , 6, 798-807		8
474	Ischemic postconditioning during primary percutaneous coronary intervention: the effects of postconditioning on myocardial reperfusion in patients with ST-segment elevation myocardial infarction (POST) randomized trial. <i>Circulation</i> , 2013 , 128, 1889-96	16.7	140
473	Impact of coronary microvascular function on long-term cardiac mortality in patients with acute ST-segment-elevation myocardial infarction. 2013 , 6, 207-15		63
472	Randomized study to assess the effect of thrombus aspiration on flow area in patients with ST-elevation myocardial infarction: an optical frequency domain imaging study--TROFI trial. 2013 , 34, 1050-60		89
471	Late Clinical Follow-up of Patients with Complex Coronary Lesions Treated with the New MGuard [®] Bare-Metal Stent. 2013 , 21, 23-29		
470	Predictors of Failed Thrombus Aspiration in Patients Undergoing Primary Percutaneous Coronary Intervention. 2013 , 21, 234-239		
469	Relationship of plasma neuropeptide Y with angiographic, electrocardiographic and coronary physiology indices of reperfusion during ST elevation myocardial infarction. 2013 , 99, 1198-203		31
468	Patients with microvascular obstruction after primary percutaneous coronary intervention show a gp91phox (NOX2) mediated persistent oxidative stress after reperfusion. 2013 , 2, 379-88		14
467	Soluble endothelial protein C receptor levels and protein C activity in patients with acute ST-segment elevation myocardial infarction. 2013 , 24, 209-16		5
466	Mean platelet volume is associated with poor postinterventional myocardial blush grade in patients with ST-segment elevation myocardial infarction. 2013 , 24, 285-9		13
465	Impact of altitude on predicting midterm outcome in patients with ST elevation myocardial infarction. 2013 , 19, 382-8		2
464	Spontaneous and procedural plaque embolisation in native coronary arteries: pathophysiology, diagnosis, and prevention. 2013 , 2013, 364247		9
463	Is postdilatation with a noncompliant balloon necessary after coronary stent deployment during primary angioplasty?. 2013 , 26, 325-31		14
462	MGuard mesh-covered stent for treatment of ST-segment elevation myocardial infarction with high thrombus burden despite manual aspiration. 2013 , 26, 1-7		8

461	The efficacy and safety of pharmacoinvasive therapy with prourokinase for acute ST-segment elevation myocardial infarction patients with expected long percutaneous coronary intervention-related delay. 2013 , 31, 285-90	5
460	Is the difference in outcome between men and women treated by primary percutaneous coronary intervention age dependent? Gender difference in STEMI stratified on age. 2013 , 2, 334-41	86
459	Vaporizing thrombus with excimer laser before coronary stenting improves myocardial reperfusion in acute coronary syndrome. 2013 , 77, 1445-52	25
458	Effect of statin pretreatment on myocardial perfusion in patients undergoing primary percutaneous coronary intervention: a systematic review and meta-analysis. 2013 , 36, E17-24	6
457	Microvascular Obstruction After Primary Percutaneous Coronary Intervention: Pathogenesis, Diagnosis and Prognostic Significance. 2013 , 11, 245-262	1
456	Myocardial "No-Reflow" Prevention. 2013 , 11, 263-277	
455	Primary percutaneous coronary intervention: devices to prevent no-reflow phenomenon. 2013 , 5, 289-300	2
454	Intracoronary fixed dose of nitroprusside via thrombus aspiration catheter for the prevention of the no-reflow phenomenon following primary percutaneous coronary intervention in acute myocardial infarction. 2013 , 6, 479-484	24
453	No reflow: what's in a name?. 2013 , 19, 4517-8	1
452	Clopidogrel pretreatment in primary percutaneous coronary intervention: prevalence of high on-treatment platelet reactivity and impact on preprocedural patency of the infarct-related artery. 2013 , 110, 110-7	17
451	Immediate and midterm follow-up results of excimer laser application in complex percutaneous coronary interventions: Report from a single center experience. 2013 , 5, 10-5	3
450	Efficacy of combination treatment with intracoronary abciximab and aspiration thrombectomy on myocardial perfusion in patients with ST-segment elevation myocardial infarction undergoing primary coronary stenting. 2014 , 55, 606-16	18
449	Persistent enhanced platelet activation in patients with acute myocardial infarction and coronary microvascular obstruction: clinical implications. 2014 , 111, 122-30	14
448	Intra-Coronary Administration of Tacrolimus Prior to First-Balloon Inflation Attenuates Infarct Size and Improves Left Ventricular Function in Patients with ST-segment Elevation Myocardial Infarction (COAT-STEMI) Undergoing Primary Coronary Intervention. 2014 , 04,	
447	Effect of High Dose Rosuvastatin Loading before Primary Percutaneous Coronary Intervention on Infarct Size in Patients with ST-Segment Elevation Myocardial Infarction. 2014 , 44, 76-81	23
446	[Procedural aspects in primary PCI: arterial access, stent selection, thrombectomy and treatment of non-culprit lesions]. 2014 , 39, 685-91	
445	The REFLO-STEMI trial comparing intracoronary adenosine, sodium nitroprusside and standard therapy for the attenuation of infarct size and microvascular obstruction during primary percutaneous coronary intervention: study protocol for a randomised controlled trial. 2014 , 15, 371	12
444	Does the Inhibition of the Final Common Pathway of Platelet Aggregation Reduce the No-Reflow Phenomenon During Primary Percutaneous Coronary Intervention? Tirofiban no Infarto Agudo do miocárdio e a não Reperfução (TIARA). 2014 , 22, 81-86	1

443	Time of symptom onset and value of myocardial blush and infarct size on prognosis in patients with ST-elevation myocardial infarction. 2014 , 31, 797-806	9
442	Catheter Aspiration after Every Stage during Primary Percutaneous Angioplasty; ADMIT Trial. 2014 , 23, 29-40	2
441	A Randomized Comparison between Everolimus-Eluting Stent and Cobalt Chromium Stent in Patients with Acute ST-Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention Using Routine Intravenous Eptifibatide: The X-MAN (Xience vs. Multi-Link Stent in Acute Myocardial Infarction) Trial, A Pilot Study. 2014 , 23, 93-100	
440	Effect of intracoronary nitroprusside in preventing no reflow phenomenon during primary percutaneous coronary intervention: a meta-analysis. 2014 , 27, 356-64	21
439	The randomized physiologic assessment of thrombus aspiration in patients with acute ST-segment elevation myocardial infarction trial (PATA STEMI): study rationale and design. 2014 , 27, 341-7	4
438	Prospective study of effects of endogenous estrogens on myocardial no-reflow risk in postmenopausal women with acute myocardial infarction. 2014 , 27, 437-43	5
437	Clinical profile and impact of family history of premature coronary artery disease on clinical outcomes of patients undergoing primary percutaneous coronary intervention for ST-elevation myocardial infarction: analysis from the HORIZONS-AMI Trial. 2014 , 15, 375-80	8
436	Length of left anterior descending coronary artery determines prognosis in acute anterior wall myocardial infarction. 2014 , 84, 316-20	11
435	Sequential evaluation of coronary flow patterns after primary angioplasty in acute anterior ST-elevation myocardial infarction predicts recovery of left ventricular systolic function. 2014 , 31, 644-653	1
434	Left anterior descending coronary artery flow after primary angioplasty in acute anterior ST-elevation myocardial infarction: how much flow is needed for left ventricular functional recovery?. 2014 , 3, 223-8	1
433	Neutrophil to lymphocyte ratio in acute ST-segment elevation myocardial infarction. 2014 , 348, 37-42	15
432	Clinical importance of hyperemic coronary blood flow (thrombolysis in myocardial infarction-intravenous flow) after primary percutaneous coronary intervention. 2014 , 25, 665-70	
431	Cytochrome c release in acute myocardial infarction predicts poor prognosis and myocardial reperfusion on contrast-enhanced magnetic resonance imaging. 2014 , 25, 66-72	10
430	CMD in Obstructive CAD. 2014 , 145-180	
429	Angiographic and Clinical Impact of Successful Manual Thrombus Aspiration in Diabetic Patients Undergoing Primary PCI. 2014 , 2014, 263926	6
428	TASTE: To aspirate, or not to aspirate, remains a question. 2014 , 2014, 37-9	1
427	Effect of ischemic postconditioning on microvascular obstruction in reperfused myocardial infarction. Results of a randomized study in patients and of an experimental model in swine. 2014 , 175, 138-46	27
426	Design and rationale of the TOTAL trial: a randomized trial of routine aspiration Thrombectomy with percutaneous coronary intervention (PCI) versus PCI ALone in patients with ST-elevation myocardial infarction undergoing primary PCI. 2014 , 167, 315-321.e1	58

425	Thrombus burden and myocardial damage during primary percutaneous coronary intervention. 2014 , 113, 1449-56	35
424	Diaphragm border detection in coronary X-ray angiographies: New method and applications. 2014 , 38, 296-305	8
423	Efficacy of early intensive rosuvastatin therapy in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention (ROSEMARY Study). 2014 , 114, 29-35	10
422	Impact of advanced age on myocardial perfusion, distal embolization, and mortality patients with ST-segment elevation myocardial infarction treated by primary angioplasty and glycoprotein IIb-IIIa inhibitors. 2014 , 29, 15-20	15
421	The value of plasma D-dimer level on admission in predicting no-reflow after primary percutaneous coronary intervention and long-term prognosis in patients with acute ST segment elevation myocardial infarction. 2014 , 38, 339-47	26
420	Chronic metformin treatment is associated with reduced myocardial infarct size in diabetic patients with ST-segment elevation myocardial infarction. 2014 , 28, 163-71	41
419	Impact of nonculprit vessel myocardial perfusion on outcomes of patients undergoing percutaneous coronary intervention for acute coronary syndromes: analysis from the ACUTY trial (Acute Catheterization and Urgent Intervention Triage Strategy). 2014 , 7, 266-75	2
418	Impact of cigarette smoking on extent of coronary artery disease and prognosis of patients with non-ST-segment elevation acute coronary syndromes: an analysis from the ACUTY Trial (Acute Catheterization and Urgent Intervention Triage Strategy). 2014 , 7, 372-9	34
417	Early change in invasive measures of microvascular function can predict myocardial recovery following PCI for ST-elevation myocardial infarction. 2014 , 35, 1971-80	52
416	Prognostic value of angiographic lesion complexity in patients with acute coronary syndromes undergoing percutaneous coronary intervention (from the acute catheterization and urgent intervention triage strategy trial). 2014 , 114, 1638-45	6
415	Pre-angiography total ST-segment resolution is not a reliable predictor of an open infarct-related artery. 2014 , 25, 826-30	1
414	Impact of myocardial reperfusion status on QT dispersion after successful recanalization of the infarct-related artery in acute myocardial infarction. 2014 , 27, 252-9	5
413	Coronary thrombus composition: links with inflammation, platelet and endothelial markers. 2014 , 237, 555-61	54
412	Impact of microvascular obstruction on the assessment of coronary flow reserve, index of microcirculatory resistance, and fractional flow reserve after ST-segment elevation myocardial infarction. 2014 , 64, 1894-904	99
411	Coronary thrombi neovascularization in patients with ST-elevation myocardial infarction - clinical and angiographic implications. 2014 , 134, 1038-45	4
410	Effect of microvascular obstruction and intramyocardial hemorrhage by CMR on LV remodeling and outcomes after myocardial infarction: a systematic review and meta-analysis. 2014 , 7, 940-52	143
409	Impact of overweight on myocardial infarct size in patients undergoing primary percutaneous coronary intervention: a magnetic resonance imaging study. 2014 , 235, 570-5	9
408	Myocardial tissue perfusion predicts the evolution of fragmented QRS in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2014 , 19, 454-61	13

407	Impact of delay to reperfusion on reperfusion success, infarct size, and clinical outcomes in patients with ST-segment elevation myocardial infarction: the INFUSE-AMI Trial (INFUSE-Anterior Myocardial Infarction). 2014 , 7, 733-40	38
406	Impact of white blood cell count on myocardial salvage, infarct size, and clinical outcomes in patients undergoing primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: a magnetic resonance imaging study. 2014 , 30, 129-36	15
405	Development of a theory for generating regional cardiac perfusion images during coronary angiography in the coronary angiography lab. 2014 , 30, 9-19	6
404	Microvascular obstruction in patients with non-ST-elevation myocardial infarction: a contrast-enhanced cardiac magnetic resonance study. 2014 , 30, 1087-95	12
403	Intracoronary thrombolysis in patients with ST-segment elevation myocardial infarction presenting with massive intraluminal thrombus and failed aspiration. 2014 , 3, 229-36	24
402	Association between local interleukin-6 levels and slow flow/microvascular dysfunction. 2014 , 37, 475-82	11
401	Angiographic evolution of myocardial perfusion impairment in a single case of Tako-Tsubo like syndrome. 2014 , 172, e82-3	
400	Assessment of coronary blood flow in the cardiac catheterization laboratory. 2014 , 39, 159-84	1
399	Outcomes in patients with ST-segment elevation acute myocardial infarction treated with clopidogrel versus prasugrel (from the INFUSE-AMI trial). 2014 , 113, 1457-60	32
398	Comparison of the frequencies of myocardial edema determined by cardiac magnetic resonance in diabetic versus nondiabetic patients having percutaneous coronary intervention for ST elevation myocardial infarction. 2014 , 113, 607-12	9
397	Relation of ST-segment elevation before and after percutaneous transluminal coronary angioplasty to left ventricular area at risk, myocardial infarct size, and systolic function. 2014 , 113, 593-600	14
396	Use of electrocardiogram indices of myocardial ischemia for risk stratification and decision making of reperfusion strategies. 2014 , 47, 520-4	2
395	The effect of thrombus aspiration during primary percutaneous coronary intervention on clinical outcome in daily clinical practice. 2014 , 111, 165-71	14
394	Relationships of elevated systemic pentraxin-3 levels with high-risk coronary plaque components and impaired myocardial perfusion after percutaneous coronary intervention in patients with ST-elevation acute myocardial infarction. 2014 , 78, 159-69	16
393	Postconditioning reduces enzymatic infarct size and improves microvascular reperfusion in patients with ST-segment elevation myocardial infarction. 2014 , 129, 250-7	17
392	The 'MAP strategy' (Maximum aspiration of atherothrombus and adjunctive glycoprotein IIb/IIIa inhibitor utilization combined with prolonged inflation of balloon/stent) for preventing no-reflow in patients with ST-segment elevation myocardial infarction undergoing percutaneous coronary intervention: A retrospective analysis of seventy-one cases. 2015 , 67 Suppl 3, S43-6	4
391	A risk score for no reflow in patients with ST-segment elevation myocardial infarction after primary percutaneous coronary intervention. 2015 , 38, 208-15	20
390	Association of Intravascular Ultrasound- and Optical Coherence Tomography-Assessed Coronary Plaque Morphology With Periprocedural Myocardial Injury in Patients With Stable Angina Pectoris. 2015 , 79, 1944-53	15

- 389 Matrix Metalloproteinase-9 in the Culprit Coronary Artery and Myocardial No-Reflow. **2015**, 350, 352-6 2
- 388 Association of lymphocyte-to-monocyte ratio with the no-reflow phenomenon in patients who underwent a primary percutaneous coronary intervention for ST-elevation myocardial infarction. **2015**, 26, 706-12 38
- 387 The impact of initial and residual thrombus burden on the no-reflow phenomenon in patients with ST-segment elevation myocardial infarction. **2015**, 26, 245-53 9
- 386 Cardiac troponin elevation pattern in patients undergoing a primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: characterization and relationship with cardiovascular events during hospitalization. **2015**, 26, 503-9 10
- 385 QRS duration: a novel marker of microvascular reperfusion as assessed by myocardial blush grade in ST elevation myocardial infarction patients undergoing a primary percutaneous intervention. **2015**, 26, 583-6 6
- 384 Effects of prasugrel pretreatment on angiographic myocardial perfusion parameters in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. **2015**, 26, 665-70 3
- 383 Effect of Intravenous Administration of Liposomal Prostaglandin E1 on Microcirculation in Patients with ST Elevation Myocardial Infarction Undergoing Primary Percutaneous Intervention. **2015**, 128, 1147-50 8
- 382 Adenosina intracoronaria durante la angioplastia primaria para la preservaci3n del flujo coronario. **2015**, 34, 93-99
- 381 Effects of high-dose atorvastatin pretreatment in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention: a cardiac magnetic resonance study. **2015**, 30, 435-41 3
- 380 The MGuard coronary stent: safety, efficacy, and clinical utility. **2015**, 11, 533-9 8
- 379 The Relationship among N-Terminal Pro-B-Type Natriuretic Peptide, High-Sensitivity C-Reactive Protein and Infarct Size in Patients with Acute ST-Elevation Myocardial Infarction. **2015**, 45, 285-93 7
- 378 Coronary Microcirculation: A Clinical Perspective. **2015**, 639-654
- 377 Treatment of Acute ST-Elevation Myocardial Infarction. **2015**, 505-532
- 376 Quantification of manual thrombus removal in patients with acute coronary syndromes: a study exploiting serial frequency domain-optical coherence tomography. **2015**, 16, 204-12 4
- 375 Doppler-derived intracoronary physiology indices predict the occurrence of microvascular injury and microvascular perfusion deficits after angiographically successful primary percutaneous coronary intervention. **2015**, 8, e001786 42
- 374 Hypotestosteronemia is frequent in ST-elevation myocardial infarction patients and is associated with coronary microvascular obstruction. **2015**, 22, 855-63 4
- 373 Impact of Nonalcoholic Fatty Liver Disease on Myocardial Perfusion in Nondiabetic Patients Undergoing Primary Percutaneous Coronary Intervention for ST-Segment Elevation Myocardial Infarction. **2015**, 116, 1810-4 24
- 372 Effect of Ischemia Duration and Door-to-Balloon Time on Myocardial Perfusion in ST-Segment Elevation Myocardial Infarction: An Analysis From HORIZONS-AMI Trial (Harmonizing Outcomes with Revascularization and Stents in Acute Myocardial Infarction). **2015**, 8, 1966-1974 41

371	Postconditioning attenuates early ventricular arrhythmias in patients with high-risk ST-segment elevation myocardial infarction. 2015 , 65, 459-65	9
370	Diagnosis of coronary microvascular dysfunction - Present status. 2015 , 67, 552-60	9
369	Remote ischemic conditioning. 2015 , 65, 177-95	391
368	Utility of peak creatine kinase-MB measurements in predicting myocardial infarct size, left ventricular dysfunction, and outcome after first anterior wall acute myocardial infarction (from the INFUSE-AMI trial). 2015 , 115, 563-70	44
367	Association of mean platelet volume with angiographic thrombus burden and short-term mortality in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2015 , 85 Suppl 1, 724-33	14
366	Abnormal left ventricular contractile response to exercise in the absence of obstructive coronary artery disease is associated with resting left ventricular long-axis dysfunction. 2015 , 28, 95-105	4
365	Implications of ventricular arrhythmia "bursts" with normal epicardial flow, myocardial blush, and ST-segment recovery in anterior ST-elevation myocardial infarction reperfusion: a biosignature of direct myocellular injury "downstream of downstream". 2015 , 4, 51-9	20
364	Development and Validation of a Distal Embolization Risk Score During Primary Angioplasty in ST-Elevation Myocardial Infarction. 2015 , 116, 1172-8	2
363	Invasive assessment of the coronary microcirculation in the catheter laboratory. 2015 , 199, 141-9	8
362	Epicardial adipose tissue thickness is associated with myocardial infarction and impaired coronary perfusion. 2015 , 15, 224-31	13
361	Is Aspiration Thrombectomy Beneficial in Patients Undergoing Primary Percutaneous Coronary Intervention? Meta-Analysis of Randomized Trials. 2015 , 8, e002258	52
360	Thrombus aspiration in patients with ST elevation myocardial infarction: meta-analysis of 16 randomized trials. 2015 , 15, 175-87	3
359	A Combined Optical Coherence Tomography and Intravascular Ultrasound Study on Plaque Rupture, Plaque Erosion, and Calcified Nodule in Patients With ST-Segment Elevation Myocardial Infarction: Incidence, Morphologic Characteristics, and Outcomes After Percutaneous Coronary Intervention. 2015 , 8, 1166-1176	150
358	Aspiration Thrombectomy for Treatment of ST-segment Elevation Myocardial Infarction: a Meta-analysis of 26 Randomized Trials in 11,943 Patients. 2015 , 68, 746-52	4
357	Impairment of pulmonary function and changes in the right cardiac structure of pneumoconiotic coal workers in China. 2015 , 28, 62-70	6
356	Novel insights into an "old" phenomenon: the no reflow. 2015 , 187, 273-80	76
355	Usefulness of the Left Anterior Descending Artery Wrapping Around the Left Ventricular Apex to Predict Adverse Clinical Outcomes in Patients With Anterior Wall ST-Segment Elevation Myocardial Infarction (an INFUSE-AMI Substudy). 2015 , 115, 1389-95	15
354	Effect of ischemic postconditioning on myocardial salvage in patients undergoing primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: cardiac magnetic resonance substudy of the POST randomized trial. 2015 , 31, 629-37	19

353	Development of a method for automated and stable myocardial perfusion measurement using coronary X-ray angiography images. 2015 , 31, 905-14	4
352	Intracoronary epinephrine in the treatment of refractory no-reflow after primary percutaneous coronary intervention: a retrospective study. 2015 , 15, 10	21
351	Soluble interleukin 6 receptor levels are associated with reduced myocardial reperfusion after percutaneous coronary intervention for acute myocardial infarction. 2015 , 73, 207-12	11
350	Use and impact of thrombectomy in primary percutaneous coronary intervention for acute myocardial infarction with persistent ST-segment elevation: results of the prospective ALKK PCI-registry. 2015 , 104, 803-11	7
349	Zero-Flow Pressure Measured Immediately After Primary Percutaneous Coronary Intervention for ST-Segment Elevation Myocardial Infarction Provides the Best Invasive Index for Predicting the Extent of Myocardial Infarction at 6 Months: An OxAMI Study (Oxford Acute Myocardial Infarction). 2015 , 6, 1410-1421	35
348	Usefulness of the Left Anterior Descending Coronary Artery Wrapping Around the Left Ventricular Apex to Predict Adverse Clinical Outcomes in Patients With Anterior Wall ST-Segment Elevation Myocardial Infarction (from the Harmonizing Outcomes With Revascularization and Stents in Acute Myocardial Infarction Trial). 2015 , 116, 1158-65	12
347	Effects of liraglutide on left ventricular function in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2015 , 170, 845-54	76
346	Evaluation of Microvascular Disease and Clinical Outcomes. 2015 , 4, 443-457	
345	How does coronary stent implantation impact on the status of the microcirculation during primary percutaneous coronary intervention in patients with ST-elevation myocardial infarction?. 2015 , 36, 3165-77	60
344	Alternative stents in ST-segment elevation myocardial infarction: improving the efficacy of primary percutaneous coronary intervention. 2015 , 11, 347-57	1
343	Adenosine as an Adjunct Therapy in ST Elevation Myocardial Infarction Patients: Myth or Truth?. 2015 , 29, 481-93	7
342	Impact of n-3 polyunsaturated fatty acids in predicting ischemia/reperfusion injury and progression of myocardial damage after reperfusion in patients with ST-segment elevation acute myocardial infarction. 2015 , 66, 101-7	9
341	Effect of Diltiazem on Coronary Artery Flow and Myocardial Perfusion in Patients With Isolated Coronary Artery Ectasia and Either Stable Angina Pectoris or Positive Myocardial Ischemic Stress Test. 2015 , 116, 1199-203	4
340	The no-reflow phenomenon: State of the art. 2015 , 108, 661-74	90
339	Trombectomía por aspiración para el tratamiento del infarto agudo de miocardio con elevación del segmento ST: un metanálisis de 26 ensayos aleatorizados con 11.943 pacientes. 2015 , 68, 746-752	12
338	Effects of tirofiban on platelet activation and endothelial function in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2015 , 71, 135-42	11
337	Impact of thrombus burden on outcomes after standard versus mesh-covered stents in acute myocardial infarction (from the MGuard for acute ST elevation reperfusion trial). 2015 , 115, 161-6	16
336	Treating thrombus in the coronary arteries. 2015 , 40, 60-5	2

335	Real world comparison of the MGuard Stent versus the bare metal stent for ST elevation myocardial infarction (the REWARD-MI study). 2015 , 85, E1-9	11
334	Coronary microvascular dysfunction: mechanisms and functional assessment. 2015 , 12, 48-62	262
333	Baseline SYNTAX score and long-term outcome in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2015 , 21, 712-9	17
332	Efficacy of Intra-aortic Balloon Pump before versus after Primary Percutaneous Coronary Intervention in Patients with Cardiogenic Shock from ST-elevation Myocardial Infarction. 2016 , 129, 1400-5	6
331	Pathophysiological aspects and management workflow of coronary microvascular obstruction in ST-segment elevation myocardial infarction. 2016 , 10, 10	
330	Thrombus aspiration in acute myocardial infarction: concepts, clinical trials, and current guidelines. 2016 , 27, 233-43	3
329	Effect of a combination of antiplatelet and antithrombotic pretreatment on myocardial perfusion in patients with an acute ST-segment elevation myocardial infarction undergoing a primary percutaneous coronary intervention. 2016 , 27, 580-5	
328	Manual Thrombus Aspiration and the Improved Survival of Patients With Unstable Angina Pectoris Treated With Percutaneous Coronary Intervention (30 Months Follow-Up). 2016 , 95, e2919	4
327	Impact of oral P2Y12 inhibitors on residual thrombus burden and reperfusion indexes in patients with ST-segment elevation myocardial infarction. 2016 , 17, 701-6	3
326	Impact of thrombus burden on procedural and mid-term outcomes after primary percutaneous coronary intervention. 2016 , 27, 169-75	5
325	Concordance of angiographic and electrocardiographic indexes of microvascular obstruction: myocardial haemorrhage role. 2016 , 17, 382-91	2
324	Prognostic utility of myocardial blush grade after PCI in patients with NSTEMI-ACS: Analysis from the ACUITY trial. 2016 , 88, 215-24	5
323	Ischemic Postconditioning After Routine Thrombus Aspiration During Primary Percutaneous Coronary Intervention: Rationale and Design of the POCconditioning Rotterdam Trial. 2016 , 88, 508-514	2
322	Association of mean platelet volume with impaired myocardial reperfusion and short-term mortality in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2016 , 27, 5-12	7
321	Plasma DPP4 activity is associated with no-reflow and major bleeding events in Chinese PCI-treated STEMI patients. 2016 , 6, 39412	5
320	Effects of Liraglutide on Reperfusion Injury in Patients With ST-Segment-Elevation Myocardial Infarction. 2016 , 9,	58
319	INNOVATION Study (Impact of Immediate Stent Implantation Versus Deferred Stent Implantation on Infarct Size and Microvascular Perfusion in Patients With ST-Segment-Elevation Myocardial Infarction). 2016 , 9,	20
318	Impact of the origin of the collateral feeding donor artery on short-term mortality in ST-elevation myocardial infarction with comorbid chronic total occlusion. 2016 , 218, 158-163	11

317	Impact of myocardial staining on In-hospital outcome after primary percutaneous coronary intervention in the Padua Registry on ST-elevation myocardial infarction. 2016 , 212, 352-4	
316	Manual Aspiration Thrombectomy in Acute Myocardial Infarction: A Clinical Experience. 2016 , 25, 20-8	
315	Non-contrast cardiac CT immediately after percutaneous coronary intervention: does it predict the risk of left ventricular remodeling in patients with ST-elevation myocardial infarction?. 2016 , 32 Suppl 1, 147-54	
314	Myocardial blush and microvascular reperfusion following manual thrombectomy during percutaneous coronary intervention for ST elevation myocardial infarction: insights from the TOTAL trial. 2016 , 37, 1891-8	26
313	The Coronary Circulation as a Target of Cardioprotection. 2016 , 118, 1643-58	142
312	Beneficial effect of early infusion of landiolol, a very short-acting beta-1 adrenergic receptor blocker, on reperfusion status in acute myocardial infarction. 2016 , 221, 321-6	5
311	Increased Platelet-leukocyte Aggregates Are Associated With Myocardial No-reflow in Patients With ST Elevation Myocardial Infarction. 2016 , 352, 261-6	26
310	Shock Index as a Predictor of Myocardial Injury in ST-segment Elevation Myocardial Infarction. 2016 , 352, 574-581	11
309	Does Residual Thrombus After Aspiration Thrombectomy Affect the Outcome of Primary PCI in Patients With ST-Segment Elevation Myocardial Infarction?: An Optical Coherence Tomography Study. 2016 , 9, 2002-2011	33
308	Complications of Percutaneous Coronary Intervention. 2016 ,	3
307	Reducing Microvascular Dysfunction in Revascularized Patients with ST-Elevation Myocardial Infarction by Off-Target Properties of Ticagrelor versus Prasugrel. Rationale and Design of the REDUCE-MVI Study. 2016 , 9, 249-256	12
306	No-Reflow Phenomenon. 2016 , 177-183	
305	Aspiration thrombectomy and intracoronary tirofiban in ST-segment elevation myocardial infarction : Combination treatment for patients undergoing primary percutaneous coronary intervention. 2016 , 41, 732-740	6
304	Serious infections among unselected patients with ST-elevation myocardial infarction treated with contemporary primary percutaneous coronary intervention. 2016 , 181, 52-59	17
303	Current Considerations of Thrombectomy for Acute Myocardial Infarction. 2016 , 1, 265-272	4
302	Coronary Plaque Characteristics Associated With Reduced TIMI (Thrombolysis in Myocardial Infarction) Flow Grade in Patients With ST-Segment-Elevation Myocardial Infarction: A Combined Optical Coherence Tomography and Intravascular Ultrasound Study. 2016 , 9,	10
301	Impact of optical coherence tomography- and coronary angiography-assessed neointimal tissue characteristics on occurrence of periprocedural myonecrosis in patients with in-stent restenosis. 2016 , 32, 1483-94	3
300	Comparative Prognostic Utility of Indexes of Microvascular Function Alone or in Combination in Patients With an Acute ST-Segment-Elevation Myocardial Infarction. <i>Circulation</i> , 2016 , 134, 1833-1847	16.7 89

299	Remote ischaemic preconditioning reduces myocardial ischaemic reperfusion injury in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2016 , 71, 596-603	20
298	Association of serum cystatin C levels with myocardial perfusion and cardiac functional recovery in patients with anterior wall ST elevation myocardial infarction treated with primary coronary intervention. 2016 , 31, 1456-66	7
297	Influence of microvascular dysfunction on regional myocardial deformation post-acute myocardial infarction: insights from a novel angiographic index for assessing myocardial tissue-level reperfusion. 2016 , 32, 711-9	4
296	Chronic ischemic mitral regurgitation and papillary muscle infarction detected by late gadolinium-enhanced cardiac magnetic resonance imaging in patients with ST-segment elevation myocardial infarction. 2016 , 105, 981-991	14
295	Left atrial volume index as a predictor of left ventricular remodeling in patients with anterior STEMI treated with primary PCI. 2016 , 68, 17-22	0
294	Effect of admission glycometabolic state on clinical outcome in non diabetic subjects with acute st segment elevation myocardial infarction. 2016 , 4, 73-78	2
293	Upstream clopidogrel, prasugrel, or ticagrelor for patients treated with primary angioplasty: Results of an angiographic randomized pilot study. 2016 , 87, 1187-93	3
292	The relationship between admission monocyte HDL-C ratio with short-term and long-term mortality among STEMI patients treated with successful primary PCI. 2016 , 27, 176-84	31
291	Adiponectin and ischemia-reperfusion injury in ST segment elevation myocardial infarction. 2016 , 5, 71-6	16
290	Toll-like receptor-4 is upregulated in plaque debris of patients with acute coronary syndrome more than Toll-like receptor-2. 2016 , 31, 1-5	15
289	B-type natriuretic peptide as a predictor of ischemia/reperfusion injury immediately after myocardial reperfusion in patients with ST-segment elevation acute myocardial infarction. 2016 , 5, 62-70	5
288	Reperfusion Therapy for Acute Myocardial Infarction. 2016 ,	
287	Mechanical Reperfusion for STEMI. 2016 ,	
286	Impact of residual platelet reactivity on reperfusion in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2016 , 5, 475-86	12
285	A protective role of early collateral blood flow in patients with ST-segment elevation myocardial infarction. 2016 , 171, 56-63	27
284	Comparison of Outcome of Patients With ST-Segment Elevation Myocardial Infarction and Complete Versus Incomplete ST-Resolution Before Primary Percutaneous Coronary Intervention. 2016 , 117, 1735-40	9
283	Effects of liraglutide on no-reflow in patients with acute ST-segment elevation myocardial infarction. 2016 , 208, 109-14	23
282	Ventricular arrhythmia burst is an independent indicator of larger infarct size even in optimal reperfusion in STEMI. 2016 , 49, 345-52	5

281	Thrombus aspiration in acute myocardial infarction. 2016 , 13, 418-28	27
280	Frame counting improves the assessment of post-reperfusion microvascular patency by TIMI myocardial perfusion grade: Evidence from cardiac magnetic resonance imaging. 2016 , 203, 360-6	13
279	Coronary microvascular obstruction in acute myocardial infarction. 2016 , 37, 1024-33	201
278	Effect of Hematologic Parameters on Microvascular Reperfusion in Patients With ST-Segment Elevation Myocardial Infarction Treated With Primary Percutaneous Coronary Intervention. 2016 , 67, 151-6	8
277	Fenómeno de no-reflow coronario: revisión actualizada de un fenómeno antiguo. 2017 , 52, 155-161	
276	Coronary flow improvement following unsuccessful primary percutaneous coronary intervention in ST-elevation myocardial infarction with diffuse ectatic coronary artery. 2017 , 6, 623-631	7
275	Assessing MICRO-vascular resistances via IMR to predict outcome in STEMI patients with multivessel disease undergoing primary PCI (AMICRO): Rationale and design of a prospective multicenter clinical trial. 2017 , 187, 37-44	0
274	Invasive Assessment of the Coronary Microcirculation in Reperused ST-Segment-Elevation Myocardial Infarction Patients: Where Do We Stand?. 2017 , 10,	28
273	A review of strategies for infarct size reduction during acute myocardial infarction. 2017 , 18, 374-383	7
272	Clinical outcome and correlates of coronary microvascular obstruction in latecomers after acute myocardial infarction. 2017 , 236, 30-35	11
271	Targeting the dominant mechanism of coronary microvascular dysfunction with intracoronary physiology tests. 2017 , 33, 1041-1059	24
270	Clinical Significance of Postinfarct Fever in ST-Segment Elevation Myocardial Infarction: A Cardiac Magnetic Resonance Imaging Study. 2017 , 6,	5
269	P wave peak time; a novel electrocardiographic parameter in the assessment of coronary no-reflow. 2017 , 50, 584-590	11
268	Relationship between R-wave peak time and no-reflow in ST elevation myocardial infarction treated with a primary percutaneous coronary intervention. 2017 , 28, 326-331	6
267	Identification of High-Risk Patients After ST-Segment-Elevation Myocardial Infarction: Comparison Between Angiographic and Magnetic Resonance Parameters. 2017 , 10, e005841	17
266	Role of deferred stenting in patients with ST elevation myocardial infarction treated with primary percutaneous coronary intervention: A systematic review and meta-analysis. 2017 , 30, 264-273	15
265	Collateral filling efficiency of comorbid chronic total occlusion segment on short-term mortality in ST-elevation myocardial infarction. 2017 , 230, 346-352	2
264	Early peripheral endothelial dysfunction predicts myocardial infarct extension and microvascular obstruction in patients with ST-elevation myocardial infarction. 2017 , 36, 731-742	9

263	High-sensitivity C-reactive protein and long term reperfusion success of primary percutaneous intervention in ST-elevation myocardial infarction. 2017 , 248, 51-56	12
262	The influence of coronary plaque morphology assessed by optical coherence tomography on final microvascular function after stenting in patients with ST-elevation myocardial infarction. 2017 , 28, 198-208	3
261	Anatomical and physiological complications related to left ventricular apical aneurysm. 2017 , 4, 70-75	
260	The Coronary Circulation in Acute Coronary Syndromes. 2017 , 99-110	
259	Invasive Assessment of the Coronary Microvasculature: The Index of Microcirculatory Resistance. 2017 , 10,	46
258	Early peripheral endothelial dysfunction predicts myocardial infarct extension and microvascular obstruction in patients with ST-elevation myocardial infarction. 2017 , 36, 731-742	3
257	Prognostic value of clinical, echocardiographic and angiographic indicators in patients with large anterior ST-segment elevation myocardial infarction as a first acute coronary event. 2017 , 18, 946-953	2
256	Efficacy and safety of intracoronary verapamil versus sodium nitroprusside for the prevention of microvascular obstruction during primary percutaneous coronary intervention for ST-segment elevation myocardial infarction. 2017 , 28, 11-16	4
255	A prospective evaluation of a standardized strategy for the use of a polymeric everolimus-eluting bioresorbable scaffold in ST-segment elevation myocardial infarction: Rationale and design of the BVS STEMI STRATEGY-IT study. 2017 , 89, 1129-1138	7
254	Indications and outcomes of excimer laser coronary atherectomy: Efficacy and safety for thrombotic lesions-The ULTRAMAN registry. 2017 , 69, 314-319	29
253	Comparison of the Infarct Size between the Loading of Ticagrelor and Clopidogrel in Patients with Acute Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. 2017 , 47, 705-713	5
252	Symptom-to-balloon time and myocardial blush grade are predictors of left ventricular remodelling after successful primary percutaneous coronary intervention. 2017 , 28, 186-190	1
251	QRS Complex Duration is a Marker of Reperfusion in Patients Presenting with Acute ST Segment Elevation Myocardial Infarction. 2017 , 05,	
250	Glycemic variability is associated with myocardial damage in nondiabetic patients with ST-elevation myocardial infarction. 2018 , 7, 47-53	5
249	Elevated serum uric acid affects myocardial reperfusion and infarct size in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2018 , 19, 240-246	10
248	Clinical impact of direct stenting and interaction with thrombus aspiration in patients with ST-segment elevation myocardial infarction undergoing percutaneous coronary intervention: Thrombectomy Trialists Collaboration. 2018 , 39, 2472-2479	15
247	Usefulness of The C-Reactive Protein/Albumin Ratio for Predicting No-Reflow in ST-elevation myocardial infarction treated with primary percutaneous coronary intervention. 2018 , 48, e12928	32
246	Markers of Reperfusion and Long-Term (8-Year) Prognosis after Primary Percutaneous Coronary Intervention. 2018 , 122, 39-46	4

245	Cost assessment of treatment of acute myocardial infarction and angiographically visible coronary thrombus. 2018 , 7, 471-481	
244	A Randomized Double-Blind Placebo-Controlled Study Comparing Intracoronary Versus Intravenous Abciximab in Patients With ST-Elevation Myocardial Infarction Undergoing Transradial Rescue Percutaneous Coronary Intervention After Failed Thrombolysis. 2018 , 122, 47-53	1
243	Outcomes of direct stenting in patients with ST-elevated myocardial infarction. 2018 , 43, 447-454	2
242	Invasive Assessment for Microcirculation. 2018 , 303-310	
241	Intramyocardial dissecting hematoma in anterior wall ST elevation myocardial infarction: impact on left ventricular remodeling and prognosis. 2018 , 34, 201-210	8
240	Prasugrel versus clopidogrel for residual thrombus burden in patients with ST-segment elevation myocardial infarction: an optical coherence tomography study. 2018 , 29, 663-669	3
239	A Novel Concept in the Progression of Acute Coronary Syndrome: Acute Coronary Continuum (ACC). 2018 , 07,	
238	Effect of hemorheological parameters on myocardial injury after primary or elective percutaneous coronary intervention. 2018 , 29, 638-646	2
237	Fractional Flow Reserve Versus Angiographically-Guided Coronary Artery Bypass Grafting. 2018 , 72, 2732-2743	47
236	Primary PCI versus pharmacoinvasive strategy for ST elevation myocardial infarction. 2018 , 21, 87-93	5
235	Dedicated Thrombus-Containing Stent Platforms. 2018 , 285-302	
234	Acute Myocardial Infarction: STEMI and NSTEMI. 2018 , 123-146	0
233	Correlation of QRS duration with myocardial blush grade as a marker of myocardial reperfusion in primary percutaneous coronary intervention. 2018 , 70 Suppl 3, S359-S364	3
232	Coronary Thrombolysis Waveforms After Acute Reperused ST-Segment-Elevation Myocardial Infarction: Relation to Microvascular Obstruction and Prognosis. 2018 , 7, e008957	4
231	The clinical significance of echo-attenuated plaque in stable angina pectoris compared with acute coronary syndromes: A combined intravascular ultrasound and optical coherence tomography study. 2018 , 270, 1-6	3
230	Right ventricular function in patients presenting with non-ST-segment elevation myocardial infarction undergoing an invasive approach. 2018 , 70, 149-153	1
229	Takotsubo Cardiomyopathy: One More Angiographic Evidence of Microvascular Dysfunction. 2018 , 2018, 5281485	8
228	Intravascular Ultrasound and Angioscopy Assessment of Coronary Plaque Components in Chronic Totally Occluded Lesions. 2018 , 82, 2032-2040	4

227	Incidence and Significance of Spontaneous ST Segment Re-elevation After Reperfused Anterior Acute Myocardial Infarction - Relationship With Infarct Size, Adverse Remodeling, and Events at 1 Year. 2018 , 82, 1379-1386	6
226	Epidemiology of Coronary Microvascular Obstruction. 2018 , 53-68	
225	Angiography and Electrocardiography (ECG) for the Assessment of Coronary Microvascular Obstruction. 2018 , 109-126	
224	Invasive Assessment of Coronary Microvascular Obstruction. 2018 , 127-153	
223	Impact of total ischemic time on manual thrombus aspiration benefit during primary percutaneous coronary intervention. 2018 , 204, 34-42	2
222	No-Reflow Phenomenon. 2018 , 1007-1015	
221	Thrombus aspiration in late presenters with ST-elevation myocardial infarction: A single-center randomized trial. 2018 , 31, 632-639	1
220	Microvascular perfusion in infarcted and remote myocardium after successful primary PCI: angiographic and CMR findings. 2019 , 29, 941-950	5
219	Early versus delayed invasive strategy in patients with non-ST-elevation acute coronary syndrome and concomitant congestive heart failure. 2019 , 74, 320-327	1
218	Pathophysiology, Diagnosis, and Management of the No-Reflow Phenomenon. 2019 , 33, 589-597	19
217	Assessment of right ventricular function after successful revascularization for acute anterior myocardial infarction without right ventricular infarction by echocardiography. 2019 , 31, 261-268	3
216	The role of Glucagon-Like Peptide 1 Loading on periprocedural myocardial infarction During elective PCI (GOLD-PCI study): A randomized, placebo-controlled trial. 2019 , 215, 41-51	4
215	Recurrence of angina after ST-segment elevation myocardial infarction: the role of coronary microvascular obstruction. 2019 , 2048872619880661	
214	Coronary microvascular obstruction: the new frontier in cardioprotection. 2019 , 114, 45	106
213	ROUTINELY AVAILABLE BIOMARKERS AS LONG-TERM PREDICTORS OF DEVELOPING SYSTOLIC DYSFUNCTION IN COMPLETELY REVASCULARIZED PATIENTS WITH ACUTE ST ELEVATION MYOCARDIAL INFARCTION. 2019 , 58, 95-102	2
212	Tissue Doppler-derived atrial dyssynchrony predicts new-onset atrial fibrillation during hospitalization for ST-elevation myocardial infarction. 2019 , 36, 1799-1805	1
211	Fragmented QRS and QRS Duration As a Marker of Myocardial Reperfusion Measured by Myocardial Blush Grade in Reperfusion Therapy: Systematic Review and Meta-Analysis. 2019 , 28, 255-261	0
210	Relationship between the ST-Segment Resolution and Microvascular Dysfunction in Patients Who Underwent Primary Percutaneous Coronary Intervention. 2019 , 2019, 8695065	2

209	Impact of intracoronary contrast injection pressure on reperfusion during primary percutaneous coronary intervention in acute ST-segment elevation myocardial infarction: A prospective randomized pilot study. 2019 , 24, 100412	1
208	Thrombus aspiration catheter improve the myocardial reperfusion of STEMI patients with high thrombus load during the emergency PCI operation. 2019 , 14, 172	5
207	Clinical and procedural predictors of suboptimal myocardial reperfusion in primary percutaneous coronary intervention. 2019 , 23, 100357	3
206	Comparison of 0.9-mm and 1.4-mm catheters in excimer laser coronary angioplasty for acute myocardial infarction. 2019 , 34, 1747-1754	5
205	Prognostic efficacy of C-reactive protein/albumin ratio in ST elevation myocardial infarction. 2019 , 53, 83-90	38
204	Efficacy of IntraCoronary Erythropoietin Delivery BEfore Reperfusion-Gauging Infarct Size in Patients with Acute ST-segment Elevation Myocardial Infarction (ICEBERG). 2019 , 60, 255-263	6
203	Optimal strategy of primary percutaneous coronary intervention for acute myocardial infarction due to unprotected left main coronary artery occlusion (OPTIMAL): study protocol for a randomised controlled trial. 2019 , 20, 162	2
202	Significado clíñico de los cambios recíprocos del segmento ST en pacientes con IAMCEST: estudio de imagen con resonancia magnética cardíaca. 2019 , 72, 120-129	1
201	Incremental Value of Coronary Microcirculation Resistive Reserve Ratio in Predicting the Extent of Myocardial Infarction in Patients with STEMI. Insights from the Oxford Acute Myocardial Infarction (OxAMI) Study. 2019 , 20, 1148-1155	14
200	Season and myocardial injury in patients with ST-segment elevation myocardial infarction: A cardiac magnetic resonance imaging study. 2019 , 14, e0211807	2
199	Does pre-angiography Total ST-segment resolution reliably predict spontaneous reperfusion of the infarct-related artery in patients with acute myocardial infarction?. 2019 , 19, 264	1
198	Current Perspectives in Facilitated Angioplasty. 2019 , 26, e208-e212	2
197	Stent malapposition, strut coverage and atherothrombotic prolapse after percutaneous coronary interventions in ST-segment elevation myocardial infarction. 2019 , 20, 122-130	4
196	Effect of coronary collateral circulation on the prognosis of elderly patients with acute ST-segment elevation myocardial infarction treated with underwent primary percutaneous coronary intervention. 2019 , 98, e16502	3
195	The coronary circulation in acute myocardial ischaemia/reperfusion injury: a target for cardioprotection. 2019 , 115, 1143-1155	77
194	Plasma interleukin 6 levels are associated with cardiac function after ST-elevation myocardial infarction. 2019 , 108, 612-621	31
193	Clinical Significance of Reciprocal ST-segment Changes in Patients With STEMI: A Cardiac Magnetic Resonance Imaging Study. 2019 , 72, 120-129	1
192	Impact of elevated HbA1c on long-term mortality in patients presenting with acute myocardial infarction in daily clinical practice: insights from a 'real world' prospective registry of the Zwolle Myocardial Infarction Study Group. 2020 , 9, 616-625	3

191	Effects of Intracoronary Nicorandil on Myocardial Microcirculation and Clinical Outcomes in Patients with Acute Myocardial Infarction: A Meta-Analysis of Randomized Controlled Trials. 2020 , 20, 191-198	3
190	Residual atherothrombotic burden after primary percutaneous coronary intervention and myocardial reperfusion-An optical frequency domain imaging study. 2020 , 96, 91-97	3
189	Short-term safety and long-term benefits of stent postdilation after primary percutaneous coronary intervention: Results of a cohort study. 2020 , 95, 1249-1256	3
188	Microcirculation. 2020 ,	1
187	Pathophysiology and diagnosis of coronary microvascular dysfunction in ST-elevation myocardial infarction. 2020 , 116, 787-805	36
186	Mitral annular calcification predicted major cardiovascular events in patients presented with acute coronary syndrome and underwent percutaneous coronary intervention. 2020 , 75, 767-773	
185	State of the Art: No-Reflow Phenomenon. 2020 , 38, 563-573	9
184	Plasma calprotectin was associated with platelet activation and no-reflow phenomenon in acute coronary syndrome. 2020 , 20, 443	1
183	High Thrombus Burden in Patients With COVID-19 Presenting With ST-Segment Elevation Myocardial Infarction. 2020 , 76, 1168-1176	116
182	Predictive accuracy of lymphocyte-to-monocyte ratio and monocyte-to-high-density-lipoprotein-cholesterol ratio in determining the slow flow/no-reflow phenomenon in patients with non-ST-elevated myocardial infarction. 2020 , 31, 518-526	9
181	Safety and efficacy of intracoronary prourokinase administration in patients with high thrombus burden. 2020 , 31, 493-499	6
180	No-Reflow after PPCI-A Predictor of Short-Term Outcomes in STEMI Patients. 2020 , 9,	7
179	Association of microRNA-224-3p and microRNA-155-5p expressions with plasma long pentraxin 3 concentration and coronary microvascular obstruction following primary angioplasty for acute ST-segment elevation myocardial infarction. 2020 , 13, 499	1
178	Serum uric acid may modulate the inflammatory response after primary percutaneous coronary intervention in patients with ST-elevation myocardial infarction. 2020 , 21, 337-339	6
177	Efficacy and safety of intracoronary prourokinase during percutaneous coronary intervention in treating ST-segment elevation myocardial infarction patients: a randomized, controlled study. 2020 , 20, 308	5
176	Prolonged enoxaparin therapy compared with standard-of-care antithrombotic therapy in opiate-treated patients undergoing primary percutaneous coronary intervention. 2021 , 32, 555-559	2
175	Coronary Microvascular Dysfunction: An Update. 2020 , 16, 43-49	
174	Impact of Thrombus Aspiration on Clinical Outcomes in Korean Patients with ST Elevation Myocardial Infarction. 2020 , 56, 36-43	1

173	Assessment of Myocardial Perfusion by Angiographic Methods in Tortuous Coronary Arteries. 2020 , 71, 616-620	1
172	Relation of collateral circulation with reciprocal changes in patients with acute ST-elevation myocardial infarction. 2020 , 60, 36-43	1
171	Coronary slow flow is associated with a worse clinical outcome in patients with Takotsubo syndrome. 2020 , 106, 923-930	21
170	Predictive values of D-dimer for the long-term prognosis of acute ST-segment elevation infarction: A retrospective study in southwestern China. 2020 , 99, e19724	2
169	A Scoring System to Predict No-Reflow Phenomenon in Elective Percutaneous Coronary Intervention: The RECOVER Score. 2021 , 46, 100676	
168	Efficacy and safety of intracoronary epinephrine versus conventional treatments alone in STEMI patients with refractory coronary no-reflow during primary PCI: The RESTORE observational study. 2021 , 97, 602-611	7
167	MiR-98-5p promotes ischemia/reperfusion-induced microvascular dysfunction by targeting NGF and is a potential biomarker for microvascular reperfusion. 2021 , 28, e12657	5
166	Residual platelet reactivity after pre-treatment with ticagrelor prior to primary percutaneous coronary intervention is associated with suboptimal myocardial reperfusion. 2019 , 2048872619860214	
165	Prognostic role of neutrophil-to-lymphocyte ratio in patients with ST-elevation myocardial infarction undergoing to pharmaco-invasive strategy. 2021 , 34, 99-99	1
164	Clinical and echocardiographic predictors of left ventricular remodeling following anterior acute myocardial infarction. 2021 , 76, e2732	
163	Low Ankle-Brachial Index is Associated with Microvascular Coronary Obstruction After Primary PCI. 2021 , 17, 23-32	
162	Gensini Score and Thrombus Burden Add Predictive Value to the SYNTAX Score in Detecting No-Reflow after Myocardial Infarction. 2021 , 116, 466-472	0
161	Predictors of Microvascular Reperfusion After Myocardial Infarction. 2021 , 23, 21	1
160	Angiographic predictors of outcome in myocardial infarction patients presenting with cardiogenic shock: a CULPRIT-SHOCK angiographic substudy. 2021 , 16, e1237-e1244	3
159	Pathophysiology, Diagnosis, and Management of Coronary No-Reflow Phenomenon. 2021 , 30, 15-21	3
158	Pressure-bounded coronary flow reserve to assess the extent of microvascular dysfunction in patients with ST-elevation acute myocardial infarction. 2021 , 16, 1434-1443	0
157	Coronary Microvascular Dysfunction Assessed by Pressure Wire and CMR After STEMI Predicts Long-Term Outcomes. 2021 , 14, 1948-1959	6
156	Patients with COVID-19 who experience a myocardial infarction have complex coronary morphology and high in-hospital mortality: Primary results of a nationwide angiographic study. 2021 , 98, E370-E378	4

155	Impact of a micro-net mesh technology covering stent on coronary microvascular dysfunction in patients with high thrombus burden. 2021,	
154	Clinical outcome of patients with ST-elevation myocardial infarction and angiographic evidence of coronary artery ectasia. 2021,	2
153	Impact of Acute Insulin Resistance on Myocardial Blush in Non-Diabetic Patients Undergoing Primary Percutaneous Coronary Intervention. 2021, 8, 647366	
152	Predictors of no-reflow phenomenon following percutaneous coronary intervention for ST-segment elevation myocardial infarction. 2021, 70, 136-142	2
151	Associations of Lipoprotein(a) With Coronary Atherosclerotic Burden and All-Cause Mortality in Patients With ST-Segment Elevation Myocardial Infarction Treated With Primary Percutaneous Coronary Intervention. 2021, 8, 638679	0
150	SARS-COV-2 colonizes coronary thrombus and impairs heart microcirculation bed in asymptomatic SARS-CoV-2 positive subjects with acute myocardial infarction. 2021, 25, 217	15
149	Manual der Arbeitsgruppe Interventionelle Kardiologie (AGIK) der Deutschen Gesellschaft für Kardiologie – Herz- und Kreislaufforschung e. V. (DGK). 2021, 15, 370-403	1
148	Myocardial preservation during primary percutaneous intervention: It's time to rethink?. 2021, 73, 395-403	
147	Adenosine-Induced Coronary Steal Is Observed in Patients Presenting With ST-Segment-Elevation Myocardial Infarction. 2021, 10, e019899	1
146	COVID-19 and Acute Coronary Syndromes: From Pathophysiology to Clinical Perspectives. 2021, 2021, 4936571	6
145	The Predictive Value of Baseline Target Lesion SYNTAX Score for No-Reflow during Urgent Percutaneous Coronary Intervention in Acute Myocardial Infarction. 2021, 2021, 9987265	0
144	Acute Coronary Syndromes (ACS)-Unravelling Biology to Identify New Therapies-The Microcirculation as a Frontier for New Therapies in ACS. 2021, 10,	1
143	Go With the Flow: Another Method to Measure Coronary Microvascular Function. 2021, 14, 1685-1687	
142	Delayed endovascular surgery in patients with acute ST-segment elevation myocardial infarction due to massive culprit arterial thrombosis in the prevention of slow/no-reflow phenomenon. 2021, 20, 2761	1
141	Protective Effect of Shexiang Baixin Pill on Myocardial Ischemia/Reperfusion Injury in Patients With STEMI. 2021, 12, 721011	0
140	Safety and efficacy of a platelet glycoprotein Ib inhibitor for patients with non-ST segment elevation myocardial infarction: A phase Ib/IIa study. 2021, 41, 828-836	1
139	Successful Intracoronary Thrombolysis in Acute ST Elevation Myocardial Infarction Patient With Uncorrected Tetralogy of Fallot. 2021, 12, 90-93	0
138	Prognostic value and clinical predictors of intramyocardial hemorrhage measured by CMR T2* sequences in STEMI. 2021, 37, 1735-1744	0

137	Outcome of manual thrombus aspiration for patients undergoing primary PCI for acute STEMI showing large thrombus burden. 2021 , 73, 8	0
136	Treatment of Acute ST-Elevation Myocardial Infarction. 2007 , 963-977	1
135	Quantitative Tool for the Assessment of Myocardial Perfusion during X-Ray Angiographic Procedures. 2009 , 124-133	3
134	Coronary Atherosclerosis: Acute Coronary Syndromes. 2002 , 746-779	1
133	Role of perilipin 2 in microvascular obstruction in patients with ST-elevation myocardial infarction. 2021 , 10, 633-642	1
132	Microvascular resistance of the culprit coronary artery in acute ST-elevation myocardial infarction. 2016 , 1, e85768	31
131	Protective effect of injection in patients with acute myocardial infarction at a high risk of no-reflow during primary percutaneous coronary intervention. 2019 , 16, 406-413	5
130	The Value of Combining Plasma D-Dimer and Endothelin-1 Levels to Predict No-Reflow After Percutaneous Coronary Intervention of ST-Segment Elevation in Acute Myocardial Infarction Patients with a Type 2 Diabetes Mellitus History. 2018 , 24, 3549-3556	10
129	Effect of Metformin Treatment on Lipoprotein Subfractions in Non-Diabetic Patients with Acute Myocardial Infarction: A Glycometabolic Intervention as Adjunct to Primary Coronary Intervention in ST Elevation Myocardial Infarction (GIPS-III) Trial. 2016 , 11, e0145719	9
128	Morphine Does Not Affect Myocardial Salvage in ST-Segment Elevation Myocardial Infarction. 2017 , 12, e0170115	14
127	Assessment of the relationship between reperfusion success and T-peak to T-end interval in patients with ST elevation myocardial infarction treated with percutaneous coronary intervention. 2018 , 19, 50-57	2
126	Pharmacological prevention of peri-, and post-procedural myocardial injury in percutaneous coronary intervention. 2008 , 4, 223-30	5
125	Coronary microvascular dysfunction. 2020 , 68, 153-163	3
124	The REFLO-STEMI (REperfusion Facilitated by LOcal adjunctive therapy in ST-Elevation Myocardial Infarction) trial: a randomised controlled trial comparing intracoronary administration of adenosine or sodium nitroprusside with control for attenuation of microvascular obstruction during primary percutaneous coronary intervention. 2016 , 3, 1-18	7
123	Terminal QRS complex distortion on the admission electrocardiogram in anterior acute myocardial infarction and association with residual flow and infarct size after primary angioplasty. 2005 , 20, 21-5	12
122	Assessment of No-Reflow Phenomenon by Myocardial Blush Grade and Pulsed Wave Tissue Doppler Imaging in Patients with Acute Coronary Syndrome. 2014 , 24, 52-56	1
121	Effect of ischemic postconditioning during primary percutaneous coronary intervention for patients with ST-segment elevation myocardial infarction: A single-center cross-sectional study. 2019 , 22, 347-352	2
120	Quantitative myocardial blush grade reserve during pharmacologic hyperaemia: a way to perform a real wireless fractional flow reserve measurement in patients with coronary artery disease and intermediate coronary lesions. 2017 , 12, e2219-e2227	2

119	Microvascular dysfunction following ST-elevation myocardial infarction and its recovery over time. 2017 , 13, e578-e584	5
118	ST peak during percutaneous coronary intervention serves as an early prognostic predictor in patients with ST-segment elevation myocardial. 2014 , 10, 466-74	3
117	A tool for predicting the outcome of reperfusion in ST-elevation myocardial infarction using age, thrombotic burden and index of microcirculatory resistance (ATI score). 2016 , 12, 1223-1230	20
116	Ongoing tirofiban in myocardial infarction evaluation (On-TIME) 2 trial: rationale and study design. 2007 , 3, 371-80	21
115	Systematic primary aspiration in acute myocardial percutaneous intervention: a multicentre randomised controlled trial of the export aspiration catheter. 2008 , 4, 222-8	55
114	Impact of dye injection on intracoronary pressure. 2009 , 5, 272-6	3
113	Incidence, predictors and prognostic importance of bleeding after primary PCI for ST-elevation myocardial infarction. 2010 , 6, 106-111	21
112	Coronary thrombectomy, technical comparison of two systems on a laboratory bench: the impact of bends, angles and thrombus age. 2011 , 6, 729-34	4
111	Operator dependence of outcome after primary percutaneous coronary intervention. 2011 , 6, 760-7	10
110	Abortion of myocardial infarction by primary angioplasty mainly depends on preprocedural TIMI flow. 2011 , 6, 854-9	3
109	A new electrocardiographic marker of myocardial reperfusion in patients with acute ST-segment elevation myocardial infarction treated with primary percutaneous intervention: the value of QRS duration. 2012 , 7, 1406-12	4
108	Impact of perfusion restoration at epicardial and tissue levels on markers of myocardial necrosis and clinical outcome of patients with acute myocardial infarction. 2011 , 7, 128-35	6
107	A simple nomogram for early prediction of myocardial reperfusion after pre-hospital thrombolysis. 2011 , 7, 248-55	3
106	Impact of infarct-related artery patency before primary PCI on outcome in patients with ST-segment elevation myocardial infarction: the HORIZONS-AMI trial. 2013 , 8, 1307-14	26
105	Time-related impact of distal embolisation on myocardial perfusion and survival among patients undergoing primary angioplasty with glycoprotein IIb/IIIa inhibitors: insights from the EGYPT cooperation. 2012 , 8, 470-6	3
104	Benefits of optimising coronary flow before stenting in primary percutaneous coronary intervention for ST-elevation myocardial infarction: insights from INFUSE-AMI. 2014 , 9, 1195-201	14
103	Ruptured plaque and large plaque burden are risks of distal embolisation during percutaneous coronary intervention: evaluation by angioscopy and virtual histology intravascular ultrasound imaging. 2013 , 9, 235-42	11
102	Impact of bifurcation target lesion on angiographic, electrocardiographic, and clinical outcomes of patients undergoing primary percutaneous coronary intervention (from the Harmonizing Outcomes With Revascularization and Stents in Acute Myocardial Infarction [HORIZONS-AMI] trial). 2013 , 9, 817-23	13

101	Quantitative myocardial blush score (QuBE) allows the prediction of heart failure development in long-term follow-up in patients with ST-segment elevation myocardial infarction: Proof of concept study. 2019 , 26, 322-332	4
100	Diabetes mellitus and glucose as predictors of mortality in primary coronary percutaneous intervention. 2014 , 103, 323-30	7
99	Regional QT Interval Dispersion as an Early Predictor of Reperfusion in Patients with Acute Myocardial Infarction after Fibrinolytic Therapy. 2019 , 112, 20-29	4
98	Predictors of in-hospital mortality in patients with ST-segment elevation myocardial infarction undergoing pharmacoinvasive treatment. 2013 , 68, 1516-20	16
97	Increased neutrophil-to-lymphocyte ratio predicts persistent coronary no-flow after wire insertion in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2015 , 70, 34-40	14
96	The Association between Monocyte HDL-C Ratio and SYNTAX Score and SYNTAX Score II in STEMI Patients Treated with Primary PCI. 2018 , 34, 23-30	13
95	Effect of Ranolazine in Patients with Chest Pain and Normal Coronaries- A Hospital Based Study. 2017 , 11, OC14-OC16	4
94	Primer per k�an koroner giri�m uygulanan ST-segment y�seklikli miyokard infarkt� hastalarda platelet/lenfosit oranı ile no-reflow olu�mu arasındaki ili�i. 2021 , 46, 1441-1448	
93	Cardiac Rupture: Pathobiology, Diagnosis, Medical Management, and Surgical Intervention. 2004 , 597-628	
92	The No-Reflow Phenomenon. 2005 , 78-91	
91	Acute Myocardial Infarction. 2006 , 611-646	
90	Abrupt vessel closure and no-reflow. 2006 , 85-101	
89	Angiographic Assessment of Myocardial Perfusion Using Correlation Analysis. 2007 , 409-413	
88	Impact of PercuSurge GuardWire device on prevention and reduction of recovery time from complete heart block in patients with acute inferior myocardial infarction undergoing primary percutaneous coronary intervention. 2007 , 48, 35-44	3
87	Use of a thrombus extraction catheter (Thrombuster II(R)) in an acute myocardial infarction. 2008 , 3, 529-31	
86	Successful Myocardial Reperfusion. 2008 , 29-41	
85	Therapies Targeted at Preserving Microvascular Integrity and Preventing Reperfusion Injury. 2009 , 135-167	
84	Primary Coronary Intervention: The Technical Approach. 2009 , 47-63	

- 83 Devices for thrombectomy and distal emboli protection during percutaneous coronary interventions. **2009**, 15, 153-161
- 82 The Assessment of Myocardial Reperfusion and Its Clinical Significance in Acute Myocardial Infarction. **2009**, 223-240
- 81 Evaluation of myocardial perfusion. **2009**, 24-32
- 80 Redefining the Success of Mechanical Reperfusion. **2010**, 203-214
- 79 Use of Thrombectomy Devices in Primary Percutaneous Interventions for ST-elevation Myocardial Infarction - An Update. **2014**, 9, 102-107 1
- 78 Research Techniques. **2011**, 336-362
- 77 Interventional X-Ray Coronary Angiography. **2013**, 279-301
- 76 Coronary No-Reflow Following Percutaneous Coronary Intervention. **2013**, 1-26
- 75 Prozedurale Komplikationen. **2013**, 21-82
- 74 Optimal Angiographic Technique and Quantitative Analysis. **2014**, 9-22
- 73 Study of the possible medical and medication explanatory factors of angiographic outcomes in patients with acute ST elevation myocardial infarction undergoing primary percutaneous intervention. **2014**, 3, 186 1
- 72 Evaluation of Medical Therapies and Intravascular Devices with Quantitative Coronary Angiography. **2014**, 41-49
- 71 Microcirculation-Coronary: A Clinical Perspective. **2014**, 1-17
- 70 [Relationship between early myocardial reperfusion assessed by videodensitometry and late left ventricular function. Results following invasive treatment of acute myocardial infarction]. **2014**, 155, 187-93
- 69 Coronary No-Reflow Following Percutaneous Coronary Intervention. **2015**, 1865-1884
- 68 Interventional Strategies in Thrombus Management for ST Elevation Myocardial Infarction. **2015**, 10, 35-38 1
- 67 Qualitative and Quantitative Coronary Angiography. **2015**, 47-74
- 66 In-Lab Upfront Use of Tirofiban May Reduce the Occurrence of No-Reflow During Primary Percutaneous Coronary Intervention. A Pilot Randomized Study. **2016**, 107, 403-410 4

65	New Catheter-based Calibration Method for 2D X-ray Angiography Myocardial Perfusion. 2016 , 136, 1305-1311	
64	Acute Hyperglycemia an Independent Risk Factor for Contrast-Induced Nephropathy in Patients Undergoing Primary Percutaneous Coronary Intervention for STEMI. 2017 , 9,	0
63	Left Heart Catheterization. 2018 , 39-57	
62	The no-reflow phenomenon: mechanisms and therapeutic targets. 2018 , 17, 5-12	2
61	[Deferred versus immediate stenting in patients with ST - segment elevation myocardial infarction and residual large thrombus burden reclassified in the culprit lesion]. 2018 , 88, 432-440	0
60	Clinical outcome of intracoronary versus intravenous high-dose bolus administration of tirofiban in diabetic patients undergoing primary percutaneous coronary intervention. 2019 , 30, 285-289	1
59	Role of Coronary Microcirculation in No-Reflow Phenomenon in Myocardial Infarction with ST Segment Elevation. 2020 , 121-139	
58	Beneficial effects of early administration of recombinant human B-type natriuretic peptide in ST-elevation myocardial infarction patients receiving percutaneous coronary intervention treatment. 2019 , 60, 621-625	0
57	[The Clinical Value of no-Reflow Phenomenon in the Routing Operation of The Regional Vascular Center]. 2020 , 60, 146-150	
56	Effectiveness of Different P2Y12 Inhibitors on Coronary Flow in Patients with ST-Elevation Myocardial Infarction. 2020 , 6, 91-97	
55	LONG-TERM CONSEQUENCES OF PRIMARY PERCUTANEOUS CORONARY INTERVENTION IN PATIENTS WITH ST-ELEVATION MYOCARDIAL INFARCTION. 2020 , 73, 662-667	
54	Effect of Intravenous Abciximab on Coronary Flow Improvement After Re-vascularization in Primary Coronary Intervention and Short Term Impact. 2020 , 74, 265-269	1
53	Statin therapy and mortality in patients with ST-segment elevation myocardial infarction treated with primary angioplasty. 2004 , 12, 271-278	2
52	The NVVC guidelines for the management of patients with ST-elevation acute coronary syndromes (STE-ACS). 2002 , 10, 125-135	
51	Angiographic determinants of infarct size after successful percutaneous intervention for acute ST-elevation myocardial infarction: the impact of distal embolisation. 2002 , 10, 353-359	2
50	Rationale of the REPARATOR study: A randomised trial with serial cardiac MRI follow-up testing the ability of atorvastatin to reduce reperfusion damage after primary PCI for acute MI. 2006 , 14, 95-99	5
49	The relation between myocardial blush grade and myocardial contrast echocardiography: which one is a better predictor of myocardial damage?. 2010 , 18, 25-30	6
48	Modern risk stratification in coronary heart disease. 2011 , 4, 377-86	13

47	Restoration of microcirculatory patency after myocardial infarction: results of current coronary interventional strategies and techniques. 2012 , 39, 342-50	5
46	Manual vs mechanical thrombectomy during PCI for STEMI: a comprehensive direct and adjusted indirect meta-analysis of randomized trials. 2013 , 3, 146-57	8
45	High tenascin-C levels cause inadequate myocardial blush grade in patients with acute myocardial infarction. 2015 , 8, 2554-61	3
44	Evaluation of Short- and Long-Term Efficacy of Combined Intracoronary Administration of High-Dose Adenosine and Tirofiban during Primary Percutaneous Coronary Intervention. 2016 , 32, 640-648	3
43	Cutoff Value of Admission N-Terminal Pro-Brain Natriuretic Peptide Which Predicts Poor Myocardial Perfusion after Primary Percutaneous Coronary Intervention for ST-Segment-Elevation Myocardial Infarction. 2016 , 32, 649-655	2
42	Can thrombectomy and catheters used increase angiographically visible distal embolization in ST elevation myocardial infarction?. 2016 , 1, e139-e144	1
41	Effect of Selective Thrombus Aspiration on Serum Lipoprotein-Associated Phospholipase A2 in Patients with ST-Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention with High Thrombus Burden. 2018 , 34, 233-241	4
40	Limitations of Quantitative Blush Evaluator (QuBE) as myocardial perfusion assessment method on digital coronary angiograms. 2018 , 3, 394-400	
39	Intra-Coronary Administration of Tacrolimus Improves Myocardial Perfusion and Left Ventricular Function in Patients with ST-Segment Elevation Myocardial Infarction (COAT-STEMI) Undergoing Primary Percutaneous Coronary Intervention. 2021 , 37, 239-253	
38	Clinical Significance of Serum Lactate in Acute Myocardial Infarction: A Cardiac Magnetic Resonance Imaging Study. 2021 , 10,	2
37	Effects of Remote Ischemic Conditioning on Late Infarct Size and Left Ventricular Function in Patients with ST-segment Elevation Myocardial Infarction. 1,	
36	Adjunctive Catheter-Directed Thrombolysis during Primary PCI for ST-Segment Elevation Myocardial Infarction with High Thrombus Burden.. 2022 , 11,	1
35	Pathophysiology, Diagnosis, and Management of Coronary No-Reflow Phenomenon.	
34	A randomized controlled clinical trial of prolonged balloon inflation during stent deployment strategy in primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: a pilot study.. 2022 , 22, 30	0
33	Effect of Coronary Thrombus Aspiration in Non ST Elevation Acute Coronary Syndrome Patients on Three-Year Survival- Does it add any Benefit?. 2022 , 33197211053404	1
32	Administration of intracoronary adenosine before stenting for the prevention of no-reflow in patients with ST-elevation myocardial infarction.. 2022 , 56, 23-27	0
31	Risk Factors for No-Reflow in Patients with ST-Elevation Myocardial Infarction Who Underwent Percutaneous Coronary Intervention: A Case-Control Study.. 2022 , 2022, 3482518	
30	A Narrative Review of the Classical and Modern Diagnostic Methods of the No-Reflow Phenomenon.. 2022 , 12,	0

29	Modern atlas of invasive coronary angiography views: a practical approach for fellows and young interventionalists.. 2021 , 1	1
28	"No-Reflow" Phenomenon: A Contemporary Review.. 2022 , 11,	1
27	Systemic Immune-Inflammation Index as a Predictor of Microvascular Dysfunction in Patients With Cardiac Syndrome X.. 2022 , 33197221087777	3
26	Can thrombectomy and catheters used increase angiographically visible distal embolization in ST elevation myocardial infarction?. 2016 , 1, 139-144	1
25	Intracoronary pharmacological therapy versus aspiration thrombectomy in STEMI (IPAT-STEMI): A systematic review and meta-analysis of randomized trials.. 2022 , 17, e0263270	2
24	The Systemic Immune-Inflammation Index Predicts Impaired Myocardial Perfusion and Short-Term Mortality in ST-Segment Elevation Myocardial Infarction Patients. 000331972211068	0
23	Risk assessment score of no-reflow phenomenon in predicting myocardial perfusion disorders by contrast echocardiography in ST-segment elevation myocardial infarction patients after endovascular revascularization. 2022 , 37, 57-64	
22	Use of a Cutting Balloon Reduces the Incidence of Distal Embolism in Acute Coronary Syndrome Requiring Predilatation Before Stenting. 2022 ,	
21	Prognostic Value of Myocardial Blush Grade in ST-elevation MI: A Systematic Review and Meta-analysis. 17,	0
20	Rationale and design of the TACTICS registry: Optical coherence tomography guided primary percutaneous coronary intervention for patients with acute coronary syndrome. 2022 ,	0
19	IntraCoronary Artery Retrograde Thrombolysis vs. Thrombus Aspiration in ST-Segment Elevation Myocardial Infarction: Study Protocol for a Randomized Controlled Trial. 9,	0
18	Effects of Loading-Dose Statins Combined with PCSK9 Inhibitor Pre-Treatment before Primary Percutaneous Coronary Intervention on the Short-Term Prognosis in Patients with ST-Segment Elevation Myocardial Infarction. 2022 ,	0
17	Machine learning assisted reflectance spectral characterisation of coronary thrombi correlates with microvascular injury in patients with ST-segment elevation acute coronary syndrome. 9,	0
16	Correlation between endothelial dysfunction and occurrence of no-reflow in patients undergoing post-thrombolysis early invasive percutaneous intervention for ST-elevation myocardial infarction. 2022 , 74,	0
15	Effect of direct stenting on microvascular dysfunction during percutaneous coronary intervention in acute myocardial infarction: a randomized pilot study. 2022 , 50, 030006052211278	0
14	Association Between History of Polymerase Chain Reaction-verified COVID-19 Infection and Outcomes of Subsequent ST-Elevation Myocardial Infarction. 000331972211399	0
13	Effect Produced by a Cyclooctyne Derivative on Both Infarct Area and Left Ventricular Pressure via Calcium Channel Activation.	0
12	One-year results from the Assessing MICRO-vascular resistances via IMR to predict outcome in ST-elevation myocardial infarction patients with multivessel disease undergoing primary PCI (AMICRO) trial. 9,	0

- 11 Benefits of excimer laser coronary angioplasty over thrombus aspiration therapy for patients with acute coronary syndrome and thrombolysis in myocardial infarction flow grade 0. **2023**, 38, ○
- 10 Adjunctive Pharmacologic Agents and Mechanical Devices in Primary Percutaneous Coronary Intervention. **2010**, 39, 230-236 ○
- 9 Low Quantitative Blush Evaluator score predicts larger infarct size and reduced left ventricular systolic function in patients with STEMI regardless of diabetes status. **2023**, 13, ○
- 8 Growth Differentiation Factor 15 as a Predictor of the No-Reflow Phenomenon in Patients with ST-Segment Elevation Myocardial Infarction. **2023**, 12, 245 ○
- 7 RDW as A Predictor for No-Reflow Phenomenon in DM Patients with ST-Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. **2023**, 12, 807 ○
- 6 Predictors of adverse diastolic remodeling in non-diabetic patients presenting with ST-elevation myocardial infarction. **2023**, 23, ○
- 5 Coronary Microcirculation: The Next Frontier in the Management of STEMI. **2023**, 12, 1602 ○
- 4 Sustained Mechanical Aspiration Thrombectomy for High Thrombus Burden Coronary Vessel Occlusion: The Multicenter CHEETAH Study. **2023**, 16, ○
- 3 Evaluation of Biological Activity Exerted by Dibenzo[b,e]Thiophene-11(6H)-One on Left Ventricular Pressure Using an Isolated Rat Heart Model. ○
- 2 Angiography-Derived and Sensor-Wire Methods to Assess Coronary Microvascular Dysfunction in Patients With Acute Myocardial Infarction. **2023**, ○
- 1 Impact of maintenance dose of eptifibatide in patients with ST-segment elevation myocardial infarction who underwent primary percutaneous coronary intervention. **2023**, 75, ○