

# CITATION REPORT

List of articles citing

## RanGTP-regulated interactions of CRM1 with nucleoporins and a shuttling DEAD-box helicase

DOI: 10.1128/mcb.19.9.6276

Molecular and Cellular Biology, 1999, 19, 6276-85.

**Source:** <https://exaly.com/paper-pdf/30111569/citation-report.pdf>

**Version:** 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| #   | Paper  | IF  | Citations |
|-----|--|-----|-----------|
| 187 | Transport of proteins and RNAs in and out of the nucleus. <b>1999</b> , 99, 677-90   |     | 645       |
| 186 | Transport between the cell nucleus and the cytoplasm. <b>1999</b> , 15, 607-60   |     | 1685      |
| 185 | The nuclear pore complex: a protein machine bridging the nucleus and cytoplasm. <b>2000</b> , 12, 361-71   |     | 214       |
| 184 | Influenza A virus NS2 protein mediates vRNP nuclear export through NES-independent interaction with hCRM1. <i>EMBO Journal</i> , <b>2000</b> , 19, 6751-8  | 13  | 185       |
| 183 | The C-terminal domain of TAP interacts with the nuclear pore complex and promotes export of specific CTE-bearing RNA substrates. <b>2000</b> , 6, 136-58   |     | 268       |
| 182 | Identification of two novel RanGTP-binding proteins belonging to the importin beta superfamily. <i>Journal of Biological Chemistry</i> , <b>2000</b> , 275, 40163-8  | 5.4 | 42        |
| 181 | Nuclear export of the DEAD box An3 protein by CRM1 is coupled to An3 helicase activity. <i>Journal of Biological Chemistry</i> , <b>2000</b> , 275, 11561-8  | 5.4 | 32        |
| 180 | Nup50, a nucleoplasmically oriented nucleoporin with a role in nuclear protein export. <i>Molecular and Cellular Biology</i> , <b>2000</b> , 20, 5619-30   | 4.8 | 102       |
| 179 | Yeast Ran-binding protein 1 (Yrb1) shuttles between the nucleus and cytoplasm and is exported from the nucleus via a CRM1 (XPO1)-dependent pathway. <i>Molecular and Cellular Biology</i> , <b>2000</b> , 20, 4295-308 | 4.8 | 50        |
| 178 | A comparison of the activity, sequence specificity, and CRM1-dependence of different nuclear export signals. <b>2000</b> , 256, 213-24   |     | 362       |
| 177 | PHAX, a mediator of U snRNA nuclear export whose activity is regulated by phosphorylation. <b>2000</b> , 101, 187-98   |     | 266       |
| 176 | Rev protein and its cellular partners. <b>2000</b> , 48, 251-98  |     | 57        |
| 175 | A role for the basic patch and the C terminus of RanGTP in regulating the dynamic interactions with importin beta, CRM1 and RanBP1. <b>2001</b> , 305, 231-43  |     | 16        |
| 174 | Qualitative highly divergent nuclear export signals can regulate export by the competition for transport cofactors in vivo. <b>2001</b> , 2, 544-55  |     | 23        |
| 173 | Protein export from the nucleus. <b>2001</b> , 2, 684-9  |     | 65        |
| 172 | RanBP3 influences interactions between CRM1 and its nuclear protein export substrates. <b>2001</b> , 2, 926-32   |     | 99        |
| 171 | Caspases mediate nucleoporin cleavage, but not early redistribution of nuclear transport factors and modulation of nuclear permeability in apoptosis. <b>2001</b> , 8, 495-505   |     | 89        |

|     |  |     |     |
|-----|--|-----|-----|
| 170 | Nucleocytoplasmic transport: Ran, beta and beyond. <b>2001</b> , 11, 497-503   |     | 190 |
| 169 | Nuclear pores and nuclear assembly. <b>2001</b> , 13, 363-75   |     | 212 |
| 168 | NXT1 is necessary for the terminal step of Crm1-mediated nuclear export. <i>Journal of Cell Biology</i> , <b>2001</b> , 152, 141-55  | 7.3 | 56  |
| 167 | Influenza B and C virus NEP (NS2) proteins possess nuclear export activities. <i>Journal of Virology</i> , <b>2001</b> , 75, 7375-83   | 6.6 | 51  |
| 166 | CRM1 mediates the export of ADAR1 through a nuclear export signal within the Z-DNA binding domain. <i>Molecular and Cellular Biology</i> , <b>2001</b> , 21, 7862-71   | 4.8 | 116 |
| 165 | Stimulation of nuclear export and inhibition of nuclear import by a Ran mutant deficient in binding to Ran-binding protein 1. <i>Journal of Biological Chemistry</i> , <b>2001</b> , 276, 14524-31                                     | 5.4 | 29  |
| 164 | Calreticulin Is a receptor for nuclear export. <i>Journal of Cell Biology</i> , <b>2001</b> , 152, 127-40  | 7.3 | 227 |
| 163 | The N-terminal nuclear export sequence of IkappaBalpha is required for RanGTP-dependent binding to CRM1. <i>Journal of Biological Chemistry</i> , <b>2001</b> , 276, 23599-606   | 5.4 | 26  |
| 162 | Cofactor requirements for nuclear export of Rev response element (RRE)- and constitutive transport element (CTE)-containing retroviral RNAs. An unexpected role for actin. <i>Journal of Cell Biology</i> , <b>2001</b> , 152, 895-910 | 7.3 | 183 |
| 161 | Nuclear pore complexes: dynamics in unexpected places. <i>Journal of Cell Biology</i> , <b>2001</b> , 154, 17-20   | 7.3 | 25  |
| 160 | The nuclear export receptor Xpo1p forms distinct complexes with NES transport substrates and the yeast Ran binding protein 1 (Yrb1p). <i>Molecular Biology of the Cell</i> , <b>2001</b> , 12, 539-49                                  | 3.5 | 53  |
| 159 | The balance of nuclear import and export determines the intracellular distribution and function of tomato heat stress transcription factor HsfA2. <i>Molecular and Cellular Biology</i> , <b>2001</b> , 21, 1759-68                    | 4.8 | 108 |
| 158 | Ran-binding protein 3 is a cofactor for Crm1-mediated nuclear protein export. <i>Journal of Cell Biology</i> , <b>2001</b> , 153, 1391-402   | 7.3 | 119 |
| 157 | NXT1 (p15) is a crucial cellular cofactor in TAP-dependent export of intron-containing RNA in mammalian cells. <i>Molecular and Cellular Biology</i> , <b>2001</b> , 21, 2545-54   | 4.8 | 91  |
| 156 | In vitro analysis of nuclear transport mediated by the C-terminal shuttle domain of Tap. <i>Journal of Biological Chemistry</i> , <b>2001</b> , 276, 42355-63  | 5.4 | 32  |
| 155 | Reconstitution of nuclear protein export in isolated nuclear envelopes. <i>Journal of Cell Biology</i> , <b>2002</b> , 158, 849-54   | 7.3 | 14  |
| 154 | Members of the tristetraprolin family of tandem CCCH zinc finger proteins exhibit CRM1-dependent nucleocytoplasmic shuttling. <i>Journal of Biological Chemistry</i> , <b>2002</b> , 277, 11606-13                                     | 5.4 | 80  |
| 153 | A synthetic HIV-1 Rev inhibitor interfering with the CRM1-mediated nuclear export. <b>2002</b> , 99, 14440-5   |     | 103 |

|     |  |     |     |
|-----|--|-----|-----|
| 152 | The cytoplasmic filaments of the nuclear pore complex are dispensable for selective nuclear protein import. <i>Journal of Cell Biology</i> , <b>2002</b> , 158, 63-77  | 7.3 | 161 |
| 151 | Interaction between parvovirus NS2 protein and nuclear export factor Crm1 is important for viral egress from the nucleus of murine cells. <i>Journal of Virology</i> , <b>2002</b> , 76, 3257-66   | 6.6 | 53  |
| 150 | Analysis of nuclear protein import and export in vitro using fluorescent cargoes. <i>Methods in Molecular Biology</i> , <b>2002</b> , 189, 231-45  | 1.4 | 5   |
| 149 | Deciphering networks of protein interactions at the nuclear pore complex. <b>2002</b> , 1, 930-46  |     | 61  |
| 148 | Ran GTPase cycle and importins alpha and beta are essential for spindle formation and nuclear envelope assembly in living <i>Caenorhabditis elegans</i> embryos. <i>Molecular Biology of the Cell</i> , <b>2002</b> , 13, 4355-70        | 3.5 | 157 |
| 147 | The NS2 proteins of parvovirus minute virus of mice are required for efficient nuclear egress of progeny virions in mouse cells. <i>Journal of Virology</i> , <b>2002</b> , 76, 10307-19   | 6.6 | 60  |
| 146 | Karyopherin beta 2B participates in mRNA export from the nucleus. <b>2002</b> , 99, 14195-9  |     | 30  |
| 145 | Steady-state nuclear localization of exportin-t involves RanGTP binding and two distinct nuclear pore complex interaction domains. <i>Molecular and Cellular Biology</i> , <b>2002</b> , 22, 5708-20                                     | 4.8 | 35  |
| 144 | Fluorescence resonance energy transfer biosensors that detect Ran conformational changes and a Ran x GDP-importin-beta -RanBP1 complex in vitro and in intact cells. <i>Journal of Biological Chemistry</i> , <b>2002</b> , 277, 30121-7 | 5.4 | 31  |
| 143 | Nuclear export of 5S rRNA-containing ribonucleoprotein complexes requires CRM1 and the RanGTPase cycle. <i>European Journal of Cell Biology</i> , <b>2002</b> , 81, 549-56   | 6.1 | 13  |
| 142 | GTPase Protocols. <b>2002</b> ,  |     | 0   |
| 141 | The C-terminal extension of the small GTPase Ran is essential for defining the GDP-bound form. <b>2002</b> , 318, 583-93   |     | 18  |
| 140 | PBX1 nuclear export is regulated independently of PBX-MEINOX interaction by PKA phosphorylation of the PBC-B domain. <i>EMBO Journal</i> , <b>2003</b> , 22, 89-99   | 13  | 48  |
| 139 | Coordinated nuclear export of 60S ribosomal subunits and NMD3 in vertebrates. <i>EMBO Journal</i> , <b>2003</b> , 22, 2841-51  | 13  | 95  |
| 138 | Monocytic cells hyperacetylate chromatin protein HMGB1 to redirect it towards secretion. <i>EMBO Journal</i> , <b>2003</b> , 22, 5551-60   | 13  | 929 |
| 137 | Functional analysis of the interaction of the human immunodeficiency virus type 1 Rev nuclear export signal with its cofactors. <b>2003</b> , 314, 591-600   |     | 13  |
| 136 | Nucleocytoplasmic transport: navigating the channel. <b>2003</b> , 4, 127-35   |     | 118 |
| 135 | Ratjadones inhibit nuclear export by blocking CRM1/exportin 1. <b>2003</b> , 286, 321-31   |     | 74  |

|     |   |     |     |
|-----|---|-----|-----|
| 134 | A multifunctional domain in human CRM1 (exportin 1) mediates RanBP3 binding and multimerization of human T-cell leukemia virus type 1 Rex protein. <i>Molecular and Cellular Biology</i> , <b>2003</b> , 23, 8751-61                | 4.8 | 26  |
| 133 | CRM1/Ran-mediated nuclear export of p27(Kip1) involves a nuclear export signal and links p27 export and proteolysis. <i>Molecular Biology of the Cell</i> , <b>2003</b> , 14, 201-13  | 3.5 | 158 |
| 132 | A temperature-sensitive mutant of the mammalian RNA helicase, DEAD-BOX X isoform, DBX, defective in the transition from G1 to S phase. <b>2003</b> , 134, 71-82   |     | 30  |
| 131 | A nuclear kinesin-like protein interacts with and stimulates the activity of the leucine-rich nuclear export signal of the human immunodeficiency virus type 1 rev protein. <i>Journal of Virology</i> , <b>2003</b> , 77, 7236-43  | 6.6 | 10  |
| 130 | The strategy for coupling the RanGTP gradient to nuclear protein export. <b>2003</b> , 100, 1717-22   |     | 35  |
| 129 | Nuclear export inhibitors and kinase inhibitors identified using a MAPK-activated protein kinase 2 redistribution screen. <b>2004</b> , 2, 7-20   |     | 53  |
| 128 | The dynamic association of RCC1 with chromatin is modulated by Ran-dependent nuclear transport. <i>Molecular Biology of the Cell</i> , <b>2004</b> , 15, 245-55   | 3.5 | 22  |
| 127 | CRM1 and Ran are present but a NES-CRM1-RanGTP complex is not required in Balbiani ring mRNP particles from the gene to the cytoplasm. <i>Journal of Cell Science</i> , <b>2004</b> , 117, 1553-66                                  | 5.3 | 9   |
| 126 | The death effector domain protein PEA-15 prevents nuclear entry of ERK2 by inhibiting required interactions. <i>Journal of Biological Chemistry</i> , <b>2004</b> , 279, 12840-7  | 5.4 | 67  |
| 125 | Nup358/RanBP2 attaches to the nuclear pore complex via association with Nup88 and Nup214/CAN and plays a supporting role in CRM1-mediated nuclear protein export. <i>Molecular and Cellular Biology</i> , <b>2004</b> , 24, 2373-84 | 4.8 | 133 |
| 124 | Nucleo-cytoplasmic shuttling of Axin, a negative regulator of the Wnt-beta-catenin Pathway. <i>Journal of Biological Chemistry</i> , <b>2004</b> , 279, 5263-7  | 5.4 | 71  |
| 123 | Supraphysiological nuclear export signals bind CRM1 independently of RanGTP and arrest at Nup358. <i>EMBO Journal</i> , <b>2004</b> , 23, 3643-52   | 13  | 77  |
| 122 | Structural basis for the assembly of a nuclear export complex. <b>2004</b> , 432, 872-7   |     | 158 |
| 121 | Aberrant intracellular localization of SET-CAN fusion protein, associated with a leukemia, disorganizes nuclear export. <b>2004</b> , 111, 501-7  |     | 45  |
| 120 | Requirement of DDX3 DEAD box RNA helicase for HIV-1 Rev-RRE export function. <b>2004</b> , 119, 381-92  |     | 410 |
| 119 | Nuclear export of signal recognition particle RNA in mammalian cells. <b>2004</b> , 313, 351-5  |     | 25  |
| 118 | Human DDX3Y, the Y-encoded isoform of RNA helicase DDX3, rescues a hamster temperature-sensitive ET24 mutant cell line with a DDX3X mutation. <b>2004</b> , 300, 213-22   |     | 60  |
| 117 | Architecture of CRM1/Exportin1 suggests how cooperativity is achieved during formation of a nuclear export complex. <b>2004</b> , 16, 761-75  |     | 107 |

|     |  |     |     |
|-----|--|-----|-----|
| 116 | NUP98 fusion in human leukemia: dysregulation of the nuclear pore and homeodomain proteins. <b>2005</b> , 82, 21-7   |     | 44  |
| 115 | Leucine-rich nuclear-export signals: born to be weak. <b>2005</b> , 15, 121-4  |     | 275 |
| 114 | The human T-cell leukemia virus Rex protein. <b>2005</b> , 10, 431-45  |     | 59  |
| 113 | Yeast poly(A)-binding protein Pab1 shuttles between the nucleus and the cytoplasm and functions in mRNA export. <b>2005</b> , 11, 517-31   |     | 91  |
| 112 | Proteins of the S100 family regulate the oligomerization of p53 tumor suppressor. <b>2005</b> , 102, 4735-40   |     | 149 |
| 111 | Detailed mapping of the nuclear export signal in the Rous sarcoma virus Gag protein. <i>Journal of Virology</i> , <b>2005</b> , 79, 8732-41  | 6.6 | 35  |
| 110 | Kinetic and molecular analysis of nuclear export factor CRM1 association with its cargo in vivo. <i>Molecular and Cellular Biology</i> , <b>2005</b> , 25, 728-39                              | 4.8 | 55  |
| 109 | Belle is a Drosophila DEAD-box protein required for viability and in the germ line. <i>Developmental Biology</i> , <b>2005</b> , 277, 92-101   | 3.1 | 88  |
| 108 | Nuclear import in viral infections. <i>Current Topics in Microbiology and Immunology</i> , <b>2005</b> , 285, 109-38   | 3.3 | 28  |
| 107 | The nucleoporin Nup214 sequesters CRM1 at the nuclear rim and modulates NFkappaB activation in Drosophila. <i>Journal of Cell Science</i> , <b>2006</b> , 119, 4409-19                         | 5.3 | 55  |
| 106 | Divergent roles of the DEAD-box protein BS-PL10, the urochordate homologue of human DDX3 and DDX3Y proteins, in colony astogeny and ontogeny. <b>2006</b> , 235, 1508-21                       |     | 38  |
| 105 | Multiple importins function as nuclear transport receptors for the Rev protein of human immunodeficiency virus type 1. <i>Journal of Biological Chemistry</i> , <b>2006</b> , 281, 20883-20890 | 5.4 | 67  |
| 104 | Mouse Disabled1 (DAB1) is a nucleocytoplasmic shuttling protein. <i>Journal of Biological Chemistry</i> , <b>2006</b> , 281, 38951-65  | 5.4 | 32  |
| 103 | Nup214 is required for CRM1-dependent nuclear protein export in vivo. <i>Molecular and Cellular Biology</i> , <b>2006</b> , 26, 6772-85  | 4.8 | 121 |
| 102 | A novel Ded1-like RNA helicase interacts with the Y-box protein ctYB-1 in nuclear mRNP particles and in polysomes. <i>Journal of Biological Chemistry</i> , <b>2006</b> , 281, 14263-72        | 5.4 | 20  |
| 101 | Nup214-Nup88 nucleoporin subcomplex is required for CRM1-mediated 60 S preribosomal nuclear export. <i>Journal of Biological Chemistry</i> , <b>2006</b> , 281, 19378-86                       | 5.4 | 61  |
| 100 | Novel interaction of the 60S ribosomal subunit export adapter Nmd3 at the nuclear pore complex. <i>Journal of Biological Chemistry</i> , <b>2007</b> , 282, 14028-37                           | 5.4 | 8   |
| 99  | The adapter importin-alpha provides flexible control of nuclear import at the expense of efficiency. <b>2007</b> , 3, 118  |     | 23  |

|    |  |      |     |
|----|--|------|-----|
| 98 | Nuclear traffic of influenza virus proteins and ribonucleoprotein complexes. <i>Virus Research</i> , <b>2007</b> , 124, 12-21  | 6.4  | 160 |
| 97 | SET-CAN, the product of the t(9;9) in acute undifferentiated leukemia, causes expansion of early hematopoietic progenitors and hyperproliferation of stomach mucosa in transgenic mice. <b>2007</b> , 171, 654-66                |      | 24  |
| 96 | The DEAD box RNA helicase VBH-1 is required for germ cell function in <i>C. elegans</i> . <b>2007</b> , 45, 533-46   |      | 26  |
| 95 | Intracellular bacterial growth is controlled by a kinase network around PKB/AKT1. <b>2007</b> , 450, 725-30  |      | 243 |
| 94 | Homodimerization antagonizes nuclear export of survivin. <b>2007</b> , 8, 1495-502   |      | 35  |
| 93 | CRM1-mediated nuclear export: to the pore and beyond. <b>2007</b> , 17, 193-201  |      | 270 |
| 92 | Inhibition of the CRM1-mediated nucleocytoplasmic transport by N-azolyacrylates: structure-activity relationship and mechanism of action. <b>2008</b> , 16, 9487-97  |      | 51  |
| 91 | Beta-catenin can bind directly to CRM1 independently of adenomatous polyposis coli, which affects its nuclear localization and LEF-1/beta-catenin-dependent gene expression. <b>2008</b> , 32, 394-400                           |      | 4   |
| 90 | Deciphering the assembly pathway of Sm-class U snRNPs. <b>2008</b> , 582, 1997-2003  |      | 84  |
| 89 | A solution to limited genomic capacity: using adaptable binding surfaces to assemble the functional HIV Rev oligomer on RNA. <b>2008</b> , 31, 824-34  |      | 92  |
| 88 | Entamoeba histolytica EhDEAD1 is a conserved DEAD-box RNA helicase with ATPase and ATP-dependent RNA unwinding activities. <i>Gene</i> , <b>2008</b> , 414, 19-31  | 3.8  | 12  |
| 87 | The DEAD-box RNA helicase DDX3 associates with export messenger ribonucleoproteins as well as tip-associated protein and participates in translational control. <i>Molecular Biology of the Cell</i> , <b>2008</b> , 19, 3847-58 | 3.5  | 163 |
| 86 | Human DDX3 functions in translation and interacts with the translation initiation factor eIF3. <i>Nucleic Acids Research</i> , <b>2008</b> , 36, 4708-18   | 20.1 | 128 |
| 85 | STRADalpha regulates LKB1 localization by blocking access to importin-alpha, and by association with Crm1 and exportin-7. <i>Molecular Biology of the Cell</i> , <b>2008</b> , 19, 1614-26                                       | 3.5  | 81  |
| 84 | Hormone-dependent nuclear export of estradiol receptor and DNA synthesis in breast cancer cells. <i>Journal of Cell Biology</i> , <b>2008</b> , 182, 327-40  | 7.3  | 68  |
| 83 | A supraphysiological nuclear export signal is required for parvovirus nuclear export. <i>Molecular Biology of the Cell</i> , <b>2008</b> , 19, 2544-52   | 3.5  | 31  |
| 82 | The Nup358-RanGAP complex is required for efficient importin alpha/beta-dependent nuclear import. <i>Molecular Biology of the Cell</i> , <b>2008</b> , 19, 2300-10   | 3.5  | 107 |
| 81 | The current understanding of Ded1p/DDX3 homologs from yeast to human. <i>RNA Biology</i> , <b>2009</b> , 6, 17-20  | 4.8  | 70  |

|    |   |      |     |
|----|---|------|-----|
| 80 | Identification and functional dissection of localization signals within ataxin-3. <i>Neurobiology of Disease</i> , <b>2009</b> , 36, 280-92   | 7.5  | 35  |
| 79 | Karyopherin binding interactions and nuclear import mechanism of nuclear pore complex protein Tpr. <i>BMC Cell Biology</i> , <b>2009</b> , 10, 74   |      | 26  |
| 78 | P68 RNA helicase is a nucleocytoplasmic shuttling protein. <i>Cell Research</i> , <b>2009</b> , 19, 1388-400  | 24.7 | 35  |
| 77 | Comparative proteomic analyses of the nuclear envelope and pore complex suggests a wide range of heretofore unexpected functions. <i>Journal of Proteomics</i> , <b>2009</b> , 72, 56-70                              | 3.9  | 22  |
| 76 | The <i>C. elegans</i> sex determination gene <i>laf-1</i> encodes a putative DEAD-box RNA helicase. <i>Developmental Biology</i> , <b>2009</b> , 330, 358-67  | 3.1  | 15  |
| 75 | Nuclear export complexes in the frame. <i>Current Opinion in Structural Biology</i> , <b>2010</b> , 20, 247-52  | 8.1  | 61  |
| 74 | NES consensus redefined by structures of PKI-type and Rev-type nuclear export signals bound to CRM1. <i>Nature Structural and Molecular Biology</i> , <b>2010</b> , 17, 1367-76                                       | 17.6 | 180 |
| 73 | An allosteric mechanism to displace nuclear export cargo from CRM1 and RanGTP by RanBP1. <i>EMBO Journal</i> , <b>2010</b> , 29, 2002-13  | 13   | 77  |
| 72 | The anti-inflammatory prostaglandin 15-deoxy-delta(12,14)-PGJ2 inhibits CRM1-dependent nuclear protein export. <i>Journal of Biological Chemistry</i> , <b>2010</b> , 285, 22202-10                                   | 5.4  | 39  |
| 71 | Venezuelan equine Encephalitis virus capsid protein forms a tetrameric complex with CRM1 and importin alpha/beta that obstructs nuclear pore complex function. <i>Journal of Virology</i> , <b>2010</b> , 84, 4158-71 | 6.6  | 79  |
| 70 | Casein-kinase-II-dependent phosphorylation of PPARgamma provokes CRM1-mediated shuttling of PPARgamma from the nucleus to the cytosol. <i>Journal of Cell Science</i> , <b>2010</b> , 123, 192-201                    | 5.3  | 34  |
| 69 | Inhibition of CRM1-mediated nuclear export of transcription factors by leukemogenic NUP98 fusion proteins. <i>Journal of Biological Chemistry</i> , <b>2010</b> , 285, 16248-57                                       | 5.4  | 42  |
| 68 | Host factors mediating HIV-1 replication. <i>Virus Research</i> , <b>2011</b> , 161, 101-14   | 6.4  | 60  |
| 67 | Biogenesis of spliceosomal small nuclear ribonucleoproteins. <i>Wiley Interdisciplinary Reviews RNA</i> , <b>2011</b> , 2, 718-31   | 9.3  | 96  |
| 66 | Patterning of the <i>Drosophila</i> oocyte by a sequential translation repression program involving the d4EHP and Belle translational repressors. <i>RNA Biology</i> , <b>2011</b> , 8, 904-12                        | 4.8  | 20  |
| 65 | Ran-dependent nuclear export mediators: a structural perspective. <i>EMBO Journal</i> , <b>2011</b> , 30, 3457-74   | 13   | 149 |
| 64 | Calcium-calmodulin kinase I cooperatively regulates nucleocytoplasmic shuttling of CCT by accessing a nuclear export signal. <i>Molecular Biology of the Cell</i> , <b>2012</b> , 23, 2755-69                         | 3.5  | 8   |
| 63 | Virus-producing cells determine the host protein profiles of HIV-1 virion cores. <i>Retrovirology</i> , <b>2012</b> , 9, 65   | 3.6  | 52  |



|    |  |      |     |
|----|--|------|-----|
| 62 | The nucleoporin-like protein NLP1 (hCG1) promotes CRM1-dependent nuclear protein export. <i>Journal of Cell Science</i> , <b>2012</b> , 125, 144-54  | 5.3  | 20  |
| 61 | Role of Nucleocytoplasmic RNA Transport during the Life Cycle of Retroviruses. <i>Frontiers in Microbiology</i> , <b>2012</b> , 3, 179   | 5.7  | 16  |
| 60 | Structural basis for the nuclear export activity of Importin13. <i>EMBO Journal</i> , <b>2013</b> , 32, 899-913  | 13   | 28  |
| 59 | Structural determinants and mechanism of mammalian CRM1 allostery. <i>Structure</i> , <b>2013</b> , 21, 1350-60  | 5.2  | 16  |
| 58 | Phosphorylation of Crm1 by CDK1-cyclin-B promotes Ran-dependent mitotic spindle assembly. <i>Journal of Cell Science</i> , <b>2013</b> , 126, 3417-28  | 5.3  | 30  |
| 57 | Several phenylalanine-glycine motives in the nucleoporin Nup214 are essential for binding of the nuclear export receptor CRM1. <i>Journal of Biological Chemistry</i> , <b>2013</b> , 288, 3952-63                             | 5.4  | 25  |
| 56 | Phosphorylation of human chromosome maintenance 1 mediates association with 14-3-3 proteins. <i>Animal Cells and Systems</i> , <b>2013</b> , 17, 186-195   | 2.3  | 1   |
| 55 | The DEAD-box helicase Ded1 from yeast is an mRNP cap-associated protein that shuttles between the cytoplasm and nucleus. <i>Nucleic Acids Research</i> , <b>2014</b> , 42, 10005-22  | 20.1 | 32  |
| 54 | Atomic basis of CRM1-cargo recognition, release and inhibition. <i>Seminars in Cancer Biology</i> , <b>2014</b> , 27, 52-61  | 12.7 | 90  |
| 53 | The Ded1/DDX3 subfamily of DEAD-box RNA helicases. <i>Critical Reviews in Biochemistry and Molecular Biology</i> , <b>2014</b> , 49, 343-60  | 8.7  | 96  |
| 52 | RNA Export through the NPC in Eukaryotes. <i>Genes</i> , <b>2015</b> , 6, 124-49   | 4.2  | 76  |
| 51 | Formation of a Trimeric Xpo1-Ran[GTP]-Ded1 Exportin Complex Modulates ATPase and Helicase Activities of Ded1. <i>PLoS ONE</i> , <b>2015</b> , 10, e0131690   | 3.7  | 7   |
| 50 | Spatial regulation of cytoplasmic snRNP assembly at the cellular level. <i>Journal of Experimental Botany</i> , <b>2015</b> , 66, 7019-30  | 7    | 5   |
| 49 | Eukaryotic Ribosome Assembly and Nuclear Export. <i>International Review of Cell and Molecular Biology</i> , <b>2015</b> , 319, 107-40   | 6    | 70  |
| 48 | Structural and Functional Characterization of CRM1-Nup214 Interactions Reveals Multiple FG-Binding Sites Involved in Nuclear Export. <i>Cell Reports</i> , <b>2015</b> , 13, 690-702   | 10.6 | 57  |
| 47 | Components and regulation of nuclear transport processes. <i>FEBS Journal</i> , <b>2015</b> , 282, 445-62  | 5.7  | 146 |
| 46 | The RanBP2/RanGAP1*SUMO1/Ubc9 SUMO E3 ligase is a disassembly machine for Crm1-dependent nuclear export complexes. <i>Nature Communications</i> , <b>2016</b> , 7, 11482   | 17.4 | 47  |
| 45 | Leukemia-Associated Nup214 Fusion Proteins Disturb the XPO1-Mediated Nuclear-Cytoplasmic Transport Pathway and Thereby the NF- $\kappa$ B Signaling Pathway. <i>Molecular and Cellular Biology</i> , <b>2016</b> , 36, 1820-35 | 4.8  | 26  |

|    |   |      |    |
|----|---|------|----|
| 44 | Principles of 60S ribosomal subunit assembly emerging from recent studies in yeast. <i>Biochemical Journal</i> , <b>2017</b> , 474, 195-214   | 3.8  | 55 |
| 43 | PL10 DEAD-Box Protein is Expressed during Germ Cell Differentiation in the Reptile <i>Podarcis sicula</i> (Family Lacertidae). <i>Journal of Experimental Zoology Part B: Molecular and Developmental Evolution</i> , <b>2017</b> , 328, 433-448                        | 1.8  | 3  |
| 42 | Nuclear Export Signal Masking Regulates HIV-1 Rev Trafficking and Viral RNA Nuclear Export. <i>Journal of Virology</i> , <b>2017</b> , 91,  | 6.6  | 24 |
| 41 | Characterization of the Es-DDX52 involved in the spermatogonial mitosis and spermatid differentiation in Chinese mitten crab ( <i>Eriocheir sinensis</i> ). <i>Gene</i> , <b>2018</b> , 646, 106-119  | 3.8  | 5  |
| 40 | mRNP Transport in Eukaryotes. mRNP Export from the Nucleus. <i>Molecular Genetics, Microbiology and Virology</i> , <b>2018</b> , 33, 182-186  | 0.4  | 1  |
| 39 | Nuclear-Cytoplasmic Transport. <i>Nucleic Acids and Molecular Biology</i> , <b>2018</b> ,   |      |    |
| 38 | Plant snRNP Biogenesis: A Perspective from the Nucleolus and Cajal Bodies. <i>Frontiers in Plant Science</i> , <b>2017</b> , 8, 2184  | 6.2  | 15 |
| 37 | Exportin Crm1 is repurposed as a docking protein to generate microtubule organizing centers at the nuclear pore. <i>ELife</i> , <b>2018</b> , 7,  | 8.9  | 10 |
| 36 | Investigating nucleo-cytoplasmic shuttling of the human DEAD-box helicase DDX3. <i>European Journal of Cell Biology</i> , <b>2018</b> , 97, 501-511   | 6.1  | 13 |
| 35 | Distinct RanBP1 nuclear export and cargo dissociation mechanisms between fungi and animals. <i>ELife</i> , <b>2019</b> , 8,   | 8.9  | 6  |
| 34 | Gonadotropin Regulation Testicular RNA Helicase, Two Decades of Studies on Its Structure Function and Regulation From Its Discovery Opens a Window for Development of a Non-hormonal Oral Male Contraceptive. <i>Frontiers in Endocrinology</i> , <b>2019</b> , 10, 576 | 5.7  | 4  |
| 33 | Exportin-1-Dependent Nuclear Export of DEAD-box Helicase DDX3X is Central to its Role in Antiviral Immunity. <i>Cells</i> , <b>2019</b> , 8,  | 7.9  | 7  |
| 32 | NUP214 in Leukemia: It's More than Transport. <i>Cells</i> , <b>2019</b> , 8,   | 7.9  | 20 |
| 31 | The coming-of-age of nucleocytoplasmic transport in motor neuron disease and neurodegeneration. <i>Cellular and Molecular Life Sciences</i> , <b>2019</b> , 76, 2247-2273   | 10.3 | 16 |
| 30 | A nuclear lamina-chromatin-Ran GTPase axis modulates nuclear import and DNA damage signaling. <i>Aging Cell</i> , <b>2019</b> , 18, e12851  | 9.9  | 16 |
| 29 | NXF1 and CRM1 nuclear export pathways orchestrate nuclear export, translation and packaging of murine leukaemia retrovirus unspliced RNA. <i>RNA Biology</i> , <b>2020</b> , 17, 528-538  | 4.8  | 8  |
| 28 | Nuclear Import Receptors Directly Bind to Arginine-Rich Dipeptide Repeat Proteins and Suppress Their Pathological Interactions. <i>Cell Reports</i> , <b>2020</b> , 33, 108538  | 10.6 | 25 |
| 27 | SKIP-HOPS recruits TBC1D15 for a Rab7-to-Arl8b identity switch to control late endosome transport. <i>EMBO Journal</i> , <b>2020</b> , 39, e102301  | 13   | 23 |

|    |  |      |    |
|----|--|------|----|
| 26 | The Identification of Potential Members of the Ded1/DDX3 Subfamily of DEAD-Box RNA Helicases from the Protozoan Parasite and Their Analyses in Yeast. <i>Genes</i> , <b>2021</b> , 12, | 4.2  | 1  |
| 25 | DDX3X: structure, physiologic functions and cancer. <i>Molecular Cancer</i> , <b>2021</b> , 20, 38   | 42.1 | 22 |
| 24 | A nuclear export sequence promotes CRM1-dependent targeting of the nucleoporin Nup214 to the nuclear pore complex. <i>Journal of Cell Science</i> , <b>2021</b> , 134,                 | 5.3  | 1  |
| 23 | The nuclear lamina is a hub for the nuclear function of Jacob. <i>Molecular Brain</i> , <b>2021</b> , 14, 9  | 4.5  | 2  |
| 22 | The Molecular Mechanisms of mRNA Export. <b>2005</b> , 161-174   |      | 1  |
| 21 | Direct and Indirect Roles of Ran-GTP in Nuclear Export of RNAS in Higher Eukaryotes. <b>2001</b> , 59-83   |      | 2  |
| 20 | On the Effects of Leukemogenic Nucleoporin Fusion Proteins on Nucleocytoplasmic Transport and Gene Expression. <i>Nucleic Acids and Molecular Biology</i> , <b>2018</b> , 223-248      |      | 1  |
| 19 | Exportin-mediated nuclear export of proteins and ribonucleoproteins. <i>Results and Problems in Cell Differentiation</i> , <b>2002</b> , 35, 67-91                                     | 1.4  | 37 |
| 18 | Constitutive transport element-mediated nuclear export. <i>Current Topics in Microbiology and Immunology</i> , <b>2001</b> , 259, 77-93  | 3.3  | 25 |
| 17 | Targeted CRM1-inhibition perturbs leukemogenic NUP214 fusion proteins and exerts anti-cancer effects in leukemia cell lines with NUP214 rearrangements.                                |      | 1  |
| 16 | members only encodes a Drosophila nucleoporin required for Rel protein import and immune response activation. <i>Genes and Development</i> , <b>2000</b> , 14, 1945-1957               | 12.6 | 35 |
| 15 | Insights into the function of the CRM1 cofactor RanBP3 from the structure of its Ran-binding domain. <i>PLoS ONE</i> , <b>2011</b> , 6, e17011   | 3.7  | 15 |
| 14 | Molecular Characterization and Functional Analysis of Annulate Lamellae Pore Complexes in Nuclear Transport in Mammalian Cells. <i>PLoS ONE</i> , <b>2015</b> , 10, e0144508           | 3.7  | 10 |
| 13 | Importin- $\beta$ modulates the permeability of the nuclear pore complex in a Ran-dependent manner. <i>ELife</i> , <b>2015</b> , 4,  | 8.9  | 69 |
| 12 | The export receptor Crm1 forms a dimer to promote nuclear export of HIV RNA. <i>ELife</i> , <b>2014</b> , 3, e04121  | 8.9  | 46 |
| 11 | Chromatin-bound CRM1 recruits SET-Nup214 and NPM1c onto clusters causing aberrant expression in leukemia cells. <i>ELife</i> , <b>2019</b> , 8,  | 8.9  | 12 |
| 10 | A nuclear role for the DEAD-box protein Dbp5 in tRNA export. <i>ELife</i> , <b>2019</b> , 8,   | 8.9  | 6  |
| 9  | Nuclear export mediated by the Rev/Rex class of retroviral Trans-activator proteins. <i>Current Topics in Microbiology and Immunology</i> , <b>2001</b> , 259, 55-76                   | 3.3  | 7  |

- 8 Ran GTPASE Regulation of the CRM1-Dependent Export Pathway. **2001**, 15-31
- 7 Human T-Cell Leukemia Virus Type 2 (HTLV-2) Biology and Pathogenesis. **2012**, 647-675
- 6 Exportin Crm1 is repurposed as a docking protein to generate microtubule organizing centers at the nuclear pore.
- 5 mRNA transport in eukaryotes. mRNA export from the nucleus. *Molekuliarnaia Genetika, Mikrobiologiya I Virusologiya*, **2018**, 36, 25 0.3
- 4 Distinct RanBP1 nuclear export and cargo dissociation mechanisms between fungi and animals.
- 3 Targeted CRM1-inhibition perturbs leukemogenic NUP214 fusion proteins and exerts anti-cancer effects in leukemia cell lines with rearrangements. *Oncotarget*, **2020**, 11, 3371-3386 3.3 0
- 2 members only encodes a Drosophila nucleoporin required for rel protein import and immune response activation. *Genes and Development*, **2000**, 14, 1945-57 12.6 65
- 1 Binding Affinity Measurement of Nuclear Export Signal Peptides to Their Exporter CRM1.. *Methods in Molecular Biology*, **2022**, 2502, 245-256 1.4