Comparison of models for defining nearshore flatfish nursery areas in Alaskan waters

DOI: 10.1046/j.1365-2419.1999.00087.x
Fisheries Oceanography, 1999, 8, 50-67.

#	Paper	IF	Citations
53	Associations between flatfish abundance and surficial sediments in the eastern Bering Sea. *Canadian Journal of Fisheries and Aquatic Sciences*, **2000**, 57, 2410-2419	2.4	75
52	Sediment preferences and size-specific distribution of young-of-the-year Pacific halibut in an Alaska nursery. *Journal of Fish Biology*, **2002**, 61, 540-559	1.9	25
51	Patterns in use of estuarine habitat by juvenile English sole (Pleuronectes vetulus) in four Eastern North Pacific estuaries. *Estuaries and Coasts*, **2003**, 26, 1142-1154		21
50	Relationships between size-specific sediment preferences and burial capabilities in juveniles of two Alaska flatfishes. *Journal of Experimental Marine Biology and Ecology*, **2003**, 282, 85-101	2.1	45
49	Biological structures and bottom type influence habitat choices made by Alaska flatfishes. *Journal of Experimental Marine Biology and Ecology*, **2003**, 292, 43-59	2.1	63
48	Quantitative description of habitat suitability for the juvenile common sole (Solea solea, L.) in the Bay of Biscay (France) and the contribution of different habitats to the adult population. *Journal of Sea Research*, **2003**, 50, 139-149	1.9	130
47	Estimating limits to the spatial extent and suitability of sole (Solea solea) nursery grounds in the Dover Strait. *Journal of Sea Research*, **2003**, 50, 151-165	1.9	44
46	Shallow water predation risk for a juvenile flatfish (winter flounder; Pseudopleuronectes americanus, Walbaum) in a northwest Atlantic estuary. *Journal of Experimental Marine Biology and Ecology*, **2004**, 304, 137-157	2.1	50
45	Growth, feeding and distribution of the solenette Buglossidium luteum with particular reference to its habitat preference. *Journal of Sea Research*, **2004**, 51, 211-217	1.9	18
44	The Planktonic Stages of Flatfishes: Physical and Biological Interactions in Transport Processes. 94-119		17
43	Applying the basin model: Assessing habitat suitability of young-of-the-year demersal fishes on the New York Bight continental shelf. *Continental Shelf Research*, **2006**, 26, 1551-1570	2.4	9
42	Using habitat suitability index and particle dispersion models for early detection of marine invaders. **2006**, 16, 1377-90		44
41	The distribution of life cycle stages of two deep-water pleuronectids, Dover sole (Microstomus pacificus) and rex sole (Glyptocephalus zachirus), at the northern extent of their range in the Gulf of Alaska. *Journal of Sea Research*, **2007**, 57, 198-208	1.9	7
40	Flatfish-habitat associations in Alaska nursery grounds: Use of continuous video records for multi-scale spatial analysis. *Journal of Sea Research*, **2007**, 57, 137-150	1.9	46
39	Habitat suitability for juvenile common sole (Solea solea, L.) in the Bay of Biscay (France): A quantitative description using indicators based on epibenthic fauna. *Journal of Sea Research*, **2007**, 57, 126-136	1.9	46
38	Using decision trees to predict benthic communities within and near the German Exclusive Economic Zone (EEZ) of the North Sea. *Environmental Monitoring and Assessment*, **2008**, 136, 313-25	3.1	22
37	Berechnung einer landschaftsökologischen Raumgliederung Europas. *Environmental Sciences Europe*, **2008**, 20, 25-35	5	6

36	Elemental chemistry of left and right sagittal otoliths in a marine fish Hippoglossus stenolepis displaying cranial asymmetry. *Journal of Fish Biology*, **2008**, 73, 870-887	1.9	11
35	Ocean transport paths for the early life history stages of offshore-spawning flatfishes: a case study in the Gulf of Alaska. *Fish and Fisheries*, **2008**, 9, 44-66	6	38
34	Modelling the spatial distribution of plaice (Pleuronectes platessa), sole (Solea solea) and thornback ray (Raja clavata) in UK waters for marine management and planning. *Journal of Sea Research*, **2009**, 61, 258-267	1.9	58
33	Fine-scale population genetic structure in Alaskan Pacific halibut (Hippoglossus stenolepis). *Conservation Genetics*, **2010**, 11, 999-1012	2.6	14
32	Stream Condition in Piedmont Streams with Restored Riparian Buffers in the Chesapeake Bay Watershed1. *Journal of the American Water Resources Association*, **2010**, 46, 473-485	2.1	20
31	Habitat distribution model for European flounder juveniles in the Venice lagoon. *Journal of Sea Research*, **2010**, 64, 133-144	1.9	21
30	Application of CART in ecological landscape mapping: Two case studies. *Ecological Indicators*, **2011**, 11, 115-122	5.8	24
29	Updated analysis of flatfish recruitment response to climate variability and ocean conditions in the Eastern Bering Sea. *Deep-Sea Research Part II: Topical Studies in Oceanography*, **2013**, 94, 157-164	2.3	26
28	Modeled connectivity between northern rock sole (Lepidopsetta polyxystra) spawning and nursery areas in the eastern Bering Sea. *Journal of Sea Research*, **2013**, 84, 2-12	1.9	24
27	Predicting estuarine use patterns of juvenile fish with Generalized Linear Models. *Estuarine, Coastal and Shelf Science*, **2013**, 120, 64-74	2.9	29
26	Effect of starvation on condition and growth of juvenile plaice Pleuronectes platessa: nursery habitat quality assessment during the settlement period. *Journal of the Marine Biological Association of the United Kingdom*, **2013**, 93, 479-488	1.1	12
25	The role of wind-forcing in the distribution of larval fish in Galway Bay, Ireland. *Journal of the Marine Biological Association of the United Kingdom*, **2013**, 93, 471-478	1.1	8
24	Nursery areas of juvenile northern rock sole (Lepidopsetta polyxystra) in the eastern Bering Sea in relation to hydrography and thermal regimes. *ICES Journal of Marine Science*, **2014**, 71, 1683-1695	2.7	17
23	Quantitative mapping of fish habitat: A useful tool to design spatialised management measures and marine protected area with fishery objectives. *Ocean and Coastal Management*, **2014**, 87, 8-19	3.9	39
22	The food limitation hypothesis for juvenile marine fish. *Fish and Fisheries*, **2015**, 16, 373-398	6	74
21	Incorporation of bomb-produced14C into fish otoliths. An example of basin-specific rates from the North Pacific Ocean. *Canadian Journal of Fisheries and Aquatic Sciences*, **2015**, 72, 879-892	2.4	7
20	Shallow-water habitat use by Bering Sea flatfishes along the central Alaska Peninsula. *Journal of Sea Research*, **2016**, 111, 37-46	1.9	8
19	Introduction to the North Pacific Research Board Gulf of Alaska Integrated Ecosystem Research Program (GOAIERP): Volume I. *Deep-Sea Research Part II: Topical Studies in Oceanography*, **2016**, 132, 1-5	2.3	5

18	Using smooth sheets to describe groundfish habitat in Alaskan waters, with specific application to two flatfishes. *Deep-Sea Research Part II: Topical Studies in Oceanography*, **2016**, 132, 210-226	2.3	7
17	Assessment of resource selection models to predict occurrence of five juvenile flatfish species (Pleuronectidae) over the continental shelf in the western Gulf of Alaska. *Journal of Sea Research*, **2016**, 111, 54-64	1.9	6
16	Effect of sand grain size on substrate preference and burial behaviour in cultured Japanese flounder juvenile, Paralichthys olivaceus. *Aquaculture Research*, **2018**, 49, 1664-1671	1.9	4
15	A full life history synthesis of Arrowtooth Flounder ecology in the Gulf of Alaska: Exposure and sensitivity to potential ecosystem change. *Journal of Sea Research*, **2018**, 142, 28-51	1.9	6
14	Comparison of the physical attributes of the central and eastern Gulf of Alaska integrated ecosystem research program inshore study sites. *Deep-Sea Research Part II: Topical Studies in Oceanography*, **2019**, 165, 280-291	2.3	6
13	Habitat suitability models for groundfish in the Gulf of Alaska. *Deep-Sea Research Part II: Topical Studies in Oceanography*, **2019**, 165, 303-321	2.3	14
12	Age validation of four rockfishes (genera Sebastes and Sebastolobus) with bomb-produced radiocarbon. *Marine and Freshwater Research*, **2020**, 71, 1355	2.2	1
11	Eddy retention and seafloor terrain facilitate cross-shelf transport and delivery of fish larvae to suitable nursery habitats. *Limnology and Oceanography*, **2020**, 65, 2800-2818	4.8	3
10	Multiple life-stage connectivity of Pacific halibut (Hippoglossus stenolepis) across the Bering Sea and Gulf of Alaska. *Fisheries Oceanography*, **2021**, 30, 174-193	2.4	1
9	Modeling nearshore fish habitats using Alaska as a regional case study. *Fisheries Research*, **2021**, 238, 105905	2.3	3
8	Hazards evaluation of a valuable vulnerable sand-wave field forage fish habitat in the marginal Central Salish Sea using a submersible. *Oceanologia*, **2021**,	2.2	1
7	The potential utility of otolith microchemistry as an indicator of nursery origins in Pacific halibut (Hippoglossus stenolepis). *Fisheries Research*, **2021**, 243, 106072	2.3	0
6	The planktonic stages of flatfishes: physical and biological interactions in transport processes. 132-170		3
5	Growth and distributional correlates of behavior in three co-occurring juvenile flatfishes. *Marine Ecology - Progress Series*, **2012**, 460, 183-193	2.6	14
4	Substrate preference and delayed settlement in northern rock sole larvae Lepidopsetta polyxystra. *Marine Ecology - Progress Series*, **2015**, 519, 183-193	2.6	10
3	Environmental Drivers of Nearshore Fish Community Composition and Size Structure in Glacially Influenced Gulf of Alaska Estuaries. *Estuaries and Coasts*,	2.8	2
2	Synthesizing integrated ecosystem research to create informed stock-specific indicators for next generation stock assessments. *Deep-Sea Research Part II: Topical Studies in Oceanography*, **2022**, 105070	2.3	
1	Small and medium-scale biorefineries: Biomass quantification and its bioeconomic potential in the Southern Coastal Territory of Bahia.		0