

CITATION REPORT

List of articles citing

**The Inventory of Depressive Symptomatology (IDS):
Clinician (IDS-C) and Self-Report (IDS-SR) ratings of
depressive symptoms**

DOI: 10.1002/mpr.79

**International Journal of Methods in Psychiatric
Research, 2000, 9, 45-59.**

Source: <https://exaly.com/paper-pdf/31306742/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
278	A comparison of alternative assessments of depressive symptom severity: a pilot study. <i>Psychiatry Research</i> , 2000 , 95, 55-65	9.9	6
277	A comparison of alternative assessments of depressive symptom severity: a pilot study. <i>Psychiatry Research</i> , 2000 , 96, 269-79	9.9	28
276	Conceptual underpinnings and empirical support for the mood spectrum. 2002 , 25, 699-712, v		54
275	Background and rationale for the sequenced treatment alternatives to relieve depression (STAR*D) study. 2003 , 26, 457-94, x		373
274	The 16-Item Quick Inventory of Depressive Symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): a psychometric evaluation in patients with chronic major depression. 2003 , 54, 573-83		2278
273	Sequenced treatment alternatives to relieve depression (STAR*D): rationale and design. 2004 , 25, 119-42		725
272	Clinical features of depressed outpatients with and without co-occurring general medical conditions in STAR*D. 2004 , 26, 421-9		83
271	The Implementation of Cognitive Therapy in STAR*D. <i>Cognitive Therapy and Research</i> , 2004 , 28, 819-833	2.7	5
270	Factors associated with chronic depressive episodes: a preliminary report from the STAR-D project. 2005 , 112, 425-33		137
269	Comorbid psychiatric disorders in depressed outpatients: demographic and clinical features. <i>Journal of Affective Disorders</i> , 2005 , 87, 43-55	6.6	185
268	Self-reported depressive symptom measures: sensitivity to detecting change in a randomized, controlled trial of chronically depressed, nonpsychotic outpatients. 2005 , 30, 405-16		96
267	Comparison of the Internal State Scale to Clinician-Administered Assessments in Patients with Bipolar Disorder and Alcohol Abuse or Dependence. 2005 , 1, 61-69		
266	Quality improvement methods as applied to a multicenter effectiveness trial--STAR*D. 2005 , 26, 95-112		6
265	Effects of 12 months of vagus nerve stimulation in treatment-resistant depression: a naturalistic study. 2005 , 58, 355-63		288
264	Vagus nerve stimulation for treatment-resistant depression: a randomized, controlled acute phase trial. 2005 , 58, 347-54		439
263	A one-year comparison of vagus nerve stimulation with treatment as usual for treatment-resistant depression. 2005 , 58, 364-73		264
262	An evaluation of the quick inventory of depressive symptomatology and the hamilton rating scale for depression: a sequenced treatment alternatives to relieve depression trial report. 2006 , 59, 493-501		300

261	Algorithms and collaborative-care systems for depression: are they effective and why? A systematic review. 2006 , 59, 1029-38		102
260	Exercise as an augmentation strategy for treatment of major depression. 2006 , 12, 205-13		120
259	Clinical and demographic factors associated with DSM-IV melancholic depression. 2006 , 18, 91-8		86
258	TREAD: Treatment with Exercise Augmentation for Depression: study rationale and design. 2006 , 3, 291-305		60
257	DSM melancholic features are unreliable predictors of ECT response: a CORE publication. 2007 , 23, 139-46		39
256	Comparison of quality of life measures in a depressed population. 2007 , 195, 219-25		27
255	Measurement-based care for refractory depression: a clinical decision support model for clinical research and practice. 2007 , 88 Suppl 2, S61-71		74
254	Assessing physicians' use of treatment algorithms: Project IMPACTS study design and rationale. 2007 , 28, 192-212		17
253	Pain in depression: STAR*D study findings. 2007 , 63, 113-22		64
252	Sertraline as a first-line treatment for cholestatic pruritus. 2007 , 45, 666-74		207
251	Clinical differences among depressed patients with and without a history of suicide attempts: findings from the STAR*D trial. <i>Journal of Affective Disorders</i> , 2007 , 97, 77-84	6.6	59
250	Influences of hormone-based contraception on depressive symptoms in premenopausal women with major depression. 2007 , 32, 843-53		61
249	Clinical vs. self-report versions of the quick inventory of depressive symptomatology in a public sector sample. <i>Journal of Psychiatric Research</i> , 2007 , 41, 239-46	5.2	40
248	Electronic source materials in clinical research: acceptability and validity of symptom self-rating in major depressive disorder. <i>Journal of Psychiatric Research</i> , 2007 , 41, 737-43	5.2	18
247	Symptom features of postpartum depression: are they distinct?. <i>Depression and Anxiety</i> , 2008 , 25, 20-6	8.4	81
246	Family history of completed suicide and characteristics of major depressive disorder: a STAR*D (sequenced treatment alternatives to relieve depression) study. <i>Journal of Affective Disorders</i> , 2008 , 108, 129-34	6.6	12
245	How much cognitive therapy, for which patients, will prevent depressive relapse?. <i>Journal of Affective Disorders</i> , 2008 , 111, 185-92	6.6	25
244	The FKBP5-gene in depression and treatment response--an association study in the Sequenced Treatment Alternatives to Relieve Depression (STAR*D) Cohort. 2008 , 63, 1103-10		218

243	Daily process methodology for measuring earlier antidepressant response. 2008 , 29, 867-77		21
242	Sex differences in depression symptoms in treatment-seeking adults: confirmatory analyses from the Sequenced Treatment Alternatives to Relieve Depression study. <i>Comprehensive Psychiatry</i> , 2008 , 49, 238-46	7.3	156
241	Antecedentes familiares de suicidio consumado y características del trastorno depresivo mayor: estudio STAR*D (sequenced treatment alternatives to relieve depression). 2008 , 15, 195-201		
240	Pharmacogenetic analysis of genes implicated in rodent models of antidepressant response: association of TREK1 and treatment resistance in the STAR(*)D study. 2008 , 33, 2810-9		89
239	Increased vesicular monoamine transporter binding during early abstinence in human methamphetamine users: Is VMAT2 a stable dopamine neuron biomarker?. 2008 , 28, 9850-6		80
238	Making the business case for enhanced depression care: the National Institute of Mental Health-harvard Work Outcomes Research and Cost-effectiveness Study. 2008 , 50, 468-75		33
237	Effectiveness Study of Venlafaxine-XR Combined with Aripiprazole for Chronic or Recurrent Major Depressive Disorder. 2009 , 43, 956-967		8
236	The epidemiology of depression in metropolitan China. <i>Psychological Medicine</i> , 2009 , 39, 735-47	6.9	121
235	Decreased binding of the D3 dopamine receptor-preferring ligand [11C]-(+)-PHNO in drug-naive Parkinson's disease. 2009 , 132, 1366-75		85
234	Genetic and clinical predictors of sexual dysfunction in citalopram-treated depressed patients. 2009 , 34, 1819-28		78
233	What predicts attrition in second step medication treatments for depression?: a STAR*D Report. 2009 , 12, 459-73		58
232	Income and attrition in the treatment of depression: a STAR*D report. <i>Depression and Anxiety</i> , 2009 , 26, 622-33	8.4	39
231	Diurnal mood variation in outpatients with major depressive disorder. <i>Depression and Anxiety</i> , 2009 , 26, 851-63	8.4	18
230	Sociodemographic, clinical, and treatment characteristics associated with worsened depression during treatment with citalopram: results of the NIMH STAR(*)D trial. <i>Depression and Anxiety</i> , 2009 , 26, 612-21	8.4	25
229	A psychometric evaluation of the clinician-rated Quick Inventory of Depressive Symptomatology (QIDS-C16) in patients with bipolar disorder. <i>International Journal of Methods in Psychiatric Research</i> , 2009 , 18, 138-46	4.3	16
228	Release bias in accessing medical records in clinical trials: a STAR*D report. <i>International Journal of Methods in Psychiatric Research</i> , 2009 , 18, 147-58	4.3	1
227	Concurrent anxiety and substance use disorders among outpatients with major depression: clinical features and effect on treatment outcome. 2009 , 99, 248-60		109
226	STAR*D: revising conventional wisdom. 2009 , 23, 627-47		172

225	Predictors of attrition during one year of depression treatment: a roadmap to personalized intervention. 2009 , 15, 113-24		22
224	Screening for major depression in private practice. 2009 , 15, 87-94		18
223	Genome-wide association study of suicidal ideation emerging during citalopram treatment of depressed outpatients. 2009 , 19, 666-74		80
222	A systematic overview of measurement-based care in the treatment of childhood and adolescent depression. 2010 , 16, 217-34		15
221	Early adverse events and attrition in selective serotonin reuptake inhibitor treatment: a suicide assessment methodology study report. 2010 , 30, 259-66		17
220	The quick inventory of depressive symptomatology (clinician and self-report versions) in patients with bipolar disorder. 2010 , 15, 367-73		19
219	Frequency and correlates of gambling problems in outpatients with major depressive disorder and bipolar disorder. 2010 , 55, 568-76		39
218	A Comparison of the QIDS-C16, QIDS-SR16, and the MADRS in an Adult Outpatient Clinical Sample. 2010 , 15, 458-68		20
217	Using the QIDS-SR16 to identify major depressive disorder in primary care medical patients. 2010 , 41, 423-31		32
216	Employer burden of mild, moderate, and severe major depressive disorder: mental health services utilization and costs, and work performance. <i>Depression and Anxiety</i> , 2010 , 27, 78-89	8.4	257
215	Screening for serious mental illness in the general population with the K6 screening scale: results from the WHO World Mental Health (WMH) survey initiative. <i>International Journal of Methods in Psychiatric Research</i> , 2010 , 19 Suppl 1, 4-22	4.3	686
214	Psychometric properties of the Quick Inventory of Depressive Symptomatology in adolescents. <i>International Journal of Methods in Psychiatric Research</i> , 2010 , 19, 185-94	4.3	75
213	Diagnostic utility of the Quick Inventory of Depressive Symptomatology (QIDS-C16 and QIDS-SR16) in the elderly. 2010 , 122, 226-34		26
212	The importance of irritability as a symptom of major depressive disorder: results from the National Comorbidity Survey Replication. 2010 , 15, 856-67		119
211	Prevalence, impairment and severity of 12-month DSM-IV major depressive episodes in Te Rau Hinengaro: New Zealand Mental Health Survey 2003/4. 2010 , 44, 750-8		7
210	Anticipated Benefits of Care (ABC): psychometrics and predictive value in psychiatric disorders. <i>Psychological Medicine</i> , 2010 , 40, 955-65	6.9	1
209	Painful physical symptoms and treatment outcome in major depressive disorder: a STAR*D (Sequenced Treatment Alternatives to Relieve Depression) report. <i>Psychological Medicine</i> , 2010 , 40, 239-51	6.9	66
208	Decreased cerebral cortical serotonin transporter binding in ecstasy users: a positron emission tomography/[¹¹ C]DASB and structural brain imaging study. 2010 , 133, 1779-97		122

207	Does comorbid substance use disorder impair recovery from major depression with SSRI treatment? An analysis of the STAR*D level one treatment outcomes. 2010 , 107, 161-70		59
206	Effect of naltrexone plus bupropion on weight loss in overweight and obese adults (COR-I): a multicentre, randomised, double-blind, placebo-controlled, phase 3 trial. 2010 , 376, 595-605		599
205	Optimal zone for bandwidth selection in semiparametric models. 2011 , 23, 701-717		14
204	Review of major measuring instruments in comorbid depression and coronary heart disease. 2011 , 35, 905-12		8
203	Introduction: Evidence-Based Practice for Major Depressive Disorder. 2011 , 1-19		
202	Residual symptoms in depressed outpatients who respond by 50% but do not remit to antidepressant medication. 2011 , 31, 180-6		123
201	The relationship between adverse events during selective serotonin reuptake inhibitor treatment for major depressive disorder and nonremission in the suicide assessment methodology study. 2011 , 31, 31-8		13
200	Psychometric considerations of depression symptom rating scales. 2011 , 1, 611-623		6
199	Weight loss with naltrexone SR/bupropion SR combination therapy as an adjunct to behavior modification: the COR-BMOD trial. 2011 , 19, 110-20		308
198	The treatment of minor depression with St. John's Wort or citalopram: failure to show benefit over placebo. <i>Journal of Psychiatric Research</i> , 2011 , 45, 931-41	5.2	37
197	International Study to Predict Optimized Treatment for Depression (iSPOT-D), a randomized clinical trial: rationale and protocol. <i>Trials</i> , 2011 , 12, 4	2.8	126
196	The Inventory Of Depressive Symptomatology (IDS-C(28)) is more sensitive to changes in depressive symptomatology than the Hamilton Depression Rating Scale (HAMD(17)) in patients with mild major, minor or subsyndromal depression. 2011 , 261, 357-67		19
195	Psychometric properties of the IDS-SR30 for the assessment of depressive symptoms in Spanish population. 2011 , 11, 131		6
194	Effectiveness, relapse prevention and mechanisms of change of cognitive therapy vs. interpersonal therapy for depression: Study protocol for a randomised controlled trial. <i>Trials</i> , 2011 , 12, 150	2.8	26
193	Stimulant reduction intervention using dosed exercise (STRIDE) - CTN 0037: study protocol for a randomized controlled trial. <i>Trials</i> , 2011 , 12, 206	2.8	37
192	Is prior course of illness relevant to acute or longer-term outcomes in depressed out-patients? A STAR*D report. <i>Psychological Medicine</i> , 2012 , 42, 1131-49	6.9	46
191	Correlates and outcomes of depressed out-patients with greater and fewer anxious symptoms: a CO-MED report. 2012 , 15, 1387-99		18
190	Applications of the Bootstrap in ROC Analysis. 2012 , 41, 865-877		4

189	Evidence-based practices in community mental health: outcome evaluation. 2012 , 39, 434-44		8
188	Baseline depression severity as a predictor of single and combination antidepressant treatment outcome: results from the CO-MED trial. 2012 , 22, 183-99		48
187	Escitalopram for severe asthma and major depressive disorder: a randomized, double-blind, placebo-controlled proof-of-concept study. 2012 , 53, 75-80		23
186	What are the clinical implications of new onset or worsening anxiety during the first two weeks of SSRI treatment for depression?. <i>Depression and Anxiety</i> , 2012 , 29, 94-101	8.4	20
185	Effect of concurrent substance use disorder on the effectiveness of single and combination antidepressant medications for the treatment of major depression: an exploratory analysis of a single-blind randomized trial. <i>Depression and Anxiety</i> , 2012 , 29, 111-22	8.4	13
184	Sensitivity to changes during antidepressant treatment: a comparison of unidimensional subscales of the Inventory of Depressive Symptomatology (IDS-C) and the Hamilton Depression Rating Scale (HAMD) in patients with mild major, minor or subsyndromal depression. 2012 , 262, 291-304		14
183	Effects of heart disease on depression treatment: results from the COMED study. 2012 , 34, 24-34		4
182	The prevalence and impact of major depressive disorder among Chinese, Malays and Indians in an Asian multi-racial population. <i>Journal of Affective Disorders</i> , 2012 , 138, 128-36	6.6	29
181	Altered slow wave activity in major depressive disorder with hypersomnia: a high density EEG pilot study. <i>Psychiatry Research - Neuroimaging</i> , 2012 , 201, 240-4	2.9	22
180	A comparative analysis between site-based and centralized ratings and patient self-ratings in a clinical trial of Major Depressive Disorder. <i>Journal of Psychiatric Research</i> , 2013 , 47, 944-54	5.2	20
179	Vagus nerve stimulation therapy randomized to different amounts of electrical charge for treatment-resistant depression: acute and chronic effects. 2013 , 6, 631-40		107
178	Prevalence of and factors related to the use of antidepressants and benzodiazepines: results from the Singapore Mental Health Study. <i>BMC Psychiatry</i> , 2013 , 13, 231	4.2	10
177	A randomized, phase 3 trial of naltrexone SR/bupropion SR on weight and obesity-related risk factors (COR-II). 2013 , 21, 935-43		381
176	Does exercise improve self-reported sleep quality in non-remitted major depressive disorder?. <i>Psychological Medicine</i> , 2013 , 43, 699-709	6.9	36
175	Using multiple methods to characterize the phenotype of individuals with a family history of major depressive disorder. <i>Journal of Affective Disorders</i> , 2013 , 150, 474-80	6.6	14
174	Talking bodies: nonverbal behavior in the assessment of depression severity. <i>Journal of Affective Disorders</i> , 2013 , 150, 1114-9	6.6	9
173	Assessing depression in obese women: an examination of two commonly-used measures. 2013 , 75, 425-30		1
172	Iowa Gambling Task scores predict future drug use in bipolar disorder outpatients with stimulant dependence. <i>Psychiatry Research</i> , 2013 , 210, 871-9	9.9	19

171	Neuropsychological testing of cognitive impairment in euthymic bipolar disorder: an individual patient data meta-analysis. 2013 , 128, 149-62		373
170	Pro-inflammatory cytokines as predictors of antidepressant effects of exercise in major depressive disorder. 2013 , 18, 1119-24		137
169	FP02 Beta Training for Drug-Resistant Depression: A New Protocol That Usually Reduces Depression and Keeps It Reduced. 2013 , 17, 198-200		11
168	Clinical and sociodemographic characteristics associated with suicidal ideation in depressed outpatients. 2013 , 58, 113-22		25
167	Sexual functioning in patients with recurrent major depressive disorder enrolled in the PREVENT study. 2013 , 201, 266-73		16
166	Models of depression. 1-6		
165	The diagnosis of mood disorders. 7-19		
164	Blended vs. face-to-face cognitive behavioural treatment for major depression in specialized mental health care: study protocol of a randomized controlled cost-effectiveness trial. <i>BMC Psychiatry</i> , 2014 , 14, 290	4.2	52
163	Association of depressive symptoms with hippocampal volume in 1936 adults. 2014 , 39, 770-9		45
162	The Dark-Side of Rule Governed Behavior: An Experimental Analysis of Problematic Rule-Following in an Adolescent Population With Depressive Symptomatology. 2014 , 38, 587-613		35
161	The Spanish version of the Quick Inventory of Depressive Symptomatology-Self-Report (QIDS-SR16): a psychometric analysis in a clinical sample. <i>Journal of Affective Disorders</i> , 2014 , 169, 189-96 ^{6.6}		10
160	A randomized controlled trial of cognitive remediation and d-cycloserine for individuals with bipolar disorder. 2014 , 2, 41		10
159	Correlates of depression in bipolar disorder. 2014 , 281, 20132320		9
158	Tablet-based screening of depressive symptoms in quito, ecuador: efficiency in primary care. 2014 , 2014, 845397		7
157	Age and task difficulty differences in dual tasking using circle tracing and serial subtraction tasks. 2014 , 26, 201-11		9
156	Frontal alpha EEG asymmetry before and after behavioral activation treatment for depression. 2014 , 99, 198-208		84
155	A randomized, double-blind, placebo-controlled trial of pregnenolone for bipolar depression. 2014 , 39, 2867-73		58
154	Mood dynamics in bipolar disorder. 2014 , 2, 11		20

153	Curcumin for the treatment of major depression: a randomised, double-blind, placebo controlled study. <i>Journal of Affective Disorders</i> , 2014 , 167, 368-75	6.6	123
152	Resting-state functional connectivity in women with Major Depressive Disorder. <i>Journal of Psychiatric Research</i> , 2014 , 59, 38-44	5.2	19
151	Magnitude of change with antidepressants and placebo in antidepressant clinical trials using structured, taped and appraised rater interviews (SIGMA-RAPS) compared to trials using traditional semi-structured interviews. 2014 , 231, 4301-7		6
150	Magnitude of placebo response and response variance in antidepressant clinical trials using structured, taped and appraised rater interviews compared to traditional rating interviews. <i>Journal of Psychiatric Research</i> , 2014 , 51, 88-92	5.2	13
149	Developing and Selecting Mental Health Measures. 2014 , 143-169		2
148	The Assessment of Depression. 2015 , 24-51		0
147	Modifying Cognitive-Behavioral Therapy for a Depressed Older Adult With Partial Sight: A Case Report. 2015 , 1, 2333721415585432		2
146	Differences in the neural correlates of affective responses in depressed and healthy women. <i>Psychiatry Research - Neuroimaging</i> , 2015 , 234, 336-45	2.9	6
145	Randomized, placebo-controlled, adjunctive study of armodafinil for bipolar I depression: implications of novel drug design and heterogeneity of concurrent bipolar maintenance treatments. 2015 , 3, 34		20
144	Novel Study Designs for Clinical Trials in Mood Disorders. 2015 , 87-104		0
143	Dual Task Performance May be a Better Measure of Cognitive Processing in Huntington's Disease than Traditional Attention Tests. 2015 , 4, 119-30		6
142	Mediators of sexual functioning and marital quality in chronically depressed adults with and without a history of childhood sexual abuse. 2015 , 12, 813-23		24
141	Effects of task difficulty during dual-task circle tracing in Huntington's disease. 2015 , 262, 268-76		11
140	Hamilton depression rating subscales to predict antidepressant treatment outcome in the early course of treatment. <i>Journal of Affective Disorders</i> , 2015 , 175, 199-208	6.6	5
139	Reduced striatal activation in females with major depression during the processing of affective stimuli. <i>Journal of Psychiatric Research</i> , 2015 , 68, 384-91	5.2	7
138	Adjunctive armodafinil for major depressive episodes associated with bipolar I disorder. <i>Journal of Affective Disorders</i> , 2015 , 181, 87-91	6.6	25
137	Improving systematic reviews on psychometric properties of measurement instruments: a letter to the Editor commenting on Reilly et al.'s (2015) review of the psychometric literature on QIDS. <i>Journal of Psychiatric Research</i> , 2015 , 62, 136-7	5.2	
136	Sleep disturbance among people with major depressive disorders (MDD) in Singapore. 2016 , 25, 492-499		12

135	Covariates of depression and high utilizers of healthcare: Impact on resource use and costs. 2016 , 85, 35-43		19
134	Atypical depressive symptoms as a predictor of treatment response to exercise in Major Depressive Disorder. <i>Journal of Affective Disorders</i> , 2016 , 200, 156-8	6.6	12
133	Randomized controlled study of early medication change for non-improvers to antidepressant therapy in major depression--The EMC trial. 2016 , 26, 705-16		30
132	Concurrent and prospective associations between HPA axis activity and depression symptoms in newlywed women. 2016 , 73, 125-132		4
131	Group schema therapy versus group cognitive behavioral therapy for social anxiety disorder with comorbid avoidant personality disorder: study protocol for a randomized controlled trial. <i>Trials</i> , 2016 , 17, 487	2.8	8
130	Mindfulness based stress reduction adapted for depressed disadvantaged women in an urban Federally Qualified Health Center. 2016 , 25, 59-67		23
129	Investigating the (cost-) effectiveness of attention bias modification (ABM) for outpatients with major depressive disorder (MDD): a randomized controlled trial protocol. <i>BMC Psychiatry</i> , 2016 , 16, 370	4.2	6
128	Tablet-based support for older adults with severe mood disorders treated in an ambulatory geriatric psychiatry setting: Protocol of a feasibility study of the eCare@Home platform. 2016 , 6, 22-28		1
127	Interaction of oxytocin level and past depression may predict postpartum depressive symptom severity. 2016 , 19, 799-808		16
126	Development and initial evaluation of blended cognitive behavioural treatment for major depression in routine specialized mental health care. 2016 , 4, 61-71		46
125	An open-label pilot study of icariin for co-morbid bipolar and alcohol use disorder. 2016 , 42, 162-7		12
124	Can Medication Free, Treatment-Resistant, Depressed Patients Who Initially Respond to TMS Be Maintained Off Medications? A Prospective, 12-Month Multisite Randomized Pilot Study. 2016 , 9, 251-7		38
123	Twice the negativity bias and half the positivity offset: Evaluative responses to emotional information in depression. 2016 , 52, 166-170		20
122	The negativity bias predicts response rate to Behavioral Activation for depression. 2016 , 52, 171-178		4
121	Microtubule associated protein 2 in bipolar depression: Impact of pregnenolone. <i>Journal of Affective Disorders</i> , 2017 , 218, 49-52	6.6	7
120	A combined marker of early non-improvement and the occurrence of melancholic features improve the treatment prediction in patients with Major Depressive Disorders. <i>Journal of Affective Disorders</i> , 2017 , 221, 184-191	6.6	4
119	Early Pubertal Timing and Childhood Family Adversity Interact to Predict Newlywed Women's Anxiety Symptoms. 2017 , 26, 591-602		2
118	The efficacy of the modified Atkins diet in North Sea Progressive Myoclonus Epilepsy: an observational prospective open-label study. 2017 , 12, 45		6

117	The Frequency of PTSD and Subthreshold PTSD among African-American Women with Depressive Symptoms in a Disadvantaged Urban Neighborhood: Pilot Study. 2017 , 4, 1069-1073		4
116	Impact of morbid obesity on treatment outcome in a clinical trial of major depressive disorder. 2017 , 3, 18-23		2
115	Continuation and maintenance treatments for persistent depressive disorder. <i>The Cochrane Library</i> , 2017 ,	5.2	2
114	An actigraphy study investigating sleep in bipolar I patients, unaffected siblings and controls. <i>Journal of Affective Disorders</i> , 2017 , 208, 248-254	6.6	10
113	Exercise is an effective treatment for positive valence symptoms in major depression. <i>Journal of Affective Disorders</i> , 2017 , 209, 188-194	6.6	30
112	Interpersonal Circumplex Profiles Of Persistent Depression: Goals, Self-Efficacy, Problems, And Effects Of Group Therapy. 2017 , 73, 595-611		20
111	Efficacy of curcumin, and a saffron/curcumin combination for the treatment of major depression: A randomised, double-blind, placebo-controlled study. <i>Journal of Affective Disorders</i> , 2017 , 207, 188-196	6.6	80
110	Early treatment response affects signal detection in a placebo-controlled depression study. 2017 , 4-6, 19-24		0
109	A review of physiological and behavioral monitoring with digital sensors for neuropsychiatric illnesses. 2018 , 39, 05TR01		57
108	Real-time fMRI neurofeedback training of the amygdala activity with simultaneous EEG in veterans with combat-related PTSD. 2018 , 19, 106-121		56
107	A longitudinal study of women's depression symptom profiles during and after the postpartum phase. <i>Depression and Anxiety</i> , 2018 , 35, 292-304	8.4	12
106	Effectiveness of smartphone-based ambulatory assessment (SBAA-BD) including a predicting system for upcoming episodes in the long-term treatment of patients with bipolar disorders: study protocol for a randomized controlled single-blind trial. <i>BMC Psychiatry</i> , 2018 , 18, 349	4.2	10
105	An augmentation study of MSI-195 (S-adenosylmethionine) in Major Depressive Disorder. <i>Journal of Psychiatric Research</i> , 2018 , 107, 86-96	5.2	6
104	Are rating scales for bipolar disorders fit for purpose?. 2018 , 213, 627-629		1
103	Accessibility and feasibility of using technology to support mindfulness practice, reduce stress and promote long term mental health. 2018 , 33, 93-99		8
102	Prevention of violent revictimization in depressed patients with an add-on internet-based emotion regulation training (iERT): study protocol for a multicenter randomized controlled trial. <i>BMC Psychiatry</i> , 2018 , 18, 29	4.2	6
101	Control charts for monitoring mood stability as a predictor of severe episodes in patients with bipolar disorder. 2018 , 6, 7		12
100	Functional connectivity of reflective and brooding rumination in depressed and healthy women. 2018 , 18, 884-901		15

99	The Promoting Activity in Cancer Survivors (PACES) trial: a multiphase optimization of strategy approach to increasing physical activity in breast cancer survivors. 2018 , 18, 744		6
98	A longitudinal approach to biological psychiatric research: The PsyCourse study. 2019 , 180, 89-102		24
97	Comparability of blinded remote and site-based assessments of response to adjunctive esketamine or placebo nasal spray in patients with treatment resistant depression. <i>Journal of Psychiatric Research</i> , 2019 , 111, 68-73	5.2	6
96	Childhood Adversity Is Associated With Increased Methylation in Healthy Individuals but Not in Bipolar Disorder Patients. <i>Frontiers in Psychiatry</i> , 2018 , 9, 743	5	7
95	Changes in mindfulness facets in a dialectical behaviour therapy skills training group program for borderline personality disorder. 2019 , 75, 958-969		7
94	Systematic Review of Symptom Assessment Measures for Use in Measurement-Based Care of Bipolar Disorders. 2019 , 70, 396-408		9
93	Acceptability of a mindfulness intervention for depressive symptoms among African-American women in a community health center: A qualitative study. 2019 , 45, 19-24		8
92	Efficacy of an internet-based self-management intervention for depression or dysthymia - a study protocol of an RCT using an active control condition. <i>BMC Psychiatry</i> , 2019 , 19, 90	4.2	4
91	A Mindfulness-Based Intervention for Low-Income African American Women with Depressive Symptoms Delivered by an Experienced Instructor Versus a Novice Instructor. 2019 , 25, 699-708		5
90	Remote assessment of disease and relapse in major depressive disorder (RADAR-MDD): a multi-centre prospective cohort study protocol. <i>BMC Psychiatry</i> , 2019 , 19, 72	4.2	44
89	A pilot study of brexpiprazole for bipolar depression. <i>Journal of Affective Disorders</i> , 2019 , 249, 315-318	6.6	15
88	Effects of Opioid Dependence on Visuospatial Memory and Its Associations With Depression and Anxiety. <i>Frontiers in Psychiatry</i> , 2019 , 10, 743	5	6
87	Comparative effectiveness of continuation and maintenance treatments for persistent depressive disorder in adults. <i>The Cochrane Library</i> , 2019 , 5, CD012855	5.2	16
86	Relationship between depressive symptom severity and amygdala volume in a large community-based sample. <i>Psychiatry Research - Neuroimaging</i> , 2019 , 283, 77-82	2.9	5
85	Predictors of the effectiveness of an early medication change strategy in patients with major depressive disorder. <i>BMC Psychiatry</i> , 2019 , 19, 24	4.2	2
84	A randomized-controlled trial of cognitive-behavioral therapy for depression with integrated techniques from emotion-focused and exposure therapies. 2019 , 29, 30-44		13
83	Patient characteristics and the therapist as predictors of depressed patients' outcome expectation over time: A multilevel analysis. 2019 , 29, 709-722		10
82	Predictors of depression and life satisfaction in visually impaired people. 2019 , 41, 1012-1023		2

81	Psychiatric assessment in Phelan-McDermid Syndrome (22q13 deletion syndrome). 2020 , 45, 54-58		9
80	Neural responses during extinction learning predict exposure therapy outcome in phobia: results from a randomized-controlled trial. 2020 , 45, 534-541		20
79	A Brain Imaging-Based Diagnostic Biomarker for Periodic Catatonia: Preliminary Evidence Using a Bayesian Approach. 2020 , 79, 352-365		5
78	Efficacy and tolerability of adjunctive modafinil/armodafinil in bipolar depression: A meta-analysis of randomized controlled trials. <i>Bipolar Disorders</i> , 2020 , 22, 109-120	3.8	9
77	Is sleep disturbance linked to short- and long-term outcomes following treatments for recurrent depression?. <i>Journal of Affective Disorders</i> , 2020 , 262, 323-332	6.6	6
76	Early score fluctuation and placebo response in a study of major depressive disorder. <i>Journal of Psychiatric Research</i> , 2020 , 121, 118-125	5.2	3
75	Group Cognitive Behavioral Analysis System of Psychotherapy (CBASP): A Pilot Study for Bipolar Depression. <i>Frontiers in Psychiatry</i> , 2020 , 11, 565681	5	1
74	Practical recommendations for the management of treatment-resistant depression with esketamine nasal spray therapy: Basic science, evidence-based knowledge and expert guidance. <i>World Journal of Biological Psychiatry</i> , 2021 , 22, 468-482	3.8	13
73	Psychotic depression in hospitalized patients: Longitudinal outcomes of psychotic vs. nonpsychotic depression among inpatients. <i>Journal of Psychiatric Research</i> , 2020 , 129, 73-79	5.2	5
72	Vasopressin Surrogate Marker Copeptin as a Potential Novel Endocrine Biomarker for Antidepressant Treatment Response in Major Depression: A Pilot Study. <i>Frontiers in Psychiatry</i> , 2020 , 11, 453	5	3
71	Effects of age on depressive symptomatology and response to antidepressant treatment in patients with major depressive disorder aged 18 to 65 years. <i>Comprehensive Psychiatry</i> , 2020 , 99, 152170	7.3	3
70	Discontinuation of long-term antidepressant use for depressive and anxiety disorders in adults. <i>The Cochrane Library</i> , 2020 ,	5.2	1
69	Fractal biomarker of activity in patients with bipolar disorder. <i>Psychological Medicine</i> , 2021 , 51, 1562-1569	6.9	4
68	CBT MobileWork ^r : User-Centered Development and Testing of a Mobile Mental Health Application for Depression. <i>Cognitive Therapy and Research</i> , 2021 , 45, 287-302	2.7	8
67	The long shadow of childhood trauma for depression in midlife: examining daily psychological stress processes as a persistent risk pathway. <i>Psychological Medicine</i> , 2021 , 1-10	6.9	1
66	Depressive Symptoms, External Stress, and Marital Adjustment: The Buffering Effect of Partner's Responsive Behavior.. <i>Social Psychological and Personality Science</i> , 2022 , 13, 220-232	4.3	2
65	Approaches for discontinuation versus continuation of long-term antidepressant use for depressive and anxiety disorders in adults. <i>The Cochrane Library</i> , 2021 , 4, CD013495	5.2	12
64	The D*Phase-study: study protocol for a pragmatic two-phased, randomised controlled (non-inferiority) trial that addresses treatment non-response and compares cognitive behavioural therapy and short-term psychodynamic supportive psychotherapy for major depression. <i>BMC Psychiatry</i> , 2021 , 21, 233	4.2	0

63	Evaluating the Clinical Feasibility of an Artificial Intelligence-Powered, Web-Based Clinical Decision Support System for the Treatment of Depression in Adults: Longitudinal Feasibility Study. <i>JMIR Formative Research</i> , 2021 , 5, e31862	2.5	0
62	Evaluating the Clinical Feasibility of an Artificial Intelligence-Powered Clinical Decision Support System: A Longitudinal Feasibility Study.		
61	The effect of esketamine in patients with treatment-resistant depression with and without comorbid anxiety symptoms or disorder. <i>Depression and Anxiety</i> , 2021 , 38, 1120-1130	8.4	0
60	Audio-Guided Mindfulness Meditation During Transcranial Magnetic Stimulation Sessions for the Treatment of Major Depressive Disorder: A Pilot Feasibility Study. <i>Frontiers in Psychology</i> , 2021 , 12, 6789314		
59	Virtual Reality Relaxation for Patients With a Psychiatric Disorder: Crossover Randomized Controlled Trial. <i>Journal of Medical Internet Research</i> , 2021 , 23, e17233	7.6	15
58	Pilot validation of blood-based biomarkers during pregnancy and postpartum in women with prior or current depression. <i>Translational Psychiatry</i> , 2021 , 11, 68	8.6	4
57	Algorithmusgestützte Behandlung. 2005 , 459-476		1
56	Quality of Life and Rating Scales of Depression. <i>Handbook of Experimental Pharmacology</i> , 2004 , 149-167	3.2	3
55	Cognitive impairment in individuals with bipolar disorder with and without comorbid alcohol and/or cocaine use disorders. <i>Journal of Affective Disorders</i> , 2020 , 272, 355-362	6.6	1
54	Gender differences in major depressive disorder: findings from the Singapore Mental Health Study. <i>Singapore Medical Journal</i> , 2017 , 58, 649-655	1.9	39
53	Dual task performance in normal aging: a comparison of choice reaction time tasks. <i>PLoS ONE</i> , 2013 , 8, e60265	3.7	30
52	The Epidemiology of Major Depressive Episode in the Iraqi General Population. <i>PLoS ONE</i> , 2015 , 10, e0131937	3.7	17
51	German Translation and Validation of the Cognitive Style Questionnaire Short Form (CSQ-SF-D). <i>PLoS ONE</i> , 2016 , 11, e0149530	3.7	4
50	Item Response Analysis of the Inventory of Depressive Symptomatology. <i>Neuropsychiatric Disease and Treatment</i> , 2006 , 2, 557-564	3.1	12
49	Cost and Effectiveness of Blended Versus Standard Cognitive Behavioral Therapy for Outpatients With Depression in Routine Specialized Mental Health Care: Pilot Randomized Controlled Trial. <i>Journal of Medical Internet Research</i> , 2019 , 21, e14261	7.6	28
48	Efficacy of a Guided Web-Based Self-Management Intervention for Depression or Dysthymia: Randomized Controlled Trial With a 12-Month Follow-Up Using an Active Control Condition. <i>Journal of Medical Internet Research</i> , 2020 , 22, e15361	7.6	9
47	Tamper-Resistant Mobile Health Using Blockchain Technology. <i>JMIR MHealth and UHealth</i> , 2017 , 5, e1115	5.5	124
46	Treatment strategies to improve and sustain remission in major depressive disorder. <i>Dialogues in Clinical Neuroscience</i> , 2008 , 10, 377-84	5.7	52

45	Turkish Validity & Reliability of The Quick Inventory of Depressive Symptomatology Adolescent Version (QIDS-A17-SR) In Comparison with The Beck Depression Inventory-II Among Late Adolescents. <i>Journal of Microbiology and Biotechnology</i> , 2016 , 26, 303-309	3.3	4
44	Multimodal assessment increases objective identification of hypersomnolence in patients referred for multiple sleep latency testing. <i>Journal of Clinical Sleep Medicine</i> , 2020 , 16, 1241-1248	3.1	1
43	The clinical and cost-effectiveness of a self-management intervention for patients with persistent depressive disorder and their partners/caregivers: study protocol of a multicenter pragmatic randomized controlled trial. <i>Trials</i> , 2021 , 22, 731	2.8	
42	Efficacy of a Guided Web-Based Self-Management Intervention for Depression or Dysthymia: Randomized Controlled Trial With a 12-Month Follow-Up Using an Active Control Condition (Preprint).		
41	Virtual Reality Relaxation for Patients With a Psychiatric Disorder: Crossover Randomized Controlled Trial (Preprint).		
40	Parkinson disease rating scales: a literature review. <i>Annals of Movement Disorders</i> , 2020 , 3, 3	0.3	1
39	Anterior cingulate cortex in individuals with depressive symptoms: A structural MRI study. <i>Psychiatry Research - Neuroimaging</i> , 2021 , 319, 111420	2.9	0
38	A randomized controlled trial of cognitive control training (CCT) as an add-on treatment for late-life depression: a study protocol. <i>BMC Psychiatry</i> , 2021 , 21, 596	4.2	
37	MTORC1 signaling as a biomarker in major depressive disorder and its pharmacological modulation by novel rapid-acting antidepressants. <i>Therapeutic Advances in Psychopharmacology</i> , 2021 , 11, 20451253211036814	4.9	0
36	Previous disorders and depression outcomes in individuals with 12-month major depressive disorder in the World Mental Health surveys. <i>Epidemiology and Psychiatric Sciences</i> , 2021 , 30, e70	5.1	2
35	Evaluating the perceived utility of an artificial intelligence-powered clinical decision support system for depression treatment using a simulation center.. <i>Psychiatry Research</i> , 2021 , 308, 114336	9.9	1
34	Process and Outcome Evaluations of Smartphone Apps for Bipolar Disorder: Scoping Review.. <i>Journal of Medical Internet Research</i> , 2022 , 24, e29114	7.6	0
33	Assessing Depressive Symptoms in Patients With Cancer Treated With Interleukin-2: A Comparison of 2 Measures.. <i>Cancer Nursing</i> , 2022 ,	2.6	
32	Routinely accessible parameters of mineralocorticoid receptor function, depression subtypes and response prediction: a post-hoc analysis from the Early Medication Change trial in major depressive disorder.. <i>World Journal of Biological Psychiatry</i> , 2022 , 1-17	3.8	
31	Pediatric Sleep Disturbances and Symptoms of Depression in Older Adults.. <i>International Journal of Aging and Human Development</i> , 2022 , 914150221077950	1.8	0
30	Remote Assessment of Disease and Relapse in Major Depressive Disorder (RADAR-MDD): recruitment, retention, and data availability in a longitudinal remote measurement study.. <i>BMC Psychiatry</i> , 2022 , 22, 136	4.2	2
29	Allostatic load as a predictor of response to repetitive transcranial magnetic stimulation in treatment resistant depression: Research protocol and hypotheses. <i>Comprehensive Psychoneuroendocrinology</i> , 2022 , 10, 100133	1.1	
28	Suicidal ideation in remitted major depressive disorder predicts recurrence.. <i>Journal of Psychiatric Research</i> , 2022 , 151, 65-72	5.2	0

27 Table_1.docx. 2019,

26 The association of mindfulness and depression stigma among African American women participants in a mindfulness-based intervention: A pilot study.. *Transcultural Psychiatry*, 2022, 13634615221076709 ²⁻⁷

25 Association Between Self-rating Depression Scores and Total Ghrelin and Adipokine Serum Levels in a Large Population-Based Sample. *Frontiers in Psychiatry*, 2022, 13, 5

24 Predictors of engagement with remote sensing technologies for symptom measurement in Major Depressive Disorder.. *Journal of Affective Disorders*, 2022, 310, 106-115 6.6 1

23 The impact of a history of child abuse on cognitive performance: a cross-sectional study in older patients with a depressive, anxiety, or somatic symptom disorder.. *BMC Geriatrics*, 2022, 22, 377 4.1

22 Protocol for Rhapsody: a longitudinal observational study examining the feasibility of speech phenotyping for remote assessment of neurodegenerative and psychiatric disorders. *BMJ Open*, 2022, 12, e061193 3 0

21 Montgomery-Åberg Depression Rating Scale factors in treatment-resistant depression at onset of treatment: Derivation, replication, and change over time during treatment with esketamine. *International Journal of Methods in Psychiatric Research*, 4.3 0

20 The Application of Exposure Principles to the Treatment of Depression. 2022, 317-345 0

19 Central monitoring of depression and anxiety symptoms reduces placebo responses in depression clinical trials: A post hoc exploratory analysis of data from the phase III CCT-004 trial of vortioxetine. 0

18 Associations of negative affective biases and depressive symptoms in a community-based sample. 1-10 0

17 Study protocol for evaluating the clinical efficacy and neurobiological correlates of sequential treatment with tDCS primed iTBS and ECT in treatment-resistant depression. 7, 242 0

16 Cognitive profiles in bipolar I disorder and associated risk factors: Using Wechsler adult intelligence scaleIV. 13, 0

15 The benefits of physical exercise on state anxiety: Exploring possible mechanisms. 2022, 100478 0

14 Delayed affective recovery to daily-life stressors signals a risk for depression. 2022, 0

13 Components of Behavioral Activation Therapy for Depression Engage Specific Reinforcement Learning Mechanisms in a Pilot Study. 2022, 6, 238 0

12 Gamification improves antidepressant effects of cognitive control training: A pilot trial. 4, 0

11 Detecting Impending Symptom Transitions Using Early-Warning Signals in Individuals Receiving Treatment for Depression. 216770262211370 1

10 Comparative Effectiveness of a Mindfulness-Based Intervention (M-Body) on Depressive Symptoms: Study Protocol of a Randomized Controlled Trial in a Federally Qualified Health Center (FQHC). 0

- 9 The association between persistent cognitive difficulties and depression and functional outcomes in people with major depressive disorder. 1-11
- 8 Alternative metrics for characterizing longer-term clinical outcomes in difficult-to-treat depression: I. Association with change in quality of life. 1-13
- 7 Efficacy of celecoxib add-on treatment for immuno-metabolic depression: Protocol of the INFLAMED double-blind placebo-controlled randomized controlled trial. **2023**, 27, 100585
- 6 A feasibility trial of conjoint magnetic seizure therapy and dialectical behavior therapy for suicidal patients with borderline personality disorder and treatment-resistant depression. **2023**, 1, 45-54
- 5 Meta-analysis of the effects of adjuvant drugs in co-occurring bipolar and substance use disorder. **2023**,
- 4 Comparative effectiveness of a mindfulness-based intervention (M-Body) on depressive symptoms: study protocol of a randomized controlled trial in a Federally Qualified Health Center (FQHC). **2023**, 24,
- 3 Multidimensional variability in ecological assessments predicts two clusters of suicidal patients. **2023**, 13,
- 2 Deep phenotyping towards precision psychiatry of first-episode depression The Brain Drugs-Depression cohort. **2023**, 23,
- 1 A critical evaluation of QIDS-SR-16 using data from a trial of psilocybin therapy versus escitalopram treatment for depression. 026988112311678