

# CITATION REPORT

List of articles citing

Partially unfolded states of beta(2)-microglobulin and amyloid formation in vitro

DOI: 10.1021/bi000276j

Biochemistry, 2000, 39, 8735-46.

**Source:** <https://exaly.com/paper-pdf/31577498/citation-report.pdf>

**Version:** 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
3 <sup>16</sup>	Kidney dialysis-associated amyloidosis: a molecular role for copper in fiber formation. <b>2001</b> , 309, 339-45		151
3 <sup>15</sup>	Beta(2)-microglobulin and its deamidated variant, N17D form amyloid fibrils with a range of morphologies in vitro. <b>2001</b> , 313, 559-71		180
3 <sup>14</sup>	Detection of two partially structured species in the folding process of the amyloidogenic protein beta 2-microglobulin. <b>2001</b> , 307, 379-91		110
3 <sup>13</sup>	Apolipoprotein E inhibits the depolymerization of beta 2-microglobulin-related amyloid fibrils at a neutral pH. <i>Biochemistry</i> , <b>2001</b> , 40, 8499-507	3.2	66
3 <sup>12</sup>	Crystal structures of the yeast prion Ure2p functional region in complex with glutathione and related compounds. <i>Biochemistry</i> , <b>2001</b> , 40, 13564-73	3.2	61
3 <sup>11</sup>	Anion shielding of electrostatic repulsions in transthyretin modulates stability and amyloidosis: insight into the chaotrope unfolding dichotomy. <i>Biochemistry</i> , <b>2001</b> , 40, 11453-9	3.2	75
3 <sup>10</sup>	Role of the single disulphide bond of beta(2)-microglobulin in amyloidosis in vitro. <b>2001</b> , 10, 1775-84		69
3 <sup>09</sup>	Protein aggregation. <b>2001</b> , 39, 1065-75		23
3 <sup>08</sup>	A partially structured species of beta 2-microglobulin is significantly populated under physiological conditions and involved in fibrillogenesis. <b>2001</b> , 276, 46714-21		124
3 <sup>07</sup>	Conformational intermediate of the amyloidogenic protein beta 2-microglobulin at neutral pH. <b>2001</b> , 276, 32657-62		44
3 <sup>06</sup>	Conformational Abs recognizing a generic amyloid fibril epitope. <b>2002</b> , 99, 1485-90		296
3 <sup>05</sup>	Sequence-dependent denaturation energetics: A major determinant in amyloid disease diversity. <b>2002</b> , 99 Suppl 4, 16427-32		283
3 <sup>04</sup>	Investigation of a peptide responsible for amyloid fibril formation of beta 2-microglobulin by achromobacter protease I. <b>2002</b> , 277, 1310-5		109
3 <sup>03</sup>	Studies of the aggregation of mutant proteins in vitro provide insights into the genetics of amyloid diseases. <b>2002</b> , 99 Suppl 4, 16419-26		246
3 <sup>02</sup>	Elucidation of the molecular mechanism during the early events in immunoglobulin light chain amyloid fibrillation. Evidence for an off-pathway oligomer at acidic pH. <b>2002</b> , 277, 12666-79		84
3 <sup>01</sup>	A designed system for assessing how sequence affects alpha to beta conformational transitions in proteins. <b>2002</b> , 277, 10150-5		79
3 <sup>00</sup>	Crystal structure of monomeric human beta-2-microglobulin reveals clues to its amyloidogenic properties. <b>2002</b> , 99, 9771-6		166

299	Conformation of beta 2-microglobulin amyloid fibrils analyzed by reduction of the disulfide bond. <b>2002</b> , 277, 21554-60		58
298	Cleaved beta 2-microglobulin partially attains a conformation that has amyloidogenic features. <b>2002</b> , 277, 11184-9		40
297	Formation of a copper specific binding site in non-native states of beta-2-microglobulin. <i>Biochemistry</i> , <b>2002</b> , 41, 10646-56	3.2	95
296	Amyloid fibril formation of the mouse V(L) domain at acidic pH. <i>Biochemistry</i> , <b>2002</b> , 41, 3389-95	3.2	22
295	Insights into the origin of the tendency of the PI3-SH3 domain to form amyloid fibrils. <b>2002</b> , 322, 1147-58		38
294	Acid Denaturation of alpha1-antitrypsin: characterization of a novel mechanism of serpin polymerization. <b>2002</b> , 324, 859-70		72
293	Capillary electrophoresis investigation of a partially unfolded conformation of beta(2)-microglobulin. <b>2002</b> , 23, 918-25		51
292	The structural basis for amyloid formation by plasma apolipoproteins: a review. <b>2002</b> , 31, 2-8		76
291	Towards an understanding of amyloidogenesis. <b>2002</b> , 9, 323-5		73
290	Picture story. Locked but not loaded. <b>2002</b> , 9, 325		
289	Structural properties of an amyloid precursor of beta(2)-microglobulin. <b>2002</b> , 9, 326-31		167
288	Mapping the core of the beta(2)-microglobulin amyloid fibril by H/D exchange. <b>2002</b> , 9, 332-6		290
287	Thermal denaturation of Bungarus fasciatus acetylcholinesterase: Is aggregation a driving force in protein unfolding?. <b>2002</b> , 11, 2022-32		16
286	The yeast prion Ure2p retains its native alpha-helical conformation upon assembly into protein fibrils in vitro. <b>2002</b> , 21, 2903-11		118
285	AFM and STM study of beta-amyloid aggregation on graphite. <b>2003</b> , 97, 73-9		65
284	Effect of pH on the aggregate formation of a non-amyloid component (1-13). <b>2003</b> , 9, 177-86		8
283	Amyloidogenesis: historical and modern observations point to heparan sulfate proteoglycans as a major culprit. <b>2003</b> , 10, 67-79		128
282	Mutations in the B1 domain of protein G that delay the onset of amyloid fibril formation in vitro. <b>2003</b> , 12, 567-76		13

281	Dissolution of beta2-microglobulin amyloid fibrils by dimethylsulfoxide. <b>2003</b> , 134, 159-64		89
280	Role of the C-terminal 28 residues of beta2-microglobulin in amyloid fibril formation. <i>Biochemistry</i> , <b>2003</b> , 42, 13536-40	3.2	32
279	An amyloid-like C-terminal domain of thrombospondin-1 displays CD47 agonist activity requiring both VVM motifs. <i>Biochemistry</i> , <b>2003</b> , 42, 10001-11	3.2	28
278	The intact human acetylcholinesterase C-terminal oligomerization domain is alpha-helical in situ and in isolation, but a shorter fragment forms beta-sheet-rich amyloid fibrils and protofibrillar oligomers. <i>Biochemistry</i> , <b>2003</b> , 42, 10863-73	3.2	22
277	pH-dependent amyloid and protofibril formation by the ABri peptide of familial British dementia. <b>2003</b> , 333, 1003-23		80
276	Amyloid-forming peptides from beta2-microglobulin-Insights into the mechanism of fibril formation in vitro. <b>2003</b> , 325, 249-57		140
275	Hierarchical assembly of beta2-microglobulin amyloid in vitro revealed by atomic force microscopy. <b>2003</b> , 330, 785-97		206
274	A systematic investigation into the effect of protein destabilisation on beta 2-microglobulin amyloid formation. <b>2003</b> , 330, 943-54		128
273	Role of the N and C-terminal strands of beta 2-microglobulin in amyloid formation at neutral pH. <b>2003</b> , 330, 935-41		68
272	Effects of salt concentration on association of the amyloid protofilaments of hen egg white lysozyme studied by time-resolved neutron scattering. <b>2003</b> , 331, 21-8		41
271	A general model for amyloid fibril assembly based on morphological studies using atomic force microscopy. <b>2003</b> , 85, 1135-44		263
270	The native-like conformation of Ure2p in fibrils assembled under physiologically relevant conditions switches to an amyloid-like conformation upon heat-treatment of the fibrils. <b>2003</b> , 141, 132-42		65
269	Assembly of the yeast prion Ure2p into protein fibrils. Thermodynamic and kinetic characterization. <b>2003</b> , 278, 30199-205		31
268	Amyloid-like fibril formation in an all beta-barrel protein. Partially structured intermediate state(s) is a precursor for fibril formation. <b>2003</b> , 278, 17701-9		51
267	An amyloid-forming segment of beta2-microglobulin suggests a molecular model for the fibril. <b>2004</b> , 101, 10584-9		209
266	Properties of some variants of human beta2-microglobulin and amyloidogenesis. <b>2004</b> , 279, 9176-89		59
265	Co-populated conformational ensembles of beta2-microglobulin uncovered quantitatively by electrospray ionization mass spectrometry. <b>2004</b> , 279, 27069-77		63
264	Amyloid fibril formation by lens crystallin proteins and its implications for cataract formation. <b>2004</b> , 279, 3413-9		140

263	A mechanical unfolding intermediate in an actin-crosslinking protein. <b>2004</b> , 11, 81-5		179
262	Differences in aggregation properties of three site-specific mutants of recombinant human stefin B. <b>2004</b> , 13, 63-70		17
261	Increase in the conformational flexibility of beta 2-microglobulin upon copper binding: a possible role for copper in dialysis-related amyloidosis. <b>2004</b> , 13, 797-809		78
260	Amyloid fibrils from the viewpoint of protein folding. <b>2004</b> , 61, 511-524		119
259	Separation of beta2-microglobulin conformers by high-field asymmetric waveform ion mobility spectrometry (FAIMS) coupled to electrospray ionisation mass spectrometry. <b>2004</b> , 18, 2229-34		50
258	Conformational constraints for amyloid fibrillation: the importance of being unfolded. <b>2004</b> , 1698, 131-53		817
257	Structural characterization of the fibrillar form of the yeast <i>Saccharomyces cerevisiae</i> prion Ure2p. <i>Biochemistry</i> , <b>2004</b> , 43, 5022-32	3.2	52
256	beta2-microglobulin H31Y variant 3D structure highlights the protein natural propensity towards intermolecular aggregation. <b>2004</b> , 335, 1051-64		35
255	Cystatin forms a tetramer through structural rearrangement of domain-swapped dimers prior to amyloidogenesis. <b>2004</b> , 336, 165-78		74
254	Core and heterogeneity of beta2-microglobulin amyloid fibrils as revealed by H/D exchange. <b>2004</b> , 338, 559-71		94
253	Phospholipid catalysis of diabetic amyloid assembly. <b>2004</b> , 341, 1175-87		290
252	Amyloid fibril formation by a partially structured intermediate state of alpha-chymotrypsin. <b>2004</b> , 342, 321-31		176
251	Specific Binding of .BETA.2-Microglobulin with Trypan Blue. <b>2005</b> , 51, 702-707		1
250	Towards an understanding of the structural molecular mechanism of beta(2)-microglobulin amyloid formation in vitro. <b>2005</b> , 1753, 51-63		47
249	The three-dimensional structure of beta2 microglobulin: results from X-ray crystallography. <b>2005</b> , 1753, 85-91		12
248	Structural studies reveal that the diverse morphology of beta(2)-microglobulin aggregates is a reflection of different molecular architectures. <b>2005</b> , 1753, 108-20		36
247	Interactions of charged ligands with beta(2)-microglobulin conformers in affinity capillary electrophoresis. <b>2005</b> , 1753, 131-40		16
246	Proteomics of beta2-microglobulin amyloid fibrils. <b>2005</b> , 1753, 23-33		34

245	Structural stability of amyloid fibrils of beta(2)-microglobulin in comparison with its native fold. <b>2005</b> , 1753, 64-75		30
244	From chance to frequent encounters: origins of beta2-microglobulin fibrillogenesis. <b>2005</b> , 1753, 92-9		36
243	Studies on unfolded beta-microglobulin at C-terminal in dialysis-related amyloidosis. <b>2005</b> , 67, 314-20		9
242	The importance of sequence diversity in the aggregation and evolution of proteins. <b>2005</b> , 438, 878-81		263
241	Search of ligands for the amyloidogenic protein beta2-microglobulin by capillary electrophoresis and other techniques. <b>2005</b> , 26, 4055-63		15
240	Amyloid fibril formation by the CAD domain of caspase-activated DNase. <b>2005</b> , 79, 39-47		6
239	Prediction of "hot spots" of aggregation in disease-linked polypeptides. <b>2005</b> , 5, 18		148
238	Amyloid fibril formation by bovine cytochrome c. <b>2005</b> , 19, 199-205		27
237	Lipid Modulators of Protein Misfolding and Aggregation. 111-129		
236	β2-Microglobulin. 667-695		5
235	Lysozyme. 635-656		5
234	Seeding-dependent maturation of beta2-microglobulin amyloid fibrils at neutral pH. <b>2005</b> , 280, 12012-8		60
233	Structure of the prion Ure2p in protein fibrils assembled in vitro. <b>2005</b> , 280, 37149-58		26
232	Stereospecific amyloid-like fibril formation by a peptide fragment of beta2-microglobulin. <i>Biochemistry</i> , <b>2005</b> , 44, 157-64	3.2	58
231	Characterization of two distinct beta2-microglobulin unfolding intermediates that may lead to amyloid fibrils of different morphology. <i>Biochemistry</i> , <b>2005</b> , 44, 16098-107	3.2	36
230	Native state kinetic stabilization as a strategy to ameliorate protein misfolding diseases: a focus on the transthyretin amyloidoses. <b>2005</b> , 38, 911-21		231
229	Unfolding, aggregation, and seeded amyloid formation of lysine-58-cleaved beta 2-microglobulin. <i>Biochemistry</i> , <b>2005</b> , 44, 4397-407	3.2	55
228	Dynamics in the unfolded state of beta2-microglobulin studied by NMR. <b>2005</b> , 346, 279-94		88

227	Amyloid-like properties of bacterial inclusion bodies. <b>2005</b> , 347, 1025-37		191
226	A toy model for predicting the rate of amyloid formation from unfolded protein. <b>2005</b> , 351, 195-205		38
225	Competing pathways determine fibril morphology in the self-assembly of beta2-microglobulin into amyloid. <b>2005</b> , 351, 850-64		299
224	A single disulfide bond differentiates aggregation pathways of beta2-microglobulin. <b>2005</b> , 354, 473-82		47
223	Sequence determinants of protein aggregation: tools to increase protein solubility. <b>2005</b> , 4, 11		75
222	Amyloid formation of a protein in the absence of initial unfolding and destabilization of the native state. <b>2005</b> , 89, 4234-44		60
221	Structural and Conformational Prerequisites of Amyloidogenesis. <b>2006</b> , 1-20		4
220	Relative Importance of Hydrophobicity, Net Charge, and Secondary Structure Propensities in Protein Aggregation. <b>2006</b> , 43-59		13
219	A statistical-mechanical theory of fibril formation in dilute protein solutions. <b>2006</b> , 90, 3134-45		28
218	A single mutation induces amyloid aggregation in the alpha-spectrin SH3 domain: analysis of the early stages of fibril formation. <b>2006</b> , 356, 453-68		47
217	A molecular dynamics approach to the structural characterization of amyloid aggregation. <b>2006</b> , 357, 1306-21		82
216	Direct observation of oligomeric species formed in the early stages of amyloid fibril formation using electrospray ionisation mass spectrometry. <b>2006</b> , 364, 9-19		130
215	Investigating the structural properties of amyloid-like fibrils formed in vitro from beta2-microglobulin using limited proteolysis and electrospray ionisation mass spectrometry. <b>2006</b> , 20, 1628-36		55
214	A native to amyloidogenic transition regulated by a backbone trigger. <b>2006</b> , 13, 202-8		177
213	Molecular chaperones and the assembly of the prion Sup35p, an in vitro study. <b>2006</b> , 25, 822-33		96
212	A systematic study of the effect of physiological factors on beta2-microglobulin amyloid formation at neutral pH. <i>Biochemistry</i> , <b>2006</b> , 45, 2311-21	3,2	111
211	Simulation of pH-dependent edge strand rearrangement in human beta-2 microglobulin. <b>2006</b> , 15, 200-7		18
210	Insights into the architecture of the Ure2p yeast protein assemblies from helical twisted fibrils. <b>2006</b> , 15, 2481-7		16

209	Characterization of the native and fibrillar conformation of the human Nalpha-acetyltransferase ARD1. <b>2006</b> , 15, 1968-76	16
208	Amyloid fibril formation by a domain of rat cell adhesion molecule. <b>2006</b> , 44, 241-9	6
207	Motional properties of unfolded ubiquitin: a model for a random coil protein. <b>2006</b> , 35, 175-86	49
206	Micro-heterogeneity and aggregation in beta2-microglobulin solutions: effects of temperature, pH, and conformational variant addition. <b>2006</b> , 35, 439-45	26
205	Collagen plays an active role in the aggregation of beta2-microglobulin under physiopathological conditions of dialysis-related amyloidosis. <b>2006</b> , 281, 16521-9	122
204	A systematic screen of beta(2)-microglobulin and insulin for amyloid-like segments. <b>2006</b> , 103, 4079-82	115
203	Aggregation of natively folded proteins: a theoretical approach. <b>2007</b> , 19, 285221	4
202	Protein Misfolding, Aggregation, and Conformational Diseases. <b>2007</b> ,	13
201	Investigation into the role of macrophages in the formation and degradation of beta2-microglobulin amyloid fibrils. <b>2007</b> , 282, 29691-700	27
200	Biochemical and functional analysis of the assembly of full-length Sup35p and its prion-forming domain. <b>2007</b> , 282, 1679-86	45
199	The mechanism of the amyloidogenic conversion of T7 endonuclease I. <b>2007</b> , 282, 14968-74	7
198	Dimethylsulfoxide-quenched hydrogen/deuterium exchange method to study amyloid fibril structure. <b>2007</b> , 1768, 1886-99	42
197	Hydrogen/deuterium exchange mass spectrometric analysis of conformational changes accompanying the assembly of the yeast prion Ure2p into protein fibrils. <b>2007</b> , 369, 1113-25	20
196	Production and characterization of RNA aptamers specific for amyloid fibril epitopes. <b>2007</b> , 282, 34500-9	37
195	Visualization and classification of amyloid beta supramolecular assemblies. <i>Biochemistry</i> , <b>2007</b> , 46, 15009-17	78
194	Structure and aggregation mechanism of beta(2)-microglobulin (83-99) peptides studied by molecular dynamics simulations. <b>2007</b> , 93, 3353-62	17
193	Formation of structured polymers upon controlled denaturation of beta-lactoglobulin with different chaotropes. <b>2007</b> , 86, 57-72	33
192	Peptide fibrillization. <b>2007</b> , 46, 8128-47	524



191	Bildung von Peptidfibrillen. <b>2007</b> , 119, 8274-8295	64
190	Specific glycosaminoglycans promote unseeded amyloid formation from beta2-microglobulin under physiological conditions. <b>2007</b> , 72, 174-81	48
189	A simple algorithm locates beta-strands in the amyloid fibril core of alpha-synuclein, Abeta, and tau using the amino acid sequence alone. <b>2007</b> , 16, 906-18	90
188	Monitoring copopulated conformational states during protein folding events using electrospray ionization-ion mobility spectrometry-mass spectrometry. <b>2007</b> , 18, 2180-90	114
187	Sulfonated molecules that bind a partially structured species of beta2-microglobulin also influence refolding and fibrillogenesis. <b>2008</b> , 29, 1502-10	16
186	Cu(II) organizes beta-2-microglobulin oligomers but is released upon amyloid formation. <b>2008</b> , 17, 748-59	35
185	Conformational characterization of oligomeric intermediates and aggregates in beta-lactoglobulin heat aggregation. <b>2001</b> , 10, 1312-8	109
184	Identification and functional characterization of an aggregation domain in long myosin light chain kinase. <b>2008</b> , 275, 2489-500	5
183	A regulatable switch mediates self-association in an immunoglobulin fold. <b>2008</b> , 15, 965-71	80
182	Protein folding and misfolding on surfaces. <b>2008</b> , 9, 2515-42	57
181	PrP(106-126) does not interact with membranes under physiological conditions. <b>2008</b> , 95, 1877-89	70
180	The in vivo and in vitro aggregation properties of globular proteins correlate with their conformational stability: the SH3 case. <b>2008</b> , 378, 1116-31	52
179	Folding versus aggregation: polypeptide conformations on competing pathways. <b>2008</b> , 469, 100-17	305
178	Interactive sequences in the molecular chaperone, human alphaB crystallin modulate the fibrillation of amyloidogenic proteins. <b>2008</b> , 40, 954-67	32
177	Early stages of misfolding and association of beta2-microglobulin: insights from infrared spectroscopy and dynamic light scattering. <i>Biochemistry</i> , <b>2008</b> , 47, 6895-906	3-2 41
176	A common beta-sheet architecture underlies in vitro and in vivo beta2-microglobulin amyloid fibrils. <b>2008</b> , 283, 17279-86	48
175	Amyloid fibrils: abnormal protein assembly. <b>2008</b> , 2, 112-7	295
174	Molecular chaperones and the assembly of the prion Ure2p in vitro. <b>2008</b> , 283, 15732-9	30

173	The structure of a folding intermediate provides insight into differences in immunoglobulin amyloidogenicity. <b>2008</b> , 105, 13373-8		34
172	Structure, formation and propagation of amyloid fibrils. <b>2008</b> , 14, 3205-18		29
171	Molecular dynamics simulations to investigate the relationship between the structural stability and amyloidogenesis of the wild-type and N-terminal hexapeptide deletion $\Delta$ 6 $\beta$ -microglobulin. <b>2009</b> , 35, 755-765		3
170	Kinetic analysis of amyloid formation in the presence of heparan sulfate: faster unfolding and change of pathway. <b>2009</b> , 284, 29921-34		51
169	A comprehensive model for packing and hydration for amyloid fibrils of beta2-microglobulin. <b>2009</b> , 284, 2169-75		49
168	A circulating beta 2-microglobulin intermediate in hemodialysis patients. <b>2009</b> , 111, c173-81		12
167	Metal binding sheds light on mechanisms of amyloid assembly. <b>2009</b> , 3, 1-4		35
166	Single-molecule mechanical unfolding of amyloidogenic beta2-microglobulin: the force-spectroscopy approach. <b>2009</b> , 10, 1471-7		5
165	Disorder-to-order conformational transitions in protein structure and its relationship to disease. <b>2009</b> , 330, 105-20		26
164	Tuning the conformational properties of the prion peptide. <b>2009</b> , 76, 213-25		10
163	Amyloid formation by globular proteins under native conditions. <b>2009</b> , 5, 15-22		671
162	The solution structure of human beta2-microglobulin reveals the prodromes of its amyloid transition. <b>2002</b> , 11, 487-99		132
161	Mechanism of lysophosphatidic acid-induced amyloid fibril formation of beta(2)-microglobulin in vitro under physiological conditions. <i>Biochemistry</i> , <b>2009</b> , 48, 5689-99	3.2	27
160	Copper binding to beta-2-microglobulin and its pre-amyloid oligomers. <i>Biochemistry</i> , <b>2009</b> , 48, 9871-81	3.2	42
159	Delineating the conformational elements responsible for Cu(2+)-induced oligomerization of beta-2 microglobulin. <i>Biochemistry</i> , <b>2009</b> , 48, 6610-7	3.2	16
158	A beta2-microglobulin cleavage variant fibrillates at near-physiological pH. <b>2009</b> , 381, 187-91		9
157	A generic mechanism of beta2-microglobulin amyloid assembly at neutral pH involving a specific proline switch. <b>2009</b> , 386, 1312-26		73
156	Agitation and high ionic strength induce amyloidogenesis of a folded PDZ domain in native conditions. <b>2009</b> , 96, 2289-98		30

155	Synthetic lipid vesicles recruit native-like aggregates and affect the aggregation process of the prion Ure2p: insights on vesicle permeabilization and charge selectivity. <b>2009</b> , 96, 3319-30		15
154	Topological investigation of amyloid fibrils obtained from beta2-microglobulin. <b>2002</b> , 11, 2362-9		49
153	The role of disulfide bond in the amyloidogenic state of beta(2)-microglobulin studied by heteronuclear NMR. <b>2002</b> , 11, 2218-29		80
152	A kinetic study of beta-lactoglobulin amyloid fibril formation promoted by urea. <b>2002</b> , 11, 2417-26		208
151	The toxicity of prion protein fragment PrP(106-126) is not mediated by membrane permeabilization as shown by a M112W substitution. <i>Biochemistry</i> , <b>2009</b> , 48, 4198-208	3.2	30
150	beta(2)-microglobulin: from physiology to amyloidosis. <b>2009</b> , 16, 151-73		46
149	Structural integrity of beta-sheet assembly. <b>2009</b> , 37, 671-6		39
148	Tertiary butanol induced amyloidogenesis of hen egg white lysozyme (HEWL) is facilitated by aggregation-prone alkali-induced molten globule like conformational state. <b>2009</b> , 16, 56-60		16
147	The Sheet Structure pH Dependence of the Core Fragments of $\beta$ -Microglobulin Amyloid Fibrils. <b>2010</b> , 83, 495-504		11
146	Effects of a lipid environment on the fibrillogenic pathway of the N-terminal polypeptide of human apolipoprotein A-I, responsible for in vivo amyloid fibril formation. <b>2010</b> , 39, 1289-99		21
145	Binding mode of Thioflavin T and other molecular probes in the context of amyloid fibrils-current status. <b>2010</b> , 3, 1-18		438
144	Why Proteins Misfold. <b>2010</b> , 1-20		
143	Hemodialysis-Related Amyloidosis. <b>2010</b> , 347-379		2
142	Genetically Engineered Polypeptides as a Model of Intrinsically Disordered Fibrillogenic Proteins: Deep UV Resonance Raman Spectroscopic Study. <b>2010</b> , 253-302		2
141	Fast membrane association is a crucial factor in the peptide pep-1 translocation mechanism: a kinetic study followed by surface plasmon resonance. <b>2010</b> , 94, 314-22		24
140	A de novo designed 11 kDa polypeptide: model for amyloidogenic intrinsically disordered proteins. <b>2010</b> , 93, 607-18		8
139	Structure and assembly properties of the N-terminal domain of the prion Ure2p in isolation and in its natural context. <i>PLoS ONE</i> , <b>2010</b> , 5, e9760	3.7	13
138	Stacked sets of parallel, in-register beta-strands of beta2-microglobulin in amyloid fibrils revealed by site-directed spin labeling and chemical labeling. <b>2010</b> , 285, 17137-47		54

137	TAB2 scaffolds TAK1 and NLK in repressing canonical Wnt signaling. <b>2010</b> , 285, 13397-404		25
136	Molten globule precursor states are conformationally correlated to amyloid fibrils of human beta-2-microglobulin. <b>2010</b> , 132, 9223-5		40
135	Magic angle spinning NMR analysis of beta2-microglobulin amyloid fibrils in two distinct morphologies. <b>2010</b> , 132, 10414-23		73
134	[Membrane roughness: A relevant concept in haemodialysis]. <b>2010</b> , 6, 158-61		1
133	Structure of the preamyloid dimer of beta-2-microglobulin from covalent labeling and mass spectrometry. <i>Biochemistry</i> , <b>2010</b> , 49, 1522-32	3.2	55
132	Differences in the molecular structure of beta(2)-microglobulin between two morphologically different amyloid fibrils. <i>Biochemistry</i> , <b>2010</b> , 49, 742-51	3.2	20
131	Structural insights into the pre-amyloid tetramer of $\beta$ 2-microglobulin from covalent labeling and mass spectrometry. <i>Biochemistry</i> , <b>2011</b> , 50, 6711-22	3.2	41
130	Structure and dynamics of oligomeric intermediates in $\beta$ 2-microglobulin self-assembly. <b>2011</b> , 101, 1238-47		23
129	Cyclodextrin, a novel therapeutic tool for suppressing amyloidogenic transthyretin misfolding in transthyretin-related amyloidosis. <b>2011</b> , 437, 35-42		22
128	Reversible heat-induced dissociation of $\beta$ 2-microglobulin amyloid fibrils. <i>Biochemistry</i> , <b>2011</b> , 50, 3211-20	3.2	49
127	$\beta$ 2-microglobulin forms three-dimensional domain-swapped amyloid fibrils with disulfide linkages. <b>2011</b> , 18, 49-55		93
126	Large proteins have a great tendency to aggregate but a low propensity to form amyloid fibrils. <i>PLoS ONE</i> , <b>2011</b> , 6, e16075	3.7	41
125	Qualitative and quantitative multiplexed proteomic analysis of complex yeast protein fractions that modulate the assembly of the yeast prion Sup35p. <i>PLoS ONE</i> , <b>2011</b> , 6, e23659	3.7	2
124	An anti-A $\beta$ amyloid $\beta$ single-chain variable fragment prevents amyloid fibril formation and cytotoxicity by withdrawing A $\beta$ oligomers from the amyloid pathway. <b>2011</b> , 437, 25-34		32
123	A mutation within the C-terminal domain of Sup35p that affects [PSI <sup>+</sup> ] prion propagation. <b>2011</b> , 81, 640-58		27
122	Amyloid structure: conformational diversity and consequences. <b>2011</b> , 80, 557-85		205
121	$\beta$ synuclein propagates from mouse brain to grafted dopaminergic neurons and seeds aggregation in cultured human cells. <b>2011</b> , 121, 715-25		616
120	Characterization of $\beta$ 2-microglobulin conformational intermediates associated to different fibrillation conditions. <b>2011</b> , 46, 734-41		12

119	Removal of intact $\beta$ -microglobulin at neutral pH by using seed-conjugated polymer beads prepared with $\beta$ -microglobulin-derived peptide (58-67). <b>2011</b> , 27, 521-9	7
118	Expression in <i>E. coli</i> and purification of the fibrillogenic fusion proteins TTR-sfGFP and $\beta$ M-sfGFP. <b>2011</b> , 41, 337-49	11
117	Binding with nucleic acids or glycosaminoglycans converts soluble protein oligomers to amyloid. <b>2012</b> , 287, 736-747	43
116	Structural Elements Regulating Interactions in the Early Stages of Fibrillogenesis: A Human Calcitonin Model System. <b>2012</b> , 351-388	1
115	Kinetic intermediates of amyloid fibrillation studied by hydrogen exchange methods with nuclear magnetic resonance. <b>2012</b> , 1824, 1307-23	18
114	Polymorphism of $\beta$ -microglobulin amyloid fibrils manifested by ultrasonication-enhanced fibril formation in trifluoroethanol. <b>2012</b> , 287, 22827-37	36
113	Inhibiting, promoting, and preserving stability of functional protein fibrils. <b>2012</b> , 8, 876-895	112
112	Distinguishing crystal-like amyloid fibrils and glass-like amorphous aggregates from their kinetics of formation. <b>2012</b> , 109, 14446-51	200
111	The transthyretin amyloidoses: from delineating the molecular mechanism of aggregation linked to pathology to a regulatory-agency-approved drug. <b>2012</b> , 421, 185-203	227
110	Protein aggregation: mechanisms and functional consequences. <b>2012</b> , 44, 1541-54	111
109	The small heat-shock protein $\beta$ -crystallin uses different mechanisms of chaperone action to prevent the amorphous versus fibrillar aggregation of $\beta$ lactalbumin. <b>2012</b> , 448, 343-52	42
108	$\beta$ -microglobulin amyloidosis. <b>2012</b> , 65, 517-40	11
107	Structural features and cytotoxicity of amyloid oligomers: implications in Alzheimer's disease and other diseases with amyloid deposits. <b>2012</b> , 99, 226-45	138
106	Pathological self-aggregation of $\beta$ -microglobulin: a challenge for protein biophysics. <b>2012</b> , 65, 165-83	6
105	Mapping local structural perturbations in the native state of stefin B (cystatin B) under amyloid forming conditions. <b>2012</b> , 5, 94	6
104	Photoinduced fibrils formation of chicken egg white lysozyme under native conditions. <b>2012</b> , 80, 2501-13	27
103	The role of conformational flexibility in $\beta$ -microglobulin amyloid fibril formation at neutral pH. <b>2012</b> , 26, 1783-92	22
102	Protein aggregation and misfolding: good or evil?. <b>2012</b> , 24, 244101	13

101	Effect of heparin on conformation of the $\beta_2$ -microglobulin molecule. <b>2012</b> , 16, 159-62		2
100	Capillary electrophoretic profile of $\beta_2$ -microglobulin intermediate associated with hemodialysis. <b>2012</b> , 16, 350-4		2
99	Exploring critical determinants of protein amyloidogenesis: a review. <b>2013</b> , 19, 529-36		14
98	Amyloid fibril formation by a normally folded protein in the absence of denaturants and agitation. <b>2013</b> , 20, 226-32		5
97	Protein folding and aggregation into amyloid: the interference by natural phenolic compounds. <b>2013</b> , 14, 12411-57		132
96	Isolation, characterization, and aggregation of a structured bacterial matrix precursor. <b>2013</b> , 288, 17559-68		42
95	The molten globule of $(\beta_2)$ -microglobulin accumulated at pH 4 and its role in protein folding. <b>2013</b> , 425, 273-91		20
94	Both the cis-trans equilibrium and isomerization dynamics of a single proline amide modulate $\beta_2$ -microglobulin amyloid assembly. <b>2013</b> , 110, 20051-6		52
93	Dynamics and dimension of an amyloidogenic disordered state of human $(\beta_2)$ -microglobulin. <b>2013</b> , 42, 767-76		8
92	Oligomeric states along the folding pathways of $\beta_2$ -microglobulin: kinetics, thermodynamics, and structure. <b>2013</b> , 425, 2722-36		27
91	Amyloid formation in heterogeneous environments: islet amyloid polypeptide glycosaminoglycan interactions. <b>2013</b> , 425, 492-505		25
90	Advances in ion mobility spectrometry-mass spectrometry reveal key insights into amyloid assembly. <b>2013</b> , 1834, 1257-68		70
89	Delicate balance between functionally required flexibility and aggregation risk in a $\beta$ -rich protein. <i>Biochemistry</i> , <b>2013</b> , 52, 8843-54	3,2	18
88	Native-state heterogeneity of $(\beta_2)$ -microglobulin as revealed by kinetic folding and real-time NMR experiments. <b>2013</b> , 425, 257-72		15
87	The amyloid-cell membrane system. The interplay between the biophysical features of oligomers/fibrils and cell membrane defines amyloid toxicity. <b>2013</b> , 182, 30-43		80
86	Heparin promotes the rapid fibrillization of a peptide with low intrinsic amyloidogenicity. <i>Biochemistry</i> , <b>2013</b> , 52, 8984-92	3,2	16
85	Assessing the effect of loop mutations in the folding space of $\beta_2$ -microglobulin with molecular dynamics simulations. <b>2013</b> , 14, 17256-78		13
84	Ultrasonication: An Efficient Agitation for Accelerating the Supersaturation-Limited Amyloid Fibrillation of Proteins. <b>2013</b> , 52, 07HA01		25

83	Structure of an early native-like intermediate of $\beta$ -microglobulin amyloidogenesis. <b>2013</b> , 22, 1349-57		12
82	The extracellular chaperone haptoglobin prevents serum fatty acid-promoted amyloid fibril formation of $\beta$ -microglobulin, resistance to lysosomal degradation, and cytotoxicity. <b>2013</b> , 288, 32326-32342		12
81	Rapid generation of amyloid from native proteins in vitro. <b>2013</b> , 50869		5
80	Calcium binding to beta-2-microglobulin at physiological pH drives the occurrence of conformational changes which cause the protein to precipitate into amorphous forms that subsequently transform into amyloid aggregates. <i>PLoS ONE</i> , <b>2014</b> , 9, e95725	3.7	10
79	$\beta$ -Microglobulin amyloid fibril-induced membrane disruption is enhanced by endosomal lipids and acidic pH. <i>PLoS ONE</i> , <b>2014</b> , 9, e104492	3.7	27
78	A simulated intermediate state for folding and aggregation provides insights into $\beta$ 6 $\beta$ -microglobulin amyloidogenic behavior. <b>2014</b> , 10, e1003606		28
77	Intermediate conformation between native $\beta$ sheet and non-native $\beta$ helix is a precursor of trifluoroethanol-induced aggregation of human carbonic anhydrase-II. <b>2014</b> , 449, 126-31		10
76	Significance of unfolding thermodynamics for predicting aggregation kinetics: a case study on high concentration solutions of a multi-domain protein. <b>2014</b> , 31, 1575-87		21
75	Unique effect of Cu(II) in the metal-induced amyloid formation of $\beta$ 2-microglobulin. <i>Biochemistry</i> , <b>2014</b> , 53, 1263-74	3.2	15
74	Secondary structure in the core of amyloid fibrils formed from human $\beta$ h and its truncated variant $\beta$ 6. <b>2014</b> , 136, 6313-25		37
73	Protonation favors aggregation of lysozyme with SDS. <b>2014</b> , 10, 2591-9		81
72	Dynamical properties of $\beta$ synuclein in soluble and fibrillar forms by Quasi Elastic Neutron Scattering. <b>2014</b> , 1844, 1307-16		3
71	[Structure analysis of disease-related proteins using vibrational spectroscopy]. <b>2014</b> , 134, 1013-20		1
70	Effect of nitric oxide on conformational changes of ovalbumin accompanying self-assembly into non-disease-associated fibrils. <b>2015</b> , 47, 1-9		2
69	C-terminal unfolding of an amyloidogenic $\beta$ -microglobulin fragment: $\beta$ 6 $\beta$ -microglobulin. <b>2015</b> , 22, 54-60		5
68	Substitution of proline32 by $\beta$ methylproline preorganizes $\beta$ -microglobulin for oligomerization but not for aggregation into amyloids. <b>2015</b> , 137, 2524-35		13
67	Regulation of protein homeostasis in neurodegenerative diseases: the role of coding and non-coding genes. <b>2015</b> , 72, 4027-47		24
66	Systemic amyloidosis: lessons from $\beta$ -microglobulin. <b>2015</b> , 290, 9951-8		57

65	Sequence- and Temperature-Dependent Properties of Unfolded and Disordered Proteins from Atomistic Simulations. <b>2015</b> , 119, 14622-30	49
64	Advances in Organic Crystal Chemistry. <b>2015</b> ,	9
63	Ultrasonication-Forced Amyloid Fibrillation of Proteins. <b>2015</b> , 15-29	
62	Alpha-synuclein oligomers and fibrils originate in two distinct conformer pools: a small angle X-ray scattering and ensemble optimisation modelling study. <b>2015</b> , 11, 190-6	22
61	Historical and Current Concepts of Fibrillogenesis and In vivo Amyloidogenesis: Implications of Amyloid Tissue Targeting. <b>2016</b> , 3, 17	15
60	Denaturation induced aggregation in Icrystallin: differential action of chaotropes. <b>2016</b> , 29, 536-543	7
59	Characterization of Salt-Induced Oligomerization of Human I-Microglobulin at Low pH. <b>2016</b> , 120, 7815-23	3
58	Comparison of the aggregation of homologous I-microglobulin variants reveals protein solubility as a key determinant of amyloid formation. <b>2016</b> , 428, 631-643	12
57	Stepwise unfolding of human I-microglobulin into a disordered amyloidogenic precursor at low pH. <b>2017</b> , 46, 65-76	2
56	Probing Conformational Exchange Dynamics in a Short-Lived Protein Folding Intermediate by Real-Time Relaxation-Dispersion NMR. <b>2017</b> , 139, 1065-1068	15
55	Increased ISheet Dynamics and D-E Loop Repositioning Are Necessary for Cu(II)-Induced Amyloid Formation by I-Microglobulin. <i>Biochemistry</i> , <b>2017</b> , 56, 1095-1104	3.2 10
54	Inhibition of Human Serum Albumin Fibrillation by Two-Dimensional Nanoparticles. <b>2017</b> , 121, 5474-5482	25
53	Small molecule-mediated inhibition of I-microglobulin-based amyloid fibril formation. <b>2017</b> , 292, 10630-10638	10
52	Self-assembling peptide and protein amyloids: from structure to tailored function in nanotechnology. <b>2017</b> , 46, 4661-4708	467
51	-Stilbenoids with Extended Fluorescence Lifetimes for the Characterization of Amyloid Fibrils. <b>2017</b> , 2, 4693-4704	9
50	sw ApoMb Amyloid Aggregation under Nondenaturing Conditions: The Role of Native Structure Stability. <b>2017</b> , 113, 991-1001	5
49	Mechanism of Initiation, Association, and Formation of Amyloid Fibrils Modeled with the N-Terminal Peptide Fragment, IKYLEFIS, of Myoglobin G-Helix. <b>2017</b> , 121, 7536-7549	3
48	Melanosomal formation of PMEL core amyloid is driven by aromatic residues. <b>2017</b> , 7, 44064	19



47	Simple Model of the Effect of Solution Conditions on the Nucleation of Amyloid Fibrils. <b>2017</b> , 121, 8893-8901	7
46	Amyloidogenic Pattern Prediction of HIV-1 Proteins. <b>2017</b> , 823-895	1
45	Real-Time Observation of the Interaction between Thioflavin T and an Amyloid Protein by Using High-Sensitivity Rheo-NMR. <b>2017</b> , 18,	7
44	Comparative Study of Toluidine Blue O and Methylene Blue Binding to Lysozyme and Their Inhibitory Effects on Protein Aggregation. <b>2018</b> , 3, 2588-2601	29
43	Sodium lauryl sarcosinate (sarkosyl) modulate amyloid fibril formation in hen egg white lysozyme (HEWL) at alkaline pH: a molecular insight study. <b>2018</b> , 36, 1550-1565	12
42	Insights into the Origin of Distinct Medin Fibril Morphologies Induced by Incubation Conditions and Seeding. <b>2018</b> , 19,	3
41	Emerging Developments in Targeting Proteotoxicity in Neurodegenerative Diseases. <b>2019</b> , 33, 883-904	14
40	The Early Phase of $\beta$ m Aggregation: An Integrative Computational Study Framed on the D76N Mutant and the $\beta$ 6 Variant. <b>2019</b> , 9,	6
39	Conformational Evolution of Molecular Signatures during Amyloidogenic Protein Aggregation. <b>2019</b> , 10, 4593-4611	7
38	Investigating the Molecular Basis of the Aggregation Propensity of the Pathological D76N Mutant of Beta-2 Microglobulin: Role of the Denatured State. <b>2019</b> , 20,	2
37	Structural Heterogeneity in the Preamyloid Oligomers of $\beta$ 2-Microglobulin. <b>2020</b> , 432, 396-409	5
36	Collagen I Weakly Interacts with the $\beta$ Sheets of $\beta$ Microglobulin and Enhances Conformational Exchange To Induce Amyloid Formation. <b>2020</b> , 142, 1321-1331	8
35	Colloidal-like aggregation of a functional amyloid protein. <b>2020</b> , 22, 23286-23294	2
34	The <i>Yarrowia lipolytica</i> orthologs of Sup35p assemble into thioflavin T-negative amyloid fibrils. <b>2020</b> , 529, 533-539	3
33	The effects of mutation and modification on the structure and stability of human lysozyme: A molecular link between carbamylation and atherosclerosis. <b>2020</b> , 100, 107703	2
32	Cationic gemini surfactant stimulates amyloid fibril formation in bovine liver catalase at physiological pH. A biophysical study.. <b>2020</b> , 10, 43751-43761	3
31	The Early Phase of $\beta$ -Microglobulin Aggregation: Perspectives From Molecular Simulations. <b>2020</b> , 7, 578433	3
30	Effect of TMAO and Urea on Dimers and Tetramers of Amyloidogenic Heptapeptides (FGAILSS). <b>2020</b> , 5, 26986-26998	1

- 29 Measuring the Energy Barrier of the Structural Change That Initiates Amyloid Formation. **2020**, 92, 4731-4735 3
- 28 Epigallocatechin-3-gallate Inhibits Cu(II)-Induced  $\beta$ -Microglobulin Amyloid Formation by Binding to the Edge of Its  $\beta$ -Sheets. *Biochemistry*, **2020**, 59, 1093-1103 3,2 2
- 27 Conformational Priming of RepA-WH1 for Functional Amyloid Conversion Detected by NMR Spectroscopy. **2020**, 28, 336-347.e4 4
- 26 Functional amyloids. **2020**, 263-282 1
- 25 Proline isomerization effects in the amyloidogenic protein  $\beta$ -microglobulin. **2021**, 23, 356-367 2
- 24 Identification of critical amino acid residues in the regulatory N-terminal domain of PMEL. **2021**, 11, 7730 1
- 23 AlphaFold and the amyloid landscape. **2021**, 433, 167059 10
- 22 Nicking and fragmentation are responsible for  $\beta$ -lactalbumin amyloid fibril formation at acidic pH and elevated temperature. **2021**, 30, 1919-1934 4
- 21  $\beta$ -Microglobulin triggers NLRP3 inflammasome activation in tumor-associated macrophages to promote multiple myeloma progression. **2021**, 54, 1772-1787.e9 8
- 20 Spontaneous Fibrillation of Bovine Serum Albumin at Physiological Temperatures Promoted by Hydrolysis-Prone Ionic Liquids. **2021**, 37, 10319-10329 1
- 19 From uncovering the mechanism of transthyretin aggregation to the drug tafamidis for ameliorating neurodegeneration and cardiomyopathy. **2021**, 65-103 1
- 18  $\beta$ -Microglobulin and Dialysis-Related Amyloidosis. **2007**, 217-239 2
- 17 Protein Misfolding and Toxicity in Dialysis-Related Amyloidosis. **2012**, 377-405 2
- 16 Optimized fast mixing device for real-time NMR applications. *Journal of Magnetic Resonance*, **2017**, 281, 125-129 3 9
- 15 SDS can be utilized as an amyloid inducer: a case study on diverse proteins. *PLoS ONE*, **2012**, 7, e29694 3,7 103
- 14 Molecular Insight into Human Lysozyme and Its Ability to Form Amyloid Fibrils in High Concentrations of Sodium Dodecyl Sulfate: A View from Molecular Dynamics Simulations. *PLoS ONE*, **2016**, 11, e0165213 3,7 21
- 13 Effect of UV Light on Amyloidogenic Proteins: Nucleation and Fibril Extension. **2009**, 267-288
- 12 Real-Time Observation of Amyloid Fibril Growth by Total Internal Reflection Fluorescence Microscopy. **2009**, 289-299

- 11 Conformational Priming of RepA-WH1 for Functional Amyloid Conversion Detected by NMR Spectroscopy.
- 10 Collagen I weakly interacts with the  $\beta$ -sheets of  $\beta$ -microglobulin and enhances conformational exchange to induce amyloid formation.
- 9 Plant Natriuretic Peptide Active Domain Forms Amyloid-like Fibrils in a pH-Dependent Manner.. *Plants*, **2021**, 11, 4-5
- 8 Structure and Polymorphism of Amyloid and Amyloid-Like Aggregates. *Biochemistry (Moscow)*, **2022**, 87, 450-463 2.9
- 7 Molecular Insights into the Inhibition of Dialysis-Related  $\beta$ m Amyloidosis Orchestrated by a Bispidine Peptidomimetic Analogue. *Biochemistry*, 3.2
- 6 The Possible Mechanism of Amyloid Transformation Based on the Geometrical Parameters of Early-Stage Intermediate in Silico Model for Protein Folding. **2022**, 23, 9502 ○
- 5 Recent developments in the treatment of amyloidosis. **2022**, 47-64 ○
- 4 Amyloid fibrillation of the glaucoma associated myocilin protein is inhibited by epicatechin gallate (ECG). **2022**, 12, 29469-29481 ○
- 3 The residual structure of acid-denatured  $\beta$  -microglobulin is relevant to an ordered fibril morphology. ○
- 2 Inhibition of lysozyme amyloid fibrillation by curcumin-conjugated silver nanoparticles: A multispectroscopic molecular level study. **2022**, 121156 ○
- 1 Application of nanofibrous protein for the purification of contaminated water as a next generational sorption technology: a review. **2023**, 34, 232004 ○