

CITATION REPORT

List of articles citing

Arterial hypertension, microalbuminuria, and risk of ischemic heart disease

DOI: 10.1161/01.hyp.35.4.898
Hypertension, 2000, 35, 898-903.

Source: <https://exaly.com/paper-pdf/31986638/citation-report.pdf>

Version: 2024-04-26

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
359	Strokes Associated with Nephrotic Syndrome and Other Renal Diseases. 469-477		
358	Dissociation between microalbuminuria and common carotid thickness in essential hypertensive men. 2000 , 14, 831-5		14
357	Ramipril delays progression of hypertensive renal disease more than amlodipine. 2001 , 5, 143-144		
356	Microalbuminuria Is Associated with Increased Risk of Death in Patients with Myocardial Infarction. 2001 , 1, 305-309		3
355	Current World Literature. 2001 , 10, 271-293		
354	Utilization of the non-covalent fluorescent dye, NanoOrange, as a potential clinical diagnostic tool. Nanomolar human serum albumin quantitation. 2001 , 754, 345-56		50
353	Non-diabetic microalbuminuria, endothelial dysfunction and cardiovascular disease. 2001 , 6, 257-64		73
352	Microalbuminuria, cardiovascular, and renal risk in primary hypertension. 2002 , 13 Suppl 3, S169-72		47
351	Microalbuminuria is an integrated marker of subclinical organ damage in primary hypertension. 2002 , 16, 399-404		55
350	Impact of an ACE inhibitor and calcium antagonist on microalbuminuria and lipid subfractions in type 2 diabetes: a randomised, multi-centre pilot study. 2002 , 16, 185-91		33
349	Microalbuminuria is associated with unfavourable cardiac geometric adaptations in essential hypertensive subjects. 2002 , 16, 249-54		26
348	Microalbuminuria, an integrated marker of cardiovascular risk in essential hypertension. 2002 , 16, 79-89		99
347	Microalbuminuria in essential hypertension. 2002 , 16 Suppl 1, S74-7		13
346	Blood pressure and cardiac autonomic function in relation to risk factors and treatment perspectives in Type 1 diabetes. 2002 , 3, 222-42		8
345	Factors related to the occurrence of microalbuminuria during antihypertensive treatment in essential hypertension. <i>Hypertension</i> , 2002 , 39, 794-8	8.5	43
344	Microalbuminuria identifies overall cardiovascular risk in essential hypertension: an artificial neural network-based approach. <i>Journal of Hypertension</i> , 2002 , 20, 1315-21	1.9	50
343	Microalbuminuria in hypertensive patients with electrocardiographic left ventricular hypertrophy: the LIFE study. <i>Journal of Hypertension</i> , 2002 , 20, 405-12	1.9	123

342	Primary renal disease as a cardiovascular risk factor. 2002 , 324, 138-45	9
341	Prevention of post-transplant cardiovascular disease--report and recommendations of an ad hoc group. 2002 , 2, 491-500	63
340	Urine albumin/creatinine ratio and echocardiographic left ventricular structure and function in hypertensive patients with electrocardiographic left ventricular hypertrophy: the LIFE study. Losartan Intervention for Endpoint Reduction. 2002 , 143, 319-26	107
339	Relationships between new risk factors and circadian blood pressure variation in untreated subjects with essential hypertension. 2002 , 15, 600-4	31
338	Microalbuminúria em pacientes diabéticos tipo 1: prevalúncia e fatores associados. 2002 , 46, 640-647	0
337	Association between high-normal albuminuria and risk factors for cardiovascular and renal disease in essential hypertensive men. 2002 , 40, 1-8	73
336	The impact of antihypertensive drug groups on urinary albumin excretion in a non-diabetic population. 2002 , 53, 31-6	14
335	Microalbuminuria in hypertension. 2003 , 5, 208-14	21
334	Association of dietary protein intake and microalbuminuria in healthy adults: Third National Health and Nutrition Examination Survey. 2003 , 41, 580-7	61
333	Microalbuminuria and all-cause mortality in 2,089 apparently healthy individuals: a 4.4-year follow-up study. The Nord-Trúdelag Health Study (HUNT), Norway. 2003 , 42, 466-73	195
332	Cigarette smoking: an important renal risk factor - far beyond carcinogenesis. 2002 , 1, 137-55	31
331	Blood pressure independent association of microalbuminuria and left ventricular hypertrophy in hypertensive men. 2003 , 254, 76-84	34
330	Effects of fenofibrate on albuminuria in patients with hypertriglyceridemia and/or hyperuricemia: a multicenter, randomized, double-blind, placebo-controlled, crossover study. 2003 , 64, 434-46	10
329	Corticosteroid use and its association with microalbuminuria in the adult population. 2003 , 16, 349-53	9
328	A comparison of the prevalence of the metabolic syndrome using two proposed definitions. 2003 , 26, 575-81	419
327	Preventing renal dysfunction in patients with hypertension: clinical implications from the early AASK Trial results. 2003 , 3, 193-200	1
326	Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. <i>Hypertension</i> , 2003 , 42, 1206-52	8.5 10207
325	Mild renal dysfunction and subclinical cardiovascular damage in primary hypertension. <i>Hypertension</i> , 2003 , 42, 14-8	8.5 59

324	ACE inhibitor intervention in Type 1 diabetes with low grade microalbuminuria. 2003 , 4, 17-26		4
323	Microalbuminuria is closely related to impaired arterial elasticity in untreated patients with essential hypertension. 2003 , 93, c106-11		24
322	Microalbuminuria and all-cause mortality in treated hypertensive individuals: does sex matter? The Nord-Trøndelag Health Study (HUNT), Norway. 2003 , 108, 2783-9		52
321	Microalbuminuria. <i>Journal of Hypertension</i> , 2003 , 21, 1229-1233	1.9	9
320	Is it time to measure microalbuminuria in hypertension?. <i>Journal of Hypertension</i> , 2003 , 21, 1213-20	1.9	35
319	Nitric oxide in diabetes mellitus. <i>Journal of Hypertension</i> , 2003 , 21, 1081-3	1.9	4
318	2003 European Society of Hypertension-European Society of Cardiology guidelines for the management of arterial hypertension. <i>Journal of Hypertension</i> , 2003 , 21, 1011-53	1.9	2526
317	Microalbuminuria as a Predictor of Cardiovascular Risk in Initial Screening of the Hypertensive Patient. 2003 , 3, 2-3		
316	Microalbuminuria and cardiovascular events in elderly hypertensive patients without previous cardiovascular complications. <i>Hypertension Research</i> , 2003 , 26, 603-8	4.7	11
315	Recent insights in the development of organ damage caused by hypertension. 2004 , 59, 369-81		11
314	The metabolic syndrome is associated with advanced vascular damage in patients with coronary heart disease, stroke, peripheral arterial disease or abdominal aortic aneurysm. 2004 , 25, 342-8		178
313	Microalbuminuria independently predicts all-cause and cardiovascular mortality in a British population: The European Prospective Investigation into Cancer in Norfolk (EPIC-Norfolk) population study. 2004 , 33, 189-98		138
312	Very low levels of microalbuminuria are associated with increased risk of coronary heart disease and death independently of renal function, hypertension, and diabetes. 2004 , 110, 32-5		502
311	A prospective study of microalbuminuria and incident coronary heart disease and its prognostic significance in a British population: the EPIC-Norfolk study. 2004 , 159, 284-93		76
310	Usefulness of microalbuminuria in cardiovascular risk stratification of essential hypertensive patients. 2004 , 96, c123-30		11
309	The role of oxidative stress-altered lipoprotein structure and function and microinflammation on cardiovascular risk in patients with minor renal dysfunction. 2004 , 15, 538-48		138
308	Estimating laboratory precision of urinary albumin excretion and other urinary measures in the International Study on Macronutrients and Blood Pressure. 2004 , 160, 287-94		11
307	New European guidelines for management of hypertension: what is relevant for the nephrologist. 2004 , 19, 524-8		0

306	Evaluation of measures of urinary albumin excretion in epidemiologic studies. 2004 , 160, 1122-31		63
305	Cardiovascular and renal risk assessment as a guide for treatment in primary hypertension. 2004 , 15 Suppl 1, S34-6		5
304	Relationship between albumin excretion rate and aortic stiffness in untreated essential hypertensive patients. 2004 , 256, 22-9		54
303	Relation of urinary albumin excretion to coronary heart disease and low renal function: role of blood pressure. 2004 , 65, 2290-7		16
302	Albuminuria and cardiovascular risk in hypertensive patients with left ventricular hypertrophy: the LIFE Study. 2004 , S56-8		18
301	The case for blood pressure control in risk groups. 2004 , 58, 844-9		4
300	Polymorphisms of the angiotensinogen gene and the outcome of microalbuminuria in essential hypertension: a 3-year follow-up study. 2004 , 18, 25-31		5
299	Guidelines for management of hypertension: report of the fourth working party of the British Hypertension Society, 2004-BHS IV. 2004 , 18, 139-85		868
298	Creatinine clearance and signs of end-organ damage in primary hypertension. 2004 , 18, 511-6		15
297	Hypertension and microalbuminuria: the new detrimental duo. 2004 , 13, 198-211		28
296	[Relevance of chronic kidney disease and cardiovascular prognosis in arterial hypertension]. 2004 , 123, 143-8		1
295	Inter-relationships of microalbuminuria with the other surrogates of the atherosclerotic cardiovascular disease in hypertensive subjects. 2004 , 17, 470-6		42
294	Microalbuminuria and cardiovascular risk. 2004 , 17, 986-93		98
293	Cardiovascular implications of albuminuria. <i>Journal of Clinical Hypertension</i> , 2004 , 6, 13-7	2.3	15
292	Microalbuminuria: definition, detection, and clinical significance. <i>Journal of Clinical Hypertension</i> , 2004 , 6, 2-7	2.3	76
291	On the importance of estimating renal function for cardiovascular risk assessment. <i>Journal of Hypertension</i> , 2004 , 22, 1635-9	1.9	16
290	Optimizing global risk evaluation in primary hypertension: the role of microalbuminuria and cardiovascular ultrasonography. <i>Journal of Hypertension</i> , 2004 , 22, 907-13	1.9	32
289	Does albuminuria predict cardiovascular outcome on treatment with losartan versus atenolol in hypertension with left ventricular hypertrophy? A LIFE substudy. <i>Journal of Hypertension</i> , 2004 , 22, 1805-11	1.9	99

288	Treatment of hypertension in patients with diabetes. 2004 , 164, 1850-7		59
287	2003 European Society of Hypertension-European Society of Cardiology Guidelines for the Management of Arterial Hypertension. 2004 , 4, 6-51		
286	What is the evidence that microalbuminuria is a predictor of cardiovascular disease events?. 2005 , 14, 271-6		76
285	Microalbuminuria, renal function and development of sustained hypertension: a longitudinal study in the early stage of hypertension. <i>Journal of Hypertension</i> , 2005 , 23, 175-82	1.9	36
284	What is the evidence that microalbuminuria is a predictor of cardiovascular disease events?. 2005 , 4, 375-380		
283	Controlled-release nifedipine and candesartan low-dose combination therapy in patients with essential hypertension: the NICE Combi (Nifedipine and Candesartan Combination) Study. <i>Journal of Hypertension</i> , 2005 , 23, 445-53	1.9	57
282	Protective effects of an angiotensin II receptor blocker and a long-acting calcium channel blocker against cardiovascular organ injuries in hypertensive patients. <i>Hypertension Research</i> , 2005 , 28, 351-9	4.7	22
281	Minor abnormalities of renal function: a situation requiring integrated management of cardiovascular risk. 2005 , 19, 429-37		8
280	Prevention of chronic kidney and vascular disease: toward global health equity--the Bellagio 2004 Declaration. 2005 , S1-6		59
279	The kidney, a cardiovascular risk marker, and a new target for therapy. 2005 , S25-9		71
278	Diagnostic significance of transferrinuria and albumin-specific dipstick testing in primary care patients with elevated office blood pressure. 2005 , 19, 205-9		9
277	Microalbuminuria and C-reactive protein: similar messengers of cardiovascular risk?. 2005 , 7, 379-84		23
276	The Relationship between Microalbuminuria and Coronary Artery Stenosis or Inflammatory Markers in Patients with Angina Pectoris. 2005 , 35, 49		1
275	Impact of target organ damage assessment in the evaluation of global risk in patients with essential hypertension. 2005 , 16 Suppl 1, S89-91		14
274	Brachial-Ankle Pulse Wave Velocity as a Novel Measure of Arterial Stiffness: Present Evidences and Perspectives. 2005 , 1, 223-234		16
273	Role of microalbuminuria in the assessment of cardiovascular risk in essential hypertension. 2005 , 16 Suppl 1, S39-41		17
272	Low-grade albuminuria and incidence of cardiovascular disease events in nonhypertensive and nondiabetic individuals: the Framingham Heart Study. 2005 , 112, 969-75		563
271	New definition of microalbuminuria in hypertensive subjects: association with incident coronary heart disease and death. <i>Hypertension</i> , 2005 , 46, 33-7	8.5	83

270	Reduction in albuminuria translates to reduction in cardiovascular events in hypertensive patients: losartan intervention for endpoint reduction in hypertension study. <i>Hypertension</i> , 2005 , 45, 198-202	8.5	436
269	Early changes in 24-hour ambulatory blood pressure are associated with high normal albumin excretion rate in children with type 1 diabetes mellitus. 2005 , 18, 879-85		11
268	Modest salt reduction reduces blood pressure and urine protein excretion in black hypertensives: a randomized control trial. <i>Hypertension</i> , 2005 , 46, 308-12	8.5	127
267	Clinical use and pathogenetic basis of laboratory tests for the evaluation of primary arterial hypertension. 2005 , 42, 393-452		4
266	Renal manifestations of the metabolic syndrome. 2005 , 20, 861-4		19
265	Urinary albumin and cardiovascular profile in the middle-aged population. 2005 , 25, 367-71		7
264	Associations of microalbuminuria and blood pressure with carotid, aortic and femoral atheromatous plaques in elderly Finns. 2005 , 69, 262-71		4
263	True resistant hypertension: is it possible to be recognized in the office?. 2005 , 18, 1534-40		71
262	Pediatric hypertension: diagnosis, evaluation, management, and treatment for the primary care physician. 2005 , 35, 262-94		19
261	2007 Guidelines for the management of arterial hypertension: The Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2007 , 28, 1462-536		1418
260	Diabetic hypertension. 2006 , 2, 25-36		4
259	Brachial-ankle pulse wave velocity and microalbuminuria. <i>Hypertension Research</i> , 2006 , 29, 469-70	4.7	1
258	Microalbuminuria as an early marker for cardiovascular disease. 2006 , 17, 2100-5		258
257	Albumin-to-creatinine ratio predicts change in ambulatory blood pressure in normotensive persons: a 7.5-year prospective study. 2006 , 19, 220-6		9
256	Microalbuminuria, blood pressure load, and systemic vascular permeability in primary hypertension. 2006 , 19, 1183-9		20
255	Total cardiovascular risk: a new treatment concept. <i>Journal of Hypertension</i> , 2006 , 24, S17-24	1.9	20
254	Microalbuminuria and endothelial dysfunction: emerging targets for primary prevention of end-organ damage. 2006 , 47 Suppl 2, S151-62; discussion S172-6		81
253	Alpha-adducin and angiotensin-converting enzyme polymorphisms in hypertension: evidence for a joint influence on albuminuria. <i>Journal of Hypertension</i> , 2006 , 24, 931-7	1.9	16

252	Update on microalbuminuria as a biomarker in renal and cardiovascular disease. 2006 , 15, 631-6	56
251	Target organ damage and cardiovascular complications in patients with hypertension and type 2 diabetes in Spain: a cross-sectional study. 2006 , 5, 23	29
250	Endothelial dysfunction: a comprehensive appraisal. 2006 , 5, 4	277
249	Very low level of microalbuminuria is associated with increased risk of death in subjects with cardiovascular or cerebrovascular diseases. 2006 , 260, 231-7	58
248	Glomerular hyperfiltration predicts the development of microalbuminuria in stage 1 hypertension: the HARVEST. 2006 , 70, 578-84	71
247	ESH-ESC guidelines for the management of hypertension. 2006 , 31, 331-8	41
246	Microalbuminuria as a target to improve cardiovascular and renal outcomes. 2006 , 47, 927-46	71
245	Microalbuminuria and chronic kidney disease as risk factors for cardiovascular disease. 2006 , 21, 2366-74	85
244	Urinary albumin excretion and the renin-angiotensin system in cardiovascular risk management. 2006 , 7, 2505-20	2
243	Microalbuminuria In Primary Hypertension. 2006 , 2, 11-19	2
242	Does albuminuria predict cardiovascular outcomes on treatment with losartan versus atenolol in patients with diabetes, hypertension, and left ventricular hypertrophy? The LIFE study. 2006 , 29, 595-600	88
241	Chronic kidney disease as a situation of high added risk in hypertensive patients. 2006 , 17, S136-40	12
240	The Diabetic Kidney. 2006 ,	1
239	Prevalence of abnormal urinary albumin excretion rate in hypertensive patients with impaired fasting glucose and its association with cardiovascular disease. 2006 , 17, S178-88	21
238	Microalbuminuria and risk for cardiovascular disease: Analysis of potential mechanisms. 2006 , 17, 2106-11	305
237	Urinary albumin excretion is associated with impaired flow- and nitroglycerin-mediated brachial artery dilatation in hypertensive adults. 2007 , 21, 231-8	56
236	Occlusion therapy for amblyopia. 2007 , 335, 678-9	5
235	Reform of the coroner system and death certification. 2007 , 335, 680-1	8

234	Encouraging children and adolescents to be more active. 2007 , 335, 677-8		17
233	Chapter 1: Introduction. 2007 , 14, S2-S113		
232	Albuminuria: pathophysiology, epidemiology and clinical relevance of an emerging marker for cardiovascular disease. 2007 , 3, 519-24		12
231	2007 Guidelines for the Management of Arterial Hypertension: The Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). <i>Journal of Hypertension</i> , 2007 , 25, 1105-87	1.9	3825
230	Reliability of urinary albumin, total protein, and creatinine assays after prolonged storage: the Family Investigation of Nephropathy and Diabetes. 2007 , 2, 1156-62		18
229	Insights into the Relationship Between Hypertension and Albuminuria. 2007 , 3, 29-37		6
228	Mental health in disaster settings. 2007 , 335, 679-80		11
227	Diagnosing left ventricular hypertrophy in arterial hypertension. 2007 , 335, 681-2		8
226	Association of cardiovascular risk factors with microalbuminuria in hypertensive individuals: the i-SEARCH global study. <i>Journal of Hypertension</i> , 2007 , 25, 2317-24	1.9	61
225	Lack of association between endothelial nitric oxide synthase gene polymorphisms, microalbuminuria and endothelial dysfunction in hypertensive men. <i>Journal of Hypertension</i> , 2007 , 25, 1389-95	1.9	20
224	2007 ESH-ESC Guidelines for the management of arterial hypertension: the task force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2007 , 16, 135-232		247
223	Microalbuminuria is a risk factor for cerebral small vessel disease in community-based elderly subjects. 2007 , 255, 27-34		85
222	[ESH/ESC 2007 Guidelines for the management of arterial hypertension]. 2007 , 60, 968.e1-94		29
221	Changes in self-monitored pulse pressure correlate with improvements in B-type natriuretic Peptide and urinary albumin in treated hypertensive patients. 2007 , 20, 1268-75		10
220	Microalbuminuria associated with systolic blood pressure and arterial compliance in Chinese metabolic syndrome patients. 2007 , 120, 1395-1399		4
219	The association between metabolic syndrome, microalbuminuria and impaired renal function in the general population: impact on cardiovascular disease and mortality. 2007 , 262, 470-8		65
218	Microalbuminuria screening in patients with hypertension: recommendations for clinical practice. 2008 , 62, 97-108		30
217	La microalbuminuria como factor pronóstico en el enfermo cardiovascular. 2007 , 7, 31A-43A		

216	Taking the pressure off type 2 diabetes mellitus: implementing hypertension guidelines. 2007 , 22, 159-65	1
215	Effect of microalbuminuria lowering on regression of left ventricular hypertrophy in children and adolescents with essential hypertension. 2007 , 28, 27-33	78
214	Inhibition of the renin-angiotensin system and chronic kidney disease. 2008 , 40, 1015-25	19
213	Low-grade urinary albumin excretion in normotensive/non-diabetic obstructive sleep apnea patients. 2008 , 12, 217-22	21
212	Relation of left ventricular hypertrophy to microalbuminuria and C-reactive protein in children and adolescents with essential hypertension. 2008 , 29, 580-4	46
211	What is the mechanism of microalbuminuria in diabetes: a role for the glomerular endothelium?. 2008 , 51, 714-25	235
210	Association of obstructive sleep apnea with urinary albumin excretion in essential hypertension: a cross-sectional study. 2008 , 52, 285-93	51
209	[Prevalence and clinical characteristics of microalbuminuria in the Spanish hypertensive population]. 2008 , 130, 201-5	6
208	Renal Filtration, Transport, and Metabolism of Albumin and Albuminuria. 2008 , 2081-2112	3
207	Secondary hypertension: current diagnosis and treatment. 2008 , 124, 6-21	47
206	Microalbuminuria and global myocardial function in patients with essential hypertension [correction of hypertensive]. 2008 , 126, 268-72	4
205	Cardiovascular sequelae of preeclampsia/eclampsia: a systematic review and meta-analyses. 2008 , 156, 918-30	617
204	Severe, very early onset preeclampsia: subsequent pregnancies and future parental cardiovascular health. 2008 , 140, 171-7	57
203	Complicaciones microvasculares. Nefropatía diabética. 2008 , 10, 1125-1129	
202	Overview of the i-SEARCH Global Study : Cardiovascular Risk Factors and Microalbuminuria in Hypertensive Individuals. 2008 , 15, 217-24	4
201	Association of microalbuminuria with brachial-ankle pulse wave velocity: the Ohasama study. 2008 , 21, 413-8	39
200	Combination of the electrocardiographic strain pattern and albuminuria for the prediction of new-onset heart failure in hypertensive patients: the LIFE study. 2008 , 21, 273-9	16
199	Prevalence of albuminuria and cardiovascular risk profile in a referred cohort of patients with type 2 diabetes: an Asian perspective. 2008 , 10, 397-403	23

198	Microalbuminuria in type 2 diabetes and hypertension: a marker, treatment target, or innocent bystander?. 2008 , 31 Suppl 2, S194-201		104
197	Endothelial dysfunction in normal and abnormal glucose metabolism. 2008 , 45, 17-43		52
196	Vascular permeability, blood pressure, and organ damage in primary hypertension. <i>Hypertension Research</i> , 2008 , 31, 873-9	4.7	11
195	Association of low-grade inflammation and platelet activation in patients with hypertension with microalbuminuria. 2008 , 114, 449-55		22
194	Soluble CD40L and its role in essential hypertension: diagnostic and therapeutic implications. 2008 , 8, 194-202		16
193	Inappropriate left ventricular mass is associated with microalbuminuria independently of left ventricular hypertrophy in primary hypertension. <i>Journal of Hypertension</i> , 2008 , 26, 345-50	1.9	11
192	Evidence for no global effect of metabolic syndrome per se on early hypertensive sequelae. <i>Journal of Hypertension</i> , 2008 , 26, 773-9	1.9	12
191	Early detection and management of the high-risk patient with elevated blood pressure. 2008 , 4, 289-96		19
190	MICROSCOPIC POLYANGIITIS AND POLYARTERITIS NODOSA. 311-330		
189	NEPHROTIC SYNDROME AND OTHER RENAL DISEASES AND STROKE. 391-400		1
188	The relationship between proteinuria and coronary risk: a systematic review and meta-analysis. 2008 , 5, e207		148
187	Abdominal obesity is associated with microalbuminuria and an elevated cardiovascular risk profile in patients with hypertension. 2009 , 5, 577-85		42
186	Atrial fibrillation and heart rate independently correlate to microalbuminuria in hypertensive patients. 2009 , 30, 1364-71		36
185	Proteinuria and clinical outcomes in hypertensive patients. 2009 , 22, 1137-47		9
184	Reappraisal of European guidelines on hypertension management: a European Society of Hypertension Task Force document. <i>Journal of Hypertension</i> , 2009 , 27, 2121-58	1.9	1004
183	Determinants of platelet activation in hypertensives with microalbuminuria. 2009 , 46, 922-7		15
182	Biochemical and histopathological effects of dietary oxidized cholesterol in rats. 2009 , 29, 715-23		16
181	Fixed-dose manidipine/delapril versus losartan/hydrochlorothiazide in hypertensive patients with type 2 diabetes and microalbuminuria. 2009 , 26, 313-24		11

180	Microalbuminuria and obesity: impact on cardiovascular disease and mortality. 2009 , 71, 40-5		25
179	Chronic kidney disease: the forgotten cardiovascular risk factor. 2009 , 63, 964-5		1
178	A history of preeclampsia identifies women who have underlying cardiovascular risk factors. 2009 , 200, 58.e1-8		146
177	Reappraisal of European guidelines on hypertension management: a European Society of Hypertension Task Force document. 2009 , 18, 308-47		186
176	Cardiovascular Endocrinology. 2009 ,		2
175	Cardiovascular Risk in Hypertensive Patients with Renal Dysfunction. 2009 , 16, 13-20		1
174	Risk stratification with the risk chart from the European Society of Hypertension compared with SCORE in the general population. <i>Journal of Hypertension</i> , 2009 , 27, 2351-7	1.9	29
173	Albuminuria in heart failure: what do we really know?. 2009 , 24, 148-54		16
172	Effects of angiotensin II receptor blockers on diabetic nephropathy. <i>Journal of Hypertension</i> , 2009 , 27, S15-21	1.9	23
171	The assessment of pulmonary diffusing capacity in diabetes mellitus with regard to microalbuminuria. 2009 , 48, 1939-43		18
170	End-organ dysfunction and cardiovascular outcomes: the role of the microcirculation. 2009 , 116, 175-90		49
169	Combined effect of albuminuria and estimated glomerular filtration rate on cardiovascular events and all-cause mortality in uncomplicated hypertensive patients. <i>Journal of Hypertension</i> , 2010 , 28, 848-55 ^{1.9}		20
168	Kidney disease after preeclampsia: a systematic review and meta-analysis. 2010 , 55, 1026-39		141
167	Poor blood pressure and urinary albumin excretion responses to home blood pressure-based antihypertensive therapy in depressive hypertensive patients. <i>Journal of Clinical Hypertension</i> , 2010 , 12, 345-9	2.3	2
166	Urine albumin excretion, within normal range, reflects increasing prevalence of metabolic syndrome in patients with essential hypertension. <i>Journal of Clinical Hypertension</i> , 2010 , 12, 597-602	2.3	14
165	VEGF-A gene promoter polymorphisms and microvascular complications in patients with essential hypertension. 2010 , 43, 1090-5		16
164	La protection vasculaire chez l'insuffisant rénal. 2010 , 22, 237-245		
163	Racial/ethnic differences in microalbuminuria among adults with prehypertension and hypertension: National Health and Nutrition Examination Survey (NHANES), 1999-2006. 2010 , 23, 859-64		36

162	Microalbuminuria is a predictor of chronic renal insufficiency in patients without diabetes and with hypertension: the MAGIC study. 2010 , 5, 1099-106	39
161	Albuminuria and the risk of incident stroke and stroke types in older adults. 2010 , 75, 1343-50	47
160	The association of microalbuminuria with mortality in patients with acute myocardial infarction. A ten-year follow-up study. 2010 , 5, e2	1
159	Hypertension in pregnancy: a comprehensive update. 2010 , 18, 178-89	51
158	The G protein B subunit C825T polymorphism is associated with microalbuminuria in hypertensive women and cardiovascular disease in hypertensive men. 2010 , 23, 1114-20	3
157	Selection on albuminuria enhances the efficacy of screening for cardiovascular risk factors. 2010 , 25, 3560-8	13
156	Spontaneous intracerebral hemorrhage. 2010 , 30, 555-64	79
155	Initial clinical encounter with the patient with established hypertension. 2010 , 28, 587-95	3
154	The contribution of early nephropathy to cardiovascular risk. 2010 , 28, 427-36	3
153	The relationship between microalbuminuria and the presence and extent of coronary atherosclerosis. 2010 , 61, 184-91	9
152	[Subclinical and established kidney disease in recently diagnosed hypertensive patients]. 2010 , 134, 246-53	0
151	Urinary albumin excretion among nondipper hypertensive patients is closely related with the pattern of nondipping. 2010 , 4, 196-202	7
150	Subclinical organ damage and cardiovascular risk prediction. 2010 , 19, 132-9	8
149	Microalbuminuria is not associated with endothelial dysfunction and coronary atherosclerosis in patients with acute coronary syndromes. 2010 , 32, 659-65	5
148	Persistent microalbuminuria after treatment with renin-angiotensin axis blockers: causes and results of treatment intensification. 2011 , 12, 333-9	7
147	The new concept of total cardiovascular risk management. 2011 , 5, 169-174	
146	Increased B-type natriuretic peptide and decreased proteinuria might reflect decreased capillary leakage and is associated with a better outcome in patients with severe burns. 2011 , 15, R161	10
145	Prognostic importance of baseline and serial changes in microalbuminuria in patients with resistant hypertension. 2011 , 216, 199-204	29

144	Novel Therapies of Diabetic Nephropathy. 2011 , 3, 23-35		1
143	Antihypertensive Therapy Considering the Prevention of Vascular Aging. 2011 , 17, 85		3
142	Relation of albuminuria to angiographically determined coronary arterial narrowing in patients with and without type 2 diabetes mellitus and stable or suspected coronary artery disease. 2011 , 107, 1144-8		32
141	Clinical practice: proteinuria. 2011 , 170, 15-20		17
140	What is the ideal blood pressure goal for patients with stage III or higher chronic kidney disease?. 2011 , 13, 492-501		10
139	Prevalence of target organ damage in hypertensive subjects attending primary care: C.V.P.C. study (epidemiological cardio-vascular study in primary care). 2011 , 12, 75		14
138	The characteristics of new semi-quantitative method for diagnosing proteinuria by using random urine samples. 2011 , 25, 14-9		3
137	Association of serum bilirubin levels with albuminuria in patients with essential hypertension. 2011 , 44, 859-63		5
136	Differences in MetS marker prevalence between black African and Caucasian teachers from the North West Province: Sympathetic Activity and Ambulatory Blood Pressure in Africans (SABPA) Study. 2011 , 16, 49-56		7
135	Arterial stiffness, ambulatory blood pressure and low-grade albuminuria in non-diabetic African and Caucasian men: the SABPA study. <i>Hypertension Research</i> , 2011 , 34, 862-8	4.7	13
134	Premenopausal women have increased risk of hypertensive target organ damage compared with men of similar age. 2011 , 20, 1175-81		26
133	Renal scintigraphy predicts global cardiovascular risk in hypertensive subjects with normal serum creatinine levels. 2011 , 20, 387-93		2
132	Angiotensin-converting enzyme inhibitors, angiotensin receptor blockers and combined therapy in patients with micro- and macroalbuminuria and other cardiovascular risk factors: a systematic review of randomized controlled trials. 2011 , 26, 2827-47		72
131	Urinary albumin excretion is associated with nocturnal systolic blood pressure in resistant hypertensives. <i>Hypertension</i> , 2011 , 57, 556-60	8.5	41
130	Physical activity is inversely associated with microalbuminuria in hypertensive patients at high cardiovascular risk: data from I-SEARCH. 2012 , 19, 1066-73		9
129	Microalbuminuria is independently associated with deep or infratentorial brain microbleeds in hypertensive adults. 2012 , 25, 430-6		36
128	Exposure to bisphenol A is associated with low-grade albuminuria in Chinese adults. 2012 , 81, 1131-9		52
127	Prognostic factors in resistant hypertension: implications for cardiovascular risk stratification and therapeutic management. 2012 , 10, 735-45		15

126	Angiogenesis and hypertension: the dual role of anti-hypertensive and anti-angiogenic therapies. 2012 , 10, 479-93		22
125	Cardiovascular risk assessment beyond Systemic Coronary Risk Estimation: a role for organ damage markers. <i>Journal of Hypertension</i> , 2012 , 30, 1056-64	1.9	63
124	Thresholds for pulse wave velocity, urine albumin creatinine ratio and left ventricular mass index using SCORE, Framingham and ESH/ESC risk charts. <i>Journal of Hypertension</i> , 2012 , 30, 1928-36	1.9	45
123	Oxidative stress impairs endothelial function in nondipper hypertensive patients. 2012 , 30, 85-92		21
122	A proposal for improving the KDIGO renal disease risk table. 2012 , 44, 1431-3		
121	Comparison of microalbuminuria with hs-CRP and low density lipoprotein levels in nondiabetic, nonhypertensive myocardial infarction patients. 2012 , 3, 287-9		4
120	Vascular heterogeneity in the kidney. 2012 , 32, 145-55		58
119	Effects of up-titration of candesartan versus candesartan plus amlodipine on kidney function in type 2 diabetic patients with albuminuria. 2012 , 26, 214-9		2
118	Guideline adherence in cardiovascular risk assessment and analysis in 15,000 hypertensive German patients in real life: results of the Prospective 3A Registry. <i>Journal of Clinical Hypertension</i> , 2012 , 14, 496-501	2.3	2
117	[Determination of urinary albumin excretion in arterial hypertension]. 2012 , 212, 172-8		6
116	Cardiovascular disease in chronic kidney disease: untying the Gordian knot. 2013 , 67, 14-31		37
115	Intensive treatment of persistent microalbuminuria: determinants of treatment resistance. 2013 , 8, 302-309		
114	Guía de práctica clínica de la ESH/ESC para el manejo de la hipertensión arterial (2013). 2013 , 30, 4-91		7
113	Creatinine and insulin predict cardiac mass in drug-naïve hypertensive patients. 2013 , 167, 519-24		6
112	Guía de práctica clínica de la ESH/ESC para el manejo de la hipertensión arterial (2013). 2013 , 66, 880.e1-880.e64		13
111	Approach to a case of secondary hypertension. 2013 , 2, 62-70		
110	2013 ESH/ESC Guidelines for the Management of Arterial Hypertension. 2013 , 22, 193-278		286
109	Increased frequency of microalbuminuria in patients receiving statins. 2013 , 8, 257-262		1

108	2013 ESH/ESC guidelines for the management of arterial hypertension: the Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2013 , 34, 2159-219		3400
107	Prevalence of abnormal urinary albumin excretion in elderly people: a Spanish survey. 2013 , 45, 553-60		9
106	Clinical and imaging parameters to predict cardiovascular outcome in asymptomatic subjects. 2013 , 29, 1595-602		14
105	Titration of telmisartan, but not addition of amlodipine, reduces urine albumin in diabetic patients treated with telmisartan-diuretic. <i>Journal of Hypertension</i> , 2013 , 31, 186-91	1.9	2
104	Preclinical cardiorenal interrelationships in essential hypertension. 2013 , 3, 38-47		9
103	Urinary albumin excretion at follow-up predicts cardiovascular outcomes in subjects with resistant hypertension. 2013 , 26, 1148-54		9
102	Kidney-protective effects of azelnidipine versus a diuretic in combination with olmesartan in hypertensive patients with diabetes and albuminuria: a randomized study. 2013 , 28, 1802-10		5
101	Screening for albuminuria with subsequent screening for hypertension and hypercholesterolaemia identifies subjects in whom treatment is warranted to prevent cardiovascular events. 2013 , 28, 2805-15		14
100	Global cardiovascular risk assessment in the management of primary hypertension: the role of the kidney. <i>International Journal of Hypertension</i> , 2013 , 2013, 542646	2.4	3
99	Renal Effects of DPP-4 Inhibitors: A Focus on Microalbuminuria. 2013 , 2013, 895102		32
98	Regression of target organ damage. <i>Journal of Hypertension</i> , 2013 , 31, 1535-6	1.9	1
97	2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). <i>Journal of Hypertension</i> , 2013 , 31, 1281-357	1.9	3363
96	Obesity is a strong determinant of hypertensive target organ damage in young-to-middle-age patients. 2013 , 37, 224-9		12
95	De Gruyter. 2013 , 21,		1
94	Proteinuria and its relation to cardiovascular disease. <i>International Journal of Nephrology and Renovascular Disease</i> , 2013 , 7, 13-24	2.5	47
93	COPD and microalbuminuria: a 12-year follow-up study. 2014 , 43, 1042-50		19
92	Association between albuminuria, atherosclerotic plaques, elevated pulse wave velocity, age, risk category and prognosis in apparently healthy individuals. <i>Journal of Hypertension</i> , 2014 , 32, 1034-41; discussion 1041	1.9	8
91	Brachial flow-mediated dilation predicts subclinical target organ damage progression in essential hypertensive patients: a 3-year follow-up study. <i>Journal of Hypertension</i> , 2014 , 32, 2393-400; discussion 2400	1.9	13

90	Serum uric acid concentration and asymptomatic hyperuricemia with subclinical organ damage in general population. 2014 , 65, 634-40		15
89	Renal disease and left atrial remodeling predict atrial fibrillation in patients with cardiovascular risk factors. 2014 , 175, 90-5		21
88	Impact of the albumin to creatinine ratio and the coronary artery state on vascular events. 2014 , 113, 1616-20		6
87	Blood pressure, albuminuria and renal dysfunction: the chicken or egg dilemma. 2014 , 29, 1453-5		14
86	Matrix metalloproteinase 9-mediated shedding of syndecan 4 in response to tumor necrosis factor α a contributor to endothelial cell glycocalyx dysfunction. 2014 , 28, 4686-99		84
85	Effective risk stratification in patients with moderate cardiovascular risk using albuminuria and atherosclerotic plaques in the carotid arteries. <i>Journal of Hypertension</i> , 2015 , 33, 1563-70	1.9	9
84	Discordant associations of lipid parameters with albuminuria and chronic kidney disease: a population-based study. 2015 , 14, 152		14
83	Left ventricular dilatation and subclinical renal damage in primary hypertension. <i>Journal of Hypertension</i> , 2015 , 33, 605-11; discussion 611	1.9	8
82	Body mass index (BMI) is associated with microalbuminuria in Chinese hypertensive patients. 2015 , 12, 1998-2008		10
81	Statins can improve proteinuria and glomerular filtration rate loss in chronic kidney disease patients, further reducing cardiovascular risk. Fact or fiction?. 2015 , 16, 1449-61		33
80	Is albuminuria a myocardial infarction risk equivalent for atherothrombotic events?. 2015 , 240, 21-5		10
79	Glycated haemoglobin A1c is associated with low-grade albuminuria in Chinese adults. 2015 , 5, e007429		8
78	Prognostic value of combined target-organ damage in patients with essential hypertension. 2015 , 28, 127-34		13
77	Use of Stem Cell Suspensions containing Separated Fetal Stem Cells in Complex Treatment of Patients with Essential Hypertension. 2016 , 6,		
76	ASSESSMENT OF MICROALBUMINURIA IN ESSENTIAL HYPERTENSIVES AND ITS RESPONSE TO ANGIOTENSIN-CONVERTING ENZYME INHIBITOR THERAPY. 2016 , 9, 32		4
75	Red cell distribution width and end-organ damage in patients with systo-diastolic hypertension. 2016 , 12, 319-25		5
74	Analytical Evaluation of the DiaSys Albumin in Urine/CSF FS Kit for Urine Albumin Measurement Using a JEOL BioMajesty JCA-BM6010/C Analyzer. 2016 , 6, 64		
73	Ambulatory versus clinic blood pressure in predicting overall subclinical target organ damage progression in essential hypertensive patients: a 3-year follow-up study. <i>Blood Pressure Monitoring</i> , 2016 , 21, 319-326	1.3	4

72	Association between atrial function assessed by 2D-speckle tracking echocardiography and albuminuria in patients with type 2 diabetes and coronary artery disease. 2016 , 44, 561-570		4
71	Residual small artery impairment in hypertensive patients with normal albumin-creatinine ratio. 2016 , 50, 167-71		1
70	Microalbuminuria predicts the recurrence of cardiovascular events in patients with essential hypertension. <i>Journal of Hypertension</i> , 2016 , 34, 646-53	1.9	12
69	Relationship Between Dyslipidemia and Albuminuria in Hypertensive Adults: A Nationwide Population-Based Study. 2016 , 95, e3224		7
68	[Comparison of the cardiovascular predictive value of MDRD and CKD-EPI in estimating chronic kidney disease]. 2016 , 42, 11-8		
67	Cardiovascular risk factors and incident albuminuria in screen-detected type 2 diabetes. 2017 , 33, e2877		2
66	Guidelines on the management of arterial hypertension and related comorbidities in Latin America. <i>Journal of Hypertension</i> , 2017 , 35, 1529-1545	1.9	36
65	3-year risk prediction of Coronary Heart Disease in hypertension patients: A preliminary study. 2017 , 2017, 1182-1185		4
64	Albuminuria and kidney function as prognostic marker of left ventricular mass among South Asians with hypertension. 2017 , 11, 811-822.e2		2
63	Most advisable strategy in search of asymptomatic target organ damage in hypertensive patients. 2017 , 34, 149-156		4
62	The micro-albuminuria is a new predictor of atrial fibrillation in hypertensive patients. 2017 , 7, 159-162		1
61	Two-year changes in proteinuria and risk for myocardial infarction in patients with hypertension: a prospective cohort study. <i>Journal of Hypertension</i> , 2017 , 35, 2295-2302	1.9	5
60	Urine Albumin Excretion as a Marker of Acute Glycemic Changes in Isolated Postprandial Hyperglycemia. 2017 , 9, 36-41		2
59	Gender-Specific Associations between Low Skeletal Muscle Mass and Albuminuria in the Middle-Aged and Elderly Population. <i>International Journal of Medical Sciences</i> , 2017 , 14, 1054-1064	3.7	2
58	Association between common carotid artery diameter and target organ damage in essential hypertension. <i>Journal of Hypertension</i> , 2018 , 36, 537-543	1.9	8
57	[Usefulness of patient education in antihypertensive treatment compliance in black Africans]. <i>Annales De Cardiologie Et DpAngiologie</i> , 2018 , 67, 9-13	0.5	1
56	Proteinuria as a Marker for Cardiovascular Disease. <i>Indian Journal of Cardiovascular Disease in Women WINCARS</i> , 2018 , 03, 215-216	0.1	
55	The ratio of AGE to sRAGE independently associated with albuminuria in hypertensive patients. <i>BMC Endocrine Disorders</i> , 2018 , 18, 84	3.3	2

54	Relationships between blood pressure measurements and target organ damage: Data from the Korea women's chest pain registry. <i>Journal of Clinical Hypertension</i> , 2018 , 20, 1724-1730	2.3	3
53	Interaction between Albuminuria and Treatment Group Outcomes for Patients with Renal Artery Stenosis: The NITER Study. <i>Journal of Vascular and Interventional Radiology</i> , 2018 , 29, 966-970	2.4	4
52	Accuracy Evaluation of a Tetrabromophenolphthalein Ethyl Ester Colorimetric Assay for Urinary Albumin. <i>Journal of applied laboratory medicine</i> , 2019 , 4, 201-213	2	2
51	Number of parity is associated with low-grade albuminuria in middle-aged and elderly Chinese women. <i>BMC Women's Health</i> , 2019 , 19, 117	2.9	1
50	Long noncoding RNA UCA1 inhibits ischaemia/reperfusion injury induced cardiomyocytes apoptosis via suppression of endoplasmic reticulum stress. <i>Genes and Genomics</i> , 2019 , 41, 803-810	2.1	13
49	Relation of short-term blood pressure variability to early renal effects in hypertensive patients with controlled blood pressure. <i>Blood Pressure Monitoring</i> , 2019 , 24, 221-224	1.3	9
48	Aortic-flow propagation velocity is associated with proteinuria and left ventricular hypertrophy in newly diagnosed hypertensive patients. <i>Echocardiography</i> , 2019 , 36, 328-335	1.5	1
47	Low-grade albuminuria in pulmonary arterial hypertension. <i>Pulmonary Circulation</i> , 2019 , 9, 20458940188245646	2.5	6
46	Association of High Levels of Spot Urine Protein with High Blood Pressure, Mean Arterial Pressure and Pulse Pressure with the Development of Diabetic Chronic Kidney Dysfunction or Failure among Diabetic Patients. Statistical Regression Modeling to Predict Diabetic Proteinuria. <i>Current Diabetes Reviews</i> , 2019 , 15, 486-496	2.7	4
45	Blood Pressure Control, Left Ventricular Hypertrophy and Treatment Practice Among Hypertensive Patients in Ethiopia. <i>International Journal of General Medicine</i> , 2020 , 13, 903-916	2.3	
44	Prevalence of Albuminuria in Cardiology and Endocrinology Departments and Its Influencing Factors: A Multicenter, Real-World Evidence Study in China. <i>International Journal of Hypertension</i> , 2020 , 2020, 1231593	2.4	1
43	An Observational Registry to Assess Urinary Albumin Evolution in Saudi Hypertensive Patients with the Current Treatment Local algorithm: Results of the RATIONAL Study. <i>International Journal of Nephrology and Renovascular Disease</i> , 2020 , 13, 75-83	2.5	
42	Association between Hypertension, Antihypertensive Drugs, and Osteoporosis in Postmenopausal Syrian Women: A Cross-Sectional Study. <i>Advances in Medicine</i> , 2020 , 2020, 7014212	2.3	3
41	Relationship Between Vascular Aging and Left Ventricular Concentric Geometry in Community-Dwelling Elderly: The Northern Shanghai Study. <i>Clinical Interventions in Aging</i> , 2020 , 15, 853-863	2.4	2
40	Association of Microalbuminuria with High HbA1C levels in Melanesian Adults with Diabetes of at-least 1 Year Duration.		
39	Association of microalbuminuria with left ventricular dysfunction in type 2 diabetes mellitus. <i>The Egyptian Journal of Internal Medicine</i> , 2021 , 33,	0.5	
38	Association of proteinuria and hypertension with incident atrial fibrillation in an elderly population: nationwide data from a community-based elderly cohort. <i>Journal of Hypertension</i> , 2022 , 40, 128-135	1.9	0
37	Treatment of Hypertension in the Elderly Patient With Diabetes. 2005 , 451-469		1

36	Microalbuminuria and arterial stiffness in a general population: the Shimanami Health Promoting Program (J-SHIP) study. <i>Hypertension Research</i> , 2004 , 27, 471-7	4.7	71
35	Low-Grade Albuminuria Is Associated with Metabolic Syndrome and Its Components in Middle-Aged and Elderly Chinese Population. <i>PLoS ONE</i> , 2013 , 8, e65597	3.7	21
34	Prevalence and control of hypertension and albuminuria in South Korea: focus on obesity and abdominal obesity in the Korean National Health and Nutrition Examination Survey, 2011-2012. <i>PLoS ONE</i> , 2014 , 9, e111179	3.7	4
33	Serum gamma - glutamyltransferase is associated with albuminuria: a population-based study. <i>PLoS ONE</i> , 2014 , 9, e114970	3.7	12
32	Moderately increased albuminuria is an independent risk factor of cardiovascular events in the general Japanese population under 75 years of age: the Watari study. <i>PLoS ONE</i> , 2015 , 10, e0123893	3.7	15
31	Funduscopy examination has limited benefit for management of hypertension. <i>International Heart Journal</i> , 2007 , 48, 187-94	1.8	9
30	The association of silent coronary artery disease and metabolic syndrome in Chinese with type 2 diabetes mellitus. <i>Review of Diabetic Studies</i> , 2004 , 1, 18-28	3.6	4
29	Microalbuminuria in pediatric patients with hypertension. <i>Health</i> , 2013 , 05, 40-46	0.4	2
28	The relationship between serum albumin levels and 24-h ambulatory blood pressure monitoring recordings in non-diabetic essential hypertensive patients. <i>Clinics</i> , 2016 , 71, 257-63	2.3	5
27	The Heart and Endocrine Diseases. 2001 , 381-389		
26	Arterial Hypertension in Diabetes: Etiology and Treatment.		
25	Increased Levels of Urinary Albumin: A Cardiovascular Risk Factor and a Target for Treatment. 2010 , 105-116		
24	Microalbuminuria in White and Black Hypertensive Nondiabetic Brazilian Patients. <i>World Journal of Cardiovascular Diseases</i> , 2014 , 04, 236-243	0	
23	MICRO-ALBUMINIURIA AS A PREDICTOR OF LEFT VENTRICULAR HYPERTROPHY IN NON-DIABETIC HYPERTENSIVES. <i>Journal of Evolution of Medical and Dental Sciences</i> , 2015 , 4, 17023-17026	0.1	
22	Postmenopausal Hypertension and Coronary Artery Disease Risk. 2016 , 329-350		
21	Follow-Up of the Hypertensive Patients with Cardiovascular Disease. 2016 , 261-277		
20	RELATIONSHIP OF MICROALBUMINURIA WITH ACUTE MYOCARDIAL INFARCTION IN NONDIABETIC NORMOTENSIVE PATIENTS. <i>Journal of Evidence Based Medicine and Healthcare</i> , 2016 , 3, 4104-4108	0	
19	Atherosclerotic Cardiovascular Disease Short-Term Risk Estimate among Civilian Licensed Aircrew. <i>World Journal of Cardiovascular Diseases</i> , 2019 , 09, 92-108	0	2

18	Relationship between Hypertension and Microalbuminuria according to Obesity Status in Prediabetes. <i>Korean Journal of Health Promotion</i> , 2019 , 19, 202	0.4	1
17	Urine Microalbumin in Obese Comparing with Healthy Children. <i>Caspian Journal of Health Research</i> , 2020 , 5, 8-11	0.1	
16	A STUDY ON ASSOCIATION OF MICROALBUMINURIA WITH ISCHAEMIC HEART DISEASE IN NON-DIABETIC SUBJECTS. 2020 , 31-32		
15	Microalbuminuria and Chronic Kidney Disease as Cardiovascular Risk Factors. 2009 , 149-167		
14	Major Risks Indicators for Diabetic Kidney Disease. 2006 , 351-357		
13	On redefining hypertension. <i>Texas Heart Institute Journal</i> , 2010 , 37, 439-41	0.8	
12	Efficacy of pentoxifylline for reduction of proteinuria in type II diabetic patients. <i>Caspian Journal of Internal Medicine</i> , 2011 , 2, 309-13	1	7
11	Microalbuminuria in untreated prehypertension and hypertension without diabetes. <i>International Journal of Clinical and Experimental Medicine</i> , 2014 , 7, 3420-9		4
10	The relationship between urinary albumin to creatinine ratio and all-cause mortality in the elderly population in the Chinese community: a 10-year follow-up study.. <i>BMC Nephrology</i> , 2022 , 23, 16	2.7	2
9	What is the impact of microvascular complications of diabetes on severe COVID-19?. <i>Microvascular Research</i> , 2021 , 140, 104310	3.7	2
8	Hypertension-mediated organ damage in masked hypertension.. <i>Journal of Hypertension</i> , 2022 , 40,	1.9	0
7	Development of a Risk Model for Predicting Microalbuminuria in the Chinese Population Using Machine Learning Algorithms.. <i>Frontiers in Medicine</i> , 2022 , 9, 775275	4.9	
6	The HARVEST. Looking for optimal management of young people with stage 1 hypertension. <i>Panminerva Medica</i> , 2021 ,	2	0
5	Analysis of cadmium accumulation in community adults and its correlation with low-grade albuminuria.. <i>Science of the Total Environment</i> , 2022 , 155210	10.2	0
4	Home Blood Pressure-based Guidance Did not Increase Anti-albuminuric Effects on Diagnostic Provision of Microalbuminuria in School Workers: A Miyagi Karoshi Prevention Study. 2022 ,		0
3	From Structural to Functional Hypertension Mediated Target Organ Damage—A Long Way to Heart Failure with Preserved Ejection Fraction. 2022 , 11, 5377		0
2	Lipid Parameters and the Development of Chronic Kidney Disease: A Prospective Cohort Study in Middle-Aged and Elderly Chinese Individuals. 2023 , 15, 112		1
1	Effects of sodium-glucose cotransporter 2 inhibitors, mineralocorticoid receptor antagonists, and their combination on albuminuria in diabetic patients.		1

