

Methacrylamidohistidine in affinity ligands for immob chromatography of human serum albumin

Biotechnology and Bioprocess Engineering
6, 402-409

DOI: 10.1007/bf02932321

Citation Report

#	ARTICLE	IF	CITATIONS
1	Metal-chelated polyamide hollow fibers for human serum albumin separation. <i>Journal of Applied Polymer Science</i> , 2002, 86, 3346-3354.	1.3	11
2	Dye Affinity Hollow Fibers for Albumin Purification. <i>Macromolecular Bioscience</i> , 2004, 4, 84-91.	2.1	25
3	IgG purification by negative chromatography in amine-based ligands: A comparison of l-lysine and poly-l-lysine. <i>Process Biochemistry</i> , 2011, 46, 2277-2285.	1.8	18
4	Utilizing a tripeptide conjugated fluorescent hybrid nanoparticles as a fluorescence sensor for the determination of copper ions. <i>Sensors and Actuators A: Physical</i> , 2012, 175, 15-18.	2.0	15
5	Investigation of the effects of Zn ²⁺ , Ca ²⁺ and Na ⁺ ions on the interaction between zonisamide and human serum albumin (HSA) by spectroscopic methods. <i>Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy</i> , 2014, 122, 48-54.	2.0	17
6	Aromatic ligands for plasmid deoxyribonucleic acid chromatographic analysis and purification: An overview. <i>Journal of Chromatography A</i> , 2014, 1327, 1-13.	1.8	11
7	Overview of Albumin and Its Purification Methods. <i>Advanced Pharmaceutical Bulletin</i> , 2016, 6, 495-507.	0.6	96
8	Whole cell based microcontact imprinted capacitive biosensor for the detection of <i>Escherichia coli</i> . <i>Biosensors and Bioelectronics</i> , 2017, 87, 807-815.	5.3	136
9	<scp>l</scp>-Histidine-Derived Smart Antifouling Biohybrid with Multistimuli Responsivity. <i>Biomacromolecules</i> , 2021, 22, 3941-3949.	2.6	9