

CITATION REPORT

List of articles citing

A lipopolysaccharide-deficient mutant of *Neisseria meningitidis* elicits attenuated cytokine release by human macrophages and signals via toll-like receptor (TLR) 2 but not via TLR4/MD2

DOI: 10.1086/317647

Journal of Infectious Diseases, 2001, 183, 89-96.

Source: <https://exaly.com/paper-pdf/33362987/citation-report.pdf>

Version: 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
148	Invited review: Bacterial lipopolysaccharides and innate immunity. 2001 , 7, 167-202		123
147	Toll receptors and sepsis. 2001 , 7, 371-5		20
146	Mannose-binding lectin regulates the inflammatory response of human professional phagocytes to <i>Neisseria meningitidis</i> serogroup B. <i>Journal of Infectious Diseases</i> , 2001 , 184, 1152-62	7	161
145	Interaction of <i>Neisseria meningitidis</i> with human dendritic cells. <i>Infection and Immunity</i> , 2001 , 69, 6912-22	3.7	45
144	A functional polymorphism of toll-like receptor 4 is not associated with likelihood or severity of meningococcal disease. <i>Journal of Infectious Diseases</i> , 2001 , 184, 640-2	7	173
143	Invited review: <i>Neisseria meningitidis</i> lipopolysaccharides in human pathology. 2001 , 7, 401-420		16
142	Cutting edge: Immune stimulation by neisserial porins is toll-like receptor 2 and MyD88 dependent. <i>Journal of Immunology</i> , 2002 , 168, 1533-7	5.3	261
141	The role of toll-like receptor (TLR) 2 and TLR4 in the host defense against disseminated candidiasis. <i>Journal of Infectious Diseases</i> , 2002 , 185, 1483-9	7	388
140	Platelet-platelet and platelet-leukocyte interactions induced by outer membrane vesicles from <i>N. meningitidis</i> . 2002 , 13, 91-9		31
139	Lipooligosaccharide and polysaccharide capsule: virulence factors of <i>Neisseria meningitidis</i> that determine meningococcal interaction with human dendritic cells. <i>Infection and Immunity</i> , 2002 , 70, 2454-62	3.7	66
138	The clinical relevance of endotoxin in human sepsis: a critical analysis. 2002 , 8, 473-476		24
137	Meningococcal disease: treatment and prevention. 2002 , 34, 624-34		14
136	The class A macrophage scavenger receptor is a major pattern recognition receptor for <i>Neisseria meningitidis</i> which is independent of lipopolysaccharide and not required for secretory responses. <i>Infection and Immunity</i> , 2002 , 70, 5346-54	3.7	231
135	Current concepts in the role of the host response in <i>Neisseria meningitidis</i> septic shock. 2002 , 15, 247-52		40
134	Toll-like receptors. <i>Critical Care Medicine</i> , 2002 , 30, S1-S11	1.4	210
133	Bench-to-bedside review: genetic influences on meningococcal disease. 2002 , 6, 60-5		11
132	Initial responses to endotoxins and Gram-negative bacteria. 2002 , 323, 59-72		264

131	Lipopolysaccharide endotoxins. 2002 , 71, 635-700		3259
130	[Physiopathology and curative treatment of meningococcal diseases: current aspects]. 2002 , 50, 613-9		1
129	Non-LPS components of Chlamydia pneumoniae stimulate cytokine production through Toll-like receptor 2-dependent pathways. 2002 , 32, 1188-95		96
128	Responsiveness of intestinal epithelial cell lines to lipopolysaccharide is correlated with Toll-like receptor 4 but not Toll-like receptor 2 or CD14 expression. 2003 , 18, 25-32		97
127	Systemic responsiveness to lipopolysaccharide and polymorphisms in the toll-like receptor 4 gene in human beings. 2003 , 112, 923-9		122
126	The dual role of LBP and CD14 in response to Gram-negative bacteria or Gram-negative compounds. 2003 , 9, 381-384		17
125	Identification of meningococcal LPS as a major monocyte activator in IL-10 depleted shock plasmas and CSF by blocking the CD14-TLR4 receptor complex. 2003 , 9, 155-163		17
124	TOLL-like receptors linking innate and adaptive immune response. 2003 , 91, 1-12		300
123	The role of human dendritic cells in meningococcal and listerial meningitis. <i>International Journal of Medical Microbiology</i> , 2003 , 293, 241-9	3.7	19
122	Toll-like receptors and their function in innate and adaptive immunity. 2003 , 130, 180-92		127
121	CD14- and Toll-like receptor-dependent activation of bladder epithelial cells by lipopolysaccharide and type 1 piliated Escherichia coli. <i>Infection and Immunity</i> , 2003 , 71, 1470-80	3.7	123
120	Activation of toll-like receptor 2 (TLR2) and TLR4/MD2 by Neisseria is independent of capsule and lipooligosaccharide (LOS) sialylation but varies widely among LOS from different strains. <i>Infection and Immunity</i> , 2003 , 71, 3901-8	3.7	58
119	Transforming growth factor-beta, interleukin (IL)-18, and IL-12: effect on the clinical course and complications of Plasmodium falciparum malaria. <i>Journal of Infectious Diseases</i> , 2003 , 187, 338-9; author reply 339-40	7	7
118	The immunopathogenesis of meningococcal disease. 2003 , 56, 417-22		20
117	Lessons from an LPS-deficient Neisseria meningitidis mutant. 2003 , 9, 124-128		16
116	Lipooligosaccharide-deficient Neisseria meningitidis shows altered pilus-associated characteristics. <i>Infection and Immunity</i> , 2003 , 71, 155-62	3.7	54
115	Activation of human dendritic cells is modulated by components of the outer membranes of Neisseria meningitidis. <i>Infection and Immunity</i> , 2003 , 71, 5590-7	3.7	28
114	Reply. <i>Journal of Infectious Diseases</i> , 2003 , 187, 342-343	7	9

113	The role that the functional Asp299Gly polymorphism in the toll-like receptor-4 gene plays in susceptibility to Chlamydia trachomatis-associated tubal infertility. <i>Journal of Infectious Diseases</i> , 2003 , 187, 341-2; author reply 342-3	7	41
112	Clinical value of adjunctive interleukin-2: clarifications and corrections. <i>Journal of Infectious Diseases</i> , 2003 , 187, 340; author reply 340-1	7	
111	Penicillin enhances the toll-like receptor 2-mediated proinflammatory activity of Streptococcus pneumoniae. <i>Journal of Infectious Diseases</i> , 2003 , 188, 1040-8	7	26
110	Reply. <i>Journal of Infectious Diseases</i> , 2003 , 187, 339-340	7	
109	The Lip lipoprotein from Neisseria gonorrhoeae stimulates cytokine release and NF-kappaB activation in epithelial cells in a Toll-like receptor 2-dependent manner. 2003 , 278, 46252-60		83
108	The dual role of LBP and CD14 in response to Gram-negative bacteria or Gram-negative compounds. 2003 , 9, 381-4		15
107	Lessons from an LPS-deficient Neisseria meningitidis mutant. 2003 , 9, 124-8		15
106	Identification of meningococcal LPS as a major monocyte activator in IL-10 depleted shock plasmas and CSF by blocking the CD14-TLR4 receptor complex. 2003 , 9, 155-63		16
105	Reply. <i>Journal of Infectious Diseases</i> , 2003 , 187, 340-341	7	65
104	Endotoxin as a drug target. <i>Critical Care Medicine</i> , 2003 , 31, S57-64	1.4	81
103	Teasing apart structural determinants of toxicity and adjuvanticity: implications for meningococcal vaccine development. 2004 , 10, 113-119		2
102	Lipopolysaccharides of Bacteroides fragilis, Chlamydia trachomatis and Pseudomonas aeruginosa signal via toll-like receptor 2. 2004 , 53, 735-740		147
101	Complement activation and complement-dependent inflammation by Neisseria meningitidis are independent of lipopolysaccharide. <i>Infection and Immunity</i> , 2004 , 72, 3344-9	3.7	61
100	Activation of murine macrophages by Neisseria meningitidis and IFN-gamma in vitro: distinct roles of class A scavenger and Toll-like pattern recognition receptors in selective modulation of surface phenotype. <i>Journal of Leukocyte Biology</i> , 2004 , 76, 577-84	6.5	42
99	Teasing apart structural determinants of toxicity and adjuvanticity: implications for meningococcal vaccine development. 2004 , 10, 113-9		30
98	High-level endothelial E-selectin (CD62E) cell adhesion molecule expression by a lipopolysaccharide-deficient strain of Neisseria meningitidis despite poor activation of NF-kappaB transcription factor. 2004 , 135, 85-93		18
97	Neisseria meningitidis-induced death of cerebrovascular endothelium: mechanisms triggering transcriptional activation of inducible nitric oxide synthase. 2004 , 89, 1166-74		28
96	Toll-like receptor 2 (TLR2) and TLR4 are present inside human dendritic cells, associated with microtubules and the Golgi apparatus but are not detectable on the cell surface: integrity of microtubules is required for interleukin-12 production in response to internalized bacteria. 2004 , 111, 173-8		92

95	Mannose binding lectin enhances IL-1beta and IL-10 induction by non-lipopolysaccharide (LPS) components of <i>Neisseria meningitidis</i> . 2004 , 28, 59-66		21
94	[Pathophysiological role of endotoxins, a common denominator to various diseases]. 2004 , 52, 415-22		2
93	Innate immunity and toll-like receptors: clinical implications of basic science research. 2004 , 144, 421-9		146
92	Differentiation of <i>Campylobacter coli</i> , <i>Campylobacter jejuni</i> , <i>Campylobacter lari</i> , and <i>Campylobacter upsaliensis</i> by a multiplex PCR developed from the nucleotide sequence of the lipid A gene <i>lpxA</i> . 2004 , 42, 5549-57		142
91	Human models of endotoxemia and recombinant human activated protein C. <i>Critical Care Medicine</i> , 2004 , 32, S202-8	1.4	36
90	Lipopolysaccharide heterogeneity: innate host responses to bacterial modification of lipid a structure. 2005 , 84, 584-95		159
89	Activation of human meningeal cells is modulated by lipopolysaccharide (LPS) and non-LPS components of <i>Neisseria meningitidis</i> and is independent of Toll-like receptor (TLR)4 and TLR2 signalling. <i>Cellular Microbiology</i> , 2005 , 7, 415-30	3.9	38
88	Lipooligosaccharide-independent alteration of cellular homeostasis in <i>Neisseria meningitidis</i> -infected epithelial cells. <i>Cellular Microbiology</i> , 2005 , 7, 869-85	3.9	11
87	Activation of hypoxia-inducible factor-1 in bacillary angiomatosis: evidence for a role of hypoxia-inducible factor-1 in bacterial infections. 2005 , 111, 1054-62		124
86	<i>Neisseria gonorrhoeae</i> enhances infection of dendritic cells by HIV type 1. <i>Journal of Immunology</i> , 2005 , 174, 7995-8002	5.3	58
85	Chemokine patterns in meningococcal disease. <i>Journal of Infectious Diseases</i> , 2005 , 191, 768-75	7	80
84	Differential involvement of toll-like receptors 2 and 4 in the host response to acute respiratory infections with wild-type and mutant <i>Haemophilus influenzae</i> strains. <i>Infection and Immunity</i> , 2005 , 73, 2075-82	3.7	29
83	Genetic polymorphisms in sepsis. 2005 , 6, S61-73		78
82	Single nucleotide polymorphisms of Toll-like receptors and susceptibility to infectious disease. 2005 , 5, 156-64		387
81	Meningococcal infections at the start of the 21st century. 2005 , 52, 129-62		19
80	Recognition of bacterial products by toll-like receptors. 2005 , 86, 99-119		32
79	Toll-like receptors: applications to dermatologic disease. 2006 , 54, 951-83; quiz 983-6		78
78	Expression of TOLL-like receptors (TLR) by bovine antigen-presenting cells-potential role in pathogen discrimination?. 2006 , 112, 2-11		79

77	Heterologous prime-boost strategy to overcome weak immunogenicity of two serosubtypes in hexavalent <i>Neisseria meningitidis</i> outer membrane vesicle vaccine. <i>Vaccine</i> , 2006 , 24, 1569-77	4.1	11
76	. 2006 ,		7
75	MyD88-dependent signaling affects the development of meningococcal sepsis by nonlipooligosaccharide ligands. <i>Infection and Immunity</i> , 2006 , 74, 3538-46	3.7	15
74	Cellular Immune Responses in Meningococcal Disease. 2006 , 295-320		1
73	Pathogenesis and Pathophysiology of Invasive Meningococcal Disease. 2006 , 427-480		16
72	<i>Neisseria meningitidis</i> expressing lgtB lipopolysaccharide targets DC-SIGN and modulates dendritic cell function. <i>Cellular Microbiology</i> , 2006 , 8, 316-25	3.9	73
71	Neurocysticercal antigens stimulate chemokine secretion from human monocytes via an NF-kappaB-dependent pathway. <i>Microbes and Infection</i> , 2006 , 8, 1732-40	9.3	11
70	TLR2 and TLR4 expression during bacterial infections. 2006 , 12, 4185-93		35
69	Differential role of lipooligosaccharide of <i>Neisseria meningitidis</i> in virulence and inflammatory response during respiratory infection in mice. <i>Infection and Immunity</i> , 2006 , 74, 5506-12	3.7	17
68	Meningococcal porin PorB binds to TLR2 and requires TLR1 for signaling. <i>Journal of Immunology</i> , 2006 , 176, 2373-80	5.3	128
67	Identification of <i>Neisseria meningitidis</i> nonlipopolysaccharide ligands for class A macrophage scavenger receptor by using a novel assay. <i>Infection and Immunity</i> , 2006 , 74, 5191-9	3.7	47
66	TNF-alpha-independent IL-8 expression: alterations in bacterial challenge dose cause differential human monocytic cytokine response. <i>Journal of Immunology</i> , 2006 , 177, 1314-22	5.3	13
65	Agonists of Toll-like receptors 3, 4, 7, and 9 are candidates for use as adjuvants in an outer membrane vaccine against <i>Neisseria meningitidis</i> serogroup B. <i>Infection and Immunity</i> , 2007 , 75, 5939-46	3.7	47
64	A critical contribution of both CD28 and ICOS in the adjuvant activity of <i>Neisseria meningitidis</i> H44/76 LPS and lpxL1 LPS. <i>Vaccine</i> , 2007 , 25, 4681-8	4.1	4
63	The host response to endotoxin, antilipopolysaccharide strategies, and the management of severe sepsis. <i>International Journal of Medical Microbiology</i> , 2007 , 297, 365-77	3.7	185
62	Establishment of a newly improved detection system for NF-kappaB activity. 2007 , 109, 175-81		10
61	Non-lipooligosaccharide-mediated signalling via Toll-like receptor 4 causes fatal meningococcal sepsis in a mouse model. <i>Cellular Microbiology</i> , 2007 , 9, 657-69	3.9	7
60	Regulation of Th-1 T cell-dominated immunity to <i>Neisseria meningitidis</i> within the human mucosa. <i>Cellular Microbiology</i> , 2007 , 9, 1050-61	3.9	26

59	Neisseria meningitidis type C capsular polysaccharide inhibits lipooligosaccharide-induced cell activation by binding to CD14. <i>Cellular Microbiology</i> , 2007 , 9, 1297-310	3.9	17
58	Innate-immunity cytokines induced by very small size proteoliposomes, a Neisseria-derived immunological adjuvant. 2007 , 147, 379-88		9
57	Reverse transcriptase-PCR differential display analysis of meningococcal transcripts during infection of human cells: up-regulation of priA and its role in intracellular replication. 2008 , 8, 131		16
56	The response of HEK293 cells transfected with bovine TLR2 to established pathogen-associated molecular patterns and to bacteria causing mastitis in cattle. 2008 , 125, 326-36		22
55	Investigation of lipopolysaccharide receptor expression on human monocytes after major orthopaedic surgery. 2008 , 40, 239-45		4
54	Identification of genes particularly sensitive to lipopolysaccharide (LPS) in human monocytes induced by wild-type versus LPS-deficient Neisseria meningitidis strains. <i>Infection and Immunity</i> , 2008 , 76, 2685-95	3.7	36
53	Mucosal immunity in healthy adults after parenteral vaccination with outer-membrane vesicles from Neisseria meningitidis serogroup B. <i>Journal of Infectious Diseases</i> , 2008 , 198, 731-40	7	32
52	Stages of meningococcal sepsis simulated in vitro, with emphasis on complement and Toll-like receptor activation. <i>Infection and Immunity</i> , 2008 , 76, 4183-9	3.7	35
51	Increased surface toll-like receptor 2 expression in superantigen shock. <i>Critical Care Medicine</i> , 2008 , 36, 1267-76	1.4	18
50	Effect of a single dose of tobramycin on systemic inflammatory response-induced acute kidney injury in a 6-hour porcine model *. <i>Critical Care Medicine</i> , 2009 , 37, 2782-2790	1.4	0
49	Recognition of meningococcal molecular patterns by innate immune receptors. <i>International Journal of Medical Microbiology</i> , 2009 , 299, 9-20	3.7	8
48	Toll-like receptors in bacterial meningitis. <i>Current Topics in Microbiology and Immunology</i> , 2009 , 336, 15-40	3.3	35
47	Toll-like receptors: methods and protocols.Preface. <i>Methods in Molecular Biology</i> , 2009 , 517, v-vi	1.4	
46	Toll-like Receptors: Roles in Infection and Neuropathology. <i>Current Topics in Microbiology and Immunology</i> , 2009 ,	3.3	3
45	Science and Practice of Pediatric Critical Care Medicine. 2009 ,		1
44	Effect of a single dose of tobramycin on systemic inflammatory response-induced acute kidney injury in a 6-hour porcine model. <i>Critical Care Medicine</i> , 2009 , 37, 2782-90	1.4	16
43	The tumor necrosis factor polymorphism TNF (-308) is associated with susceptibility to meningococcal sepsis, but not with lethality. <i>Critical Care Medicine</i> , 2009 , 37, 1237-43	1.4	37
42	Inflammatory and circulatory effects of the reduction of endotoxin concentration in established porcine endotoxemic shock—a model of endotoxin elimination. <i>Critical Care Medicine</i> , 2009 , 37, 1031-e4	1.4	15

41	Innate immune activation of swine intestinal epithelial cell lines (IPEC-J2 and IPI-2I) in response to LPS from <i>Salmonella typhimurium</i> . <i>Comparative Immunology, Microbiology and Infectious Diseases</i> , 2010 , 33, 161-74	2.6	78
40	Structural analysis of lipopolysaccharides from Gram-negative bacteria. <i>Biochemistry (Moscow)</i> , 2010 , 75, 383-404	2.9	50
39	Phosphoryl moieties of lipid A from <i>Neisseria meningitidis</i> and <i>N. gonorrhoeae</i> lipooligosaccharides play an important role in activation of both MyD88- and TRIF-dependent TLR4-MD-2 signaling pathways. <i>Journal of Immunology</i> , 2010 , 185, 6974-84	5.3	46
38	TLR4-mediated podosome loss discriminates gram-negative from gram-positive bacteria in their capacity to induce dendritic cell migration and maturation. <i>Journal of Immunology</i> , 2010 , 184, 1280-91	5.3	24
37	Mechanisms of polymyxin B endotoxin removal from extracorporeal blood flow: molecular interactions. <i>Contributions To Nephrology</i> , 2010 , 167, 45-54	1.6	7
36	Endotoxins and other sepsis triggers. <i>Contributions To Nephrology</i> , 2010 , 167, 14-24	1.6	149
35	Improved OMV vaccine against <i>Neisseria meningitidis</i> using genetically engineered strains and a detergent-free purification process. <i>Vaccine</i> , 2010 , 28, 4810-6	4.1	110
34	Pneumolysin-induced CXCL8 production by nasopharyngeal epithelial cells is dependent on calcium flux and MAPK activation via Toll-like receptor 4. <i>Microbes and Infection</i> , 2011 , 13, 65-75	9.3	17
33	Studies on Experimental Models. 2011 ,		
32	Nuclear receptor signaling inhibits HIV-1 replication in macrophages through multiple trans-repression mechanisms. <i>Journal of Virology</i> , 2011 , 85, 10834-50	6.6	33
31	<i>Neisseria meningitidis</i> capsular polysaccharides induce inflammatory responses via TLR2 and TLR4-MD-2. <i>Journal of Leukocyte Biology</i> , 2011 , 89, 469-80	6.5	51
30	Vigorous response of human innate functioning IgM memory B cells upon infection by <i>Neisseria gonorrhoeae</i> . <i>Journal of Immunology</i> , 2012 , 188, 4008-22	5.3	17
29	The nature of an in vivo anti-capsular polysaccharide response is markedly influenced by the composition and/or architecture of the bacterial subcapsular domain. <i>Journal of Immunology</i> , 2012 , 188, 569-77	5.3	16
28	Classification and pathogenesis of meningococcal infections. <i>Methods in Molecular Biology</i> , 2012 , 799, 21-35	1.4	60
27	Species-specific PAMP recognition by TLR2 and evidence for species-restricted interaction with Dectin-1. <i>Journal of Leukocyte Biology</i> , 2013 , 94, 449-58	6.5	34
26	Lipopolysaccharide modifications of a cholera vaccine candidate based on outer membrane vesicles reduce endotoxicity and reveal the major protective antigen. <i>Infection and Immunity</i> , 2013 , 81, 2379-93	3.7	47
25	Lipopolysaccharide-deficient <i>Acinetobacter baumannii</i> shows altered signaling through host Toll-like receptors and increased susceptibility to the host antimicrobial peptide LL-37. <i>Infection and Immunity</i> , 2013 , 81, 684-9	3.7	49
24	The biology of <i>Neisseria adhesins</i> . <i>Biology</i> , 2013 , 2, 1054-109	4.9	30

23	Hexa-acylated lipid A is required for host inflammatory response to <i>Neisseria gonorrhoeae</i> in experimental gonorrhea. <i>Infection and Immunity</i> , 2014 , 82, 184-92	3.7	19
22	<i>Neisseria meningitidis</i> elicits a pro-inflammatory response involving I κ B α in a human blood-cerebrospinal fluid barrier model. <i>Journal of Neuroinflammation</i> , 2014 , 11, 163	10.1	25
21	Activation of NOD receptors by <i>Neisseria gonorrhoeae</i> modulates the innate immune response. <i>Innate Immunity</i> , 2014 , 20, 377-89	2.7	24
20	Different cytokine production and Toll-like receptor expression induced by heat-killed invasive and carrier strains of <i>Neisseria meningitidis</i> . <i>Apmis</i> , 2014 , 122, 33-41	3.4	3
19	How the Knowledge of Interactions between Meningococcus and the Human Immune System Has Been Used to Prepare Effective <i>Neisseria meningitidis</i> Vaccines. <i>Journal of Immunology Research</i> , 2015 , 2015, 189153	4.5	15
18	Dual 3Seq using deepSuperSAGE uncovers transcriptomes of interacting <i>Salmonella enterica</i> Typhimurium and human host cells. <i>BMC Genomics</i> , 2015 , 16, 323	4.5	11
17	Outer membrane vesicles derived from <i>Salmonella</i> Typhimurium mutants with truncated LPS induce cross-protective immune responses against infection of <i>Salmonella enterica</i> serovars in the mouse model. <i>International Journal of Medical Microbiology</i> , 2016 , 306, 697-706	3.7	32
16	A critical role for ATF2 transcription factor in the regulation of E-selectin expression in response to non-endotoxin components of <i>Neisseria meningitidis</i> . <i>Cellular Microbiology</i> , 2016 , 18, 66-79	3.9	4
15	Innate immune recognition and inflammation in <i>Neisseria meningitidis</i> infection. <i>Pathogens and Disease</i> , 2017 , 75,	4.2	15
14	Neuronal Damage and Neuroinflammation, a Bridge Between Bacterial Meningitis and Neurodegenerative Diseases. <i>Frontiers in Cellular Neuroscience</i> , 2021 , 15, 680858	6.1	5
13	Severe Sepsis and Septic Shock. 2008 , 287-304		1
12	Toll-Like Receptors: Ligands, Cell-Based Models, and Readouts for Receptor Action. <i>Methods in Molecular Biology</i> , 2016 , 1390, 3-27	1.4	15
11	Ligands, cell-based models, and readouts required for Toll-like receptor action. <i>Methods in Molecular Biology</i> , 2009 , 517, 15-32	1.4	5
10	Effect of the administration rate on the biological responses to a fixed dose of endotoxin in the anesthetized pig. <i>Shock</i> , 2008 , 29, 173-80	3.4	17
9	Role of Toll-Like Receptors in Innate Immunity of the Intestine. 79-94		1
8	Differential effect of TLR2 and TLR4 on the immune response after immunization with a vaccine against <i>Neisseria meningitidis</i> or <i>Bordetella pertussis</i> . <i>PLoS ONE</i> , 2010 , 5, e15692	3.7	33
7	Death in a Lonely Place: Pathophysiology of the Dying Patient. 2008 , 45-64		
6	Genetic Polymorphisms in Critical Care and Illness. 2009 , 1-16		

5 Endotoxin Recognition. **2009**, 767-773

4 Oxidative Stress in Animal Models with Special Reference to Experimental Porcine Endotoxemia. **2011**, 497-510

3 Toll-Like Receptors and Neuroinflammation. *Advances in Neurobiology*, **2013**, 135-156 2.1

2 The Host-Pathogen Interactions and Epicellular Lifestyle of .. *Frontiers in Cellular and Infection Microbiology*, **2022**, 12, 862935 5.9 1

1 Biological functions of sialic acid as a component of bacterial endotoxin. 13, 0