CITATION REPORT List of articles citing

Association of C-reactive protein with carotid atherosclerosis in men and women: the Framingham Heart Study

DOI: 10.1161/01.atv.0000034543.78801.69 Arteriosclerosis, Thrombosis, and Vascular Biology, 2002, 22, 1662-7.

Source: https://exaly.com/paper-pdf/33648259/citation-report.pdf

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| # | Paper | IF | Citations |
|-----|---|------|-----------|
| 203 | C-reactive protein, subclinical atherosclerosis, and risk of cardiovascular events. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2002 , 22, 1512-3 | 9.4 | 38 |
| 202 | High-sensitivity C-reactive protein in high-grade carotid stenosis: risk marker for unstable carotid plaque. <i>Journal of Vascular Surgery</i> , 2003 , 38, 1018-24 | 3.5 | 89 |
| 201 | C-reactive protein, carotid intima-media thickness, and incidence of ischemic stroke in the elderly: the Cardiovascular Health Study. 2003 , 108, 166-70 | | 267 |
| 200 | C-reactive protein and coronary artery calcification: The Study of Inherited Risk of Coronary Atherosclerosis (SIRCA). <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2003 , 23, 1851-6 | 9.4 | 84 |
| 199 | Current World Literature. 2003, 14, 621-628 | | |
| 198 | Lipid metabolism: inflammatory-immune responses in atherosclerosis. 2003, 14, 337-9 | | 4 |
| 197 | Update on vascular disease in systemic lupus erythematosus. 2003 , 15, 519-27 | | 59 |
| 196 | C-reactive protein and interleukin-6 in vascular disease: culprits or passive bystanders?. 2003 , 21, 1787 | -803 | 133 |
| 195 | Current World Literature. 2003, 14, 513-527 | | |
| 194 | New risk factors for atherosclerosis and patient risk assessment. 2004 , 109, III15-9 | | 181 |
| 193 | Should C-reactive protein be added to metabolic syndrome and to assessment of global cardiovascular risk?. 2004 , 109, 2818-25 | | 508 |
| 192 | Atherosclerotic Vascular Disease Conference: Writing Group I: epidemiology. 2004 , 109, 2605-12 | | 142 |
| 191 | The relationship between C-reactive protein and carotid intima-media wall thickness in middle-aged women with polycystic ovary syndrome. 2004 , 89, 6061-7 | | 82 |
| 190 | CagA-positive Helicobacter pylori strains may influence the natural history of atherosclerotic stroke. 2004 , 63, 800-4 | | 39 |
| 189 | Surrogate markers for cardiovascular disease: structural markers. 2004 , 109, IV22-30 | | 146 |
| 188 | Monocyte count, but not C-reactive protein or interleukin-6, is an independent risk marker for subclinical carotid atherosclerosis. 2004 , 35, 1619-24 | | 165 |
| 187 | Pre-procedural plasma levels of C-reactive protein and interleukin-6 do not predict late coronary angiographic restenosis after elective stenting. 2004 , 25, 1029-35 | | 30 |

| 186 | Autoimmune and inflammatory mechanisms in atherosclerosis. 2004 , 22, 361-403 | | 344 |
|-----|---|-----|-----|
| 185 | Circulating tissue kallikrein levels correlate with severity of carotid atherosclerosis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology,</i> 2004 , 24, 1104-10 | 9.4 | 15 |
| 184 | Strong association of C-reactive protein with body mass index and 2-h post-challenge glucose in non-diabetic, non-smoker subjects without hypertension. 2004 , 21, 581-5 | | 37 |
| 183 | High-sensitivity C-reactive protein: Clinical importance. 2004 , 29, 439-493 | | 14 |
| 182 | C-reactive protein, the metabolic syndrome, and prediction of cardiovascular events in the Framingham Offspring Study. 2004 , 110, 380-5 | | 514 |
| 181 | In vivo magnetic resonance evaluation of associations between aortic atherosclerosis and both risk factors and coronary artery disease in patients referred for coronary angiography. 2004 , 148, 137-43 | | 64 |
| 180 | Inflammation, infection, and aortic valve sclerosis; Insights from the Olmsted County (Minnesota) population. 2004 , 174, 337-42 | | 44 |
| 179 | The relationship between high-sensitive C-reactive protein and pulse wave velocity in healthy Japanese men. 2004 , 174, 373-7 | | 34 |
| 178 | New and confirmatory evidence of an association between APOE genotype and baseline C-reactive protein in dyslipidemic individuals. 2004 , 177, 345-51 | | 58 |
| 177 | High-sensitivity C-reactive protein: Clinical importance. 2004 , 29, 439-493 | | 30 |
| 176 | Carotid IMT: A Surrogate Marker for Cerebrovascular Disease?. 2005 , 5, 66-73 | | О |
| 175 | Gender-specific association between pulse pressure and C-reactive protein in a Chinese population. 2005 , 19, 293-9 | | 9 |
| 174 | Comprehensive endovascular therapy for femoropopliteal arterial atherosclerotic occlusive disease. 2005 , 201, 275-96 | | 32 |
| 173 | Atherosclerosis as inflammation. 2005 , 19, 130-8 | | 71 |
| 172 | Oxidant stress, inflammation and atherogenesis. 2005 , 14, 760-4 | | 77 |
| 171 | Inflammation and Carotid ArteryRisk for Atherosclerosis Study (ICARAS). 2005 , 111, 2203-9 | | 178 |
| 170 | C-reactive protein and the 10-year incidence of coronary heart disease in older men and women: the cardiovascular health study. 2005 , 112, 25-31 | | 289 |
| 169 | C-reactive protein, interleukin-6, and soluble adhesion molecules as predictors of progressive peripheral atherosclerosis in the general population: Edinburgh Artery Study. 2005 , 112, 976-83 | | 328 |

| 168 | Is high-sensitivity C-reactive protein associated with carotid atherosclerosis in healthy Koreans?. 2005 , 12, 548-554 | | 3 |
|-----|---|-----|-----|
| 167 | Increased carotid intima-media thickness and serum inflammatory markers in obstructive sleep apnea. 2005 , 172, 625-30 | | 317 |
| 166 | Lifecourse socioeconomic position, C-reactive protein, and carotid intima-media thickness in young adults: the cardiovascular risk in Young Finns Study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2005 , 25, 2197-202 | 9.4 | 72 |
| 165 | Elevated serum C-reactive protein levels and advanced atherosclerosis in youth. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2005 , 25, 1237-43 | 9.4 | 52 |
| 164 | Defects in regulation of local immune responses resulting in atherosclerosis. 2005 , 12, 225-34 | | 13 |
| 163 | Associations of serum IL-18 levels with carotid intima-media thickness. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2005 , 25, 1458-62 | 9.4 | 97 |
| 162 | Soluble CD40L and cardiovascular risk in asymptomatic low-grade carotid stenosis. 2005 , 36, 673-5 | | 54 |
| 161 | The association of C-reactive protein levels with carotid intima-media complex thickness and plaque formation in the general population. 2005 , 36, 2138-42 | | 98 |
| 160 | Interleukin-2 levels are associated with carotid artery intima-media thickness. 2005, 180, 181-7 | | 47 |
| 159 | Brief review and critical examination of the use of hs-CRP for cardiac risk assessment with the conclusion that it is premature to use this test. 2005 , 356, 1-8 | | 14 |
| 158 | Hormone therapy, C-reactive protein, and progression of atherosclerosis: data from the Estrogen Replacement on Progression of Coronary Artery Atherosclerosis (ERA) trial. 2005 , 150, 907-11 | | 28 |
| 157 | The relationship between C-reactive protein and subclinical cardiovascular disease in the Diabetes Heart Study (DHS). 2005 , 150, 1032-8 | | 30 |
| 156 | Cardiovascular Biomarkers. 2006, | | 2 |
| 155 | Inflammatory and long-term risk markers. 2006 , 26, 553-70, v | | 8 |
| 154 | The relationship between plasma levels of oxidized and reduced thiols and early atherosclerosis in healthy adults. 2006 , 47, 1005-11 | | 180 |
| 153 | Rosiglitazone improves insulin sensitivity in nonobese subjects with impaired glucose tolerance: the role of adiponectin and C-reactive protein. 2006 , 55, 439-44 | | 16 |
| 152 | What does minor elevation of C-reactive protein signify?. 2006 , 119, 166.e17-28 | | 244 |
| 151 | Epidemiology of carotid artery atherosclerosis. 22-34 | | |

Monitoring pharmaceutical interventions with conventional ultrasound (IMT). 430-450

| 149 | High-sensitivity C-reactive protein is only weakly related to cardiovascular damage after adjustment for traditional cardiovascular risk factors. 2006 , 24, 655-61 | 32 |
|-----|---|-----|
| 148 | PAI-1 Gene 4G/5G polymorphism and risk of type 2 diabetes in a population-based sample. 2006 , 14, 753-8 | 24 |
| 147 | Rheumatoid arthritis: links with cardiovascular disease and the receptor for advanced glycation end products. 2006 , 156, 42-52 | 24 |
| 146 | Diagnostic tools for the study of vascular cognitive dysfunction in hypertension and antihypertensive drug research. 2006 , 109, 274-83 | 14 |
| 145 | C-reactive protein's place on the cardiovascular stage: prima ballerina or chorus girl?. 2006 , 24, 627-32 | 6 |
| 144 | Systemic inflammatory markers and risk of dementia. 2006 , 21, 258-62 | 64 |
| 143 | Cardiovascular risk reduction in high-risk pediatric patients: a scientific statement from the American Heart Association Expert Panel on Population and Prevention Science; the Councils on Cardiovascular Disease in the Young, Epidemiology and Prevention, Nutrition, Physical Activity and | 524 |
| 142 | Variants at the APOA5 locus, association with carotid atherosclerosis, and modification by obesity: the Framingham Study. 2006 , 47, 990-6 | 59 |
| 141 | C-Reactive Protein as a Tool for Risk Assessment in Primary Prevention. 2006 , 237-260 | |
| 140 | Correlation of coronary and cerebral atherosclerosis: difference between extracranial and intracranial arteries. 2006 , 21, 112-9 | 50 |
| 139 | Hemostatic markers of endothelial dysfunction and risk of incident type 2 diabetes: the Framingham Offspring Study. 2006 , 55, 530-7 | 138 |
| 138 | Serum chitotriosidase activity, a marker of activated macrophages, predicts new cardiovascular events independently of C-reactive protein. 2007 , 108, 297-306 | 28 |
| 137 | N-terminal pro-brain natriuretic peptide, but not high sensitivity C-reactive protein, improves cardiovascular risk prediction in the general population. 2007 , 28, 1374-81 | 109 |
| 136 | High-sensitivity C-reactive protein is not associated with carotid intima-media progression: the carotid atherosclerosis progression study. 2007 , 38, 1774-9 | 66 |
| 135 | Elevated inflammation markers in pheochromocytoma compared to other forms of hypertension. 2007 , 14, 57-64 | 30 |
| 134 | Low-grade inflammation is a risk factor for clinical stroke events in addition to silent cerebral infarcts in Japanese older hypertensives: the Jichi Medical School ABPM Study, wave 1. 2007 , 38, 911-7 | 58 |
| 133 | High-sensitivity C-reactive protein is independently associated with early carotid artery progression in women but not in men: the INVADE Study. 2007 , 38, 2881-6 | 32 |
| | | |

| 132 | Asymptomatic Carotid Artery Stenosis. 2007, | 13 |
|--------------------------|--|---------------------|
| 131 | Cardiovascular risk reduction in high-risk pediatric patients: a scientific statement from the American Heart Association Expert Panel on Population and Prevention Science; the Councils on Cardiovascular Disease in the Young, Epidemiology and Prevention, Nutrition, Physical Activity and | 41 |
| 130 | C-reactive protein, inflammatory conditions, and cardiovascular disease risk. 2007 , 120, 1054-62 | 75 |
| 129 | Effects of IL-6, adiponectin, CRP and metabolic syndrome on subclinical atherosclerosis. 2007 , 384, 99-104 | 65 |
| 128 | Both cyclooxygenase- and cytokine-mediated inflammation are associated with carotid intima-media thickness. 2007 , 38, 130-6 | 35 |
| 127 | C-reactive protein and features of metabolic syndrome in Brazilian women with previous gestational diabetes. 2007 , 78, 23-9 | 24 |
| 126 | C-reactive protein is related to extent and progression of coronary and extra-coronary atherosclerosis; results from the Rotterdam study. 2007 , 195, e195-202 | 54 |
| 125 | Intraplaque hemorrhage assessed by high-resolution magnetic resonance imaging and C-reactive protein in carotid atherosclerosis. <i>Journal of Vascular Surgery</i> , 2007 , 46, 1130-7 | 29 |
| 124 | Arterial vascular disease in women. <i>Journal of Vascular Surgery</i> , 2007 , 46, 1295-302 | 57 |
| | | |
| 123 | La Aterosclerosis como enfermedad sistfhica. 2007 , 60, 184-195 | 42 |
| 123 | La Aterosclerosis como enfermedad sistEhica. 2007 , 60, 184-195 Atherosclerosis As a Systemic Disease. 2007 , 60, 184-195 | 42 5 |
| | | |
| | Atherosclerosis As a Systemic Disease. 2007 , 60, 184-195 | |
| 122 | Atherosclerosis As a Systemic Disease. 2007, 60, 184-195 Protefia C reativa e instabilidade clitica na doenii obstrutiva de artifias cartidas. 2007, 6, 124-129 Differential association of C-reactive protein with adiposity in men and women in an Aboriginal | 5 |
| 122 121 120 | Atherosclerosis As a Systemic Disease. 2007, 60, 184-195 Protefia C reativa e instabilidade clítica na doena obstrutiva de artifias cartidas. 2007, 6, 124-129 Differential association of C-reactive protein with adiposity in men and women in an Aboriginal community in northeast Arnhem Land of Australia. 2007, 31, 103-8 A dose-response study of hormone replacement in young hypogonadal women: effects on intima | 33 |
| 122 121 120 | Atherosclerosis As a Systemic Disease. 2007, 60, 184-195 Protefla C reativa e instabilidade clílica na doeníl obstrutiva de artílias carílidas. 2007, 6, 124-129 Differential association of C-reactive protein with adiposity in men and women in an Aboriginal community in northeast Arnhem Land of Australia. 2007, 31, 103-8 A dose-response study of hormone replacement in young hypogonadal women: effects on intima media thickness and metabolism. 2007, 66, 557-64 Association of interleukin-6 and C-reactive protein with subclinical carotid atherosclerosis (the | 5 33 38 |
| 122 121 120 119 | Atherosclerosis As a Systemic Disease. 2007, 60, 184-195 Protella C reativa e instabilidade clilica na doenil obstrutiva de artilias cartidas. 2007, 6, 124-129 Differential association of C-reactive protein with adiposity in men and women in an Aboriginal community in northeast Arnhem Land of Australia. 2007, 31, 103-8 A dose-response study of hormone replacement in young hypogonadal women: effects on intima media thickness and metabolism. 2007, 66, 557-64 Association of interleukin-6 and C-reactive protein with subclinical carotid atherosclerosis (the Rancho Bernardo Study). 2007, 99, 99-102 Comparison of characteristics of future myocardial infarctions in women with baseline high versus | 5 33 38 46 |

(2009-2008)

| 114 | The elusive link between high sensitivity C-reactive protein and carotid subclinical atherosclerosis in coronary artery bypass grafting candidates: a cross-sectional study. 2008 , 6, 23 | 3 |
|-----|--|-----|
| 113 | Effects of the progestagen-only contraceptive implant Implanon on cardiovascular risk factors. 2008 , 68, 355-60 | 28 |
| 112 | Inflammatory markers and progression of subclinical atherosclerosis in healthy postmenopausal women (from the Estrogen in the Prevention of Atherosclerosis Trial). 2008 , 101, 1131-3 | 16 |
| 111 | Direct association between C-reactive protein serum levels and endothelial dysfunction in patients with claudication. 2008 , 35, 480-6 | 21 |
| 110 | C-reactive protein and the metabolic syndrome correlate differently with carotid atherosclerosis between men and women in a Taiwanese community. 2008 , 57, 1023-8 | 11 |
| 109 | Low-grade chronic inflammation is not accompanied by structural arterial injury in polycystic ovary syndrome. 2008 , 81, 179-83 | 16 |
| 108 | Carotid intima-media thickness and markers of inflammation, endothelial damage and hemostasis. 2008 , 40, 21-44 | 71 |
| 107 | Slow heart rate recovery after exercise is associated with carotid atherosclerosis. 2008, 196, 256-261 | 33 |
| 106 | Prevalence and correlates of silent cerebral infarcts in the Framingham offspring study. 2008, 39, 2929-35 | 236 |
| 105 | Adherence to the mediterranean diet is inversely associated with circulating interleukin-6 among middle-aged men: a twin study. 2008 , 117, 169-75 | 110 |
| 104 | High-sensitivity CRP is correlated with neurologic symptoms and plaque instability in patients with severe stenosis of the carotid bifurcation. 2008 , 42, 249-55 | 7 |
| 103 | Genetics of the Framingham Heart Study population. 2008 , 62, 33-65 | 66 |
| 102 | Cardiovascular risk evaluated by C-reactive protein levels in diabetic and obese Mexican subjects. 2008 , 72, 1170-4 | 8 |
| 101 | Association of soluble adhesion molecule and C-reactive protein levels with silent brain infarction in patients with and without type 2 diabetes. 2008 , 5, 106-11 | 6 |
| 100 | Cardiovascular risk prediction by N-terminal pro brain natriuretic peptide and high sensitivity C-reactive protein is affected by age and sex. 2008 , 26, 26-34 | 32 |
| 99 | Does high C-reactive protein concentration increase atherosclerosis? The Whitehall II Study. 2008 , 3, e3013 | 33 |
| 98 | Erythrocyte aggregation as an early biomarker in patients with asymptomatic carotid stenosis. 2008 , 24, 33-9 | 14 |
| 97 | Effects of pitavastatin on serum lipids and high sensitivity C-reactive protein in type 2 diabetic patients. 2009 , 16, 546-52 | 18 |

| 96 | Sex differences in inflammatory markers: what is the contribution of visceral adiposity?. 2009 , 89, 1307-14 | 136 |
|----|--|-----|
| 95 | The association between inflammatory markers and carotid atherosclerosis is sex dependent: the Tromslbtudy. 2009 , 27, 392-7 | 35 |
| 94 | Prevalence of asymptomatic carotid artery stenosis according to age and sex: systematic review and metaregression analysis. 2009 , 40, 1105-13 | 144 |
| 93 | New risk markers may change the HeartScore risk classification significantly in one-fifth of the population. 2009 , 23, 105-12 | 8 |
| 92 | Increased carotid intima-media thickness in patients with pheochromocytoma in comparison to essential hypertension. 2009 , 23, 350-8 | 13 |
| 91 | High-sensitivity C-reactive protein is not independently associated with peripheral subclinical atherosclerosis. 2009 , 60, 12-20 | 4 |
| 90 | Asthma is an independent risk for elevation of plasma C-reactive protein levels. 2009, 399, 79-82 | 21 |
| 89 | Association of the -757T>C polymorphism in the CRP gene with circulating C-reactive protein levels and carotid atherosclerosis. 2009 , 124, 458-62 | 16 |
| 88 | [C-reactive protein and carotid intima-media thickness in atherothrombotic ischemic stroke]. 2009 , 133, 496-500 | 3 |
| 87 | The relationships among serum glucose, albumin concentrations and carotid atherosclerosis in men with spinal cord injury. 2009 , 206, 528-34 | 25 |
| 86 | Carotid intima-media thickness: an early marker of cardiovascular disease. 2009 , 25, 55-61 | 22 |
| 85 | Highly sensitive C-reactive protein, body mass index, and serum lipids in HIV-infected persons receiving antiretroviral therapy: a longitudinal study. 2009 , 52, 480-7 | 29 |
| 84 | Distribution of C-reactive protein and its association with subclinical atherosclerosis in asymptomatic postmenopausal Chinese women. 2010 , 59, 1672-9 | 5 |
| 83 | Gender-specific features of plasmatic and circulating cell alterations as risk factors in cardiovascular disease. 2010 , 24, 665-74 | 7 |
| 82 | Prehypertension subtype with elevated C-reactive protein: risk of ischemic stroke in a general Japanese population. 2010 , 23, 1108-13 | 23 |
| 81 | Cilostazol reduces restenosis after carotid artery stenting. <i>Journal of Vascular Surgery</i> , 2010 , 51, 51-6 3.5 | 45 |
| 80 | Gestational diabetes and subclinical inflammation: evaluation of first year postpartum outcomes. 2011 , 94, 426-33 | 24 |
| 79 | Association of systemic inflammatory activity with coronary and carotid atherosclerosis in the very elderly. 2011 , 216, 212-6 | 37 |

| 78 | Carotid intima-media thickness as a surrogate marker of cardiovascular disease in diabetes. 2011 , 4, 23-34 | 52 |
|----|--|----|
| 77 | High-Sensitivity C-Reactive Protein and Cardiovascular Disease. 2011 , 3, 178-186 | |
| 76 | Significant correlation between visceral adiposity and high-sensitivity C-reactive protein (hs-CRP) in Japanese subjects. 2011 , 50, 2767-73 | 31 |
| 75 | [Classification of etiologic subtypes for transient ischemic attacks: clinical significance of lacunar transient ischemic attack]. 2011 , 51, 406-11 | 6 |
| 74 | Role of C-reactive protein in cerebrovascular disease: a critical review. 2011 , 9, 1565-84 | 53 |
| 73 | Increased risk of death in pediatric and adult patients with ESRD secondary to lupus. 2011 , 26, 93-8 | 46 |
| 72 | A crossover study of rosuvastatin and pitavastatin in patients with type 2 diabetes. 2011 , 28, 160-71 | 14 |
| 71 | Ultrasound imaging of carotid intima-media thickness: an office-based tool to assist physicians in cardiovascular risk assessment. 2011 , 13, 431-6 | 8 |
| 70 | Blood pressure tracking over the adult life course: patterns and correlates in the Framingham heart study. 2012 , 60, 1393-9 | 89 |
| 69 | High-sensitive C-reactive protein: universal prognostic and causative biomarker in heart disease?. 2012 , 6, 19-34 | 34 |
| 68 | The effect of chronic carbon monoxide exposure on hs-CRP, CIM thickness and PEF in furnacemen. 2012 , 37, 42-47 | |
| 67 | Evaluation of clinical variables associated with increased carotid intima-media thickness in middle-aged hypertensive women. 2012 , 2012, 257501 | 5 |
| 66 | Factors associated with carotid intima-media thickness and carotid plaques in type 2 diabetic patients. 2012 , 30, 940-7 | 25 |
| 65 | Criticial limb ischemia: epidemiology. 2012 , 8, 10-4 | 57 |
| 64 | Can premenstrual syndrome affect arterial stiffness or blood pressure?. 2012, 224, 170-6 | 15 |
| 63 | Periprocedural cilostazol treatment and restenosis after carotid artery stenting: the Retrospective Study of In-Stent Restenosis after Carotid Artery Stenting (ReSISteR-CAS). 2012 , 21, 193-9 | 25 |
| 62 | Inflammation and Atherosclerosis. 2012, | 3 |
| 61 | Angiographically confirmed stent over expansion in the internal carotid artery during stenting: incidence, predictors, and outcomes. 2012 , 54, 481-6 | 1 |

| 60 | Managing cardiovascular risk in patients with chronic inflammatory diseases. 2012 , 31, 585-94 | | 16 |
|----|--|-----|----|
| 59 | Relationship among circulating inflammatory proteins, platelet gene expression, and cardiovascular risk. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2013 , 33, 2666-73 | 9.4 | 49 |
| 58 | Atherosclerotic biomarkers and aortic atherosclerosis by cardiovascular magnetic resonance imaging in the Framingham Heart Study. 2013 , 2, e000307 | | 11 |
| 57 | The value of C-reactive protein in symptomatic versus asymptomatic carotid artery stenosis. 2013 , 123, 311-7 | | 1 |
| 56 | Inflammatory markers and blood pressure: sex differences and the effect of fat mass in the CoLaus Study. 2013 , 27, 169-75 | | 27 |
| 55 | Increased C-reactive protein associates with elevated carotid intima-media thickness in Chinese adults with normal low density lipoprotein cholesterol levels. 2013 , 20, 575-84 | | 7 |
| 54 | Biomarkers of inflammation may be of use for identification of more severe peripheral arterial occlusive disease. 2014 , 13, 182-191 | | 2 |
| 53 | Two-hour post-load plasma glucose levels are associated with carotid intima-media thickness in subjects with normal glucose tolerance. 2014 , 31, 76-83 | | 10 |
| 52 | Incidence and risk factors of significant carotid artery stenosis in asymptomatic survivors of head and neck cancer after radiotherapy. 2014 , 36, 215-9 | | 44 |
| 51 | C-reactive protein and subclinical cardiovascular disease among African-Americans: (the Jackson Heart Study). 2014 , 15, 371-6 | | 11 |
| 50 | Relationship between periodontal disease and subclinical atherosclerosis: the Dong-gu study. 2014 , 41, 262-8 | | 13 |
| 49 | Increased Hs-CRP/adiponectin ratio is associated with increase carotid intima-media thickness. 2014 , 13, 120 | | 13 |
| 48 | Carotid intima-media thickness value distributions in the Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). 2014 , 237, 227-35 | | 52 |
| 47 | Aortic arch atherosclerosis in ischaemic stroke of unknown origin affects prognosis. 2014 , 4, 92-101 | | 4 |
| 46 | Increased Soluble TNF Receptor-1 and Glutathione Peroxidase may Predict Carotid Intima Media Thickness in Females with Cushing Syndrome. 2015 , 21, 286-95 | | 4 |
| 45 | Associations between Inflammatory Markers and Subclinical Atherosclerosis in Middle-aged White, Japanese-American and Japanese Men: The ERA-JUMP Study. 2015 , 22, 590-8 | | 17 |
| 44 | High red blood cell distribution width and preclinical carotid atherosclerosis. 2015 , 20, 376-81 | | 14 |
| 43 | Modulation of C-reactive protein and plasma omega-6 fatty acid levels by phospholipase A2 gene polymorphisms following a 6-week supplementation with fish oil. 2015 , 102-103, 37-45 | | 4 |

(2021-2017)

| 42 | Long-Term Effect of Adrenalectomy on Cardiovascular Remodeling in Patients With Pheochromocytoma. 2017 , 102, 1208-1217 | 11 |
|----|--|----|
| 41 | Residential proximity to abandoned uranium mines and serum inflammatory potential in chronically exposed Navajo communities. 2017 , 27, 365-371 | 29 |
| 40 | Association Between Absolute Eosinophil Count and Complex Aortic Arch Plaque in Patients With Acute Ischemic Stroke. 2017 , 48, 1074-1076 | 8 |
| 39 | MicroRNA Signature of Cigarette Smoking and Evidence for a Putative Causal Role of MicroRNAs in Smoking-Related Inflammation and Target Organ Damage. 2017 , 10, | 28 |
| 38 | C-reactive protein in atherosclerosis - A risk marker but not a causal factor? A 13-year population-based longitudinal study: The Troms[study. 2017 , 263, 293-300 | 15 |
| 37 | No Association Between High-Sensitivity C-Reactive Protein and Carotid Intima-Media Progression: The APAC Study. 2017 , 26, 252-259 | 4 |
| 36 | Obesity and Its Influence on Mediators of Inflammation: Clinical Relevance of C-Reactive Protein in Obese Subjects. 2017 , | |
| 35 | Association of Chronic Low-grade Inflammation With Risk of Alzheimer Disease in ApoE4 Carriers. 2018 , 1, e183597 | 76 |
| 34 | ☑aluation et quantification d៊une stBose carotidienne par Ēhographie-Doppler. 2019 , 2, 204-216 | 1 |
| 33 | Effects of sex steroids on cardiovascular risk profile in transgender men under gender affirming hormone therapy. 2019 , 66, 385-392 | 1 |
| 32 | Management of Patients with Asymptomatic and Symptomatic Carotid Artery Disease: Update on Anti-Thrombotic Therapy. 2019 , 119, 576-585 | 6 |
| 31 | Catecholamines Induce Left Ventricular Subclinical Systolic Dysfunction: A Speckle-Tracking Echocardiography Study. 2019 , 11, | 4 |
| 30 | Association between neutrophil to lymphocyte ratio and carotid artery wall thickness in healthy pregnant women. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2020 , 255, 98-10 ⁴⁻⁴ | 4 |
| 29 | Evaluation of single nucleotide polymorphisms in 6 candidate genes and carotid intima-media thickness in community-dwelling residents. 2020 , 15, e0230715 | 1 |
| 28 | Is multiple sclerosis a risk factor for atherosclerosis?. 2021 , 48, 99-103 | 6 |
| 27 | Future stroke risk in the chronic phase of post-percutaneous coronary intervention. 2021 , 16, e0251253 | |
| 26 | Analysis of the levels of inflammatory parameters in persons over the age of 90. 2021 , 148, 111278 | |
| 25 | C-reactive protein in adult Samoans: Population variation and physiological correlates. 2021 , e23646 | |

| 24 | Association between C-reactive protein and carotid plaque in mild-to-moderate idiopathic pulmonary fibrosis. 2021 , 16, 1529-1539 | 5 |
|----------------------|--|----|
| 23 | B-Mode Ultrasound: A Noninvasive Method for Assessing Atherosclerosis. 2007 , 1783-1796 | 2 |
| 22 | Effects of sex steroids on cardiovascular risk profile in transgender men under gender affirming hormone therapy. 2019 , 66, 385-392 | 10 |
| 21 | Is high-sensitivity C-reactive protein associated with carotid atherosclerosis in healthy Koreans?. 2005 , 12, 548-54 | 11 |
| 20 | Association of general and abdominal obesities and metabolic syndrome with subclinical atherosclerosis in asymptomatic Chinese postmenopausal women. 2008 , 15, 185-92 | 21 |
| 19 | The association between circulating inflammatory markers and metabolic syndrome in Korean rural adults. 2008 , 41, 413-8 | 13 |
| 18 | [C-reactive protein and carotid intima-media thickness in a population of middle-aged Koreans]. 2009 , 42, 29-34 | 5 |
| 17 | The Effects of Omega-3 and Vitamin E Co-supplementation on Carotid Intima-media Thickness and Inflammatory Factors in Patients with Polycystic Ovary Syndrome. 2018 , 33, 473-479 | 12 |
| 16 | Impact of C-reactive protein on the likelihood of carotid atherosclerosis in Japanese adults. 2006 , 13, 175-82 | 8 |
| 15 | Serum Biomarkers in Prediction of Stroke Risk and Outcome. 2004 , 257-278 | |
| | | |
| 14 | [Relationship between C-reactive protein and features of the metabolic syndrome in military pilots in the Serbia and Montenegro]. 2005 , 62, 811-9 | |
| 14 | | 1 |
| | in the Serbia and Montenegro]. 2005, 62, 811-9 [The relationship between C-reactive protein and subclinical carotid arteriosclerosis in military | 1 |
| 13 | in the Serbia and Montenegro]. 2005, 62, 811-9 [The relationship between C-reactive protein and subclinical carotid arteriosclerosis in military pilots]. 2009, 66, 465-71 Cardiovascular Risk Prediction by Measurement of Arterial Elastic Properties and Wall Thickness. | |
| 13 | in the Serbia and Montenegro]. 2005, 62, 811-9 [The relationship between C-reactive protein and subclinical carotid arteriosclerosis in military pilots]. 2009, 66, 465-71 Cardiovascular Risk Prediction by Measurement of Arterial Elastic Properties and Wall Thickness. 2009, 399-421 | |
| 13 12 11 | in the Serbia and Montenegro]. 2005, 62, 811-9 [The relationship between C-reactive protein and subclinical carotid arteriosclerosis in military pilots]. 2009, 66, 465-71 Cardiovascular Risk Prediction by Measurement of Arterial Elastic Properties and Wall Thickness. 2009, 399-421 Future Directions of Atherosclerosis Research and Translation into Clinical Application. 2012, 613-623 Predictive value of C-reactive protein and carotid intimal medial thickness in acute ischemic stroke. | 0 |
| 13 12 11 10 | [The relationship between C-reactive protein and subclinical carotid arteriosclerosis in military pilots]. 2009, 66, 465-71 Cardiovascular Risk Prediction by Measurement of Arterial Elastic Properties and Wall Thickness. 2009, 399-421 Future Directions of Atherosclerosis Research and Translation into Clinical Application. 2012, 613-623 Predictive value of C-reactive protein and carotid intimal medial thickness in acute ischemic stroke. 2019, 55, | 0 |

CITATION REPORT

| 6 | Ethnicity and sex modify the association of serum c-reactive protein with microalbuminuria. 2008 , 18, 324-9 | | 5 |
|---|---|-----|---|
| 5 | Associations between Adverse Childhood Experiences and the novel inflammatory marker glycoprotein acetyls in two generations of the Avon Longitudinal Study of Parents and Children birth cohort. 2021 , 100, 112-120 | | 1 |
| 4 | Prognostic indicators of persistent carotid intima-media thickness increase in postpartum period in a population of normotensive women with gestational diabetes mellitus <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2021 , 269, 47-54 | 2.4 | O |
| 3 | Periodontitis is associated with higher subclinical atherosclerosis in patients with systemic lupus erythematosus <i>Journal of Periodontal Research</i> , 2022 , | 4.3 | O |
| 2 | Lack of association between cognitive impairment and systemic inflammation in asymptomatic carotid stenosis <i>Journal of Vascular Surgery</i> , 2021 , | 3.5 | 1 |
| 1 | Aort Kapak Sklerozunun 🗓 🗹 Emesinde Yeni Hematolojik 🖺 flamatuar Parametrelerin 🖰 emi. 2022 , 7, 37-41 | | |