

CITATION REPORT

List of articles citing

Elevated production of interleukin-18 is associated with renal disease in patients with systemic lupus erythematosus

DOI: 10.1046/j.1365-2249.2002.01989.x

Clinical and Experimental Immunology, 2002, 130, 345-51.

Source: <https://exaly.com/paper-pdf/34642381/citation-report.pdf>

Version: 2024-04-10

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
62	IL-18 cDNA vaccination protects mice from spontaneous lupus-like autoimmune disease. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 14181-6	11.5	101
61	Blockade of IL-18 receptor signaling delays the onset of autoimmune disease in MRL-Faslpr mice. <i>Journal of Immunology</i> , 2004 , 173, 5312-8	5.3	52
60	Up-regulation of IL-18 and predominance of a Th1 immune response is a hallmark of lupus nephritis. <i>Clinical and Experimental Immunology</i> , 2004 , 138, 171-8	6.2	100
59	Biomarkers in systemic lupus erythematosus: II. Markers of disease activity. <i>Arthritis and Rheumatism</i> , 2004 , 50, 2048-65		120
58	Nitric oxide and glomerulonephritis. <i>Seminars in Nephrology</i> , 2004 , 24, 324-32	4.8	23
57	Biomarkers for systemic lupus erythematosus: a review and perspective. <i>Current Opinion in Rheumatology</i> , 2005 , 17, 543-9	5.3	53
56	Th1 cytokines in the pathogenesis of lupus nephritis: the role of IL-18. <i>Autoimmunity Reviews</i> , 2005 , 4, 542-8	13.6	57
55	Interleukin-18 receptor signaling is not required for autoantibody production and end-organ disease in murine lupus. <i>Arthritis and Rheumatism</i> , 2005 , 52, 984-6		12
54	Immunothérapies du lupus systémique : innovations et perspectives. <i>Revue Du Rhumatisme (Edition Française)</i> , 2005 , 72, 572-591	0.1	1
53	Aberrant production of soluble costimulatory molecules CTLA-4, CD28, CD80 and CD86 in patients with systemic lupus erythematosus. <i>Rheumatology</i> , 2005 , 44, 989-94	3.9	100
52	Plasma interleukin-18 levels in chronic renal failure and continuous ambulatory peritoneal dialysis. <i>Blood Purification</i> , 2005 , 23, 144-8	3.1	25
51	Elevation of plasma osteopontin concentration is correlated with disease activity in patients with systemic lupus erythematosus. <i>Rheumatology</i> , 2005 , 44, 602-6	3.9	76
50	Immunomodulatory effects of lingzhi and san-miao-san supplementation on patients with rheumatoid arthritis. <i>Immunopharmacology and Immunotoxicology</i> , 2006 , 28, 197-200	3.2	20
49	Blockade of TLR9 agonist-induced type I interferons promotes inflammatory cytokine IFN-gamma and IL-17 secretion by activated human PBMC. <i>Cytokine</i> , 2006 , 35, 235-46	4	31
48	Interleukin-18 overexpression as a hallmark of the activity of autoimmune inflammatory myopathies. <i>Clinical and Experimental Immunology</i> , 2006 , 146, 21-31	6.2	46
47	Elevation of plasma interleukin-18 concentration is associated with insulin levels in patients with systemic lupus erythematosus. <i>Lupus</i> , 2006 , 15, 207-12	2.6	9
46	Immunomodulatory effects of a traditional Chinese medicine with potential antiviral activity: a self-control study. <i>The American Journal of Chinese Medicine</i> , 2006 , 34, 13-21	6	60

45	Raised plasma concentration and ex vivo production of inflammatory chemokines in patients with systemic lupus erythematosus. <i>Annals of the Rheumatic Diseases</i> , 2006 , 65, 209-15	2.4	136
44	Serum concentration of interleukin-18 is up-regulated in patients with ANCA-associated vasculitis. <i>Autoimmunity</i> , 2007 , 40, 529-31	3	17
43	Elevation of pro-inflammatory cytokines, C-reactive protein and cardiac troponin T in chronic renal failure patients on dialysis. <i>Immunological Investigations</i> , 2007 , 36, 47-57	2.9	30
42	Cytokines in glomerulonephritis. <i>Seminars in Nephrology</i> , 2007 , 27, 275-85	4.8	20
41	Safety and efficacy of Ganoderma lucidum (lingzhi) and San Miao San supplementation in patients with rheumatoid arthritis: a double-blind, randomized, placebo-controlled pilot trial. <i>Arthritis and Rheumatism</i> , 2007 , 57, 1143-50		55
40	Type I interferons inhibition of inflammatory T helper cell responses in systemic lupus erythematosus. <i>Annals of the New York Academy of Sciences</i> , 2007 , 1108, 11-23	6.5	24
39	Association of serum nitrate and nitrite levels with longitudinal assessments of disease activity and damage in systemic lupus erythematosus and lupus nephritis. <i>Arthritis and Rheumatism</i> , 2008 , 58, 263-72		46
38	Evidence for a pathogenetic role of interleukin-18 in cutaneous lupus erythematosus. <i>Arthritis and Rheumatism</i> , 2008 , 58, 3205-15		68
37	Overexpression of interleukin-12 and T helper 1 predominance in lupus nephritis. <i>Clinical and Experimental Immunology</i> , 2008 , 154, 247-54	6.2	80
36	Hyperproduction of IL-23 and IL-17 in patients with systemic lupus erythematosus: implications for Th17-mediated inflammation in auto-immunity. <i>Clinical Immunology</i> , 2008 , 127, 385-93	9	409
35	Cytokines and their receptors as biomarkers of systemic lupus erythematosus. <i>Expert Review of Molecular Diagnostics</i> , 2008 , 8, 189-98	3.8	34
34	Interleukin-18 in urine and serum of children with idiopathic nephrotic syndrome. <i>Kidney and Blood Pressure Research</i> , 2008 , 31, 122-6	3.1	10
33	IL-18 activity in systemic lupus erythematosus. <i>Annals of the New York Academy of Sciences</i> , 2009 , 1173, 301-9	6.5	52
32	Increased spontaneous but decreased mitogen-stimulated expression and excretion of interleukin 18 by mononuclear cells in patients with active systemic lupus erythematosus. <i>Journal of Rheumatology</i> , 2009 , 36, 1910-6	4.1	3
31	Association of interleukin-18 promoter polymorphisms with WHO pathological classes and serum IL-18 levels in Chinese patients with lupus nephritis. <i>Lupus</i> , 2009 , 18, 29-37	2.6	29
30	Association of IL-18 gene polymorphism (-137C) with arthritis manifestations in SLE: combined effect with IFN gamma gene polymorphism (+874A). <i>Clinical Rheumatology</i> , 2009 , 28, 219-23	3.9	21
29	Activation profile of intracellular mitogen-activated protein kinases in peripheral lymphocytes of patients with systemic lupus erythematosus. <i>Journal of Clinical Immunology</i> , 2009 , 29, 738-46	5.7	26
28	IL-18 105 A>C polymorphism contributes to renal manifestations in patients with SLE. <i>Rheumatology International</i> , 2009 , 30, 187-91	3.6	12

27	Decreased expression of T lymphocyte co-stimulatory molecule CD26 on invariant natural killer T cells in systemic lupus erythematosus. <i>Immunological Investigations</i> , 2009 , 38, 350-64	2.9	17
26	Expressions of IL-18 and its binding protein in peripheral blood leukocytes and kidney tissues of lupus nephritis patients. <i>Clinical Rheumatology</i> , 2010 , 29, 717-21	3.9	22
25	High circulating levels of free interleukin-18 in patients with active SLE in the presence of elevated levels of interleukin-18 binding protein. <i>Journal of Autoimmunity</i> , 2010 , 34, 121-6	15.5	62
24	Serum interleukin-18 binding protein increases with behavior different from IL-18 in patients with diabetic nephropathy. <i>Diabetes Research and Clinical Practice</i> , 2011 , 92, e66-9	7.4	4
23	Retinal self-antigen induces a predominantly Th1 effector response in Axl and Mertk double-knockout mice. <i>Journal of Immunology</i> , 2011 , 187, 4178-86	5.3	26
22	Association of IL-18 polymorphisms with rheumatoid arthritis and systemic lupus erythematosus in Asian populations: a meta-analysis. <i>BMC Medical Genetics</i> , 2012 , 13, 107	2.1	16
21	Serum cytokine profile in patients with active lupus nephritis. <i>Cytokine</i> , 2012 , 60, 410-6	4	65
20	Interleukin-18 contributes more closely to the progression of diabetic nephropathy than other diabetic complications. <i>Acta Diabetologica</i> , 2012 , 49, 111-7	3.9	46
19	Elevated interleukin-18 and skewed Th1:Th2 immune response in lupus nephritis. <i>Rheumatology International</i> , 2012 , 32, 223-9	3.6	16
18	IL-18 in inflammatory and autoimmune disease. <i>Cellular and Molecular Life Sciences</i> , 2013 , 70, 4795-808	10.3	90
17	The role of cytokines in the pathogenesis of systemic lupus erythematosus - from bench to bedside. <i>Nephrology</i> , 2013 , 18, 243-55	2.2	68
16	Serum interleukin-18 levels in patients with systemic lupus erythematosus: Relation with disease activity and lupus nephritis. <i>Egyptian Rheumatologist</i> , 2013 , 35, 45-51	1	15
15	Interleukin-12 is associated with arterial stiffness in healthy individuals. <i>American Journal of Hypertension</i> , 2013 , 26, 159-62	2.3	23
14	Systemic Lupus Erythematosus and Psoriasis: Comparison of Immunopathogenesis. <i>Psoriasis Forum</i> , 2013 , 19a, 165-175		
13	Association between interleukin 18 polymorphisms and alopecia areata in Koreans. <i>Journal of Interferon and Cytokine Research</i> , 2014 , 34, 349-53	3.5	9
12	Serum interleukin-18 and interleukin-10 levels in systemic lupus erythematosus: correlation with SLEDAI score and disease activity parameters. <i>Egyptian Rheumatology and Rehabilitation</i> , 2014 , 41, 160-166	0.5	4
11	Role of renal biomarkers as predictors of acute kidney injury in cardiac surgery. <i>Asian Cardiovascular and Thoracic Annals</i> , 2014 , 22, 234-41	0.6	13
10	Elevation of Interleukin-18 Correlates With Cardiovascular, Cerebrovascular, and Peripheral Vascular Events: A Cohort Study of Hemodialysis Patients. <i>Medicine (United States)</i> , 2015 , 94, e1836	1.8	6

9	Th-17 related regulatory network in the pathogenesis of Arab patients with systemic lupus erythematosus and lupus nephritis. <i>International Journal of Rheumatic Diseases</i> , 2016 , 19, 512-20	2.3	10
8	Serum IL-18 as biomarker in predicting long-term renal outcome among pediatric-onset systemic lupus erythematosus patients. <i>Medicine (United States)</i> , 2016 , 95, e5037	1.8	17
7	Cytokine targets in lupus nephritis: Current and future prospects. <i>Clinical Immunology</i> , 2019 , 206, 42-52	9	13
6	High circulating levels of interleukin-18 binding protein indicate the severity of glomerular involvement in systemic lupus erythematosus. <i>Modern Rheumatology</i> , 2012 , 22, 73-9	3.3	6
5	Correlation of plasma interleukin-18 concentration and severity of renal involvement and disease activity in systemic lupus erythematosus. <i>Journal of Nephropathology</i> , 2016 , 5, 28-33	0.6	17
4	Effects of Interleukin Families Polymorphisms on Systemic Lupus Erythematosus: Focus on Interleukin-1. <i>Gene, Cell and Tissue</i> , 2018 , In Press,	0.6	1
3	Enhanced Activity of NLRP3 Inflammasome in the Lung of Patients with Anti-Synthetase Syndrome. 2023 , 12, 60		0
2	Construction of molecular subgroups in childhood systemic lupus erythematosus using bioinformatics. 2022 , 101, e32274		0
1	Interleukin-18 Promoter Gene Polymorphisms in Chinese Patients With Systemic Lupus Erythematosus: Association With CC Genotype at Position 807. 2007 , 36, 91-95		1