

CITATION REPORT

List of articles citing

Separation methods for estimating octanol-water partition coefficients

DOI: 10.1016/j.jchromb.2003.08.032

Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2003, 797, 3-19.

Source: <https://exaly.com/paper-pdf/35097857/citation-report.pdf>

Version: 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
205	High-throughput multiplexed capillary electrophoresis in drug discovery. 2004 , 9, 1072-80		52
204	Determination of liquid-liquid partition coefficients by separation methods. 2004 , 1037, 3-14		198
203	Determination of acid dissociation constants by capillary electrophoresis. 2004 , 1037, 445-54		168
202	Column selection for liquid chromatographic estimation of the k'_w hydrophobicity parameter. 2004 , 1042, 37-48		41
201	Classification of drugs in absorption classes using the classification and regression trees (CART) methodology. 2005 , 39, 91-103		106
200	Prediction of gastro-intestinal absorption using multivariate adaptive regression splines. 2005 , 39, 1021-30		31
199	Determination of lipophilicity by reversed-phase high-performance liquid chromatography. Influence of 1-octanol in the mobile phase. 2005 , 1091, 51-9		81
198	Evaluation of a Semi-Empirical Approach to Correlation of Retention with Octanol-Water Partition Coefficients by Use of Adjustable Hydrogen-Bonding Indicator Variables. 2005 , 61, 479-492		5
197	Recent advances in micellar electrokinetic chromatography. 2005 , 26, 719-34		76
196	Quantitative structure-retention relationships for ionic and non-ionic compounds in biopartitioning micellar chromatography. 2005 , 19, 155-68		17
195	The molecular retention mechanism in reversed-phase liquid chromatography of meso-ionic compounds by quantitative structure-retention relationships (QSRR). 2005 , 2, 1691-700		7
194	Capillary electrophoresis technologies for screening in drug discovery. 2005 , 2, 171-8		18
193	High-throughput physicochemical and in vitro ADMET screening. 2005 , 3, 83-100		14
192	Presentation of a structurally diverse and commercially available drug data set for correlation and benchmarking studies. 2006 , 49, 6660-71		40
191	Chromatographic estimation of drug disposition properties by means of immobilized artificial membranes (IAM) and C18 columns. 2006 , 49, 4861-70		78
190	Classification tree models for the prediction of blood-brain barrier passage of drugs. 2006 , 46, 1410-9		49
189	Planar chromatography. 2006 , 78, 3841-52		19

188	. 2006 ,	23
187	Electromigration Separation Techniques in Pharmaceutical Analysis. 351-371	
186	Selectivity Characterization of Pseudostationary Phases Using the Solvation Parameter Model. 55-78	1
185	Heteroscedasticity of retention factor and adequate modeling in micellar liquid chromatography. 2006 , 576, 229-38	14
184	Models and objective functions for the optimisation of selectivity in reversed-phase liquid chromatography. 2006 , 579, 125-45	97
183	Synthesis and comparative molecular field analysis (CoMFA) of argentatin B derivatives as growth inhibitors of human cancer cell lines. 2006 , 14, 1889-901	13
182	Comparative evaluation of three high-performance liquid chromatography-based Kow estimation methods for highly hydrophobic organic compounds: polybrominated diphenyl ethers and hexabromocyclododecane. 2006 , 25, 2018-27	51
181	Experimental studies of environmental processes: A practical course in environmental chemistry. 2006 , 13, 435-40	2
180	Correlations Between Phenols-Micelles Partition Coefficients and Several Molecular Descriptors. An Approach to Predict the Phenols Behaviour in MSPME. 2006 , 63, 167-174	8
179	Investigation of copper-azamacrocyclic complexes by high-performance liquid chromatography. 2006 , 20, 37-47	4
178	Recent applications of microemulsion electrokinetic chromatography. 2006 , 27, 60-75	72
177	Chromatographic Approaches for Measuring Log P. 2007 , 331-355	7
176	The Good, the Bad and the Ugly of Distribution Coefficients: Current Status, Views and Outlook. 2007 , 407-437	6
175	Regression method of the hydrophobicity ruler approach for determining octanol/water partition coefficients of very hydrophobic compounds. 2007 , 66, 1086-93	7
174	Tungstate-Based Inorganic/Organic Hybrid Nanobelts/Nanotubes with Lamellar Mesostructures: Synthesis, Characterization, and Formation Mechanism. 2007 , 19, 1808-1815	55
173	Chemical speciation of copper(II) diaminediamide derivative of pentacycloundecane--a potential anti-inflammatory agent. 2007 , 1140-9	15
172	Structure-Retention Correlation in Liquid Chromatography for Pharmaceutical Applications. 2007 , 30, 761-789	16
171	A system for LogD screening of 96-well plates using a water-plug aspiration/injection method combined with high-performance liquid chromatography-mass spectrometry. 2007 , 79, 8312-5	51

170	A study of the lipophilicity of amide derivatives of (1,2,3,4-tetrahydroisoquinolin-2-yl)- β -hydroxybutyric acid by use of RP-TLC and calculation. 2007 , 20, 101-106	8
169	. 2007 ,	6
168	Evaluation of boosted regression trees (BRTs) and two-step BRT procedures to model and predict blood-brain barrier passage. 2007 , 21, 280-291	11
167	MEKC: an update focusing on practical aspects. 2007 , 28, 174-92	65
166	Structure-retention relationship study of diastereomeric (Z)- and (E)-2-alkylidene-4-oxothiazolidines. 2007 , 30, 2241-8	11
165	Evaluation of chromatographic descriptors for the prediction of gastro-intestinal absorption of drugs. 2007 , 1138, 190-202	33
164	Counter-current chromatographic estimation of hydrophobicity of Z-(cis) and E-(trans) enalapril and kinetics of cis/trans isomerization. 2007 , 1157, 101-7	11
163	Polymer/water partition coefficients of extended range measured by using organic modifiers in the aqueous phase. 2007 , 48, 7523-7530	8
162	Relationship between physicochemical properties, lipophilicity parameters, and local anesthetic activity of dibasic esters of phenylcarbamic acid. 2007 , 61,	6
161	High-throughput method for lipophilicity measurement. 2007 , 79, 1043-9	28
160	QSRR: quantitative structure-(chromatographic) retention relationships. 2007 , 107, 3212-46	372
159	Micellar solid-phase microextraction for determining partition coefficients of substituted polycyclic aromatic hydrocarbons in micellar media: possible prediction of hydrocarbon-micelle behaviour. 2007 , 387, 2271-81	5
158	Chemometric characterization of (chromatographic) lipophilicity parameters of newly synthesized s-triazine derivatives. 2008 , 22, 195-202	17
157	An advanced application of the quantitative structure-activity relationship concept in electrokinetic chromatography of metal complexes. 2008 , 29, 827-34	13
156	In vitro and in vivo studies of N,N'-bis[2(2-pyridyl)-methyl]pyridine-2,6-dicarboxamide/copper(II) and rheumatoid arthritis. 2008 , 27, 453-464	15
155	Quantitative structure-retention (property) relationships in micellar electrokinetic chromatography. 2008 , 1182, 1-24	54
154	Effect of n-octanol in the mobile phase on lipophilicity determination by reversed-phase high-performance liquid chromatography on a modified silica column. 2008 , 1209, 111-9	34
153	Determination of binding constants by affinity capillary electrophoresis, electrospray ionization mass spectrometry and phase-distribution methods. 2008 , 27, 738-748	120

152	High-throughput log P determination by ultraperformance liquid chromatography: a convenient tool for medicinal chemists. 2008 , 51, 396-9	28
151	Influence of lipophilia and of the vehicle used in the transdermal absorption of novel benzimidazole compounds with possible anti-HIV activity. 2008 , 13, 127-33	4
150	Quantitative structure-property relationships for octanol-water partition coefficients of polybrominated diphenyl ethers. 2008 , 72, 1602-1606	59
149	LogD: lipophilicity for ionisable compounds. 2008 , 72, 1401-1408	136
148	Hydrophobicity Parameter (log Kow) Estimation for Some Phenolic Compounds of Pharmaceutical Interest from Retention Studies with Mobile Phase Composition in Reversed-Phase Liquid Chromatography. 2008 , 43, 147-163	12
147	Characterization of the GEMINI C18 Column: Lipophilicity Measurement and LSER. 2008 , 31, 2198-2218	10
146	RPTLC determination of logPof structurally diverse neutral compounds. 2008 , 21, 143-149	12
145	RPTLC retention indices of basic and neutral drugs as surrogates of octanol-water distribution coefficients. Effect of buffer constituents and pH. 2009 , 22, 217-224	6
144	Normal-phase thin-layer chromatography of some angiotensin converting enzyme (ACE) inhibitors and their metabolites. 2009 , 74, 677-688	10
143	Application of Okik's Equation for Description of Solute Retention in RP HPLC and Calculation of Retention Factor in Water. 2009 , 32, 2779-2794	8
142	High throughput screening of physicochemical properties and in vitro ADME profiling in drug discovery. 2009 , 12, 315-29	72
141	Chromatography methods in investigation of lipophilicity of the biological active substances. 2009 , 63, 33-37	
140	High-throughput phase-distribution method to determine drug-cyclodextrin binding constants. 2009 , 98, 229-38	7
139	Analytical tools for the physicochemical profiling of drug candidates to predict absorption/distribution. 2009 , 394, 707-29	63
138	Fast log P determination by ultra-high-pressure liquid chromatography coupled with UV and mass spectrometry detections. 2009 , 394, 1919-30	31
137	Prediction of chemical absorption into and through the skin from cosmetic and dermatological formulations. 2009 , 160, 80-91	40
136	Improvement of the high-speed logD assay using an injection marker for the water plug aspiration/injection method. 2009 , 1216, 2984-8	12
135	In vitro and in vivo studies of the dermally absorbed Cu(II) complexes of N5O2 donor ligands □ Potential anti-inflammatory drugs. 2009 , 362, 125-135	15

134	Retention models for isocratic and gradient elution in reversed-phase liquid chromatography. 2009 , 1216, 1737-55	114
133	Determination of solute descriptors by chromatographic methods. 2009 , 652, 32-53	187
132	Lipophilicity of basic drugs measured by hydrophilic interaction chromatography. 2009 , 52, 3416-9	25
131	In Silico Tools and In Vitro HTS Approaches to Determine Lipophilicity During the Drug Discovery Process. 2009 , 91-116	
130	Optimization of Chromatographic Systems for Determination of Lipophilicity for Selected Isoquinoline Alkaloids. 2009 , 32, 2265-2280	5
129	Relationship between brain tissue partitioning and microemulsion retention factors of CNS drugs. 2009 , 52, 1693-700	44
128	Novel melphalan and chlorambucil derivatives of 2,2,6,6-tetramethyl-1-piperidinyloxy radicals: synthesis, characterization, and biological evaluation in vitro. 2010 , 58, 332-5	4
127	Unusual Retention Behavior of Some Cationic-Type Aldoximes Used as AChE Reactivators Under Ion-Pairing Liquid Chromatographic Mechanism. 2010 , 43, 1267-1276	6
126	High-throughput log P determination by MEEKC coupled with UV and MS detections. 2010 , 31, 952-64	25
125	Hydrophobicity/hydrophilicity descriptors obtained from extrapolated chromatographic retention data as modeling tools for biological distribution: application to some oxime-type acetylcholinesterase reactivators. 2010 , 52, 508-16	17
124	Characterization of Ascentis RP-Amide column: Lipophilicity measurement and linear solvation energy relationships. 2010 , 1217, 65-74	20
123	Properties of subcritical water as an eluent for reversed-phase liquid chromatography--disruption of the hydrogen-bond network at elevated temperature and its consequences. 2010 , 1217, 5769-75	13
122	. 2010 ,	3
121	Chromatography in the Millennium Perspectives. 2010 , 363-395	
120	QSAR in Chromatography: Quantitative Structure Retention Relationships (QSRRs). 2010 , 223-259	4
119	Recent Advances in QSAR Studies. 2010 ,	82
118	Application of successive projections algorithm (SPA) as a variable selection in a QSPR study to predict the octanol/water partition coefficients (Kow) of some halogenated organic compounds. 2010 , 2, 758	18
117	Advances in LC platforms for drug discovery. 2010 , 5, 475-89	20

116	Studies on log Po/w of quinoxaline di-N-oxides: a comparison of RP-HPLC experimental and predictive approaches. 2011 , 16, 7893-908	6
115	4-fluoro-2-methoxyphenol, an apocynin analog with enhanced inhibitory effect on leukocyte oxidant production and phagocytosis. 2011 , 660, 445-53	14
114	Recent application of analytical methods to phase I and phase II drugs development: a review. 2012 , 26, 283-300	27
113	Evaluation of the membrane permeability (PAMPA and skin) of benzimidazoles with potential cannabinoid activity and their relation with the Biopharmaceutics Classification System (BCS). 2011 , 12, 573-8	13
112	A novel method for the investigation of liquid/liquid distribution coefficients and interface permeabilities applied to the water-octanol-drug system. 2011 , 28, 2140-6	12
111	Lipophilicity of kinetically labile metal complexes through the example of antidiabetic Zn(II) and VO(IV) compounds. 2011 , 54, 1073-81	29
110	A new high-throughput method utilizing porous silica-based nano-composites for the determination of partition coefficients of drug candidates. 2011 , 34, 2505-12	2
109	The use of phospholipid modified column for the determination of lipophilic properties in high performance liquid chromatography. 2011 , 1218, 1211-8	13
108	Orthogonal chromatographic descriptors for modelling Caco-2 drug permeability. 2012 , 50, 175-83	5
107	Simultaneous multiple-development HPTLC quantification of water- and oil-soluble sunscreens. 2012 , 25, 344-348	7
106	Micellar electrokinetic chromatography: a new simple tool for the analysis of synthetic cannabinoids in herbal blends and for the rapid estimation of their logP values. 2012 , 1267, 198-205	27
105	Lipophilicity screening of novel drug-like compounds and comparison to clog P. 2012 , 1258, 161-7	19
104	RP TLC ASSAY OF THE LIPOPHILICITY OF NEW AZATHIOPRINE ANALOGS. 2012 , 35, 1686-1696	2
103	A spontaneous gold(I)-azide alkyne cycloaddition reaction yields gold-peptide bioconjugates which overcome cisplatin resistance in a p53-mutant cancer cell line. 2012 , 3, 2062	82
102	Molecular interaction fields based descriptors to interpret and compare chromatographic indexes. 2012 , 1252, 84-9	21
101	Comparative solution equilibrium studies of anticancer gallium(III) complexes of 8-hydroxyquinoline and hydroxy(thio)pyrone ligands. <i>Journal of Inorganic Biochemistry</i> , 2012 , 117, 189-97 ⁴⁻²	43
100	Complex-Formation Ability of Salicylaldehyde Thiosemicarbazone towards ZnII, CuII, FeII, FeIII and GaIII Ions. 2012 , 2012, 4036-4047	37
99	Quantification of Lipophilicity of 1,2,4-Triazoles Using Micellar Chromatography. 2012 , 75, 449-456	22

98	Determination of descriptors for fragrance compounds by gas chromatography and liquid-liquid partition. 2012 , 1235, 159-65	27
97	Exploring the interplay of physicochemical properties, membrane permeability and giardicidal activity of some benzimidazole derivatives. 2012 , 52, 193-204	26
96	A comparative study of the lipophilicity of 1,2,4-triazoles by reversed-phase and micellar TLC and OPLC. 2013 , 26, 153-159	5
95	Quantitative Structure Property (Retention) Relationships in Liquid Chromatography. 2013 , 385-405	3
94	Estimation of the environmental properties of compounds from chromatographic measurements and the solvation parameter model. 2013 , 1317, 85-104	120
93	Reversed-phase thin-layer chromatography technique for the comparison of the lipophilicity of selected non-steroidal anti-inflammatory drugs. 2013 , 85, 132-7	25
92	The Block Relevance (BR) analysis to aid medicinal chemists to determine and interpret lipophilicity. 2013 , 4, 1376	20
91	Determination of lipophilicity of γ -butyrolactone derivatives with anticonvulsant and analgesic activity using micellar electrokinetic chromatography. 2013 , 34, 3079-85	5
90	Solution equilibrium studies on anticancer ruthenium(II)- <i>p</i> -cymene complexes of 3-hydroxy-2(1H)-pyridones. 2013 , 734, 38-44	19
89	Hydrophilic Interaction Chromatography (HILIC) for Drug Discovery. 2013 , 169-193	1
88	^#.^t11: Statistical Evaluation and Validation of Quantitative Methods of Drug Analysis. 2013 , 216-221	
87	Critical review and interpretation of environmental data for volatile methylsiloxanes: partition properties. 2014 , 48, 11748-59	50
86	Retention-property relationships of 1,2,4-triazoles by micellar and reversed-phase liquid chromatography. 2014 , 37, 1419-28	13
85	Evaluation of the lipophilicity of selected sunscreens--a chemometric analysis of thin-layer chromatographic retention data. 2014 , 37, 3074-81	14
84	Evaluation of the three-phase equilibrium method for measuring temperature dependence of internally consistent partition coefficients (K(OW), K(OA), and K(AW)) for volatile methylsiloxanes and trimethylsilanol. 2014 , 33, 2702-10	38
83	Lipophilicity indices derived from the liquid chromatographic behavior observed under bimodal retention conditions (reversed phase/hydrophilic interaction): application to a representative set of pyridinium oximes. 2014 , 122, 172-9	11
82	Hydroxypyridinecarboxylic acid derivatives influencing metal ion levels in the brain: Equilibrium complexation studies with Cu(II) and Zn(II). 2014 , 67, 481-489	3
81	Introduction of polar groups on the naphthalene scaffold of molecular tongs inhibiting wild-type and mutated HIV-1 protease dimerization. 2014 , 5, 719-727	1

80	Synthesis and biological evaluation of glucosyl curcuminoids. 2014 , 347, 834-9	8
79	Chromatographic behavior of new antiepileptic active compounds on different reversed-phase materials. 2014 , 1338, 188-96	5
78	Solution speciation of potential anticancer metal complexes of salicylaldehyde semicarbazone and its bromo derivative. 2014 , 67, 242-252	26
77	Quantification of octanol/water partition coefficients of several aldehydes in a bubble column using selected ion flow tube mass spectrometry. 2014 , 367, 22-28	6
76	Lipophilicity of amyloid β -peptide 12-28 and 25-35 to unravel their ability to promote hydrophobic and electrostatic interactions. 2015 , 495, 179-185	6
75	Towards better understanding of lipophilicity: assessment of in silico and chromatographic logP measures for pharmaceutically important compounds by nonparametric rankings. 2015 , 115, 183-91	37
74	Confocal Raman Microscopy for in Situ Measurement of Octanol-Water Partitioning within the Pores of Individual C18-Functionalized Chromatographic Particles. 2015 , 87, 5340-7	12
73	Recent advances in lipophilicity measurement by reversed-phase high-performance liquid chromatography. 2015 , 68, 28-36	50
72	Exploring experimental and computational markers of cyclic peptides: Charting islands of permeability. 2015 , 97, 202-13	60
71	Lipophilicity study of fifteen fluoroquinolones by reversed-phase thin-layer chromatography. 2015 , 7, 3841-3848	8
70	Estimation of alkane-water logP for neutral, acidic, and basic compounds using an alkylated polystyrene-divinylbenzene high-performance liquid chromatography column. 2015 , 1417, 21-9	4
69	Carrier Mediated Distribution System (CAMDIS): a new approach for the measurement of octanol/water distribution coefficients. 2015 , 68, 68-77	18
68	Zone fluidics for measurement of octanol-water partition coefficient of drugs. 2015 , 860, 1-7	11
67	Electrophoretic Approaches for Log P Determination of Small Molecules and Polymers. 2016 , 1-25	
66	Multivariate assessment of lipophilicity scales-computational and reversed phase thin-layer chromatographic indices. 2016 , 127, 81-93	46
65	Solution equilibria and the X-ray structure of Cu(ii) complexation with 3-amino-N-(pyridin-2-ylmethyl)propanamide, a pseudo-mimic of human serum albumin. 2016 , 45, 17010-17019 ⁵	
64	Design, synthesis and evaluation of diphenyl ether analogues as antitubercular agents. 2016 , 6, 110571-110582	
63	Effects of polar β -receptor agonists designed for the periphery on ATP-induced Ca ²⁺ release from keratinocytes. 2016 , 7, 317-326	16

62	Limits of rapid log P determination methods for highly lipophilic and flexible compounds. 2016 , 915, 90-101	5
61	A Fast Chromatographic Method for Estimating Lipophilicity and Ionization in Nonpolar Membrane-Like Environment. 2016 , 13, 1100-10	23
60	High-throughput determination of octanol/water partition coefficients using a shake-flask method and novel two-phase solvent system. 2016 , 117, 338-44	20
59	Applications of the solvation parameter model in reversed-phase liquid chromatography. 2017 , 1486, 2-19	41
58	Medicating Patients During Extracorporeal Membrane Oxygenation: The Evidence is Building. 2017 , 389-400	
57	Lipophilicity estimation of statins as a decisive physicochemical parameter for their hepato-selectivity using reversed-phase thin layer chromatography. 2017 , 142, 7-14	25
56	Consensus-based comparison of chromatographic and computationally estimated lipophilicity of benzothiepine[3,2-c]pyridine derivatives as potential antifungal drugs. 2017 , 40, 2089-2096	13
55	DAPNe with micro-capillary separatory chemistry-coupled to MALDI-MS for the analysis of polar and non-polar lipid metabolism in one cell. 2017 , 28, 918-928	10
54	P and H NMR Studies of the Molecular Organization of Lipids in the Parallel Artificial Membrane Permeability Assay. 2017 , 14, 284-295	6
53	Physicochemical Parameters Affecting the Electrospray Ionization Efficiency of Amino Acids after Acylation. 2017 , 89, 9159-9166	23
52	Quantitative structure property (retention) relationships in liquid chromatography. 2017 , 553-572	3
51	Medicating patients during extracorporeal membrane oxygenation: the evidence is building. 2017 , 21, 66	56
50	Phospholipid bilayer affinities and solvation characteristics by electrokinetic chromatography with a nanodisc pseudostationary phase. 2017 , 38, 738-746	4
49	Permeation characteristics of hypericin across Caco-2 monolayers in the presence of single flavonoids, defined flavonoid mixtures or Hypericum extract matrix. 2019 , 71, 58-69	7
48	Comparative evaluation of new dihydropyrimidine and dihydropyridine derivatives perturbing mitotic spindle formation. 2018 , 10, 2395-2410	1
47	Confocal Raman Microscopy for in Situ Measurement of Phospholipid-Water Partitioning into Model Phospholipid Bilayers within Individual Chromatographic Particles. 2018 , 90, 7048-7055	6
46	Inhibitory Influence of Panax notoginseng Saponins on Aspirin Hydrolysis in Human Intestinal Caco-2 Cells. 2018 , 23,	10
45	Application of Capillary Electromigration Methods for Physicochemical Measurements. 2018 , 547-591	4

44	Evaluation of Hydrophobicity for Fatty Acids Using Reversed-phase Thin Layer Chromatography. 2019 , 68, 665-670	1
43	Simultaneous Determination and Quantification of Triterpene Saponins from Seeds Using UPLC-PDA-QTOF-MS/MS. 2019 , 24,	13
42	Drug Disposition and Pharmacotherapy in Neonatal ECMO: From Fragmented Data to Integrated Knowledge. 2019 , 7, 360	24
41	State of the art and prospects of methods for determination of lipophilicity of chemical compounds. 2019 , 113, 54-73	23
40	A New Straightforward Method for Lipophilicity (logP) Measurement using ¹⁹ F NMR Spectroscopy. 2019 ,	4
39	Reversed phase HPLC with high temperature ethanol/water mobile phases as a green alternative method for the estimation of octanol/water partition coefficients. 2019 , 1601, 243-254	8
38	Experimental lipophilicity for beyond Rule of 5 compounds. 2019 , 1, FDD10	11
37	Moderate to Severe Acute Respiratory Distress Syndrome Management Strategies: A Narrative Review. 2019 , 32, 347-360	2
36	Ventilatory and Pharmacotherapeutic Strategies for Management of Adult Patients on Extracorporeal Life Support. 2019 , 39, 355-368	4
35	Prediction of the Blood-Brain Barrier Permeability Using RP-18 Thin Layer Chromatography. 2019 , 17, 43-56	10
34	Study of pH and temperature effect on lipophilicity of catechol-containing antioxidants by reversed phase liquid chromatography. 2019 , 145, 380-387	6
33	Provitamin supramolecular polymer micelle with pH responsiveness to control release, bioavailability enhancement and potentiation of cytotoxic efficacy. 2019 , 173, 85-93	11
32	Design, synthesis and evaluation of antitubercular activity of Triclosan analogues. 2019 , 12, 3316-3323	15
31	Design, Synthesis, Biological Evaluation and Molecular Dynamic Simulation Studies of Diphenyl Ether Derivatives as Antitubercular and Antibacterial Agents. 2020 , 5, 201-210	2
30	LogP determination for highly lipophilic hydrogen-bonding anion receptor molecules. 2020 , 1132, 123-133	5
29	Multiphase Behavior of Tetraphenylethylene Derivatives with Different Polarities at High Pressures. 2020 , 124, 7263-7271	3
28	Determination of physicochemical properties of small molecules by reversed-phase liquid chromatography. 2020 , 1626, 461427	9
27	Experimental determination of phase ratio of C8 columns employing retention factors and octane-mobile phase partition coefficients of homologous series of linear alkylbenzenes. 2020 , 1634, 461668	4

26	In Vitro Antioxidant and Insulin Mimetic Activities of Heteroleptic Oxovanadium(IV) Complexes with Thiosemicarbazones and Naproxen. 2020 , 5, 6245-6254	3
25	Versatile in silico modeling of partition coefficients of organic compounds in polydimethylsiloxane using linear and nonlinear methods. 2020 , 399, 123012	7
24	Synthesis, structural characterization, in vitro DNA binding, and antitumor activity properties of Ru(II) compounds containing 2(2,6-dimethoxypyridine-3-yl)-1H-imidazo(4,5-f)[1, 10]phenanthroline. 2020 , 39, 760-791	3
23	Design, synthesis, evaluation, and molecular dynamic simulation of triclosan mimic diphenyl ether derivatives as antitubercular and antibacterial agents. 2020 , 31, 983-998	6
22	High-Throughput Liquid-Liquid Extractions with Nanoliter Volumes. 2020 , 92, 3189-3197	17
21	Acyl Derivatives of Eudesmanolides To Boost their Bioactivity: An Explanation of Behavior in the Cell Membrane Using a Molecular Dynamics Approach. 2021 , 16, 1297-1307	1
20	Continuous and automated slug flow nanoextraction for rapid partition coefficient measurement. 2021 , 146, 5722-5731	1
19	Assessment of Lipophilicity Descriptors of Selected NSAIDs Obtained at Different TLC Stationary Phases. 2021 , 13,	2
18	A thiol-responsive and self-immolative podophyllotoxin prodrug for cancer therapy. 2021 , 71, 153044	1
17	Highly Hydrophilic and Lipophilic Derivatives of Bile Salts. 2021 , 22,	1
16	Predicting Octanol/Water Partition Coefficients of Fluorinated Drug-Like Molecules: A Combined Experimental and Theoretical Study. 2020 , 73, 677	4
15	Analytical Utility of Liposomes: From Past to Present. 2016 , 1-82	2
14	Standardized reversed-phase thin-layer chromatographic study of the lipophilicity of five anti-diabetic thiazolidinediones. 2007 , 20, 261-265	6
13	Bio-organometallic Peptide Conjugates: Recent Advances in Their Synthesis and Prospects for Biomedical Application. 2020 , 24, 2508-2523	3
12	Chromatographic evaluation of lipophilicity for angiotensin II AT1 receptor antagonists. 2008 , 21, 23-29	
11	Chapter 2 Solubility Determinations for Pharmaceutical API. 2017 , 19-84	
10	Estimation of lipophilicity data for derivatives of alkandiamine-N,NEdi-2-(3-cyclohexyl) propanoic acid with potential antineoplastic activity, by UHPLC-MS method. 2017 , 402-409	
9	Evaluation of a Semi-Empirical Approach to Correlation of Retention with Octanol-Water Partition Coefficients by Use of Adjustable Hydrogen-Bonding Indicator Variables. 2005 , 61, 479	

8	Lipophilicity Examination of Some ACE inhibitors and Hydrochlorothiazide on Cellulose in RP Thin-Layer Chromatography. 2012 , 11, 763-70		8
7	CHAPTER 5. Biophysics of Flavour Perception. 2022 , 109-136		
6	A DFT study of the chemical reactivity properties, spectroscopy and bioactivity scores of bioactive flavonols. 2022 , 1210, 113658		1
5	The coordination modes of (thio)semicarbazone copper(II) complexes strongly modulate the solution chemical properties and mechanism of anticancer activity.. <i>Journal of Inorganic Biochemistry</i> , 2022 , 231, 111786	4.2	6
4	Review on development of ionic liquids in lignocellulosic biomass refining. <i>Journal of Molecular Liquids</i> , 2022 , 359, 119326	6	1
3	Recent advances for estimating environmental properties for small molecules from chromatographic measurements and the solvation parameter model. 2023 , 1687, 463682		1
2	Red-emissive Dual-state Fluorogenic Probe for Wash-free Imaging of Lipid Droplets in Living Cells and Fatty Liver Tissues. 2023 , 18,		0
1	Optimizing pharmacotherapy regimens in adult patients receiving extracorporeal membrane oxygenation: A narrative review for clinical pharmacists.		0