Comparison of bioelectrical impedance analysis and due the assessment of appendicular body composition in an

European Journal of Clinical Nutrition 57, 1068-1072 DOI: 10.1038/sj.ejcn.1601643

Citation Report

#	Article	IF	CITATIONS
1	Cross-calibration of eight-polar bioelectrical impedance analysis versus dual-energy X-ray absorptiometry for the assessment of total and appendicular body composition in healthy subjects aged 21-82 years. Annals of Human Biology, 2003, 30, 380-391.	1.0	284
2	Bioelectrical impedance analysis?part I: review of principles and methods. Clinical Nutrition, 2004, 23, 1226-1243.	5.0	2,089
3	Bioelectrical impedance analysis—part II: utilization in clinical practice. Clinical Nutrition, 2004, 23, 1430-1453.	5.0	1,643
4	The Validity of Bioelectrical Impedance Models in Clinical Populations. Nutrition in Clinical Practice, 2004, 19, 433-446.	2.4	175
5	Improvement of Nutritional Status as Assessed by Multifrequency BIA During 15 Weeks of Refeeding in Adolescent Girls with Anorexia Nervosa. Journal of Nutrition, 2004, 134, 3026-3030.	2.9	44
6	Body water distribution in severe obesity and its assessment from eight-polar bioelectrical impedance analysis. European Journal of Clinical Nutrition, 2005, 59, 155-160.	2.9	137
7	Accuracy of eight-polar bioelectrical impedance analysis for the assessment of total and appendicular body composition in peritoneal dialysis patients. European Journal of Clinical Nutrition, 2005, 59, 932-937.	2.9	54
8	Phase angle is a predictor of basal metabolic rate in female patients with anorexia nervosa. Physiological Measurement, 2005, 26, S145-S152.	2.1	24
9	Sarcopenia is predictive of nosocomial infection in care of the elderly. British Journal of Nutrition, 2006, 96, 895-901.	2.3	208
10	Evaluation of air-displacement plethysmography and bioelectrical impedance analysis vs dual-energy X-ray absorptiometry for the assessment of fat-free mass in elderly subjects. European Journal of Clinical Nutrition, 2008, 62, 1282-1286.	2.9	14
11	Body composition changes in female adolescents with anorexia nervosa. American Journal of Clinical Nutrition, 2009, 89, 1005-1010.	4.7	35
12	Cross-validation of bioelectrical impedance analysis for the assessment of body composition in a representative sample of 6- to 13-year-old children. European Journal of Clinical Nutrition, 2009, 63, 619-626.	2.9	73
13	Crossâ€calibration of multiâ€frequency bioelectrical impedance analysis with eightâ€point tactile electrodes and dualâ€energy Xâ€ray absorptiometry for assessment of body composition in healthy children aged 6–18â€fyears. Pediatrics International, 2009, 51, 263-268.	0.5	119
14	Underweight patients with anorexia nervosa: Comparison of bioelectrical impedance analysis using five equations to dual X-ray absorptiometry. Clinical Nutrition, 2011, 30, 746-752.	5.0	29
15	Anorexia nervosa and nutritional assessment: contribution of body composition measurements. Nutrition Research Reviews, 2011, 24, 39-45.	4.1	22
16	Comparison of three segmental multifrequency bioelectrical impedance techniques in healthy adults. Annals of Human Biology, 2012, 39, 468-478.	1.0	9
17	New Application of Bioelectrical Impedance Analysis by the Back Propagation Artificial Neural Network Mathematically Predictive Model of Tissue Composition in the Lower Limbs of Elderly People. International Journal of Gerontology, 2012, 6, 20-26.	0.6	10
18	Bioelectrical Impedance Analysis in a Mathematical Model for Estimating Fat-free Mass in Multiple Segments in Elderly Taiwanese Males. International Journal of Gerontology, 2012, 6, 273-277.	0.6	9

CITATION REPORT

#	Article	IF	CITATIONS
19	Development and Validation of a Predictive Equation for Lean Body Mass in Children and Adolescents. Annals of Human Biology, 2012, 39, 171-182.	1.0	59
20	Development and Evaluation of a Bio-ion Measurement System on Acupoints for Meridian Diagnosis. JAMS Journal of Acupuncture and Meridian Studies, 2013, 6, 110-118.	0.7	5
21	Validity of Standing Posture Eight-electrode Bioelectrical Impedance to Estimate Body Composition in Taiwanese Elderly. International Journal of Gerontology, 2014, 8, 137-142.	0.6	11
22	Body composition in young female eating-disorder patients with severe weight loss and controls: evidence from the four-component model and evaluation of DXA. European Journal of Clinical Nutrition, 2015, 69, 1330-1335.	2.9	13
23	Changes in lean and skeletal muscle body mass in adult females with anorexia nervosa before and after weight restoration. Clinical Nutrition, 2017, 36, 170-178.	5.0	20
24	Prevalence of sarcopenia in communityâ€dwelling older people of Mexico City using the EGWSOP (European Working Group on Sarcopenia in Older People) diagnostic criteria. JCSM Clinical Reports, 2017, 2, 1-9.	1.3	5
25	Prediction of body composition in anorexia nervosa: Results from a retrospective study. Clinical Nutrition, 2018, 37, 1670-1674.	5.0	19
26	Comparison of body composition assessment by DXA and BIA according to the body mass index: A retrospective study on 3655 measures. PLoS ONE, 2018, 13, e0200465.	2.5	168
27	Validity of Bioimpedance Equations to Evaluate Fat-Free Mass and Muscle Mass in Severely Malnourished Anorectic Patients. Journal of Clinical Medicine, 2020, 9, 3664.	2.4	6
28	Body composition in subjects with anorexia nervosa: bioelectrical impedance analysis and dual-energy X-ray absorptiometry. Eating and Weight Disorders, 2012, 17, e298-303.	2.5	7
29	Estimation of segmental fat free mass in Taiwanese elderly females by bioelectrical impedance analysis with new mathematical model. African Journal of Biotechnology, 2011, 10, .	0.6	1
30	Comparison of Dual-Energy X-ray Absorptiometry and Bioelectrical Impedance Analysis in the Assessment of Body Composition in Women with Anorexia Nervosa upon Admission and Discharge from an Inpatient Specialist Unit. International Journal of Environmental Research and Public Health, 2021. 18. 11388.	2.6	8
31	Accuracy of bioimpedance equations for measuring body composition in a cohort of 2134 patients with obesity. Clinical Nutrition, 2022, 41, 2013-2024.	5.0	3