

A NEW NEMATODE INFECTION OF MAN

JAMA - Journal of the American Medical Association

72, 567

DOI: [10.1001/jama.1919.02610080033010](https://doi.org/10.1001/jama.1919.02610080033010)

Citation Report

#	ARTICLE	IF	CITATIONS
1	A Mouse Oxyurid, <i>Syphacia obvelata</i> , as a Parasite of Man. <i>Journal of Parasitology</i> , 1919, 6, 89.	0.7	21
2	Institutional Hookworm Disease in a Non-Endemic Region. <i>The Journal of Hygiene</i> , 1923, 22, 77-88.	0.9	4
3	The mass diagnosis of ankylostome infestation (parts VIII to XIII). <i>Transactions of the Royal Society of Tropical Medicine and Hygiene</i> , 1924, 18, 278-310.	1.8	2
4	Pseudoparasites in the Faeces of Man. <i>Journal of the Royal Society of Medicine</i> , 1926, 19, 14-18.	0.1	1
5	Inheritance of Resistance to the Wartelle Race of Root-knot Nematode in Soybeans 1. <i>Crop Science</i> , 1976, 16, 783-785.	1.8	5
6	Species and distribution of Heteroderidae and Meloidogynidae (Nematoda: Tylenchida) in Canada. <i>Canadian Journal of Plant Pathology</i> , 1986, 8, 170-184.	1.4	7
7	Interspecific differences in the fine structure of the body cuticle of Trichodoridae Thorne, 1935 (Nematoda: Diphtherophorina) and review of anchoring structures of the epidermis. <i>Nematology</i> , 2001, 3, 525-533.	0.6	15
8	Meloidogyne brasiliensis n. sp. (Nematoda: Meloidogynidae), a root-knot nematode parasitising tomato cv. Rossol in Brazil. <i>Nematology</i> , 2002, 4, 629-643.	0.6	7
9	Characterisation of Meloidogyne spp. (Tylenchida: Meloidogynidae) from coffee plantations in Central America and Brazil. <i>Nematology</i> , 2004, 6, 193-204.	0.6	18
10	Literary Collections in Tang Dynasty China. <i>T'oung Pao</i> , 2007, 93, 1-52.	0.1	4
11	Screening of <i>Tagetes patula</i> L. on different populations of Meloidogyne. <i>Crop Protection</i> , 2008, 27, 96-100.	2.1	23
12	Additional information on <i>Meloidogyne inornata</i> Lordello, 1956 (Tylenchida: Meloidogynidae) and its characterisation as a valid species. <i>Nematology</i> , 2008, 10, 123-136.	0.6	12
13	Meloidogyne phaseoli n. sp. (Nematoda: Meloidogynidae), a root-knot nematode parasitising bean in Brazil. <i>Nematology</i> , 2008, 10, 525-538.	0.6	6
14	Meloidogyne pisi n. sp. (Nematoda: Meloidogynidae), a root-knot nematode parasitising pea in Brazil. <i>Nematology</i> , 2008, 10, 479-493.	0.6	6
15	Taxonomy of Meloidogyne (Nematoda: Heteroderidae) with descriptions of four new species*. <i>Transactions of the Zoological Society of London</i> , 1968, 31, 263-401.	2.6	77
16	Histopathological changes induced by <i>Meloidogyne incognita</i> in some ornamental plants. <i>Crop Protection</i> , 2014, 65, 216-220.	2.1	12
17	A new root-knot nematode <i>Meloidogyne spartelensis</i> n. sp. (Nematoda: Meloidogynidae) in Northern Morocco. <i>European Journal of Plant Pathology</i> , 2015, 143, 25-42.	1.7	16
18	Passage of Meloidogyne Eggs in Human Stool: Forgotten, but Not Gone. <i>Journal of Clinical Microbiology</i> , 2015, 53, 1458-1459.	3.9	11

#	ARTICLE	IF	CITATIONS
19	Comparison of concomitant and sequential inoculation of <i>Steinernema</i> sp. in the management of root-knot (<i>Meloidogyne incognita</i>) nematode infecting eggplant (<i>Solanum melongena</i>). <i>Archives of Phytopathology and Plant Protection</i> , 2015, 48, 277-287.	1.3	0
20	Infection by <i>Meloidogyne javanica</i> does not breakdown resistance to the defoliating pathotype of <i>Verticillium dahliae</i> in selected clones of wild olive. <i>Scientia Horticulturae</i> , 2016, 199, 149-157.	3.6	10
21	Fungal root endophytes of tomato from Kenya and their nematode biocontrol potential. <i>Mycological Progress</i> , 2016, 15, 1.	1.4	43
22	Evaluation of nematicides for southern root-knot nematode management in lima bean. <i>Crop Protection</i> , 2017, 96, 151-157.	2.1	54
23	<i>Meloidogyne aberrans</i> sp. nov. (Nematoda: Meloidogynidae), a new root-knot nematode parasitizing kiwifruit in China. <i>PLoS ONE</i> , 2017, 12, e0182627.	2.5	16
24	Morphological and morphometrical identification of <i>Meloidogyne</i> populations from various crop production areas in South Africa with emphasis on <i>M. enterolobii</i> . <i>Zootaxa</i> , 2019, 4658, zootaxa.4658.2.3.	0.5	7
25	< i>Meloidogyne daklakensis</i> n. sp. (Nematoda: Meloidogynidae), a new root-knot nematode associated with Robusta coffee (< i>Coffea canephora</i> Pierre ex A. Froehner) in the Western Highlands, Vietnam. <i>Journal of Helminthology</i> , 2019, 93, 242-254.	1.0	22
26	Intestinal helminthic parasites of rodents in the central region of Iran: first report of a capillariid nematode from <i>Dryomys nitedula</i> . <i>BMC Research Notes</i> , 2020, 13, 461.	1.4	5
27	Degree of resistance of < i>Solanum torvum</i> cultivars to < i>Mi-1.2</i>-virulent and avirulent isolates of < i>Meloidogyne incognita</i>, < i>Meloidogyne javanica</i>, and < i>Meloidogyne luci</i>. <i>Journal of Nematology</i> , 2021, 53, 1-7.	0.9	12
28	First report and new molecular and morphological characterizations of root-knot nematode, < i>Meloidogyne javanica</i>, infecting ginger and long coriander in Vietnam. <i>Journal of Nematology</i> , 2021, 53, 1-8.	0.9	3
29	First report of root-knot nematode, < i>Meloidogyne incognita</i> on calendula in Turkey. <i>Journal of Nematology</i> , 2021, 53, 1-5.	0.9	2
30	Nematodes Associated with Stone Fruits and Their Management Strategies. , 2021, , 423-436.	1	
31	Biodisinfestation With Agricultural By-Products Developed Long-Term Suppressive Soils Against <i>Meloidogyne incognita</i> in Lettuce Crop. <i>Frontiers in Sustainable Food Systems</i> , 2021, 5, .	3.9	4
32	The weed <i>Amaranthus palmeri</i> is a host of <i>Meloidogyne incognita</i> in Uruguay. <i>Rhizosphere</i> , 2021, 19, 100400.	3.0	3
33	Taxonomy, identification and principal species.. , 2009, , 55-97.		171
34	Quantitative Field Testing <i>Heterodera glycines</i> from Metagenomic DNA Samples Isolated Directly from Soil under Agronomic Production. <i>PLoS ONE</i> , 2014, 9, e89887.	2.5	9
35	Reproduction of < i>Meloidogyne incognita</i> Race 3 on Flue-cured Tobacco Homozygous for < i>Rk1</i> and/or < i>Rk2</i> Resistance Genes. <i>Journal of Nematology</i> , 2016, 48, 79-86.	0.9	6
36	New Hosts and Records in Portugal for the Root-Knot Nematode < i>Meloidogyne luci</i>. <i>Journal of Nematology</i> , 2019, 51, 1-4.	0.9	15

#	ARTICLE	IF	CITATIONS
37	First report of cultivated Cretan mountain tea (<i>Sideritis syriaca</i>) as a host of <i>Meloidogyne hapla</i> and <i>M. javanica</i> in Crete, with some additional records on the occurrence of <i>Meloidogyne</i> species in Greece. <i>Journal of Nematology</i> , 2019, 51, 1-4.	0.9	4
38	First Report of <i>Scutellonema brachyurus</i> (Steiner, 1938) Andrassy, 1958 and Occurrence of <i>Meloidogyne incognita</i> (Kofoid & White, 1919) Chitwood, 1949 in Belgium. <i>Journal of Nematology</i> , 2019, 51, 1-6.	0.9	3
39	Oxyurid nematodes of pet rodents in Slovakia - a neglected zoonotic threat. <i>Brazilian Journal of Veterinary Parasitology</i> , 2020, 29, e014319.	0.7	0
40	First report of southern root-knot nematode, <i>Meloidogyne incognita</i> , infecting <i>Brassica nigra</i> in Peru. <i>Journal of Nematology</i> , 2020, 52, 1-13.	0.9	1
41	First report of southern root-knot nematode, <i>Meloidogyne incognita</i> , infecting pomegranate, <i>Punica granatum</i> , in Peru. <i>Journal of Nematology</i> , 2020, 52, 1-3.	0.9	1
42	Chenopodium album is a weed host of <i>Meloidogyne incognita</i> (Nematoda: Meloidogynidae) in Peru. <i>Journal of Nematology</i> , 2020, 52, 1-4.	0.9	0
43	Vertebrates as uninfected disseminators of helminth eggs and larvae. <i>Advances in Parasitology</i> , 2022, 115, 45-170.	3.2	2
44	Evaluation of bacterial formulations as potential biocontrol agents against the southern root-knot nematode, <i>Meloidogyne incognita</i> . <i>Egyptian Journal of Biological Pest Control</i> , 2022, 32, .	1.8	6
45	Integrative taxonomy of the aggressive pest <i>Meloidogyne enterolobii</i> and molecular phylogeny of <i>Meloidogyne</i> spp. based on five gene regions. <i>Australasian Plant Pathology</i> , 2022, 51, 345-358.	1.0	6
46	Türkiye'de Melisada (<i>Melissa officinalis</i> L.) Bulunan Kükür Nematodu <i>Meloidogyne arenaria</i> 'nın Ölk Kaydı. Kahramanmaraş Sıratı Ümmetü'l-Mam Aöeniversitesi Tarım Ve Doğa Dergisi, 2022, 25, 176-179.	0.7	0
47	<i>Belonolaimus longicaudatus</i> Host Status and Pathogenicity on Sweetpotato. <i>Journal of Nematology</i> , 2022, 54, .	0.9	0
48	Agricultural weeds maintain populations of <i>Meloidogyne</i> spp. root-knot nematodes. <i>Rhizosphere</i> , 2022, 23, 100576.	3.0	0
49	A quantitative and qualitative assessment of sugar beet genotype resistance to root-knot nematode, <i>Meloidogyne incognita</i> . <i>Frontiers in Plant Science</i> , 0, 13, .	3.6	2
51	Galls induced by a root-knot nematode in <i>Petroselinum crispum</i> (Mill.): impacts on host development, histology, and cell wall dynamics. <i>Protoplasma</i> , 0, , .	2.1	1
52	<i>Meloidogyne paramali</i> n. sp. (Nematoda: Meloidogyninae) and First Report of <i>M. marylandi</i> in maple and yacca tree from Japan. <i>Journal of Nematology</i> , 2023, 55, .	0.9	0
53	Plant parasitic nematodes associated with cucumber cultivation in Lorestan province of Iran. <i>Plant Pathology Science</i> , 2022, 11, 73-82.	0.2	0
54	Efficiency of Vinasse Application on Root-Knot Nematodes in Soybean. <i>Agronomy</i> , 2023, 13, 2719.	3.0	0
55	Detection and molecular characterization of the northern root-knot nematode, <i>Meloidogyne hapla</i> , infesting a tree tomato field in Ecuador. <i>Australasian Plant Disease Notes</i> , 2023, 18, .	0.7	0

#	ARTICLE	IF	CITATIONS
56	Role of Phenolics in Establishing Mycorrhizal Association in Plants for Management of Biotic Stress. , 2024, , 35-74.		0