

CITATION REPORT

List of articles citing

Family presence during cardiopulmonary resuscitation and invasive procedures: practices of critical care and emergency nurses

DOI: 10.1067/men.2003.100

Journal of Emergency Nursing, 2003, 29, 208-21.

Source: <https://exaly.com/paper-pdf/35386387/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
662	Family member presence during resuscitation in the emergency department: An Australian perspective. 2004 , 16, 295-308		20
661	Emergency: family-centered care in the ED. 2004 , 104, 59-62		4
660	Report of the National Consensus Conference on Family Presence During Pediatric Cardiopulmonary Resuscitation and Procedures. <i>Pediatric Emergency Care</i> , 2005 , 21, 787-91	1.4	42
659	With you until the end: family presence during failed resuscitation. 2005 , 10, 204-7		15
658	A European survey of critical care nurses' attitudes and experiences of having family members present during cardiopulmonary resuscitation. <i>International Journal of Nursing Studies</i> , 2005 , 42, 557-68	5.8	82
657	Family presence during invasive procedures and resuscitation in the emergency department: a review of the literature. <i>Journal of Emergency Nursing</i> , 2005 , 31, 58-72; quiz 119	1.3	33
656	Practice Alert: Family Presence During CPR and Invasive Procedures. 2005 , 24, 49-51		2
655	The public's attitude and perception concerning witnessed cardiopulmonary resuscitation. 2006 , 34, 2925-8		42
654	Family presence during resuscitation: A randomised controlled trial of the impact of family presence. <i>Australasian Emergency Nursing Journal</i> , 2006 , 8, 139-147		64
653	Report of the National Consensus Conference on Family Presence During Pediatric Cardiopulmonary Resuscitation and Procedures. <i>Journal of Emergency Nursing</i> , 2006 , 32, 23-9	1.3	43
652	Effects of family presence during resuscitation and invasive procedures in a pediatric emergency department. <i>Journal of Emergency Nursing</i> , 2006 , 32, 225-33	1.3	123
651	Narratives about resuscitation--family members differ about presence. <i>European Journal of Cardiovascular Nursing</i> , 2006 , 5, 68-74	3.3	31
650	Should family members be present during cardiopulmonary resuscitation? A review of the literature. 2007 , 24, 311-7		48
649	The introduction of family presence evidence-based practice into a baccalaureate nursing curriculum. 2007 , 30, 364-71		7
648	Family experiences during resuscitation at a children's hospital emergency department. 2007 , 21, 217-25		53
647	Parent presence during complex invasive procedures and cardiopulmonary resuscitation: a systematic review of the literature. 2007 , 120, 842-54		97
646	Family presence during CPR: a study of the experiences and opinions of Turkish critical care nurses. <i>International Journal of Nursing Studies</i> , 2007 , 44, 83-92	5.8	57

645	Paediatric critical care nurses' attitudes and experiences of parental presence during cardiopulmonary resuscitation: a European survey. <i>International Journal of Nursing Studies</i> , 2007 , 44, 1238-49	5.8	47
644	Emergency nurses' current practices and understanding of family presence during CPR. <i>Journal of Emergency Nursing</i> , 2007 , 33, 433-40	1.3	54
643	Part B: A survey of staff attitudes immediately post-resuscitation to Family Presence during resuscitation. <i>Australasian Emergency Nursing Journal</i> , 2008 , 11, 114-122		19
642	Family presence during resuscitation and invasive procedures: the nurse experience. 2009 , 19, 1431-42		29
641	The views and preferences of resuscitated and non-resuscitated patients towards family witnessed resuscitation: a qualitative study. <i>International Journal of Nursing Studies</i> , 2009 , 46, 12-21	5.8	54
640	A study of Turkish critical care nurses' perspectives regarding family-witnessed resuscitation. 2009 , 18, 2907-15		36
639	Family presence during resuscitation: Canadian critical care nurses' perspectives. 2009 , 41, 233-40		40
638	Partners in care: implementing a policy on family member passengers. 2009 , 28, 31-6		1
637	Effects of family-witnessed resuscitation after trauma prior to hospitalization. <i>Journal of Trauma Nursing: the Official Journal of the Society of Trauma Nurses</i> , 2010 , 17, 11-8	1	13
636	Health care providers' attitudes regarding family presence during resuscitation of adults: an integrated review of the literature. <i>Clinical Nurse Specialist</i> , 2010 , 24, 161-74	0.6	26
635	Family presence during trauma activations and medical resuscitations in a pediatric emergency department: an evidence-based practice project. <i>Journal of Emergency Nursing</i> , 2010 , 36, 115-21	1.3	25
634	Health care providers' evaluations of family presence during resuscitation. <i>Journal of Emergency Nursing</i> , 2010 , 36, 524-33	1.3	27
633	Family Presence During Adult Resuscitation. <i>Advanced Emergency Nursing Journal</i> , 2010 , 32, 46-58	0.8	15
632	Ventilator-associated pneumonia caused by <i>Pseudomonas aeruginosa</i> and respiratory colonization by <i>Candida</i> spp. <i>Critical Care</i> , 2010 , 14, P87	10.8	78
631	Abstracts of the 30th International Symposium on Intensive Care and Emergency Medicine. Brussels, Belgium. March 9-12, 2010. <i>Critical Care</i> , 2010 , 14 Suppl 1, P1-602	10.8	
630	Sodium 4-phenylbutylate protects against myocardial ischemia-reperfusion injury by reducing unfolded protein response-mediated apoptosis in mice. <i>Critical Care</i> , 2010 , 14, P2	10.8	78
629	Time-dependent effects of intravenous H ₂ S during long-term, resuscitated porcine hemorrhagic shock. <i>Critical Care</i> , 2010 , 14, P3	10.8	1
628	An ovine intensive care model of septic shock. <i>Critical Care</i> , 2010 , 14, P4	10.8	1

627	Effects of temperature and H ₂ S inhalation on glucose metabolism in murine resuscitated septic shock. <i>Critical Care</i> , 2010 , 14, P5	10.8	78
626	Mitochondrial respiration and cytochrome c inhibition by sulfide in peritoneal macrophages in vitro: effects of temperature and pH. <i>Critical Care</i> , 2010 , 14, P6	10.8	2
625	Alterations of caspase 9 mRNA gene expression in survivors and nonsurvivors of severe sepsis. <i>Critical Care</i> , 2010 , 14, P7	10.8	78
624	The TLR4 antagonist CRX-526 reduces LPS-induced leukocyte activation and improves capillary perfusion of the rat intestine. <i>Critical Care</i> , 2010 , 14, P8	10.8	78
623	Effects of lipopolysaccharide on isolated muscle mitochondrial respiration are dose and time dependent. <i>Critical Care</i> , 2010 , 14, P9	10.8	78
622	Clinical severity and local inflammatory responses in animal models of sepsis. <i>Critical Care</i> , 2010 , 14, P10	10.8	78
621	Adenosine increases during human experimental endotoxemia, but does not influence the immune response and subsequent organ injury. <i>Critical Care</i> , 2010 , 14, P11	10.8	78
620	Bacterial load plays a crucial role for survival in experimental peritonitis and modulates immunoparalysis of monocytes. <i>Critical Care</i> , 2010 , 14, P12	10.8	78
619	Decreased whole blood TNF α production capacity after acute alcohol exposure and LPS stimulation ex vivo. <i>Critical Care</i> , 2010 , 14, P13	10.8	3
618	Leukocyte immunophenotyping: methodological aspects. <i>Critical Care</i> , 2010 , 14, P14	10.8	78
617	Phenotypical analysis of peripheral human T lymphocytes in early sepsis. <i>Critical Care</i> , 2010 , 14, P15	10.8	78
616	CD64, a marker of leucocyte activation kinetics after uncomplicated cardiac surgery. <i>Critical Care</i> , 2010 , 14, P16	10.8	78
615	Interferon-gamma reverses sepsis-induced immunoparalysis of monocytes in vitro. <i>Critical Care</i> , 2010 , 14, P17	10.8	78
614	Early hydrocortisone treatment counteracts circulatory derangement but not cytokine response in porcine endotoxemia. <i>Critical Care</i> , 2010 , 14, P18	10.8	78
613	CD40L is selectively expressed on platelets from thrombocytopenic septic patients. <i>Critical Care</i> , 2010 , 14, P19	10.8	1
612	Neuronal NOS-inhibition in the setting of septic cardiomyopathy. <i>Critical Care</i> , 2010 , 14, P20	10.8	78
611	Urinary albumin excretion is elevated in sepsis, but does not correlate with circulating VEGF-A levels. <i>Critical Care</i> , 2010 , 14, P21	10.8	1
610	High mobility group box protein-1 preconditions isolated rat hearts and protects human pulmonary and renal cells against hypoxic injury. <i>Critical Care</i> , 2010 , 14, P22	10.8	78

609	Intravenous immunoglobulins prevent breakdown of the blood-brain barrier in experimental sepsis. <i>Critical Care</i> , 2010 , 14, P24	10.8	1
608	Early use of immunoglobulin in septic shock. <i>Critical Care</i> , 2010 , 14, P25	10.8	1
607	Low levels of immunoglobulin G in patients with sepsis or septic shock: a signum mali ominis?. <i>Critical Care</i> , 2010 , 14, P26	10.8	2
606	Background factors in patients receiving immunoglobulin administration and changes in sepsis markers. <i>Critical Care</i> , 2010 , 14, P27	10.8	78
605	C1-esterase inhibitor (C1INH) implication in systemic inflammation in sepsis. <i>Critical Care</i> , 2010 , 14, P28	10.8	1
604	Prognostic value of B-type natriuretic peptide in critically ill patients with new onset of fever: preliminary study. <i>Critical Care</i> , 2010 , 14, P29	10.8	78
603	Microalbuminuria evaluated as a biomarker in patients with septic shock. <i>Critical Care</i> , 2010 , 14, P30	10.8	78
602	Evaluation of automated hematologic VCS parameters in severe sepsis and septic shock: a case-control study. <i>Critical Care</i> , 2010 , 14, P31	10.8	1
601	Use of CD64 for the diagnosis of sepsis: a case-control study. <i>Critical Care</i> , 2010 , 14, P32	10.8	1
600	Endotoxin assay with endotoxin scattering photometry is one of the diagnostic markers of sepsis. <i>Critical Care</i> , 2010 , 14, P33	10.8	78
599	Immature platelet fraction as predictive index of sepsis. <i>Critical Care</i> , 2010 , 14, P34	10.8	1
598	Serial estimation of cytokine markers and their correlation with mortality in elderly patients with sepsis. <i>Critical Care</i> , 2010 , 14, P35	10.8	78
597	Influence of TIMP-1/MMP-9 ratio on the severity and mortality in sepsis. <i>Critical Care</i> , 2010 , 14, P36	10.8	78
596	Cytokine response in severe sepsis: predicting and modelling the course of illness. <i>Critical Care</i> , 2010 , 14, P37	10.8	78
595	Time course of IL-6 and LBP, candidate biomarkers of sepsis in surgical critical care. <i>Critical Care</i> , 2010 , 14, P38	10.8	78
594	Prognostic value of melatonin in patients with sepsis: a comparison of survivors and nonsurvivors. <i>Critical Care</i> , 2010 , 14, P39	10.8	1
593	Serum C-reactive protein as a prognostic variable in elective surgery ICU patients: especially valuable following esophagectomy. <i>Critical Care</i> , 2010 , 14, P40	10.8	78
592	Role of procalcitonin as a diagnostic and prognostic marker of infectious diseases in the emergency department: preliminary data. <i>Critical Care</i> , 2010 , 14, P41	10.8	78

591	Evaluation of procalcitonin at James Paget Intensive Care Unit. <i>Critical Care</i> , 2010 , 14, P42	10.8	78
590	Procalcitonin facing the postoperative culture results after liver transplantation. <i>Critical Care</i> , 2010 , 14, P43	10.8	78
589	Efficiency diagnostic and advantages of procalcitonin and C-reactive protein in the early diagnosis of sepsis. <i>Critical Care</i> , 2010 , 14, P45	10.8	78
588	Biomarkers of sepsis in burn patients. <i>Critical Care</i> , 2010 , 14, P46	10.8	1
587	Age-related differences of outcomes in patients hospitalized with community-acquired pneumonia are not explained by differences in immune response. <i>Critical Care</i> , 2010 , 14, P47	10.8	78
586	A multicentre study of bacteraemia using a new commercial universal 16S rDNA PCR test. <i>Critical Care</i> , 2010 , 14, P49	10.8	78
585	Time to antibiotics in sepsis. <i>Critical Care</i> , 2010 , 14, P50	10.8	4
584	Utility of an antibiotic guideline in hospital-associated infections. <i>Critical Care</i> , 2010 , 14, P51	10.8	78
583	Assessment of antibiotic prescriptions in two French intensive care units. <i>Critical Care</i> , 2010 , 14, P52	10.8	78
582	Occurrence of alert pathogens in the clinical materials and consumption of antibiotics in the ICU, in the years 2007 and 2008. <i>Critical Care</i> , 2010 , 14, P53	10.8	78
581	Treatment cost of secondary peritonitis in Germany: a comparative study of medical cost incurred for tigecycline therapy and standard regimens. <i>Critical Care</i> , 2010 , 14, P54	10.8	78
580	Efficacy and safety of once daily dosing of colistin to critically ill patients. <i>Critical Care</i> , 2010 , 14, P56	10.8	78
579	A comparison of the effect on renal function of flucloxacillin and vancomycin antibiotic prophylaxis in cardiac surgery patients. <i>Critical Care</i> , 2010 , 14, P57	10.8	78
578	Microdialysis study of meropenem cerebral distribution in patients with acute brain injury. <i>Critical Care</i> , 2010 , 14, P58	10.8	78
577	Effects of two regimens of antibiotic prophylaxis on colonizing flora of the respiratory tract in patients with cardiovascular surgery. <i>Critical Care</i> , 2010 , 14, P59	10.8	78
576	Prevalence of infections among patients admitted to ICUs in Nordic countries and the Netherlands in comparison with Mediterranean countries: a report from the EPIC II study. <i>Critical Care</i> , 2010 , 14, P60	10.8	78
575	Risk factors for the development of carbapenem-resistant <i>Klebsiella pneumoniae</i> infections in critically ill patients. <i>Critical Care</i> , 2010 , 14, P61	10.8	78
574	Outcome of unimicrobial versus polymicrobial sepsis. <i>Critical Care</i> , 2010 , 14, P62	10.8	2

573	De-escalation practice pattern in an Indian intensive care unit. <i>Critical Care</i> , 2010 , 14, P63	10.8	1
572	Incidence, identification of multidrug-resistant pathogens and impact on the outcome after cardiac surgery. <i>Critical Care</i> , 2010 , 14, P64	10.8	2
571	Surveillance of ICU-acquired infections in Belgium: 2008 reference data. <i>Critical Care</i> , 2010 , 14, P65	10.8	78
570	Risk of acquiring drug-resistant (MDR) Gram-negative isolates with previous exposure to antibiotic. <i>Critical Care</i> , 2010 , 14, P66	10.8	78
569	Ozone fumigation successfully controlled and eradicated multidrug-resistant <i>Acinetobacter baumannii</i> from an intensive care unit. <i>Critical Care</i> , 2010 , 14, P67	10.8	0
568	Viral infections in the ICU: should we search for them?. <i>Critical Care</i> , 2010 , 14, P68	10.8	78
567	Oseltamivir dosing with haemofiltration. <i>Critical Care</i> , 2010 , 14, P69	10.8	6
566	Efficacy and utility of a protocol for pre-emptive antimycotic therapy. <i>Critical Care</i> , 2010 , 14, P70	10.8	2
565	Amphotericin B used as a continuous infusion is safer. <i>Critical Care</i> , 2010 , 14, P71	10.8	78
564	Anidulafungin compared with fluconazole therapy in critically ill patients. <i>Critical Care</i> , 2010 , 14, P72	10.8	5
563	Increase in systolic blood pressure and improvement in laboratory parameters following polymyxin B-immobilized fiber treatment in septic shock. <i>Critical Care</i> , 2010 , 14, P73	10.8	78
562	Selection of acute blood purification therapy according to lipid mediator adsorption and blood purification in patients with septic shock. <i>Critical Care</i> , 2010 , 14, P74	10.8	78
561	Prognostic assessment in community-acquired pneumonia by pneumonia severity scores and biomarkers. <i>Critical Care</i> , 2010 , 14, P75	10.8	78
560	Current progress in pneumonia research. <i>Critical Care</i> , 2010 , 14, P76	10.8	78
559	Bacterial colonization and infections of the lower respiratory tract in an interdisciplinary ICU. <i>Critical Care</i> , 2010 , 14, P77	10.8	78
558	Rehydration does not affect pulmonary immune responses to influenza or susceptibility to secondary bacterial pneumonia. <i>Critical Care</i> , 2010 , 14, P78	10.8	78
557	Bacteremic nosocomial pneumonia cases from the ATTAIN studies. <i>Critical Care</i> , 2010 , 14, P79	10.8	78
556	Late ventilator-associated pneumonia: analysis of baseline characteristics and clinical outcomes in the ATTAIN studies. <i>Critical Care</i> , 2010 , 14, P80	10.8	1

555	On the diagnosis of acute respiratory distress syndrome in nosocomial pneumonias. <i>Critical Care</i> , 2010 , 14, P81	10.8	78
554	Genetic variability at the surfactant proteins A and D in susceptibility and severity of pneumonia. <i>Critical Care</i> , 2010 , 14, P82	10.8	1
553	Successful strategy to reduce ventilator-associated pneumonia. <i>Critical Care</i> , 2010 , 14, P83	10.8	78
552	Ventilator-associated pneumonia rate and ventilator bundle compliance in a district general hospital. <i>Critical Care</i> , 2010 , 14, P84	10.8	78
551	Oral chlorhexidine to prevent nosocomial pneumonia in critically ill patients: a systematic review and meta-analysis. <i>Critical Care</i> , 2010 , 14, P85	10.8	78
550	Impact of routine multiple site decontamination in intubated patients on ICU-acquired infections. <i>Critical Care</i> , 2010 , 14, P86	10.8	78
549	Time course of RT-PCR positivity in H1N1-induced ARDS. <i>Critical Care</i> , 2010 , 14, P88	10.8	78
548	Novel swine influenza A: most frequent ultrasonographic lung patterns. <i>Critical Care</i> , 2010 , 14, P89	10.8	78
547	Outcome and prognostic factors in patients with HIV-negative pneumocystis pneumonia requiring mechanical ventilation. <i>Critical Care</i> , 2010 , 14, P90	10.8	78
546	Acute respiratory failure from pandemic influenza A (H1N1) in an intensive care unit in southern Brazil. <i>Critical Care</i> , 2010 , 14, P91	10.8	4
545	Extracorporeal membrane oxygenation for influenza A (H1N1): experience in a regional referral center. <i>Critical Care</i> , 2010 , 14, P92	10.8	78
544	Budd-Chiari syndrome complicated by abdominal compartment syndrome: evidence of central hypovolaemia?. <i>Critical Care</i> , 2010 , 14, P93	10.8	78
543	Adenosine signalling has a protective role in intestinal ischemia and reperfusion injury. <i>Critical Care</i> , 2010 , 14, P94	10.8	78
542	Threshold levels of extreme body surface area that may cause a misrepresentation of cardiac performance. <i>Critical Care</i> , 2010 , 14, P96	10.8	78
541	Cardiac output measurement is feasible in the presence of left-to-right shunt with ultrasound dilution method: a validation study in lambs. <i>Critical Care</i> , 2010 , 14, P97	10.8	2
540	The PAPIKAS trial: a comparative clinical trial of pulmonary catheter versus the PiCCO device during therapy of patients with acute heart failure and cardiogenic shock. <i>Critical Care</i> , 2010 , 14, P98	10.8	1
539	Bioreactance versus PiCCOTD/PC in critically ill septic shock patients. <i>Critical Care</i> , 2010 , 14, P99	10.8	1
538	NICOM vs LiDCOplus during changes in cardiac output in critically ill patients. <i>Critical Care</i> , 2010 , 14, P100	10.8	78

537	PulseCO consistency: variation in calibration factor over 24 and 48 hours. <i>Critical Care</i> , 2010 , 14, P102	10.8	5
536	Comparison of cardiac output values; pulse contour methods against thermodilution technique. <i>Critical Care</i> , 2010 , 14, P105	10.8	78
535	Continuous cardiac output measurement: effect of time. <i>Critical Care</i> , 2010 , 14, P106	10.8	78
534	A comparison in cardiac output data: a random effects model for repeated measures. <i>Critical Care</i> , 2010 , 14, P107	10.8	78
533	In vivo validation of a new transpulmonary thermodilution method to assess global end-diastolic volume and extravascular lung water. <i>Critical Care</i> , 2010 , 14, P108	10.8	78
532	Cardiac index derived from arterial pressure waveform: uncalibrated analysis vs once-only calibrated analysis. <i>Critical Care</i> , 2010 , 14, P109	10.8	78
531	Cardiac output monitoring during abdominal aortic cross clamping: a comparison between Vigileo/FloTrac system and transoesophageal Doppler. <i>Critical Care</i> , 2010 , 14, P110	10.8	78
530	Stroke volume index assessment using two minimal invasive devices during hemodynamic postoperative optimization. <i>Critical Care</i> , 2010 , 14, P111	10.8	78
529	Non-invasive cardiac output monitoring in children: clinical validation. <i>Critical Care</i> , 2010 , 14, P112	10.8	2
528	Global end-diastolic volume and its correlation to cardiac index inside and outside normal values. <i>Critical Care</i> , 2010 , 14, P114	10.8	78
527	Relationship of stroke volume variation, pulse pressure variation and global end-diastolic volume in patients undergoing brain surgery. <i>Critical Care</i> , 2010 , 14, P115	10.8	78
526	The value of pulse pressure variation to predict volume response in patients ventilated with low VT. <i>Critical Care</i> , 2010 , 14, P116	10.8	1
525	Pre-ejection period is a reliable parameter to estimate cardiac preload in a hemorrhagic shock. <i>Critical Care</i> , 2010 , 14, P117	10.8	78
524	Predictive value of pulse pressure variation for fluid responsiveness after maneuver to change tidal volume in patients with protective ventilatory strategy. <i>Critical Care</i> , 2010 , 14, P118	10.8	1
523	Does stroke volume increase after a fluid challenge? A study on the management of patients undergoing major head and neck free flap surgery: preliminary data. <i>Critical Care</i> , 2010 , 14, P119	10.8	1
522	Optimizing stroke volume and oxygen delivery in elective abdominal aortic surgery. <i>Critical Care</i> , 2010 , 14, P120	10.8	78
521	Oxygen saturation determined from the mouth is an early indicator of central hypovolemia in humans. <i>Critical Care</i> , 2010 , 14, P121	10.8	78
520	Plethysmographic indices as predictors of fluid responsiveness: a meta-analysis. <i>Critical Care</i> , 2010 , 14, P122	10.8	1

519	Maintenance of cardiac index within normal range is associated with mortality reduction in patients undergoing major urological surgery. <i>Critical Care</i> , 2010 , 14, P123	10.8	2
518	Effect of a preload challenge on peripheral perfusion in critically ill patients. <i>Critical Care</i> , 2010 , 14, P124	10.8	78
517	Distensibility index of inferior vena cava diameter in ventilated septic and trauma patients with shock. <i>Critical Care</i> , 2010 , 14, P125	10.8	1
516	Aortic dP/dtmax accurately reflects left ventricular contractility when effective preload independence is achieved. <i>Critical Care</i> , 2010 , 14, P126	10.8	78
515	Cardiac cycle efficiency: a new index for cardiac work estimation tested during aortic valve plasty. <i>Critical Care</i> , 2010 , 14, P127	10.8	1
514	Nonocclusive mesenteric ischemia following cardiothoracic surgery. <i>Critical Care</i> , 2010 , 14, P128	10.8	78
513	Etiology, comorbidity and prognosis of hospitalized patients with congestive heart failure. <i>Critical Care</i> , 2010 , 14, P129	10.8	1
512	Factors associated with intensive care morbidity following surgical repair of tetralogy of fallot. <i>Critical Care</i> , 2010 , 14, P130	10.8	78
511	Sildenafil attenuates pulmonary vascular remodeling and upregulates Kv1.5 mRNA expression in pulmonary hypertension secondary to left-to-right shunt in rats. <i>Critical Care</i> , 2010 , 14, P131	10.8	78
510	Validation of echocardiographic indices of right ventricular afterload: an experimental, open pericardium pig model. <i>Critical Care</i> , 2010 , 14, P132	10.8	78
509	Activation of apoptotic pathways in experimental acute afterload-induced right ventricular failure. <i>Critical Care</i> , 2010 , 14, P133	10.8	78
508	High incidence of severely prolonged QT interval after cardiac surgery. <i>Critical Care</i> , 2010 , 14, P134	10.8	2
507	Endothelial protein C receptor expression after cardiopulmonary bypass in adult cardiac surgical patients. <i>Critical Care</i> , 2010 , 14, P135	10.8	78
506	Urapidyl for hypertension control in severe pre-eclampsia: a comparative study with nicardipine. <i>Critical Care</i> , 2010 , 14, P136	10.8	2
505	Comparison of the effects of furosemide, captopril and lorazepam on noncomplicated hypertensive patients in the emergency department. <i>Critical Care</i> , 2010 , 14, P137	10.8	78
504	Incidence, risk factors and outcome of hypertensive crises in critically ill patients: a retrospective survey. <i>Critical Care</i> , 2010 , 14, P138	10.8	78
503	Comparative study to evaluate blood pressure levels measured by an invasive method versus a non-invasive method in patients in an intensive care unit. <i>Critical Care</i> , 2010 , 14, P139	10.8	78
502	Chronic atrial fibrillation in intensive care unit treated with electrical cardioversion: is it appropriate?. <i>Critical Care</i> , 2010 , 14, P140	10.8	78

501	Applanation tonometry application in ICU patients with acute cardiogenic edema. <i>Critical Care</i> , 2010 , 14, P141	10.8	78
500	A better prognosis for septic patients with left ventricular dysfunction?. <i>Critical Care</i> , 2010 , 14, P142	10.8	78
499	Microvascular effects of pulsatile versus nonpulsatile perfusion during cardiopulmonary bypass. <i>Critical Care</i> , 2010 , 14, P143	10.8	78
498	Microcirculation and intravascular coagulopathy in patients with severe sepsis and septic shock. <i>Critical Care</i> , 2010 , 14, P144	10.8	78
497	Prevalence and prognostic significance of cardiac abnormalities in the ICU: an echocardiographic study. <i>Critical Care</i> , 2010 , 14, P145	10.8	78
496	BNP and NT-proBNP in patients with acute myocardial infarction complicated by cardiogenic shock: results from the IABP Shock trial. <i>Critical Care</i> , 2010 , 14, P146	10.8	6
495	Understanding vital-sign abnormalities in critical care patients. <i>Critical Care</i> , 2010 , 14, P147	10.8	78
494	Tissue microdialysis in critically ill septic patients: associations with sepsis severity and mortality. <i>Critical Care</i> , 2010 , 14, P148	10.8	1
493	Delayed increase in inflammation and thenar VO ₂ in marathon runners. <i>Critical Care</i> , 2010 , 14, P149	10.8	78
492	Value of peripheral tissue oxygen tension in animal experimental models of haemorrhagic and septic shock. <i>Critical Care</i> , 2010 , 14, P150	10.8	78
491	Incidence of a low tissue oxygen saturation in a mixed population of critically ill patients. <i>Critical Care</i> , 2010 , 14, P151	10.8	2
490	Use of near-infrared spectroscopy during a vascular occlusion test to assess the microcirculatory response during fluid responsiveness. <i>Critical Care</i> , 2010 , 14, P152	10.8	1
489	Influence of mechanical ventilation on intestinal microcirculation and inflammation in septic rats. <i>Critical Care</i> , 2010 , 14, P153	10.8	78
488	ScvO ₂ and Pcv-aCO ₂ as complementary tools for goal-directed therapy during high-risk surgery. <i>Critical Care</i> , 2010 , 14, P154	10.8	3
487	Prognostic value of central venous-arterial pCO ₂ difference in severe sepsis and septic shock patients. <i>Critical Care</i> , 2010 , 14, P155	10.8	78
486	Veno-arterial carbon dioxide gradient at the early stage of septic shock. <i>Critical Care</i> , 2010 , 14, P156	10.8	2
485	Venous-arterial PCO ₂ difference as an early predictor of organ dysfunction in the ICU. <i>Critical Care</i> , 2010 , 14, P157	10.8	78
484	Lactate and ScvO ₂ combinations do not predict mortality in ICU patients. <i>Critical Care</i> , 2010 , 14, P158	10.8	78

483	Early changes in lactate and central venous saturation combinations do not predict subsequent changes in organ performance. <i>Critical Care</i> , 2010 , 14, P159	10.8	78
482	Regional lactate and oxygen saturation gradients: preliminary experimental data. <i>Critical Care</i> , 2010 , 14, P160	10.8	78
481	Metformin increases platelet lactate production by inhibiting mitochondrial function. <i>Critical Care</i> , 2010 , 14, P162	10.8	1
480	Artifactual lactic acidosis from point of care analysis: mind the gap. <i>Critical Care</i> , 2010 , 14, P163	10.8	78
479	Physicochemical acid-base parameters and mortality in 1,070 ICU patients: a retrospective cohort study. <i>Critical Care</i> , 2010 , 14, P165	10.8	78
478	Accuracy of predicting arterial blood gases following mechanical ventilator adjustment. <i>Critical Care</i> , 2010 , 14, P166	10.8	78
477	Use of capnography and the availability of airway equipment on UK intensive care units. <i>Critical Care</i> , 2010 , 14, P167	10.8	78
476	Combined oximetry and transcutaneous capnography is a reliable non-invasive monitoring in patients after cardiac surgery. <i>Critical Care</i> , 2010 , 14, P168	10.8	1
475	Computerized lung sound monitoring to assess effectiveness of physiotherapy and secretion removal: a feasibility study. <i>Critical Care</i> , 2010 , 14, P169	10.8	3
474	Analysis of sound waves recorded with the VRI pulmonary acoustic monitoring system in critically ill mechanically ventilated patients. <i>Critical Care</i> , 2010 , 14, P170	10.8	78
473	Lung tissue properties obtained by optical alveolar elastometry. <i>Critical Care</i> , 2010 , 14, P171	10.8	1
472	Combined monitoring of functional residual capacity and compliance may avoid hyperinflation and cardiac depression in ARDS. <i>Critical Care</i> , 2010 , 14, P172	10.8	78
471	Electrical impedance tomography at two thoracic levels provides detailed information about ventilation distribution in the cranial to caudal direction. <i>Critical Care</i> , 2010 , 14, P173	10.8	78
470	Changes in regional ventilation determined by volume and pressure-controlled ventilation as assessed by electrical impedance tomography in an experimental model of acid-induced acute lung injury. <i>Critical Care</i> , 2010 , 14,	10.8	78
469	Changes in lung tidal distribution assessed by electrical impedance tomography in a model of hemorrhagic shock with endotoxemia and resuscitation. <i>Critical Care</i> , 2010 , 14,	10.8	78
468	Estimating the intratidal compliance profile with the new gliding-SLICE method. <i>Critical Care</i> , 2010 , 14, P176	10.8	1
467	Atelectasis and overdistension are detected by heartbeat-induced disturbances in the pressure-volume loop. <i>Critical Care</i> , 2010 , 14, P177	10.8	78
466	Transpulmonary pressure and ventilation distribution measured with EIT during a PEEP trial in porcine acute lung injury. <i>Critical Care</i> , 2010 , 14, P178	10.8	1

465	A prospective pilot study of the effect of neutrophil elastase on the pulmonary vascular permeability in patients with pneumonia. <i>Critical Care</i> , 2010 , 14, P179	10.8	78
464	Evaluating the fibroproliferative response to ventilator-induced lung injury. <i>Critical Care</i> , 2010 , 14, P180	10.8	78
463	Effects of severe hemorrhage on pulmonary mechanics in ventilated pigs with ARDS. <i>Critical Care</i> , 2010 , 14, P181	10.8	78
462	Higher versus lower positive end-expiratory pressure in acute lung injury and acute respiratory distress syndrome: systematic review and individual patient data meta-analysis. <i>Critical Care</i> , 2010 , 14, P182	10.8	1
461	Effect of different recruitment maneuvers on bacterial translocation. <i>Critical Care</i> , 2010 , 14, P183	10.8	78
460	Nonlinear recruitment model with viscoelastic component fit respiratory mechanics in ARDS. <i>Critical Care</i> , 2010 , 14, P184	10.8	2
459	Lung recruitment and PEEP response in ARDS-related H1N1 virus patients. <i>Critical Care</i> , 2010 , 14, P185	10.8	78
458	Maximal recruitment strategy minimizes tidal recruitment in severe ARDS: a CT scan study. <i>Critical Care</i> , 2010 , 14, P186	10.8	78
457	Mechanical ventilation with different alveolar pressures improves alveolar recruitment: a model study. <i>Critical Care</i> , 2010 , 14, P187	10.8	78
456	Decreasing tidal volume from 6 to 4 ml/kg: feasibility and effects on repeated opening and closing. <i>Critical Care</i> , 2010 , 14, P188	10.8	78
455	Barriers to providing lung-protective ventilation to patients with ALI/ARDS. <i>Critical Care</i> , 2010 , 14, P189	10.8	1
454	Influence of aetiology of acute respiratory distress syndrome and early differences in oxygenation on outcome. <i>Critical Care</i> , 2010 , 14, P190	10.8	78
453	Acute lung injury in TBI patients with SIRS: role of vascular endothelial damage. <i>Critical Care</i> , 2010 , 14, P191	10.8	1
452	Mechanical ventilation aggravates transfusion-related acute lung injury induced by MHC class I antibodies. <i>Critical Care</i> , 2010 , 14, P192	10.8	78
451	Adaptive support ventilation may be inappropriate for patients with ALI/ARDS after recruitment: an observational study. <i>Critical Care</i> , 2010 , 14, P193	10.8	78
450	Receptor for advanced glycation end products is associated with systemic and organ-specific severity of acute lung injury and acute respiratory distress syndrome. <i>Critical Care</i> , 2010 , 14, P194	10.8	1
449	Mild hypothermia was protective in a physiological model of ventilator-induced lung injury by reducing inflammation but not by reducing the respiratory rate. <i>Critical Care</i> , 2010 , 14, P195	10.8	78
448	Effect of induced hypothermia on respiratory parameters in mechanically ventilated patients. <i>Critical Care</i> , 2010 , 14, P196	10.8	78

447	Extracorporeal life support service in a regional referral center: the Florence experience. <i>Critical Care</i> , 2010 , 14, P197	10.8	78
446	Interhospital ground transportation of severe acute respiratory distress syndrome patients on extracorporeal membrane oxygenation: Monza's experience. <i>Critical Care</i> , 2010 , 14, P198	10.8	78
445	Effects of hypertonic saline on a pig model of acute lung injury induced by hydrochloric acid instillation. <i>Critical Care</i> , 2010 , 14,	10.8	78
444	Conventional mechanical ventilation can injury intact lungs in severe trauma patients. <i>Critical Care</i> , 2010 , 14, P200	10.8	1
443	Protective effect of methylprednisolone on ventilator-induced diaphragm dysfunction is dose dependent. <i>Critical Care</i> , 2010 , 14, P201	10.8	78
442	Alveolar recruitment with non-invasive mechanical ventilation (C-PAP) in patients with nonobstructive respiratory failure. <i>Critical Care</i> , 2010 , 14, P202	10.8	78
441	N-acetylcysteine attenuates ventilator-induced diaphragm dysfunction in rats. <i>Critical Care</i> , 2010 , 14, P203	10.8	78
440	In vitro muscle contraction force measurements on isolated and entire rat diaphragms. <i>Critical Care</i> , 2010 , 14, P204	10.8	1
439	Useful of implementation of neurally adjusted ventilatory assist in critically ill patients. <i>Critical Care</i> , 2010 , 14, P205	10.8	1
438	Twenty-four-hour neurally adjusted ventilatory assist and volume cycled ventilation have similar effects on respiratory neuromuscular function in a porcine model of fecal peritonitis. <i>Critical Care</i> , 2010 , 14, P206	10.8	78
437	Effects of conventional and closed-loop neurally adjusted ventilatory assist in intubated spinal cord injured patients. <i>Critical Care</i> , 2010 , 14, P207	10.8	78
436	A level of neurally adjusted ventilatory assist corresponds to low levels of continuous positive airway pressure and pressure support ventilation in patients preparing to be extubated from mechanical ventilation. <i>Critical Care</i> , 2010 , 14, P208	10.8	78
435	Towards partially automated ventilation: adapting decision-making according to medical preferences. <i>Critical Care</i> , 2010 , 14, P209	10.8	78
434	Postoperative pulmonary complications in adult elective surgery patients in the US: severity, outcomes and resources use. <i>Critical Care</i> , 2010 , 14, P210	10.8	10
433	Increased duration of mechanical ventilation is associated with decreased diaphragmatic force. <i>Critical Care</i> , 2010 , 14, P211	10.8	1
432	Haemodynamics and volhaemic stage in patients with respiratory disorders to thermal injury. <i>Critical Care</i> , 2010 , 14, P212	10.8	78
431	Invasive mechanical ventilation in cancer patients: prior non-invasive ventilation is a poor prognostic factor. <i>Critical Care</i> , 2010 , 14, P213	10.8	78
430	Laryngeal angioedema: do we need a new treatment algorithm in the ER?. <i>Critical Care</i> , 2010 , 14, P214	10.8	78

429	Comparison of two insertion techniques of the ProSeal laryngeal mask airway: standard versus 90° rotation in pediatric patients. <i>Critical Care</i> , 2010 , 14, P215	10.8	78
428	Endotracheal intubation in the critically ill: complication rate and use of intubation aids. <i>Critical Care</i> , 2010 , 14, P216	10.8	78
427	Intensity and adequacy of laryngoscope light in seven intensive care units. <i>Critical Care</i> , 2010 , 14, P217	10.8	78
426	Airtraq: new device in patients at increased risk for difficult tracheal intubation. <i>Critical Care</i> , 2010 , 14, P218	10.8	78
425	Evaluation of a new tracheostomy technique for morbidly obese patients on an intensive care unit. <i>Critical Care</i> , 2010 , 14, P219	10.8	2
424	Our percutaneous tracheostomy experiences using the rotation-dilatation screw method. <i>Critical Care</i> , 2010 , 14, P220	10.8	78
423	Single-step vs balloon dilatation tracheostomy: a pilot study on 20 tracheostomies. <i>Critical Care</i> , 2010 , 14, P221	10.8	78
422	Recruitment maneuver for prevention of alveolar derecruitment during percutaneous dilation tracheostomy. <i>Critical Care</i> , 2010 , 14, P222	10.8	78
421	An audit of timing of tracheostomy formation in neurosurgical patients: how long should we wait?. <i>Critical Care</i> , 2010 , 14, P223	10.8	78
420	Percutaneous dilational tracheotomy in adult burn patients. <i>Critical Care</i> , 2010 , 14, P224	10.8	2
419	Safety of percutaneous tracheostomy in trauma patients. <i>Critical Care</i> , 2010 , 14, P225	10.8	3
418	Safety of balloon dilatational tracheostomy with the Ciaglia Blue Dolphin device. <i>Critical Care</i> , 2010 , 14, P226	10.8	78
417	Influence of the cuff pressure on the swallowing reflex in tracheostomized ICU patients. <i>Critical Care</i> , 2010 , 14, P227	10.8	78
416	Microaspirations during mechanical ventilation: polyurethane versus polyvinyl cuffed endotracheal tubes. <i>Critical Care</i> , 2010 , 14, P228	10.8	78
415	Do endotracheal tubes prevent microaspiration?. <i>Critical Care</i> , 2010 , 14, P229	10.8	4
414	A cuff -leak pressure test is a simple method for assessing severe laryngeal edema in postoperative patients. <i>Critical Care</i> , 2010 , 14, P230	10.8	78
413	Code critical airway teams improves patient safety. <i>Critical Care</i> , 2010 , 14, P231	10.8	3
412	A prospective evaluation of hemodynamic indexes to predict weaning from mechanical ventilation in patients after cardiac surgery. <i>Critical Care</i> , 2010 , 14, P232	10.8	78

411	Rapid shallow breathing index predicts readiness to wean?. <i>Critical Care</i> , 2010 , 14, P233	10.8	78
410	Increasing pressure support during vibrocompression is useful during respiratory therapy?. <i>Critical Care</i> , 2010 , 14, P234	10.8	78
409	Predicting success in weaning from mechanical ventilation in patients with and without cardiac disease. <i>Critical Care</i> , 2010 , 14, P235	10.8	78
408	Implementation of a standardised weaning protocol in patients with prolonged mechanical ventilation in a post-acute care ICU. <i>Critical Care</i> , 2010 , 14, P236	10.8	0
407	Comparison of pressure control and pressure support modes for non-invasive mechanical ventilation in acute hypercapnic respiratory failure. <i>Critical Care</i> , 2010 , 14, P237	10.8	1
406	Non-invasive CPAP with face mask: comparison between different devices. <i>Critical Care</i> , 2010 , 14, P238	10.8	1
405	Effects of non-invasive mechanical ventilation on hemodynamic and gas exchanges parameters after cardiac surgery in patients with $300 > PaO_2 / FiO_2 > 150$. <i>Critical Care</i> , 2010 , 14, P239	10.8	78
404	Non-invasive positive pressure ventilation in burns. <i>Critical Care</i> , 2010 , 14, P240	10.8	78
403	Non-invasive mechanical ventilation in the weaning process: continuous hemodynamic monitoring. <i>Critical Care</i> , 2010 , 14, P241	10.8	78
402	Influence of severe obesity on non-invasive ventilation strategies and responses in patients with acute hypercapnic respiratory failure in ICU. <i>Critical Care</i> , 2010 , 14, P242	10.8	1
401	A simple predictive scoring system for prolonged mechanical ventilation in severe sepsis and septic shock. <i>Critical Care</i> , 2010 , 14, P243	10.8	78
400	Simplified approach to ICU severity scoring with MPM and EUROSCORE. <i>Critical Care</i> , 2010 , 14, P244	10.8	78
399	Validation of a computerised system to calculate the sequential organ failure assessment score. <i>Critical Care</i> , 2010 , 14, P245	10.8	3
398	Sequential Organ Failure Assessment scores and mortality. <i>Critical Care</i> , 2010 , 14, P246	10.8	78
397	Bacteremia, APACHE II and need for renal replacement therapy are independent predictors of mortality in critically ill transplanted patients with cirrhosis. <i>Critical Care</i> , 2010 , 14, P247	10.8	78
396	External validation of six mortality prediction systems for intensive care unit patients. <i>Critical Care</i> , 2010 , 14, P248	10.8	78
395	Prognostic accuracy of severity score for prolonged ICU length of stay >72 hours in general surgical ICU: a prospective study. <i>Critical Care</i> , 2010 , 14, P249	10.8	78
394	Perioperative scores to predict mortality in surgical oncologic patients: a review of 1,362 cases. <i>Critical Care</i> , 2010 , 14,	10.8	78

393	Scoring patients with a suspected infection in the emergency department (ED): comparison of the ED-specific MEDS score with APACHE II and SOFA score. <i>Critical Care</i> , 2010 , 14, P251	10.8	1
392	Validation of the new proposed Emergency Trauma Score (EMTRAS). <i>Critical Care</i> , 2010 , 14, P252	10.8	2
391	A new severity score for community-acquired pneumonia: PARB score. <i>Critical Care</i> , 2010 , 14, P253	10.8	2
390	Modified Early Warning Score and identification of patients with severe sepsis. <i>Critical Care</i> , 2010 , 14, P254	10.8	78
389	Impact of a systematic MEWS introduction on preoperative and postoperative evaluation in urgent/emergency surgery. <i>Critical Care</i> , 2010 , 14, P255	10.8	2
388	Clinical predictors of physiological deterioration. <i>Critical Care</i> , 2010 , 14, P256	10.8	78
387	Modified Early Warning Scores: inaccurate summation or inaccurate assignment of score?. <i>Critical Care</i> , 2010 , 14, P257	10.8	8
386	Analysis of Modified Early Warning System scores and intraoperative factors on the incidence of sepsis and septic shock after elective major surgery. <i>Critical Care</i> , 2010 , 14, P258	10.8	2
385	Temporal variability of medical emergency team calls. <i>Critical Care</i> , 2010 , 14, P259	10.8	78
384	Medical emergency team use and outcomes in an Australian private hospital. <i>Critical Care</i> , 2010 , 14, P260	10.8	1
383	Continuously improving patient safety by a rapid response system. <i>Critical Care</i> , 2010 , 14, P261	10.8	78
382	Role of critical care outreach to provide the intention to treat of hematological disorders: a pilot study. <i>Critical Care</i> , 2010 , 14, P262	10.8	78
381	Score to door time: a benchmarking tool for rapid response systems. <i>Critical Care</i> , 2010 , 14, P263	10.8	1
380	A comparison of injected lidocaine 2% and no analgesia for peripheral intravenous cannula insertion in the emergency department. <i>Critical Care</i> , 2010 , 14, P264	10.8	78
379	High-sensitive troponin T measurement for patients with acute chest pain: improvement of diagnostics?. <i>Critical Care</i> , 2010 , 14, P265	10.8	78
378	Correlation of lung comet score with decreased gas exchange in patients with acute pulmonary edema: a pilot study. <i>Critical Care</i> , 2010 , 14, P267	10.8	78
377	Magnitude of first medical contact in the long-term prognosis of patients treated with STEMI. <i>Critical Care</i> , 2010 , 14, P268	10.8	78
376	Ultrasound-guided hematoma block and fracture reduction: a new way to go forward. <i>Critical Care</i> , 2010 , 14, P269	10.8	2

375	Pulmonary edema in the prehospital setting: the importance of early treatment. <i>Critical Care</i> , 2010 , 14, P270	10.8	78
374	Efficacy of bedside ultrasonography in early diagnosis of patients admitted to the emergency department with dyspnea. <i>Critical Care</i> , 2010 , 14, P271	10.8	78
373	Auscultation in flight: comparison of amplified and traditional stethoscopes. <i>Critical Care</i> , 2010 , 14, P272	10.8	1
372	Performance of ventilators at simulated altitude: study of fraction of inspired oxygen. <i>Critical Care</i> , 2010 , 14, P273	10.8	78
371	Aeromedical evacuation of the French Army: survey of the past 10 years. <i>Critical Care</i> , 2010 , 14, P274	10.8	1
370	Comparison of two disaster drills' management performed by trained and not-trained students: key times evaluation. <i>Critical Care</i> , 2010 , 14, P275	10.8	78
369	Acute severe asthma: performance of ventilator at simulated altitude. <i>Critical Care</i> , 2010 , 14, P276	10.8	78
368	Validation of a new French-language triage algorithm: the ELISA scale. <i>Critical Care</i> , 2010 , 14, P277	10.8	1
367	Examining the suitability of medical admissions to the emergency short stay ward of a large UK hospital. <i>Critical Care</i> , 2010 , 14, P278	10.8	3
366	A study on reliability of a new triage emergency method among nursing students. <i>Critical Care</i> , 2010 , 14, P279	10.8	78
365	Aggression and violence against staff of ambulance services. <i>Critical Care</i> , 2010 , 14, P280	10.8	78
364	Analysis of the disease spectrum of treated patients after Wenchuan earthquake. <i>Critical Care</i> , 2010 , 14, P281	10.8	78
363	Large-scale implementation of a critical care surge capacity management program. <i>Critical Care</i> , 2010 , 14, P282	10.8	78
362	Standard X-rays for the victims of severe trauma: time for a change. <i>Critical Care</i> , 2010 , 14, P283	10.8	78
361	Focused assessment with sonography for trauma in the prehospital setting. <i>Critical Care</i> , 2010 , 14, P284	10.8	0
360	Clinicopathogenetic aspects of systemic inflammatory response development in polytrauma. <i>Critical Care</i> , 2010 , 14, P285	10.8	78
359	Diagnostic and prognostic significance of serum apoptosis markers in the patients with severe trauma injury. <i>Critical Care</i> , 2010 , 14, P286	10.8	78
358	Predictors of mechanical ventilation after burn injury. <i>Critical Care</i> , 2010 , 14, P287	10.8	78

357	Impact of regional imaging guidelines for head injury and unconscious polytrauma patients. <i>Critical Care</i> , 2010 , 14, P288	10.8	78
356	Blunt vascular injuries of the carotid and vertebral arteries: should we screen the asymptomatic trauma patients at high risk? Preliminary results of a prospective cohort study. <i>Critical Care</i> , 2010 , 14, P289	10.8	78
355	Effect of 8.4% sodium bicarbonate on raised intracranial pressure after traumatic brain injury. <i>Critical Care</i> , 2010 , 14, P290	10.8	78
354	Hyponatremia in pediatric patients with severe traumatic brain injury. <i>Critical Care</i> , 2010 , 14, P291	10.8	78
353	Use of transcranial Doppler in patients with severe traumatic brain injury. <i>Critical Care</i> , 2010 , 14, P292	10.8	2
352	Effect of hyperbaric oxygen on cerebral oxygen saturation in patients with sub-acute traumatic brain injury. <i>Critical Care</i> , 2010 , 14, P294	10.8	78
351	Brain biomarkers and brain tissue oxygenation: changes and correlations following severe head injury. <i>Critical Care</i> , 2010 , 14, P295	10.8	78
350	Cortex and deep grey matter have different sensitivities to hypoxia-hypotension and traumatic brain injury: a combined PtIO ₂ and microdialysis study in rats. <i>Critical Care</i> , 2010 , 14, P296	10.8	78
349	Diagnosis of carotid and vertebral artery injury in major trauma with head injury. <i>Critical Care</i> , 2010 , 14, P297	10.8	78
348	Decompressive craniectomy in severe intracranial hypertension after brain injury: early or late?. <i>Critical Care</i> , 2010 , 14, P298	10.8	78
347	Optic nerve ultrasound for detecting intracranial hypertension. <i>Critical Care</i> , 2010 , 14, P300	10.8	1
346	Measurement of neuronal specific enolase as a predictor of outcome in patients suffering from severe traumatic brain injury. <i>Critical Care</i> , 2010 , 14, P301	10.8	78
345	Hyperphosphorylated neurofilament NF-H, S100 β and NSE as indicators of severity of brain injury in children. <i>Critical Care</i> , 2010 , 14, P302	10.8	78
344	A simple hospital discharge score predicts Glasgow Outcome Scale at 12 months in patients with traumatic brain injury. <i>Critical Care</i> , 2010 , 14, P303	10.8	78
343	Outcome of aggressive treatment for blunt and penetrating traumatic cardiac arrest. <i>Critical Care</i> , 2010 , 14, P304	10.8	78
342	Using flow-rate limiting bag-valve-mask device without training. <i>Critical Care</i> , 2010 , 14, P305	10.8	78
341	Out-of-hospital cardiac arrest and resource utilisation in the ICU. <i>Critical Care</i> , 2010 , 14, P306	10.8	78
340	Comparison of external cardiac compression on the floor and in a dental chair. <i>Critical Care</i> , 2010 , 14, P307	10.8	78

339	Time factors rather than type and/or nature of CPR modify the prognosis of out-of-hospital cardiac arrests witnessed by bystanders. <i>Critical Care</i> , 2010 , 14, P309	10.8	78
338	Use of the laryngeal tube by paramedics during prehospital resuscitation: an observational study. <i>Critical Care</i> , 2010 , 14, P310	10.8	78
337	The i-gel supraglottic airway as an alternative in the preclinical emergency medical aid? A comparative study on a manikin. <i>Critical Care</i> , 2010 , 14, P311	10.8	78
336	Retention of skills during emergency airway management training: a 12-month follow-up. <i>Critical Care</i> , 2010 , 14, P312	10.8	2
335	Factors correlated with the advanced life support simulation performance of senior medical students. <i>Critical Care</i> , 2010 , 14, P313	10.8	78
334	Aging augments unwillingness to the initial three links in chains of survival. <i>Critical Care</i> , 2010 , 14, P314	10.8	78
333	Prediction of the success of cardiac resuscitation: a pattern classification approach based on ECG spectral and temporal features. <i>Critical Care</i> , 2010 , 14, P315	10.8	78
332	Dispatcher-assisted telephone cardiopulmonary resuscitation using a French compression-only protocol: performance of volunteers with or without prior life support training. <i>Critical Care</i> , 2010 , 14, P316	10.8	78
331	CPR team members' attitudes towards family presence during cardiopulmonary resuscitation. <i>Critical Care</i> , 2010 , 14, P317	10.8	78
330	Prehospital therapeutic hypothermia induced with cold infusions improves haemodynamic stability in nonshockable cardiac arrest patients. <i>Critical Care</i> , 2010 , 14, P318	10.8	1
329	Trans-nasal cooling during CPR: a single-center experience. <i>Critical Care</i> , 2010 , 14, P319	10.8	1
328	Intra-arrest cooling using a novel intra-nasal cooling method for immediate induction of therapeutic hypothermia in Germany. <i>Critical Care</i> , 2010 , 14, P320	10.8	78
327	Therapeutic hypothermia after cardiac arrest: implementation in UK intensive care units. <i>Critical Care</i> , 2010 , 14, P321	10.8	78
326	Hospital-wide therapeutic hypothermia protocol significantly reduces the time to induce hypothermia but does not improve outcomes. <i>Critical Care</i> , 2010 , 14, P322	10.8	78
325	Induced hypothermia after cardiac arrest for 12 hours: single-centre experience. <i>Critical Care</i> , 2010 , 14, P323	10.8	78
324	Survival following adult cardiac arrest in intensive care units: a 5-year retrospective analysis. <i>Critical Care</i> , 2010 , 14, P324	10.8	78
323	Effects of bed height and time on the quality of chest compressions during cardiopulmonary resuscitation. <i>Critical Care</i> , 2010 , 14, P325	10.8	78
322	New results about early therapeutic hypothermia in sudden cardiac death and following return of spontaneous circulation. <i>Critical Care</i> , 2010 , 14, P326	10.8	78

321	Infection rates among cooled patients post cardiac arrest. <i>Critical Care</i> , 2010 , 14, P327	10.8	78
320	Strategies to implement a therapeutic hypothermia program. <i>Critical Care</i> , 2010 , 14, P328	10.8	78
319	A comparison of core and tympanic temperature measurement in the critically ill. <i>Critical Care</i> , 2010 , 14, P329	10.8	1
318	Can we predict neurological prognoses with computed tomography just after resuscitation?. <i>Critical Care</i> , 2010 , 14, P330	10.8	78
317	Unintended consequences of technology improvement. <i>Critical Care</i> , 2010 , 14, P331	10.8	78
316	Bispectral Index Scale alterations in brain death patients. <i>Critical Care</i> , 2010 , 14, P332	10.8	78
315	Prognosis in patients presenting with nontraumatic coma. <i>Critical Care</i> , 2010 , 14, P333	10.8	2
314	Dynamic changes in cerebral oxygenation by two methods during cardiac surgery and postoperative cognitive decline. <i>Critical Care</i> , 2010 , 14, P334	10.8	1
313	Acetazolamide-induced cerebrovascular reactivity is impaired in sepsis-associated encephalopathy. <i>Critical Care</i> , 2010 , 14, P335	10.8	1
312	Epidemiological analysis of patients with cerebral aneurysms submitted to embolization at S3 Hospital. <i>Critical Care</i> , 2010 , 14, P336	10.8	78
311	Frequency/prevalence analysis of risk factors on aneurysmal subarachnoid hemorrhage. <i>Critical Care</i> , 2010 , 14, P337	10.8	1
310	Monitoring regional brain oxygen saturation in patients with clipped aneurysms of the anterior circle of Willis. <i>Critical Care</i> , 2010 , 14, P338	10.8	78
309	Early prognosis value of the first 24 hours mean plasma S100 β protein level after subarachnoid aneurysmal hemorrhage. <i>Critical Care</i> , 2010 , 14, P339	10.8	78
308	Altered aortic stiffness associated with stress cardiomyopathy in patients with aneurysmal subarachnoid haemorrhage. <i>Critical Care</i> , 2010 , 14, P340	10.8	78
307	Effect of packed red blood cell transfusion on cerebral oxygenation and metabolism after subarachnoid hemorrhage. <i>Critical Care</i> , 2010 , 14, P341	10.8	1
306	Comparison of surgical or endovascular treated aneurysmal SAH patients with a special emphasis on cerebral vasospasm. <i>Critical Care</i> , 2010 , 14, P342	10.8	78
305	Prediction of cerebral vasospasm in patients with aneurysmal subarachnoid hemorrhage using jugular bulb oximetry monitoring: preliminary results. <i>Critical Care</i> , 2010 , 14, P343	10.8	0
304	Magnesium use on prophylaxis of vasospasm morbidity and mortality rate in subarachnoid hemorrhage. <i>Critical Care</i> , 2010 , 14, P344	10.8	78

303	Effects of simvastatin in prevention of vasospasm in nontraumatic subarachnoid hemorrhage: preliminary data. <i>Critical Care</i> , 2010 , 14, P345	10.8	78
302	Addition of new criteria to Sequential Organ Failure Assessment for patients with subarachnoid hemorrhage. <i>Critical Care</i> , 2010 , 14, P346	10.8	78
301	Cardiac output augmentation with fluid resuscitation improves brain tissue oxygenation after severe brain injury. <i>Critical Care</i> , 2010 , 14, P347	10.8	78
300	Multicentre experience of using ICM+ for investigations of cerebrovascular dynamics with near-infrared spectroscopy. <i>Critical Care</i> , 2010 , 14, P348	10.8	2
299	Cerebral oxygen saturation and cerebral blood flow are relatively stable during single-lung ventilation, if normocapnia is maintained. <i>Critical Care</i> , 2010 , 14, P349	10.8	78
298	Factors affecting the transient hyperaemic response test of the cerebral autoregulation. <i>Critical Care</i> , 2010 , 14, P350	10.8	78
297	Infection associated with the external ventricular derivation catheter: preliminary data. <i>Critical Care</i> , 2010 , 14, P351	10.8	78
296	fMRI default mode connectivity in vegetative state and locked-in syndrome patients. <i>Critical Care</i> , 2010 , 14, P352	10.8	78
295	Incidence of cerebral salt wasting syndrome: comparison in subarachnoid haemorrhagic compared with head trauma. <i>Critical Care</i> , 2010 , 14, P353	10.8	78
294	Sodium dysbalances in neurointensive care: results of a 5-year prospective study. <i>Critical Care</i> , 2010 , 14, P354	10.8	78
293	Impact of hypernatremia on patients with severe traumatic brain injury. <i>Critical Care</i> , 2010 , 14, P355	10.8	5
292	Prevalence and clinical setting of SIADH in neurosurgery: comparison between subarachnoid haemorrhage and head trauma. <i>Critical Care</i> , 2010 , 14, P356	10.8	78
291	Risk factors for venous thromboembolism in Japanese patients with severe blunt trauma. <i>Critical Care</i> , 2010 , 14, P357	10.8	78
290	Safety of heparin prophylaxis for venous thromboembolism in patients with neurotrauma. <i>Critical Care</i> , 2010 , 14, P358	10.8	78
289	Prevalence and incidence of proximal deep vein thrombosis in critically ill patients. <i>Critical Care</i> , 2010 , 14, P359	10.8	78
288	Trauma patients with prophylactic IVC filter and chemical prophylaxis: a descriptive study. <i>Critical Care</i> , 2010 , 14, P360	10.8	78
287	Clinical signs and risk factors of acute deep vein thrombosis of lower extremities in patients after abdominal surgery for cancer: efficiency and safety of different types of anticoagulant therapy. <i>Critical Care</i> , 2010 , 14, P361	10.8	78
286	Multidrug sedation as a risk factor for deep vein thrombosis in the intensive care unit. <i>Critical Care</i> , 2010 , 14, P362	10.8	78

285	New method diagnostics for coagulation disorders after surgery. <i>Critical Care</i> , 2010 , 14, P363	10.8	78
284	Thrombocytopenia in surgical ICU patients. <i>Critical Care</i> , 2010 , 14, P364	10.8	78
283	Platelet count as a prognostic marker in intensive care. <i>Critical Care</i> , 2010 , 14, P365	10.8	1
282	66 Evaluation of the particle immunofiltration anti-platelet factor 4 rapid assay in MICU patients with thrombocytopenia. <i>Critical Care</i> , 2010 , 14, P366	10.8	78
281	Heparin-induced thrombocytopenia incidence in the ICU: preliminary results. <i>Critical Care</i> , 2010 , 14, P367	10.8	78
280	Treatment of the coagulation deficit during extracorporeal circulation in a porcine model by a prothrombin complex concentrate. <i>Critical Care</i> , 2010 , 14, P368	10.8	78
279	Fibrinogen in dilutional coagulopathy: a dose study in pigs. <i>Critical Care</i> , 2010 , 14, P369	10.8	78
278	Tranexamic acid reduces blood loss in postpartum haemorrhage. <i>Critical Care</i> , 2010 , 14, P370	10.8	6
277	Regional variation of the off-licence use of rFVIIa for patients with uncontrolled haemorrhage in England. <i>Critical Care</i> , 2010 , 14, P371	10.8	78
276	Early use of factor VIIa in cardiac surgery is associated with lower rates of blood transfusions without impact on outcome. <i>Critical Care</i> , 2010 , 14,	10.8	78
275	Low-frequency hemoviscoelastography in monitoring coagulation disorders after abdominal surgery for cancer. <i>Critical Care</i> , 2010 , 14, P373	10.8	78
274	Reliability of the HemoCue [®] hemoglobinometer in critically ill patients. <i>Critical Care</i> , 2010 , 14, P374	10.8	78
273	Effect of transfusion, leukocyte-depleted blood product on onset of new septic shock and mortality in septic shock. <i>Critical Care</i> , 2010 , 14, P375	10.8	78
272	Transfusion Requirements After Cardiac Surgery (TRACS) study: preliminary data of a randomized controlled clinical trial. <i>Critical Care</i> , 2010 , 14,	10.8	3
271	Analysis of red cell transfusion practices in patients without active haemorrhage over a 12-month period in a UK intensive care unit. <i>Critical Care</i> , 2010 , 14, P377	10.8	78
270	Effects of red blood cell transfusion in central venous saturation and lactate levels in patients with septic shock. <i>Critical Care</i> , 2010 , 14, P378	10.8	78
269	Massive transfusion in paediatric trauma: a single-centre experience. <i>Critical Care</i> , 2010 , 14, P379	10.8	78
268	Cytokine levels evaluation during acute isovolemic anemia. <i>Critical Care</i> , 2010 , 14,	10.8	78

267	Haemodynamic alterations in community-acquired sepsis at early stage. <i>Critical Care</i> , 2010 , 14, P381	10.8	3
266	Epidemiology of severe sepsis in India: an update. <i>Critical Care</i> , 2010 , 14, P382	10.8	18
265	Systemic inflammatory response syndrome is reduced by inline filtration in intensive care patients. <i>Critical Care</i> , 2010 , 14, P383	10.8	78
264	Review of central venous catheter-related infections in neurointensive care patients in a tertiary referral centre. <i>Critical Care</i> , 2010 , 14, P384	10.8	78
263	Clinical impact of diagnosis methods for catheter-related bloodstream infection in ICU patients. <i>Critical Care</i> , 2010 , 14, P385	10.8	78
262	Catheter-related bacteremia in a jugular site with tracheostomy and in a femoral site. <i>Critical Care</i> , 2010 , 14, P386	10.8	78
261	A comparative study between conventional and antiseptic impregnated central venous catheters in neurocritical patients. <i>Critical Care</i> , 2010 , 14, P387	10.8	78
260	Comparison between peripherally inserted central venous catheters and short-term central venous catheter in patients discharged from the ICU. <i>Critical Care</i> , 2010 , 14, P388	10.8	1
259	Vacuum-assisted closure of laparostomy, an efficient strategy for complicated intra-abdominal infection. <i>Critical Care</i> , 2010 , 14, P389	10.8	78
258	Comparison between left ventricular tissue Doppler strain imaging and the sequential organ failure assessment score (SOFA score) in patients with severe sepsis or septic shock. <i>Critical Care</i> , 2010 , 14, P390	10.8	78
257	Systolic and diastolic function and survival in severe sepsis and septic shock: clinical, echocardiographic, biochemical correlation. <i>Critical Care</i> , 2010 , 14, P391	10.8	78
256	Frequency and clinical spectrum of myocardial dysfunction in severe sepsis and septic shock. <i>Critical Care</i> , 2010 , 14, P392	10.8	78
255	Effects of nitroglycerin on intestinal microcirculatory blood flow in patients with abdominal sepsis. <i>Critical Care</i> , 2010 , 14, P393	10.8	78
254	Diastolic dysfunction in septic shock is an independent predictor of elevated B-type natriuretic peptide and hospital mortality. <i>Critical Care</i> , 2010 , 14, P394	10.8	1
253	Cardiac-specific troponin I and cardiovascular dysfunction in post-traumatic sepsis. <i>Critical Care</i> , 2010 , 14, P395	10.8	78
252	Both systolic and diastolic functions are impaired in critically ill patients with community-acquired sepsis. <i>Critical Care</i> , 2010 , 14, P396	10.8	78
251	Selective V1a agonism reduces vascular leakage and cardiopulmonary dysfunction in methicillin-resistant <i>Staphylococcus aureus</i> sepsis. <i>Critical Care</i> , 2010 , 14, P397	10.8	2
250	Sepsis resuscitation bundle compliance in septic shock: improves survival and yet better late than never. <i>Critical Care</i> , 2010 , 14, P398	10.8	78

249	Effect of myocardial preservation of early goal-directed therapy on severe sepsis/septic shock patients in the ICU. <i>Critical Care</i> , 2010 , 14, P399	10.8	1
248	Impact of the sepsis bundle strategy on the outcomes of the patients suffering from severe sepsis and septic shock. <i>Critical Care</i> , 2010 , 14, P400	10.8	78
247	Impact of Surviving Sepsis Campaign implementation in Brazil. <i>Critical Care</i> , 2010 , 14, P401	10.8	78
246	Effect of synthetic vasopressin in septic shock using esophageal Doppler. <i>Critical Care</i> , 2010 , 14, P402	10.8	78
245	Effects of sepsis resuscitation bundle in cirrhotic patients with septic shock. <i>Critical Care</i> , 2010 , 14, P403	10.8	78
244	Outcomes and incidence of bleeding events associated with drotrecogin alfa: a single-centre experience of 440 patients. <i>Critical Care</i> , 2010 , 14, P404	10.8	78
243	Activated protein C: cost matters!. <i>Critical Care</i> , 2010 , 14, P405	10.8	78
242	Outcome of patients receiving two or more infusions of activated protein C: a single-centre experience. <i>Critical Care</i> , 2010 , 14, P406	10.8	78
241	Drotrecogin alfa (activated): a single-centre experience of 110 patients. <i>Critical Care</i> , 2010 , 14, P407	10.8	78
240	Efficacy of recombinant human soluble thrombomodulin in disseminated intravascular coagulation. <i>Critical Care</i> , 2010 , 14, P408	10.8	78
239	Clinical experience with lipopolysaccharide adsorber in cancer patients with severe sepsis and septic shock. <i>Critical Care</i> , 2010 , 14, P409	10.8	78
238	Apheresis of activated leukocytes with an immobilized polymyxin B filter. <i>Critical Care</i> , 2010 , 14, P410	10.8	0
237	Therapeutic blood purification using functionalized core/shell nanomagnets. <i>Critical Care</i> , 2010 , 14, P411	10.8	78
236	Coupled plasma filtration and adsorption in septic shock: a multicentric experience. <i>Critical Care</i> , 2010 , 14, P412	10.8	2
235	Patients admitted to the ICU for extra-abdominal disease and operated on for emergency laparotomy have significant survival. <i>Critical Care</i> , 2010 , 14, P413	10.8	78
234	Can ICU admission be predicted?. <i>Critical Care</i> , 2010 , 14, P414	10.8	78
233	Unsuitable for ICU: what happens next?. <i>Critical Care</i> , 2010 , 14, P415	10.8	78
232	Admission pattern and outcome in a UK weaning centre. <i>Critical Care</i> , 2010 , 14, P416	10.8	78

231	Referral to intensive care: who and when?. <i>Critical Care</i> , 2010 , 14, P417	10.8	78
230	Acute drug overdose in an ICU: 15 years experience. <i>Critical Care</i> , 2010 , 14, P418	10.8	3
229	Prognostic factors for very short-term mortality in severe sepsis. <i>Critical Care</i> , 2010 , 14, P419	10.8	78
228	Is sepsis mortality rate driven by differences in Brazilian healthcare systems?. <i>Critical Care</i> , 2010 , 14, P420	10.8	78
227	Prognoses indicators of HIV patients admitted to the ICU. <i>Critical Care</i> , 2010 , 14, P421	10.8	78
226	Outcomes of cancer patients admitted to intensive care units with severe acute kidney injury. <i>Critical Care</i> , 2010 , 14, P422	10.8	78
225	Is intensive care beneficial for patients with haematologic malignancies?. <i>Critical Care</i> , 2010 , 14, P423	10.8	78
224	Postoperative mortality in the Netherlands: identifying high-risk surgery. <i>Critical Care</i> , 2010 , 14, P424	10.8	0
223	Outcomes of 1,000 patients with cancer admitted to the intensive care unit: a Brazilian prospective study. <i>Critical Care</i> , 2010 , 14,	10.8	1
222	Trend in outcome of critically ill patients with co-morbidity: a 15-year study. <i>Critical Care</i> , 2010 , 14, P426	10.8	78
221	Clinical profile, maternal and fetal mortality analysis of severe gestosis admitted to the ICU. <i>Critical Care</i> , 2010 , 14, P427	10.8	78
220	Obstetric critical care in developing countries: a comparison between two intensive care units. <i>Critical Care</i> , 2010 , 14, P428	10.8	78
219	Co-morbidity, demographic and social factors as predictors of outcome from intensive care: an observational cohort study. <i>Critical Care</i> , 2010 , 14, P429	10.8	78
218	C-Reactive protein and albumin as prognostic markers in intensive care. <i>Critical Care</i> , 2010 , 14, P430	10.8	1
217	Statin therapy on admission to intensive care and outcome: an observational cohort study. <i>Critical Care</i> , 2010 , 14, P431	10.8	78
216	How did you sleep in the ICU?. <i>Critical Care</i> , 2010 , 14, P432	10.8	1
215	Factors associated with longer stay on a surgical intensive care unit. <i>Critical Care</i> , 2010 , 14, P433	10.8	78
214	Social support: a significant factor for health-related quality of life after critical illness?. <i>Critical Care</i> , 2010 , 14, P434	10.8	78

213	Quality of the evaluations of clinical studies in adult critical care and emergency medicine on a specialized Internet site. <i>Critical Care</i> , 2010, 14, P435	10.8	0
212	Functional status and quality of life in patients suffering severe cranioencephalic trauma at the time of discharge from the intensive care unit and 1 year after. <i>Critical Care</i> , 2010, 14, P436	10.8	2
211	Long-term (>6 years) quality of life after surgical intensive care admission. <i>Critical Care</i> , 2010, 14, P437	10.8	78
210	Children's self-reported health-related quality of life after intensive care treatment. <i>Critical Care</i> , 2010, 14, P438	10.8	4
209	Activities of daily living (Barthel Index) at discharge from the intensive care unit. <i>Critical Care</i> , 2010, 14, P439	10.8	4
208	A comparison of Short Form 36 and Hospital Anxiety and Depression Scale based on patients 1 month after discharge from the intensive care unit. <i>Critical Care</i> , 2010, 14, P440	10.8	78
207	Post-traumatic stress disorder after surgical ICU admission. <i>Critical Care</i> , 2010, 14, P441	10.8	78
206	Health-related quality of life before intensive care. <i>Critical Care</i> , 2010, 14, P442	10.8	78
205	Nursing staff development: strategies for success in a unique critical care unit in a developing country. <i>Critical Care</i> , 2010, 14, P443	10.8	78
204	Implementing the Leapfrog standard in a developing country. <i>Critical Care</i> , 2010, 14, P444	10.8	78
203	Impact of thiopental use on cerebral perfusion pressure in severe brain trauma: an example of caregivers' adaptation. <i>Critical Care</i> , 2010, 14, P445	10.8	78
202	Knowledge of ICU nurses regarding selected care areas commonly guided by protocols. <i>Critical Care</i> , 2010, 14, P446	10.8	78
201	Pharmacist clinical interventions in the ICU. <i>Critical Care</i> , 2010, 14, P447	10.8	5
200	Adverse drug events in the critical care unit. <i>Critical Care</i> , 2010, 14, P448	10.8	78
199	Evaluation of appropriateness of intensive care unit admissions: the GIVITI's StART approach. <i>Critical Care</i> , 2010, 14, P449	10.8	1
198	Noise in the critical care environment. <i>Critical Care</i> , 2010, 14, P450	10.8	1
197	Allergy documentation and transfer within critical care. <i>Critical Care</i> , 2010, 14, P451	10.8	78
196	Use of an anonymous medication incident reporting system on a critical care unit. <i>Critical Care</i> , 2010, 14, P452	10.8	1

195	Three years experience of a quality improvement programme. <i>Critical Care</i> , 2010 , 14, P453	10.8	78
194	Improving team work during bedside rounds: using daily goals and best practices. <i>Critical Care</i> , 2010 , 14, P454	10.8	78
193	Stress induced by simulation-based training in anesthesiologist residents. <i>Critical Care</i> , 2010 , 14, P455	10.8	78
192	Impact of an education program in respiratory therapy quality of care. <i>Critical Care</i> , 2010 , 14, P456	10.8	78
191	Use of the FAST HUG checklist in Mexican critically ill patients. <i>Critical Care</i> , 2010 , 14, P457	10.8	78
190	Bay rooms vs single-bed rooms in intensive care unit nosocomial infections: a case-control study. <i>Critical Care</i> , 2010 , 14, P458	10.8	2
189	Impact of quality improvement process on healthcare-associated infection in the ICU in a tertiary care hospital in India. <i>Critical Care</i> , 2010 , 14, P459	10.8	78
188	Teaching airway resuscitation to novices: a randomized controlled simulator mannequin study. <i>Critical Care</i> , 2010 , 14, P460	10.8	78
187	Impact of ultrasound guidance in central venous catheterization by inexperienced trainees: a prospective observational study. <i>Critical Care</i> , 2010 , 14, P461	10.8	78
186	Performance analysis of an ICU according to GiViTI's StART approach. <i>Critical Care</i> , 2010 , 14, P462	10.8	78
185	Simulation training at the point of care to decrease the incidence of airway complications on the intensive care unit. <i>Critical Care</i> , 2010 , 14, P463	10.8	78
184	Management of needle stick injuries from incapacitated patients. <i>Critical Care</i> , 2010 , 14, P464	10.8	78
183	Stress and fatigue among anaesthesia and intensive care doctors in Lithuania. <i>Critical Care</i> , 2010 , 14, P465	10.8	78
182	Implementation of daily goals in the ICU reduces length of ICU stay and errors of omission in patient care. <i>Critical Care</i> , 2010 , 14, P466	10.8	1
181	Use of single patient rooms to decrease acquisition of antibiotic-resistant bacteria in the ICU. <i>Critical Care</i> , 2010 , 14, P467	10.8	78
180	Time to first review of new admissions to critical care by the consultant intensivist. <i>Critical Care</i> , 2010 , 14, P468	10.8	78
179	Analysis of critical care referrals. <i>Critical Care</i> , 2010 , 14, P469	10.8	2
178	Communication with intensive care patients. <i>Critical Care</i> , 2010 , 14, P470	10.8	78

177	Quality of transfers of critically ill patients within the hospital. <i>Critical Care</i> , 2010 , 14, P471	10.8	78
176	Are bariatric patients putting a strain on intensive care beds?. <i>Critical Care</i> , 2010 , 14, P472	10.8	78
175	Five-year evolution of drug prescribing in a university adult intensive care unit. <i>Critical Care</i> , 2010 , 14, P473	10.8	78
174	A semiquantitative assessment of the effects of delayed discharge from the intensive care unit of survivors of critical illness. <i>Critical Care</i> , 2010 , 14, P474	10.8	78
173	Robotics telepresence in acute care facilities during influenza AH1N1 outbreak in Mexico. <i>Critical Care</i> , 2010 , 14, P475	10.8	78
172	Association between out of hours discharge from the ICU and subsequent readmission. <i>Critical Care</i> , 2010 , 14, P476	10.8	3
171	Sequential Organ Failure Assessment in pandemic planning. <i>Critical Care</i> , 2010 , 14, P477	10.8	2
170	Failure mode and effect analysis in interhospital transport of critically ill patients: a tool to design checklists for handovers in the mobile intensive care unit. <i>Critical Care</i> , 2010 , 14, P478	10.8	78
169	Selective neurogenic blockades and perioperative immune reactivity in patients undergoing lung resection. <i>Critical Care</i> , 2010 , 14, P479	10.8	78
168	Adverse effects of beta adrenergic blockers during spinal anesthesia with 0.5% bupivacaine. <i>Critical Care</i> , 2010 , 14, P480	10.8	2
167	Remifentanyl for postoperative analgo-sedation: effects on glycaemia and HOMA score. <i>Critical Care</i> , 2010 , 14, P481	10.8	78
166	Adding ultralow-dose oral naltrexone to morphine: a double-blind, randomized clinical trial. <i>Critical Care</i> , 2010 , 14, P482	10.8	78
165	Four-day sedation using sevoflurane: fluoride and renal function. <i>Critical Care</i> , 2010 , 14, P483	10.8	78
164	Beneficial effect of sevoflurane on endotoxin-induced pulmonary hypertension. <i>Critical Care</i> , 2010 , 14, P484	10.8	1
163	Does etomidate affect prognosis in septic shock patients treated with hydrocortisone?. <i>Critical Care</i> , 2010 , 14, P485	10.8	78
162	Use of methyl-naltrexone for the treatment of opioid-induced constipation in critical care patients. <i>Critical Care</i> , 2010 , 14, P486	10.8	78
161	Safety and efficacy of a nurse-driven sedation protocol with daily sedation interruption in the ICU. <i>Critical Care</i> , 2010 , 14, P487	10.8	78
160	Effects of propofol and midazolam on indocyanine green elimination assessed with LIMON for early sepsis patients. <i>Critical Care</i> , 2010 , 14, P488	10.8	78

159	Assessment of correlation between Bispectral Index monitorization and sedation scales used in the intensive care unit. <i>Critical Care</i> , 2010 , 14, P489	10.8	0
158	Sedation and analgesia: early results from a web-based survey of UK practice. <i>Critical Care</i> , 2010 , 14, P490	10.8	4
157	Severe alcohol withdrawal syndrome in ICU: is propofol a safe option?. <i>Critical Care</i> , 2010 , 14, P491	10.8	1
156	Agitation and pain during physiotherapy in intensive care. <i>Critical Care</i> , 2010 , 14, P492	10.8	1
155	Pain evaluation in the intensive care unit: listen to the patient, a new approach!. <i>Critical Care</i> , 2010 , 14, P493	10.8	0
154	Pharmacokinetics of orally administered melatonin in critically ill patients. <i>Critical Care</i> , 2010 , 14, P494	10.8	78
153	Manifestations and clinical course of delirium tremens in the ICU. <i>Critical Care</i> , 2010 , 14, P495	10.8	1
152	Comparison of two methods for diagnosing delirium conducted in a mixed medical ICU and the concordance between two independent nurse assessors. <i>Critical Care</i> , 2010 , 14, P496	10.8	78
151	Quetiapine in prolonged ICU delirium. <i>Critical Care</i> , 2010 , 14, P497	10.8	2
150	PREDICT, Prediction of Delirium in ICU Patients: development and validation of a prediction model. <i>Critical Care</i> , 2010 , 14, P498	10.8	2
149	Hospital anxiety depression scale in our surgical ICU. <i>Critical Care</i> , 2010 , 14, P499	10.8	1
148	Fluid therapy change from synthetic colloids to only crystalloids in patients with severe sepsis. <i>Critical Care</i> , 2010 , 14, P500	10.8	78
147	Effects of balanced and unbalanced colloid and crystalloid solutions on renal microvascular perfusion in endotoxemic rats. <i>Critical Care</i> , 2010 , 14, P501	10.8	78
146	Effects of balanced and unbalanced colloid and crystalloid solutions on renal oxygenation in a rat model of hemorrhagic shock and resuscitation. <i>Critical Care</i> , 2010 , 14, P502	10.8	78
145	Effect of rapidly infused crystalloids on acid-base status of dehydrated patients in the emergency department. <i>Critical Care</i> , 2010 , 14, P503	10.8	78
144	Impact of saline-based tetrastarch vs balanced tetrastarch and crystalloid solutions on hyperchloraemic acidosis in endotoxaemic sheep. <i>Critical Care</i> , 2010 , 14, P504	10.8	78
143	Are different hydroxyethyl starch solutions an independent risk factor for acute kidney injury in surgical critically ill patients? A retrospective analysis of 3,591 patients. <i>Critical Care</i> , 2010 , 14, P505	10.8	78
142	Role of early continuous venovenous hemodiafiltration in septic shock and multiorgan failure. <i>Critical Care</i> , 2010 , 14, P506	10.8	78

141	Effect of intravenous H2S on porcine aortic occlusion-induced systemic inflammation and kidney ischemia/reperfusion injury. <i>Critical Care</i> , 2010 , 14, P507	10.8	78
140	Survival improvement in pigs with liver failure and superimposed sepsis by a new liver support system (Hepa Wash [®]). <i>Critical Care</i> , 2010 , 14, P508	10.8	1
139	Increased glomerular filtration in the critically ill patient receiving anti-infective treatment. <i>Critical Care</i> , 2010 , 14, P509	10.8	4
138	Therapeutic plasma exchange in children with acute liver failure: assessment of laboratory parameters. <i>Critical Care</i> , 2010 , 14, P510	10.8	2
137	New markers of inflammation-induced renal injury subside when endotoxin tolerance develops in humans as measured by urine proteomics. <i>Critical Care</i> , 2010 , 14, P511	10.8	78
136	Extraction and purification of 1-hydroxymidazolam glucuronide from ultrafiltrate. <i>Critical Care</i> , 2010 , 14, P512	10.8	78
135	Continuous renal replacement therapy: are we getting the dose right?. <i>Critical Care</i> , 2010 , 14, P513	10.8	78
134	Use of CRRT-CVVH in a hemato-oncological ICU in patients treated with defibrotide for veno-occlusive disease in patients after allogeneic hematopoietic stem cell transplantation: single-center experience. <i>Critical Care</i> , 2010 , 14, P514	10.8	1
133	Comparison of the efficacy and safety of two regional citrate anticoagulation protocols using acid citrate dextrose A or Prismocitrate 10/2, in patients with acute renal failure undergoing continuous venovenous haemodiafiltration. <i>Critical Care</i> , 2010 , 14, P515	10.8	2
132	Addition of prostacyclin to heparin does not prolong circuit life during continuous haemofiltration. <i>Critical Care</i> , 2010 , 14, P516	10.8	78
131	Recurrent early filter clotting in regional citrate anticoagulated continuous venovenous hemodialysis due to undetected antibodies to heparin-platelet factor 4 complexes. <i>Critical Care</i> , 2010 , 14, P517	10.8	1
130	Pilot study on a regional citrate anticoagulated continuous venovenous hemodiafiltration protocol with variable treatment dose. <i>Critical Care</i> , 2010 , 14, P518	10.8	1
129	Characteristics and outcome of patients treated with renal replacement therapy: single-center, 2-year experience. <i>Critical Care</i> , 2010 , 14, P519	10.8	78
128	Renal function in living related kidney donors above the age of 70. <i>Critical Care</i> , 2010 , 14, P520	10.8	78
127	Augmented renal clearance in traumatic brain injury. <i>Critical Care</i> , 2010 , 14, P521	10.8	2
126	Pressure-controlled hemorrhagic shock in mice: a new model of acute kidney injury. <i>Critical Care</i> , 2010 , 14, P522	10.8	78
125	Continuous urinary electrolyte measurement in a swine model of mechanical ventilation. <i>Critical Care</i> , 2010 , 14, P523	10.8	78
124	Impact of positive fluid balance on survival in critically ill cancer patients. <i>Critical Care</i> , 2010 , 14, P524	10.8	78

123	Fluid accumulation has an adverse effect on outcome of ICU patients with acute kidney injury. <i>Critical Care</i> , 2010 , 14, P525	10.8	78
122	Significant microalbuminuria, on admission to the ICU, may indicate AKI in patients with sepsis. <i>Critical Care</i> , 2010 , 14, P526	10.8	78
121	Neutrophil gelatinase-associated lipocalin as an early indicator for postoperative renal failure. <i>Critical Care</i> , 2010 , 14, P527	10.8	78
120	Neutrophil gelatinase-associated lipocalin as an early indicator for mortality in adult critically ill patients with acute kidney injury. <i>Critical Care</i> , 2010 , 14, P528	10.8	1
119	Urinary and plasma NGAL levels reflect the severity of acute kidney injury. <i>Critical Care</i> , 2010 , 14, P529	10.8	78
118	Economic impact and cost-effectiveness of urine neutrophil gelatinase-associated lipocalin following cardiac surgery. <i>Critical Care</i> , 2010 , 14, P530	10.8	2
117	Neutrophil gelatinase-associated lipocalin predicts acute kidney injury in critically ill patients. <i>Critical Care</i> , 2010 , 14, P531	10.8	78
116	Study of serum vancomycin levels in different modalities of hemodialysis in Indian patients. <i>Critical Care</i> , 2010 , 14, P532	10.8	78
115	RIFLE criteria for acute kidney dysfunction following liver transplantation: incidence and risk factors. <i>Critical Care</i> , 2010 , 14, P533	10.8	78
114	Prophylactic use of laxative for constipation in critically ill patients. <i>Critical Care</i> , 2010 , 14, P534	10.8	1
113	Study of intra-abdominal hypertension in the medical intensive care unit. <i>Critical Care</i> , 2010 , 14, P535	10.8	78
112	Intra-abdominal pressure values of emergency department intensive care unit patients and clinical outcomes. <i>Critical Care</i> , 2010 , 14, P536	10.8	78
111	Advanced haemodynamic assessment of patients with liver disease and abdominal compartment syndrome. <i>Critical Care</i> , 2010 , 14, P537	10.8	78
110	Molecular adsorbent recirculating system in hepatic failure patients with a ventricular assist device. <i>Critical Care</i> , 2010 , 14, P538	10.8	78
109	Molecular adsorbent recirculating system in patients with acute liver failure: prospective follow-up. <i>Critical Care</i> , 2010 , 14, P539	10.8	78
108	A study of patients with cirrhosis admitted to nontransplant general intensive care in the UK: prevalence, case mix, outcomes and evaluation of critical illness and disease-specific scoring systems. <i>Critical Care</i> , 2010 , 14, P540	10.8	1
107	Acute liver failure: a European perspective. <i>Critical Care</i> , 2010 , 14, P541	10.8	78
106	Intracranial pressure monitoring in acute liver failure: a review of 10 years experience. <i>Critical Care</i> , 2010 , 14, P542	10.8	8

105	Microdialysis catheters in liver transplants detect vascular complications and rejection. <i>Critical Care</i> , 2010 , 14, P543	10.8	78
104	Drug-induced hepatic cholestasis in the ICU: a case-control study. <i>Critical Care</i> , 2010 , 14, P544	10.8	78
103	Diagnosis of acute cholecystitis in septic patients. <i>Critical Care</i> , 2010 , 14, P545	10.8	78
102	Bleeding and pneumonia in intensive care unit patients given PPIs and H2RAs for prevention of stress ulcer: a meta analysis. <i>Critical Care</i> , 2010 , 14, P546	10.8	78
101	Influence of septic shock on energy expenditure in critically ill ICU patients, estimated with indirect calorimetry. <i>Critical Care</i> , 2010 , 14, P547	10.8	78
100	Evaluation of a novel motility capsule for gastric emptying in a porcine model of acute lung injury. <i>Critical Care</i> , 2010 , 14, P548	10.8	78
99	Preliminary results on the validation of a novel method for measuring intra-abdominal pressure and gastric residual volume. <i>Critical Care</i> , 2010 , 14, P549	10.8	78
98	Monitoring enteral nutrition efficacy in ICU patients: is there a place for the Brix value-derived fraction of enteral formula in gastric residual volume?. <i>Critical Care</i> , 2010 , 14, P550	10.8	0
97	Initial gastric volume, nasogastric tube and pneumonia incidence in trauma patients. <i>Critical Care</i> , 2010 , 14, P551	10.8	78
96	Achieving nutritional targets in the ICU. <i>Critical Care</i> , 2010 , 14, P552	10.8	78
95	Evaluation of studies of nutritional support team: 2 years' experience. <i>Critical Care</i> , 2010 , 14, P553	10.8	78
94	Are we feeding our critically ill patients appropriately?. <i>Critical Care</i> , 2010 , 14, P554	10.8	78
93	Effect of increased enteral nutrition on ICU-acquired infections: a multicenter observational study. <i>Critical Care</i> , 2010 , 14, P555	10.8	78
92	Impact of early enteral nutrition up to goal on the prognosis of patients with different severity of illness in the ICU. <i>Critical Care</i> , 2010 , 14, P556	10.8	78
91	Benefit of supplemental parenteral nutrition in the critically ill patient? Results of a multicenter observational study. <i>Critical Care</i> , 2010 , 14, P557	10.8	2
90	Role of fiber diet formula in surgical critically ill septic patients. <i>Critical Care</i> , 2010 , 14, P558	10.8	78
89	Effect of oral supplementation with immunonutrients on radiation-induced hematologic toxicity in cancer patients. <i>Critical Care</i> , 2010 , 14, P559	10.8	1
88	Impact of multichamber bag parenteral nutrition (OliClinomel) upon inflammatory markers (EPICOS Study). <i>Critical Care</i> , 2010 , 14, P560	10.8	78

87	Effect of n-3 fatty acids on cerebral markers and the inflammatory response in sepsis. <i>Critical Care</i> , 2010 , 14, P561	10.8	78
86	Effects of an olive oil-based lipid emulsion (ClinOleic) versus LCT/MCT-based lipid emulsions upon the inflammatory markers of critically ill patients (EPICOS Study). <i>Critical Care</i> , 2010 , 14, P562	10.8	78
85	Does a fish-oil-containing lipid emulsion improve liver function in comparison with a soybean oil lipid emulsion?. <i>Critical Care</i> , 2010 , 14, P565	10.8	78
84	Therapy of compartmentalization of the proinflammatory and anti-inflammatory cytokines in sepsis. <i>Critical Care</i> , 2010 , 14, P566	10.8	1
83	Intensive insulin therapy and brief hypoglycemia do not increase neurological injury markers in critically ill children. <i>Critical Care</i> , 2010 , 14, P567	10.8	78
82	A cluster-controlled implementation project of intensive insulin therapy: effects on blood glucose control and incidence of severe hypoglycemia. <i>Critical Care</i> , 2010 , 14, P568	10.8	78
81	Incidence of hypoglycemia in NICE sugar not so nice after consideration of the treatment period. <i>Critical Care</i> , 2010 , 14, P569	10.8	78
80	Comparison of a planned versus emergent approach implementation of tight glycaemic control therapy on an ICU: an experimental design. <i>Critical Care</i> , 2010 , 14, P570	10.8	78
79	Coagulation profile in hyperglycemic septic shock patients. <i>Critical Care</i> , 2010 , 14, P571	10.8	78
78	Impact of glycaemic control on survival in critical care patients with sepsis. <i>Critical Care</i> , 2010 , 14, P572	10.8	1
77	Therapeutic impact on intravenous insulin dosing with three new point-of-care glucose meters in an adult ICU. <i>Critical Care</i> , 2010 , 14, P573	10.8	78
76	Significance of blood glucose variability in the early phase for predicting survival in acutely ill patients with glucose intolerance. <i>Critical Care</i> , 2010 , 14, P574	10.8	78
75	Blood glucose amplitude variation: effects of intensive insulin therapy and relative association with mortality. <i>Critical Care</i> , 2010 , 14, P575	10.8	78
74	Prevention of fatal and disabling neuroglycopenia: a comparison of arterial blood sampling methods. <i>Critical Care</i> , 2010 , 14, P576	10.8	78
73	Impact of real-time continuous glucose monitoring on glucose variability in critically ill patients. <i>Critical Care</i> , 2010 , 14, P577	10.8	78
72	Automated blood glucose monitor is insensitive to hematocrit variation and common saccharide interferences. <i>Critical Care</i> , 2010 , 14, P578	10.8	78
71	Performance of an automated blood glucose monitor in the OR/ICU. <i>Critical Care</i> , 2010 , 14, P579	10.8	78
70	ICU glucose monitoring measured in plasma using mid-infrared spectroscopy. <i>Critical Care</i> , 2010 , 14, P580	10.8	1

69	Validation of HbA1c as a diagnostic marker for diabetes in the critically ill. <i>Critical Care</i> , 2010 , 14, P581	10.8	1
68	Effect of intravenous glucocorticoids on the efficiency of tight glycaemic control. <i>Critical Care</i> , 2010 , 14, P582	10.8	78
67	Chronic critical illness: high levels of estrogens and mortality. <i>Critical Care</i> , 2010 , 14, P583	10.8	78
66	Estrogen decreases inflammation and apoptotic signaling in the heart following severe burn injury. <i>Critical Care</i> , 2010 , 14, P584	10.8	78
65	Dexamethasone treatment during cardiac surgery and postoperative lactate levels. <i>Critical Care</i> , 2010 , 14, P585	10.8	78
64	Risk factors for hypertriglyceridemia in the intensive care unit: an exploratory study. <i>Critical Care</i> , 2010 , 14, P586	10.8	78
63	Septic patients show significantly lower 1,25-dihydroxy vitamin D levels than trauma patients. <i>Critical Care</i> , 2010 , 14, P587	10.8	1
62	High prevalence of severe vitamin D deficiency in intensive care patients. <i>Critical Care</i> , 2010 , 14, P588	10.8	3
61	Vitamin D, parathyroid hormone and serum calcium levels and their association with hospital mortality in critically ill patients. <i>Critical Care</i> , 2010 , 14, P589	10.8	5
60	Effect of restoration of 25OHD3 status in prolonged critical illness on serum innate immunity parameters LL-37 and sCD163. <i>Critical Care</i> , 2010 , 14, P590	10.8	10
59	Alterations in adipose tissue during critical illness: an adaptive and protective response?. <i>Critical Care</i> , 2010 , 14, P591	10.8	1
58	Impact of collection method and sample handling on measured levels of circulating ACTH. <i>Critical Care</i> , 2010 , 14, P592	10.8	1
57	Prognostic value of serum zinc levels in critically ill patients. <i>Critical Care</i> , 2010 , 14, P593	10.8	1
56	High stress during admission predicts the impact of a PICU follow-up clinic on parents' psychological outcome. <i>Critical Care</i> , 2010 , 14, P594	10.8	1
55	Patient's contact person in the intensive care unit: is it really important?. <i>Critical Care</i> , 2010 , 14, P595	10.8	78
54	Evaluating the quality of communication with patients' relatives in critical care. <i>Critical Care</i> , 2010 , 14, P596	10.8	78
53	Family satisfaction on the intensive care unit. <i>Critical Care</i> , 2010 , 14, P597	10.8	78
52	Survey into bereavement of family members of patients who died in the intensive care unit. <i>Critical Care</i> , 2010 , 14, P598	10.8	78

51	Exploration of the needs and experiences of family members of a patient being admitted to a neurosurgery intensive care unit in Shiraz, Iran. <i>Critical Care</i> , 2010 , 14, P599	10.8	78
50	Deciding the end of life for patients with disorders of consciousness: a European survey. <i>Critical Care</i> , 2010 , 14, P600	10.8	78
49	Evaluation of a new limitations and withdrawal of care framework order form in the intensive care unit. <i>Critical Care</i> , 2010 , 14, P601	10.8	78
48	Examination of the utility of serum lactate and base deficit in hemorrhagic shock. <i>Critical Care</i> , 2010 , 14, P161	10.8	78
47	Response of NICOM stroke volume to passive leg raising to predict fluid responsiveness in critically ill patients with spontaneous breathing activity. <i>Critical Care</i> , 2010 , 14, P113	10.8	2
46	[Assessment of health care providers on family presence during resuscitation]. <i>Enfermería Clínica</i> , 2011 , 21, 227-9	0.6	1
45	Family presence preference when patients are receiving resuscitation in an accident and emergency department. <i>Journal of Advanced Nursing</i> , 2011 , 67, 56-67	3.1	29
44	An introduction to the multisystem model of knowledge integration and translation. <i>Advances in Nursing Science</i> , 2011 , 34, 29-38	2.1	20
43	Challenges in conducting research after family presence during resuscitation. <i>Journal of Trauma Nursing: the Official Journal of the Society of Trauma Nurses</i> , 2012 , 19, 190-4	1	3
42	Family engagement regarding the critically ill patient. <i>Surgical Clinics of North America</i> , 2012 , 92, 1637-47	3.7	14
41	"It's just what we do": a qualitative study of emergency nurses working with well-established family presence protocol. <i>Journal of Emergency Nursing</i> , 2012 , 38, 329-34	1.3	28
40	Parental presence during child resuscitation: a critical review of a research article. <i>Journal of Perioperative Practice</i> , 2012 , 22, 63-6	0.4	1
39	Attitudes, implementation and practice of family presence during resuscitation (FPDR): a quantitative literature review. <i>International Emergency Nursing</i> , 2013 , 21, 26-34	2.4	33
38	Family-centered care in the pediatric intensive care unit. <i>Pediatric Clinics of North America</i> , 2013 , 60, 761-72	3.8	77
37	Assessing health professionals' perceptions of family presence during resuscitation: a replication study. <i>International Emergency Nursing</i> , 2013 , 21, 17-25	2.4	28
36	Facilitated family presence at resuscitation: effectiveness of a nursing student toolkit. <i>Nurse Education Today</i> , 2013 , 33, 1258-63	3.7	10
35	Implemented family presence protocols: an integrative review. <i>Journal of Nursing Care Quality</i> , 2013 , 28, 281-8	1.7	12
34	Changes in provider perceptions of family presence during resuscitation. <i>Clinical Nurse Specialist</i> , 2013 , 27, 239-44	0.6	9

33	Integrative review: nurses' and physicians' experiences and attitudes towards inpatient-witnessed resuscitation of an adult patient. <i>Journal of Advanced Nursing</i> , 2014 , 70, 957-74	3.1	30
32	Resuscitation team perceptions of family presence during CPR. <i>Advanced Emergency Nursing Journal</i> , 2014 , 36, 325-34	0.8	16
31	Family presence during resuscitation (FPDR): Perceived benefits, barriers and enablers to implementation and practice. <i>International Emergency Nursing</i> , 2014 , 22, 69-74	2.4	38
30	Australian Emergency Department health professionals' reasons to invite or not invite Family Witnessed Resuscitation: a qualitative perspective. <i>International Emergency Nursing</i> , 2014 , 22, 18-24	2.4	16
29	Factors associated with experiences and attitudes of healthcare professionals towards family-witnessed resuscitation: a cross-sectional study. <i>Journal of Advanced Nursing</i> , 2015 , 71, 2595-608	3.1	12
28	Family presence during resuscitation: A Canadian Critical Care Society position paper. <i>Canadian Respiratory Journal</i> , 2015 , 22, 201-5	2.1	37
27	Family presence during resuscitation (FPDR): A survey of emergency personnel in Victoria, Australia. <i>Australasian Emergency Nursing Journal</i> , 2015 , 18, 98-105		16
26	Predictors of public support for family presence during cardiopulmonary resuscitation: A population based study. <i>International Journal of Nursing Studies</i> , 2015 , 52, 1064-70	5.8	23
25	Inviting family to be present during cardiopulmonary resuscitation: Impact of education. <i>Nurse Education in Practice</i> , 2016 , 16, 274-9	3.2	12
24	Family presence during resuscitation: Validation of the riskBenefit and self-confidence scales for student nurses. <i>Journal of Research in Nursing</i> , 2016 , 21, 306-322	0.9	6
23	When Someone Dies in the Emergency Department: Perspectives of Emergency Nurses. <i>Journal of Emergency Nursing</i> , 2016 , 42, 207-12	1.3	18
22	Psychometric properties of the critical care family needs inventory-emergency department. <i>Applied Nursing Research</i> , 2017 , 33, 113-120	1.8	2
21	Predicting medical professionals' intention to allow family presence during resuscitation: A cross sectional survey. <i>International Journal of Nursing Studies</i> , 2017 , 70, 11-16	5.8	6
20	Family Presence During Trauma Resuscitation: Family Members' Attitudes, Behaviors, and Experiences. <i>American Journal of Critical Care</i> , 2017 , 26, 229-239	1.7	21
19	Family Presence in the Intensive Care Unit and Emergency Department: A Metasynthesis. <i>Journal of Family Theory and Review</i> , 2017 , 9, 201-218	2.3	5
18	Factors influencing medical staff's intentions to implement family-witnessed cardiopulmonary resuscitation: A cross-sectional, multihospital survey. <i>European Journal of Cardiovascular Nursing</i> , 2017 , 16, 492-501	3.3	5
17	Effect of Family Presence on Advanced Trauma Life Support Task Performance During Pediatric Trauma Team Evaluation. <i>Pediatric Emergency Care</i> , 2017 ,	1.4	2
16	Family presence during resuscitation: A concise narrative review. <i>Trends in Anaesthesia and Critical Care</i> , 2017 , 15, 12-16	0.4	8

15	Should Family be Allowed During Resuscitation. 2017 ,		2
14	Owner-witnessed cardiopulmonary resuscitation in small animal emergency veterinary medicine. <i>Journal of the American Veterinary Medical Association</i> , 2018 , 253, 1032-1037	1	2
13	A survey of emergency department staff's opinions and experiences of family presence during invasive procedures and resuscitation. <i>Emergency Nurse</i> , 2019 , 27, 13-19	0.7	3
12	Family Presence in the Adult ICU During Bedside Procedures. <i>Journal of Intensive Care Medicine</i> , 2019 , 34, 587-593	3.3	2
11	Experiences and Opinions of Patients and Their Relatives to Family Presence During Adult Resuscitation in Poland: Quantitative Research. <i>Patient Preference and Adherence</i> , 2020 , 14, 227-234	2.4	1
10	Family Presence During Resuscitation: A Critical Review of the Literature. <i>American Journal of Critical Care</i> , 2005 , 14, 494-511	1.7	72
9	Nurses' Perceptions of Their Self-confidence and the Benefits and Risks of Family Presence During Resuscitation. <i>American Journal of Critical Care</i> , 2008 , 17, 101-111	1.7	60
8	Impact of a Multifaceted Intervention on Nurses' and Physicians' Attitudes and Behaviors Toward Family Presence During Resuscitation. <i>Critical Care Nurse</i> , 2007 , 27, 52-61	1.6	72
7	My experience of being present during resuscitation. <i>Critical Care Nurse</i> , 2007 , 27, 17-17	1.6	1
6	Family Presence During Resuscitation. <i>Journal of Continuing Education in Nursing</i> , 2008 , 39, 530-531	0.9	1
5	Knowledge, Attitude, and Practice of Healthcare Professionals Regarding Family Presence During Resuscitation: An Interventional Study in a Tertiary Care Setting, Karachi, Pakistan. <i>Connect the World of Critical Care Nursing</i> , 2019 , 13, 46-55	0.3	0
4	Parental presence during pediatric emergency procedures: finding answers in an Asian context. <i>Clinical and Experimental Emergency Medicine</i> , 2019 , 6, 340-344	1.7	3
3	Family Presence and Audience Effects on Cardiopulmonary Resuscitation: Perceptions of Pre-hospital Emergency Caregivers in Turkey.		
2	Parent Participation in Morbidity and Mortality Review: Parent and Physician Perspectives. <i>Journal of Patient Experience</i> , 2022 , 9, 237437352211026	1.3	
1	General Approach to the Pediatric Patient. 2013 , 113-116.e1		0