

CITATION REPORT

List of articles citing

Genome-wide insertional mutagenesis of *Arabidopsis thaliana*

DOI: 10.1126/science.1086391
Science, 2003, 301, 653-7.

Source: <https://exaly.com/paper-pdf/35532798/citation-report.pdf>

Version: 2024-04-25

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
2303	Abstract.		
2302	Progressive akinetic mutism in cerebellar hemangioblastoma with "normal-pressure hydrocephalus". 1969 , 19, 32-6		10
2301	The prognostic significance of thymic germinal center proliferation in myasthenia gravis. 1973 , 23, 69-72		14
2300	An Arabidopsis thaliana T-DNA mutagenized population (GABI-Kat) for flanking sequence tag-based reverse genetics. 2003 , 53, 247-59		597
2299	Regulation of flowering time by histone acetylation in Arabidopsis. <i>Science</i> , 2003 , 302, 1751-4	33.3	385
2298	Empirical analysis of transcriptional activity in the Arabidopsis genome. <i>Science</i> , 2003 , 302, 842-6	33.3	782
2297	Plant responses to ethylene gas are mediated by SCF(EBF1/EBF2)-dependent proteolysis of EIN3 transcription factor. 2003 , 115, 667-77		689
2296	Site-specific integration of Agrobacterium tumefaciens T-DNA via double-stranded intermediates. 2003 , 133, 1011-23		121
2295	VARICOSE, a WD-domain protein, is required for leaf blade development. 2003 , 130, 6577-88		44
2294	IBR5, a dual-specificity phosphatase-like protein modulating auxin and abscisic acid responsiveness in Arabidopsis. 2003 , 15, 2979-91		130
2293	Agrobacterium tumefaciens and the plant: the David and Goliath of modern genetics. 2003 , 133, 948-55		39
2292	Generation and analysis of end sequence database for T-DNA tagging lines in rice. 2003 , 133, 2040-7		213
2291	Interaction of Arabidopsis Membrane Proteins with SPIRAL1. 2004 , 21, 261-267		
2290	The role of SEUSS in auxin response and floral organ patterning. 2004 , 131, 4697-707		61
2289	A homolog of prokaryotic thiol disulfide transporter CcdA is required for the assembly of the cytochrome b6f complex in Arabidopsis chloroplasts. 2004 , 279, 32474-82		79
2288	Genome-wide analysis of Arabidopsis pentatricopeptide repeat proteins reveals their essential role in organelle biogenesis. 2004 , 16, 2089-103		956
2287	Genetics of the biogenesis and dynamics of the photosynthetic machinery in eukaryotes. 2004 , 16, 1650-60		42

2286	Chlamydomonas and Arabidopsis. A dynamic duo. 2004 , 135, 607-10	57
2285	Arabidopsis CAND1, an unmodified CUL1-interacting protein, is involved in multiple developmental pathways controlled by ubiquitin/proteasome-mediated protein Degradation. 2004 , 16, 1870-82	105
2284	Arabidopsis to rice. Applying knowledge from a weed to enhance our understanding of a crop species. 2004 , 135, 622-9	68
2283	Identification of the Arabidopsis palmitoyl-monogalactosyldiacylglycerol delta7-desaturase gene FAD5, and effects of plastidial retargeting of Arabidopsis desaturases on the fad5 mutant phenotype. 2004 , 136, 4237-45	75
2282	Arabidopsis HMA2, a divalent heavy metal-transporting P(1B)-type ATPase, is involved in cytoplasmic Zn ²⁺ homeostasis. 2004 , 136, 3712-23	171
2281	The phytochrome-interacting transcription factor, PIF3, acts early, selectively, and positively in light-induced chloroplast development. 2004 , 101, 16091-8	244
2280	Towards an understanding of photosynthetic acclimation. 2005 , 56, 435-47	407
2279	Circadian rhythms of ethylene emission in Arabidopsis. 2004 , 136, 3751-61	132
2278	Interaction of Polycomb-group proteins controlling flowering in Arabidopsis. 2004 , 131, 5263-76	384
2277	A mechanism related to the yeast transcriptional regulator Paf1c is required for expression of the Arabidopsis FLC/MAF MADS box gene family. 2004 , 16, 2940-53	142
2276	Type-A Arabidopsis response regulators are partially redundant negative regulators of cytokinin signaling. 2004 , 16, 658-71	503
2275	Arabidopsis thaliana DNA gyrase is targeted to chloroplasts and mitochondria. 2004 , 101, 7821-6	124
2274	PIL5, a phytochrome-interacting basic helix-loop-helix protein, is a key negative regulator of seed germination in Arabidopsis thaliana. 2004 , 16, 3045-58	324
2273	Reentry of the Ethylene MPK6 Module. 2004 , 16, 3169-3173	54
2272	GCR1 can act independently of heterotrimeric G-protein in response to brassinosteroids and gibberellins in Arabidopsis seed germination. 2004 , 135, 907-15	148
2271	ADL2a, like ADL2b, is involved in the control of higher plant mitochondrial morphology. 2004 , 55, 783-5	88
2270	Phosphorylation of 1-aminocyclopropane-1-carboxylic acid synthase by MPK6, a stress-responsive mitogen-activated protein kinase, induces ethylene biosynthesis in Arabidopsis. 2004 , 16, 3386-99	635
2269	Members of the Toc159 import receptor family represent distinct pathways for protein targeting to plastids. 2004 , 15, 3379-92	164

2268	Interchangeable functions of Arabidopsis PIROGI and the human WAVE complex subunit SRA1 during leaf epidermal development. 2004 , 131, 4345-55	104
2267	Arabidopsis ETA2, an apparent ortholog of the human cullin-interacting protein CAND1, is required for auxin responses mediated by the SCF(TIR1) ubiquitin ligase. 2004 , 16, 1883-97	92
2266	Deletion of the chloroplast-localized Thylakoid formation1 gene product in Arabidopsis leads to deficient thylakoid formation and variegated leaves. 2004 , 136, 3594-604	140
2265	PIN-FORMED1 and PINOID regulate boundary formation and cotyledon development in Arabidopsis embryogenesis. 2004 , 131, 5021-30	202
2264	Disparate roles for the regulatory A subunit isoforms in Arabidopsis protein phosphatase 2A. 2004 , 16, 709-22	88
2263	Large-scale mutagenesis of the yeast genome using a Tn7-derived multipurpose transposon. 2004 , 14, 1975-86	43
2262	A transcriptional coactivator, AtGIF1, is involved in regulating leaf growth and morphology in Arabidopsis. 2004 , 101, 13374-9	253
2261	Nuclear SDH2-1 and SDH2-2 genes, encoding the iron-sulfur subunit of mitochondrial complex II in Arabidopsis, have distinct cell-specific expression patterns and promoter activities. 2004 , 136, 4072-87	59
2260	P-type ATPase heavy metal transporters with roles in essential zinc homeostasis in Arabidopsis. 2004 , 16, 1327-39	532
2259	Extensive phenotypic variation in early flowering mutants of Arabidopsis. 2004 , 135, 201-11	31
2258	IAR4, a gene required for auxin conjugate sensitivity in Arabidopsis, encodes a pyruvate dehydrogenase E1alpha homolog. 2004 , 135, 989-99	31
2257	Inactivation of the clpC1 gene encoding a chloroplast Hsp100 molecular chaperone causes growth retardation, leaf chlorosis, lower photosynthetic activity, and a specific reduction in photosystem content. 2004 , 136, 4114-26	115
2256	Genome-wide analysis of spatial gene expression in Arabidopsis flowers. 2004 , 16, 1314-26	201
2255	NAPP and PIRP encode subunits of a putative wave regulatory protein complex involved in plant cell morphogenesis. 2004 , 16, 2335-49	87
2254	A novel NADPH thioredoxin reductase, localized in the chloroplast, which deficiency causes hypersensitivity to abiotic stress in Arabidopsis thaliana. 2004 , 279, 43821-7	268
2253	mRNA deadenylation by PARN is essential for embryogenesis in higher plants. 2004 , 10, 1200-14	68
2252	Loss of plastidic lysophosphatidic acid acyltransferase causes embryo-lethality in Arabidopsis. 2004 , 45, 503-10	92
2251	The geminivirus nuclear shuttle protein is a virulence factor that suppresses transmembrane receptor kinase activity. 2004 , 18, 2545-56	150

2250	Ice-cap. A high-throughput method for capturing plant tissue samples for genotype analysis. 2004 , 135, 1162-9	14
2249	FRIGIDA-related genes are required for the winter-annual habit in Arabidopsis. 2004 , 101, 3281-5	125
2248	Calcium/calmodulin up-regulates a cytoplasmic receptor-like kinase in plants. 2004 , 279, 42552-9	72
2247	Natural genetic variation in Arabidopsis identifies BREVIS RADIX, a novel regulator of cell proliferation and elongation in the root. 2004 , 18, 700-14	185
2246	Identification of genes required for embryo development in Arabidopsis. 2004 , 135, 1206-20	378
2245	Cloning and functional characterization of a phospholipid:diacylglycerol acyltransferase from Arabidopsis. 2004 , 135, 1324-35	217
2244	The Cell Wall Navigator database. A systems-based approach to organism-unrestricted mining of protein families involved in cell wall metabolism. 2004 , 136, 3003-8; discussion 3001	62
2243	Molecular analysis of high-copy insertion sites in maize. 2004 , 32, e54	65
2242	Lipid utilization, gluconeogenesis, and seedling growth in Arabidopsis mutants lacking the glyoxylate cycle enzyme malate synthase. 2004 , 279, 42916-23	92
2241	AtNAP7 is a plastidic SufC-like ATP-binding cassette/ATPase essential for Arabidopsis embryogenesis. 2004 , 101, 9143-8	110
2240	Blue light-induced transcription of plastid-encoded psbD gene is mediated by a nuclear-encoded transcription initiation factor, AtSig5. 2004 , 101, 3304-9	103
2239	An Arabidopsis indole-3-butyric acid-response mutant defective in PEROXIN6, an apparent ATPase implicated in peroxisomal function. 2004 , 101, 1786-91	114
2238	A novel protein for photosystem I biogenesis. 2004 , 279, 10243-51	72
2237	Peroxisomal Acyl-CoA synthetase activity is essential for seedling development in Arabidopsis thaliana. 2004 , 16, 394-405	186
2236	Nuclear protein phosphatases with Kelch-repeat domains modulate the response to brassinosteroids in Arabidopsis. 2004 , 18, 448-60	289
2235	UPS1 and UPS2 from Arabidopsis mediate high affinity transport of uracil and 5-fluorouracil. 2004 , 279, 44817-24	49
2234	SGS3 and SGS2/SDE1/RDR6 are required for juvenile development and the production of trans-acting siRNAs in Arabidopsis. 2004 , 18, 2368-79	711
2233	Functional reclassification of the putative cinnamyl alcohol dehydrogenase multigene family in Arabidopsis. 2004 , 101, 1455-60	186

2232	Molecular characterization of two highly homologous receptor-like kinase genes, RLK902 and RKL1, in <i>Arabidopsis thaliana</i> . 2004 , 68, 1935-41	25
2231	Plant proteins that interact with VirB2, the <i>Agrobacterium tumefaciens</i> pilin protein, mediate plant transformation. 2004 , 16, 3148-67	100
2230	Regulation of flowering time in <i>Arabidopsis</i> by K homology domain proteins. 2004 , 101, 12759-64	128
2229	Mre11 deficiency in <i>Arabidopsis</i> is associated with chromosomal instability in somatic cells and Spo11-dependent genome fragmentation during meiosis. 2004 , 16, 1968-78	163
2228	The SHINE clade of AP2 domain transcription factors activates wax biosynthesis, alters cuticle properties, and confers drought tolerance when overexpressed in <i>Arabidopsis</i> . 2004 , 16, 2463-80	539
2227	Functional specialization amongst the <i>Arabidopsis</i> Toc159 family of chloroplast protein import receptors. 2004 , 16, 2059-77	154
2226	The gravitropism defective 2 mutants of <i>Arabidopsis</i> are deficient in a protein implicated in endocytosis in <i>Caenorhabditis elegans</i> . 2004 , 136, 3095-103; discussion 3002	67
2225	Versatile gene-specific sequence tags for <i>Arabidopsis</i> functional genomics: transcript profiling and reverse genetics applications. 2004 , 14, 2176-89	265
2224	<i>Arabidopsis</i> FHY3/FAR1 gene family and distinct roles of its members in light control of <i>Arabidopsis</i> development. 2004 , 136, 4010-22	88
2223	Reserve mobilization in the <i>Arabidopsis</i> endosperm fuels hypocotyl elongation in the dark, is independent of abscisic acid, and requires PHOSPHOENOLPYRUVATE CARBOXYKINASE1. 2004 , 16, 2705-18	195
2222	RHM2 is involved in mucilage pectin synthesis and is required for the development of the seed coat in <i>Arabidopsis</i> . 2004 , 134, 286-95	113
2221	MIPS <i>Arabidopsis thaliana</i> Database (MAtDB): an integrated biological knowledge resource for plant genomics. 2004 , 32, D373-6	74
2220	<i>Arabidopsis</i> CYP707As encode (+)-abscisic acid 8'-hydroxylase, a key enzyme in the oxidative catabolism of abscisic acid. 2004 , 134, 1439-49	414
2219	<i>Cryptococcus neoformans</i> virulence gene discovery through insertional mutagenesis. 2004 , 3, 420-9	161
2218	T-DNA locus structure in a large population of soybean plants transformed using the <i>Agrobacterium</i> -mediated cotyledonary-node method. 2004 , 2, 289-300	39
2217	Mutants for photosystem I subunit D of <i>Arabidopsis thaliana</i> : effects on photosynthesis, photosystem I stability and expression of nuclear genes for chloroplast functions. 2004 , 37, 839-52	99
2216	An outer envelope membrane component of the plastid protein import apparatus plays an essential role in <i>Arabidopsis</i> . 2004 , 38, 93-106	96
2215	<i>Arabidopsis</i> Yellow Stripe-Like2 (YSL2): a metal-regulated gene encoding a plasma membrane transporter of nicotianamine-metal complexes. 2004 , 39, 403-14	272

2214	Roles of the AtErcc1 protein in recombination. 2004 , 39, 334-42	44
2213	High throughput T-DNA insertion mutagenesis in rice: a first step towards in silico reverse genetics. 2004 , 39, 450-64	217
2212	A compendium of methods useful for characterizing Arabidopsis pollen mutants and gametophytically-expressed genes. 2004 , 39, 761-75	190
2211	Establishing gene function by mutagenesis in Arabidopsis thaliana. 2004 , 39, 682-96	114
2210	Nfu2: a scaffold protein required for [4Fe-4S] and ferredoxin iron-sulphur cluster assembly in Arabidopsis chloroplasts. 2004 , 40, 101-11	87
2209	The ubiquitin ligase XBAT32 regulates lateral root development in Arabidopsis. 2004 , 40, 996-1006	77
2208	AtMYB32 is required for normal pollen development in Arabidopsis thaliana. 2004 , 40, 979-95	222
2207	Arabidopsis vacuolar H-ATPase subunit E isoform 1 is required for Golgi organization and vacuole function in embryogenesis. 2005 , 41, 125-32	72
2206	Genome-wide analysis of glucose-6-phosphate dehydrogenases in Arabidopsis. 2005 , 41, 243-56	110
2205	An Arabidopsis NPR1-like gene, NPR4, is required for disease resistance. 2005 , 41, 304-18	112
2204	Molecular and functional characterization of Arabidopsis Cullin 3A. 2005 , 41, 386-99	76
2203	In vivo studies on the roles of Tic110, Tic40 and Hsp93 during chloroplast protein import. 2005 , 41, 412-28	152
2202	Identification of Arabidopsis thaliana transformants without selection reveals a high occurrence of silenced T-DNA integrations. 2005 , 41, 464-77	68
2201	Differing requirements for the Arabidopsis Rad51 paralogs in meiosis and DNA repair. 2005 , 41, 533-45	121
2200	A collection of 11 800 single-copy Ds transposon insertion lines in Arabidopsis. 2004 , 37, 897-905	183
2199	NAD - new roles in signalling and gene regulation in plants. 2004 , 163, 31-44	101
2198	MACROALGAL CANDIDATES FOR GENOMICS1. 2004 , 40, 26-33	61
2197	Systematic genome-wide screens of gene function. 2004 , 5, 11-22	261

2196	Maximizing the potential of functional genomics. 2004 , 5, 190-201	79
2195	The Arabidopsis homologue of Xrcc3 plays an essential role in meiosis. 2004 , 23, 439-49	115
2194	The Arabidopsis cytochrome P450 CYP707A encodes ABA 8'-hydroxylases: key enzymes in ABA catabolism. 2004 , 23, 1647-56	683
2193	Molecular analysis of telomere fusions in Arabidopsis: multiple pathways for chromosome end-joining. 2004 , 23, 2304-13	145
2192	A proteoglycan mediates inductive interaction during plant vascular development. 2004 , 429, 873-8	264
2191	Rapid BDNF-induced retrograde synaptic modification in a developing retinotectal system. 2004 , 429, 878-83	94
2190	Plant biology: good neighbours. 2004 , 430, 302-4	8
2189	Solar physics: hidden magnetism. 2004 , 430, 304-5	15
2188	The AtNACK1/HINKEL and STUD/TETRASPORE/AtNACK2 genes, which encode functionally redundant kinesins, are essential for cytokinesis in Arabidopsis. 2004 , 9, 1199-211	101
2187	Agrobacterium T-DNA integration: molecules and models. 2004 , 20, 375-83	225
2186	Integrating signals in stomatal development. 2004 , 7, 26-32	75
2185	Conservation and diversity in flower land. 2004 , 7, 84-91	133
2184	The ethylene signaling pathway: new insights. 2004 , 7, 40-9	729
2183	Genomics applications to biotech traits: a revolution in progress?. 2004 , 7, 226-30	38
2182	Genomics tools for QTL analysis and gene discovery. 2004 , 7, 132-6	103
2181	Network building: transcriptional circuits in the root. 2004 , 7, 582-8	33
2180	A global view of defense gene expression regulation--a highly interconnected signaling network. 2004 , 7, 506-11	120
2179	RNA silencing in plants: a shortcut to functional analysis. 2004 , 72, 65-73	25

2178	Plant development: slowing root growth naturally. 2004 , 14, R395-6	5
2177	MicroRNA regulation of NAC-domain targets is required for proper formation and separation of adjacent embryonic, vegetative, and floral organs. 2004 , 14, 1035-46	54 ^o
2176	Arabidopsis GNARLED encodes a NAP125 homolog that positively regulates ARP2/3. 2004 , 14, 1405-9	82
2175	Helical growth of the Arabidopsis mutant tortifolia1 reveals a plant-specific microtubule-associated protein. 2004 , 14, 1515-21	104
2174	BYPASS1 negatively regulates a root-derived signal that controls plant architecture. 2004 , 14, 1739-46	80
2173	Phosphate transport in Arabidopsis: Pht1;1 and Pht1;4 play a major role in phosphate acquisition from both low- and high-phosphate environments. 2004 , 39, 629-42	55 ^o
2172	Structure of the hypothetical protein At3g17210 from Arabidopsis thaliana. 2004 , 28, 397-400	8
2171	Rice mutant resources for gene discovery. 2004 , 54, 325-34	185
2170	Transcripts of MYB-like genes respond to phosphorous and nitrogen deprivation in Arabidopsis. 2004 , 219, 1003-9	46
2169	Structural and functional analysis of rice genome. 2004 , 83, 79-99	46
2168	Silencing of the pollen-specific gene NTP303 and its family members in tobacco affects in vivo pollen tube growth and results in male sterile plants. 2004 , 55, 715-26	21
2167	Root cap specific expression of an endo-beta-1,4-D-glucanase (cellulase): a new marker to study root development in Arabidopsis. 2004 , 56, 309-23	49
2166	Functional and regulatory analysis of the Arabidopsis thaliana CAX2 cation transporter. 2004 , 56, 959-71	76
2165	Scaffolds, levers, rods and springs: diverse cellular functions of long coiled-coil proteins. 2004 , 61, 1996-2009	99
2164	Isochore structures in the genome of the plant Arabidopsis thaliana. 2004 , 59, 227-38	21
2163	RNAi for plant functional genomics. 2004 , 5, 240-4	42
2162	Transgene integration in plants: poking or patching holes in promiscuous genomes?. 2004 , 15, 126-31	59
2161	The plant-specific database. Classification of Arabidopsis proteins based on their phylogenetic profile. 2004 , 135, 1888-92	15

2160	Conserved MYC transcription factors play a key role in jasmonate signaling both in tomato and Arabidopsis. 2004 , 18, 1577-91		448
2159	The essential basic helix-loop-helix protein FIT1 is required for the iron deficiency response. 2004 , 16, 3400-12		565
2158	JASMONATE-INSENSITIVE1 encodes a MYC transcription factor essential to discriminate between different jasmonate-regulated defense responses in Arabidopsis. 2004 , 16, 1938-50		925
2157	PAF1-complex-mediated histone methylation of FLOWERING LOCUS C chromatin is required for the vernalization-responsive, winter-annual habit in Arabidopsis. 2004 , 18, 2774-84		228
2156	Stomatal development and pattern controlled by a MAPKK kinase. <i>Science</i> , 2004 , 304, 1494-7	33:3	417
2155	Della proteins and gibberellin-regulated seed germination and floral development in Arabidopsis. 2004 , 135, 1008-19		431
2154	The role of [Delta]1-pyrroline-5-carboxylate dehydrogenase in proline degradation. 2004 , 16, 3413-25		176
2153	AMPDB: the Arabidopsis Mitochondrial Protein Database. 2005 , 33, D605-10		50
2152	Genetic models in pathogenesis. 2004 , 38, 347-63		49
2151	Plant genomics: the third wave. 2004 , 5, 443-77		59
2150	The ethylene signaling pathway. <i>Science</i> , 2004 , 306, 1513-5	33:3	168
2149	Antagonistic interaction between abscisic acid and jasmonate-ethylene signaling pathways modulates defense gene expression and disease resistance in Arabidopsis. 2004 , 16, 3460-79		849
2148	The BDGP gene disruption project: single transposon insertions associated with 40% of Drosophila genes. 2004 , 167, 761-81		668
2147	Flagellin is not a major defense elicitor in <i>Ralstonia solanacearum</i> cells or extracts applied to Arabidopsis thaliana. 2004 , 17, 696-706		90
2146	Convergence of signaling pathways in the control of differential cell growth in Arabidopsis. 2004 , 7, 193-204		253
2145	The plant exon finder: a tool for precise detection of exons using a T-DNA-based tagging approach. 2004 , 338, 267-73		2
2144	Translation termination in Arabidopsis thaliana: characterisation of three versions of release factor 1. 2004 , 341, 219-25		15
2143	Identification and functional analysis of Arabidopsis proteins that interact with resistance gene product RPS2 in yeast. 2004 , 65, 257-267		7

2142	Endogenous trans-acting siRNAs regulate the accumulation of Arabidopsis mRNAs. 2004 , 16, 69-79	671
2141	Toward a systems approach to understanding plant cell walls. <i>Science</i> , 2004 , 306, 2206-11	333 931
2140	Generation of Arabidopsis thaliana plants with complex N-glycans lacking beta1,2-linked xylose and core alpha1,3-linked fucose. 2004 , 561, 132-6	246
2139	TILLING. Traditional mutagenesis meets functional genomics. 2004 , 135, 630-6	286
2138	The Ins and Outs of Nondestructive Cell-to-Cell and Systemic Movement of Plant Viruses. 2004 , 23, 195-250	177
2137	TELOMERASE ACTIVATOR1 induces telomerase activity and potentiates responses to auxin in Arabidopsis. 2004 , 16, 2910-22	41
2136	Control of male gametophyte development. 2004 , 16 Suppl, S142-53	427
2135	Target-selected gene inactivation in zebrafish. 2004 , 77, 69-90	42
2134	The recessive epigenetic swellmap mutation affects the expression of two step II splicing factors required for the transcription of the cell proliferation gene STRUWWELPETER and for the timing of cell cycle arrest in the Arabidopsis leaf. 2005 , 17, 1994-2008	51
2133	Integration of Genomic and Phenotypic Data. 2005 , 83-97	1
2132	T-DNA insertional mutagenesis in Arabidopsis: a tool for functional genomics. 2005 , 8,	12
2131	Applications of Genomics in Plant Biology. 59-80	1
2130	Genetic analysis of developmentally regulated resistance to downy mildew (<i>Hyaloperonospora parasitica</i>) in Arabidopsis thaliana. 2005 , 18, 1226-34	44
2129	Ustilago maydis Infection of the Nonnatural Host Arabidopsis thaliana. 2005 , 95, 480-8	22
2128	The Arabidopsis inositol 1,3,4-trisphosphate 5/6 kinase, AtItpk-1, is involved in plant photomorphogenesis under red light conditions, possibly via interaction with COP9 signalosome. 2005 , 43, 947-54	16
2127	AtHsfA2 modulates expression of stress responsive genes and enhances tolerance to heat and oxidative stress in Arabidopsis. 2005 , 48, 540-50	77
2126	Hydrogen peroxide-induced cell death in Arabidopsis: transcriptional and mutant analysis reveals a role of an oxoglutarate-dependent dioxygenase gene in the cell death process. 2005 , 57, 181-8	88
2125	AtEnSEMBL. 2007 , 406, 213-27	4

2124	Arabidopsis ETO1 specifically interacts with and negatively regulates type 2 1-aminocyclopropane-1-carboxylate synthases. 2005 , 5, 14	125
2123	Chlorophyll breakdown in senescent Arabidopsis leaves. Characterization of chlorophyll catabolites and of chlorophyll catabolic enzymes involved in the degreening reaction. 2005 , 139, 52-63	233
2122	What is base excision repair good for?: knockout mutants for FPG and OGG glycosylase genes in Arabidopsis. 2005 , 123, 227-232	12
2121	Rapid generation of plant traits via regulation of DNA mismatch repair. 2005 , 3, 399-407	12
2120	Novel gene functions required for melanization of the human pathogen <i>Cryptococcus neoformans</i> . 2005 , 57, 1381-96	173
2119	A novel isoform of glucan, water dikinase phosphorylates pre-phosphorylated alpha-glucans and is involved in starch degradation in Arabidopsis. 2005 , 41, 595-605	161
2118	An alternative tandem affinity purification strategy applied to Arabidopsis protein complex isolation. 2005 , 41, 767-78	200
2117	Mutations in Arabidopsis acyl-CoA oxidase genes reveal distinct and overlapping roles in beta-oxidation. 2005 , 41, 859-74	88
2116	A role of Toc33 in the protochlorophyllide-dependent plastid import pathway of NADPH:protochlorophyllide oxidoreductase (POR) A. 2005 , 42, 1-12	33
2115	BAS1 and SOB7 act redundantly to modulate Arabidopsis photomorphogenesis via unique brassinosteroid inactivation mechanisms. 2005 , 42, 23-34	144
2114	Functional genomics by integrated analysis of metabolome and transcriptome of Arabidopsis plants over-expressing an MYB transcription factor. 2005 , 42, 218-35	747
2113	Functional importance of the family 1 glucosyltransferase UGT72B1 in the metabolism of xenobiotics in Arabidopsis thaliana. 2005 , 42, 556-66	56
2112	Two sesquiterpene synthases are responsible for the complex mixture of sesquiterpenes emitted from Arabidopsis flowers. 2005 , 42, 757-71	262
2111	The plant Mo-hydroxylases aldehyde oxidase and xanthine dehydrogenase have distinct reactive oxygen species signatures and are induced by drought and abscisic acid. 2005 , 42, 862-76	129
2110	An important role of phosphatidic acid in ABA signaling during germination in Arabidopsis thaliana. 2005 , 43, 107-17	124
2109	NPH4/ARF7 and ARF19 promote leaf expansion and auxin-induced lateral root formation. 2005 , 43, 118-30	312
2108	The Botany Array Resource: e-Northerns, Expression Angling, and promoter analyses. 2005 , 43, 153-63	587
2107	Ex vitro composite plants: an inexpensive, rapid method for root biology. 2005 , 43, 449-57	166

2106	UPF3 suppresses aberrant spliced mRNA in Arabidopsis. 2005 , 43, 530-40	117
2105	The transcription factor ATAF2 represses the expression of pathogenesis-related genes in Arabidopsis. 2005 , 43, 745-57	234
2104	Sucrose rescues seedling establishment but not germination of Arabidopsis mutants disrupted in peroxisomal fatty acid catabolism. 2005 , 43, 861-72	138
2103	Secondary metabolites influence Arabidopsis/Botrytis interactions: variation in host production and pathogen sensitivity. 2005 , 44, 25-36	225
2102	Steady-state transposon mutagenesis in inbred maize. 2005 , 44, 52-61	184
2101	Genetic analysis of two Arabidopsis DNA polymerase epsilon subunits during early embryogenesis. 2005 , 44, 223-36	55
2100	Regulation of plant defense responses in Arabidopsis by EDR2, a PH and START domain-containing protein. 2005 , 44, 245-57	79
2099	RNA interference of Arabidopsis beta-amylase8 prevents maltose accumulation upon cold shock and increases sensitivity of PSII photochemical efficiency to freezing stress. 2005 , 44, 730-43	176
2098	Requirement of aminoacyl-tRNA synthetases for gametogenesis and embryo development in Arabidopsis. 2005 , 44, 866-78	100
2097	Identification of a novel protein, CRR7, required for the stabilization of the chloroplast NAD(P)H dehydrogenase complex in Arabidopsis. 2005 , 44, 1036-44	42
2096	Regulating transgenic crops sensibly: lessons from plant breeding, biotechnology and genomics. 2005 , 23, 439-44	200
2095	Prevention of early flowering by expression of FLOWERING LOCUS C requires methylation of histone H3 K36. 2005 , 7, 1256-60	226
2094	Developmental specificity of auxin response by pairs of ARF and Aux/IAA transcriptional regulators. 2005 , 24, 1874-85	303
2093	An Arabidopsis aspartic protease functions as an anti-cell-death component in reproduction and embryogenesis. 2005 , 6, 282-8	110
2092	Cell adhesion in Arabidopsis thaliana is mediated by ECTOPICALLY PARTING CELLS 1--a glycosyltransferase (GT64) related to the animal exostosins. 2005 , 43, 384-97	35
2091	Genetic analysis of Arabidopsis GATA transcription factor gene family reveals a nitrate-inducible member important for chlorophyll synthesis and glucose sensitivity. 2005 , 44, 680-92	144
2090	Ethylene signalling and response pathway: a unique signalling cascade with a multitude of inputs and outputs. 2005 , 123, 195-206	67
2089	The vacuolar Ca ²⁺ -activated channel TPC1 regulates germination and stomatal movement. 2005 , 434, 404-8	430

2088	The map-based sequence of the rice genome. 2005 , 436, 793-800	2923
2087	Ca ²⁺ /calmodulin is critical for brassinosteroid biosynthesis and plant growth. 2005 , 437, 741-5	115
2086	SERRATE coordinates shoot meristem function and leaf axial patterning in Arabidopsis. 2005 , 437, 1022-6	183
2085	Two types of replication protein A in seed plants. 2005 , 272, 3270-81	34
2084	Integrating genetic information into plant breeding programmes: how will we produce varieties from molecular variation, using bioinformatics?. 2005 , 146, 223-237	17
2083	Arabidopsis SOI33/AtENT8 Gene Encodes a Putative Equilibrative Nucleoside Transporter That Is Involved in Cytokinin Transport In Planta. 2005 , 47, 588-603	63
2082	Two meiotic crossover classes cohabit in Arabidopsis: one is dependent on MER3, whereas the other one is not. 2005 , 15, 692-701	149
2081	Partially redundant functions of Arabidopsis DICER-like enzymes and a role for DCL4 in producing trans-acting siRNAs. 2005 , 15, 1494-500	480
2080	MOS2, a protein containing G-patch and KOW motifs, is essential for innate immunity in Arabidopsis thaliana. 2005 , 15, 1936-42	73
2079	Transcriptional analysis and functional genomics in wheat. 2005 , 41, 149-163	23
2078	TILLING without a plough: a new method with applications for reverse genetics. 2005 , 8, 211-5	101
2077	The Arabidopsis Mei2 homologue AML1 binds AtRaptor1B, the plant homologue of a major regulator of eukaryotic cell growth. 2005 , 5, 2	42
2076	Callose (beta-1,3 glucan) is essential for Arabidopsis pollen wall patterning, but not tube growth. 2005 , 5, 22	158
2075	Distributed robustness versus redundancy as causes of mutational robustness. 2005 , 27, 176-88	216
2074	Genes, enzymes and regulation of arginine biosynthesis in plants. 2005 , 43, 729-45	190
2073	The inositol oxygenase gene family of Arabidopsis is involved in the biosynthesis of nucleotide sugar precursors for cell-wall matrix polysaccharides. 2005 , 221, 243-54	103
2072	Arabidopsis emb175 and other ppr knockout mutants reveal essential roles for pentatricopeptide repeat (PPR) proteins in plant embryogenesis. 2005 , 221, 424-36	117
2071	SKS6, a multicopper oxidase-like gene, participates in cotyledon vascular patterning during Arabidopsis thaliana development. 2005 , 222, 652-66	36

2070	Isogene specific oligo arrays reveal multifaceted changes in gene expression during grape berry (<i>Vitis vinifera</i> L.) development. 2005 , 222, 832-47	213
2069	Disruption of <i>RCI2A</i> leads to over-accumulation of Na ⁺ and increased salt sensitivity in <i>Arabidopsis thaliana</i> plants. 2005 , 222, 1001-9	47
2068	Physiological characterisation of <i>Arabidopsis</i> mutants affected in the expression of the putative regulatory protein PII. 2005 , 223, 28-39	52
2067	Genomics of plant cell wall biogenesis. 2005 , 221, 747-51	83
2066	Four distinct classes of proteins as interaction partners of the PABC domain of <i>Arabidopsis thaliana</i> Poly(A)-binding proteins. 2005 , 272, 651-65	32
2065	Frequency and character of alternative somatic recombination fates of paralogous genes during T-DNA integration. 2005 , 274, 91-102	2
2064	<i>Agrobacterium</i> T-DNA integration in <i>Arabidopsis</i> is correlated with DNA sequence compositions that occur frequently in gene promoter regions. 2005 , 5, 240-53	39
2063	Structural and functional characterization of <i>AtPTR3</i> , a stress-induced peptide transporter of <i>Arabidopsis</i> . 2005 , 11, 226-36	32
2062	A double mutant for the <i>CYP85A1</i> and <i>CYP85A2</i> Genes of <i>Arabidopsis</i> exhibits a Brassinosteroid dwarf phenotype. 2005 , 48, 237-244	37
2061	Reciprocal regulation of <i>Arabidopsis</i> <i>UGT78D2</i> and <i>BANYULS</i> is critical for regulation of the metabolic flux of anthocyanidins to condensed tannins in developing seed coats. 2005 , 48, 356-370	29
2060	<i>Agrobacterium</i> -Mediated Transformation of Common Bermudagrass (<i>Cynodon dactylon</i>). 2005 , 83, 223-229	22
2059	Enhancing gene targeting efficiency in higher plants: rice is on the move. 2005 , 14, 1-14	18
2058	Insertional mutagenesis in <i>Populus</i> : relevance and feasibility. 2005 , 1, 135-142	17
2057	Reverse genetic approaches for functional genomics of rice. 2005 , 59, 111-23	92
2056	<i>AINTEGUMENTA</i> -like (<i>AIL</i>) genes are expressed in young tissues and may specify meristematic or division-competent states. 2005 , 57, 613-28	144
2055	An annotation update via cDNA sequence analysis and comprehensive profiling of developmental, hormonal or environmental responsiveness of the <i>Arabidopsis</i> <i>AP2/EREBP</i> transcription factor gene family. 2005 , 59, 853-68	140
2054	Two callose synthases, <i>GSL1</i> and <i>GSL5</i> , play an essential and redundant role in plant and pollen development and in fertility. 2005 , 58, 333-49	139
2053	Expression of <i>MADS</i> -box genes during the embryonic phase in <i>Arabidopsis</i> . 2005 , 58, 89-107	51

2052	Genome-wide expression profiling and identification of gene activities during early flower development in Arabidopsis. 2005 , 58, 401-19	63
2051	NADK2, an Arabidopsis chloroplastic NAD kinase, plays a vital role in both chlorophyll synthesis and chloroplast protection. 2005 , 59, 553-64	103
2050	Genome-wide introgression lines and their use in genetic and molecular dissection of complex phenotypes in rice (<i>Oryza sativa</i> L.). 2005 , 59, 33-52	123
2049	Redundant and distinct functions of the ABA response loci ABA-INSENSITIVE(ABI)5 and ABRE-BINDING FACTOR (ABF)3. 2005 , 59, 253-67	154
2048	The Arabidopsis AtRaptor genes are essential for post-embryonic plant growth. 2005 , 3, 12	117
2047	Proteome mapping of mature pollen of Arabidopsis thaliana. 2005 , 5, 4864-84	207
2046	Genetic Programming of Crops to Sustain or Increase Yields under Reduced Irrigation. 2005 , 1	
2045	. 2005 ,	13
2044	Inferring hypotheses on functional relationships of genes: Analysis of the Arabidopsis thaliana subtilase gene family. 2005 , 1, e40	122
2043	Complete maturation of the plastid protein translocation channel requires a type I signal peptidase. 2005 , 171, 425-30	87
2042	PGP4, an ATP binding cassette P-glycoprotein, catalyzes auxin transport in Arabidopsis thaliana roots. 2005 , 17, 2922-39	288
2041	Mutations in the Arabidopsis phosphoinositide phosphatase gene SAC9 lead to overaccumulation of PtdIns(4,5)P2 and constitutive expression of the stress-response pathway. 2005 , 138, 686-700	124
2040	AUXIN RESPONSE FACTOR1 and AUXIN RESPONSE FACTOR2 regulate senescence and floral organ abscission in Arabidopsis thaliana. 2005 , 132, 4563-74	396
2039	Ethylene signaling pathway. 2005 , 2005, cm3	11
2038	The Arabidopsis plastidic glucose 6-phosphate/phosphate translocator GPT1 is essential for pollen maturation and embryo sac development. 2005 , 17, 760-75	163
2037	SEC8, a subunit of the putative Arabidopsis exocyst complex, facilitates pollen germination and competitive pollen tube growth. 2005 , 138, 2005-18	149
2036	Dissection of Arabidopsis ADP-RIBOSYLATION FACTOR 1 function in epidermal cell polarity. 2005 , 17, 525-36	347
2035	The RPN1 subunit of the 26S proteasome in Arabidopsis is essential for embryogenesis. 2005 , 17, 2723-37	62

2034	Psb29, a conserved 22-kD protein, functions in the biogenesis of Photosystem II complexes in <i>Synechocystis</i> and <i>Arabidopsis</i> . 2005 , 17, 2768-81	88
2033	AtCCMH, an essential component of the c-type cytochrome maturation pathway in <i>Arabidopsis</i> mitochondria, interacts with apocytochrome c. 2005 , 102, 16113-8	77
2032	The <i>Arabidopsis</i> genome: a foundation for plant research. 2005 , 15, 1632-42	89
2031	Two homologous ATP-binding cassette transporter proteins, AtMDR1 and AtPGP1, regulate <i>Arabidopsis</i> photomorphogenesis and root development by mediating polar auxin transport. 2005 , 138, 949-64	137
2030	A resource of 5,814 dissociation transposon-tagged and sequence-indexed lines of <i>Arabidopsis</i> transposed from start loci on chromosome 5. 2005 , 46, 1149-53	52
2029	The involvement of gibberellin 20-oxidase genes in phytochrome-regulated petiole elongation of <i>Arabidopsis</i> . 2005 , 138, 1106-16	142
2028	Detoxification and transcriptome response in <i>Arabidopsis</i> seedlings exposed to the allelochemical benzoxazolin-2(3H)-one. 2005 , 280, 21867-81	139
2027	A Sec14p-nodulin domain phosphatidylinositol transfer protein polarizes membrane growth of <i>Arabidopsis thaliana</i> root hairs. 2005 , 168, 801-12	182
2026	Class III homeodomain-leucine zipper gene family members have overlapping, antagonistic, and distinct roles in <i>Arabidopsis</i> development. 2005 , 17, 61-76	530
2025	The BLADE ON PETIOLE genes act redundantly to control the growth and development of lateral organs. 2005 , 132, 2203-13	160
2024	Functional association of <i>Arabidopsis</i> CAX1 and CAX3 is required for normal growth and ion homeostasis. 2005 , 138, 2048-60	157
2023	The critical role of <i>Arabidopsis</i> electron-transfer flavoprotein:ubiquinone oxidoreductase during dark-induced starvation. 2005 , 17, 2587-600	191
2022	Functional genomic analysis of the AUXIN/INDOLE-3-ACETIC ACID gene family members in <i>Arabidopsis thaliana</i> . 2005 , 17, 3282-300	265
2021	MicroRNA-directed regulation of <i>Arabidopsis</i> AUXIN RESPONSE FACTOR17 is essential for proper development and modulates expression of early auxin response genes. 2005 , 17, 1360-75	682
2020	Functional genomics of eukaryotic photosynthesis using insertional mutagenesis of <i>Chlamydomonas reinhardtii</i> . 2005 , 137, 545-56	162
2019	BLADE-ON-PETIOLE-dependent signaling controls leaf and floral patterning in <i>Arabidopsis</i> . 2005 , 17, 1434-48	216
2018	Pollen development and fertilization in <i>Arabidopsis</i> is dependent on the MALE GAMETOGENESIS IMPAIRED ANTHERS gene encoding a type V P-type ATPase. 2005 , 19, 2757-69	77
2017	An inducible targeted tagging system for localized saturation mutagenesis in <i>Arabidopsis</i> . 2005 , 137, 3-12	28

2016	TWIN SISTER OF FT (TSF) acts as a floral pathway integrator redundantly with FT. 2005 , 46, 1175-89	338	
2015	Loss of the CONSTITUTIVE PHOTOMORPHOGENIC9 signalosome subunit 5 is sufficient to cause the cop/det/fus mutant phenotype in Arabidopsis. 2005 , 17, 1967-78	79	
2014	Localized hormone fluxes and early haustorium development in the hemiparasitic plant <i>Triphysaria versicolor</i> . 2005 , 138, 1469-80	49	
2013	Identification and functional characterization of Arabidopsis PEROXIN4 and the interacting protein PEROXIN22. 2005 , 17, 3422-35	100	
2012	Dissecting the ethylene pathway of Arabidopsis. 2005 , 3, 372-81	18	
2011	The Arabidopsis peroxisomal targeting signal type 2 receptor PEX7 is necessary for peroxisome function and dependent on PEX5. 2005 , 16, 573-83	121	
2010	SWI3 subunits of putative SWI/SNF chromatin-remodeling complexes play distinct roles during Arabidopsis development. 2005 , 17, 2454-72	100	
2009	BASIC PENTACYSSTEINE1, a GA binding protein that induces conformational changes in the regulatory region of the homeotic Arabidopsis gene SEEDSTICK. 2005 , 17, 722-9	86	
2008	Arabidopsis peroxisomal citrate synthase is required for fatty acid respiration and seed germination. 2005 , 17, 2037-48	157	
2007	Reinforcement of silencing at transposons and highly repeated sequences requires the concerted action of two distinct RNA polymerases IV in Arabidopsis. 2005 , 19, 2030-40	308	
2006	A basic helix-loop-helix transcription factor in Arabidopsis, MYC2, acts as a repressor of blue light-mediated photomorphogenic growth. 2005 , 17, 1953-66	202	
2005	RIN13 is a positive regulator of the plant disease resistance protein RPM1. 2005 , 17, 1016-28	22	
2004	A central role of Arabidopsis thaliana ovate family proteins in networking and subcellular localization of 3-aa loop extension homeodomain proteins. 2005 , 102, 4908-12	195	
2003	Identification of genes required for cellulose synthesis by regression analysis of public microarray data sets. 2005 , 102, 8633-8	479	
2002	AtREC8 and AtSCC3 are essential to the monopolar orientation of the kinetochores during meiosis. 2005 , 118, 4621-32	197	
2001	Auxin response factors ARF6 and ARF8 promote jasmonic acid production and flower maturation. 2005 , 132, 4107-18	464	
2000	Positional signaling mediated by a receptor-like kinase in Arabidopsis. <i>Science</i> , 2005 , 307, 1111-3	33.3	176
1999	Multiple type-B response regulators mediate cytokinin signal transduction in Arabidopsis. 2005 , 17, 3007-18	315	

1998	The use of microarrays to study the anaerobic response in Arabidopsis. 2005 , 96, 661-8	51
1997	New perspectives in flooding research: the use of shade avoidance and Arabidopsis thaliana. 2005 , 96, 533-40	44
1996	Auxin and light control of adventitious rooting in Arabidopsis require ARGONAUTE1. 2005 , 17, 1343-59	278
1995	A plasma membrane H ⁺ -ATPase is required for the formation of proanthocyanidins in the seed coat endothelium of Arabidopsis thaliana. 2005 , 102, 2649-54	169
1994	Uncoupling of the functions of the Arabidopsis VIP1 protein in transient and stable plant genetic transformation by Agrobacterium. 2005 , 102, 5733-8	122
1993	Site preferences of insertional mutagenesis agents in Arabidopsis. 2005 , 137, 168-75	45
1992	Arabidopsis RAD51C gene is important for homologous recombination in meiosis and mitosis. 2005 , 139, 896-908	83
1991	DISTORTED3/SCAR2 is a putative arabidopsis WAVE complex subunit that activates the Arp2/3 complex and is required for epidermal morphogenesis. 2005 , 17, 502-24	110
1990	Roles of Arabidopsis AtREV1 and AtREV7 in translesion synthesis. 2005 , 138, 870-81	52
1989	Folate metabolism in plants: an Arabidopsis homolog of the mammalian mitochondrial folate transporter mediates folate import into chloroplasts. 2005 , 280, 34823-31	96
1988	The membrane-anchored BOTRYTIS-INDUCED KINASE1 plays distinct roles in Arabidopsis resistance to necrotrophic and biotrophic pathogens. 2006 , 18, 257-73	315
1987	FRIGIDA-ESSENTIAL 1 interacts genetically with FRIGIDA and FRIGIDA-LIKE 1 to promote the winter-annual habit of Arabidopsis thaliana. 2005 , 132, 5471-8	74
1986	Molecular genetics using T-DNA in rice. 2005 , 46, 14-22	107
1985	A putative nucleoporin 96 is required for both basal defense and constitutive resistance responses mediated by suppressor of npr1-1, constitutive 1. 2005 , 17, 1306-16	185
1984	The plant VirE2 interacting protein 1. a molecular link between the Agrobacterium T-complex and the host cell chromatin?. 2005 , 138, 1318-21	62
1983	The Arabidopsis PEX12 gene is required for peroxisome biogenesis and is essential for development. 2005 , 139, 231-9	92
1982	A molecular-genetic study of the Arabidopsis Toc75 gene family. 2005 , 138, 715-33	98
1981	A plant-specific protein essential for blue-light-induced chloroplast movements. 2005 , 139, 101-14	83

1980	Identification of three urease accessory proteins that are required for urease activation in <i>Arabidopsis</i> . 2005 , 139, 1155-62	55
1979	Genetic and chemical analyses of the action mechanisms of sirtinol in <i>Arabidopsis</i> . 2005 , 102, 3129-34	74
1978	The evolutionarily conserved TOUGH protein is required for proper development of <i>Arabidopsis thaliana</i> . 2005 , 17, 2473-85	22
1977	Auxin response factors mediate <i>Arabidopsis</i> organ asymmetry via modulation of KANADI activity. 2005 , 17, 2899-910	373
1976	Maize rough sheath2 and its <i>Arabidopsis</i> orthologue ASYMMETRIC LEAVES1 interact with HIRA, a predicted histone chaperone, to maintain <i>knox</i> gene silencing and determinacy during organogenesis. 2005 , 17, 2886-98	167
1975	IRREGULAR TRICHOME BRANCH1 in <i>Arabidopsis</i> encodes a plant homolog of the actin-related protein2/3 complex activator Scar/WAVE that regulates actin and microtubule organization. 2005 , 17, 2314-26	83
1974	Control of root cap formation by MicroRNA-targeted auxin response factors in <i>Arabidopsis</i> . 2005 , 17, 2204-16	619
1973	Expression of <i>Arabidopsis</i> MIRNA genes. 2005 , 138, 2145-54	523
1972	Functional genomic analysis of the AUXIN RESPONSE FACTOR gene family members in <i>Arabidopsis thaliana</i> : unique and overlapping functions of ARF7 and ARF19. 2005 , 17, 444-63	699
1971	HOS10 encodes an R2R3-type MYB transcription factor essential for cold acclimation in plants. 2005 , 102, 9966-71	156
1970	Combining experimental and predicted datasets for determination of the subcellular location of proteins in <i>Arabidopsis</i> . 2005 , 139, 598-609	116
1969	Analysis of mutations induced by carbon ions in <i>Arabidopsis thaliana</i> . 2005 , 56, 587-96	123
1968	PIRLs: a novel class of plant intracellular leucine-rich repeat proteins. 2005 , 46, 913-22	36
1967	Cellular sterol ester synthesis in plants is performed by an enzyme (phospholipid:sterol acyltransferase) different from the yeast and mammalian acyl-CoA:sterol acyltransferases. 2005 , 280, 34626-34	43
1966	Functional and developmental impact of cytosolic protein N-terminal methionine excision in <i>Arabidopsis</i> . 2005 , 137, 623-37	60
1965	An indole-3-acetic acid carboxyl methyltransferase regulates <i>Arabidopsis</i> leaf development. 2005 , 17, 2693-704	218
1964	MYB98 is required for pollen tube guidance and synergid cell differentiation in <i>Arabidopsis</i> . 2005 , 17, 2981-92	211
1963	Looking for invisible phenotypes in cell wall mutants of <i>Arabidopsis thaliana</i> . 2005 , 139, 80-83	12

1962	Organ-specific expression of Arabidopsis genome during development. 2005 , 138, 80-91	146
1961	The Arabidopsis R2R3 MYB proteins FOUR LIPS and MYB88 restrict divisions late in the stomatal cell lineage. 2005 , 17, 2754-67	157
1960	STN8 protein kinase in Arabidopsis thaliana is specific in phosphorylation of photosystem II core proteins. 2005 , 280, 33679-86	173
1959	Higher plant plastids and cyanobacteria have folate carriers related to those of trypanosomatids. 2005 , 280, 38457-63	76
1958	ATM and ATR make distinct contributions to chromosome end protection and the maintenance of telomeric DNA in Arabidopsis. 2005 , 19, 2111-5	45
1957	Tripeptidyl peptidase II. An oligomeric protease complex from Arabidopsis. 2005 , 138, 1046-57	51
1956	Mutants of Arabidopsis lacking a chloroplastic isoamylase accumulate phyto glycogen and an abnormal form of amylopectin. 2005 , 138, 184-95	145
1955	Loss-of-function mutations in chitin responsive genes show increased susceptibility to the powdery mildew pathogen Erysiphe cichoracearum. 2005 , 138, 1027-36	171
1954	Protein geranylgeranyltransferase I is involved in specific aspects of abscisic acid and auxin signaling in Arabidopsis. 2005 , 139, 722-33	59
1953	Pathogen-responsive expression of glycosyltransferase genes UGT73B3 and UGT73B5 is necessary for resistance to Pseudomonas syringae pv tomato in Arabidopsis. 2005 , 139, 1890-901	125
1952	Repressor- and activator-type ethylene response factors functioning in jasmonate signaling and disease resistance identified via a genome-wide screen of Arabidopsis transcription factor gene expression. 2005 , 139, 949-59	448
1951	Arabidopsis has two redundant Cullin3 proteins that are essential for embryo development and that interact with RBX1 and BTB proteins to form multisubunit E3 ubiquitin ligase complexes in vivo. 2005 , 17, 1180-95	124
1950	D-GLYCERATE 3-KINASE, the last unknown enzyme in the photorespiratory cycle in Arabidopsis, belongs to a novel kinase family. 2005 , 17, 2413-20	110
1949	The Arabidopsis thaliana SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASES1 and 2 control male sporogenesis. 2005 , 17, 3337-49	252
1948	Peroxisomal localization of a myosin XI isoform in Arabidopsis thaliana. 2005 , 46, 782-9	82
1947	The TIP GROWTH DEFECTIVE1 S-acyl transferase regulates plant cell growth in Arabidopsis. 2005 , 17, 2554-63	98
1946	Characterisation of a T-DNA-tagged gene of Arabidopsis thaliana that regulates gibberellin metabolism and flowering time. 2005 , 32, 923-932	3
1945	RNR1, a 3'-5' exoribonuclease belonging to the RNR superfamily, catalyzes 3' maturation of chloroplast ribosomal RNAs in Arabidopsis thaliana. 2005 , 33, 2751-63	97

1944	The bromodomain protein GTE6 controls leaf development in Arabidopsis by histone acetylation at ASYMMETRIC LEAVES1. 2005 , 19, 2245-54	41
1943	Integration of flowering signals in winter-annual Arabidopsis. 2005 , 137, 149-56	237
1942	Arabidopsis tic110 is essential for the assembly and function of the protein import machinery of plastids. 2005 , 17, 1482-96	109
1941	Interactions between the cell cycle and embryonic patterning in Arabidopsis uncovered by a mutation in DNA polymerase epsilon. 2005 , 17, 3362-77	69
1940	The elongata mutants identify a functional Elongator complex in plants with a role in cell proliferation during organ growth. 2005 , 102, 7754-9	132
1939	Photosystem II core phosphorylation and photosynthetic acclimation require two different protein kinases. 2005 , 437, 1179-82	376
1938	Arabidopsis cytokinin receptor mutants reveal functions in shoot growth, leaf senescence, seed size, germination, root development, and cytokinin metabolism. 2006 , 18, 40-54	696
1937	Induction of protein secretory pathway is required for systemic acquired resistance. <i>Science</i> , 2005 , 308, 1036-40	333 441
1936	A Link between ethylene and auxin uncovered by the characterization of two root-specific ethylene-insensitive mutants in Arabidopsis. 2005 , 17, 2230-42	385
1935	Transcriptional profile of the Arabidopsis root quiescent center. 2005 , 17, 1908-25	257
1934	New subunits NDH-M, -N, and -O, encoded by nuclear genes, are essential for plastid Ndh complex functioning in higher plants. 2005 , 17, 219-32	157
1933	Vitamin B6 biosynthesis in higher plants. 2005 , 102, 13687-92	168
1932	Dissecting the beta-aminobutyric acid-induced priming phenomenon in Arabidopsis. 2005 , 17, 987-99	295
1931	Purification of the COP9 signalosome from porcine spleen, human cell lines, and Arabidopsis thaliana plants. 2005 , 398, 468-81	9
1930	Characterization of an Arabidopsis enzyme family that conjugates amino acids to indole-3-acetic acid. 2005 , 17, 616-27	719
1929	A novel phosphatidylcholine-hydrolyzing phospholipase C induced by phosphate starvation in Arabidopsis. 2005 , 280, 7469-76	190
1928	The potassium transporter AtHAK5 functions in K(+) deprivation-induced high-affinity K(+) uptake and AKT1 K(+) channel contribution to K(+) uptake kinetics in Arabidopsis roots. 2005 , 137, 1105-14	363
1927	Identification of novel genes in Arabidopsis involved in secondary cell wall formation using expression profiling and reverse genetics. 2005 , 17, 2281-95	598

1926	D-type cyclins activate division in the root apex to promote seed germination in Arabidopsis. 2005 , 102, 15694-9	126
1925	Structure of the higher plant light harvesting complex I: in vivo characterization and structural interdependence of the Lhca proteins. 2005 , 44, 3065-73	62
1924	Targeted mutagenesis using zinc-finger nucleases in Arabidopsis. 2005 , 102, 2232-7	339
1923	Functional genomics of protein kinases in plants. 2005 , 3, 362-71	22
1922	Ethylene signal transduction. 2005 , 95, 901-15	317
1921	An Arabidopsis homolog of RAPTOR/KOG1 is essential for early embryo development. 2005 , 326, 844-50	125
1920	A reference map of the Arabidopsis thaliana mature pollen proteome. 2005 , 337, 1257-66	123
1919	A new class of transcription factors mediates brassinosteroid-regulated gene expression in Arabidopsis. 2005 , 120, 249-59	539
1918	microRNA-directed phasing during trans-acting siRNA biogenesis in plants. 2005 , 121, 207-21	1783
1917	AtNTRB is the major mitochondrial thioredoxin reductase in Arabidopsis thaliana. 2005 , 579, 337-42	96
1916	Selective involvement of members of the eukaryotic initiation factor 4E family in the infection of Arabidopsis thaliana by potyviruses. 2005 , 579, 1167-71	103
1915	MDR-like ABC transporter AtPGP4 is involved in auxin-mediated lateral root and root hair development. 2005 , 579, 5399-5406	172
1914	Purification and characterization of topoisomerase IIA from Arabidopsis thaliana. 2005 , 168, 1023-1033	7
1913	POL and related phosphatases are dosage-sensitive regulators of meristem and organ development in Arabidopsis. 2005 , 285, 272-84	72
1912	Tall tales from sly dwarves: novel functions of gibberellins in plant development. 2005 , 10, 123-9	140
1911	Insertional mutagenesis: a Swiss Army knife for functional genomics of Medicago truncatula. 2005 , 10, 229-35	103
1910	A rapid and versatile combined DNA/RNA extraction protocol and its application to the analysis of a novel DNA marker set polymorphic between Arabidopsis thaliana ecotypes Col-0 and Landsberg erecta. 2005 , 1, 4	53
1909	The Arabidopsis Genome and Its Use in Cereal Genomics. 2004 , 515-534	

1908	Circadian genetics in the model higher plant, <i>Arabidopsis thaliana</i> . 2005 , 393, 23-35	32
1907	The AIP2 E3 ligase acts as a novel negative regulator of ABA signaling by promoting ABI3 degradation. 2005 , 19, 1532-43	320
1906	<i>Arabidopsis</i> ethylene signaling pathway. 2005 , 2005, cm4	34
1905	Ethylene biosynthesis and signaling: an overview. 2005 , 72, 399-430	44
1904	Cereal Genomics. 2005 ,	3
1903	<i>Agrobacterium</i> Protocols. 2006 ,	2
1902	Mechanisms for nitric oxide synthesis in plants. 2006 , 57, 471-8	266
1901	TOPLESS regulates apical embryonic fate in <i>Arabidopsis</i> . <i>Science</i> , 2006 , 312, 1520-3	333 381
1900	<i>Arabidopsis</i> Protocols. 2006 ,	2
1899	Gene identification and cloning by molecular marker mapping. 2006 , 323, 115-26	18
1898	Will you let me use your nucleus? How <i>Agrobacterium</i> gets its T-DNA expressed in the host plant cell. 2006 , 84, 333-45	35
1897	A systems approach to understanding root development. 2006 , 84, 695-701	3
1896	KNAT6: an <i>Arabidopsis</i> homeobox gene involved in meristem activity and organ separation. 2006 , 18, 1900-7	138
1895	Plant Metabolomics. 2006 ,	29
1894	Inexpensive fine mapping and positional cloning in plants using visible, mapped transgenes. 2006 , 84, 179-188	1
1893	Transformation-induced mutations in transgenic plants: analysis and biosafety implications. 2006 , 23, 209-37	58
1892	Identification of primary target genes of phytochrome signaling. Early transcriptional control during shade avoidance responses in <i>Arabidopsis</i> . 2006 , 141, 85-96	107
1891	Blind homologous R2R3 Myb genes control the pattern of lateral meristem initiation in <i>Arabidopsis</i> . 2006 , 18, 586-97	189

1890	Plastid cues posttranscriptionally regulate the accumulation of key enzymes of the methylerythritol phosphate pathway in Arabidopsis. 2006 , 141, 75-84	72
1889	Maturation of arabidopsis seeds is dependent on glutathione biosynthesis within the embryo. 2006 , 141, 446-55	210
1888	A pathogen-inducible endogenous siRNA in plant immunity. 2006 , 103, 18002-7	400
1887	Organization and function of the actin cytoskeleton in developing root cells. 2006 , 252, 219-64	21
1886	Arabidopsis SAMT1 defines a plastid transporter regulating plastid biogenesis and plant development. 2006 , 18, 3088-105	85
1885	Cell biology of the Arabidopsis nuclear siRNA pathway for RNA-directed chromatin modification. 2006 , 71, 473-80	29
1884	Brassinosteroid induction of AtACS4 encoding an auxin-responsive 1-aminocyclopropane-1-carboxylate synthase 4 in Arabidopsis seedlings. 2006 , 126, 060217072449002-???	4
1883	The AtNFXL1 gene encodes a NF-X1 type zinc finger protein required for growth under salt stress. 2006 , 580, 4851-6	42
1882	Photodamaged D1 protein is degraded in Arabidopsis mutants lacking the Deg2 protease. 2006 , 580, 6929-32	46
1881	Arabidopsis thaliana floral dip transformation method. 2006 , 343, 87-103	122
1880	Forward genetic screening of insertional mutants. 2006 , 323, 147-61	2
1879	Molecular mechanisms of ethylene signaling in Arabidopsis. 2006 , 2, 165-73	55
1878	Pollen Development, a Genetic and Transcriptomic View. 15-45	21
1877	Screening and Analysis of Pollen Tube Mutations. 243-263	
1876	Arabidopsis mitochondria have two basic amino acid transporters with partially overlapping specificities and differential expression in seedling development. 2006 , 1757, 1277-83	43
1875	Methylation of tRNA ^{Asp} by the DNA methyltransferase homolog Dnmt2. <i>Science</i> , 2006 , 311, 395-8	33-3 797
1874	Genome-wide analysis of the ERF gene family in Arabidopsis and rice. 2006 , 140, 411-32	1395
1873	Arabidopsis TEBICHI, with helicase and DNA polymerase domains, is required for regulated cell division and differentiation in meristems. 2006 , 18, 879-92	81

1872	The accumulation of oleosins determines the size of seed oilbodies in Arabidopsis. 2006 , 18, 1961-74	320
1871	A gain-of-function mutation in the Arabidopsis pleiotropic drug resistance transporter PDR9 confers resistance to auxinic herbicides. 2006 , 142, 63-74	123
1870	CYP707A1 and CYP707A2, which encode abscisic acid 8'-hydroxylases, are indispensable for proper control of seed dormancy and germination in Arabidopsis. 2006 , 141, 97-107	375
1869	Expression of the Arabidopsis high-affinity hexose transporter STP13 correlates with programmed cell death. 2006 , 580, 2381-7	77
1868	Profound redox sensitivity of peptidyl-prolyl isomerase activity in Arabidopsis thylakoid lumen. 2006 , 580, 3671-6	42
1867	Arabidopsis RNA-binding protein UBA2a relocalizes into nuclear speckles in response to abscisic acid. 2006 , 580, 4160-5	38
1866	Elucidation of the beta-carotene hydroxylation pathway in Arabidopsis thaliana. 2006 , 580, 4718-22	58
1865	Functional analysis of a calcium-binding transcription factor involved in plant salt stress signaling. 2006 , 580, 5251-6	55
1864	Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts Agrobacterium-mediated transformation. 2006 , 125, 749-60	1321
1863	Regulatory functions of nuclear hexokinase1 complex in glucose signaling. 2006 , 127, 579-89	330
1862	Molecular evolution of the AP2 subfamily. 2006 , 366, 256-65	149
1861	The pleiotropic Arabidopsis frd mutation with altered coordination of chloroplast biogenesis, cell size and differentiation, organ size and number. 2006 , 382, 88-99	5
1860	Atmnd1-delta1 is sensitive to gamma-irradiation and defective in meiotic DNA repair. 2006 , 5, 455-64	35
1859	Systems biology of circadian rhythms: an outlook. 2006 , 21, 507-18	17
1858	Arabidopsis GLUTAMINE-RICH PROTEIN23 is essential for early embryogenesis and encodes a novel nuclear PPR motif protein that interacts with RNA polymerase II subunit III. 2006 , 18, 815-30	118
1857	MEKK1 is required for MPK4 activation and regulates tissue-specific and temperature-dependent cell death in Arabidopsis. 2006 , 281, 36969-76	224
1856	An Arabidopsis glutathione peroxidase functions as both a redox transducer and a scavenger in abscisic acid and drought stress responses. 2006 , 18, 2749-66	379
1855	FUNCTIONAL GENOMICS OF STRESS TOLERANCE. 2006 , 301-334	3

1854	Uncovering a hidden distributed architecture behind scale-free transcriptional regulatory networks. 2006 , 360, 204-12	61
1853	AGO1 homeostasis entails coexpression of MIR168 and AGO1 and preferential stabilization of miR168 by AGO1. 2006 , 22, 129-36	272
1852	PEPPER, a novel K-homology domain gene, regulates vegetative and gynoecium development in Arabidopsis. 2006 , 289, 346-59	36
1851	Pollen-specific pectin methylesterase involved in pollen tube growth. 2006 , 294, 83-91	156
1850	Analysis of T-DNA insertion site distribution patterns in Arabidopsis thaliana reveals special features of genes without insertions. 2006 , 87, 645-52	33
1849	Characterization of T-DNA integration in creeping bentgrass. 2006 , 170, 225-237	14
1848	Harvesting the potential of induced biological diversity. 2006 , 11, 71-9	43
1847	Cloned sequence repertoires for small- and large-scale biology. 2006 , 11, 133-41	30
1846	Licensed to kill: the lifestyle of a necrotrophic plant pathogen. 2006 , 11, 247-53	502
1845	Plant genetic and molecular responses to water deficit. 2006 , 1, 617	2
1844	CDPKs CPK6 and CPK3 function in ABA regulation of guard cell S-type anion- and Ca(2+)-permeable channels and stomatal closure. 2006 , 4, e327	446
1843	Thale Cress (Arabidopsis thaliana) Genome. 2006 ,	1
1842	AGROBACTERIUM-HOST INTERACTIONS: BIOLOGY AND BIOTECHNOLOGY. 2006 , 129-142	
1841	Model Host Systems: Tools for Comprehensive Analysis of Host-Pathogen Interactions. 2006 , 445-456	
1840	Functional annotation of the Arabidopsis P450 superfamily based on large-scale co-expression analysis. 2006 , 34, 1192-8	27
1839	Arabidopsis thaliana RGXT1 and RGXT2 encode Golgi-localized (1,3)-alpha-D-xylosyltransferases involved in the synthesis of pectic rhamnogalacturonan-II. 2006 , 18, 2593-607	106
1838	AtMND1 is required for homologous pairing during meiosis in Arabidopsis. 2006 , 7, 24	29
1837	Cytokinin affects circadian-clock oscillation in a phytochrome B- and Arabidopsis response regulator 4-dependent manner. 2006 , 127, 277-292	25

1836	The BIG gene is involved in regulation of sulfur deficiency-responsive genes in <i>Arabidopsis thaliana</i> . 2006 , 129, 351-363	18
1835	Genomic-assisted identification of genes involved in secondary growth in <i>Arabidopsis</i> utilising transcript profiling of poplar wood-forming tissues. 2006 , 129, 415-428	16
1834	Gene Discovery and Functional Analyses in the Model Plant <i>Arabidopsis</i> . 2006 , 48, 5-14	8
1833	An alpha-(1,4)-amylase is essential for alpha-(1,3)-glucan production and virulence in <i>Histoplasma capsulatum</i> . 2006 , 62, 970-83	96
1832	Over-expression of different aldehyde dehydrogenase genes in <i>Arabidopsis thaliana</i> confers tolerance to abiotic stress and protects plants against lipid peroxidation and oxidative stress. 2006 , 29, 1033-48	243
1831	The CLAVATA1-related BAM1, BAM2 and BAM3 receptor kinase-like proteins are required for meristem function in <i>Arabidopsis</i> . 2006 , 45, 1-16	287
1830	Analysis of the root-hair morphogenesis transcriptome reveals the molecular identity of six genes with roles in root-hair development in <i>Arabidopsis</i> . 2006 , 45, 83-100	163
1829	Generation of a flanking sequence-tag database for activation-tagging lines in japonica rice. 2006 , 45, 123-32	283
1828	Functional analysis of <i>Arabidopsis</i> NCED6 and NCED9 genes indicates that ABA synthesized in the endosperm is involved in the induction of seed dormancy. 2006 , 45, 309-19	323
1827	The WAG1 and WAG2 protein kinases negatively regulate root waving in <i>Arabidopsis</i> . 2006 , 45, 752-64	108
1826	<i>Arabidopsis</i> CDKA;1, a cdc2 homologue, controls proliferation of generative cells in male gametogenesis. 2006 , 45, 819-31	166
1825	The <i>Arabidopsis thaliana</i> multifunctional protein gene (MFP2) of peroxisomal beta-oxidation is essential for seedling establishment. 2006 , 45, 930-41	84
1824	Peroxisredoxin Q of <i>Arabidopsis thaliana</i> is attached to the thylakoids and functions in context of photosynthesis. 2006 , 45, 968-81	139
1823	Characterization of three members of the <i>Arabidopsis</i> carotenoid cleavage dioxygenase family demonstrates the divergent roles of this multifunctional enzyme family. 2006 , 45, 982-93	265
1822	Microarray analysis of the transcriptome as a stepping stone towards understanding biological systems: practical considerations and perspectives. 2006 , 45, 630-50	72
1821	CYP707A3, a major ABA 8'-hydroxylase involved in dehydration and rehydration response in <i>Arabidopsis thaliana</i> . 2006 , 46, 171-82	233
1820	An <i>Arabidopsis</i> cell wall-associated kinase required for invertase activity and cell growth. 2006 , 46, 307-16	138
1819	<i>Arabidopsis</i> myrosinases TGG1 and TGG2 have redundant function in glucosinolate breakdown and insect defense. 2006 , 46, 549-62	293

1818	AtNAP, a NAC family transcription factor, has an important role in leaf senescence. 2006 , 46, 601-12	572
1817	The N-terminal ATPase AT-hook-containing region of the Arabidopsis chromatin-remodeling protein SPLAYED is sufficient for biological activity. 2006 , 46, 685-99	23
1816	Transcriptional regulators of stamen development in Arabidopsis identified by transcriptional profiling. 2006 , 46, 984-1008	250
1815	Light activates the degradation of PIL5 protein to promote seed germination through gibberellin in Arabidopsis. 2006 , 47, 124-39	279
1814	Phosphatidylglycerol biosynthesis in chloroplasts of Arabidopsis mutants deficient in acyl-ACP glycerol-3- phosphate acyltransferase. 2006 , 47, 296-309	83
1813	UPF1 is required for nonsense-mediated mRNA decay (NMD) and RNAi in Arabidopsis. 2006 , 47, 480-9	167
1812	A trial of phenome analysis using 4000 Ds-insertional mutants in gene-coding regions of Arabidopsis. 2006 , 47, 640-51	96
1811	CUL4 associates with DDB1 and DET1 and its downregulation affects diverse aspects of development in Arabidopsis thaliana. 2006 , 47, 591-603	116
1810	A role for a flavin-containing mono-oxygenase in resistance against microbial pathogens in Arabidopsis. 2006 , 47, 629-39	76
1809	The pbs1-3 mutation in a putative dual-specificity protein tyrosine phosphatase gene provokes hypersensitive responses to abscisic acid in Arabidopsis thaliana. 2006 , 47, 711-9	48
1808	Mutation of SAD2, an importin beta-domain protein in Arabidopsis, alters abscisic acid sensitivity. 2006 , 47, 776-87	73
1807	AtEXO70A1, a member of a family of putative exocyst subunits specifically expanded in land plants, is important for polar growth and plant development. 2006 , 48, 54-72	206
1806	Arabidopsis SPO11-2 functions with SPO11-1 in meiotic recombination. 2006 , 48, 206-16	155
1805	An Arabidopsis chloroplast-targeted Hsp101 homologue, APG6, has an essential role in chloroplast development as well as heat-stress response. 2006 , 48, 249-60	62
1804	AtCAT6, a sink-tissue-localized transporter for essential amino acids in Arabidopsis. 2006 , 48, 414-26	92
1803	Regulation of hormone metabolism in Arabidopsis seeds: phytochrome regulation of abscisic acid metabolism and abscisic acid regulation of gibberellin metabolism. 2006 , 48, 354-66	319
1802	Arabidopsis WRKY33 transcription factor is required for resistance to necrotrophic fungal pathogens. 2006 , 48, 592-605	589
1801	Negative regulation of defense responses in Arabidopsis by two NPR1 paralogs. 2006 , 48, 647-56	152

1800	Yeast complementation reveals a role for an <i>Arabidopsis thaliana</i> late embryogenesis abundant (LEA)-like protein in oxidative stress tolerance. 2006 , 48, 743-56	80
1799	Association of <i>Arabidopsis</i> topoisomerase IIA cleavage sites with functional genomic elements and T-DNA loci. 2006 , 48, 697-709	11
1798	Isolation and characterization of the RAD54 gene from <i>Arabidopsis thaliana</i> . 2006 , 48, 827-42	75
1797	Non conservation of the meiotic function of the Ski8/Rec103 homolog in <i>Arabidopsis</i> . 2006 , 11, 615-22	39
1796	'Cisgenic' as a product designation. 2006 , 24, 1329-31; author reply 1331-3	12
1795	Reply to 'Cisgenic' as a product designation. 2006 , 24, 1331-1333	4
1794	Intracellular trafficking and proteolysis of the <i>Arabidopsis</i> auxin-efflux facilitator PIN2 are involved in root gravitropism. 2006 , 8, 249-56	471
1793	Moving forward in reverse: genetic technologies to enable genome-wide phenomic screens in <i>Arabidopsis</i> . 2006 , 7, 524-36	197
1792	AtSNX1 defines an endosome for auxin-carrier trafficking in <i>Arabidopsis</i> . 2006 , 443, 106-9	288
1791	AtSufE is an essential activator of plastidic and mitochondrial desulfurases in <i>Arabidopsis</i> . 2006 , 25, 900-9	74
1790	Reduced meiotic crossovers and delayed prophase I progression in AtMLH3-deficient <i>Arabidopsis</i> . 2006 , 25, 1315-23	102
1789	bZIP10-LSD1 antagonism modulates basal defense and cell death in <i>Arabidopsis</i> following infection. 2006 , 25, 4400-11	191
1788	A homologue of the breast cancer-associated gene BARD1 is involved in DNA repair in plants. 2006 , 25, 4326-37	38
1787	The role of AtMSH2 in homologous recombination in <i>Arabidopsis thaliana</i> . 2006 , 7, 100-5	64
1786	Characterization of a novel putative zinc finger gene MIF1: involvement in multiple hormonal regulation of <i>Arabidopsis</i> development. 2006 , 45, 399-422	76
1785	Distinct and overlapping roles of two gibberellin 3-oxidases in <i>Arabidopsis</i> development. 2006 , 45, 804-18	225
1784	Transgressive segregation reveals two <i>Arabidopsis</i> TIR-NB-LRR resistance genes effective against <i>Leptosphaeria maculans</i> , causal agent of blackleg disease. 2006 , 46, 218-30	71
1783	AtMBD9: a protein with a methyl-CpG-binding domain regulates flowering time and shoot branching in <i>Arabidopsis</i> . 2006 , 46, 282-96	48

1782	Phytochrome-mediated agravitropism in Arabidopsis hypocotyls requires GIL1 and confers a fitness advantage. 2006 , 46, 641-8	19
1781	Impaired sucrose induction1 encodes a conserved plant-specific protein that couples carbohydrate availability to gene expression and plant growth. 2006 , 46, 1045-58	31
1780	Cloning of two splice variants of the rice PTS1 receptor, OsPex5pL and OsPex5pS, and their functional characterization using pex5-deficient yeast and Arabidopsis. 2006 , 47, 457-66	30
1779	Functional and expression analysis of Arabidopsis SPA genes during seedling photomorphogenesis and adult growth. 2006 , 47, 577-90	69
1778	Expression profiling and mutant analysis reveals complex regulatory networks involved in Arabidopsis response to Botrytis infection. 2006 , 48, 28-44	217
1777	The protein kinase genes MAP3K epsilon 1 and MAP3K epsilon 2 are required for pollen viability in Arabidopsis thaliana. 2006 , 48, 193-205	31
1776	PDX1 is essential for vitamin B6 biosynthesis, development and stress tolerance in Arabidopsis. 2006 , 48, 933-46	121
1775	The Arabidopsis SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASE1 protein complex includes BRASSINOSTEROID-INSENSITIVE1. 2006 , 18, 626-38	215
1774	AtGRP7 is involved in the regulation of abscisic acid and stress responses in Arabidopsis. 2006 , 11, 526-35	60
1773	Deletion of core components of the plastid protein import machinery causes differential arrest of embryo development in Arabidopsis thaliana. 2006 , 8, 18-30	44
1772	Diverse functions and molecular properties emerging for CAX cation/H ⁺ exchangers in plants. 2006 , 8, 419-29	96
1771	Evaluation of a non-targeted "omic" approach in the safety assessment of genetically modified plants. 2006 , 8, 662-72	16
1770	A map of KNAT gene expression in the Arabidopsis root. 2006 , 60, 1-20	72
1769	OsEIL1, a rice homolog of the Arabidopsis EIN3 regulates the ethylene response as a positive component. 2006 , 61, 141-52	81
1768	Plant coenzyme A biosynthesis: characterization of two pantothenate kinases from Arabidopsis. 2006 , 61, 629-42	25
1767	Development of a virus-induced gene-silencing system for functional analysis of the RPS2-dependent resistance signalling pathways in Arabidopsis. 2006 , 62, 223-32	36
1766	The putative SWI/SNF complex subunit BRAHMA activates flower homeotic genes in Arabidopsis thaliana. 2006 , 62, 291-304	98
1765	State transitions revisited-a buffering system for dynamic low light acclimation of Arabidopsis. 2006 , 62, 779-93	71

1764	BOLITA, an Arabidopsis AP2/ERF-like transcription factor that affects cell expansion and proliferation/differentiation pathways. 2006 , 62, 825-43	65
1763	The Arabidopsis stearyl-acyl carrier protein-desaturase family and the contribution of leaf isoforms to oleic acid synthesis. 2007 , 63, 257-71	167
1762	Biotechnology approaches to overcome biotic and abiotic stress constraints in legumes. 2006 , 147, 1-24	179
1761	Isolation and characterization of a novel semi-lethal Arabidopsis thaliana mutant of gene for pentatricopeptide (PPR) repeat-containing protein. 2006 , 128, 395-407	15
1760	Chromosome synapsis in Arabidopsis: analysis of the transverse filament protein ZYP1 reveals novel functions for the synaptonemal complex. 2006 , 115, 212-9	42
1759	Two cell wall associated peroxidases from Arabidopsis influence root elongation. 2006 , 223, 965-74	129
1758	Progress report: ethylene signaling and responses. 2006 , 223, 387-91	35
1757	AtAAH encodes a protein with allantoate amidohydrolase activity from Arabidopsis thaliana. 2006 , 223, 1108-13	43
1756	UDP-sugar pyrophosphorylase is essential for pollen development in Arabidopsis. 2006 , 224, 520-32	70
1755	ABA-responsive RNA-binding proteins are involved in chloroplast and stromule function in Arabidopsis seedlings. 2006 , 224, 900-14	42
1754	Involvement of AtLAC15 in lignin synthesis in seeds and in root elongation of Arabidopsis. 2006 , 224, 1185-96	127
1753	Evidence for a role of AtCAD 1 in lignification of elongating stems of Arabidopsis thaliana. 2006 , 225, 23-39	70
1752	In vitro-mutagenesis of NADPH:protochlorophyllide oxidoreductase B: two distinctive protochlorophyllide binding sites participate in enzyme catalysis and assembly. 2006 , 275, 540-52	15
1751	Coordination of cell proliferation and cell expansion in the control of leaf size in Arabidopsis thaliana. 2006 , 119, 37-42	166
1750	Sequence stability of the T-DNA - plant junctions in tissue culture in Arabidopsis transgenic lines. 2006 , 25, 1362-8	6
1749	The ATL gene family from Arabidopsis thaliana and Oryza sativa comprises a large number of putative ubiquitin ligases of the RING-H2 type. 2006 , 62, 434-45	88
1748	Unintended consequences of plant transformation: a molecular insight. 2006 , 47, 277-86	86
1747	Two kinesins are involved in the spatial control of cytokinesis in Arabidopsis thaliana. 2006 , 16, 888-94	109

1746	Molecular characterization of a bHLH transcription factor involved in Arabidopsis abscisic acid-mediated response. 2006 , 1759, 191-4	27
1745	Arabidopsis thaliana--a model organism to study plant peroxisomes. 2006 , 1763, 1382-91	64
1744	Application of TILLING in plant improvement. 2006 , 33, 957-64	8
1743	A case of promiscuity: Agrobacterium's endless hunt for new partners. 2006 , 22, 29-37	154
1742	Root development--branching into novel spheres. 2006 , 9, 66-71	27
1741	New FATB thioesterases from a high-laurate Cuphea species: Functional and complementation analyses. 2006 , 108, 979-990	10
1740	How ethylene works in the reproductive organs of higher plants: a signaling update from the third millennium. 2006 , 1, 231-42	17
1739	Altered Expression of PERK Receptor Kinases in Arabidopsis Leads to Changes in Growth and Floral Organ Formation. 2006 , 1, 251-60	21
1738	Use of microarray analysis to dissect the plant defense response. 2007 , 354, 121-30	3
1737	Using information from public Arabidopsis databases to aid research. 2006 , 323, 187-211	1
1736	Characterization of the three Arabidopsis thaliana RAD21 cohesins reveals differential responses to ionizing radiation. 2006 , 57, 971-83	34
1735	The role of AtMUS81 in DNA repair and its genetic interaction with the helicase AtRecQ4A. 2006 , 34, 4438-48	83
1734	Molecular characterization and phylogeny of U2AF35 homologs in plants. 2006 , 140, 624-36	37
1733	The Arabidopsis flavin-dependent monooxygenase FMO1 is an essential component of biologically induced systemic acquired resistance. 2006 , 141, 1666-75	188
1732	Adenosine kinase modulates root gravitropism and cap morphogenesis in Arabidopsis. 2006 , 142, 564-73	50
1731	Increased level of polyploidy1, a conserved repressor of CYCLINA2 transcription, controls endoreduplication in Arabidopsis. 2006 , 18, 2452-68	57
1730	Identification of components in disease-resistance signaling in Arabidopsis by map-based cloning. 2007 , 354, 69-78	10
1729	The Arabidopsis histidine phosphotransfer proteins are redundant positive regulators of cytokinin signaling. 2006 , 18, 3073-87	313

1728	PCR-based screening for insertional mutants. 2006 , 323, 163-72	8
1727	A molecular portrait of Arabidopsis meiosis. 2006 , 4, e0095	44
1726	SUPPRESSOR OF FRI 4 encodes a nuclear-localized protein that is required for delayed flowering in winter-annual Arabidopsis. 2006 , 133, 4699-707	55
1725	Generation of composite plants using Agrobacterium rhizogenes. 2006 , 343, 155-67	26
1724	Optimisation of tobacco rattle virus-induced gene silencing in Arabidopsis. 2006 , 33, 347-355	56
1723	Transcriptome profiling, molecular biological, and physiological studies reveal a major role for ethylene in cotton fiber cell elongation. 2006 , 18, 651-64	424
1722	Arabidopsis INOSITOL TRANSPORTER4 mediates high-affinity H ⁺ symport of myoinositol across the plasma membrane. 2006 , 141, 565-77	63
1721	ABA-hypersensitive germination3 encodes a protein phosphatase 2C (AtPP2CA) that strongly regulates abscisic acid signaling during germination among Arabidopsis protein phosphatase 2Cs. 2006 , 140, 115-26	284
1720	Arabidopsis thaliana cyclic nucleotide gated channel 3 forms a non-selective ion transporter involved in germination and cation transport. 2006 , 57, 791-800	167
1719	The mutational consequences of plant transformation. 2006 , 2006, 25376	92
1718	The SCABRA3 nuclear gene encodes the plastid RpoTp RNA polymerase, which is required for chloroplast biogenesis and mesophyll cell proliferation in Arabidopsis. 2006 , 141, 942-56	117
1717	The ENHANCER OF SHOOT REGENERATION 2 gene in Arabidopsis regulates CUP-SHAPED COTYLEDON 1 at the transcriptional level and controls cotyledon development. 2006 , 47, 1443-56	119
1716	Characterization of the plant-specific BREVIS RADIX gene family reveals limited genetic redundancy despite high sequence conservation. 2006 , 140, 1306-16	46
1715	BIGYIN, an orthologue of human and yeast FIS1 genes functions in the control of mitochondrial size and number in Arabidopsis thaliana. 2006 , 57, 1275-80	82
1714	Functional profiling reveals that only a small number of phytochrome-regulated early-response genes in Arabidopsis are necessary for optimal deetiolation. 2006 , 18, 2157-71	81
1713	Arabidopsis CUP-SHAPED COTYLEDON3 regulates postembryonic shoot meristem and organ boundary formation. 2006 , 18, 2946-57	237
1712	Arabidopsis REGULATOR OF AXILLARY MERISTEMS1 controls a leaf axil stem cell niche and modulates vegetative development. 2006 , 18, 598-611	148
1711	Ethylene-dependent and -independent pathways controlling floral abscission are revealed to converge using promoter::reporter gene constructs in the ida abscission mutant. 2006 , 57, 3627-37	49

1710	Analysis of the Arabidopsis <i>rsr4-1/pdx1-3</i> mutant reveals the critical function of the PDX1 protein family in metabolism, development, and vitamin B6 biosynthesis. 2006 , 18, 1722-35	66
1709	Branched-chain aminotransferase4 is part of the chain elongation pathway in the biosynthesis of methionine-derived glucosinolates in Arabidopsis. 2006 , 18, 2664-79	132
1708	A cytosolic Arabidopsis D-xylulose kinase catalyzes the phosphorylation of 1-deoxy-D-xylulose into a precursor of the plastidial isoprenoid pathway. 2006 , 142, 441-57	42
1707	Double knockouts of phospholipases Dzeta1 and Dzeta2 in Arabidopsis affect root elongation during phosphate-limited growth but do not affect root hair patterning. 2006 , 140, 761-70	162
1706	Locus-specific control of DNA methylation by the Arabidopsis SUVH5 histone methyltransferase. 2006 , 18, 1166-76	223
1705	FZL, an FZO-like protein in plants, is a determinant of thylakoid and chloroplast morphology. 2006 , 103, 6759-64	93
1704	The cell surface leucine-rich repeat receptor for AtPep1, an endogenous peptide elicitor in Arabidopsis, is functional in transgenic tobacco cells. 2006 , 103, 10104-9	309
1703	Tic21 is an essential translocon component for protein translocation across the chloroplast inner envelope membrane. 2006 , 18, 2247-57	135
1702	Desensitization of GSTF8 induction by a prior chemical treatment is long lasting and operates in a tissue-dependent manner. 2006 , 142, 245-53	15
1701	Enhancement of abscisic acid sensitivity and reduction of water consumption in Arabidopsis by combined inactivation of the protein phosphatases type 2C ABI1 and HAB1. 2006 , 141, 1389-99	197
1700	SUGAR-DEPENDENT1 encodes a patatin domain triacylglycerol lipase that initiates storage oil breakdown in germinating Arabidopsis seeds. 2006 , 18, 665-75	303
1699	Arabidopsis HAP2 (GCS1) is a sperm-specific gene required for pollen tube guidance and fertilization. 2006 , 133, 4761-9	232
1698	G-protein-coupled receptor 1, G-protein Galpha-subunit 1, and prephenate dehydratase 1 are required for blue light-induced production of phenylalanine in etiolated Arabidopsis. 2006 , 140, 844-55	101
1697	Plastid movement impaired 2, a new gene involved in normal blue-light-induced chloroplast movements in Arabidopsis. 2006 , 141, 1328-37	62
1696	A basic leucine zipper transcription factor, G-box-binding factor 1, regulates blue light-mediated photomorphogenic growth in Arabidopsis. 2006 , 281, 22190-22199	76
1695	Arabidopsis orthologs of maize chloroplast splicing factors promote splicing of orthologous and species-specific group II introns. 2006 , 142, 1656-63	93
1694	The HVE/CAND1 gene is required for the early patterning of leaf venation in Arabidopsis. 2006 , 133, 3755-66	52
1693	A role for chromatin remodeling in regulation of CUC gene expression in the Arabidopsis cotyledon boundary. 2006 , 133, 3223-30	79

1692	A redox-regulated chloroplast protein phosphatase binds to starch diurnally and functions in its accumulation. 2006 , 103, 9732-7	99
1691	The Arabidopsis SUPPRESSOR OF AUXIN RESISTANCE proteins are nucleoporins with an important role in hormone signaling and development. 2006 , 18, 1590-603	130
1690	AHK5 histidine kinase regulates root elongation through an ETR1-dependent abscisic acid and ethylene signaling pathway in Arabidopsis thaliana. 2007 , 48, 375-80	53
1689	A PERK-like receptor kinase interacts with the geminivirus nuclear shuttle protein and potentiates viral infection. 2006 , 80, 6648-56	70
1688	Chromatin assembly factor CAF-1 is required for cellular differentiation during plant development. 2006 , 133, 4163-72	94
1687	Regulation of flowering time by Arabidopsis MSI1. 2006 , 133, 1693-702	76
1686	Gamma-tubulin is essential for microtubule organization and development in Arabidopsis. 2006 , 18, 1412-25	131
1685	An Arabidopsis Fip1 homolog interacts with RNA and provides conceptual links with a number of other polyadenylation factor subunits. 2006 , 281, 176-86	35
1684	Gain-of-function phenotypes of many CLAVATA3/ESR genes, including four new family members, correlate with tandem variations in the conserved CLAVATA3/ESR domain. 2006 , 140, 1331-44	160
1683	ARABINAN DEFICIENT 1 is a putative arabinosyltransferase involved in biosynthesis of pectic arabinan in Arabidopsis. 2006 , 140, 49-58	210
1682	SCARFACE encodes an ARF-GAP that is required for normal auxin efflux and vein patterning in Arabidopsis. 2006 , 18, 1396-411	110
1681	Auxin biosynthesis by the YUCCA flavin monooxygenases controls the formation of floral organs and vascular tissues in Arabidopsis. 2006 , 20, 1790-9	803
1680	Genetic basis for natural variation in seed vitamin E levels in Arabidopsis thaliana. 2006 , 103, 18834-41	61
1679	BRICK1/HSPC300 functions with SCAR and the ARP2/3 complex to regulate epidermal cell shape in Arabidopsis. 2006 , 133, 1091-100	81
1678	Kinase and phosphatase: the cog and spring of the circadian clock. 2006 , 250, 47-72	30
1677	The transcription factors WRKY11 and WRKY17 act as negative regulators of basal resistance in Arabidopsis thaliana. 2006 , 18, 3289-302	331
1676	Involvement of a glycerol-3-phosphate dehydrogenase in modulating the NADH/NAD ⁺ ratio provides evidence of a mitochondrial glycerol-3-phosphate shuttle in Arabidopsis. 2006 , 18, 422-41	112
1675	The Arabidopsis zinc finger-homeodomain genes encode proteins with unique biochemical properties that are coordinately expressed during floral development. 2006 , 140, 1095-108	103

1674	Quantitative profiling of Arabidopsis polar glycerolipids in response to phosphorus starvation. Roles of phospholipases D zeta1 and D zeta2 in phosphatidylcholine hydrolysis and digalactosyldiacylglycerol accumulation in phosphorus-starved plants. 2006 , 142, 750-61	186
1673	The protein phosphatase AtPP2CA negatively regulates abscisic acid signal transduction in Arabidopsis, and effects of abh1 on AtPP2CA mRNA. 2006 , 140, 127-39	212
1672	A novel plant-specific family gene, ROOT PRIMORDIUM DEFECTIVE 1, is required for the maintenance of active cell proliferation. 2006 , 140, 591-602	34
1671	Separation of Arabidopsis pollen tetrads is regulated by QUARTET1, a pectin methylesterase gene. 2006 , 142, 1004-13	180
1670	Type-B ARR transcription factors, ARR10 and ARR12, are implicated in cytokinin-mediated regulation of protoxylem differentiation in roots of Arabidopsis thaliana. 2007 , 48, 84-96	154
1669	EARLY IN SHORT DAYS 1 (ESD1) encodes ACTIN-RELATED PROTEIN 6 (AtARP6), a putative component of chromatin remodelling complexes that positively regulates FLC accumulation in Arabidopsis. 2006 , 133, 1241-52	106
1668	Polycomb-group proteins repress the floral activator AGL19 in the FLC-independent vernalization pathway. 2006 , 20, 1667-78	146
1667	AtATG genes, homologs of yeast autophagy genes, are involved in constitutive autophagy in Arabidopsis root tip cells. 2006 , 47, 1641-52	141
1666	The LEAFY target LMI1 is a meristem identity regulator and acts together with LEAFY to regulate expression of CAULIFLOWER. 2006 , 133, 1673-82	114
1665	Loss of function of COBRA, a determinant of oriented cell expansion, invokes cellular defence responses in Arabidopsis thaliana. 2006 , 57, 2923-36	48
1664	Vacuolar invertase regulates elongation of Arabidopsis thaliana roots as revealed by QTL and mutant analysis. 2006 , 103, 2994-9	149
1663	Characterization of 43 non-protein-coding mRNA genes in Arabidopsis, including the MIR162a-derived transcripts. 2006 , 140, 1192-204	103
1662	DAWDLE, a forkhead-associated domain gene, regulates multiple aspects of plant development. 2006 , 141, 932-41	43
1661	Mutations in the microRNA complementarity site of the INCURVATA4 gene perturb meristem function and adaxialize lateral organs in arabidopsis. 2006 , 141, 607-19	77
1660	Receptor signal output mediated by the ETR1 N terminus is primarily subfamily I receptor dependent. 2006 , 142, 492-508	72
1659	The Arabidopsis-me12-like genes play a role in meiosis and vegetative growth in Arabidopsis. 2006 , 18, 545-59	67
1658	CONSTITUTIVELY PHOTOMORPHOGENIC1 is required for the UV-B response in Arabidopsis. 2006 , 18, 1975-90	289
1657	Arabidopsis NRP1 and NRP2 encode histone chaperones and are required for maintaining postembryonic root growth. 2006 , 18, 2879-92	105

1656	Mutant identification and characterization of the laccase gene family in Arabidopsis. 2006 , 57, 2563-9	126
1655	The Arabidopsis thaliana PARTING DANCERS gene encoding a novel protein is required for normal meiotic homologous recombination. 2006 , 17, 1331-43	63
1654	A Golgi-localized hexose transporter is involved in heterotrimeric G protein-mediated early development in Arabidopsis. 2006 , 17, 4257-69	66
1653	A PHD finger protein involved in both the vernalization and photoperiod pathways in Arabidopsis. 2006 , 20, 3244-8	161
1652	GA4 is the active gibberellin in the regulation of LEAFY transcription and Arabidopsis floral initiation. 2006 , 18, 2172-81	268
1651	A bacterial virulence protein suppresses host innate immunity to cause plant disease. <i>Science</i> , 2006 , 313, 220-3	333 382
1650	Control of leaf vascular patterning by polar auxin transport. 2006 , 20, 1015-27	586
1649	Cytochrome P450 CYP710A encodes the sterol C-22 desaturase in Arabidopsis and tomato. 2006 , 18, 1008-22	115
1648	Mutants of Arabidopsis lacking starch branching enzyme II substitute plastidial starch synthesis by cytoplasmic maltose accumulation. 2006 , 18, 2694-709	86
1647	PsbR, a missing link in the assembly of the oxygen-evolving complex of plant photosystem II. 2006 , 281, 145-50	98
1646	Arabidopsis reversibly glycosylated polypeptides 1 and 2 are essential for pollen development. 2006 , 142, 1480-92	86
1645	AtGRXcp, an Arabidopsis chloroplastic glutaredoxin, is critical for protection against protein oxidative damage. 2006 , 281, 26280-8	119
1644	AtIREG2 encodes a tonoplast transport protein involved in iron-dependent nickel detoxification in Arabidopsis thaliana roots. 2006 , 281, 25532-40	169
1643	A pentatricopeptide repeat protein facilitates the trans-splicing of the maize chloroplast rps12 pre-mRNA. 2006 , 18, 2650-63	227
1642	Developmentally controlled farnesylation modulates AtNAP1;1 function in cell proliferation and cell expansion during Arabidopsis leaf development. 2006 , 142, 1412-26	51
1641	A second thylakoid membrane-localized Alb3/Oxal/YidC homologue is involved in proper chloroplast biogenesis in Arabidopsis thaliana. 2006 , 281, 16632-42	60
1640	Plant-Pathogen Interactions. 2006 ,	2
1639	AtALMT1, which encodes a malate transporter, is identified as one of several genes critical for aluminum tolerance in Arabidopsis. 2006 , 103, 9738-43	420

1638	Metabolomics and Gene Identification in Plant Natural Product Pathways. 2006 , 243-259	3
1637	Physical and functional interactions between pathogen-induced Arabidopsis WRKY18, WRKY40, and WRKY60 transcription factors. 2006 , 18, 1310-26	541
1636	The AUXIN RESPONSE FACTOR 2 gene of Arabidopsis links auxin signalling, cell division, and the size of seeds and other organs. 2006 , 133, 251-61	403
1635	ETHYLENE-INSENSITIVE5 encodes a 5'→3' exonuclease required for regulation of the EIN3-targeting F-box proteins EBF1/2. 2006 , 103, 13286-93	148
1634	CONSTANS and the CCAAT box binding complex share a functionally important domain and interact to regulate flowering of Arabidopsis. 2006 , 18, 2971-84	415
1633	Arabidopsis LHT1 is a high-affinity transporter for cellular amino acid uptake in both root epidermis and leaf mesophyll. 2006 , 18, 1931-46	239
1632	The Arabidopsis major intrinsic protein NIP5;1 is essential for efficient boron uptake and plant development under boron limitation. 2006 , 18, 1498-509	526
1631	Arabidopsis Hsa32, a novel heat shock protein, is essential for acquired thermotolerance during long recovery after acclimation. 2006 , 140, 1297-305	208
1630	REVERSION-TO-ETHYLENE SENSITIVITY1, a conserved gene that regulates ethylene receptor function in Arabidopsis. 2006 , 103, 7917-22	171
1629	The balance between the MIR164A and CUC2 genes controls leaf margin serration in Arabidopsis. 2006 , 18, 2929-45	414
1628	RACK1 mediates multiple hormone responsiveness and developmental processes in Arabidopsis. 2006 , 57, 2697-708	107
1627	AUXIN RESPONSE FACTOR8 is a negative regulator of fruit initiation in Arabidopsis. 2006 , 18, 1873-86	203
1626	Evidence for distinct mechanisms of starch granule breakdown in plants. 2006 , 281, 12050-9	115
1625	Manipulation of Arabidopsis fatty acid amide hydrolase expression modifies plant growth and sensitivity to N-acyl ethanolamines. 2006 , 103, 12197-202	61
1624	Multiple heme oxygenase family members contribute to the biosynthesis of the phytochrome chromophore in Arabidopsis. 2006 , 140, 856-68	94
1623	An Arabidopsis basic helix-loop-helix leucine zipper protein modulates metal homeostasis and auxin conjugate responsiveness. 2006 , 174, 1841-57	83
1622	Downregulation of ClpR2 leads to reduced accumulation of the ClpPRS protease complex and defects in chloroplast biogenesis in Arabidopsis. 2006 , 18, 1704-21	100
1621	A fluoroorotic acid-resistant mutant of Arabidopsis defective in the uptake of uracil. 2006 , 57, 3563-73	8

1620	Cloning and characterization of Arabidopsis and Brassica juncea flavin-containing amine oxidases. 2006 , 57, 4155-69	15
1619	Trans-acting siRNA-mediated repression of ETTIN and ARF4 regulates heteroblasty in Arabidopsis. 2006 , 133, 2973-81	281
1618	Partially redundant functions of two SET-domain polycomb-group proteins in controlling initiation of seed development in Arabidopsis. 2006 , 103, 13244-9	111
1617	Identification of a peroxisomal acyl-activating enzyme involved in the biosynthesis of jasmonic acid in Arabidopsis. 2006 , 281, 33511-20	152
1616	An ordered, nonredundant library of Pseudomonas aeruginosa strain PA14 transposon insertion mutants. 2006 , 103, 2833-8	707
1615	The Arabidopsis thaliana MND1 homologue plays a key role in meiotic homologous pairing, synapsis and recombination. 2006 , 119, 2486-96	67
1614	Global analysis of della direct targets in early gibberellin signaling in Arabidopsis. 2007 , 19, 3037-57	464
1613	The Arabidopsis irregular xylem8 mutant is deficient in glucuronoxylan and homogalacturonan, which are essential for secondary cell wall integrity. 2007 , 19, 237-55	219
1612	Identification of acyltransferases required for cutin biosynthesis and production of cutin with suberin-like monomers. 2007 , 104, 18339-44	274
1611	MYB98 positively regulates a battery of synergid-expressed genes encoding filiform apparatus localized proteins. 2007 , 19, 2557-68	100
1610	MEKK1 is required for flg22-induced MPK4 activation in Arabidopsis plants. 2007 , 143, 661-9	248
1609	AtERF14, a member of the ERF family of transcription factors, plays a nonredundant role in plant defense. 2007 , 143, 400-9	153
1608	DNA demethylation in the Arabidopsis genome. 2007 , 104, 6752-7	376
1607	Two Arabidopsis genes (IPMS1 and IPMS2) encode isopropylmalate synthase, the branchpoint step in the biosynthesis of leucine. 2007 , 143, 970-86	69
1606	Arabidopsis CAP1 - a key regulator of actin organisation and development. 2007 , 120, 2609-18	54
1605	Effects of mutations in Arabidopsis FtsZ1 on plastid division, FtsZ ring formation and positioning, and FtsZ filament morphology in vivo. 2007 , 48, 775-91	53
1604	Pollen tetrad-based visual assay for meiotic recombination in Arabidopsis. 2007 , 104, 3913-8	146
1603	Arabidopsis thaliana squalene epoxidase 1 is essential for root and seed development. 2007 , 282, 17002-13	68

1602	A cluster of disease resistance genes in Arabidopsis is coordinately regulated by transcriptional activation and RNA silencing. 2007 , 19, 2929-39	152
1601	Profile of Joseph R. Ecker. 2007 , 104, 3023-3025	78
1600	A small ATPase protein of Arabidopsis, TGD3, involved in chloroplast lipid import. 2007 , 282, 35945-53	107
1599	TRICHOMELESS1 regulates trichome patterning by suppressing GLABRA1 in Arabidopsis. 2007 , 134, 3873-82	134
1598	A heteromeric plastidic pyruvate kinase complex involved in seed oil biosynthesis in Arabidopsis. 2007 , 19, 2006-22	147
1597	Arabidopsis relatives of the human lysine-specific Demethylase1 repress the expression of FWA and FLOWERING LOCUS C and thus promote the floral transition. 2007 , 19, 2975-87	172
1596	PIL5, a phytochrome-interacting bHLH protein, regulates gibberellin responsiveness by binding directly to the GAI and RGA promoters in Arabidopsis seeds. 2007 , 19, 1192-208	328
1595	The Arabidopsis ATNRT2.7 nitrate transporter controls nitrate content in seeds. 2007 , 19, 1590-602	154
1594	ASY1 mediates AtDMC1-dependent interhomolog recombination during meiosis in Arabidopsis. 2007 , 21, 2220-33	184
1593	A nitrite transporter associated with nitrite uptake by higher plant chloroplasts. 2007 , 48, 1022-35	89
1592	Identification of two protein kinases required for abscisic acid regulation of seed germination, root growth, and gene expression in Arabidopsis. 2007 , 19, 485-94	482
1591	Arabidopsis BRANCHED1 acts as an integrator of branching signals within axillary buds. 2007 , 19, 458-72	507
1590	The Significance of Biology for Human Development: A Developmental Psychobiological Systems View. 2007 ,	8
1589	Methylation of gibberellins by Arabidopsis GAMT1 and GAMT2. 2007 , 19, 32-45	182
1588	GreenPhylDB: a database for plant comparative genomics. 2008 , 36, D991-8	64
1587	The central cell plays a critical role in pollen tube guidance in Arabidopsis. 2007 , 19, 3563-77	136
1586	Inositol polyphosphate 5-phosphatases 1 and 2 are required for regulating seedling growth. 2007 , 143, 1408-17	51
1585	The auxin-regulated AP2/EREBP gene PUCHI is required for morphogenesis in the early lateral root primordium of Arabidopsis. 2007 , 19, 2156-68	141

1584	Genetic characterization of mutants resistant to the antiauxin p-chlorophenoxyisobutyric acid reveals that AAR3, a gene encoding a DCN1-like protein, regulates responses to the synthetic auxin 2,4-dichlorophenoxyacetic acid in Arabidopsis roots. 2007 , 145, 773-85	38
1583	Identification of Arabidopsis cyclase-associated protein 1 as the first nucleotide exchange factor for plant actin. 2007 , 18, 3002-14	62
1582	A novel major facilitator superfamily protein at the tonoplast influences zinc tolerance and accumulation in Arabidopsis. 2007 , 143, 1705-19	177
1581	NAC transcription factors, NST1 and NST3, are key regulators of the formation of secondary walls in woody tissues of Arabidopsis. 2007 , 19, 270-80	576
1580	Telomere dynamics and fusion of critically shortened telomeres in plants lacking DNA ligase IV. 2007 , 35, 6490-500	60
1579	Mutational evidence that the Arabidopsis MAP kinase MPK6 is involved in anther, inflorescence, and embryo development. 2007 , 58, 2181-91	88
1578	An extra-plastidial alpha-glucan, water dikinase from Arabidopsis phosphorylates amylopectin in vitro and is not necessary for transient starch degradation. 2007 , 58, 3949-60	26
1577	Zip4/Spo22 is required for class I CO formation but not for synapsis completion in Arabidopsis thaliana. 2007 , 3, e83	126
1576	Distinct modulations of the hexokinase1-mediated glucose response and hexokinase1-independent processes by HYS1/CPR5 in Arabidopsis. 2007 , 58, 3239-48	25
1575	The GCR1, GPA1, PRN1, NF-Y signal chain mediates both blue light and abscisic acid responses in Arabidopsis. 2007 , 143, 1590-600	161
1574	STRESS RESPONSE SUPPRESSOR1 and STRESS RESPONSE SUPPRESSOR2, two DEAD-box RNA helicases that attenuate Arabidopsis responses to multiple abiotic stresses. 2007 , 145, 814-30	137
1573	Two Arabidopsis threonine aldolases are nonredundant and compete with threonine deaminase for a common substrate pool. 2006 , 18, 3564-75	69
1572	An ABC transporter gene of Arabidopsis thaliana, AtWBC11, is involved in cuticle development and prevention of organ fusion. 2007 , 48, 1790-802	125
1571	GABI-Kat SimpleSearch: an Arabidopsis thaliana T-DNA mutant database with detailed information for confirmed insertions. 2007 , 35, D874-8	42
1570	Transcriptional profiling of the Arabidopsis embryo. 2007 , 143, 924-40	109
1569	A heat-inducible transcription factor, HsfA2, is required for extension of acquired thermotolerance in Arabidopsis. 2007 , 143, 251-62	393
1568	SEF, a new protein required for flowering repression in Arabidopsis, interacts with PIE1 and ARP6. 2007 , 143, 893-901	99
1567	Arabidopsis nucleolin affects plant development and patterning. 2007 , 144, 173-86	83

1566	Mitochondrial complex II Is essential for gametophyte development in Arabidopsis. 2007 , 143, 1534-46	85
1565	Deficiency in phosphatidylserine decarboxylase activity in the psd1 psd2 psd3 triple mutant of Arabidopsis affects phosphatidylethanolamine accumulation in mitochondria. 2007 , 144, 904-14	59
1564	Genetic dissection of histidine biosynthesis in Arabidopsis. 2007 , 144, 890-903	58
1563	The GH3 acyl adenylase family member PBS3 regulates salicylic acid-dependent defense responses in Arabidopsis. 2007 , 144, 1144-56	158
1562	Identification of the photorespiratory 2-phosphoglycolate phosphatase, PGLP1, in Arabidopsis. 2007 , 144, 1580-6	74
1561	Induction of isoforms of tetrapyrrole biosynthetic enzymes, AtHEMA2 and AtFC1, under stress conditions and their physiological functions in Arabidopsis. 2007 , 144, 1039-51	52
1560	Mitochondrial protein lipoylation does not exclusively depend on the mtKAS pathway of de novo fatty acid synthesis in Arabidopsis. 2007 , 145, 41-8	32
1559	Global expression profiling applied to the analysis of Arabidopsis stamen development. 2007 , 145, 747-62	95
1558	yucca6, a dominant mutation in Arabidopsis, affects auxin accumulation and auxin-related phenotypes. 2007 , 145, 722-35	115
1557	Distinct functions for the two PsbP-like proteins PPL1 and PPL2 in the chloroplast thylakoid lumen of Arabidopsis. 2007 , 145, 668-79	119
1556	The cytochrome P450 enzyme CYP96A15 is the midchain alkane hydroxylase responsible for formation of secondary alcohols and ketones in stem cuticular wax of Arabidopsis. 2007 , 145, 653-67	195
1555	Arabidopsis INOSITOL TRANSPORTER2 mediates H ⁺ symport of different inositol epimers and derivatives across the plasma membrane. 2007 , 145, 1395-407	50
1554	Functional diversification of acyl-coenzyme A oxidases in jasmonic acid biosynthesis and action. 2007 , 143, 812-24	142
1553	Arabidopsis inositol polyphosphate 6-/3-kinase (AtIpk2beta) is involved in axillary shoot branching via auxin signaling. 2007 , 144, 942-51	41
1552	The nuclear pore protein AtTPR is required for RNA homeostasis, flowering time, and auxin signaling. 2007 , 144, 1383-90	80
1551	A putative CCAAT-binding transcription factor is a regulator of flowering timing in Arabidopsis. 2007 , 145, 98-105	116
1550	Arabidopsis potential calcium sensors regulate nitric oxide levels and the transition to flowering. 2007 , 2, 446-54	55
1549	The Protein Phosphatases and Protein Kinases of Arabidopsis thaliana. 2007 , 5, e0106	41

1548	The <i>Arabidopsis thaliana</i> homolog of yeast BRE1 has a function in cell cycle regulation during early leaf and root growth. 2007 , 19, 417-32	123
1547	Genetic analysis of SUMOylation in <i>Arabidopsis</i> : conjugation of SUMO1 and SUMO2 to nuclear proteins is essential. 2007 , 145, 119-34	208
1546	Safe genetically engineered plants. 2007 , 19, 395005	4
1545	Stomatal development and patterning are regulated by environmentally responsive mitogen-activated protein kinases in <i>Arabidopsis</i> . 2007 , 19, 63-73	587
1544	R1R2R3-Myb proteins positively regulate cytokinesis through activation of KNOLLE transcription in <i>Arabidopsis thaliana</i> . 2007 , 134, 1101-10	133
1543	SUBA: the <i>Arabidopsis</i> Subcellular Database. 2007 , 35, D213-8	364
1542	Detached and attached <i>Arabidopsis</i> leaf assays reveal distinctive defense responses against hemibiotrophic <i>Colletotrichum</i> spp. 2007 , 20, 1308-19	74
1541	The receptor-like kinase SERK3/BAK1 is a central regulator of innate immunity in plants. 2007 , 104, 12217-22	784
1540	Insertional mutagenesis as a tool to study genes/functions in <i>Chlamydomonas</i> . 2007 , 616, 77-89	26
1539	<i>Arabidopsis</i> PLDzeta2 regulates vesicle trafficking and is required for auxin response. 2007 , 19, 281-95	171
1538	A conserved carboxylesterase is a SUPPRESSOR OF AVRBS1-ELICITED RESISTANCE in <i>Arabidopsis</i> . 2007 , 19, 688-705	49
1537	Functional definition of outer membrane proteins involved in preprotein import into mitochondria. 2007 , 19, 3739-59	117
1536	The DELLA domain of GA INSENSITIVE mediates the interaction with the GA INSENSITIVE DWARF1A gibberellin receptor of <i>Arabidopsis</i> . 2007 , 19, 1209-20	336
1535	Characterization of the VIER F-BOX PROTEINE genes from <i>Arabidopsis</i> reveals their importance for plant growth and development. 2007 , 19, 1163-78	44
1534	CYP703 is an ancient cytochrome P450 in land plants catalyzing in-chain hydroxylation of lauric acid to provide building blocks for sporopollenin synthesis in pollen. 2007 , 19, 1473-87	271
1533	BIN4, a novel component of the plant DNA topoisomerase VI complex, is required for endoreduplication in <i>Arabidopsis</i> . 2007 , 19, 3655-68	90
1532	Characterization of <i>Arabidopsis thaliana</i> SufE2 and SufE3: functions in chloroplast iron-sulfur cluster assembly and Nad synthesis. 2007 , 282, 18254-18264	64
1531	KEEP ON GOING, a RING E3 ligase essential for <i>Arabidopsis</i> growth and development, is involved in abscisic acid signaling. 2006 , 18, 3415-28	302

1530	MONODEHYDROASCORBATE REDUCTASE4 is required for seed storage oil hydrolysis and postgerminative growth in Arabidopsis. 2007 , 19, 1376-87	88
1529	Role of the MPN subunits in COP9 signalosome assembly and activity, and their regulatory interaction with Arabidopsis Cullin3-based E3 ligases. 2007 , 19, 564-81	107
1528	Repression of the LEAFY COTYLEDON 1/B3 regulatory network in plant embryo development by VP1/ABSCISIC ACID INSENSITIVE 3-LIKE B3 genes. 2007 , 143, 902-11	179
1527	The Arabidopsis BEL1-LIKE HOMEODOMAIN proteins SAW1 and SAW2 act redundantly to regulate KNOX expression spatially in leaf margins. 2007 , 19, 2719-35	105
1526	Role of SVP in the control of flowering time by ambient temperature in Arabidopsis. 2007 , 21, 397-402	442
1525	A core subunit of the RNA-processing/degrading exosome specifically influences cuticular wax biosynthesis in Arabidopsis. 2007 , 19, 904-13	91
1524	VIM1, a methylcytosine-binding protein required for centromeric heterochromatinization. 2007 , 21, 267-77	143
1523	The acyltransferase GPAT5 is required for the synthesis of suberin in seed coat and root of Arabidopsis. 2007 , 19, 351-68	261
1522	PIC1, an ancient permease in Arabidopsis chloroplasts, mediates iron transport. 2007 , 19, 986-1006	196
1521	Arabidopsis Teosinte Branched1-like 1 regulates axillary bud outgrowth and is homologous to monocot Teosinte Branched1. 2007 , 48, 667-77	162
1520	CLASP localizes in two discrete patterns on cortical microtubules and is required for cell morphogenesis and cell division in Arabidopsis. 2007 , 120, 4416-25	97
1519	LITTLE NUCLEI genes affecting nuclear morphology in Arabidopsis thaliana. 2007 , 19, 2793-803	158
1518	Roles of Arabidopsis cyclin-dependent kinase C complexes in cauliflower mosaic virus infection, plant growth, and development. 2007 , 19, 1388-402	63
1517	The essential nature of sphingolipids in plants as revealed by the functional identification and characterization of the Arabidopsis LCB1 subunit of serine palmitoyltransferase. 2006 , 18, 3576-93	122
1516	Pollen lethality: a phenomenon in Arabidopsis RNA interference plants. 2007 , 145, 330-3	19
1515	The gene MACCHI-BOU 4/ENHANCER OF PINOID encodes a NPH3-like protein and reveals similarities between organogenesis and phototropism at the molecular level. 2007 , 134, 3849-59	81
1514	A family of microRNAs present in plants and animals. 2006 , 18, 3355-69	117
1513	An analysis of univalent segregation in meiotic mutants of Arabidopsis thaliana: a possible role for synaptonemal complex. 2007 , 175, 505-11	26

1512	Cell-fate switch of synergid to egg cell in <i>Arabidopsis eostre</i> mutant embryo sacs arises from misexpression of the BEL1-like homeodomain gene BLH1. 2007 , 19, 3578-92	184
1511	Ubiquitin lysine 63 chain forming ligases regulate apical dominance in <i>Arabidopsis</i> . 2007 , 19, 1898-911	80
1510	Mutational loss of the prohibitin AtPHB3 results in an extreme constitutive ethylene response phenotype coupled with partial loss of ethylene-inducible gene expression in <i>Arabidopsis</i> seedlings. 2007 , 58, 2237-48	44
1509	Nitrite acts as a transcriptome signal at micromolar concentrations in <i>Arabidopsis</i> roots. 2007 , 145, 1735-45	111
1508	SAD2, an importin -like protein, is required for UV-B response in <i>Arabidopsis</i> by mediating MYB4 nuclear trafficking. 2007 , 19, 3805-18	131
1507	NUCLEAR PORE ANCHOR, the <i>Arabidopsis</i> homolog of Tpr/Mlp1/Mlp2/megator, is involved in mRNA export and SUMO homeostasis and affects diverse aspects of plant development. 2007 , 19, 1537-48	148
1506	Phytochemical genomics in <i>Arabidopsis thaliana</i> : A case study for functional identification of flavonoid biosynthesis genes. 2007 , 79, 811-823	76
1505	Vitamer levels, stress response, enzyme activity, and gene regulation of <i>Arabidopsis</i> lines mutant in the pyridoxine/pyridoxamine 5'-phosphate oxidase (PDX3) and the pyridoxal kinase (SOS4) genes involved in the vitamin B6 salvage pathway. 2007 , 145, 985-96	60
1504	Characterization of AtALMT1 expression in aluminum-inducible malate release and its role for rhizotoxic stress tolerance in <i>Arabidopsis</i> . 2007 , 145, 843-52	150
1503	The phosphatase laforin crosses evolutionary boundaries and links carbohydrate metabolism to neuronal disease. 2007 , 178, 477-88	127
1502	Hawaiian skirt: an F-box gene that regulates organ fusion and growth in <i>Arabidopsis</i> . 2007 , 144, 1370-82	64
1501	Temporary expression of the TAF10 gene and its requirement for normal development of <i>Arabidopsis thaliana</i> . 2007 , 48, 134-46	18
1500	Using genomics to study legume seed development. 2007 , 144, 562-74	121
1499	The HECATE genes regulate female reproductive tract development in <i>Arabidopsis thaliana</i> . 2007 , 134, 3593-601	177
1498	Two closely related RecQ helicases have antagonistic roles in homologous recombination and DNA repair in <i>Arabidopsis thaliana</i> . 2007 , 104, 18836-41	115
1497	Genetic evidence for three unique components in primary cell-wall cellulose synthase complexes in <i>Arabidopsis</i> . 2007 , 104, 15566-71	403
1496	NPY1, a BTB-NPH3-like protein, plays a critical role in auxin-regulated organogenesis in <i>Arabidopsis</i> . 2007 , 104, 18825-9	111
1495	<i>Arabidopsis</i> mutants lacking long chain base phosphate lyase are fumonisin-sensitive and accumulate trihydroxy-18:1 long chain base phosphate. 2007 , 282, 28195-206	60

1494	Ethylene-induced Arabidopsis hypocotyl elongation is dependent on but not mediated by gibberellins. 2007 , 58, 4269-81	55
1493	The CCCH-type zinc finger proteins AtSZF1 and AtSZF2 regulate salt stress responses in Arabidopsis. 2007 , 48, 1148-58	144
1492	A tale of two templates: automatically resolving double traces has many applications, including efficient PCR-based elucidation of alternative splices. 2007 , 17, 212-8	12
1491	GIGANTEA acts in blue light signaling and has biochemically separable roles in circadian clock and flowering time regulation. 2007 , 143, 473-86	130
1490	The Arabidopsis DESPERADO/AtWBC11 transporter is required for cutin and wax secretion. 2007 , 145, 1345-60	239
1489	New gateways to discovery. 2007 , 145, 1100-9	44
1488	Components of the Arabidopsis mRNA decapping complex are required for early seedling development. 2007 , 19, 1549-64	99
1487	Nitrate efflux at the root plasma membrane: identification of an Arabidopsis excretion transporter. 2007 , 19, 3760-77	138
1486	Inactivation of thioredoxin reductases reveals a complex interplay between thioredoxin and glutathione pathways in Arabidopsis development. 2007 , 19, 1851-65	196
1485	VPS9a, the common activator for two distinct types of Rab5 GTPases, is essential for the development of Arabidopsis thaliana. 2007 , 19, 3504-15	108
1484	Phytochrome-Interacting Factors. 81-105	4
1483	The iron-responsive element (IRE)/iron-regulatory protein 1 (IRP1)-cytosolic aconitase iron-regulatory switch does not operate in plants. 2007 , 405, 523-31	57
1482	Arabidopsis WRKY70 is required for full RPP4-mediated disease resistance and basal defense against Hyaloperonospora parasitica. 2007 , 20, 120-8	158
1481	TLP18.3, a novel thylakoid lumen protein regulating photosystem II repair cycle. 2007 , 406, 415-25	81
1480	Heterotrimeric G protein gamma subunits provide functional selectivity in Gbetagamma dimer signaling in Arabidopsis. 2007 , 19, 1235-50	143
1479	Participation of endomembrane cation/H ⁺ exchanger AtCHX20 in osmoregulation of guard cells. 2007 , 144, 82-93	83
1478	Dual lipid modification of Arabidopsis Ggamma-subunits is required for efficient plasma membrane targeting. 2007 , 143, 1119-31	66
1477	Somatic cytokinesis and pollen maturation in Arabidopsis depend on TPLATE, which has domains similar to coat proteins. 2006 , 18, 3502-18	90

1476	Unique, shared, and redundant roles for the Arabidopsis SWI/SNF chromatin remodeling ATPases BRAHMA and SPLAYED. 2007 , 19, 403-16	123
1475	Tandem gene arrays: a challenge for functional genomics. 2007 , 12, 203-10	23
1474	The role of the SCRAMBLED receptor-like kinase in patterning the Arabidopsis root epidermis. 2007 , 302, 118-31	96
1473	C2H2 zinc finger-SET histone methyltransferase is a plant-specific chromatin modifier. 2007 , 303, 259-69	56
1472	Combinations of WOX activities regulate tissue proliferation during Arabidopsis embryonic development. 2007 , 309, 306-16	120
1471	Distinct and overlapping roles of single-repeat MYB genes in root epidermal patterning. 2007 , 311, 566-78	129
1470	An efficient method for producing an indexed, insertional-mutant library in rice. 2007 , 89, 532-40	14
1469	Arabidopsis cytochrome P450 monooxygenase 71A13 catalyzes the conversion of indole-3-acetaldoxime in camalexin synthesis. 2007 , 19, 2039-52	259
1468	A systematic analysis of T-DNA insertion events in Magnaporthe oryzae. 2007 , 44, 1050-64	52
1467	The retromer protein VPS29 links cell polarity and organ initiation in plants. 2007 , 130, 1057-70	189
1466	Genome-wide high-resolution mapping of exosome substrates reveals hidden features in the Arabidopsis transcriptome. 2007 , 131, 1340-53	258
1465	RPK1 and TOAD2 are two receptor-like kinases redundantly required for arabidopsis embryonic pattern formation. 2007 , 12, 943-56	115
1464	The evolutionarily conserved Arabidopsis thaliana F-box protein AtFBP7 is required for efficient translation during temperature stress. 2007 , 392, 106-16	39
1463	A nucleosome interaction module is required for normal function of Arabidopsis thaliana BRAHMA. 2007 , 373, 240-50	44
1462	Redox regulation and antioxidative defence in Arabidopsis leaves viewed from a systems biology perspective. 2007 , 129, 229-48	33
1461	Transcript profiling in host-pathogen interactions. 2007 , 45, 329-69	136
1460	MIKC* MADS-protein complexes bind motifs enriched in the proximal region of late pollen-specific Arabidopsis promoters. 2007 , 143, 447-60	90
1459	RNA Silencing and Its Application in Functional Genomics. 2007 , 291-332	1

1458	The conserved cysteine-rich domain of a tesmin/TSO1-like protein binds zinc in vitro and TSO1 is required for both male and female fertility in <i>Arabidopsis thaliana</i> . 2007 , 58, 3657-70	48
1457	Regulation of telomerase in <i>Arabidopsis</i> by BT2, an apparent target of TELOMERASE ACTIVATOR1. 2007 , 19, 23-31	59
1456	T-DNA Insertion Mutants as a Resource for Rice Functional Genomics. 2007 , 181-221	10
1455	MYC2 differentially modulates diverse jasmonate-dependent functions in <i>Arabidopsis</i> . 2007 , 19, 2225-45	722
1454	Rice Functional Genomics. 2007 ,	
1453	MADS-complexes regulate transcriptome dynamics during pollen maturation. 2007 , 8, R249	82
1452	Genetic subtraction profiling identifies genes essential for <i>Arabidopsis</i> reproduction and reveals interaction between the female gametophyte and the maternal sporophyte. 2007 , 8, R204	113
1451	Transposable elements in fish functional genomics: technical challenges and perspectives. 2007 , 8 Suppl 1, S6	4
1450	Beyond Structural Genomics for Plant Science. 2007 , 77-161	13
1449	Transgenic Microalgae as Green Cell Factories. 2007 ,	11
1448	Ethylene Biosynthesis and Signaling: A Puzzle Yet to be Completed. 2007 , 125-145	1
1447	Tomato. 2007 , 1-125	13
1446	Efficient genome-wide mutagenesis of zebrafish genes by retroviral insertions. 2007 , 104, 12428-33	92
1445	A plant porphyria related to defects in plastid import of protochlorophyllide oxidoreductase A. 2007 , 104, 2019-23	38
1444	Localization of members of the gamma-glutamyl transpeptidase family identifies sites of glutathione and glutathione S-conjugate hydrolysis. 2007 , 144, 1715-32	81
1443	SCABP8/CBL10, a putative calcium sensor, interacts with the protein kinase SOS2 to protect <i>Arabidopsis</i> shoots from salt stress. 2007 , 19, 1415-31	361
1442	Induced mutants in common bean (<i>Phaseolus vulgaris</i>), and their potential use in nutrition quality breeding and gene discovery. 2007 , 55, 191-200	11
1441	AtMEK1 mediates stress-induced gene expression of CAT1 catalase by triggering H ₂ O ₂ production in <i>Arabidopsis</i> . 2007 , 58, 2969-81	140

1440	Arabidopsis FIERY1, XRN2, and XRN3 are endogenous RNA silencing suppressors. 2007 , 19, 3451-61	221
1439	The catalytically active tyrosine residues of both SPO11-1 and SPO11-2 are required for meiotic double-strand break induction in Arabidopsis. 2007 , 19, 3090-9	95
1438	Genetic analysis reveals functional redundancy and the major target genes of the Arabidopsis miR159 family. 2007 , 104, 16371-6	243
1437	A WD40 domain cyclophilin interacts with histone H3 and functions in gene repression and organogenesis in Arabidopsis. 2007 , 19, 2403-16	53
1436	Dual regulation role of GH3.5 in salicylic acid and auxin signaling during Arabidopsis-Pseudomonas syringae interaction. 2007 , 145, 450-64	219
1435	The Arabidopsis EIN3 binding F-Box proteins EBF1 and EBF2 have distinct but overlapping roles in ethylene signaling. 2007 , 19, 509-23	202
1434	Omics-based identification of Arabidopsis Myb transcription factors regulating aliphatic glucosinolate biosynthesis. 2007 , 104, 6478-83	556
1433	Multilevel interactions between ethylene and auxin in Arabidopsis roots. 2007 , 19, 2169-85	416
1432	Auxin synthesized by the YUCCA flavin monooxygenases is essential for embryogenesis and leaf formation in Arabidopsis. 2007 , 19, 2430-9	457
1431	Transcriptome Analysis of Plant Drought and Salt Stress Response. 2007 , 261-283	5
1430	Ethylene. 2007 ,	
1429	Salicylic acid-, jasmonic acid- and ethylene-mediated regulation of plant defense signaling. 2007 , 28, 55-83	48
1428	Phenotypic diversity and altered environmental plasticity in Arabidopsis thaliana with reduced Hsp90 levels. 2007 , 2, e648	121
1427	Towards the identification of flower-specific genes in Citrus spp. 2007 , 30, 761-768	3
1426	Biochemistry and molecular biology of Arabidopsis-aphid interactions. 2007 , 29, 871-83	114
1425	The future of metabolic phytochemistry: larger numbers of metabolites, higher resolution, greater understanding. 2007 , 68, 2861-80	89
1424	Characterization of a BAHD acyltransferase responsible for producing the green leaf volatile (Z)-3-hexen-1-yl acetate in Arabidopsis thaliana. 2007 , 49, 194-207	156
1423	A plasmodesmata-associated beta-1,3-glucanase in Arabidopsis. 2007 , 49, 669-82	222

1422	Arabidopsis ESK1 encodes a novel regulator of freezing tolerance. 2007 , 49, 786-99	120
1421	Identification of an arsenic tolerant double mutant with a thiol-mediated component and increased arsenic tolerance in phyA mutants. 2007 , 49, 1064-75	25
1420	QQT proteins colocalize with microtubules and are essential for early embryo development in Arabidopsis. 2007 , 50, 615-26	20
1419	Identification of a flavin-monooxygenase as the S-oxygenating enzyme in aliphatic glucosinolate biosynthesis in Arabidopsis. 2007 , 50, 902-10	186
1418	Genome-wide analysis of Agrobacterium T-DNA integration sites in the Arabidopsis genome generated under non-selective conditions. 2007 , 51, 779-91	159
1417	Two genes in Arabidopsis thaliana encoding GDP-L-galactose phosphorylase are required for ascorbate biosynthesis and seedling viability. 2007 , 52, 673-89	305
1416	Arabidopsis eIF3e is regulated by the COP9 signalosome and has an impact on development and protein translation. 2008 , 53, 300-11	33
1415	An Arabidopsis homolog of the bacterial peptidoglycan synthesis enzyme MurE has an essential role in chloroplast development. 2008 , 53, 924-34	65
1414	Root tip contact with low-phosphate media reprograms plant root architecture. 2007 , 39, 792-6	444
1413	An adapter ligation-mediated PCR method for high-throughput mapping of T-DNA inserts in the Arabidopsis genome. 2007 , 2, 2910-7	96
1412	SKB1-mediated symmetric dimethylation of histone H4R3 controls flowering time in Arabidopsis. 2007 , 26, 1934-41	98
1411	AtPRD1 is required for meiotic double strand break formation in Arabidopsis thaliana. 2007 , 26, 4126-37	77
1410	Interaction of shade avoidance and auxin responses: a role for two novel atypical bHLH proteins. 2007 , 26, 4756-67	163
1409	A novel plant gene essential for meiosis is related to the human CtIP and the yeast COM1/SAE2 gene. 2007 , 26, 5061-70	81
1408	The Arabidopsis Spontaneous Cell Death1 gene, encoding a zeta-carotene desaturase essential for carotenoid biosynthesis, is involved in chloroplast development, photoprotection and retrograde signalling. 2007 , 17, 458-70	91
1407	Disruption of phytoene desaturase gene results in albino and dwarf phenotypes in Arabidopsis by impairing chlorophyll, carotenoid, and gibberellin biosynthesis. 2007 , 17, 471-82	231
1406	Arabidopsis AtBECLIN 1/AtAtg6/AtVps30 is essential for pollen germination and plant development. 2007 , 17, 249-63	96
1405	Transcription factor control of asymmetric cell divisions that establish the stomatal lineage. 2007 , 445, 537-40	367

1404	A type III effector ADP-ribosylates RNA-binding proteins and quells plant immunity. 2007 , 447, 284-8	292
1403	Positive darwinian selection at the imprinted MEDEA locus in plants. 2007 , 448, 349-52	124
1402	A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence. 2007 , 448, 497-500	1260
1401	The JAZ family of repressors is the missing link in jasmonate signalling. 2007 , 448, 666-71	1576
1400	Emerging trends in the functional genomics of the abiotic stress response in crop plants. 2007 , 5, 361-80	181
1399	Bacterial non-host resistance: interactions of Arabidopsis with non-adapted Pseudomonas syringae strains. 2007 , 131, 448-61	44
1398	Biological systems of the host cell involved in Agrobacterium infection. 2007 , 9, 9-20	120
1397	Genome-wide analysis of T-DNA integration into the chromosomes of Magnaporthe oryzae. 2007 , 66, 371-82	69
1396	Involvement of the xyloglucan endotransglycosylase/hydrolases encoded by celery XTH1 and Arabidopsis XTH33 in the phloem response to aphids. 2007 , 30, 187-201	60
1395	Increased expression of the MYB-related transcription factor, PHR1, leads to enhanced phosphate uptake in Arabidopsis thaliana. 2007 , 30, 1499-512	204
1394	Development of an Arabidopsis thaliana-based bioassay for investigating seed colonization by mycotoxigenic Aspergillus species. 2007 , 56, 848-854	
1393	Metabolite profiling of plant carotenoids using the matrix-assisted laser desorption ionization time-of-flight mass spectrometry. 2007 , 49, 552-64	110
1392	A SNARE-protein has opposing functions in penetration resistance and defence signalling pathways. 2007 , 49, 302-12	117
1391	The Arabidopsis PsbO2 protein regulates dephosphorylation and turnover of the photosystem II reaction centre D1 protein. 2007 , 49, 528-39	87
1390	The ubiquitin pathway is required for innate immunity in Arabidopsis. 2007 , 49, 540-51	83
1389	ABA induction of miR159 controls transcript levels of two MYB factors during Arabidopsis seed germination. 2007 , 49, 592-606	567
1388	Dissecting the biosynthetic pathway for the bypass1 root-derived signal. 2007 , 49, 619-28	58
1387	A lysine-rich arabinogalactan protein in Arabidopsis is essential for plant growth and development, including cell division and expansion. 2007 , 49, 629-40	75

1386	Arabidopsis thaliana nicotinate/nicotinamide mononucleotide adenylyltransferase (AtNMNAT) is required for pollen tube growth. 2007 , 49, 694-703	46
1385	EDM2 is required for RPP7-dependent disease resistance in Arabidopsis and affects RPP7 transcript levels. 2007 , 49, 829-39	82
1384	Non-random distribution of T-DNA insertions at various levels of the genome hierarchy as revealed by analyzing 13 804 T-DNA flanking sequences from an enhancer-trap mutant library. 2007 , 49, 947-59	98
1383	Characterization of the extracellular gamma-glutamyl transpeptidases, GGT1 and GGT2, in Arabidopsis. 2007 , 49, 865-77	99
1382	Nicotinamidase participates in the salvage pathway of NAD biosynthesis in Arabidopsis. 2007 , 49, 1020-9	77
1381	Structural and functional characterization of ferredoxin-NADP+-oxidoreductase using knock-out mutants of Arabidopsis. 2007 , 49, 1041-52	81
1380	Alanine aminotransferase catalyses the breakdown of alanine after hypoxia in Arabidopsis thaliana. 2007 , 49, 1108-21	186
1379	A mutation in NLA, which encodes a RING-type ubiquitin ligase, disrupts the adaptability of Arabidopsis to nitrogen limitation. 2007 , 50, 320-37	218
1378	Structural analogs of ABA reveal novel features of ABA perception and signaling in Arabidopsis. 2007 , 50, 414-28	76
1377	Overexpression of CRK13, an Arabidopsis cysteine-rich receptor-like kinase, results in enhanced resistance to Pseudomonas syringae. 2007 , 50, 488-99	101
1376	Differential regulation of closely related R2R3-MYB transcription factors controls flavonol accumulation in different parts of the Arabidopsis thaliana seedling. 2007 , 50, 660-77	669
1375	Receptor-like protein kinase 2 (RPK 2) is a novel factor controlling anther development in Arabidopsis thaliana. 2007 , 50, 751-66	147
1374	WRINKLED1 specifies the regulatory action of LEAFY COTYLEDON2 towards fatty acid metabolism during seed maturation in Arabidopsis. 2007 , 50, 825-38	319
1373	The Arabidopsis ABA-deficient mutant aba4 demonstrates that the major route for stress-induced ABA accumulation is via neoxanthin isomers. 2007 , 50, 810-24	173
1372	Members of the PHO1 gene family show limited functional redundancy in phosphate transfer to the shoot, and are regulated by phosphate deficiency via distinct pathways. 2007 , 50, 982-94	137
1371	Arabidopsis GH3-LIKE DEFENSE GENE 1 is required for accumulation of salicylic acid, activation of defense responses and resistance to Pseudomonas syringae. 2007 , 51, 234-46	96
1370	Induction of phosphatidylinositol 3-kinase-mediated endocytosis by salt stress leads to intracellular production of reactive oxygen species and salt tolerance. 2007 , 51, 185-97	224
1369	A role for the AtMTP11 gene of Arabidopsis in manganese transport and tolerance. 2007 , 51, 198-210	188

1368	Mutations in CHLOROPLAST RNA BINDING provide evidence for the involvement of the chloroplast in the regulation of the circadian clock in Arabidopsis. 2007 , 51, 551-62	47
1367	Members of the plant NIMA-related kinases are involved in organ development and vascularization in poplar, Arabidopsis and rice. 2007 , 51, 575-88	34
1366	Salt tolerance (STO), a stress-related protein, has a major role in light signalling. 2007 , 51, 563-74	104
1365	Characterization of seed-specific benzoyloxyglucosinolate mutations in Arabidopsis thaliana. 2007 , 51, 1062-76	84
1364	Toc64/OEP64 is not essential for the efficient import of proteins into chloroplasts in Arabidopsis thaliana. 2007 , 52, 53-68	66
1363	The Arabidopsis AtPNG1 gene encodes a peptide: N-glycanase. 2007 , 52, 94-104	38
1362	AtDUR3 represents the major transporter for high-affinity urea transport across the plasma membrane of nitrogen-deficient Arabidopsis roots. 2007 , 52, 30-40	82
1361	Function of plastidial pyruvate kinases in seeds of Arabidopsis thaliana. 2007 , 52, 405-19	100
1360	PRMT11: a new Arabidopsis MBD7 protein partner with arginine methyltransferase activity. 2007 , 52, 210-22	31
1359	Phosphorylation of threonine 161 in plant cyclin-dependent kinase A is required for cell division by activation of its associated kinase. 2007 , 52, 435-48	34
1358	The rice nuclear gene, VIRESCENT 2, is essential for chloroplast development and encodes a novel type of guanylate kinase targeted to plastids and mitochondria. 2007 , 52, 512-27	96
1357	Characterization of Arabidopsis ABCG11/WBC11, an ATP binding cassette (ABC) transporter that is required for cuticular lipid secretion. 2007 , 52, 485-98	296
1356	Conditional oxidative stress responses in the Arabidopsis photorespiratory mutant cat2 demonstrate that redox state is a key modulator of daylength-dependent gene expression, and define photoperiod as a crucial factor in the regulation of H ₂ O ₂ -induced cell death. 2007 , 52, 640-57	327
1355	The Arabidopsis TALE homeobox gene ATH1 controls floral competency through positive regulation of FLC. 2007 , 52, 899-913	45
1354	Genetic characterization reveals no role for the reported ABA receptor, GCR2, in ABA control of seed germination and early seedling development in Arabidopsis. 2007 , 52, 1001-13	98
1353	The Arabidopsis her1 mutant implicates GABA in E-2-hexenal responsiveness. 2008 , 53, 197-213	72
1352	Arabidopsis extra-large G proteins (XLGs) regulate root morphogenesis. 2008 , 53, 248-63	83
1351	Subcellular co-localization of Arabidopsis RTE1 and ETR1 supports a regulatory role for RTE1 in ETR1 ethylene signaling. 2008 , 53, 275-86	110

1350	Arabidopsis endo-1,4-beta-glucanases are involved in the formation of root syncytia induced by <i>Heterodera schachtii</i> . 2008 , 53, 336-51	54
1349	Loss of the vacuolar cation channel, AtTPC1, does not impair Ca ²⁺ signals induced by abiotic and biotic stresses. 2008 , 53, 287-99	147
1348	ARL2, ARG1 and PIN3 define a gravity signal transduction pathway in root statocytes. 2008 , 53, 380-92	140
1347	The gibberellin biosynthetic genes AtGA20ox1 and AtGA20ox2 act, partially redundantly, to promote growth and development throughout the Arabidopsis life cycle. 2008 , 53, 488-504	258
1346	Phytochromes and cryptochromes regulate the differential growth of Arabidopsis hypocotyls in both a PGP19-dependent and a PGP19-independent manner. 2008 , 53, 516-29	68
1345	Histone H2A.Z and homologues of components of the SWR1 complex are required to control immunity in Arabidopsis. 2008 , 53, 475-87	166
1344	The AT-hook-containing proteins SOB3/AHL29 and ESC/AHL27 are negative modulators of hypocotyl growth in Arabidopsis. 2008 , 54, 1-14	54
1343	GASSHO1 and GASSHO2 encoding a putative leucine-rich repeat transmembrane-type receptor kinase are essential for the normal development of the epidermal surface in Arabidopsis embryos. 2008 , 54, 30-42	81
1342	The use of the zinc-fluorophore, Zinpyr-1, in the study of zinc homeostasis in Arabidopsis roots. 2007 , 174, 39-45	95
1341	The Chloroplast Outer Envelope Membrane: The Edge of Light and Excitement. 2007 , 49, 1100-1111	37
1340	GAMT2 Encodes a Methyltransferase of Gibberellic Acid That is Involved in Seed Maturation and Germination in Arabidopsis. 2007 , 49, 368-381	14
1339	Molecular characterisation of the STRUBBELIG-RECEPTOR FAMILY of genes encoding putative leucine-rich repeat receptor-like kinases in Arabidopsis thaliana. 2007 , 7, 16	59
1338	Discovery of chemically induced mutations in rice by TILLING. 2007 , 7, 19	312
1337	Functional analysis of Arabidopsis WRKY25 transcription factor in plant defense against <i>Pseudomonas syringae</i> . 2007 , 7, 2	153
1336	Mutations in a plastid-localized elongation factor G alter early stages of plastid development in Arabidopsis thaliana. 2007 , 7, 37	38
1335	Combining classifiers to predict gene function in Arabidopsis thaliana using large-scale gene expression measurements. 2007 , 8, 358	26
1334	Sequence-indexed mutations in maize using the UniformMu transposon-tagging population. 2007 , 8, 116	91
1333	Effect of chromatin upon Agrobacterium T-DNA integration and transgene expression. 2007 , 1769, 410-21	46

1332	A Novel ROP/RAC effector links cell polarity, root-meristem maintenance, and vesicle trafficking. 2007 , 17, 947-52	193
1331	The NTT gene is required for transmitting-tract development in carpels of Arabidopsis thaliana. 2007 , 17, 1101-8	115
1330	Tolerance of dividing cells to replication stress in UVB-irradiated Arabidopsis roots: requirements for DNA translesion polymerases eta and zeta. 2007 , 6, 1341-58	63
1329	Disease induction by human microbial pathogens in plant-model systems: potential, problems and prospects. 2007 , 12, 167-73	18
1328	T-DNA-induced mutations in transgenic plants. 2007 , 43, 1-11	8
1327	Mosaic expression pattern of the nptII gene in transgenic tobacco plants Nu 21. 2007 , 43, 780-790	7
1326	A guide to choosing vectors for transformation of the plastid genome of higher plants. 2007 , 145, 1201-10	49
1325	Construction and behavior of engineered minichromosomes in maize. 2007 , 104, 8924-9	125
1324	In the grip of algal genomics. 2007 , 616, 54-76	19
1323	A mutation in the AtPRP4 splicing factor gene suppresses seed development in Arabidopsis. 2007 , 9, 447-52	8
1322	Microhomologies between T-DNA ends and target sites often occur in inverted orientation and may be responsible for the high frequency of T-DNA-associated inversions. 2007 , 26, 617-30	12
1321	The Arabidopsis thaliana AT PRP39-1 gene, encoding a tetratricopeptide repeat protein with similarity to the yeast pre-mRNA processing protein PRP39, affects flowering time. 2007 , 26, 1357-66	35
1320	LeERF1 positively modulated ethylene triple response on etiolated seedling, plant development and fruit ripening and softening in tomato. 2007 , 26, 1999-2008	95
1319	Expression and functional analysis of ZmDWF4, an ortholog of Arabidopsis DWF4 from maize (Zea mays L.). 2007 , 26, 2091-9	43
1318	Molecular Basis of the Ethylene Signaling and Response Pathway in Arabidopsis. 2007 , 26, 106-117	46
1317	The negative influence of N-mediated TMV resistance on yield in tobacco: linkage drag versus pleiotropy. 2007 , 115, 169-78	25
1316	Characterization of T-DNA insertion patterns in the genome of rice blast fungus Magnaporthe oryzae. 2007 , 51, 233-43	40
1315	Expression profiles as detailed snapshots of biological states. 2007 , 16, 399-403	2

1314	RNA interference for wheat functional gene analysis. 2007 , 16, 689-701	68
1313	Aconitase plays a role in regulating resistance to oxidative stress and cell death in Arabidopsis and Nicotiana benthamiana. 2007 , 63, 273-87	108
1312	Functionality of the beta/six site-specific recombination system in tobacco and Arabidopsis: a novel tool for genetic engineering of plant genomes. 2007 , 63, 545-56	20
1311	A large-scale collection of phenotypic data describing an insertional mutant population to facilitate functional analysis of rice genes. 2007 , 63, 625-35	121
1310	IBR3, a novel peroxisomal acyl-CoA dehydrogenase-like protein required for indole-3-butyric acid response. 2007 , 64, 59-72	76
1309	Functional analysis of Arabidopsis genes involved in mitochondrial iron-sulfur cluster assembly. 2007 , 64, 225-40	44
1308	Molecular characterization of two Arabidopsis thaliana glycosyltransferase mutants, rra1 and rra2, which have a reduced residual arabinose content in a polymer tightly associated with the cellulosic wall residue. 2007 , 64, 439-51	76
1307	The HvNAC6 transcription factor: a positive regulator of penetration resistance in barley and Arabidopsis. 2007 , 65, 137-50	117
1306	DORNROSCHEN-LIKE, an AP2 gene, is necessary for stamen emergence in Arabidopsis. 2007 , 65, 219-32	39
1305	Transcript profiling of the hypomethylated hog1 mutant of Arabidopsis. 2007 , 65, 571-86	23
1304	The bHLH-type transcription factor AtAIB positively regulates ABA response in Arabidopsis. 2007 , 65, 655-65	87
1303	The WiscDsLox T-DNA collection: an arabidopsis community resource generated by using an improved high-throughput T-DNA sequencing pipeline. 2007 , 120, 157-65	120
1302	N-terminal domain including conserved flg22 is required for flagellin-induced hypersensitive cell death in Arabidopsis thaliana. 2007 , 73, 281-285	15
1301	Arabidopsis thaliana myosin XIX is involved in root hair as well as trichome morphogenesis on stems and leaves. 2007 , 230, 193-202	86
1300	Segregation distortion in Arabidopsis gametophytic factor 1 (gfa1) mutants is caused by a deficiency of an essential RNA splicing factor. 2007 , 20, 87-97	23
1299	pgd1, an Arabidopsis thaliana deletion mutant, is defective in pollen germination. 2007 , 20, 137-149	4
1298	Alpha-tocopherol may influence cellular signaling by modulating jasmonic acid levels in plants. 2007 , 225, 681-91	82
1297	AtPTR3, a wound-induced peptide transporter needed for defence against virulent bacterial pathogens in Arabidopsis. 2007 , 225, 1431-45	56

1296	Arabidopsis thaliana GH3.9 influences primary root growth. 2007 , 226, 21-34	66
1295	Expression and localization of AtAGP18, a lysine-rich arabinogalactan-protein in Arabidopsis. 2007 , 226, 169-79	24
1294	The amino acid permease AAP8 is important for early seed development in Arabidopsis thaliana. 2007 , 226, 805-13	101
1293	Nitration of gamma-tocopherol in plant tissues. 2007 , 226, 1311-22	37
1292	Osmotically stress-regulated the expression of glutathione peroxidase 3 in Arabidopsis. 2007 , 52, 127-130	9
1291	Characterization of three Arabidopsis AP2/EREBP family transcription factors involved in ABA sensitivity, freeze and salt tolerance. 2007 , 52, 1746-1753	2
1290	AtERF38 (At2g35700), an AP2/ERF family transcription factor gene from Arabidopsis thaliana, is expressed in specific cell types of roots, stems and seeds that undergo suberization. 2008 , 46, 1051-61	23
1289	Pantothenate synthetase is essential but not limiting for pantothenate biosynthesis in Arabidopsis. 2008 , 66, 1-14	15
1288	Altered gravitropic response, amyloplast sedimentation and circumnutation in the Arabidopsis shoot gravitropism 5 mutant are associated with reduced starch levels. 2008 , 67, 57-69	45
1287	The genetic locus At1g73660 encodes a putative MAPKKK and negatively regulates salt tolerance in Arabidopsis. 2008 , 67, 125-34	39
1286	The microRNA regulated SBP-box genes SPL9 and SPL15 control shoot maturation in Arabidopsis. 2008 , 67, 183-95	347
1285	Stay-green protein, defective in Mendel's green cotyledon mutant, acts independent and upstream of pheophorbide a oxygenase in the chlorophyll catabolic pathway. 2008 , 67, 243-56	78
1284	Characterisation of 3' transgene insertion site and derived mRNAs in MON810 YieldGard maize. 2008 , 67, 271-81	41
1283	A phosphatidylinositol phosphate-specific myo-inositol polyphosphate 5-phosphatase required for seedling growth. 2008 , 67, 375-88	26
1282	Sucrose induction of Arabidopsis miR398 represses two Cu/Zn superoxide dismutases. 2008 , 67, 403-17	200
1281	A small plant-specific protein family of ABI five binding proteins (AFPs) regulates stress response in germinating Arabidopsis seeds and seedlings. 2008 , 67, 643-58	101
1280	Arabidopsis DEMETER-LIKE proteins DML2 and DML3 are required for appropriate distribution of DNA methylation marks. 2008 , 67, 671-81	171
1279	The Arabidopsis thaliana carboxyl-terminal domain phosphatase-like 2 regulates plant growth, stress and auxin responses. 2008 , 67, 683-97	39

1278	An ArabidopsisSUPERMAN-like gene,AtZFP12, Expressed at Shoot Organ Boundaries Suppresses cell Growth.. 2008 , 51, 413-417	2
1277	The Fruits of Tropical Plant Genomics. 2008 , 1, 3-19	7
1276	Morphological characterisation and genetic analysis of a bi-pistil mutant (bip) in Medicago truncatula Gaertn.. 2008 , 21, 133-141	8
1275	New insights on sucrose metabolism: evidence for an active A/N-Inv in chloroplasts uncovers a novel component of the intracellular carbon trafficking. 2008 , 227, 795-807	66
1274	Redirection of the phenylpropanoid pathway to feruloyl malate in Arabidopsis mutants deficient for cinnamoyl-CoA reductase 1. 2008 , 227, 943-56	144
1273	The role of JAR1 in Jasmonoyl-L: -isoleucine production during Arabidopsis wound response. 2008 , 227, 1221-32	172
1272	PRD, an Arabidopsis AINTEGUMENTA-like gene, is involved in root architectural changes in response to phosphate starvation. 2008 , 228, 511-22	34
1271	The long-term response to fluctuating light quality is an important and distinct light acclimation mechanism that supports survival of Arabidopsis thaliana under low light conditions. 2008 , 228, 573-87	70
1270	Transcript profiling demonstrates absence of dosage compensation in Arabidopsis following loss of a single RPL23a paralog. 2008 , 228, 627-40	19
1269	SAZ, a new SUPERMAN-like protein, negatively regulates a subset of ABA-responsive genes in Arabidopsis. 2008 , 279, 183-92	22
1268	Homologous recombination-mediated cloning and manipulation of genomic DNA regions using Gateway and recombineering systems. 2008 , 8, 88	18
1267	Gene duplications, robustness and evolutionary innovations. 2008 , 30, 367-73	88
1266	Biochemical and genetic characterization of Arabidopsis flavanone 3beta-hydroxylase. 2008 , 46, 833-43	66
1265	ASY1 coordinates early events in the plant meiotic recombination pathway. 2008 , 120, 302-12	36
1264	Evolutionary linkage between eukaryotic cytokinesis and chloroplast division by dynamin proteins. 2008 , 105, 15202-7	67
1263	Light regulation of CaS, a novel phosphoprotein in the thylakoid membrane of Arabidopsis thaliana. 2008 , 275, 1767-77	121
1262	Dual control of nuclear EIN3 by bifurcate MAPK cascades in C2H4 signalling. 2008 , 451, 789-95	392
1261	T-DNA-mediated transfer of Agrobacterium tumefaciens chromosomal DNA into plants. 2008 , 26, 1015-7	51

1260	Enzymatic, but not non-enzymatic, 1O ₂ -mediated peroxidation of polyunsaturated fatty acids forms part of the EXECUTER1-dependent stress response program in the flu mutant of <i>Arabidopsis thaliana</i> . 2008 , 54, 236-48	103
1259	Identification of a novel cis-regulatory element for UV-B-induced transcription in <i>Arabidopsis</i> . 2008 , 54, 402-14	47
1258	Deletion of the chloroplast-localized AtTerC gene product in <i>Arabidopsis thaliana</i> leads to loss of the thylakoid membrane and to seedling lethality. 2008 , 55, 428-42	25
1257	A role for inositol hexakisphosphate in the maintenance of basal resistance to plant pathogens. 2008 , 56, 638-52	112
1256	CLB19, a pentatricopeptide repeat protein required for editing of rpoA and clpP chloroplast transcripts. 2008 , 56, 590-602	212
1255	Overexpression of the non-canonical Aux/IAA genes causes auxin-related aberrant phenotypes in <i>Arabidopsis</i> . 2008 , 133, 397-405	79
1254	Sterol-dependent endocytosis mediates post-cytokinetic acquisition of PIN2 auxin efflux carrier polarity. 2008 , 10, 237-44	268
1253	Whole-plant mineral partitioning throughout the life cycle in <i>Arabidopsis thaliana</i> ecotypes Columbia, Landsberg erecta, Cape Verde Islands, and the mutant line ysl1ysl3. 2008 , 177, 389-405	119
1252	An illustrated gardener's guide to transgenic <i>Arabidopsis</i> field experiments. 2008 , 180, 545-555	17
1251	The proton pump interactor (Ppi) gene family of <i>Arabidopsis thaliana</i> : expression pattern of Ppi1 and characterisation of knockout mutants for Ppi1 and 2. 2008 , 10, 237-49	14
1250	The plant WNK gene family and regulation of flowering time in <i>Arabidopsis</i> . 2008 , 10, 548-62	66
1249	Transcriptional regulation of plant senescence: from functional genomics to systems biology. 2008 , 10 Suppl 1, 99-109	18
1248	<i>Agrobacterium</i> -mediated transformation of the temperate grass <i>Brachypodium distachyon</i> (genotype Bd21) for T-DNA insertional mutagenesis. 2008 , 6, 236-45	99
1247	Immune subversion by chromatin manipulation: a 'new face' of host-bacterial pathogen interaction. 2008 , 10, 1582-90	15
1246	Relative contribution of AtHAK5 and AtAKT1 to K ⁺ uptake in the high-affinity range of concentrations. 2008 , 134, 598-608	147
1245	Expression differences for genes involved in lignin, glutathione and sulphate metabolism in response to cadmium in <i>Arabidopsis thaliana</i> and the related Zn/Cd-hyperaccumulator <i>Thlaspi caerulescens</i> . 2008 , 31, 301-24	253
1244	The disturbance of small RNA pathways enhanced abscisic acid response and multiple stress responses in <i>Arabidopsis</i> . 2008 , 31, 562-74	93
1243	The lack of alternative oxidase at low temperature leads to a disruption of the balance in carbon and nitrogen metabolism, and to an up-regulation of antioxidant defence systems in <i>Arabidopsis thaliana</i> leaves. 2008 , 31, 1190-202	108

1242	Identification and characterization of Cor413im proteins as novel components of the chloroplast inner envelope. 2008 , 31, 1470-83	46
1241	Responses of Brassica oleracea cultivars to infestation by the aphid Brevicoryne brassicae: an ecological and molecular approach. 2008 , 31, 1592-605	58
1240	Adequate phenylalanine synthesis mediated by G protein is critical for protection from UV radiation damage in young etiolated Arabidopsis thaliana seedlings. 2008 , 31, 1756-70	23
1239	Specialization of CDC27 function in the Arabidopsis thaliana anaphase-promoting complex (APC/C). 2008 , 53, 78-89	65
1238	The GRV2/RME-8 protein of Arabidopsis functions in the late endocytic pathway and is required for vacuolar membrane flow. 2008 , 53, 29-41	65
1237	Duplicated P5CS genes of Arabidopsis play distinct roles in stress regulation and developmental control of proline biosynthesis. 2008 , 53, 11-28	478
1236	26S proteasome regulatory particle mutants have increased oxidative stress tolerance. 2008 , 53, 102-14	129
1235	Armadillo repeat-containing kinesins and a NIMA-related kinase are required for epidermal-cell morphogenesis in Arabidopsis. 2008 , 53, 157-71	85
1234	Degradation of the cyclin-dependent kinase inhibitor KRP1 is regulated by two different ubiquitin E3 ligases. 2008 , 53, 705-16	87
1233	The first zygotic division in Arabidopsis requires de novo transcription of thymidylate kinase. 2008 , 53, 776-89	38
1232	Regulation of the anthocyanin biosynthetic pathway by the TTG1/bHLH/Myb transcriptional complex in Arabidopsis seedlings. 2008 , 53, 814-27	965
1231	LZF1, a HY5-regulated transcriptional factor, functions in Arabidopsis de-etiolation. 2008 , 54, 205-19	121
1230	The AAE14 gene encodes the Arabidopsis o-succinylbenzoyl-CoA ligase that is essential for phylloquinone synthesis and photosystem-I function. 2008 , 54, 272-83	52
1229	Large-scale insertional mutagenesis using the Tnt1 retrotransposon in the model legume Medicago truncatula. 2008 , 54, 335-47	344
1228	Reduced activity of Arabidopsis thaliana HMT2, a methionine biosynthetic enzyme, increases seed methionine content. 2008 , 54, 310-20	50
1227	AtMKK1 mediates ABA-induced CAT1 expression and H ₂ O ₂ production via AtMPK6-coupled signaling in Arabidopsis. 2008 , 54, 440-51	310
1226	A NIMA-related protein kinase suppresses ectopic outgrowth of epidermal cells through its kinase activity and the association with microtubules. 2008 , 54, 829-44	38
1225	Arabidopsis PCFS4, a homologue of yeast polyadenylation factor Pcf11p, regulates FCA alternative processing and promotes flowering time. 2008 , 54, 899-910	61

1224	Dual function of Arabidopsis glucan synthase-like genes GSL8 and GSL10 in male gametophyte development and plant growth. 2008 , 54, 911-23	80
1223	Generation of a 3D indexed Petunia insertion database for reverse genetics. 2008 , 54, 1105-14	38
1222	AGL23, a type I MADS-box gene that controls female gametophyte and embryo development in Arabidopsis. 2008 , 54, 1037-48	99
1221	The oomycete response gene LURP1 is required for defense against <i>Hyaloperonospora parasitica</i> in Arabidopsis thaliana. 2008 , 55, 53-64	42
1220	RLM3, a TIR domain encoding gene involved in broad-range immunity of Arabidopsis to necrotrophic fungal pathogens. 2008 , 55, 188-200	69
1219	CUL1 regulates TOC1 protein stability in the Arabidopsis circadian clock. 2008 , 55, 568-79	36
1218	AtCYP38 ensures early biogenesis, correct assembly and sustenance of photosystem II. 2008 , 55, 639-51	87
1217	The Arabidopsis sweetie mutant is affected in carbohydrate metabolism and defective in the control of growth, development and senescence. 2008 , 55, 665-86	53
1216	Ethylene signaling in Arabidopsis involves feedback regulation via the elaborate control of EBF2 expression by EIN3. 2008 , 55, 821-31	110
1215	Cysteine proteases XCP1 and XCP2 aid micro-autolysis within the intact central vacuole during xylogenesis in Arabidopsis roots. 2008 , 56, 303-315	118
1214	Vein patterning screens and the defectively organized tributaries mutants in Arabidopsis thaliana. 2008 , 56, 251-263	58
1213	Getting the most out of publicly available T-DNA insertion lines. 2008 , 56, 665-77	47
1212	Efficient virus-induced gene silencing in Arabidopsis using a 'one-step' TYMV-derived vector. 2008 , 56, 678-90	84
1211	The DDF1 transcriptional activator upregulates expression of a gibberellin-deactivating gene, GA2ox7, under high-salinity stress in Arabidopsis. 2008 , 56, 613-26	238
1210	Transcriptional regulation by an NAC (NAM-ATAF1,2-CUC2) transcription factor attenuates ABA signalling for efficient basal defence towards <i>Blumeria graminis</i> f. sp. hordei in Arabidopsis. 2008 , 56, 867-80	172
1209	Non-identical contributions of two membrane-bound cpSRP components, cpFtsY and Alb3, to thylakoid biogenesis. 2008 , 56, 1007-17	42
1208	Identification, cloning and characterization of sis7 and sis10 sugar-insensitive mutants of Arabidopsis. 2008 , 8, 104	23
1207	RACK1 genes regulate plant development with unequal genetic redundancy in Arabidopsis. 2008 , 8, 108	58

1206	ESKIMO1 is a key gene involved in water economy as well as cold acclimation and salt tolerance. 2008 , 8, 125	51
1205	Overexpression of mtDNA-associated AtWhy2 compromises mitochondrial function. 2008 , 8, 42	33
1204	Articulation of three core metabolic processes in Arabidopsis: fatty acid biosynthesis, leucine catabolism and starch metabolism. 2008 , 8, 76	65
1203	The plant Apolipoprotein D ortholog protects Arabidopsis against oxidative stress. 2008 , 8, 86	80
1202	AtGRX4, an Arabidopsis chloroplastic monothiol glutaredoxin, is able to suppress yeast grx5 mutant phenotypes and respond to oxidative stress. 2008 , 582, 848-54	43
1201	NADPH-dependent thioredoxin reductase and 2-Cys peroxiredoxins are needed for the protection of Mg-protoporphyrin monomethyl ester cyclase. 2008 , 582, 2773-8	79
1200	A conserved DYW domain of the pentatricopeptide repeat protein possesses a novel endoribonuclease activity. 2008 , 582, 4163-8	66
1199	The peroxin loss-of-function mutation abstinence by mutual consent disrupts male-female gametophyte recognition. 2008 , 18, 63-8	108
1198	SHOC1, an XPF endonuclease-related protein, is essential for the formation of class I meiotic crossovers. 2008 , 18, 1432-7	53
1197	T-DNA insertion, plasmid rescue and integration analysis in the model mycorrhizal fungus <i>Laccaria bicolor</i> . 2008 , 1, 258-69	30
1196	Characterization of Arabidopsis calreticulin mutants in response to calcium and salinity stresses. 2008 , 18, 1219-1224	7
1195	Functional analysis of lactate dehydrogenase during hypoxic stress in Arabidopsis. 2008 , 35, 131-140	35
1194	Trehalose-6-phosphate synthase/phosphatase regulates cell shape and plant architecture in Arabidopsis. 2008 , 146, 97-107	101
1193	PlantDB - a versatile database for managing plant research. 2008 , 4, 1	49
1192	RNA Interference. 207-225	
1191	TOPLESS mediates auxin-dependent transcriptional repression during Arabidopsis embryogenesis. <i>Science</i> , 2008 , 319, 1384-6	33-3 547
1190	Mutagenesis and High-Throughput Functional Genomics in Cereal Crops: Current Status. 2008 , 98, 357-414	3
1189	UTILdb, a <i>Pisum sativum</i> in silico forward and reverse genetics tool. 2008 , 9, R43	122

1188	Plant Growth Signaling. 2008 ,	2
1187	TOR Signaling in Plants. 2007 , 243-259	
1186	Endophytic actinobacteria induce defense pathways in Arabidopsis thaliana. 2008 , 21, 208-18	256
1185	Agrobacterium: From Biology to Biotechnology. 2008 ,	34
1184	Agrobacterium Tumefaciens-Mediated Transformation: Patterns of T-Dna Integration Into the Host Genome. 2008 , 441-481	11
1183	PPR336 is associated with polysomes in plant mitochondria. 2008 , 375, 626-36	60
1182	COP1 and ELF3 control circadian function and photoperiodic flowering by regulating GI stability. 2008 , 32, 617-30	273
1181	Unexpected silencing effects from T-DNA tags in Arabidopsis. 2008 , 13, 4-6	88
1180	Identifying essential genes in Arabidopsis thaliana. 2008 , 13, 483-91	176
1179	Ethylene biosynthesis and signaling in rice. 2008 , 175, 32-42	82
1178	The dual-and-opposing-effect of ethylene on the negative gravitropism of Arabidopsis inflorescence stem and light-grown hypocotyls. 2008 , 175, 71-86	11
1177	Melon, an alternative model plant for elucidating fruit ripening. 2008 , 175, 121-129	47
1176	Ethylene-regulated transcription and crosstalk with jasmonic acid. 2008 , 175, 18-23	29
1175	Mapping of the phosphorylation sites on the phototropic signal transducer, NPH3. 2008 , 174, 626-633	32
1174	The Arabidopsis SDG4 contributes to the regulation of pollen tube growth by methylation of histone H3 lysines 4 and 36 in mature pollen. 2008 , 315, 355-68	80
1173	Systems approaches to identifying gene regulatory networks in plants. 2008 , 24, 81-103	85
1172	Comparative mutant analysis of Arabidopsis ABCC-type ABC transporters: AtMRP2 contributes to detoxification, vacuolar organic anion transport and chlorophyll degradation. 2008 , 49, 557-69	55
1171	The histone methyltransferase SDG8 regulates shoot branching in Arabidopsis. 2008 , 373, 659-64	75

1170	A complex containing PGRL1 and PGR5 is involved in the switch between linear and cyclic electron flow in Arabidopsis. 2008 , 132, 273-85	397
1169	The importance of being essential: EMBRYO-DEFECTIVE genes in Arabidopsis. 2008 , 331, 726-36	35
1168	Differential expression of WOX genes mediates apical-basal axis formation in the Arabidopsis embryo. 2008 , 14, 867-76	274
1167	A link between RNA metabolism and silencing affecting Arabidopsis development. 2008 , 14, 854-66	328
1166	The basic helix-loop-helix transcription factor PIF5 acts on ethylene biosynthesis and phytochrome signaling by distinct mechanisms. 2007 , 19, 3915-29	117
1165	Comparative genomics and reverse genetics analysis reveal indispensable functions of the serine acetyltransferase gene family in Arabidopsis. 2008 , 20, 2484-96	98
1164	Plant simple sequence repeats: distribution, variation, and effects on gene expression. 2008 , 51, 79-90	16
1163	Mutations in SUPPRESSOR OF VARIATION1, a factor required for normal chloroplast translation, suppress var2-mediated leaf variegation in Arabidopsis. 2008 , 20, 1786-804	111
1162	Arabidopsis stromal 70-kD heat shock proteins are essential for plant development and important for thermotolerance of germinating seeds. 2008 , 146, 1231-41	198
1161	T-DNA Tagging Lines. 2008 , 95-106	
1160	A membrane-tethered transcription factor defines a branch of the heat stress response in Arabidopsis thaliana. 2008 , 105, 16398-403	198
1159	Antagonistic jacalin-related lectins regulate the size of ER body-type beta-glucosidase complexes in Arabidopsis thaliana. 2008 , 49, 969-80	69
1158	EMB2473/MIRO1, an Arabidopsis Miro GTPase, is required for embryogenesis and influences mitochondrial morphology in pollen. 2008 , 20, 589-601	56
1157	High temperature-induced abscisic acid biosynthesis and its role in the inhibition of gibberellin action in Arabidopsis seeds. 2008 , 146, 1368-85	303
1156	Dual roles of the nuclear cap-binding complex and SERRATE in pre-mRNA splicing and microRNA processing in Arabidopsis thaliana. 2008 , 105, 8795-800	297
1155	Putrescine is involved in Arabidopsis freezing tolerance and cold acclimation by regulating abscisic acid levels in response to low temperature. 2008 , 148, 1094-105	269
1154	The Arabidopsis thaliana type I Isopentenyl Diphosphate Isomerases are targeted to multiple subcellular compartments and have overlapping functions in isoprenoid biosynthesis. 2008 , 20, 677-96	110
1153	Sphingolipid long-chain base hydroxylation is important for growth and regulation of sphingolipid content and composition in Arabidopsis. 2008 , 20, 1862-78	123

1152	Functions of chloroplastic adenylate kinases in Arabidopsis. 2008 , 146, 492-504	35
1151	MYB72 is required in early signaling steps of rhizobacteria-induced systemic resistance in Arabidopsis. 2008 , 146, 1293-304	206
1150	A distinct endosomal Ca ²⁺ /Mn ²⁺ pump affects root growth through the secretory process. 2008 , 147, 1675-89	87
1149	Arabidopsis mitogen-activated protein kinase kinases MKK1 and MKK2 have overlapping functions in defense signaling mediated by MEKK1, MPK4, and MKS1. 2008 , 148, 212-22	221
1148	Two Arabidopsis ADP-glucose pyrophosphorylase large subunits (APL1 and APL2) are catalytic. 2008 , 148, 65-76	69
1147	Two new clock proteins, LWD1 and LWD2, regulate Arabidopsis photoperiodic flowering. 2008 , 148, 948-59	79
1146	A naturally occurring mutation in an Arabidopsis accession affects a beta-D-galactosidase that increases the hydrophilic potential of rhamnogalacturonan I in seed mucilage. 2007 , 19, 3990-4006	103
1145	The Arabidopsis phytochrome-interacting factor PIF7, together with PIF3 and PIF4, regulates responses to prolonged red light by modulating phyB levels. 2008 , 20, 337-52	261
1144	Arabidopsis SCARs function interchangeably to meet actin-related protein 2/3 activation thresholds during morphogenesis. 2008 , 20, 995-1011	60
1143	The Arabidopsis Phosphatidylinositol Phosphate 5-Kinase PIP5K3 is a key regulator of root hair tip growth. 2008 , 20, 367-80	164
1142	The microtubule plus-end binding protein EB1 functions in root responses to touch and gravity signals in Arabidopsis. 2008 , 20, 396-410	74
1141	A SNARE complex unique to seed plants is required for protein storage vacuole biogenesis and seed development of Arabidopsis thaliana. 2008 , 20, 3006-21	156
1140	A mutant impaired in the production of plastome-encoded proteins uncovers a mechanism for the homeostasis of isoprenoid biosynthetic enzymes in Arabidopsis plastids. 2008 , 20, 1303-15	137
1139	Histone H2B monoubiquitination in the chromatin of FLOWERING LOCUS C regulates flowering time in Arabidopsis. 2008 , 20, 2586-602	176
1138	The ABC subfamily B auxin transporter AtABCB19 is involved in the inhibitory effects of N-1-naphthylphthalamic acid on the phototropic and gravitropic responses of Arabidopsis hypocotyls. 2008 , 49, 1250-5	71
1137	Role of Arabidopsis CHL27 protein for photosynthesis, chloroplast development and gene expression profiling. 2008 , 49, 1350-63	52
1136	Arabidopsis thaliana has a set of J proteins in the endoplasmic reticulum that are conserved from yeast to animals and plants. 2008 , 49, 1547-62	58
1135	NPY genes and AGC kinases define two key steps in auxin-mediated organogenesis in Arabidopsis. 2008 , 105, 21017-22	112

1134	L-galactono-1,4-lactone dehydrogenase is required for the accumulation of plant respiratory complex I. 2008 , 283, 32500-5	116
1133	The POK/AtVPS52 protein localizes to several distinct post-Golgi compartments in sporophytic and gametophytic cells. 2008 , 59, 3087-98	21
1132	Artificial trans-acting siRNAs confer consistent and effective gene silencing. 2008 , 147, 543-51	52
1131	Reduced V-ATPase activity in the trans-Golgi network causes oxylipin-dependent hypocotyl growth inhibition in Arabidopsis. 2008 , 20, 1088-100	102
1130	Root hair defective4 encodes a phosphatidylinositol-4-phosphate phosphatase required for proper root hair development in Arabidopsis thaliana. 2008 , 20, 381-95	127
1129	Degradation of sphingoid long-chain base 1-phosphates (LCB-1Ps): functional characterization and expression of AtDPL1 encoding LCB-1P lyase involved in the dehydration stress response in Arabidopsis. 2008 , 49, 1758-63	30
1128	An exocyst complex functions in plant cell growth in Arabidopsis and tobacco. 2008 , 20, 1330-45	213
1127	Transcriptional profiling of mature Arabidopsis trichomes reveals that NOECK encodes the MIXTA-like transcriptional regulator MYB106. 2008 , 148, 1583-602	163
1126	F-actin organization and pollen tube tip growth in Arabidopsis are dependent on the gametophyte-specific Armadillo repeat protein ARO1. 2008 , 20, 2798-814	31
1125	Degradation of a polyadenylated rRNA maturation by-product involves one of the three RRP6-like proteins in Arabidopsis thaliana. 2008 , 28, 3038-44	72
1124	SHORT HYPOCOTYL IN WHITE LIGHT1, a serine-arginine-aspartate-rich protein in Arabidopsis, acts as a negative regulator of photomorphogenic growth. 2008 , 147, 169-78	13
1123	Genetic analysis reveals that C19-GA 2-oxidation is a major gibberellin inactivation pathway in Arabidopsis. 2008 , 20, 2420-36	217
1122	Identification and characterization of Arabidopsis indole-3-butyric acid response mutants defective in novel peroxisomal enzymes. 2008 , 180, 237-51	104
1121	ChromDB: the chromatin database. 2008 , 36, D298-302	66
1120	A bifunctional locus (BIO3-BIO1) required for biotin biosynthesis in Arabidopsis. 2008 , 146, 60-73	39
1119	The Arabidopsis AtOPT3 protein functions in metal homeostasis and movement of iron to developing seeds. 2008 , 146, 589-601	177
1118	The MADS-domain transcriptional regulator AGAMOUS-LIKE15 promotes somatic embryo development in Arabidopsis and soybean. 2008 , 146, 1663-72	108
1117	A novel role for protein farnesylation in plant innate immunity. 2008 , 148, 348-57	50

1116	Arabidopsis transcriptome reveals control circuits regulating redox homeostasis and the role of an AP2 transcription factor. 2008 , 148, 2050-8	74
1115	Functional and physiological characterization of Arabidopsis INOSITOL TRANSPORTER1, a novel tonoplast-localized transporter for myo-inositol. 2008 , 20, 1073-87	62
1114	Calmodulin7 plays an important role as transcriptional regulator in Arabidopsis seedling development. 2008 , 20, 1747-59	91
1113	Medicago truncatula as a model for understanding plant interactions with other organisms, plant development and stress biology: past, present and future. 2008 , 35, 253-264	65
1112	Arabidopsis ATG6 is required to limit the pathogen-associated cell death response. 2008 , 4, 20-7	150
1111	Arabidopsis DDB1-CUL4 ASSOCIATED FACTOR1 forms a nuclear E3 ubiquitin ligase with DDB1 and CUL4 that is involved in multiple plant developmental processes. 2008 , 20, 1437-55	109
1110	Interaction of the WD40 domain of a myo-inositol polyphosphate 5-phosphatase with SnRK1 links inositol, sugar, and stress signaling. 2008 , 148, 1868-82	66
1109	NIP6;1 is a boric acid channel for preferential transport of boron to growing shoot tissues in Arabidopsis. 2008 , 20, 2860-75	225
1108	Role of Arabidopsis RAP2.4 in regulating light- and ethylene-mediated developmental processes and drought stress tolerance. 2008 , 1, 42-57	130
1107	Phylogeny, gene structures, and expression patterns of the ERF gene family in soybean (<i>Glycine max</i> L.). 2008 , 59, 4095-107	182
1106	The coenzyme a biosynthetic enzyme phosphopantetheine adenylyltransferase plays a crucial role in plant growth, salt/osmotic stress resistance, and seed lipid storage. 2008 , 148, 546-56	33
1105	Enhanced tolerance to oxidative stress in transgenic Arabidopsis plants expressing proteins of unknown function. 2008 , 148, 280-92	83
1104	Further evidence for the mandatory nature of polysaccharide debranching for the aggregation of semicrystalline starch and for overlapping functions of debranching enzymes in Arabidopsis leaves. 2008 , 148, 1309-23	68
1103	SCREAM/ICE1 and SCREAM2 specify three cell-state transitional steps leading to arabidopsis stomatal differentiation. 2008 , 20, 1775-85	338
1102	BSKs mediate signal transduction from the receptor kinase BRI1 in Arabidopsis. <i>Science</i> , 2008 , 321, 557-60, 3	456
1101	Plastid signals remodel light signaling networks and are essential for efficient chloroplast biogenesis in Arabidopsis. 2007 , 19, 3944-60	160
1100	New connections across pathways and cellular processes: industrialized mutant screening reveals novel associations between diverse phenotypes in Arabidopsis. 2008 , 146, 1482-500	70
1099	Amino acid sequence of bacterial microbe-associated molecular pattern flg22 is required for virulence. 2008 , 21, 1165-74	61

1098	Overexpression of AtMYB44 enhances stomatal closure to confer abiotic stress tolerance in transgenic Arabidopsis. 2008 , 146, 623-35	464
1097	The EPIP peptide of INFLORESCENCE DEFICIENT IN ABSCISSION is sufficient to induce abscission in arabidopsis through the receptor-like kinases HAESA and HAESA-LIKE2. 2008 , 20, 1805-17	216
1096	Prefoldin 6 is required for normal microtubule dynamics and organization in Arabidopsis. 2008 , 105, 18064-9	46
1095	Arginase-negative mutants of Arabidopsis exhibit increased nitric oxide signaling in root development. 2008 , 147, 1936-46	137
1094	The transcript and metabolite networks affected by the two clades of Arabidopsis glucosinolate biosynthesis regulators. 2008 , 148, 2021-49	155
1093	An Arabidopsis protein phosphorylated in response to microbial elicitation, AtPHOS32, is a substrate of MAP kinases 3 and 6. 2008 , 283, 10493-9	61
1092	Identification of the wax ester synthase/acyl-coenzyme A: diacylglycerol acyltransferase WSD1 required for stem wax ester biosynthesis in Arabidopsis. 2008 , 148, 97-107	238
1091	Two leucine-rich repeat receptor kinases mediate signaling, linking cell wall biosynthesis and ACC synthase in Arabidopsis. 2008 , 20, 3065-79	226
1090	Two distinct interacting classes of nuclear envelope-associated coiled-coil proteins are required for the tissue-specific nuclear envelope targeting of Arabidopsis RanGAP. 2008 , 20, 1639-51	85
1089	Knocking out cytosolic cysteine synthesis compromises the antioxidant capacity of the cytosol to maintain discrete concentrations of hydrogen peroxide in Arabidopsis. 2008 , 147, 562-72	73
1088	SRA-domain proteins required for DRM2-mediated de novo DNA methylation. 2008 , 4, e1000280	120
1087	An exocyst vesicle tethering complex in plants. 2008 , 20, 1188	2
1086	Allelic mutant series reveal distinct functions for Arabidopsis cycloartenol synthase 1 in cell viability and plastid biogenesis. 2008 , 105, 3163-8	80
1085	The high affinity K ⁺ transporter AtHAK5 plays a physiological role in planta at very low K ⁺ concentrations and provides a caesium uptake pathway in Arabidopsis. 2008 , 59, 595-607	215
1084	A genome-wide functional investigation into the roles of receptor-like proteins in Arabidopsis. 2008 , 147, 503-17	190
1083	The glycerophosphoryl diester phosphodiesterase-like proteins SHV3 and its homologs play important roles in cell wall organization. 2008 , 49, 1522-35	68
1082	Topoisomerase 3alpha and RMI1 suppress somatic crossovers and are essential for resolution of meiotic recombination intermediates in Arabidopsis thaliana. 2008 , 4, e1000285	70
1081	Double fertilization in Arabidopsis thaliana involves a polyspermy block on the egg but not the central cell. 2008 , 1, 611-9	57

1080	The MADS domain protein DIANA acts together with AGAMOUS-LIKE80 to specify the central cell in <i>Arabidopsis</i> ovules. 2008 , 20, 2088-101	97
1079	IMPa-4, an <i>Arabidopsis</i> importin alpha isoform, is preferentially involved in agrobacterium-mediated plant transformation. 2008 , 20, 2661-80	87
1078	Three SRA-domain methylcytosine-binding proteins cooperate to maintain global CpG methylation and epigenetic silencing in <i>Arabidopsis</i> . 2008 , 4, e1000156	126
1077	A 125 kDa RNase E/G-like protein is present in plastids and is essential for chloroplast development and autotrophic growth in <i>Arabidopsis</i> . 2008 , 59, 2597-610	32
1076	NAD(H)-dependent glutamate dehydrogenase is essential for the survival of <i>Arabidopsis thaliana</i> during dark-induced carbon starvation. 2008 , 59, 667-80	126
1075	Mutations in AtPS1 (<i>Arabidopsis thaliana</i> parallel spindle 1) lead to the production of diploid pollen grains. 2008 , 4, e1000274	94
1074	The <i>Arabidopsis</i> P4-ATPase ALA3 localizes to the golgi and requires a beta-subunit to function in lipid translocation and secretory vesicle formation. 2008 , 20, 658-76	106
1073	The AGL62 MADS domain protein regulates cellularization during endosperm development in <i>Arabidopsis</i> . 2008 , 20, 635-47	192
1072	A cytosolic pathway for the conversion of hydroxypyruvate to glycerate during photorespiration in <i>Arabidopsis</i> . 2008 , 20, 2848-59	160
1071	Transcription factors relevant to auxin signalling coordinate broad-spectrum metabolic shifts including sulphur metabolism. 2008 , 59, 2831-46	40
1070	The <i>Arabidopsis</i> BLAP75/Rmi1 homologue plays crucial roles in meiotic double-strand break repair. 2008 , 4, e1000309	42
1069	The <i>Arabidopsis</i> cytochrome P450 CYP86A1 encodes a fatty acid omega-hydroxylase involved in suberin monomer biosynthesis. 2008 , 59, 2347-60	175
1068	Purification, molecular cloning, and characterization of glutathione S-transferases (GSTs) from pigmented <i>Vitis vinifera</i> L. cell suspension cultures as putative anthocyanin transport proteins. 2008 , 59, 3621-34	166
1067	Agrobacterium-mediated transformation of the haploid liverwort <i>Marchantia polymorpha</i> L., an emerging model for plant biology. 2008 , 49, 1084-91	209
1066	Histone arginine methylation is required for vernalization-induced epigenetic silencing of FLC in winter-annual <i>Arabidopsis thaliana</i> . 2008 , 105, 411-6	98
1065	SOMNUS, a CCCH-type zinc finger protein in <i>Arabidopsis</i> , negatively regulates light-dependent seed germination downstream of PIL5. 2008 , 20, 1260-77	210
1064	Activated expression of an <i>Arabidopsis</i> HD-START protein confers drought tolerance with improved root system and reduced stomatal density. 2008 , 20, 1134-51	263
1063	The <i>Arabidopsis</i> MUM2 gene encodes a beta-galactosidase required for the production of seed coat mucilage with correct hydration properties. 2007 , 19, 4007-21	119

1062	Exportin1 genes are essential for development and function of the gametophytes in <i>Arabidopsis thaliana</i> . 2008 , 180, 1493-500	36
1061	Association of the <i>Agrobacterium</i> T-DNA-protein complex with plant nucleosomes. 2008 , 105, 15429-34	56
1060	Efficient and high-throughput vector construction and <i>Agrobacterium</i> -mediated transformation of <i>Arabidopsis thaliana</i> suspension-cultured cells for functional genomics. 2008 , 49, 242-50	40
1059	Di- and tri- but not monomethylation on histone H3 lysine 36 marks active transcription of genes involved in flowering time regulation and other processes in <i>Arabidopsis thaliana</i> . 2008 , 28, 1348-60	235
1058	Subclade of flavin-monooxygenases involved in aliphatic glucosinolate biosynthesis. 2008 , 148, 1721-33	123
1057	<i>Arabidopsis thaliana</i> class II poly(A)-binding proteins are required for efficient multiplication of turnip mosaic virus. 2008 , 89, 2339-2348	42
1056	<i>Arabidopsis</i> SOMATIC EMBRYOGENESIS RECEPTOR KINASE proteins serve brassinosteroid-dependent and -independent signaling pathways. 2008 , 148, 611-9	147
1055	Two CRM protein subfamilies cooperate in the splicing of group IIB introns in chloroplasts. 2008 , 14, 2319-32	53
1054	XAP5 CIRCADIAN TIMEKEEPER coordinates light signals for proper timing of photomorphogenesis and the circadian clock in <i>Arabidopsis</i> . 2008 , 20, 1244-59	44
1053	Autophagy protein 6 (ATG6) is required for pollen germination in <i>Arabidopsis thaliana</i> . 2008 , 4, 339-48	70
1052	Functional characterization of <i>Arabidopsis</i> calreticulin1a: a key alleviator of endoplasmic reticulum stress. 2008 , 49, 912-24	55
1051	<i>Arabidopsis</i> SMG7 protein is required for exit from meiosis. 2008 , 121, 2208-16	66
1050	Sex-biased lethality or transmission of defective transcription machinery in <i>Arabidopsis</i> . 2008 , 180, 207-18	36
1049	A versatile transposon-based activation tag vector system for functional genomics in cereals and other monocot plants. 2008 , 146, 189-99	52
1048	Serine palmitoyltransferase, a key enzyme for de novo synthesis of sphingolipids, is essential for male gametophyte development in <i>Arabidopsis</i> . 2008 , 146, 1322-32	69
1047	Integrative approaches for mining transcriptional regulatory programs in <i>Arabidopsis</i> . 2008 , 7, 264-74	12
1046	Maternal control of male-gamete delivery in <i>Arabidopsis</i> involves a putative GPI-anchored protein encoded by the LORELEI gene. 2008 , 20, 3038-49	137
1045	<i>Arabidopsis</i> bZIP60 is a proteolysis-activated transcription factor involved in the endoplasmic reticulum stress response. 2008 , 20, 3107-21	221

1044	Minor antenna proteins CP24 and CP26 affect the interactions between photosystem II subunits and the electron transport rate in grana membranes of Arabidopsis. 2008 , 20, 1012-28	149
1043	Contribution of the GABA shunt to hypoxia-induced alanine accumulation in roots of Arabidopsis thaliana. 2008 , 49, 92-102	142
1042	SUGAR-DEPENDENT6 encodes a mitochondrial flavin adenine dinucleotide-dependent glycerol-3-phosphate dehydrogenase, which is required for glycerol catabolism and post germinative seedling growth in Arabidopsis. 2008 , 148, 519-28	33
1041	The ATG12-conjugating enzyme ATG10 is essential for autophagic vesicle formation in Arabidopsis thaliana. 2008 , 178, 1339-53	210
1040	The transcription factor MtSERF1 of the ERF subfamily identified by transcriptional profiling is required for somatic embryogenesis induced by auxin plus cytokinin in Medicago truncatula. 2008 , 146, 1622-36	118
1039	Potential sites of bioactive gibberellin production during reproductive growth in Arabidopsis. 2008 , 20, 320-36	175
1038	pur4 mutations are lethal to the male, but not the female, gametophyte and affect sporophyte development in Arabidopsis. 2008 , 147, 650-60	16
1037	A homolog of ScRAD5 is involved in DNA repair and homologous recombination in Arabidopsis. 2008 , 146, 1786-96	41
1036	Semidominant mutations in reduced epidermal fluorescence 4 reduce phenylpropanoid content in Arabidopsis. 2008 , 178, 2237-51	29
1035	A CRM domain protein functions dually in group I and group II intron splicing in land plant chloroplasts. 2007 , 19, 3864-75	68
1034	Web-based Arabidopsis functional and structural genomics resources. 2008 , 6, e0118	9
1033	Dual effects of miR156-targeted SPL genes and CYP78A5/KLUH on plastochron length and organ size in Arabidopsis thaliana. 2008 , 20, 1231-43	370
1032	Chloroplast Fe(III) chelate reductase activity is essential for seedling viability under iron limiting conditions. 2008 , 105, 10619-24	139
1031	Physiological roles of the beta-substituted alanine synthase gene family in Arabidopsis. 2008 , 146, 310-20	138
1030	Two class XI myosins function in organelle trafficking and root hair development in Arabidopsis. 2008 , 146, 1109-16	196
1029	Two GATA transcription factors are downstream effectors of floral homeotic gene action in Arabidopsis. 2008 , 147, 707-18	93
1028	LEUNIG_HOMOLOG and LEUNIG perform partially redundant functions during Arabidopsis embryo and floral development. 2008 , 147, 672-81	48
1027	AtCCX3 is an Arabidopsis endomembrane H ⁺ -dependent K ⁺ transporter. 2008 , 148, 1474-86	51

1026	AGL61 interacts with AGL80 and is required for central cell development in Arabidopsis. 2008 , 148, 259-68	87
1025	Characterization and biological function of the ISOCHORISMATE SYNTHASE2 gene of Arabidopsis. 2008 , 147, 1279-87	255
1024	In planta analysis of the cell cycle-dependent localization of AtCDC48A and its critical roles in cell division, expansion, and differentiation. 2008 , 148, 246-58	61
1023	AtPTR1 and AtPTR5 transport dipeptides in planta. 2008 , 148, 856-69	136
1022	Redundant requirement for a pair of PROTEIN ARGININE METHYLTRANSFERASE4 homologs for the proper regulation of Arabidopsis flowering time. 2008 , 148, 490-503	40
1021	Functional characterization of the Arabidopsis AtSUC2 Sucrose/H ⁺ symporter by tissue-specific complementation reveals an essential role in phloem loading but not in long-distance transport. 2008 , 148, 200-11	151
1020	The DEG15 serine protease cleaves peroxisomal targeting signal 2-containing proteins in Arabidopsis. 2008 , 148, 1847-56	63
1019	A gamma-glutamyl transpeptidase-independent pathway of glutathione catabolism to glutamate via 5-oxoproline in Arabidopsis. 2008 , 148, 1603-13	113
1018	LEAFY COTYLEDON1 is a key regulator of fatty acid biosynthesis in Arabidopsis. 2008 , 148, 1042-54	269
1017	Comparative analyses of Arabidopsis complex glycan1 mutants and genetic interaction with staurosporin and temperature sensitive3a. 2008 , 148, 1354-67	51
1016	Arabidopsis CLP1-SIMILAR PROTEIN3, an ortholog of human polyadenylation factor CLP1, functions in gametophyte, embryo, and postembryonic development. 2008 , 148, 2059-69	28
1015	Identification of a xylogalacturonan xylosyltransferase involved in pectin biosynthesis in Arabidopsis. 2008 , 20, 1289-302	100
1014	S-nitrosylation of peroxiredoxin II E promotes peroxynitrite-mediated tyrosine nitration. 2007 , 19, 4120-30	283
1013	NAI2 is an endoplasmic reticulum body component that enables ER body formation in Arabidopsis thaliana. 2008 , 20, 2529-40	50
1012	A heterocomplex of iron superoxide dismutases defends chloroplast nucleoids against oxidative stress and is essential for chloroplast development in Arabidopsis. 2008 , 20, 3148-62	201
1011	Disrupting two Arabidopsis thaliana xylosyltransferase genes results in plants deficient in xyloglucan, a major primary cell wall component. 2008 , 20, 1519-37	312
1010	A class I ADP-ribosylation factor GTPase-activating protein is critical for maintaining directional root hair growth in Arabidopsis. 2008 , 147, 1659-74	38
1009	Functional analysis of a predicted flavonol synthase gene family in Arabidopsis. 2008 , 147, 1046-61	162

1008	Genetically engineered crops for biofuel production: regulatory perspectives. 2008 , 25, 331-61	9
1007	Cellulose binding protein from the parasitic nematode <i>Heterodera schachtii</i> interacts with <i>Arabidopsis</i> pectin methylesterase: cooperative cell wall modification during parasitism. 2008 , 20, 3080-93	162
1006	<i>Pseudomonas syringae</i> elicits emission of the terpenoid (E,E)-4,8,12-trimethyl-1,3,7,11-tridecatetraene in <i>Arabidopsis</i> leaves via jasmonate signaling and expression of the terpene synthase TPS4. 2008 , 21, 1482-97	31
1005	Regulation of floral organ abscission in <i>Arabidopsis thaliana</i> . 2008 , 105, 15629-34	228
1004	Expression and developmental function of the 3-ketoacyl-ACP synthase2 gene in <i>Arabidopsis thaliana</i> . 2008 , 83, 143-52	9
1003	An inversion identified in <i>acl1-1</i> mutant functions as an enhancer of the <i>acl1-1</i> phenotype. 2008 , 83, 293-300	1
1002	Discovery of ADP-ribosylation and other plant defense pathway elements through expression profiling of four different <i>Arabidopsis</i> - <i>Pseudomonas</i> R-avr interactions. 2008 , 21, 646-57	47
1001	Diverse roles for chloroplast stromal and thylakoid-bound ascorbate peroxidases in plant stress responses. 2008 , 412, 275-85	142
1000	In vivo quantitative relationship between plastid division proteins FtsZ1 and FtsZ2 and identification of ARC6 and ARC3 in a native FtsZ complex. 2008 , 412, 367-78	48
999	Poplars. 2008 , 1-34	
998	Recent patents on agrobacterium-mediated gene and protein transfer, for research and biotechnology. 2008 , 2, 69-81	9
997	Identification, biochemical characterization, and subcellular localization of allantoate amidohydrolases from <i>Arabidopsis</i> and soybean. 2008 , 146, 418-30	76
996	MATERNALLY EXPRESSED PAB C-TERMINAL, a novel imprinted gene in <i>Arabidopsis</i> , encodes the conserved C-terminal domain of polyadenylate binding proteins. 2008 , 20, 2387-98	88
995	<i>Arabidopsis</i> VIP6/ELF8, the homolog of CTR9 component of the transcriptional complex PAF1, is essential for plant development. 2008 , 25, 447-455	3
994	The maternally expressed WRKY transcription factor TTG2 controls lethality in interploidy crosses of <i>Arabidopsis</i> . 2008 , 6, 2707-20	92
993	Parasites lead to evolution of robustness against gene loss in host signaling networks. 2008 , 4, 202	20
992	Characterization of PAP1-upregulated Glutathione S-transferase genes in <i>Arabidopsis thaliana</i> . 2008 , 25, 191-196	20
991	. 2008 ,	2

990	Transposon-based activation tagging in cereals. 2009 , 36, 915-921	12
989	Metal A and metal B sites of nuclear RNA polymerases Pol IV and Pol V are required for siRNA-dependent DNA methylation and gene silencing. 2009 , 4, e4110	40
988	AGO1 homeostasis involves differential production of 21-nt and 22-nt miR168 species by MIR168a and MIR168b. 2009 , 4, e6442	82
987	Integrating the genetic and physical maps of <i>Arabidopsis thaliana</i> : identification of mapped alleles of cloned essential (EMB) genes. 2009 , 4, e7386	40
986	Characterization of genes encoding poly(A) polymerases in plants: evidence for duplication and functional specialization. 2009 , 4, e8082	19
985	A Multiparent Advanced Generation Inter-Cross to fine-map quantitative traits in <i>Arabidopsis thaliana</i> . 2009 , 5, e1000551	421
984	Generation of "backbone" free, low transgene copy plants by launching T-DNA from the <i>Agrobacterium</i> chromosome. 2009 ,	
983	Genome-wide identification, structure and expression studies, and mutant collection of 22 early nodulin-like protein genes in <i>Arabidopsis</i> . 2009 , 73, 2452-9	44
982	Highly boron deficiency-tolerant plants generated by enhanced expression of NIP5;1, a boric acid channel. 2009 , 50, 58-66	84
981	<i>Arabidopsis</i> mutant deficient in 3 abscisic acid-activated protein kinases reveals critical roles in growth, reproduction, and stress. 2009 , 106, 8380-5	565
980	Male gametophyte development: a molecular perspective. 2009 , 60, 1465-78	229
979	GUN4 is required for posttranslational control of plant tetrapyrrole biosynthesis. 2009 , 2, 1198-210	72
978	The <i>Arabidopsis</i> GRF-INTERACTING FACTOR gene family performs an overlapping function in determining organ size as well as multiple developmental properties. 2009 , 151, 655-68	136
977	Bell-like homeodomain selectively regulates the high-irradiance response of phytochrome A. 2009 , 106, 13624-9	22
976	Assembly of an interactive correlation network for the <i>Arabidopsis</i> genome using a novel heuristic clustering algorithm. 2010 , 152, 29-43	135
975	Omics and bioinformatics: an essential toolbox for systems analyses of plant functions beyond 2010. 2009 , 50, 1177-80	19
974	Replication stress leads to genome instabilities in <i>Arabidopsis</i> DNA polymerase delta mutants. 2009 , 21, 2700-14	28
973	Histone acetylation, VERNALIZATION INSENSITIVE 3, FLOWERING LOCUS C, and the vernalization response. 2009 , 2, 724-737	50

972	Abscisic acid represses growth of the Arabidopsis embryonic axis after germination by enhancing auxin signaling. 2009 , 21, 2253-68	107
971	Differential recruitment of WOX transcription factors for lateral development and organ fusion in Petunia and Arabidopsis. 2009 , 21, 2269-83	153
970	Genetic resources for maize cell wall biology. 2009 , 151, 1703-28	117
969	Evidence for base excision repair of oxidative DNA damage in chloroplasts of Arabidopsis thaliana. 2009 , 284, 17006-17012	41
968	Arabidopsis SBP-box genes SPL10, SPL11 and SPL2 control morphological change in association with shoot maturation in the reproductive phase. 2009 , 50, 2133-45	184
967	LATERAL ORGAN FUSION1 and LATERAL ORGAN FUSION2 function in lateral organ separation and axillary meristem formation in Arabidopsis. 2009 , 136, 2423-32	97
966	Interaction of the Tobacco mosaic virus replicase protein with a NAC domain transcription factor is associated with the suppression of systemic host defenses. 2009 , 83, 9720-30	81
965	REVEILLE1, a Myb-like transcription factor, integrates the circadian clock and auxin pathways. 2009 , 106, 16883-8	158
964	Identification of plant cell wall mutants by means of a forward chemical genetic approach using hydrolases. 2009 , 106, 14699-704	104
963	The calmodulin-related calcium sensor CML42 plays a role in trichome branching. 2009 , 284, 31647-57	71
962	ARABIDOPSIS DEHISCENCE ZONE POLYGALACTURONASE1 (ADPG1), ADPG2, and QUARTET2 are Polygalacturonases required for cell separation during reproductive development in Arabidopsis. 2009 , 21, 216-33	260
961	CYP86B1 is required for very long chain omega-hydroxyacid and alpha, omega -dicarboxylic acid synthesis in root and seed suberin polyester. 2009 , 150, 1831-43	123
960	A chloroplastic UDP-glucose pyrophosphorylase from Arabidopsis is the committed enzyme for the first step of sulfolipid biosynthesis. 2009 , 21, 892-909	174
959	Functional analysis of transcription factors in Arabidopsis. 2009 , 50, 1232-48	200
958	Peptidyl-prolyl isomerase activity in chloroplast thylakoid lumen is a dispensable function of immunophilins in Arabidopsis thaliana. 2009 , 50, 1801-14	32
957	Methyl salicylate production and jasmonate signaling are not essential for systemic acquired resistance in Arabidopsis. 2009 , 21, 954-71	185
956	NADPH Thioredoxin reductase C controls the redox status of chloroplast 2-Cys peroxiredoxins in Arabidopsis thaliana. 2009 , 2, 298-307	71
955	HOS3, an ELO-like gene, inhibits effects of ABA and implicates a S-1-P/ceramide control system for abiotic stress responses in Arabidopsis thaliana. 2009 , 2, 138-51	37

954	Instrumentation and Software for Analysis of Arabidopsis Circadian Leaf Movement. 2009 , 1, 22-25	3
953	Accumulation of flavonoids in an ntra ntrb mutant leads to tolerance to UV-C. 2009 , 2, 249-58	44
952	Silencing of a plant gene by transcriptional interference. 2009 , 37, 3739-46	17
951	Redox regulation in photosynthetic organisms: signaling, acclimation, and practical implications. 2009 , 11, 861-905	1030
950	Novel nuclear-encoded subunits of the chloroplast NAD(P)H dehydrogenase complex. 2009 , 284, 905-12	48
949	ARABIDOPSIS TRITHORAX-RELATED7 is required for methylation of lysine 4 of histone H3 and for transcriptional activation of FLOWERING LOCUS C. 2009 , 21, 3257-69	149
948	Whirly proteins maintain plastid genome stability in Arabidopsis. 2009 , 106, 14693-8	152
947	The mediator complex subunit PFT1 is a key regulator of jasmonate-dependent defense in Arabidopsis. 2009 , 21, 2237-52	246
946	Arabidopsis LON2 is necessary for peroxisomal function and sustained matrix protein import. 2009 , 151, 1354-65	75
945	Uncovering small RNA-mediated responses to phosphate deficiency in Arabidopsis by deep sequencing. 2009 , 151, 2120-32	520
944	KAGIANA: an excel-based tool for retrieving summary information on Arabidopsis genes. 2009 , 50, 173-7	10
943	AtAMT1;4, a pollen-specific high-affinity ammonium transporter of the plasma membrane in Arabidopsis. 2009 , 50, 13-25	72
942	The signal peptide peptidase is required for pollen function in Arabidopsis. 2009 , 149, 1289-301	16
941	Histone H2B deubiquitination is required for transcriptional activation of FLOWERING LOCUS C and for proper control of flowering in Arabidopsis. 2009 , 149, 1196-204	80
940	MYB108 acts together with MYB24 to regulate jasmonate-mediated stamen maturation in Arabidopsis. 2009 , 149, 851-62	156
939	Temporal global expression data reveal known and novel salicylate-impacted processes and regulators mediating powdery mildew growth and reproduction on Arabidopsis. 2009 , 149, 1435-51	54
938	The short-rooted phenotype of the brevis radix mutant partly reflects root abscisic acid hypersensitivity. 2009 , 149, 1917-28	52
937	Triple loss of function of protein phosphatases type 2C leads to partial constitutive response to endogenous abscisic acid. 2009 , 150, 1345-55	207

936	CYP704B1 is a long-chain fatty acid omega-hydroxylase essential for sporopollenin synthesis in pollen of Arabidopsis. 2009 , 151, 574-89	224
935	Truncation of a protein disulfide isomerase, PDIL2-1, delays embryo sac maturation and disrupts pollen tube guidance in Arabidopsis thaliana. 2008 , 20, 3300-11	57
934	Arabidopsis ASA1 is important for jasmonate-mediated regulation of auxin biosynthesis and transport during lateral root formation. 2009 , 21, 1495-511	253
933	The association of the Arabidopsis actin-related protein2/3 complex with cell membranes is linked to its assembly status but not its activation. 2009 , 151, 2095-109	43
932	AINTEGUMENTA and AINTEGUMENTA-LIKE6 act redundantly to regulate Arabidopsis floral growth and patterning. 2009 , 150, 1916-29	126
931	Chloroplast NADPH-thioredoxin reductase interacts with photoperiodic development in Arabidopsis. 2009 , 149, 1261-76	127
930	Disruption of poly(ADP-ribosyl)ation mechanisms alters responses of Arabidopsis to biotic stress. 2010 , 152, 267-80	100
929	Postembryonic seedling lethality in the sterol-deficient Arabidopsis cyp51A2 mutant is partially mediated by the composite action of ethylene and reactive oxygen species. 2010 , 152, 192-205	30
928	Functional characterization of a higher plant sphingolipid Delta4-desaturase: defining the role of sphingosine and sphingosine-1-phosphate in Arabidopsis. 2009 , 149, 487-98	92
927	Establishing RNA interference as a reverse-genetic approach for gene functional analysis in protoplasts. 2009 , 149, 642-52	80
926	VTC4 is a bifunctional enzyme that affects myoinositol and ascorbate biosynthesis in plants. 2009 , 150, 951-61	109
925	Identification and functional characterization of inhibitor-3, a regulatory subunit of protein phosphatase 1 in plants. 2009 , 150, 144-56	25
924	Loss of 26S proteasome function leads to increased cell size and decreased cell number in Arabidopsis shoot organs. 2009 , 150, 178-89	88
923	Functional characterization of the Arabidopsis beta-ketoacyl-coenzyme A reductase candidates of the fatty acid elongase. 2009 , 150, 1174-91	146
922	AtBXL1 encodes a bifunctional beta-D-xylosidase/alpha-L-arabinofuranosidase required for pectic arabinan modification in Arabidopsis mucilage secretory cells. 2009 , 150, 1219-34	103
921	AMR1, an Arabidopsis gene that coordinately and negatively regulates the mannose/l-galactose ascorbic acid biosynthetic pathway. 2009 , 150, 942-50	77
920	Arabidopsis methionine gamma-lyase is regulated according to isoleucine biosynthesis needs but plays a subordinate role to threonine deaminase. 2009 , 151, 367-78	70
919	The role of oxophytodienoate reductases in the detoxification of the explosive 2,4,6-trinitrotoluene by Arabidopsis. 2009 , 151, 253-61	60

9 ¹⁸	Plant SMU-1 and SMU-2 homologues regulate pre-mRNA splicing and multiple aspects of development. 2009 , 151, 1498-512	29
9 ¹⁷	Abnormal physiological and molecular mutant phenotypes link chloroplast polynucleotide phosphorylase to the phosphorus deprivation response in Arabidopsis. 2009 , 151, 905-24	37
9 ¹⁶	The ABC transporter PXA1 and peroxisomal beta-oxidation are vital for metabolism in mature leaves of Arabidopsis during extended darkness. 2009 , 21, 2733-49	127
9 ¹⁵	DORNROSCHEN is a direct target of the auxin response factor MONOPTEROS in the Arabidopsis embryo. 2009 , 136, 1643-51	120
9 ¹⁴	The Arabidopsis nitrate transporter NRT1.7, expressed in phloem, is responsible for source-to-sink remobilization of nitrate. 2009 , 21, 2750-61	203
9 ¹³	Conservation of lotus and Arabidopsis basic helix-loop-helix proteins reveals new players in root hair development. 2009 , 151, 1175-85	79
9 ¹²	Multiple mechanism-mediated retention of a defective brassinosteroid receptor in the endoplasmic reticulum of Arabidopsis. 2008 , 20, 3418-29	147
9 ¹¹	Arabidopsis plants harbouring a mutation in AtSUC2, encoding the predominant sucrose/proton symporter necessary for efficient phloem transport, are able to complete their life cycle and produce viable seed. 2009 , 104, 1121-8	47
9 ¹⁰	Penetration of the stigma and style elicits a novel transcriptome in pollen tubes, pointing to genes critical for growth in a pistil. 2009 , 5, e1000621	258
9 ⁰⁹	Gibberellin acts through jasmonate to control the expression of MYB21, MYB24, and MYB57 to promote stamen filament growth in Arabidopsis. 2009 , 5, e1000440	266
9 ⁰⁸	DETORQUEO, QUIRKY, and ZERZAUST represent novel components involved in organ development mediated by the receptor-like kinase STRUBBELIG in Arabidopsis thaliana. 2009 , 5, e1000355	59
9 ⁰⁷	CHD3 proteins and polycomb group proteins antagonistically determine cell identity in Arabidopsis. 2009 , 5, e1000605	124
9 ⁰⁶	A link among DNA replication, recombination, and gene expression revealed by genetic and genomic analysis of TEBICHI gene of Arabidopsis thaliana. 2009 , 5, e1000613	40
9 ⁰⁵	Loss of chloroplast protease SPPA function alters high light acclimation processes in Arabidopsis thaliana L. (Heynh.). 2009 , 60, 1715-27	21
9 ⁰⁴	Repression of flowering by the miR172 target SMZ. 2009 , 7, e1000148	320
9 ⁰³	Arabidopsis thaliana T-DNA mutants implicate GAUT genes in the biosynthesis of pectin and xylan in cell walls and seed testa. 2009 , 2, 1000-14	99
9 ⁰²	Two Prp19-like U-box proteins in the MOS4-associated complex play redundant roles in plant innate immunity. 2009 , 5, e1000526	95
9 ⁰¹	A high throughput genetic screen identifies new early meiotic recombination functions in Arabidopsis thaliana. 2009 , 5, e1000654	105

900	A combinatorial interplay among the 1-aminocyclopropane-1-carboxylate isoforms regulates ethylene biosynthesis in <i>Arabidopsis thaliana</i> . 2009 , 183, 979-1003	208
899	Interplay between ethylene, ETP1/ETP2 F-box proteins, and degradation of EIN2 triggers ethylene responses in <i>Arabidopsis</i> . 2009 , 23, 512-21	238
898	The low-oxygen-induced NAC domain transcription factor ANAC102 affects viability of <i>Arabidopsis</i> seeds following low-oxygen treatment. 2009 , 149, 1724-38	114
897	In vivo studies on the roles of Tic55-related proteins in chloroplast protein import in <i>Arabidopsis thaliana</i> . 2009 , 2, 1397-409	22
896	DDBJ dealing with mass data produced by the second generation sequencer. 2009 , 37, D16-8	27
895	Regulation of membrane trafficking and organ separation by the NEVERSHED ARF-GAP protein. 2009 , 136, 1909-18	89
894	DLA-based strategies for cloning insertion mutants: cloning the gl4 locus of maize using Mu transposon tagged alleles. 2009 , 183, 1215-25	31
893	Quantitative analysis of ER body morphology in an <i>Arabidopsis</i> mutant. 2009 , 50, 2015-22	23
892	Mutation of the membrane-associated M1 protease APM1 results in distinct embryonic and seedling developmental defects in <i>Arabidopsis</i> . 2009 , 21, 1693-721	41
891	Roles for <i>Arabidopsis</i> CAMTA transcription factors in cold-regulated gene expression and freezing tolerance. 2009 , 21, 972-84	440
890	Helical growth of the <i>Arabidopsis</i> mutant <i>tortifolia2</i> does not depend on cell division patterns but involves handed twisting of isolated cells. 2009 , 21, 2090-106	66
889	Chloroplast ribonucleoprotein CP31A is required for editing and stability of specific chloroplast mRNAs. 2009 , 106, 6002-7	91
888	<i>Arabidopsis</i> FAB1/PIKfyve proteins are essential for development of viable pollen. 2009 , 151, 1812-22	64
887	The cytosolic protein response as a subcomponent of the wider heat shock response in <i>Arabidopsis</i> . 2009 , 21, 642-54	151
886	The gene controlling the indole glucosinolate modifier1 quantitative trait locus alters indole glucosinolate structures and aphid resistance in <i>Arabidopsis</i> . 2009 , 21, 985-99	160
885	The ferroportin metal efflux proteins function in iron and cobalt homeostasis in <i>Arabidopsis</i> . 2009 , 21, 3326-38	244
884	A search for factors influencing etioplast-chloroplast transition. 2009 , 106, 12201-6	15
883	Gene duplication and hypermutation of the pathogen Resistance gene SNC1 in the <i>Arabidopsis</i> bal variant. 2009 , 183, 1227-34	55

882	Peroxisome-associated matrix protein degradation in Arabidopsis. 2009 , 106, 4561-6	85
881	The paralogous genes RADICAL-INDUCED CELL DEATH1 and SIMILAR TO RCD ONE1 have partially redundant functions during Arabidopsis development. 2009 , 151, 180-98	58
880	Arabidopsis STN7 kinase provides a link between short- and long-term photosynthetic acclimation. 2009 , 21, 2402-23	212
879	AtnMat2, a nuclear-encoded maturase required for splicing of group-II introns in Arabidopsis mitochondria. 2009 , 15, 2299-311	99
878	MAP kinase phosphatase1 and protein tyrosine phosphatase1 are repressors of salicylic acid synthesis and SNC1-mediated responses in Arabidopsis. 2009 , 21, 2884-97	170
877	The Arabidopsis thaliana double-stranded RNA binding protein DRB1 directs guide strand selection from microRNA duplexes. 2009 , 15, 2219-35	164
876	A 1-megadalton translocation complex containing Tic20 and Tic21 mediates chloroplast protein import at the inner envelope membrane. 2009 , 21, 1781-97	82
875	The STRUCTURAL MAINTENANCE OF CHROMOSOMES 5/6 complex promotes sister chromatid alignment and homologous recombination after DNA damage in Arabidopsis thaliana. 2009 , 21, 2688-99	77
874	Web-queryable large-scale data sets for hypothesis generation in plant biology. 2009 , 21, 1034-51	98
873	Members of the LBD family of transcription factors repress anthocyanin synthesis and affect additional nitrogen responses in Arabidopsis. 2009 , 21, 3567-84	365
872	Agrobacterium tumefaciens VirC2 enhances T-DNA transfer and virulence through its C-terminal ribbon-helix-helix DNA-binding fold. 2009 , 106, 9643-8	27
871	Disruption of the pollen-expressed FERONIA homologs ANXUR1 and ANXUR2 triggers pollen tube discharge. 2009 , 136, 3279-88	220
870	VAJ/GFA1/CLO is involved in the directional control of floral organ growth. 2009 , 50, 515-27	15
869	RNA silencing is required for Arabidopsis defence against Verticillium wilt disease. 2009 , 60, 591-602	153
868	Understanding synergy in genetic interactions. 2009 , 25, 368-76	90
867	An archived activation tagged population of Arabidopsis thaliana to facilitate forward genetics approaches. 2009 , 9, 101	52
866	A modified TILLING approach to detect induced mutations in tetraploid and hexaploid wheat. 2009 , 9, 115	226
865	Characterization of WRKY co-regulatory networks in rice and Arabidopsis. 2009 , 9, 120	112

864	Salt-dependent regulation of a CNG channel subfamily in Arabidopsis. 2009 , 9, 140	68
863	A high-density collection of EMS-induced mutations for TILLING in Landsberg erecta genetic background of Arabidopsis. 2009 , 9, 147	36
862	The extracellular EXO protein mediates cell expansion in Arabidopsis leaves. 2009 , 9, 20	52
861	Effective carbon partitioning driven by exotic phloem-specific regulatory elements fused to the Arabidopsis thaliana AtSUC2 sucrose-proton symporter gene. 2009 , 9, 7	26
860	Characterization and isolation of a T-DNA tagged banana promoter active during in vitro culture and low temperature stress. 2009 , 9, 77	23
859	TILLING in the two-rowed barley cultivar 'Barke' reveals preferred sites of functional diversity in the gene HvHox1. 2009 , 2, 258	88
858	The signaling peptide EPF2 controls asymmetric cell divisions during stomatal development. 2009 , 19, 864-9	268
857	Dosage-dependent deregulation of an AGAMOUS-LIKE gene cluster contributes to interspecific incompatibility. 2009 , 19, 1128-32	102
856	ANXUR1 and 2, sister genes to FERONIA/SIRENE, are male factors for coordinated fertilization. 2009 , 19, 1327-31	206
855	AtAzg1 and AtAzg2 comprise a novel family of purine transporters in Arabidopsis. 2009 , 583, 481-6	34
854	SGS3 and RDR6 interact and colocalize in cytoplasmic SGS3/RDR6-bodies. 2009 , 583, 1261-6	124
853	AtCYP20-2 is an auxiliary protein of the chloroplast NAD(P)H dehydrogenase complex. 2009 , 583, 2355-8	39
852	AtGRP5, a vacuole-located glycine-rich protein involved in cell elongation. 2009 , 230, 253-65	32
851	Over-expression of CONSTANS-LIKE 5 can induce flowering in short-day grown Arabidopsis. 2009 , 230, 481-91	129
850	Transcriptional activation of the maize endosperm transfer cell-specific gene BETL1 by ZmMRP-1 is enhanced by two C2H2 zinc finger-containing proteins. 2009 , 230, 807-18	19
849	Diversity in subcellular targeting of the PP2A B'eta subfamily members. 2009 , 230, 935-45	37
848	Gene expression profiling of ozone-treated Arabidopsis abi1td insertional mutant: protein phosphatase 2C ABI1 modulates biosynthesis ratio of ABA and ethylene. 2009 , 230, 1003-17	35
847	The application of expression analysis in elucidating the eukaryotic elongation factor one alpha gene family in Arabidopsis thaliana. 2009 , 281, 391-405	18

846	Functional characterization of the Arabidopsis bHLH92 transcription factor in abiotic stress. 2009 , 282, 503-16	106
845	LAP3, a novel plant protein required for pollen development, is essential for proper exine formation. 2009 , 22, 167-77	49
844	The extracellular lipase EXL4 is required for efficient hydration of Arabidopsis pollen. 2009 , 22, 197-204	79
843	Incidence and post-pollination mechanisms of nonrandom mating in Arabidopsis thaliana. 2009 , 22, 257-62	14
842	Cohesin gene defects may impair sister chromatid alignment and genome stability in Arabidopsis thaliana. 2009 , 118, 591-605	38
841	Inactivation of genes, encoding tocopherol biosynthetic pathway enzymes, results in oxidative stress in outdoor grown Arabidopsis thaliana. 2009 , 47, 384-90	57
840	Characterization of T-DNA insertions in transgenic grapevines obtained by Agrobacterium-mediated transformation. 2009 , 24, 305-320	22
839	BIM1, a bHLH protein involved in brassinosteroid signalling, controls Arabidopsis embryonic patterning via interaction with DORNROSCHEN and DORNROSCHEN-LIKE. 2009 , 69, 57-68	69
838	Functional characterization of Arabidopsis NaCl-inducible WRKY25 and WRKY33 transcription factors in abiotic stresses. 2009 , 69, 91-105	413
837	Identification of two Arabidopsis genes encoding a peroxisomal oxidoreductase-like protein and an acyl-CoA synthetase-like protein that are required for responses to pro-auxins. 2009 , 69, 503-15	39
836	A protein related to prokaryotic UMP kinases is involved in psaA/B transcript accumulation in Arabidopsis. 2009 , 69, 517-28	10
835	The RNA-binding proteins CSP41a and CSP41b may regulate transcription and translation of chloroplast-encoded RNAs in Arabidopsis. 2009 , 69, 541-52	58
834	AtMKK1 and AtMPK6 are involved in abscisic acid and sugar signaling in Arabidopsis seed germination. 2009 , 70, 725-36	42
833	Antagonism between abscisic acid and ethylene in Arabidopsis acts in parallel with the reciprocal regulation of their metabolism and signaling pathways. 2009 , 71, 61-80	94
832	Arabidopsis thaliana encodes a bacterial-type heterodimeric isopropylmalate isomerase involved in both Leu biosynthesis and the Met chain elongation pathway of glucosinolate formation. 2009 , 71, 227-39	50
831	Deletion of an organellar peptidosome PreP affects early development in Arabidopsis thaliana. 2009 , 71, 497-508	29
830	Identification and functional analysis of two ZIP metal transporters of the hyperaccumulator Thlaspi caerulescens. 2009 , 325, 79-95	36
829	Glyphosate Resistance as a Versatile Selection Marker for Arabidopsis Transformation. 2009 , 27, 132-138	1

828	Quantitative analysis of the relationship between induction kinetics of chlorophyll fluorescence and function of genes in the cyanobacterium <i>Synechocystis</i> sp. PCC 6803. 2009 , 101, 47-58	7
827	Identification and Functional Analysis of Phosphoproteins Regulated by Auxin in <i>Arabidopsis</i> Roots. 2009 , 52, 65-72	4
826	Involvement of an R2R3-MYB transcription factor gene <i>AtMYB118</i> in embryogenesis in <i>Arabidopsis</i> . 2009 , 28, 337-46	33
825	Functional analysis of an <i>Arabidopsis</i> transcription factor <i>WRKY25</i> in heat stress. 2009 , 28, 683-93	133
824	High frequency <i>Agrobacterium tumefaciens</i> -mediated plant transformation induced by ammonium nitrate. 2009 , 28, 737-57	22
823	A transformation booster sequence (TBS) from <i>Petunia hybrida</i> functions as an enhancer-blocking insulator in <i>Arabidopsis thaliana</i> . 2009 , 28, 1095-104	32
822	Analysis of cytosolic and plastidic serine acetyltransferase mutants and subcellular metabolite distributions suggests interplay of the cellular compartments for cysteine biosynthesis in <i>Arabidopsis</i> . 2009 , 32, 349-367	122
821	Three novel subunits of <i>Arabidopsis</i> chloroplastic NAD(P)H dehydrogenase identified by bioinformatic and reverse genetic approaches. 2009 , 57, 207-19	71
820	The E2 ubiquitin-conjugating enzymes, <i>AtUBC1</i> and <i>AtUBC2</i> , play redundant roles and are involved in activation of <i>FLC</i> expression and repression of flowering in <i>Arabidopsis thaliana</i> . 2009 , 57, 279-88	122
819	The essential gene <i>EMB1611</i> maintains shoot apical meristem function during <i>Arabidopsis</i> development. 2009 , 57, 579-92	4
818	Systematic approaches to using the FOX hunting system to identify useful rice genes. 2009 , 57, 883-94	99
817	<i>VH1/BRL2</i> receptor-like kinase interacts with vascular-specific adaptor proteins <i>VIT</i> and <i>VIK</i> to influence leaf venation. 2009 , 57, 1000-14	53
816	Comparative analysis of leaf-type ferredoxin-NADP oxidoreductase isoforms in <i>Arabidopsis thaliana</i> . 2009 , 57, 1103-15	54
815	The timely deposition of callose is essential for cytokinesis in <i>Arabidopsis</i> . 2009 , 58, 13-26	89
814	<i>BTB</i> and <i>TAZ</i> domain scaffold proteins perform a crucial function in <i>Arabidopsis</i> development. 2009 , 58, 109-21	69
813	The <i>Arabidopsis</i> gene <i>YS1</i> encoding a DYW protein is required for editing of <i>rpoB</i> transcripts and the rapid development of chloroplasts during early growth. 2009 , 58, 82-96	155
812	<i>FIERY1</i> regulates light-mediated repression of cell elongation and flowering time via its 3'(2'),5'-bisphosphate nucleotidase activity. 2009 , 58, 208-19	48
811	A BAHD acyltransferase is expressed in the tapetum of <i>Arabidopsis</i> anthers and is involved in the synthesis of hydroxycinnamoyl spermidines. 2009 , 58, 246-59	140

810	Fusarium oxysporum hijacks COI1-mediated jasmonate signaling to promote disease development in Arabidopsis. 2009 , 58, 927-39	220
809	Interplay between SCARECROW, GA and LIKE HETEROCHROMATIN PROTEIN 1 in ground tissue patterning in the Arabidopsis root. 2009 , 58, 1016-27	85
808	The plant adherin AtSCC2 is required for embryogenesis and sister-chromatid cohesion during meiosis in Arabidopsis. 2009 , 59, 1-13	24
807	Identification of a novel E3 ubiquitin ligase that is required for suppression of premature senescence in Arabidopsis. 2009 , 59, 39-51	103
806	ATHB4, a regulator of shade avoidance, modulates hormone response in Arabidopsis seedlings. 2009 , 59, 266-77	94
805	VERNALIZATION INSENSITIVE 3 (VIN3) is required for the response of Arabidopsis thaliana seedlings exposed to low oxygen conditions. 2009 , 59, 576-87	48
804	Calcium-dependent protein kinases regulate polarized tip growth in pollen tubes. 2009 , 59, 528-39	146
803	Xylella fastidiosa subsp. pauca and fastidiosa Colonize Arabidopsis Systemically and Induce Anthocyanin Accumulation in Infected Leaves. 2019 , 109, 225-232	5
802	ELF18-INDUCED LONG NONCODING RNA evicts fibrillarin from mediator subunit to enhance PATHOGENESIS-RELATED GENE1 (PR1) expression. 2019 , 221, 2067-2079	38
801	CBL-interacting protein kinase 25 contributes to root meristem development. 2019 , 70, 133-147	18
800	EFFECTOR OF TRANSCRIPTION factors are novel plant-specific regulators associated with genomic DNA methylation in Arabidopsis. 2019 , 221, 261-278	4
799	Analysis of potential redundancy among Arabidopsis 6-phosphogluconolactonase isoforms in peroxisomes. 2020 , 71, 823-836	7
798	Thioredoxin h2 contributes to the redox regulation of mitochondrial photorespiratory metabolism. 2020 , 43, 188-208	20
797	Monitoring Ethylene in Plants: Genetically Encoded Reporters and Biosensors. 2020 , 4, 1900260	6
796	Biotechnological mechanism for improving plant remobilization of phosphorus during leaf senescence. 2020 , 18, 470-478	1
795	Genome-wide identification of EMBRYO-DEFECTIVE (EMB) genes required for growth and development in Arabidopsis. 2020 , 226, 306-325	54
794	Lack of FIBRILLIN6 in Arabidopsis thaliana affects light acclimation and sulfate metabolism. 2020 , 225, 1715-1731	6
793	Blurring the boundaries between cereal crops and model plants. 2020 , 228, 1721-1727	16

792	NAP is involved in GA-mediated chlorophyll degradation and leaf senescence by interacting with DELLAs in Arabidopsis. 2020 , 39, 75-87	12
791	The mismatch repair protein MSH6 regulates somatic recombination in Arabidopsis thaliana. 2020 , 87, 102789	3
790	YODA-HSP90 Module Regulates Phosphorylation-Dependent Inactivation of SPEECHLESS to Control Stomatal Development under Acute Heat Stress in Arabidopsis. 2020 , 13, 612-633	34
789	A Kinase and a Glycosylase Catabolize Pseudouridine in the Peroxisome to Prevent Toxic Pseudouridine Monophosphate Accumulation. 2020 , 32, 722-739	8
788	The Arabidopsis Receptor Kinase IRK Is Polarized and Represses Specific Cell Divisions in Roots. 2020 , 52, 183-195.e4	12
787	PEROXIREDOXIN Q stimulates the activity of the chloroplast 16:1 FATTY ACID DESATURASE4. 2020 , 102, 718-729	11
786	Decrease of Arabidopsis PAO activity entails increased RBOH activity, ROS content and altered responses to Pseudomonas. 2020 , 292, 110372	6
785	The Acetate Pathway Supports Flavonoid and Lipid Biosynthesis in Arabidopsis. 2020 , 182, 857-869	19
784	Reverse genetics-based biochemical studies of the ribosomal exit tunnel constriction region in eukaryotic ribosome stalling: spatial allocation of the regulatory nascent peptide at the constriction. 2020 , 48, 1985-1999	3
783	Inositol hexakisphosphate biosynthesis underpins PAMP-triggered immunity to Pseudomonas syringae pv. tomato in Arabidopsis thaliana but is dispensable for establishment of systemic acquired resistance. 2020 , 21, 376-387	4
782	Overexpression of a Rice Monosaccharide Transporter Gene (OsMST6) Confers Enhanced Tolerance to Drought and Salinity Stress in Arabidopsis thaliana. 2020 , 38, 151-164	9
781	The effector GpRbp-1 of Globodera pallida targets a nuclear HECT E3 ubiquitin ligase to modulate gene expression in the host. 2020 , 21, 66-82	4
780	A PXY-Mediated Transcriptional Network Integrates Signaling Mechanisms to Control Vascular Development in Arabidopsis. 2020 , 32, 319-335	44
779	Nitrate-inducible NIGT1 proteins modulate phosphate uptake and starvation signalling via transcriptional regulation of SPX genes. 2020 , 102, 448-466	20
778	Overexpression of a developing xylem cDNA library in transgenic poplar generates high mutation rate specific to wood formation. 2020 , 18, 1434-1443	3
777	The Earth BioGenome project: opportunities and challenges for plant genomics and conservation. 2020 , 102, 222-229	17
776	The Past, Present, and Future of Maize Improvement: Domestication, Genomics, and Functional Genomic Routes toward Crop Enhancement. 2020 , 1, 100010	34
775	Diversity of genetic lesions characterizes new Arabidopsis flavonoid pigment mutant alleles from T-DNA collections. 2020 , 291, 110335	6

774	Growth and development of AtMSH7 mutants in Arabidopsis thaliana. 2020 , 146, 329-336	6
773	NatB-Mediated N-Terminal Acetylation Affects Growth and Biotic Stress Responses. 2020 , 182, 792-806	25
772	PP7L is essential for MAIL1-mediated transposable element silencing and primary root growth. 2020 , 102, 703-717	3
771	CrRLK1L receptor-like kinases HERK1 and ANJEA are female determinants of pollen tube reception. 2020 , 21, e48466	29
770	Polar recruitment of RLD by LAZY1-like protein during gravity signaling in root branch angle control. 2020 , 11, 76	36
769	Development and screening of EMS mutants with altered seed oil content or fatty acid composition in Brassica napus. 2020 , 104, 1410-1422	5
768	Organelle movement and apical accumulation of secretory vesicles in pollen tubes of Arabidopsis thaliana depend on class XI myosins. 2020 , 104, 1685-1697	3
767	Discovery of a Family of Mixed Lineage Kinase Domain-like Proteins in Plants and Their Role in Innate Immune Signaling. 2020 , 28, 813-824.e6	21
766	ARMADILLO REPEAT ONLY proteins confine Rho GTPase signalling to polar growth sites. 2020 , 6, 1275-1288	7
765	The Cyclin CYCA3;4 Is a Postprophase Target of the APC/C E3-Ligase Controlling Formative Cell Divisions in Arabidopsis. 2020 , 32, 2979-2996	10
764	An efficient protocol for Agrobacterium-mediated genetic transformation of Antirrhinum majus. 2020 , 142, 527-536	2
763	Heterologous Expression of the and Genes from Restored the Hypersensitive Response and Resistance of Mutant to Bacterial Pathogen. 2020 , 9,	2
762	DREAM complex suppresses DNA methylation maintenance genes and precludes DNA hypermethylation. 2020 , 6, 942-956	16
761	Involvement of Arabidopsis BIG protein in cell death mediated by Myo-inositol homeostasis. 2020 , 10, 11268	1
760	Formate dehydrogenase takes part in molybdenum and iron homeostasis and affects dark-induced senescence in plants. 2020 , 15, 386-397	3
759	Touch-induced seedling morphological changes are determined by ethylene-regulated pectin degradation. 2020 , 6,	6
758	The Arabidopsis L-Type Amino Acid Transporter 5 (LAT5/PUT5) Is Expressed in the Phloem and Alters Seed Nitrogen Content When Knocked Out. 2020 , 9,	1
757	: A Sequence-Indexed Resource of Transposon-Induced Maize Mutations for Functional Genomics Studies. 2020 , 184, 620-631	9

756	Two chloroplast thioredoxin systems differentially modulate photosynthesis in Arabidopsis depending on light intensity and leaf age. 2020 , 104, 718-734	7
755	PUB11-Dependent Ubiquitination of the Phospholipid Flippase ALA10 Modifies ALA10 Localization and Affects the Pool of Linolenic Phosphatidylcholine. 2020 , 11, 1070	4
754	An Artificial Conversion of Roots into Organs with Shoot Stem Characteristics by Inducing Two Transcription Factors. 2020 , 23, 101332	2
753	Thiol Redox Regulation of Plant Carbonic Anhydrase. 2020 , 10,	9
752	AtPPRT3, a novel E3 ubiquitin ligase, plays a positive role in ABA signaling. 2020 , 39, 1467-1478	3
751	Roles for the chloroplast-localized pentatricopeptide repeat protein 30 and the 'mitochondrial' transcription termination factor 9 in chloroplast quality control. 2020 , 104, 735-751	9
750	PI4K α Interacts with E3 Ligase MIEL1 to Regulate Auxin Metabolism and Root Development. 2020 , 184, 933-944	2
749	The Critical Role of <i>AtHsp101</i> and Genes in Arabidopsis Phosphate Compensation Network. 2020 , 11, 565865	5
748	<i>AtHsp101</i> research sets course of action for the genetic improvement of crops against heat stress. 2020 , 29, 715-732	4
747	An ethyl methanesulfonate-induced neutral mutant-bridging method efficiently identifies spontaneously mutated genes in rice. 2020 , 104, 1129-1141	2
746	Regulation of Expression by bHLH and WRKY Transcription Factors Helps to Confer Increased Salt Tolerance to Plants. 2020 , 11, 1311	11
745	ABSCISIC ACID-DEFICIENT4 Has an Essential Function in Both cis-Violaxanthin and cis-Neoxanthin Synthesis. 2020 , 184, 1303-1316	11
744	Key role of the motor protein Kinesin 13B in the activity of homeodomain-leucine zipper I transcription factors. 2020 , 71, 6282-6296	1
743	COLD REGULATED 27 and 28 are targets of CONSTITUTIVELY PHOTOMORPHOGENIC 1 and negatively affect phytochrome B signalling. 2020 , 104, 1038-1053	4
742	Iron-Sulfur Cluster Protein NITROGEN FIXATION S-LIKE1 and Its Interactor FRATAXIN Function in Plant Immunity. 2020 , 184, 1532-1548	4
741	The peptidyl-prolyl isomerases FKBP15-1 and FKBP15-2 negatively affect lateral root development by repressing the vacuolar invertase VIN2 in Arabidopsis. 2020 , 252, 52	2
740	Regulation of a Cytochrome P450 Gene by WRKY33 Transcription Factor Controls Apoplastic Barrier Formation in Roots to Confer Salt Tolerance. 2020 , 184, 2199-2215	24
739	The Salt Tolerance Related Protein (STRP) Mediates Cold Stress Responses and Abscisic Acid Signalling in Arabidopsis. 2020 , 11, 1251	2

738	Construction of Mutant Library Provides Genetic Resources for Cotton Germplasm Improvement. 2020 , 21,	3
737	A natriuretic peptide from <i>Arabidopsis thaliana</i> (AtPNP-A) can modulate catalase 2 activity. 2020 , 10, 19632	2
736	Role of Splicing factor SF1 in Temperature-Responsive Alternative Splicing of pre-mRNA. 2020 , 11, 596354	10
735	Functional Interchangeability of Nucleotide Sugar Transporters URGT1 and URGT2 Reveals That and Cell Wall Chemotypes Depend on Their Spatio-Temporal Expression. 2020 , 11, 594544	2
734	Participates in the Defense Response of Pepper to through Regulating the Expression of Defense-Related Genes. 2020 , 21,	3
733	The Tapetal Major Facilitator NPF2.8 Is Required for Accumulation of Flavonol Glycosides on the Pollen Surface in. 2020 , 32, 1727-1748	11
732	GAL4/GFP enhancer-trap lines for identification and manipulation of cells and tissues in developing <i>Arabidopsis</i> leaves. 2020 , 249, 1127-1146	5
731	TRIPP Is a Plant-Specific Component of the <i>Arabidopsis</i> TRAPP1 Membrane Trafficking Complex with Important Roles in Plant Development. 2020 , 32, 2424-2443	10
730	CpARF2 and CpEIL1 interact to mediate auxin-ethylene interaction and regulate fruit ripening in papaya. 2020 , 103, 1318-1337	20
729	Exocyst mutants suppress pollen tube growth and cell wall structural defects of hydroxyproline O-arabinosyltransferase mutants. 2020 , 103, 1399-1419	9
728	FRA1 Kinesin Modulates the Lateral Stability of Cortical Microtubules through Cellulose Synthase-Microtubule Uncoupling Proteins. 2020 , 32, 2508-2524	8
727	The Lipid Flippases ALA4 and ALA5 Play Critical Roles in Cell Expansion and Plant Growth. 2020 , 182, 2111-2125	7
726	Identification, deployment, and transferability of quantitative trait loci from genome-wide association studies in plants. 2020 , 24, 100145	7
725	Insertional mutagenesis of <i>Brachypodium distachyon</i> using the Tnt1 retrotransposable element. 2020 , 103, 1924-1936	4
724	STARCH SYNTHASE5, a Noncanonical Starch Synthase-Like Protein, Promotes Starch Granule Initiation in <i>Arabidopsis</i> . 2020 , 32, 2543-2565	21
723	The isoleucic acid triad: distinct impacts on plant defense, root growth, and formation of reactive oxygen species. 2020 , 71, 4258-4270	6
722	The <i>Arabidopsis</i> Protein CGL20 Is Required for Plastid 50S Ribosome Biogenesis. 2020 , 182, 1222-1238	5
721	Maize and Genes Have Overlapping and Distinct Functions in Cuticular Lipid Deposition. 2020 , 183, 840-853	5

720	Regulation of Leaf Longevity by DML3-Mediated DNA Demethylation. 2020 , 13, 1149-1161	6
719	An impedance-coupled microfluidic device for single-cell analysis of primary cell wall regeneration. 2020 , 165, 112374	8
718	Non-Catalytic Subunits Facilitate Quaternary Organization of Plastidic Acetyl-CoA Carboxylase. 2020 , 182, 756-775	7
717	Endogenous Biosynthesis of S-Nitrosoglutathione From Nitro-Fatty Acids in Plants. 2020 , 11, 962	8
716	An iron (II) dependent oxygenase performs the last missing step of plant lysine catabolism. 2020 , 11, 2931	2
715	Adaptive reduction of male gamete number in the selfing plant <i>Arabidopsis thaliana</i> . 2020 , 11, 2885	9
714	PWR/HDA9/ABI4 Complex Epigenetically Regulates ABA Dependent Drought Stress Tolerance in. 2020 , 11, 623	21
713	<i>Arabidopsis</i> Modulates Pathogen Defense and Tolerance to Oxidative Stress. 2020 , 11, 703	3
712	A simple, non-toxic method for separating seeds based on density, and its application in isolating seed oil mutants. 2020 , 8, e11332	2
711	The <i>Arabidopsis</i> GPI-Anchored LTPg5 Encoded by Has a Role in Resistance against a Diverse Range of Pathogens. 2020 , 21,	3
710	RETICULON-LIKE4 (RTNLB4) Protein Participates in Infection and VirB2 Peptide-Induced Plant Defense Response. 2020 , 21,	1
709	RNA Interference-Based Screen Reveals Concerted Functions of MEKK2 and CRCK3 in Plant Cell Death Regulation. 2020 , 183, 331-344	4
708	The Circadian Clock Influences the Long-Term Water Use Efficiency of <i>Arabidopsis</i> . 2020 , 183, 317-330	18
707	<i>Arabidopsis</i> ROOT PHOTOTROPISM2 Is a Light-Dependent Dynamic Modulator of Phototropin1. 2020 , 32, 2004-2019	9
706	The Position and Complex Genomic Architecture of Plant T-DNA Insertions Revealed by 4SEE. 2020 , 21,	2
705	Analytical Techniques for Determining the Role of Domain of Unknown Function 579 Proteins in the Synthesis of -Methylated Plant Polysaccharides. 2020 , 25, 345-355	2
704	Integrated multi-omics framework of the plant response to jasmonic acid. 2020 , 6, 290-302	59
703	Vitamin B6 Is Under a Tight Balance During Disease Development by on Different Cultivars of Potato and on Mutants. 2020 , 11, 875	3

702	Ubiquitin-Conjugating Enzymes UBC7, UBC13, and UBC14 Are Required in Plant Responses to Multiple Stress Conditions. 2020 , 9,	3
701	A <i>Leymus chinensis</i> histidine-rich Ca-binding protein binds Ca/Zn and suppresses abscisic acid signaling in <i>Arabidopsis</i> . 2020 , 252, 153209	2
700	De Novo Shoot Regeneration Controlled by HEN1 and TCP3/4 in <i>Arabidopsis</i> . 2020 , 61, 1600-1613	6
699	Molecular Cloning and Functional Characterization of and Transcription Factors from Wintersweet (<i>L.</i>). 2020 , 9,	9
698	<i>Arabidopsis</i> cysteine-rich trans-membrane module (CYSTM) small proteins play a protective role mainly against heat and UV stresses. 2020 , 47, 195-202	1
697	The SWI/SNF ATP-Dependent Chromatin Remodeling Complex in <i>Arabidopsis</i> Responds to Environmental Changes in Temperature-Dependent Manner. 2020 , 21,	5
696	Functional Aspects of Early Light-Induced Protein (ELIP) Genes from the Desiccation-Tolerant Moss. 2020 , 21,	2
695	Sulfur Deficiency-Induced Glucosinolate Catabolism Attributed to Two β Glucosidases, BGLU28 and BGLU30, is Required for Plant Growth Maintenance under Sulfur Deficiency. 2020 , 61, 803-813	15
694	The Regulates Dark-Induced Senescence and Plays Contrasting Roles in Defense Against Bacterial and Fungal Pathogens. 2020 , 33, 754-766	11
693	Light Activates the Translational Regulatory Kinase GCN2 via Reactive Oxygen Species Emanating from the Chloroplast. 2020 , 32, 1161-1178	18
692	CmCYC2-like transcription factors may interact with each other or bind to the promoter to regulate floral symmetry development in <i>Chrysanthemum morifolium</i> . 2020 , 103, 159-171	6
691	Two <i>Arabidopsis</i> Chloroplast GrpE Homologues Exhibit Distinct Biological Activities and Can Form Homo- and Hetero-Oligomers. 2019 , 10, 1719	0
690	The TIR-NB-LRR pair DSC1 and WRKY19 contributes to basal immunity of <i>Arabidopsis</i> to the root-knot nematode <i>Meloidogyne incognita</i> . 2020 , 20, 73	9
689	An importin-beta-like protein mediates lignin-modification-induced dwarfism in <i>Arabidopsis</i> . 2020 , 102, 1281-1293	12
688	A malectin domain kinesin functions in pollen and seed development in <i>Arabidopsis</i> . 2020 , 71, 1828-1841	4
687	Rediscovery of Genetic and Genomic Resources for Future Food Security. 2020 ,	3
686	Calcium channel CNGC19 mediates basal defense signaling to regulate colonization by <i>Piriformospora indica</i> in <i>Arabidopsis</i> roots. 2020 , 71, 2752-2768	19
685	Effects of <i>Metaxenia</i> on Stone Cell Formation in Pear (<i>Pyrus bretschneideri</i>) Based on Transcriptomic Analysis and Functional Characterization of the Lignin-Related Gene PbC4H2. 2020 , 11, 53	0

684	Gene Type and Mutation Position Influence Responses in Root Traits across Nutrient Environments. 2020 , 181, 210-223	3
683	PARC6 Is Critical for Plastid Morphogenesis in Pavement, Trichome, and Guard Cells in Leaf Epidermis. 2019 , 10, 1665	4
682	Epistatic Transcription Factor Networks Differentially Modulate Growth and Defense. 2020 , 214, 529-541	7
681	Functional dissection of the DORNREICHEN-LIKE enhancer 2 during embryonic and phyllotactic patterning. 2020 , 251, 90	3
680	STRESS INDUCED FACTOR 2 Regulates Arabidopsis Stomatal Immunity through Phosphorylation of the Anion Channel SLAC1. 2020 , 32, 2216-2236	6
679	High-Throughput CRISPR/Cas9 Mutagenesis Streamlines Trait Gene Identification in Maize. 2020 , 32, 1397-1413	68
678	mTERF8, a Member of the Mitochondrial Transcription Termination Factor Family, Is Involved in the Transcription Termination of Chloroplast Gene. 2020 , 182, 408-423	13
677	The growth of Arabidopsis primary root is repressed by several and diverse amino acids through auxin-dependent and independent mechanisms and MPK6 kinase activity. 2021 , 302, 110717	3
676	Endogenous indole-3-acetamide levels contribute to the crosstalk between auxin and abscisic acid, and trigger plant stress responses in Arabidopsis. 2021 , 72, 459-475	10
675	SAW homeodomain transcription factors regulate initiation of leaf margin serrations. 2021 , 72, 1738-1747	0
674	Silencing of FOREVER YOUNG FLOWER-Like Genes from Phalaenopsis Orchids Promotes Flower Senescence and Abscission. 2021 , 62, 111-124	3
673	The Arabidopsis leucine-rich repeat receptor-like kinase MIK2 is a crucial component of early immune responses to a fungal-derived elicitor. 2021 , 229, 3453-3466	16
672	The transcriptional repressors VAL1 and VAL2 recruit PRC2 for genome-wide Polycomb silencing in Arabidopsis. 2021 , 49, 98-113	20
671	Approaching the genetic dissection of indirect adventitious organogenesis process in tomato explants. 2021 , 302, 110721	0
670	A novel zinc transporter essential for Arabidopsis zinc and iron-dependent growth. 2021 , 256, 153296	6
669	EARLY RESPONSE TO DEHYDRATION 7 Remodels Cell Membrane Lipid Composition during Cold Stress in Arabidopsis. 2021 , 62, 80-91	6
668	Application of genomics and transcriptomics to accelerate development of clubroot resistant canola. 2021 , 43, 189-208	5
667	CRISPR/Cas-Mediated Abiotic Stress Tolerance in Crops. 2021 , 177-211	2

- 666 Piriformospora indica employs host-derived putrescine for growth promotion in plants.
- 665 Coordinated cytokinin signaling and auxin biosynthesis mediates arsenate-induced root growth inhibition. **2021**, 185, 1166-1181 7
- 664 Meiotic chromosome axis remodelling is critical for meiotic recombination in Brassica rapa. **2021**, 72, 3012-3027 7
- 663 Phytochrome B inhibits darkness-induced hypocotyl adventitious root formation by stabilizing IAA14 and suppressing ARF7 and ARF19. **2021**, 105, 1689-1702 2
- 662 Easymap: a user-friendly software package for rapid mapping by sequencing of point mutations and large insertions.
- 661 Local jasmonic acid cues drive systemic acquired resistance signal generation.
- 660 Functional relationships of three NFU proteins in the biogenesis of chloroplastic iron-sulfur clusters. **2021**, 5, e00303 1
- 659 The Arabidopsis NOT4A E3 ligase promotes PGR3 expression and regulates chloroplast translation. **2021**, 12, 251 2
- 658 The CLASSY family controls tissue-specific DNA methylation patterns in Arabidopsis. 4
- 657 Identification of tung tree FATB as a promoter of 18:3 fatty acid accumulation through hydrolyzing 18:0-ACP. **2021**, 145, 143-154 1
- 656 Reported differences in the flg22 response of the null mutation of AtRGS1 correlates with fixed genetic variation in the background of Col-0 isolates. **2021**, 16, 1878685 2
- 655 Chemokine-like MDL proteins modulate flowering time and innate immunity in plants. **2021**, 296, 100611 0
- 654 Effects of Natural Variations on Biofortification. **2021**, 115-138 1
- 653 Development of -based activation tags for crop insertional mutagenesis. **2021**, 5, e00300 1
- 652 Review on forward and reverse genetics in plant breeding. **2021**, 14, 127-135 3
- 651 Arabidopsis Ca²⁺-ATPases 1, 2, and 7 in the endoplasmic reticulum contribute to growth and pollen fitness. **2021**, 185, 1966-1985 10
- 650 slim shady is a novel allele of PHYTOCHROME B present in the T-DNA line SALK_015201.
- 649 Analysis of sticky generative cell mutants reveals that suppression of callose deposition in the generative cell is necessary for generative cell internalization and differentiation in Arabidopsis. **2021**, 106, 228-244 1

648	Arabidopsis Mitochondrial Transcription Termination Factor mTERF2 Promotes Splicing of Group IIB Introns. 2021 , 10,	5
647	A Centrifuge-Based Method for Identifying Novel Genetic Traits That Affect Root-Substrate Adhesion in. 2021 , 12, 602486	1
646	Opposing functions of the plant TOPLESS gene family during SNC1-mediated autoimmunity. 2021 , 17, e1009026	4
645	Yield maintenance under drought is orchestrated by the qDTY12.1-encoded DECUSSATE gene of rice through a network with other flowering-associated genes across the genetic background.	1
644	Singlet oxygen-dependent chloroplast degradation is independent of macroautophagy in the Arabidopsis ferrochelatase two mutant.	
643	Discovery of photosynthesis genes through whole-genome sequencing of acetate-requiring mutants of <i>Chlamydomonas reinhardtii</i> .	
642	The mitochondrial AAA protease FTSH3 regulates Complex I abundance by promoting its disassembly. 2021 , 186, 599-610	1
641	AtFAHD1a: A New Player Influencing Seed Longevity and Dormancy in Arabidopsis?. 2021 , 22,	3
640	Genome-Wide Analysis of Tubulin Gene Family in Cassava and Expression of Family Member during Various Stress. 2021 , 10,	0
639	Large scale genomic rearrangements in selected Arabidopsis thaliana T-DNA lines are caused by T-DNA insertion mutagenesis.	3
638	Construction of a Full-Length cDNA Over-Expressing Library to Identify Valuable Genes from. 2021 , 22,	1
637	Sucrose triggers a novel signaling cascade promoting <i>Bacillus subtilis</i> rhizosphere colonization. 2021 , 15, 2723-2737	17
636	Uncovering a novel function of the CCR4-NOT complex in phytochrome A-mediated light signalling in plants. 2021 , 10,	6
635	Can genetic engineering-based methods for gene function identification be eclipsed by genome editing in plants? A comparison of methodologies. 2021 , 296, 485-500	0
634	The ATXN2 Orthologs and , Act Redundantly to In-Fluence Developmental Pathways throughout the Life Cycle of. 2021 , 22,	
633	Genome-wide analysis of AP2/ERF superfamily in lotus (<i>Nelumbo nucifera</i>) and the association between NnADAP and rhizome morphology. 2021 , 22, 171	7
632	An Arabidopsis Oxalyl-CoA Decarboxylase, AtOXC, Is Important for Oxalate Catabolism in Plants. 2021 , 22,	3
631	The Phosphofructokinase Isoform AtPFK5 Is a Novel Target of Plastidic Thioredoxin-f-Dependent Redox Regulation. 2021 , 10,	1

630	The Sequential Action of MIDA9/PP2C.D1, PP2C.D2, and PP2C.D5 Is Necessary to Form and Maintain the Hook After Germination in the Dark. 2021 , 12, 636098	0
629	Expansion and Molecular Characterization of AP2/ERF Gene Family in Wheat (L.). 2021 , 12, 632155	5
628	Ribosome biogenesis factor OLI2 and its interactor BRX1-2 are associated with morphogenesis and lifespan extension in. 2021 , 38, 117-125	
627	The TOR-EIN2 axis mediates nuclear signalling to modulate plant growth. 2021 , 591, 288-292	21
626	WRKY9 transcription factor regulates cytochrome P450 genes CYP94B3 and CYP86B1, leading to increased root suberin and salt tolerance in Arabidopsis. 2021 , 172, 1673-1687	5
625	RNA-seq-based identification of potential resistance genes against the soybean cyst nematode (Heterodera glycines) HG Type 1.2.3.5.7 in Dongnong L-10 2021 , 114, 101627	3
624	Turgor regulation defect 1 proteins play a conserved role in pollen tube reproductive innovation of the angiosperms. 2021 , 106, 1356-1365	0
623	FASCICLIN-LIKE 18 Is a New Player Regulating Root Elongation in. 2021 , 12, 645286	1
622	Genome-Wide Analysis of the RAV Transcription Factor Genes in Rice Reveals Their Response Patterns to Hormones and Virus Infection. 2021 , 13,	0
621	Spatial and temporal localization of SPIRRIG and WAVE/SCAR reveal roles for these proteins in actin-mediated root hair development. 2021 , 33, 2131-2148	2
620	A Case of Gene Fragmentation in Plant Mitochondria Fixed by the Selection of a Compensatory Restorer of Fertility-Like PPR Gene. 2021 , 38, 3445-3458	0
619	An Argon-Ion-Induced Pale Green Mutant of Exhibiting Rapid Disassembly of Mesophyll Chloroplast Grana. 2021 , 10,	0
618	tRNA-Dependent Import of a Transit Sequence-Less Aminoacyl-tRNA Synthetase (LeuRS2) into the Mitochondria of Arabidopsis. 2021 , 22,	0
617	The contribution of PIP2-type aquaporins to photosynthetic response to increased vapour pressure deficit. 2021 , 72, 5066-5078	1
616	mRNA surveillance complex PELOTA-HBS1 regulates phosphoinositide-dependent protein kinase1 and plant growth. 2021 , 186, 2003-2020	3
615	A single-cell analysis of the Arabidopsis vegetative shoot apex. 2021 , 56, 1056-1074.e8	26
614	MES7 Modulates Seed Germination via Regulating Salicylic Acid Content in Arabidopsis. 2021 , 10,	0
613	CRISPR screens in plants: approaches, guidelines, and future prospects. 2021 , 33, 794-813	10

612	Arabidopsis phosphatidylinositol 4-phosphate 5-kinase genes PIP5K7, PIP5K8, and PIP5K9 are redundantly involved in root growth adaptation to osmotic stress. 2021 , 106, 913-927	2
611	Isolation of Arabidopsis extracellular ATP binding proteins by affinity proteomics and identification of PHOSPHOLIPASE[̳-LIKE] as an extracellular protein essential for fumonisin[B1 toxicity. 2021 , 106, 1387-1400	2
610	Easymap: A User-Friendly Software Package for Rapid Mapping-by-Sequencing of Point Mutations and Large Insertions. 2021 , 12, 655286	2
609	PRAT Proteins Operate in Organellar Protein Import and Export in. 2021 , 10,	0
608	A Genome-Scale TF-DNA Interaction Network of Transcriptional Regulation of Arabidopsis Primary and Specialized Metabolism.	
607	The chemical compound 'Heatin' stimulates hypocotyl elongation and interferes with the Arabidopsis NIT1-subfamily of nitrilases. 2021 , 106, 1523-1540	1
606	A combination of genome-wide and transcriptome-wide association studies reveals genetic elements leading to male sterility during high temperature stress in cotton. 2021 , 231, 165-181	4
605	RAD54 is essential for RAD51-mediated repair of meiotic DSB in Arabidopsis. 2021 , 17, e1008919	3
604	HsfA1d promotes hypocotyl elongation under chilling via enhancing expression of ribosomal protein genes in Arabidopsis. 2021 , 231, 646-660	0
603	Genome-wide identification of the BASS gene family in four Gossypium species and functional characterization of GhBASSs against salt stress. 2021 , 11, 11342	3
602	Rapid identification of mutations caused by fast neutron bombardment in Medicago truncatula. 2021 , 17, 62	2
601	PGRL2 triggers degradation of PGR5 in the absence of PGRL1. 2021 , 12, 3941	5
600	A MYB4-MAN3-Mannose-MNB1 signaling cascade regulates cadmium tolerance in Arabidopsis. 2021 , 17, e1009636	2
599	Protocol for screening and expression studies of T-DNA and tagging-based insertional mutants in. 2021 , 11, 332	
598	is a novel allele of present in the T-DNA line SALK_015201. 2021 , 5, e00326	0
597	Ferroportin 3 is a dual-targeted mitochondrial/chloroplast iron exporter necessary for iron homeostasis in Arabidopsis. 2021 , 107, 215-236	7
596	A network of stress-related genes regulates hypocotyl elongation downstream of selective auxin perception. 2021 , 187, 430-445	1
595	Two evolutionary distinct effectors from a nematode and virus target RanGAP1 and 2 via the WPP domain to promote disease.	0

594	Three β -Glucuronosyltransferase Genes Involved in Arabinogalactan Biosynthesis Function in Arabidopsis Growth and Development. 2021 , 10,	1
593	Transcription-coupled and epigenome-encoded mechanisms direct H3K4 methylation.	0
592	Is Critical for Chloroplast Development in Leaf Epidermal Guard Cells. 2021 , 10,	0
591	Expanding the range of CRISPR/Cas9-directed genome editing in soybean. 1	0
590	Comparison of metabolites and variety authentication of Amomum tsao-ko and Amomum paratsao-ko using GC-MS and NIR spectroscopy. 2021 , 11, 15200	2
589	Local regulation of auxin transport in root-apex transition zone mediates aluminium-induced Arabidopsis root-growth inhibition. 2021 , 108, 55-66	3
588	-Triggered Cell Enlargement and Loss of Cellular Integrity in Root Systems Are Mediated by Pectin Demethylation. 2021 , 12, 711838	1
587	TRANVIA (TVA) facilitates cellulose synthase trafficking and delivery to the plasma membrane. 2021 , 118,	2
586	The core autophagy machinery is not required for chloroplast singlet oxygen-mediated cell death in the Arabidopsis thaliana plastid ferrochelatase two mutant. 2021 , 21, 342	6
585	High-throughput insertional mutagenesis reveals novel targets for enhancing lipid accumulation in Nannochloropsis oceanica. 2021 , 66, 239-258	13
584	Guard cell endomembrane Ca-ATPases underpin a 'carbon memory' of photosynthetic assimilation that impacts on water-use efficiency. 2021 , 7, 1301-1313	6
583	The Arabidopsis Iron-Sulfur (Fe-S) Cluster Gene Plays a Role in Host and Nonhost Disease Resistance by Accumulation of Defense-Related Metabolites. 2021 , 22,	0
582	Differential degradation of RNA species by autophagy-related pathways in Arabidopsis. 2021 , 72, 6867-6881	1
581	Adaptation to coastal soils through pleiotropic boosting of ion and stress hormone concentrations in wild Arabidopsis thaliana. 2021 , 232, 208-220	2
580	FERONIA receptor kinase-regulated reactive oxygen species mediate self-incompatibility in Brassica rapa. 2021 , 31, 3004-3016.e4	13
579	THESEUS1 modulates cell wall stiffness and abscisic acid production in Arabidopsis thaliana.	0
578	Large scale genomic rearrangements in selected Arabidopsis thaliana T-DNA lines are caused by T-DNA insertion mutagenesis. 2021 , 22, 599	5
577	Combinations of maternal-specific repressive epigenetic marks in the endosperm control seed dormancy. 2021 , 10,	2

576	Heterologous Complementation of SPO11-1 and -2 Depends on the Splicing Pattern. 2021 , 22,	0
575	A competitive PCR-based method to detect a single copy of T-DNA insertion in transformants. 2021 , 173, 1179-1188	0
574	Arabidopsis RAB8A, RAB8B and RAB8D Proteins Interact with Several RTNLB Proteins and are Involved in the Agrobacterium tumefaciens Infection Process. 2021 , 62, 1572-1588	1
573	ERF1 delays flowering through direct inhibition of FLOWERING LOCUS T expression in Arabidopsis. 2021 , 63, 1712-1723	5
572	Sulfur deficiency-induced genes affect seed protein accumulation and composition under sulfate deprivation. 2021 , 187, 2419-2434	5
571	SET DOMAIN GROUP 721 protein functions in saline-alkaline stress tolerance in the model rice variety Kitaake. 2021 , 19, 2576-2588	1
570	CLCd and CLCf act redundantly at the trans-Golgi network/early endosome and prevent acidification of the Golgi stack. 2021 , 134,	0
569	The DREAM complex represses growth in response to DNA damage in. 2021 , 4,	4
568	MuWU: Mutant-seq library analysis and annotation. 2021 ,	
567	Crosstalk between the chloroplast protein import and SUMO systems revealed through genetic and molecular investigation in. 2021 , 10,	1
566	A Role for Inositol Pyrophosphates in the Metabolic Adaptations to Low Phosphate in. 2021 , 11,	1
565	A Course-Based Undergraduate Research Experience in CRISPR-Cas9 Experimental Design to Support Reverse Genetic Studies in. 2021 , 22,	
564	Discovery of photosynthesis genes through whole-genome sequencing of acetate-requiring mutants of Chlamydomonas reinhardtii. 2021 , 17, e1009725	4
563	Redundant roles of four ZIP family members in zinc homeostasis and seed development in Arabidopsis thaliana. 2021 , 108, 1162-1173	2
562	The APC/CCCS52A2 E3-Ligase Complex targeted PKN1 Protein is a Plant Lineage-Specific Driver of Cell Division.	
561	Genome-wide Excision Repair Map of Cyclobutane Pyrimidine Dimers in Arabidopsis and the Roles of CSA1 and CSA2 Proteins in Transcription-coupled Repair. 2021 ,	
560	Cytosolic phosphofructokinases are important for sugar homeostasis in leaves of Arabidopsis thaliana. 2021 ,	1
559	Sterol Glucosyltransferases Tailor Polysaccharide Accumulation in Arabidopsis Seed Coat Epidermal Cells. 2021 , 10,	1

558	A BTB-TAZ protein is required for gene activation by Cauliflower mosaic virus 35S multimerized enhancers. 2021 ,	0
557	Myosin XI-B is involved in the transport of vesicles and organelles in pollen tubes of <i>Arabidopsis thaliana</i> . 2021 , 108, 1145-1161	1
556	AtPng1 knockout mutant of <i>Arabidopsis thaliana</i> shows a juvenile phenotype, morpho-functional changes, altered stress response and cell wall modifications. 2021 , 167, 11-21	0
555	AtCGL160 recruits chloroplast coupling factor 1.	
554	Insight into ripening-associated transcription factors in tomato: A review. 2021 , 288, 110363	2
553	Knockout of FER decreases cadmium concentration in roots of <i>Arabidopsis thaliana</i> by inhibiting the pathway related to iron uptake. 2021 , 798, 149285	1
552	The PELOTA-HBS1 Complex Orchestrates mRNA Translation Surveillance and PDK1-mediated Plant Growth and Development.	
551	Model Plants for Understanding Evolution. 1-9	2
550	Nuclear-encoded sigma factor 6 (SIG6) is involved in the block of greening response in <i>Arabidopsis thaliana</i> . 2020 , 107, 329-338	3
549	Approaches for identification of fungal genes essential for plant disease. 2004 , 26, 85-103	3
548	Mechanisms of T-DNA integration. 2008 , 395-440	10
547	Transposon Resources for Forward and Reverse Genetics in Maize. 2009 , 561-584	23
546	Cell Wall Genomics in the Recombinogenic Moss <i>Physcomitrella patens</i> . 2011 , 241-261	4
545	The Role of Nucleus- and Chloroplast-Encoded Factors in the Synthesis of the Photosynthetic Apparatus. 2007 , 145-165	8
544	Biotechnology Approaches To Engineering Drought Tolerant Crop. 2007 , 333-357	7
543	Transformation as a Tool for Genetic Analysis in <i>Populus</i> . 2010 , 113-133	6
542	Natural Variation in <i>Arabidopsis thaliana</i> . 2011 , 123-151	3
541	Identification of <i>Arabidopsis</i> knockout lines for genes of interest. 2014 , 1110, 347-62	5

540	Brachypodium distachyon. 2015 , 1223, 17-33	14
539	A user's guide to the Arabidopsis T-DNA insertion mutant collections. 2015 , 1284, 323-42	52
538	Multipurpose transposon insertion libraries for large-scale analysis of gene function in yeast. 2008 , 416, 117-29	6
537	Northern blotting techniques for small RNAs. 2010 , 631, 87-107	10
536	Analysing cellulose biosynthesis with confocal microscopy. 2011 , 715, 141-52	2
535	Chloroplast phenomics: systematic phenotypic screening of chloroplast protein mutants in Arabidopsis. 2011 , 775, 161-85	9
534	Tailor-made mutations in Arabidopsis using zinc finger nucleases. 2014 , 1062, 193-209	7
533	Arabidopsis database and stock resources. 2014 , 1062, 65-96	9
532	A growth quantification assay for Hyaloperonospora arabidopsidis isolates in Arabidopsis thaliana. 2014 , 1127, 145-58	8
531	Transposon Tagging and Reverse Genetics. 2009 , 143-159	10
530	Integration of Agrobacterium T-DNA in Plant Cells. 2009 , 157-186	3
529	Self-Incompatibility in the Brassicaceae. 2014 , 245-254	1
528	Functional Analysis of the Synechococcus elongatus PCC 7942 Genome. 2012 , 119-137	9
527	Molecular Approaches for Harvesting Natural Diversity for Crop Improvement. 2020 , 67-169	6
526	T-protein is present in large excess over the other proteins of the glycine cleavage system in leaves of Arabidopsis. 2018 , 247, 41-51	18
525	An Arabidopsis DISEASE RELATED NONSPECIFIC LIPID TRANSFER PROTEIN 1 is required for resistance against various phytopathogens and tolerance to salt stress. 2020 , 753, 144802	8
524	Non-specific activities of the major herbicide-resistance gene BAR. 2017 , 3, 937-945	23
523	CRISPR-Cas9-mediated induction of heritable chromosomal translocations in Arabidopsis. 2020 , 6, 638-645	53

522	Arabidopsis 4-COUMAROYL-COA LIGASE 8 contributes to the biosynthesis of the benzenoid ring of coenzyme Q in peroxisomes. 2019 , 476, 3521-3532	10
521	Fine tuning chloroplast movements through physical interactions between phototropins. 2016 , 67, 4963-78	16
520	Arabidopsis ECERIFERUM2-LIKEs Are Mediators of Condensing Enzyme Function. 2021 , 61, 2126-2138	5
519	Anno genominis XX: 20 years of Arabidopsis genomics. 2021 , 33, 832-845	5
518	Transport of UDP-rhamnose by URG2, URG4, and URG6 modulates rhamnogalacturonan-I length. 2021 , 185, 914-933	3
517	The Sequence of 1504 Mutants in the Model Rice Variety Kitaake Facilitates Rapid Functional Genomic Studies.	1
516	A centrifuge-based method for identifying novel genetic traits that affect root-substrate adhesion in Arabidopsis thaliana.	2
515	RAD54 is essential for RAD51-mediated repair of meiotic DSB in Arabidopsis.	1
514	Fibrillin 2 interacts with other proteins to protect photosystem II against abiotic stress in Arabidopsis thaliana.	1
513	LRX Proteins play a crucial role in pollen grain and pollen tube cell wall development.	2
512	Arabidopsis inositol phosphate kinases, IPK1 and ITPK1, constitute a metabolic pathway in maintaining phosphate homeostasis.	2
511	A variably imprinted epiallele impacts seed development.	2
510	Identification of the Calmodulin-dependent NAD ⁺ kinase sustaining the elicitor-induced oxidative burst in plants.	1
509	PBS3 and EPS1 complete salicylic acid biosynthesis from isochorismate in Arabidopsis.	8
508	Plant mixed lineage kinase domain-like proteins limit biotrophic pathogen growth.	6
507	The ALBA RNA-binding proteins function redundantly to promote growth and flowering in Arabidopsis.	1
506	Light activates the translational regulatory GCN2 kinase via reactive oxygen species emanating from the chloroplast.	1
505	The contribution of PIP2-type aquaporins to photosynthesis in response to increased vapour pressure deficit.	1

504	Auxin and Flower Development: A Blossoming Field. 2021 , 13,	5
503	Trimethylguanosine Synthase1 (TGS1) Is Essential for Chilling Tolerance. 2017 , 174, 1713-1727	7
502	Methylation of protein phosphatase 2A-Influence of regulators and environmental stress factors. 2017 , 40, 2347-2358	12
501	Effects of Arabidopsis Ku80 deletion on the integration of the left border of T-DNA into plant chromosomal DNA via Agrobacterium tumefaciens. 2020 , 95, 173-182	1
500	Embryo-lethal phenotypes in early abp1 mutants are due to disruption of the neighboring BSM gene. 2015 , 4, 1104	31
499	EIN3 and ORE1 Accelerate Degreening during Ethylene-Mediated Leaf Senescence by Directly Activating Chlorophyll Catabolic Genes in Arabidopsis. 2015 , 11, e1005399	177
498	MET18 Connects the Cytosolic Iron-Sulfur Cluster Assembly Pathway to Active DNA Demethylation in Arabidopsis. 2015 , 11, e1005559	33
497	Chloroplast RNA-Binding Protein RBD1 Promotes Chilling Tolerance through 23S rRNA Processing in Arabidopsis. 2016 , 12, e1006027	28
496	PARAQUAT TOLERANCE3 Is an E3 Ligase That Switches off Activated Oxidative Response by Targeting Histone-Modifying PROTEIN METHYLTRANSFERASE4b. 2016 , 12, e1006332	17
495	NUCLEAR FACTOR Y, Subunit C (NF-YC) Transcription Factors Are Positive Regulators of Photomorphogenesis in Arabidopsis thaliana. 2016 , 12, e1006333	36
494	The RTR Complex Partner RMI2 and the DNA Helicase RTEL1 Are Both Independently Involved in Preserving the Stability of 45S rDNA Repeats in Arabidopsis thaliana. 2016 , 12, e1006394	22
493	T-DNA-genome junctions form early after infection and are influenced by the chromatin state of the host genome. 2017 , 13, e1006875	21
492	POWERDRESS-mediated histone deacetylation is essential for thermomorphogenesis in Arabidopsis thaliana. 2018 , 14, e1007280	54
491	NIGT1 family proteins exhibit dual mode DNA recognition to regulate nutrient response-associated genes in Arabidopsis. 2020 , 16, e1009197	4
490	A systems biology approach identifies a R2R3 MYB gene subfamily with distinct and overlapping functions in regulation of aliphatic glucosinolates. 2007 , 2, e1322	255
489	DICER-LIKE2 plays a primary role in transitive silencing of transgenes in Arabidopsis. 2008 , 3, e1755	126
488	Heat stress enhances the accumulation of polyadenylated mitochondrial transcripts in Arabidopsis thaliana. 2008 , 3, e2889	19
487	Crosstalks between myo-inositol metabolism, programmed cell death and basal immunity in Arabidopsis. 2009 , 4, e7364	82

486	Dual localized AtHscB involved in iron sulfur protein biogenesis in Arabidopsis. 2009 , 4, e7662	26
485	ATL9, a RING zinc finger protein with E3 ubiquitin ligase activity implicated in chitin- and NADPH oxidase-mediated defense responses. 2010 , 5, e14426	71
484	Trans-repression of gene activity upstream of T-DNA tagged RLK902 links Arabidopsis root growth inhibition and downy mildew resistance. 2011 , 6, e19028	4
483	Structure-function analysis of STRUBBELIG, an Arabidopsis atypical receptor-like kinase involved in tissue morphogenesis. 2011 , 6, e19730	38
482	Altered germination and subcellular localization patterns for PUB44/SAUL1 in response to stress and phytohormone treatments. 2011 , 6, e21321	35
481	The stromal processing peptidase of chloroplasts is essential in Arabidopsis, with knockout mutations causing embryo arrest after the 16-cell stage. 2011 , 6, e23039	30
480	Arabidopsis ovate family proteins, a novel transcriptional repressor family, control multiple aspects of plant growth and development. 2011 , 6, e23896	77
479	A fasciclin-like arabinogalactan-protein (FLA) mutant of Arabidopsis thaliana, fla1, shows defects in shoot regeneration. 2011 , 6, e25154	60
478	Carbon dynamics, development and stress responses in Arabidopsis: involvement of the APL4 subunit of ADP-glucose pyrophosphorylase (starch synthesis). 2011 , 6, e26855	12
477	Cell expansion-mediated organ growth is affected by mutations in three EXIGUA genes. 2012 , 7, e36500	24
476	Arabidopsis thaliana AUCSIA-1 regulates auxin biology and physically interacts with a kinesin-related protein. 2012 , 7, e41327	13
475	High-to-low CO ₂ acclimation reveals plasticity of the photorespiratory pathway and indicates regulatory links to cellular metabolism of Arabidopsis. 2012 , 7, e42809	66
474	Mediator subunit18 controls flowering time and floral organ identity in Arabidopsis. 2013 , 8, e53924	41
473	Arabidopsis DEAD-box RNA helicase UAP56 interacts with both RNA and DNA as well as with mRNA export factors. 2013 , 8, e60644	41
472	The TMK subfamily of receptor-like kinases in Arabidopsis display an essential role in growth and a reduced sensitivity to auxin. 2013 , 8, e60990	55
471	Loss of the Arabidopsis thaliana P _{ATPase} ALA3 reduces adaptability to temperature stresses and impairs vegetative, pollen, and ovule development. 2013 , 8, e62577	23
470	Elemental concentrations in the seed of mutants and natural variants of Arabidopsis thaliana grown under varying soil conditions. 2013 , 8, e63014	15
469	Arabidopsis TRANSCURVATA1 encodes NUP58, a component of the nucleopore central channel. 2013 , 8, e67661	16

468	Antagonistic regulation of flowering time through distinct regulatory subunits of protein phosphatase 2A. 2013 , 8, e67987	23
467	Evolution of chloroplast J proteins. 2013 , 8, e70384	24
466	Time- and cost-efficient identification of T-DNA insertion sites through targeted genomic sequencing. 2013 , 8, e70912	25
465	Analysis of essential Arabidopsis nuclear genes encoding plastid-targeted proteins. 2013 , 8, e73291	36
464	Mu-seq: sequence-based mapping and identification of transposon induced mutations. 2013 , 8, e77172	35
463	GreenGate---a novel, versatile, and efficient cloning system for plant transgenesis. 2013 , 8, e83043	178
462	Efficient gene silencing mediated by tobacco rattle virus in an emerging model plant physalis. 2014 , 9, e85534	12
461	Development of a Cucumis sativus TILLinG platform for forward and reverse genetics. 2014 , 9, e97963	29
460	Metabolome analysis of Arabidopsis thaliana roots identifies a key metabolic pathway for iron acquisition. 2014 , 9, e102444	96
459	RNA-seq analysis of the effect of kanamycin and the ABC transporter AtWBC19 on Arabidopsis thaliana seedlings reveals changes in metal content. 2014 , 9, e109310	5
458	Pale-green phenotype of atl31atl6 double mutant leaves is caused by disruption of 5-aminolevulinic acid biosynthesis in Arabidopsis thaliana. 2015 , 10, e0117662	14
457	Alanine aminotransferase variants conferring diverse NUE phenotypes in Arabidopsis thaliana. 2015 , 10, e0121830	22
456	Functional roles of three cutin biosynthetic acyltransferases in cytokinin responses and skotomorphogenesis. 2015 , 10, e0121943	5
455	Genome-Wide Analysis of the AP2/ERF Transcription Factors Family and the Expression Patterns of DREB Genes in Moso Bamboo (<i>Phyllostachys edulis</i>). 2015 , 10, e0126657	68
454	The Arabidopsis KH-Domain RNA-Binding Protein ESR1 Functions in Components of Jasmonate Signalling, Unlinking Growth Restraint and Resistance to Stress. 2015 , 10, e0126978	35
453	Decreased Expression of a Gene Caused by a T-DNA Insertion in an Adjacent Gene in Arabidopsis. 2016 , 11, e0147911	4
452	The Early-Acting Peroxin PEX19 Is Redundantly Encoded, Farnesylated, and Essential for Viability in Arabidopsis thaliana. 2016 , 11, e0148335	11
451	AtGRP3 Is Implicated in Root Size and Aluminum Response Pathways in Arabidopsis. 2016 , 11, e0150583	19

450	Structure of a Berberine Bridge Enzyme-Like Enzyme with an Active Site Specific to the Plant Family Brassicaceae. 2016 , 11, e0156892	20
449	Functional Characterization of CYP94-Genes and Identification of a Novel Jasmonate Catabolite in Flowers. 2016 , 11, e0159875	30
448	PIF4 and ELF3 Act Independently in Arabidopsis thaliana Thermoresponsive Flowering. 2016 , 11, e0161791	11
447	Homologous Recombination Defective Arabidopsis Mutants Exhibit Enhanced Sensitivity to Abscisic Acid. 2017 , 12, e0169294	7
446	Molecular and Functional Characterization of GR2-R1 Event Based Backcross Derived Lines of Golden Rice in the Genetic Background of a Mega Rice Variety Swarna. 2017 , 12, e0169600	24
445	SnRK1 activates autophagy via the TOR signaling pathway in Arabidopsis thaliana. 2017 , 12, e0182591	98
444	Sugarcane Mosaic Virus-Based Gene Silencing in. 2017 , 15, 260-267	3
443	Glucose uptake to guard cells via STP transporters provides carbon sources for stomatal opening and plant growth. 2020 , 21, e49719	19
442	Molecular characteristics, recombinant expression and activity detection of OsGSTL1 from rice. 2014 , 57, 70-76	1
441	Variation in GFP Gene Expression in Arabidopsis thaliana Monoclonal Cell Lines. 2019 , 35, 58-67	2
440	Accumulation of the Auxin Precursor Indole-3-Acetamide Curtails Growth through the Repression of Ribosome-Biogenesis and Development-Related Transcriptional Networks. 2021 , 22,	3
439	[Progresses on plant AP2/ERF transcription factors]. 2012 , 34, 835-47	9
438	Expressions of Three Members of ACS Gene Family Induced by Ethephon Relationship of Ethylene Production and Sugar Accumulation in Sugarcane Stalks. 2008 , 34, 418-422	5
437	The Response of AtCER to Oxidative Stress in Arabidopsis thaliana. 2010 , 31, 326-334	2
436	Plant and microbial research seeks biofuel production from lignocellulose. 2009 , 63, 178-184	4
435	The Potential Wheat Signaling Pathways in Response to Abiotic Stress. 2007 , 2, 295-302	3
434	Phenotypic Characterization of the Arabidopsis ufm1 (Ubiquitin Fold Modifier) Gene Involved in Seed Development. 2016 , 12, 10-19	1
433	Investigation of the Effect of AtWIN1/SHN1 Overexpression on Poplar Trees. 2017 , 12, 1-13	1

432	Analysis of junction between T-DNA and plant genome in insect resistance GM Chinese cabbage. 2008 , 35, 101-108	3
431	Progress in Genetic Manipulation of the Brassicaceae. 2012 , 39, 1-12	2
430	Message in a Bottle: Chemical Biology of Induced Disease Resistance in Plants. 2008 , 24, 245-268	34
429	A versatile method to prevent transcriptional gene silencing in <i>Arabidopsis thaliana</i> . 2011 , 28, 515-519	3
428	ATL54, a RING-H2 domain protein selected by a gene co-expression network analysis, is associated with secondary cell wall formation in <i>Arabidopsis</i> . 2013 , 30, 169-177	9
427	<i>Arabidopsis</i> glycerol-3-phosphate permease 4 is localized in the plastids and involved in the accumulation of seed oil. 2014 , 31, 159-165	5
426	Expression analysis of gene trap lines and mapping of donor loci for Dissociation transposition in <i>Arabidopsis</i> . 2007 , 24, 467-479	5
425	ARTRA: a new database of the <i>Arabidopsis</i> transcriptome and gene-specific sequences for microarray probes and RNAi triggers. 2009 , 26, 161-165	1
424	Accurate timekeeping is controlled by a cycling activator in <i>Arabidopsis</i> . 2013 , 2, e00473	119
423	The transcriptional regulator BZR1 mediates trade-off between plant innate immunity and growth. 2013 , 2, e00983	144
422	Combining growth-promoting genes leads to positive epistasis in <i>Arabidopsis thaliana</i> . 2014 , 3, e02252	27
421	Natural genetic variation in <i>Arabidopsis thaliana</i> defense metabolism genes modulates field fitness. 2015 , 4,	64
420	PCH1 integrates circadian and light-signaling pathways to control photoperiod-responsive growth in <i>Arabidopsis</i> . 2016 , 5, e13292	53
419	The MBD7 complex promotes expression of methylated transgenes without significantly altering their methylation status. 2017 , 6,	12
418	Integration of two RAB5 groups during endosomal transport in plants. 2018 , 7,	16
417	Drought adaptation in by extensive genetic loss-of-function. 2018 , 7,	35
416	Dynamic ubiquitination determines transcriptional activity of the plant immune coactivator NPR1. 2019 , 8,	14
415	Genetic transformation of , a resource for studying chytrid biology and evolutionary cell biology. 2020 , 9,	18

414	Oomycete small RNAs bind to the plant RNA-induced silencing complex for virulence. 2020 , 9,	40
413	auxin biosynthetic genes are required for Arabidopsis shade avoidance. 2016 , 4, e2574	41
412	Recombination between T-DNA insertions to cause chromosomal deletions in is a rare phenomenon. 2018 , 6, e5076	4
411	TRANSPARENT TESTA GLABRA 1 participates in flowering time regulation in. 2020 , 8, e8303	4
410	Multiple immunity-related genes control susceptibility of to the parasitic weed. 2020 , 8, e9268	5
409	Cinnamoyl coA:NADP oxidoreductase like 1 regulates abscisic acid response by modulating phaseic acid homeostasis in Arabidopsis thaliana. 2021 ,	1
408	Evolutionarily informed machine learning enhances the power of predictive gene-to-phenotype relationships. 2021 , 12, 5627	4
407	Genome-wide identification, phylogeny and expression analysis of AP2/ERF transcription factors family in sweet potato. 2021 , 22, 748	2
406	A FLASH pipeline for arrayed CRISPR library construction and the gene function discovery of rice receptor-like kinases. 2021 ,	3
405	Arabidopsis PLD α 1 and PLD α 2 localize to post-Golgi membrane compartments in a partially overlapping manner. 2021 , 108, 31	
404	Structural Maintenance of Chromosomes 5/6 Complex Is Necessary for Tetraploid Genome Stability in. 2021 , 12, 748252	3
403	The ARRE RING-Type E3 Ubiquitin Ligase Negatively Regulates Cuticular Wax Biosynthesis in by Controlling ECERIFERUM1 and ECERIFERUM3 Protein Levels. 2021 , 12, 752309	0
402	Arabidopsis bioinformatics: tools and strategies. 2021 ,	1
401	Control of shoot stem cell homeostasis by two antagonistic CLE peptide signalling pathways. 2021 , 10,	10
400	Arabidopsis fibrillin 1-2 subfamily exerts their functions via specific protein-protein interactions. 2021 ,	1
399	Identification of binding interaction between Curcumin derivative and PERK13 Proline rich receptor like protein kinase protein using In silico docking techniques Π o help plants tackle salt water.	
398	Specific interaction of an RNA-binding protein with the 3'-UTR of its target mRNA is critical to oomycete sexual reproduction. 2021 , 17, e1010001	1
397	A dedicated flavin-dependent monooxygenase catalyzes the hydroxylation of demethoxyubiquinone into ubiquinone (coenzyme Q) in Arabidopsis. 2021 , 297, 101283	2

- 396 Isolation of Arabidopsis mutants that have altered sensitivity to staurosporine and ABA. **2005**, 22, 303-310
- 395 [Host factors and their relevance to virus infection in plants]. **2006**, 56, 155-63
- 394 The Study of Plant Natural Product Biosynthesis in the Pangenomics and Genomics Eras. **2006**, 203-220
- 393 Transcription Factors Regulating Plant Defense Responses. **2006**, 159-205
- 392 Knockout of AtFKBP13 Proves Degeneration of Prolyl Isomerase Activity in the Thylakoid Lumen of Arabidopsis thaliana. **2008**, 1129-1132
- 391 Structural and Functional Characterization of Leaf-Type Ferredoxin-NADP⁺-Oxidoreductase Isoforms in Arabidopsis thaliana. **2008**, 933-936
- 390 Intracellular Transport of Agrobacterium T-DNA. **2008**, 365-394
- 389 Molecular and Genetic Analyses of Transgenic Plants. **2008**, 385-401
- 388 Using reverse genetics to develop small knockout collections for specific biological questions. **2009**, 495, 1-10 1
- 387 Genome Wide Approaches in Natural Product Research. **2009**, 475-503
- 386 Unravelling Gene Function Through Mutagenesis. **2010**, 437-467
- 385 Phenome analysis of root development in Arabidopsis. **2010**, 27, 345-347
- 384 Rice functional genomics using T-DNA mutants. **2010**, 37, 133-143 1
- 383 Roadmap of Genomics Research in the 21st Century. **2010**, 571-582
- 382 Agrobacterium tumefaciens-Mediated Transformation of Ectomycorrhizal Fungi. **2011**, 123-141
- 381 Screening or selection for chloroplast biogenesis mutants of Arabidopsis, following chemical or insertional mutagenesis. **2011**, 774, 3-18 1
- 380 Functional Genomics Dissection of Photosynthetic Mechanisms in Arabidopsis thaliana. **2012**, 177-198 1
- 379 Construction of the RNAi vector for Citrus ERF1 gene. **2011**, 37, 385-390

- 378 A one-time inducible transposon to create knockout mutants in rice. **2012**, 847, 369-77
- 377 Functional Genomics and Transcriptomics in Musa. **2012**, 156-180
- 376 Comparative Photosynthetic Analyses of Three Widely Used Arabidopsis Ecotypes. **2013**, 583-585
- 375 Do State Transitions Control CEF1 in Higher Plants?. **2013**, 286-289 1
- 374 Target Identification of Biologically Active Small Molecules. 233-246
- 373 The Use of Small Molecules to Dissect Cell Wall Biosynthesis and Manipulate the Cortical Cytoskeleton. 203-217
- 372 Arabidopsis PIP5K2 Is Involved in Lateral Root Development Through Regulating Auxin Accumulation. **2014**, 29-43
- 371 Arabidopsis PIP5K2 Is Involved in Root Gravitropism Through Regulation of Polar Auxin Transport. **2014**, 45-62
- 370 Forward and Reverse Genetics: The LORE1 Retrotransposon Insertion Mutants. **2014**, 221-227 1
- 369 Molecular Life Sciences. **2014**, 1-15
- 368 Arabidopsis PIP5K2 Is Involved in Salt Tolerance. **2014**, 63-77
- 367 Organization and function of shoot apical meristem affecting growth and development in plants. **2014**, 41, 180-193
- 366 Veränderungen im Genom: Mutationen. **2015**, 399-458
- 365 Genetic Modification of the Pigment Optical Density. **2015**, 69-82
- 364 An efficient gene targeting system using homologous recombination in plants. **2015**, 42, 154-160
- 363 ELF3 polyQ variation in Arabidopsis thaliana reveals PIF4-independent role in thermoresponsive flowering. 0
- 362 Interplay of vacuolar transporters for coupling primary and secondary active transport. **2016**, 3, 479-500 0
- 361 PARAQUAT TOLERANCE3 is an E3 ligase and acts as a negative regulator of oxidative stress response.

- 360 1Chapter 1 Molecular Approaches to the Study of Plant Development. **2016**, 173-196
- 359 Plant functional genomics: Approaches and applications. **2016**, 157-186 2
- 358 Phenotyping Zebrafish. **2016**, 75-95
- 357 INTRODUCTION. **2016**, 10-11
- 356 The variable ELF3 polyglutamine tract mediates complex epistatic interactions in *Arabidopsis thaliana*.
- 355 NUCLEAR FACTOR Y, subunit C (NF-YC) transcription factors are positive regulators of photomorphogenesis in *Arabidopsis thaliana*. 2
- 354 The Pollen Membrane Proteome. **2017**, 293-318
- 353 T-DNA integration is rapid and influenced by the chromatin state of the host genome.
- 352 Discovery and reduction of nonspecific activities of the major herbicide-resistance gene BAR.
- 351 Characterization and Expression Analysis of an AP2 Gene from *Ginkgo biloba*. **2017**, 16, 108-115
- 350 Massive variation of short tandem repeats with functional consequences across strains of *Arabidopsis thaliana*. 1
- 349 Flavonol rhamnosylation indirectly modifies the cell wall defects of RHAMNOSE BIOSYNTHESIS1 mutants by altering rhamnose flux.
- 348 POWERDRESS-mediated histone deacetylation is essential for thermomorphogenesis in *Arabidopsis thaliana*.
- 347 CLASP stabilizes microtubule plus ends created by serving to drive cortical array reorientation. 0
- 346 PWO1 interacts with PcG proteins and histones to regulate *Arabidopsis* flowering and development.
- 345 The mutant line contains a loss of function allele in the gene caused by a remnant inactivated selection marker—a cautionary tale. **2017**, 5, e4134
- 344 A molecular network for functional versatility of HECATE transcription factors.
- 343 Mapping and dynamics of regulatory DNA in maturing seeds.

342 Molecular Life Sciences. **2018**, 905-918

341 Adaptive Reduction of Male Gamete Number in a Selfing Species.

340 The complex architecture of plant transgene insertions.

1

339 Arabidopsis thaliana as an Experimental Organism. 1-9

338 PII1: a protein involved in starch initiation that determines granule number and size in Arabidopsis chloroplast.

337 General Themes of Molecular Stress Physiology. **2019**, 9-55

0

336 Drought adaptation in nature by extensive genetic loss-of-function.

335 Effects of green seaweed extract on Arabidopsis early development suggest roles for hormone signalling in plant responses to algal fertilisers.

334 CrRLK1L receptor-like kinases HERCULES RECEPTOR KINASE 1 and ANJEA are female determinants of pollen tube reception.

333 New Drosophila long-term memory genes revealed by assessing computational function prediction methods.

3

332 A seed resource for screening functionally redundant genes and isolation of new mutants impaired in CO₂ and ABA responses.

331 A role for the auxin precursor anthranilic acid in root gravitropism via regulation of PIN-FORMED protein polarity and relocalization in Arabidopsis.

330 Positive feedback and feedforward loops between PERIANTHIA, WUSCHEL-RELATED HOMEBOX5 and GRF-INTERACTING FACTOR 1 modulate gene expression and function in the Arabidopsis root.

329 Tissue morphogenesis mediated by the Arabidopsis receptor kinase STRUBBELIG involves a clathrin-dependent process.

328 Hypermorphic SERK1 mutations function via a SOBIR1 pathway to activate floral abscission signaling.

327 Arabidopsis TRM5 encodes a nuclear-localised bifunctional tRNA guanine and inosine-N1-methyltransferase that is important for growth.

326 Invasive DNA elements modify nuclear architecture by KNOT-Linked Silencing in plants.

325 Encyclopedia of Animal Cognition and Behavior. **2019**, 1-7

- 324 The striking flower-in-flower phenotype of *Arabidopsis thaliana* Nossen (No-0) is caused by a novel LEAFY allele. 1
- 323 Development of Flax (*Linum usitatissimum* L.) Mutant Populations for Forward and Reverse Genetics. **2019**, 145-165
- 322 A Novel Conversion of Roots into Organs with Shoot Stem Characteristics by Inducing Two Transcription Factors.
- 321 *Arabidopsis* *ml03* mutant plants exhibit spontaneous callose deposition and signs of early leaf senescence. 1
- 320 Functional characterization of genes mediating cell wall metabolism and responses to plant cell wall integrity impairment.
- 319 The circadian clock contributes to the long-term water use efficiency of *Arabidopsis*. 0
- 318 A Musashi-Related Protein is Essential for Gametogenesis in *Arabidopsis*. 1
- 317 Epistatic Transcription Factor Networks Differentially Modulate *Arabidopsis* Growth and Defense.
- 316 *Arabidopsis* glutathione reductase 2 is indispensable in plastids, while mitochondrial glutathione is safeguarded by additional reduction and transport systems. 0
- 315 Proteomic studies of the *Arabidopsis* TRAPP complexes reveal conserved organization and a novel plant-specific component with a role in plant development.
- 314 FRA1 modulates cortical microtubule localization of CMU proteins.
- 313 Nuclear encoded photosynthesis genes are specifically controlled by the NuA4 complex.
- 312 Analysis of the *Arabidopsis* organellar rhomboid At1g74140 transcript population uncovered splicing patterns different from its close relative At1g74130. 8, 1925
- 311 The TIFY Gene Family in Wheat and its Progenitors: Genome-wide Identification, Evolution and Expression Analysis. **2019**, 20, 371-388 2
- 310 *Arabidopsis* ROOT PHOTOTROPISM2 is a Light-Dependent Dynamic Modulator of Phototropin1.
- 309 CYCA3;4 Is a Post-Prophase Target of the APC/CCCS52A2 E3-Ligase Controlling Formative Cell Divisions in *Arabidopsis*.
- 308 Hormonal regulation of the BRC1-dependent strigolactone transcriptome involved in shoot branching responses. 1
- 307 Rerouting of ribosomal proteins into splicing in plant organelles.

- 306 Oposing functions of the plant TOPLESS gene family during SNC1-mediated autoimmunity.
- 305 Regulation of CYP94B1 by WRKY33 controls apoplastic barrier formation in the roots leading to salt tolerance.
- 304 Perturbations of the AMI1 IAM-amidohydrolase expression trigger plant stress responses in *Arabidopsis thaliana*.
- 303 Characterization of β Glutamyl Peptidases and β Glutamyl Cyclotransferases for Glutathione Degradation in *Arabidopsis*.
- 302 Specific PP2A Catalytic Subunits Are a Prerequisite for Positive Growth Effects in *Arabidopsis* Co-Cultivated with and. **2020**, 10,
- 301 Elucidation of genomic organizations of transgenic soybean plants through de novo genome assembly with short paired-end reads. **2021**, 41, 1 3
- 300 The nucleolar protein SAHY1 is involved in pre-rRNA processing and normal plant growth. **2021**, 185, 1039-1058 0
- 299 The nucellus: between cell elimination and sugar transport. **2021**, 185, 478-490 4
- 298 Functional and evolutionary analysis of the *Arabidopsis* 4R-MYB protein SNAPc4 as part of the SNAP complex. **2021**, 185, 1002-1020 2
- 297 An iron (II) dependent oxygenase performs the last missing step of plant lysine catabolism.
- 296 GAL4/GFP enhancer-trap lines for identification and manipulation of cells and tissues in developing *Arabidopsis* leaves. 0
- 295 Plastid gene expression is required for singlet oxygen-induced chloroplast degradation and cell death.
- 294 The *Arabidopsis* NOT4A E3 ligase coordinates PGR3 expression to regulate chloroplast protein translation.
- 293 Development of mPing-based Activation Tags for Crop Insertional Mutagenesis.
- 292 Crosstalk between chloroplast protein import and the SUMO system revealed through genetic and molecular investigation.
- 291 Ferroportin 3 is a dual-targeted mitochondrial/chloroplast iron exporter necessary for iron homeostasis in *Arabidopsis*.
- 290 Case studies for transcriptional profiling. **2007**, 97, 87-97
- 289 Regulatory Mechanisms for Floral Organ Identity Specification in *Arabidopsis thaliana*. **2006**, 533-547

- 288 The use of functional genomics to understand components of plant metabolism and the regulation occurring at molecular, cellular and whole plant levels. **2007**, 1-26
- 287 Describing Epigenomic Information in Arabidopsis. **2009**, 163-175
- 286 The interplay of HSP90s with YDA regulates main body axis formation during early embryogenesis in Arabidopsis.
- 285 Ubiquitylome Analysis Reveals a Central Role for the Ubiquitin-Proteasome System in Plant Innate Immunity.
- 284 Induction of GLUTAMINE DUMPER1 reveals a link between amino acid export, abscisic acid, and immune responses.
- 283 Combinations of maternal-specific repressive epigenetic marks in the endosperm control seed dormancy.
- 282 Spatial and Temporal Localization of SPIRRIG and WAVE/SCAR Reveal Roles for These Proteins in Actin-Mediated Root Hair Development. 0
- 281 Bioinformatic Tools in Arabidopsis Research. **2021**, 2200, 25-89 2
- 280 BRI1 Signaling in the Root is Mediated through the SERK1 and SERK3 Co-receptors. **2020**, 43,
- 279 The Arabidopsis transcription factors AtPHL1 and AtHB23 act together promoting carbohydrate transport from pedicel-silique nodes to seeds.. **2022**, 315, 111133 0
- 278 Heat-induced transposition leads to rapid drought resistance of Arabidopsis thaliana. 0
- 277 CFM6 is an Essential CRM Protein Required for the Splicing of nad5 Transcript in Arabidopsis Mitochondria. **2021**, 0
- 276 Comparative Transcriptomic Analysis Provides Novel Insights into the Blanched Stem of. **2021**, 10, 2
- 275 A genome-scale TF-DNA interaction network of transcriptional regulation of Arabidopsis primary and specialized metabolism. **2021**, 17, e10625 1
- 274 The Role of DNA Topoisomerase Binding Protein 1 (TopBP1) in Genome Stability in .. **2021**, 10, 2
- 273 Piriformospora indica recruits host-derived putrescine for growth promotion in plants. **2021**, 2
- 272 Asymmetric distribution of extracellular matrix proteins in seed coat epidermal cells of Arabidopsis is determined by polar secretion. **2021**, 5, e360
- 271 Gain-of-function mutations of AtNHX1 suppress sos1 salt sensitivity and improve salt tolerance in Arabidopsis. **2021**, 1, 1 0

- 270 Initiation of cytosolic plant purine nucleotide catabolism involves a monospecific xanthosine monophosphate phosphatase. **2021**, 12, 6846 0
- 269 Heterochromatin is a quantitative trait associated with spontaneous epiallele formation. **2021**, 12, 6958 1
- 268 Identification, evolutionary analysis and functional diversification of RAV gene family in cotton (*G. hirsutum* L.). **2021**, 255, 14 0
- 267 Transient gene expression system in zoysiagrass leaf mesophyll protoplasts. 1 0
- 266 Role of the mismatch repair protein MSH7 in Arabidopsis adaptation to acute salt stress. **2021**, 169, 280-290 0
- 265 The CLASSY family controls tissue-specific DNA methylation patterns in Arabidopsis.. **2022**, 13, 244 3
- 264 A Transposon-Based Activation Tag System for Functional Genomics in Cereals: Detection of Mping-Based Activation Tag Mobilization in Wheat. **2022**, 195-207
- 263 Abscisic Acid Drives Stomatal Closure through Increases in Hydrogen Peroxide in Distinct Subcellular Compartments Including Mitochondria. 0
- 262 Arabidopsis CIA2 and CIL have distinct and overlapping functions in regulating chloroplast and flower development.. **2022**, 6, e380 0
- 261 Disruption of the N-Acetyltransferase NatB Causes Sensitivity to Reductive Stress in .. **2021**, 12, 799954
- 260 NuA4 and H2A.Z control environmental responses and autotrophic growth in Arabidopsis.. **2022**, 13, 277 3
- 259 Phloem iron remodels root development in response to ammonium as the major nitrogen source.. **2022**, 13, 561 1
- 258 A mitochondrial ADXR-ADX-P450 electron transport chain is essential for maternal gametophytic control of embryogenesis in .. **2022**, 119, 1 1
- 257 CFI 25 Subunit of Cleavage Factor I is Important for Maintaining the Diversity of 3' UTR Lengths in Arabidopsis thaliana (L.) Heynh.. **2022**, 0
- 256 The SNF5-type protein BUSHY regulates seed germination via the gibberellin pathway and is dependent on HUB1 in Arabidopsis.. **2022**, 255, 34 0
- 255 Systematic histone H4 replacement in Arabidopsis thaliana reveals a role for H4R17 in regulating flowering time. 0
- 254 Cold Tolerance is Governed by Diverse Genetic Mechanisms Including Those Regulated by NB-LRR-type Receptor Proteins in Arabidopsis. 0
- 253 Mutations in TIC100 impair and repair chloroplast protein import and impact retrograde signalling. 0

- 252 Arabidopsis pavement cell shape formation involves spatially confined ROPGAP regulators.. **2022**, 3
- 251 Contrasting gene-level signatures of selection with reproductive fitness.. **2021**,
- 250 Overexpression mutants reveal a role for a chloroplast MPD protein in regulation of reactive oxygen species during chilling in Arabidopsis.. **2022**, 0
- 249 Genome-wide identification and expression profiling analysis of maize AP2/ERF superfamily genes reveal essential roles in abiotic stress tolerance.. **2022**, 23, 125 2
- 248 Chloroplast Protein Tic55 Involved in Dark-Induced Senescence through AtbHLH/AtWRKY-ANAC003 Controlling Pathway of .. **2022**, 13, 0
- 247 Arabidopsis GOLD36/MVP1/ERMO3 is required for powdery mildew penetration resistance and proper targeting of the PEN3 transporter.. **2022**,
- 246 Unraveling the Dynamic Integration of Auxin, Brassinosteroid and Gibberellin in Early Shade-Induced Hypocotyl Elongation. 1
- 245 Study of natural diversity in response to a key pathogenicity regulator of *Ralstonia solanacearum* reveals new susceptibility genes in *Arabidopsis thaliana*.. **2021**, 2
- 244 Tnt1 Insertional Mutagenesis in *Medicago truncatula* for Gene Function Analysis. **2022**, 23-33
- 243 Protoplast for Gene Functional Analysis in Arabidopsis.. **2022**, 2464, 29-47
- 242 An inosine triphosphate pyrophosphatase safeguards nucleic acids from aberrant purine nucleotides and prevents a constitutive salicylic acid response.
- 241 Trends in Arabidopsis Research post genome sequencing- A Scientometric study.
- 240 Dynamics of Etiolation Monitored by Seedling Morphology, Carotenoid Composition, Antioxidant Level, and Photoactivity of Protochlorophyllide in .. **2021**, 12, 772727
- 239 The Structure of the PEX4-PEX22 Peroxin Complex-Insights Into Ubiquitination at the Peroxisomal Membrane.. **2022**, 10, 838923 0
- 238 Identification of Mutants With Increased Susceptibility to .. **2022**, 13, 799142 0
- 237 Complex Changes in Membrane Lipids Associated with the Modification of Autophagy in .. **2022**, 12, 0
- 236 Evolutionarily conserved bacterial effectors hijack abscisic acid signaling to induce an aqueous environment in the apoplast.. **2022**, 4
- 235 Comparison between Germinated Seed and Isolated Microspore EMS Mutagenesis in Chinese Cabbage (*Brassica rapa* L. ssp. *pekinensis*). **2022**, 8, 232 0

- 234 The DC1 domain protein BINUCLEATE POLLEN is required for pollen development in Arabidopsis.
- 233 Interplay between Genotype, Plant Growth and Rhizosphere Colonization by Phytobeneficial Phenazine-Producing .. **2022**, 10, 0
- 232 The APAF1_C/WD40 repeat domain-encoding gene from the sea lettuce *Ulva mutabilis* sheds light on the evolution of NB-ARC domain-containing proteins in green plants.. **2022**, 255, 76 1
- 231 The phytochrome interacting proteins ERF55 and ERF58 repress light-induced seed germination in *Arabidopsis thaliana*.. **2022**, 13, 1656 0
- 230 Genetically-determined variations in photosynthesis indicate roles for specific fatty acid species in chilling responses.. **2022**, 1
- 229 The nitrate-inducible NAC transcription factor NAC056 controls nitrate assimilation and promotes lateral root growth in *Arabidopsis thaliana*.. **2022**, 18, e1010090 2
- 228 HPCA1 is required for systemic ROS and calcium cell-to-cell signaling and plant acclimation to stress. 1
- 227 Redundant PLETHORA activity promotes development of early embryonic cell lineages in *Arabidopsis*.
- 226 Insights on suppression of bacterial leaf spot by *Bacillus* cyclic lipopeptides via induced resistance in *Arabidopsis thaliana*. 1
- 225 Genetics as a key to improving crop photosynthesis.. **2022**, 1
- 224 Using CRISPR-Kill for organ specific cell elimination by cleavage of tandem repeats.. **2022**, 13, 1502 0
- 223 CRISPR-Cas9-mediated chromosome engineering in *Arabidopsis thaliana*.. **2022**, 0
- 222 Jasmonate Signaling Pathway Modulates Plant Defense, Growth, and Their Trade-Offs.. **2022**, 23, 2
- 221 Transcriptome, metabolome and suppressor analysis reveal an essential role for the ubiquitin-proteasome system in seedling chloroplast development.. **2022**, 22, 183 1
- 220 SEC1A is a major *Arabidopsis* Sec1/Munc18 gene in vesicle trafficking during pollen tube tip growth.. **2022**, 0
- 219 ATL8, a RING E3 ligase, modulates root growth and phosphate homeostasis in *Arabidopsis*.. **2022**, 179, 90-99 1
- 218 Creation of cotton mutant library based on linear electron accelerator radiation mutation.. **2022**, 30, 101228 0
- 217 Advances in gene editing without residual transgenes in plants. **2021**, 4

216	UV-B Irradiation Results in Changes in Elongation, Cell Cycle, and Decreased Endoreduplication Mediated by miR5642. 2021,	2
215	InMut-finder: a software tool for insertion identification in mutagenesis using Nanopore long reads.. 2021, 22, 908	0
214	Telomerase Interaction Partners-Insight from Plants.. 2021, 23,	1
213	Functional Differentiation among the Arabidopsis Phosphatidylinositol 4-Phosphate 5-Kinase Genes PIP5K1, PIP5K2 and PIP5K3.. 2022,	0
212	Genome-Wide Analysis of AP2/ERF Superfamily Genes in Contrasting Wheat Genotypes Reveals Heat Stress-Related Candidate Genes.. 2022, 13, 853086	0
211	The Dawn of Plant Molecular Biology: How Three Key Methodologies Paved the Way.. 2022, 2, e417	2
210	The Role of Nucleus- and Chloroplast-Encoded Factors in the Synthesis of the Photosynthetic Apparatus. 2007, 145-165	
209	Image_1.tif. 2019,	
208	Image_2.tif. 2019,	
207	Image_3.tif. 2019,	
206	Image_4.tif. 2019,	
205	Table_1.xlsx. 2019,	
204	Table_2.xlsx. 2019,	
203	Table_3.xlsx. 2019,	
202	Data_Sheet_1.doc. 2019,	
201	Data_Sheet_1.XLSX. 2018,	
200	Presentation_1.PDF. 2018,	
199	Presentation_1.pdf. 2018,	

198 Data_Sheet_1.PDF. **2019**,

197 Data_Sheet_2.PDF. **2019**,

196 Data_Sheet_3.PDF. **2019**,

195 Table_1.pdf. **2019**,

194 Table_2.pdf. **2019**,

193 Data_Sheet_1.pdf. **2018**,

192 Data_Sheet_1.docx. **2019**,

191 Presentation_1.zip. **2019**,

190 Presentation_1.pdf. **2018**,

189 DataSheet_1.pdf. **2020**,

188 DataSheet_2.pdf. **2020**,

187 DataSheet_3.pdf. **2020**,

186 DataSheet_4.pdf. **2020**,

185 DataSheet_5.pdf. **2020**,

184 Video_1.mov. **2020**,

183 Video_2.mov. **2020**,

182 DataSheet_1.pdf. **2020**,

181 Image_1.TIF. **2020**,

180 Image_2.TIF. 2020,

179 Image_3.TIF. 2020,

178 Table_1.DOCX. 2020,

177 Data_Sheet_1.DOCX. 2018,

176 Image_1.tif. 2020,

175 Image_2.tif. 2020,

174 Image_3.tif. 2020,

173 Image_4.tif. 2020,

172 Table_1.DOCX. 2020,

171 Data_Sheet_1.docx. 2020,

170 Image_1.TIF. 2020,

169 Image_10.TIF. 2020,

168 Image_11.TIF. 2020,

167 Image_2.TIF. 2020,

166 Image_3.TIF. 2020,

165 Image_4.TIF. 2020,

164 Image_5.TIF. 2020,

163 Image_6.TIF. 2020,

162 Image_7.TIF. 2020,

161 Image_8.TIF. 2020,

160 Image_9.TIF. 2020,

159 Image_1.TIF. 2018,

158 Image_2.tif. 2018,

157 Image_3.tif. 2018,

156 Image_4.TIF. 2018,

155 Image_5.pdf. 2018,

154 Table_1.pdf. 2018,

153 Table_2.pdf. 2018,

152 Table_3.pdf. 2018,

151 Presentation_1.pptx. 2020,

150 Table_1.xlsx. 2020,

149 Image_1.pdf. 2018,

148 Image_1.tif. 2020,

147 Image_2.tif. 2020,

146 Image_3.tif. 2020,

145 Table_1.xlsx. 2020,

144 DataSheet_1.pdf. **2020**,

143 Presentation_1.pptx. **2019**,

142 Table_1.xlsx. **2019**,

141 Image_1.jpg. **2019**,

140 Image_2.jpg. **2019**,

139 Image_3.jpg. **2019**,

138 Image_4.jpg. **2019**,

137 Table_1.docx. **2019**,

136 Data_Sheet_1.pdf. **2019**,

135 DataSheet_1.pdf. **2020**,

134 Table_1.DOCX. **2019**,

133 Data_Sheet_1.docx. **2020**,

132 Data_Sheet_1.pdf. **2020**,

131 Table_1.pdf. **2020**,

130 Efficient Generation of CRISPR/Cas9-Based Mutants Supported by Fluorescent Seed Selection in Different Arabidopsis Accessions.. **2022**, 2484, 161-182 ○

129 Robustness of organ morphology is associated with modules of co-expressed genes related to plant cell wall. ○

128 PILS proteins provide a homeostatic feedback on auxin signaling output.

127 Transcriptome Analysis Points to BES1 as a Transducer of Strigolactone Effects on Drought Memory in Arabidopsis thaliana.. **2022**, ○

- 126 Increased branching independent of strigolactone in cytokinin oxidase 2-overexpressing tomato is mediated by reduced auxin transport. **2022**, 2, 0
- 125 Extracellular ATP targets Arabidopsis RIBONUCLEASE 1 to suppress mycotoxin stress-induced cell death.. **2022**,
- 124 Nuclear Movement in Growing Arabidopsis Root Hairs Involves Both Actin Filaments and Microtubules.. **2022**, 1
- 123 Full Issue PDF.. **2022**, 35, 368-438
- 122 The Kelch-F-box protein SMALL AND GLOSSY LEAVES 1 (SAGL1) negatively influences salicylic acid biosynthesis in Arabidopsis thaliana by promoting the turn-over of transcription factor SYSTEMIC ACQUIRED RESISTANCE DEFICIENT 1 (SARD1).. **2022**, 1
- 121 Plant Growth-Promoting rhizobacterium Pseudomonas sp. CM11 Specifically Induces Lateral Roots.. **2022**, 0
- 120 Genome-wide Identification and Characterization of the Strawberry (Fragaria Vesca) FvAP2/ERF Gene Family in Abiotic Stress. 0
- 119 MAMP-elicited changes in amino acid transport activity contribute to restricting bacterial growth.. **2022**, 0
- 118 Regulatory interaction of BcWRKY33A and BcHSA4A promotes salt tolerance in non-heading Chinese cabbage [Brassica campestris (syn. Brassica rapa) ssp. chinensis]. 0
- 117 Neophaseic acid catabolism in 9'-hydroxylation pathway of abscisic acid in Arabidopsis thaliana.. **2022**, 100340 1
- 116 Conjugation. **2022**, 1619-1625
- 115 COR27/28 Regulate the Evening Transcriptional Activity of the RVE8-LNK1/2 Circadian Complex.
- 114 Mutations in the chloroplast inner envelope protein TIC100 impair and repair chloroplast protein import and impact retrograde signalling. 1
- 113 Overlapping functions of YDA and MAPKKK3/MAPKKK5 upstream of MPK3/MPK6 in plant immunity and growth/development. 0
- 112 Non-cell-autonomous HSC70.1 chaperone displays homeostatic feed-back regulation by binding its own mRNA.
- 111 Ovule initiation: the essential step controlling offspring number in Arabidopsis. 0
- 110 Lipidomic Analysis of Arabidopsis T-DNA Insertion Lines Leads to Identification and Characterization of C-Terminal Alterations in FATTY ACID DESATURASE 6. 0
- 109 The Indole-3-Acetamide-Induced Arabidopsis Transcription Factor MYB74 Decreases Plant Growth and Contributes to the Control of Osmotic Stress Responses. 13, 1

108	Experimentally heat-induced transposition increases drought tolerance in <i>Arabidopsis thaliana</i> .	1
107	Photoreceptors regulate gene expression of <i>Arabidopsis thaliana</i> grown with biophilic LED-sourced lighting systems. 2022 , 17, e0269868	1
106	Random Mutagenesis as a Promising Tool for Microalgal Strain Improvement towards Industrial Production. 2022 , 20, 440	1
105	Characterization of an autonomous pathway complex that promotes flowering in <i>Arabidopsis</i> .	1
104	A multiplexed, three-dimensional pooling and next-generation sequencing strategy for creating barcoded mutant arrays: construction of a <i>Schizosaccharomyces pombe</i> transposon insertion library.	
103	PILS proteins provide a homeostatic feedback on auxin signaling output. 2022 , 149,	0
102	<i>Medicago truncatula</i> resources to study legume biology and symbiotic nitrogen fixation. 2022 ,	1
101	High-throughput Identification of Novel Heat Tolerance Genes via Genome-wide Pooled Mutant Screens in the Model Green Alga <i>Chlamydomonas reinhardtii</i> .	0
100	Gibberellins promote polar auxin transport to regulate stem cell fate decisions in cambium.	
99	Construction and analysis of a <i>Noccaea caerulescens</i> TILLING population. 2022 , 22,	1
98	Systematic histone H4 replacement in <i>Arabidopsis thaliana</i> reveals a role for H4R17 in regulating flowering time.	0
97	Spatial consistency of cell growth direction during organ morphogenesis requires CELLULOSE-SYNTHASE INTERACTIVE1.	
96	A Role for Allantoate Amidohydrolase (AtAAH) in the Germination of <i>Arabidopsis Thaliana</i> Seeds.	
95	DEFECTIVELY ORGANIZED TRIBUTARIES 5 is not required for leaf venation patterning in <i>Arabidopsis thaliana</i> .	
94	AIG2A and AIG2B limit the activation of salicylic acid-regulated defenses by tryptophan-derived secondary metabolism in <i>Arabidopsis</i> .	0
93	The <i>Arabidopsis</i> INNER NO OUTER (INO) gene acts exclusively and quantitatively in regulation of ovule outer integument development.	
92	The calcium sensor CBL7 is required for <i>Serendipita indica</i> -induced growth stimulation in <i>Arabidopsis thaliana</i> , controlling defense against the endophyte and K ⁺ homeostasis in the symbiosis.	
91	Transcription-coupled and epigenome-encoded mechanisms direct H3K4 methylation. 2022 , 13,	0

90	Glutathione degradation activity of Eglutamyl peptidase 1 manifests its dual roles in primary and secondary sulfur metabolism in Arabidopsis.	1
89	The unconventional prefoldin RPB5 interactor mediates the gravitropic response by modulating cytoskeleton organization and auxin transport in Arabidopsis.	
88	Synergistic action of the Arabidopsis spliceosome components PRP39a and SmD1 in promoting post-transcriptional transgene silencing.	0
87	Assessment of functional relevance of genes associated with local temperature variables in Arabidopsis thaliana.	
86	Multiple pathways mediate chloroplast singlet oxygen stress signaling.	
85	Expression and Functional Analyses of the WIP Gene Family in Arabidopsis. 2022 , 11, 2010	
84	HPCA1 is required for systemic reactive oxygen species and calcium cell-to-cell signaling and plant acclimation to stress.	3
83	Reverse to Forward Genetic Screen Spots the C-Terminus of Plastidial Desaturase FAD6.	
82	The DC1 domain protein BINUCLEATE POLLEN is required for pollen development in Arabidopsis thaliana.	0
81	The regulation of Alternaria alternata resistance by LRR-RK4 through ERF109, defensin19 and phytoalexin scopoletin in Nicotiana attenuata. 2022 , 323, 111414	
80	Grape CIPK18 acts as a positive regulator of CBF cold signaling pathway by modulating ROS homeostasis. 2022 , 203, 105063	0
79	Genetic factors governing bacterial virulence and host plant susceptibility during Agrobacterium infection. 2022 ,	0
78	A single amino acid transporter controls the uptake of priming-inducing beta-amino acids and the associated tradeoff between induced resistance and plant growth.	0
77	ZAT10 plays dual roles in cadmium uptake and detoxification in Arabidopsis. 13,	0
76	The root-specific NtR12 promoter-based expression of RIP increased the resistance against bacterial wilt disease in tobacco.	0
75	DEFECTIVELY ORGANIZED TRIBUTARIES 5 is not required for leaf venation patterning in Arabidopsis thaliana.	0
74	Differential effects of root-level exposure to triazine xenobiotics on root development plasticity in Arabidopsis thaliana. 2022 , 44,	1
73	High-throughput detection of T-DNA insertion sites for multiple transgenes in complex genomes. 2022 , 23,	0

72	CGL160-mediated recruitment of the coupling factor CF1 is required for efficient thylakoid ATP synthase assembly, photosynthesis, and chloroplast development in Arabidopsis.	0
71	Multiple pathways mediate chloroplast singlet oxygen stress signaling.	0
70	P58IPK facilitates plant recovery from ER stress by enhancing protein synthesis.	0
69	Trait Improvement of Solanaceae Fruit Crops for Vertical Farming by Genome Editing.	0
68	Cytokinin modulates sulfur and glutathione metabolic network.	1
67	Loss of LEUCINE CARBOXYL METHYLTRANSFERASE 1 interferes with metal homeostasis in Arabidopsis and enhances susceptibility to environmental stresses. 2022 , 279, 153843	0
66	Agrobacterium. 2022 ,	0
65	Plethora of QTLs found in Arabidopsis thaliana reveals complexity of genetic variation for photosynthesis in dynamic light conditions.	0
64	A plastid nucleoside kinase is involved in inosine salvage and control of purine nucleotide biosynthesis.	1
63	m6A modification of U6 snRNA modulates usage of two major classes of pre-mRNA 5' splice site. 11,	1
62	The intracellular immune receptor like gene SNC1 is an enhancer of effector-triggered immunity in Arabidopsis.	0
61	Plant Experimental Resources. 2022 , 246-252	0
60	The root-knot nematode effector MiMSP32 targets host 12-oxophytodienoate reductase 2 (OPR2) to regulate plant susceptibility.	0
59	Discovery of endogenous nitroxyl as a new redox player in Arabidopsis thaliana.	0
58	Non-cell-autonomous HSC70 .1 chaperone displays homeostatic feed-back regulation by binding its own mRNA.	0
57	High-throughput identification of novel heat tolerance genes via genome-wide pooled mutant screens in the model green alga Chlamydomonas reinhardtii.	0
56	The vacuolar sugar transporter Early Response to Dehydration 6-Like4 regulates fructose signaling and plant growth.	0
55	An inosine triphosphate pyrophosphatase safeguards plant nucleic acids from aberrant purine nucleotides.	0

54	Characterization of γ -glutamyltransferase- and phytochelatin synthase-mediated catabolism of glutathione and glutathione S-conjugates in <i>Arabidopsis thaliana</i> . 2022 , 39, 381-389	1
53	UV Radiation Induces Specific Changes in the Carotenoid Profile of <i>Arabidopsis thaliana</i> . 2022 , 12, 1879	1
52	Intergenic splicing-stimulated transcriptional readthrough is suppressed by nonsense-mediated mRNA decay in <i>Arabidopsis</i> . 2022 , 5,	0
51	Abscisic Acid Increases Hydrogen Peroxide in Mitochondria to Facilitate Stomatal Closure.	0
50	The LHT1 amino acid transporter limits <i>Pseudomonas simiae</i> growth in the rhizosphere, an essential function for bacteria-mediated plant growth promotion.	0
49	Analysis of BRCT5 domain-containing proteins reveals a new component of DNA damage repair in <i>Arabidopsis</i> . 13,	0
48	Transcriptomic reprogramming in a susceptible <i>Phaseolus vulgaris</i> L. variety during <i>Pseudomonas syringae</i> attack: The key role of homogalacturonan methylation.	0
47	Reduced Expression of PRX2/ATPRX1, PRX8, PRX35, and PRX73 Affects Cell Elongation, Vegetative Growth, and Vasculature Structures in <i>Arabidopsis thaliana</i> . 2022 , 11, 3353	0
46	The <i>Arabidopsis</i> LHT1 Amino Acid Transporter Contributes to <i>Pseudomonas simiae</i> -Mediated Plant Growth Promotion by Modulating Bacterial Metabolism in the Rhizosphere. 2023 , 12, 371	0
45	Identification of multiple novel genetic mechanisms that regulate chilling tolerance in <i>Arabidopsis</i> . 13,	0
44	<i>Arabidopsis</i> TRB proteins function in H3K4me3 demethylation by recruiting JMJ14.	0
43	Involvement of the Membrane Nanodomain Protein, AtFlot1, in Vesicular Transport of Plasma Membrane H ⁺ -ATPase in <i>Arabidopsis thaliana</i> under Salt Stress. 2023 , 24, 1251	0
42	Light induction of somatic embryogenesis in <i>Arabidopsis</i> is regulated by PHYTOCHROME E. 2023 , 195, 163-169	0
41	Peroxisomal protein phosphatase PP2A-B β interacts with and piggybacks SINA-like 10 E3 ligase into peroxisomes. 2023 , 644, 34-39	0
40	The intracellular auxin homeostasis regulators PIN5 and PIN8 have a divergent membrane topology in <i>Arabidopsis thaliana</i> root cells.	0
39	A simple and cost-effective method for studying anoxia tolerance in plants.	0
38	BBX24 interacts with JAZ3 to promote growth by reducing DELLA activity in shade avoidance..	0
37	The non-coding RNA SValka locus produces a cis- natural antisense transcript that negatively regulates the expression of CBF1 and biomass production at normal temperatures. 2023 , 100551	0

- 36 FKF1 Interacts with CHUP1 and Regulates Chloroplast Movement in Arabidopsis. **2023**, 12, 542 ○
- 35 GoSAMTs are required for pectin methyl-esterification and mucilage release in seed coat epidermal cells. 14, ○
- 34 Generation, storage, and utilizations of mutant libraries. **2023**, 403-420 ○
- 33 UGT76F1 glycosylates an isomer of the C7 -necic acid component of pyrrolizidine alkaloids in Arabidopsis thaliana. ○
- 32 A molecular framework underlying low-nitrogen-induced early leaf senescence in Arabidopsis thaliana. **2023**, 16, 756-774 ○
- 31 Deficiency of the Arabidopsis mismatch repair MSH6 attenuates Pseudomonas syringae invasion. **2023**, 332, 111713 ○
- 30 pin2 mutant agravitropic root phenotype is conditional and nutrient-sensitive. **2023**, 329, 111606 ○
- 29 MKKK20 works as an upstream triple-kinase of MKK3-MPK6-MYC2 module in Arabidopsis seedling development. **2023**, 26, 106049 ○
- 28 Comprehensive analysis of cucumber RAV Family genes and functional characterization of CsRAV1 in salt and ABA tolerance in cucumber. 14, ○
- 27 PAT1-type GRAS-domain proteins control regeneration by activating DOF3.4 to drive cell proliferation in Arabidopsis roots. **2023**, 35, 1513-1531 ○
- 26 The atrzf1 Mutation of the Novel RING-Type E3 Ubiquitin Ligase Increases Proline Contents and Enhances Drought Tolerance in Arabidopsis. **2021**, 59, 15-28 ○
- 25 Genome-wide analysis of mutations induced by carbon ion beam irradiation in cotton. 14, ○
- 24 Biological insights from multi-omics analysis strategies: Complex pleiotropic effects associated with autophagy. 14, ○
- 23 m6A mRNA modification promotes chilling tolerance and modulates gene translation efficiency in Arabidopsis. ○
- 22 The Arabidopsis INNER NO OUTER (INO) gene acts exclusively and quantitatively in regulation of ovule outer integument development. **2023**, 7, ○
- 21 Functional genomics of Chlamydomonas reinhardtii. **2023**, 65-84 ○
- 20 Deregulation of β -carotene desaturase in Arabidopsis and tomato exposes a unique carotenoid-derived redundant regulation of floral meristem identity and function. ○
- 19 Canonical and ETT-mediated auxin signalling pathways synergistically promote gynoecium development in Arabidopsis thaliana. ○

- 18 Insights into the function of the chloroplastic ribosome-associated GTPase HflX in *Arabidopsis thaliana*. ○
- 17 A gene silencing screen uncovers diverse tools for targeted gene repression in *Arabidopsis*. **2023**, 9, 460-472 ○
- 16 Induction of T-DNA amplification by retrotransposon-derived sequences. ○
- 15 *Arabidopsis thaliana* BBX14 is a target of GLK1 and involved in high-light acclimation, photomorphogenesis and GUN-type retrograde signaling. ○
- 14 Roles of microbiota in autoimmunity in *Arabidopsis*. ○
- 13 Genetic manipulation of microRNAs: approaches and limitations. ○
- 12 *Arabidopsis* cell suspension culture and RNA sequencing reveal regulatory networks underlying plant programmed cell death. ○
- 11 Sensory circuitry controls cytosolic calcium-mediated phytochrome B phototransduction. **2023**, 186, 1230-1243. e14 ○
- 10 Multi-Knock[®] multi-targeted genome-scale CRISPR toolbox to overcome functional redundancy in plants. **2023**, 9, 572-587 ○
- 9 *Arabidopsis* TRB proteins function in H3K4me3 demethylation by recruiting JMJ14. **2023**, 14, ○
- 8 Analysis of *Arabidopsis venosa4-0* supports the role of VENOSA4 in dNTP homeostasis. ○
- 7 Further insight into the involvement of PII1 in starch granule initiation in *Arabidopsis* leaf chloroplasts. ○
- 6 Crop Improvement Through Induced Genetic Diversity and Mutation Breeding: Challenges and Opportunities. **2023**, 293-300 ○
- 5 Gibberellins promote polar auxin transport to regulate stem cell fate decisions in cambium. **2023**, 9, 631-644 ○
- 4 COLD REGULATED GENE 27 and 28 antagonize the transcriptional activity of the RVE8/LNK1/LNK2 circadian complex. ○
- 3 Clathrin Light Chains are essential in negative regulation of cell death and immunity in *Arabidopsis* through interacting with autophagy pathway. ○
- 2 Auxin-dependent regulation of cell division rates governs root thermomorphogenesis. ○
- 1 Influence of a phyA Mutation on Polyamine Metabolism in *Arabidopsis* Depends on Light Spectral Conditions. **2023**, 12, 1689 ○

