

# CITATION REPORT

List of articles citing

**Cross-calibration of eight-polar bioelectrical impedance analysis versus dual-energy X-ray absorptiometry for the assessment of total and appendicular body composition in healthy subjects aged 21-82 years**

**DOI: 10.1080/0301446031000095211**

**Annals of Human Biology, 2003, 30, 380-91.**

**Source:** <https://exaly.com/paper-pdf/35646727/citation-report.pdf>

**Version:** 2024-04-23

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| #   | Paper   | IF | Citations |
|-----|---|----|-----------|
| 257 | Body water distribution in severe obesity and its assessment from eight-polar bioelectrical impedance analysis. <b>2005</b> , 59, 155-60  |    | 110       |
| 256 | Accuracy of eight-polar bioelectrical impedance analysis for the assessment of total and appendicular body composition in peritoneal dialysis patients. <b>2005</b> , 59, 932-7                           |    | 50        |
| 255 | Interleukin-1 gene cluster polymorphisms are associated with nutritional status and inflammation in patients with end-stage renal disease. <b>2005</b> , 23, 384-93                                       |    | 21        |
| 254 | Role of interleukin-1beta in the development of malnutrition in chronic renal failure patients. <b>2005</b> , 23, 275-81  |    | 8         |
| 253 | A white blood cell count in the normal concentration range is independently related to cardiorespiratory fitness in apparently healthy Korean men. <b>2005</b> , 54, 1448-52                              |    | 13        |
| 252 | Within-day variability of whole-body and segmental bioelectrical impedance in a standing position. <b>2006</b> , 60, 938-41   |    | 18        |
| 251 | Insulin resistance and muscle wasting in non-diabetic end-stage renal disease patients. <b>2007</b> , 22, 2554-62   |    | 31        |
| 250 | Effect of protein and energy intakes on body composition in non-diabetic maintenance-hemodialysis patients. <b>2007</b> , 53, 410-8   |    | 11        |
| 249 | Quantification of Spinning bike performance during a standard 50-minute class. <b>2007</b> , 25, 421-9  |    | 29        |
| 248 | Body fat distribution and insulin resistance: beyond obesity in nonalcoholic fatty liver disease among overweight men. <b>2007</b> , 26, 321-6  |    | 47        |
| 247 | Birth size, adult body composition and muscle strength in later life. <b>2007</b> , 31, 1392-9  |    | 160       |
| 246 | Relative contribution of body composition to bone mineral density at different sites in men and women of South Korea. <b>2007</b> , 25, 165-71  |    | 76        |
| 245 | Evaluation of air-displacement plethysmography and bioelectrical impedance analysis vs dual-energy X-ray absorptiometry for the assessment of fat-free mass in elderly subjects. <b>2008</b> , 62, 1282-6 |    | 11        |
| 244 | The mediating role of C-reactive protein and handgrip strength between obesity and walking limitation. <b>2008</b> , 56, 462-9  |    | 85        |
| 243 | Different pattern of fluid loss from the lower extremities in normohydrated and overhydrated stage 5 chronic-kidney-disease patients after haemodialysis. <b>2008</b> , 13, 109-15                        |    | 3         |
| 242 | Validity and reliability of body composition analysers in children and adults. <b>2008</b> , 100, 859-65  |    | 117       |
| 241 | Effects of a short-term resistance program using elastic bands versus weight machines for sedentary middle-aged women. <b>2008</b> , 22, 1441-8   |    | 112       |

|     |  |     |
|-----|--|-----|
| 240 | Ability of new octapolar bioimpedance spectroscopy analyzers to predict 4-component-model percentage body fat in Hispanic, black, and white adults. <b>2008</b> , 87, 332-8  | 127 |
| 239 | Body composition and fitness during strength and/or endurance training in older men. <b>2008</b> , 40, 950-8   | 68  |
| 238 | Agreements between indirect calorimetry and prediction equations of resting energy expenditure in end-stage renal disease patients on continuous ambulatory peritoneal dialysis. <b>2008</b> , 49, 255-64                                      | 12  |
| 237 | Body mass index during childhood and adult body composition in men and women aged 56-70 y. <b>2008</b> , 87, 1769-75   | 85  |
| 236 | Japanese study to organize proper lifestyle modifications for metabolic syndrome (J-STOP-Mets): design and method. <b>2008</b> , 4, 415-20   | 6   |
| 235 | Body composition in psychotic disorders: a general population survey. <b>2009</b> , 39, 801-10   | 51  |
| 234 | Body composition at high altitude: a randomized placebo-controlled trial of dietary carbohydrate supplementation. <b>2009</b> , 90, 1193-202   | 33  |
| 233 | Validity of segmental multiple-frequency bioelectrical impedance analysis to estimate body composition of adults across a range of body mass indexes. <b>2009</b> , 25, 25-32  | 144 |
| 232 | Study design and methods of the Ansan Geriatric Study (AGE study). <b>2009</b> , 9, 10   | 34  |
| 231 | Effects of aquatic resistance training on health and fitness in postmenopausal women. <b>2009</b> , 106, 113-22  | 90  |
| 230 | Cross-validation of bioelectrical impedance analysis for the assessment of body composition in a representative sample of 6- to 13-year-old children. <b>2009</b> , 63, 619-26   | 56  |
| 229 | New method of predicting dry weight using bioelectrical impedance analysis in haemodialysis patients. <b>2009</b> , 14, 705-11   | 3   |
| 228 | Cross-calibration of multi-frequency bioelectrical impedance analysis with eight-point tactile electrodes and dual-energy X-ray absorptiometry for assessment of body composition in healthy children aged 6-18 years. <b>2009</b> , 51, 263-8 | 88  |
| 227 | Adiposity parameters and cognitive function in the elderly: application of "Jolly Fat" hypothesis to cognition. <b>2009</b> , 49, e133-e138  | 39  |
| 226 | Oxygen uptake for cycling in relation to body composition: a pilot study. <b>2009</b> , 25, 544-51   | 2   |
| 225 | Physical activity and body composition changes during military service. <b>2009</b> , 41, 1735-42  | 23  |
| 224 | Negative association between metabolic syndrome and bone mineral density in Koreans, especially in men. <b>2010</b> , 86, 350-8  | 71  |
| 223 | Efficacy of Biodanza for treating women with fibromyalgia. <b>2010</b> , 16, 1191-200  | 24  |

|     |  |     |
|-----|--|-----|
| 222 | Temporal parameters of the foot roll-over during walking: influence of obesity and sarcopenic obesity on postmenopausal women. <b>2010</b> , 67, 178-85  | 19  |
| 221 | Analysis of the body composition of Spanish women with fibromyalgia. <b>2011</b> , 7, 7-12   |     |
| 220 | [Analysis of the body composition of Spanish women with fibromyalgia]. <b>2011</b> , 7, 7-12   | 9   |
| 219 | Assessment of body composition using bioelectrical impedance analysis in Prader-Willi syndrome. <b>2011</b> , 110, 719-23  | 6   |
| 218 | Pain and functional capacity in female fibromyalgia patients. <b>2011</b> , 12, 1667-75  | 51  |
| 217 | Accuracy of direct segmental multi-frequency bioimpedance analysis in the assessment of total body and segmental body composition in middle-aged adult population. <b>2011</b> , 30, 610-5                               | 339 |
| 216 | Reliability and accuracy of segmental bioelectrical impedance analysis for assessing muscle and fat mass in older Europeans: a comparison with dual-energy X-ray absorptiometry. <b>2011</b> , 111, 1879-87              | 37  |
| 215 | Does a 3-month multidisciplinary intervention improve pain, body composition and physical fitness in women with fibromyalgia?. <b>2011</b> , 45, 1189-95   | 44  |
| 214 | Research letter. Body composition in subtypes of depression--a population-based survey. <b>2011</b> , 41, 1113-7   | 3   |
| 213 | The effects of androgen deprivation therapy on lipid metabolism and body composition in Japanese patients with prostate cancer. <b>2011</b> , 41, 577-81   | 23  |
| 212 | Are there gender differences in quality of life and symptomatology between fibromyalgia patients?. <b>2012</b> , 6, 314-9  | 16  |
| 211 | Effectiveness of a tai-chi training and detraining on functional capacity, symptomatology and psychological outcomes in women with fibromyalgia. <b>2012</b> , 2012, 614196  | 26  |
| 210 | Body composition analysis in critically ill survivors: a comparison of bioelectrical impedance spectroscopy devices. <b>2012</b> , 36, 306-15  | 18  |
| 209 | Racial differences in the association between adiposity measures and the risk of hepatitis C-related liver disease. <b>2012</b> , 46, 779-88   | 2   |
| 208 | Effects of aquatic and dry land resistance training devices on body composition and physical capacity in postmenopausal women. <b>2012</b> , 32, 185-95  | 40  |
| 207 | North Korean refugee health in South Korea (NORNS) study: study design and methods. <b>2012</b> , 12, 172  | 26  |
| 206 | Obesity in young-adult Nigerians: variations in prevalence determined by anthropometry and bioelectrical impedance analysis, and the development of % body fat prediction equations. <b>2012</b> , 5, 22                 | 14  |
| 205 | New Application of Bioelectrical Impedance Analysis by the Back Propagation Artificial Neural Network Mathematically Predictive Model of Tissue Composition in the Lower Limbs of Elderly People. <b>2012</b> , 6, 20-26 | 7   |

|     |   |     |    |
|-----|---|-----|----|
| 204 | Bioelectrical Impedance Analysis in a Mathematical Model for Estimating Fat-free Mass in Multiple Segments in Elderly Taiwanese Males. <b>2012</b> , 6, 273-277   |     | 8  |
| 203 | Urban-rural differences in physical performance and health status among older Japanese community-dwelling women. <b>2012</b> , 3, 127-131   |     | 6  |
| 202 | Birth-weight and resting metabolic rate in adulthood - sex-specific differences. <b>2012</b> , 44, 296-303  |     | 10 |
| 201 | Development and validation of a predictive equation for lean body mass in children and adolescents. <i>Annals of Human Biology</i> , <b>2012</b> , 39, 171-82   | 1.7 | 51 |
| 200 | The utility of visceral fat level by bioelectrical impedance analysis in the screening of metabolic syndrome - the results of the Kyushu and Okinawa Population Study (KOPS). <b>2012</b> , 19, 462-70                  |     | 42 |
| 199 | Higher serum testosterone is associated with increased risk of advanced hepatitis C-related liver disease in males. <b>2012</b> , 55, 759-68  |     | 54 |
| 198 | A comparison of dual energy X-ray absorptiometry and bioelectrical impedance analysis to measure total and segmental body composition in healthy young adults. <b>2012</b> , 112, 589-95                                |     | 73 |
| 197 | Body composition changes by DXA, BIA and skinfolds during exercise training in women. <b>2013</b> , 113, 2331-41  |     | 23 |
| 196 | Physical inactivity and insufficient dietary intake are associated with the frequency of sarcopenia in patients with compensated viral liver cirrhosis. <b>2013</b> , 43, 1264-75                                       |     | 77 |
| 195 | Assessing Body Composition. <b>2013</b> , 88-101  |     |    |
| 194 | Water versus land-based multimodal exercise program effects on body composition in breast cancer survivors: a controlled clinical trial. <b>2013</b> , 21, 521-30   |     | 34 |
| 193 | Fibromyalgia's key symptoms in normal-weight, overweight, and obese female patients. <b>2013</b> , 14, 268-276  |     | 21 |
| 192 | Effect of fasting in Ramadan on body composition and nutritional intake: a prospective study. <b>2013</b> , 26 Suppl 1, 97-104  |     | 75 |
| 191 | A warm water pool-based exercise program decreases immediate pain in female fibromyalgia patients: uncontrolled clinical trial. <b>2013</b> , 34, 600-5   |     | 11 |
| 190 | Cut-off value of body fat in association with metabolic syndrome in Thai peri- and postmenopausal women. <b>2013</b> , 16, 393-7  |     | 14 |
| 189 | Direct segmental multi-frequency bioelectrical impedance analysis is useful to evaluate sarcopenia. <b>2013</b> , 13, 2506-7  |     | 24 |
| 188 | Retrospective analysis of safety and efficacy of insulin-to-liraglutide switch in Japanese type 2 diabetes: A caution against inappropriate use in patients with reduced $\beta$ cell function. <b>2013</b> , 4, 585-94 |     | 23 |
| 187 | Relative skeletal muscle mass is associated with development of metabolic syndrome. <b>2013</b> , 37, 458-64  |     | 59 |

|     |   |       |
|-----|---|-------|
| 186 | A Two-Stage Regression Using Bioimpedance and Temperature for Hydration Assessment During Sports. <b>2014</b> ,   | 2     |
| 185 | Arterial stiffness is associated with low skeletal muscle mass in Japanese community-dwelling older adults. <b>2014</b> , 14 Suppl 1, 109-14  | 56    |
| 184 | Body composition measurements using bioimpedance analysis in peritoneal dialysis patients are affected by the presence of dialysate. <b>2014</b> , 19, 727-31   | 6     |
| 183 | Fat mass by bioelectrical impedance analysis is not associated with increased risk of Barrett esophagus. <b>2014</b> , 48, 218-23   | 21    |
| 182 | Vitamin D insufficiency is associated with lower physical function in patients with heart failure and diabetes. <b>2014</b> , 2014, 320930  | 9     |
| 181 | The importance of visceral fat mass in obese pregnant women and relation with pregnancy outcomes. <b>2014</b> , 7, 22-5   | 6     |
| 180 | Body composition in 18- to 88-year-old adults--comparison of multifrequency bioimpedance and dual-energy X-ray absorptiometry. <b>2014</b> , 22, 101-9  | 64    |
| 179 | Metabolically-healthy obesity and coronary artery calcification. <b>2014</b> , 63, 2679-86  | 178   |
| 178 | Concentrations of chromium, selenium, and copper in the hair of viscerally obese adults are associated with insulin resistance. <b>2014</b> , 158, 152-7  | 29    |
| 177 | Top 10 research questions related to body composition. <b>2014</b> , 85, 38-48  | 7     |
| 176 | Accuracy of quantitative magnetic resonance and eight-electrode bioelectrical impedance analysis in normal weight and obese women. <b>2014</b> , 33, 471-7  | 21    |
| 175 | Cardiorespiratory fitness and body composition in postmenopausal women. <b>2014</b> , 43, 139-48  | 10    |
| 174 | Measurement of Skeletal Muscle mass of Japanese Men and Women Aged 18-84 by Bioelectrical Impedance Analysis Focusing on the Difference in Measured Values Produced by Different Equipment. <i>Rigakuryoho Kagaku</i> , <b>2015</b> , 30, 265-271 | 0.1 0 |
| 173 | Sarcopenic obesity and physical performance in middle aged women: a cross-sectional study in Northeast Brazil. <b>2016</b> , 16, 43   | 41    |
| 172 | Differences in sedentary time and physical activity between female patients with fibromyalgia and healthy controls: the al-Bdalus project. <b>2015</b> , 67, 3047-57  | 38    |
| 171 | Comparisons of Predictive Equations for Resting Energy Expenditure in Patients with Cerebral Infarct during Acute Care. <b>2015</b> , 24, 1879-85   | 16    |
| 170 | Association of visceral fat area with chronic kidney disease and metabolic syndrome risk in the general population: analysis using multi-frequency bioimpedance. <b>2015</b> , 40, 223-30   | 29    |
| 169 | Identification of peripheral arterial disease in diabetic patients and its association with quality of life, physical activity and body composition. <b>2015</b> , 14, 46-54  | 6     |

|     |   |     |     |
|-----|---|-----|-----|
| 168 | Inflammation is associated to volume status in peritoneal dialysis patients. <b>2015</b> , 37, 935-40   |     | 3   |
| 167 | Genetic Variants in Interleukin-28B Are Associated with Diabetes and Diabetes-Related Complications in Patients with Chronic Hepatitis C Virus Infection. <b>2015</b> , 60, 2030-7  |     | 3   |
| 166 | Vitamin D status is a determinant of skeletal muscle mass in obesity according to body fat percentage. <b>2015</b> , 31, 801-6  |     | 22  |
| 165 | Assessment of fat-free mass from bioelectrical impedance analysis in obese women with Prader-Willi syndrome. <i>Annals of Human Biology</i> , <b>2015</b> , 42, 538-42  | 1.7 | 11  |
| 164 | Motivational interviewing with parents for obesity: an RCT. <b>2015</b> , 135, e644-52  |     | 32  |
| 163 | Association of dual-task walking performance and leg muscle quality in healthy children. <b>2015</b> , 15, 2  |     | 20  |
| 162 | Agreement Between Omron 306 and Biospace InBody 720 Bioelectrical Impedance Analyzers (BIA) in Children and Adolescents. <b>2015</b> , 86 Suppl 1, S58-65   |     | 10  |
| 161 | Age at menarche and non-alcoholic fatty liver disease. <b>2015</b> , 62, 1164-70  |     | 37  |
| 160 | The influence of body mass index on survival in breast cancer patients. <b>2015</b> , 15, e117-23   |     | 9   |
| 159 | Relationship of sitting time and physical activity with non-alcoholic fatty liver disease. <b>2015</b> , 63, 1229-37  |     | 116 |
| 158 | Triiodothyronine Levels Are Independently Associated with Metabolic Syndrome in Euthyroid Middle-Aged Subjects. <b>2016</b> , 31, 311-9   |     | 21  |
| 157 | Insulin resistance increases loss of antibody to hepatitis B surface antigen in nondiabetic healthy adults. <b>2016</b> , 23, 889-896   |     | 2   |
| 156 | The association of total and central body fat with pain, fatigue and the impact of fibromyalgia in women; role of physical fitness. <b>2016</b> , 20, 811-21  |     | 12  |
| 155 | The discordance between subjectively and objectively measured physical function in women with fibromyalgia: association with catastrophizing and self-efficacy cognitions. The al-Badalus project. <i>Disability and Rehabilitation</i> , <b>2018</b> , 40, 329-337 | 2.4 | 26  |
| 154 | Metabolically Healthy Obesity and the Development of Nonalcoholic Fatty Liver Disease. <b>2016</b> , 111, 1133-40   |     | 107 |
| 153 | Age at Menarche and Gallstone Disease in Middle-Aged Women. <b>2016</b> , 23, 1304-13   |     | 2   |
| 152 | Relationship Between Low Relative Muscle Mass and Coronary Artery Calcification in Healthy Adults. <b>2016</b> , 36, 1016-21  |     | 52  |
| 151 | Metabolically healthy obesity is associated with an increased risk of diabetes independently of nonalcoholic fatty liver disease. <b>2016</b> , 24, 1996-2003   |     | 19  |

|     |  |     |     |
|-----|--|-----|-----|
| 150 | Metabolically Healthy Obesity and Development of Chronic Kidney Disease: A Cohort Study. <b>2016</b> , 164, 305-12   |     | 161 |
| 149 | The amino acid-rich elemental diet Elental <sup>®</sup> preserves lean body mass during chemo- or chemoradiotherapy for esophageal cancer. <b>2016</b> , 36, 1093-100  |     | 20  |
| 148 | Association of Physical Fitness with Depression in Women with Fibromyalgia. <b>2016</b> , 17, 1542-52  |     | 17  |
| 147 | Cut-off points to identify sarcopenia according to European Working Group on Sarcopenia in Older People (EWGSOP) definition. <b>2016</b> , 35, 1557-1563   |     | 191 |
| 146 | Association of physical fitness and fatness with cognitive function in women with fibromyalgia. <b>2016</b> , 34, 1731-9   |     | 6   |
| 145 | A Temperature-Based Bioimpedance Correction for Water Loss Estimation During Sports. <b>2016</b> , 20, 1477-1484   |     | 4   |
| 144 | Association of triiodothyronine levels with future development of metabolic syndrome in euthyroid middle-aged subjects: a 6-year retrospective longitudinal study. <i>European Journal of Endocrinology</i> , <b>2017</b> , 176, 443-452 | 6.5 | 6   |
| 143 | Low muscle mass and risk of type 2 diabetes in middle-aged and older adults: findings from the KoGES. <b>2017</b> , 60, 865-872  |     | 81  |
| 142 | Development of Nephrolithiasis in Asymptomatic Hyperuricemia: A Cohort Study. <b>2017</b> , 70, 173-181  |     | 33  |
| 141 | Sarcopenia is associated with an increased risk of advanced colorectal neoplasia. <b>2017</b> , 32, 557-565  |     | 18  |
| 140 | Association Between Low Relative Muscle Mass and the Risk of Colorectal Neoplasms. <b>2017</b> , 51, e83-e89   |     | 7   |
| 139 | Obesity and Risk of Nonalcoholic Fatty Liver Disease: A Comparison of Bioelectrical Impedance Analysis and Conventionally-Derived Anthropometric Measures. <i>Clinical Gastroenterology and Hepatology</i> , <b>2017</b> , 15, 1965-1967 | 6.9 | 7   |
| 138 | Time-motion analysis and patterns of salivary cortisol during different judo championship phases. <b>2017</b> , 13, 419-426  |     | 1   |
| 137 | Serum uric acid is positively associated with pulmonary function in Korean health screening examinees. <b>2017</b> , 27, 1057-1065   |     | 14  |
| 136 | Contribution of branched-chain amino acids to purine nucleotide cycle: a pilot study. <b>2017</b> , 71, 587-593  |     | 4   |
| 135 | Independent and joint associations of physical activity and fitness with fibromyalgia symptoms and severity: The al-Bidal project. <b>2017</b> , 35, 1565-1574   |     | 10  |
| 134 | Fat-free mass index predicts survival in patients with idiopathic pulmonary fibrosis. <b>2017</b> , 22, 480-485  |     | 21  |
| 133 | The Associations between Circulating Bile Acids and the Muscle Volume in Patients with Non-alcoholic Fatty Liver Disease (NAFLD). <b>2017</b> , 56, 755-762  |     | 17  |

|     |   |      |
|-----|---|------|
| 132 | Relative muscle mass and the risk of incident type 2 diabetes: A cohort study. <i>PLoS ONE</i> , <b>2017</b> , 12, e0188650   | 41   |
| 131 | A Randomized Controlled Trial of Nurse Coaching vs. Herbal Supplementation for Weight Reduction in Soldiers. <i>Military Medicine</i> , <b>2017</b> , 182, 274-280                                | 13 5 |
| 130 | DIFERENCIAS ESPACIO-TEMPORALES DE LA LOCOMOCIÓN EN ADULTOS VARONES CON NORMOPESO Y SOBREPESO. <b>2017</b> , 23, 8-11  |      |
| 129 | Heterogeneity of gestational diabetes (GDM) and long-term risk of diabetes and metabolic syndrome: findings from the RADIEL study follow-up. <b>2018</b> , 55, 493-501                            | 23   |
| 128 | Positive Effects of a Short-Term Intense Elastic Resistance Training Program on Body Composition and Physical Functioning in Overweight Older Women. <b>2018</b> , 20, 321-334                    | 16   |
| 127 | Low relative muscle mass and left ventricular diastolic dysfunction in middle-aged adults. <b>2018</b> , 255, 118-123   | 12   |
| 126 | Sitting time, physical activity and the risk of lower urinary tract symptoms: a cohort study. <b>2018</b> , 122, 293-299  | 8    |
| 125 | Does obesity attenuate the beneficial cardiovascular effects of cardiorespiratory fitness?. <b>2018</b> , 272, 21-26  | 3    |
| 124 | Relationships among peak expiratory flow rate, body composition, physical function, and sarcopenia in community-dwelling older adults. <b>2018</b> , 30, 331-340                                  | 20   |
| 123 | Non-Discriminant Relationships between Leg Muscle Strength, Mass and Gait Performance in Healthy Young and Old Adults. <b>2018</b> , 64, 11-18  | 13   |
| 122 | The Obesity Factor: How Cardiorespiratory Fitness is Estimated More Accurately in People with Obesity. <b>2018</b> , 26, 291-298  | 5    |
| 121 | Pre-pregnancy overweight or obesity and gestational diabetes as predictors of body composition in offspring twenty years later: evidence from two birth cohort studies. <b>2018</b> , 42, 872-879 | 34   |
| 120 | Relationship between skeletal muscle mass and lung function in Korean adults without clinically apparent lung disease. <b>2018</b> , 97, e12281   | 26   |
| 119 | Contrast in the circadian behaviors of an electrodermal activity and bioimpedance spectroscopy. <b>2018</b> , 35, 1413-1422   | 3    |
| 118 | Dapagliflozin Reduces Fat Mass without Affecting Muscle Mass in Type 2 Diabetes. <b>2018</b> , 25, 467-476  | 40   |
| 117 | Association between appendicular skeletal muscle mass and depressive symptoms: Review of the cardiovascular and metabolic diseases etiology research center cohort. <b>2018</b> , 238, 8-15       | 11   |
| 116 | Body composition as a predictor of physical performance in older age: A ten-year follow-up of the Helsinki Birth Cohort Study. <b>2018</b> , 77, 163-168  | 10   |
| 115 | Effect of Tofogliflozin on Body Composition and Glycemic Control in Japanese Subjects with Type 2 Diabetes Mellitus. <b>2018</b> , 2018, 6470137  | 18   |

|     |   |        |
|-----|---|--------|
| 114 | Nutritional status of outpatients with chronic stable heart failure based on serum amino acid concentration. <b>2018</b> , 72, 458-465  | 15     |
| 113 | Low muscle quality in Japanese type 2 diabetic patients with visceral fat accumulation. <b>2018</b> , 17, 112   | 27     |
| 112 | Reduced leptin level is independent of fat mass changes and hunger scores from high-intensity intermittent plus strength training. <b>2018</b> , 58, 1045-1051  | 3      |
| 111 | Metformin in Pregnancy: Mechanisms and Clinical Applications. <b>2018</b> , 19,   | 33     |
| 110 | The SGLT2 Inhibitor Dapagliflozin Significantly Improves the Peripheral Microvascular Endothelial Function in Patients with Uncontrolled Type 2 Diabetes Mellitus. <b>2018</b> , 57, 2147-2156              | 41     |
| 109 | Visceral fat mass as a novel risk factor for predicting gestational diabetes in obese pregnant women. <b>2018</b> , 11, 121-125   | 9      |
| 108 | Association of objectively measured physical activity and sedentary time with health-related quality of life in women with fibromyalgia: The al-Bdalu project. <b>2019</b> , 8, 258-266                     | 9      |
| 107 | Substituting Sedentary Time With Physical Activity in Fibromyalgia and the Association With Quality of Life and Impact of the Disease: The al-Bdalu Project. <b>2019</b> , 71, 281-289                      | 9      |
| 106 | Decline in lung function is associated with elevated lipoprotein (a) in individuals without clinically apparent disease: A cross-sectional study. <b>2019</b> , 24, 68-75                                   | 2      |
| 105 | Associations between Skeletal Muscle and Myocardium in Aging: A Syndrome of "Cardio-Sarcopenia"?. <b>2019</b> , 67, 2568-2573   | 18     |
| 104 | Clinical features, treatment and outcomes of colon and rectum mucosa-associated lymphoid tissue (MALT) lymphoma: Literature reviews published in English between 1993 and 2017. <b>2019</b> , 30, v440-v441 |        |
| 103 | Consumption of dietary supplements to support weight reduction in adults according to sociodemographic background, body mass index, waist-hip ratio, body fat and physical activity. <b>2019</b> , 38, 31   | 7      |
| 102 | Comparative assessment of skeletal muscle mass using computerized tomography and bioelectrical impedance analysis in critically ill patients. <b>2019</b> , 38, 2747-2755                                   | 22     |
| 101 | A Study of Using Tetrapolar and Eight-Polar Electrode System to Detect Fat by Bioelectrical Impedance Analysis (BIA) Technique. <b>2019</b> ,   | 0      |
| 100 | Lean body mass and risk of type 2 diabetes - a Danish cohort study. <b>2019</b> , 18, 445-451   | 8      |
| 99  | Viscus fat area contributes to the Framingham 10-year general cardiovascular disease risk in patients with type 2 diabetes mellitus. <b>2019</b> , 220, 69-75   | 1      |
| 98  | Decreased lung function is associated with risk of developing non-alcoholic fatty liver disease: A longitudinal cohort study. <i>PLoS ONE</i> , <b>2019</b> , 14, e0208736                                  | 3.7 17 |
| 97  | Assessment of fat-free mass from bioelectrical impedance analysis in men and women with Prader-Willi syndrome: cross-sectional study. <b>2019</b> , 70, 645-649   | 5      |

|    |  |     |    |
|----|--|-----|----|
| 96 | Association of Sarcopenia With Performance on Multiple Cognitive Domains: Results From the ELSA-Brasil Study. <b>2019</b> , 74, 1805-1811  |     | 11 |
| 95 | Dry Electrode-Based Body Fat Estimation System with Anthropometric Data for Use in a Wearable Device. <b>2019</b> , 19,  |     | 2  |
| 94 | SPA-Based Methods for the Quantitative Estimation of the Soil Salt Content in Saline-Alkali Land from Field Spectroscopy Data: A Case Study from the Yellow River Irrigation Regions. <b>2019</b> , 11, 967                          |     | 16 |
| 93 | A Prospective Cohort Study of Vitamin D Supplementation in AD Soldiers: Preliminary Findings. <i>Military Medicine</i> , <b>2019</b> , 184, 498-505  | 1.3 | 3  |
| 92 | Serum concentration of thyroid hormones in abnormal and euthyroid ranges and breast cancer risk: A cohort study. <b>2019</b> , 145, 3257-3266  |     | 22 |
| 91 | Association of Patterns of Moderate-to-Vigorous Physical Activity Bouts With Pain, Physical Fatigue, and Disease Severity in Women With Fibromyalgia: the al-Bidalus Project. <b>2019</b> , 100, 1234-1242.e1                        |     | 12 |
| 90 | Predictors of Discordance in the Assessment of Skeletal Muscle Mass between Computed Tomography and Bioimpedance Analysis. <b>2019</b> , 8,  |     | 4  |
| 89 | Association of mineral content outside of bone with coronary artery calcium and 1-year cardiovascular prognosis in maintenance hemodialysis patients. <b>2019</b> , 43, 988-1001   |     | 5  |
| 88 | High-Risk Human Papillomavirus Infection and the Risk of Cardiovascular Disease in Korean Women. <b>2019</b> , 124, 747-756  |     | 11 |
| 87 | A Cross-Sectional Study of the Association between Arterial Stiffness and Sarcopenia in Chinese Community-Dwelling Elderly Using the Asian Working Group for Sarcopenia Criteria. <b>2019</b> , 23, 195-201                          |     | 20 |
| 86 | Depression is Associated With Sarcopenia Due to Low Muscle Strength: Results From the ELSA-Brasil Study. <b>2019</b> , 20, 1641-1646   |     | 22 |
| 85 | Low Skeletal Muscle Mass Predicts Incident Dipstick Albuminuria in Korean Adults without Chronic Kidney Disease: A Prospective Cohort Study. <b>2019</b> , 141, 105-111  |     | 7  |
| 84 | Association between rosacea severity and relative muscle mass: A cross-sectional study. <b>2019</b> , 46, 11-17  |     | 3  |
| 83 | Associations of Fat and Lean Body Mass with Circulating Amino Acids in Older Men and Women. <b>2020</b> , 75, 885-891  |     | 6  |
| 82 | Transgenerational effects of maternal obesity and gestational diabetes on offspring body composition and left ventricle mass: the Finnish Gestational Diabetes Prevention Study (RADIEL) 6-year follow-up. <b>2020</b> , 37, 147-156 |     | 4  |
| 81 | Assessment of Skin Properties in Chronic Lymphedema: Measurement of Skin Stiffness, Percentage Water Content, and Transepidermal Water Loss. <i>Lymphatic Research and Biology</i> , <b>2020</b> , 18, 212-218                       | 2.3 | 2  |
| 80 | Equation models developed with bioelectric impedance analysis tools to assess muscle mass: A systematic review. <b>2020</b> , 35, 47-62  |     | 16 |
| 79 | The prevalence of sarcopenic obesity and its association with cognitive performance in type 2 diabetes in Singapore. <b>2020</b> , 39, 2274-2281   |     | 17 |

|    |  |       |
|----|--|-------|
| 78 | Association Between Peritonitis and Low Muscle Mass in Peritoneal Dialysis Patients. <b>2020</b> , 30, 341-346   | 2     |
| 77 | Relation between Lifestyle and Body Composition among Young Females in Serbia of 18-29 Years of Age. <b>2020</b> ,   |       |
| 76 | Low relative skeletal muscle mass predicts incident hypertension in Korean men: a prospective cohort study. <b>2020</b> , 38, 2223-2229  | 3     |
| 75 | Menopausal Transition, Body Mass Index, and Prevalence of Mammographic Dense Breasts in Middle-Aged Women. <b>2020</b> , 9,  | 1     |
| 74 | Measuring maternal body composition by biomedical impedance can predict risk for gestational diabetes mellitus: a retrospective study among 22,223 women. <b>2020</b> , 1-8  | 4     |
| 73 | Associations Between Nutrition Markers and Muscle Mass on Bioimpedance Analysis in Patients Receiving Parenteral Nutrition. <b>2021</b> , 45, 1089-1099  | 2     |
| 72 | Lean body mass is not beneficial, but may be detrimental for glucose tolerance - Splitting body mass index according to body composition. <b>2020</b> , 14, 747-752  | 1     |
| 71 | Lifestyle and glycemic health 5 years postpartum in obese and non-obese high diabetes risk women. <b>2020</b> , 57, 1453-1462  | 2     |
| 70 | The Comparison of Segmental Multifrequency Bioelectrical Impedance Analysis and Dual-Energy X-ray Absorptiometry for Estimating Fat Free Mass and Percentage Body Fat in an Ambulatory Population. <b>2021</b> , 45, 1231-1238 | 9     |
| 69 | Body image in COPD and its relation with physical activity levels, lung function and body composition: An observational study. <b>2020</b> , 14, 1182-1190   | 0     |
| 68 | Adaptations to Swimming Training in Athletes with Down Syndrome. <i>International Journal of Environmental Research and Public Health</i> , <b>2020</b> , 17,  | 4.6 1 |
| 67 | Physical activity is favorably associated with arterial stiffness in patients with obesity and elevated metabolic risk. <b>2020</b> , 74, e13563   | 3     |
| 66 | Sedentary Time Accumulated in Bouts is Positively Associated with Disease Severity in Fibromyalgia: The Al-Badlus Project. <b>2020</b> , 9,  | 2     |
| 65 | Thyroid Function and Sarcopenia: Results from the ELSA-Brasil Study. <b>2020</b> , 68, 1545-1553   | 9     |
| 64 | Body composition and changes in health-related quality of life in older age: a 10-year follow-up of the Helsinki Birth Cohort Study. <b>2020</b> , 29, 2039-2050   | 9     |
| 63 | Effects of volume status on body composition in incident peritoneal dialysis patients. <b>2020</b> , 74, 633-641   | 4     |
| 62 | Resistance training using different elastic components offers similar gains on muscle strength to weight machine equipment in Individuals with COPD: A randomized controlled trial. <b>2020</b> , 1-14                         | 4     |
| 61 | Skeletal muscle mass and risk of advanced adenoma in surveillance colonoscopy. <b>2020</b> , 35, 2088-2095   | 1     |

|    |  |      |     |
|----|--|------|-----|
| 60 | Prevalence and factors associated with motoric cognitive risk syndrome in community-dwelling older Chinese: a cross-sectional study. <b>2020</b> , 27, 1137-1145   |      | 6   |
| 59 | Effects of Body Weight vs. Lean Body Mass on Wingate Anaerobic Test Performance in Endurance Athletes. <b>2020</b> , 41, 545-551   |      | 1   |
| 58 | The prospective relationship between low muscle mass and thyroid hormones among 198'069 euthyroid men and women; comparing different definitions of low muscle mass. <b>2021</b> , 75, e13710                                      |      | 1   |
| 57 | Extracellular-to-intracellular water ratios are associated with functional disability levels in patients with knee osteoarthritis: results from the Nagahama Study. <i>Clinical Rheumatology</i> , <b>2021</b> , 40, 2889-2896     | 3.9  | 3   |
| 56 | Is active commuting associated with sedentary behaviour and physical activity in women with fibromyalgia? The al-Bidalus project. <i>Disability and Rehabilitation</i> , <b>2021</b> , 1-9   | 2.4  | 0   |
| 55 | A Comparison of Supine Versus Stand-on Bioimpedance Devices to Assess Breast Cancer-Related Lymphedema. <i>Lymphatic Research and Biology</i> , <b>2021</b> ,  | 2.3  | 1   |
| 54 | Variants in NEB and RIF1 genes on chr2q23 are associated with skeletal muscle index in Koreans: genome-wide association study. <i>Scientific Reports</i> , <b>2021</b> , 11, 2333  | 4.9  | 1   |
| 53 | Low muscle mass is associated with progression of chronic kidney disease and albuminuria - An 8-year longitudinal study in Asians with Type 2 Diabetes. <i>Diabetes Research and Clinical Practice</i> , <b>2021</b> , 174, 108777 | 7.4  | 2   |
| 52 | Weight Change and the Development of Nonalcoholic Fatty Liver Disease in Metabolically Healthy Overweight Individuals. <i>Clinical Gastroenterology and Hepatology</i> , <b>2021</b> ,   | 6.9  | 1   |
| 51 | Body Fat Mass, Percent Body Fat, Fat-Free Mass, and Skeletal Muscle Mass Reference Curves for Czech Children Aged 6-11 Years. <i>Children</i> , <b>2021</b> , 8,   | 2.8  | 0   |
| 50 | Association between skeletal muscle mass and mammographic breast density. <i>Scientific Reports</i> , <b>2021</b> , 11, 16785  | 4.9  | 0   |
| 49 | Total and regional fat-to-muscle mass ratio measured by bioelectrical impedance and risk of incident type 2 diabetes. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , <b>2021</b> ,  | 10.3 | 2   |
| 48 | Both lean and fat body mass associate with blood pressure. <i>European Journal of Internal Medicine</i> , <b>2021</b> , 91, 40-44  | 3.9  | 6   |
| 47 | A high lean body mass is not protecting from type 2 diabetes in the presence of a high body fat mass. <i>Diabetes and Metabolism</i> , <b>2021</b> , 47, 101219  | 5.4  | 5   |
| 46 | Novel Phototherapy Kiosk Shows Promise as a Treatment Option for Low Vitamin D. <i>Military Medicine</i> , <b>2021</b> , 186, 722-728  | 1.3  | 1   |
| 45 | Proposal of a normative table for body fat percentages of Brazilian young adults through bioimpedanciometry. <i>Journal of Exercise Rehabilitation</i> , <b>2018</b> , 14, 974-979   | 1.8  | 10  |
| 44 | Regular exercise is associated with a reduction in the risk of NAFLD and decreased liver enzymes in individuals with NAFLD independent of obesity in Korean adults. <i>PLoS ONE</i> , <b>2012</b> , 7, e46819                      | 3.7  | 128 |
| 43 | The correlates of body composition with heart rate recovery after step test: an exploratory study of Malaysian adolescents. <i>PLoS ONE</i> , <b>2013</b> , 8, e82893  | 3.7  | 12  |

|    |  |     |    |
|----|--|-----|----|
| 42 | Non-alcoholic fatty liver disease and the development of nephrolithiasis: A cohort study. <i>PLoS ONE</i> , <b>2017</b> , 12, e0184506   | 3.7 | 8  |
| 41 | Impact of Dapagliflozin Therapy on Renal Protection and Kidney Morphology in Patients With Uncontrolled Type 2 Diabetes Mellitus. <i>Journal of Clinical Medicine Research</i> , <b>2018</b> , 10, 466-477                     | 2.9 | 11 |
| 40 | Initial Acute Decline in Estimated Glomerular Filtration Rate After Sodium-Glucose Cotransporter-2 Inhibitor in Patients With Chronic Kidney Disease. <i>Journal of Clinical Medicine Research</i> , <b>2020</b> , 12, 724-733 | 3.9 | 1  |
| 39 | Physical Function and Cardio-Ankle Vascular Index in Elderly Heart Failure Patients. <i>International Heart Journal</i> , <b>2020</b> , 61, 769-775  | 1.8 | 5  |
| 38 | Capacidade de avaliação de testes de desempenho físico para identificar baixa massa muscular em mulheres de meia-idade e idosas. <i>Revista Brasileira De Geriatria E Gerontologia</i> , <b>2020</b> , 23,                     | 0.8 | 1  |
| 37 | Fatty liver is an independent risk factor for gallbladder polyps. <i>World Journal of Gastroenterology</i> , <b>2020</b> , 26, 6979-6992   | 5.6 | 8  |
| 36 | Obesity Determined by Different Measures and its Impact on the Health-Related Quality of Life of Young-Adult Nigerians. <i>Asian Journal of Clinical Nutrition</i> , <b>2015</b> , 7, 64-75                                    | 0   | 1  |
| 35 | Association of appendicular skeletal muscle mass with carotid intima-media thickness according to body mass index in Korean adults. <i>Epidemiology and Health</i> , <b>2018</b> , 40, e2018049                                | 5.6 | 9  |
| 34 | Peritoneal-Membrane Characteristics and Hypervolemia Management in Peritoneal Dialysis: A Randomized Control Trial. <i>Membranes</i> , <b>2021</b> , 11,   | 3.8 |    |
| 33 | Association between Muscle Mass Deficits and the Non-Alcoholic Fatty Liver Disease Fibrosis Score in Adults without Central Obesity. <i>Korean Journal of Family Practice</i> , <b>2021</b> , 11, 345-351                      | 0.4 |    |
| 32 | Correlation of drinking water nutritional element levels with body composition of women aged 55-70 years living in Batman province. <i>Dicle Medical Journal</i> , <b>2017</b> , 44, 99-99                                     | 0   |    |
| 31 | Correlation between the Cross-sectional Area of the Multifidus Muscle at Each Vertebral Level and Body Composition. <i>Rigakuryoho Kagaku</i> , <b>2018</b> , 33, 733-737  | 0.1 | 1  |
| 30 | Association of Age at Menarche With Left Ventricular Diastolic Dysfunction in Middle-Aged Women. <i>Circulation Journal</i> , <b>2018</b> , 82, 708-714  | 2.9 | 1  |
| 29 | BIA?????????. <i>The Japanese Journal of SURGICAL METABOLISM and NUTRITION</i> , <b>2019</b> , 53, 123-130   | 0   | 0  |
| 28 | Association of Serum Amino Acid Concentration With Loss of Skeletal Muscle Mass After 1 Year in Cardiac Rehabilitation Center Patients. <i>Circulation Reports</i> , <b>2019</b> , 1, 456-461                                  | 0.7 | 1  |
| 27 | Comparison of two bioelectrical impedance analysis instruments for determining body composition in adolescent girls. <i>International Journal of Body Composition Research</i> , <b>2006</b> , 4, 153-160                      |     | 19 |
| 26 | Cross-calibration of pencil-beam (DPX-NT) and fan-beam (QDR-4500C) dual-energy X-ray absorptiometry for sarcopenia. <i>Nagoya Journal of Medical Science</i> , <b>2015</b> , 77, 647-52  | 0.7 | 2  |
| 25 | Using nutritional assessment to predict gender-specific risks of sarcopenia among the elderly receiving health check-up in Taipei. <i>Journal of Geriatric Cardiology</i> , <b>2016</b> , 13, 261-6                            | 1.7 |    |

|    |  |     |   |
|----|--|-----|---|
| 24 | [Clinical guidelines «Obesity in children»]. <i>Problemy Endokrinologii</i> , <b>2021</b> , 67, 67-83  | 0   | 5 |
| 23 | Impact of muscle mass on blood glucose level. <i>Journal of Basic and Clinical Physiology and Pharmacology</i> , <b>2021</b> ,   | 1.6 |   |
| 22 | Association of sleep disturbances with sarcopenia and its defining components: the ELSA-Brasil study. <i>Brazilian Journal of Medical and Biological Research</i> , <b>2021</b> , 54, e11539                 | 2.8 | 0 |
| 21 | Excess Weight and Body Fat Percentage Associated with Waist Circumference as a Cardiometabolic Risk Factor in University Students.. <i>Scientifica</i> , <b>2022</b> , 2022, 1310030                         | 2.6 | 1 |
| 20 | Relationship between Seasonal Changes in Food Intake and Energy Metabolism, Physical Activity, and Body Composition in Young Japanese Women.. <i>Nutrients</i> , <b>2022</b> , 14,                           | 6.7 | 0 |
| 19 | Serum 25-hydroxy vitamin D and the risk of low muscle mass in young and middle-aged Korean adults.. <i>European Journal of Endocrinology</i> , <b>2022</b> ,   | 6.5 | 0 |
| 18 | Comparison of the effects of empagliflozin and glimepiride on endothelial function in patients with type 2 diabetes: A randomized controlled study.. <i>PLoS ONE</i> , <b>2022</b> , 17, e0262831            | 3.7 | 1 |
| 17 | Bicycle Touring 480 km in Seven Days: Effects on Body Composition and Physical Fitness-A Case Study.. <i>International Journal of Environmental Research and Public Health</i> , <b>2022</b> , 19,           | 4.6 |   |
| 16 | Concerns in using Segment Impedances for Estimation Body Composition in 8-polar Bioimpedance Measurement System. <b>2022</b> ,   |     |   |
| 15 | Amino acid profile of skeletal muscle loss in type 2 diabetes: results from a 7-year longitudinal study in Asians.. <i>Diabetes Research and Clinical Practice</i> , <b>2022</b> , 109803                    | 7.4 | 0 |
| 14 | The effect of metabolic health and obesity on lung function: A cross sectional study of 114,143 participants from Kangbuk Samsung Health Study.. <i>PLoS ONE</i> , <b>2022</b> , 17, e0266885                | 3.7 | 0 |
| 13 | Adiposity-Related Predictors of Vascular Aging From a Life Course Perspective-Findings From the Helsinki Birth Cohort Study.. <i>Frontiers in Cardiovascular Medicine</i> , <b>2022</b> , 9, 865544          | 5.4 |   |
| 12 | The importance of muscle mass in predicting intradialytic hypotension in patients undergoing maintenance hemodialysis.. <i>Kidney Research and Clinical Practice</i> , <b>2022</b> ,                         | 3.6 | 0 |
| 11 | Influence of acute water ingestion and prolonged standing on raw bioimpedance and subsequent body fluid and composition estimates. <i>Journal of Electrical Bioimpedance</i> , <b>2022</b> , 13, 10-20       | 1.5 | 1 |
| 10 | Metabolically healthy and unhealthy obesity and risk of vasomotor symptoms in premenopausal women: cross-sectional and cohort studies. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , | 3.7 | 1 |
| 9  | Effect of body posture on segmental multifrequency bioimpedance variables.   |     | 0 |
| 8  | Sex-specific associations between gut microbiota and skeletal muscle mass in a population-based study.   |     | 0 |
| 7  | Influence of the nutritional status on facial morphology in young Japanese women. <b>2022</b> , 12,  |     | 0 |

- 6 Body composition in late midlife as a predictor of accelerated age-associated deficit accumulation from late midlife into old age: a longitudinal birth cohort study. ○
- 5 Age-specific and sex-specific associations of visceral adipose tissue mass and fat-to-muscle mass ratio with risk of mortality. ○
- 4 Gradual Strength Training Improves Sleep Quality, Physical Function and Pain in Women with Fibromyalgia. **2022**, 19, 15662 ○
- 3 Effects of a 14-week community health program of exercise and learning/education in older adults: A single-arm pre-post comparison study. **2023**, 51, 1-8 ○
- 2 Influence of Upper-Extremity and Lower-Extremity Resistance Exercise on Segmental Body Composition and Body Fluid Estimates. **2022**, Publish Ahead of Print, ○
- 1 The impact of insulin resistance on the association between metabolic syndrome and lung function: the Kangbuk Samsung Health Study. **2023**, 15, ○