CITATION REPORT List of articles citing

Educational and organizational interventions to improve the management of depression in primary care: a systematic review

DOI: 10.1001/jama.289.23.3145 JAMA - Journal of the American Medical Association, 2003, 289, 3145-51.

Source: https://exaly.com/paper-pdf/35935840/citation-report.pdf

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper IF	Citations
549	Quality improvements in clinical therapeutics. 2003 , 16, 725-729	
548	Achieving long-term optimal outcomes in geriatric depression and anxiety. 2003, 8, 27-34	13
547	Web and telecounselling in Australia. 2004 , 180, 604-5	11
546	Can care management enhance integration of primary and specialty care?. 2004, 329, 605	8
545	Self-help: Empowering service users or aiding cash strapped mental health services?. 2004 , 13, 117-123	31
544	Effects of a primary care-based intervention on violent behavior and injury in children. 2004, 114, e392-9	73
543	Re-engineering systems for the treatment of depression in primary care: cluster randomised controlled trial. 2004 , 329, 602	190
542	Effectiveness of screening and treatment for depression in ambulatory indigent patients. 2004 , 19, 78-84	29
541	Depression with atypical features in a sample of primary care outpatients: prevalence, specific characteristics and consequences. 2004 , 83, 237-42	21
540	Antidepressant drug use in Lombardy, Italy: a population-based study. 2004, 83, 169-75	44
539	Use of brief depression screening tools in primary care: consideration of heterogeneity in performance in different patient groups. 2004 , 26, 190-8	103
538	Differences in the indicators of depressive symptoms among a community sample of African-American and Caucasian older adults. 2004 , 40, 309-31	29
537	Building concordant relationships with patients starting antidepressant medication. 2004 , 55, 241-6	56
536	Recognizing and treating depression in patients with diabetes. 2004 , 4, 119-25	83
535	Costs and quality in the treatment of acute depression in primary care: a comparison between England, Germany and Switzerland. 2004 , 19, 201-8	5
534	The 13- and 20-item Hopkins Symptom Checklist Depression Scale: psychometric properties in primary care patients with minor depression or dysthymia. 2004 , 34, 37-50	40
533	Which organizational characteristics are associated with increased management of depression using antidepressants in US nursing homes?. <i>Medical Care</i> , 2004 , 42, 992-1000	46

[2006-2004]

532	Disease management programs for depression: a systematic review and meta-analysis of randomized controlled trials. <i>Medical Care</i> , 2004 , 42, 1211-21	160
531	NICE, but will they help people with depression? The new National Institute for Clinical Excellence depression guidelines. 2005 , 186, 177-8	21
530	Effect of comorbid chronic diseases on prevalence and odds of depression in adults with diabetes. 2005 , 67, 46-51	74
529	Training in neuropsychiatry: is it time to reintegrate into mainstream psychiatry?. 2005 , 29, 361-364	6
528	Mental health in general practice: assessment of current state and future needs. 2005 , 22, 83-86	31
527	Assessing the implementation of the chronic care model in quality improvement collaboratives. 2005 , 40, 978-96	146
526	The development and implementation of an educational intervention on first episode psychosis for primary care. 2005 , 39, 1006-14	20
525	District nurses' involvement and attitudes to mental health problems: a three-area cross-sectional study. 2005 , 14, 976-85	39
524	Telephonic case-finding of major depression in a Medicaid chronic disease management program for diabetes and heart failure. 2005 , 27, 338-43	21
523	Feasibility and acceptability of a telephone psychotherapy program for depressed adults treated in primary care. 2005 , 27, 400-10	29
522	Overt and covert barriers to the integration of primary and specialist mental health care. 2005 , 61, 1785-94	57
521	Case management for the treatment of patients with major depression in general practicesrationale, design and conduct of a cluster randomized controlled trialPRoMPT (PRimary care Monitoring for depressive Patient's Trial) [ISRCTN66386086]study protocol. 2005 , 5, 101	28
520	Depression. 2005 , 50, 265-273	6
519	General Practitioners' opinions on their practice in mental health and their collaboration with mental health professionals. 2005 , 6, 18	44
518	General practitioners' conceptions about treatment of depression and factors that may influence their practice in this area. A postal survey. 2005 , 6, 21	13
517	Perioperative beta-blockade: guidelines and practice in New Zealand. 2005 , 33, 645-50	4
516	Remission in depressed geriatric primary care patients: a report from the PROSPECT study. 2005 , 162, 718-24	157
515	The potential impact of the recovery movement on family interventions for schizophrenia: opportunities and obstacles. 2006 , 32, 451-63	112

514	Enhancing end-of-life care in nursing homes: Palliative Care Educational Resource Team (PERT) program. 2005 , 8, 556-66	43
513	A randomized effectiveness trial of cognitive-behavioral therapy and medication for primary care panic disorder. 2005 , 62, 290-8	193
512	Stepped care in psychological therapies: access, effectiveness and efficiency. Narrative literature review. 2005 , 186, 11-7	609
511	Role of specialists in common chronic diseases. 2005 , 330, 651-3	29
510	Depression in the elderly. 2005 , 365, 1961-70	1098
509	From establishing an evidence-based practice to implementation in real-world settings: IMPACT as a case study. 2005 , 28, 1079-92	32
508	Screening and case finding instruments for depression. 2005 , CD002792	102
507	Evaluation and Treatment of Depression (Part I). 2005 , 13, 295-306	7
506	Evaluation and Treatment of Depression in Managed Care (Part II). 2005, 13, 307-316	2
505	Improving women's health during internatal periods: developing an evidenced-based approach to addressing maternal depression in pediatric settings. 2006 , 15, 692-703	35
504	[Programmes against depression]. 2006 , 32, 9-15	2
503	Disease-focused or integrated treatment: diabetes and depression. 2006 , 90, 627-46	19
502	Collaborative care for depression in primary care. Making sense of a complex intervention: systematic review and meta-regression. 2006 , 189, 484-93	293
501	Adherence to treatment in mood disorders. 2006 , 19, 44-9	33
500	Do patient requests for antidepressants enhance or hinder physicians' evaluation of depression? A randomized controlled trial. <i>Medical Care</i> , 2006 , 44, 1107-13	12
499	Evaluation of a Train-the-Trainer Program to Enhance Hospice and Palliative Care in Nursing Homes. 2006 , 8, 42-49	23
498	Management of the cardinal features of andropause. 2006 , 13, 145-60	19
497	Organizational interventions to implement improvements in patient care: a structured review of reviews. 2006 , 1, 2	147

(2006-2006)

496	What should non-US behavioral health systems learn from the USA?: US behavior health services trends in the 1980s and 1990s. 2006 , 60, 261-70	3
495	Medical patients' attitudes toward emotional problems and their treatment. What do they really want?. 2006 , 21, 39-45	59
494	Depression in primary care: encouragement and caution for the business case. 2006 , 21, 1125-7	6
493	Increased antidepressant use and fewer suicides in Jthland county, Sweden, after a primary care educational programme on the treatment of depression. 2006 , 114, 159-67	59
492	An educational programme for peer review groups to improve treatment of chronic heart failure and diabetes mellitus type 2 in general practice. 2006 , 12, 613-21	23
491	Realigning clinical and economic incentives to support depression management within a medicaid population: the Colorado access experience. 2006 , 33, 26-33	7
490	Realigning economic incentives for depression care at UCSF. 2006 , 33, 34-8	5
489	Implementation of guideline-based care for depression in primary care. 2006 , 33, 43-53	24
488	Evaluation of "depression in primary care" innovations. 2006 , 33, 86-91	12
487	Toward the implementation of mental health consumer provider services. 2006 , 33, 176-95	96
486	Improving general medical care for persons with mental and addictive disorders: systematic review. 2006 , 28, 145-53	132
485	Pebbles in a pond: NIMH grants stimulate improvements in primary care treatment of depression. 2006 , 28, 185-8	10
484	Developing a U.K. protocol for collaborative care: a qualitative study. 2006 , 28, 296-305	54
483	A systematic review of complex system interventions designed to increase recovery from depression in primary care. 2006 , 6, 88	126
482	Ethnography's role in assisting mental health research and clinical practice. 2006, 62, 859-79	16
481	Dissemination of evidence-based mental health interventions: importance to the trauma field. 2006 , 19, 611-23	25
480	Should we screen for depression?. 2006 , 332, 1027-30	138

478	Raising the awareness of primary care providers about postpartum depression. 2006 , 27, 59-73	17
477	Challenges to implementing and sustaining comprehensive mental health service programs. 2006 , 29, 195-218	30
476	Identification and management of depression in primary care settings. A meta-review of evidence. 2006 , 15, 276-283	14
475	The logic behind a multimethod intervention to improve adherence to clinical practice guidelines in a nationwide network of primary care practices. 2006 , 29, 65-88	45
474	Challenges in the treatment of anxiety disorders: beyond guidelines. 2006 , 10 Suppl 3, 18-24	1
473	Costs and consequences of enhanced primary care for depression: systematic review of randomised economic evaluations. 2006 , 189, 297-308	134
472	A taxonomy for disease management: a scientific statement from the American Heart Association Disease Management Taxonomy Writing Group. 2006 , 114, 1432-45	2 10
471	In-Home Cognitive Behavior Therapy for a Depressed Mother in a Home Visitation Program. 2007 , 6, 161-180	23
47°	Guidelines for Adolescent Depression in Primary Care (GLAD-PC): II. Treatment and ongoing management. 2007 , 120, e1313-26	179
469	Telephone screening, outreach, and care management for depressed workers and impact on clinical and work productivity outcomes: a randomized controlled trial. <i>JAMA - Journal of the American</i> 27.4 <i>Medical Association</i> , 2007 , 298, 1401-11	274
468	Care management of poststroke depression: a randomized, controlled trial. 2007, 38, 998-1003	101
467	Reducing health disparities in depressive disorders outcomes between non-Hispanic Whites and ethnic minorities: a call for pragmatic strategies over the life course. 2007 , 64, 157S-94S	51
466	Antidepressant treatment of veterans with Parkinson's disease and depression: analysis of a national sample. 2007 , 20, 161-5	38
465	Factors associated with integrating self-management support into primary care. 2007 , 33 Suppl 6, 126S-131S	12
464	Guidelines for Adolescent Depression in Primary Care (GLAD-PC): I. Identification, assessment, and initial management. 2007 , 120, e1299-312	150
463	Collaborative care for depression and anxiety problems. 2007,	6
462	Treatment adequacy for anxiety and depressive disorders in six European countries. 2007, 190, 172-3	121

460	Models of care for treating late-life depression in primary care. 2007 , 3, 67-75	2
459	Primary care physicians' discussion of emotional distress and patient satisfaction. 2007 , 37, 331-45	11
458	Acc et continuit des soins pour les troubles mentaux les plus frquents : les consultations de soins partag du RPSM 78. 2007 , 165, 741-745	1
457	Randomized trials with concurrent economic evaluations reported unrepresentatively large clinical effect sizes. 2007 , 60, 781-6	10
456	[Improving the management of depression in primary care: review and prospects]. 2007, 33, 552-60	4
455	Adherence to antidepressant treatment. 2007 , 7, 57-62	38
454	Reducing the societal burden of depression: a review of economic costs, quality of care and effects of treatment. 2007 , 25, 7-24	141
453	Primary care - mental health collaboration: an example of assessing usual practice and potential barriers. 2007 , 21, 207-16	29
452	Best practice case management for improved medical adherence. 2007, 44, 161-77	10
451	Perfil sintom l ico de los pacientes con depresili tratados en atencili primaria. Un estudio epidemioligico. 2007 , 14, 47-52	1
450	New challenges in family interventions for schizophrenia. 2007 , 7, 33-43	16
449	Introduction. 2007 , 12, 3-3	
448	Increased access to evidence-based primary mental health care: will the implementation match the rhetoric?. 2007 , 187, 100-3	46
447	Introduction. 2007 , 12, 3-3	11
446	Current practices in depression care. 2007 , 27 Suppl 1, S9-17	6
445	How quality improvement interventions can address disparities in depression. 2007 , 27 Suppl 1, S33-9	4
444	Improving depression care for ethnic and racial minorities: a concept for an intervention that integrates CME planning with improvement strategies. 2007 , 27 Suppl 1, S65-74	3
443	The effects of a shared decision-making intervention in primary care of depression: a cluster-randomized controlled trial. 2007 , 67, 324-32	266

442	Depression and the nature of Trinidadian family practice: a cross-sectional study. 2007, 8, 25	11
441	Depression and anxiety, an indicated prevention (DIP) protocol in homes for the elderly: feasibility and (cost) effectiveness of a stepped care programme. 2007 , 7, 6	13
440	Primary care provider preferences for working with a collaborative support team. 2007 , 2, 16	9
439	Depressed patients' perceptions of depression treatment decision-making. 2007 , 10, 62-74	36
438	Effects of a nationwide programme: interventions to reduce perceived barriers to collaboration and to increase structural one-on-one contact. 2007 , 13, 860-6	8
437	Teaching junior doctors to manage patients who somatise: is it possible in an afternoon?. 2007 , 41, 995-1001	14
436	The comparison of effectiveness of two modalities of mental health nurse follow-up programmes for female outpatients with depression in Taipei, Taiwan. 2007 , 16, 1141-50	8
435	The methodological quality of economic evaluations of guideline implementation into clinical practice: a systematic review of empiric studies. 2007 , 10, 305-16	47
434	Assessment of an enhanced program for depression management in primary care: a cluster randomized controlled trial. The INDI project (Interventions for Depression Improvement). 2007 , 7, 253	17
433	Systematic review of multifaceted interventions to improve depression care. 2007 , 29, 91-116	196
432	Design of the Coordinated Anxiety Learning and Management (CALM) study: innovations in collaborative care for anxiety disorders. 2007 , 29, 379-87	80
431	Major depression in individuals with chronic medical disorders: prevalence, correlates and association with health resource utilization, lost productivity and functional disability. 2007 , 29, 409-16	420
430	Controlled trial of problem-solving therapy and consultation-liaison for common mental disorders in general medical settings in Taiwan. 2007 , 29, 402-8	10
429	[Changes in the frequency of suicidal behaviour after a 2-year intervention campaign]. 2007 , 78, 272-6, 278-80, 282	6
428	[Innovative care models for treating depression]. 2007, 78 Suppl 3, 585-94; quiz 595	7
427	Learning by doing: a novel approach to improving general practitioners' diagnostic skills for common mental disorders. 2007 , 119, 117-23	7
426	Increase in observed mental health difficulties one year after acute coronary syndrome: general practitioner survey. 2007 , 176, 205-9	1
425	Failure to recognize depression in primary care: issues and challenges. 2007 , 22, 701-3	36

(2008-2008)

424	How well do GPs fulfill their educator role in consultations for depression and anxiety?. 2008 , 73, 389-95	4
423	Impact of shared mental health care in the general population on subjects' perceptions of mental health status. 2008 , 43, 113-20	3
422	Ambulante psychotherapeutische Versorgung depressiver Patienten. 2008, 53, 260-267	1
421	Family advocacy, support and education in children's mental health: results of a national survey. 2008 , 35, 73-83	29
420	Models in the delivery of depression care: a systematic review of randomised and controlled intervention trials. 2008 , 9, 25	56
419	Long term GP opinions and involvement after a consultation-liaison intervention for mental health problems. 2008 , 9, 41	9
418	Developing a national dissemination plan for collaborative care for depression: QUERI Series. 2008 , 3, 59	33
417	A multicenter study of major depressive disorder among emergency department patients in Latin-American countries. 2008 , 25, E199-204	11
416	Strategies for facilitating change in alcohol and other drugs (AOD) professional practice: a systematic review of the effectiveness of reminders and feedback. 2008 , 27, 548-58	9
415	Improving the role of nursing in the treatment of depression in primary care in Spain. 2008, 44, 248-58	14
415 414	Improving the role of nursing in the treatment of depression in primary care in Spain. 2008, 44, 248-58 The effects of quality improvement for depression in primary care at nine years: results from a randomized, controlled group-level trial. 2008, 43, 1952-74	14 31
	The effects of quality improvement for depression in primary care at nine years: results from a	
414	The effects of quality improvement for depression in primary care at nine years: results from a randomized, controlled group-level trial. 2008 , 43, 1952-74	31
414	The effects of quality improvement for depression in primary care at nine years: results from a randomized, controlled group-level trial. 2008, 43, 1952-74 Discussion of depression in follow-up medical visits with older patients. 2008, 56, 16-22 Patient education and group counselling to improve the treatment of depression in primary care: a	31
414 413 412	The effects of quality improvement for depression in primary care at nine years: results from a randomized, controlled group-level trial. 2008, 43, 1952-74 Discussion of depression in follow-up medical visits with older patients. 2008, 56, 16-22 Patient education and group counselling to improve the treatment of depression in primary care: a randomized controlled trial. 2008, 105, 235-40 Prevalence and rates of recognition of depressive disorders in internal medicine outpatient	31 12 20
414 413 412 411	The effects of quality improvement for depression in primary care at nine years: results from a randomized, controlled group-level trial. 2008, 43, 1952-74 Discussion of depression in follow-up medical visits with older patients. 2008, 56, 16-22 Patient education and group counselling to improve the treatment of depression in primary care: a randomized controlled trial. 2008, 105, 235-40 Prevalence and rates of recognition of depressive disorders in internal medicine outpatient departments of 23 general hospitals in Shenyang, China. 2008, 110, 46-54 Improving the prevention and management of chronic disease in low-income and middle-income	31 12 20 50
414 413 412 411 410	The effects of quality improvement for depression in primary care at nine years: results from a randomized, controlled group-level trial. 2008, 43, 1952-74 Discussion of depression in follow-up medical visits with older patients. 2008, 56, 16-22 Patient education and group counselling to improve the treatment of depression in primary care: a randomized controlled trial. 2008, 105, 235-40 Prevalence and rates of recognition of depressive disorders in internal medicine outpatient departments of 23 general hospitals in Shenyang, China. 2008, 110, 46-54 Improving the prevention and management of chronic disease in low-income and middle-income countries: a priority for primary health care. 2008, 372, 940-9 Treatment of depressive symptoms and attitudes towards treatment options in a representative	31 12 20 50 382

406	General practitioners understanding of depression in young people: qualitative study. 2008, 9, 269	19
405	Care management for depression in primary care practice: findings from the RESPECT-Depression trial. 2008 , 6, 30-7	46
404	Depression in primary care: three key challenges. 2008 , 84, 545-8	12
403	Opportunities for improving management of advanced chronic kidney disease. 2008 , 23, 184-92	7
402	Development and evaluation of a Nursing Assistant Computerized Education Programme. 2008, 14, 502-9	13
401	Screening and case-finding instruments for depression: a meta-analysis. 2008 , 178, 997-1003	312
400	Management of chronic hepatitis C in veterans: the potential of integrated care models. 2008, 103, 1810-23	46
399	Battling depression. 2008 , 178, 1023-4	6
398	Women's care-seeking experiences after referral for postpartum depression. 2008, 18, 1161-73	80
397	Collaborative care for depression in UK primary care: a randomized controlled trial. 2008, 38, 279-87	120
396	Research Priorities for P atient-Centred Mental Health Services: Findings From a National Consultation. 2008 , 13, 33-43	2
395	Organising health care services for persons with an intellectual disability. 2008, CD007492	36
394	Inducing sustainable improvement in depression care in primary care practices. 2008, 34, 247-55	17
393	Consultation liaison in primary practice for mental health problems. 2008,	4
392	Educational interventions. 116-131	
391	Collaborative mental health and primary care for bipolar disorder. 2008, 14 Suppl 2, 55-64	5
390	Gaps in depression care: why primary care physicians should hone their depression screening, diagnosis, and management skills. 2008 , 50, 451-8	35
389	Whither DSM and ICD, Chapter V?. 2008 , 13, 4-15	18

The time dimension:. 129-140

387	The central role of staff for better mental health care. 153-164		
386	Psychiatry in primary care. 479-497		17
385	BIBLIOTHERAPY AS AN ADJUNCTIVE TREATMENT. 2008 , 7-39		2
384	Knowledge transfer and improvement of primary and ambulatory care for patients with anxiety. 2008 , 53, 277-93		16
383	A Quality Improvement Initiative for Depression: Finally, a Model for use in R ealFamily Physician Time. 2008 , 27, 191-199		
382	Coordinated care in the management of patients with unexplained physical symptoms: depression is a key issue. 2008 , 188, S133-7		15
381	Plenty of activity but little outcome data: a review of the "grey literature" on primary care anxiety and depression programs in Australia. 2008 , 188, S103-6		3
380	Recurrence rates in Ontario physicians monitored for major depression and bipolar disorder. 2009 , 54, 777-82		6
379	[Which are the recent clinical findings regarding the association between depression and suicide?]. 2009 , 31 Suppl 1, S18-25		21
378	Primary care contact prior to suicide in individuals with mental illness. <i>British Journal of General Practice</i> , 2009 , 59, 825-32	1.6	81
377	Disease Management Programmes for Major Depression: Making the Financial Case. 2009 , 15, 679-725		
376	Simulated physician learning intervention to improve safety and quality of diabetes care: a randomized trial. 2009 , 32, 585-90		36
375	Improving outpatient care of depression by implementing practice guidelines: a controlled clinical trial. 2009 , 21, 29-36		13
374	Cross-national analysis of the associations among mental disorders and suicidal behavior: findings from the WHO World Mental Health Surveys. 2009 , 6, e1000123		482
373	Reducing suicidal ideation and depression in older primary care patients: 24-month outcomes of the PROSPECT study. 2009 , 166, 882-90		258
372	Collaborative Depression Trial (CADET): multi-centre randomised controlled trial of collaborative care for depressionstudy protocol. 2009 , 9, 188		27
371	Researching the mental health needs of hard-to-reach groups: managing multiple sources of evidence. 2009 , 9, 226		31

370	The Indiana Cancer Pain and Depression (INCPAD) trial Design of a telecare management intervention for cancer-related symptoms and baseline characteristics of study participants. 2009 , 31, 240-53	57
369	Adherence to evidence-based guidelines for depression and anxiety disorders is associated with recording of the diagnosis. 2009 , 31, 460-9	56
368	Clinic-level process of care for depression in primary care settings. 2009 , 36, 144-58	7
367	The applications of PROs in clinical practice: what are they, do they work, and why?. 2009 , 18, 115-23	413
366	Translating evidence to practice: two stories from the field. 2009 , 16, 47-57	21
365	The train is leaving the station: is psychology aboard?. 2009 , 16, 1-3	3
364	[Evaluation of Germany's sixth national health target entitled "Depressive illnesses - prevention, early diagnosis, sustainable treatment"]. 2009 , 52, 897-904	8
363	Managing high-acuity-depressed adults in primary care. 2009 , 21, 362-70	2
362	Is it clinically and cost effective to screen for postnatal depression: a systematic review of controlled clinical trials and economic evidence. 2009 , 116, 1019-27	46
361	[Current and lifetime depression as a risk factor for hospital admission in patients older than 74 years attended in an emergency department]. 2009 , 44, 305-10	3
360	[Family doctors and psychiatrists and the patient with depression: the need to re-adjust health care approaches and organisational dynamics]. 2009 , 41, 33-40	2
359	The International Study on General Practitioners and Early Psychosis (IGPS). 2009 , 108, 182-90	27
358	Screening for depression in high-risk groups: prospective cohort study in general practice. 2009 , 194, 399-403	34
357	The bright side of being blue: depression as an adaptation for analyzing complex problems. 2009 , 116, 620-54	358
356	Improving adherence in mood disorders: the struggle against relapse, recurrence and suicide risk. 2009 , 9, 985-1004	67
355	An Ultra-Brief Screening Scale for Anxiety and Depression: The PHQA. 2009, 50, 613-621	152
354	Objective: remission of depression in primary care The Oreon Study. 2009 , 19, 169-76	21
353	[Patient safety and a culture of responsibility in ambulatory care: strategies for improving practice]. 2009 , 103, 510-4	

352	Clinical diagnosis of depression in primary care: a meta-analysis. 2009 , 374, 609-19	714
351	Reducing psychosocial and behavioral pregnancy risk factors: results of a randomized clinical trial among high-risk pregnant african american women. 2009 , 99, 1053-61	62
350	REDIRECT: cluster randomised controlled trial of GP training in first-episode psychosis. <i>British Journal of General Practice</i> , 2009 , 59, e183-90	59
349	Organizational factors and depression management in community-based primary care settings. 2009 , 4, 84	14
348	System factors affect the recognition and management of posttraumatic stress disorder by primary care clinicians. <i>Medical Care</i> , 2009 , 47, 686-94	24
347	Gatekeeper training as a preventative intervention for suicide: a systematic review. 2009 , 54, 260-8	233
346	On-site mental health workers delivering psychological therapy and psychosocial interventions to patients in primary care: effects on the professional practice of primary care providers. 2009 , CD000532	36
345	Positive screens for psychiatric disorders in primary care: a long-term follow-up of patients who were not in treatment. 2010 , 61, 151-9	27
344	Which physician and practice characteristics are associated with adherence to evidence-based guidelines for depressive and anxiety disorders?. <i>Medical Care</i> , 2010 , 48, 240-8	52
343	Initiation of Primary Care-Mental Health Integration programs in the VA Health System: associations with psychiatric diagnoses in primary care. <i>Medical Care</i> , 2010 , 48, 843-51	80
342	A review of the costs associated with depression and treatment noncompliance: the potential benefits of online support. 2010 , 25, 288-96	26
341	Twelve evidence-based principles for implementing self-management support in primary care. 2010 , 36, 561-70	125
340	Development of a patient-report measure of psychotherapy for depression. 2010 , 37, 245-53	11
339	Mind and body reunited: improving care at the behavioral and primary healthcare interface. 2010 , 37, 529-42	24
338	Organizational readiness in specialty mental health care. 2010 , 25 Suppl 1, 27-31	22
337	Implementation of a family intervention for individuals with schizophrenia. 2010 , 25 Suppl 1, 32-7	24
336	Assessing and treating pain in hospices: current state of evidence-based practices. 2010 , 39, 803-19	53
335	General practitioners' opinions on how to improve treatment of mental disorders in primary health care. Interviews with one hundred Norwegian general practitioners. 2010 , 10, 35	26

334	Suicide prevention in primary care: General practitioners' views on service availability. 2010 , 3, 246	23
333	Falling through the net - black and minority ethnic women and perinatal mental healthcare: health professionals' views. 2010 , 32, 17-25	45
332	What is the role of consultation-liaison psychiatry in the management of depression in primary care? A systematic review and meta-analysis. 2010 , 32, 246-54	29
331	Guideline-concordant antidepressant use among patients with major depressive disorder. 2010 , 32, 360-7	16
330	Research article: Antidepressant use among Asians in the United States. 2010 , 27, 46-55	14
329	Improving child protection in the emergency department: a systematic review of professional interventions for health care providers. 2010 , 17, 117-25	28
328	Evidence-based treatment for patients with obsessive disorder: questions and controversies. 249-259	
327	The treatment of anxiety disorders in primary care. 348-355	
326	Prodromal schizophrenia in primary care: a randomised sensitisation study. <i>British Journal of General Practice</i> , 2010 , 60, e353-9	6
325	Linking evidence to practice. 31-54	
325 324	Linking evidence to practice. 31-54 Long-term clinical outcomes of care management for chronically depressed primary care patients: a report from the depression in primary care project. 2010 , 8, 387-96	18
	Long-term clinical outcomes of care management for chronically depressed primary care patients: a	18
324	Long-term clinical outcomes of care management for chronically depressed primary care patients: a report from the depression in primary care project. 2010 , 8, 387-96	
324	Long-term clinical outcomes of care management for chronically depressed primary care patients: a report from the depression in primary care project. 2010 , 8, 387-96 Personalized medicine for depression: can we match patients with treatments?. 2010 , 167, 1445-55 A case study of early experience with implementation of collaborative care in the Veterans Health	228
3 ² 4 3 ² 3 3 ² 2	Long-term clinical outcomes of care management for chronically depressed primary care patients: a report from the depression in primary care project. 2010 , 8, 387-96 Personalized medicine for depression: can we match patients with treatments?. 2010 , 167, 1445-55 A case study of early experience with implementation of collaborative care in the Veterans Health Administration. 2010 , 13, 331-7	228
3 ² 4 3 ² 3 3 ² 2 3 ² 1	Long-term clinical outcomes of care management for chronically depressed primary care patients: a report from the depression in primary care project. 2010 , 8, 387-96 Personalized medicine for depression: can we match patients with treatments?. 2010 , 167, 1445-55 A case study of early experience with implementation of collaborative care in the Veterans Health Administration. 2010 , 13, 331-7 The Mental Health Nurse Incentive Program: the benefits from a client perspective. 2010 , 31, 646-53	228 14 18
324 323 322 321 320	Long-term clinical outcomes of care management for chronically depressed primary care patients: a report from the depression in primary care project. 2010, 8, 387-96 Personalized medicine for depression: can we match patients with treatments?. 2010, 167, 1445-55 A case study of early experience with implementation of collaborative care in the Veterans Health Administration. 2010, 13, 331-7 The Mental Health Nurse Incentive Program: the benefits from a client perspective. 2010, 31, 646-53 Suicide victims' last contact with the primary care physician: report from Slovenia. 2010, 56, 280-7	228141812

316	Randomized trial of a computer-tailored intervention for patients with depression. 2011, 26, 77-89	29
315	Prevalencia del trastorno depresivo en pacientes que consultan un servicio de urgencias en Bogot□ 2011 , 40, 22-37	2
314	Effectiveness of guided self-help for depression and anxiety disorders in primary care: a pragmatic randomized controlled trial. 2011 , 187, 113-20	21
313	Improving outcomes for patients with depression by enhancing antidepressant therapy with non-pharmacological interventions: a systematic review of reviews. 2011 , 125, 357-67	29
312	Supporting the couple with conception challenges. 2011 , 36, 38-45	2
311	Beliefs and attitudes of French family practitioners toward depression: the impact of training in mental health. 2011 , 41, 107-22	11
310	New Zealand general practice nurses' roles in mental health care. 2011 , 58, 225-33	7
309	Primary care clinician adherence to guidelines for the management of chronic musculoskeletal pain: results from the study of the effectiveness of a collaborative approach to pain. 2011 , 12, 1490-501	29
308	Effectiveness of interventions to improve antidepressant medication adherence: a systematic review. 2011 , 65, 954-75	81
307	Outcomes for depression and anxiety in primary care and details of treatment: a naturalistic longitudinal study. 2011 , 11, 180	21
306	Optimal duration of combined psychotherapy and pharmacotherapy for patients with moderate and severe depression: a meta-analysis. 2011 , 131, 24-36	22
305	Positive impact of a family practice-based depression case management on patient's self-management. 2011 , 33, 23-8	12
304	OB CARESThe Obstetric Clinics and Resources Study: providers' perceptions of addressing perinatal depressiona qualitative study. 2011 , 33, 267-78	17
303	Zelfhulp voor depressie en angststoornissen. 2011 , 54, 488-490	
302	Depression and the onset of chronic illness in older adults: a 12-year prospective study. 2011 , 38, 373-82	32
301	Talking about depression: a qualitative study of barriers to managing depression in people with long term conditions in primary care. 2011 , 12, 10	106
300	A mobile phone application for the assessment and management of youth mental health problems in primary care: a randomised controlled trial. 2011 , 12, 131	105
299	Reducing depression in older home care clients: design of a prospective study of a nurse-led interprofessional mental health promotion intervention. 2011 , 11, 50	8

298	Implementing collaborative care for depression treatment in primary care: a cluster randomized evaluation of a quality improvement practice redesign. 2011 , 6, 121	53
297	Screening for distress and depression in cancer settings: 10 lessons from 40 years of primary-care research. 2011 , 20, 572-84	100
296	Implementing the NICE guideline for schizophrenia recommendations for psychological therapies: a qualitative analysis of the attitudes of CMHT staff. 2011 , 18, 48-59	67
295	Lay health worker led intervention for depressive and anxiety disorders in India: impact on clinical and disability outcomes over 12 months. 2011 , 199, 459-66	156
294	An illness perception model of primary care patients' help seeking for depression. 2011 , 21, 1495-507	18
293	Does integrated care improve treatment for depression? A systematic review. 2011 , 34, 113-25	42
292	New Jersey's efforts to improve postpartum depression care did not change treatment patterns for women on medicaid. 2011 , 30, 293-301	53
291	[Collaborative care for depression in general practice: overview of systematic reviews]. 2011 , 20, 91-8	
290	Perspective: Physician education: a promising strategy to prevent adolescent suicide. 2011 , 86, 342-7	31
289	Use of an Internet portal to improve community-based pediatric ADHD care: a cluster randomized trial. 2011 , 128, e1201-8	55
288	Patient-reported care coordination: associations with primary care continuity and specialty care use. 2011 , 9, 323-9	46
287	Psychiatry and primary care integration: challenges and opportunities. 2011 , 23, 28-30	5
286	Managing child and adolescent mental health problems in primary care: taking the leap from knowledge to practice. 2011 , 12, 301-9	12
285	Treatment of late-life mental disorders in primary care: we can do a better job. 2011 , 23, 274-85	9
284	A compilation of strategies for implementing clinical innovations in health and mental health. 2012 , 69, 123-57	482
283	Evaluating Depression Care Management in a Community Setting: Main Outcomes for a Medicaid HMO Population with Multiple Medical and Psychiatric Comorbidities. 2012 , 2012, 769298	3
282	A randomized trial to reduce the prevalence of depression and self-harm behavior in older primary care patients. 2012 , 10, 347-56	41
281	Depression Program in Primary Health Care. 2012 , 41, 38-47	15

280	Effect of routine mental health screening in a low-resource pediatric primary care population. 2012 , 51, 359-65	15
279	Collaborative care for depression and anxiety problems. 2012 , 10, CD006525	45 ¹
278	Recognising and responding to adolescent depression in general practice: developing and implementing the Therapeutic Identification of Depression in Young people (TIDY) programme. 2012 , 17, 482-94	14
277	Simplifying care for complex patients. 2012 , 10, 3-5	20
276	Addressing methodological challenges in implementing the nursing home pain management algorithm randomized controlled trial. 2012 , 9, 634-44	20
275	Stepped care: a method to deliver increased access to psychological therapies. 2012 , 57, 210-5	37
274	A qualitative evaluation of barriers to care for trauma-related mental health problems among low-income minorities in primary care. 2012 , 200, 438-43	9
273	Impact of a work-focused intervention on the productivity and symptoms of employees with depression. 2012 , 54, 128-35	41
272	Early dropout from psychotherapy for depression with group- and network-model therapists. 2012 , 39, 440-7	10
271	Rethinking the mental health treatment skills of primary care staff: a framework for training and research. 2012 , 39, 489-502	19
270	['Optimal care for depression': Freiburg model of integrated care for depressive disorders]. 2012 , 106, 625-30	2
269	Effectiveness of a multi-component programme for managing depression in primary care: a cluster randomized trial. The INDI project. 2012 , 142, 297-305	28
268	Collaborative Interventions for Circulation and Depression (COINCIDE): study protocol for a cluster randomized controlled trial of collaborative care for depression in people with diabetes and/or coronary heart disease. 2012 , 13, 139	22
267	[Views of patients diagnosed with depression and cared for by general practitioners and psychiatrists]. 2012 , 44, 595-602	O
266	Depression and diabetes: the role and impact of models of health care systems. 2012 , 142 Suppl, S80-8	12
265	Depression screening and mental health outcomes in children and adolescents: a systematic review protocol. 2012 , 1, 58	14
264	Therapeutic identification of depression in young people: lessons from the introduction of a new technique in general practice. <i>British Journal of General Practice</i> , 2012 , 62, e174-82	13
263	Collaborative care approaches for people with severe mental illness. 2012 ,	5

262	Does GP training in depression care affect patient outcome? - A systematic review and meta-analysis. 2012 , 12, 10	48
261	Yield of practice-based depression screening in VA primary care settings. 2012 , 27, 331-8	27
260	What is the role of general practice in addressing youth mental health? A discussion paper. 2012 , 181, 189-97	7
259	Barriers and facilitators to optimize function and prevent disability worsening: a content analysis of a nurse home visit intervention. 2012 , 68, 80-93	13
258	Use of the theory of planned behaviour to assess factors influencing the identification of individuals at ultra-high risk for psychosis in primary care. 2012 , 6, 265-75	5
257	The Swiss Early Psychosis Project SWEPP: a national network. 2012 , 6, 106-11	11
256	Implementing a knowledge application program for anxiety and depression in community-based primary mental health care: a multiple case study research protocol. 2013 , 8, 26	7
255	Comparison of high and low intensity contact between secondary and primary care to detect people at ultra-high risk for psychosis: study protocol for a theory-based, cluster randomized controlled trial. 2013 , 14, 222	9
254	Primary care clinicians' recognition and management of depression: a model of depression care in real-world primary care practice. 2013 , 28, 1430-9	6
253	Continuity of care: an Italian clinical experience. 2013 , 8, 595-9	4
253 252	Continuity of care: an Italian clinical experience. 2013 , 8, 595-9 Effectiveness of collaborative care for depression in Italy. A randomized controlled trial. 2013 , 35, 579-86	26
252	Effectiveness of collaborative care for depression in Italy. A randomized controlled trial. 2013 , 35, 579-86 Psychosis 101: evaluating a training programme for northern and remote youth mental health	26
252 251	Effectiveness of collaborative care for depression in Italy. A randomized controlled trial. 2013 , 35, 579-86 Psychosis 101: evaluating a training programme for northern and remote youth mental health service providers. 2013 , 7, 442-50 Increasing antiviral treatment through integrated hepatitis C care: a randomized multicenter trial.	26
252251250	Effectiveness of collaborative care for depression in Italy. A randomized controlled trial. 2013 , 35, 579-86 Psychosis 101: evaluating a training programme for northern and remote youth mental health service providers. 2013 , 7, 442-50 Increasing antiviral treatment through integrated hepatitis C care: a randomized multicenter trial. 2013 , 35, 97-107 Monitoring and managing metabolic effects of antipsychotics: a cluster randomized trial of an	26 2 26
252251250249	Effectiveness of collaborative care for depression in Italy. A randomized controlled trial. 2013, 35, 579-86 Psychosis 101: evaluating a training programme for northern and remote youth mental health service providers. 2013, 7, 442-50 Increasing antiviral treatment through integrated hepatitis C care: a randomized multicenter trial. 2013, 35, 97-107 Monitoring and managing metabolic effects of antipsychotics: a cluster randomized trial of an intervention combining evidence-based quality improvement and external facilitation. 2013, 8, 120 Better together? a naturalistic qualitative study of inter-professional working in collaborative care	26 2 26 13
252 251 250 249 248	Effectiveness of collaborative care for depression in Italy. A randomized controlled trial. 2013, 35, 579-86 Psychosis 101: evaluating a training programme for northern and remote youth mental health service providers. 2013, 7, 442-50 Increasing antiviral treatment through integrated hepatitis C care: a randomized multicenter trial. 2013, 35, 97-107 Monitoring and managing metabolic effects of antipsychotics: a cluster randomized trial of an intervention combining evidence-based quality improvement and external facilitation. 2013, 8, 120 Better together? a naturalistic qualitative study of inter-professional working in collaborative care for co-morbid depression and physical health problems. 2013, 8, 110 Depression care management for Chinese Americans in primary care: a feasibility pilot study. 2013,	26 2 26 13 51

(2014-2013)

244	Evaluation of a mental health training intervention for multidisciplinary teams in primary care in Brazil: a pre- and posttest study. 2013 , 35, 304-8	31
243	Screening for depression in patients with myocardial infarction by general practitioners. 2013 , 20, 800-6	15
242	Is treatment of depression cost-effective in people with diabetes? A systematic review of the economic evidence. 2013 , 29, 384-91	17
241	Depression screening and management among adolescents in primary care: factors associated with best practice. 2013 , 52, 557-67	16
240	Barriers to improving primary care of depression: perspectives of medical group leaders. 2013 , 23, 805-14	25
239	Common mental disorder severity and its association with treatment contact and treatment intensity for mental health problems. 2013 , 43, 2203-13	40
238	The effects of care management on depression treatment in a psychiatric clinic: a randomized controlled trial. 2013 , 28, 1023-30	2
237	Adolescent classroom education on knowledge and attitudes about deceased organ donation: a systematic review. 2013 , 17, 119-28	32
236	Practice-based versus telemedicine-based collaborative care for depression in rural federally qualified health centers: a pragmatic randomized comparative effectiveness trial. 2013 , 170, 414-25	193
235	Educational theories. 2013 , 298-307	1
235	Educational theories. 2013, 298-307 Depression in primary care: current and future challenges. 2013, 58, 442-8	97
234	Depression in primary care: current and future challenges. 2013 , 58, 442-8	
234	Depression in primary care: current and future challenges. 2013 , 58, 442-8 Behavioral Emergencies for the Emergency Physician. 53-59 Collaborative stepped care v. care as usual for common mental disorders: 8-month, cluster	97
234233232	Depression in primary care: current and future challenges. 2013, 58, 442-8 Behavioral Emergencies for the Emergency Physician. 53-59 Collaborative stepped care v. care as usual for common mental disorders: 8-month, cluster randomised controlled trial. 2013, 203, 132-9	97
234233232231	Depression in primary care: current and future challenges. 2013, 58, 442-8 Behavioral Emergencies for the Emergency Physician. 53-59 Collaborative stepped care v. care as usual for common mental disorders: 8-month, cluster randomised controlled trial. 2013, 203, 132-9 Educating family physicians to recognize and manage depression: where are we now?. 2013, 58, 449-55 Clinical implementation of a new antibiotic prophylaxis regimen for percutaneous endoscopic	97 39 12
234233232231230	Depression in primary care: current and future challenges. 2013, 58, 442-8 Behavioral Emergencies for the Emergency Physician. 53-59 Collaborative stepped care v. care as usual for common mental disorders: 8-month, cluster randomised controlled trial. 2013, 203, 132-9 Educating family physicians to recognize and manage depression: where are we now?. 2013, 58, 449-55 Clinical implementation of a new antibiotic prophylaxis regimen for percutaneous endoscopic gastrostomy. BMJ Open, 2013, 3, Perspective: Reconsidering the focus on "outcomes research" in medical education: a cautionary	97 39 12 4

226	Does primary care mental health resourcing affect the use and costs of secondary psychiatric services?. <i>International Journal of Environmental Research and Public Health</i> , 2014 , 11, 8743-54	4.6	6
225	Interpersonal Counseling (IPC) for Depression in Primary Care. 2014 , 68, 359-83		30
224	The challenges of human resources in low- and middle-income countries. 117-125		
223	Setting up integrated mental health systems. 144-151		
222	Collaborative treatment of late-life depression in primary care (GermanIMPACT): study protocol of a cluster-randomized controlled trial. 2014 , 15, 351		15
221	Routine depression screening in an MS clinic and its association with provider treatment recommendations and related treatment outcome. 2014 , 21, 347-55		1
220	12-month outcomes of community engagement versus technical assistance to implement depression collaborative care: a partnered, cluster, randomized, comparative effectiveness trial. 2014 , 161, S23-34		40
219	Including nurses in care models for older people with mild to moderate depression: an integrative review. 2014 , 23, 911-26		6
218	Strategies for guideline implementation in primary care focusing on patients with cardiovascular disease: a systematic review. 2014 , 31, 247-66		27
217	CASPER plus (CollAborative care in Screen-Positive EldeRs with major depressive disorder): study protocol for a randomised controlled trial. 2014 , 15, 451		12
216	Educational interventions for general practitioners to identify and manage depression as a suicide risk factor in young people: a systematic review and meta-analysis protocol. 2014 , 3, 145		11
215	Integration of depression and primary care: barriers to adoption. 2014 , 5, 67-73		18
214	The Affordable Care Act and integrated care. 2014 , 13, 44-86		25
213	Effective treatment for depression in patients with cancer. 2014 , 384, 1076-8		9
212	Do depressed and anxious men do groups? What works and what are the barriers to help seeking?. 2014 , 15, 287-301		12
211	Different answers to different questions: exploring clinical decision making by general practitioners and psychiatrists about depressed patients. 2014 , 36, 425-30		6
210	[An experience of collaboration between primary health care and mental health care in La Ribera Department of Health (Valencia, Spain)]. 2014 , 28, 405-7		1
209	Evaluation of GPs diagnostic knowledge and treatment practice in detection and treatment of early schizophrenia: a French postal survey in Brittany. 2014 , 49, 69-77		3

208	Prevention of depression in older age. 2014 , 79, 136-41		43
207	Development and evaluation of culturally sensitive psychosocial interventions for under-served people in primary care. 2014 , 14, 217		18
206	[Patient education of depression in primary care]. 2014 , 43, 883-5		
205	Listening to depression and suicide risk in medical students: the Healer Education Assessment and Referral (HEAR) Program. 2014 , 38, 547-53		49
204	An interprofessional nurse-led mental health promotion intervention for older home care clients with depressive symptoms. 2014 , 14, 62		31
203	Overview of evidence-based practice and translation science. 2014 , 49, 269-74		7
202	Collaborative care for depression in European countries: a systematic review and meta-analysis. 2014 , 77, 247-63		32
201	Participation in and impact of a depression care management program targeting low-income minority patients in an urban community-based clinic. 2014 , 36, 778-90		2
200	Study protocol for a randomized, controlled, superiority trial comparing the clinical and cost-effectiveness of integrated online mental health assessment-referral-care in pregnancy to usual prenatal care on prenatal and postnatal mental health and infant health and development: the		28
199	Integrated Maternal Psychosocial Assessment to Care Trial (IMPACT). 2014, 15, 72 Diabetes and depression in general practice: meeting the challenge of managing comorbidity. British Journal of General Practice, 2014, 64, 386-7	1.6	2
198	Tobacco use screening and treatment by outpatient psychiatrists before and after release of the American Psychiatric Association treatment guidelines for nicotine dependence. 2014 , 104, 90-5		32
197	Directions for future patient-centered and comparative effectiveness research for people with serious mental illness in a learning mental health care system. 2014 , 40 Suppl 1, S1-S94		19
196	Diagnoses of psychiatric disorders in hypothetical patients by non-psychiatric physicians in Japan. 2014 , 47, 65-74		2
195	Suicidal behaviour in mood disorderswho, when, and why?. 2014 , 59, 120-30		173
194	Achieving change in primary careeffectiveness of strategies for improving implementation of complex interventions: systematic review of reviews. <i>BMJ Open</i> , 2015 , 5, e009993	3	113
193	Study protocol for "Study of Practices Enabling Implementation and Adaptation in the Safety Net (SPREAD-NET)": a pragmatic trial comparing implementation strategies. 2015 , 10, 144		9
192	Comprehensive technology-assisted training and supervision program to enhance depression management in primary care in Santiago, Chile: study protocol for a cluster randomized controlled trial. 2015 , 16, 311		3
191	Consultation liaison in primary care for people with mental disorders. 2015 , CD007193		30

190	Prevalence, Comorbidity, and Prognosis of Mental Health Among US Veterans. 2015, 105, 2564-9	159
189	Continuum of care for persons with common mental health disorders in Nunavik: a descriptive study. 2015 , 74, 27186	10
188	Participation in Training for Depression Care Quality Improvement: A Randomized Trial of Community Engagement or Technical Support. 2015 , 66, 831-9	23
187	Mental and physical long-term conditions in the UK: spanning the boundaries. 2015 , 20, 190; 192-3	4
186	Promoting professional behaviour change in healthcare: what interventions work, and why? A theory-led overview of systematic reviews. <i>BMJ Open</i> , 2015 , 5, e008592	230
185	Clinical effectiveness and cost-effectiveness of tailored intensive liaison between primary and secondary care to identify individuals at risk of a first psychotic illness (the LEGs study): a cluster-randomised controlled trial. 2015 , 2, 984-93	35
184	COSMOSimproving the quality of life in nursing home patients: protocol for an effectiveness-implementation cluster randomized clinical hybrid trial. 2015 , 10, 131	38
183	Is there a case for mental health promotion in the primary care setting? A systematic review. 2015 , 76 Suppl, S5-11	7
182	Development and evaluation of an educational intervention in youth mental health for primary care practitioners. 2015 , 32, 137-146	3
181	Conceptualizations of postpartum depression by public-sector health care providers in Mexico. 2015 , 25, 551-68	6
180	A refined compilation of implementation strategies: results from the Expert Recommendations for Implementing Change (ERIC) project. 2015 , 10, 21	1234
179	Training Health-Care Professionals for the Assessment and Management of Perinatal Depression and Anxiety. 2015 , 210-227	1
178	Evaluation of the role of training in the implementation of a depression screening and treatment protocol in 2 academic outpatient internal medicine clinics utilizing the electronic medical record. 2015 , 30, 359-66	9
177	Evaluation of knowledge and attitudes among primary care physicians in Cavan-Monaghan as 'gatekeepers-in-waiting' for the introduction of Carepath for Overcoming Psychosis Early (COPE). 2015 , 9, 141-50	7
176	Chile: Acceptability of a Training Program for Depression Management in Primary Care. 2016 , 7, 853	1
175	The effect of knowledge and expectations on adherence to and persistence with antidepressants. 2016 , 10, 761-8	13
174	Identifying Maternal Depression in Pediatric Primary Care: Changes Over a Decade. 2016 , 37, 113-20	29
173	The Impact of Community Engagement on Health, Social, and Utilization Outcomes in Depressed, Impoverished Populations: Secondary Findings from a Randomized Trial. 2016 , 29, 325-38	17

172	The Rise of Primary Care Physicians in the Provision of US Mental Health Care. 2016, 41, 559-83	26
171	Evaluation of Mindfulness-Based Cognitive Therapy to Reduce Psychological Distress and to Promote Well-Being: A Pilot Study in a Primary Health Care Setting. 2016 , 6, 215824401666954	3
170	Collaborative care for anxiety disorders in primary care: a systematic review and meta-analysis. 2016 , 17, 62	36
169	Association of antidepressant treatment with emergency admission to medical units for patients 65 years or older. 2016 , 9, 210-218	
168	Irish General Practitioner referrals to psychological therapies. 2016 , 33, 73-80	2
167	Pain Management Algorithms for Implementing Best Practices in Nursing Homes: Results of a Randomized Controlled Trial. 2016 , 17, 348-56	11
166	Collaborative nurse-led self-management support for primary care patients with anxiety, depressive or somatic symptoms: Cluster-randomised controlled trial (findings of the SMADS study). 2016 , 63, 101-111	29
165	Improving the Capacity to Treat Depression using Talking Therapies. 2016 , 181-204	3
164	A qualitative study of perceived needs and factors associated with the quality of care for common mental disorders in patients with chronic diseases: the perspective of primary care clinicians and patients. 2016 , 17, 134	21
163	Organising healthcare services for persons with an intellectual disability. 2016 , 4, CD007492	22
162	Organisation of health services for preventing and treating pressure ulcers. 2016,	3
161	Receipt of pharmacotherapy and psychotherapy among a nationally representative US sample of privately insured adults with depression: associations with insurance plan arrangements and provider specialty. 2016 , 7, 53-62	2
160	The Relevance of the Affordable Care Act for Improving Mental Health Care. 2016, 12, 515-42	37
159	Treatment approaches of palliative medicine specialists for depression in the palliative care setting: findings from a qualitative, in-depth interview study. 2016 , 6, 186-93	2
158	Validation of the self-reporting questionnaire (SRQ 20) in British Pakistani and White European population in the United Kingdom. 2016 , 189, 392-6	5
157	Association of antidepressant treatment with emergency admission to medical units for patients 65 years or older. 2016 , 9, 210-218	3
156	Cross-Sectional and Longitudinal Association Between Trust in Physician and Depressive Symptoms Among U.S. Community-Dwelling Chinese Older Adults. 2017 , 72, S125-S130	10
155	Managing Depression among Diverse Older Adults in Primary Care: The BRIGHTEN Program. 2017 , 40, 88-96	3

154	Recognition of depression by primary care clinicians in rural Ethiopia. 2017, 18, 56		25
153	Integration of behavioral health and primary care: current knowledge and future directions. 2017 , 40, 69-84		44
152	Do measures of depressive symptoms function differently in people with spinal cord injury versus primary care patients: the CES-D, PHQ-9, and PROMIS-D. 2017 , 26, 139-148		17
151	Global Shifts in Cardiovascular Disease, the Epidemiologic Transition, and Other Contributing Factors: Toward a New Practice of Global Health Cardiology. 2017 , 35, 1-12		26
150	Random or predictable?: Adoption patterns of chronic care management practices in physician organizations. 2017 , 12, 106		5
149	German general practitioners' self-reported management of patients with chronic depression. 2017 , 17, 401		2
148	Embedding a Social Work-Led Behavioral Health Program in a Primary Care System: A 2012-2018 Case Study. 2017 , 23 Suppl 6 Suppl, Gulf Region , S40-S46		1
147	Guidelines for Adolescent Depression in Primary Care (GLAD-PC): Part I. Practice Preparation, Identification, Assessment, and Initial Management. 2018 , 141,		164
146	Die pdagogische Dimension in den Gesundheitsberufen. 2018 , 26, 30-32		
145	Interventions to improve screening and appropriate referral of patients with cancer for psychosocial distress: systematic review. <i>BMJ Open</i> , 2018 , 8, e017959	3	31
145 144		3	28
	psychosocial distress: systematic review. <i>BMJ Open</i> , 2018 , 8, e017959 Clinical effectiveness of care managers in collaborative care for patients with depression in	3	
144	psychosocial distress: systematic review. <i>BMJ Open</i> , 2018 , 8, e017959 Clinical effectiveness of care managers in collaborative care for patients with depression in Swedish primary health care: a pragmatic cluster randomized controlled trial. 2018 , 19, 28 Strategies to translate knowledge related to common musculoskeletal conditions into	3	28
144	Clinical effectiveness of care managers in collaborative care for patients with depression in Swedish primary health care: a pragmatic cluster randomized controlled trial. 2018, 19, 28 Strategies to translate knowledge related to common musculoskeletal conditions into physiotherapy practice: a systematic review. 2018, 104, 1-8 Associations Among Depressive Symptoms, Wellness, Patient Involvement, Provider Cultural Competency, and Treatment Nonadherence: A Pilot Study Among Community Patients Seen at a	3	28
144 143 142	Clinical effectiveness of care managers in collaborative care for patients with depression in Swedish primary health care: a pragmatic cluster randomized controlled trial. 2018, 19, 28 Strategies to translate knowledge related to common musculoskeletal conditions into physiotherapy practice: a systematic review. 2018, 104, 1-8 Associations Among Depressive Symptoms, Wellness, Patient Involvement, Provider Cultural Competency, and Treatment Nonadherence: A Pilot Study Among Community Patients Seen at a University Medical Center. 2018, 54, 138-148 Maintained effect of a training program on attitudes towards depression in family physicians. 2018,	3	28 23 2
144 143 142	Clinical effectiveness of care managers in collaborative care for patients with depression in Swedish primary health care: a pragmatic cluster randomized controlled trial. 2018, 19, 28 Strategies to translate knowledge related to common musculoskeletal conditions into physiotherapy practice: a systematic review. 2018, 104, 1-8 Associations Among Depressive Symptoms, Wellness, Patient Involvement, Provider Cultural Competency, and Treatment Nonadherence: A Pilot Study Among Community Patients Seen at a University Medical Center. 2018, 54, 138-148 Maintained effect of a training program on attitudes towards depression in family physicians. 2018, 35, 61-66 The rationale, design, and methods of a randomized, controlled trial to evaluate the effectiveness of collaborative telecare in preserving function among patients with late stage cancer and	3	28 23 2
144 143 142 141 140	Clinical effectiveness of care managers in collaborative care for patients with depression in Swedish primary health care: a pragmatic cluster randomized controlled trial. 2018, 19, 28 Strategies to translate knowledge related to common musculoskeletal conditions into physiotherapy practice: a systematic review. 2018, 104, 1-8 Associations Among Depressive Symptoms, Wellness, Patient Involvement, Provider Cultural Competency, and Treatment Nonadherence: A Pilot Study Among Community Patients Seen at a University Medical Center. 2018, 54, 138-148 Maintained effect of a training program on attitudes towards depression in family physicians. 2018, 35, 61-66 The rationale, design, and methods of a randomized, controlled trial to evaluate the effectiveness of collaborative telecare in preserving function among patients with late stage cancer and hematologic conditions. 2018, 64, 254-264 Development of a 2-h suicide prevention program for medical staff including nurses and medical	3	28 23 2 3

136 Teaching and Learning about Mental Health. 439-458

135	Undertaking Mental Health Research in Primary Care. 459-472	
134	What Is Primary Care Mental Health?. 1-11	1
133	Collaborative Care and Stepped Care. 401-419	
132	Management of treatment-resistant depression in primary care: a mixed-methods study. <i>British Journal of General Practice</i> , 2018 , 68, e673-e681	7
131	Protocol of an ongoing randomized controlled trial of care management for comorbid depression and hypertension: the Chinese Older Adult Collaborations in Health (COACH) study. 2018 , 18, 124	10
130	Cost-effectiveness of a care manager collaborative care programme for patients with depression in primary care: economic evaluation of a pragmatic randomised controlled study. <i>BMJ Open</i> , 2018 , 8, e024741	9
129	Elusive search for effective provider interventions: a systematic review of provider interventions to increase adherence to evidence-based treatment for depression. 2018 , 13, 99	9
128	Helping patients help themselves: A systematic review of self-management support strategies in primary health care practice. 2019 , 14, e0220116	94
127	Care managers can be useful for patients with depression but their role must be clear: a qualitative study of GPs' experiences. 2019 , 37, 273-282	7
126	Achieving universal health coverage for mental disorders. 2019 , 366, l4516	13
125	Collaborative care vs consultation liaison for depression and anxiety disorders in general practice: study protocol for two randomized controlled trials (the Danish Collabri Flex trials). 2019 , 20, 607	5
124	Lifelong Learning for Professional Development in Psychiatry: Pedagogy, Innovations, and Maintenance of Certification. 2019 , 42, 425-437	3
123	Comorbidity Profiles Identified in Older Primary Care Patients Who Attempt Suicide. 2019 , 67, 2553-2559	9
122	Registered nurses experiences of managing depressive symptoms at care centres for older people: a qualitative descriptive study. 2019 , 18, 43	1
121	Evaluation of the impacts of a district-level mental health care plan on contact coverage, detection and individual outcomes in rural Uganda: a mixed methods approach. 2019 , 13, 63	5
120	Is care really shared? A systematic review of collaborative care (shared care) interventions for adult cancer patients with depression. 2019 , 19, 120	12
119	Depression in primary care: part 2-management. 2019 , 365, l835	19

118	WHO-mhGAP Training in Mexico: Increasing Knowledge and Readiness for the Identification and Management of Depression and Suicide Risk in Primary Care. 2019 , 50, 558-566		9
117	Facilitators, barriers and ethical values related to the coordination of return-to-work among employees on sick leave due to common mental disorders: a protocol for a qualitative study (the CORE-project). <i>BMJ Open</i> , 2019 , 9, e032463	3	2
116	Pharmacy-based management for depression in adults. 2019 , 12, CD013299		10
115	Newly educated care managers' experiences of providing care for persons with stress-related mental disorders in the clinical primary care context. 2019 , 14, e0224929		3
114	Treatment adherence as a mediator of blood pressure control in Chinese older adults with depression. 2019 , 34, 432-438		5
113	Unmet social support needs among older adults with cancer. 2019 , 125, 473-481		31
112	Matter over mind: How mental health symptom presentations shape diagnostic outcomes. 2020 , 24, 755-772		2
111	Not seeing the forest for the trees: a systematic review of comprehensive distress management programs and implementation strategies. 2020 , 14, 220-231		6
110	Implementing primary healthcare-based measurement, advice and treatment for heavy drinking and comorbid depression at the municipal level in three Latin American countries: final protocol for a quasiexperimental study (SCALA study). <i>BMJ Open</i> , 2020 , 10, e038226	3	10
109	Psychological interventions for posttraumatic stress disorder involving primary care physicians: systematic review and Meta-analysis of randomized controlled trials. 2020 , 21, 176		
108	Efficacy of guided internet-assisted intervention on depression reduction among educational technology students of Nigerian universities. 2020 , 99, e18774		5
107	The Effect of Psychological First Aid Training on Knowledge and Understanding about Psychosocial Support Principles: A Cluster-Randomized Controlled Trial. <i>International Journal of Environmental Research and Public Health</i> , 2020 , 17,	4.6	30
106	Does early identification of high work related stress affect pharmacological treatment of primary care patients? - analysis of Swedish pharmacy dispensing data in a randomised control study. 2020 , 21, 70		2
105	Building a platform for translational research in chronic noncommunicable diseases to address population health: lessons from NHLBI supported CRONICAS in Peru. 2015 , 10, 13-9		2
104	Breast cancer endocrine therapy adherence in health professional shortage areas: Unique effects on patients with mental illness. 2021 , 140, 110294		O
103	Facilitators and Barriers to Implementing Standardized Outcome Measurement for Children With Cleft Lip and Palate. 2021 , 58, 7-18		2
102	Transforming Population-Based Depression Care: a Quality Improvement Initiative Using Remote, Centralized Care Management. 2021 , 36, 333-340		2
101	Exploring the Role of the Internet, Care Quality and Communication in Shaping Mental Health: An Analysis of the Health Information National Trends Survey. 2021 , PP,		O

100 Rural and Remote Psychiatry. 2021, 327-350

99	Effectiveness of therapeutic milieu intervention on inpatients with depressive disorder: A feasibility study from North India. 2021 , 57, 1604-1615		2
98	Association of Depression with Uncontrolled Hypertension in Primary Care Setting: A Cross-Sectional Study in Less-Developed Northwest China. 2021 , 2021, 6652228		1
97	Factors Differentiating Rural and Urban Population in Determining Anxiety and Depression in Patients with Chronic Cardiovascular Disease: A Pilot Study. <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18,	4.6	O
96	Implementation of a care manager organisation and its association with antidepressant medication patterns: a register-based study of primary care centres in Sweden. <i>BMJ Open</i> , 2021 , 11, e044959	3	1
95	Perceived need for treatment and non-utilization of outpatient psychotherapy in old age: two cohorts of a nationwide survey. 2021 , 21, 442		3
94	Association of Team-Based Care and Continuity of Care with Hospitalizations for Veterans with Comorbid Mental and Physical Health Conditions. 2021 , 1		O
93	Association between contact with a general practitioner and depressive symptoms during the COVID-19 pandemic and lockdown: a large community-based study in Hangzhou, China. <i>BMJ Open</i> , 2021 , 11, e052383	3	1
92	Cost-effectiveness of a care manager collaborative care programme for patients with depression in primary care: 12-month economic evaluation of a pragmatic randomised controlled trial. 2021 , 19, 52		O
91	Primary care-based screening and management of depression amongst heavy drinking patients: Interim secondary outcomes of a three-country quasi-experimental study in Latin America. 2021 , 16, e0255594		O
90	Prevalence and predictive nomogram of depression among hypertensive patients in primary care: A cross-sectional study in less developed Northwest China. 2021 , 100, e24422		
89	Organisation of health services for preventing and treating pressure ulcers. 2018 , 12, CD012132		4
88	Depression. 269-308		4
87	Behandlungspotenziale in der allgemeinEztlichen Versorgung. 2006 , 405-421		2
86	Depression and Work Performance: The Work and Health Initiative Study. 2011 , 103-120		8
85	How Behavioral Healthcare Informatics Systems Interface with Medical Informatics Systems: A Work in Progress. 2010 , 195-204		2
84	Promoting Treatment Engagement with Specific Populations: Depression. 2017, 151-170		O
83	Achieving Mental Health Equity: Collaborative Care. 2020 , 43, 501-510		4

82	A systematic review of elderly suicide prevention programs. 2011 , 32, 88-98		190
81	Treatment of depression and PTSD in primary care clinics serving uninsured low-income mostly Latina/o immigrants: A naturalistic prospective evaluation. 2019 , 25, 579-589		3
80	Mental health treatment dose and annual healthcare costs in patients with cancer and major depressive disorder. 2018 , 37, 1035-1040		12
79	Treatment of attempted suicide and suicidal patients in primary care. 2009 , 464-470		1
78	Counterpoint: chronic illness and primary care. <i>Medical Care</i> , 2011 , 49, 973-5	.1	7
77	Changing profiles of service sectors used for mental health care in the United States. 2006 , 163, 1187-98		148
76	Scaling-up primary health care-based prevention and management of alcohol use disorder at the municipal level in middle-income countries in Latin America: Background and pre-protocol for a three-country quasi-experimental study. <i>F1000Research</i> , 2017 , 6, 311	.6	8
75	How to decrease suicide rates in both genders? An effectiveness study of a community-based intervention (EAAD). 2013 , 8, e75081		38
74	Effectiveness of telephone-based aftercare case management for adult patients with unipolar depression compared to usual care: A randomized controlled trial. 2017 , 12, e0186967		5
73	Views of Young People in Rural Australia on SPARX, a Fantasy World Developed for New Zealand Youth With Depression. 2014 , 2, e3		30
72	Childcare service centers' preferences and intentions to use a web-based program to implement healthy eating and physical activity policies and practices: a cross-sectional study. 2015 , 17, e108		19
71	Comparative Effectiveness of a Technology-Facilitated Depression Care Management Model in Safety-Net Primary Care Patients With Type 2 Diabetes: 6-Month Outcomes of a Large Clinical Trial. 2018 , 20, e147		15
70	An integrated web-based mental health intervention of assessment-referral-care to reduce stress, anxiety, and depression in hospitalized pregnant women with medically high-risk pregnancies: a feasibility study protocol of hospital-based implementation. 2015 , 4, e9		24
69	The Feasibility and Acceptability of a Web-Based Alcohol Management Intervention in Community Sports Clubs: A Cross-Sectional Study. 2017 , 6, e123		1
68	Clinical effectiveness and cost-effectiveness of collaborative care for depression in UK primary care (CADET): a cluster randomised controlled trial. 2016 , 20, 1-192		30
67	Increasing equity of access to high-quality mental health services in primary care: a mixed-methods study. 2013 , 1, XXXX-XXXX		18
66	Qualitative study of depression management in primary care: GP and patient goals, and the value of listening. <i>British Journal of General Practice</i> , 2007 , 57, 872-9	.6	58
65	Comment amliorer la coopfation entre mflecins gfifalistes et psychiatres?. 2014 , 90, 359		2

64	A mixed-methods approach to investigating the adoption of evidence-based pain practices in nursing homes. 2014 , 40, 52-60		11
63	Pilot Study of Implementation of an Internet-Based Depression Prevention Intervention (CATCH-IT) for Adolescents in 12 US Primary Care Practices: Clinical and Management/Organizational Behavioral Perspectives. <i>primary care companion for CNS disorders, The</i> , 2013 , 15,	1.2	17
62	Clinical effectiveness of using an integrated model to treat depressive symptoms in veterans affairs primary care clinics and its impact on health care utilization. <i>primary care companion for CNS disorders, The</i> , 2011 , 13,	1.2	Ο
61	Collaboration Between Mental Health Professionals and Family Physicians: A Survey of New Jersey Family Physicians. 2005 , 7, 12-14		6
60	Randomized trial of the effect of research design and publication characteristics on physician change. 2006 , 8, 32-8		3
59	Mental health practice and attitudes of family physicians can be changed!. 2013, 17, 14-7		17
58	Outcomes of mental health pharmacist-managed electronic consults at a Veterans Affairs health care system. <i>Mental Health Clinician</i> , 2017 , 7, 131-136	1.6	7
57	Depression: We've Come a Long Way!. 2003 , 5, 151-152		
56	Prevention and Intervention in Primary Care. 2004 , 313-325		
55	Billing for the Evaluation and Treatment of Adult Depression by the Primary Care Clinician. 2004 , 6, 21-	26	1
54	Physical treatments. 2004 , 289-296		
53	Psychological treatment centres: the case against. 2006 , 332, 1154-5		1
52	Collaborative care to improve treatment of co-occurring diabetes mellitus and psychiatric disorders. 2007 , 219-238		
51	The Prediction of Occupational Disability Related to Depressive and Anxiety Disorders. 2008 , 371-384		1
5 0			
50	Early detection of mental ill-health, harmful stress and suicidal behaviours. 2009 , 544-550		
49	Early detection of mental ill-health, harmful stress and suicidal behaviours. 2009 , 544-550 Psychopharmacotherapy efficacy in patients with disorders of depressive spectrum depending on some peculiarities of prehospital period. <i>Bulletin of Siberian Medicine</i> , 2009 , 8, 81-84	0.4	
	Psychopharmacotherapy efficacy in patients with disorders of depressive spectrum depending on	0.4	

46	Depression in primary care: Chile's Programme for Screening, Diagnosis and Comprehensive Treatment of Depression. <i>International Psychiatry: Bulletin of the Board of International Affairs of the Royal College of Psychiatrists</i> , 2011 , 8, 6-7		2
45	Role of the general practitioner in the care of patients recently discharged from the hospital after a first psychotic episode: influence of length of stay. <i>primary care companion for CNS disorders, The</i> , 2011 , 13,	1.2	1
44	Depressive Disorders. 2013 , 749-756		
43	Mental Health Outreach at Community Behavior Health Centers. <i>Health</i> , 2014 , 06, 531-540	0.4	
42	SEMIOTICS, DIAGNOSTICS AND THERAPY TACTICS OF DEPRESSIVE DISORDERS IN CLINICAL MEDICINE. <i>Bulletin of Siberian Medicine</i> , 2014 , 13, 5-24	0.4	2
41	[Clinical/dynamic and epidemiological aspects of depressive disorders in the protracted emergency situation]. Zhurnal Nevrologii I Psikhiatrii Imeni S S Korsakova, 2015 , 115, 65-69	0.4	
40	Depression Management in the Medical Home. 2015 , 139-155		
39	Evaluation of a depression case management service: Improving depression care and impact on suicide assessment. <i>Mental Health Clinician</i> , 2015 , 5, 197-201	1.6	
38	Klinisches Management. 2017 , 69-80		1
37	Comparative Effectiveness of a Technology-Facilitated Depression Care Management Model in Safety-Net Primary Care Patients With Type 2 Diabetes: 6-Month Outcomes of a Large Clinical Trial (Preprint).		
36	Scaling-up primary health care-based prevention and management of alcohol use disorder at the municipal level in middle-income countries in Latin America: Background and pre-protocol for a three-country quasi-experimental study. <i>F1000Research</i> , 6, 311	3.6	2
35	Sozialpsychiatrie als Versorgungsforschung. 2018 , 87-110		
34	Scaling-up primary health care-based prevention and management of heavy drinking at the municipal level in middle-income countries in Latin America: Background and protocol for a three-country quasi-experimental study. <i>F1000Research</i> , 2017 , 6, 311	3.6	8
33	Assessment of the Antidepressant Side Effects Occurrence in Patients Treated in Primary Care. <i>Materia Socio-medica</i> , 2020 , 32, 131-134	0.9	1
32	Clinical effectiveness and cost effectiveness of individual mental health workers colocated within primary care practices: a systematic literature review. <i>BMJ Open</i> , 2020 , 10, e042052	3	
31	Evaluation of knowledge and competency among nurses after a brief suicide prevention educational program: A pilot study. <i>Journal of Family Medicine and Primary Care</i> , 2020 , 9, 6018-6022	1.5	1
30	Rural and Remote Psychiatry. 2020 , 1-24		
29	Theoretical Bases of Interventions to Improve Use of Medicines and Tests. 2020 , 29-57		

FrBerkennung und Awareness. 2006, 371-386 28 7 A Study Examining Practice Style Traits of Physical Therapists: Do They Explain Practice Behaviors 27 and Knowledge Translation?. Computerised patient-specific guidelines for management of common mental disorders in primary 26 1.6 19 care: a randomised controlled trial. British Journal of General Practice, 2004, 54, 832-7 Frequency of consultations and general practitioner recognition of psychological symptoms. British 1.6 25 Journal of General Practice, 2004, 54, 838-43 Self-help books for depression: how can practitioners and patients make the right choice?. British 1.6 67 24 Journal of General Practice, 2005, 55, 387-92 Interventions to improve management of anxiety disorders in general practice: a systematic review. 1.6 16 23 British Journal of General Practice, 2005, 55, 867-74 Designing CIS to improve decisions in depression disease management: a discourse analysis of front 22 0.7 line practice. **2006**, 564-8 Patients' experiences of receiving collaborative care for the treatment of depression in the UK: a 21 17 qualitative investigation. Mental Health in Family Medicine, 2008, 5, 95-104 Primary care, public health, and mental health. Preventing Chronic Disease, 2010, 7, A04 20 3.7 4 Depression in adults: psychological treatments and care pathways. Clinical Evidence, 2007, 2007, 19 A meta-analysis of interventions to improve care for chronic illnesses. American Journal of Managed 18 280 2.1 Care, 2005, 11, 478-88 Training teachers to teach mental health skills to staff in primary care settings in a vast, 17 under-populated area. Mental Health in Family Medicine, 2012, 9, 219-24 16 Geriatric depression assessment by rural primary care physicians. Rural and Remote Health, 2009, 9, 11801.3 5 Evaluation of a primary care adult mental health service: Year 2. Mental Health in Family Medicine, 15 4 2013, 10, 53-9 A Qualitative Study on Incentives and Disincentives for Care of Common Mental Disorders in 1.1 14 5 Ontario Family Health Teams. *Healthcare Policy*, **2016**, 12, 84-96 Depression in primary care: Chile's Programme for Screening, Diagnosis and Comprehensive Treatment of Depression. International Psychiatry: Bulletin of the Board of International Affairs of 13 the Royal College of Psychiatrists, 2011, 8, 6-7 A psychological intervention by community pharmacies to prevent depression in adults with subthreshold depression and long-term conditions: the CHEMIST pilot RCT. Public Health Research, 12 1.7 O 2022, 10, 1-160 Trauma-informed Collaborative Care for African American Primary Care Patients in Federally 11 3.1 Qualified Health Centers: A Pilot Randomized Trial.. Medical Care, 2022, 60, 232-239

Care management of depression: treatment of depression in primary care and the need for a multidisciplinary approach. 24-45

9	Anatomy of a review. 55-72	
8	Strategies to improve suicide prevention. 357-365	O
7	Consumer Perspectives on Anxiety Management in Australian General Practice <i>International Journal of Environmental Research and Public Health</i> , 2022 , 19,	4.6
6	Effectiveness of integrated care for older adults with depression and hypertension in rural China: a cluster randomized controlled trial.	
5	Effectiveness of integrated care for older adults with depression and hypertension in rural China: A cluster randomized controlled trial. 2022 , 19, e1004019	O
4	Response to commentaries on Improving access to psychological therapy: The Doncaster demonstration site organisational model 2008 , 1, 47-51	O
3	A theory-informed, rapid cycle approach to identifying and adapting strategies to promote sustainability: optimizing depression treatment in primary care clinics seeking to sustain collaborative care (The Transform DepCare Study). 2023 , 4,	O
2	Associations of Discrimination, Low Social Support, and Limited English Proficiency with Depression in South Asian Immigrants.	О
1	A Randomized Controlled Trial of the Effectiveness of Cognitive Behavior Therapy for Major Depression in Bangladesh.	O