

Assessment of estuarine water-quality indicators using Initial results from Tampa Bay, FL

Remote Sensing of Environment

93, 423-441

DOI: [10.1016/j.rse.2004.08.007](https://doi.org/10.1016/j.rse.2004.08.007)

Citation Report

#	ARTICLE	IF	CITATIONS
1	MODIS detects a devastating algal bloom in Paracas Bay, Peru. <i>Eos</i> , 2004, 85, 465-472.	0.1	74
2	Application of satellite data for integrated assessment of water quality in the Pearl River Estuary, China. , 0, , .		0
3	Using MODIS medium-resolution bands to monitor harmful algal blooms. , 2005, , .		3
4	Absorption and scattering properties of water body in Taihu Lake, China: absorption. <i>International Journal of Remote Sensing</i> , 2006, 27, 4277-4304.	1.3	68
5	Remote sensing of the water quality of shallow lakes: A mixture modelling approach to quantifying phytoplankton in water characterized by high suspended sediment. <i>International Journal of Remote Sensing</i> , 2006, 27, 1521-1537.	1.3	161
6	Mapping the concentrations of total suspended matter in Lake Taihu, China, using Landsat TM data. <i>International Journal of Remote Sensing</i> , 2006, 27, 1177-1191.	1.3	68
7	Light availability in the coastal ocean: impact on the distribution of benthic photosynthetic organisms and their contribution to primary production. <i>Biogeosciences</i> , 2006, 3, 489-513.	1.3	246
8	Operational monitoring of suspended matter distribution using MODIS images and numerical modelling. <i>Advances in Space Research</i> , 2006, 38, 2182-2188.	1.2	52
9	Quantitative modeling of suspended sediment in middle Changjiang River from modis. <i>Chinese Geographical Science</i> , 2006, 16, 79-82.	1.2	41
10	Developing a Monitoring Support Model for Chlorophyll a Concentration in the James River Estuary Using MODIS Images. , 2007, , .		1
11	Ocean color remote sensing of turbid plumes in the southern California coastal waters during storm events. , 2007, , .		0
12	On the absorption of light in the Orinoco River plume. <i>Continental Shelf Research</i> , 2007, 27, 1447-1464.	0.9	28
13	Monitoring variation of water turbidity and related environmental factors in Poyang Lake National Nature Reserve, China. , 2007, , .		5
14	Colored dissolved organic matter in Tampa Bay, Florida. <i>Marine Chemistry</i> , 2007, 104, 98-109.	0.9	104
15	Extending the MODIS 1 km ocean colour atmospheric correction to the MODIS 500 m bands and 500 m chlorophyll-a estimation towards coastal and estuarine monitoring. <i>Remote Sensing of Environment</i> , 2007, 107, 521-532.	4.6	24
16	Remotely sensed assessment of water quality levels in the Pearl River Estuary, China. <i>Marine Pollution Bulletin</i> , 2007, 54, 1267-1272.	2.3	56
17	Satellite-based monitoring of tropical seagrass vegetation: current techniques and future developments. <i>Hydrobiologia</i> , 2007, 591, 59-71.	1.0	33
18	Baroclinic tidal flows and inundation processes in Cook Inlet, Alaska: numerical modeling and satellite observations. <i>Ocean Dynamics</i> , 2007, 57, 205-221.	0.9	43

#	ARTICLE	IF	CITATIONS
19	Monitoring turbidity in Tampa Bay using MODIS/Aqua 250-m imagery. Remote Sensing of Environment, 2007, 109, 207-220.	4.6	252
20	Determination of Environmental Quality of a Drinking Water Reservoir by Remote Sensing, GIS and Regression Analysis. Water, Air, and Soil Pollution, 2008, 194, 275-285.	1.1	28
21	A GIS-based WFD oriented typology of shallow micro-tidal soft bottom using wave exposure and turbidity mapping. Estuarine, Coastal and Shelf Science, 2008, 78, 27-37.	0.9	10
22	Development and application of a remote sensing-based salinity prediction model for a large estuarine lake in the US Gulf of Mexico coast. Journal of Hydrology, 2008, 360, 184-194.	2.3	45
23	Origin and Offshore Extent of Floating Algae in Olympic Sailing Area. Eos, 2008, 89, 302-303.	0.1	121
24	Ocean Color Reveals Sand Ridge Morphology on the West Florida Shelf. IEEE Geoscience and Remote Sensing Letters, 2008, 5, 443-447.	1.4	7
25	Comparison of Turbidity Measurement by MODIS and AVHRR Images. , 2008, , .		2
26	Remote sensing data assimilation in environmental models. , 2008, , .		1
27	Quantitative analysis of suspended sediment in the Yangtze River estuary from MODIS images. , 2008, , .		1
28	Study on the atmospheric correction for high spatial resolution bands of MODIS. , 2008, , .		2
29	Study on the quantitative model of suspended sediment from MODIS in the Yangtze River. , 2008, , .		0
30	Optical Algorithms at Satellite Wavelengths for Total Suspended Matter in Tropical Coastal Waters. Sensors, 2008, 8, 4165-4185.	2.1	102
31	Water Quality Determination of KÄ¼ÄÄ¼ÄSekmece Lake, Turkey by Using Multispectral Satellite Data. Scientific World Journal, The, 2009, 9, 1215-1229.	0.8	26
32	Exploring the potential of MODIS visible and thermal channels in monitoring and assessing the impact of desalination plant discharges in the Arabian Gulf. , 2009, , .		0
33	Use of Reflectance Ratios as a Proxy for Coastal Water Constituent Monitoring in the Pearl River Estuary. Sensors, 2009, 9, 656-673.	2.1	9
34	Measurement of water colour using AVIRIS imagery to assess the potential for an operational monitoring capability in the Pamlico Sound Estuary, USA. International Journal of Remote Sensing, 2009, 30, 3291-3314.	1.3	30
35	Conditioning of reflectance signals by linear diffusion for improving narrow-band ratio-based remote-sensing bottom depth retrieval in shallow coastal waters. Journal of Applied Remote Sensing, 2009, 3, 033539.	0.6	0
36	Linear algorithm for salinity distribution modelling from MODIS data. , 2009, , .		6

#	ARTICLE	IF	CITATIONS
37	Monitoring the impact of backflow and dredging on water clarity using MODIS images of Poyang Lake, China. <i>Hydrological Processes</i> , 2009, 23, 342-350.	1.1	27
38	A novel ocean color index to detect floating algae in the global oceans. <i>Remote Sensing of Environment</i> , 2009, 113, 2118-2129.	4.6	683
39	Environmental modelling of Omerli catchment area in Istanbul, Turkey using remote sensing and GIS techniques. <i>Environmental Monitoring and Assessment</i> , 2009, 153, 323-332.	1.3	12
40	Application of MODIS data in monitoring suspended sediment of Taihu Lake, China. <i>Chinese Journal of Oceanology and Limnology</i> , 2009, 27, 614-620.	0.7	27
41	Validation of satellite ocean color primary products at optically complex coastal sites: Northern Adriatic Sea, Northern Baltic Proper and Gulf of Finland. <i>Remote Sensing of Environment</i> , 2009, 113, 2574-2591.	4.6	154
42	Remote sensing assessment of sediment re-suspension during Hurricane Frances in Apalachicola Bay, USA. <i>Remote Sensing of Environment</i> , 2009, 113, 2670-2681.	4.6	58
43	Characterization of mesoscale spatio-temporal patterns and variability of remotely sensed Chl <i>a</i> and SST in the Interior Sea of Chiloe (41.4°-43.5°S). <i>International Journal of Remote Sensing</i> , 2009, 30, 1521-1536.	1.3	10
44	Error decomposition and estimation of inherent optical properties. <i>Applied Optics</i> , 2009, 48, 4947.	2.1	34
45	Automatic Registration of High-Resolution Multispectral Imageries from Band-Reconfigurable Imaging System on Board Unmanned Airship. , 2009, , .		2
46	One Cost-Effective High-Resolution Multispectral Imaging System for Quick-Response to Time-Critical Applications. , 2009, , .		0
47	Least square algorithm for sea surface salinity retrieving from MODIS satellite data. , 2009, , .		1
48	Light scattering by particles suspended in the sea: The role of particle size and density. <i>Continental Shelf Research</i> , 2009, 29, 1748-1755.	0.9	76
49	Sediment plume response to surface melting and supraglacial lake drainages on the Greenland ice sheet. <i>Journal of Glaciology</i> , 2009, 55, 1072-1082.	1.1	58
50	Assessment of water quality in Sheyang Estuary (China) using hyperspectral data. <i>Proceedings of SPIE</i> , 2009, , .	0.8	1
51	Ocean color response to an episode of heavy rainfall in the lagoon of New Caledonia. , 2009, , .		7
52	Retrieval of total suspended matter concentration in the Yellow and East China Seas from MODIS imagery. <i>Remote Sensing of Environment</i> , 2010, 114, 392-403.	4.6	211
53	Satellite Estimates of Wide-Range Suspended Sediment Concentrations in Changjiang (Yangtze) Estuary Using MERIS Data. <i>Estuaries and Coasts</i> , 2010, 33, 1420-1429.	1.0	166
54	Inherent optical properties and satellite retrieval of chlorophyll concentration in the lagoon and open ocean waters of New Caledonia. <i>Marine Pollution Bulletin</i> , 2010, 61, 503-518.	2.3	57

#	ARTICLE	IF	CITATIONS
55	Short-term variability of suspended sediment and phytoplankton in Tampa Bay, Florida: Observations from a coastal oceanographic tower and ocean color satellites. <i>Estuarine, Coastal and Shelf Science</i> , 2010, 89, 62-72.	0.9	61
56	MODIS imagery of turbid plumes in San Diego coastal waters during rainstorm events. <i>Remote Sensing of Environment</i> , 2010, 114, 332-344.	4.6	58
57	Calibration and validation of a generic multisensor algorithm for mapping of total suspended matter in turbid waters. <i>Remote Sensing of Environment</i> , 2010, 114, 854-866.	4.6	533
58	Development of the Metropolitan Water Availability Index (MWAI) and short-term assessment with multi-scale remote sensing technologies. <i>Journal of Environmental Management</i> , 2010, 91, 1397-1413.	3.8	9
59	Time-series MODIS Image-based Retrieval and Distribution Analysis of Total Suspended Matter Concentrations in Lake Taihu (China). <i>International Journal of Environmental Research and Public Health</i> , 2010, 7, 3545-3560.	1.2	35
60	An Investigation on Water Quality of Darlik Dam Drinking Water using Satellite Images. <i>Scientific World Journal</i> , The, 2010, 10, 1293-1306.	0.8	8
61	Quantitative monitoring of inland water using remote sensing of the upper reaches of the Huangpu River, China. <i>International Journal of Remote Sensing</i> , 2010, 31, 2471-2492.	1.3	6
62	Full Issue in PDF / Numéro complet en PDF. <i>Canadian Journal of Remote Sensing</i> , 2010, 36, 437-630.	1.1	8
63	Remote sensing of chlorophyll- <i>a</i> concentrations of the Pearl River Estuary from MODIS land bands. <i>International Journal of Remote Sensing</i> , 2010, 31, 4625-4633.	1.3	16
64	AERONET-OC: an overview. <i>Canadian Journal of Remote Sensing</i> , 2010, 36, 488-497.	1.1	32
65	Moderate Resolution Imaging Spectroradiometer (MODIS) observations of cyanobacteria blooms in Taihu Lake, China. <i>Journal of Geophysical Research</i> , 2010, 115, .	3.3	280
66	Remote-sensing reflectance characteristics of highly turbid estuarine waters – a comparative experiment of the Yangtze River and the Yellow River. <i>International Journal of Remote Sensing</i> , 2010, 31, 2639-2654.	1.3	109
67	Estimating turbidity and total suspended matter in the Adour River plume (South Bay of Biscay) using MODIS 250-m imagery. <i>Continental Shelf Research</i> , 2010, 30, 379-392.	0.9	204
68	Modelling Sea Surface Salinity from MODIS Satellite Data. <i>Lecture Notes in Computer Science</i> , 2010, , 545-556.	1.0	12
69	Dynamics of the turbidity plume in the Guadalquivir estuary (SW Spain): A remote sensing approach. , 2011, , .		3
70	A study of sediment transport in a shallow estuary using MODIS imagery and particle tracking simulation. <i>International Journal of Remote Sensing</i> , 2011, 32, 6653-6671.	1.3	27
71	Remote sensing analysis of rainstorm effects on sediment concentrations in Apalachicola Bay, USA. <i>Ecological Informatics</i> , 2011, 6, 147-155.	2.3	27
72	Estimation of cyanobacterial pigments in a freshwater lake using OCM satellite data. <i>Remote Sensing of Environment</i> , 2011, 115, 3409-3423.	4.6	43

#	ARTICLE	IF	CITATIONS
73	Relationships between suspended mineral concentrations and red-waveband reflectances in moderately turbid shelf seas. <i>Remote Sensing of Environment</i> , 2011, 115, 3719-3730.	4.6	35
74	A Multi-Sensor Approach to Examining the Distribution of Total Suspended Matter (TSM) in the Albemarle-Pamlico Estuarine System, NC, USA. <i>Remote Sensing</i> , 2011, 3, 962-974.	1.8	38
75	An enhanced MODIS remote sensing model for detecting rainfall effects on sediment plume in the coastal waters of Apalachicola Bay. <i>Marine Environmental Research</i> , 2011, 72, 265-272.	1.1	35
76	Temporal and spatial variability of chlorophyll a concentration in Lake Taihu using MODIS time-series data. <i>Hydrobiologia</i> , 2011, 661, 235-250.	1.0	48
77	Integrating satellite imagery and geostatistics of point samples for monitoring spatio-temporal changes of total suspended solids in bay waters: application to Tien Yen Bay (Northern Vietnam). <i>Frontiers of Earth Science</i> , 2011, 5, 305.	0.9	6
78	A simple method for mapping bathymetry over turbid coastal waters from MODIS data: possibilities and limitations. <i>International Journal of Remote Sensing</i> , 2011, 32, 7575-7590.	1.3	6
79	Retrieving seasonal sea surface salinity from MODIS satellite data using a Box-Jenkins algorithm. , 2011, , .		7
80	Development of clarity model for Caspian Sea using MERIS data. <i>Proceedings of SPIE</i> , 2011, , .	0.8	0
81	Impact of sub-pixel variations on ocean color remote sensing products. <i>Optics Express</i> , 2012, 20, 20844.	1.7	34
82	Estimating Secchi depth by simulated HSI of HJ-1A from in situ hyperspectral data: A case study in Sishili Bay, China. , 2012, , .		1
84	Tracing offshore low-salinity plumes in the Northeastern Gulf of Mexico during the summer season by use of multispectral remote-sensing data. <i>Journal of Oceanography</i> , 2012, 68, 743-760.	0.7	16
85	Water Quality Monitoring Using Remote Sensing and an Artificial Neural Network. <i>Water, Air, and Soil Pollution</i> , 2012, 223, 4875-4887.	1.1	96
86	Characterization of turbidity in Florida's Lake Okeechobee and Caloosahatchee and St. Lucie Estuaries using MODIS-Aqua measurements. <i>Water Research</i> , 2012, 46, 5410-5422.	5.3	46
87	Quantifying and Mapping Ecosystem Services Supplies and Demands: A Review of Remote Sensing Applications. <i>Environmental Science & Technology</i> , 2012, 46, 8529-8541.	4.6	112
88	The development of a new optical total suspended matter algorithm for the Chesapeake Bay. <i>Remote Sensing of Environment</i> , 2012, 119, 243-254.	4.6	93
89	Phytoplankton bloom during the northeast monsoon in the Luzon Strait bordering the Kuroshio. <i>Remote Sensing of Environment</i> , 2012, 124, 38-48.	4.6	33
90	Human induced turbidity changes in Poyang Lake between 2000 and 2010: Observations from MODIS. <i>Journal of Geophysical Research</i> , 2012, 117, .	3.3	116
91	Current Advances in Uncertainty Estimation of Earth Observation Products of Water Quality. , 2012, , .		0

#	ARTICLE	IF	CITATIONS
92	Atmospheric Correction and Vicarious Calibration of Oceansat-1 Ocean Color Monitor (OCM) Data in Coastal Case 2 Waters. <i>Remote Sensing</i> , 2012, 4, 1716-1740.	1.8	24
93	Comparative Analysis of Four Models to Estimate Chlorophyll-a Concentration in Case-2 Waters Using MODerate Resolution Imaging Spectroradiometer (MODIS) Imagery. <i>Remote Sensing</i> , 2012, 4, 2373-2400.	1.8	48
94	Assessment of total suspended sediment concentrations in Poyang Lake using HJ-1A/1B CCD imagery. <i>Chinese Journal of Oceanology and Limnology</i> , 2012, 30, 295-304.	0.7	29
95	Use of EO-1 Advanced Land Imager (ALI) multispectral image data and real-time field sampling for water quality mapping in the Hirfanlı Dam Lake, Turkey. <i>Environmental Science and Pollution Research</i> , 2013, 20, 5416-5424.	2.7	2
96	Optical variability along a river plume gradient: Implications for management and remote sensing. <i>Estuarine, Coastal and Shelf Science</i> , 2013, 131, 149-161.	0.9	15
97	Inherent and apparent optical properties of the complex estuarine waters of Tampa Bay: What controls light?. <i>Estuarine, Coastal and Shelf Science</i> , 2013, 117, 54-69.	0.9	41
98	Using geostationary satellite ocean color data to map the diurnal dynamics of suspended particulate matter in coastal waters. <i>Remote Sensing of Environment</i> , 2013, 133, 225-239.	4.6	210
99	Comparison of MODIS-based models for retrieving suspended particulate matter concentrations in Poyang Lake, China. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2013, 24, 63-72.	1.4	39
100	Monitoring estuarine water quality using satellite imagery. The Mondego river estuary (Portugal) as a case study. <i>Ocean and Coastal Management</i> , 2013, 72, 13-21.	2.0	13
101	Satellite observations of turbidity in the Dead Sea. <i>Journal of Geophysical Research: Oceans</i> , 2013, 118, 3146-3160.	1.0	30
102	Phytoplankton composition indicators for the assessment of eutrophication in marine waters: Present state and challenges within the European directives. <i>Marine Pollution Bulletin</i> , 2013, 66, 7-16.	2.3	85
103	Evaluation of chlorophyll-a remote sensing algorithms for an optically complex estuary. <i>Remote Sensing of Environment</i> , 2013, 129, 75-89.	4.6	152
104	Analyzing the effects of particle size on remotely sensed spectra: a study on optical properties and spectral similarity scale of suspended particulate matters in water. <i>Applied Optics</i> , 2013, 52, 879.	0.9	9
105	A hybrid approach to estimate chromophoric dissolved organic matter in turbid estuaries from satellite measurements: A case study for Tampa Bay. <i>Optics Express</i> , 2013, 21, 18849.	1.7	27
106	Atmospheric correction and inherent optical property estimation in the southwest New Caledonia lagoon using AVNIR-2 high-resolution data. <i>Applied Optics</i> , 2013, 52, 182.	0.9	21
107	Long-term statistical assessment of Aqua-MODIS aerosol optical depth over coastal regions: bias characteristics and uncertainty sources. <i>Tellus, Series B: Chemical and Physical Meteorology</i> , 2022, 65, 20805.	0.8	38
108	Modeling Methods for Identifying Critical Source Areas of Bacteria: Recent Developments and Future Perspectives. <i>Water Environment Research</i> , 2013, 85, 259-269.	1.3	2
109	The Application of GOCI Data Based Remote Sensing Technology in the Field of Sediment Reserch. <i>Applied Mechanics and Materials</i> , 2013, 303-306, 786-789.	0.2	0

#	ARTICLE	IF	CITATIONS
110	Evaluation of spatio-temporal variations in chlorophyll- <i>a</i> in Lake Naivasha, Kenya: remote-sensing approach. <i>International Journal of Remote Sensing</i> , 2013, 34, 8142-8155.	1.3	17
111	Regional algorithms for remote-sensing estimates of total suspended matter in the Beaufort Sea. <i>International Journal of Remote Sensing</i> , 2013, 34, 6562-6576.	1.3	21
112	Using remotely sensed suspended sediment concentration variation to improve management of Poyang Lake, China. <i>Lake and Reservoir Management</i> , 2013, 29, 47-60.	0.4	51
113	Ecosystem modelling for ecosystem-based management of bivalve aquaculture sites in data-poor environments. <i>Aquaculture Environment Interactions</i> , 2013, 4, 117-133.	0.7	26
114	Assessment of MERIS ocean color data products for European seas. <i>Ocean Science</i> , 2013, 9, 521-533.	1.3	40
115	Study of the variation of schistosomiasis risk in Lake Poyang in the People's Republic of China using multiple space-borne sensors for monitoring and modelling. <i>Geospatial Health</i> , 2014, 8, 353.	0.3	5
116	Assessment of Total Suspended Sediment Distribution under Varying Tidal Conditions in Deep Bay: Initial Results from HJ-1A/1B Satellite CCD Images. <i>Remote Sensing</i> , 2014, 6, 9911-9929.	1.8	24
117	Improving Sediment Transport Prediction by Assimilating Satellite Images in a Tidal Bay Model of Hong Kong. <i>Water (Switzerland)</i> , 2014, 6, 642-660.	1.2	14
118	Simulation of Tsunami Impact on Sea Surface Salinity along Banda Aceh Coastal Waters, Indonesia. , 2014, , .		3
119	Variability of particulate organic carbon in inland waters observed from MODIS Aqua imagery. <i>Environmental Research Letters</i> , 2014, 9, 084011.	2.2	56
120	Atmospheric correction of HJ-1 CCD imagery over turbid lake waters. <i>Optics Express</i> , 2014, 22, 7906.	1.7	17
121	Developing MODIS-based retrieval models of suspended particulate matter concentration in Dongting Lake, China. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2014, 32, 46-53.	1.4	27
122	Modeling suspended sediment distribution patterns of the Amazon River using MODIS data. <i>Remote Sensing of Environment</i> , 2014, 147, 232-242.	4.6	117
123	Assessment of suspended solids in the Guadalquivir estuary using new DEIMOS-1 medium spatial resolution imagery. <i>Remote Sensing of Environment</i> , 2014, 146, 148-158.	4.6	35
124	Monitoring spatio-temporal variability of the Adour River turbid plume (Bay of Biscay, France) with MODIS 250-m imagery. <i>Continental Shelf Research</i> , 2014, 74, 35-49.	0.9	64
125	Influence of the Three Gorges Dam on total suspended matters in the Yangtze Estuary and its adjacent coastal waters: Observations from MODIS. <i>Remote Sensing of Environment</i> , 2014, 140, 779-788.	4.6	151
126	A Validation Study of an Improved SWIR Iterative Atmospheric Correction Algorithm for MODIS-Aqua Measurements in Lake Taihu, China. <i>IEEE Transactions on Geoscience and Remote Sensing</i> , 2014, 52, 4686-4695.	2.7	23
127	Influence of a red band-based water classification approach on chlorophyll algorithms for optically complex estuaries. <i>Remote Sensing of Environment</i> , 2014, 155, 289-302.	4.6	39

#	ARTICLE	IF	CITATIONS
128	Remote sensing of selected water-quality indicators with the hyperspectral imager for the coastal ocean (HICO) sensor. <i>International Journal of Remote Sensing</i> , 2014, 35, 2927-2962.	1.3	64
129	Performance evaluation of normalized difference chlorophyll index in northern Gulf of Mexico estuaries using the Hyperspectral Imager for the Coastal Ocean. <i>GIScience and Remote Sensing</i> , 2014, 51, 175-198.	2.4	29
130	Characterization of spatial and temporal variation of suspended sediments in the Yellow and East China Seas using satellite ocean color data. <i>GIScience and Remote Sensing</i> , 2014, 51, 212-226.	2.4	34
131	A Novel Algorithm to Estimate Algal Bloom Coverage to Subpixel Resolution in Lake Taihu. <i>IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing</i> , 2014, 7, 3060-3068.	2.3	54
132	Mapping bottom depth and albedo in coastal waters of the South China Sea islands and reefs using Landsat TM and ETM+ data. <i>International Journal of Remote Sensing</i> , 2014, 35, 4156-4172.	1.3	21
133	Harbour dredging and fish mortality in an aquaculture zone: assessment of changes in suspended particulate matter using multi-sensor remote-sensing data. <i>International Journal of Remote Sensing</i> , 2014, 35, 4383-4398.	1.3	6
134	Challenges and opportunities for geostationary ocean colour remote sensing of regional seas: A review of recent results. <i>Remote Sensing of Environment</i> , 2014, 146, 63-76.	4.6	69
135	Evaluating suitability of MODIS-Terra images for reproducing historic sediment concentrations in water bodies: Lake Tana, Ethiopia. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2014, 26, 286-297.	1.4	44
136	Assimilation of MODIS Chlorophyll-a Data Into a Coupled Hydrodynamic-Biological Model of Taihu Lake. <i>IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing</i> , 2014, 7, 1623-1631.	2.3	11
137	An EOF-Based Algorithm to Estimate Chlorophyll a Concentrations in Taihu Lake from MODIS Land-Band Measurements: Implications for Near Real-Time Applications and Forecasting Models. <i>Remote Sensing</i> , 2014, 6, 10694-10715.	1.8	59
138	An Adaptive Model to Monitor Chlorophyll-a in Inland Waters in Southern Quebec Using Downscaled MODIS Imagery. <i>Remote Sensing</i> , 2014, 6, 6446-6471.	1.8	16
139	Simulation of tsunami effects on sea surface salinity using MODIS satellite data. <i>IOP Conference Series: Earth and Environmental Science</i> , 2014, 18, 012004.	0.2	0
140	WACODI: A generic algorithm to derive the intrinsic color of natural waters from digital images. <i>Limnology and Oceanography: Methods</i> , 2015, 13, 697-711.	1.0	29
141	Remote Sensing of Suspended Sediment Over Gulf of Martaban. <i>Ekologia</i> , 2015, 34, .	0.2	2
142	Observations of hysteresis in the annual exchange circulation of a large microtidal estuary. <i>Journal of Geophysical Research: Oceans</i> , 2015, 120, 2904-2919.	1.0	4
143	Estuarine turbidity maxima revisited. <i>Developments in Sedimentology</i> , 2015, 68, 49-109.	0.5	18
144	Fourteen-Year Record (2000–2013) of the Spatial and Temporal Dynamics of Floating Algae Blooms in Lake Chaohu, Observed from Time Series of MODIS Images. <i>Remote Sensing</i> , 2015, 7, 10523-10542.	1.8	99
145	Monitoring for Spatiotemporal Estuarine Chlorophyll Using MODIS and In Situ Characteristics. <i>Journal of Environmental Engineering, ASCE</i> , 2015, 141, 04015003.	0.7	0

#	ARTICLE	IF	CITATIONS
146	Remote Sensing of Spatiotemporal Variation of Apparent Optical Properties in Bohai Sea. IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, 2015, 8, 1176-1184.	2.3	15
147	MODIS Observations of Human-Induced Changes in the Mesopotamian Marshes in Iraq. Wetlands, 2015, 35, 31-40.	0.7	23
148	Using remote sensing to monitor the influence of river discharge on watershed outlets and adjacent coral Reefs: Magdalena River and Rosario Islands, Colombia. International Journal of Applied Earth Observation and Geoinformation, 2015, 38, 204-215.	1.4	23
149	Towards a practical remote-sensing model of suspended sediment concentrations in turbid waters using MERIS measurements. International Journal of Remote Sensing, 2015, 36, 3875-3889.	1.3	11
150	Long-term remote monitoring of total suspended matter concentration in Lake Taihu using 250m MODIS-Aqua data. Remote Sensing of Environment, 2015, 164, 43-56.	4.6	197
151	Northern Gulf of Mexico estuarine coloured dissolved organic matter derived from MODIS data. International Journal of Remote Sensing, 2015, 36, 2219-2237.	1.3	13
152	Evaluation of stream water quality data generated from MODIS images in modeling total suspended solid emission to a freshwater lake. Science of the Total Environment, 2015, 523, 170-177.	3.9	29
153	A model for partitioning the light absorption coefficient of natural waters into phytoplankton, nonalgal particulate, and colored dissolved organic components: A case study for the Chesapeake Bay. Journal of Geophysical Research: Oceans, 2015, 120, 2601-2621.	1.0	34
154	Enhanced satellite remote sensing of coastal waters using spatially improved bio-optical products from SNPP-VIIRS. Remote Sensing of Environment, 2015, 165, 53-63.	4.6	17
155	Study on the seasonal migration of surface suspended sediment in the Taiwan Strait based on remote sensing. , 2015, , .		0
156	Smart Information Reconstruction via Time-Space-Spectrum Continuum for Cloud Removal in Satellite Images. IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, 2015, 8, 1898-1912.	2.3	27
157	Abundance, distribution, and fluxes of dissolved organic carbon (DOC) in four small sub-tropical rivers of the Tampa Bay Estuary (Florida, USA). Applied Geochemistry, 2015, 63, 550-562.	1.4	26
158	Sediment plumes induced by the Port of Miami dredging: Analysis and interpretation using Landsat and MODIS data. Remote Sensing of Environment, 2015, 170, 328-339.	4.6	41
159	Habitat and water quality variables as predictors of community composition in an Indonesian coral reef: a multi-taxon study in the Spermonde Archipelago. Science of the Total Environment, 2015, 537, 139-151.	3.9	43
160	Assessing the potential of MODIS/Terra version 5 images to improve near shore lake bathymetric surveys. International Journal of Applied Earth Observation and Geoinformation, 2015, 36, 13-21.	1.4	11
161	Estimating wide range Total Suspended Solids concentrations from MODIS 250-m imageries: An improved method. ISPRS Journal of Photogrammetry and Remote Sensing, 2015, 99, 58-69.	4.9	85
162	Spatio-temporal Variability of Chlorophyll-A in the Coastal Zone of Northeastern Brazil. Estuaries and Coasts, 2015, 38, 72-83.	1.0	7
163	On the Potential of Robust Satellite Techniques Approach for SPM Monitoring in Coastal Waters: Implementation and Application over the Basilicata Ionian Coastal Waters Using MODIS-Aqua. Remote Sensing, 2016, 8, 922.	1.8	16

#	ARTICLE	IF	CITATIONS
164	Pre-melt-season sediment plume variability at Jökulsárlón, Iceland, a preliminary evaluation using in-situ spectroradiometry and satellite imagery. <i>Annals of Glaciology</i> , 2016, 57, 39-46.	2.8	9
165	A Semi-Analytic Model for Estimating Total Suspended Sediment Concentration in Turbid Coastal Waters of Northern Western Australia Using MODIS-Aqua 250 m Data. <i>Remote Sensing</i> , 2016, 8, 556.	1.8	39
166	A Sensor Web and Web Service-Based Approach for Active Hydrological Disaster Monitoring. <i>ISPRS International Journal of Geo-Information</i> , 2016, 5, 171.	1.4	10
167	Development of a Semi-Analytical Algorithm for the Retrieval of Suspended Particulate Matter from Remote Sensing over Clear to Very Turbid Waters. <i>Remote Sensing</i> , 2016, 8, 211.	1.8	92
168	A Quantitative Comparison of Total Suspended Sediment Algorithms: A Case Study of the Last Decade for MODIS and Landsat-Based Sensors. <i>Remote Sensing</i> , 2016, 8, 810.	1.8	28
169	A MODIS-Based Retrieval Model of Suspended Particulate Matter Concentration for the Two Largest Freshwater Lakes in China. <i>Sustainability</i> , 2016, 8, 832.	1.6	7
170	A Comprehensive Review on Water Quality Parameters Estimation Using Remote Sensing Techniques. <i>Sensors</i> , 2016, 16, 1298.	2.1	581
171	Monitoring water turbidity and surface suspended sediment concentration of the Bagre Reservoir (Burkina Faso) using MODIS and field reflectance data. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2016, 52, 243-251.	1.4	39
172	Spaceborne and airborne sensors in water quality assessment. <i>International Journal of Remote Sensing</i> , 2016, 37, 3143-3180.	1.3	33
173	Error analysis on Green Tide monitoring using MODIS data in the Yellow Sea based on GF-1 WFV data. , 2016, , .		1
174	A simple correction method for the MODIS surface reflectance product over typical inland waters in China. <i>International Journal of Remote Sensing</i> , 2016, 37, 6076-6096.	1.3	45
175	A novel MODIS algorithm to estimate chlorophyll a concentration in eutrophic turbid lakes. <i>Ecological Indicators</i> , 2016, 69, 138-151.	2.6	31
176	Nutrient reduction magnifies the impact of extreme weather on cyanobacterial bloom formation in large shallow Lake Taihu (China). <i>Water Research</i> , 2016, 103, 302-310.	5.3	131
177	Spatial and seasonal shifts in bloom-forming cyanobacteria in Lake Chaohu: Patterns and driving factors. <i>Phycological Research</i> , 2016, 64, 44-55.	0.8	88
178	Remote sensing observations and numerical studies of a super typhoon-induced suspended sediment concentration variation in the East China Sea. <i>Ocean Modelling</i> , 2016, 104, 187-202.	1.0	26
179	Retrieval of total suspended matter concentration from Gaofen-1 Wide Field Imager (WFI) multispectral imagery with the assistance of Terra MODIS in turbid water – case in Deep Bay. <i>International Journal of Remote Sensing</i> , 2016, 37, 3400-3413.	1.3	17
180	Island building in the South China Sea: detection of turbidity plumes and artificial islands using Landsat and MODIS data. <i>Scientific Reports</i> , 2016, 6, 33194.	1.6	24
181	Satellite analysis to identify changes and drivers of CyanoHABs dynamics in Lake Taihu. <i>Water Science and Technology: Water Supply</i> , 2016, 16, 1451-1466.	1.0	19

#	ARTICLE	IF	CITATIONS
182	Evaluation of the suitability of MODIS, OLCI and OLI for mapping the distribution of total suspended matter in the Barra Bonita Reservoir (Tiet�a River, Brazil). <i>Remote Sensing Applications: Society and Environment</i> , 2016, 4, 68-82.	0.8	14
183	Remote monitoring of sediment dynamics in a coastal lagoon: Long-term spatio-temporal variability of suspended sediment in Chilika. <i>Estuarine, Coastal and Shelf Science</i> , 2016, 170, 155-172.	0.9	40
184	Coral reefs chronically exposed to river sediment plumes in the southwestern Caribbean: Rosario Islands, Colombia. <i>Science of the Total Environment</i> , 2016, 553, 316-329.	3.9	50
185	Remote sensing of the impacts of construction in coastal waters on suspended particulate matter concentration – the case of the Yangtze River delta, China. <i>International Journal of Remote Sensing</i> , 2016, 37, 2132-2147.	1.3	15
186	Light absorption properties of the New York/New Jersey Harbor Estuary. <i>Hydrobiologia</i> , 2016, 766, 173-188.	1.0	0
187	Variability of Yellow River turbid plume detected with satellite remote sensing during water-sediment regulation. <i>Continental Shelf Research</i> , 2017, 135, 74-85.	0.9	27
188	Climate- and human-induced changes in suspended particulate matter over Lake Hongze on short and long timescales. <i>Remote Sensing of Environment</i> , 2017, 192, 98-113.	4.6	133
189	A study of a matching pixel by pixel (MPP) algorithm to establish an empirical model of water quality mapping, as based on unmanned aerial vehicle (UAV) images. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2017, 58, 213-224.	1.4	50
190	Remote Sensing Studies of Suspended Sediment Concentration Variations in a Coastal Bay During the Passages of Atmospheric Cold Fronts. <i>IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing</i> , 2017, 10, 2608-2622.	2.3	10
191	MODIS observations of cyanobacterial risks in a eutrophic lake: Implications for long-term safety evaluation in drinking-water source. <i>Water Research</i> , 2017, 122, 455-470.	5.3	107
192	Land adjacency effects on <sc>MODIS A</sc>qua topâ€ofâ€atmosphere radiance in the shortwave infrared: <sc>S</sc>tatistical assessment and correction. <i>Journal of Geophysical Research: Oceans</i> , 2017, 122, 4802-4818.	1.0	45
193	Improving river water quality monitoring using satellite data products and a genetic algorithm processing approach. <i>Sustainability of Water Quality and Ecology</i> , 2017, 9-10, 88-114.	2.0	19
194	Retrieving Total Suspended Matter in Tropical Reservoirs Within a Cascade System With Widely Differing Optical Properties. <i>IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing</i> , 2017, 10, 5495-5512.	2.3	8
195	Estimating sea surface salinity in the northern Gulf of Mexico from satellite ocean color measurements. <i>Remote Sensing of Environment</i> , 2017, 201, 115-132.	4.6	62
196	Uncertainties and applications of satellite-derived coastal water quality products. <i>Progress in Oceanography</i> , 2017, 159, 45-72.	1.5	74
197	A Hybrid EOF Algorithm to Improve MODIS Cyanobacteria Phycocyanin Data Quality in a Highly Turbid Lake: Bloom and Nonbloom Condition. <i>IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing</i> , 2017, 10, 4430-4444.	2.3	19
198	Differential Impact of Anniversary-Severe Cyclones on the Water Quality of a Tropical Coastal Lagoon. <i>Estuaries and Coasts</i> , 2017, 40, 317-342.	1.0	14
199	Using Satellite Observations to Characterize the Response of Estuarine Turbidity Maxima to External Forcing. <i>Estuaries and Coasts</i> , 2017, 40, 343-358.	1.0	22

#	ARTICLE	IF	CITATIONS
200	Wind and rainfall regulation of the diffuse attenuation coefficient in large, shallow lakes from long-term MODIS observations using a semianalytical model. <i>Journal of Geophysical Research D: Atmospheres</i> , 2017, 122, 6748-6763.	1.2	16
201	Variability of apparent and inherent optical properties of sediment-laden waters in large river basins – lessons from in situ measurements and bio-optical modeling. <i>Optics Express</i> , 2017, 25, A283.	1.7	30
202	A MODIS-Based Novel Method to Distinguish Surface Cyanobacterial Scums and Aquatic Macrophytes in Lake Taihu. <i>Remote Sensing</i> , 2017, 9, 133.	1.8	64
203	Turbidity in Apalachicola Bay, Florida from Landsat 5 TM and Field Data: Seasonal Patterns and Response to Extreme Events. <i>Remote Sensing</i> , 2017, 9, 367.	1.8	28
204	Assessment of Chlorophyll-a Remote Sensing Algorithms in a Productive Tropical Estuarine-Lagoon System. <i>Remote Sensing</i> , 2017, 9, 516.	1.8	43
205	Application of the Support Vector Regression Method for Turbidity Assessment with MODIS on a Shallow Coral Reef Lagoon (Voh-Kon-Pouembout, New Caledonia). <i>Water (Switzerland)</i> , 2017, 9, 737.	1.2	9
206	Spatio-Temporal Variability in a Turbid and Dynamic Tidal Estuarine Environment (Tasmania, Australia): An Assessment of MODIS Band 1 Reflectance. <i>ISPRS International Journal of Geo-Information</i> , 2017, 6, 320.	1.4	9
207	Application of Landsat 8 for Monitoring Impacts of Wastewater Discharge on Coastal Water Quality. <i>Frontiers in Marine Science</i> , 0, 4, .	1.2	44
208	Impact of the spatial resolution of satellite remote sensing sensors in the quantification of total suspended sediment concentration: A case study in turbid waters of Northern Western Australia. <i>PLoS ONE</i> , 2017, 12, e0175042.	1.1	38
209	A Landsat-based model for retrieving total suspended solids concentration of estuaries and coasts in China. <i>Geoscientific Model Development</i> , 2017, 10, 4347-4365.	1.3	42
210	A satellite remote-sensing multi-index approach to discriminate pelagic <i>Sargassum</i> in the waters of the Yucatan Peninsula, Mexico. <i>International Journal of Remote Sensing</i> , 2018, 39, 3608-3627.	1.3	37
211	Development of a MODIS data based algorithm for retrieving nearshore sea surface salinity along the northern Gulf of Mexico coast. <i>International Journal of Remote Sensing</i> , 2018, 39, 3497-3511.	1.3	12
212	Can MODIS Land Reflectance Products be Used for Estuarine and Inland Waters?. <i>Water Resources Research</i> , 2018, 54, 3583-3601.	1.7	20
213	Monitoring algal blooms in drinking water reservoirs using the Landsat-8 Operational Land Imager. <i>International Journal of Remote Sensing</i> , 2018, 39, 2818-2846.	1.3	22
214	Changes in water quality following opening and closure of a bar-built estuary (Pescadero, Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 182 Td	0.9	5
215	Applying geographic information systems and remote sensing for water quality assessment of mangrove forest. <i>Acta Ecologica Sinica</i> , 2018, 38, 135-143.	0.9	12
216	Using VIIRS/NPP and MODIS/Aqua data to provide a continuous record of suspended particulate matter in a highly turbid inland lake. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2018, 64, 256-265.	1.4	26
217	A novel cross-satellite based assessment of the spatio-temporal development of a cyanobacterial harmful algal bloom. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2018, 66, 69-81.	1.4	46

#	ARTICLE	IF	CITATIONS
218	Can Multispectral Information Improve Remotely Sensed Estimates of Total Suspended Solids? A Statistical Study in Chesapeake Bay. <i>Remote Sensing</i> , 2018, 10, 1393.	1.8	28
219	A study on the evaluation of water-bloom using image processing. <i>Environmental Science and Pollution Research</i> , 2018, 25, 36775-36780.	2.7	15
220	Remote Sensing of Chlorophyll-a in Case II Waters: A Novel Approach With Improved Accuracy Over Widely Implemented Turbid Water Indices. <i>Journal of Geophysical Research: Oceans</i> , 2018, 123, 8138-8158.	1.0	14
221	Characterizing a Sea Turtle Developmental Habitat Using Landsat Observations of Surface-Pelagic Drift Communities in the Eastern Gulf of Mexico. <i>IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing</i> , 2018, 11, 3646-3659.	2.3	11
222	Remote Sensing of Lakes' Water Environment. , 2018, , 249-277.		5
223	Applicability of Earth Observation chlorophyll-a data in assessment of water status via MERIS " With implications for the use of OLCI sensors. <i>Remote Sensing of Environment</i> , 2018, 212, 273-287.	4.6	31
224	Application of HySpex hyperspectral images for verification of a two-dimensional hydrodynamic model. <i>European Journal of Remote Sensing</i> , 2018, 51, 637-649.	1.7	5
225	Assessment of trophic state and water quality of coastal-inland lakes based on Fuzzy Inference System. <i>Journal of Great Lakes Research</i> , 2018, 44, 1010-1025.	0.8	17
226	Variations of transparency derived from GOCI in the Bohai Sea and the Yellow Sea. <i>Optics Express</i> , 2018, 26, 12191.	1.7	25
227	Comparison of two atmospheric correction approaches applied to MODIS measurements over North American waters. <i>Remote Sensing of Environment</i> , 2018, 216, 442-455.	4.6	21
228	Water Quality Drivers in 11 Gulf of Mexico Estuaries. <i>Remote Sensing</i> , 2018, 10, 255.	1.8	11
229	Retrieval of total suspended matter concentrations from high resolution WorldView-2 imagery: a case study of inland rivers. <i>IOP Conference Series: Earth and Environmental Science</i> , 2018, 121, 032036.	0.2	5
230	Impacts of 40 years of land cover change on water quality in Tampa Bay, Florida. <i>Cogent Geoscience</i> , 2018, 4, 1422956.	0.6	6
231	Evaluation of Unified Algorithms for Remote Sensing of Chlorophyll-a and Turbidity in Lake Shinji and Lake Nakaumi of Japan and the Vaal Dam Reservoir of South Africa under Eutrophic and Ultra-Turbid Conditions. <i>Water (Switzerland)</i> , 2018, 10, 618.	1.2	19
232	MERIS observations of phytoplankton phenology in the Baltic Sea. <i>Science of the Total Environment</i> , 2018, 642, 447-462.	3.9	7
233	A harmonized image processing workflow using Sentinel-2/MSI and Landsat-8/OLI for mapping water clarity in optically variable lake systems. <i>Remote Sensing of Environment</i> , 2019, 231, 111284.	4.6	89
234	Indirect Assessment of Sedimentation in Hydropower Dams Using MODIS Remote Sensing Images. <i>Remote Sensing</i> , 2019, 11, 314.	1.8	16
235	Remote Sensing Estimation of Lake Total Phosphorus Concentration Based on MODIS: A Case Study of Lake Hongze. <i>Remote Sensing</i> , 2019, 11, 2068.	1.8	31

#	ARTICLE	IF	CITATIONS
236	Enhancing the accuracy of retrieving quantities of turbidity and total suspended solids using Landsat-8-based-principal component analysis technique. <i>Journal of Spatial Science</i> , 2019, , 1-20.	1.0	7
237	Market Demand Forecasting Model of Urban Underground Comprehensive Pipe Gallery PPP Project. , 2019, , .		0
238	Estimation of Chlorophyll-a Concentrations in a Highly Turbid Eutrophic Lake Using a Classification-Based MODIS Land-Band Algorithm. <i>IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing</i> , 2019, 12, 3769-3783.	2.3	21
239	Reconstruction of sea-land interactions between terrestrial vegetation cover and water quality constituents in the Mattapoissett Harbor area during the 1991 Hurricane Bob event. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2019, 83, 101929.	1.4	0
240	Ensemble-Based Systems to Monitor Algal Bloom With Remote Sensing. <i>IEEE Transactions on Geoscience and Remote Sensing</i> , 2019, 57, 7955-7971.	2.7	13
241	High-frequency observation of floating algae from AHI on Himawari-8. <i>Remote Sensing of Environment</i> , 2019, 227, 151-161.	4.6	37
242	Preliminary Assessment of Turbidity and Chlorophyll Impact on Bathymetry Derived from Sentinel-2A and Sentinel-3A Satellites in South Florida. <i>Remote Sensing</i> , 2019, 11, 645.	1.8	78
243	A Near-Infrared Band-Based Algorithm for Suspended Sediment Estimation for Turbid Waters Using the Experimental Tiangong 2 Moderate Resolution Wide-Wavelength Imager. <i>IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing</i> , 2019, 12, 774-787.	2.3	9
244	Remote Sensing Estimation of Sea Surface Salinity from GOCI Measurements in the Southern Yellow Sea. <i>Remote Sensing</i> , 2019, 11, 775.	1.8	18
245	Long-term distribution patterns of remotely sensed water quality variables in Pearl River Delta, China. <i>Estuarine, Coastal and Shelf Science</i> , 2019, 221, 90-103.	0.9	21
246	Developing a Community of Practice for Applied Uses of Future PACE Data to Address Marine Food Security Challenges. <i>Frontiers in Earth Science</i> , 2019, 7, .	0.8	14
247	Correlation of Remotely Sensed Surface Reflectance With Forcing Variables in Six Different Estuaries. <i>Journal of Geophysical Research: Oceans</i> , 2019, 124, 9439-9461.	1.0	3
248	Chlorophyll-a estimation from remote sensing data of sea surface temperature and aerosol optical thickness through a shallow neural network. , 2019, , .		0
249	Satellite-based monitoring of contrasting characteristics of suspended sediment discharged from the Red and the Ma river systems along the northern coast of Vietnam. <i>International Journal of Sediment Research</i> , 2019, 34, 191-204.	1.8	5
251	The Salinity Structure of the Amazon River Plume Drives Spatiotemporal Variation of Oceanic Primary Productivity. <i>Journal of Geophysical Research G: Biogeosciences</i> , 2019, 124, 147-165.	1.3	27
252	Comparative analysis of four semi-analytical models for estimating chlorophyll-a concentration in case-2 waters using field hyperspectral reflectance. <i>International Journal of Remote Sensing</i> , 2020, 41, 584-594.	1.3	2
253	Drivers to spatial and temporal dynamics of column integrated phytoplankton biomass in the shallow lake of Chaohu, China. <i>Ecological Indicators</i> , 2020, 109, 105812.	2.6	15
254	Quantitative modeling of cyanobacterial concentration using MODIS imagery in the Southern Caspian Sea. <i>Journal of Great Lakes Research</i> , 2020, 46, 1251-1261.	0.8	8

#	ARTICLE	IF	CITATIONS
255	The effects of freshwater flow and salinity on turbidity and dissolved oxygen in a shallow Macrotidal estuary: A case study of Portsmouth Harbour. <i>Ocean and Coastal Management</i> , 2020, 191, 105179.	2.0	16
256	Opportunities for seagrass research derived from remote sensing: A review of current methods. <i>Ecological Indicators</i> , 2020, 117, 106560.	2.6	62
257	Spatiotemporal pattern of gypsum blooms in the Salton Sea, California, during 2000-2018. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2020, 89, 102090.	1.4	7
258	Quantification of lake clarity in China using Landsat OLI imagery data. <i>Remote Sensing of Environment</i> , 2020, 243, 111800.	4.6	74
259	Impacts of a Major Mississippi River Freshwater Diversion on Suspended Sediment Plume Kinematics in Lake Pontchartrain, a Semi-enclosed Gulf of Mexico Estuary. <i>Estuaries and Coasts</i> , 2021, 44, 704-721.	1.0	8
260	Using Satellite Remote Sensing to Study the Effect of Sand Excavation on the Suspended Sediment in the Hong Kong-Zhuhai-Macau Bridge Region. <i>Water (Switzerland)</i> , 2021, 13, 435.	1.2	5
261	Wind Effects for Floating Algae Dynamics in Eutrophic Lakes. <i>Remote Sensing</i> , 2021, 13, 800.	1.8	18
262	Potential for High Fidelity Global Mapping of Common Inland Water Quality Products at High Spatial and Temporal Resolutions Based on a Synthetic Data and Machine Learning Approach. <i>Frontiers in Environmental Science</i> , 2021, 9, .	1.5	17
263	Coastal water bathymetry for critical zone management using regression tree models from Gaofen-6 imagery. <i>Ocean and Coastal Management</i> , 2021, 204, 105522.	2.0	9
264	Retrieval of suspended sediment concentration of the Chilika Lake, India using Landsat-8 OLI satellite data. <i>Environmental Earth Sciences</i> , 2021, 80, 1.	1.3	8
265	Estimation of suspended matter concentration in manwan reservoir, lancang river using remotely sensed small satellite constellation for environment and disaster monitoring and forecasting (HJ-1A/1B), charge coupled device (CCD) data. <i>International Journal of Remote Sensing</i> , 2021, 42, 5232-5252.	1.3	1
266	Optimized remote sensing estimation of the lake algal biomass by considering the vertically heterogeneous chlorophyll distribution: Study case in Lake Chaohu of China. <i>Science of the Total Environment</i> , 2021, 771, 144811.	3.9	17
267	Estimation of Suspended Sediment Concentration from Remote Sensing and In Situ Measurement over Lake Tana, Ethiopia. <i>Advances in Civil Engineering</i> , 2021, 2021, 1-17.	0.4	5
268	Remote Detection of Cyanobacterial Blooms and Chlorophyll-a Analysis in a Eutrophic Reservoir Using Sentinel-2. <i>Sustainability</i> , 2021, 13, 8570.	1.6	14
269	Adaptive Threshold Model in Google Earth Engine: A Case Study of <i>Ulva prolifera</i> Extraction in the South Yellow Sea, China. <i>Remote Sensing</i> , 2021, 13, 3240.	1.8	16
270	Spatiotemporal variability of secchi depths of the North Arabian Gulf over the last two decades. <i>Estuarine, Coastal and Shelf Science</i> , 2021, 260, 107487.	0.9	8
271	Quantifying Turbidity Variation for Lakes in Daqing of Northeast China Using Landsat Images From 1984 to 2018. <i>IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing</i> , 2021, 14, 8884-8897.	2.3	3
272	Examining Material Transport in Dynamic Coastal Environments: An Integrated Approach Using Field Data, Remote Sensing and Numerical Modeling. <i>Coastal Research Library</i> , 2014, , 333-364.	0.2	1

#	ARTICLE	IF	CITATIONS
273	High-Performance Ocean Color Monte Carlo Simulation in the Geo-info Project. Lecture Notes in Computer Science, 2010, , 370-379.	1.0	5
274	Cyanobacterial Blooms as an Indicator of Environmental Degradation in Waters and Their Monitoring Using Satellite Remote Sensing. Structure and Function of Mountain Ecosystems in Japan, 2016, , 71-85.	0.1	1
275	Integrated monitoring of lakes' turbidity in Wuhan, China during the COVID-19 epidemic using multi-sensor satellite observations. International Journal of Digital Earth, 2021, 14, 443-463.	1.6	11
276	Ocean color as a proxy to predict sea surface salinity in the Banda Sea. IOP Conference Series: Earth and Environmental Science, 0, 618, 012037.	0.2	1
277	Evaluating ocean-color algorithms to remotely sense the surface suspended particle matter in the northeast Aegean Sea, Greece. , 2018, , .		1
278	Review of Selected Moderate-Resolution Imaging Spectroradiometer Algorithms, Data Products and Applications. Taylor & Francis Series in Remote Sensing Applications, 2011, , 31-55.	0.0	4
279	Coupled approach for radiometric calibration and parameter retrieval to improve SPM estimations in turbid inland/coastal waters. Optics Express, 2020, 28, 5567.	1.7	6
280	Detection and Characterization of Low Temperature Peat Fires during the 2015 Fire Catastrophe in Indonesia Using a New High-Sensitivity Fire Monitoring Satellite Sensor (FireBird). PLoS ONE, 2016, 11, e0159410.	1.1	43
281	Development of Regional TSS Algorithm over Penang using Modis Terra (250 M) Surface Reflectance Product. Ekologia, 2016, 35, 289-294.	0.2	2
282	Remote Sensing of Temporal and Spatial Variations of Suspended Sediment Concentration in Bahmanshir Estuary, Iran. Indian Journal of Science and Technology, 2013, 6, 1-10.	0.5	12
283	Cyanobacteria bloom monitoring with remote sensing in Lake Taihu. Hupo Kexue/Journal of Lake Sciences, 2008, 20, 145-152.	0.3	37
284	The NIR peak of the reflectance spectrum associated with chlorophyll in the pool waters: preliminary results. Hupo Kexue/Journal of Lake Sciences, 2009, 21, 228-233.	0.3	2
285	Time-series MODIS images-based retrieval and change analysis of suspended sediment concentration during flood period in Lake Poyang. Hupo Kexue/Journal of Lake Sciences, 2009, 21, 288-297.	0.3	3
286	Suitability of the retrieval models for estimating chlorophyll concentration in Lake Taihu. Hupo Kexue/Journal of Lake Sciences, 2015, 27, 150-162.	0.3	4
287	Temporal and spatial distribution of algal blooms in Lake Chaohu, 2000-2015. Hupo Kexue/Journal of Lake Sciences, 2017, 29, 276-284.	0.3	17
289	Monitoring turbidity in the Ionical coast during extreme events by applying a Robust Satellite Technique (RST) to MODIS imagery. , 2011, , .		1
290	Remote Quantification of Total Suspended Matter through Empirical Approaches for Inland Waters. Journal of Environmental Informatics, 2014, 23, 23-36.	6.0	13
291	Liquid crystal tunable filters in detecting water pollution. , 2014, , .		1

#	ARTICLE	IF	CITATIONS
292	Extracting A-L Relationship for Urmia Lake, Iran Using MODIS NDVI/NDWI Indices. Journal of Hydrogeology and Hydrologic Engineering, 2018, 07, .	0.1	3
293	Análisis multisensor para el estudio de los patrones de turbidez en el estuario del Guadalquivir. Revista De Teledeteccion, 2016, , 1.	0.6	5
295	Hydrologic controls on coastal suspended sediment plumes around the Greenland Ice Sheet. Cryosphere, 2012, 6, 1-19.	1.5	56
297	Quantitative Retrieval of Water Quality by Remote Sensing in Taihu Lake Based on MODIS data. Geo-information Science, 2010, 11, 104-110.	0.1	0
299	Water Quality Monitoring in Potable Water Reservoirs of Cities by Remote Sensing and GIS Techniques. , 2012, , .		0
300	Remote Sensing Retrieval of Turbidity Near Radial Sand Ridges Area in the South Yellow Sea Using HJ-1A/B CCD Imagery. Lecture Notes in Electrical Engineering, 2012, , 121-127.	0.3	0
303	Application of Advanced Remote Sensing Techniques to Improve Modeling Estuary Water Quality. Coastal Research Library, 2014, , 295-313.	0.2	1
304	Suitability of MODIS 250 m resolution band data for quantitative mapping of cyanobacterial blooms. Estonian Journal of Ecology, 2006, 55, 318.	0.5	12
305	An operational algorithm to estimate chlorophyll-a concentrations in Lake Chaohu from MODIS imagery. Hupo Kexue/Journal of Lake Sciences, 2015, 27, 1140-1150.	0.3	2
306	Can Watershed Models Aid in Determining Historic Lake Sediment Concentrations in Data-Scarce Areas?. Springer Geography, 2016, , 819-833.	0.3	0
307	Estimating chlorophyll-A concentration in the Caspian Sea from MODIS images using artificial neural networks. Environmental Engineering Research, 2020, 25, 515-521.	1.5	2
308	Long-Term Analysis of Water Quality in Chilika Lagoon and Application of Bio-optical Models for Cyclone Impact Assessment. Wetlands: Ecology, Conservation and Management, 2020, , 165-202.	0.0	0
309	Sentinel-2 and Landsat-8 Observations for Harmful Algae Blooms in a Small Eutrophic Lake. Remote Sensing, 2021, 13, 4479.	1.8	13
310	Analyzing the distribution and variation of Suspended Particulate Matter (SPM) in the Yellow River Estuary (YRE) using Landsat 8 OLI. Regional Studies in Marine Science, 2021, 48, 102064.	0.4	1
313	Spatial and seasonal variability of chlorophyll <i>a</i> in different-sized lakes across eastern China. Inland Waters, 2022, 12, 205-214.	1.1	4
314	Machine Learning Based Long-Term Water Quality in the Turbid Pearl River Estuary, China. Journal of Geophysical Research: Oceans, 2022, 127, .	1.0	15
315	Modeling Secchi Disk Depth Over the North Arabian Gulf Waters Using MODIS and MERIS Images. PFG - Journal of Photogrammetry, Remote Sensing and Geoinformation Science, 2022, 90, 177-189.	0.7	2
316	A deep learning method for cyanobacterial harmful algae blooms prediction in Taihu Lake, China. Harmful Algae, 2022, 113, 102189.	2.2	30

#	ARTICLE	IF	CITATIONS
317	Development of remote sensing algorithm for total phosphorus concentration in eutrophic lakes: Conventional or machine learning?. <i>Water Research</i> , 2022, 215, 118213.	5.3	28
318	Remote sensing of total suspended matter concentration in lakes across China using Landsat images and Google Earth Engine. <i>ISPRS Journal of Photogrammetry and Remote Sensing</i> , 2022, 187, 61-78.	4.9	26
319	Monitoring of Turbidity Variation in the Ukai Reservoir, Gujarat, INDIA, during 1993-2018 using Landsat Series of Dataset. <i>International Journal of Scientific Research in Science and Technology</i> , 2021, , 236-251.	0.1	1
320	Decline of suspended particulate matter concentrations in Lake Taihu from 1984 to 2020: observations from Landsat TM and OLI. <i>Optics Express</i> , 2022, 30, 22572.	1.7	6
321	Prediction of Total Phosphorus Concentration in Macrophytic Lakes Using Chlorophyll-Sensitive Bands: A Case Study of Lake Baiyangdian. <i>Remote Sensing</i> , 2022, 14, 3077.	1.8	6
322	Turbidity dynamics of large lakes and reservoirs in northeastern China in response to natural factors and human activities. <i>Journal of Cleaner Production</i> , 2022, 368, 133148.	4.6	7
323	Remote sensing of chlorophyll-a concentrations in coastal oceans of the Greater Bay Area in China: Algorithm development and long-term changes. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2022, 112, 102922.	0.9	3
324	MODIS-Landsat fusion-based single-band algorithms for TSS and turbidity estimation in an urban-waste-dominated river reach. <i>Water Research</i> , 2022, 224, 119082.	5.3	7
325	Sea Surface Salinity Inversion Model for Changjiang Estuary and Adjoining Sea Area with SMAP and MODIS Data Based on Machine Learning and Preliminary Application. <i>Remote Sensing</i> , 2022, 14, 5358.	1.8	0
326	Remote Estimation of the Chlorophyll-a Concentration in Lake Dianshan, China Using High-Spatial-Resolution Satellite Imagery. <i>Journal of the Indian Society of Remote Sensing</i> , 2022, 50, 2465-2477.	1.2	1
327	Evaluating and Optimizing VIIRS Retrievals of Chlorophyll-a and Suspended Particulate Matter in Turbid Lakes Using a Machine Learning Approach. <i>IEEE Transactions on Geoscience and Remote Sensing</i> , 2022, 60, 1-17.	2.7	4
328	Mysterious increases of whiting events in the Bahama Banks. <i>Remote Sensing of Environment</i> , 2023, 285, 113389.	4.6	6
329	Adaptive optical algorithms with differentiation of water bodies based on varying composition of suspended particulate matter: A case study for estimating the particulate organic carbon concentration in the western Arctic seas. <i>Remote Sensing of Environment</i> , 2023, 286, 113360.	4.6	6
330	Using MODIS data to track the long-term variations of dissolved oxygen in Lake Taihu. <i>Frontiers in Environmental Science</i> , 0, 10, .	1.5	3
331	Overview of the Application of Remote Sensing in Effective Monitoring of Water Quality Parameters. <i>Remote Sensing</i> , 2023, 15, 1938.	1.8	22
332	Algal biomass mapping of eutrophic lakes using a machine learning approach with MODIS images. <i>Science of the Total Environment</i> , 2023, 880, 163357.	3.9	4
334	Tracer study of CDOM for nitrogen and phosphorus pollution in the offshore Leizhou Peninsula, China. <i>Ecological Indicators</i> , 2023, 148, 110019.	2.6	2
335	HY1C/D-CZI Noctiluca scintillans Bloom Recognition Network Based on Hybrid Convolution and Self-Attention. <i>Remote Sensing</i> , 2023, 15, 1757.	1.8	3

#	ARTICLE	IF	CITATIONS
352	UAV and satellite remote sensing for inland water quality assessments: a literature review. Environmental Monitoring and Assessment, 2024, 196, .	1.3	0