

CITATION REPORT

List of articles citing

Determination of sets of solute descriptors from chromatographic measurements

DOI: 10.1016/j.chroma.2003.12.004
Journal of Chromatography A, 2004, 1037, 29-47.

Source: <https://exaly.com/paper-pdf/37159472/citation-report.pdf>

Version: 2024-04-29

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
776	Systematic Selection of Internal Standard with Similar Chemical and UV Properties to Drug to Be Quantified in Serum Samples. <i>Chromatographia</i> , 2004 , 60, 391-397	2.1	5
775	Appraisal of an Empirical Model for Simulation of Retention from Structure in Temperature-Programmed Gas Chromatography. <i>Chromatographia</i> , 2004 , 60, 663-672	2.1	5
774	Insights into the retention mechanism of neutral organic compounds on polar chemically bonded stationary phases in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2004 , 1052, 199-204	4.5	22
773	Correlation and prediction of partition coefficients between the gas phase and water, and the solvents dodecane and undecane. 2004 , 28, 1538		39
772	Characterisation of the water-isopropyl myristate system. <i>International Journal of Pharmaceutics</i> , 2005 , 294, 121-8	6.5	29
771	Solubility of 9-fluorenone, thianthrene and xanthene in organic solvents. 2005 , 232, 113-121		59
770	Predicting the equilibrium partitioning of organic compounds using just one linear solvation energy relationship (LSER). 2005 , 233, 19-22		130
769	Comments regarding predicting the equilibrium partitioning of organic compounds using just one linear solvation energy relationship (LSER) 2005 , 237, 224-226		22
768	Effects of modifiers in subcritical fluid chromatography on retention with porous graphitic carbon. <i>Journal of Chromatography A</i> , 2005 , 1087, 64-76	4.5	45
767	System constants of synthesized poly(methyl-3,3,3-trifluoropropyl) siloxanes. <i>Journal of Chromatography A</i> , 2005 , 1100, 208-17	4.5	17
766	Evaluation of a Semi-Empirical Approach to Correlation of Retention with Octanol-Water Partition Coefficients by Use of Adjustable Hydrogen-Bonding Indicator Variables. <i>Chromatographia</i> , 2005 , 61, 479-492	2.1	5
765	System Maps for XTerra MS C18: Effect of Solvent Type on Selectivity in Reversed-Phase Liquid Chromatography. <i>Chromatographia</i> , 2005 , 61, 587-593	2.1	22
764	Contribution of Steric Repulsion to Retention on an Octadecylsiloxane-Bonded Silica Stationary Phase in Reversed-Phase Liquid Chromatography. <i>Chromatographia</i> , 2005 , 62, 553-561	2.1	34
763	Using comprehensive two-dimensional gas chromatography retention indices to estimate environmental partitioning properties for a complete set of diesel fuel hydrocarbons. 2005 , 77, 7172-82		90
762	Correlation of the Solubility Behavior of Crystalline 1-Nitronaphthalene in Organic Solvents With the Abraham Solvation Parameter Model. <i>Journal of Solution Chemistry</i> , 2005 , 34, 1121-1133	1.8	24
761	SOLUBILITY BEHAVIOR OF CRYSTALLINE POLYCYCLIC AROMATIC HYDROCARBONS (PAHS): PREDICTION OF FLUORENE SOLUBILITIES IN ORGANIC SOLVENTS WITH THE ABRAHAM SOLVATION PARAMETER MODEL. 2005 , 25, 313-326		13
760	Solvation parameters for the 209 PCBs: calculation of physicochemical properties. 2005 , 7, 295-301		42

759	Models for the adsorption of organic compounds at gas-water interfaces. 2005 , 7, 577-80		15
758	Comment on "model for the adsorption of organic compounds at gas-water interfaces" by C. F. Poole, JEM, 2005, 7, 577. 2005 , 7, 1105-6; author reply 1107-9		2
757	Solubility of crystalline nonelectrolyte solutes in organic solvents: Mathematical correlation of ibuprofen solubilities with the Abraham solvation parameter model. <i>Physics and Chemistry of Liquids</i> , 2005 , 43, 261-268	1.5	20
756	The estimation of physicochemical properties of methyl and other alkyl naphthalenes. 2005 , 7, 445-9		6
755	Characterisation of the water/o-nitrophenyl octyl ether system in terms of the partition of nonelectrolytes and of ions. 2005 , 7, 2418-22		33
754	Solubility of crystalline nonelectrolyte solutes in organic solvents: mathematical correlation of 4-chloro-3-nitrobenzoic acid and 2-chloro-5-nitrobenzoic acid solubilities with the Abraham solvation parameter model. <i>Physics and Chemistry of Liquids</i> , 2005 , 43, 351-360	1.5	56
753	Octanol/water partition of ionic species, including 544 cations. 2005 , 70, 2633-40		47
752	The electrostatic origin of Abraham's solute polarity parameter. 2005 , 109, 7564-73		56
751	A concentration-dependent multi-term linear free energy relationship for sorption of organic compounds to soils based on the hexadecane dilute-solution reference state. <i>Environmental Science & Technology</i> , 2005 , 39, 8817-28	10.3	48
750	Air to blood distribution of volatile organic compounds: a linear free energy analysis. 2005 , 18, 904-11		67
749	Polyparameter linear free energy relationships for estimating the equilibrium partition of organic compounds between water and the natural organic matter in soils and sediments. <i>Environmental Science & Technology</i> , 2005 , 39, 913-24	10.3	224
748	Investigating Unused Hydrogen Bond Acceptors Using Known and Hypothetical Crystal Polymorphism. 2005 , 5, 983-993		46
747	High-throughput physicochemical and in vitro ADMET screening. 2005 , 3, 83-100		14
746	Chemical toxicity correlations for several fish species based on the Abraham solvation parameter model. 2005 , 18, 1497-505		72
745	Air to muscle and blood/plasma to muscle distribution of volatile organic compounds and drugs: linear free energy analyses. 2006 , 19, 801-8		53
744	Prediction of partitioning between complex organic mixtures and water: application of polyparameter linear free energy relationships. <i>Environmental Science & Technology</i> , 2006 , 40, 536-45	10.3	35
743	Comments on "solvation parameters. 2. A simplified molecular topology to generate easily optimized values". <i>Journal of Chemical Information and Modeling</i> , 2006 , 46, 1879-81	6.1	8
742	Gas to olive oil partition coefficients: a linear free energy analysis. <i>Journal of Chemical Information and Modeling</i> , 2006 , 46, 1735-41	6.1	33

741	Sorption equilibrium of a wide spectrum of organic vapors in Leonardite humic acid: modeling of experimental data. <i>Environmental Science & Technology</i> , 2006 , 40, 5374-9	10.3	72
740	The partition behavior of fluorotelomer alcohols and olefins. <i>Environmental Science & Technology</i> , 2006 , 40, 3572-7	10.3	90
739	. 2006 ,		23
738	Selectivity Characterization of Pseudostationary Phases Using the Solvation Parameter Model. 55-78		1
737	Correlation of Minimum Inhibitory Concentrations Toward Oral Bacterial Growth Based on the Abraham Model. 2006 , 25, 912-920		7
736	Hydrogen bonding, steric effects and thermodynamics of partitioning. 2006 , 19, 242-248		22
735	Air to brain, blood to brain and plasma to brain distribution of volatile organic compounds: linear free energy analyses. 2006 , 41, 494-502		50
734	Air to fat and blood to fat distribution of volatile organic compounds and drugs: linear free energy analyses. 2006 , 41, 1430-8		46
733	Characterisation of stationary phases in subcritical fluid chromatography by the solvation parameter model. II. Comparison tools. <i>Journal of Chromatography A</i> , 2006 , 1110, 191-9	4.5	60
732	Characterisation of stationary phases in subcritical fluid chromatography with the solvation parameter model IV. Aromatic stationary phases. <i>Journal of Chromatography A</i> , 2006 , 1115, 233-45	4.5	88
731	Insights into the retention mechanism on an octadecylsiloxane-bonded silica stationary phase (HyPURITY C18) in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2006 , 1115, 133-41	4.5	32
730	Solute-solvent interactions in micellar liquid chromatography. Characterization of hybrid micellar systems of sodium dodecyl sulfate-pentanol. <i>Journal of Chromatography A</i> , 2006 , 1117, 47-55	4.5	24
729	Application of the solvation parameter model to poly(methylcyanopropylsiloxane) stationary phases. <i>Journal of Chromatography A</i> , 2006 , 1122, 230-41	4.5	14
728	Evaluation of the separation characteristics of application-specific (pesticides and dioxins) open-tubular columns for gas chromatography. <i>Journal of Chromatography A</i> , 2006 , 1128, 228-35	4.5	15
727	Evaluation of the separation characteristics of application-specific (volatile organic compounds) open-tubular columns for gas chromatography. <i>Journal of Chromatography A</i> , 2006 , 1134, 284-90	4.5	10
726	Revised solute descriptors for characterizing retention properties of open-tubular columns in gas chromatography and their application to a carborane-siloxane copolymer stationary phase. <i>Journal of Chromatography A</i> , 2006 , 1104, 299-312	4.5	42
725	Model for the distribution of neutral organic compounds between n-hexane and acetonitrile. <i>Journal of Chromatography A</i> , 2006 , 1104, 82-90	4.5	32
724	Characterisation of stationary phases in subcritical fluid chromatography with the solvation parameter model. III. Polar stationary phases. <i>Journal of Chromatography A</i> , 2006 , 1110, 200-13	4.5	113

723	Characterization of stationary phases in subcritical fluid chromatography by the solvation parameter model. I. Alkylsiloxane-bonded stationary phases. <i>Journal of Chromatography A</i> , 2006 , 1110, 181-90	4.5	88
722	The chemical interpretation and practice of linear solvation energy relationships in chromatography. <i>Journal of Chromatography A</i> , 2006 , 1126, 143-94	4.5	410
721	Comparative analysis of solvation and selectivity in room temperature ionic liquids using the Abraham linear free energy relationship. 2006 , 8, 906		124
720	Insights into the Retention Mechanism on a Pentafluorophenylpropylsiloxane-Bonded Silica Stationary Phase (Discovery HS F5) in RP-LC. <i>Chromatographia</i> , 2006 , 63, 407-417	2.1	30
719	Partitioning properties of linear and branched ethers: Determination of linear solvation energy relationship (LSER) descriptors. 2006 , 246, 143-152		16
718	Correlating toxicities of organic compounds to select protozoa using the Abraham model. 2006 , 369, 109-18		21
717	Correlation of the toxicity of organic compounds to tadpoles using the Abraham model. 2006 , 371, 99-109		37
716	The analysis of solvation in ionic liquids and organic solvents using the Abraham linear free energy relationship. 2006 , 81, 1441-1446		87
715	Separation characteristics of phenyl-containing stationary phases for gas chromatography based on silarylene-siloxane copolymer chemistries. 2006 , 29, 211-7		18
714	Distribution of neutral organic compounds between n-heptane and methanol or N,N-dimethylformamide. 2006 , 29, 2158-65		26
713	A data base for partition of volatile organic compounds and drugs from blood/plasma/serum to brain, and an LFER analysis of the data. 2006 , 95, 2091-100		109
712	Partition coefficient correlations for transfer of solutes from gas phase and water to room temperature ionic liquids. <i>Physics and Chemistry of Liquids</i> , 2007 , 45, 241-249	1.5	11
711	A quantitative structure activity analysis on the relative sensitivity of the olfactory and the nasal trigeminal chemosensory systems. 2007 , 32, 711-9		47
710	Chemical toxicity correlations for several protozoas, bacteria, and water fleas based on the Abraham solvation parameter model. 2007 , 6, 165-174		36
709	Modelling and field application of the Chemcatcher passive sampler calibration data for the monitoring of hydrophobic organic pollutants in water. 2007 , 145, 895-904		84
708	Henry's Law constants or air to water partition coefficients for 1,3,5-triazines by an LFER method. 2007 , 9, 234-9		15
707	Comment on "systematic investigation of the sorption properties of polyurethane foams for organic vapors". 2007 , 79, 6891-3		14
706	Limiting partition coefficients of solutes in biphasic trihexyltetradecylphosphonium chloride ionic liquid-supercritical CO ₂ system: measurement and LSER-based correlation. 2007 , 111, 7620-5		16

705	Alternative approaches for modeling gas-particle partitioning of semivolatile organic chemicals: model development and comparison. <i>Environmental Science & Technology</i> , 2007 , 41, 1272-8	10.3	78
704	Characterization of room-temperature ionic liquids by the Abraham model with cation-specific and anion-specific equation coefficients. <i>Journal of Chemical Information and Modeling</i> , 2007 , 47, 1123-9	6.1	69
703	Predicting dermal permeability of biocides in commercial cutting fluids using a LSER approach. 2007 , 175, 34-43		14
702	Linear free energy relationship correlation of the distribution of solutes between water and sodium dodecyl sulfate (SDS) micelles and between gas and SDS micelles. <i>Journal of Chemical Information and Modeling</i> , 2007 , 47, 1808-17	6.1	31
701	Enthalpy of solvation correlations for gaseous solutes dissolved in water and in 1-octanol based on the Abraham model. <i>Journal of Chemical Information and Modeling</i> , 2007 , 47, 115-21	6.1	95
700	In Silico Models to Predict Brain Uptake. 2007 , 745-766		6
699	Evaluation of the separation characteristics of application-specific (fatty acid methyl esters) open-tubular columns for gas chromatography. 2007 , 30, 740-5		16
698	Distribution model for Folch partition. 2007 , 30, 2326-31		17
697	Enthalpy of solvation correlations for gaseous solutes dissolved in dimethyl sulfoxide and propylene carbonate based on the Abraham model. 2007 , 459, 17-25		43
696	Air to liver partition coefficients for volatile organic compounds and blood to liver partition coefficients for volatile organic compounds and drugs. 2007 , 42, 743-51		54
695	Partition of compounds from gas to water and from gas to physiological saline at 310 K: Linear free energy relationships. 2007 , 251, 93-109		65
694	Distribution of neutral organic compounds between n-heptane and fluorine-containing alcohols. <i>Journal of Chromatography A</i> , 2007 , 1143, 276-83	4.5	34
693	Combined supercritical fluid chromatographic methods for the characterization of octadecylsiloxane-bonded stationary phases. <i>Journal of Chromatography A</i> , 2007 , 1149, 345-57	4.5	30
692	Quantitative structure-(chromatographic) retention relationships. <i>Journal of Chromatography A</i> , 2007 , 1158, 273-305	4.5	302
691	Characterization of the retention behavior of organic and pharmaceutical drug molecules on an immobilized artificial membrane column with the Abraham model. <i>Journal of Chromatography A</i> , 2007 , 1160, 235-45	4.5	30
690	Could linear solvation energy relationships give insights into chiral recognition mechanisms? 2. Characterization of macrocyclic glycopeptide stationary phases. <i>Journal of Chromatography A</i> , 2007 , 1166, 70-8	4.5	21
689	Could linear solvation energy relationships give insights into chiral recognition mechanisms? 1. Pi-pi and charge interaction in the reversed versus the normal phase mode. <i>Journal of Chromatography A</i> , 2007 , 1166, 61-9	4.5	32
688	Determination of descriptors for organosilicon compounds by gas chromatography and non-aqueous liquid-liquid partitioning. <i>Journal of Chromatography A</i> , 2007 , 1169, 179-92	4.5	99

687	Characterization of the sorption of gaseous and organic solutes onto polydimethyl siloxane solid-phase microextraction surfaces using the Abraham model. <i>Journal of Chromatography A</i> , 2007 , 1175, 162-73	4.5	83
686	Quantitative structure-retention relationships of pesticides in reversed-phase high-performance liquid chromatography. 2007 , 582, 235-42		44
685	Modelling of retention of pesticides in reversed-phase high-performance liquid chromatography: quantitative structure-retention relationships based on solute quantum-chemical descriptors and experimental (solvatochromic and spin-probe) mobile phase descriptors. 2007 , 593, 140-51		26
684	Investigation of retention behaviour of non-steroidal anti-inflammatory drugs in high-performance liquid chromatography by using quantitative structure-retention relationships. 2007 , 601, 68-76		38
683	Characterisation of stationary phases in supercritical fluid chromatography with the solvation parameter model V. Elaboration of a reduced set of test solutes for rapid evaluation. <i>Journal of Chromatography A</i> , 2007 , 1169, 205-19	4.5	35
682	Blood or plasma to skin distribution of drugs: a linear free energy analysis. <i>International Journal of Pharmaceutics</i> , 2007 , 329, 129-34	6.5	25
681	Enthalpy of Solvation Correlations for Gaseous Solutes Dissolved in Benzene and in Alkane Solvents Based on the Abraham Model. 2007 , 26, 881-888		36
680	Relationship between the structure of compounds and the effect of Pluronic L61 on their permeation through lipid membranes. 2007 , 49, 1034-1044		
679	QSRR: quantitative structure-(chromatographic) retention relationships. 2007 , 107, 3212-46		372
678	Insights into the Retention Mechanisms on Perfluorohexylpropylsiloxane-Bonded (Fluophase-RP) and Octadecylsiloxane-Bonded (Betasil C18) Stationary Phases Based on the Same Silica Substrate in RP-LC. <i>Chromatographia</i> , 2007 , 65, 127-139	2.1	29
677	Comparison of the Separation Characteristics of the Organic-Inorganic Hybrid Octadecyl Stationary Phases XTerra MS C18 and XBridge C18 and Shield RP18 in RPLC. <i>Chromatographia</i> , 2007 , 66, 453-460	2.1	22
676	PEG-linked geminal dicationic ionic liquids as selective, high-stability gas chromatographic stationary phases. 2007 , 389, 2265-75		139
675	Enthalpy of Solvation Correlations for Gaseous Solutes Dissolved in Toluene and Carbon Tetrachloride Based on the Abraham Model. <i>Journal of Solution Chemistry</i> , 2007 , 36, 947-966	1.8	39
674	Enthalpy of solvation correlations for gaseous solutes dissolved in chloroform and 1,2-dichloroethane based on the Abraham model. 2007 , 258, 191-198		34
673	Prediction of gas to water partition coefficients from 273 to 373K using predicted enthalpies and heat capacities of hydration. 2007 , 262, 97-110		33
672	Correlation and prediction of partition coefficients for solute transfer to 1,2-dichloroethane from both water and from the gas phase. 2008 , 273, 78-86		33
671	Progress and perspectives of quantitative structure-activity relationships used for ecological risk assessment of toxic organic compounds. 2008 , 51, 593		23
670	Comparison of solubility of gases and vapours in wet and dry alcohols, especially octan-1-ol. 2008 , 21, 823-832		26

669	Enthalpy of Solvation Correlations For Gaseous Solutes Dissolved in Linear Alkanes (C5-C16) Based on the Abraham Model. 2008 , 27, 179-186		25
668	Characterization of the Partitioning of Gaseous Solutes Into Humic Acid with the Abraham Model and Temperature-Independent Equation Coefficients. 2008 , 27, 483-491		14
667	Enthalpy of Solvation Correlations for Gaseous Solutes Dissolved in Alcohol Solvents based on the Abraham Model. 2008 , 27, 627-635		30
666	Mathematical Correlations for Gas-to-Olive Oil, Gas-to-Saline Solution, and Saline Solution-to-Olive Oil Partition Coefficients Based on the Goss Modified Abraham Model. 2008 , 27, 890-900		16
665	Correlation of Human and Animal Air-to-Blood Partition Coefficients With a Single Linear Free Energy Relationship Model. 2008 , 27, 1130-1139		15
664	Effect of anesthetic structure on inhalation anesthesia: implications for the mechanism. 2008 , 97, 2373-84		20
663	The orthogonal character of stationary phases for gas chromatography. 2008 , 31, 1118-23		43
662	Overview of the retention in subcritical fluid chromatography with varied polarity stationary phases. 2008 , 31, 1238-51		66
661	Correlation and prediction of partition coefficient between the gas phase and water, and the solvents dry methyl acetate, dry and wet ethyl acetate, and dry and wet butyl acetate. 2008 , 270, 30-44		60
660	Separation characteristics of wall-coated open-tubular columns for gas chromatography. <i>Journal of Chromatography A</i> , 2008 , 1184, 254-80	4-5	156
659	Cluster and principal component analysis for Kov _{ls} ' retention indices on apolar and polar stationary phases in gas chromatography. <i>Journal of Chromatography A</i> , 2008 , 1177, 175-82	4-5	16
658	Quantitative structure-retention (property) relationships in micellar electrokinetic chromatography. <i>Journal of Chromatography A</i> , 2008 , 1182, 1-24	4-5	54
657	Application of linear solvation energy relationships to a custom-made polyaniline solid-phase microextraction fiber and three commercial fibers. <i>Journal of Chromatography A</i> , 2008 , 1188, 108-17	4-5	8
656	A unified classification of stationary phases for packed column supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2008 , 1191, 21-39	4-5	138
655	Combination of artificial neural network technique and linear free energy relationship parameters in the prediction of gradient retention times in liquid chromatography. <i>Journal of Chromatography A</i> , 2008 , 1190, 241-52	4-5	32
654	Solute descriptors for characterizing retention properties of open-tubular columns of different selectivity in gas chromatography at intermediate temperatures. <i>Journal of Chromatography A</i> , 2008 , 1195, 136-45	4-5	49
653	Investigation of retention behavior of drug molecules in supercritical fluid chromatography using linear solvation energy relationships. <i>Journal of Chromatography A</i> , 2008 , 1206, 186-95	4-5	32
652	Enthalpy of solvation correlations for gaseous solutes dissolved in dibutyl ether and ethyl acetate. 2008 , 470, 67-76		29

651	Micellar versus hydro-organic reversed-phase liquid chromatography: a solvation parameter-based perspective. <i>Journal of Chromatography A</i> , 2008 , 1182, 176-96	4.5	23
650	Orthogonal screening system of columns for supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2008 , 1203, 105-13	4.5	62
649	System Maps for RP-LC on an Octadecylsiloxane-Bonded Silica Stationary Phase (SunFire C18). <i>Chromatographia</i> , 2008 , 68, 11-17	2.1	26
648	Comparison of the Separation Characteristics of the Organic/Inorganic Hybrid Stationary Phases XBridge C8 and Phenyl and XTerra Phenyl in RP-LC. <i>Chromatographia</i> , 2008 , 68, 491-500	2.1	23
647	Hydrogen-bond acidic polymers for chemical vapor sensing. 2008 , 108, 726-45		180
646	Interactions between spiropyrans and room-temperature ionic liquids: photochromism and solvatochromism. 2008 , 112, 7530-6		109
645	Absorption or adsorption? Insights from molecular probes n-alkanes and cycloalkanes into modes of sorption by environmental solid matrices. <i>Environmental Science & Technology</i> , 2008 , 42, 3989-95 ^{10.3}		34
644	Concentration-detection functions for the odor of homologous n-acetate esters. 2008 , 95, 658-67		35
643	Modeling and prediction (correction) of partition coefficients of bile acids and their derivatives by multivariate regression methods. <i>Talanta</i> , 2008 , 75, 651-7	6.2	28
642	Evaluating coal tar-water partitioning coefficient estimation methods and solute-solvent molecular interactions in tar phase. 2008 , 73, 532-8		22
641	Solvation parameters for mercury and mercury(II) compounds: calculation of properties of environmental interest. 2008 , 10, 435-42		12
640	Computation methodology for determining Abraham solute descriptors from limited experimental data by combining Abraham model and Goss-modified Abraham model correlations. <i>Physics and Chemistry of Liquids</i> , 2008 , 46, 574-585	1.5	17
639	Compound-specific factors influencing sorption nonlinearity in natural organic matter. <i>Environmental Science & Technology</i> , 2008 , 42, 5897-903	10.3	38
638	Correlation and Prediction of Partition Coefficients From the Gas Phase and from Water to Alkan-1-ols. <i>Industrial & Engineering Chemistry Research</i> , 2008 , 47, 3990-3995	3.9	42
637	Partitioning of polynuclear aromatic hydrocarbons into water from biodiesel fuel mixtures. 2008 , 5, 435		5
636	Selectivity of calixarene-bonded silica-phases in HPLC: description of special characteristics with a multiple term linear equation at two different pH-values. 2008 , 24, 1157-64		9
635	Characterisation of room temperature ionic liquid chromatographic stationary phases by combining experimental retention factor and partition coefficient data into a single model. <i>Physics and Chemistry of Liquids</i> , 2009 , 47, 74-83	1.5	9
634	Prediction of biomagnification factors for some organochlorine compounds using linear free energy relationship parameters and artificial neural networks. 2009 , 20, 453-65		9

633	Chapter 3 Theoretical Considerations. 2009 , 55, 49-76		1
632	A prediction model for blood-brain barrier permeation and analysis on its parameter biologically. 2009 , 95, 280-7		31
631	Chiral recognition mechanisms with macrocyclic glycopeptide selectors. 2009 , 21, 167-75		63
630	Modelling of partition constants: linear solvation energy relationships or PLS regression?. 2009 , 23, 254-262		6
629	Etched succinate-functionalized silica hydride stationary phase for open-tubular CEC. 2009 , 30, 3855-62		16
628	Exploring enantioselective molecular recognition mechanisms with chemoinformatic techniques. 2009 , 32, 1566-84		43
627	Prediction of immobilized artificial membrane-liquid chromatography retention of some drugs from their molecular structure descriptors and LFER parameters. 2009 , 32, 3395-402		4
626	Artificial neural network modelling of retention of pesticides in various octadecylsiloxane-bonded reversed-phase columns and water-acetonitrile mobile phase. 2009 , 646, 47-61		11
625	Prediction of convulsant activity of gases and vapors. 2009 , 44, 885-90		6
624	Linear Free Energy Relationship Correlation of The Distribution of Solutes Between Water And Cetytrimethylammonium Bromide (CTAB) Micelles. 2009 , 28, 72-88		26
623	Enthalpy of solvation correlations for organic solutes and gases dissolved in N,N-dimethylformamide and tert-butanol. 2009 , 144, 23-31		28
622	Enthalpy of solvation correlations for organic solutes and gases dissolved in acetonitrile and acetone. 2009 , 484, 65-69		28
621	Solvation parameter model of comprehensive two-dimensional gas chromatography separations. <i>Journal of Chromatography A</i> , 2009 , 1216, 1650-7	4-5	34
620	Retention mechanism of high-performance liquid chromatographic enantioseparation on macrocyclic glycopeptide-based chiral stationary phases. <i>Journal of Chromatography A</i> , 2009 , 1216, 1845-60	4-5	91
619	Retention models for isocratic and gradient elution in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2009 , 1216, 1737-55	4-5	114
618	Partitioning behaviour of organic compounds between ionic liquids and supercritical fluids. <i>Journal of Chromatography A</i> , 2009 , 1216, 1861-80	4-5	52
617	Inverse gas chromatography as a source of physiochemical data. <i>Journal of Chromatography A</i> , 2009 , 1216, 1551-66	4-5	189
616	Application of the solvation parameter model in method development for analysis of residual solvents in pharmaceuticals. <i>Journal of Chromatography A</i> , 2009 , 1216, 86-91	4-5	3

615	Extension of the system constants database for open-tubular columns: system maps at low and intermediate temperatures for four new columns. <i>Journal of Chromatography A</i> , 2009 , 1216, 1640-9	4.5	30
614	Theoretical characterization of gas-liquid chromatographic stationary phases with quantum chemical descriptors. <i>Journal of Chromatography A</i> , 2009 , 1216, 2540-7	4.5	12
613	Partition coefficients of organic compounds in new imidazolium based ionic liquids using inverse gas chromatography. <i>Journal of Chromatography A</i> , 2009 , 1216, 4775-86	4.5	73
612	Characterization of bipartitioning micellar chromatography system using monolithic column by linear solvation energy relationship and application to predict blood-brain barrier penetration. <i>Journal of Chromatography A</i> , 2009 , 1216, 5190-8	4.5	20
611	Possibility of predicting separations in supercritical fluid chromatography with the solvation parameter model. <i>Journal of Chromatography A</i> , 2009 , 1216, 5600-7	4.5	23
610	Prediction of milk/plasma concentration ratios of drugs and environmental pollutants. 2009 , 44, 2452-8		22
609	Correlation and prediction of solute transfer to chloroalkanes from both water and the gas phase. 2009 , 281, 144-162		45
608	Development of Abraham model correlations for solvation characteristics of linear alcohols. 2009 , 286, 170-174		47
607	Beyond physical properties--application of Abraham descriptors and LFER analysis in agrochemical research. 2009 , 17, 4153-9		25
606	Determination of solute descriptors by chromatographic methods. 2009 , 652, 32-53		187
605	Strategy for reduced calibration sets to develop quantitative structure-retention relationships in high-performance liquid chromatography. 2009 , 652, 180-8		19
604	Determination of descriptors for semivolatile organosilicon compounds by gas chromatography and non-aqueous liquid-liquid partition. <i>Journal of Chromatography A</i> , 2009 , 1216, 7882-8	4.5	57
603	The influence of the eluent composition on the retention of derivatives of some aromatic heterocyclic compounds in reversed-phase high-performance liquid chromatography. 2009 , 83, 1223-1229		4
602	LFERs for soil organic carbon-water distribution coefficients (K _{oc}) at environmentally relevant sorbate concentrations. <i>Environmental Science & Technology</i> , 2009 , 43, 3094-100	10.3	60
601	Linear Free Energy Relationship Correlations for Room Temperature Ionic Liquids: Revised Cation-Specific and Anion-Specific Equation Coefficients for Predictive Applications Covering a Much Larger Area of Chemical Space. <i>Industrial & Engineering Chemistry Research</i> , 2009 , 48, 4145-4154	3.9	64
600	Distribution of organic solutes in biphasic 1-n-butyl-3-methylimidazolium methyl sulfate-supercritical CO ₂ system. 2009 , 113, 9520-6		16
599	Activity coefficients at infinite dilution of organic solutes in the ionic liquid 1-ethyl-3-methyl-imidazolium nitrate. 2009 , 113, 4323-32		47
598	Environmentally Relevant Properties of All 209 Polychlorinated Biphenyl Congeners for Modeling Their Fate in Different Natural and Climatic Conditions. 2009 , 54, 1189-1213		37

597	Important role of reaction field in photodegradation of deca-bromodiphenyl ether: theoretical and experimental investigations of solvent effects. 2009 , 76, 1486-90		51
596	Partition of compounds from water and from air into amides. 2009 , 33, 2034-2043		53
595	The hydrogen bond acidity and other descriptors for oximes. 2009 , 33, 76-81		11
594	Systematic investigation of the sorption properties of Tenax TA, Chromosorb 106, Porapak N, and Carbpak F. 2009 , 81, 3017-21		18
593	Bibliography. 2009 , 1-241		1
592	The partition of compounds from water and from air into wet and dry ketones. 2009 , 33, 568-573		68
591	Partition of compounds from water and from air into the wet and dry monohalobenzenes. 2009 , 33, 1685		38
590	Models for the sorption of volatile organic compounds by diesel soot and atmospheric aerosols. 2009 , 11, 815-22		11
589	Factors Affecting the Interpretation of Selectivity on Synergi Reversed-Phase Columns. <i>Chromatographia</i> , 2010 , 71, 185-193	2.1	18
588	Trends in the adsorption of volatile organic compounds in a large-pore metal-organic framework, IRMOF-1. 2010 , 26, 11319-29		67
587	Retention and selectivity properties of carbamate pesticides on novel polar-embedded stationary phases. 2010 , 397, 2513-24		13
586	Development of Abraham model correlations for solvation characteristics of secondary and branched alcohols. 2010 , 288, 121-127		41
585	Comment on Development of Correlations for Describing Solute Transfer into Acyclic Alcohol Solvents Based on the Abraham Model and Fragment-Specific Equation Coefficients 2010 , 291, 212-213		4
584	Partitioning of organochlorine pesticides from water to polyethylene passive samplers. 2010 , 158, 2511-7		73
583	Modeling and predicting competitive sorption of organic compounds in soil. 2010 , 29, 2676-84		8
582	Multiple-column RP-HPLC retention modelling based on solvatochromic or theoretical solute descriptors. 2010 , 33, 155-66		19
581	Model for the partition of neutral compounds between n-heptane and formamide. 2010 , 33, 1167-73		21
580	Characterization of calixarene-bonded stationary phases. 2010 , 33, 2930-42		8

579	Selectivity of calixarene-bonded silica phases in HPLC: Description of special characteristics with a multiple term linear equation at different methanol concentrations. 2010 , 33, 2943-55		5
578	Prediction of solubility of drugs and other compounds in organic solvents. 2010 , 99, 1500-15		217
577	Water-solvent partition coefficients and Delta Log P values as predictors for blood-brain distribution; application of the Akaike information criterion. 2010 , 99, 2492-501		17
576	Extraction of organic compounds with room temperature ionic liquids. <i>Journal of Chromatography A</i> , 2010 , 1217, 2268-86	4-5	392
575	Solute-solvent interactions in micellar electrokinetic chromatography: VII. Characterization of sodium cholate-sodium deoxycholate mixed-micellar systems. <i>Journal of Chromatography A</i> , 2010 , 1217, 1701-8	4-5	7
574	Evaluation of ternary mobile phases for reversed-phase liquid chromatography: effect of composition on retention mechanism. <i>Journal of Chromatography A</i> , 2010 , 1217, 5957-64	4-5	15
573	Partition coefficients of macrocyclic crown ethers in water-organic biphasic system: DFT/COSMO-RS approach. 2010 , 288, 111-120		14
572	Development of correlations for describing solute transfer into acyclic alcohol solvents based on the Abraham model and fragment-specific equation coefficients. 2010 , 288, 139-144		23
571	Mathematical correlations for describing solute transfer into functionalized alkane solvents containing hydroxyl, ether, ester or ketone solvents. 2010 , 298, 48-53		28
570	Activity coefficients at infinite dilution of organic solutes in the ionic liquid 1-ethyl-3-methylimidazolium methanesulfonate. 2010 , 299, 198-206		37
569	. 2010 ,		3
568	Memory-efficient fully coupled filtering approach for observational model building. 2010 , 21, 680-6		2
567	Chromatography in the Millennium Perspectives. 2010 , 363-395		
566	A simple QSPR model to predict aqueous solubility of drugs. 2010 , 20, 467-476		14
565	System selectivity cube: a 3D visualization tool for comparing the selectivity of gas chromatography, supercritical-fluid chromatography, high-pressure liquid chromatography, and micellar electrokinetic capillary chromatography systems. 2010 , 82, 6251-8		7
564	Equations for the transfer of neutral molecules and ionic species from water to organic phases. 2010 , 75, 1006-15		102
563	Solute descriptors for phenoxide anions and their use to establish correlations of rates of reaction of anions with iodomethane. 2010 , 75, 3021-6		57
562	Hydrogen bonding between solutes in solvents octan-1-ol and water. 2010 , 75, 7651-8		8

561	Characterization of Adsorption Mechanisms of Volatile Organic Compounds with Montmorillonite at Different Levels of Relative Humidity via a Linear Solvation Energy Relationship Approach. 2010 , 55, 5766-5770		9
560	Estimating the hydrogen bond energy. 2010 , 114, 9529-36		234
559	The biological and toxicological activity of gases and vapors. 2010 , 24, 357-62		27
558	Predicting solvation energies for kinetic modeling. 2010 , 106, 211		44
557	Chiral Recognition Mechanisms in Enantiomers Separations: A General View. 2010 , 1-32		13
556	Comment on "Important role of reaction field in photodegradation of deca-bromodiphenyl ether: Theoretical and experimental investigations of solvent effects" by Q. Xie, J. Chen, J. Shao, C. Chen, H. Zhao, C. Hao [Chemosphere 76(11) (2009) 1486-1490]. 2010 , 80, 676-8; author reply 679-80		1
555	QSAR in Chromatography: Quantitative StructureRetention Relationships (QSRRs). 2010 , 223-259		4
554	Recent Advances in QSAR Studies. 2010 ,		82
553	Comparing linear free energy relationships for organic chemicals in soils: effects of soil and solute properties. <i>Environmental Science & Technology</i> , 2010 , 44, 6971-7	10.3	12
552	The transfer of neutral molecules, ions and ionic species from water to wet octanol. 2010 , 12, 13182-8		57
551	The transfer of neutral molecules, ions and ionic species from water to ethylene glycol and to propylene carbonate; descriptors for pyridinium cations. 2010 , 34, 2298		64
550	Infinite-Dilution Activity Coefficients of Volatile Organic Compounds in Cyclic Olefin Copolymers. 2011 , 56, 1255-1260		1
549	The lipophilicity and hydrogen bond strength of pyridine-N-oxides and protonated pyridine-N-oxides. 2011 , 35, 930		22
548	Predicting sorption of pesticides and other multifunctional organic chemicals to soil organic carbon. <i>Environmental Science & Technology</i> , 2011 , 45, 1313-9	10.3	83
547	Equilibrium partition coefficients of diverse polar and nonpolar organic compounds to polyoxymethylene (POM) passive sampling devices. <i>Environmental Science & Technology</i> , 2011 , 45, 10124-32	10.3	80
546	Serum albumin binding of structurally diverse neutral organic compounds: data and models. 2011 , 24, 2293-301		91
545	Development and use of polyethylene passive samplers to detect triclosans and alkylphenols in an urban estuary. <i>Environmental Science & Technology</i> , 2011 , 45, 2270-7	10.3	48
544	Interactions of Volatile Organic Compounds with the Ionic Liquid 1-Butyl-1-methylpyrrolidinium Dicyanamide. 2011 , 56, 4909-4918		42

543	Linear Solvation Energy Relationships Used To Evaluate Sorption Mechanisms of Volatile Organic Compounds with One Organomontmorillonite under Different Humidities. 2011 , 56, 4950-4955		9
542	A systematic investigation of quaternary ammonium ions as asymmetric phase-transfer catalysts. Application of quantitative structure activity/selectivity relationships. 2011 , 76, 4337-57		93
541	Models for liquid-liquid partition in the system formamide-organic solvent and their use for estimating descriptors for organic compounds. <i>Talanta</i> , 2011 , 83, 1118-25	6.2	20
540	Quality Control in Pharmaceuticals: Residual Solvents Testing and Analysis. 2011 ,		1
539	Multivariate visualization of chromatographic systems. 2011 ,		2
538	The transfer of neutral molecules, ions and ionic species from water to benzonitrile; comparison with nitrobenzene. 2011 , 526, 22-28		27
537	Thermodynamics and activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid 1-butyl-1-methylpyrrolidinium tetracyanoborate. 2011 , 43, 1810-1817		74
536	Abraham model correlations for solute partitioning into o-xylene, m-xylene and p-xylene from both water and the gas phase. 2011 , 308, 64-71		21
535	Estimation of Hansen solubility parameters using multivariate nonlinear QSPR modeling with COSMO screening charge density moments. 2011 , 309, 8-14		47
534	Solubility and rate of dissolution for Miscanthus in hydrophilic ionic liquids. 2011 , 309, 89-96		41
533	Abraham model correlations for transfer of neutral molecules and ions to sulfolane. 2011 , 309, 30-35		22
532	PTR-MS measurements and analysis of models for the calculation of Henry's law constants of monosulfides and disulfides. 2011 , 83, 311-7		22
531	Supercritical fluid chromatography in research laboratories: design, development and implementation of an efficient generic screening for exploiting this technique in the achiral environment. <i>Journal of Chromatography A</i> , 2011 , 1218, 8551-60	4.5	39
530	Linear Solvation Energy Relationships as classifiers in non-target analysis--a capillary liquid chromatography approach. <i>Journal of Chromatography A</i> , 2011 , 1218, 8192-6	4.5	29
529	Ionic liquids: predictions of physicochemical properties with experimental and/or DFT-calculated LFER parameters to understand molecular interactions in solution. 2011 , 115, 6040-50		49
528	Partition coefficients of ketones, phenols, aliphatic and aromatic acids, and esters in n-hexane/nitromethane. 2011 , 9, 813-824		3
527	Regularities of the sorption of cycloalkenyl-substituted thiophenes and 2,2'-bithiophenes from water-acetonitrile solutions on hexadecyl silica gel under conditions of high-performance liquid chromatography. 2011 , 85, 1440-1449		3
526	Partition Coefficients and Solubilities of Compounds in the Water-Ethanol Solvent System. <i>Journal of Solution Chemistry</i> , 2011 , 40, 1279-1290	1.8	45

525	Correlation of the Solubilizing Abilities of Hexyl(trimethyl)ammonium bis((Trifluoromethyl)sulfonyl)imide, 1-Propyl-1-methylpiperidinium bis((Trifluoromethyl)sulfonyl)imide, and 1-Butyl-1-methyl-pyrrolidinium Thiocyanate. <i>Journal of Solution Chemistry</i> , 2011 , 40, 2000-2022	1.8	24
524	Models for Liquid-Liquid Partition in the System Ethylene Glycol/Organic Solvent and Their Use for Estimating Descriptors for Organic Compounds. <i>Chromatographia</i> , 2011 , 73, 941-951	2.1	25
523	Separation Characteristics of Ionic Liquids Grafted Polymethylsiloxanes Stationary Phases for Capillary GC. <i>Chromatographia</i> , 2011 , 74, 717-724	2.1	14
522	Predicting the partitioning of biological compounds between room-temperature ionic liquids and water by means of the solvation-parameter model. 2011 , 399, 2807-20		16
521	Diffusion coefficients in ethanol and in water at 298 K: Linear free energy relationships. 2011 , 303, 45-55		53
520	Comparison of theoretical and experimental models for characterizing solvent properties using reversed phase liquid chromatography. 2011 , 702, 37-44		9
519	Enthalpy of solvation correlations for organic solutes and gases dissolved in 1-propanol and tetrahydrofuran. 2011 , 519, 103-113		21
518	Prediction models for the Abraham hydrogen bond donor strength: comparison of semi-empirical, ab initio, and DFT methods. 2011 , 24, 1072-1080		25
517	Ionic liquid stationary phases for gas chromatography. 2011 , 34, 888-900		173
516	The permeation of neutral molecules, ions, and ionic species through membranes: brain permeation as an example. 2011 , 100, 1690-701		43
515	Linear free energy relationship analysis of retention factors in cerasome electrokinetic chromatography intended for predicting drug skin permeation. 2011 , 100, 3105-3113		26
514	Chromatographic selectivity triangles. <i>Journal of Chromatography A</i> , 2011 , 1218, 556-86	4.5	39
513	Insights into chiral recognition mechanisms in supercritical fluid chromatography. I. Non-enantiospecific interactions contributing to the retention on tris-(3,5-dimethylphenylcarbamate) amylose and cellulose stationary phases. <i>Journal of Chromatography A</i> , 2011 , 1218, 2019-32	4.5	40
512	High temperature and highly selective stationary phases of ionic liquid bonded polysiloxanes for gas chromatography. <i>Journal of Chromatography A</i> , 2011 , 1218, 833-41	4.5	38
511	Fundamental challenges and opportunities for preparative supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2011 , 1218, 1037-114	4.5	179
510	Models for liquid-liquid partition in the system propylene carbonate-organic solvent and their use for estimating descriptors for organic compounds. <i>Journal of Chromatography A</i> , 2011 , 1218, 809-16	4.5	28
509	Characterization of new R-naphthylethyl cyclodextrin 6 chiral stationary phase and its comparison with R-naphthylethyl β -cyclodextrin-based column. <i>Journal of Chromatography A</i> , 2011 , 1218, 1393-8	4.5	34
508	Retention mechanism of a cholesterol-coated C18 stationary phase: van't Hoff and Linear Solvation Energy Relationships (LSER) approaches. <i>Journal of Chromatography A</i> , 2011 , 1218, 2936-43	4.5	15

507	Investigations on the chromatographic behaviour of zwitterionic stationary phases used in hydrophilic interaction chromatography. <i>Journal of Chromatography A</i> , 2011 , 1218, 5939-63	4.5	134
506	Chromatographic characterization of hydrophilic interaction liquid chromatography stationary phases: hydrophilicity, charge effects, structural selectivity, and separation efficiency. <i>Journal of Chromatography A</i> , 2011 , 1218, 5903-19	4.5	144
505	Multi-variable retention modelling in reversed-phase high-performance liquid chromatography based on the solvation method: a comparison between curvilinear and artificial neural network regression. 2011 , 690, 35-46		13
504	Models for liquid-liquid partition in the system dimethyl sulfoxide-organic solvent and their use for estimating descriptors for organic compounds. <i>Journal of Chromatography A</i> , 2011 , 1218, 4525-36	4.5	18
503	Activity coefficients at infinite dilution measurements for organic solutes and water in the ionic liquid 1-ethyl-3-methylimidazolium tetracyanoborate. 2011 , 43, 1050-1057		93
502	Effect-Directed Analysis of Complex Environmental Contamination. <i>Handbook of Environmental Chemistry</i> , 2011 ,	0.8	12
501	Sorption of Anthropogenic Organic Compounds to Airborne Particles. 2011 , 113-147		
500	Experimental and predicted solubilities of 3,4-dichlorobenzoic acid in select organic solvents and in binary aqueous-ethanol mixtures. <i>Physics and Chemistry of Liquids</i> , 2012 , 50, 324-335	1.5	30
499	Determination of Abraham model solute descriptors for benzoin based on measured solubility ratios. <i>Physics and Chemistry of Liquids</i> , 2012 , 50, 254-265	1.5	31
498	An algorithm for 353 odor detection thresholds in humans. 2012 , 37, 207-18		44
497	Activity Coefficients at Infinite Dilution for Organic Compounds Dissolved in 1-Alkyl-1-methylpyrrolidinium Bis(trifluoromethylsulfonyl)imide Ionic Liquids Having Six-, Eight-, and Ten-Carbon Alkyl Chains. 2012 , 57, 3510-3518		59
496	Partial solvation parameters and mixture thermodynamics. 2012 , 116, 7302-21		36
495	Determination of Descriptors for Plasticizers by Chromatography and Liquid-Liquid Partition. <i>Chromatographia</i> , 2012 , 75, 1135-1146	2.1	18
494	Correlation of the Solubilizing Abilities of 1-Butyl-1-methylpiperidinium Bis(trifluoromethylsulfonyl)imide and 1-Butyl-1-methylpyrrolidinium Tetracyanoborate. <i>Journal of Solution Chemistry</i> , 2012 , 41, 1165-1184	1.8	24
493	Determination of partition coefficients of refrigerants by gas liquid chromatographic headspace analysis. <i>Journal of Chromatography A</i> , 2012 , 1265, 144-8	4.5	2
492	Compounds for expanding the descriptor space for characterizing separation systems. <i>Journal of Chromatography A</i> , 2012 , 1266, 124-30	4.5	20
491	Easy and accurate calculation of programmed temperature gas chromatographic retention times by back-calculation of temperature and hold-up time profiles. <i>Journal of Chromatography A</i> , 2012 , 1263, 179-88	4.5	15
490	Linear solvation energy relationships as classifiers in non-target analysis - a gas chromatographic approach. <i>Journal of Chromatography A</i> , 2012 , 1264, 95-103	4.5	8

489	Partitioning of neutral organic compounds to structural proteins. <i>Environmental Science & Technology</i> , 2012 , 46, 12697-703	10.3	64
488	The hydrogen bond properties of water from 273 K to 573 K; equations for the prediction of gas-water partition coefficients. 2012 , 14, 7433-40		13
487	Linear free-energy relationships for water/hexadec-1-ene and water/deca-1,9-diene partitions, and for permeation through lipid bilayers; comparison of permeation systems. 2012 , 36, 1798		23
486	Sorption interactions of volatile organic compounds with organoclays under different humidities by using linear solvation energy relationships. 2012 , 18, 329-336		10
485	Solute Partitioning Between 1-n-Butyl-3-methylimidazolium Trifluoromethanesulfonate Ionic Liquid and Supercritical CO ₂ . 2012 , 57, 1064-1071		7
484	Experimentally determined soil organic matter-water sorption coefficients for different classes of natural toxins and comparison with estimated numbers. <i>Environmental Science & Technology</i> , 2012 , 46, 6118-26	10.3	34
483	Salting-out effect in aqueous NaCl solutions: trends with size and polarity of solute molecules. <i>Environmental Science & Technology</i> , 2012 , 46, 1496-503	10.3	151
482	Stationary phases for packed-column supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2012 , 1250, 157-71	4.5	68
481	Characterization and use of hydrophilic interaction liquid chromatography type stationary phases in supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2012 , 1250, 182-95	4.5	62
480	Multi-component trace analysis of organic xenobiotics in surface water containing suspended particular matter by solid phase extraction/gas chromatography-mass spectrometry. <i>Journal of Chromatography A</i> , 2012 , 1249, 181-9	4.5	19
479	Inverse gas chromatography and partial solvation parameters. <i>Journal of Chromatography A</i> , 2012 , 1251, 194-207	4.5	17
478	A principal component analysis approach for developing retention models in liquid chromatography. <i>Journal of Chromatography A</i> , 2012 , 1251, 134-140	4.5	5
477	Gas solvent and water solvent partition coefficients of the tetraphenyl compounds of group (IV). 2012 , 36, 626-631		5
476	Principles and Practice of Solid-Phase Extraction. 2012 , 273-297		6
475	Classification and Selection of Open-Tubular Columns for Analytical Separations. 2012 , 137-159		4
474	Chemometric methods to classify stationary phases for achiral packed column supercritical fluid chromatography. 2012 , 26, 52-65		26
473	Solvent classification for chromatography and extraction. <i>Journal of Planar Chromatography - Modern TLC</i> , 2012 , 25, 190-199	0.9	26
472	Influence of Solvent Nature on the Solubility of Halogenated Alkanes. 2012 , 57, 1945-1952		1

471	Equations for the Partition of Neutral Molecules, Ions and Ionic Species from Water to Water-Ethanol Mixtures. <i>Journal of Solution Chemistry</i> , 2012 , 41, 730-740	1.8	31
470	Occurrence of residual water within disk-based solid-phase extraction and its effect on GC-MS measurement of organic extracts of environmental samples. 2012 , 403, 2541-52		8
469	Stationary phase selection and comprehensive two-dimensional gas chromatographic analysis of trace biodiesel in petroleum-based fuel. <i>Journal of Chromatography A</i> , 2012 , 1226, 103-9	4.5	26
468	Determination of descriptors for fragrance compounds by gas chromatography and liquid-liquid partition. <i>Journal of Chromatography A</i> , 2012 , 1235, 159-65	4.5	27
467	Study of chemical selectivity of molecular binary mixed micelles of sodium 10-undecenyl sulfate and sodium N-undecenyl leucinate using linear solvation energy relationships model. <i>Journal of Chromatography A</i> , 2012 , 1236, 207-14	4.5	2
466	Desorption kinetics with PTR-MS: Isothermal differential desorption kinetics from a heterogeneous inlet surface at ambient pressure and a new concept for compound identification. 2012 , 314, 33-41		7
465	The effect of ionized species on microsomal binding. 2012 , 47, 202-5		35
464	Interactions of volatile organic compounds with the ionic liquid 1-ethyl-3-methylimidazolium tetracyanoborate. 2012 , 47, 100-108		33
463	Determination of LFER descriptors of 30 cations of ionic liquids--progress in understanding their molecular interaction potentials. 2012 , 13, 780-7		11
462	Correlation of the Solubilizing Abilities of 1-Butyl-1-methyl-pyrrolidinium Tris(pentafluoroethyl)trifluorophosphate, 1-Butyl-1-methylpyrrolidinium Triflate and 1-Methoxyethyl-1-methylmorpholinium Tris(pentafluoroethyl)trifluorophosphate. <i>Journal of Solution Chemistry</i> , 2013 , 42, 772-799	1.8	20
461	Quantitative structure/eluent-retention relationships in reversed-phase high-performance liquid chromatography based on the solvatochromic method. 2013 , 405, 755-66		1
460	On the solubility of nicotinic acid and isonicotinic acid in water and organic solvents. 2013 , 61, 74-78		9
459	Estimation of extraction properties of new imidazolidine anion based ionic liquids on the basis of activity coefficient at infinite dilution measurements. 2013 , 118, 242-254		33
458	Evaluation of Triethylamine as a Counter Solvent in Totally Organic Biphasic Liquid-Liquid Partition Systems. <i>Chromatographia</i> , 2013 , 76, 1031-1039	2.1	11
457	Physicochemical properties and activity coefficients at infinite dilution for organic solutes and water in a novel bicyclic guanidinium superbases-derived protic ionic liquid. 2013 , 58, 62-69		31
456	Evaluation of the suitability of chromatographic systems to predict human skin permeation of neutral compounds. <i>European Journal of Pharmaceutical Sciences</i> , 2013 , 50, 557-68	5.1	16
455	IUPAC-NIST Solubility Data Series. 99. Solubility of Benzoic Acid and Substituted Benzoic Acids in Both Neat Organic Solvents and Organic Solvent Mixtures. <i>Journal of Physical and Chemical Reference Data</i> , 2013 , 42, 033103	4.3	6
454	Determination of solvation descriptors for terpene hydrocarbons from chromatographic measurements. <i>Journal of Chromatography A</i> , 2013 , 1293, 133-41	4.5	20

453	Characterization of stationary phases based on polysiloxanes thermally immobilized onto silica and metalized silica using supercritical fluid chromatography with the solvation parameter model. <i>Journal of Chromatography A</i> , 2013 , 1315, 176-87	4.5	13
452	Quantitative Structure Property (Retention) Relationships in Liquid Chromatography. 2013 , 385-405		3
451	Estimation of the environmental properties of compounds from chromatographic measurements and the solvation parameter model. <i>Journal of Chromatography A</i> , 2013 , 1317, 85-104	4.5	120
450	Comparative characterization of hydrophilic interaction liquid chromatography columns by linear solvation energy relationships. <i>Journal of Chromatography A</i> , 2013 , 1273, 73-94	4.5	71
449	"Retention projection" enables reliable use of shared gas chromatographic retention data across laboratories, instruments, and methods. 2013 , 85, 11650-7		18
448	Application of the LSER method for investigation of intermolecular interactions of porous polymer surface with organic molecules. 2013 , 49, 639-641		4
447	Main interactions and influences of the chromatographic parameters in HILIC separations. 2013 , 51, 684-93		99
446	Descriptors for the Prediction of Partition Coefficients and Solubilities of Organophosphorus Compounds. 2013 , 48, 884-897		29
445	Activity Coefficients at Infinite Dilution for Organic Solutes Dissolved in Three 1-Alkyl-1-methylpyrrolidinium Bis(trifluoromethylsulfonyl)imide Ionic Liquids Bearing Short Linear Alkyl Side Chains of Three to Five Carbons. 2013 , 58, 2210-2218		61
444	Prediction of chromatographic retention time in high-resolution anti-doping screening data using artificial neural networks. 2013 , 85, 10330-7		49
443	Models for Liquid-Liquid Partition in the System Ethanolamine-Organic Solvent and Their Use for Estimating Descriptors for Organic Compounds. <i>Chromatographia</i> , 2013 , 76, 157-164	2.1	13
442	A critique of Abraham and Acree's correlation for deca-1,9-diene/water partition coefficients. 2013 , 37, 283-285		2
441	Totally organic biphasic solvent systems for extraction and descriptor determinations. 2013 , 36, 96-109		30
440	Mobile Phases and Their Properties. 2013 , 363-447		6
439	HPLC Analysis. 2013 , 465-519		2
438	A correlation for 1,9-decadiene/water partition coefficients. 2013 , 102, 136-44		12
437	GC for flavonoids analysis: Past, current, and prospective trends. 2013 , 36, 20-36		31
436	Study of organic compounds-water interactions by partition in aqueous two-phase systems. <i>Journal of Chromatography A</i> , 2013 , 1322, 97-104	4.5	16

435	Analysis of immobilized artificial membrane retention factors for both neutral and ionic species. <i>Journal of Chromatography A</i> , 2013 , 1298, 44-9	4.5	25
434	The use of biopartitioning micellar chromatography and immobilized artificial membrane column for in silico and in vitro determination of blood-brain barrier penetration of phenols. <i>Journal of Chromatography A</i> , 2013 , 1286, 127-36	4.5	13
433	COSMO-RS for the prediction of the retention behavior in micellar liquid chromatography based on partition coefficients of non-dissociated and dissociated solutes. <i>Journal of Chromatography A</i> , 2013 , 1273, 66-72	4.5	9
432	Characterization of five chemistries and three particle sizes of stationary phases used in supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2013 , 1319, 148-59	4.5	63
431	Quantitative Structure Retention Relationships as Useful Tool to Characterize Chromatographic Systems and Their Potential to Simulate Biological Processes. <i>Chromatographia</i> , 2013 , 76, 211-226	2.1	23
430	Polymer-polymer miscibility and partial solvation parameters. 2013 , 54, 1621-1638		31
429	An extensible framework for capturing solvent effects in computer generated kinetic models. 2013 , 117, 2955-70		23
428	The Determination of Abraham Descriptors and Their Application to Crop Protection Research. 2013 , 273-305		11
427	An insight into the use of dimethylphenyl carbamate cyclofructan 7 chiral stationary phase in supercritical fluid chromatography: the basic comparison with HPLC. 2013 , 36, 1711-9		28
426	Application of Abraham solvation parameters for surface tension prediction of mono-solvents and solvent mixtures at various temperatures. 2013 , 178, 44-56		4
425	General model for estimating partition coefficients to organisms and their tissues using the biological compositions and polyparameter linear free energy relationships. <i>Environmental Science & Technology</i> , 2013 , 47, 6630-9	10.3	89
424	Effects of mobile phase composition on retention and selectivity in achiral supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2013 , 1302, 152-62	4.5	65
423	Infinite dilution activity coefficients of volatile organic compounds in two ionic liquids composed of the tris(pentafluoroethyl)trifluorophosphate ([FAP]) anion and a functionalized cation. 2013 , 65, 53-64		11
422	Characterisation of capillary ionic liquid columns for gas chromatography-mass spectrometry analysis of fatty acid methyl esters. 2013 , 803, 166-73		65
421	Hydration Free Energies of Multifunctional Nitroaromatic Compounds. 2013 , 9, 2774-85		18
420	Quantitative Structure Retention Relationship Models in an Analytical Quality by Design Framework: Simultaneously Accounting for Compound Properties, Mobile-Phase Conditions, and Stationary-Phase Properties. <i>Industrial & Engineering Chemistry Research</i> , 2013 , 52, 12269-12284	3.9	21
419	Interactions of volatile organic compounds with the ionic liquids 1-butyl-1-methylpyrrolidinium tetracyanoborate and 1-butyl-1-methylpyrrolidinium bis(oxalato)borate. 2013 , 57, 344-354		23
418	Experimental determination of polyparameter linear free energy relationship (pp-LFER) substance descriptors for pesticides and other contaminants: new measurements and recommendations. <i>Environmental Science & Technology</i> , 2013 , 47, 14204-14	10.3	54

4 ¹⁷	Polyethylene-water partitioning coefficients for parent- and alkylated-polycyclic aromatic hydrocarbons and polychlorinated biphenyls. <i>Environmental Science & Technology</i> , 2013 , 47, 6943-50 ^{10.3}	48
4 ¹⁶	Interaction Mechanisms and Predictive Model for the Sorption of Aromatic Compounds onto Nonionic Resins. 2013 , 117, 17707-17715	23
4 ¹⁵	Summation Solute Hydrogen Bonding Acidity Values for Hydroxyl Substituted Flavones Determined by NMR Spectroscopy. 2013 , 8, 1934578X1300800	1
4 ¹⁴	Physical Ionic Liquid/Polysiloxane Mixtures for Tuning the Polarity and the Selectivity of the Polysiloxane Stationary Phase for GC Analysis. <i>Chromatographia</i> , 2014 , 77, 1671-1681	2.1 3
4 ¹³	Abraham model correlations describing the solubilising ability of peanut oil. <i>Physics and Chemistry of Liquids</i> , 2014 , 52, 792-803	1.5 5
4 ¹²	Withdrawn: Using multidimensional gas chromatography to define a carbon number-functionality grid for characterizing oxidation of organic aerosol. 2014 ,	
4 ¹¹	Correlations between chromatographic parameters and bioactivity predictors of potential herbicides. 2014 , 52, 676-84	5
4 ¹⁰	Sorption mechanism and predictive models for removal of cationic organic contaminants by cation exchange resins. <i>Environmental Science & Technology</i> , 2014 , 48, 14572-81	10.3 10
4 ⁰⁹	What experimental factors influence the accuracy of retention projections in gas chromatography-mass spectrometry?. <i>Journal of Chromatography A</i> , 2014 , 1373, 179-89	4.5 5
4 ⁰⁸	Insights into chiral recognition mechanisms in supercritical fluid chromatography V. Effect of the nature and proportion of alcohol mobile phase modifier with amylose and cellulose tris-(3,5-dimethylphenylcarbamate) stationary phases. <i>Journal of Chromatography A</i> , 2014 , 1373, 197-210	4.5 44
4 ⁰⁷	Quantum mechanical estimation of Abraham hydrogen bond parameters using 1:1 donor-acceptor complexes. 2014 , 27, 783-793	6
4 ⁰⁶	A practical methodology to measure unbiased gas chromatographic retention factor vs. temperature relationships. <i>Journal of Chromatography A</i> , 2014 , 1374, 207-215	4.5 5
4 ⁰⁵	From Molecular Structures of Ionic Liquids to Predicted Retention of Fatty Acid Methyl Esters in Comprehensive Two-Dimensional Gas Chromatography. 2014 , 79, 790-797	24
4 ⁰⁴	Descriptors for the Prediction of Partition Coefficients of 8-Hydroxyquinoline and its Derivatives. 2014 , 49, 2135-2141	15
4 ⁰³	Abraham model correlations for describing solute transfer into ionic liquid solvents: calculation of ion-specific equation coefficients for the 4,5-dicyano-2-(trifluoromethyl)imidazolide anion. <i>Physics and Chemistry of Liquids</i> , 2014 , 52, 777-791	1.5 25
4 ⁰²	On the solubility of quercetin. 2014 , 197, 157-159	26
4 ⁰¹	Novel imidazolium-embedded and imidazolium-spaced octadecyl stationary phases for reversed phase liquid chromatography. <i>Talanta</i> , 2014 , 126, 177-84	6.2 43
4 ⁰⁰	Amino acid/water interactions study: a new amino acid scale. 2014 , 32, 959-68	34

399	Comparative assessment of achiral stationary phases for high throughput analysis in supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2014 , 1332, 73-81	4.5	20
398	IUPAC-NIST Solubility Data Series. 102. Solubility of Nonsteroidal Anti-inflammatory Drugs (NSAIDs) in Neat Organic Solvents and Organic Solvent Mixtures. <i>Journal of Physical and Chemical Reference Data</i> , 2014 , 43, 023102	4.3	28
397	Gas chromatography on wall-coated open-tubular columns with ionic liquid stationary phases. <i>Journal of Chromatography A</i> , 2014 , 1357, 87-109	4.5	109
396	Linear solvation energy relationships as classifier in non-target analysis--an approach for isocratic liquid chromatography. <i>Journal of Chromatography A</i> , 2014 , 1324, 96-103	4.5	6
395	Applications of polyparameter linear free energy relationships in environmental chemistry. <i>Environmental Science & Technology</i> , 2014 , 48, 12477-91	10.3	186
394	Quantitative analysis of molecular interaction potentials of ionic liquid anions using multi-functionalized stationary phases in HPLC. 2014 , 15, 2351-8		8
393	Sorption and Quantitative Structure-Activity Relationship (QSAR). 2014 , 85-118		1
392	An NMR method for the quantitative assessment of intramolecular hydrogen bonding; application to physicochemical, environmental, and biochemical properties. 2014 , 79, 11075-83		60
391	Characterization of novel metallocarborane-based sorbents by linear solvation energy relationships. <i>Journal of Chromatography A</i> , 2014 , 1371, 220-6	4.5	5
390	Mapping environmental partitioning properties of nonpolar complex mixtures by use of GC/GC. <i>Environmental Science & Technology</i> , 2014 , 48, 6814-26	10.3	25
389	Predicting partition coefficients of Polyfluorinated and organosilicon compounds using polyparameter linear free energy relationships (PP-LFERs). <i>Environmental Science & Technology</i> , 2014 , 48, 2776-84	10.3	33
388	Interaction of organic compounds and boron clusters with new silica matrices containing the phosphatidylcholine headgroup. 2014 , 6, 3045-3055		7
387	Responses of polar organic compounds to different ionic environments in aqueous media are interrelated. 2014 , 16, 23347-54		11
386	Organic Contaminant Sorption and Diffusion in Engineered Clays with Organoclay Additives-Understanding Barrier Performance at the Molecular Level. 2014 ,		1
385	Using multidimensional gas chromatography to group secondary organic aerosol species by functionality. 2014 , 96, 310-321		7
384	in vitro and in silico determination of oral, jejenum and Caco-2 human absorption of fatty acids and polyphenols. Micellar liquid chromatography. <i>Talanta</i> , 2014 , 130, 265-73	6.2	14
383	The prediction of blood-tissue partitions, water-skin partitions and skin permeation for agrochemicals. 2014 , 70, 1130-7		19
382	Reconstruction of adsorption potential in Polanyi-based models and application to various adsorbents. <i>Environmental Science & Technology</i> , 2014 , 48, 6772-9	10.3	10

381	Understanding and modeling removal of anionic organic contaminants (AOCs) by anion exchange resins. <i>Environmental Science & Technology</i> , 2014 , 48, 7494-502	10.3	20
380	Determination of descriptors for polycyclic aromatic hydrocarbons and related compounds by chromatographic methods and liquid-liquid partition in totally organic biphasic systems. <i>Journal of Chromatography A</i> , 2014 , 1361, 240-54	4.5	16
379	Ionic liquids as stationary phases in gas chromatography--an LSER investigation of six commercial phases and some applications. 2014 , 406, 5347-58		12
378	Novel Concepts and Challenges of Flavor Microencapsulation and Taste Modification. 2014 , 421-442		3
377	Investigation of active pharmaceutical ingredient loss in pharmaceutical compounding of capsules. 2014 , 96, 68-76		6
376	The solubility of liquid and solid compounds in dry octan-1-ol. 2014 , 103, 26-34		17
375	Determination of a solvent hydrogen-bond acidity scale by means of the solvatochromism of pyridinium-N-phenolate betaine dye 30 and PCM-TD-DFT calculations. 2014 , 118, 4605-14		41
374	Multidimensional gas chromatography beyond simple volatiles separation. 2014 , 50, 8819-33		50
373	Correlations between the zeta potentials of silica hydride-based stationary phases, analyte retention behaviour and their ionic interaction descriptors. 2014 , 817, 48-60		32
372	Human intestinal absorption--neutral molecules and ionic species. 2014 , 103, 1956-1966		20
371	Cross-column prediction of gas-chromatographic retention indices of saturated esters. <i>Journal of Chromatography A</i> , 2014 , 1355, 269-77	4.5	9
370	Prediction of sorption of aromatic and aliphatic organic compounds by carbon nanotubes using poly-parameter linear free-energy relationships. 2014 , 59, 295-303		51
369	Molecular Thermodynamics of Hydrogen-Bonded Systems. 2015 , 145-198		
368	Application of two methods of calculation of solvation descriptor L to estimate C5 -C7 alkenes retention. 2015 , 38, 2447-54		0
367	Green sample-preparation methods using room-temperature ionic liquids for the chromatographic analysis of organic compounds. 2015 , 71, 144-156		60
366	Deduction of Physicochemical Properties from Solubilities: 2,4-Dihydroxybenzophenone, Biotin, and Caprolactam as Examples. 2015 , 60, 1440-1446		23
365	Thermodynamic characteristics of adsorption of organic molecules on pyrimidinophane. 2015 , 64, 800-805		3
364	The transfer of neutral molecules from water and from the gas phase to solvents acetophenone and aniline. 2015 , 212, 301-306		19

363	Organic Carbon/Water and Dissolved Organic Carbon/Water Partitioning of Cyclic Volatile Methylsiloxanes: Measurements and Polyparameter Linear Free Energy Relationships. <i>Environmental Science & Technology</i> , 2015 , 49, 12161-8	10.3	22
362	Physicochemical and biochemical properties for the dialkyl phthalates. 2015 , 119, 871-880		15
361	Prediction of the zeta potentials and ionic descriptors of a silica hydride stationary phase with mobile phases of different pH and ionic strength. 2015 , 859, 79-86		19
360	Automatic Mechanism and Kinetic Model Generation for Gas- and Solution-Phase Processes: A Perspective on Best Practices, Recent Advances, and Future Challenges. 2015 , 47, 199-231		80
359	Evaluation of differences between Chiralpak IA and Chiralpak AD-RH amylose-based chiral stationary phases in reversed-phase high-performance liquid chromatography. 2015 , 38, 711-9		15
358	The many faces of packed column supercritical fluid chromatography--a critical review. <i>Journal of Chromatography A</i> , 2015 , 1382, 2-46	4.5	285
357	Abraham model correlations for solute transfer into tributyl phosphate from both water and the gas phase. <i>Physics and Chemistry of Liquids</i> , 2015 , 53, 10-24	1.5	44
356	Determination of Abraham model solute descriptors for three dichloronitrobenzenes from measured solubilities in organic solvents. <i>Physics and Chemistry of Liquids</i> , 2015 , 53, 163-173	1.5	19
355	A System Map for the Ionic Liquid Stationary Phase 1,9-Di(3-vinylimidazolium)nonane Bis(trifluoromethylsulfonyl)imide. <i>Chromatographia</i> , 2015 , 78, 81-88	2.1	13
354	Order diagram: a universal and versatile approach for system comparison and classification: application to solvent properties. <i>Journal of Chromatography A</i> , 2015 , 1389, 49-64	4.5	22
353	The influence of chlorinated aromatics structure on their adsorption characteristics on activated carbon to tackle chemical spills in drinking water source. 2015 , 9, 138-146		14
352	Abraham model correlations for describing solute transfer into 2-butoxyethanol from both water and the gas phase at 298 K. 2015 , 209, 196-202		35
351	Abraham model correlations for solute transfer into 2-methoxyethanol from water and from the gas phase. 2015 , 209, 738-744		36
350	Predicting storage-lipid water partitioning of organic solutes from molecular structure. <i>Environmental Science & Technology</i> , 2015 , 49, 5538-45	10.3	9
349	In silico model for predicting soil organic carbon normalized sorption coefficient (K(OC)) of organic chemicals. 2015 , 119, 438-444		31
348	Studies on the hydrogen bond acidity, and other descriptors and properties for hydroxyflavones and hydroxyisoflavones. 2015 , 208, 363-372		14
347	Chemically diverse sensor arrays based on electrochemically copolymerized pyrrole and styrene derivatives. 2015 , 215, 510-517		7
346	Abraham model correlations for describing solute transfer into diisopropyl ether. <i>Physics and Chemistry of Liquids</i> , 2015 , 53, 25-37	1.5	46

- 345 Partitioning of Aromatic Constituents into Water from Jet Fuels. **2015**, 69, 153-62
- 344 Characterization and classification of stationary phases in HPLC and SFC $\bar{\Delta}$ review. **2015**, 886, 1-15 51
- 343 Determination of Abraham model solute descriptors for the monomeric and dimeric forms of trans-cinnamic acid using measured solubilities from the Open Notebook Science Challenge. **2015**, 9, 11 17
- 342 Comprehensive Two-Dimensional Gas Chromatography/Mass Spectrometry Analysis of Different Types of Vegetable Oils. **2015**, 92, 783-790 4
- 341 In silico prediction of linear free energy relationship descriptors of neutral and ionic compounds. **2015**, 5, 80634-80642 19
- 340 Sum of ranking differences to rank stationary phases used in packed column supercritical fluid chromatography. *Journal of Chromatography A*, **2015**, 1409, 241-50 4.5 23
- 339 Using water-solvent systems to estimate in vivo blood-tissue partition coefficients. **2015**, 9, 58 3
- 338 An attempt to estimate ionic interactions with phenyl and pentafluorophenyl stationary phases in supercritical fluid chromatography. *Journal of Chromatography A*, **2015**, 1412, 126-38 4.5 32
- 337 Partial solvation parameters and the equation-of-state approach. **2015**, 406, 101-115 12
- 336 LSER model for organic compounds adsorption by single-walled carbon nanotubes: Comparison with multi-walled carbon nanotubes and activated carbon. **2015**, 206, 652-60 33
- 335 Abraham model correlations for estimating solute transfer of neutral molecules into anhydrous acetic acid from water and from the gas phase. **2015**, 212, 16-22 11
- 334 Is there an intramolecular hydrogen bond in 2-halophenols? A theoretical and spectroscopic investigation. **2015**, 17, 25151-9 25
- 333 Analysis of partitioning of organic compounds and proteins in aqueous polyethylene glycol-sodium sulfate aqueous two-phase systems in terms of solute-solvent interactions. *Journal of Chromatography A*, **2015**, 1415, 1-10 4.5 28
- 332 Assessment of the orthogonality in two-dimensional separation systems using criteria defined by the maximal information coefficient. *Journal of Chromatography A*, **2015**, 1415, 108-14 4.5 11
- 331 Comments concerning "A possible simplification of the Goss-modified Abraham solvation equation". **2015**, 138, 1058-61 1
- 330 Are Crystallinity Parameters Critical for Drug Solubility Prediction?. *Journal of Solution Chemistry*, **2015**, 44, 2297-2315 1.8 11
- 329 Polymer selection for SAW sensor array based electronic noses by fuzzy c-means clustering of partition coefficients: Model studies on detection of freshness and spoilage of milk and fish. **2015**, 209, 751-769 43
- 328 Solvent Extraction. **2015**, 131-189 3

327	Analysis of the solubility of betaine: calculation of descriptors and physicochemical properties. 2015 , 387, 1-4		6
326	Comparison of the performance of different silica hydride particles for the solid-phase extraction of non-volatile analytes from dark chocolate with analysis by gas chromatography-quadrupole mass spectrometry. 2015 , 174, 434-9		20
325	A simple method for estimating in vitro air-tissue and in vivo blood-tissue partition coefficients. 2015 , 120, 188-91		29
324	Solvent Selection and Method Development. 2015 , 313-350		2
323	Sorption of the monoterpenes α -pinene and limonene to carbonaceous geosorbents including biochar. 2015 , 119, 881-888		20
322	Sorption of Hydrophobic Organic Compounds to Plastics in the Marine Environment: Equilibrium. <i>Handbook of Environmental Chemistry</i> , 2016 , 185-204	0.8	26
321	Polarizable polysiloxane stationary phase containing a cyano unit attached to an aromatic side group for highly selective separation of H-bonding and aromatic analytes. 2016 , 6, 109786-109792		5
320	System maps for retention of small neutral compounds on a biphenylsiloxane-bonded silica stationary phase in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2016 , 1478, 68-74	4.5	6
319	Determination of Abraham model solute descriptors for isophthalic acid from experimental solubility data in organic solvents at 298 K. <i>Physics and Chemistry of Liquids</i> , 2016 , 54, 747-757	1.5	14
318	Recent advances in development and characterization of stationary phases for hydrophilic interaction chromatography. 2016 , 81, 23-33		77
317	The factors that influence solubility in perfluoroalkane solvents. 2016 , 421, 59-66		10
316	Chromatographic methods enabling the characterization of stationary phases and retention prediction in high-performance liquid chromatography and supercritical fluid chromatography. 2016 , 39, 115-31		20
315	Use and practice of achiral and chiral supercritical fluid chromatography in pharmaceutical analysis and purification. 2016 , 39, 212-33		82
314	Equations for water-triolein partition coefficients for neutral species; comparison with other water-solvent partitions, and environmental and toxicological processes. 2016 , 154, 48-54		14
313	Is the solvation parameter model or its adaptations adequate to account for ionic interactions when characterizing stationary phases for drug impurity profiling with supercritical fluid chromatography?. 2016 , 924, 9-20		13
312	Experimental determination of solvent-water partition coefficients and Abraham parameters for munition constituents. 2016 , 161, 429-437		6
311	System maps for retention of small neutral compounds on a superficially porous particle column in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2016 , 1468, 250-256	4.5	15
310	Deconvoluting the effects of buffer salt concentration in hydrophilic interaction chromatography on a zwitterionic stationary phase. <i>Journal of Chromatography A</i> , 2016 , 1461, 92-7	4.5	28

309	Thermodynamics and limiting activity coefficients measurements for organic solutes and water in the ionic liquid 1-dodecyl-3-methylimidazolium bis(trifluoromethylsulfonyl) imide. 2016 , 103, 76-85		30
308	Comparison of liquid and supercritical fluid chromatography mobile phases for enantioselective separations on polysaccharide stationary phases. <i>Journal of Chromatography A</i> , 2016 , 1467, 463-472	4.5	46
307	Commentary on Measurement and Correlation of the Solubility of Telmisartan (Form A) in Nine Different Solvents from 277.85 to 338.35 K <i>Journal of Solution Chemistry</i> , 2016 , 45, 1902-1905	1.8	3
306	Inverse liquid chromatography as a tool for characterisation of the surface layer of ceramic biomaterials. <i>Journal of Chromatography A</i> , 2016 , 1468, 116-125	4.5	3
305	Predicting solvent-water partitioning of charged organic species using quantum-chemically estimated Abraham pp-LFER solute parameters. 2016 , 164, 634-642		3
304	Rapid characterization of commercial polysorbate 80 by ultra-high performance supercritical fluid chromatography combined with quadrupole time-of-flight mass spectrometry. <i>Journal of Chromatography A</i> , 2016 , 1465, 190-6	4.5	13
303	Solvation descriptors for porphyrins (porphines). 2016 , 40, 9945-9950		6
302	Evaluation of an amide-based stationary phase for supercritical fluid chromatography. 2016 , 39, 3469-76		4
301	Micronization of levofloxacin by supercritical antisolvent precipitation. 2016 , 10, 1201-1210		7
300	Linear modeling of the soil-water partition coefficient normalized to organic carbon content by reversed-phase thin-layer chromatography. <i>Journal of Chromatography A</i> , 2016 , 1458, 136-44	4.5	8
299	Equations for the Partition of Neutral Molecules, Ions and Ionic Species from Water to Water-Methanol Mixtures. <i>Journal of Solution Chemistry</i> , 2016 , 45, 861-874	1.8	28
298	Solvatochromic parameters of imidazolium-, hydroxyammonium-, pyridinium- and phosphonium-based room temperature ionic liquids. 2016 , 213, 107-114		23
297	Concept model of the formation process of humic acid-kaolin complexes deduced by trichloroethylene sorption experiments and various characterizations. 2016 , 151, 116-23		7
296	Development of Abraham model correlations for enthalpies of solvation of organic solutes dissolved in 1,3-dioxolane. <i>Physics and Chemistry of Liquids</i> , 2016 , 54, 786-796	1.5	9
295	An improved classification of stationary phases for ultra-high performance supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2016 , 1440, 212-228	4.5	90
294	Development of Abraham model correlations for predicting enthalpies of solvation of nonionic solutes dissolved in formamide. <i>Physics and Chemistry of Liquids</i> , 2016 , 54, 313-324	1.5	11
293	Development of Abraham model expressions for predicting the enthalpies of solvation of solutes dissolved in acetic acid. <i>Physics and Chemistry of Liquids</i> , 2016 , 54, 141-154	1.5	12
292	Advances in immobilized artificial membrane (IAM) chromatography for novel drug discovery. 2016 , 11, 473-88		34

291	Rational Design of Ionic Liquids for Lipid Processing. 2016 , 153-203		1
290	High-throughput logPo/w determination from UHPLC measurements: Revisiting the chromatographic hydrophobicity index. 2016 , 127, 26-31		8
289	Quantitative structure-retention relationship of selected imidazoline derivatives on β -acid glycoprotein column. 2016 , 127, 101-11		11
288	Modelling for antimicrobial activities of ionic liquids towards <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> and <i>Candida albicans</i> using linear free energy relationship descriptors. <i>Journal of Hazardous Materials</i> , 2016 , 311, 168-75	12.8	29
287	An assessment of air quality reflecting the chemosensory irritation impact of mixtures of volatile organic compounds. 2016 , 86, 84-91		13
286	PCDD/F and PCB water column partitioning examination using natural organic matter and black carbon partition coefficient models. 2016 , 23, 6322-33		0
285	Comparison of different models predicting the phospholipid-membrane water partition coefficients of charged compounds. 2016 , 144, 382-91		42
284	Characterization of ceramic hydroxyapatite surface by inverse liquid chromatography in aquatic systems. <i>Talanta</i> , 2016 , 147, 44-9	6.2	8
283	Determination of the solubilising character of 2-methoxyethyl-(dimethyl)ethylammonium tris(pentafluoroethyl)trifluorophosphate based on the Abraham solvation parameter model. <i>Physics and Chemistry of Liquids</i> , 2016 , 54, 110-126	1.5	15
282	Abraham model correlations for describing solute transfer into anhydrous 1,2-propylene glycol for neutral and ionic species. <i>Physics and Chemistry of Liquids</i> , 2016 , 54, 1-13	1.5	35
281	Descriptors for ions and ion-pairs for use in linear free energy relationships. <i>Journal of Chromatography A</i> , 2016 , 1430, 2-14	4.5	43
280	Applications of the solvation parameter model in reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2017 , 1486, 2-19	4.5	41
279	Abraham model ion-specific equation coefficients for the 1-butyl-2,3-dimethylimidazolium and 4-cyano-1-butylpyridinium cations calculated from measured gas-to-liquid partition coefficient data. <i>Physics and Chemistry of Liquids</i> , 2017 , 55, 218-237	1.5	7
278	Development of polyparameter linear free energy relationship models for octanol-air partition coefficients of diverse chemicals. 2017 , 19, 300-306		5
277	Metabolomic Strategies Involving Mass Spectrometry Combined with Liquid and Gas Chromatography. 2017 , 965, 77-98		17
276	Descriptors for ferrocene and some substituted ferrocenes. 2017 , 232, 325-331		23
275	An equation for the prediction of human skin permeability of neutral molecules, ions and ionic species. <i>International Journal of Pharmaceutics</i> , 2017 , 521, 259-266	6.5	21
274	Combination of partial least squares regression and design of experiments to model the retention of pharmaceutical compounds in supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2017 , 1491, 182-194	4.5	21

273	Recent advances in stationary phases and understanding of retention in hydrophilic interaction chromatography. A review. 2017 , 967, 12-32		171
272	Assessment of the chromatographic lipophilicity of eight cephalosporins on different stationary phases. <i>European Journal of Pharmaceutical Sciences</i> , 2017 , 101, 115-124	5.1	13
271	Modeling Bisolute Adsorption of Aromatic Compounds Based on Adsorbed Solution Theories. <i>Environmental Science & Technology</i> , 2017 , 51, 5552-5562	10.3	3
270	3D-QSAR predictions for Cyclodextrin binding constants using quantum mechanically based descriptors. 2017 , 169, 693-699		9
269	A simple and facile NMR method for the determination of hydrogen bonding by amide NH protons in protein models and other compounds. 2017 , 41, 6064-6066		11
268	High selective water/butan-1-ol separation on investigation of limiting activity coefficients with [P 8,8,8,8][NTf 2] ionic liquid. 2017 , 449, 1-9		21
267	Gas-solvent and water-solvent partition of trans -stilbene at 298 K. 2017 , 238, 58-61		32
266	Baseline toxicity and ion-trapping models to describe the pH-dependence of bacterial toxicity of pharmaceuticals. 2017 , 19, 901-916		19
265	Redefining solubility parameters: Bulk and surface properties from unified molecular descriptors. 2017 , 111, 207-220		16
264	RP-HPLC Analytical Columns. 2017 , 279-328		9
263	Solvents, Buffers, and Additives Used in the Mobile Phase. 2017 , 393-450		2
262	Separation of binary mixtures hexane/hex-1-ene, cyclohexane/cyclohexene and ethylbenzene/styrene based on limiting activity coefficients. 2017 , 110, 227-236		26
261	Quantitative structure-property relationships for predicting sorption of pharmaceuticals to sewage sludge during waste water treatment processes. 2017 , 579, 1512-1520		23
260	Hydrogen-Bond Donicity in DMSO and Gas Phase and Its Dependence on Brønsted Acidity. 2017 , 121, 357-369		10
259	Partition constant database for totally organic biphasic systems. <i>Journal of Chromatography A</i> , 2017 , 1527, 18-32	4.5	12
258	Estimation of Solvation Quantities from Experimental Thermodynamic Data: Development of the Comprehensive CompSol Databank for Pure and Mixed Solutes. <i>Journal of Physical and Chemical Reference Data</i> , 2017 , 46, 033102	4.3	20
257	The New Method for Correlation and Prediction of Thermophysical Properties of Fluids. Critical Temperature. 2017 , 62, 3723-3731		4
256	Graphene Nanoplatelet-Polymer Chemiresistive Sensor Arrays for the Detection and Discrimination of Chemical Warfare Agent Simulants. 2017 , 2, 1669-1678		29

255	A system map for the ionic liquid stationary phase 1,12-di(triisopropylphosphonium)dodecane bis(trifluoromethylsulfonyl)imide for gas chromatography. <i>Journal of Chromatography A</i> , 2017 , 1525, 138-144	4.5	6
254	Estimation of descriptors for hydrogen-bonding compounds from chromatographic and liquid-liquid partition measurements. <i>Journal of Chromatography A</i> , 2017 , 1526, 13-22	4.5	14
253	Evaluating Nonlinear Sorption of Four Substituted Phenols to Agriculture Soils Using Expanded Polyparameter Linear Free Energy Relationship. 2017 , 228, 1		2
252	System map for the ionic liquid stationary phase tri(triisopropylphosphoniumhexanamido)triethylamine bis(trifluoromethylsulfonyl)imide for gas chromatography. <i>Journal of Chromatography A</i> , 2017 , 1524, 210-214	4.5	12
251	Advances in Hydrophilic Interaction Liquid Chromatography. 2017 , 39-87		2
250	Toward a Simple Predictive Molecular Thermodynamic Model for Bulk Phases and Interfaces. <i>Industrial & Engineering Chemistry Research</i> , 2017 , 56, 10900-10910	3.9	13
249	Phase Ratio and Equilibrium Constant in RP-HPLC Obtained from Octanol/Water Partition Constant Through Solvophobic Theory. <i>Chromatographia</i> , 2017 , 80, 1491-1500	2.1	3
248	Study of benzyl- or cyclohexyl-functionalized ionic liquids using inverse gas chromatography. 2017 , 242, 550-559		28
247	Quantum Chemically Estimated Abraham Solute Parameters Using Multiple Solvent-Water Partition Coefficients and Molecular Polarizability. <i>Environmental Science & Technology</i> , 2017 , 51, 9887-9898	10.3	10
246	Solvent selection in liquid chromatography. 2017 , 343-373		2
245	Quantitative structure property (retention) relationships in liquid chromatography. 2017 , 553-572		3
244	Reversed-phase liquid chromatography. 2017 , 91-123		6
243	Determination of molar refractions and Abraham descriptors for tris(acetylacetonato)chromium(III), tris(acetylacetonato)iron(III) and tris(acetylacetonato)cobalt(III). 2017 , 41, 14259-14265		8
242	Chiral surfaces: The many faces of chiral recognition. 2017 , 32, 94-107		22
241	Separation of water/butan-1-ol mixtures based on limiting activity coefficients with phosphonium-based ionic liquid. 2017 , 113, 183-191		23
240	System Maps for Retention of Small Neutral Compounds on a Superficially Porous Ethyl-Bridged, Octadecylsiloxane-Bonded Silica Stationary Phase in Reversed-Phase Liquid Chromatography. <i>Chromatographia</i> , 2017 , 80, 1279-1286	2.1	10
239	The correlation and prediction of infinite dilution activity coefficients of compounds in water at 298.15 K. 2017 , 449, 117-129		9
238	Liquid chromatography with room temperature ionic liquids. <i>Journal of Planar Chromatography - Modern TLC</i> , 2017 , 30, 97-105	0.9	11

237	Ion-specific equation coefficient version of the Abraham model for ionic liquid solvents: determination of coefficients for tributylethylphosphonium, 1-butyl-1-methylmorpholinium, 1-allyl-3-methylimidazolium and octyltriethylammonium cations. <i>Physics and Chemistry of Liquids</i> , 2017 , 55, 358-385	1.5	31
236	Separation of binary mixtures based on gamma infinity data using [EMIM][TCM] ionic liquid and modelling of thermodynamic functions. 2017 , 225, 382-390		29
235	Descriptors for Pentane-2,4-dione and Its Derivatives. <i>Journal of Solution Chemistry</i> , 2017 , 46, 1625-1638	1.8	7
234	Abraham Model Correlations for Triethylene Glycol Solvent Derived from Infinite Dilution Activity Coefficient, Partition Coefficient and Solubility Data Measured at 298.15 K. <i>Journal of Solution Chemistry</i> , 2017 , 46, 2249-2267	1.8	10
233	The Evaluation of Pro-Cognitive and Antiamnestic Properties of Berberine and Magnoflorine Isolated from Barberry Species by Centrifugal Partition Chromatography (CPC), in Relation to QSAR Modelling. 2017 , 18,		14
232	Practical Approaches to Column Selection for Supercritical Fluid Chromatography. 2017 , 57-101		
231	Column Characterization. 2017 , 103-125		
230	Insights into the Retention Mechanism of Small Neutral Compounds on Octylsiloxane-Bonded and Diisobutyloctadecylsiloxane-Bonded Silica Stationary Phases in Reversed-Phase Liquid Chromatography. <i>Chromatographia</i> , 2018 , 81, 373-385	2.1	9
229	Sorption of organic compounds by aged polystyrene microplastic particles. 2018 , 236, 218-225		223
228	Linear solvation energy relationship (LSER) characterization of the normal phase retention mechanism on hypercrosslinked polystyrenes. <i>Journal of Chromatography A</i> , 2018 , 1543, 40-47	4.5	1
227	Physicochemical property guidelines for modern agrochemicals. 2018 , 74, 1979		34
226	Estimating system parameters for solvent-water and plant cuticle-water using quantum chemically estimated Abraham solute parameters. 2018 , 20, 813-821		4
225	Descriptors for the α -dicarboxylic acids from oxalic acid to sebacic acid. 2018 , 467, 17-24		6
224	Validation and updating of QSAR models for partitioning coefficients of ionic liquids in octanol-water and development of a new LFER model. 2018 , 633, 920-928		14
223	Thermodynamic study of molecular interaction-selectivity in separation processes based on limiting activity coefficients. 2018 , 121, 112-120		16
222	Characterization of Retention Mechanisms in Mixed-Mode HPLC with a Bimodal Reversed-Phase/Cation-Exchange Stationary Phase. <i>Chromatographia</i> , 2018 , 81, 387-399	2.1	9
221	Infinite Dilution Activity Coefficients and Gas-to-Liquid Partition Coefficients of Organic Solutes Dissolved in 1-Benzylpyridinium Bis(Trifluoromethylsulfonyl)Imide and 1-Cyclohexylmethyl-1-Methylpyrrolidinium Bis(Trifluoromethylsulfonyl)Imide. <i>Journal of Solution Chemistry</i> , 2018 , 47, 308-335	1.8	25
220	Characterizing Sorption and Permeation Properties of Membrane Filters Used for Aquatic Integrative Passive Samplers. <i>Environmental Science & Technology</i> , 2018 , 52, 2118-2125	10.3	27

219	Thermodynamic characterization of polymers. 2018 , 136, 47-61		12
218	Chromatographic efficiency of polar capillary columns applied for the analysis of fatty acid methyl esters by gas chromatography. 2018 , 41, 1582-1592		3
217	Screening tools for the bioconcentration potential of monovalent organic ions in fish. 2018 , 20, 845-853		12
216	Partition of Neutral Molecules and Ions from Water to -Nitrophenyl Octyl Ether and of Neutral Molecules from the Gas Phase to -Nitrophenyl Octyl Ether. <i>Journal of Solution Chemistry</i> , 2018 , 47, 293-307	1.8	6
215	A system map for the ionic liquid stationary phase 1,12-di(triisopropylphosphonium)dodecane bis(trifluoromethylsulfonyl)imide trifluoromethanesulfonate for gas chromatography. <i>Journal of Chromatography A</i> , 2018 , 1559, 164-169	4-5	9
214	Updated Abraham model correlations for correlating solute transfer into dry butanone and dry cyclohexanone solvents. <i>Physics and Chemistry of Liquids</i> , 2018 , 56, 571-583	1.5	27
213	Structural properties governing drug-plasma protein binding determined by high-performance liquid chromatography method. 2018 , 149, 16-21		12
212	Illustration of the calculation of solute descriptors for maltol from published solubility data. <i>Physics and Chemistry of Liquids</i> , 2018 , 56, 403-409	1.5	2
211	The correlation and prediction of the temperature variation of infinite dilution activity coefficients of compounds in water. 2018 , 455, 1-5		4
210	Biorelevant Drug Solubility Enhancement Modeled by a Linear Solvation Energy Relationship. 2018 , 107, 503-506		16
209	Insights into the Retention Mechanism for Small Neutral Compounds on Silica-Based Phenyl Phases in Reversed-Phase Liquid Chromatography. <i>Chromatographia</i> , 2018 , 81, 225-238	2.1	9
208	Effects of structure on the performance of latex nanoparticles as a pseudostationary phase in electrokinetic chromatography. 2018 , 1000, 293-302		1
207	Determination of Abraham model solute descriptors for monomeric 3,4,5-trimethoxybenzoic acid from experimental solubility data in organic solvents measured at 298.2 K. <i>Physics and Chemistry of Liquids</i> , 2018 , 56, 381-390	1.5	16
206	Recent advances in mixed-mode chromatographic stationary phases. 2019 , 42, 89-129		36
205	Preconcentration of Organic Substances on Low-Polar Adsorbents in the Flow Systems of Analysis. <i>Journal of Analytical Chemistry</i> , 2018 , 73, 1030-1042	1.1	1
204	A new method for calculating normal boiling point of liquids. 2018 , 67, 1823-1830		
203	An Alternative Method for Correlation and Evaluation of Mutual Diffusion Coefficients of Solutes in Organic Solvents at Infinite Dilution. <i>Journal of Solution Chemistry</i> , 2018 , 47, 1479-1491	1.8	2
202	Protonation of polyaniline-coated silica stationary phase affects the retention behavior of neutral hydrophobic solutes in reversed-phase capillary liquid chromatography. 2018 , 41, 2886		8

201	Descriptors for Cyclooctasulfur: Estimation of Water-Solvent Partition Coefficients, Solubilities in Solvents, and Physicochemical Properties. 2018 , 3, 5516-5521		8
200	Monolithic rod columns for HPLC based on divinylbenzene-styrene copolymer with 1-vinylimidazole and 4-vinylpyridine. 2018 , 41, 458-466		6
199	Comparative Estimation of the Selectivity Parameters for Monolith HPLC Columns with Organic Sorbents. 2018 , 92, 1615-1620		
198	Linear free energy relationship analysis of permeability across polydimethylsiloxane (PDMS) membranes and comparison with human skin permeation in vitro. <i>European Journal of Pharmaceutical Sciences</i> , 2018 , 123, 524-530	5.1	5
197	Predicting the skin-permeating components of externally-applied medicinal herbs: application of a newly constructed linear free-energy relationship equation for human skin permeation. 2018 , 42, 11930-11943 ²		
196	Characterization of stationary phases based on monosubstituted benzene retention indices using correspondence factor analysis and linear solvation energy relationships in RPLC. 2018 , 15, 2295-2305		2
195	Chromatographic test methods for characterizing alkylsiloxane-bonded silica columns for reversed-phase liquid chromatography. 2018 , 1092, 207-219		21
194	Determination of Abraham model solute descriptors for 4-tert-butylbenzoic acid from experimental solubility data in organic mono-solvents. <i>Physics and Chemistry of Liquids</i> , 2019 , 57, 445-452 ^{1.5}		8
193	Infinite dilution activity coefficients and gas-to-liquid partition coefficients of organic solutes dissolved in 1-sec-butyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide and in 1-tert-butyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide. <i>Physics and Chemistry of Liquids</i> , 2019 , 57, 453-472	1.5	21
192	Determination of Abraham model solute descriptors for o-acetoacetanilide based on experimental solubility data in organic mono-solvents. <i>Physics and Chemistry of Liquids</i> , 2019 , 57, 528-535 ^{1.5}		12
191	Chemoresistive materials for electronic nose: Progress, perspectives, and challenges. 2019 , 1, 289-316		71
190	Evaluation of separation properties of stationary phases in supercritical fluid chromatography; deazapurine nucleosides case study. 2019 , 150, 104137		7
189	Skin-permeating components of <i>Lonicera japonica</i> flos: a comprehensive study from observations and model computations. 2019 , 43, 12538-12547		0
188	Characterization of three macrocyclic glycopeptide stationary phases in supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2019 , 1604, 460485	4.5	18
187	Mechanistic studies of congener-specific adsorption and bioaccumulation of polycyclic aromatic hydrocarbons and phthalates in soil by novel QSARs. 2019 , 179, 108838		5
186	A comparison of the solvation power of the green solvent 2,2,5,5-tetramethyloxolane versus toluene via partition coefficients. 2019 , 240, 118175		4
185	A new method for the determination of Henry's law constants (air-water-partition coefficients). 2019 , 502, 112300		11
184	Gas chromatography system constant database for 52 wall-coated, open-tubular columns covering the temperature range 60-140°C. <i>Journal of Chromatography A</i> , 2019 , 1604, 460482	4.5	10

183	Effects of high concentrations of mobile phase additives on retention and separation mechanisms on a teicoplanin aglycone stationary phase in supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2019 , 1604, 460494	4.5	16
182	Development of Abraham model expressions for predicting the standard molar enthalpies of vaporization of organic compounds at 298.15 K. 2019 , 681, 178372		13
181	Determination of the hydrogen-bond acidity and basicity for un-dissociated hydrazoic acid, isocyanic acid and isothiocyanic acid. 2019 , 294, 111666		0
180	Atmospheric histories, growth rates and solubilities in seawater and other natural waters of the potential transient tracers HCFC-22, HCFC-141b, HCFC-142b, HFC-134a, HFC-125, HFC-23, PFC-14 and PFC-116. 2019 , 15, 33-60		6
179	Design, synthesis and evaluation of a series of alkylsiloxane-bonded stationary phases for expanded supercritical fluid chromatography separations. <i>Journal of Chromatography A</i> , 2019 , 1593, 127-134	4.5	2
178	Hybrid QSPR models for the prediction of the free energy of solvation of organic solute/solvent pairs. 2019 , 21, 13706-13720		19
177	110th Anniversary: From Solubility Parameters to Predictive Equation-of-State Modeling. <i>Industrial & Engineering Chemistry Research</i> , 2019 , 58, 12787-12800	3.9	7
176	Reversed-phase liquid chromatography system constant database over an extended mobile phase composition range for 25 siloxane-bonded silica-based columns. <i>Journal of Chromatography A</i> , 2019 , 1600, 112-126	4.5	16
175	Solvent and temperature effects on the solubility of syringic, vanillic or veratric acids: Experimental, modeling and solid phase studies. 2019 , 289, 111089		16
174	GC/MS and DFT studies of S,S-dialkyl methylphosphonothioloselenoates related to Schedule 2.B.04 of Chemical Weapons Convention. 2019 , 40, 516-528		4
173	Reversed phase HPLC with high temperature ethanol/water mobile phases as a green alternative method for the estimation of octanol/water partition coefficients. <i>Journal of Chromatography A</i> , 2019 , 1601, 243-254	4.5	8
172	Developing Predictive Models for Carrying Ability of Micro-Plastics towards Organic Pollutants. 2019 , 24,		12
171	Comprehensive two-dimensional gas-chromatography-based property estimation to assess the fate and behavior of complex mixtures: A case study of vehicle engine oil. 2019 , 669, 739-745		5
170	Determination of Some Physicochemical Properties of Mebendazole with RPLC Method. 2019 , 64, 2736-2741		1
169	System Maps for the Retention of Neutral Compounds on an Electrostatic-Shielded Reversed-Phase Column. <i>Chromatographia</i> , 2019 , 82, 799-808	2.1	5
168	Characterization of the solubilizing ability of tetraalkylammonium ionic liquids containing a pendant alkyl chain bearing a basic N,N-dimethylamino or N,N-dimethylaminoethoxy functionality. 2019 , 283, 380-390		14
167	Prediction of water solubility and Setschenow coefficients by tree-based regression strategies. 2019 , 282, 401-406		4
166	QSAR characterization of new synthesized hydantoin with antiproliferative activity. 2019 , 33, e4539		1

165	Solvation Descriptors for Zwitterionic α -Aminoacids; Estimation of Water-Solvent Partition Coefficients, Solubilities, and Hydrogen-Bond Acidity and Hydrogen-Bond Basicity. 2019 , 4, 2883-2892		8
164	Characterization of sorption properties of high-density polyethylene using the poly-parameter linearfree-energy relationships. 2019 , 248, 312-319		15
163	The Poisson-Boltzmann model for implicit solvation of electrolyte solutions: Quantum chemical implementation and assessment via Sechenov coefficients. 2019 , 151, 224111		20
162	Thermodynamic characterization of ionic liquids. 2019 , 277, 10-21		9
161	Descriptors for the hydrogen halides, their solution properties and hydrogen- bonding acidity and basicity: Comparison of the latter with gas phase data. 2019 , 275, 667-673		4
160	Drug supersaturation during formulation digestion, including real-time analytical approaches. 2019 , 142, 50-61		18
159	Partial Solvation Parameters of Drugs as a New Thermodynamic Tool for Pharmaceuticals. 2019 , 11,		8
158	Development of Abraham model IL-specific correlations for N-triethyl(octyl)ammonium bis(fluorosulfonyl)imide and 1-butyl-3-methylpyrrolidinium bis(fluorosulfonyl)imide. <i>Physics and Chemistry of Liquids</i> , 2019 , 57, 733-745	1.5	6
157	Influence of Solvent Effects on Retention of Small Molecules in Reversed-Phase Liquid Chromatography. <i>Chromatographia</i> , 2019 , 82, 49-64	2.1	21
156	Abraham model correlations for enthalpies of solvation of organic solutes dissolved in methyl acetate and octane. <i>Physics and Chemistry of Liquids</i> , 2020 , 58, 18-30	1.5	7
155	Core concepts and milestones in the development of solid-phase extraction. 2020 , 1-36		3
154	Polyethersulfone as suitable passive sampler for waterborne hydrophobic organic compounds - Laboratory calibration and field test in the Sosiani river, Kenya. 2020 , 699, 134056		3
153	Use of HPLC retention to investigate new P descriptors designed to represent ion-ion interactions. 2020 , 43, 83-93		1
152	Descriptors for terpene esters from chromatographic and partition measurements: Estimation of human odor detection thresholds. <i>Journal of Chromatography A</i> , 2020 , 1609, 460428	4.5	3
151	Solubility of 4-methyl-3-nitrobenzoic acid in organic mono-solvents: calculation of Abraham model solute descriptors. <i>Physics and Chemistry of Liquids</i> , 2020 , 58, 782-791	1.5	2
150	Characterization of mesoporous aluminosilicate materials by means of inverse liquid chromatography. <i>Journal of Chromatography A</i> , 2020 , 1610, 460544	4.5	4
149	Solvent Selection for Liquid-Phase Extraction. 2020 , 45-89		2
148	Predicting low density polyethylene-water partition coefficients based on pp-LFER and QSPR models using molecular descriptors. 2020 , 506, 112374		4



147	Classification of biphasic solvent systems according to Abraham descriptors for countercurrent chromatography. <i>Journal of Chromatography A</i> , 2020 , 1617, 460820	4.5	5
146	Wayne State University experimental descriptor database for use with the solvation parameter model. <i>Journal of Chromatography A</i> , 2020 , 1617, 460841	4.5	19
145	Development of pp-LFER and QSPR models for predicting the diffusion coefficients of hydrophobic organic compounds in LDPE. 2020 , 190, 110179		5
144	Preparation and comparison of three zwitterionic stationary phases for hydrophilic interaction liquid chromatography. 2020 , 43, 1071-1079		2
143	Totally Organic Biphasic Systems. 2020 , 265-288		
142	Evaluation of a series of phenyl-type stationary phases in supercritical fluid chromatography with the linear solvation energy relationship model and its application to the separation of phenolic compounds. <i>Journal of Chromatography A</i> , 2020 , 1614, 460700	4.5	7
141	Room-temperature-ionic-liquid coated graphitized carbons for selective preconcentration of polar vapors. <i>Journal of Chromatography A</i> , 2020 , 1609, 460486	4.5	5
140	[DCA]-based ionic liquids for the extraction of sulfur and nitrogen compounds from fuels: Activity coefficients at infinite dilution. 2020 , 507, 112424		11
139	Selection of calibration compounds for selectivity evaluation of wall-coated, open-tubular columns for gas chromatography by the solvation parameter model. <i>Journal of Chromatography A</i> , 2020 , 1629, 461500	4.5	5
138	Application of HPLC measurements for the determination of physicochemical and biomimetic properties to model in vivo drug distribution in support of early drug discovery. 2020 , 8, 667-758		2
137	Selectivity evaluation of core-shell silica columns for reversed-phase liquid chromatography using the solvation parameter model. <i>Journal of Chromatography A</i> , 2020 , 1634, 461692	4.5	10
136	Concepts, selectivity options and experimental design approaches in multidimensional and comprehensive two-dimensional gas chromatography. 2020 , 130, 115995		13
135	Estimation of heat capacities of gases, liquids and solids, and heat capacities of vaporization and of sublimation of organic chemicals at 298.15 K. 2020 , 317, 113969		5
134	Determination of physicochemical properties of small molecules by reversed-phase liquid chromatography. <i>Journal of Chromatography A</i> , 2020 , 1626, 461427	4.5	9
133	Selection of calibration compounds for selectivity evaluation of siloxane-bonded silica columns for reversed-phase liquid chromatography by the solvation parameter model. <i>Journal of Chromatography A</i> , 2020 , 1633, 461652	4.5	6
132	Predicting Sorbent-Air Partition Coefficients for Terpenoids at Multiple Temperatures. <i>Industrial & Engineering Chemistry Research</i> , 2020 , 59,	3.9	1
131	Uptake/release of organic contaminants by microplastics: A critical review of influencing factors, mechanistic modeling, and thermodynamic prediction methods. 2020 , 1-45		3
130	Temperature-dependent vapor/liquid equilibria and solvation free energy estimation from minimal data. 2020 , 66, e16976		5

129	Yolk-Water Partitioning of Neutral Organic Compounds in the Model Organism <i>Danio rerio</i> . 2020 , 39, 1506-1516		1
128	Microgeometry-independent equation for measuring infinite dilution activity coefficients using gas-liquid chromatography with static-wall-coated open-tubular columns. <i>Journal of Chromatography A</i> , 2020 , 1624, 461264	4.5	1
127	Versatile modelling of polyoxymethylene-water partition coefficients for hydrophobic organic contaminants using linear and nonlinear approaches. 2020 , 728, 138881		5
126	Differentiation of ring-substituted regioisomers of cathinone analogs by supercritical fluid chromatography. 2020 , 1, 22		1
125	Applications of Machine Learning to In Silico Quantification of Chemicals without Analytical Standards. <i>Journal of Chemical Information and Modeling</i> , 2020 , 60, 2718-2727	6.1	12
124	VOCs adsorption on activated carbon with initial water vapor contents: Adsorption mechanism and modified characteristic curves. 2020 , 731, 139184		9
123	Estimation of vapor pressures of liquid and solid organic and organometallic compounds at 298.15 K. 2020 , 519, 112595		9
122	Estimation of Vapor Pressures of Solvent + Salt Systems with Quadratic Solvation Relationships. <i>Journal of Solution Chemistry</i> , 2020 , 49, 559-582	1.8	
121	Evaluation of the solvation parameter model as a quantitative structure-retention relationship model for gas and liquid chromatography. <i>Journal of Chromatography A</i> , 2020 , 1626, 461308	4.5	9
120	Does phase ratio in reversed phase high performance liquid chromatography vary with temperature?. <i>Journal of Chromatography A</i> , 2020 , 1620, 461023	4.5	6
119	Towards a predictive thermodynamic description of sorption processes in polymers: The synergy between theoretical EoS models and vibrational spectroscopy. 2020 , 140, 100525		17
118	The assessment of intramolecular hydrogen bonding in ortho-substituted anilines by an NMR method. 2020 , 315, 113730		1
117	Solid-liquid phase equilibrium of trans-cinnamic acid, p-coumaric acid and ferulic acid in water and organic solvents: Experimental and modelling studies. 2020 , 521, 112747		6
116	Hesperetin Solubility in Aqueous Co-solvent Mixtures of Methanol and Ethanol: Solute Descriptors, Solvent Effect and Preferential Solvation Analysis. <i>Journal of Solution Chemistry</i> , 2020 , 49, 179-194	1.8	0
115	Biodegradation and Ecotoxicity of Branched Alcohol Ethoxylates: Application of the Target Lipid Model and Implications for Environmental Classification. 2020 , 23, 383-403		7
114	Commercial Applications of Ionic Liquids. <i>Green Chemistry and Sustainable Technology</i> , 2020 ,	1.1	27
113	Comments on "Classification of biphasic solvent systems according to Abraham descriptors for countercurrent chromatography". <i>Journal of Chromatography A</i> , 2020 , 1618, 460889	4.5	0
112	Incorporation of Imidazolium Ionic Liquids in GC Stationary Phases via the SolGel Process. <i>Chromatographia</i> , 2020 , 83, 439-449	2.1	0

111	Predicting the Blood-Brain Barrier Permeability of New Drug-Like Compounds via HPLC with Various Stationary Phases. 2020 , 25,		4
110	Property Estimation of Per- and Polyfluoroalkyl Substances: A Comparative Assessment of Estimation Methods. 2020 , 39, 775-786		12
109	Estimation of enthalpies of sublimation of organic, organometallic and inorganic compounds. 2020 , 515, 112575		8
108	Abraham model correlations for describing dissolution of organic solutes and inorganic gases in dimethyl carbonate. <i>Physics and Chemistry of Liquids</i> , 2021 , 59, 181-195	1.5	5
107	Prediction of Analyte Retention Time in Liquid Chromatography. 2021 , 93, 228-256		29
106	The solubility parameters of carbon dioxide and ionic liquids: Are they an enigma?. 2021 , 527, 112828		3
105	BTEX compounds leachates from cigarette butts into water environment: A primary study. 2021 , 269, 116185		12
104	Descriptors for High-Energy Nitro Compounds; Estimation of Thermodynamic, Physicochemical and Environmental Properties. 2021 , 46, 267-279		1
103	Abraham model correlations for describing solute transfer processes into diethyl carbonate. <i>Physics and Chemistry of Liquids</i> , 2021 , 59, 26-39	1.5	7
102	Column Selection and Optimization for Comprehensive Two-Dimensional Gas Chromatography: A Review. 2021 , 51, 183-202		7
101	Solvent extraction. 2021 , 191-279		
100	Cellobiose as a Model Carbohydrate for Predicting Solubilities in Nonaqueous Solvents. <i>Industrial & Engineering Chemistry Research</i> , 2021 , 60, 1859-1871	3.9	2
99	Properties of the tert-butyl halide solvolysis transition states. 2021 , 23, 3311-3320		1
98	Predicting the adsorption of organic pollutants on boron nitride nanosheets via in silico techniques: DFT computations and QSAR modeling. 2021 , 8, 795-805		5
97	Characterization of stationary phases in supercritical fluid chromatography including exploration of shape selectivity. <i>Journal of Chromatography A</i> , 2021 , 1639, 461923	4.5	5
96	Congener-specific partition properties of chlorinated paraffins evaluated with COSMOtherm and gas chromatographic retention indices. 2021 , 11, 4426		1
95	Performance of smoothly clipped absolute deviation as a variable selection method in the artificial neural network-based QSAR studies. 2021 , 35, e3338		2
94	Descriptors for fluorotelomere alcohols. Calculation of physicochemical properties. <i>Physics and Chemistry of Liquids</i> , 1-6	1.5	1

93	Application of polymethyl[4-(2,3-difluoro-4-hydroxyphenoxy)butyl] siloxane in surface acoustic wave gas sensors for dimethyl methylphosphonate detection. 2021 , 329, 129216		11
92	Relook on the Linear Free Energy Relationships Describing the Partitioning Behavior of Diverse Chemicals for Polyethylene Water Passive Samplers. 2021 , 6, 5221-5232		1
91	Use of Partition Coefficients in a Hexane-Acetonitrile System in the GC-MS Analysis of Polyaromatic Hydrocarbons in the Example of Delayed Coking Gas Oils. 2021 , 6, 9910-9919		4
90	Descriptors for adamantane and some of its derivatives. 2021 , 325, 114894		0
89	Equations for the Correlation and Prediction of Partition Coefficients of Neutral Molecules and Ionic Species in the Water-Isopropanol Solvent System. <i>Journal of Solution Chemistry</i> , 2021 , 50, 458-472	1.8	4
88	Descriptors for vitamin K3 (menadione); calculation of biological and physicochemical properties. 2021 , 330, 115707		4
87	Solvation parameter model: Tutorial on its application to separation systems for neutral compounds. <i>Journal of Chromatography A</i> , 2021 , 1645, 462108	4.5	10
86	Development and Application of a Novel Semi-quantification Approach in LC-QToF-MS Analysis of Natural Products. 2021 , 32, 1412-1423		2
85	Determination of lipid-water partition coefficient of neutral and ionic drugs by liposome electrokinetic chromatography. 2021 , 42, 1436-1449		1
84	Determination of physicochemical properties of ionic liquids by gas chromatography. <i>Journal of Chromatography A</i> , 2021 , 1644, 461964	4.5	5
83	Safer Solvent Blends for Food, Dye, and Environmental Analyses Using Reversed-Phase High Performance Liquid Chromatography. <i>Chromatographia</i> , 2021 , 84, 769	2.1	3
82	Descriptors for Edaravone; studies on its structure, and prediction of properties. 2021 , 332, 115821		0
81	In silico prediction of polyethylene-aqueous and air partition coefficients of organic contaminants using linear and nonlinear approaches. 2021 , 289, 112437		0
80	A Comprehensive Non-targeted Analysis Study of the Prenatal Exposome. <i>Environmental Science & Technology</i> , 2021 , 55, 10542-10557	10.3	5
79	Determination of Water-Solvent Partition Coefficients for Fluorescein: Evaluation of Descriptors for the Lactone Form and Prediction of Properties. <i>Journal of Solution Chemistry</i> , 2021 , 50, 1027-1035	1.8	
78	Linear solvation energy relationships in the determination of phospholipids by supercritical fluid chromatography. 2021 , 173, 105206		4
77	Empirical regressions between system parameters and solute descriptors of polyparameter linear free energy relationships (PPLFERs) for predicting solvent-air partitioning. 2021 , 540, 113035		0
76	Development of Predictive Expressions for Infinite Dilution Activity Coefficients, Molar Solubilities and Partition Coefficients for Solutes Dissolved in 2-Pyrrolidone Based on the Abraham Solvation Parameter Model. <i>Journal of Solution Chemistry</i> , 1	1.8	

75	Prediction of Chameleonic Efficiency. <i>ChemMedChem</i> , 2021 , 16, 2669-2685	3.7	1
74	Development of Abraham model correlations for solute transfer into cyclopentanol from both water and the gas phase based on measured solubility ratios. <i>Physics and Chemistry of Liquids</i> , 1-10	1.5	2
73	Modeling polyurethane foam (PUF)-air partition coefficients for persistent organic pollutants using linear and non-linear chemometric methods. <i>Journal of Environmental Chemical Engineering</i> , 2021 , 9, 105615	6.8	
72	Predicting Leachables Solubilization in Polysorbate 80 Solutions by a Linear Solvation Energy Relationship (LSER). <i>Pharmaceutical Research</i> , 2021 , 38, 1549-1561	4.5	1
71	Abraham model description of the solubilising properties of the isopropyl acetate organic mono-solvent. <i>Physics and Chemistry of Liquids</i> , 1-13	1.5	3
70	Synthesis of octadecylamine-derived carbon dots and application in reversed phase/hydrophilic interaction liquid chromatography. <i>Journal of Chromatography A</i> , 2021 , 1656, 462548	4.5	1
69	Inverse gas chromatographic evaluation of polysiloxanes containing phenolic and fluorophenolic functional groups for use in gas sensors. <i>Talanta</i> , 2021 , 234, 122711	6.2	1
68	Prediction models with multiple machine learning algorithms for POPs: The calculation of PDMS-air partition coefficient from molecular descriptor. <i>Journal of Hazardous Materials</i> , 2022 , 423, 127037	12.8	2
67	Column classification and structure-retention relationships. 2021 , 165-190		
66	Design and Information Content of Arrays of Sorption-Based Vapor Sensors Using Solubility Interactions and Linear Solvation Energy Relationships. 2009 , 193-218		4
65	Fate and Transfer of Semivolatile Organic Compounds in a Multi-Compartment Environment. <i>Environmental Science and Engineering</i> , 2010 , 277-307	0.2	3
64	Separation Techniques in Effect-Directed Analysis. <i>Handbook of Environmental Chemistry</i> , 2011 , 83-118	0.8	3
63	Columns and Stationary Phases. 2014 , 59-116		2
62	Designing of Novel 6(H)-1,3,4-Thiadiazine Derivatives as MMP12 Inhibitors: A MLR and Docking Approach. <i>American Journal of Pharmacological Sciences</i> , 2013 , 1, 29-34	0	4
61	An interphase model for retention in liquid chromatography. <i>Journal of Planar Chromatography - Modern TLC</i> , 2015 , 28, 98-105	0.9	14
60	Comprehensive structural characterisation of the newly emerged synthetic cannabimimetics Cumyl-BC[2.2.1]HpMeGaClone, Cumyl-BC[2.2.1]HpMINACA, and Cumyl-BC[2.2.1]HpMICA featuring a norbornyl methyl side chain. <i>Forensic Chemistry</i> , 2021 , 26, 100371	2.8	1
59	IUPAC-NIST Solubility Data Series. 105. Solubility of Solid Alkanoic Acids, Alkenoic Acids, Alkanedioic Acids, and Alkenedioic Acids Dissolved in Neat Organic Solvents, Organic Solvent Mixtures, and Aqueous Organic Solvent Mixtures. I. Alkanoic Acids. <i>Journal of Physical and Chemical Reference Data</i> , 2021 , 50, 043103	4.3	1
58	Building Structure-Property Predictive Models Using Data Assimilation. <i>Lecture Notes in Computer Science</i> , 2006 , 64-73	0.9	

57	Deposition from the Atmosphere to Water and Soils with Aerosol Particles and Precipitation. 2010 , 103-135		
56	  <i>Russian Journal of Physical Chemistry A</i> , 2018 , 1344-1350	1.5	
55	Chapter 3. Conventional Gas Chromatography: Basic Principles and Instrumental Aspects. <i>Food Chemistry, Function and Analysis</i> , 2019 , 83-130	0.6	
54	Gas Chromatography Columns Using Ionic Liquids as Stationary Phase. <i>Green Chemistry and Sustainable Technology</i> , 2020 , 131-165	1.1	2
53	Solvent Selection for Environmental Organic Chemistry ExperimentsBasic Concepts. <i>Journal of Environmental Chemistry</i> , 2020 , 30, 145-150	0.3	
52	Olfactory Sensing Using Quartz Crystal Microbalances with Radio-Frequency Sputtered Organic Films Based on Phenomenological Gas-Sorption Dynamics. 170-182		
51	Evaluation of a Semi-Empirical Approach to Correlation of Retention with Octanol-Water Partition Coefficients by Use of Adjustable Hydrogen-Bonding Indicator Variables. <i>Chromatographia</i> , 2005 , 61, 479	2.1	
50	Mathematical Modelling in Analytical Chromatography: Problems and Solutions. <i>Journal of Analytical Chemistry</i> , 2021 , 76, 1233-1244	1.1	
49	Kinetics Assessment of the Homogeneously Catalyzed Hydroformylation of Ethylene on an Rh Catalyst. <i>Industrial & Engineering Chemistry Research</i> , 2021 , 60, 16665-16681	3.9	0
48	Abraham Model Descriptors for Melatonin; Prediction of Solution, Biological and Thermodynamic Properties. <i>Journal of Solution Chemistry</i> , 1	1.8	1
47	Abraham model solute descriptors: effect of structural features on the calculated numerical descriptor values for vanillin and select derivatives. <i>Physics and Chemistry of Liquids</i> , 1-13	1.5	0
46	Group Contribution and Machine Learning Approaches to Predict Abraham Solute Parameters, Solvation Free Energy, and Solvation Enthalpy.. <i>Journal of Chemical Information and Modeling</i> , 2022 ,	6.1	6
45	Combining Experimental Sorption Parameters with QSAR to Predict Neonicotinoid and Transformation Product Sorption to Carbon Nanotubes and Granular Activated Carbon.. <i>ACS ES&T Water</i> , 2022 , 2, 247-258		0
44	Structural Effects on the Hydrogen-Bonding Descriptors of the Solvation Parameter Model. <i>Journal of Solution Chemistry</i> , 1	1.8	2
43	Linear Solvation Energy Relationships (LSERs) for Robust Prediction of Partition Coefficients between Low Density Polyethylene and Water Part II: Model Evaluation and Benchmarking.. <i>European Journal of Pharmaceutical Sciences</i> , 2022 , 172, 106138	5.1	
42	Investigation of partition coefficients and fingerprints of atmospheric gas- and particle-phase intermediate volatility and semi-volatile organic compounds using pixel-based approaches.. <i>Journal of Chromatography A</i> , 2022 , 1665, 462808	4.5	2
41	Linear Solvation Energy Relationships (LSERs) for Robust Prediction of Partition Coefficients between Low Density Polyethylene and Water Part I: Experimental Partition Coefficients and Model Calibration.. <i>European Journal of Pharmaceutical Sciences</i> , 2022 , 172, 106137	5.1	3
40	OctanolWater Partition Coefficients and Aqueous Solubility Data of Monoterpenoids: Experimental, Modeling, and Environmental Distribution. <i>Industrial & Engineering Chemistry Research</i> , 2022 , 61, 3154-3167	3.9	2

39	Descriptors for some compounds with pharmacological activity; calculation of properties.. <i>International Journal of Pharmaceutics</i> , 2022 , 617, 121597	6.5	0
38	An exploration into the use of Hansen solubility parameters for modelling reversed-phase chromatographic separations. <i>Journal of Analytical Science and Technology</i> , 2022 , 13,	3.4	
37	Prediction of heel build-up on activated carbon using machine learning.. <i>Journal of Hazardous Materials</i> , 2022 , 433, 128747	12.8	1
36	Quantum Chemical Calculation and Evaluation of Partition Coefficients for Classical and Emerging Environmentally Relevant Organic Compounds.. <i>Environmental Science & Technology</i> , 2021 ,	10.3	2
35	Applicability Domain of Polyparameter Linear Free Energy Relationship Models Evaluated by Leverage and Prediction Interval Calculation.. <i>Environmental Science & Technology</i> , 2022 ,	10.3	1
34	Applications of the solvation parameter model in thin-layer chromatography. <i>Journal of Planar Chromatography - Modern TLC</i> , 1	0.9	0
33	Temperature and indoor environments. <i>Indoor Air</i> , 2022 , 32,	5.4	0
32	In Silico Package Solutions for Deriving Values of Solute Parameters in Linear Solvation Energy Relationships. <i>SSRN Electronic Journal</i> ,	1	1
31	Hydrogen-Bonding Interactions in Polymer/Organic Solvent Mixtures. <i>Macromolecules</i> , 2022 , 55, 4578-4588	9.5	2
30	Mobile phases and their properties. 2022 , 207-269		
29	Reversed-phase HPLC. 2022 , 341-419		
28	Selectivity Map of Stationary Phases: A Graphical Method for Systematizing and Searching for Conditions for the Gas Chromatographic Separation of Polar Substances. <i>Journal of Analytical Chemistry</i> , 2022 , 77, 898-906	1.1	
27	Abraham Solvation Parameter Model: Calculation of L Solute Descriptors for Large C11 to C42 Methylated Alkanes from Measured Gas/Liquid Chromatographic Retention Data. <i>Liquids</i> , 2022 , 2, 85-105		1
26	Abraham model correlations for solute transfer into cyclopentanone. <i>Physics and Chemistry of Liquids</i> , 1-13	1.5	1
25	Acoustic Wave Sensors for Detection of Blister Chemical Warfare Agents and Their Simulants. <i>Sensors</i> , 2022 , 22, 5607	3.8	2
24	Ionic liquid and octadecylamine co-derived carbon dots for multi-mode high performance liquid chromatography. 2022 , 182, 107932		0
23	Activity coefficients at infinite dilution via a perturbation method of NRHB model. 2022 , 262, 118043		2
22	Strategies for structure elucidation of small molecules based on LC/MS/MS data from complex biological samples. 2022 , 20, 5085-5097		0

21	Novel concepts and challenges of flavor microencapsulation. 2023 , 427-450	0
20	Solvent selection for chemical reactions toward optimal thermodynamic and kinetic performances: Group contribution and COSMO-based modeling. 2023 , 564, 113623	0
19	Development of Abraham Model Correlations for Solute Transfer into the tert-Butyl Acetate Mono-Solvent and Updated Equations for Both Ethyl Acetate and Butyl Acetate. 2022 , 2, 258-288	2
18	Correlation of Surface Tension of Mono-Solvents at Various Temperatures. 2022 , 2, 378-387	0
17	Volatility and Nonspecific van der Waals Interaction Properties of Per- and Polyfluoroalkyl Substances (PFAS): Evaluation Using Hexadecane/Air Partition Coefficients.	1
16	Models Used to Predict Chemical Bioaccumulation in Fish from In Vitro Biotransformation Rates Require Accurate Estimates of Blood-Water Partitioning and Chemical Volume of Distribution.	0
15	Solid-liquid equilibrium solubility prediction of sulfanilamide in four binary solvent mixtures by using pure solvents solubility data from 278.15 to 318.15 K with the Abraham solvation parameter model, Yalkowsky Log-Linear and extended log-linear solubility thermodynamic models. 2022 , 368, 120634	0
14	Prediction of organic contaminant rejection by nanofiltration and reverse osmosis membranes using interpretable machine learning models. 2023 , 857, 159348	0
13	Recent advances for estimating environmental properties for small molecules from chromatographic measurements and the solvation parameter model. 2023 , 1687, 463682	1
12	The surface site interaction point approach to non-covalent interactions. 2022 , 51, 10064-10082	1
11	Linear Solvation Energy Relationships (LSER) and Equation-of-State Thermodynamics: On the Extraction of Thermodynamic Information from the LSER Database. 2023 , 3, 66-89	0
10	In silico package models for deriving values of solute parameters in linear solvation energy relationships. 1-17	0
9	Solvent selection and method development for thin-layer chromatography. 2023 , 47-80	0
8	Determination of Abraham Model Solute Descriptors for 62 Additional C10 through C13 Methyl- and Ethyl-Branched Alkanes. 2023 , 3, 118-131	0
7	Linear Free-Energy Relationships and Solvation Thermodynamics: The Thermodynamic Basis of LFER Linearity. 2023 , 62, 2989-3000	0
6	The influence of descriptor database selection on the solvation parameter model for separation processes. 2023 , 1692, 463851	0
5	A machine learning based approach for estimating site-specific partition coefficient K _d of organic compounds: Application to nonionic pesticides. 2023 , 323, 121297	0
4	Modeling the Solubility of Monoterpenoids with Hybrid and Predictive Thermodynamic Tools. 2023 , 62, 5326-5335	0

- 3 Selectivity evaluation of extraction systems. **2023**, 1695, 463939 ○
- 2 Electron ionization mass spectrometry fragmentation routes of Chemical Weapons Convention-related organoarsenic compounds: Electron ionization and density functional theory studies. **2023**, 37, ○
- 1 Comprehensive Two-Dimensional Gas Chromatography as a Bioanalytical Platform for Drug Discovery and Analysis. **2023**, 15, 1121 ○