New interdisciplinary cognitive-behavioural-nutritions 5-year follow-up study

Eating and Weight Disorders 9, 29-34

DOI: 10.1007/bf03325042

Citation Report

#	Article	IF	CITATIONS
1	Interest of pronostic score for optimal clinical management of obese patients. Diabetes and Metabolism, 2003, 29, 418-423.	1.4	4
2	Effect of Orlistat in Obese Patients with Binge Eating Disorder. Obesity, 2005, 13, 1701-1708.	4.0	94
3	Interdisciplinary rehabilitation in morbidly obese subjects: an observational pilot study. Monaldi Archives for Chest Disease, 2006, 65, 89-95.	0.3	7
4	Dietary and Body Weight Control: Therapeutic Education, Motivational Interviewing and Cognitive-Behavioral Approaches for Long-Term Weight Loss Maintenance., 2006, 11, 127-137.		5
5	Une éducation thérapeutique en quatre dimensions pour diminuer la résistance au changement des patients obÃ"ses. Medecine Des Maladies Metaboliques, 2007, 1, 69-76.	0.1	5
6	Home-centred physical fitness programme in morbidly obese individuals: a randomized controlled trial. Clinical Rehabilitation, 2008, 22, 940-950.	1.0	17
7	Treatment Alternatives for Overweight and Obesity: The Role of Online Interventions. Behaviour Change, 2008, 25, 1-14.	0.6	17
8	Helping Clients Think Through Their Causal Models. Quality Management in Health Care, 2008, 17, 66-79.	0.4	1
9	Adding cognitive behavioural treatment to either low-carbohydrate or low-fat diets: differential short-term effects. British Journal of Nutrition, 2009, 102, 1847-1853.	1.2	26
10	Efficacy of therapeutic patient education in chronic diseases and obesity. Patient Education and Counseling, 2010, 79, 283-286.	1.0	139
11	New psycho-pedagogic approach to obesity treatment: A 5-year follow-up. Patient Education and Counseling, 2010, 79, 333-337.	1.0	21
12	Psychological factors influencing weight loss maintenance: An integrative literature review. Journal of the American Academy of Nurse Practitioners, 2011, 23, 592-601.	1.4	56
13	Multi-Factorial Approach Associated with a New â€~On/Off' Orlistat® Use in a Weight Loss Maintenance Programme: 4 Years Follow-Up. Obesity Facts, 2011, 4, 191-196.	1.6	2
14	Do Obese Patients after Weight Loss Become Metabolically Normal?. Obesity Facts, 2011, 4, 218-221.	1.6	6
15	Using cognitiveâ€behavioural techniques to improve exclusive breastfeeding in a lowâ€literacy disadvantaged population. Maternal and Child Nutrition, 2012, 8, 57-71.	1.4	29
16	If it does not significantly change HbA _{1c} levels why should we waste time on it? A plea for the prioritization of psychological wellâ€being in people with diabetes. Diabetic Medicine, 2015, 32, 155-163.	1.2	70
17	Sustainable Lifestyle Changeâ€"Participatory Design of Support Together with Persons with Obesity in the Third Age. International Journal of Environmental Research and Public Health, 2016, 13, 1248.	1.2	4
18	Dance therapy combined with patient education improves quality of life of persons with obesity: A pilot feasibility study for a randomised controlled trial. Obesity Research and Clinical Practice, 2017, 11, 79-87.	0.8	13

#	Article	IF	CITATIONS
19	Bariatric surgery versus lifestyle interventions for severe obesity: 5â€year changes in body weight, risk factors and comorbidities. Clinical Obesity, 2017, 7, 183-190.	1.1	19
20	The challenge of keeping it off, a descriptive systematic review of highâ€quality, followâ€up studies of obesity treatments. Obesity Reviews, 2020, 21, e12949.	3.1	76
21	L'efficacité médico-économique de l'éducation thérapeutique chez des patients obà ses. Education Therapeutique Du Patient, 2009, 1, 57-62.	0.5	7
22	Effects of Lifestyle Interventions That Include a Physical Activity Component in Class II and III Obese Individuals: A Systematic Review and Meta-Analysis. PLoS ONE, 2015, 10, e0119017.	1.1	98
24	The vital link between chronic disease and depressive disorders. Preventing Chronic Disease, 2005, 2, A14.	1.7	163
25	Understanding weight regain after a nutritional weight loss intervention: Systematic review and meta-analysis. Clinical Nutrition ESPEN, 2022, 49, 138-153.	0.5	15