Fluoxetine-induced change in rat brain expression of br varies depending on length of treatment

Neuroscience 128, 597-604

DOI: 10.1016/j.neuroscience.2004.06.054

Citation Report

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Chronic fluoxetine administration inhibits extracellular signal-regulated kinase 1/2 phosphorylation in rat brain. Journal of Neurochemistry, 2005, 93, 1551-1560. | 2.1 | 98 |
| 2 | New drug targets for depression and anxiety: Is the peptides era arriving?. Drug Development Research, 2005, 65, 93-96. | 1.4 | 2 |
| 3 | Chronic administration of the delta opioid receptor agonist (+)BW373U86 and antidepressants on behavior in the forced swim test and BDNF mRNA expression in rats. Psychopharmacology, 2005, 183, 31-40. | 1.5 | 30 |
| 4 | Possible involvement of brain-derived neurotrophic factor in the antinociceptive effect of antidepressants in neuropathic pain. Medical Hypotheses, 2005, 65, 530-533. | 0.8 | 9 |
| 5 | Possible involvement of the BDNF-dependent pathway in treatment-emergent suicidality or decreased response to antidepressants. Medical Hypotheses, 2005, 65, 942-946. | 0.8 | 19 |
| 6 | Sequential changes in BDNF mRNA expression and synaptic levels of AMPA receptor subunits in rat hippocampus after chronic antidepressant treatment. Neuropharmacology, 2005, 49, 1178-1188. | 2.0 | 85 |
| 7 | Signaling Pathways Regulating Gene Expression, Neuroplasticity, and Neurotrophic Mechanisms in the Action of Antidepressants: A Critical Overview. Pharmacological Reviews, 2006, 58, 115-134. | 7.1 | 270 |
| 8 | A Neurotrophic Model for Stress-Related Mood Disorders. Biological Psychiatry, 2006, 59, 1116-1127. | 0.7 | 2,873 |
| 9 | Peripheral immune activation by lipopolysaccharide decreases neurotrophins in the cortex and hippocampus in rats. Brain, Behavior, and Immunity, 2006, 20, 64-71. | 2.0 | 206 |
| 10 | Lithium and antidepressants: Potential agents for the treatment of Rett syndrome. Medical Hypotheses, 2006, 67, 626-629. | 0.8 | 17 |
| 11 | Antidepressants reverse corticosterone-mediated decrease in brain-derived neurotrophic factor expression: Differential regulation of specific exons by antidepressants and corticosterone. Neuroscience, 2006, 139, 1017-1029. | 1.1 | 180 |
| 12 | Chronic fluoxetine induces region-specific changes in translation factor eIF4E and eEF2 activity in the rat brain. European Journal of Neuroscience, 2006, 23, 2814-2818. | 1.2 | 31 |
| 13 | Chronic fluoxetine upregulates activity, protein and mRNA levels of cytosolic phospholipase A2 in rat frontal cortex. Pharmacogenomics Journal, 2006, 6, 413-420. | 0.9 | 40 |
| 14 | Repeated electroconvulsive stimuli increase brain-derived neurotrophic factor in ACTH-treated rats. European Journal of Pharmacology, 2006, 529, 114-121. | 1.7 | 44 |
| 15 | Peptidic delta opioid receptor agonists produce antidepressant-like effects in the forced swim test and regulate BDNF mRNA expression in rats. Brain Research, 2006, 1069, 172-181. | 1.1 | 87 |
| 16 | Biphasic change in BDNF gene expression following antidepressant drug treatment explained by differential transcript regulation. Brain Research, 2006, 1106, 12-20. | 1.1 | 62 |
| 17 | Multi-target strategies for the improved treatment of depressive states: Conceptual foundations and neuronal substrates, drug discovery and therapeutic application. , 2006, 110, 135-370. | | 483 |
| 18 | Chronic treatment with fluoxetine up-regulates cellular BDNF mRNA expression in rat dopaminergic regions. International Journal of Neuropsychopharmacology, 2006, 9, 307. | 1.0 | 103 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Metabotropic Glutamate Receptor Subtype 7 Ablation Causes Dysregulation of the HPA Axis and Increases Hippocampal BDNF Protein Levels: Implications for Stress-Related Psychiatric Disorders. Neuropsychopharmacology, 2006, 31, 1112-1122. | 2.8 | 122 |
| 20 | Chronic Duloxetine Treatment Induces Specific Changes in the Expression of BDNF Transcripts and in the Subcellular Localization of the Neurotrophin Protein. Neuropsychopharmacology, 2007, 32, 2351-2359. | 2.8 | 110 |
| 22 | Brain-Derived Neurotrophic Factor and Major Depression. Psychopharm Review: Timely Reports in Psychopharmacology and Device-based Therapies, 2007, 42, 59-66. | 0.1 | 0 |
| 23 | Psychopharm Review: CME QUIZ. Psychopharm Review: Timely Reports in Psychopharmacology and Device-based Therapies, 2007, 42, 66. | 0.1 | 0 |
| 24 | AMPA Receptor Potentiators: Application for Depression and Parkinsons Disease. Current Drug Targets, 2007, 8, 603-620. | 1.0 | 93 |
| 25 | A Clinical View of BDNF-TrkB Signaling in the Treatment of Major Depression. Current Signal Transduction Therapy, 2007, 2, 186-189. | 0.3 | 1 |
| 26 | Wheel running and fluoxetine antidepressant treatment have differential effects in the hippocampus and the spinal cord. Neuroscience, 2007, 144, 1033-1044. | 1.1 | 66 |
| 27 | Methylphenidate regulates activity regulated cytoskeletal associated but not brain-derived neurotrophic factor gene expression in the developing rat striatum. Neuroscience, 2007, 144, 969-984. | 1.1 | 50 |
| 28 | Acute onset by 5-HT6-receptor activation on rat brain brain-derived neurotrophic factor and activity-regulated cytoskeletal-associated protein mRNA expression. Neuroscience, 2007, 147, 778-785. | 1.1 | 61 |
| 29 | Effects of fluoxetine on ischemic cells and expressions in BDNF and some antioxidants in the gerbil hippocampal CA1 region induced by transient ischemia. Experimental Neurology, 2007, 204, 748-758. | 2.0 | 68 |
| 30 | Measurements of brain-derived neurotrophic factor: Methodological aspects and demographical data. Brain Research Bulletin, 2007, 73, 143-149. | 1.4 | 178 |
| 31 | Antidepressant like effects of piperine in chronic mild stress treated mice and its possible mechanisms. Life Sciences, 2007, 80, 1373-1381. | 2.0 | 207 |
| 32 | A microarray gene expression study of the molecular pharmacology of lithium carbonate on mouse brain mRNA to understand the neurobiology of mood stabilization and treatment of bipolar affective disorder. Pharmacogenetics and Genomics, 2007, 17, 605-617. | 0.7 | 133 |
| 33 | Antidepressant-Like Effects of the Histone Deacetylase Inhibitor, Sodium Butyrate, in the Mouse. Biological Psychiatry, 2007, 62, 55-64. | 0.7 | 462 |
| 34 | Regulation of Serotonin-1A Receptor Function in Inducible Brain-Derived Neurotrophic Factor Knockout Mice After Administration of Corticosterone. Biological Psychiatry, 2007, 62, 521-529. | 0.7 | 45 |
| 36 | Is it time to reassess the BDNF hypothesis of depression?. Molecular Psychiatry, 2007, 12, 1079-1088. | 4.1 | 369 |
| 38 | BDNF Level in the Rat Prefrontal Cortex Increases Following Chronic but Not Acute Treatment with Duloxetine, a Dual Acting Inhibitor of Noradrenaline and Serotonin Re-uptake. Cellular and Molecular Neurobiology, 2008, 28, 457-468. | 1.7 | 51 |
| 39 | S32006, a novel 5-HT2C receptor antagonist displaying broad-based antidepressant and anxiolytic properties in rodent models. Psychopharmacology, 2008, 199, 549-568. | 1.5 | 109 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 40 | Changes in rat hippocampal CA1 synapses following imipramine treatment. Hippocampus, 2008, 18, 631-639. | 0.9 | 48 |
| 41 | Chronic Administration of Ketamine Elicits Antidepressantâ€Like Effects in Rats without Affecting Hippocampal Brainâ€Derived Neurotrophic Factor Protein Levels. Basic and Clinical Pharmacology and Toxicology, 2008, 103, 502-506. | 1.2 | 101 |
| 42 | Evidence of biologic epistasis between BDNF and SLC6A4 and implications for depression. Molecular Psychiatry, 2008, 13, 709-716. | 4.1 | 216 |
| 43 | Brainâ€derived neurotropic factor and neurogenesis in the adult rat dentate gyrus: interactions with corticosterone. European Journal of Neuroscience, 2008, 27, 2493-2500. | 1.2 | 59 |
| 44 | Time-dependent biphasic modulation of human BDNF by antidepressants in neuroblastoma cells. BMC Neuroscience, 2008, 9, 61. | 0.8 | 25 |
| 45 | Fluoxetine-induced proliferation and differentiation of neural progenitor cells isolated from rat postnatal cerebellum. Biochemical Pharmacology, 2008, 76, 391-403. | 2.0 | 37 |
| 46 | Differential regulation of central BDNF protein levels by antidepressant and non-antidepressant drug treatments. Brain Research, 2008, 1211, 37-43. | 1.1 | 173 |
| 47 | Temporal expression of brain-derived neurotrophic factor (BDNF) mRNA in the rat hippocampus after treatment with selective and mixed monoaminergic antidepressants. European Journal of Pharmacology, 2008, 578, 114-122. | 1.7 | 67 |
| 48 | Plasma and serum brain-derived neurotrophic factor (BDNF) in depressed patients during 1Âyear of antidepressant treatments. Journal of Affective Disorders, 2008, 105, 279-283. | 2.0 | 183 |
| 49 | Brain-derived neurotrophic factor and its receptor tropomyosin-related kinase B in the mechanism of action of antidepressant therapies. , 2008, 117, 30-51. | | 173 |
| 50 | Nutritional Tryptophan Restriction and the Role of Serotonin in Development and Plasticity of Central Visual Connections. NeuroImmunoModulation, 2008, 15, 170-175. | 0.9 | 24 |
| 51 | The differential regulation of BDNF and TrkB levels in juvenile rats after four days of escitalopram and desipramine treatment. Neuropharmacology, 2008, 54, 251-257. | 2.0 | 42 |
| 52 | Acute administration of ketamine induces antidepressant-like effects in the forced swimming test and increases BDNF levels in the rat hippocampus. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2008, 32, 140-144. | 2.5 | 377 |
| 53 | Acute tryptophan depletion does not alter central or plasma brain-derived neurotrophic factor in the rat. European Neuropsychopharmacology, 2008, 18, 317-322. | 0.3 | 12 |
| 54 | Relationship between antidepressants and IGF-1 system in the brain: Possible role in cognition. European Neuropsychopharmacology, 2008, 18, 431-438. | 0.3 | 35 |
| 55 | Maintenance Treatment with Fluoxetine is Necessary to Sustain Normal Levels of Synaptic Markers in an Experimental Model of Depression: Correlation with Behavioral Response. Neuropsychopharmacology, 2008, 33, 1896-1908. | 2.8 | 71 |
| 56 | Serum Concentrations of Nerve Growth Factor and Brain-Derived Neurotrophic Factor in Depressed Patients before and after Antidepressant Treatment. Pharmacopsychiatry, 2008, 41, 66-71. | 1.7 | 115 |
| 57 | Genetic Regulation of Behavioral and Neuronal Responses to Fluoxetine. Neuropsychopharmacology, 2008, 33, 1312-1322. | 2.8 | 83 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 58 | Brain-derived neurotrophic factor: role in depression and suicide. Neuropsychiatric Disease and Treatment, 2009, 5, 433. | 1.0 | 276 |
| 59 | Early induction of CREB activation and CREB-regulating signalling by antidepressants. International Journal of Neuropsychopharmacology, 2009, 12, 1367. | 1.0 | 40 |
| 60 | Ketamine induces rapid onset of antidepressant action: neurophysiological biomarkers as predictors of effect. Biomarkers in Medicine, 2009, 3, 5-8. | 0.6 | 3 |
| 61 | Changes in BDNF serum levels in patients with major depression disorder (MDD) after 6 months treatment with sertraline, escitalopram, or venlafaxine. Journal of Psychiatric Research, 2009, 43, 247-254. | 1.5 | 212 |
| 62 | Early raise of BDNF in hippocampus suggests induction of posttranscriptional mechanisms by antidepressants. BMC Neuroscience, 2009, 10, 48. | 0.8 | 53 |
| 63 | Effects of escitalopram on the regulation of brainâ€derived neurotrophic factor and nerve growth factor protein levels in a rat model of chronic stress. Journal of Neuroscience Research, 2009, 87, 2551-2560. | 1.3 | 42 |
| 64 | Inhibition of serotonin and norepinephrine reuptake and inhibition of phosphodiesterase by multi-target inhibitors as potential agents for depression. Bioorganic and Medicinal Chemistry, 2009, 17, 6890-6897. | 1.4 | 30 |
| 65 | Targeting the Hippocampal Mossy Fiber Synapse for the Treatment of Psychiatric Disorders. Molecular Neurobiology, 2009, 39, 24-36. | 1.9 | 63 |
| 66 | Depression and antidepressants: molecular and cellular aspects. Cellular and Molecular Life Sciences, 2009, 66, 2985-3008. | 2.4 | 70 |
| 67 | Antiapoptotic and neurotrophic effects of antidepressants: A review of clinical and experimental studies. Brain Research Bulletin, 2009, 79, 248-257. | 1.4 | 76 |
| 68 | Synergistic effects of dehydroepiandrosterone and fluoxetine on proliferation of progenitor cells in the dentate gyrus of the adult male rat. Neuroscience, 2009, 158, 1644-1651. | 1.1 | 34 |
| 69 | Venlafaxine treatment stimulates expression of brain-derived neurotrophic factor protein in frontal cortex and inhibits long-term potentiation in hippocampus. Neuroscience, 2009, 162, 1411-1419. | 1.1 | 36 |
| 70 | Antidepressant effects of ginseng total saponins in the forced swimming test and chronic mild stress models of depression. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2009, 33, 1417-1424. | 2.5 | 194 |
| 71 | Acute harmine administration induces antidepressive-like effects and increases BDNF levels in the rat hippocampus. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2009, 33, 1425-1430. | 2.5 | 109 |
| 72 | Chronic imipramine downregulates cyclic AMP signaling in rat hippocampus. NeuroReport, 2009, 20, 307-311. | 0.6 | 9 |
| 73 | Chronic administration of harmine elicits antidepressant-like effects and increases BDNF levels in rat hippocampus. Journal of Neural Transmission, 2010, 117, 1131-1137. | 1.4 | 85 |
| 74 | Regulation of brain-derived neurotrophic factor (BDNF) in the chronic unpredictable stress rat model and the effects of chronic antidepressant treatment. Journal of Psychiatric Research, 2010, 44, 808-816. | 1.5 | 137 |
| 75 | Chronic antidepressant administration alleviates frontal and hippocampal BDNF deficits in CUMS rat. Brain Research, 2010, 1366, 141-148. | 1.1 | 114 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 76 | Research in people with psychosis risk syndrome: a review of the current evidence and future directions. Journal of Child Psychology and Psychiatry and Allied Disciplines, 2010, 51, 390-431. | 3.1 | 167 |
| 77 | Depressionâ€resistant endophenotype in mice overexpressing cannabinoid CB ₂ receptors. British Journal of Pharmacology, 2010, 160, 1773-1784. | 2.7 | 169 |
| 78 | The Roles of BDNF in the Pathophysiology of Major Depression and in Antidepressant Treatment. Psychiatry Investigation, 2010, 7, 231. | 0.7 | 364 |
| 79 | A Role for p11 in the Antidepressant Action of Brain-Derived Neurotrophic Factor. Biological Psychiatry, 2010, 68, 528-535. | 0.7 | 83 |
| 80 | Neurochemical and behavioural effects of acute and chronic memantine administration in rats: Further support for NMDA as a new pharmacological target for the treatment of depression?. Brain Research Bulletin, 2010, 81, 585-589. | 1.4 | 97 |
| 81 | Promoter specific alterations of brain-derived neurotrophic factor mRNA in schizophrenia. Neuroscience, 2010, 169, 1071-1084. | 1.1 | 87 |
| 82 | Exposure of mice to long-light: A new animal model to study depression. European Neuropsychopharmacology, 2010, 20, 802-812. | 0.3 | 12 |
| 83 | Neurotrophic Factors and Antidepressant Action: Recent Advances. Modern Problems of Pharmacopsychiatry, 2010, , 199-223. | 2.5 | 3 |
| 84 | Multiple Sclerosis and Depression: A Neuroimmunological Perspective. NeuroImmune Biology, 2010, , 269-307. | 0.2 | 1 |
| 85 | Potential role of the 5-HT6 receptor in depression and anxiety: an overview of preclinical data. Pharmacological Reports, 2010, 62, 564-577. | 1.5 | 106 |
| 86 | NO-SSRIs: Nitric Oxide Chimera Drugs Incorporating a Selective Serotonin Reuptake Inhibitor. ACS Medicinal Chemistry Letters, 2011, 2, 656-661. | 1.3 | 12 |
| 87 | Antidepressant Treatments Change 5-HT2C Receptor mRNA Expression in Rat Prefrontal/Frontal Cortex and Hippocampus. Neuropsychobiology, 2011, 63, 160-168. | 0.9 | 38 |
| 88 | Chronic antidepressant treatments induce a time-dependent up-regulation of AMPA receptor subunit protein levels. Neurochemistry International, 2011, 59, 896-905. | 1.9 | 61 |
| 89 | The molecular interaction between the glutamatergic, noradrenergic, dopaminergic and serotoninergic systems informs a detailed genetic perspective on depressive phenotypes. Progress in Neurobiology, 2011, 94, 418-460. | 2.8 | 61 |
| 90 | In animal models, psychosocial stress-induced (neuro)inflammation, apoptosis and reduced neurogenesis are associated to the onset of depression. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2011, 35, 744-759. | 2.5 | 369 |
| 91 | New Therapeutic Strategy for Mood Disorders. Current Medicinal Chemistry, 2011, 18, 4284-4298. | 1.2 | 16 |
| 92 | Dose-related effects of venlafaxine on pCREB and brain-derived neurotrophic factor (BDNF) in the hippocampus of the rat by chronic unpredictable stress. Acta Neuropsychiatrica, 2011, 23, 20-30. | 1.0 | 5 |
| 93 | Strain differences in the neurochemical response to chronic restraint stress in the rat: Relevance to depression. Pharmacology Biochemistry and Behavior, 2011, 97, 690-699. | 1.3 | 74 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 94 | Eszopiclone facilitation of the antidepressant efficacy of fluoxetine using a social defeat stress model. Pharmacology Biochemistry and Behavior, 2011, 99, 648-658. | 1.3 | 7 |
| 95 | Effects of GABAB ligands alone and in combination with paroxetine on hippocampal BDNF gene expression. European Journal of Pharmacology, 2011, 671, 33-38. | 1.7 | 14 |
| 96 | Protective Effects of Flavonoid Extract from Apocynum venetum Leaves Against Corticosterone-Induced Neurotoxicity in PC12 Cells. Cellular and Molecular Neurobiology, 2011, 31, 421-428. | 1.7 | 31 |
| 97 | Chronic ketamine use increases serum levels of brain-derived neurotrophic factor. Psychopharmacology, 2011, 215, 143-148. | 1.5 | 57 |
| 98 | The Effects of Fluoxetine Treatment in a Chronic Mild Stress Rat Model on Depression-Related Behavior, Brain Neurotrophins and ERK Expression. Journal of Molecular Neuroscience, 2011, 45, 246-255. | 1.1 | 106 |
| 99 | CART Peptides Regulate Psychostimulants and May be Endogenous Antidepressants. Current Neuropharmacology, 2011, 9, 12-16. | 1.4 | 12 |
| 100 | Behavioral Pharmacology: Potential Antidepressant and Anxiolytic Properties. International Review of Neurobiology, 2011, 96, 49-71. | 0.9 | 12 |
| 101 | Neurotrophins Role in Depression Neurobiology: A Review of Basic and Clinical Evidence. Current Neuropharmacology, 2011, 9, 530-552. | 1.4 | 130 |
| 102 | Dissociation of antidepressant-like activity of escitalopram and nortriptyline on behaviour and hippocampal BDNF expression in female rats. Journal of Psychopharmacology, 2011, 25, 1378-1387. | 2.0 | 22 |
| 103 | The impact of chronic imipramine treatment on amino acid concentrations in the hippocampus of mice. Nutritional Neuroscience, 2012, 15, 26-33. | 1.5 | 7 |
| 104 | Modulation of neuroplasticity pathways and antidepressant-like behavioural responses following the short-term (3 and 7 days) administration of the 5-HT4 receptor agonist RS67333. International Journal of Neuropsychopharmacology, 2012, 15, 631-643. | 1.0 | 76 |
| 105 | Interaction between the BDNF gene Val/66/Met polymorphism and morning cortisol levels as a predictor of depression in adult women. British Journal of Psychiatry, 2012, 201, 313-319. | 1.7 | 27 |
| 106 | Ginsenoside Rb3 exerts antidepressant-like effects in several animal models. Journal of Psychopharmacology, 2012, 26, 697-713. | 2.0 | 49 |
| 107 | Vascular Endothelial Growth Factor-dependent Spinogenesis Underlies Antidepressant-like Effects of Enriched Environment. Journal of Biological Chemistry, 2012, 287, 40938-40955. | 1.6 | 27 |
| 108 | Molecular adaptation to chronic antidepressant treatment: evidence for a more rapid response to the novel α2-adrenoceptor antagonist/5-HT-noradrenaline reuptake inhibitor (SNRI), S35966, compared to the SNRI, venlafaxine. International Journal of Neuropsychopharmacology, 2012, 15, 617-629. | 1.0 | 14 |
| 109 | Acute running stimulates hippocampal dopaminergic neurotransmission in rats, but has no influence on brain-derived neurotrophic factor. Journal of Applied Physiology, 2012, 112, 535-541. | 1.2 | 59 |
| 110 | Early intervention with fluoxetine reverses abnormalities in the serotonergic system and behavior of rats exposed prenatally to dexamethasone. Neuropharmacology, 2012, 63, 292-300. | 2.0 | 53 |
| 111 | The genetics of selective serotonin reuptake inhibitors. , 2012, 136, 375-400. | | 38 |

ARTICLE IF CITATIONS # Preclinical discovery of duloxetine for the treatment of depression. Expert Opinion on Drug 112 2.5 9 Discovery, 2012, 7, 745-755. Analysis of region-specific changes in gene expression upon treatment with citalopram and desipramine reveals temporal dynamics in response to antidepressant drugs at the transcriptome 1.5 level. Psychopharmacology, 2012, 223, 281-297. Brain-derived neurotrophic factor and exercise in fibromyalgia syndrome patients: a mini review. 114 1.5 41 Rheumatology International, 2012, 32, 2593-2599. lmipramine treatment reverses depressiveâ€like behavior in alloxanâ€diabetic rats. Diabetes/Metabolism Research and Reviews, 2012, 28, 139-144. Subchronic treatment with fluoxetine and ketanserin increases hippocampal brainâ€derived neurotrophic factor, βâ€catenin and antidepressantâ€like effects. British Journal of Pharmacology, 2012, 117 2.7 55 165, 1046-1057. Fluoxetine rescues impaired hippocampal neurogenesis in a transgenic A53T synuclein mouse model. European Journal of Neuroscience, 2012, 35, 10-19. 1.2 Effects of repeated 5-HT6 receptor stimulation on BDNF gene expression and cell survival. 119 1.0 5 Neuroscience Letters, 2013, 553, 211-215. Icaritin opposes the development of social aversion after defeat stress via increases of GR mRNA and 120 1.2 44 BDNF mRNA in mice. Behavioural Brain Research, 2013, 256, 602-608. Essential oil of Perilla frutescens-induced change in hippocampal expression of brain-derived 121 neurotrophic factor in chronic unpredictable mild stress in mice. Journal of Ethnopharmacology, 2.0 41 2013, 147, 245-253. Involvement of Brain-Derived Neurotrophic Factor in Late-Life Depression. American Journal of Geriatric Psychiatry, 2013, 21, 433-449. Activity-dependent, stress-responsive BDNF signaling and the quest for optimal brain health and 123 1.1 130 resilience throughout the lifespan. Neuroscience, 2013, 239, 228-240. Antidepressant-like effect of macranthol isolated from Illicium dunnianum tutch in mice. European 124 29 Journal of Pharmacology, 2013, 707, 112-119. Tetrandrine exerts antidepressant-like effects in animal models: Role of brain-derived neurotrophic 125 1.2 43 factor. Behavioural Brain Research, 2013, 238, 79-85. Lithium augmentation of the effects of desipramine in a mouse model of treatment-resistant depression: A role for hippocampal cell proliferation. Neuroscience, 2013, 228, 36-46. 1.1 Antidepressant-like effect of oleanolic acid in mice exposed to the repeated forced swimming test. 127 2.0 24 Journal of Psychopharmacology, 2013, 27, 459-468. The role of 5-hydroxytryptamine receptor subtypes in the regulation of <i>brain-derived neurotrophic 1.2 factor </i> gene expression. Journal of Pharmacy and Pharmacology, 2013, 66, 53-61. Captodiamine, a putative antidepressant, enhances hypothalamic BDNF expression in vivo by synergistic 129 5-HT_{2c} receptor antagonism and sigma-1 receptor agonism. Journal of 2.0 16 Psychopharmacology, 2013, 27, 930-939. Ketamine alters behavior and decreases BDNF levels in the rat brain as a function of time after drug administration. Revista Brasileira De Psiquiatria, 2013, 35, 262-266.

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 131 | Long-Term Adaptive Changes Induced by Antidepressants: From Conventional to Novel Therapies. , 0, , . | | 0 |
| 132 | Fluoxetine Induces Proliferation and Inhibits Differentiation of Hypothalamic Neuroprogenitor Cells In Vitro. PLoS ONE, 2014, 9, e88917. | 1.1 | 11 |
| 133 | Contingency-based emotional resilience: effort-based reward training and flexible coping lead to adaptive responses to uncertainty in male rats. Frontiers in Behavioral Neuroscience, 2014, 8, 124. | 1.0 | 29 |
| 134 | Effect of fluoxetine on disease progression in a mouse model of ALS. Journal of Neurophysiology, 2014, 111, 2164-2176. | 0.9 | 19 |
| 135 | Fluoxetine treatment ameliorates depression induced by perinatal arsenic exposure via a neurogenic mechanism. NeuroToxicology, 2014, 44, 98-109. | 1.4 | 24 |
| 136 | Effects of negative stressors on DNA methylation in the brain: Implications for mood and anxiety disorders. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 2014, 165, 541-554. | 1.1 | 32 |
| 137 | Exercise as an add-on strategy for the treatment of major depressive disorder: a systematic review. CNS Spectrums, 2014, 19, 496-508. | 0.7 | 74 |
| 138 | High baseline BDNF serum levels and early psychopathological improvement are predictive of treatment outcome in major depression. Psychopharmacology, 2014, 231, 2955-2965. | 1.5 | 56 |
| 139 | The Antidepressant Fluoxetine Mobilizes Vesicles to the Recycling Pool of Rat Hippocampal Synapses During High Activity. Molecular Neurobiology, 2014, 49, 916-930. | 1.9 | 12 |
| 140 | BDNF signaling is necessary for the antidepressant-like effect of naringenin. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2014, 48, 135-141. | 2.5 | 64 |
| 141 | Antidepressants that inhibit both serotonin and norepinephrine reuptake impair long-term potentiation in hippocampus. Psychopharmacology, 2014, 231, 4429-4441. | 1.5 | 14 |
| 142 | Effects of antidepressant drug exposure on gene expression in the developing cerebral cortex. Synapse, 2014, 68, 209-220. | 0.6 | 10 |
| 143 | Chronic social instability stress enhances vulnerability of BDNF response to LPS in the limbic structures of female rats: A protective role of antidepressants. Neuroscience Research, 2014, 88, 74-83. | 1.0 | 22 |
| 144 | Roles of olfactory system dysfunction in depression. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2014, 54, 26-30. | 2.5 | 75 |
| 145 | New insight in expression, transport, and secretion of brain-derived neurotrophic factor: Implications in brain-related diseases. World Journal of Biological Chemistry, 2014, 5, 409. | 1.7 | 132 |
| 146 | Long-term consequences of chronic fluoxetine exposure on the expression of myelination-related genes in the rat hippocampus. Translational Psychiatry, 2015, 5, e642-e642. | 2.4 | 24 |
| 147 | Neuropeptide Trefoil Factor 3 Reverses Depressive-Like Behaviors by Activation of BDNF-ERK-CREB Signaling in Olfactory Bulbectomized Rats. International Journal of Molecular Sciences, 2015, 16, 28386-28400. | 1.8 | 29 |
| 148 | A critical evaluation of the activity-regulated cytoskeleton-associated protein (Arc/Arg3.1)'s putative role in regulating dendritic plasticity, cognitive processes, and mood in animal models of depression. Frontiers in Neuroscience, 2015, 9, 279. | 1.4 | 65 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 149 | Epigenetic mechanisms underlying the role of brain-derived neurotrophic factor in depression and response to antidepressants. Journal of Experimental Biology, 2015, 218, 21-31. | 0.8 | 82 |
| 150 | Tricyclic antidepressant treatment evokes regional changes in neurotrophic factors over time within the intact and degenerating nigrostriatal system. Experimental Neurology, 2015, 266, 11-21. | 2.0 | 22 |
| 151 | Antidepressant-like effect of zinc is dependent on signaling pathways implicated in BDNF modulation. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2015, 59, 59-67. | 2.5 | 36 |
| 152 | Is serotonin an upper or a downer? The evolution of the serotonergic system and its role in depression and the antidepressant response. Neuroscience and Biobehavioral Reviews, 2015, 51, 164-188. | 2.9 | 214 |
| 153 | Antidepressant activity of fluoxetine in the zinc deficiency model in rats involves the NMDA receptor complex. Behavioural Brain Research, 2015, 287, 323-330. | 1.2 | 27 |
| 154 | Fluoxetine Maintains a State of Heightened Responsiveness to Motor Training Early After Stroke in a Mouse Model. Stroke, 2015, 46, 2951-2960. | 1.0 | 75 |
| 155 | Brain-derived neurotrophic factor correlates with functional and cognitive impairment in non-disabled older individuals. Experimental Gerontology, 2015, 72, 129-137. | 1.2 | 17 |
| 156 | Chronic Amitriptyline Treatment Attenuates Nigrostriatal Degeneration and Significantly Alters Trophic Support in a Rat Model of Parkinsonism. Neuropsychopharmacology, 2015, 40, 874-883. | 2.8 | 21 |
| 157 | Effect of using fluoxetine at different time windows on neurological functional prognosis after ischemic stroke. Restorative Neurology and Neuroscience, 2016, 34, 177-187. | 0.4 | 21 |
| 158 | Sleep disturbance as detected by actigraphy in pre-pubertal juvenile monkeys receiving therapeutic doses of fluoxetine. Neurotoxicology and Teratology, 2016, 55, 1-7. | 1.2 | 16 |
| 159 | Peripheral fibroblast metabolic pathway alterations in juvenile rhesus monkeys undergoing long-term fluoxetine administration. European Neuropsychopharmacology, 2016, 26, 1110-1118. | 0.3 | 11 |
| 161 | The absence of 5-HT4 receptors modulates depression- and anxiety-like responses and influences the response of fluoxetine in olfactory bulbectomised mice: Adaptive changes in hippocampal neuroplasticity markers and 5-HT1A autoreceptor. Neuropharmacology, 2016, 111, 47-58. | 2.0 | 48 |
| 162 | Blunted response of hippocampal AMPK associated with reduced neurogenesis in older versus younger mice. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2016, 71, 57-65. | 2.5 | 12 |
| 163 | Effects of Fluoxetine on Neural Functional Prognosis after Ischemic Stroke: A Randomized Controlled Study in China. Journal of Stroke and Cerebrovascular Diseases, 2016, 25, 761-770. | 0.7 | 27 |
| 164 | Chronic stress, hippocampus and parvalbumin-positive interneurons: what do we know so far?. Reviews in the Neurosciences, 2016, 27, 397-409. | 1.4 | 49 |
| 165 | Gestational stress induces depressive-like and anxiety-like phenotypes through epigenetic regulation of BDNF expression in offspring hippocampus. Epigenetics, 2016, 11, 150-162. | 1.3 | 76 |
| 166 | Piperine potentiates the effects of trans-resveratrol on stress-induced depressive-like behavior: involvement of monoaminergic system and cAMP-dependent pathway. Metabolic Brain Disease, 2016, 31, 837-848. | 1.4 | 21 |
| 167 | Hippocampal BDNF in physiological conditions and social isolation. Reviews in the Neurosciences, 2017, 28, 675-692. | 1.4 | 88 |
| | | | |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 168 | Translatable Models of Brain and Cognitive Reserve. , 2017, , 79-104. | | 1 |
| 170 | Expression of <scp>BDNF</scp> and trkB in the hippocampus of a rat genetic model of vulnerability (Roman lowâ€avoidance) and resistance (Roman highâ€avoidance) to stressâ€induced depression. Brain and Behavior, 2017, 7, e00861. | 1.0 | 31 |
| 171 | Effect of co-administration of memantine and sertraline on the antidepressant-like activity and brain-derived neurotrophic factor (BDNF) levels in the rat brain. Brain Research Bulletin, 2017, 128, 29-33. | 1.4 | 28 |
| 172 | Computational Model of Antidepressant Response Heterogeneity as Multi-pathway Neuroadaptation. Frontiers in Pharmacology, 2017, 8, 925. | 1.6 | 10 |
| 173 | Immediate-Early Genes Modulation by Antipsychotics: Translational Implications for a Putative Gateway to Drug-Induced Long-Term Brain Changes. Frontiers in Behavioral Neuroscience, 2017, 11, 240. | 1.0 | 35 |
| 174 | Antidepressant-Like Effect of Lipid Extract ofChanna striatusin Postpartum Model of Depression in Rats. Evidence-based Complementary and Alternative Medicine, 2017, 2017, 1-15. | 0.5 | 7 |
| 175 | Zinc in the Monoaminergic Theory of Depression: Its Relationship to Neural Plasticity. Neural Plasticity. Plasticity, 2017, 2017, 1-18. | 1.0 | 58 |
| 176 | 5-HT2A Receptors and BDNF Regulation: Implications for Psychopathology. , 2018, , 395-438. | | 7 |
| 177 | Early Selective Serotonin Reuptake Inhibitors for Recovery after Stroke: A Meta-Analysis and Trial Sequential Analysis. Journal of Stroke and Cerebrovascular Diseases, 2018, 27, 1178-1189. | 0.7 | 42 |
| 178 | The potential benefit of combined versus monotherapy of coenzyme Q10 and fluoxetine on depressive-like behaviors and intermediates coupled to Gsk-3β in rats. Toxicology and Applied Pharmacology, 2018, 340, 39-48. | 1.3 | 15 |
| 179 | Chronic antidepressant-like effect of EMD386088, a partial 5-HT6 receptor agonist, in olfactory bulbectomy model may be connected with BDNF and/or CREB signalling pathway. Pharmacological Reports, 2018, 70, 1047-1056. | 1.5 | 5 |
| 180 | The effect of exercise on resting concentrations of peripheral brain-derived neurotrophic factor (BDNF) in major depressive disorder: A meta-analysis. Journal of Psychiatric Research, 2018, 105, 123-131. | 1.5 | 34 |
| 181 | Agomelatine-induced modulation of brain-derived neurotrophic factor (BDNF) in the rat hippocampus. Life Sciences, 2018, 210, 177-184. | 2.0 | 61 |
| 182 | Effects of Fluoxetine on Poststroke Dysphagia: A Clinical Retrospective Study. Journal of Stroke and Cerebrovascular Diseases, 2018, 27, 3320-3327. | 0.7 | 11 |
| 183 | Citalopram in first episode schizophrenia: The DECIFER trial. Schizophrenia Research, 2019, 208, 331-337. | 1.1 | 15 |
| 184 | Interactive effect of 5-HTTLPR and BDNF polymorphisms on amygdala intrinsic functional connectivity and anxiety. Psychiatry Research - Neuroimaging, 2019, 285, 1-8. | 0.9 | 12 |
| 185 | Repeated fluoxetine treatment induces long-lasting neurotrophic changes in the medial prefrontal cortex of adult rats. Behavioural Brain Research, 2019, 365, 114-124. | 1.2 | 26 |
| 186 | Antidepressant Effect of Blue Light on Depressive Phenotype in Light-Deprived Male Rats. Journal of Neuropathology and Experimental Neurology, 2020, 79, 1344-1353. | 0.9 | 1 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 187 | Fluoxetine modulates neuronal activity in stress-related limbic areas of adult rats subjected to the chronic social isolation. Brain Research Bulletin, 2020, 163, 95-108. | 1.4 | 8 |
| 188 | Is there a relationship between morphological and functional platelet changes and depressive disorder?. CNS Spectrums, 2020, , 1-34. | 0.7 | 3 |
| 189 | Astrocytic GABAB Receptors in Mouse Hippocampus Control Responses to Behavioral Challenges through Astrocytic BDNF. Neuroscience Bulletin, 2020, 36, 705-718. | 1.5 | 16 |
| 190 | Effect of pharmacological manipulations on Arc function. Current Research in Pharmacology and Drug Discovery, 2021, 2, 100013. | 1.7 | 6 |
| 191 | Possible Modulatory Role of ARC Gene Variants in Mood Disorders. Clinical Psychopharmacology and Neuroscience, 2021, 19, 46-52. | 0.9 | 2 |
| 192 | Glatiramer acetate attenuates depressive/anxiety-like behaviors and cognitive deficits induced by post-weaning social isolation in male mice. Psychopharmacology, 2021, 238, 2121-2132. | 1.5 | 10 |
| 193 | Creatine and Alpha-Lipoic Acid Antidepressant-Like Effect Following Cyclosporine A Administration. Turkish Journal of Pharmaceutical Sciences, 2022, 19, 196-201. | 0.6 | 3 |
| 194 | Adult Neural Progenitor Cells and Cell Replacement Therapy for Huntington Disease. Pancreatic Islet Biology, 2011, , 299-314. | 0.1 | 1 |
| 195 | Effects of adolescent administration of fluoxetine on novel object recognition memory, anxiety-like behaviors, and hippocampal brain-derived neurotrophic factor level. Life Sciences, 2020, 260, 118338. | 2.0 | 21 |
| 196 | Neurologic links between epilepsy and depression in women. Neurology, 2006, 66, S13-22. | 1.5 | 26 |
| 197 | The Roles of BDNF, pCREB and Wnt3a in the Latent Period Preceding Activation of Progenitor Cell Mitosis in The Adult Dentate Gyrus by Fluoxetine. PLoS ONE, 2010, 5, e13652. | 1.1 | 65 |
| 198 | Increased cellular turnover in response to fluoxetine in neuronal precursors derived from human embryonic stem cells. International Journal of Developmental Biology, 2010, 54, 707-715. | 0.3 | 14 |
| 199 | Beneficial effect of fluoxetine treatment aganist psychological stress is mediated by increasing BDNF expression in selected brain areas. Oncotarget, 2017, 8, 69527-69537. | 0.8 | 19 |
| 200 | The Monoaminergic Tripartite Synapse: A Putative Target for Currently Available Antidepressant Drugs. Current Drug Targets, 2013, 14, 1277-1294. | 1.0 | 42 |
| 201 | Assessing the Neuronal Serotonergic Target-based Antidepressant Stratagem: Impact of In Vivo Interaction Studies and Knockout Models. Current Neuropharmacology, 2008, 6, 215-234. | 1.4 | 9 |
| 202 | Neural Mechanisms of Exercise: Anti-Depression, Neurogenesis, and Serotonin Signaling. CNS and Neurological Disorders - Drug Targets, 2015, 14, 1307-1311. | 0.8 | 45 |
| 203 | Ketamina, da anestesia ao uso abusivo. Revista Neurociencias, 2010, 18, 227-237. | 0.0 | 8 |
| 204 | Antidepressant-Like Effects of Central BDNF Administration in Mice of Antidepressant Sensitive Catalepsy (ASC) Strain. Chinese Journal of Physiology, 2012, 55, 284-93. | 0.4 | 14 |

| # | Article | IF | Citations |
|-----|--|-------------------|---------------|
| 205 | Time of Day and Length of Antidepressant Drug Administration Influence Brain-Derived Neurotrophic Factor and TrkB Levels in Rat Brain. The Open Pharmacology Journal, 2009, 3, 1-8. | 0.4 | 0 |
| 206 | Antidepressant Drugs and Phosphodiesterases. , 2016, , 805-823. | | 0 |
| 207 | Imaging Genetics of Depression. , 2016, , 209-222. | | 0 |
| 209 | Even neural stem cells get the blues: evidence for a molecular link between modulation of adult neurogenesis and depression. Gene Expression, 2008, 14, 183-93. | 0.5 | 24 |
| 210 | A salience hypothesis of stress in PTSD. European Journal of Neuroscience, 2021, 54, 8029-8051. | 1.2 | 1 |
| 211 | Antidepressant-like Effects of Combined Fluoxetine and Zinc Treatment in Mice Exposed to Chronic Restraint Stress Are Related to Modulation of Histone Deacetylase. Molecules, 2022, 27, 22. | 1.7 | 5 |
| 213 | Differential effects of fluoxetine on the phototactic behavior of 3 amphipod species (Crustacea;) Tj ETQq0 0 0 rg | BT /Overlo 2.0 | ck 10 Tf 50 5 |
| 214 | Single and Repeated Exposure to Cannabidiol Differently Modulate BDNF Expression and Signaling in the Cortico-Striatal Brain Network. Biomedicines, 2022, 10, 1853. | 1.4 | 8 |
| 215 | Gαs, adenylyl cyclase, and their relationship to the diagnosis and treatment of depression. Frontiers in Pharmacology, 0, 13, . | 1.6 | 5 |
| 216 | Chronic escitalopram in healthy volunteers has specific effects on reinforcement sensitivity: a double-blind, placebo-controlled semi-randomised study. Neuropsychopharmacology, 2023, 48, 664-670. | 2.8 | 12 |
| 217 | Neonatal treatment with fluoxetine alters locomotor activity and the cortical glial/neuron index in rats with cerebral palsy. Canadian Journal of Physiology and Pharmacology, 0, , . | 0.7 | 0 |

| 218 | Translatable Models of Brain and Cognitive Reserve. Contemporary Clinical Neuroscience, 2023, , 93-119. | 0.3 | 0 |
|-----|---|-----|---|
| | | | |