THE CARBONIFEROUS GONIATITES OF THE NORTH O

Proceedings of the Yorkshire Geological Society 20, 40-124

DOI: 10.1144/pygs.20.1.40

Citation Report

#	Article	IF	CITATIONS
1	Some recent stratigraphical work in its bearing on South Pembrokeshire problems Proceedings of the Geologists Association, 1933, 44, 217-225.	1.1	9
2	Late Paleozoic Crustal Movements of Europe and North America. AAPG Bulletin, 1935, 19, .	1.5	2
3	An outline of the geology of Ashover, Derbyshire. Proceedings of the Geologists Association, 1946, 57, 117-IN4.	1.1	1
4	The lower carboniferous rocks of ireland. Geological Journal, 1951, 1, 113-147.	1.3	18
5	Neue Faunen aus dem Namur des nordöstlichen Spaniens. Palaontologische Zeitschrift, 1951, 24, 184-193.	1.6	5
6	The Namurian of the Lancaster Fells. Quarterly Journal of the Geological Society of London, 1953, 109, 423-454.	0.5	26
7	The beds above the Carboniferous Limestone in North-West County Clare, Eire. Quarterly Journal of the Geological Society of London, 1953, 109, 259-283.	0.5	40
8	XIVâ€"Scottish Carboniferous Goniatites. Transactions of the Royal Society of Edinburgh, 1954, 62, 527-602.	0.3	56
9	The Upper Carboniferous Rocks of the Ingleton Coalfield. Quarterly Journal of the Geological Society of London, 1954, 110, 231-265.	0.5	13
10	CURRENT-BEDDING DIRECTIONS IN SANDSTONES OF LOWER <i>RET1CULOCERAS</i> AGE IN THE MILLSTONE GRIT OF WHARFEDALE, YORKSHIRE. Proceedings of the Yorkshire Geological Society, 1955, 30, 115-132.	0.3	12
11	X.â€" The Mode of Life of Certain Goniatites . Transactions of the Geological Society of Glasgow, 1957, 22, 169-186.	0.2	6
12	THE YOREDALE SERIES OF UPPER WENSLEYDALE AND ADJACENT PARTS OF NORTH-WEST YORKSHIRE. Proceedings of the Yorkshire Geological Society, 1957, 31, 91-148.	0.3	41
13	Namurian Stage Names. Geological Magazine, 1958, 95, 85-86.	1.5	1
14	Dinantian, Namurian and Westphalian rocks of the district south-west of Barnstaple, North Devon. Quarterly Journal of the Geological Society of London, 1959, 115, 261-290.	0.5	20
15	THE BASE OF THE NAMURIAN AND OF THE MILLSTONE GRIT IN NORTH-EASTERN ENGLAND. Proceedings of the Yorkshire Geological Society, 1962, 33, 341-362.	0.3	28
16	Radiolarian Nature of the Thicker-shelled Goniatite Faunal Phase in some Namurian Limestone †Bullions'. Nature, 1964, 201, 697-699.	27.8	5
17	A Review of the Factors Affecting the Sedimentation of the Millstone Grit (Namurian) in the Central Pennines. Developments in Sedimentology, 1964, , 340-346.	0.5	20
18	THE CARBONIFEROUS SUCCESSION IN THE KIRKBY MALZEARD AREA, YORKSHIRE. Proceedings of the Yorkshire Geological Society, 1965, 35, 203-227.	0.3	9

#	Article	IF	Citations
19	Viséan ammonoids from South Urals. International Geology Review, 1966, 8, 1067-1077.	2.1	2
20	Relation of upper visean sedimentology to the bowland shale overlap in Yorkshire, England. Sedimentary Geology, 1967, 1, 117-136.	2.1	1
21	Dinantian and namurian stratigraphy east and south-east of leek, north staffordshire. Proceedings of the Geologists Association, 1969, 80, 145-IN1.	1.1	5
22	RATES OF SEDIMENTATION IN THE UPPER CARBONIFEROUS OF BRITAIN. Lethaia, 1970, 3, 1-9.	1.4	14
23	William Sawney Bisat, 1886-1973. Biographical Memoirs of Fellows of the Royal Society, 1974, 20, 27-40.	0.1	2
24	THE HASLINGDEN FLAGS (NAMURIAN G1) OF SOUTH-EAST LANCASHIRE: BAR FINGER SANDS IN THE PENNINE BASIN. Proceedings of the Yorkshire Geological Society, 1975, 40, 431-458.	0.3	34
25	Lower Pennsylvanian (Morrowan) Ammonoids from the North American Midcontinent. Journal of Paleontology, 1980, 54, 1-56.	0.8	42
26	Newsletter on Carboniferous Stratigraphy. Lethaia, 1980, 13, 78-78.	1.4	9
27	Ammonoids and the correlation of the Lower Carboniferous rocks of eastern Australia. Alcheringa, 1983, 7, 75-123.	1.2	20
28	Morphology and morphologic diversity of mid-Carboniferous (Namurian) ammonoids in time and space. Paleobiology, 1984, 10, 195-228.	2.0	105
30	Recent growth of nautiloid and ammonite taxonomy. Palaontologische Zeitschrift, 1989, 63, 281-296.	1.6	4
31	Early fill of the Western Irish Namurian Basin: a complex relationship between turbidites and deltas. Basin Research, 1991, 3, 223-242.	2.7	60
32	Dinantian (Lower Carboniferous) biostratigraphy and chronostratigraphy in the British Isles. Journal of the Geological Society, 1993, 150, 427-446.	2.1	96
33	Re-evaluation of two Early Pennsylvanian (Middle Namurian) ammonoids and their bearing on mid-Carboniferous correlations. Journal of Paleontology, 1993, 67, 993-999.	0.8	2
34	The Late Namurian genusCancelloceras (Carboniferous Ammonoidea) and its distribution. Palaontologische Zeitschrift, 1995, 69, 353-376.	1.6	3
35	Discrimination of regionally extensive coals in the Upper Carboniferous of the Pennine Basin, UK using high resolution sequence stratigraphic concepts. Geological Society Special Publication, 1995, 82, 79-97.	1.3	10
36	Dinantian (Lower Carboniferous) biostratigraphy and chronostratigraphy in the British Isles. Geological Society Memoir, 1995, 16, 105-124.	1.7	1
37	New developments in the Late Carboniferous geology of the central Pennines, northern England: a review. Proceedings of the Yorkshire Geological Society, 1996, 51, 81-86.	0.3	6

#	Article	IF	CITATIONS
38	MID-CARBONIFEROUS AMMONOID BIOSTRATIGRAPHY, SOUTHERN NYE COUNTY, NEVADA: IMPLICATIONS OF THE FIRST NORTH AMERICANHOMOCERAS. Journal of Paleontology, 2001, 75, 1-31.	0.8	8
39	A well-preserved early Namurian ammonoid fauna with <i>Cravenoceras leion</i> Bisat 1930 from Backdale Mine, Hassop, Derbyshire, England. Proceedings of the Yorkshire Geological Society, 2002, 54, 111-119.	0.3	8
40	Comparative palynostratigraphy of the early Arnsbergian (Namurian) sequences between Great Britain and northwest China. Review of Palaeobotany and Palynology, 2002, 118, 227-238.	1.5	5
41	W. S. Bisat (1886–1973): his life and influence on Carboniferous stratigraphy. Proceedings of the Geologists Association, 2004, 115, 371-377.	1.1	O
42	A revised palynozonation of British Namurian deposits and comparisons with eastern Europe. Micropaleontology, 2004, 50, 89.	1.0	5
43	Ammonoid assemblages from the Asbian B _{2b} (Early Carboniferous: Mississippian) buildups of the Peak District, England. Proceedings of the Yorkshire Geological Society, 2006, 56, 111-150.	0.3	9
44	Correlation of Mississippian (Upper Viséan) foraminiferan, conodont, miospore and ammonoid zonal schemes, and correlation with the Asbian–Brigantian boundary in northwest Ireland. Geological Journal, 2006, 41, 221-241.	1.3	24
45	Tournaisian and Visean ammonoid stratigraphy in North Africa. Neues Jahrbuch Fur Geologie Und Palaontologie - Abhandlungen, 2007, 243, 127-148.	0.4	45
46	The Early Carboniferous (Mississippian) ammonoids from the Chebket el Hamra (Jerada Basin,) Tj ETQq0 0 0 rgBT	/Overlock	10 Tf 50 422
47	Biostratigraphic zonation and correlation of Mississippian rocks in Western Europe: some case studies in the late Viséan/Serpukhovian. Geological Journal, 2008, 43, 209-240.	1.3	68
47	Biostratigraphic zonation and correlation of Mississippian rocks in Western Europe: some case studies in the late VisÃ@an/Serpukhovian. Geological Journal, 2008, 43, 209-240. Paleontology and Microfacies of the Serpukhovian in the Verkhnyaya Kardailovka Section, South Urals, Russia: potential candidate for the GSSP for the Visean-Serpukhovian boundary. Newsletters on Stratigraphy, 2009, 43, 165-193.	1.3	68
	Paleontology and Microfacies of the Serpukhovian in the Verkhnyaya Kardailovka Section, South Urals, Russia: potential candidate for the GSSP for the Visean-Serpukhovian boundary. Newsletters on	1.2	49
48	Paleontology and Microfacies of the Serpukhovian in the Verkhnyaya Kardailovka Section, South Urals, Russia: potential candidate for the GSSP for the Visean-Serpukhovian boundary. Newsletters on Stratigraphy, 2009, 43, 165-193.	1.2	49
48 49	Paleontology and Microfacies of the Serpukhovian in the Verkhnyaya Kardailovka Section, South Urals, Russia: potential candidate for the GSSP for the Visean-Serpukhovian boundary. Newsletters on Stratigraphy, 2009, 43, 165-193. The ammonoids from the Argiles de Teguentour of Oued Temertasset (early Late Tournaisian; Mouydir,) Tj ETQq1 The ammonoids from the Argiles de Timimoun of Timimoun (Early and Middle VisÃ@an; Gourara, Algeria).	1.2 10.78431	49 4_rgBT /Ove
48 49 50	Paleontology and Microfacies of the Serpukhovian in the Verkhnyaya Kardailovka Section, South Urals, Russia: potential candidate for the GSSP for the Visean-Serpukhovian boundary. Newsletters on Stratigraphy, 2009, 43, 165-193. The ammonoids from the Argiles de Teguentour of Oued Temertasset (early Late Tournaisian; Mouydir,) Tj ETQq1 The ammonoids from the Argiles de Timimoun of Timimoun (Early and Middle VisÃ@an; Gourara, Algeria). Fossil Record, 2010, 13, 215-278. Submarine channel response to intrabasinal tectonics: The influence of lateral tilt. AAPG Bulletin,	1.2 1.0.78431 0.5	49 4 rgBT /Ove
48 49 50 51	Paleontology and Microfacies of the Serpukhovian in the Verkhnyaya Kardailovka Section, South Urals, Russia: potential candidate for the GSSP for the Visean-Serpukhovian boundary. Newsletters on Stratigraphy, 2009, 43, 165-193. The ammonoids from the Argiles de Teguentour of Oued Temertasset (early Late Tournaisian; Mouydir,) Tj ETQq1 The ammonoids from the Argiles de Timimoun of Timimoun (Early and Middle VisÃ@an; Gourara, Algeria). Fossil Record, 2010, 13, 215-278. Submarine channel response to intrabasinal tectonics: The influence of lateral tilt. AAPG Bulletin, 2010, 94, 189-219. The Serpukhovian and Bashkirian (Carboniferous, Namurian and basal Westphalian) faunas of	1.2 1.0.78431 0.5	49 4 ₂₁ gBT /Ove 14 48
48 49 50 51 52	Paleontology and Microfacies of the Serpukhovian in the Verkhnyaya Kardailovka Section, South Urals, Russia: potential candidate for the GSSP for the Visean-Serpukhovian boundary. Newsletters on Stratigraphy, 2009, 43, 165-193. The ammonoids from the Argiles de Teguentour of Oued Temertasset (early Late Tournaisian; Mouydir,) Tj ETQq1 The ammonoids from the Argiles de Timimoun of Timimoun (Early and Middle VisÃ@an; Gourara, Algeria). Fossil Record, 2010, 13, 215-278. Submarine channel response to intrabasinal tectonics: The influence of lateral tilt. AAPG Bulletin, 2010, 94, 189-219. The Serpukhovian and Bashkirian (Carboniferous, Namurian and basal Westphalian) faunas of northern England. Proceedings of the Yorkshire Geological Society, 2011, 58, 143-165.	1.2 1.0.78431 0.5	49 4 ₂₁ gBT /Ove 14 48

#	Article	IF	CITATIONS
56	Biostratigraphy and lithostratigraphy of the Mid-Carboniferous boundary beds in the Muradymovo section (South Urals, Russia). Geological Magazine, 2014, 151, 269-298.	1.5	30
57	Controls on deltaic sedimentation in glacio-eustatic cycles of late Marsdenian (Namurian R) Tj ETQq1 1 0.784314 of the Yorkshire Geological Society, 2014, 60, 63-84.	rgBT /Ove 0.3	rlock 10 Tf 4
58	Paleozoic Ammonoid Biostratigraphy. Topics in Geobiology, 2015, , 299-328.	0.5	13
59	Gastropods from the Carboniferous (Namurian) of Congleton Edge, Cheshire, <scp>UK</scp> . Papers in Palaeontology, 2016, 2, 399-438.	1.5	5
60	Problems correlating the late Brigantian–Arnsbergian Western European substages within northern England. Geological Journal, 2016, 51, 817-840.	1.3	31
61	Mid-Carboniferous ammonoids from the Shannon Basin, western Ireland: identification of crushed material. Swiss Journal of Palaeontology, 2016, 135, 75-85.	1.7	6
62	The prospectivity of a potential shale gas play: An example from the southern Pennine Basin (central) Tj ETQq0 0 0	rgBT /Ov	erlock 10 Tf
63	Goniatites sphaericus (Sowerby, 1814), the archetype of Palaeozoic ammonoids: a case of decreasing phenotypic variation through ontogeny. Palaontologische Zeitschrift, 2017, 91, 337-352.	1.6	12
64	Controls on amorphous organic matter type and sulphurization in a Mississippian black shale. Review of Palaeobotany and Palynology, 2019, 268, 1-18.	1.5	20
65	Total organic carbon in the Bowland-Hodder Unit of the southern Widmerpool Gulf: a discussion. Journal of Petroleum Science and Engineering, 2019, 178, 1194-1202.	4.2	7
66	Foraminifers of the Viséan–Serpukhovian boundary interval in Western Palaeotethys: a review. Lethaia, 2019, 52, 260-284.	1.4	18
67	Utility of crinoid columnals in palaeontology illustrated by a new species: Clare Shale Formation (Carboniferous), Doolin, County Clare, western Ireland. Proceedings of the Geologists Association, 2019, 130, 696-700.	1.1	4
68	From marine bands to hybrid flows: Sedimentology of a Mississippian black shale. Sedimentology, 2020, 67, 261-304.	3.1	20
69	A Mississippian black shale record of redox oscillation in the Craven Basin, UK. Palaeogeography, Palaeoclimatology, Palaeoecology, 2020, 538, 109423.	2.3	11
70	Bilinguites â€" a cosmopolitan Late Carboniferous ammonoid genus. Palaeoworld, 2021, 30, 689-707.	1.1	3
71	The contribution of publications of the Yorkshire Geological Society to the understanding of the geological development of the Carboniferous Pennine Basin, northern England. Proceedings of the Yorkshire Geological Society, 2020, 63, 1-32.	0.3	7
72	Microconchus cravenensisn. sp.: a giant among microconchid tubeworms. Journal of Paleontology, 2020, 94, 1051-1058.	0.8	6
73	Quantitative palynological analysis of the E2a and E2b goniatite biozones (Arnsbergian, Mississippian) in mudstones from the Southern Pennine Basin (U.K.). Review of Palaeobotany and Palynology, 2020, 276, 104187.	1.5	3

#	ARTICLE	IF	Citations
74	The succession of the mid-Bashkirian ammonoids Cancelloceras and Gastrioceras in North China. Palaeoworld, 2021, 30, 72-94.	1.1	5
75	The holotype of Psephodus minutus, Wellburn, 1901 (chondrichthyes, cochliodontiformes) is a gastropod steinkern. Proceedings of the Yorkshire Geological Society, 2021, 63, pygs2020-004.	0.3	1
76	SIGNIFICANCE FOR INTERNATIONAL CORRELATION OF THE PERAPERTÃS FORMATION IN NORTHERN PALENCIA, CANTABRIAN MOUNTAINS. TECTONIC/STRATIGRAPHIC CONTEXT AND DESCRIPTION OF MISSISSIPPIAN AND UPPER BASHKIRIAN GONIATITES. Spanish Journal of Paleontology, 2021, 22, 127.	0.1	3
77	The Carboniferous chronostratigraphic scale: history, status and prospectus. Geological Society Special Publication, 2022, 512, 19-48.	1.3	6
78	Earth science collections of the Centre Grégoire Fournier (Maredsous) with comments on Middle Devonian–Carboniferous brachiopods and trilobites from southern Belgium. Geologica Belgica, 2021, 24, 33-68.	1,1	8
79	Carboniferous ammonoid genozones. Geological Society Special Publication, 2022, 512, 633-693.	1.3	3
80	The Carboniferous Period., 2020,, 811-874.		45
81	THE GEOLOGY OF THE KEASDEN AREA, WEST OF SETTLE, YORKSHIRE. Proceedings of the Yorkshire Geological Society, 1956, 30, 331-352.	0.3	11
82	MARINE BANDS OF ARNSBERGIAN AGE (NAMURIAN) IN THE SOUTH-EASTERN PORTION OF THE ASKRIGG BLOCK, YORKSHIRE. Proceedings of the Yorkshire Geological Society, 1959, 32, 45-67.	0.3	11
83	THE MILLSTONE GRIT SERIES OF COLSTERDALE AND NEIGHBOURHOOD, YORKSHIRE. Proceedings of the Yorkshire Geological Society, 1960, 32, 429-452.	0.3	11
84	Kinderscoutian and Marsdenian successions in the Bradup and Hag Farm boreholes, near Ilkley, west Yorkshire. Proceedings of the Yorkshire Geological Society, 1996, 51, 115-125.	0.3	16
85	High-resolution sequence stratigraphy in the early Marsdenian (Namurian, Carboniferous) of the central Pennines and adjacent areas. Proceedings of the Yorkshire Geological Society, 1996, 51, 127-140.	0.3	17
86	Paleosalinity in Upper Visean Yoredale Formation of England-Geochemical Method for Locating Porosity. AAPG Bulletin, 1964, 48, .	1.5	3
87	The Serpukhovian in Britain: use of foraminiferal assemblages for dating and correlating. Journal of the Geological Society, 2021, 178, jgs2020-170.	2.1	10
88	An integrated sequence stratigraphic analysis of the early Marsdenian substage of the Millstone Grit Group, Central Pennines, UK. Proceedings of the Yorkshire Geological Society, 2023, 64, .	0.3	0
89	Cross-Basin Mo and U Analysis of the Upper Mississippian Bowland Shale UK. Geological Society Special Publication, 2024, 534, .	1.3	O