

CITATION REPORT

List of articles citing

Trigonometrically fitted RungeKutta methods for the numerical solution of the Schrdinger equation

DOI: 10.1007/s10910-004-1470-8

Journal of Mathematical Chemistry, 2005, 37, 281-293.

Source: <https://exaly.com/paper-pdf/38496188/citation-report.pdf>

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
98	Stability and phase-lag analysis of explicit Runge-Kutta methods with variable coefficients for oscillatory problems. <i>Computer Physics Communications</i> , 2005 , 173, 115-130	4.2	40
97	Trigonometrically fitted and exponentially fitted symplectic methods for the numerical integration of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2006 , 40, 257-267	2.1	71
96	The numerical solution of the radial Schrödinger equation via a trigonometrically fitted family of seventh algebraic order Predictor-Corrector methods. <i>Journal of Mathematical Chemistry</i> , 2006 , 40, 269-293	2.1	66
95	A Family of Exponentially-fitted Runge-Kutta Methods with Exponential Order Up to Three for the Numerical Solution of the Schrödinger Equation. <i>Journal of Mathematical Chemistry</i> , 2007 , 41, 79-100	2.1	81
94	New Second-order Exponentially and Trigonometrically Fitted Symplectic Integrators for the Numerical Solution of the Time-independent Schrödinger Equation. <i>Journal of Mathematical Chemistry</i> , 2007 , 42, 535-545	2.1	51
93	A family of four-step trigonometrically-fitted methods and its application to the schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2008 , 44, 447-466	2.1	51
92	Closed Newton-Cotes trigonometrically-fitted formulae of high order for the numerical integration of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2008 , 44, 483-499	2.1	52
91	Numerical methods in chemistry. <i>Chemical Modelling</i> , 350-487	2	1
90	A new methodology for the development of numerical methods for the numerical solution of the Schrödinger equation. <i>Chemical Modelling</i> , 210-509	2	
89	Conditions for Trigonometrically Fitted Runge-Kutta Methods. 2009 ,		
88	A Modified Zero Dispersion and Zero Dissipation RKN Method for the Numerical Solution of the Radial Schrödinger Equation. 2009 ,		1
87	Trends of the bonding effect on the performance of DFT methods in electric properties calculations: a pattern recognition and metric space approach on some XY ₂ (X = O, S and Y = H, O, F, S, Cl) molecules. <i>Journal of Computational Chemistry</i> , 2010 , 31, 412-20	3.5	14
86	Numerical multistep methods for the efficient solution of quantum mechanics and related problems. <i>Physics Reports</i> , 2009 , 482-483, 1-240	27.7	67
85	A family of two-stage two-step methods for the numerical integration of the Schrödinger equation and related IVPs with oscillating solution. <i>Journal of Mathematical Chemistry</i> , 2009 , 45, 1102-1129	2.1	61
84	A family of Runge-Kutta methods with zero phase-lag and derivatives for the numerical solution of the Schrödinger equation and related problems. <i>Journal of Mathematical Chemistry</i> , 2009 , 46, 1158-1171	2.1	7
83	Two optimized symmetric eight-step implicit methods for initial-value problems with oscillating solutions. <i>Journal of Mathematical Chemistry</i> , 2009 , 46, 604-620	2.1	87
82	A new methodology for the development of numerical methods for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2009 , 46, 621-651	2.1	2

81	A new methodology for the construction of numerical methods for the approximate solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2009 , 46, 652-691	2.1	1
80	High order multistep methods with improved phase-lag characteristics for the integration of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2009 , 46, 692-725	2.1	2
79	High order phase fitted multistep integrators for the Schrödinger equation with improved frequency tolerance. <i>Journal of Mathematical Chemistry</i> , 2009 , 46, 1009-1049	2.1	1
78	A new Numerov-type method for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2009 , 46, 981-1007	2.1	97
77	P-stability, Trigonometric-fitting and the numerical solution of the radial Schrödinger equation. <i>Computer Physics Communications</i> , 2009 , 180, 1072-1085	4.2	9
76	High order closed Newton-Cotes trigonometrically-fitted formulae for the numerical solution of the Schrödinger equation. <i>Applied Mathematics and Computation</i> , 2009 , 209, 137-151	2.7	124
75	A semi-discretization method based on quartic splines for solving one-space-dimensional hyperbolic equations. <i>Applied Mathematics and Computation</i> , 2009 , 210, 508-514	2.7	13
74	On the efficiency of Newton and Broyden numerical methods in the investigation of the regular polygon problem of (N+1) bodies. <i>Applied Mathematics and Computation</i> , 2009 , 212, 100-112	2.7	14
73	The Use of Phase-Lag and Amplification Error Derivatives in the Numerical Integration of ODEs with Oscillating Solutions. 2009 ,		1
72	A modified phase-fitted and amplification-fitted Runge-Kutta-Nyström method for the numerical solution of the radial Schrödinger equation. <i>Journal of Molecular Modeling</i> , 2010 , 16, 1339-46	2	16
71	Exponentially and Trigonometrically Fitted Methods for the Solution of the Schrödinger Equation. <i>Acta Applicandae Mathematicae</i> , 2010 , 110, 1331-1352	1.1	137
70	An optimized explicit Runge-Kutta method with increased phase-lag order for the numerical solution of the Schrödinger equation and related problems. <i>Journal of Mathematical Chemistry</i> , 2010 , 47, 315-330	2.1	64
69	A new two-step hybrid method for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2010 , 47, 871-890	2.1	85
68	Two-step high order hybrid explicit method for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2010 , 48, 224-252	2.1	75
67	Multistep methods with vanished phase-lag and its first and second derivatives for the numerical integration of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2010 , 48, 1092-1143	2.1	30
66	A family of eight-step methods with vanished phase-lag and its derivatives for the numerical integration of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2011 , 49, 711-764	2.1	23
65	A hybrid method with phase-lag and derivatives equal to zero for the numerical integration of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2011 , 49, 1330-1356	2.1	36
64	A family of ten-step methods with vanished phase-lag and its first derivative for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2011 , 49, 1843-1888	2.1	11

63	A two-step method with vanished phase-lag and its first two derivatives for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2011 , 49, 2486-2518	2.1	43
62	A new symmetric linear eight-step method with fifth trigonometric order for the efficient integration of the Schrödinger equation. <i>Applied Mathematics Letters</i> , 2011 , 24, 1468-1472	3.5	6
61	A NEW METHODOLOGY FOR THE CONSTRUCTION OF OPTIMIZED RUNGE-KUTTA METHODS. <i>International Journal of Modern Physics C</i> , 2011 , 22, 623-634	1.1	44
60	A new high order two-step method with vanished phase-lag and its derivatives for the numerical integration of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2012 , 50, 2351-2373	2.1	5
59	New open modified trigonometrically-fitted Newton-Cotes type multilayer symplectic integrators for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2012 , 50, 782-804	2.1	5
58	High order closed Newton-Cotes exponentially and trigonometrically fitted formulae as multilayer symplectic integrators and their application to the radial Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2012 , 50, 1224-1261	2.1	71
57	A new hybrid two-step method with vanished phase-lag and its first and second derivatives for the numerical solution of the Schrödinger equation and related problems. <i>Journal of Mathematical Chemistry</i> , 2012 , 50, 1861-1881	2.1	6
56	A parametric symmetric linear four-step method for the efficient integration of the Schrödinger equation and related oscillatory problems. <i>Journal of Computational and Applied Mathematics</i> , 2012 , 236, 3880-3889	2.4	113
55	A new family of symmetric linear four-step methods for the efficient integration of the Schrödinger equation and related oscillatory problems. <i>Applied Mathematics and Computation</i> , 2012 , 218, 5370-5382	2.7	101
54	A new four-step hybrid type method with vanished phase-lag and its first derivatives for each level for the approximate integration of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2013 , 51, 2542-2571	2.1	26
53	A new optimized symmetric 8-step semi-embedded predictor-corrector method for the numerical solution of the radial Schrödinger equation and related orbital problems. <i>Journal of Mathematical Chemistry</i> , 2013 , 51, 1914-1937	2.1	42
52	A new four-step Runge-Kutta type method with vanished phase-lag and its first, second and third derivatives for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2013 , 51, 1418-1445	2.1	7
51	A new modified embedded 5(4) pair of explicit Runge-Kutta methods for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2013 , 51, 937-953	2.1	6
50	New high order multiderivative explicit four-step methods with vanished phase-lag and its derivatives for the approximate solution of the Schrödinger equation. Part I: Construction and theoretical analysis. <i>Journal of Mathematical Chemistry</i> , 2013 , 51, 194-226	2.1	68
49	New optimized explicit modified RKN methods for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2013 , 51, 390-411	2.1	4
48	High order four-step hybrid method with vanished phase-lag and its derivatives for the approximate solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2013 , 51, 532-555	2.1	7
47	A New Trigonometrically Fitted Two-Derivative Runge-Kutta Method for the Numerical Solution of the Schrödinger Equation and Related Problems. <i>Journal of Applied Mathematics</i> , 2013 , 2013, 1-9	1.1	2
46	EXPONENTIALLY FITTED TWO-DERIVATIVE RUNGE-KUTTA METHODS FOR THE SCHRÖDINGER EQUATION. <i>International Journal of Modern Physics C</i> , 2013 , 24, 1350073	1.1	10

45	A hybrid type four-step method with vanished phase-lag and its first, second and third derivatives for each level for the numerical integration of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2014 , 52, 2334-2379	2.1	55
44	A new explicit hybrid four-step method with vanished phase-lag and its derivatives. <i>Journal of Mathematical Chemistry</i> , 2014 , 52, 1690-1716	2.1	52
43	Trigonometrically fitted high-order predictor-corrector method with phase-lag of order infinity for the numerical solution of radial Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2014 , 52, 1870-1894	2.1	8
42	A new embedded 5(3) pair of modified Runge-Kutta-Nyström methods for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2014 , 52, 1081-1098	2.1	1
41	A Runge-Kutta type four-step method with vanished phase-lag and its first and second derivatives for each level for the numerical integration of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2014 , 52, 917-947	2.1	59
40	An explicit four-step method with vanished phase-lag and its first and second derivatives. <i>Journal of Mathematical Chemistry</i> , 2014 , 52, 833-855	2.1	64
39	An explicit linear six-step method with vanished phase-lag and its first derivative. <i>Journal of Mathematical Chemistry</i> , 2014 , 52, 1895-1920	2.1	22
38	A family of explicit linear six-step methods with vanished phase-lag and its first derivative. <i>Journal of Mathematical Chemistry</i> , 2014 , 52, 2087-2118	2.1	55
37	Runge-Kutta type methods with special properties for the numerical integration of ordinary differential equations. <i>Physics Reports</i> , 2014 , 536, 75-146	27.7	13
36	Efficient low computational cost hybrid explicit four-step method with vanished phase-lag and its first, second, third and fourth derivatives for the numerical integration of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2015 , 53, 1808-1834	2.1	53
35	A new explicit four-step method with vanished phase-lag and its first and second derivatives. <i>Journal of Mathematical Chemistry</i> , 2015 , 53, 402-429	2.1	25
34	A high algebraic order multistage explicit four-step method with vanished phase-lag and its first, second, third, fourth and fifth derivatives for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2015 , 53, 1915-1942	2.1	47
33	A predictor-corrector explicit four-step method with vanished phase-lag and its first, second and third derivatives for the numerical integration of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2015 , 53, 685-717	2.1	47
32	Exponentially fitted TDRK pairs for the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2015 , 53, 1470-1487	2.1	
31	A high algebraic order predictor-corrector explicit method with vanished phase-lag and its first, second, third and fourth derivatives for the numerical solution of the Schrödinger equation and related problems. <i>Journal of Mathematical Chemistry</i> , 2015 , 53, 1495-1522	2.1	48
30	Explicit Runge-Kutta method with trigonometrically-fitted for solving first order ODEs. 2016 ,		
29	A family of embedded explicit six-step methods with vanished phase-lag and its derivatives for the numerical integration of the Schrödinger equation: development and theoretical analysis. <i>Journal of Mathematical Chemistry</i> , 2016 , 54, 1159-1186	2.1	16
28	A family of two stages tenth algebraic order symmetric six-step methods with vanished phase-lag and its first derivatives for the numerical solution of the radial Schrödinger equation and related problems. <i>Journal of Mathematical Chemistry</i> , 2016 , 54, 1835-1862	2.1	15

27	A new eight algebraic order embedded explicit six-step method with vanished phase-lag and its first, second, third and fourth derivatives for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2016 , 54, 1696-1727	2.1	14
26	An implicit symmetric linear six-step methods with vanished phase-lag and its first, second, third and fourth derivatives for the numerical solution of the radial Schrödinger equation and related problems. <i>Journal of Mathematical Chemistry</i> , 2016 , 54, 1010-1040	2.1	17
25	Family of symmetric linear six-step methods with vanished phase-lag and its derivatives and their application to the radial Schrödinger equation and related problems. <i>Journal of Mathematical Chemistry</i> , 2016 , 54, 466-502	2.1	17
24	Three stages symmetric six-step method with eliminated phase-lag and its derivatives for the solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2017 , 55, 1213-1235	2.1	11
23	An efficient six-step method for the solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2017 , 55, 1521-1547	2.1	9
22	High order computationally economical six-step method with vanished phase-lag and its derivatives for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2017 , 55, 987-1013	2.1	10
21	A generator of families of two-step numerical methods with free parameters and minimal phase-lag. <i>Journal of Mathematical Chemistry</i> , 2017 , 55, 1808-1832	2.1	33
20	A new two stages tenth algebraic order symmetric six-step method with vanished phase-lag and its first and second derivatives for the solution of the radial Schrödinger equation and related problems. <i>Journal of Mathematical Chemistry</i> , 2017 , 55, 105-131	2.1	14
19	New two stages high order symmetric six-step method with vanished phase-lag and its first, second and third derivatives for the numerical solution of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2017 , 55, 503-531	2.1	12
18	Two stages six-step method with eliminated phase-lag and its first, second, third and fourth derivatives for the approximation of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2017 , 55, 961-986	2.1	12
17	Exponentially fitted multi-derivative linear methods for the resonant state of the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2017 , 55, 223-237	2.1	1
16	Phase-Fitted and Amplification-Fitted Higher Order Two-Derivative Runge-Kutta Method for the Numerical Solution of Orbital and Related Periodical IVPs. <i>Mathematical Problems in Engineering</i> , 2017 , 2017, 1-11	1.1	5
15	A new three-stages six-step finite difference pair with optimal phase properties for second order initial and/or boundary value problems with periodical and/or oscillating solutions. <i>Journal of Mathematical Chemistry</i> , 2018 , 56, 1280-1312	2.1	2
14	Two-derivative Runge-Kutta methods with increased phase-lag and dissipation order for the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2018 , 56, 1924-1934	2.1	2
13	A new six-step algorithm with improved properties for the numerical solution of second order initial and/or boundary value problems. <i>Journal of Mathematical Chemistry</i> , 2018 , 56, 1206-1233	2.1	3
12	New three-stages symmetric six-step finite difference method with vanished phase-lag and its derivatives up to sixth derivative for second order initial and/or boundary value problems with periodical and/or oscillating solutions. <i>Journal of Mathematical Chemistry</i> , 2018 , 56, 2267-2301	2.1	1
11	A new eight-order symmetric two-step multiderivative method for the numerical solution of second-order IVPs with oscillating solutions. <i>Numerical Algorithms</i> , 2018 , 77, 95-109	2.1	12
10	A multistep method with optimal properties for second order differential equations. <i>Journal of Mathematical Chemistry</i> , 2018 , 56, 1-29	2.1	9

9	A new multistep finite difference pair for the Schrödinger equation and related problems. <i>Journal of Mathematical Chemistry</i> , 2018 , 56, 656-686	2.1	3
8	Modified two-derivative Runge-Kutta methods for the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2018 , 56, 799-812	2.1	4
7	Exponentially-fitted fourth-order explicit modified Runge-Kutta type method for solving third-order ODEs. <i>Journal of Physics: Conference Series</i> , 2018 , 1132, 012016	0.3	1
6	Exponentially Fitted and Trigonometrically Fitted Explicit Modified Runge-Kutta Type Methods for Solving $y'' = f(x, y, y')$. <i>Journal of Applied Mathematics</i> , 2018 , 2018, 1-19	1.1	1
5	New four-stages symmetric six-step method with improved phase properties for second order problems with periodical and/or oscillating solutions. <i>Journal of Mathematical Chemistry</i> , 2018 , 56, 2898-2928	2.1	2
4	Trigonometrically fitted two-step Obrechhoff linear methods for the Schrödinger equation. <i>Journal of Mathematical Chemistry</i> , 2019 , 57, 1543-1556	2.1	
3	Trigonometrically fitted fifth-order explicit two-derivative Runge-Kutta method with FSAL property. <i>Journal of Physics: Conference Series</i> , 2019 , 1294, 032009	0.3	1
2	New multiple stages multistep method with best possible phase properties for second order initial/boundary value problems. <i>Journal of Mathematical Chemistry</i> , 2019 , 57, 834-857	2.1	2
1	A general solution procedure for the scaled boundary finite element method via shooting technique. <i>Computer Methods in Applied Mechanics and Engineering</i> , 2021 , 384, 113996	5.7	2