

CITATION REPORT

List of articles citing

Modified glomerular filtration rate estimating equation for Chinese patients with chronic kidney disease

DOI: 10.1681/asn.2006040368

Journal of the American Society of Nephrology: JASN, 2006, 17, 2937-44.

Source: <https://exaly.com/paper-pdf/39531091/citation-report.pdf>

Version: 2024-04-27

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
1435	Low dosages of bromocriptine added to levodopa in Parkinsons disease. 1985 , 35, 199-206		65
1434	Exploration de la fonction glomÉulaire rÉale (II). Estimation du dÉbit de filtration glomÉulaire. 2006 , 1, 1-12		
1433	ÉpidÉmiologie de l'insuffisance rÉale terminale traitÉe par dialyse. 2006 , 1, 1-12		
1432	Comparison of 99mTc-DTPA renal dynamic imaging with modified MDRD equation for glomerular filtration rate estimation in Chinese patients in different stages of chronic kidney disease. 2007 , 22, 417-23		59
1431	Community-based screening for chronic kidney disease among populations older than 40 years in Beijing. 2007 , 22, 1093-9		88
1430	Evaluation of the modification of diet in renal disease study equation in a large diverse population. <i>Journal of the American Society of Nephrology: JASN</i> , 2007 , 18, 2749-57	12.7	412
1429	Improved GFR estimation by combined creatinine and cystatin C measurements. 2007 , 72, 1535-42		83
1428	How common is early chronic kidney disease? A background paper prepared for the UK Consensus Conference on early chronic kidney disease. 2007 , 22 Suppl 9, ix8-18		6
1427	The usefulness of the International Diabetes Federation and the National Cholesterol Education Programs Adult Treatment Panel III definitions of the metabolic syndrome in predicting coronary heart disease in subjects with type 2 diabetes. 2007 , 30, 1206-11		101
1426	Carotid intima media thickness predicts cardiovascular diseases in Chinese predialysis patients with chronic kidney disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2007 , 18, 1966-72	12.7	79
1425	Chronic kidney disease the promise and the perils. <i>Journal of the American Society of Nephrology: JASN</i> , 2007 , 18, 2803-5	12.7	37
1424	An epidemic of chronic kidney disease: fact or fiction?. 2008 , 23, 1117-21		128
1423	Bibliography. Current world literature. Dialysis and transplantation. 2007 , 16, 596-616		
1422	Understanding estimated glomerular filtration rate: implications for identifying chronic kidney disease. 2007 , 16, 242-9		40
1421	Bibliography. Current world literature. Renal immunology and pathology. 2007 , 16, 272-83		
1420	Metabolic syndrome and chronic kidney disease in a Chinese population aged 40 years and older. 2007 , 82, 822-7		43
1419	Impacts of chronic kidney disease and albuminuria on associations between coronary heart disease and its traditional risk factors in type 2 diabetic patients - the Hong Kong diabetes registry. 2007 , 6, 37		28

1418	Interactive effect of retinopathy and macroalbuminuria on all-cause mortality, cardiovascular and renal end points in Chinese patients with Type 2 diabetes mellitus. 2007 , 24, 741-6	32
1417	Aberrant sialylation of serum IgA1 was associated with prognosis of patients with IgA nephropathy. 2007 , 125, 268-74	22
1416	Association between carotid artery intima-media thickness and early-stage CKD in a Chinese population. 2007 , 49, 786-92	57
1415	Testing for chronic kidney disease: a position statement from the National Kidney Foundation. 2007 , 50, 169-80	210
1414	Modification of the Modification of Diet in Renal Disease (MDRD) Study equation for Japan. 2007 , 50, 927-37	290
1413	Selection of Appropriate Glomerular Filtration Rate Estimating Equation. 2007 , 9, 15-22	
1412	Dual Therapy with ACE Inhibitors and Angiotensin II Receptor Blockers in Proteinuric IgA Nephropathy Patients. 2007 , 9, 89-93	1
1411	Modified end-stage renal disease risk score for Chinese type 2 diabetic patients--the Hong Kong Diabetes Registry. 2007 , 50, 1348-50	16
1410	Estimation of glomerular filtration rate by the MDRD study equation modified for Japanese patients with chronic kidney disease. 2007 , 11, 41-50	442
1409	Prevalence of chronic kidney disease (CKD) in the Japanese general population predicted by the MDRD equation modified by a Japanese coefficient. 2007 , 11, 156-163	223
1408	Bias and precision of estimated glomerular filtration rate in children. 2007 , 22, 167-9	14
1407	Association of kidney function with inflammatory and procoagulant markers in a diverse cohort: a cross-sectional analysis from the Multi-Ethnic Study of Atherosclerosis (MESA). 2008 , 9, 9	95
1406	Development and validation of a risk score for hospitalization for heart failure in patients with Type 2 diabetes mellitus. 2008 , 7, 9	37
1405	Risk factors in V-shaped risk associations with all-cause mortality in type 2 diabetes-The Hong Kong Diabetes Registry. 2008 , 24, 238-46	46
1404	Natural history of immunoglobulin A nephropathy and predictive factors of prognosis: a long-term follow up of 204 cases in China. 2008 , 13, 242-6	74
1403	Association between waist-to-hip ratio and chronic kidney disease in the elderly. 2008 , 38, 402-6	16
1402	The NCEP-ATPIII but not the IDF criteria for the metabolic syndrome identify Type 2 diabetic patients at increased risk of chronic kidney disease. 2008 , 25, 1419-25	18
1401	Reliability of the MDRD method for estimating glomerular filtration rate in relation to gender, body mass index and extracellular fluid volume. 2008 , 38, 486-93	6

1400	The metabolic syndrome increases cardiovascular mortality in Taiwanese elderly. 2008 , 38, 469-75	20
1399	Prevalence of chronic kidney disease in population-based studies: systematic review. 2008 , 8, 117	601
1398	Development and validation of a total coronary heart disease risk score in type 2 diabetes mellitus. 2008 , 101, 596-601	89
1397	Prevalence and factors associated with CKD: a population study from Beijing. 2008 , 51, 373-84	208
1396	Oral calcitriol for the treatment of persistent proteinuria in immunoglobulin A nephropathy: an uncontrolled trial. 2008 , 51, 724-31	78
1395	CKD in the elderly--old questions and new challenges: World Kidney Day 2008. 2008 , 51, 353-7	45
1394	Mortality risks for all causes and cardiovascular diseases and reduced GFR in a middle-aged working population in Taiwan. 2008 , 52, 1051-60	66
1393	Variability and risk factors for kidney disease progression and death following attainment of stage 4 CKD in a referred cohort. 2008 , 52, 661-71	201
1392	CKD in the elderly. 2008 , 52, 803; author reply 803-4	12
1391	Type 2 diabetes mellitus and its renal complications in relation to apolipoprotein E gene polymorphism. 2008 , 152, 134-42	10
1390	Erectile dysfunction predicts coronary heart disease in type 2 diabetes. 2008 , 51, 2045-50	158
1389	Urinary IL-18 and NGAL as early predictive biomarkers in contrast-induced nephropathy after coronary angiography. 2008 , 108, c176-81	176
1388	Influence of preoperative allograft function (effective renal plasma flow) on the short-term outcome following living donor kidney transplantation. 2008 , 40, 2108-11	4
1387	Investigation of the prevalence of CKD in 13,383 Chinese hospitalised adult patients. 2008 , 387, 128-32	17
1386	Gender-specific reference value of urine albumin-creatinine ratio in healthy Chinese adults: results of the Beijing CKD survey. 2008 , 398, 125-9	14
1385	Thresholds of risk factors for ischemic stroke in type 2 diabetic patients with and without albuminuria: a non-linear approach. 2008 , 110, 701-9	8
1384	Monitoring renal function and limitations of renal function tests. 2008 , 38, 32-46	119
1383	Implementation of the routine reporting of eGFR in Australia and New Zealand. 2008 , 241, 23-9	16

1382	Predicting values of lipids and white blood cell count for all-site cancer in type 2 diabetes. 2008 , 15, 597-607	33
1381	Screening for CKD with eGFR: doubts and dangers. 2008 , 3, 1563-8	116
1380	Outcomes of stage 3-5 chronic kidney disease before end-stage renal disease at a single center in Taiwan. 2008 , 109, c109-18	49
1379	The global burden of chronic kidney disease: how valid are the estimates?. 2008 , 110, c39-46; discussion c47	54
1378	Reference intervals of common serum analytes of Hong Kong Chinese. 2008 , 61, 632-6	18
1377	Delayed severe pneumonia in mycophenolate mofetil-treated patients with IgA nephropathy. 2008 , 23, 2868-72	26
1376	Stepwise increase in the prevalence of isolated systolic hypertension with the stages of chronic kidney disease. 2008 , 23, 3895-900	25
1375	Predictivity of survival according to different equations for estimating renal function in community-dwelling elderly subjects. 2009 , 24, 1197-205	40
1374	Comparison of the prevalence of chronic kidney disease among different ethnicities: Beijing CKD survey and American NHANES. 2009 , 24, 1220-6	22
1373	The prevalence of hematologic and metabolic abnormalities during chronic kidney disease stages in different ethnic groups. 2008 , 74, 108-14	9
1372	Antineutrophil cytoplasmic autoantibody-associated vasculitis in older patients. 2008 , 87, 203-209	85
1371	Peroxisome proliferator-activated receptor-gamma gene polymorphism and risk of cardiovascular disease in patients with diabetic nephropathy. 2008 , 28, 715-22	16
1370	Metabolic syndrome predicts new onset of chronic kidney disease in 5,829 patients with type 2 diabetes: a 5-year prospective analysis of the Hong Kong Diabetes Registry. 2008 , 31, 2357-61	134
1369	Long-term outcomes of patients with propylthiouracil-induced anti-neutrophil cytoplasmic auto-antibody-associated vasculitis. 2008 , 47, 1515-20	41
1368	Prevalence and risk factors associated with chronic kidney disease in an adult population from southern China. 2009 , 24, 1205-12	98
1367	Additive interaction of hyperglycemia and albuminuria on risk of ischemic stroke in type 2 diabetes: Hong Kong Diabetes Registry. 2008 , 31, 2294-300	24
1366	Development and validation of an all-cause mortality risk score in type 2 diabetes. 2008 , 168, 451-7	79
1365	Tacrolimus for the treatment of systemic lupus erythematosus with pure class V nephritis. 2008 , 47, 1678-81	97

1364	Aldose reductase genotypes and cardiorenal complications: an 8-year prospective analysis of 1,074 type 2 diabetic patients. 2008 , 31, 2148-53	21
1363	Natriuretic peptides in chronic kidney disease. 2008 , 3, 1644-51	64
1362	Chronic kidney disease prevalence estimates among racial/ethnic groups: the Multi-Ethnic Study of Atherosclerosis. 2008 , 3, 1391-7	37
1361	Epidemiology, major outcomes, risk factors, prevention and management of chronic kidney disease in China. 2008 , 28, 1-7	58
1360	Effects of albuminuria and renal dysfunction on development of dyslipidaemia in type 2 diabetes--the Hong Kong Diabetes Registry. 2008 , 23, 2834-40	13
1359	Complement deposition in renal histopathology of patients with ANCA-associated pauci-immune glomerulonephritis. 2009 , 24, 1247-52	81
1358	Independent associations between low-density lipoprotein cholesterol and cancer among patients with type 2 diabetes mellitus. 2008 , 179, 427-37	67
1357	Longitudinal analysis of performance of estimated glomerular filtration rate as renal function declines in chronic kidney disease. 2009 , 24, 109-16	24
1356	Is Frequency of Patient Physician Clinic Contact Important in Peritoneal Dialysis Patients?. 2009 , 29, 83-89	8
1355	Drug Dosage Adjustment Using Renal Estimation Equations: A Review of the Literature. 2009 , 44, 577-584	2
1354	Asian leadership in chronic kidney disease. 2009 , 24 Suppl, S3-6	5
1353	The prevalence of chronic kidney disease (CKD) and the associated factors to CKD in urban Korea: a population-based cross-sectional epidemiologic study. 2009 , 24 Suppl, S11-21	68
1352	The international realities of live donor kidney transplantation. 2009 , 75, 1003-5	0
1351	Glomerular filtration rate measurement and prediction equations. 2009 , 47, 1023-32	46
1350	Effect of interactions between C peptide levels and insulin treatment on clinical outcomes among patients with type 2 diabetes mellitus. 2009 , 180, 919-26	15
1349	Improvement of awareness, treatment and control of hypertension among chronic kidney disease patients in China from 1999 to 2005. 2009 , 32, 444-9	12
1348	Dissecting and refining the staging of chronic kidney disease. 2009 , 75, 1009-14	74
1347	Estimated glomerular filtration rate: time for a performance review?. 2009 , 75, 1001-3	20

1346	Interaction between variants of two glycosyltransferase genes in IgA nephropathy. 2009 , 76, 190-8	19
1345	Circulating neutrophil gelatinase-associated lipocalin: a useful biomarker for assessing disease activity of ANCA-associated vasculitis. 2009 , 48, 355-8	19
1344	Reply. 2009 , 24, 2953-2954	1
1343	Disease chronicity and activity predict subclinical left ventricular systolic dysfunction in patients with systemic lupus erythematosus. 2009 , 95, 980-7	25
1342	Evaluation of the Modification of Diet in Renal Disease equation (eGFR) against simultaneous, dual-marker multi-sample measurements of glomerular filtration rate. 2009 , 46, 58-64	4
1341	Low LDL cholesterol, albuminuria, and statins for the risk of cancer in type 2 diabetes: the Hong Kong diabetes registry. 2009 , 32, 1826-32	37
1340	Relation of adiponectin and high-sensitivity C-reactive protein to pulse-wave velocity and N-terminal pro-B-type natriuretic peptide in the general population. 2009 , 103, 1411-6	36
1339	Combination therapy of prednisone and ACE inhibitor versus ACE-inhibitor therapy alone in patients with IgA nephropathy: a randomized controlled trial. 2009 , 53, 26-32	148
1338	Current status and future perspectives for CKD testing. 2009 , 53, S17-26	40
1337	Revised equations for estimated GFR from serum creatinine in Japan. 2009 , 53, 982-92	4244
1336	GFR estimation in Japan and China: what accounts for the difference?. 2009 , 53, 932-5	73
1335	Report of the Asian Forum of Chronic Kidney Disease Initiative (AFCKDI) 2007. "Current status and perspective of CKD in Asia": diversity and specificity among Asian countries. 2009 , 13, 249-56	20
1334	Changes of serum and urine neutrophil gelatinase-associated lipocalin in type-2 diabetic patients with nephropathy: one year observational follow-up study. 2009 , 36, 45-51	63
1333	Phenotype-genotype interactions on renal function in type 2 diabetes: an analysis using structural equation modelling. 2009 , 52, 1543-53	20
1332	Prevalence of chronic kidney disease in Thai adults: a national health survey. 2009 , 10, 35	38
1331	Long-term prognosis of Chinese patients with a lacunar infarct associated with small vessel disease: a five-year longitudinal study. 2009 , 4, 81-8	20
1330	Carotid vascular remodelling in patients with autosomal dominant polycystic kidney disease. 2009 , 14, 113-7	10
1329	Association of interleukin-18 promoter polymorphism and atherosclerotic diseases in Chinese patients with diabetic nephropathy. 2009 , 14, 606-12	11

1328	Concise semiquantitative histological scoring system for immunoglobulin A nephropathy. 2009 , 14, 597-605	19
1327	The choice of estimating equations for glomerular filtration rate significantly affects the prevalence of chronic kidney disease in a multi-ethnic population during health screening. 2009 , 14, 588-96	11
1326	Detection and prevention of chronic kidney disease in Indonesia: initial community screening. 2009 , 14, 669-74	20
1325	Prediction of chronic allograft damage index of renal allografts using serum level of plasminogen activator inhibitor-1. 2009 , 23, 206-12	11
1324	A comparison of prediction equations for estimating glomerular filtration rate in Chinese potential living kidney donors. 2009 , 23, 469-75	13
1323	The Joint Asia Diabetes Evaluation (JADE) Program: a web-based program to translate evidence to clinical practice in Type 2 diabetes. 2009 , 26, 693-9	39
1322	Effects of systolic and diastolic blood pressures on incident coronary heart disease and all-cause death in Chinese women with Type 2 diabetes: the Hong Kong Diabetes Registry. 2009 , 1, 90-8	3
1321	Determination of reference intervals for creatinine and evaluation of creatinine-based estimating equation for Chinese patients with chronic kidney disease. 2009 , 403, 87-91	17
1320	Chronic kidney disease epidemic: cost and health care implications in China. 2009 , 29, 483-6	24
1319	Funci3 renal en el anciano: el pago del tiempo. 2009 , 26, 2-6	1
1318	Measured GFR as a confirmatory test for estimated GFR. <i>Journal of the American Society of Nephrology: JASN</i> , 2009 , 20, 2305-13	12.7 381
1317	Serum uric acid is independently associated with metabolic syndrome in subjects with and without a low estimated glomerular filtration rate. 2009 , 36, 1691-8	40
1316	Combination patterns of cardiovascular risks and sequelae at different stage of hypertension in natural hypertensive population in Beijing. 2009 , 31, 142-55	2
1315	Central or peripheral systolic or pulse pressure: which best relates to target organs and future mortality?. 2009 , 27, 461-7	308
1314	Comparison of drug-eluting stents and coronary artery bypass grafting for the treatment of multivessel coronary artery disease in patients with chronic kidney disease. 2009 , 73, 1228-34	41
1313	Covariate-adjusted nonlinear regression. 2009 , 37,	75
1312	Genetic variants of the protein kinase C-beta 1 gene and development of end-stage renal disease in patients with type 2 diabetes. 2010 , 304, 881-9	49
1311	Tacrolimus combined with corticosteroids in treatment of nephrotic idiopathic membranous nephropathy: a multicenter randomized controlled trial. 2010 , 339, 233-8	83

1310	Urinary neutrophil gelatinase-associated lipocalin: A potential biomarker for predicting rapid progression of drug-induced chronic tubulointerstitial nephritis. 2010 , 339, 537-42	14
1309	Does obesity attenuate the effect of metabolic syndrome on chronic kidney disease in patients with coronary artery disease? Report from China heart survey. 2010 , 74, 462-7	5
1308	Comparison of different antithrombotic regimens for patients with atrial fibrillation undergoing drug-eluting stent implantation. 2010 , 74, 701-8	78
1307	Prevalence and risk factors associated with chronic kidney disease in a Uygur adult population from Urumqi. 2010 , 30, 604-10	9
1306	ASCI 2010 contrast media guideline for cardiac imaging: a report of the Asian Society of Cardiovascular Imaging cardiac computed tomography and cardiac magnetic resonance imaging guideline working group. 2010 , 26, 203-12	10
1305	Clinical characteristics and outcomes of rural patients with ESRD in Guangxi, China: one dialysis center experience. 2010 , 42, 195-204	15
1304	Supportive Management in Patients With End-stage Renal Disease: Local Experience in Hong Kong. 2010 , 12, 31-36	2
1303	Association of hepatitis C and B virus infection with CKD in an endemic area in Taiwan: a cross-sectional study. 2010 , 56, 23-31	87
1302	Estimating GFR using the CKD Epidemiology Collaboration (CKD-EPI) creatinine equation: more accurate GFR estimates, lower CKD prevalence estimates, and better risk predictions. 2010 , 55, 622-7	551
1301	Burden of CKD, proteinuria, and cardiovascular risk among Chinese, Mongolian, and Nepalese participants in the International Society of Nephrology screening programs. 2010 , 56, 915-27	51
1300	From design to implementation--the Joint Asia Diabetes Evaluation (JADE) program: a descriptive report of an electronic web-based diabetes management program. 2010 , 10, 26	37
1299	Assessing glomerular filtration rate (GFR) in elderly Chinese patients with chronic kidney disease (CKD): a comparison of various predictive equations. 2010 , 51, 13-20	40
1298	Gender differences in age-related decline in glomerular filtration rates in healthy people and chronic kidney disease patients. 2010 , 11, 20	40
1297	Evaluation of erectile dysfunction and associated cardiovascular risk using structured questionnaires in Chinese type 2 diabetic men. 2010 , 33, 853-60	19
1296	Independent predictive roles of eotaxin Ala23Thr, paraoxonase 2 Ser311Cys and beta-adrenergic receptor Trp64Arg polymorphisms on cardiac disease in Type 2 Diabetes--an 8-year prospective cohort analysis of 1297 patients. 2010 , 27, 376-83	19
1295	Relationship between beta1-adrenergic receptor polymorphisms and cardiovascular disease in patients with diabetic nephropathy. 2010 , 15, 242-7	
1294	Prevalence and risk factors associated with chronic kidney disease in adults over 40 years: a population study from Central China. 2010 , 15, 354-61	35
1293	Current status of estimated glomerular filtration rate (eGFR) equations for Asians and an approach to create a common eGFR equation. 2010 , 15 Suppl 2, 45-8	22

1292	Association of IRF5 gene polymorphisms and lupus nephritis in a Chinese population. 2010 , 15, 710-3	25
1291	Poster Abstracts. 2010 , 15, 62-144	2
1290	A logistic regression model for microalbuminuria prediction in overweight male population. 2010 ,	1
1289	Ethnic coefficients for glomerular filtration rate estimation by the Modification of Diet in Renal Disease study equations in the Korean population. 2010 , 25, 1616-25	88
1288	Low dietary sodium intake increases the death risk in peritoneal dialysis. 2010 , 5, 240-7	46
1287	Prevalence of CKD in northeastern Italy: results of the INCIPE study and comparison with NHANES. 2010 , 5, 1946-53	50
1286	Suitability of the IDMS-traceable MDRD equation method to estimate GFR in early postoperative renal transplant recipients. 2010 , 114, c108-17	8
1285	Ultrasonic characterization (integrated backscatter) of myocardial tissue in patients with autosomal dominant polycystic kidney disease. 2010 , 114, c288-94	1
1284	The immunopathological spectrum of crescentic glomerulonephritis: a survey of 106 patients in a single Chinese center. 2010 , 116, c65-74	16
1283	Gout: an independent risk factor for all-cause and cardiovascular mortality. 2010 , 49, 141-6	151
1282	Prevalence of chronic kidney disease in a rural Chinese adult population: the Handan Eye Study. 2010 , 114, c295-302	11
1281	Wave reflection and arterial stiffness in the prediction of 15-year all-cause and cardiovascular mortalities: a community-based study. 2010 , 55, 799-805	284
1280	Associations of hyperglycemia and insulin usage with the risk of cancer in type 2 diabetes: the Hong Kong diabetes registry. 2010 , 59, 1254-60	128
1279	Level of kidney function correlates with cognitive decline. 2010 , 32, 117-21	31
1278	Impact of the clinical conditions at dialysis initiation on mortality in incident haemodialysis patients: a national cohort study in Taiwan. 2010 , 25, 2616-24	121
1277	The effect of the World Kidney Day campaign on the awareness of chronic kidney disease and the status of risk factors for cardiovascular disease and renal progression. 2010 , 25, 413-9	16
1276	Differences in progression of CKD and mortality amongst Caucasian, Oriental Asian and South Asian CKD patients. 2010 , 25, 3663-72	62
1275	Association of risk factors for cardiovascular disease and glomerular filtration rate: a community-based study of 4,925 adults in Beijing. 2010 , 25, 3924-31	29

1274	Application of radial basis function neural network to estimate glomerular filtration rate in Chinese patients with chronic kidney disease. 2010 ,	6
1273	Combination of urinary kidney injury molecule-1 and interleukin-18 as early biomarker for the diagnosis and progressive assessment of acute kidney injury following cardiopulmonary bypass surgery: a prospective nested case-control study. 2010 , 15, 332-9	56
1272	White blood cell count and renin-angiotensin system inhibitors for the risk of cancer in type 2 diabetes. 2010 , 87, 117-25	13
1271	Association of statin use and development of renal dysfunction in type 2 diabetes--the Hong Kong Diabetes Registry. 2010 , 88, 227-33	18
1270	Use of sulphonylurea and cancer in type 2 diabetes-The Hong Kong Diabetes Registry. 2010 , 90, 343-51	69
1269	New insights in glomerular filtration rate formulas and chronic kidney disease classification. 2010 , 411, 1341-7	13
1268	A prediction model for the risk of incident chronic kidney disease. 2010 , 123, 836-846.e2	64
1267	Gout and risk of non-alcoholic fatty liver disease. 2010 , 39, 466-71	36
1266	Implication of CKD-EPI equation to estimate glomerular filtration rate in Chinese patients with chronic kidney disease. 2011 , 33, 859-65	16
1265	Significant correlation between left ventricular systolic and diastolic dysfunction and decreased glomerular filtration rate. 2011 , 33, 977-82	5
1264	Prognostic impact of body mass index in patients undergoing coronary artery bypass surgery. 2011 , 97, 648-54	26
1263	Evaluation of GFR measurement method as an explanation for differences among GFR estimation equations. 2011 , 58, 496-8	21
1262	Nondiabetic older adults with untreated hypertension in Taiwan: Treatment implication in elderly hypertension. 2011 , 2, 58-61	1
1261	Acute-on-chronic kidney injury at hospital discharge is associated with long-term dialysis and mortality. 2011 , 80, 1222-30	127
1260	Metabolic syndrome and chronic kidney disease in a rural Chinese population. 2011 , 412, 1983-8	13
1259	Associations of intra-renal arterial resistance index with chronic kidney disease and carotid intima-media thickness in type 2 diabetes mellitus. 2011 , 92, e37-40	2
1258	Low HDL cholesterol, metformin use, and cancer risk in type 2 diabetes: the Hong Kong Diabetes Registry. 2011 , 34, 375-80	56
1257	A multicenter study of the association of serum uric acid, serum creatinine, and diuretic use in hypertensive patients. 2011 , 148, 325-30	19

1256	Association between high-sensitivity cardiac troponin T and predicted cardiovascular risks in a community-based population. 2011 , 149, 253-256	6
1255	Plasma asymmetric dimethylarginine predicts death and major adverse cardiovascular events in individuals referred for coronary angiography. 2011 , 153, 135-40	60
1254	Evaluation of the Chronic Kidney Disease Epidemiology Collaboration equation for estimating the glomerular filtration rate in multiple ethnicities. 2011 , 79, 555-62	288
1253	Predictive role of polymorphisms in interleukin-5 receptor alpha-subunit, lipoprotein lipase, integrin A2 and nitric oxide synthase genes on ischemic stroke in type 2 diabetes--an 8-year prospective cohort analysis of 1327 Chinese patients. 2011 , 215, 130-5	18
1252	Serum fetuin-A is correlated with metabolic syndrome in middle-aged and elderly Chinese. 2011 , 216, 180-6	62
1251	Is chronic kidney disease associated with a high ankle brachial index in adults at high cardiovascular risk?. 2011 , 18, 224-30	9
1250	Serum fetuin-A associates with type 2 diabetes and insulin resistance in Chinese adults. 2011 , 6, e19228	50
1249	Urinary podocyte-associated mRNA profile in various stages of diabetic nephropathy. 2011 , 6, e20431	67
1248	Association of low glomerular filtration rate with the incidence of stroke in patients following complete coronary revascularization. 2011 , 75, 2372-8	5
1247	Inaccuracies in estimated glomerular filtration rate in one Australian renal centre. 2011 , 16, 486-94	3
1246	Estimated glomerular filtration rate: which equation is best and how should it be used?. 2011 , 16, 483-5	1
1245	Short-term outcomes of induction therapy with tacrolimus versus cyclophosphamide for active lupus nephritis: A multicenter randomized clinical trial. 2011 , 57, 235-44	139
1244	Clinical features and outcomes of anti-glomerular basement membrane disease in older patients. 2011 , 57, 575-82	40
1243	GFR estimating equations in a multiethnic Asian population. 2011 , 58, 56-63	176
1242	Comparisons of technetium-99m diethylenetriaminepentaacetic acid plasma clearance and renal dynamic imaging with inulin clearance. 2011 , 58, 1043-5	11
1241	A comparison of clinical outcomes of Chinese sirolimus-eluting stents versus foreign sirolimus-eluting stents for the treatment of coronary artery disease. 2011 , 19, 418-22	1
1240	Hyperuricemia and metabolic syndrome: associations with chronic kidney disease. 2011 , 30, 323-30	18
1239	Obesity and chronic kidney disease in patients with chronic heart failure: an insight from the China Heart Survey. 2011 , 15, 522-8	5

1238	The association of dimethylarginine dimethylaminohydrolase 1 gene polymorphism with type 2 diabetes: a cohort study. 2011 , 10, 16	17
1237	Low triglyceride and nonuse of statins is associated with cancer in type 2 diabetes mellitus: the Hong Kong Diabetes Registry. 2011 , 117, 862-71	17
1236	Normoalbuminaemia is associated with IgA nephropathy in primary glomerulopathy with nephrotic-range proteinuria in Chinese patients. 2011 , 26, 1247-52	11
1235	Prevalence and risk factors of chronic kidney disease: a population study in the Tibetan population. 2011 , 26, 1592-9	62
1234	Cancer-specific mortality in chronic kidney disease: longitudinal follow-up of a large cohort. 2011 , 6, 1121-8	97
1233	Relationship of urinary bisphenol A concentration to risk for prevalent type 2 diabetes in Chinese adults: a cross-sectional analysis. 2011 , 155, 368-74	96
1232	Pathologic predictors of renal outcome and therapeutic efficacy in IgA nephropathy: validation of the oxford classification. 2011 , 6, 2175-84	117
1231	CPVL/CHN2 genetic variant is associated with diabetic retinopathy in Chinese type 2 diabetic patients. 2011 , 60, 3085-9	27
1230	Assessment of the CKD-EPI equation to estimate glomerular filtration rate in adults from a Chinese CKD population. 2011 , 39, 2273-80	21
1229	Effect of blood sample type on the measurement of advanced oxidation protein products as a biomarker of inflammation and oxidative stress in hemodialysis patients. 2011 , 16, 129-35	10
1228	C-reactive protein and the risk of contrast-induced acute kidney injury in patients undergoing percutaneous coronary intervention. 2011 , 34, 203-10	53
1227	Regular use of nephrotoxic medications is an independent risk factor for chronic kidney disease--results from a Chinese population study. 2011 , 26, 1916-23	10
1226	The need for robust validation for MDRD-based glomerular filtration rate estimation in various CKD populations. 2011 , 26, 2780-5	70
1225	Mortality and renal function decline among a community-based Chinese population with normal or mildly impaired renal function. 2011 , 26, 2847-52	12
1224	Reply. 2011 , 26, 2718-2719	
1223	Polymorphisms in the nonmuscle myosin heavy chain 9 gene (MYH9) are associated with the progression of IgA nephropathy in Chinese. 2011 , 26, 2544-9	27
1222	Associations of renal vascular resistance with albuminuria in adolescents and young adults. 2011 , 26, 3943-9	6
1221	Prognostic significance of glomerular filtration rate estimated by the Japanese equation among patients who underwent complete coronary revascularization. 2011 , 34, 378-83	6

1220	Estimating glomerular filtration rate in Asian subjects: where do we stand?. 2011 , 80, 439-40	18
1219	The contribution of chronic kidney disease to the global burden of major noncommunicable diseases. 2011 , 80, 1258-70	784
1218	A new tool to detect kidney disease in Chinese type 2 diabetes patients: comparison of EZSCAN with standard screening methods. 2011 , 13, 937-43	25
1217	The clinician and estimation of glomerular filtration rate by creatinine-based formulas: current limitations and quo vadis. 2011 , 6, 937-50	73
1216	Pulsatile hemodynamics and clinical outcomes in acute heart failure. 2011 , 24, 775-82	28
1215	Risk of acute kidney injury after exposure to gadolinium-based contrast in patients with renal impairment. 2011 , 33, 758-64	20
1214	Asymmetric dimethylarginine and clinical outcomes in chronic kidney disease. 2011 , 6, 1566-72	59
1213	Evaluation of serum creatinine- and cystatin C-based equations for the estimation of glomerular filtration rate in a Chinese population. 2012 , 46, 223-31	12
1212	Tacrolimus improves the proteinuria remission in patients with refractory IgA nephropathy. 2012 , 35, 312-20	34
1211	Gender difference in the association of hyperuricemia with chronic kidney disease in southern China. 2012 , 36, 98-106	5
1210	The association between plasma uric acid and renal function decline in a Chinese population-based cohort. 2012 , 27, 1836-9	29
1209	Induction therapy with autologous mesenchymal stem cells in living-related kidney transplants: a randomized controlled trial. 2012 , 307, 1169-77	431
1208	SLICC/ACR damage index independently associated with left ventricular diastolic dysfunction in patients with systemic lupus erythematosus. 2012 , 21, 1057-62	20
1207	Association between nonalcoholic fatty liver disease (NAFLD) and osteoporotic fracture in middle-aged and elderly Chinese. 2012 , 97, 2033-8	96
1206	Predictive role of multilocus genetic polymorphisms in cardiovascular disease and inflammation-related genes on chronic kidney disease in Type 2 diabetes--an 8-year prospective cohort analysis of 1163 patients. 2012 , 27, 190-6	11
1205	Effect of multi-dimensional education on disease progression in pre-dialysis patients in China. 2012 , 34, 47-52	10
1204	Imbalance of regulatory T cells to Th17 cells in IgA nephropathy. 2012 , 72, 221-9	54
1203	Excessive wave reflections on admission predict post-discharge events in patients hospitalized due to acute heart failure. 2012 , 14, 1348-55	24

1202	The impact of CKD identification in large countries: the burden of illness. 2012 , 27 Suppl 3, iii32-8	74
1201	Exposure to bisphenol A is associated with low-grade albuminuria in Chinese adults. 2012 , 81, 1131-9	52
1200	Ethnic factors of the glomerular filtration rate estimating equation. 2012 , 81, 799; author reply 799-800	3
1199	Estimating glomerular filtration rates by use of both cystatin C and standardized serum creatinine avoids ethnicity coefficients in Asian patients with chronic kidney disease. 2012 , 58, 450-7	25
1198	Re-evaluation of the histopathologic classification of ANCA-associated glomerulonephritis: a study of 121 patients in a single center. 2012 , 27, 2343-9	90
1197	Clinical features of Chinese coronary heart disease patients with chronic kidney disease. 2012 , 34, 985-90	2
1196	Serum cystatin C is an early biomarker for assessment of renal function in burn patients. 2012 , 50, 667-71	11
1195	Association of reduced renal function with hepatitis B virus infection and elevated alanine aminotransferase. 2012 , 7, 1561-6	12
1194	Aristolochic acid nephropathy: variation in presentation and prognosis. 2012 , 27, 292-8	72
1193	Predicting outcomes after myocardial infarction by using the Chronic Kidney Disease Epidemiology Collaboration equation in comparison with the Modification of Diet in Renal Disease study equation: results from the Korea Acute Myocardial Infarction Registry. 2012 , 27, 3868-74	10
1192	Comparisons between validated estimated glomerular filtration rate equations and isotopic glomerular filtration rate in HIV patients. 2012 , 26, 1781-8	30
1191	Outcome of lamivudine-resistant chronic hepatitis B after up to 5 years of combination therapy with adefovir. 2012 , 17, 1255-62	15
1190	Urinary angiotensinogen excretion and ambulatory blood pressure. 2012 , 30, 2000-6	16
1189	Estimating Equations for Glomerular Filtration Rate in the Era of Creatinine Standardization. 2012 , 156, 785	286
1188	Renal protection of losartan 50 mg in normotensive Chinese patients with nondiabetic chronic kidney disease. 2012 , 60, 1041-7	9
1187	Prevalence of chronic kidney disease in China: a cross-sectional survey. 2012 , 379, 815-22	1170
1186	Urinary mRNA expression of CCN2/CCN3 as a noninvasive marker for monitoring glomerular structure changes in nondiabetic chronic kidney disease. 2012 , 17, 714-20	3
1185	Synergistic effects of low LDL cholesterol with other factors for the risk of cancer in type 2 diabetes: the Hong Kong Diabetes Registry. 2012 , 49 Suppl 1, S185-93	11

1184	Diabetes and hypertension have become leading causes of CKD in Chinese elderly patients: a comparison between 1990-1991 and 2009-2010. 2012 , 44, 1269-76	23
1183	Impact of acute kidney injury on one-year survival after surgery for aortic dissection. 2012 , 94, 1407-12	52
1182	A decade after the KDOQI CKD guidelines: a perspective from China. 2012 , 60, 727-8	4
1181	Chronic kidney disease and its association with mortality and hospitalization in Chinese nursing home older residents: a 3-year prospective cohort study. 2012 , 13, 782-7	10
1180	Prevalence and associated comorbidities of moderate to severe chronic renal impairment in Chinese nursing home older adults. 2012 , 13, 630-3	14
1179	Asymmetric dimethylarginine predicts clinical outcomes in ischemic chronic heart failure. 2012 , 225, 504-10	40
1178	Advanced oxidation protein products as prognostic biomarkers for recovery from acute kidney injury after coronary artery bypass grafting. 2012 , 17, 507-12	15
1177	In addition to malnutrition and renal function impairment, anemia is associated with hyponatremia in the elderly. 2012 , 55, 77-81	10
1176	Prevalence of hyperuricemia and its association with antihypertensive treatment in hypertensive patients in Taiwan. 2012 , 156, 41-6	21
1175	Association of the PPARG Pro12Ala polymorphism with type 2 diabetes and incident coronary heart disease in a Hong Kong Chinese population. 2012 , 97, 483-91	23
1174	Association between blood glucose levels and high-sensitivity cardiac troponin T in an overt cardiovascular disease-free community-based study. 2012 , 97, 139-45	19
1173	Plasma Uric Acid is Associated with Postinfarction Cardiac Remodeling in Elderly with Old Myocardial Infarction. 2012 , 6, 174-176	1
1172	Chronic kidney disease predicts poor prognosis in patients with stable premature coronary artery disease. 2012 , 23, 716-9	7
1171	Acute kidney injury and bilateral symmetrical enlargement of the kidneys as first presentation of B-cell lymphoblastic lymphoma. 2012 , 60, 1044-8	9
1170	C5a and its receptors in human anti-neutrophil cytoplasmic antibody (ANCA)-associated vasculitis. 2012 , 14, R140	66
1169	U-curve association between timing of renal replacement therapy initiation and in-hospital mortality in postoperative acute kidney injury. 2012 , 7, e42952	34
1168	Acute kidney injury biomarkers for patients in a coronary care unit: a prospective cohort study. 2012 , 7, e32328	37
1167	A pilot trial assessing urinary gene expression profiling with an mRNA array for diabetic nephropathy. 2012 , 7, e34824	21

1166	Association of glomerular filtration rate with high-sensitivity cardiac troponin T in a community-based population study in Beijing. 2012 , 7, e38218	15
1165	Predicting progression of IgA nephropathy: new clinical progression risk score. 2012 , 7, e38904	87
1164	High prevalence and associated risk factors for impaired renal function and urinary abnormalities in a rural adult population from southern China. 2012 , 7, e47100	19
1163	Chronic kidney disease and automatic reporting of estimated glomerular filtration rate: new developments and revised recommendations. 2012 , 197, 224-5	95
1162	Serum small-dense LDL abnormalities in chronic renal disease patients. 2012 , 69, 99-102	20
1161	The impact of different GFR estimating equations on the prevalence of CKD and risk groups in a Southeast Asian cohort using the new KDIGO guidelines. 2012 , 13, 1	30
1160	Comparison of the predictive value of EuroSCORE, SYNTAX score, and clinical SYNTAX score for outcomes of patients undergoing percutaneous coronary intervention for unprotected left main coronary artery disease. 2012 , 80, 222-30	17
1159	Chronic kidney disease and the risk of stent thrombosis after percutaneous coronary intervention with drug-eluting stents. 2012 , 80, 361-7	28
1158	Predictive performance of eGFR equations in South Africans of African and Indian ancestry compared with ¹²⁵ I-Tc-DTPA imaging. 2012 , 44, 847-55	32
1157	Importance and benefits of dietary sodium restriction in the management of chronic kidney disease patients: experience from a single Chinese center. 2012 , 44, 549-56	22
1156	Comparison of clinicopathological features between children and adults with IgA nephropathy. 2012 , 27, 1293-300	9
1155	Frequency of established cardiovascular disease and its risk factors in Chinese patients with systemic lupus erythematosus. 2012 , 31, 669-75	18
1154	Chronic kidney disease prevention--a challenge for Asian countries: report of the Third Asian Forum of Chronic Kidney Disease Initiatives. 2012 , 16, 187-94	4
1153	Prevalence and risk factors of chronic kidney disease in first-degree relatives of chronic kidney disease patients in Southern China. 2012 , 17, 123-30	6
1152	Oral calcitriol for reduction of proteinuria in patients with IgA nephropathy: a randomized controlled trial. 2012 , 59, 67-74	56
1151	Association between GFR, proteinuria, and adverse outcomes among White, Chinese, and South Asian individuals in Canada. 2012 , 59, 390-9	24
1150	Association between preoperative allograft function (effective renal plasma flow) and the change in glomerular filtration rate among living-donor kidney transplant recipients. 2012 , 44, 248-53	4
1149	Long-term outcomes of living kidney donors over the past 28 years in a single center in Taiwan. 2012 , 44, 39-42	9

1148	Urinary mRNA markers of epithelial-mesenchymal transition correlate with progression of diabetic nephropathy. 2012 , 76, 657-64	22
1147	Comparison of prediction equations to estimate glomerular filtration rate in Chinese patients with chronic kidney disease. 2012 , 42, e59-67	13
1146	Urinary tubular biomarkers in short-term type 2 diabetes mellitus patients: a cross-sectional study. 2012 , 41, 82-8	91
1145	Common genetic variants of the human uromodulin gene regulate transcription and predict plasma uric acid levels. 2013 , 83, 733-40	26
1144	Associations of RBP4 with lipid metabolism and renal function in diabetes mellitus. 2013 , 115, 831-837	2
1143	A multicenter, randomized, double-blind, parallel-group, placebo-controlled study of the effects of qili qiangxin capsules in patients with chronic heart failure. 2013 , 62, 1065-1072	171
1142	Using a multi-staged strategy based on machine learning and mathematical modeling to predict genotype-phenotype risk patterns in diabetic kidney disease: a prospective case-control cohort analysis. 2013 , 14, 162	29
1141	Association between family members of dialysis patients and chronic kidney disease: a multicenter study in China. 2013 , 14, 19	9
1140	Questionnaire survey evaluating disease-related knowledge for 149 primary gout patients and 184 doctors in South China. 2013 , 32, 1633-40	11
1139	A higher glomerular filtration rate predicts low risk of developing chronic kidney disease in living kidney donors. 2013 , 37, 923-9	8
1138	Assessment of glomerular filtration rate in elderly patients with chronic kidney disease. 2013 , 45, 1475-82	17
1137	Improvement and influencing factors of blood pressure control by nephrologist referral in chronic kidney disease patients in China: a cohort study. 2013 , 45, 1345-53	4
1136	Impact of anemia on contrast-induced nephropathy (CIN) in patients undergoing percutaneous coronary interventions. 2013 , 45, 1065-70	21
1135	Association of handgrip strength with chronic diseases and multimorbidity: a cross-sectional study. 2013 , 35, 929-41	103
1134	Effect of far infrared therapy on arteriovenous fistula maturation: an open-label randomized controlled trial. 2013 , 62, 304-11	41
1133	An odd application of artificial vascular graft in the kidney transplantation of a patient with severe deep vein thrombosis: a case report. 2013 , 45, 2819-21	
1132	Impact of elevated serum glycated albumin levels on contrast-induced acute kidney injury in diabetic patients with moderate to severe renal insufficiency undergoing coronary angiography. 2013 , 167, 369-73	5
1131	Prevalence of chronic kidney disease across levels of glycemia among adults in Pudong New Area, Shanghai, China. 2013 , 14, 253	16

1130	The ambulatory arterial stiffness index and target-organ damage in Chinese patients with chronic kidney disease. 2013 , 14, 257	6
1129	Long-term prognostic impact of cystatin C on acute coronary syndrome octogenarians with diabetes mellitus. 2013 , 12, 157	11
1128	Ethnic differences in creatinine kinetics in a New Zealand end-stage kidney disease cohort. 2013 , 18, 222-8	1
1127	Chronic Kidney Disease: Pathophysiology and the Influence of Dietary Protein. 2013 , 3021-3072	
1126	Prevalence of anemia in Chinese nursing home older adults: implication of age and renal impairment. 2013 , 13, 591-6	9
1125	Renal transporters in drug development. 2013 , 53, 503-29	226
1124	Long-term data on corticosteroids and mycophenolate mofetil treatment in lupus nephritis. 2013 , 52, 480-6	39
1123	Association between aristolochic acid and CKD: a cross-sectional survey in China. 2013 , 61, 918-22	15
1122	Detection of viral antigens in renal tissue of glomerulonephritis patients without serological evidence of hepatitis B virus and hepatitis C virus infection. 2013 , 17, e535-8	19
1121	Efficacy of trivalent seasonal influenza vaccination in reducing mortality and hospitalization in Chinese nursing home older adults. 2013 , 14, 889-94	10
1120	Sequential changes in renal function and the risk of stroke and death in patients with atrial fibrillation. 2013 , 168, 4678-84	39
1119	The efficacy of influenza vaccination is reduced in nursing home older adults with moderate to severe renal impairment. 2013 , 14, 133-6	1
1118	Predictors for outcomes in patients with severe ANCA-associated glomerulonephritis who were dialysis-dependent at presentation: a study of 89 cases in a single Chinese center. 2013 , 42, 515-21	26
1117	Proteomic analysis of adverse outcomes in patients with acute coronary syndromes. 2013 , 416, 60-6	2
1116	A comparison of the performances of an artificial neural network and a regression model for GFR estimation. 2013 , 62, 1109-15	17
1115	Measurement of renal function in patients with chronic kidney disease. 2013 , 76, 504-15	31
1114	Relation of estimated glomerular filtration rate and body mass index to mortality in non-dialysis patients with coronary artery disease: a report from the ET-CHD registry, 1997-2003. 2013 , 62, 144-50	3
1113	Treatment of early immunoglobulin A nephropathy by angiotensin-converting enzyme inhibitor. 2013 , 126, 162-8	16

1112	Plasma levels and diagnostic value of catestatin in patients with heart failure. 2013 , 46, 20-5	32
1111	Relation of renal dysfunction to the increased risk of stroke and death in female patients with atrial fibrillation. 2013 , 168, 1502-8	22
1110	Reduced number and impaired function of circulating endothelial progenitor cells in patients with abdominal aortic aneurysm. 2013 , 168, 1070-7	22
1109	Chronic kidney disease and long-term outcomes of myocardial infarction. 2013 , 167, 2490-5	7
1108	Prospective monitoring N-terminal pro-brain natriuretic peptide during natural progression of atherosclerotic renal artery stenosis in elderly. 2013 , 56, 395-9	2
1107	Endovascular treatment of renal artery aneurysms and renal arteriovenous fistulas. 2013 , 57, 765-70	53
1106	Chapter 1: Definition and classification of CKD. 2013 , 3, 19-62	565
1105	Increased plasma sVCAM-1 is associated with severity in IgA nephropathy. 2013 , 14, 21	14
1104	HbA1c and the diagnosis of diabetes and prediabetes in a middle-aged and elderly Han population from northwest China (HbA1c). 2013 , 5, 282-90	18
1103	References. 2013 , 3, 136-150	66
1102	Remote ischemic preconditioning reduces myocardial injury in patients undergoing coronary stent implantation. 2013 , 29, 1084-9	67
1101	Chronic kidney disease: global dimension and perspectives. 2013 , 382, 260-72	2280
1100	Prevalence, clinical predictors, and prognostic impact of chronic renal insufficiency in very old Chinese patients with coronary artery disease. 2013 , 25, 385-91	2
1099	Enhancers and attenuators of risk associations of chronic hepatitis B virus infection with hepatocellular carcinoma in type 2 diabetes. 2013 , 20, 161-71	16
1098	Association between percent-free prostate-specific antigen and glomerular filtration rate in transrectal ultrasound-guided biopsy-proven patients with prostate-specific antigen levels ranging from 4 to 10 ng/ml. 2013 , 31, 313-8	
1097	The applicability of eGFR equations to different populations. 2013 , 9, 513-22	96
1096	Con: Should we abandon the use of the MDRD equation in favour of the CKD-EPI equation?. 2013 , 28, 1396-403; discussion 403	42
1095	Pro: Estimating GFR using the chronic kidney disease epidemiology collaboration (CKD-EPI) 2009 creatinine equation: the time for change is now. 2013 , 28, 1390-6	27

1094	Evaluation of the Chronic Kidney Disease Epidemiology Collaboration equation for estimating glomerular filtration rate in the Chinese population. 2013 , 28, 641-51	119
1093	Activation of the interleukin-4/signal transducer and activator of transcription 6 signaling pathway and homeodomain-interacting protein kinase 2 production by tonsillar mononuclear cells in IgA nephropathy. 2013 , 38, 321-32	23
1092	Urinary TNF- α and NGAL are correlated with the progression of nephropathy in patients with type 2 diabetes. 2013 , 6, 1482-1488	27
1091	Is blood pressure load associated, independently of blood pressure level, with target organ damage?. 2013 , 31, 1812-8	26
1090	Association of central pulse pressure with contrast-induced nephropathy and clinical outcomes in patients undergoing coronary intervention. 2013 , 31, 2187-94	9
1089	Association of retinopathy with coronary atherosclerosis determined by coronary 64-slice multidetector computed tomography angiography in type 2 diabetes. 2013 , 10, 161-8	9
1088	GFR estimating equations: getting closer to the truth?. 2013 , 8, 1414-20	46
1087	Homocysteine and C-reactive protein as useful surrogate markers for evaluating CKD risk in adults. 2013 , 37, 402-13	10
1086	Chronic kidney disease: an independent risk factor of all-cause mortality for elderly Chinese patients with chronic heart failure. 2012 , 9, 355-60	2
1085	Prevalence and risk factors of CKD in Chinese patients with periodontal disease. 2013 , 8, e70767	11
1084	Central obesity, C-reactive protein and chronic kidney disease: a community-based cross-sectional study in southern China. 2013 , 37, 392-401	21
1083	Blood pressure in relation to interactions between sodium dietary intake and renal handling. 2013 , 62, 719-25	10
1082	The impact of hyperglycemic emergencies on the kidney and liver. 2013 , 2013, 967097	9
1081	Interaction between PLA2R1 and HLA-DQA1 variants associates with anti-PLA2R antibodies and membranous nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2013 , 24, 1323-9	12.7 101
1080	Risk association of HbA1c variability with chronic kidney disease and cardiovascular disease in type 2 diabetes: prospective analysis of the Hong Kong Diabetes Registry. 2013 , 29, 384-90	92
1079	Reduced kidney function in acute coronary syndrome patients with undiagnosed diabetes or pre-diabetes. 2013 , 18, 263-8	3
1078	Plasma levels and potential roles of cathepsin in patients with coronary heart disease. 2013 , 47, 217-24	20
1077	Derivation and validation of a prediction score for acute kidney injury in patients hospitalized with acute heart failure in a Chinese cohort. 2013 , 18, 489-96	19

1076	Individuals of Pacific Asian origin with IgA nephropathy have an increased risk of progression to end-stage renal disease. 2013 , 84, 1017-24	87
1075	Predictive factors for worse pathological outcomes of upper tract urothelial carcinoma: experience from a nationwide high-volume centre in China. 2013 , 112, 917-24	47
1074	Outcomes of tacrolimus therapy in adults with refractory membranous nephrotic syndrome: a prospective, multicenter clinical trial. 2013 , 345, 81-7	16
1073	Predictive value of serum uric acid levels on mortality in acute coronary syndrome patients with chronic kidney disease after drug-eluting stent implantation. 2013 , 125, 204-12	13
1072	Association of increased serum glycated albumin levels with low coronary collateralization in type 2 diabetic patients with stable angina and chronic total occlusion. 2013 , 12, 165	28
1071	Serum surfactant protein D is associated with the prognosis in patients with chronic kidney disease. 2013 , 14, 461-5	9
1070	Contemporary management and attainment of cholesterol targets for patients with dyslipidemia in China. 2013 , 8, e47681	13
1069	Acoustic radiation force impulse imaging for noninvasive evaluation of renal parenchyma elasticity: preliminary findings. 2013 , 8, e68925	77
1068	Prognostic abilities of different calculation formulas for the glomerular filtration rate in elderly Chinese patients with coronary artery disease. 2013 , 8, 229-37	6
1067	Estimating glomerular filtration rates in elderly Chinese patients with chronic kidney disease: performance of six modified formulae developed in Asian populations. 2013 , 8, 899-904	4
1066	Is the Chronic Kidney Disease Epidemiology Collaboration creatinine-cystatin C equation useful for glomerular filtration rate estimation in the elderly?. 2013 , 8, 1387-91	17
1065	The ability of NT-proBNP to detect chronic heart failure and predict all-cause mortality is higher in elderly Chinese coronary artery disease patients with chronic kidney disease. 2013 , 8, 409-17	9
1064	FCGR2B and FCRLB gene polymorphisms associated with IgA nephropathy. 2013 , 8, e61208	18
1063	MicroRNA-21 and risk of severe acute kidney injury and poor outcomes after adult cardiac surgery. 2013 , 8, e63390	64
1062	Synergistic effect of mesangial cell-induced CXCL1 and TGF- β in promoting podocyte loss in IgA nephropathy. 2013 , 8, e73425	29
1061	Relationships between HDL-C, hs-CRP, with central arterial stiffness in apparently healthy people undergoing a general health examination. 2013 , 8, e81778	20
1060	The safety and short-term efficacy of aliskiren in the treatment of immunoglobulin a nephropathy--a randomized cross-over study. 2013 , 8, e62736	13
1059	Comparison of the efficacy of rosuvastatin versus atorvastatin in preventing contrast induced nephropathy in patient with chronic kidney disease undergoing percutaneous coronary intervention. 2014 , 9, e111124	14

1058	Association of renal biochemical parameters with left ventricular diastolic dysfunction in a community-based elderly population in China: a cross-sectional study. 2014, 9, e88638	9
1057	Association of serum irisin with metabolic syndrome in obese Chinese adults. 2014, 9, e94235	103
1056	Modification of diet in renal disease (MDRD) study and CKD epidemiology collaboration (CKD-EPI) equations for Taiwanese adults. 2014, 9, e99645	36
1055	Association of metabolic syndrome with decreased glomerular filtration rate among 75,468 Chinese adults: a cross-sectional study. 2014, 9, e113450	16
1054	Chronic kidney disease screening methods and its implication for Malaysia: an in depth review. 2015, 7, 96-109	8
1053	Association of uric acid with metabolic syndrome in men, premenopausal women and postmenopausal women. 2014, 11, 2899-910	28
1052	Serum lipid profiles, lipid ratios and Chronic Kidney Disease in a Chinese population. 2014, 11, 7622-35	11
1051	Health-Related Quality of Life in Chinese Patients with Type 2 Diabetes: An Analysis of the Joint Asia Diabetes Evaluation (JADE) Program. 2014, 05,	2
1050	Different types of atrial fibrillation, renal function, and mortality in elderly Chinese patients with coronary artery disease. 2014, 9, 301-8	3
1049	Comparison of Ten Creatinine-based Equations for Estimation of Glomerular Filtration Rate in Chinese Postmenopausal Women with Normal or Mildly Reduced Renal Function. 2014, 03,	
1048	Association between resting heart rate and N-terminal pro-brain natriuretic peptide in a community-based population study in Beijing. 2015, 10, 55-60	1
1047	Cystatin C Versus Creatinine- Based Definition of Renal Dysfunction for Predicting Poor Coronary Collateralization in Type 2 Diabetic Patients with Stable Coronary Artery Disease. 2014, 5,	
1046	Th1/Th2 polarization in tonsillar lymphocyte form patients with IgA nephropathy. 2014, 36, 407-12	19
1045	Relationship between prehypertension and incidence of chronic kidney disease in a general population: a prospective analysis in central south China. 2014, 46, 2183-9	8
1044	Detecting diabetic risk using the oral glucose tolerance test in Chinese patients with hypertension: a cross-sectional study. 2014, 37, 82-7	6
1043	Modified glomerular filtration rate-estimating equations developed in asiatic population for chinese patients with type 2 diabetes. 2014, 2014, 521071	13
1042	Prevalence of chronic kidney disease among adults in a rural community in South India: Results from the kidney disease screening (KIDS) project. 2014, 24, 214-21	36
1041	Free triiodothyronine concentrations are inversely associated with microalbuminuria. 2014, 2014, 959781	11

1040	Severe hypoglycemia identifies vulnerable patients with type 2 diabetes at risk for premature death and all-site cancer: the Hong Kong diabetes registry. 2014 , 37, 1024-31	52
1039	Prospective study on the incidences of cardiovascular-renal complications in Chinese patients with young-onset type 1 and type 2 diabetes. 2014 , 37, 149-57	70
1038	Lack of association between TLR4 genetic polymorphisms and diabetic nephropathy in a Chinese population. 2014 , 2014, 704167	2
1037	Preliminary study of Huai Qi Huang granules delay the development of primary glomerular diseases in human. 2014 , 36, 1407-10	7
1036	Nomogram predicting renal insufficiency after nephroureterectomy for upper tract urothelial carcinoma in the Chinese population: exclusion of ineligible candidates for adjuvant chemotherapy. 2014 , 2014, 529186	14
1035	Clinical analysis of the relationship between cystatin C and metabolic syndrome in the elderly. 2014 , 33, 411-6	11
1034	Metabolic syndrome and chronic kidney disease in a Southern Chinese population. 2014 , 19, 325-31	17
1033	Evolving renorrhaphy technique for retroperitoneal laparoscopic partial nephrectomy: single-surgeon series. 2014 , 21, 865-73	10
1032	Low serum potassium level is associated with nonalcoholic fatty liver disease and its related metabolic disorders. 2014 , 80, 348-55	19
1031	Long-term data on tacrolimus treatment in lupus nephritis. 2014 , 53, 2232-7	37
1030	Clinical analysis of the PADUA and the RENAL scoring systems for renal neoplasms: a retrospective study of 245 patients undergoing laparoscopic partial nephrectomy. 2014 , 21, 40-4	24
1029	Correlation between comprehensive evaluation of coronary artery lesion severity and long-term clinical outcomes in Chinese octogenarians with acute coronary syndrome. 2014 , 23, 1125-31	3
1028	CD4 (+)CD 25 (+)Treg cells and IgA nephropathy patients with tonsillectomy: a clinical and pathological study. 2014 , 46, 2361-9	11
1027	Opposite association between diabetes, dyslipidemia, and hepatocellular carcinoma mortality in the middle-aged and elderly. 2014 , 59, 2207-15	34
1026	Leflunomide in addition to steroids improves proteinuria and renal function in adult Henoch-Schoenlein nephritis with nephrotic proteinuria. 2014 , 19, 94-100	9
1025	Prevalence and risk factors associated with prehypertension in Shunde District, southern China. 2014 , 4, e006551	9
1024	Kidneys from standard-criteria donors with different severities of terminal acute kidney injury. 2014 , 46, 3335-8	9
1023	Association between metabolically unhealthy overweight/obesity and chronic kidney disease: the role of inflammation. 2014 , 40, 423-30	20

1022	WASP neuronet activated by bipolar-sigmoid functions and applied to glomerular-filtration-rate estimation. 2014,	2
1021	Wave reflections, arterial stiffness, and orthostatic hypotension. 2014, 27, 1446-55	15
1020	Association between exposure to hepatitis B virus and chronic kidney disease in China. 2014, 42, 1178-84	11
1019	Prevalence of chronic kidney disease and associated factors among the Chinese population in Taian, China. 2014, 15, 205	18
1018	Hypoglycaemia, chronic kidney disease and death in type 2 diabetes: the Hong Kong diabetes registry. 2014, 14, 48	21
1017	A comparison of different diagnostic criteria of acute kidney injury in critically ill patients. 2014, 18, R144	149
1016	Dynamic change of glomerular filtration rate in the early stage is associated with kidney allograft status: a preliminary report. 2014, 19, 72	
1015	Staging and risk factors of chronic kidney disease of outpatients in Shanghai. 2014, 36, 1018-22	2
1014	Genetic variants for type 2 diabetes and new-onset cancer in Chinese with type 2 diabetes. 2014, 103, 328-37	13
1013	Fluorescent polarity probes for identifying bovine serum albumin: Amplification effect of para-substituted benzene. 2014, 103, 1-8	32
1012	Effects of tea consumption on renal function in a metropolitan Chinese population: the Guangzhou biobank cohort study. 2014, 24, 26-31	3
1011	Metabolic syndrome and chronic kidney disease in general Chinese adults: results from the 2007-08 China National Diabetes and Metabolic Disorders Study. 2014, 430, 115-20	26
1010	Improved measurement of the glomerular filtration rate from Tc-99m DTPA scintigraphy in patients following nephrectomy. 2014, 24, 413-22	8
1009	Prevalence of chronic kidney disease and its association with metabolic diseases: a cross-sectional survey in Zhejiang province, Eastern China. 2014, 15, 36	38
1008	Prevention of contrast-induced nephropathy with prostaglandin E1 in high-risk patients undergoing percutaneous coronary intervention. 2014, 46, 781-6	18
1007	Risk factors and treatment outcomes of new contralateral upper urinary urothelial carcinoma after nephroureterectomy: the experiences of a large Chinese center. 2014, 140, 477-85	23
1006	The eGFR-C study: accuracy of glomerular filtration rate (GFR) estimation using creatinine and cystatin C and albuminuria for monitoring disease progression in patients with stage 3 chronic kidney disease--prospective longitudinal study in a multiethnic population. 2014, 15, 13	28
1005	Predictors of treatment resistance and relapse in antineutrophil cytoplasmic antibody-associated vasculitis: a study of 439 cases in a single Chinese center. 2014, 66, 1920-6	44

1004	Short-term rosuvastatin therapy for prevention of contrast-induced acute kidney injury in patients with diabetes and chronic kidney disease. 2014 , 63, 62-70	159
1003	Premature mortality and comorbidities in young-onset diabetes: a 7-year prospective analysis. 2014 , 127, 616-24	80
1002	Serum potassium level is associated with metabolic syndrome: a population-based study. 2014 , 33, 521-7	20
1001	Associations between bone mineral density and subclinical atherosclerosis: a cross-sectional study of a Chinese population. 2014 , 99, 469-77	32
1000	Correlation of Cystatin-C and radionuclidic measurement method of glomerular filtration rate in patients with lung cancer receiving cisplatin treatment. 2014 , 36, 1043-50	10
999	Effect of remote ischemic preconditioning in the elderly patients with coronary artery disease with diabetes mellitus undergoing elective drug-eluting stent implantation. 2014 , 65, 660-6	48
998	Serum cystatin C and cerebral microbleeds in patients with acute cerebral stroke. 2014 , 21, 268-73	12
997	Estimation of GFR in South Asians: a study from the general population in Pakistan. 2014 , 63, 49-58	49
996	LDL cholesterol as a novel risk factor for contrast-induced acute kidney injury in patients undergoing percutaneous coronary intervention. 2014 , 237, 453-9	15
995	Age and gender differences in the relationship between hepatitis C infection and all stages of Chronic kidney disease. 2014 , 21, 706-15	27
994	Apparent diffusion coefficient measurements of bilateral kidneys at 3 T MRI: effects of age, gender, and laterality in healthy adults. 2014 , 69, e491-6	7
993	Renin angiotensin system inhibitors may attenuate low LDL cholesterol-related cancer risk in type 2 diabetes. 2014 , 30, 415-23	4
992	Status of non-steroidal anti-inflammatory drugs use and its association with chronic kidney disease: a cross-sectional survey in China. 2014 , 19, 655-60	13
991	Wave reflections, arterial stiffness, heart rate variability and orthostatic hypotension. 2014 , 37, 1056-61	14
990	Urinary serum- and glucocorticoid-inducible kinase SGK1 reflects renal injury in patients with immunoglobulin A nephropathy. 2014 , 19, 307-17	7
989	Clinical analysis of the relationship between cystatin C and metabolic syndrome in the elderly. 2014 , 33, 411-416	5
988	Outcomes in older adults with stage 5 chronic kidney disease: comparison of peritoneal dialysis and conservative management. 2014 , 69, 308-14	37
987	The association between anti-plasminogen antibodies and disease activity in ANCA-associated vasculitis. 2014 , 53, 300-6	14

986	Different interventions in preventing contrast-induced nephropathy after percutaneous coronary intervention. 2014 , 46, 1801-7	19
985	Urinary kidney injury molecule-1 is related to pathologic involvement in IgA nephropathy with normotension, normal renal function and mild proteinuria. 2014 , 15, 107	9
984	Association between sleep quality and cardiovascular damage in pre-dialysis patients with chronic kidney disease. 2014 , 15, 131	20
983	Prevalence of baseline chronic kidney disease in 2,769 Chinese patients with renal cancer: nephron-sparing treatment is still underutilized. 2014 , 32, 1027-31	2
982	Survival rates and risk factors for mortality in systemic lupus erythematosus patients in a Chinese center. 2014 , 33, 947-53	22
981	Derivation and validation of a risk score for contrast-induced nephropathy after cardiac catheterization in Chinese patients. 2014 , 18, 892-8	29
980	Chronic kidney disease after autologous stem cell transplantation: analysis of a single center experience. 2014 , 46, 1403-10	1
979	Association between metabolic syndrome and osteoporotic fracture in middle-aged and elderly Chinese peoples. 2014 , 70, 1297-303	14
978	The revised Lund-Malm GFR estimating equation outperforms MDRD and CKD-EPI across GFR, age and BMI intervals in a large Swedish population. 2014 , 52, 815-24	94
977	Low-grade albuminuria is associated with peripheral artery disease in Chinese diabetic patients. 2014 , 232, 285-8	15
976	Pattern and risk factors of intravesical recurrence after nephroureterectomy for upper tract urothelial carcinoma: a large Chinese center experience. 2014 , 113, 820-7	25
975	Additive effect of aldose reductase Z-4 microsatellite polymorphism and glycaemic control on cataract development in type 2 diabetes. 2014 , 28, 147-51	7
974	Comparison of associations of outcomes after stroke with estimated GFR using Chinese modifications of the MDRD study and CKD-EPI creatinine equations: results from the China National Stroke Registry. 2014 , 63, 59-67	26
973	Association between waist-to-height ratio and chronic kidney disease in the Taiwanese population. 2014 , 44, 645-52	5
972	Elevated soluble VEGF receptor sFlt-1 correlates with endothelial injury in IgA nephropathy. 2014 , 9, e101779	25
971	Original article. Cost-effectiveness of annual microalbuminuria screening in Thai diabetics. 2014 , 8, 371-379	6
970	Asymmetric dimethylarginine and long-term adverse cardiovascular events in patients with type 2 diabetes: relation with the glycemic control. 2014 , 13, 156	23
969	Novel application of stereotactic ablative radiotherapy using CyberKnife for early-stage renal cell carcinoma in patients with pre-existing chronic kidney disease: Initial clinical experiences. 2014 , 8, 355-360	17

968	Validation of methods to control for immortal time bias in a pharmacoepidemiologic analysis of renin-angiotensin system inhibitors in type 2 diabetes. 2014 , 24, 267-73	17
967	Urinary interleukin-18 as an early indicator to predict contrast-induced nephropathy in patients undergoing percutaneous coronary intervention. 2014 , 8, 1263-1266	23
966	An integrative study of the genetic, social and environmental determinants of chronic kidney disease characterized by tubulointerstitial damages in the North Central Region of Sri Lanka. 2014 , 56, 28-38	73
965	Association of N-terminal pro-B-type natriuretic peptide with contrast-induced nephropathy and long-term outcomes in patients with chronic kidney disease and relative preserved left ventricular function. 2015 , 94, e358	8
964	Efficacy of Shenzhuo formula on diabetic kidney disease: a retrospective study. 2015 , 35, 528-36	9
963	Lower Barthel Index Is Associated with Higher Risk of Hospitalization-Requiring Pneumonia in Long-Term Care Facilities. 2015 , 236, 281-8	8
962	Predictive role of preoperative hydronephrosis on poor pathological outcomes and prognosis in upper tract urothelial carcinoma patients: Experience from a nationwide high-volume center in China. 2015 , 10, 3113-3122	9
961	Antihypertensive treatment improves left ventricular diastolic function in patients with chronic kidney disease. 2015 , 9, 1702-1708	
960	Association of elevated glycosylated hemoglobin A1c with hyperfiltration in a middle-aged and elderly Chinese population with prediabetes or newly diagnosed diabetes: a cross-sectional study. 2015 , 15, 47	9
959	Thyroid function modifies the association between ratio of triglyceride to high-density lipoprotein cholesterol and renal function: a multicenter cross-sectional study. 2015 , 5, 11052	
958	Common Carotid Artery Stiffness Is Associated with Left Ventricular Structure and Function and Predicts First Hospitalization for Acute Heart Failure. 2014 , 2, 18-28	6
957	Use of Dual-Source Computed Tomography to Evaluate Renal Cortical Perfusion in Patients With Essential Hypertension Without Diabetes: Preliminary Results. 2015 , 39, 473-8	
956	Kidneys from Older Living Donors Provide Excellent Short and Intermediate Outcomes--A Single China Center's Experience. 2015 , 99, e81-8	9
955	Discordant associations of lipid parameters with albuminuria and chronic kidney disease: a population-based study. 2015 , 14, 152	14
954	Plasma complement factor H is associated with disease activity of patients with ANCA-associated vasculitis. 2015 , 17, 129	29
953	Impact of the Serum Uric Acid Level on Subclinical Atherosclerosis in Middle-aged and Elderly Chinese. 2015 , 22, 823-32	17
952	Association of visceral and total body fat with hypertension and prehypertension in a middle-aged Chinese population. 2015 , 33, 1555-62	35
951	Clinical Features and Outcomes in Patients With Membranous Nephropathy and Crescent Formation. 2015 , 94, e2294	8

950	Prevalence and risk factors associated with chronic kidney disease in a Zhuang ethnic minority area in China. 2015 , 20, 807-13	3
949	The Clinicopathologic Spectrum of Rapidly Progressive Glomerulonephritis Based on Glomerular Immune Deposition and Antineutrophil Cytoplasmic Antibody. 2015 , 23, 704-10	4
948	Validation of the Fatty Liver Index for Nonalcoholic Fatty Liver Disease in Middle-Aged and Elderly Chinese. 2015 , 94, e1682	81
947	Prevalence, awareness and treatment of chronic kidney disease among middle-aged and elderly: The China Health and Retirement Longitudinal Study. 2015 , 20, 474-84	27
946	The predictive capacity and additional prognostic power of N-terminal pro-B-type natriuretic peptide in Chinese elderly with chronic heart failure. 2015 , 10, 359-65	6
945	The Association between Glomerular Hyperfiltration and Left Ventricular Structure and Function in Patients with Primary Aldosteronism. 2015 , 12, 369-77	5
944	Body mass index (BMI) is associated with microalbuminuria in Chinese hypertensive patients. 2015 , 12, 1998-2008	10
943	Hyperhomocysteinemia is an independent predictor of long-term clinical outcomes in Chinese octogenarians with acute coronary syndrome. 2015 , 10, 1467-74	15
942	Role of immunosuppressive therapy and predictors of therapeutic effectiveness and renal outcome in IgA nephropathy with proteinuria. 2015 , 11, 332-9	2
941	Combined biomarker analysis for risk of acute kidney injury in patients with ST-segment elevation myocardial infarction. 2015 , 10, e0125282	28
940	Individualizing pharmacotherapy in patients with renal impairment: the validity of the Modification of Diet in Renal Disease formula in specific patient populations with a glomerular filtration rate below 60 ml/min. A systematic review. 2015 , 10, e0116403	9
939	Nighttime Systolic Blood-Pressure Load Is Correlated with Target-Organ Damage Independent of Ambulatory Blood-Pressure Level in Patients with Non-Diabetic Chronic Kidney Disease. 2015 , 10, e0131546	16
938	Renal Function in Chinese HIV-Positive Individuals following Initiation of Antiretroviral Therapy. 2015 , 10, e0135462	4
937	Serum Cystatin C Reflects Angiographic Coronary Collateralization in Stable Coronary Artery Disease Patients with Chronic Total Occlusion. 2015 , 10, e0137253	9
936	Genetic Variation in miR-146a Is Not Associated with Susceptibility to IgA Nephropathy in Adults from a Chinese Han Population. 2015 , 10, e0139554	8
935	Hypoglycaemia, Abnormal Lipids, and Cardiovascular Disease among Chinese with Type 2 Diabetes. 2015 , 2015, 862896	4
934	Chaperonin-containing t-complex protein-1 subunit is a possible biomarker for the phase of glomerular hyperfiltration of diabetic nephropathy. 2015 , 2015, 548101	5
933	Effects of Long-Term Statin Therapy in Coronary Artery Disease Patients with or without Chronic Kidney Disease. 2015 , 2015, 252564	4

932	Contrast-induced nephropathy following chronic total occlusion percutaneous coronary intervention in patients with chronic kidney disease. 2015 , 25, 2274-81	10
931	Relationship between serum irisin levels and urinary albumin excretion in patients with type 2 diabetes. 2015 , 29, 384-9	25
930	Uric Acid is independently associated with diabetic kidney disease: a cross-sectional study in a Chinese population. 2015 , 10, e0129797	35
929	Clinicopathologic characteristics and outcomes of renal thrombotic microangiopathy in anti-neutrophil cytoplasmic autoantibody-associated glomerulonephritis. 2015 , 10, 750-8	19
928	Nocturnal Hypertension Correlates Better With Target Organ Damage in Patients With Chronic Kidney Disease than a Nondipping Pattern. 2015 , 17, 792-801	21
927	Novel risk scoring for pre-procedural prediction of contrast-induced nephropathy and poor long-term outcomes among patients with chronic total occlusion undergoing percutaneous coronary intervention. 2015 , 17, C34-C41	6
926	High prevalence of isolated nocturnal hypertension in Chinese patients with chronic kidney disease. 2015 , 4, e002025	23
925	Borderline ankle-brachial index is associated with increased prevalence of micro- and macrovascular complications in type 2 diabetes: A cross-sectional analysis of 12,772 patients from the Joint Asia Diabetes Evaluation Program. 2015 , 12, 334-41	15
924	Retinal Vessel Diameter and Chronic Kidney Disease in Rural China: A Cross-Sectional Study. 2015 , 94, e2076	10
923	Serum Fetuin-A Associated With Fatty Liver Index, Early Indicator of Nonalcoholic Fatty Liver Disease: A Strobe-Compliant Article. 2015 , 94, e1517	9
922	Multimarker Analysis for New Biomarkers in Relation to Central Arterial Stiffness and Hemodynamics in a Chinese Community-Dwelling Population. 2015 , 66, 950-6	11
921	Elevated serum uric acid after injury correlates with the early acute kidney in severe burns. 2015 , 41, 1724-1731	12
920	Influence of the Interval Between Renal Computed Tomographic Angiography and Kidney Transplantation on Allograft Function: A Prospective, Randomized, Controlled Trial. 2015 , 47, 2822-6	
919	Clinical Association of Cystatin C and Atrial Fibrillation in Chinese Elderly. 2015 , 9, 146-150	1
918	Hai-yan Wang, MD, 1937-2014. 2015 , 65, A16-7	1
917	Disparate assessment of clinic blood pressure and ambulatory blood pressure in differently aged patients with chronic kidney disease. 2015 , 183, 54-62	9
916	[Exploration of renal glomerular function: estimation of glomerular filtration rate]. 2015 , 11, 54-67	0
915	Cumulative effects of variants identified by genome-wide association studies in IgA nephropathy. 2014 , 4, 4904	13

914	Novel urinary biomarkers in pre-diabetic nephropathy. 2015 , 19, 895-900	36
913	Severe pneumonia in mycophenolate mofetil combined with low-dose corticosteroids-treated patients with immunoglobulin A nephropathy. 2015 , 31, 42-6	9
912	The association of leptin and homocysteine with renal function impairment in a population of Taiwanese adults. 2015 , 34, 943-50	9
911	Ethnic disparity in prevalence of diabetic kidney disease in an Asian primary healthcare cluster. 2015 , 20, 216-23	14
910	Association between body mass index and cause-specific mortality as well as hospitalization in frail Chinese older adults. 2015 , 15, 72-9	6
909	Evaluation of renal function in the elderly, not as easy as it seems: a review. 2015 , 27, 397-401	7
908	Clinical characteristics of IgA nephropathy associated with low complement 4 levels. 2015 , 37, 424-32	7
907	The Association between Serum Uric Acid Levels and the Prevalence of Vulnerable Atherosclerotic Carotid Plaque: A Cross-sectional Study. 2015 , 5, 10003	26
906	Noninvasive renal denervation for resistant hypertension using high-intensity focused ultrasound. 2015 , 66, e22-5	15
905	The cutoffs and performance of glycated hemoglobin for diagnosing diabetes and prediabetes in a young and middle-aged population and in an elderly population. 2015 , 109, 238-45	11
904	Impact of body composition on estimated glomerular filtration rate in relatively healthy adults in Taiwan. 2015 , 69, 34-9	6
903	Circulating fibroblast growth factor 21 levels predict progressive kidney disease in subjects with type 2 diabetes and normoalbuminuria. 2015 , 100, 1368-75	64
902	Visceral adiposity index, hypertriglyceridemic waist phenotype and chronic kidney disease in a southern Chinese population: a cross-sectional study. 2015 , 47, 1387-96	13
901	Pharmacokinetics, Pharmacodynamics, and Safety of Single-Dose Canagliflozin in Healthy Chinese Subjects. 2015 , 37, 1483-1492.e1	13
900	Human neutrophil peptide 1-3, a component of the neutrophil extracellular trap, as a potential biomarker of lupus nephritis. 2015 , 18, 533-40	11
899	Neuropeptide Y levels are associated with nutritional status and cardiovascular events in adults with chronic kidney disease. 2015 , 69, 717-21	2
898	A Study of the Applicability of GFR Evaluation Equations for an Elderly Chinese Population. 2015 , 19, 693-701	5
897	Association of serum irisin and body composition with chronic kidney disease in obese Chinese adults: a cross-sectional study. 2015 , 16, 16	19

896	Contralateral upper tract urothelial carcinoma after nephroureterectomy: the predictive role of DNA methylation. 2015 , 34, 5	13
895	Presence of Concomitant Non-muscle-invasive Bladder Cancer in Chinese Patients with Upper Tract Urothelial Carcinoma: Risk Factors, Characteristics, and Predictive Value. 2015 , 22, 2789-98	12
894	Associations between biomarkers of renal function with cerebral microbleeds in hypertensive patients. 2015 , 28, 739-45	17
893	Non-HDL-cholesterol to HDL-cholesterol ratio as an independent risk factor for the development of chronic kidney disease. 2015 , 25, 582-7	22
892	Decreased urine uric acid excretion is an independent risk factor for chronic kidney disease but not for carotid atherosclerosis in hospital-based patients with type 2 diabetes: a cross-sectional study. 2015 , 14, 36	13
891	Lipoprotein (a) is a risk factor for coronary artery disease in Chinese Han ethnic population modified by some traditional risk factors: A cross-sectional study of 3462 cases and 6125 controls. 2015 , 451, 278-86	23
890	Relationship between β cell function, metabolic control, and microvascular complications in type 2 diabetes mellitus. 2015 , 17, 29-34	12
889	Chronic kidney disease among overweight and obesity with and without metabolic syndrome in an urban Chinese cohort. 2015 , 16, 85	39
888	Insulin resistance is not independently associated with chronic kidney disease in Chinese population: A population-based cross-sectional study. 2015 , 448, 232-7	7
887	Urinary vimentin mRNA as a potential novel biomarker of renal fibrosis. 2015 , 309, F514-22	24
886	Renal function-adjusted safe contrast volume to prevent contrast-induced nephropathy and poor long-term outcomes in patients with chronic total occlusions undergoing cardiac catheterization. 2015 , 17, C17-C25	2
885	Thrombolysis in chinese ischemic stroke patients with renal dysfunction. 2015 , 3, 101-6	5
884	Associations between pentraxin 3 and severity of coronary artery disease. 2015 , 5, e007123	16
883	Association between diabetic retinopathy and subclinical atherosclerosis in China: Results from a community-based study. 2015 , 12, 366-72	11
882	Perceived stress status and sympathetic nervous system activation in young male patients with coronary artery disease in China. 2015 , 26, 726-30	9
881	Decreased urine uric acid excretion is associated with diabetic retinopathy but not with lower limb atherosclerosis in hospitalized patients with type 2 diabetes. 2015 , 242, 13-8	11
880	Intermittent high-volume hemofiltration promotes remission in steroid-resistant idiopathic nephrotic syndrome. 2015 , 37, 966-73	3
879	Glycated haemoglobin A1c is associated with low-grade albuminuria in Chinese adults. 2015 , 5, e007429	8

878	Tissue transcriptome-driven identification of epidermal growth factor as a chronic kidney disease biomarker. 2015 , 7, 316ra193		202
877	Risk factors for mortality in Chinese patients on continuous ambulatory peritoneal dialysis. 2015 , 35, 199-205		15
876	Depression in Chinese patients with type 2 diabetes: associations with hyperglycemia, hypoglycemia, and poor treatment adherence. 2015 , 7, 800-8		70
875	Incidence, characteristics, treatment strategies, and oncologic outcomes of synchronous bilateral upper tract urothelial carcinoma in the Chinese population. 2015 , 33, 66.e1-11		20
874	Variants in Complement Factor H and Complement Factor H-Related Protein Genes, CFHR3 and CFHR1, Affect Complement Activation in IgA Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2015 , 26, 1195-204	12.7	90
873	Serum creatinine levels and risk of metabolic syndrome in a middle-aged and older Chinese population. 2015 , 440, 177-82		8
872	Relationship between TG/HDL-C ratio and metabolic syndrome risk factors with chronic kidney disease in healthy adult population. 2015 , 34, 874-80		25
871	Chronic Kidney Disease in the Elderly □Who Has it? Who Does One Treat and How are They to be Treated?. 2015 , 843-853		
870	Whether Warfarin Therapy is Associated with Damage on Renal Function in Chinese Patients with Nonvalvular Atrial Fibrillation. 2016 , 129, 1135-9		1
869	Mineral and Bone Disorder and Its Association with Cardiovascular Parameters in Chinese Patients with Chronic Kidney Disease. 2016 , 129, 2275-80		6
868	Protective effect of tartary buckwheat on renal function in type 2 diabetics: a randomized controlled trial. 2016 , 12, 1721-1727		10
867	"One-Time" versus Staged Multivessel Intervention in Intermediate to Very High-Risk Patients with Non-ST-Segment Elevation Acute Coronary Syndromes. 2016 , 46, 774-783		3
866	The prognostic value of the plasma N-terminal pro-brain natriuretic peptide level on all-cause death and major cardiovascular events in a community-based population. 2016 , 11, 245-53		14
865	Association of hemoglobin with ankle-brachial index in general population. 2016 , 71, 375-80		1
864	Short-term rosuvastatin therapy prevents contrast-induced acute kidney injury in female patients with diabetes and chronic kidney disease: a subgroup analysis of the TRACK-D study. 2016 , 8, 1000-6		4
863	Comparison of CCR, Cockcroft-Gault and Mdrd Formula for the Estimation of Glomerular Filtration Rate. 2016 , 33, 207-212		
862	Analyzing Dynamic Changes of Laboratory Indexes in Patients with Acute Heart Failure Based on Retrospective Study. 2016 , 2016, 7496061		
861	Correlation of visceral adiposity index with chronic kidney disease in the People's Republic of China: to rediscover the new clinical potential of an old indicator for visceral obesity. 2016 , 12, 489-94		8

860	Visceral Adiposity Index and Lipid Accumulation Product Index: Two Alternate Body Indices to Identify Chronic Kidney Disease among the Rural Population in Northeast China. 2016 , 13,	33
859	The Association of Hypertriglyceridemic Waist Phenotype with Chronic Kidney Disease and Its Sex Difference: A Cross-Sectional Study in an Urban Chinese Elderly Population. 2016 , 13,	6
858	High Prevalence of Hyperhomocysteinemia and Its Association with Target Organ Damage in Chinese Patients with Chronic Kidney Disease. 2016 , 8,	17
857	All Components of Metabolic Syndrome Are Associated with Microalbuminuria in a Chinese Population. 2016 , 11, e0157303	9
856	The Relationship of Diabetes and Smoking Status to Hepatocellular Carcinoma Mortality. 2016 , 95, e2699	12
855	Effects of glomerular filtration rate estimating equations derived from different reference methods on staging and long term mortality risks of chronic kidney disease in a Southeast Asian cohort. 2016 , 21, 678-86	1
854	Comparison of Acarbose and Metformin on Albumin Excretion in Patients With Newly Diagnosed Type 2 Diabetes: A Randomized Controlled Trial. 2016 , 95, e3247	12
853	Podocyte Detachment Is Associated with Renal Prognosis in ANCA-Associated Glomerulonephritis: A Retrospective Cohort Study. 2016 , 95, e3294	8
852	Estimating glomerular filtration rate by serum creatinine or/and cystatin C equations: An analysis of multi-centre Chinese subjects. 2016 , 21, 372-8	17
851	Prevalence, determinants, and clinical significance of masked hypertension and white-coat hypertension in patients with chronic kidney disease. 2016 , 21, 841-50	12
850	A retrospective analysis of laparoscopic partial nephrectomy with segmental renal artery clamping and factors that predict postoperative renal function. 2016 , 118, 610-7	8
849	Treatment strategies for upper tract urothelial carcinoma (UTUC) of a solitary kidney: a single-institutional analysis of 61 cases. 2016 , 48, 1601-8	1
848	Association of NOS2 and NOS3 gene polymorphisms with susceptibility to type 2 diabetes mellitus and diabetic nephropathy in the Chinese Han population. 2016 , 68, 516-25	19
847	Kidney function is associated with severity of white matter hyperintensity in patients with acute ischemic stroke/TIA. 2016 , 16, 193	12
846	Nickel exposure and prevalent albuminuria and α -microglobulinuria: evidence from a population-based study. 2016 , 70, 437-43	8
845	Characteristics and treatment outcomes of pan-urothelial cell carcinoma: a descriptive analysis of 45 patients. 2015 , 5, 18014	5
844	Low serum magnesium levels are associated with impaired peripheral nerve function in type 2 diabetic patients. 2016 , 6, 32623	12
843	A meta-analysis of renal outcomes in living kidney donors. 2016 , 95, e3847	10

842	The Kinetics of Circulating Monocyte Subsets and Monocyte-Platelet Aggregates in the Acute Phase of ST-Elevation Myocardial Infarction: Associations with 2-Year Cardiovascular Events. 2016 , 95, e3466	30
841	Outcomes in elderly patients with end-stage renal disease: Comparison of renal replacement therapy and conservative management. 2016 , 19, 42-56	9
840	N-terminal pro-brain natriuretic peptide improves the C-ACS risk score prediction of clinical outcomes in patients with ST-elevation myocardial infarction. 2016 , 16, 255	5
839	Retinal Information is Independently Associated with Cardiovascular Disease in Patients with Type 2 diabetes. 2016 , 6, 19053	11
838	Association between serum bilirubin levels and decline in estimated glomerular filtration rate among patients with type 2 diabetes. 2016 , 30, 1255-60	14
837	Establish Albumin-creatinine Ratio Reference Value of Adults in the Rural Area of Hebei Province. 2016 , 31, 23-30	
836	Association between serum uric acid related genetic loci and diabetic kidney disease in the Chinese type 2 diabetes patients. 2016 , 30, 798-802	5
835	Combined body mass index with high-sensitivity C-reactive protein as independent predictors for chronic kidney disease in a relatively healthy population in Taiwan. 2016 , 70, 766-71	1
834	Hyponatremia and Worsening Sodium Levels Are Associated With Long-Term Outcome in Patients Hospitalized for Acute Heart Failure. 2016 , 5, e002668	28
833	Regulation of IgA Class Switch Recombination in Immunoglobulin A Nephropathy: Retinoic Acid Signaling and BATF. 2016 , 43, 179-94	8
832	Effectiveness of sulodexide might be associated with inhibition of complement system in hepatitis B virus-associated membranous nephropathy: An inspiration from a pilot trial. 2016 , 32, 96-104	4
831	[Effect of Astragali and Angelica particle on proteinuria in Chinese patients with primary glomerulonephritis]. 2016 , 36, 299-306	8
830	The relationship between albuminuria and poor clinical outcomes in patients with infective endocarditis. 2016 , 462, 28-32	1
829	Long-term outcome of patients with very small coronary artery disease: A comparison of drug-eluting and bare metal stents. 2016 , 79, 642-648	4
828	Advance care planning for 600 Chinese patients with end-stage renal disease. 2016 , 19, 19-27	5
827	Asymmetric dimethylarginine predicts the risk of contrast-induced acute kidney injury in patients undergoing cardiac catheterization. 2016 , 254, 161-166	8
826	Dose-response relationship between serum uric acid levels and risk of incident coronary heart disease in the Dongfeng-Tongji Cohort. 2016 , 224, 299-304	20
825	Standards of care for type 2 diabetes in China. 2016 , 32, 442-58	185

824	Glomerular filtration rate measured by (99m) Tc-DTPA renal dynamic imaging is significantly lower than that estimated by the CKD-EPI equation in horseshoe kidney patients. 2016 , 21, 499-505	9
823	Uric acid, renal function and risk of hypoglycaemia in Chinese type 2 diabetes patients. 2016 , 32, 875-882	6
822	Understanding the controversy surrounding the correlation between fibrinogen level and prognosis of coronary artery disease-The role of the subtypes of coronary artery disease. 2016 , 222, 968-972	2
821	Circulating periostin in relation to insulin resistance and nonalcoholic fatty liver disease among overweight and obese subjects. 2016 , 6, 37886	18
820	Determinants and Prognostic Impact of Hyperuricemia in Hospitalization for Acute Heart Failure. 2016 , 80, 404-10	19
819	Biopsy-proven kidney diseases in the elderly: clinical characteristics, renal histopathological spectrum and prognostic factors. 2016 , 44, 1092-1102	9
818	Tacrolimus combined with low-dose corticosteroids is an effective and safe therapeutic option for refractory IgA nephropathy. 2016 , 12, 1934-1938	5
817	Does N-terminal pro-brain natriuretic peptide add prognostic value to the Mehran risk score for contrast-induced nephropathy and long-term outcomes after primary percutaneous coronary intervention?. 2016 , 48, 1675-82	10
816	Hyperuricemia is associated with increased hospitalization risk and healthcare costs: Evidence from an administrative database in Italy. 2016 , 26, 951-61	9
815	Relation between admission plasma fibrinogen levels and mortality in Chinese patients with coronary artery disease. 2016 , 6, 30506	6
814	Excess Pressure Integral Predicts Long-Term All-Cause Mortality in Stable Heart Failure Patients. 2017 , 30, 271-278	10
813	Plasma selenium levels and nonalcoholic fatty liver disease in Chinese adults: a cross-sectional analysis. 2016 , 6, 37288	32
812	Community-based study on elderly CKD subjects and the associated risk factors. 2016 , 38, 1672-1676	3
811	Parity and serum lipid levels: a cross-sectional study in chinese female adults. 2016 , 6, 33831	6
810	Areca Nut Chewing and an Impaired Estimated Glomerular Filtration Rate as Significant Risk Factors for Non-Muscle-Invasive Bladder Cancer Recurrence. 2016 , 6, 29466	6
809	Levels of Serum Phosphorus and Cardiovascular Surrogate Markers. 2016 , 23, 95-104	8
808	Association of self-reported recurrent mild hypoglycemia with incident cardiovascular disease and all-cause mortality in patients with type 2 diabetes: Prospective analysis of the Joint Asia Diabetes Evaluation Registry. 2016 , 95, e5183	11
807	Comparison of Different Risk Scores for Predicting Contrast Induced Nephropathy and Outcomes After Primary Percutaneous Coronary Intervention in Patients With ST Elevation Myocardial Infarction. 2016 , 117, 1896-903	17

806	Clinical Significance of IgM and C3 Glomerular Deposition in Primary Focal Segmental Glomerulosclerosis. 2016 , 11, 1582-9	23
805	Anaphthalimide-based fluorescent probe for mercapto-containing compounds. 2016 , 27, 1776-1780	20
804	Upper Tract Urothelial Carcinomas Accompanied by Previous or Synchronous Nonmuscle-Invasive Bladder Cancer and Preoperative Hydronephrosis Might Have Worse Oncologic Outcomes After Radical Nephroureterectomy. 2016 , 14, e469-e477	9
803	Type 2 Diabetes, Diabetes Genetic Score and Risk of Decreased Renal Function and Albuminuria: A Mendelian Randomization Study. 2016 , 6, 162-170	18
802	Metabolic syndrome and osteoporotic fracture: a population-based study in China. 2016 , 16, 27	15
801	Association Between Body Mass Index and Mildly Decreased Estimated Glomerular Filtration Rate in Chinese Adults With Early Chronic Kidney Disease. 2016 , 26, 367-372	5
800	Association between FNDC5 genetic variants and proliferative diabetic retinopathy in a Chinese population. 2016 , 43, 580-2	3
799	Gender, diabetes education, and psychosocial factors are associated with persistent poor glycemic control in patients with type 2 diabetes in the Joint Asia Diabetes Evaluation (JADE) program. 2016 , 8, 109-19	13
798	Estimating GFR prior to contrast medium examinations--what the radiologist needs to know!. 2016 , 26, 425-35	4
797	Metabolic syndrome does not always play a critical role in decreased GFR. 2016 , 38, 383-9	2
796	Rare Variants in the Complement Factor H-Related Protein 5 Gene Contribute to Genetic Susceptibility to IgA Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2016 , 27, 2894-905	43
795	The efficacy of tonsillectomy on clinical remission and relapse in patients with IgA nephropathy: a randomized controlled trial. 2016 , 38, 242-8	17
794	Serum Cystatin C and the Risk of Coronary Heart Disease in Ethnic Chinese Patients With Normal Renal Function. 2016 , 47, 13-9	4
793	Association of Serum Fetuin-A Levels With the Risk of Albuminuria in Middle-Aged and Elderly Chinese. 2016 , 101, 1235-42	4
792	Association of Endothelial and Mild Renal Dysfunction With the Severity of Left Ventricular Hypertrophy in Hypertensive Patients. 2016 , 29, 501-8	9
791	Impact of Chronic Total Occlusion in a Noninfarct-related Artery on Clinical Outcomes in Patients With Acute ST-elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. 2016 , 95, e2441	10
790	Increased APRIL Expression Induces IgA1 Aberrant Glycosylation in IgA Nephropathy. 2016 , 95, e3099	41
789	High-sensitivity cardiac troponin T in geriatric inpatients. 2016 , 65, 111-5	17

788	Kidney Function and Cognitive Impairment in People Aged 80 Years and Over with Untreated Hypertension: A Cross-Sectional Survey. 2016 , 41, 70-7	5
787	IgG4 anti-phospholipase A2 receptor might activate lectin and alternative complement pathway meanwhile in idiopathic membranous nephropathy: an inspiration from a cross-sectional study. 2016 , 64, 919-30	29
786	Modifying Effect of Body Mass Index on Survival in Elderly Type 2 Diabetic Patients: Hong Kong Diabetes Registry. 2016 , 17, 276.e15-22	3
785	Plasma NT-proBNP is independently associated with albuminuria in type 2 diabetes. 2016 , 30, 669-74	4
784	Bibliometric and visualized analysis of emergy research. 2016 , 90, 285-293	85
783	Longitudinal relationship between chronic kidney disease and distribution of cerebral microbleeds in patients with ischemic stroke. 2016 , 362, 1-6	20
782	Correlation of Preoperative Renal Insufficiency With Mortality and Morbidity After Aortic Valve Replacement: A Propensity Score Matching Analysis. 2016 , 95, e2576	9
781	IntelliHealth: A medical decision support application using a novel weighted multi-layer classifier ensemble framework. 2016 , 59, 185-200	72
780	Cardiac autonomic neuropathy risk estimated by sudomotor function and arterial stiffness in Chinese subjects. 2016 , 30, 720-725	10
779	Gender-specific association between metabolic syndrome and decreased glomerular filtration rate in elderly population. 2016 , 48, 389-97	4
778	Elevated plasma tumor necrosis factor- α receptor 2 and resistin are associated with increased incidence of kidney function decline in Chinese adults. 2016 , 52, 541-9	10
777	Chronic kidney disease and cognitive impairment among the very old in China. 2016 , 28, 475-82	9
776	Is low b-factors-based apparent diffusion coefficient helpful in assessing renal dysfunction?. 2016 , 121, 6-11	3
775	Biomarkers of Renal Function in Type 2 Diabetic Patients with Cognitive Impairment. 2016 , 610, 19-23	5
774	Assessment of renal dysfunction with diffusion-weighted imaging: comparing intra-voxel incoherent motion (IVIM) with a mono-exponential model. 2016 , 57, 507-12	16
773	Screening diabetic patients for non-alcoholic fatty liver disease with controlled attenuation parameter and liver stiffness measurements: a prospective cohort study. 2016 , 65, 1359-68	271
772	Gender impact on the correlations between subclinical thyroid dysfunction and hyperuricemia in Chinese. 2016 , 35, 143-9	26
771	Renal Dysfunction, CHADS Score, and Adherence to the Anticoagulant Treatment in Nonvalvular Atrial Fibrillation. 2017 , 23, 248-254	6

770	Preliminary effects of renal denervation with saline irrigated catheter on cardiac systolic function in patients with heart failure: A Prospective, Randomized, Controlled, Pilot Study. 2017 , 89, E153-E161	28
769	Performance of the creatinine and cystatin C-based equations for estimation of GFR in Chinese patients with chronic kidney disease. 2017 , 21, 236-246	10
768	Contrast Volume to Cystatin C-Based Glomerular Filtration Ratio Predicts the Risk of Contrast-Induced Nephropathy After Cardiac Catheterization. 2017 , 68, 207-215	1
767	Interleukin 18 -607 A/C Gene Polymorphism is Associated With Susceptibility to IgA Nephropathy in a Chinese Han Population. 2017 , 25, 725-730	5
766	Increased prevalence and risk of non-alcoholic fatty liver disease in overweight and obese patients with Type 2 diabetes in South China. 2017 , 34, 505-513	18
765	Poor sleep quality is responsible for the nondipper pattern in hypertensive but not in normotensive chronic kidney disease patients. 2017 , 22, 690-698	6
764	The impact of admission neutrophil-to-platelet ratio on in-hospital and long-term mortality in patients with infective endocarditis. 2017 , 55, 899-906	8
763	Which creatinine-based estimated glomerular filtration rate equation best predicts all-cause mortality in Chinese subjects with type 2 diabetes?. 2017 , 126, 25-29	3
762	Effects of individualized exercise program on physical function, psychological dimensions, and health-related quality of life in patients with chronic kidney disease: A randomized controlled trial in China. 2017 , 23, e12519	20
761	Canada Acute Coronary Syndrome Score: A Preprocedural Risk Score for Contrast-Induced Nephropathy After Primary Percutaneous Coronary Intervention. 2017 , 68, 782-789	7
760	Association between hyperhomocysteinemia and stroke with atherosclerosis and small artery occlusion depends on homocysteine metabolism-related vitamin levels in Chinese patients with normal renal function. 2017 , 32, 859-865	15
759	Prognostic value of N-terminal prohormone brain natriuretic peptide for in-hospital and long-term outcomes in patients with infective endocarditis. 2017 , 24, 676-684	8
758	Accuracy of glomerular filtration rate equations for chronic kidney disease patients at the G3a stage: a single-center cross-sectional study. 2017 , 10, 107	2
757	Diagnostic value of renal perfusion in patients with chronic kidney disease using 3D arterial spin labeling. 2017 , 46, 589-594	24
756	Aging Kidney: Vascular Characteristics and Assessment. 2017 , 113-127	
755	Recent trends in the prevalence of chronic kidney disease: not the same old song. 2017 , 26, 187-196	25
754	Diagnostic accuracy of different methods of early detection of chronic kidney disease. 2017 , 25, 401-407	
753	Association between estimated glomerular filtration rate and outcomes in patients with diabetic foot ulcers: a 3-year follow-up study. 2017 , 177, 41-50	9

752	Performance of AHEAD Score in an Asian Cohort of Acute Heart Failure With Either Preserved or Reduced Left Ventricular Systolic Function. 2017 , 6,	16
751	Prevalence and correlates of left ventricular diastolic dysfunction and heart failure with preserved ejection fraction in elderly community residents. 2017 , 227, 820-825	3
750	Serum cystatin c is not superior to serum creatinine for early diagnosis of contrast-induced nephropathy in patients who underwent angiography. 2017 , 31,	7
749	Urine Proteome Specific for Eye Damage Can Predict Kidney Damage in Patients With Type 2 Diabetes: A Case-Control and a 5.3-Year Prospective Cohort Study. 2017 , 40, 253-260	24
748	A lower baseline glomerular filtration rate predicts high mortality and newly cerebrovascular accidents in acute ischemic stroke patients. 2017 , 96, e5868	3
747	The efficacy and safety of tacrolimus monotherapy in adult-onset nephrotic syndrome caused by idiopathic membranous nephropathy. 2017 , 39, 512-518	19
746	Serum glycated albumin, glycated hemoglobin, and arterial stiffness in a general Chinese population. 2017 , 468, 33-38	5
745	Prevalence and determinants of left ventricular geometric abnormalities in hypertensive patients: A study based on the updated classification system of left ventricular geometry. 2017 , 58, 124-132	2
744	Ascending aortic distensibility and target organ damage in primary hypertension without diabetes. 2017 , 33, 1245-1251	6
743	Cardiac Management in the Frail Elderly Patient and the Oldest Old. 2017 ,	1
742	Association of estimated glomerular filtration rate and urine albumin-to-creatinine ratio with incidence of cardiovascular diseases and mortality in chinese patients with type 2 diabetes mellitus - a population-based retrospective cohort study. 2017 , 18, 47	13
741	Association between non-invasively diagnosed hepatic steatosis and chronic kidney disease in Chinese adults on their health check-up. 2017 , 18, 229-236	11
740	Erythrocyte fatty acid composition does not influence levels of free, bioavailable, and total 25-hydroxy vitamin D. 2017 , 77, 45-52	4
739	Low testosterone and clinical outcomes in Chinese men with type 2 diabetes mellitus - Hong Kong Diabetes Registry. 2017 , 123, 97-105	15
738	Human Epididymis Protein 4 as a Potential Biomarker of Chronic Kidney Disease in Female Patients With Normal Ovarian Function. 2017 , 48, 238-243	16
737	Effects of hydration combined with Shenfu injection on contrast-induced acute kidney injury in acute coronary syndrome patients undergoing percutaneous coronary intervention. 2017 , 7, 477-481	2
736	Pharmacokinetics and Pharmacodynamics of Meloxicam in East Asian Populations: The Role of Ethnicity on Drug Response. 2017 , 6, 823-832	10
735	Oral N-acetylcysteine for prophylaxis of contrast-induced nephropathy in patients following coronary angioplasty: A meta-analysis. 2017 , 14, 1568-1576	12

734	The influence of living donor SHROOM3 and ABCB1 genetic variants on renal function after kidney transplantation. 2017 , 27, 19-26	6
733	CDKAL1 rs7756992 is associated with diabetic retinopathy in a Chinese population with type 2 diabetes. 2017 , 7, 8812	9
732	Effects of acarbose on cardiovascular and diabetes outcomes in patients with coronary heart disease and impaired glucose tolerance (ACE): a randomised, double-blind, placebo-controlled trial. 2017 , 5, 877-886	164
731	A new scoring model for the prediction of mortality in patients with acute kidney injury. 2017 , 7, 7862	13
730	Arthritis, Other Medical Illnesses and Morale Among Chinese Nonagenarians and Centenarians. 2017 , 11, 100-103	
729	Prevalence of chronic kidney disease markers: Evidence from a three-million married population with fertility desire in rural China. 2017 , 7, 2710	5
728	Serum uric acid as a simple risk factor in patients with rheumatic heart disease undergoing valve replacement surgery. 2017 , 472, 69-74	5
727	Clinical outcomes of IgA nephropathy patients with different proportions of crescents. 2017 , 96, e6190	17
726	Role of Cystatin C and glomerular filtration rate in diagnosis of kidney impairment in hepatic cirrhosis patients. 2017 , 96, e6949	12
725	Circulating Antibodies against Thrombospondin Type-I Domain-Containing 7A in Chinese Patients with Idiopathic Membranous Nephropathy. 2017 , 12, 1642-1651	51
724	Thrombocytopenia as a Preoperative Risk Assessment Tool in Patients With Rheumatic Heart Disease Undergoing Valve Replacement Surgery. 2017 , 6,	4
723	Lipoprotein-associated phospholipase A2 and oxidized low-density lipoprotein in young patients with acute coronary syndrome in China. 2017 , 7, 16092	6
722	Relatives in silent kidney disease screening (RISKS) study: A Chinese cohort study. 2017 , 22 Suppl 4, 35-42	8
721	Prognostic value of hypoalbuminemia for adverse outcomes in patients with rheumatic heart disease undergoing valve replacement surgery. 2017 , 7, 1958	4
720	Estimation of clinical parameters of chronic kidney disease by exhaled breath full-scan mass spectrometry data and iterative PCA with intensity screening algorithm. 2017 , 11, 036007	3
719	Association of high-sensitivity cardiac troponin T with mortality and cardiovascular events in a community-based prospective study in Beijing. 2017 , 7, e013431	9
718	Renal function had an independent relationship with coronary artery calcification in Chinese elderly men. 2017 , 17, 80	1
717	Strategies and cost-effectiveness evaluation of persistent albuminuria screening among high-risk population of chronic kidney disease. 2017 , 18, 135	5

716	Reduced Kidney Function, Albuminuria, and Risks for All-cause and Cardiovascular Mortality in China: A Population-based Cohort Study. 2017 , 18, 188	18
715	C-peptide is independent associated with diabetic peripheral neuropathy: a community-based study. 2017 , 9, 12	11
714	Evaluating the Efficacy, Safety, and Tolerability of Serelaxin When Added to Standard Therapy in Asian Patients With Acute Heart Failure: Design and Rationale of RELAX-AHF-ASIA Trial. 2017 , 23, 63-71	15
713	Ascending Aortic Elasticity and Related Risk Factors Study on Prehypertension Patients. 2017 , 30, 61-66	3
712	Edentulism and associated factors among community-dwelling middle-aged and elderly adults in China. 2017 , 34, 195-207	14
711	Evaluation of glomerular filtration rate by different equations in Chinese elderly with chronic kidney disease. 2017 , 49, 133-141	13
710	Prevalence of Homocysteine-Related Hypertension in Patients With Chronic Kidney Disease. 2017 , 19, 151-160	23
709	Additive prognostic value of left ventricular ejection fraction to the TIMI risk score for in-hospital and long-term mortality in patients with ST segment elevation myocardial infarction. 2017 , 43, 1-6	4
708	Association Between Serum Cystatin C Level and Total Magnetic Resonance Imaging Burden of Cerebral Small Vessel Disease in Patients With Acute Lacunar Stroke. 2017 , 26, 186-191	25
707	Retroperitoneoscopic Partial Nephrectomy for Moderately Complex Ventral Hilar Tumors: Surgical Technique and Trifecta Outcomes from a Single Institution in China. 2017 , 27, 812-817	4
706	Innovative non-invasive model for screening reduced estimated glomerular filtration rate in a working population. 2017 , 22, 892-898	2
705	Risk of glomerular filtration rate decline in patients with hypertrophic cardiomyopathy and obstructive sleep apnoea. 2017 , 7, 17399	6
704	The association between serum uric acid and blood pressure in different age groups in a healthy Chinese cohort. 2017 , 96, e8953	21
703	Hemographic indices are associated with mortality in acute heart failure. 2017 , 7, 17828	8
702	Association of Thyroid-Stimulating Hormone Levels with Microvascular Complications in Type 2 Diabetes Patients. 2017 , 23, 2715-2720	18
701	Evaluation of various equations for estimating renal function in elderly Chinese patients with type 2 diabetes mellitus. 2017 , 12, 1661-1672	7
700	Assessment of chronic kidney disease using skin texture as a key parameter: for South Indian population. 2017 , 4, 223-227	5
699	Soluble Urokinase Plasminogen Activator Receptor and the Risk of Coronary Artery Disease in Young Chinese Patients. 2017 , 2017, 4719403	5

698	Chronic Kidney Dysfunction Can Increase the Risk of Deep Vein Thrombosis after Total Hip and Knee Arthroplasty. 2017 , 2017, 8260487	11
697	Investigating Factors Associated with Depressive Symptoms of Chronic Kidney Diseases in China with Type 2 Diabetes. 2017 , 2017, 1769897	8
696	The Association of a Genetic Variant in with Diabetic Kidney Disease and Diabetic Retinopathy in a Chinese Population. 2017 , 2017, 6542689	3
695	RCHADS and RCHADSVASc schemes improved the performance of CHADS and CHADSVASc scores in death risk stratification of Chinese older patients with atrial fibrillation. 2017 , 12, 1233-1238	9
694	Comparative Performance of Creatinine-Based Estimated Glomerular Filtration Rate Equations in the Malays: A Pilot Study in Tertiary Hospital in Malaysia. 2017 , 2017, 2901581	4
693	Serum irisin levels correlated to peritoneal dialysis adequacy in nondiabetic peritoneal dialysis patients. 2017 , 12, e0176137	3
692	Segmental arterial stiffness in relation to B-type natriuretic peptide with preserved systolic heart function. 2017 , 12, e0183747	2
691	Perioperative body weight change is associated with in-hospital mortality in cardiac surgical patients with postoperative acute kidney injury. 2017 , 12, e0187280	4
690	CKD-EPI creatinine-cystatin C glomerular filtration rate estimation equation seems more suitable for Chinese patients with chronic kidney disease than other equations. 2017 , 18, 226	17
689	Predictive abilities of cardiovascular biomarkers to rapid decline of renal function in Chinese community-dwelling population: a 5-year prospective analysis. 2017 , 18, 331	2
688	Persistent hematuria in patients with antineutrophil cytoplasmic antibody-associated vasculitis during clinical remission: chronic glomerular lesion or low-grade active renal vasculitis?. 2017 , 18, 354	6
687	MDRD vs. CKD-EPI in comparison to Chromium EDTA: a cross sectional study of Malaysian CKD cohort. 2017 , 18, 363	9
686	The impact of intravenous methylprednisolone pulses on renal survival in anti-neutrophil cytoplasmic antibody associated vasculitis with severe renal injury patients: a retrospective study. 2017 , 18, 381	7
685	Association of Serum Prealbumin with Angiographic Severity in Patients with Acute Coronary Syndrome. 2017 , 23, 4041-4049	13
684	Laparoscopic radical cystectomy with novel orthotopic neobladder with bilateral isoperistaltic afferent limbs: initial experience. 2017 , 43, 57-66	6
683	Kidney function and cognitive decline in an oldest-old Chinese population. 2017 , 12, 1049-1054	11
682	Relationships of pancreatic beta-cell function with microalbuminuria and glomerular filtration rate in middle-aged and elderly population without type 2 diabetes mellitus: a Chinese community-based analysis. 2017 , 12, 753-757	2
681	Plasma Metal Concentrations and Incident Coronary Heart Disease in Chinese Adults: The Dongfeng-Tongji Cohort. 2017 , 125, 107007	84

680	Efficacy and Safety of Niaoduoqing Particles for Delaying Moderate-to-severe Renal Dysfunction: A Randomized, Double-blind, Placebo-controlled, Multicenter Clinical Study. 2017 , 130, 2402-2409	11
679	Evaluation of Serological Indicators and Glomerular Filtration Rate Equations in Chinese Cancer Patients. 2017 , 23, 2949-2960	3
678	Associations of multiple plasma metals with incident type 2 diabetes in Chinese adults: The Dongfeng-Tongji Cohort. 2018 , 237, 917-925	43
677	Prediction of Tubulointerstitial Injury in Chronic Kidney Disease Using a Non-Invasive Model: Combination of Renal Sonography and Laboratory Biomarkers. 2018 , 44, 941-948	3
676	Chinese observational prospective study of ageing population with chronic kidney disease (C-OPTION): a study protocol. 2018 , 8, e019457	7
675	Epidemiological characteristics of and risk factors for patients with postoperative acute kidney injury: a multicenter prospective study in 30 Chinese intensive care units. 2018 , 50, 1319-1328	13
674	Effects of remote ischemic preconditioning on contrast induced nephropathy after percutaneous coronary intervention in patients with acute coronary syndrome. 2018 , 97, e9579	10
673	Associations of Plasma Amino Acid and Acylcarnitine Profiles with Incident Reduced Glomerular Filtration Rate. 2018 , 13, 560-568	12
672	Glomerular Filtration Rates in Asians. 2018 , 25, 41-48	21
671	Late restoration of renal function in patients with severe ANCA-associated glomerulonephritis who were dialysis-dependent at presentation. 2018 , 37, 2143-2150	4
670	Identification of chronic kidney disease risk in relatively lean Southern Chinese: the hypertriglyceridemic waist phenotype vs. anthropometric indexes. 2018 , 23, 885-892	4
669	Significant interaction of hypertension and homocysteine on neurological severity in first-ever ischemic stroke patients. 2018 , 12, 534-541	4
668	The significance of the initial symptom in Chinese patients with upper tract urothelial carcinoma: Regular health examination is still underutilized. 2018 , 34, 511-521	3
667	Uncertainty in measurement: A review of the procedures for determining uncertainty in measurement and its use in deriving the biological variation of the estimated glomerular filtration rate. 2018 , 12, e00097	6
666	Association between QTc interval prolongation and outcomes of diabetic foot ulcers: Data from a 4-year follow-up study in China. 2018 , 138, 26-34	4
665	Synergistical action of the β adrenoceptor and fatty acid binding protein 2 polymorphisms on the loss of glomerular filtration rate in Chinese patients with type 2 diabetic nephropathy. 2018 , 50, 715-723	4
664	Prospective Comparison of Equations Based on Creatinine and Cystatin C for the Glomerular Filtration Rate Estimation in Chinese Renal Transplant Recipients. 2018 , 50, 85-91	4
663	Fatty liver index, albuminuria and the association with chronic kidney disease: a population-based study in China. 2018 , 8, e019097	10

662	Serum Ω Polyunsaturated Fatty Acids and Potential Influence Factors in Elderly Patients with Multiple Cardiovascular Risk Factors. 2018 , 8, 1102	7
661	Association of co-exposure to heavy metals with renal function in a hypertensive population. 2018 , 112, 198-206	17
660	Impact of serum albumin levels on long-term all-cause, cardiovascular, and cardiac mortality in patients with first-onset acute myocardial infarction. 2018 , 477, 89-93	16
659	Inverse association between plasma homocysteine concentrations and type 2 diabetes mellitus among a middle-aged and elderly Chinese population. 2018 , 28, 278-284	1
658	Soluble Tim-3 and Gal-9 are associated with renal allograft dysfunction in kidney transplant recipients: A cross-sectional study. 2018 , 55, 330-335	11
657	Trajectories of Circulating Monocyte Subsets After ST-Elevation Myocardial Infarction During Hospitalization: Latent Class Growth Modeling for High-Risk Patient Identification. 2018 , 11, 22-32	6
656	Existing creatinine-based equations overestimate glomerular filtration rate in Indians. 2018 , 19, 22	10
655	Comparison of clinicopathologic characteristics, epigenetic biomarkers and prognosis between renal pelvic and ureteral tumors in upper tract urothelial carcinoma. 2018 , 18, 22	5
654	Effects of statin therapy on chronic kidney disease patients with coronary artery disease. 2018 , 17, 84	3
653	Comparison of visceral, body fat indices and anthropometric measures in relation to chronic kidney disease among Chinese adults from a large scale cross-sectional study. 2018 , 19, 40	21
652	Single-marker and multi-marker approaches to appraise the relationships between biomarkers and microalbuminuria in Chinese middle-aged and elderly from communities: a cross-sectional analysis. 2018 , 19, 93	3
651	Serum carbohydrate antigen 153 and renal function in patients with type 2 diabetes mellitus. 2018 , 32, e22461	5
650	Comparison of pulse wave velocity and pulse pressure amplification in association with target organ damage in community-dwelling elderly: The Northern Shanghai Study. 2018 , 41, 372-381	9
649	Obesity, hyperhomocysteinaemia and risk of chronic kidney disease: a population-based study. 2018 , 35, 259-265	8
648	The association between plasma big endothelin-1 levels at admission and long-term outcomes in patients with atrial fibrillation. 2018 , 272, 1-7	10
647	Association between dietary patterns, cadmium intake and chronic kidney disease among adults. 2018 , 37, 276-284	64
646	T cell responses to peptides of Goodpasture autoantigen in patients with anti-glomerular basement membrane disease. 2018 , 23, 345-350	3
645	Associations of lipid parameters with insulin resistance and diabetes: A population-based study. 2018 , 37, 1423-1429	38

644	Development of a cardiovascular diseases risk prediction model and tools for Chinese patients with type 2 diabetes mellitus: A population-based retrospective cohort study. 2018 , 20, 309-318	12
643	Intra-aortic balloon pumping and thrombocytopenia in patients with acute coronary syndrome : Incidence, risk factors, and prognosis. 2018 , 43, 555-564	1
642	Effect of Chronic Total Occlusion Percutaneous Coronary Intervention on Clinical Outcomes in Elderly Patients. 2018 , 355, 174-182	10
641	Epidemiological characteristics of lower extremity arterial disease in Chinese diabetes patients at high risk: a prospective, multicenter, cross-sectional study. 2018 , 32, 150-156	20
640	Microribonucleic acid-192 as a specific biomarker for the early diagnosis of diabetic kidney disease. 2017 , 9, 602	5
639	Pharmacodynamics and pharmacokinetics of ticagrelor vs. clopidogrel in patients with acute coronary syndromes and chronic kidney disease. 2018 , 84, 88-96	19
638	Baseline characteristics and temporal differences in Acarbose Cardiovascular Evaluation (ACE) trial participants. 2018 , 199, 170-175	4
637	Lack of ethnic differences of moxifloxacin and metabolite pharmacokinetics in East Asian men. 2018 , 45, 199-214	3
636	Novel Equations for Estimating Lean Body Mass in Patients With Chronic Kidney Disease. 2018 , 28, 156-164	6
635	The effect of L-thyroxine substitution on oxidative stress in early-stage diabetic nephropathy patients with subclinical hypothyroidism: a randomized double-blind and placebo-controlled study. 2018 , 50, 97-103	5
634	Prognostic value of plasma NT-proBNP levels in very old patients with moderate renal insufficiency in China. 2018 , 51, 889-896	1
633	Effect of Diabetes on the Assessment Role of 2-Oxoglutarate to the Severity of Chronic Heart Failure. 2018 , 126, 478-486	0
632	Visceral Adiposity Index as a Predictor of Chronic Kidney Disease in a Relatively Healthy Population in Taiwan. 2018 , 28, 91-100	13
631	Association of elevated blood serum high-sensitivity C-reactive protein levels and body composition with chronic kidney disease: A population-based study in Taiwan. 2018 , 97, e11896	3
630	Glomerular Filtration Rate is Associated with Hemorrhagic Transformation in Acute Ischemic Stroke Patients without Thrombolytic Therapy. 2018 , 131, 1639-1644	7
629	Macrocytic anemia is associated with the severity of liver impairment in patients with hepatitis B virus-related decompensated cirrhosis: a retrospective cross-sectional study. 2018 , 18, 161	8
628	Serum uromodulin and progression of kidney disease in patients with chronic kidney disease. 2018 , 16, 316	18
627	Changes of Percent Body Fat as a Useful Surrogate for Risk of Declined Renal Function. 2018 , 8, 17289	2

626	Association of Blood Pressure Trajectories in Early Life with Subclinical Renal Damage in Middle Age. <i>Journal of the American Society of Nephrology: JASN</i> , 2018 , 29, 2835-2846	12.7	44
625	Is mycophenolate mofetil combined with low-dose prednisone a treatment option for advanced IgA nephropathy? A 10-year follow-up case and brief literature review. 2018 , 16, 205873921880268		
624	Aristolochic acid containing herbs induce gender-related oncological differences in upper tract urothelial carcinoma patients. 2018 , 10, 6627-6639		9
623	Female Sex as a Risk Factor for Ischemic Stroke and Systemic Embolism in Chinese Patients With Atrial Fibrillation: A Report From the China-AF Study. 2018 , 7, e009391		6
622	Low serum phosphate and magnesium levels are associated with peripheral neuropathy in patients with type 2 diabetes mellitus. 2018 , 146, 1-7		16
621	Chronic kidney disease can increase the risk of preoperative deep vein thrombosis in middle-aged and elderly patients with hip fractures. 2018 , 13, 1669-1674		5
620	Relationships of Hyperhomocysteinemia and Hyperuricemia With Metabolic Syndrome and Renal Function in Chinese Centenarians. 2018 , 9, 502		11
619	Risk factors of chronic kidney diseases in Chinese adults with type 2 diabetes. 2018 , 8, 14686		14
618	Gamma-Glutamyltransferase and Risk of Acute Coronary Syndrome in Young Chinese Patients: A Case-Control Study. 2018 , 2018, 2429160		10
617	Clinical Characteristics and Outcomes in the Very Elderly Patients Hospitalized for Acute Heart Failure: Importance of Pharmacologic Guideline Adherence. 2018 , 8, 14270		5
616	Associations of urinary epidermal growth factor and monocyte chemotactic protein-1 with kidney involvement in patients with diabetic kidney disease. 2020 , 35, 291-297		13
615	Sociodemographic associations with abnormal estimated glomerular filtration rate (eGFR) in a large Canadian city: a cross-sectional observation study. 2018 , 19, 198		6
614	Advantageous effect of telbivudine on renal function in patients with chronic hepatitis B infection. 2018 , 5, 33-37		
613	Cognitive function and 3-year mortality in the very elderly Chinese population with chronic kidney disease. 2018 , 13, 2055-2060		4
612	Effect of Donor Age on Graft Function and Pathologic Findings in Living Donor Transplantation. 2018 , 50, 2431-2435		1
611	Clinical and pathologic characteristics of pauci-immune anti-myeloperoxidase antibody associated glomerulonephritis with nephrotic range proteinuria. 2018 , 40, 554-560		6
610	The effects of AER and eGFR on outcomes of CVD in patients with T2DM in an urban community over 8 years of multifactorial treatment: the Beijing Communities Diabetes Study 18. 2018 , 14, 1537-1545		4
609	The prevalence of depressive symptoms in Chinese longevous persons and its correlation with vitamin D status. 2018 , 18, 198		14

608	Clinical implications of pathological features of primary membranous nephropathy. 2018 , 19, 215	19
607	Optimal equation for estimation of glomerular filtration rate in autosomal dominant polycystic kidney disease: influence of tolvaptan. 2018 , 22, 1213-1223	0
606	Perturbations in mitochondrial dynamics by p66Shc lead to renal tubular oxidative injury in human diabetic nephropathy. 2018 , 132, 1297-1314	20
605	Prevalence and Risk Factors for CKD: A Comparison Between the Adult Populations in China and the United States. 2018 , 3, 1135-1143	32
604	Serum Cystatin C Level Is Not a Promising Biomarker for Predicting Clinicopathological Characteristics of Bladder Urothelial Tumors. 2018 , 2018, 2617439	1
603	Prevalence of metabolic syndrome risk factors and their relationships with renal function in Chinese centenarians. 2018 , 8, 9863	3
602	Bidirectional association between nonalcoholic fatty liver disease and hypertension from the Dongfeng-Tongji cohort study. 2018 , 12, 660-670	5
601	Therapeutic drug monitoring of cycloserine and linezolid during anti-tuberculosis treatment in Beijing, China. 2018 , 22, 931-936	13
600	A pilot and comparative study between pathological and serological levels of immunoglobulin and complement among three kinds of primary glomerulonephritis. 2018 , 19, 18	6
599	Hyper-Uricemia and Gouty Access in the Adult Population of the Southeast of Gabon: Biochemical Aspects. 2018 , 6,	4
598	Association of Thyroid Function With the Estimated Glomerular Filtration Rate in a Large Chinese Euthyroid Population. 2018 , 43, 1075-1083	6
597	Serum uric acid levels and decreased estimated glomerular filtration rate in patients with type 2 diabetes: A cohort study and meta-analysis. 2018 , 34, e3046	16
596	Triglyceride to high density lipoprotein cholesterol ratio may serve as a useful predictor of major adverse coronary event in female revascularized ST-elevation myocardial infarction. 2018 , 485, 166-172	3
595	Clinicopathological Characteristics, Role of Immunosuppressive Therapy and Progression in IgA Nephropathy with Hyperuricemia. 2018 , 43, 1131-1140	7
594	Serum Corin Level Is Associated With Subsequent Decline in Renal Function in Patients With Suspected Coronary Artery Disease. 2018 , 7,	3
593	Gender-related differences in clinicopathological characteristics and renal outcomes of Chinese patients with IgA nephropathy. 2018 , 19, 31	10
592	Association of arterial stiffness and central hemodynamics with moderately reduced glomerular filtration rate in Chinese middle-aged and elderly community residents: a cross-sectional analysis. 2018 , 19, 103	3
591	Coronary Artery Bypass Surgery versus Percutaneous Coronary Intervention for Left Main Coronary Artery Disease with Chronic Kidney Disease. 2018 , 59, 279-285	9

590	Urinary epidermal growth factor predicts renal prognosis in antineutrophil cytoplasmic antibody-associated vasculitis. 2018 , 77, 1339-1344	11
589	Blood Serum Potassium and Sodium Concentration Constants: Search for Regulatory Factors. 2018 , 44, 418-423	
588	Relation of serum uric acid to asymptomatic proximal extracranial artery stenosis in a middle-aged Chinese population: a community-based cross-sectional study. 2018 , 8, e020681	3
587	Bilirubin and its changes were negatively associated with diabetic kidney disease incidence and progression: A five-yearIs cohort study based on 5323 Chinese male diabetic patients. 2018 , 32, 1012-1017	5
586	Comparison of various lipid parameters in association of target organ damage: a cohort study. 2018 , 17, 199	11
585	Relationship between high-sensitivity cardiac troponin T and the prognosis of elderly inpatients with non-acute coronary syndromes. 2018 , 13, 1091-1098	11
584	The association between time-mean serum uric acid levels and the incidence of chronic kidney disease in the general population: a retrospective study. 2018 , 19, 190	8
583	Association of various glycemic variability indices and vascular outcomes in type-2 diabetes patients: A retrospective study. 2018 , 97, e10860	6
582	Prediabetes and diabetes are both risk factors for adverse outcomes in infective endocarditis. 2018 , 35, 1499-1507	7
581	Stationary Treatment Compared with Individualized Chinese Medicine for Type 2 Diabetes Patients with Microvascular Complications: Study Protocol for a Randomized Controlled Trial. 2018 , 24, 728-733	2
580	Visceral adiposity index is associated with increased urinary albumin excretion: A population-based study. 2019 , 38, 1332-1338	12
579	The cardiothoracic ratio: a neglected preoperative risk-stratified method for patients with rheumatic heart disease undergoing valve replacement surgery. 2019 , 55, 511-517	1
578	Effect of conjugated bilirubin on clinical outcomes in infective endocarditis. 2019 , 38, 2259-2266	1
577	Perturbations of pulsatile hemodynamics and clinical outcomes in patients with acute heart failure and reduced, mid-range or preserved ejection fraction. 2019 , 14, e0220183	1
576	Complement activation products in the circulation and urine of primary membranous nephropathy. 2019 , 20, 313	22
575	Increased circulating Tfh to Tfr ratio in chronic renal allograft dysfunction: a pilot study. 2019 , 20, 26	10
574	Assessment of adiposity distribution and its association with diabetes and insulin resistance: a population-based study. 2019 , 11, 51	8
573	Moderate chronic kidney disease and left atrial enlargement independently predict thromboembolic events and mortality in elderly patients with atrial fibrillation: a retrospective single-center study. 2019 , 47, 4312-4323	2

572	Relationship between drug application and mortality rate in Chinese older coronary artery disease/chronic heart failure patients with and without low glomerular filtration rate. 2019 , 20, 44	1
571	Age, creatinine and ejection fraction (ACEF) score: a simple risk-stratified method for infective endocarditis. 2019 , 112, 900-906	6
570	Clinical Outcomes of Transmetatarsal Amputation in Patients with Diabetic Foot Ulcers Treated without Revascularization. 2019 , 10, 1465-1472	4
569	Increased cardio-ankle vascular index is independently associated with chronic kidney disease: A cross-sectional study in Chinese patients with type 2 diabetes mellitus. 2019 , 33, 623-627	1
568	Switching from allopurinol to febuxostat: efficacy and safety in the treatment of hyperuricemia in renal transplant recipients. 2019 , 41, 595-599	2
567	Association of Urinary Plasminogen-Plasmin with Edema and Epithelial Sodium Channel Activation in Patients with Nephrotic Syndrome. 2019 , 50, 92-104	10
566	Effects of aging, baseline renal function and stage of HIV infection on post-treatment changes in renal function among HIV-infected patients: a retrospective cohort study. 2019 , 20, 591-600	4
565	The Accuracy of Renal Function Measurements in Obstructive Hydronephrosis Using Dynamic Contrast-Enhanced MR Renography. 2019 , 213, 859-866	1
564	Efficacy of long-term treatment with tenofovir in Chinese nucleos(t)ide-naïve chronic hepatitis B patients regardless of baseline viral load. 2019 , 18, 260-268	1
563	Association between exposure to arsenic, nickel, cadmium, selenium, and zinc and fasting blood glucose levels. 2019 , 255, 113325	24
562	Low-Normal Thyroid Function Predicts Incident Anemia in the General Population With Euthyroid Status. 2019 , 104, 5693-5702	3
561	Prognostic Significance of Murine Double Minute 2 Expression in Tumor Cells in Upper Tract Urothelial Carcinoma: An Analysis of 341 Cases in a Large Chinese Center. 2019 , 23, 797-806	
560	Increased Expression Of SOX2 Predicts A Poor Prognosis And Promotes Malignant Phenotypes In Upper Tract Urothelial Carcinoma. 2019 , 11, 9095-9106	6
559	Prevalence and Associated Risk Factors of Chronic Kidney Disease in an Elderly Population from Eastern China. 2019 , 16,	12
558	Association of uric acid in serum and urine with subclinical renal damage: Hanzhong Adolescent Hypertension Study. 2019 , 14, e0224680	12
557	Serum haptoglobin levels are associated with renal function decline in type 2 diabetes mellitus patients in a Chinese Han population. 2019 , 156, 107865	2
556	The effect of pulse methylprednisolone induction therapy in Chinese patients with dialysis-dependent MPO-ANCA associated vasculitis. 2019 , 76, 105883	13
555	Number of parity is associated with low-grade albuminuria in middle-aged and elderly Chinese women. 2019 , 19, 117	1

554	Visit-to-Visit Variability in Total Cholesterol Correlates with the Progression of Renal Function Decline in a Chinese Community-Based Hypertensive Population. 2019 , 44, 727-742	2
553	Prevalence of kidney damage in Chinese elderly: a large-scale population-based study. 2019 , 20, 341	5
552	Decreased Expression of Urinary Mammalian Target of Rapamycin mRNA Is Related to Chronic Renal Fibrosis in IgAN. 2019 , 2019, 2424751	3
551	Outcomes of patients with anemia and renal dysfunction in hospitalized heart failure with preserved ejection fraction (from the CN-HF registry). 2019 , 25, 100415	3
550	Joint association of body mass index and central obesity with cardiovascular events and all-cause mortality in prediabetic population: A prospective cohort study. 2019 , 13, 453-461	3
549	Correlation between kidney and peripheral nerve functions in Type 2 diabetes. 2020 , 113, 173-180	2
548	Electrocardiographic left ventricular hypertrophy and mortality in an oldest-old hypertensive Chinese population. 2019 , 14, 1657-1662	5
547	Development and validation of a more accurate estimating equation for glomerular filtration rate in a Chinese population. 2019 , 95, 636-646	27
546	Outcomes and practice patterns with hemodiafiltration in Shanghai: a longitudinal cohort study. 2019 , 20, 34	2
545	Neutrophil-to-lymphocyte ratio: An effective predictor of corticosteroid response in IgA nephropathy. 2019 , 74, 105678	7
544	Relationship between serum uric acid and hypertension: a cross-sectional study in Bangladeshi adults. 2019 , 9, 9061	34
543	Association of Blood Pressure and Risk of Cardiovascular and Chronic Kidney Disease in Hong Kong Hypertensive Patients. 2019 , 74, 331-340	15
542	Genetic Variants Flanking the Gene Were Associated with Renal Function in Chinese Patients with Type 2 Diabetes. 2019 , 2019, 9387358	4
541	Derivation and validation of a prediction score for acute kidney injury secondary to acute myocardial infarction in Chinese patients. 2019 , 20, 195	5
540	Plasma metals and cardiovascular disease in patients with type 2 diabetes. 2019 , 129, 497-506	19
539	Lipid profile and renal safety of tenofovir disoproxil fumarate-based anti-retroviral therapy in HIV-infected Chinese patients. 2019 , 83, 64-71	7
538	Elevated triglycerides rather than other lipid parameters are associated with increased urinary albumin to creatinine ratio in the general population of China: a report from the REACTION study. 2019 , 18, 57	6
537	Association between serum 25-hydroxyvitamin D and diabetic kidney disease in Chinese patients with type 2 diabetes. 2019 , 14, e0214728	11

536	High serum IgA/C3 ratio better predicts a diagnosis of IgA nephropathy among primary glomerular nephropathy patients with proteinuria \geq 1 g/d: an observational cross-sectional study. 2019 , 20, 150	6
535	How to estimate renal function in patients with liver disease: choosing the most suitable equation. 2019 , 51, 677-690	4
534	HbA1C Variability Is Strongly Associated With the Severity of Peripheral Neuropathy in Patients With Type 2 Diabetes. 2019 , 13, 90	21
533	Bioinformatics-based discovery of the urinary BBOX1 mRNA as a potential biomarker of diabetic kidney disease. 2019 , 17, 59	5
532	Pioglitazone increases VEGFR3 expression and promotes activation of M2 macrophages via the peroxisome proliferator-activated receptor α 2019 , 19, 2740-2748	6
531	N-terminal pro-brain natriuretic peptide levels had an independent and added ability in the evaluation of all-cause mortality in older Chinese patients with atrial fibrillation. 2019 , 19, 56	6
530	THE ASSOCIATION OF SERUM BILIRUBIN ON KIDNEY CLINICOPATHOLOGIC FEATURES AND RENAL OUTCOME IN PATIENTS WITH DIABETIC NEPHROPATHY: A BIOPSY-BASED STUDY. 2019 , 25, 554-561	2
529	Fetuin-B Links Nonalcoholic Fatty Liver Disease to Chronic Kidney Disease in Obese Chinese Adults: A Cross-Sectional Study. 2019 , 74, 287-295	5
528	Correlation of Serum 1,5-AG with Uric Acid in Type 2 Diabetes Mellitus with Different Renal Functions. 2019 , 2019, 4353075	2
527	A randomized controlled trial on the blood pressure-lowering effect of amlodipine and nifedipine-GITS in sustained hypertension. 2019 , 21, 648-657	6
526	Diabetes-Related Complications and Mortality in Patients With Young-Onset Latent Autoimmune Diabetes: A 14-Year Analysis of the Prospective Hong Kong Diabetes Register. 2019 , 42, 1042-1050	16
525	Association between admission plasma 2-oxoglutarate levels and short-term outcomes in patients with acute heart failure: a prospective cohort study. 2019 , 25, 8	2
524	Prognostic Comparison of the Estimations of Renal Function in Patients With Acute Heart Failure. 2019 , 83, 767-774	1
523	The predictive value of the renal resistive index for contrast-induced nephropathy in patients with acute coronary syndrome. 2019 , 19, 36	5
522	Gender- and Age-Specific Associations between Visceral Obesity and Renal Function Impairment. 2019 , 12, 67-77	7
521	Current status and progression of lower urinary tract symptoms in Chinese male patients: the protocol and rationale for a nationwide, hospital-based, prospective, multicentre study. 2019 , 9, e028378	1
520	A retrospective cohort study of oral anticoagulant treatment in patients with acute coronary syndrome and atrial fibrillation. 2019 , 9, e031180	5
519	Prevalence, clinicopathological features, and prognosis in upper tract urinary carcinoma patients with severe preoperative chronic kidney disease. 2019 , 8, 641-650	3

518	Burden of CKD and Cardiovascular Disease on Life Expectancy and Health Service Utilization: a Cohort Study of Hong Kong Chinese Hypertensive Patients. <i>Journal of the American Society of Nephrology: JASN</i> , 2019 , 30, 1991-1999	12.7	6
517	Increased risk of aspirin-induced gastric mucosal erosion in elderly Chinese men harboring SLCO1B1*1b/*1b while using aspirin and an ACEI or ARB concomitantly. 2019 , 20, 183		3
516	Comparison of Renal Function Estimation Formulae for Dosing Direct Oral Anticoagulants in Patients with Atrial Fibrillation. 2019 , 8,		7
515	The association between socioeconomic status and prevalence of chronic kidney disease: A cross-sectional study among rural residents in eastern China. 2019 , 98, e14822		6
514	Increasing Time in Therapeutic Range of Tacrolimus in the First Year Predicts Better Outcomes in Living-Donor Kidney Transplantation. 2019 , 10, 2912		5
513	Diagnostic performance of serum cystatin C and complement component 1q in lupus nephritis. 2019 , 21, 267		2
512	The value of cystatin C and urinary and serum neutrophil gelatinase-associated lipocalin during the perioperative period of renal transplantation. 2019 , 8, 432-441		4
511	Epidemiology of acute kidney injury in intensive care units in Beijing: the multi-center BAKIT study. 2019 , 20, 468		21
510	PCSK9 positively correlates with plasma sdLDL in community-dwelling population but not in diabetic participants after confounder adjustment. 2019 , 98, e15062		2
509	Establishment of a predictive model for inpatient sudden cardiac death in a Chinese cardiac department population: a retrospective study. 2019 , 132, 17-24		1
508	Safety and Efficacy of Reduced Prolonged-release Tacrolimus Exposure in De Novo Kidney Transplantation: A Randomized, Open-label, Pilot Study in Asia-OPTIMIZE Study. 2019 , 5, e340		1
507	Serum PCSK6 and corin levels are not associated with cardiovascular outcomes in patients undergoing coronary angiography. 2019 , 14, e0226129		2
506	Spectrum of chronic kidney disease in China: A national study based on hospitalized patients from 2010 to 2015. 2019 , 24, 725-736		27
505	Serum immunoglobulin G provides early risk prediction in immunoglobulin A nephropathy. 2019 , 66, 13-18		7
504	Effects of Niaoduqing Particles () on Delaying Progression of Renal Dysfunction: A Post-trial, Open-Label, Follow-up Study. 2019 , 25, 168-174		2
503	Reasonable Glycemic Control Would Help Wound Healing During the Treatment of Diabetic Foot Ulcers. 2019 , 10, 95-105		26
502	Interaction between PLA2R1 and HLA-DQA1 variants contributes to the increased genetic susceptibility to membranous nephropathy in Western China. 2019 , 24, 919-925		4
501	Plasma acylcarnitines could predict prognosis and evaluate treatment of IgA nephropathy. 2019 , 16, 2		7

500	Successful endovascular treatment of chronic renal artery occlusion: a preliminary retrospective case series including 15 patients. 2019 , 51, 285-291	4
499	Combining the use of amino-terminal pro-B-type natriuretic peptide and B-type natriuretic peptide in the prognosis of hospitalized heart failure patients. 2019 , 491, 8-14	4
498	Gender impact on the correlations between GravesLhyperthyroidism and hyperuricemia in Chinese. 2019 , 188, 843-848	5
497	Interactive effects of testosterone and the androgen receptor CAG repeat length polymorphism on cardiovascular-renal events and mortality in men with diabetes. 2019 , 35, e3081	6
496	Complement deposition on renal histopathology of patients with diabetic nephropathy. 2019 , 45, 363-368	17
495	Beta 2-Microglobulin and the Severity of Coronary Stenosis in Patients With Acute Coronary Syndrome. 2019 , 28, 575-582	5
494	Renal impairment markers in type 2 diabetes patients with different types of hyperuricemia. 2019 , 10, 118-123	5
493	Elevated plasma soluble (pro)renin receptor levels are associated with left ventricular remodeling and renal function in chronic heart failure patients with reduced ejection fraction. 2019 , 111, 152-157	12
492	Associations of Amino Acid and Acylcarnitine Profiles With Incident Hyperuricemia in Middle-Aged and Older Chinese Individuals. 2020 , 72, 1305-1314	5
491	Soluble urokinase-type plasminogen activator receptor and incident end-stage renal disease in Chinese patients with chronic kidney disease. 2020 , 35, 465-470	8
490	A Novel Important Node Discovery Algorithm Based on Local Community Aggregation and Recognition in Complex Networks. 2020 , 27, 253-260	2
489	Altered DNA methylation of TRIM13 in diabetic nephropathy suppresses mesangial collagen synthesis by promoting ubiquitination of CHOP. 2020 , 51, 102582	11
488	Association of visit-to-visit variability of systolic blood pressure with cardiovascular disease, chronic kidney disease and mortality in patients with hypertension. 2020 , 38, 943-953	8
487	Using race in the estimation of glomerular filtration rates: time for a reversal?. 2020 , 29, 227-231	4
486	Plasma retinol-binding protein 4 in the first and second trimester and risk of gestational diabetes mellitus in Chinese women: a nested case-control study. 2020 , 17, 1	23
485	Anemia in centenarians: prevalence and association with kidney function. 2020 , 25, 26-33	2
484	Incidence, Risk Factors and Subsequent Prognostic Impact of New-Onset Atrial Fibrillation in Infective Endocarditis. 2020 , 84, 262-268	0
483	Association between atherogenic index of plasma and subclinical renal damage over a 12-year follow-up: Hanzhong adolescent hypertension study. 2020 , 74, 278-284	4

482	Multiple plasma metals, genetic risk and serum C-reactive protein: A metal-metal and gene-metal interaction study. 2020 , 29, 101404	12
481	An overlap of antineutrophil cytoplasmic antibody (ANCA)-associated glomerulonephritis and IgG4-related kidney disease. 2020 , 501, 12-19	3
480	Relationship between arterial stiffness and chronic kidney disease in patients with primary hypertension. 2020 , 34, 577-585	3
479	Geriatric nutrition risk index is associated with renal progression, cardiovascular events and all-cause mortality in chronic kidney disease. 2020 , 33, 783-793	3
478	Observations from a teaching hospital in Ireland: changing from MDRD to CKD-EPI eGFR in routine practice. 2021 , 74, 608-611	0
477	Associations of Triglyceride-Glucose Index and Its Derivatives with Hyperuricemia Risk: A Cohort Study in Chinese General Population. 2020 , 2020, 3214716	3
476	The plasma level of mCRP is linked to cardiovascular disease in antineutrophil cytoplasmic antibody-associated vasculitis. 2020 , 22, 228	3
475	Circulating Ceramide: A New Cardiometabolic Biomarker in Patients With Comorbid Acute Coronary Syndrome and Type 2 Diabetes Mellitus. 2020 , 11, 1104	7
474	Application of a Pharmacokinetic Model to Inform the Optimal Dose for Individualized Drug Administration. 2020 , 709-714	1
473	Variants in the RARRES2 gene are associated with serum chemerin and increase the risk of diabetic kidney disease in type 2 diabetes. 2020 , 165, 1574-1580	4
472	Novel risk models to predict acute kidney disease and its outcomes in a Chinese hospitalized population with acute kidney injury. 2020 , 10, 15636	10
471	Efficacy and safety of bivalirudin application during primary percutaneous coronary intervention in older patients with acute ST-segment elevation myocardial infarction. 2020 , 48, 300060520947942	0
470	Leptin mediate central obesity on the severity of cardiovascular autonomic neuropathy in well-controlled type 2 diabetes and prediabetes. 2020 , 18, 396	1
469	Hyperuricemia precedes non-alcoholic fatty liver disease with abdominal obesity moderating this unidirectional relationship: Three longitudinal analyses. 2020 , 311, 44-51	4
468	A high triglyceride glucose index is more closely associated with hypertension than lipid or glycemic parameters in elderly individuals: a cross-sectional survey from the Reaction Study. 2020 , 19, 112	15
467	Association of Uric Acid in Serum and Urine with Arterial Stiffness: Hanzhong Adolescent Hypertension Study. 2020 , 2020, 1638515	5
466	Age-Specific Associations Between Systolic Blood Pressure and Cardiovascular Disease: A 10-Year Diabetes Mellitus Cohort Study. 2020 , 9, e015771	4
465	Impact of estimated glomerular filtration rate on long-term clinical outcomes among Chinese patients with atrial fibrillation. 2020 , 20, 490	2

464	Serum homocysteine level is an independent predictor for hemorrhagic transformation within 24 h of intravenous thrombolysis in acute ischemic stroke. 2020 , 82, 13-19	1
463	Cystatin C-Based Renal Function in Predicting the Long-Term Outcomes of Chronic Total Occlusion After Percutaneous Coronary Intervention. 2020 , 7, 586181	2
462	Higher Serum Asprosin Level is Associated with Urinary Albumin Excretion and Renal Function in Type 2 Diabetes. 2020 , 13, 4341-4351	6
461	Neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio in evaluation of inflammation in non-dialysis patients with end-stage renal disease (ESRD). 2020 , 21, 511	3
460	The difference between nocturnal dipping status and morning blood pressure surge for target organ damage in patients with chronic kidney disease. 2020 , 22, 2025-2034	6
459	Sex-Specific Association of Serum Uric Acid Level and Change in Hyperuricemia Status with Risk of Type 2 Diabetes Mellitus: A Large Cohort Study in China. 2020 , 2020, 9637365	5
458	Associations Between Complement Components and Vitamin D and the Physical Activities of Daily Living Among a Longevous Population in Hainan, China. 2020 , 11, 1543	2
457	The association of subclinical atherosclerosis with prediabetes is stronger in people with dyslipidaemia than in those with normoglycaemia: A cross-sectional study in Chinese adults. 2020 , 14, 760-767	1
456	Associations between usual glycated haemoglobin and cardiovascular disease in patients with type 2 diabetes mellitus: A 10-year diabetes cohort study. 2020 , 22, 2325-2334	4
455	Association between diabetes mellitus and health-related quality of life among patients with chronic kidney disease: results from the Chinese Cohort Study of Chronic Kidney Disease (C-STRIDE). 2020 , 18, 266	0
454	A Diagnostic Nomogram of Pathologic Grade for Preoperative Risk Stratification in Upper Tract Urothelial Carcinoma. 2020 , 14, 1179554920927662	0
453	Coagulation parameters are associated with the prognosis of immunoglobulin a nephropathy: a retrospective study. 2020 , 21, 447	0
452	IgG deposits in the mesangium and capillary loops predict poor renal outcome in patients with IgA nephropathy: a single-center retrospective study. 2020 , 42, 902-911	1
451	Serum cystatin C is closely associated with euthyroid Hashimoto's thyroiditis in adult female Chinese patients. 2020 , 48, 300060520952973	1
450	Increased Risk of Clopidogrel-Induced Gastric Mucosal Erosion in Elderly Chinese Men Harboring the Allele. 2020 , 13, 1237-1244	0
449	Association Between Hypertriglyceridemic Waist Phenotype and Increased Urinary Albumin-Creatinine Ratio in Chinese Adults: The REACTION Study. 2020 , 13, 2965-2974	3
448	Associations between Personal PM Elemental Constituents and Decline of Kidney Function in Older Individuals: the China BAPE Study. 2020 , 54, 13167-13174	13
447	Prognostic Value of Secreted Frizzled-Related Protein 5 in Heart Failure Patients With and Without Type 2 Diabetes Mellitus. 2020 , 13, e007054	29

446	Blood Pressure is Associated with Rapid Kidney Function Decline in a Very Elderly Hypertensive Chinese Population. 2020 , 15, 1317-1323	1
445	Comparison of Novel Metabolic Indices in Estimation of Chronic Kidney Diseases in a Southern Chinese Population. 2020 , 13, 4919-4927	3
444	Does Serum Uric Acid Status Influence the Association Between Left Atrium Diameter and Atrial Fibrillation in Hypertension Patients?. 2020 , 7, 594788	1
443	Underweight in Men had a Closer Relationship with Falls than Women in Centenarians. 2020 , 24, 987-992	2
442	Mineral and bone disorder biomarkers and inflammation indexes in patients with end stage renal disease. 2020 , 9, 3938-3946	1
441	Blood group A: a risk factor for heart rupture after acute myocardial infarction. 2020 , 20, 471	0
440	The cumulative blood pressure load and target organ damage in patients with essential hypertension. 2020 , 22, 981-990	6
439	SNPs in PRKCA-HIF1A-GLUT1 are associated with diabetic kidney disease in a Chinese Han population with type 2 diabetes. 2020 , 50, e13264	4
438	Neck circumference is an independent risk factor for hyperuricemia within 3 years in women: a longitudinal study. 2020 , 39, 3757-3767	1
437	The ventilatory abnormalities and prognostic values of H FPEF score in dyspnoeic patients with preserved left ventricle systolic function. 2020 , 7, 1872-1879	2
436	Multiple Measures of Mineral Metabolism Were Associated With Renal Function in Chinese Centenarians: A Cross-Sectional Study. 2020 , 7, 120	4
435	Secreted Frizzled-Related Protein 2 and Extracellular Volume Fraction in Patients with Heart Failure. 2020 , 2020, 2563508	15
434	J-shaped relationship between serum uric acid levels and the risk of ischemic stroke in high-risk individuals: A hospital-based observational study. 2020 , 195, 105906	0
433	Carotid intraplaque neovascularization predicts atherosclerotic renal artery stenosis in patients with carotid artery stenosis. 2020 , 30, 1492-1499	1
432	Renal survival and risk factors in IgA nephropathy with crescents. 2020 , 52, 1507-1516	3
431	A novel nomogram to predict the reliability of estimated glomerular filtration rate formulae in oncology patients. 2020 , 20, 530	3
430	Admission Low-Density Lipoprotein Cholesterol Stratified by Circulating CD14++CD16+ Monocytes and Risk for Recurrent Cardiovascular Events Following ST Elevation Myocardial Infarction: Lipid Paradox Revised. 2020 , 13, 916-927	4
429	Acute kidney disease and acute kidney injury biomarkers in coronary care unit patients. 2020 , 21, 207	13

428	Comparative Performance of Creatinine-Based GFR Estimation Equations in Exceptional Longevity: The Rugao Longevity and Ageing Study. 2020 , 15, 733-742	
427	Risk stratification in patients hospitalized for acute heart failure in Asian population. 2020 , 83, 544-550	0
426	The association of hematuria on kidney clinicopathologic features and renal outcome in patients with diabetic nephropathy: a biopsy-based study. 2020 , 43, 1213-1220	5
425	Changing trajectories of serum uric acid and risk of non-alcoholic fatty liver disease: a prospective cohort study. 2020 , 18, 133	8
424	Age-specific associations of glycosylated haemoglobin variability with cardiovascular disease and mortality in patients with type 2 diabetes mellitus: A 10-year cohort study. 2020 , 22, 1316-1327	7
423	Effects of dynamic change in fetuin-A levels from the first to the second trimester on insulin resistance and gestational diabetes mellitus: a nested case-control study. 2020 , 8,	6
422	Uric acid and its correlation with hypertension in postmenopausal women: A multi-ethnic study (Observational study). 2020 , 42, 559-564	1
421	The variability of glycosylated hemoglobin is associated with renal function decline in patients with type 2 diabetes. 2020 , 11, 2040622319898370	10
420	Dysbiosis of gut microbiota in adult idiopathic membranous nephropathy with nephrotic syndrome. 2020 , 147, 104359	12
419	Alpha1-Antitrypsin in Urinary Extracellular Vesicles: A Potential Biomarker of Diabetic Kidney Disease Prior to Microalbuminuria. 2020 , 13, 2037-2048	3
418	TIR generated by continuous glucose monitoring is associated with peripheral nerve function in type 2 diabetes. 2020 , 166, 108289	18
417	Association Between Atherosclerosis and Diabetic Retinopathy in Chinese Patients with Type 2 Diabetes Mellitus. 2020 , 13, 1911-1920	3
416	Early-Onset Type 2 Diabetes as a Risk Factor for End-Stage Renal Disease in Patients With Diabetic Kidney Disease. 2020 , 17, E50	5
415	Comparison of the risk factors of hemorrhagic transformation between large artery atherosclerosis stroke and cardioembolism after intravenous thrombolysis. 2020 , 196, 106032	3
414	Association between weight status, metabolic syndrome, and chronic kidney disease among middle-aged and elderly Chinese. 2020 , 30, 2017-2026	7
413	The role of pulmonary function in patients with heart failure and preserved ejection fraction: Looking beyond chronic obstructive pulmonary disease. 2020 , 15, e0235152	4
412	Lower uric acid level may be associated with hemorrhagic transformation but not functional outcomes in patients with anterior circulation acute ischemic stroke undergoing endovascular thrombectomy. 2020 , 35, 1157-1164	4
411	Optimal Cut-Offs of Body Mass Index and Waist Circumference to Identify Obesity in Chinese Type 2 Diabetic Patients. 2020 , 13, 1899-1909	0

410	Urinary neutrophil gelatinase-associated lipocalin rapidly decreases in the first week after kidney transplantation. 2020 , 34, e23445	1
409	The association between dyslipidemia and the incidence of chronic kidney disease in the general Zhejiang population: a retrospective study. 2020 , 21, 252	5
408	Risk factors for treatment resistance and relapse of Chinese patients with MPO-ANCA-associated vasculitis. 2020 , 20, 199-206	6
407	Post-contrast acute kidney injury in a hospitalized population: short-, mid-, and long-term outcome and risk factors for adverse events. 2020 , 30, 3516-3527	7
406	Serum Fibroblast Growth Factor 19 and Total Bile Acid Concentrations Are Potential Biomarkers of Hepatocellular Carcinoma in Patients with Type 2 Diabetes Mellitus. 2020 , 2020, 1751989	4
405	HbA1C Variability Is Strongly Associated with Development of Macroalbuminuria in Normal or Microalbuminuria in Patients with Type 2 Diabetes Mellitus: A Six-Year Follow-Up Study. 2020 , 2020, 7462158	6
404	Low TSH level predicts a poor clinical outcome in patients with anterior circulation ischemic stroke after endovascular thrombectomy. 2020 , 41, 1821-1828	3
403	Association analysis of DNA methyltransferases in IgA nephropathy. 2020 , 80, 106147	2
402	Elevated triglyceride to high-density lipoprotein cholesterol (TG/HDL-C) ratio increased risk of hyperuricemia: a 4-year cohort study in China. 2020 , 68, 71-80	11
401	Dynamic Changes and Prognostic Value of Gut Microbiota-Dependent Trimethylamine-N-Oxide in Acute Ischemic Stroke. 2020 , 11, 29	12
400	Associations of apnea hypopnea index and educational attainments with microvascular complications in patients with T2DM. 2020 , 67, 363-373	7
399	Baseline serum albumin and its dynamic change is associated with type 2 diabetes risk: A large cohort study in China. 2020 , 36, e3296	5
398	Analysis of risk factors for carotid intima-media thickness in patients with type 2 diabetes mellitus in Western China assessed by logistic regression combined with a decision tree model. 2020 , 12, 8	3
397	Clustering Heatmap for Visualizing and Exploring Complex and High-dimensional Data Related to Chronic Kidney Disease. 2020 , 9,	13
396	Association of blood lipid profile with incident chronic kidney disease: A Mendelian randomization study. 2020 , 300, 19-25	12
395	Risk of chronic kidney disease defined by decreased estimated glomerular filtration rate in individuals with different prediabetic phenotypes: results from a prospective cohort study in China. 2020 , 8,	2
394	The association of neutrophil-to-lymphocyte ratio with all-cause mortality in Chinese patients with MPO-ANCA associated vasculitis. 2020 , 20, 401-408	5
393	Plasma element levels and risk of chronic kidney disease in elderly populations (≥90 Years old). 2020 , 254, 126809	6

392	Associations between estimated glomerular filtration rate and cardiac biomarkers. 2020 , 34, e23336	3
391	The impact of protein diet on bone density in people with/without chronic kidney disease: An analysis of the National Health and Nutrition Examination Survey database. 2020 , 39, 3497-3503	1
390	The association between serum uric acid levels and ischemic stroke in essential hypertension patients. 2020 , 132, 551-558	6
389	Systemic immune-inflammation index (SII) predicted clinical outcome in patients with coronary artery disease. 2020 , 50, e13230	97
388	Risk scores for predicting incident chronic kidney disease among rural Chinese people: a village-based cohort study. 2020 , 21, 120	4
387	Risk factors for subclinical renal damage and its progression: Hanzhong Adolescent Hypertension Study. 2021 , 75, 531-538	6
386	Evaluation of the Oxford classification in immunoglobulin A vasculitis with nephritis: a cohort study and meta-analysis. 2021 , 14, 516-525	1
385	Renal Sodium Handling in Relation to Environmental and Genetic Factors in Untreated Chinese. 2021 , 34, 394-403	1
384	Feasibility of the transcatheter mitral valve repair for patients with severe mitral regurgitation and endangered heart failure. 2021 , 120, 452-459	1
383	Association of multiple metals with lipid markers against different exposure profiles: A population-based cross-sectional study in China. 2021 , 264, 128505	14
382	On the Basis of Race: The Utility of a Race Factor in Estimating Glomerular Filtration. 2021 , 6, 155-166	3
381	Plasma metals and cancer incidence in patients with type 2 diabetes. 2021 , 758, 143616	4
380	Prognostic significance of uric acid change in acute ischemic stroke patients with reperfusion therapy. 2021 , 28, 1218-1224	1
379	Association between nonalcoholic fatty liver and increased low-level albuminuria in postmenopausal women in China: A cross-sectional study. 2021 , 13, 494-505	1
378	Prediction models and nomograms for 10-year risk of end-stage renal disease in Chinese type 2 diabetes mellitus patients in primary care. 2021 , 23, 897-909	3
377	Long-Term Exposure to Ambient PM and Increased Risk of CKD Prevalence in China. <i>Journal of the American Society of Nephrology: JASN</i> , 2021 , 32, 448-458	12.7 14
376	Prognostic effects of longitudinal changes in left ventricular ejection fraction with cardiac resynchronization therapy. 2021 , 8, 368-379	1
375	Effect of Combining Furosemide With Standard Hydration Therapy on Contrast-Induced Acute Kidney Injury Following Coronary Angiography or Intervention in a High-Risk Population. 2021 , 72, 138-144	1

374	Association of gonadal hormones and sex hormone binding globulin with risk of diabetes: A cohort study in middle-aged and elderly Chinese males. 2021 , 75, e14008	0
373	Validation and Comparison of Six Risk Scores for Infection in Patients With ST-Segment Elevation Myocardial Infarction Undergoing Percutaneous Coronary Intervention. 2020 , 7, 621002	1
372	Associations of Baseline and Changes in Leukocyte Counts with Incident Cardiovascular Events: The Dongfeng-Tongji Cohort Study. 2021 ,	0
371	Comorbidities and outcomes in South Asian individuals with chronic kidney disease: an observational primary care cohort. 2021 ,	2
370	Higher Apnea-Hypopnea Index (AHI) and Oxygen Desaturation Index (ODI) Were Independently Associated with Increased Risks of Hypertension in Patients with T2DM: A Cross-Sectional Study. 2021 , 2021, 8887944	2
369	The Association of Glomerular Filtration Rate With Echocardiographic Parameters in Chronic Kidney Disease. 2021 , 13, 121-129	
368	Pleural Effusion Is Associated with Severe Renal Dysfunction in Patients with Acute Pancreatitis. 2021 , 27, e928118	2
367	Modification and Validation of the Phosphate Removal Model: A Multicenter Study. 2021 , 46, 53-62	0
366	The Relationship Between Chronic Kidney Disease and the Severity and Long-Term Prognosis of Patients with Coronary Artery Disease After Drug-Eluting Stent Implantation. 2021 , 14, 399-404	
365	Predictive value of elevated alanine aminotransferase for in-hospital mortality in patients with acute myocardial infarction. 2021 , 21, 82	2
364	Bone Fragment Co-transplantation Alongside Bone Marrow Aspirate Infusion Protects Kidney Transplant Recipients. 2021 , 12, 630710	1
363	A double-blind, placebo-controlled trial on the antihypertensive treatment effect of a quadruple single-pill combination. 2021 , 23, 815-822	1
362	Urinary soluble CD90 predicts renal prognosis in patients with diabetic kidney disease. 2021 , 9, 282	2
361	A Novel Creatinine-Based Equation to Estimate Glomerular Filtration Rate in Chinese Population With Chronic Kidney Disease: Implications for DOACs Dosing in Atrial Fibrillation Patients. 2021 , 12, 615953	1
360	Associations Between Anemia, Cognitive Impairment, and All-Cause Mortality in Oldest-Old Adults: A Prospective Population-Based Cohort Study. 2021 , 8, 613426	3
359	Low-grade albuminuria is associated with high cardiovascular risk in CVD-free and normoalbuminuric Chinese adults: Results from the REACTION study. 2021 , 13, 648-660	0
358	High-normal thyroid function predicts incident non-alcoholic fatty liver disease among middle-aged and elderly euthyroid subjects. 2021 ,	5
357	Anemia among Chinese patients with chronic kidney disease and its association with quality of life - results from the Chinese cohort study of chronic kidney disease (C-STRIDE). 2021 , 22, 64	1

356	Circulating chemerin level is associated with metabolic, biochemical and haematological parameters-A population-based study. 2021 , 94, 927-939	2
355	Child-to-adult body mass index trajectories and the risk of subclinical renal damage in middle age. 2021 , 45, 1095-1104	5
354	An Evaluation of Commonly Used Surrogate Baseline Creatinine Values to Classify AKI During Acute Infection. 2021 , 6, 645-656	5
353	The Relationship Between Circulating Growth Differentiation Factor 15 Levels and Diabetic Retinopathy in Patients With Type 2 Diabetes. 2021 , 12, 627395	2
352	Lower Plasma Elabela Levels in Hypertensive Patients With Heart Failure Predict the Occurrence of Major Adverse Cardiac Events: A Preliminary Study. 2021 , 8, 638468	1
351	Impact of Baseline Neutrophil-to-Lymphocyte Ratio on Long-Term Prognosis in Patients With Atrial Fibrillation. 2021 , 72, 819-828	3
350	Optimal Cut-Off Values of Visceral Fat Area for Predicting Metabolic Syndrome Among Type 2 Diabetes Patients in Ningbo, China. 2021 , 14, 1375-1383	2
349	Elevated Plasma Big Endothelin-1 at Admission Is Associated With Poor Short-Term Outcomes in Patients With Acute Decompensated Heart Failure. 2021 , 8, 629268	4
348	Association of metabolic syndrome with the incidence of low-grade albuminuria: a cohort study in middle-aged and elderly Chinese adults. 2021 , 13, 7350-7360	0
347	Risk factors for electrocardiographic left ventricular hypertrophy in a young Chinese general population: the Hanzhong adolescent cohort study. 2021 , 21, 159	0
346	β-blocker and 1-year outcomes among patients hospitalized for heart failure with mid-range ejection fraction. 2021 ,	3
345	Feasibility of combining heart rate variability and electrochemical skin conductance as screening and severity evaluation of cardiovascular autonomic neuropathy in type 2 diabetes. 2021 , 12, 1671-1679	1
344	Prognostic significance of hypertension at the onset of lupus nephritis in Chinese patients: prevalence and clinical outcomes. 2021 ,	1
343	Pre-operative N-terminal pro-B-type natriuretic peptide for prediction of acute kidney injury after noncardiac surgery: A retrospective cohort study. 2021 , 38, 591-599	1
342	Novel equation for estimating resting energy expenditure in patients with chronic kidney disease. 2021 , 113, 1647-1656	1
341	Predictive value of CHA2DS2-VASc score combined with hs-CRP for new-onset atrial fibrillation in elderly patients with acute myocardial infarction. 2021 , 21, 175	0
340	Predictive Value of Thyroid Hormones for Incident Hyperuricemia in Euthyroid Subjects: The Tianjin Chronic Low-Grade Systemic Inflammation and Health Cohort Study. 2021 , 27, 291-297	0
339	Serum and urine ANGPTL8 expression levels are associated with hyperlipidemia and proteinuria in primary nephrotic syndrome. 2021 , 22, 130	

338	The association of baseline N-terminal pro-B-type natriuretic peptide with short and long-term prognosis following percutaneous coronary intervention in non-ST segment elevation acute coronary syndrome with multivessel coronary artery disease: a retrospective cohort study. 2021 , 21, 202	0
337	Blood lead, nutrient intake, and renal function among type 2 diabetic patients. 2021 , 28, 49063-49073	0
336	Comparative Risks of Nonsteroidal Anti-Inflammatory Drugs on CKD. 2021 , 16, 898-907	1
335	The Prevalence, Distribution, and Extent of Subclinical Atherosclerosis and Its Relation With Serum Uric Acid in Hypertension Population. 2021 , 8, 638992	3
334	Proliferative diabetic retinopathy in patients with type 2 diabetes correlates with the presence of atherosclerosis cardiovascular disease. 2021 , 13, 48	2
333	Dose-response relationship between serum 25-hydroxyvitamin D and the risk of metabolic syndrome. 2021 , 40, 1530-1536	2
332	Assessment of prognostic value of intraoperative oliguria for postoperative acute kidney injury: a retrospective cohort study. 2021 , 126, 799-807	1
331	Chronic kidney disease in acute coronary syndromes: real-world data of long-term outcomes. 2021 , 17, 1359-1369	1
330	Choosing an appropriate glomerular filtration rate estimating equation: role of body mass index. 2021 , 22, 197	0
329	Lipoprotein (a) is associated with poor long-term prognosis in patients aged 80 years and older with acute coronary syndrome. 2021 , 15, 466-476	2
328	Associations between air pollutant exposure and renal function: A prospective study of older adults without chronic kidney disease. 2021 , 277, 116750	4
327	Hypertriglyceridemic waist phenotype and risk of chronic kidney disease in community-dwelling adults aged 60 years and older in Tianjin, China: a 7-year cohort study. 2021 , 22, 182	0
326	Insulin resistance surrogates predict hypertension plus hyperuricemia. 2021 , 12, 2046-2053	10
325	Association Between Renal Dysfunction and Low HDL Cholesterol Among the Elderly in China. 2021 , 8, 644208	0
324	The association between variants in PLA2R and HLA-DQA1 and renal outcomes in patients with primary membranous nephropathy in Western China. 2021 , 14, 123	1
323	A new scoring system for predicting short-term outcomes in Chinese patients with critically-ill acute decompensated heart failure. 2021 , 21, 228	0
322	Development and Validation of a Risk Score for Predicting Post-acute Myocardial Infarction Infection in Patients Undergoing Percutaneous Coronary Intervention: Study Protocol for an Observational Study. 2021 , 8, 675142	1
321	Elevated urinary albumin concentration predicts worse clinical outcomes in hospitalized acute decompensated heart failure patients. 2021 , 8, 3037-3048	2

320	Associations Between Obesity and Kidney Disease in Chinese Men and Women With Type 2 Diabetes: A Retrospective Cohort Study. 2021 ,	0
319	The Risk Threshold for Hemoglobin A1c Associated With Albuminuria: A Population-Based Study in China. 2021 , 12, 673976	0
318	Prognostic value of D-dimer for adverse outcomes in patients with infective endocarditis: an observational study. 2021 , 21, 279	2
317	Triglyceride-Glucose Index and Extracellular Volume Fraction in Patients With Heart Failure. 2021 , 8, 704462	4
316	Relationship Between Serum Neopterin Level and Peripheral Arterial Plaque in Patients with Type 2 Diabetes. 2021 , 14, 2871-2878	0
315	Successful single kidney transplantation from pediatric donors less than or equal to 10 kg to adult recipient: a retrospective cohort study. 2021 , 10, 1618-1629	0
314	Association between consumption of ultra-processed foods and hyperuricemia: TCLSIH prospective cohort study. 2021 , 31, 1993-2003	1
313	Prognostic Value of Plasma NT-proBNP levels in Hospitalized Patients Older than 80 Years of Age in a Hospital in Beijing, China. 2021 , 116, 1027-1036	0
312	Perirenal Fat Thickness Is Significantly Associated With the Risk for Development of Chronic Kidney Disease in Patients With Diabetes. 2021 , 70, 2322-2332	4
311	Exploring the mediating role of serum retinol-binding protein 4 in the relationship between sleep quality and insulin resistance in pregnant women. 2021 , 176, 108866	1
310	Association of fluid balance trajectories with clinical outcomes in patients with septic shock: a prospective multicenter cohort study. 2021 , 8, 40	
309	Significance of serum FGF-23 for risk assessment of contrast-associated acute kidney injury and clinical outcomes in patients undergoing coronary angiography. 2021 , 16, e0254835	2
308	Association between Latent Trajectories of Fluid Balance and Clinical Outcomes in Critically Ill Patients with Acute Kidney Injury: A Prospective Multicenter Observational Study.. 2022 , 8, 82-92	
307	Impact of Serum Uric Acid Concentration on the Risk of Cardiovascular Disease: A Cohort Study Conducted in Northern China. 2021 , 117, 666-675	1
306	Systemic immune-inflammation index predicts a reduced risk of end-stage renal disease in Chinese patients with myeloperoxidase-anti-neutrophil cytoplasmic antibody-associated vasculitis: A retrospective observational study. 2021 , 22, 989	1
305	Optimal blood pressure for patients with end-stage renal disease following coronary interventions. 2021 , 23, 1622-1630	1
304	TIFA protein expression is associated with pulmonary arterial hypertension. 2021 , 11, 14140	
303	Serum albumin and albuminuria predict the progression of chronic kidney disease in patients with newly diagnosed type 2 diabetes: a retrospective study. 2021 , 9, e11735	2

302	The characteristics and risk factors of healthcare-seeking men with lower urinary tract symptoms in China: Initial report from the POInT group. 2021 , 40, 1740-1753	0
301	ALKBH1-demethylated DNA N6-methyladenine modification triggers vascular calcification via osteogenic reprogramming in chronic kidney disease. 2021 , 131,	8
300	Association of central arterial blood pressure and left ventricular hypertrophy in patients with chronic kidney disease. 2021 ,	
299	Safe Limits of Contrast Media for Contrast-Induced Nephropathy: A Multicenter Prospective Cohort Study. 2021 , 8, 701062	0
298	Association of depressive symptoms with the progression of carotid intima-media thickness in a community-based cohort in Beijing, China. 2021 , 11, e050254	
297	Sex-Related Characteristics, Quality of Care, and Short-Term Mortality Rates in Patients With Chronic Heart Failure. 2022 , 31, 77-84	0
296	Associations of Estimated Glomerular Filtration Rate with All-Cause Mortality and Cardiovascular Mortality in Patients with Diabetic Foot Osteomyelitis. 2021 , 14, 4499-4509	0
295	Association between Metabolic Syndrome and Microvascular Complications in Chinese Adults with Type 1 Diabetes Mellitus. 2021 ,	2
294	Correlation between thyroid autoantibodies and cardiovascular disease in patients with stages 3-5 chronic kidney disease. 2021 , 9, 1301	
293	Age-Specific Associations of Usual Blood Pressure Variability With Cardiovascular Disease and Mortality: 10-Year Diabetes Mellitus Cohort Study. 2021 , 10, e019026	1
292	Incorporating Post-Cessation Weight-Control Coaching into Smoking Cessation Therapy to Reduce Type 2 Diabetes Risk. 2021 , 13,	
291	Development and Validation of Creatinine-Based Estimates of the Glomerular Filtration Rate Equation from Tc-DTPA Imaging in the Malaysian Setting. 2021 , 2021, 3465472	
290	Acute Lower Respiratory Tract Infection Increased the Risk of Cardiovascular Events and All-Cause Mortality in Elderly Patients With Stable Coronary Artery Disease. 2021 , 8, 711264	
289	Blood urea nitrogen, blood urea nitrogen to creatinine ratio and incident stroke: The Dongfeng-Tongji cohort. 2021 , 333, 1-8	4
288	Nonlinear Relationship Between Macrocytic Anemia and Decompensated Hepatitis B Virus Associated Cirrhosis: A Population-Based Cross-Sectional Study. 2021 , 12, 755625	0
287	Dietary patterns and risk for hyperuricemia in the general population: Results from the TCLSIH cohort study. 2022 , 93, 111501	0
286	Association of pulsatile stress in childhood with subclinical renal damage in adults: A 30-year prospective cohort study. 2021 , 23, 1843-1851	1
285	Serum creatinine levels and risk of nonalcohol fatty liver disease in a middle-aged and older Chinese population: A cross-sectional analysis. 2021 , e3489	0

284	Comprehensive metabolomics profiling reveals common metabolic alterations underlying the four major non-communicable diseases in treated HIV infection. 2021 , 71, 103548	2
283	HbA combined with glycated albumin or 1,5-anhydroglucitol improves the efficiency of diabetes screening in a Chinese population. 2021 , e14685	2
282	Relationship Between Change in Serum Uric Acid and Ischemic Stroke in Chinese Hypertensive Patients. 2021 , 8, 717128	1
281	Comparison of cystatin C-based and creatinine-based glomerular filtration rate in the prediction of postoperative residual hypertension in aldosterone-producing adenoma patients after adrenalectomy. 2021 , 520, 147-153	0
280	Association between urinary VEGFA and renal pathology of IgA nephropathy patients. 2021 , 35, e23995	1
279	The increased miRNA-150-5p expression of the tonsil tissue in patients with IgA nephropathy may be related to the pathogenesis of disease. 2021 , 100, 108124	0
278	Clinical classification of hyperuricemia in patients with chronic kidney disease. 2021 , 53, 1665-1674	
277	Residual Risk and Its Risk Factors for Ischemic Stroke with Adherence to Guideline-Based Secondary Stroke Prevention. 2021 , 23, 51-60	4
276	Efficacy, safety and response predictors of adjuvant astragalus for diabetic kidney disease (READY): study protocol of an add-on, assessor-blind, parallel, pragmatic randomised controlled trial. 2021 , 11, e042686	4
275	Association between Plasma Osmolality and Case Fatality within 1 Year after Severe Acute Ischemic Stroke. 2021 , 62, 600-607	0
274	Reference values of glomerular filtration rate for healthy adults in southern China: a cross-sectional survey. 2021 , 12, 20406223211035287	
273	The Safety and Efficacy of Inspiratory Muscle Training for Patients With Acute Myocardial Infarction Undergoing Percutaneous Coronary Intervention: Study Protocol for a Randomized Controlled Trial. 2020 , 7, 598054	0
272	Kidney Function Tests. 2012 , 669-707	3
271	Association of plasma antimony concentration with markers of liver function in Chinese adults. 2020 , 17, 304	3
270	The impact of chronic kidney disease on long-term outcomes following semi-urgent and elective percutaneous coronary intervention. 2021 , 32, 517-525	1
269	Performance of the CKD-EPI creatinine-cystatin C glomerular filtration rate estimation equations in a multiethnic Asian population. 2014 , 55, 656-9	10
268	2018 Chinese Guidelines for Prevention and Treatment of Hypertension-A report of the Revision Committee of Chinese Guidelines for Prevention and Treatment of Hypertension. 2019 , 16, 182-241	205
267	[Association of metabolic syndrome with chronic kidney disease in premenopausal and postmenopausal women]. 2019 , 39, 861-866	1

266	Analysis of risk factors for acute kidney injury after ascending aortic replacement combined with open placement of triple-branched stent graft in type A aortic dissection: a new technique versus the traditional method in a single Chinese center. 2015 , 21, 674-80	11
265	Optimal Vitamin D Status in a Middle-Aged and Elderly Population Residing in Shanghai, China. 2017 , 23, 6001-6011	8
264	Metformin or Acarbose Treatment Significantly Reduced Albuminuria in Patients with Newly Diagnosed Type 2 Diabetes Mellitus and Low-Grade Albuminuria. 2018 , 24, 8941-8949	7
263	Clinical Features of Nephrotic Syndrome with Cerebral Hemorrhage. 2019 , 25, 2179-2185	6
262	Causes and Factors Associated with Frequent Hospitalization in Chinese Patients with Systemic Lupus Erythematosus: An Ambispective Cohort Study. 2019 , 25, 8061-8068	10
261	Association of E/ELand NT-proBNP with renal function in patients with essential hypertension. 2013 , 8, e54513	7
260	Multicenter study of creatinine- and/or cystatin C-based equations for estimation of glomerular filtration rates in Chinese patients with chronic kidney disease. 2013 , 8, e57240	25
259	Low-Grade Albuminuria Is Associated with Metabolic Syndrome and Its Components in Middle-Aged and Elderly Chinese Population. 2013 , 8, e65597	21
258	Urinary uromodulin excretion predicts progression of chronic kidney disease resulting from IgA nephropathy. 2013 , 8, e71023	25
257	Association of insulin resistance with chronic kidney disease in non-diabetic subjects with normal weight. 2013 , 8, e74058	28
256	Association of high-density lipoprotein cholesterol with the estimated glomerular filtration rate in a community-based population. 2013 , 8, e79738	15
255	Circulating serum trefoil factor 3 (TFF3) is dramatically increased in chronic kidney disease. 2013 , 8, e80271	29
254	Use of net reclassification improvement (NRI) method confirms the utility of combined genetic risk score to predict type 2 diabetes. 2013 , 8, e83093	29
253	Comparisons between the 2012 new CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration) equations and other four approved equations. 2014 , 9, e84688	28
252	Plasma homocysteine is associated with aortic arterial stiffness but not wave reflection in Chinese hypertensive subjects. 2014 , 9, e85938	18
251	Association of EZSCAN values with arterial stiffness in individuals without diabetes or cardiovascular disease. 2014 , 9, e90854	6
250	Hypertriglyceridemic waist phenotype and chronic kidney disease in a Chinese population aged 40 years and older. 2014 , 9, e92322	15
249	Relationship of arterial compliance and blood pressure with microalbuminuria and mildly decreased glomerular filtration rate: a Chinese community-based analysis. 2014 , 9, e101013	2

248	Serum gamma - glutamyltransferase is associated with albuminuria: a population-based study. 2014 , 9, e114970	12
247	Gender differences in the risk factors for endothelial dysfunction in Chinese hypertensive patients: homocysteine is an independent risk factor in females. 2015 , 10, e0118686	24
246	High prevalence of lower extremity peripheral artery disease in type 2 diabetes patients with proliferative diabetic retinopathy. 2015 , 10, e0122022	17
245	Effects of body mass index on risks for ischemic stroke, thromboembolism, and mortality in Chinese atrial fibrillation patients: a single-center experience. 2015 , 10, e0123516	20
244	Metabolic Syndrome without Diabetes or Hypertension Still Necessitates Early Screening for Chronic Kidney Disease: Information from a Chinese National Cross-Sectional Study. 2015 , 10, e0132220	5
243	The Clinical Utility of SUDOSCAN in Chronic Kidney Disease in Chinese Patients with Type 2 Diabetes. 2015 , 10, e0134981	18
242	Heart Rate Variability Is Associated with Exercise Capacity in Patients with Cardiac Syndrome X. 2016 , 11, e0144935	7
241	Circulating Level of Neutrophil Extracellular Traps Is Not a Useful Biomarker for Assessing Disease Activity in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. 2016 , 11, e0148197	34
240	Retrospective Study of Phospholipase A2 Receptor and IgG Subclasses in Glomerular Deposits in Chinese Patients with Membranous Nephropathy. 2016 , 11, e0156263	27
239	Validation of glomerular filtration rate-estimating equations in Chinese children. 2017 , 12, e0180565	7
238	The prognostic value of asymmetric dimethylarginine in patients with cardiac syndrome X. 2017 , 12, e0188995	5
237	The failure of glomerular filtration rate estimating equations among obese population. 2020 , 15, e0242447	6
236	Nighttime blood pressure decline as a predictor of renal injury in patients with hypertension: a population-based cohort study. 2019 , 11, 4310-4322	3
235	Combined efficacy of C-reactive protein and red blood cell distribution width in prognosis of patients with culture-negative infective endocarditis. 2017 , 8, 71173-71180	7
234	Changes in renal function indices in cirrhotic chronic hepatitis C patients treated with sofosbuvir-containing regimens. 2017 , 8, 90916-90924	3
233	Genetic polymorphisms in HLA-DP and STAT4 are associated with IgA nephropathy in a Southwest Chinese population. 2018 , 9, 7066-7074	5
232	Effect of CTLA-4 gene polymorphisms on long-term kidney allograft function in Han Chinese recipients. 2016 , 7, 23088-95	3
231	[Impact of diagnostic ureteroscopy and biopsy on radical nephroureterectomy of upper tract urothelial carcinoma]. 2019 , 51, 665-672	1

230	Predicting Metabolic Syndrome With Machine Learning Models Using a Decision Tree Algorithm: Retrospective Cohort Study. 2020 , 8, e17110	13
229	Comparative performance of four equations estimating glomerular filtration rate in adult Chinese diabetics. 2013 , 36, 293-7	4
228	Association of glomerular filtration rate with arterial stiffness in Chinese women with normal to mildly impaired renal function. 2012 , 9, 158-65	11
227	Efficacy and safety of combined directly acting antivirals for treatment of Chinese chronic hepatitis C patients in a real-world setting. 2017 , 23, 4072-4079	3
226	Clinicopathological features and prognosis in patients with idiopathic membranous nephropathy with hypertension. 2020 , 19, 2615-2621	2
225	Validation of the Korean coefficient for the modification of diet in renal disease study equation. 2016 , 31, 344-56	4
224	Low bone mineral density is associated with increased arterial stiffness in participants of a health records based study. 2015 , 7, 790-8	9
223	The prognostic values of beta-2 microglobulin for risks of cardiovascular events and mortality in the elderly patients with isolated systolic hypertension. 2018 , 23, 82	4
222	The association between changes in urinary albumin-to-creatinine ratio and risk of abnormal ankle-brachial index in a community-based Chinese population. 2012 , 19, 912-7	5
221	Chronic kidney disease, all-cause mortality and cardiovascular mortality among Chinese patients with established cardiovascular disease. 2010 , 17, 395-401	23
220	The impact of serum uric acid on the natural history of glomerular filtration rate: a retrospective study in the general population. 2016 , 4, e1859	6
219	Application of Modified Glomerular Filtration Rate Estimation Equations in Chinese Diabetic Patients with Chronic Kidney Diseases. 2015 , 64, 209-12	5
218	Reduced global longitudinal strain as a marker for early detection of Fabry cardiomyopathy. 2021 ,	2
217	Genetics and Clinical Characteristics of PPAR γ Variant-Induced Diabetes in a Chinese Han Population. 2021 , 12, 677130	1
216	The Mediation Effects of Aluminum in Plasma and Dipeptidyl Peptidase Like Protein 6 (DPP6) Polymorphism on Renal Function via Genome-Wide Typing Association. 2021 , 18,	
215	Dietary Approaches to Kidney Diseases. 2011 , 2170-2204	2
214	Correlation between serum uric Acid and renal function in patients with stable coronary artery disease and type 2 diabetes. 2014 , 6, 443-50	8
213	CO-MORBIDITIES AND THEIR MANAGEMENT IN PATIENTS WITH CHRONIC KIDNEY DISEASE IN A TERTIARY HOSPITAL OF KERALA. 2017 , 4, 1595-1598	

212	Accuracy of glomerular filtration rate estimation equations in patients with hematopathy. 2017 ,	
211	High Level of Lipoprotein(a) as Predictor for Recurrent Heart Failure in Patients with Chronic Heart Failure: a Cohort Study. 2019 , 113, 197-204	1
210	Parity is associated with albuminuria and chronic kidney disease: a population-based study. 2019 , 11, 11030-11039	2
209	A novel nomogram to predict the reliability of estimated glomerular filtration rate formula in oncology patients.	
208	Clinicopathological characteristics and prognosis of patients with IgA nephropathy and renal vasculitic lesions. 2021 , 22, 353	0
207	Scoring model to predict risk of chronic kidney disease in Chinese health screening examinees with type 2 diabetes. 2021 , 1	
206	Prevalence and Associated Factors of Kidney Dysfunction in Patients with Hypertension and/or Diabetes Mellitus from a Primary Care Population in Northwest China. 2021 , 14, 7567-7578	1
205	The Relationship Between Simple Renal Cysts and Renal Function in Patients With Type 2 Diabetes. 2020 , 11, 616167	0
204	The contribution of plasma uric acid to the risk of stroke in hypertensive populations. 2020 , 31, 298-303	0
203	Revealing the Decrease of Renal Cortical Perfusion in Primary Glomerular Disease and Renal Aging by Arterial Spin Labeling. 2020 , 17,	
202	Evaluation of the prognostic ability of serum uric acid for elderly acute coronary syndrome patients with diabetes mellitus: a prospective cohort study. 2021 , 22, 856-865	1
201	Health inequities and the inappropriate use of race in nephrology. 2021 ,	7
200	Underweight in Men Had a Closer Relationship with Falls than Women in Centenarians. 2020 , 24, 987-992	1
199	Sphingosine-1-phosphate in anti-neutrophil cytoplasmic antibody-associated vasculitis: coagulation-related clinical indicators and complications. 2020 , 40,	0
198	Contrast-Enhanced Ultrasound Assessment of Renal Parenchymal Perfusion in Patients with Atherosclerotic Renal Artery Stenosis to Predict Renal Function Improvement After Revascularization. 2020 , 13, 1713-1721	3
197	Ageing and the glomerular filtration rate: truths and consequences. 2009 , 120, 419-28	154
196	Rosuvastatin attenuated contrast-induced nephropathy in diabetes patients with renal dysfunction. 2015 , 8, 2342-9	9
195	Association between plasma homocysteine and progression of early nephropathy in type 2 diabetic patients. 2015 , 8, 11174-80	11

194	Comparison of Bare-Metal Stent and Drug-Eluting Stent for the Treatment of Patients Undergoing Percutaneous Coronary Intervention for Unprotected Left Main Coronary Artery Disease - Long-Term Result from a Single Center Experience. 2015 , 31, 381-9	4
193	Staged versus "one-time" multivessel intervention in elderly patients with non-ST-elevation acute coronary syndrome. 2016 , 13, 760-767	2
192	Influence of renal function on the association between homocysteine level and risk of ischemic stroke. 2017 , 9, 4553-4563	1
191	Cystatin C is Indispensable for Evaluation of Kidney Disease. 2017 , 28, 268-276	15
190	[A high level of high-density lipoprotein cholesterol is a protective factor against transplant renal artery stenosis]. 2018 , 38, 155-161	
189	[Characteristics of BK polymavirus infection in kidney transplant recipients]. 2019 , 39, 120-124	
188	[Early intervention of BK virus replication promotes stabilization of renal graft function]. 2017 , 37, 1110-1115	
187	[Clinical and pathological features in IgA nephropathy with IgG deposition in the glomerular mesangial area]. 2017 , 37, 308-311	
186	[Clinical characteristics of patients with acute renal infarction: an analysis of 52 patients in a single center]. 2019 , 51, 863-869	1
185	Influencing factors for peripheral and posterior lesions in mild non-proliferative diabetic retinopathy-the Kailuan Eye Study. 2020 , 13, 1467-1476	
184	[Correlation of type 2 diabetes and impaired glucose regulation with chronic kidney disease in middle-aged and elderly individuals]. 2020 , 40, 1457-1464	
183	Effects of sodium-glucose cotransporter 2 inhibitors on cardiovascular outcomes in elderly patients with comorbid coronary heart disease and diabetes mellitus. 2021 , 18, 440-448	
182	Mild Renal Dysfunction and Risk of Thromboembolism and Bleeding in Patients with Atrial Fibrillation: The Chinese Atrial Fibrillation Registry Study. 2021 , 1, 29-36	
181	Physical activities mediate the correlations between serum creatinine and bone mineral density in Chinese. 2021 , 524, 25-33	
180	Combined Effects of Dyslipidemia and High Adiposity on the Estimated Glomerular Filtration Rate in a Middle-Aged Chinese Population. 2021 , 14, 4513-4522	1
179	Higher aldosterone is associated with increased renal impairment risk in patients with hypertension and abnormal glucose metabolism: a longitudinal study. 2021 ,	0
178	The effect of body weight and alcohol consumption on hyperuricemia and their attributable population fractions: A National health survey in China. 2021 ,	3
177	Fibrinogen to Albumin Ratio as an Independent Risk Factor for Type 2 Diabetic Kidney Disease. 2021 , 14, 4557-4567	0

- 176 A Prospective Comparative Study on Cardiac Alterations After Surgery and Drug Treatment of Primary Aldosteronism. **2021**, 12, 770711 0
- 175 Estimation of Mycophenolic Acid Exposure in Heart Transplant Recipients by Population Pharmacokinetic and Limited Sampling Strategies. **2021**, 12, 748609
- 174 Clinic and ambulatory blood pressure in relation to the interaction between plasma advanced glycation end products and sodium dietary intake and renal handling. **2021**, 0
- 173 Utilizing reclassification to explore characteristics and prognosis of KDIGO AKI subgroups: a retrospective analysis of a multicenter prospective cohort study. **2021**, 43, 1569-1576
- 172 VLCKD: a real time safety study in obesity.. **2022**, 20, 23 3
- 171 Clinical Comparison of the Glomerular Filtration Rate Calculated from Different Renal Depths and Formulae. **2022**, 12, 698 0
- 170 Influencing factors for peripheral and posterior lesions in mild non-proliferative diabetic retinopathyThe Kailuan Eye Study. **2020**, 13, 1467-1476 0
- 169 Patients with end-stage renal disease requiring hemodialysis benefit from percutaneous coronary intervention after non-ST-segment elevation myocardial infarction.. **2022**, 1
- 168 Evaluation of the long-term prognostic ability of triglyceride-glucose index for elderly acute coronary syndrome patients: a cohort study.. **2022**, 21, 3 4
- 167 Healthcare resource utilisation for chronic kidney disease and other major non-communicable chronic diseases in China: a cross-sectional study.. **2022**, 12, e051888 0
- 166 The Effect of Body Adiposity and Alcohol Consumption on Serum Uric Acid: A Quantile Regression Analysis Based on the China National Health Survey.. **2021**, 8, 724497 0
- 165 Combined Effect of Homocysteine and Uric Acid to Identify Patients With High Risk for Subclinical Atrial Fibrillation.. **2021**, e021997 0
- 164 Effect of type 2 diabetes on coronary artery ectasia: smaller lesion diameter and shorter lesion length but similar adverse cardiovascular events.. **2022**, 21, 9 1
- 163 Mild renal insufficiency and attributable risk of adverse In-hospital outcomes in patients with Acute Coronary Syndrome from the improving care for Cardiovascular Disease in China (CCC) project.. **2022**, 23, 29 1
- 162 Association of Chinese Visceral Adiposity Index and Its Dynamic Change With Risk of Carotid Plaque in a Large Cohort in China.. **2021**, e022633 0
- 161 Association of education levels with the risk of hypertension and hypertension control: a nationwide cohort study in Chinese adults.. **2022**, 1
- 160 Association of urinary albumin:creatinine ratio with incident frailty in older populations.
- 159 A Variation in SORBS1 is Associated with Type 2 Diabetes and High-density Lipoprotein Cholesterol in Chinese Population.. **2022**, e3524

158	Cystatin C-based equations for estimating glomerular filtration rate do not require race or sex coefficients.. 2022 , 1-5	1
157	Sex Disparities in the Association of Serum Uric Acid With Kidney Stone: A Cross-Sectional Study in China.. 2022 , 9, 774351	0
156	Hyperhomocysteinemia Increases Risk of Metabolic Syndrome and Cardiovascular Death in an Elderly Chinese Community Population of a 7-Year Follow-Up Study.. 2021 , 8, 811670	3
155	Multi-scalar data integration links glomerular angiotensin-converting enzyme signaling pathway activation with progression of diabetic kidney disease.	
154	Development and Validation of a Nomogram for Predicting the Risk of Adverse Cardiovascular Events in Patients with Coronary Artery Ectasia.. 2021 , 8,	1
153	Disparate Associations of 24-h Central Aortic and Brachial Cuff Blood Pressure With Hypertension-Mediated Organ Damage and Cardiovascular Risk.. 2022 , 9, 795509	0
152	Association between Platelet to Neutrophil Ratio (PNR) and Clinical Outcomes in STEMI Patients after Successful pPCI: A Secondary Analysis Based on a Cohort Study.. 2022 , 2022, 2022657	
151	Joint Effect of Beer, Spirits Intake, and Excess Adiposity on Hyperuricemia Among Chinese Male Adults: Evidence From the China National Health Survey.. 2022 , 9, 806751	0
150	Prognostic analysis of crescentic glomerulonephritis with acute kidney injury: a single-center cohort with 5-year follow-up.. 2022 ,	0
149	Serum CTRP9 Reflects Coronary Collateralization in Nondiabetic Patients with Obstructive Coronary Artery Disease.. 2022 , 2022, 8537686	
148	Association of Serum Adipokines and Resting Energy Expenditure in Patients With Chronic Kidney Disease.. 2022 , 9, 828341	1
147	METS-IR, a novel score to evaluate insulin sensitivity, is associated with the urinary albumin-creatinine ratio in Chinese adults: A cross-sectional REACTION study.. 2022 ,	
146	Cystatin C-based estimated GFR performs best in identifying individuals with poorer survival in an unselected Chinese population: results from the China Health and Retirement Longitudinal Study (CHARLS).	
145	Serum creatinine and cystatin C-based diagnostic indices for sarcopenia in advanced non-small cell lung cancer.. 2022 ,	1
144	Paradoxical mortality of high estimated glomerular filtration rate reversed by 24-h urine creatinine excretion rate adjustment: sarcopenia matters.. 2022 ,	0
143	A study on association of serum uric acid and blood pressure in hypertensives at a tertiary care centre. 2022 , 8, 264-268	
142	Comparison of Coronary Artery Bypass Grafting and Drug-Eluting Stent Implantation in Patients With Chronic Kidney Disease: A Propensity Score Matching Study.. 2022 , 9, 802181	
141	Prognostic Role of Pulmonary Function in Patients With Heart Failure With Reduced Ejection Fraction.. 2022 , e023422	0

- 140 Efficacy and safety of Zicuiyin decoction on diabetic kidney disease: A multicenter, randomized controlled trial.. **2022**, 100, 154079 0
- 139 Prevalence, risk factors and clinical characteristics of renal dysfunction in Chinese outpatients with growth simple renal cysts. **2021**, 1
- 138 Comparison of 6 equations for estimating glomerular filtration rate in a Chinese benign hypertensive nephrosclerosis population.. **2021**, 100, e28318 1
- 137 The diagnostic role of PD-1 CXCR5 follicular helper CD8 T cell in renal allograft dysfunction.. **2021**, e24200 1
- 136 Associations of Carbonaceous Compounds and Water-Soluble Inorganic Ions in Ambient PM with Renal Function in Older Individuals: The China BAPE Study.. **2021**,
- 135 Association of genetic polymorphisms in IL-23R and IL-17A with the susceptibility to IgA nephropathy in a Chinese Han population.. **2021**, 0
- 134 Gender Differences in the Association Between Obesity Indices and Chronic Kidney Disease Among Middle-Aged and Elderly Taiwanese Population: A Community-Based Cross-Sectional Study.. **2021**, 12, 737586 0
- 133 Association Between Metabolic Syndrome and Peripheral Arterial Disease in Elderly Patients with Type 2 Diabetes.. **2021**, 14, 4783-4789 1
- 132 Gender-specific and age-specific associations of the homoeostasis model assessment for IR (HOMA-IR) with albuminuria and renal function impairment: a retrospective cross-sectional study in Southeast China.. **2021**, 11, e053649 0
- 131 A simple prediction model of hyperuricemia for use in a rural setting. **2021**, 11, 23300 0
- 130 Association Between Early Oral β -Blocker Therapy and In-Hospital Outcomes in Patients With ST-Elevation Myocardial Infarction With Mild-Moderate Heart Failure: Findings From the CCC-ACS Project.. **2022**, 9, 828614
- 129 Comparison of Risk of Target Organ Damage in Different Phenotypes of Arterial Stiffness and Central Aortic Blood Pressure.. **2022**, 9, 839875 1
- 128 Image_1.TIF. **2019**,
- 127 Table_1.DOCX. **2020**,
- 126 Table_1.DOCX. **2020**,
- 125 Table_1.DOCX. **2020**,
- 124 Data_Sheet_1.doc. **2020**,
- 123 Image_1.pdf. **2020**,

122	Association of Long-term Ambient Fine Particulate Matter (PM) and Incident CKD: A Prospective Cohort Study in China.. 2022,	0
121	Higher plasma renin activity is associated with increased kidney damage risk in patients with hypertension and glucose metabolic disorders.. 2022,	1
120	Health Risks of Chronic Exposure to Small Doses of Microcystins: An Integrative Metabolomic and Biochemical Study of Human Serum.. 2022,	1
119	Chronic kidney disease among greenhouse workers and field workers in China.. 2022, 302, 134905	0
118	Red Blood Cell Distribution Width: A Prognostic Marker in Patients With Type B Aortic Dissection Undergoing Endovascular Aortic Repair.. 2022, 9, 788476	0
117	Population pharmacokinetics of omeprazole in obese and normal-weight adults.. 2022,	1
116	Association of a novel electrolyte index, SUSPPUP, based on the measurement of fasting serum and spot urinary sodium and potassium, with prediabetes and diabetes in Chinese population.. 2022,	
115	Two Competing Guilds as a Core Microbiome Signature for Chronic Diseases.	1
114	Characteristics of acute kidney injury and its impact on outcome in patients with acute-on-chronic liver failure.. 2022, 22, 231	1
113	Early life body mass index trajectories and albuminuria in midlife: A 30-year prospective cohort study.. 2022, 48, 101420	1
112	Variability, Clearance, and Concentration of Multiple Metals and Risk of Kidney Function Abnormality: A New Integrative Metal Exposure Assessment Approach.	1
111	Role of depressive symptoms in the prognosis of heart failure and its potential clinical predictors.	0
110	Phenylacetylglutamine as a risk factor and prognostic indicator of heart failure.	0
109	Prevalence and risk factors of hyperuricaemia in non-obese Chinese: a single-centre cross-sectional study. 2022, 12, e048574	
108	Estimated glomerular filtration rate derived from different formulas and prognosis in acute coronary syndrome: Findings from the Improving Care for Cardiovascular Disease in China-Acute Coronary Syndrome Project. 2022,	
107	Metabolic Dysfunction-Associated Fatty Liver Disease Is Associated with the Risk of Incident Cardiovascular Disease: A Prospective Cohort Study in Xinjiang. 2022, 14, 2361	0
106	NRF2 Genetic Polymorphism Modifies the Association of Plasma Selenium Levels With Incident Coronary Heart Disease among Type 2 Diabetes Individuals.	0
105	Preoperative NT-proBNP and LVEF for the prediction of acute kidney injury after noncardiac surgery: a single-centre retrospective study. 2022, 22,	0

- 104 The Ratio of CD226 and TIGIT Expression in Tfh and PD-1+ICOS+Tfh Cells Are Potential Biomarkers for Chronic Antibody-Mediated Rejection in Kidney Transplantation. **2022**, 2022, 1-12
- 103 Urinary exosomes derived circRNAs as biomarkers for chronic renal fibrosis. **2022**, 54, 1966-1976 ○
- 102 Prediction model for the risk of ESKD in patients with primary FSGS.
- 101 Far-Infrared Therapy Improves Arteriovenous Fistula Patency and Decreases Plasma Asymmetric Dimethylarginine in Patients with Advanced Diabetic Kidney Disease: A Prospective Randomized Controlled Trial. **2022**, 11, 4168
- 100 Association between Perceived Salt Intake and Arterial Stiffness. **2022**, 2022, 1-7 1
- 99 Sugar-Sweetened Beverage Consumption Is Associated With More Obesity and Higher Serum Uric Acid in Chinese Male Gout Patients With Early Onset. 9, ○
- 98 Renal Denervation Reduces Blood Pressure and Improves Cardiac Function: Results from a 12-Month Study. **2022**, 2022, 1-7
- 97 Chinese Visceral Adiposity Index Is Associated With Incident Renal Damage in Patients With Hypertension and Abnormal Glucose Metabolism: A Longitudinal Study. 13, ○
- 96 Predictive Value of Glycosylated Hemoglobin for Post-operative Acute Kidney Injury in Non-cardiac Surgery Patients. 9,
- 95 Ambient ozone pollution and prevalence of chronic kidney disease: A nationwide study based on the China National survey of chronic kidney disease. **2022**, 306, 135603 ○
- 94 Clinical Characteristics of Inpatients With New-Onset Diabetes Mellitus in Eastern China: Based on Novel Clustering Analysis. 13, ○
- 93 Prevalence of hyperuricemia and the population attributable fraction of modifiable risk factors: Evidence from a general population cohort in China. 10, 1
- 92 The association between a body shape index and elevated urinary albumin:creatinine ratio in Chinese community adults. 13, ○
- 91 Comparison of three non-insulin-based insulin resistance indexes in predicting the presence and severity of coronary artery disease. 9, 1
- 90 Ticagrelor versus Clopidogrel in Patients with Severe Renal Insufficiency Undergoing PCI for Acute Coronary Syndrome. **2022**, 2022, 1-9
- 89 Triglyceride-glucose index in the prediction of adverse cardiovascular events in patients with premature coronary artery disease: a retrospective cohort study. **2022**, 21, ○
- 88 The fibrosis-4 score is associated with long-term mortality in different phenotypes of acute heart failure.
- 87 Associations between co-exposure to multiple metals and renal function: a cross-sectional study in Guangxi, China. 1

- 86 Association between sleep duration and ideal cardiovascular health in Chinese adults: results from the China health and nutrition survey.
- 85 Clinical characteristics and outcomes of chronic kidney disease in people living with HIV in a resource limited centre of central China.
- 84 High neutrophil to lymphocyte ratio with type 2 diabetes mellitus predicts poor prognosis in patients undergoing percutaneous coronary intervention: a large-scale cohort study. **2022**, 21, 0
- 83 The association between systemic inflammatory response index and new-onset atrial fibrillation in patients with ST-elevated myocardial infarction treated with percutaneous coronary intervention. 0
- 82 Association between neutrophil percentage-to-albumin ratio and 3-month functional outcome in acute ischemic stroke patients with reperfusion therapy. 13, 0
- 81 Circulating metals, leukocyte microRNAs and microRNA networks: A profiling and functional analysis in Chinese adults. **2022**, 169, 107511 0
- 80 Association between serum pyrethroid insecticide levels and incident type 2 diabetes risk: a nested case-control study in Dongfeng-Tongji cohort. **2022**, 37, 959-970 1
- 79 The prevalence of MAFLD and its association with atrial fibrillation in a nationwide health check-up population in China. 13, 0
- 78 Prognostic significance of the extent of tubulointerstitial lesions in patients with IgA nephropathy. 0
- 77 Renal fat fraction is significantly associated with the risk of chronic kidney disease in patients with type 2 diabetes. 13, 0
- 76 Effectiveness of the Risk Assessment and Management Programme for hypertension (RAMP-HT) in reducing complications and service utilization after 5 years: A population-based matched cohort study. 0
- 75 S100A8/A9, an Upregulated Host Factor in BK Virus Infection after Kidney Transplantation, Is Associated with Allograft Function Impairment. **2022**, 21, 2356-2366 1
- 74 Nomogram based on high-frequency shear wave elastography (SWE) to evaluate chronic changes after kidney transplantation. 0
- 73 Non-glomerular Tip Lesion Focal Segmental Glomerulosclerosis as a Negative Predictor in Idiopathic Membranous Nephropathy. 0
- 72 The Prognosis of Diabetic Foot Ulcer is Independent of age? A Comparative Analysis of the Characteristics of Patients with Diabetic Foot Ulcer in Different age Groups: A Cross-Sectional Study from China. 153473462211258 0
- 71 Association of exposure to multiple serum metals with the risk of chronic kidney disease in the elderly: a population-based case-control study. 1
- 70 Association of complement C3 with incident type 2 diabetes and the mediating role of BMI: A 10-year follow-up study. 0
- 69 Association between left ventricular geometry and global myocardial work in patients with heart failure with preserved ejection fraction: assessment using strain-pressure loop. 0

- 68 Association between Serum Uric Acid and Hypertension in a Large Cross-Section Study in a Chinese Population. **2022**, 9, 346 ○
- 67  **2022**, 14-23 ○
- 66 Case Report: Endovascular Treatment of Chronic Atherosclerotic Renal Artery Total Occlusions with Failed Medical Therapy. 9, ○
- 65 Length of Hospitalization-Related Differences and Associated Long-Term Prognosis of Patients with Cardiac Resynchronization Therapy: A Propensity Score-Matched Cohort. **2022**, 9, 354 ○
- 64 Influence of baseline arterial stiffness on effects of intensive compared with standard blood pressure control: a post hoc analysis of the STEP trial. **2022**, 20, ○
- 63 Impact of levosimendan on efficacy and renal function in acute heart failure according to renal function: A perspective, multi-center, real-world registry. 9, ○
- 62 Autologous wound margin point columnar full-thickness skin grafting combined with negative pressure wound therapy improves wound healing in refractory diabetic foot ulcers. ○
- 61 Risk stratification in patients with hypertrophic cardiomyopathy: Looking beyond the left side myocardial function. Publish Ahead of Print, ○
- 60 Bivalirudin vs. heparin on a background of ticagrelor and aspirin in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention: A multicenter prospective cohort study. 9, ○
- 59 Impact of chronic kidney disease on the extent and severity of coronary plaque burden in general population: evaluation by Coronary CT angiography. ○
- 58 Fat-to-Muscle Ratio Is Independently Associated with Hyperuricemia and a Reduced Estimated Glomerular Filtration Rate in Chinese Adults: The China National Health Survey. **2022**, 14, 4193 ○
- 57 Comparison of glomerular filtration rate estimating formulas among Japanese adults without kidney disease. **2022**, ○
- 56 Associations between plasma metal levels and mild renal impairment in the general population of Southern China. **2022**, 247, 114209 1
- 55 Multi-Scalar Data Integration Links Glomerular Angiopoietin-Tie Signaling Pathway Activation With Progression of Diabetic Kidney Disease. ○
- 54 Prognostic potential of liver injury in patients with dilated cardiomyopathy: a retrospective study. **2022**, 27, 1
- 53 Beneficial effect of the short-chain fatty acid propionate on vascular calcification through intestinal microbiota remodelling. **2022**, 10, ○
- 52 Performance of the renal resistive index and usual clinical indicators in predicting persistent AKI. **2022**, 44, 2028-2038 ○
- 51 Association between Serum Uric Acid Levels and the Prevalence of Heart Failure due to Acute Coronary Syndrome in Chinese Hospitalized Patients: A Cross-sectional Study. **2022**, ○

- 50 Associations of co-exposure to polycyclic aromatic hydrocarbons and metals with hyperuricemia risk in Chinese coke oven workers: Mediating roles of oxidative damage. **2023**, 318, 120891 ○
- 49 Effectiveness and Safety of DOACs in Atrial Fibrillation Patients Undergoing Catheter Ablation: Results from the China Atrial Fibrillation (China-AF) Registry. **2022**, 28, 107602962211233 ○
- 48 Association of chronic kidney disease with all-cause mortality in patients hospitalized for atrial fibrillation and impact of clinical and socioeconomic factors on this association. 9, ○
- 47 Reduced left atrial contractile strain with speckle tracking analysis predicts abnormal plasma NTproBNP in an asymptomatic community population. **2022**, 20, ○
- 46 The Effects of Indoxyl Sulfate and Oxidative Stress on the Severity of Peripheral Nerve Dysfunction in Patients with Chronic Kidney Diseases. **2022**, 11, 2350 ○
- 45 Change in Depressive Symptoms During the First Month of Discharge and 1-Year Clinical Outcomes in Patients Hospitalized for Heart Failure. **2022**, 11, ○
- 44 The association between systemic inflammatory response index and new-onset atrial fibrillation in patients with ST-elevated myocardial infarction treated with percutaneous coronary intervention. **2022**, 22, ○
- 43 Associations between plasma metal mixture exposure and risk of hypertension: A cross-sectional study among adults in Shenzhen, China. 10, ○
- 42 Angiotensin-converting enzyme 2 in human plasma and lung tissue. **2023**, 32, 6-15 ○
- 41 Correlation between fatty liver index and estimated glomerular filtration rate in an elderly population in Yangzhou, China. ○
- 40 Associations between long-term exposure to ambient fine particulate pollution with the decline of kidney function and hyperuricemia: a longitudinal cohort study. ○
- 39 Association of lipoprotein(a) with left ventricular hypertrophy in patients with new-onset acute myocardial infarction: A large cross-sectional study. **2023**, 540, 117226 ○
- 38 Association of lipoprotein(a) with coronary severity in patients with new-onset acute myocardial infarction: A large cross-sectional study. **2023**, 540, 117220 ○
- 37 Sex differences in the risk of arterial stiffness among adults with different glycemic status and modifications by age. ○
- 36 The association between liver fibrosis scores and chronic kidney disease. 10, ○
- 35 Long-Term Trajectories of High-Sensitivity C-Reactive Protein Level Among Patients with Acute Heart Failure. Volume 16, 359-371 ○
- 34 Reduced type 3 innate lymphoid cells related to worsening kidney function in renal dysfunction. 153537022211475 ○
- 33 The association among uric acid, microalbumin and estimated glomerular filtration rate in hypertensive patients: a case control study. **2023**, 23, ○

- 32 Long-term exposure to air pollution and increased risk of atrial fibrillation prevalence in China. **2023**, 378, 130-137 ○
- 31 Lipoprotein(a) distribution and its association with carotid arteriopathy in the Chinese population. **2023**, 372, 1-9 ○
- 30 Use of Race in Kidney Function Estimation: Lessons Learned and the Path Toward Health Justice. **2023**, 74, 385-400 ○
- 29 UroAngel: a single-kidney function prediction system based on computed tomography urography using deep learning. ○
- 28 A modified renal risk score for Chinese patients with antineutrophil cytoplasmic antibody-associated vasculitis. **2023**, 21, ○
- 27 Distinct Associations Between Postdischarge Cognitive Change Patterns and 1-year Outcomes in Patients Hospitalized for Heart Failure. **2023**, ○
- 26 The Association Between Metabolic Dysfunction-Associated Fatty Liver Disease and Peripheral Arterial Disease in the Chinese Population. Volume 16, 373-384 ○
- 25 Predictive value of remnant cholesterol level for all-cause mortality in heart failure patients. 10, ○
- 24 Nuclear Magnetic Resonance-Based Metabolomics Approach Revealed the Intervention Effect of Using Complementary and Alternative Medicine (CAM) by CKD Patients. **2023**, 8, 7722-7737 ○
- 23 A Novel and Noninvasive Risk Assessment Score and Its Child-to-Adult Trajectories to Screen Subclinical Renal Damage in Middle Age. **2023**, 10, 257 ○
- 22 Prevalence of chronic kidney disease in a city of Northwestern China: a cross-sectional study. ○
- 21 Insulin Resistance-Varying Associations of Adiposity Indices with Cerebral Perfusion in Older Adults: A Population-Based Study. **2023**, 27, 219-227 ○
- 20 Optimal Glomerular Filtration Rate Equations for Various Age Groups, Disease Conditions and Ethnicities in Asia: A Systematic Review. **2023**, 12, 1822 ○
- 19 Triglyceride-glucose index for predicting repeat revascularization and in-stent restenosis in patients with chronic coronary syndrome undergoing percutaneous coronary intervention. **2023**, 22, ○
- 18 Long-term cumulative high-sensitivity cardiac troponin T and mortality among patients with acute heart failure. ○
- 17 Associations between Plasma Essential Metals Levels and the Risks of All-Cause Mortality and Cardiovascular Disease Mortality among Individuals with Type 2 Diabetes. **2023**, 15, 1198 ○
- 16 Hyponatremia and lower normal serum sodium levels are associated with an increased risk of all-cause death in heart failure patients. ○
- 15 Distinct hyperuricemia trajectories are associated with different risks of incident diabetes: A prospective cohort study. **2023**, ○

- 14 Association of prognostic nutritional index level and diabetes status with the prognosis of coronary artery disease: a cohort study. **2023**, 15, ○
- 13 Quantitative thermal testing as a screening and follow-up tool for diabetic sensorimotor polyneuropathy in patients with type 2 diabetes and prediabetes. 17, ○
- 12 Metabolic dysfunction-associated fatty liver disease increased the risk of subclinical carotid atherosclerosis in China. 14, ○
- 11 Urinary Sirtuin2/Creatinine Ratio Correlates with Severity of Albuminuria and Tubular Injury in Type 2 Diabetes Patients. ○
- 10 Long-Term Exposure to Fine Particulate Constituents and Vascular Damage in a Population with Metabolic Abnormality in China. **2023**, ○
- 9 The new visceral adiposity index outperforms traditional obesity indices as a predictor of subclinical renal damage in Chinese individuals: a cross-sectional study. **2023**, 23, ○
- 8 High absolute neutrophil count with type 2 diabetes is associated with adverse outcome in patients with coronary artery disease: A large-scale cohort study. 14, ○
- 7 The association of apolipoprotein B with chronic kidney disease in the Chinese population. 14, ○
- 6 Association between Circulating Ectodysplasin A and Diabetic Kidney Disease. **2023**, 2023, 1-10 ○
- 5 Association Between Fibrosis-Related Gene Polymorphism and Long-term Allograft Outcome in Renal Transplant Recipients. ○
- 4 Association of the components of ambient fine particulate matter (PM2.5) and chronic kidney disease prevalence in China. **2023**, 339, 117885 ○
- 3 The metabolic score for insulin resistance in the prediction of major adverse cardiovascular events in patients after coronary artery bypass surgery: a multicenter retrospective cohort study. ○
- 2 Diabetes Mellitus with Mild or Moderate Kidney Dysfunction is Associated with Poor Prognosis in Patients with Coronary Artery Disease: A Large-Scale Cohort Study. **2023**, 110693 ○
- 1 Limited non-linear impact of warm ischemia time on renal functional decline after partial nephrectomy: a propensity score-matched study. ○