

IDENTIFICATION OF NOCARDIA CAVIAE (ERIKSON) N

Annals of the New York Academy of Sciences

98, 628-636

DOI: [10.1111/j.1749-6632.1962.tb30585.x](https://doi.org/10.1111/j.1749-6632.1962.tb30585.x)

Citation Report

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | ENZYMATIC STUDIES WITH PATHOGENIC FUNGI*. International Journal of Dermatology, 1965, 4, 72-79. | 0.5 | 35 |
| 2 | Investigations on the antigenic structure of actinomycetales IX. Serological classification of the Nocardiae with the polysaccharide fractions of their cell walls. Mycopathologia, 1965, 25, 173-182. | 1.3 | 7 |
| 3 | Some Criteria for the Recognition of Nocardia madurae (Vincent) Blanchard. Journal of General Microbiology, 1966, 45, 355-364. | 2.3 | 120 |
| 4 | A rapidly fatal infection caused by Nocardia caviae in a dog. The Journal of Pathology and Bacteriology, 1968, 95, 540-546. | 1.1 | 21 |
| 5 | A MEXICAN CASE OF MYCETOMA CAUSED BY STREPTOMYCES SOMALIENSIS. International Journal of Dermatology, 1968, 7, 17-22. | 0.5 | 3 |
| 6 | Studies on aerobic actinomycetes isolated from soil. Medical Mycology, 1968, 6, 140-146. | 0.3 | 19 |
| 7 | A survey of Nocardia asteroides, N. caviae and N. brasiliensis occurring in soil in India. Medical Mycology, 1968, 6, 260-266. | 0.3 | 23 |
| 8 | Studies on aerobic actinomycetes isolated from soil. Medical Mycology, 1968, 6, 192-202. | 0.3 | 1 |
| 9 | Numerical Taxonomy of the Genus Nocardia. Journal of General Microbiology, 1969, 56, 265-287. | 2.3 | 102 |
| 10 | Use of paraffin bait technique in the isolation of Nocardia asteroides from sputum. Mycopathologia, 1970, 40, 363-367. | 1.3 | 7 |
| 11 | Nocardiosis: A review. Mycopathologia, 1970, 40, 193-219. | 1.3 | 13 |
| 12 | Chapter XI Actinomycetes. Methods in Microbiology, 1971, 4, 295-334. | 0.4 | 138 |
| 13 | Nocardia caviae and Nocardia asteroides: Comparative bacteriological and mouse pathogenicity studies. Journal of Comparative Pathology, 1971, 81, 79-87. | 0.1 | 27 |
| 14 | Actinomycetoma pedis caused by Nocardia caviae in India. Medical Mycology, 1972, 10, 19-23. | 0.3 | 18 |
| 15 | Nocardiosis in a child with chronic granulomatous disease. Journal of Pediatrics, 1973, 83, 98-100. | 0.9 | 43 |
| 16 | Taxonomic Criteria for Mycobacteria and Nocardiae. Advances in Applied Microbiology, 1974, 18, 131-190. | 1.3 | 21 |
| 17 | Systemic Nocardia caviae Infection. Chest, 1974, 65, 360-362. | 0.4 | 41 |
| 18 | Mycetoma of the knee due to Nocardia caviae. Medical Mycology, 1975, 13, 170-173. | 0.3 | 8 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Actinomycotic mycetoma due to <i>Streptomyces somaliensis</i> : Report of a case in South Africa. <i>Medical Mycology</i> , 1975, 13, 316-322. | 0.3 | 2 |
| 20 | Distribution of Menaquinones in Actinomycetes and Corynebacteria. <i>Journal of General Microbiology</i> , 1977, 100, 221-230. | 2.3 | 2,030 |
| 21 | Disseminated <i>Nocardia caviae</i> infection. <i>American Journal of Medicine</i> , 1977, 62, 409-412. | 0.6 | 32 |
| 22 | Successful treatment of <i>nocardia asteroides</i> infection with minocycline in kidney transplant patients. <i>The Japanese Journal of Surgery</i> , 1978, 8, 138-144. | 0.2 | 8 |
| 23 | Chemical and Numerical Taxonomy of Strains Received as <i>Gordona aurantiaca</i> . <i>Journal of General Microbiology</i> , 1978, 109, 57-68. | 2.3 | 66 |
| 24 | Numerical Taxonomy of <i>Actinomadura</i> and Related Actinomycetes. <i>Journal of General Microbiology</i> , 1979, 112, 95-111. | 2.3 | 73 |
| 25 | Mycetoma Due to <i>Nocardia Caviae</i> . <i>International Journal of Dermatology</i> , 1980, 19, 260-262. | 0.5 | 22 |
| 26 | Laboratory evaluation of an outbreak of nocardiosis in immunocompromised hosts. <i>American Journal of Medicine</i> , 1981, 71, 928-934. | 0.6 | 81 |
| 27 | Selective Isolation and Enumeration of Actinomycetes using Rifampicin. <i>Journal of Applied Bacteriology</i> , 1981, 51, 289-297. | 1.1 | 56 |
| 28 | Numerical Classification of Some Rhodococci, Corynebacteria and Related Organisms. <i>Microbiology (United Kingdom)</i> , 1982, 128, 731-745. | 0.7 | 44 |
| 29 | Numerical and Chemical Classification of <i>Nocardia amarae</i> . <i>Microbiology (United Kingdom)</i> , 1982, 128, 1283-1297. | 0.7 | 20 |
| 30 | Numerical Classification of Sporoactinomycetes Containing meso-Diaminopimelic Acid in the Cell Wall. <i>Microbiology (United Kingdom)</i> , 1982, 128, 503-527. | 0.7 | 49 |
| 31 | The Mycolic Acids of <i>Mycobacterium chelonae</i> . <i>Microbiology (United Kingdom)</i> , 1982, 128, 817-822. | 0.7 | 29 |
| 32 | Numerical classification of <i>Rhodococcus equi</i> and related actinomycetes. <i>Journal of Applied Bacteriology</i> , 1982, 53, 199-207. | 1.1 | 48 |
| 33 | A simple method for the production of mycobactin, the lipid-soluble siderophore, from mycobacteria. <i>FEMS Microbiology Letters</i> , 1982, 15, 133-136. | 0.7 | 36 |
| 34 | <i>Streptosporangium fragile</i> sp. nov.. <i>International Journal of Systematic Bacteriology</i> , 1983, 33, 364-368. | 2.8 | 16 |
| 35 | Mycolic Acid Patterns of Representative Strains of <i>Mycobacterium fortuitum</i> , " <i>Mycobacterium peregrinum</i> " and <i>Mycobacterium smegmatis</i> . <i>Microbiology (United Kingdom)</i> , 1984, 130, 363-367. | 0.7 | 32 |
| 36 | Numerical taxonomy of cholesterol-degrading soil bacteria. <i>Journal of Applied Bacteriology</i> , 1984, 57, 429-446. | 1.1 | 20 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Aridicins, novel glycopeptide antibiotics. II. Isolation and characterization.. Journal of Antibiotics, 1985, 38, 561-571. | 1.0 | 47 |
| 38 | Cutaneous nocardiosis. Journal of the American Academy of Dermatology, 1985, 13, 125-133. | 0.6 | 123 |
| 39 | Mycolic Acid Patterns of Some Rapidly-Growing Species of Mycobacterium. Zentralblatt Fur Bakteriologie, Mikrobiologie, Und Hygiene Series A, Medical Microbiology, Infectious Diseases, Virology, Parasitology, 1985, 259, 446-460. | 0.5 | 20 |
| 40 | Kibdelins, novel glycopeptide antibiotics. I Discovery, production, and biological evaluation.. Journal of Antibiotics, 1986, 39, 1386-1394. | 1.0 | 39 |
| 41 | Mycobactin and the Competition for Iron between Mycobacterium neoaurum and M. vaccae. Microbiology (United Kingdom), 1986, 132, 839-843. | 0.7 | 1 |
| 42 | Distribution and Application of Mycobactins for the Characterization of Species within the Genus Rhodococcus. Microbiology (United Kingdom), 1986, 132, 853-856. | 0.7 | 15 |
| 43 | Characterisation of rhodococci using peptide hydrolase substrates based on 7-amino-4-methylcoumarin. FEMS Microbiology Letters, 1987, 44, 349-355. | 0.7 | 12 |
| 44 | Mycetoma due to Nocardia caviae. International Journal of Dermatology, 1987, 26, 174-177. | 0.5 | 17 |
| 45 | Histological and microbiological aspects of actinomycetoma cases in Venezuela. Revista Do Instituto De Medicina Tropical De Sao Paulo, 1988, 30, 297-304. | 0.5 | 6 |
| 46 | Saccharopolyspora gregorii and Saccharopolyspora hordei: Two New Actinomycete Species from Fodder. Microbiology (United Kingdom), 1989, 135, 2125-2139. | 0.7 | 25 |
| 47 | Primary subcutaneous abscess caused by Nocardia otitidiscaviarum. Journal of the American Academy of Dermatology, 1989, 21, 137-139. | 0.6 | 15 |
| 48 | Amycolatopsis methanolica sp. nov., a Facultatively Methylotrophic Actinomycete. International Journal of Systematic Bacteriology, 1990, 40, 194-204. | 2.8 | 77 |
| 49 | Classification and Identification of Rhodococci. Zentralblatt Fur Bakteriologie: International Journal of Medical Microbiology, 1990, 274, 299-315. | 0.5 | 39 |
| 50 | Bovine farcy: A clinico-pathological study of the disease and its aetiological agent. Journal of Comparative Pathology, 1991, 105, 287-301. | 0.1 | 16 |
| 51 | Positive selection of antibiotic-producing soil isolates. Journal of General Microbiology, 1991, 137, 2321-2329. | 2.3 | 71 |
| 52 | Rapid identification of streptomycetes by artificial neural network analysis of pyrolysis mass spectra. FEMS Microbiology Letters, 1993, 114, 115-119. | 0.7 | 27 |
| 53 | Lymphocutaneous nocardiosis caused by Nocardia caviae: The first case report from Asia. Journal of the American Academy of Dermatology, 1993, 29, 639-641. | 0.6 | 20 |
| 54 | Thin-layer chromatographic analysis of glycolipids and mycolic acids from Mycobacterium farcinogenes, Mycobacterium senegalense and related taxa. Zentralblatt Fur Bakteriologie: International Journal of Medical Microbiology, 1993, 279, 354-367. | 0.5 | 28 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | A simple chemical test to distinguish mycobacteria from other mycolic-acid-containing actinomycetes. <i>Journal of General Microbiology</i> , 1993, 139, 2203-2213. | 2.3 | 40 |
| 56 | Biosystematics and diversity amongst novel carboxydophilic actinomycetes. <i>Antonie Van Leeuwenhoek</i> , 1994, 64, 325-340. | 0.7 | 86 |
| 57 | Numerical and chemical classification of <i>Streptosporangium</i> and some related actinomycetes. <i>Antonie Van Leeuwenhoek</i> , 1994, 64, 387-429. | 0.7 | 12 |
| 58 | Differential elimination of enteric bacteria by protists in a freshwater system. <i>Journal of Applied Bacteriology</i> , 1994, 77, 476-483. | 1.1 | 35 |
| 59 | Transfer of <i>Nocardia amarae</i> Lechevalier and Lechevalier 1974 to the genus <i>Gordona</i> as <i>Gordona amarae</i> comb. nov.. <i>Letters in Applied Microbiology</i> , 1994, 19, 401-405. | 1.0 | 14 |
| 60 | Rapid characterisation and identification of mycobacteria using fluorogenic enzyme tests. <i>Zentralblatt Fur Bakteriologie: International Journal of Medical Microbiology</i> , 1994, 280, 476-487. | 0.5 | 11 |
| 61 | Primary Lymphocutaneous Nocardiosis Due to <i>Nocardia otitidiscaviarum</i> : The First Case Report from Japan. <i>Journal of Dermatology</i> , 1995, 22, 344-347. | 0.6 | 12 |
| 62 | Rapid Growth and Increased Biomass Yield of <i>Mycobacterium farcinogenes</i> and Some Related Taxa in Broth and Agar Media. <i>Zoonoses and Public Health</i> , 1995, 42, 397-404. | 1.4 | 3 |
| 63 | Primary Cutaneous <i>Nocardia otitidiscaviarum</i> Infection: Case Report and Review. <i>Clinical Infectious Diseases</i> , 1995, 20, 1266-1270. | 2.9 | 44 |
| 64 | Actinomycete diversity associated with foaming in activated sludge plants. <i>Journal of Industrial Microbiology</i> , 1996, 17, 268-280. | 0.9 | 30 |
| 65 | <i>Nocardia otitidiscaviarum</i> infection of a traumatic skin wound. <i>European Journal of Clinical Microbiology and Infectious Diseases</i> , 1997, 16, 383-384. | 1.3 | 15 |
| 66 | Novel rhodococci and other mycolate actinomycetes from the deep sea. <i>Antonie Van Leeuwenhoek</i> , 1998, 74, 27-40. | 0.7 | 102 |
| 67 | In situ detection of rhodococci associated with activated sludge foams. <i>Antonie Van Leeuwenhoek</i> , 1998, 74, 41-48. | 0.7 | 17 |
| 68 | Taxonomy and biotransformation activities of some deep-sea actinomycetes. <i>Extremophiles</i> , 1998, 2, 269-277. | 0.9 | 85 |
| 69 | <i>Saccharopolyspora spinosporotrichia</i> sp. nov., a novel actinomycete from soil. <i>International Journal of Systematic Bacteriology</i> , 1998, 48, 53-58. | 2.8 | 69 |
| 70 | <i>Amycolatopsis thermoflava</i> sp. nov., a novel soil actinomycete from Hainan Island, China. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 1999, 49, 1369-1373. | 0.8 | 44 |
| 71 | <i>Nocardia salmonicida</i> nom. rev., a fish pathogen. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 1999, 49, 833-837. | 0.8 | 85 |
| 72 | The <i>Nocardia salmonicida</i> clade, including descriptions of <i>Nocardia cummidelens</i> sp. nov., <i>Nocardia fluminea</i> sp. nov. and <i>Nocardia soli</i> sp. nov. <i>Antonie Van Leeuwenhoek</i> , 2000, 78, 367-377. | 0.7 | 69 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Genomic and phenomic differentiation of <i>Rhodococcus equi</i> and related strains. <i>Antonie Van Leeuwenhoek</i> , 2000, 78, 331-340. | 0.7 | 39 |
| 74 | Quantitative Use of Fluorescent In Situ Hybridization To Examine Relationships between Mycolic Acid-Containing Actinomycetes and Foaming in Activated Sludge Plants. <i>Applied and Environmental Microbiology</i> , 2000, 66, 1158-1166. | 1.4 | 148 |
| 75 | <i>Gordonia namibiensis</i> sp. nov., a Novel Nitrile Metabolising Actinomycete Recovered from an African Sand. <i>Systematic and Applied Microbiology</i> , 2001, 24, 510-515. | 1.2 | 35 |
| 76 | <i>Nocardia africana</i> sp. nov., a New Pathogen Isolated from Patients with Pulmonary Infections. <i>Journal of Clinical Microbiology</i> , 2001, 39, 625-630. | 1.8 | 74 |
| 77 | <i>Amycolatopsis rubida</i> sp. nov., a new <i>Amycolatopsis</i> species from soil.. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2001, 51, 1093-1097. | 0.8 | 28 |
| 78 | PRIMARY LYMPHOCUTANEOUS NOCARDIOSIS CAUSED BY <i>NOCARDIA OTITIDISCAVIARUM</i> . <i>Hand Surgery</i> , 2002, 07, 285-287. | 0.6 | 10 |
| 79 | <i>Nocardia Otitidiscaviarum</i> : An Unusual <i>Nocardia</i> Species Causing a Primary Lymphocutaneous Infectious Process in a Mildly Immunosuppressed Patient. <i>Skinmed</i> , 2002, 1, 62-64. | 0.0 | 6 |
| 80 | Dispelling the "Nocardia amarae" myth: a phylogenetic and phenotypic study of mycolic acid-containing actinomycetes isolated from activated sludge foam. <i>Water Science and Technology</i> , 2002, 46, 81-90. | 1.2 | 27 |
| 81 | Differentiation of <i>Nocardia</i> Species by PCR-Randomly Amplified Polymorphic DNA Fingerprinting. <i>Systematic and Applied Microbiology</i> , 2002, 25, 60-67. | 1.2 | 15 |
| 82 | Discrimination and taxonomy of geographically diverse strains of nitrile-metabolizing actinomycetes using chemometric and molecular sequencing techniques. <i>Environmental Microbiology</i> , 2002, 4, 262-276. | 1.8 | 38 |
| 83 | Dereplication for biotechnology screening: PyMS analysis and PCR-RFLP-SSCP (PRS) profiling of 16S rRNA genes of marine and terrestrial actinomycetes. <i>Applied Microbiology and Biotechnology</i> , 2002, 58, 77-83. | 1.7 | 36 |
| 84 | <i>Streptomyces yatensis</i> sp. nov., a novel bioactive streptomycete isolated from a New-Caledonian ultramafic soil. <i>Antonie Van Leeuwenhoek</i> , 2003, 83, 21-26. | 0.7 | 36 |
| 85 | <i>Gordonia sinesedis</i> sp. nov., a novel soil isolate. <i>Antonie Van Leeuwenhoek</i> , 2003, 83, 75-80. | 0.7 | 20 |
| 86 | <i>Tsukamurella spumae</i> sp. nov., A Novel Actinomycete Associated with Foaming in Activated Sludge Plants. <i>Systematic and Applied Microbiology</i> , 2003, 26, 367-375. | 1.2 | 41 |
| 87 | <i>Actinomadura mexicana</i> sp. nov. and <i>Actinomadura meyerii</i> sp. nov., Two Novel Soil Sporoactinomycetes. <i>Systematic and Applied Microbiology</i> , 2003, 26, 511-517. | 1.2 | 28 |
| 88 | <i>Nocardia caishijiensis</i> sp. nov., a novel soil actinomycete. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2003, 53, 999-1004. | 0.8 | 35 |
| 89 | <i>Williamsia maris</i> sp. nov., a novel actinomycete isolated from the Sea of Japan. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2004, 54, 191-194. | 0.8 | 51 |
| 90 | <i>Nocardia neocaledoniensis</i> sp. nov., a novel actinomycete isolated from a New-Caledonian brown hypermagnesian ultramafic soil. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2004, 54, 599-603. | 0.8 | 32 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | <i>Tsukamurella pseudospumae</i> sp. nov., a novel actinomycete isolated from activated sludge foam. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2004, 54, 1209-1212. | 0.8 | 71 |
| 92 | <i>Mycobacterium psychrotolerans</i> sp. nov., isolated from pond water near a uranium mine. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2004, 54, 1459-1463. | 0.8 | 29 |
| 93 | <i>Rhodococcus aetherivorans</i> sp. nov., A New Species that Contains Methyl t-butyl Ether-Degrading Actinomycetes. <i>Systematic and Applied Microbiology</i> , 2004, 27, 61-65. | 1.2 | 58 |
| 94 | Identification and Expression in <i>E. coli</i> of Novel Nitrile Hydratases from the Metagenome. <i>Engineering in Life Sciences</i> , 2004, 4, 557-562. | 2.0 | 33 |
| 95 | Diversity of cultivable actinobacteria in geographically widespread marine sediments. <i>Antonie Van Leeuwenhoek</i> , 2005, 87, 11-18. | 0.7 | 172 |
| 96 | <i>Amycolatopsis plumensis</i> sp. nov., a novel bioactive actinomycete isolated from a New-Caledonian brown hypermagnesian ultramafic soil. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2005, 55, 2057-2061. | 0.8 | 30 |
| 97 | MICROBIAL DIAGNOSIS IN A GENERAL HOSPITAL. <i>Annals of the New York Academy of Sciences</i> , 2006, 98, 647-669. | 1.8 | 9 |
| 98 | Exploration of <i>Amycolatopsis</i> diversity in soil using genus-specific primers and novel selective media. <i>Systematic and Applied Microbiology</i> , 2006, 29, 557-569. | 1.2 | 115 |
| 99 | Clinical and Laboratory Features of the <i>Nocardia</i> spp. Based on Current Molecular Taxonomy. <i>Clinical Microbiology Reviews</i> , 2006, 19, 259-282. | 5.7 | 910 |
| 100 | <i>Dermacoccus abyssi</i> sp. nov., a piezotolerant actinomycete isolated from the Mariana Trench. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2006, 56, 1233-1237. | 0.8 | 62 |
| 101 | <i>Gordonia soli</i> sp. nov., a novel actinomycete isolated from soil. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2006, 56, 2597-2601. | 0.8 | 32 |
| 102 | <i>Williamsia marianensis</i> sp. nov., a novel actinomycete isolated from the Mariana Trench. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2006, 56, 1123-1126. | 0.8 | 37 |
| 103 | <i>Dermacoccus barathri</i> sp. nov. and <i>Dermacoccus profundus</i> sp. nov., novel actinomycetes isolated from deep-sea mud of the Mariana Trench. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2006, 56, 2303-2307. | 0.8 | 45 |
| 104 | <i>Streptomyces synnematoformans</i> sp. nov., a novel actinomycete isolated from a sand dune soil in Egypt. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2007, 57, 2009-2013. | 0.8 | 22 |
| 105 | <i>Nocardia amamiensis</i> sp. nov., isolated from a sugar-cane field in Japan. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2007, 57, 1599-1602. | 0.8 | 19 |
| 106 | <i>Agrococcus casei</i> sp. nov., isolated from the surfaces of smear-ripened cheeses. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2007, 57, 92-97. | 0.8 | 40 |
| 107 | <i>Nonomuraea aegyptia</i> sp. nov., a novel actinomycete isolated from a sand dune. <i>Antonie Van Leeuwenhoek</i> , 2007, 92, 165-171. | 0.7 | 16 |
| 108 | A new preferential medium for enumeration and isolation of desert actinomycetes. <i>World Journal of Microbiology and Biotechnology</i> , 2008, 24, 1547-1552. | 1.7 | 31 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | <i>Acinetobacter soli</i> sp. nov., isolated from forest soil. <i>Journal of Microbiology</i> , 2008, 46, 396-401. | 1.3 | 67 |
| 110 | <i>Streptomyces sudanensis</i> sp. nov., a new pathogen isolated from patients with actinomycetoma. <i>Antonie Van Leeuwenhoek</i> , 2008, 93, 305-313. | 0.7 | 65 |
| 111 | <i>Dietzia papillomatosis</i> sp. nov., a novel actinomycete isolated from the skin of an immunocompetent patient with confluent and reticulated papillomatosis. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2008, 58, 68-72. | 0.8 | 68 |
| 112 | Actinomycetoma Caused by Organisms of the <i>Nocardia Asteroides</i> * Complex and Closely Related Strains. <i>Mycoses</i> , 1978, 21, 109-121. | 1.8 | 12 |
| 113 | Nocardiosis During Steroid Treatment of Autoimmune Haemolytic Anaemia. <i>Acta Medica Scandinavica</i> , 1965, 178, 221-231. | 0.0 | 13 |
| 114 | Actinobacterial diversity from marine sediments collected in Mexico. <i>Antonie Van Leeuwenhoek</i> , 2009, 95, 111-120. | 0.7 | 79 |
| 115 | Diversity of culturable actinomycetes in hyper-arid soils of the Atacama Desert, Chile. <i>Antonie Van Leeuwenhoek</i> , 2009, 95, 121-133. | 0.7 | 185 |
| 116 | Study of the diversity of culturable actinomycetes in the North Pacific and Caribbean coasts of Costa Rica. <i>Antonie Van Leeuwenhoek</i> , 2009, 96, 71-78. | 0.7 | 18 |
| 117 | Oxidation of aliphatic, branched chain, and aromatic hydrocarbons by <i>Nocardia cyriacigeorgica</i> isolated from oil-polluted sand samples collected in the Saudi Arabian Desert. <i>Journal of Basic Microbiology</i> , 2010, 50, 241-253. | 1.8 | 41 |
| 118 | Mycetoma caused by <i>Nocardia caviae</i> in the first Brazilian patient. <i>International Journal of Dermatology</i> , 2010, 49, 56-58. | 0.5 | 9 |
| 119 | <i>Pseudoclavibacter chungangensis</i> sp. nov., isolated from activated sludge. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2010, 60, 1672-1677. | 0.8 | 18 |
| 120 | <i>Lechevalieria atacamensis</i> sp. nov., <i>Lechevalieria deserti</i> sp. nov. and <i>Lechevalieria roselyniae</i> sp. nov., isolated from hyperarid soils. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2010, 60, 296-300. | 0.8 | 36 |
| 121 | <i>Bacillus chungangensis</i> sp. nov., a halophilic species isolated from sea sand. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2010, 60, 1349-1352. | 0.8 | 11 |
| 122 | <i>Nocardiopsis nikkonensis</i> sp. nov., isolated from a compost sample. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2010, 60, 2967-2971. | 0.8 | 14 |
| 123 | <i>Williamsia faeni</i> sp. nov., an actinomycete isolated from a hay meadow. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2010, 60, 2548-2551. | 0.8 | 20 |
| 124 | <i>Dietzia timorensis</i> sp. nov., isolated from soil. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2010, 60, 451-454. | 0.8 | 32 |
| 125 | Dermacozines, a new phenazine family from deep-sea dermacocci isolated from a Mariana Trench sediment. <i>Organic and Biomolecular Chemistry</i> , 2010, 8, 2352. | 1.5 | 123 |
| 126 | <i>Altererythrobacter namhicola</i> sp. nov. and <i>Altererythrobacter aestuarii</i> sp. nov., isolated from seawater. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2011, 61, 709-715. | 0.8 | 51 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 127 | <i>Paenibacillus puldeungensis</i> sp. nov., isolated from a grassy sandbank. International Journal of Systematic and Evolutionary Microbiology, 2011, 61, 670-673. | 0.8 | 23 |
| 128 | <i>Paenibacillus chungangensis</i> sp. nov., isolated from a tidal-flat sediment. International Journal of Systematic and Evolutionary Microbiology, 2011, 61, 281-285. | 0.8 | 17 |
| 129 | Diverse Metabolic Profiles of a <i>Streptomyces</i> Strain Isolated from a Hyper-arid Environment. Journal of Natural Products, 2011, 74, 1965-1971. | 1.5 | 129 |
| 130 | <i>Actinomycetospora iriomotensis</i> sp. nov., a novel actinomycete isolated from a lichen sample. Journal of Antibiotics, 2011, 64, 289-292. | 1.0 | 36 |
| 131 | <i>Nocardioides iriomotensis</i> sp. nov., an actinobacterium isolated from forest soil. International Journal of Systematic and Evolutionary Microbiology, 2011, 61, 2205-2209. | 0.8 | 17 |
| 132 | <i>Actinomycetospora rishiriensis</i> sp. nov., isolated from a lichen. International Journal of Systematic and Evolutionary Microbiology, 2011, 61, 2621-2625. | 0.8 | 26 |
| 133 | <i>Serinicoccus chungangensis</i> sp. nov., isolated from tidal flat sediment, and emended description of the genus <i>Serinicoccus</i> . International Journal of Systematic and Evolutionary Microbiology, 2011, 61, 1299-1303. | 0.8 | 20 |
| 134 | <i>Amycolatopsis thermophila</i> sp. nov. and <i>Amycolatopsis viridis</i> sp. nov., thermophilic actinomycetes isolated from arid soil. International Journal of Systematic and Evolutionary Microbiology, 2012, 62, 168-172. | 0.8 | 25 |
| 135 | <i>Chungangia koreensis</i> gen. nov., sp. nov., isolated from marine sediment. International Journal of Systematic and Evolutionary Microbiology, 2012, 62, 1914-1920. | 0.8 | 11 |
| 136 | <i>Actinomadura geliboluensis</i> sp. nov., isolated from soil. International Journal of Systematic and Evolutionary Microbiology, 2012, 62, 2011-2017. | 0.8 | 13 |
| 137 | <i>Actinoplanes rishiriensis</i> sp. nov., a novel motile actinomycete isolated by rehydration and centrifugation method. Journal of Antibiotics, 2012, 65, 249-253. | 1.0 | 10 |
| 138 | Phylum XXVI. Actinobacteria phyl. nov., 2012, , 33-2028. | | 58 |
| 139 | <i>Amycolatopsis granulosa</i> sp. nov., <i>Amycolatopsis ruanii</i> sp. nov. and <i>Amycolatopsis thermalba</i> sp. nov., thermophilic actinomycetes isolated from arid soils. International Journal of Systematic and Evolutionary Microbiology, 2012, 62, 1245-1251. | 0.8 | 23 |
| 140 | <i>Microbispora thailandensis</i> sp. nov., an actinomycete isolated from cave soil. Journal of Antibiotics, 2012, 65, 491-494. | 1.0 | 17 |
| 141 | <i>Gulosibacter chungangensis</i> sp. nov., an actinomycete isolated from a marine sediment, and emended description of the genus <i>Gulosibacter</i> . International Journal of Systematic and Evolutionary Microbiology, 2012, 62, 1055-1060. | 0.8 | 19 |
| 142 | <i>Verrucospora wenchangensis</i> sp. nov., isolated from mangrove soil. Antonie Van Leeuwenhoek, 2012, 102, 1-7. | 0.7 | 78 |
| 143 | <i>Amycolatopsis bartoniae</i> sp. nov. and <i>Amycolatopsis bullii</i> sp. nov., mesophilic actinomycetes isolated from arid Australian soils. Antonie Van Leeuwenhoek, 2012, 102, 91-98. | 0.7 | 22 |
| 144 | <i>Streptosporangium anatoliense</i> sp. nov., isolated from soil in Turkey. Antonie Van Leeuwenhoek, 2012, 102, 269-276. | 0.7 | 12 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 145 | <i>Verrucosipora maris</i> sp. nov., a novel deep-sea actinomycete isolated from a marine sediment which produces abyssomicins. <i>Antonie Van Leeuwenhoek</i> , 2012, 101, 185-193. | 0.7 | 63 |
| 146 | <i>Rhodococcus jostii</i> : a home for <i>Rhodococcus</i> strain RHA1. <i>Antonie Van Leeuwenhoek</i> , 2013, 104, 435-440. | 0.7 | 12 |
| 147 | <i>Oceanobacillus chungangensis</i> sp. nov., isolated from a sand dune. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2013, 63, 3666-3671. | 0.8 | 17 |
| 148 | <i>Methylobacterium tarhaniae</i> sp. nov., isolated from arid soil. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2013, 63, 2823-2828. | 0.8 | 24 |
| 149 | Characterisation of the first actinobacterial group isolated from a Mexican extremophile environment. <i>Antonie Van Leeuwenhoek</i> , 2013, 104, 63-70. | 0.7 | 19 |
| 150 | <i>Streptacidiphilus hamsterleyensis</i> sp. nov., isolated from a spruce forest soil. <i>Antonie Van Leeuwenhoek</i> , 2013, 104, 965-972. | 0.7 | 13 |
| 151 | <i>Bacillus songklensis</i> sp. nov., isolated from soil. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2013, 63, 4189-4195. | 0.8 | 13 |
| 152 | <i>Haloferula chungangensis</i> sp. nov., isolated from marine sediment. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2013, 63, 19-24. | 0.8 | 9 |
| 153 | <i>Sphingopyxis rigui</i> sp. nov. and <i>Sphingopyxis wooonensis</i> sp. nov., isolated from wetland freshwater, and emended description of the genus <i>Sphingopyxis</i> . <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2013, 63, 1297-1303. | 0.8 | 35 |
| 154 | <i>Nonomuraea jabiensis</i> sp. nov., isolated from arid soil. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2013, 63, 212-218. | 0.8 | 18 |
| 155 | <i>Prescottia equi</i> gen. nov., comb. nov.: a new home for an old pathogen. <i>Antonie Van Leeuwenhoek</i> , 2013, 103, 655-671. | 0.7 | 35 |
| 156 | <i>Nocardia aciditolerans</i> sp. nov., isolated from a spruce forest soil. <i>Antonie Van Leeuwenhoek</i> , 2013, 103, 1079-1088. | 0.7 | 36 |
| 157 | <i>Algoriphagus chungangensis</i> sp. nov., isolated from a tidal flat sediment. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2013, 63, 648-653. | 0.8 | 23 |
| 158 | <i>Maribacter chungangensis</i> sp. nov., isolated from a green seaweed, and emended descriptions of the genus <i>Maribacter</i> and <i>Maribacter arcticus</i> . <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2013, 63, 2553-2558. | 0.8 | 35 |
| 159 | <i>Mesoflavibacter aestuarii</i> sp. nov., a zeaxanthin-producing marine bacterium isolated from seawater. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2014, 64, 1932-1937. | 0.8 | 19 |
| 160 | <i>Sneathiella chungangensis</i> sp. nov., isolated from a marine sand, and emended description of the genus <i>Sneathiella</i> . <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2014, 64, 1468-1472. | 0.8 | 13 |
| 161 | Functional diversity of <i>Nocardia</i> in metabolism. <i>Environmental Microbiology</i> , 2014, 16, 29-48. | 1.8 | 37 |
| 162 | <i>Nocardia kroppenstedtii</i> sp. nov., an actinomycete isolated from a lung transplant patient with a pulmonary infection. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2014, 64, 751-754. | 0.8 | 17 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 163 | <i>Streptomyces hokutonensis</i> sp. nov., a novel actinomycete isolated from the strawberry root rhizosphere. <i>Journal of Antibiotics</i> , 2014, 67, 465-470. | 1.0 | 17 |
| 164 | <i>Paenibacillus doosanensis</i> sp. nov., isolated from soil. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2014, 64, 1271-1277. | 0.8 | 15 |
| 165 | Characterization of a thermotolerant laccase produced by <i>Streptomyces</i> sp. SB086. <i>Annals of Microbiology</i> , 2014, 64, 1363-1369. | 1.1 | 22 |
| 166 | Biodecolorization of a food azo dye by the deep sea <i>Dermacoccus abyssi</i> MT1.1T strain from the Mariana Trench. <i>Journal of Environmental Management</i> , 2014, 132, 155-164. | 3.8 | 20 |
| 176 | <i>Oceanobacillus arenosus</i> sp. nov., a moderately halophilic bacterium isolated from marine sand. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2015, 65, 2943-2948. | 0.8 | 12 |
| 177 | Culturable bioactive actinomycetes from the Great Indian Thar Desert. <i>Annals of Microbiology</i> , 2015, 65, 1901-1914. | 1.1 | 26 |
| 178 | <i>Amycolatopsis rhabdoformis</i> sp. nov., an actinomycete isolated from a tropical forest soil. <i>International Journal of Systematic and Evolutionary Microbiology</i> , 2015, 65, 1786-1793. | 0.8 | 8 |
| 179 | <i>Eudoraea chungangensis</i> sp. nov., isolated from an aquafarm waste water sludge. <i>Antonie Van Leeuwenhoek</i> , 2015, 107, 1009-1015. | 0.7 | 6 |
| 180 | <i>Kangiella chungangensis</i> sp. nov. isolated from a marine sand. <i>Antonie Van Leeuwenhoek</i> , 2015, 107, 1291-1298. | 0.7 | 40 |
| 181 | Mass spectrometric approaches for the identification of anthracycline analogs produced by actinobacteria. <i>Journal of Mass Spectrometry</i> , 2016, 51, 437-445. | 0.7 | 10 |
| 182 | DNA-EB in agarose gel assay: a simple methodology in the search for DNA-binders in crude extracts from actinomycetes. <i>Analytical Methods</i> , 2016, 8, 2653-2659. | 1.3 | 2 |
| 183 | Are There Any Other Compounds Isolated From <i>Dermacoccus</i> spp at All?. <i>Current Microbiology</i> , 2017, 74, 132-144. | 1.0 | 12 |
| 184 | <i>Cellulosimicrobium arenosum</i> sp. nov., Isolated from Marine Sediment Sand. <i>Current Microbiology</i> , 2018, 75, 901-906. | 1.0 | 12 |
| 185 | Diversity of culturable nocardioform actinomycetes from wastewater treatment plants in Spain and their role in the biodegradability of aromatic compounds. <i>Environmental Technology (United Kingdom)</i> 39(10):1147-1156, 2018 | 1.0 | 10 |
| 186 | <i>Rhodococcus psychrotolerans</i> sp. nov., isolated from rhizosphere of <i>Deschampsia antarctica</i> . <i>Antonie Van Leeuwenhoek</i> , 2018, 111, 629-636. | 0.7 | 16 |
| 188 | Bioactive Compounds from <i>Nocardia</i> : Biosynthesis and Production. <i>Environmental Chemistry for A Sustainable World</i> , 2019, , 49-74. | 0.3 | 1 |
| 189 | Pradimicin-IRD from <i>Amycolatopsis</i> sp. IRD-009 and its antimicrobial and cytotoxic activities. <i>Natural Product Research</i> , 2019, 33, 1713-1720. | 1.0 | 15 |
| 190 | Polyphasic taxonomic analysis of <i>Nitratireductor arenosus</i> sp. nov., isolated from sea sand. <i>FEMS Microbiology Letters</i> , 2020, 367, . | 0.7 | 2 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 191 | The Families Dietziaceae, Gordoniaceae, Nocardiaceae and Tsukamurellaceae. , 2006, , 843-888. | | 45 |
| 192 | The Family Nocardiaceae. , 2014, , 595-650. | | 47 |
| 193 | Mycetoma. , 1971, , 589-613. | | 1 |
| 194 | Mycetoma. , 1971, , 589-613. | | 2 |
| 195 | Saccharopolyspora flava sp. nov. and Saccharopolyspora thermophila sp.nov., novel actinomycetes from soil.. International Journal of Systematic and Evolutionary Microbiology, 2001, 51, 319-325. | 0.8 | 66 |
| 196 | Differentiation of Nocardioform Actinomycetes by Lysozyme Sensitivity. Journal of General Microbiology, 1978, 109, 381-384. | 2.3 | 10 |
| 197 | Hymenobacter wooonensis sp. nov., isolated from wetland freshwater. International Journal of Systematic and Evolutionary Microbiology, 2015, 65, 1871-1876. | 0.8 | 14 |
| 198 | Palleronia soli sp. nov., isolated from a soil sample on reclaimed tidal land, and emended description of the genus Palleronia. International Journal of Systematic and Evolutionary Microbiology, 2015, 65, 2516-2521. | 0.8 | 10 |
| 199 | Limibacillus halophilus gen. nov., sp. nov., a moderately halophilic bacterium in the family Rhodospirillaceae isolated from reclaimed land. International Journal of Systematic and Evolutionary Microbiology, 2015, 65, 3155-3161. | 0.8 | 14 |
| 200 | Sphingobium barthaii sp. nov., a high molecular weight polycyclic aromatic hydrocarbon-degrading bacterium isolated from cattle pasture soil. International Journal of Systematic and Evolutionary Microbiology, 2015, 65, 2919-2924. | 0.8 | 28 |
| 201 | Rhodococcus gordoniae sp. nov., an actinomycete isolated from clinical material and phenol-contaminated soil. International Journal of Systematic and Evolutionary Microbiology, 2004, 54, 407-411. | 0.8 | 32 |
| 202 | Marinivirga aestuarii gen. nov., sp. nov., a member of the family Flavobacteriaceae , isolated from marine environments, and emended descriptions of the genera Hyunsoonleella , Jejuia and Pontirhabdus and the species Hyunso. International Journal of Systematic and Evolutionary Microbiology, 2013, 63, 1524-1531. | 0.8 | 34 |
| 203 | Salegentibacter chungangensis sp. nov., isolated from a sea sand and emended description of the genus Salegentibacter. International Journal of Systematic and Evolutionary Microbiology, 2014, 64, 1514-1519. | 0.8 | 13 |
| 204 | Roseovarius aquimarinus sp. nov., a slightly halophilic bacterium isolated from seawater. International Journal of Systematic and Evolutionary Microbiology, 2015, 65, 4514-4520. | 0.8 | 17 |
| 205 | Salinimicrobium soli sp. nov., isolated from soil of reclaimed land. International Journal of Systematic and Evolutionary Microbiology, 2016, 66, 462-467. | 0.8 | 14 |
| 206 | Marimicrobium arenosum gen. nov., sp. nov., a moderately halophilic bacterium isolated from sea sand. International Journal of Systematic and Evolutionary Microbiology, 2016, 66, 856-861. | 0.8 | 15 |
| 207 | Hymenobacter coalescens sp. nov., isolated from wetland freshwater. International Journal of Systematic and Evolutionary Microbiology, 2016, 66, 3546-3551. | 0.8 | 14 |
| 208 | Maribacter arenosus sp. nov., isolated from marine sediment. International Journal of Systematic and Evolutionary Microbiology, 2016, 66, 4826-4831. | 0.8 | 17 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 209 | <i>Altererythrobacter sediminis</i> sp. nov., isolated from lagoon sediments. International Journal of Systematic and Evolutionary Microbiology, 2016, 66, 5424-5429. | 0.8 | 19 |
| 210 | <i>Tropicimonas arenosa</i> sp. nov., isolated from marine sand. International Journal of Systematic and Evolutionary Microbiology, 2016, 66, 5514-5518. | 0.8 | 10 |
| 211 | <i>Marimonas arenosa</i> gen. nov., sp. nov., isolated from sea sand. International Journal of Systematic and Evolutionary Microbiology, 2017, 67, 121-126. | 0.8 | 12 |
| 212 | <i>Nonlabens halophilus</i> sp. nov., isolated from reclaimed land. International Journal of Systematic and Evolutionary Microbiology, 2017, 67, 138-143. | 0.8 | 6 |
| 213 | <i>Thioclava arenosa</i> sp. nov., isolated from sea sand. International Journal of Systematic and Evolutionary Microbiology, 2017, 67, 1735-1739. | 0.8 | 9 |
| 214 | <i>Tessaracoccus arenae</i> sp. nov., isolated from sea sand. International Journal of Systematic and Evolutionary Microbiology, 2017, 67, 2008-2013. | 0.8 | 15 |
| 215 | <i>Hymenobacter tenuis</i> sp. nov., isolated from wastewater of an acidic water neutralization facility. International Journal of Systematic and Evolutionary Microbiology, 2017, 67, 2171-2177. | 0.8 | 7 |
| 216 | <i>Limibaculum halophilum</i> gen. nov., sp. nov., a new member of the family Rhodobacteraceae. International Journal of Systematic and Evolutionary Microbiology, 2017, 67, 3812-3818. | 0.8 | 17 |
| 217 | <i>Arthrobacter paludis</i> sp. nov., isolated from a marsh. International Journal of Systematic and Evolutionary Microbiology, 2018, 68, 47-51. | 0.8 | 13 |
| 218 | <i>Arenimonas halophila</i> sp. nov., isolated from soil. International Journal of Systematic and Evolutionary Microbiology, 2018, 68, 2188-2193. | 0.8 | 8 |
| 219 | <i>Arenibacillus arenosus</i> gen. nov., sp. nov., a member of the family Rhodobacteraceae isolated from sea sand. International Journal of Systematic and Evolutionary Microbiology, 2019, 69, 153-158. | 0.8 | 13 |
| 220 | <i>Arenibacterium halophilum</i> gen. nov., sp. nov., a halotolerant bacterium in the family Rhodobacteraceae isolated from a coastal sand dune. International Journal of Systematic and Evolutionary Microbiology, 2020, 70, 6323-6330. | 0.8 | 10 |
| 222 | <i>Nocardia, Rhodococcus, Gordonia, Actinomadura, Streptomyces</i>, and Other Aerobic Actinomycetes. , 0, , 504-535. | | 20 |
| 223 | Laboratory Identification of Clinically Important Aerobic Actinomycetes. Applied Microbiology, 1973, 25, 665-681. | 0.6 | 139 |
| 224 | <i>Nocardia caviae</i> : a Report of 13 New Isolations with Clinical Correlation. Applied Microbiology, 1974, 28, 193-198. | 0.6 | 31 |
| 225 | Role of L-forms of <i>Nocardia caviae</i> in the development of chronic mycetomas in normal and immunodeficient murine models. Infection and Immunity, 1981, 33, 893-907. | 1.0 | 38 |
| 226 | Use of CO ₂ for More Rapid Growth of the <i>Nocardia</i> Species. Journal of Clinical Microbiology, 1976, 3, 463-464. | 1.8 | 1 |
| 227 | Nocardial infection in cultured yellowtails (<i>Seriola quinqueradiata</i> and <i>S.purpurascens</i>)”II. Fish Pathology, 1968, 3, 24-33. | 0.4 | 26 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 228 | Marine Sediment Recovered <i>Salinispora</i> sp. Inhibits the Growth of Emerging Bacterial Pathogens and other Multi-Drug-Resistant Bacteria. Polish Journal of Microbiology, 2020, 69, 321-330. | 0.6 | 6 |
| 229 | Unexpected Properties of Micromonosporae from Marine Origin. Advances in Microbiology, 2015, 05, 452-456. | 0.3 | 10 |
| 230 | Transfer of <i>Tsukamurella wratislaviensis</i> Goodfellow et al. 1995 to the genus <i>Rhodococcus</i> as <i>Rhodococcus wratislaviensis</i> comb. nov... International Journal of Systematic and Evolutionary Microbiology, 2002, 52, 749-755. | 0.8 | 22 |
| 232 | The Family Dermacoccaceae. , 2014, , 301-315. | | 0 |
| 233 | The Family Tsukamurellaceae. , 2014, , 1051-1061. | | 0 |
| 234 | Taxonomy and Biotransformation Activities of Deep-Sea Actinomycetes. , 1999, , 39-54. | | 0 |
| 235 | Characterization of Actinomycetes from Smear Ripened Cheeses – A Polyphasic Approach. , 2015, , 51-101. | | 1 |
| 236 | A report of 38 unrecorded bacterial species in Korea, belonging to the phylum Actinobacteria. Journal of Species Research, 2016, 5, 223-234. | 0.1 | 0 |
| 237 | <i>Sphingomicrobium arenosum</i> sp. nov., isolated from marine sediment. International Journal of Systematic and Evolutionary Microbiology, 2018, 68, 2551-2556. | 0.8 | 7 |
| 238 | <i>Actinomadura graeca</i> sp. nov.: A novel producer of the macrocyclic antibiotic zelvomycin. PLoS ONE, 2021, 16, e0260413. | 1.1 | 7 |
| 239 | Menaquinone Composition in the Classification of <i>Actinomadura</i> and Related Taxa. Microbiology (United Kingdom), 1984, 130, 817-823. | 0.7 | 10 |
| 240 | Molecular fingerprinting of some clinically significant <i>Nocardia</i> and related strains by restriction polymorphism ribosomal RNA analyses. Turkish Journal of Biology, 0, , . | 2.1 | 0 |
| 241 | <i>Gordonia pseudamarae</i> sp. nov., a home for novel actinobacteria isolated from stable foams on activated sludge wastewater treatment plants. International Journal of Systematic and Evolutionary Microbiology, 2022, 72, . | 0.8 | 3 |