

CITATION REPORT

List of articles citing

Prevalence and factors associated with failure of liver stiffness measurement using FibroScan in a prospective study of 2114 examinations

DOI: 10.1097/00042737-200604000-00015

**European Journal of Gastroenterology and Hepatology
, 2006, 18, 411-2.**

Source: <https://exaly.com/paper-pdf/40198818/citation-report.pdf>

Version: 2024-04-29

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
313	Diagnosis of hepatic fibrosis in patients with chronic hepatitis C. 2006 , 10, 821-33		14
312	Functional hepatic imaging and disease recurrence after liver transplantation. 2006 , 12, 1744-6		3
311	[Transient elastography for diagnosing liver cirrhosis]. 2006 , 131, 2765-9		26
310	[Transient elastography (FibroScan) for the non-invasive assessment of liver fibrosis: current status and perspectives]. 2007 , 45, 387-94		8
309	P5D-4 Advances in Liver Stiffness Measurements Using Transient Elastography. 2007 ,		0
308	Nonalcoholic fatty liver disease: from pathogenesis to patient care. 2007 , 3, 458-69		69
307	[Results and place of Fibroscan in the non-invasive diagnosis of hepatic fibrosis]. 2007 , 28, 94-102		6
306	Features associated with success rate and performance of FibroScan measurements for the diagnosis of cirrhosis in HCV patients: a prospective study of 935 patients. 2007 , 46, 628-34		201
305	FibroTest and FibroScan for the prediction of hepatitis C-related fibrosis: a systematic review of diagnostic test accuracy. 2007 , 102, 2589-600		297
304	Reproducibility of transient elastography in the evaluation of liver fibrosis in patients with chronic liver disease. 2007 , 56, 968-73		596
303	Nouveaux moyens diagnostiques de la cirrhose. 2007 , 16, 563-567		
302	Assessment of hepatic fibrosis with magnetic resonance elastography. 2007 , 5, 1207-1213.e2		723
301	Ultrasound-based transient elastography for the detection of hepatic fibrosis: systematic review and meta-analysis. 2007 , 5, 1214-20		396
300	Role of liver biopsy and serum markers of liver fibrosis in non-alcoholic fatty liver disease. 2007 , 11, 25-35, viii		58
299	Elastography in hepatology. 2007 , 21, 839-42		23
298	Transient elastography for the assessment of chronic liver disease: ready for the clinic?. 2007 , 13, 4791-7		17
297	The NAFLD fibrosis score: a noninvasive system that identifies liver fibrosis in patients with NAFLD. 2007 , 45, 846-54		1757

296	Transient elastography for diagnosis of portal hypertension in liver cirrhosis: is there still a role for hepatic venous pressure gradient measurement?. 2007 , 45, 1087-90	57
295	Noninvasive diagnosis and monitoring of nonalcoholic steatohepatitis: present and future. 2007 , 46, 582-9	341
294	Transient elastography is unreliable for detection of cirrhosis in patients with acute liver damage. 2008 , 47, 592-5	381
293	Significant liver damage in patients with bleeding disorders and chronic hepatitis C: non-invasive assessment of liver fibrosis using transient elastography. 2007 , 5, 25-30	19
292	[Use of elastometry (FibroScan) for the non-invasive staging of liver fibrosis]. 2007 , 31, 524-30; quiz 500, 531-2	9
291	Transient elastography for measurement of liver stiffness measurement can detect early significant hepatic fibrosis in Japanese patients with viral and nonviral liver diseases. 2008 , 43, 720-8	59
290	Extrahepatic cholestasis increases liver stiffness (FibroScan) irrespective of fibrosis. 2008 , 48, 1718-23	431
289	Elastography assessment in patients with chronic HCV infection. 2008 , 62, 1108-12	10
288	Correlation of liver biochemistry with liver stiffness in chronic hepatitis B and development of a predictive model for liver fibrosis. 2008 , 28, 1408-16	68
287	Prospective evaluation of transient elastography for the diagnosis of hepatic fibrosis in Asians: comparison with liver biopsy and aspartate transaminase platelet ratio index. 2008 , 28, 51-61	55
286	Assessment of asymptomatic liver fibrosis in alcoholic patients using fibroscan: prospective comparison with seven non-invasive laboratory tests. 2008 , 28, 1188-98	160
285	[Non-invasive evaluation of liver fibrosis in hepatitis C]. 2008 , 32, S90-5	7
284	Transient elastography (FibroScan). 2008 , 32, 58-67	167
283	MR elastography. 2008 , 32, 68-72	31
282	Combining non-invasive methods for assessment of liver fibrosis. 2008 , 32, 73-9	11
281	Non-invasive assessment of liver fibrosis by transient elastography in post transfusional iron overload. 2008 , 80, 337-40	24
280	Noninvasive assessment of liver fibrosis: serum markers, imaging, and other modalities. 2008 , 12, 883-900, x	29
279	Noninvasive assessment of liver fibrosis and portal hypertension with transient elastography. 2008 , 134, 8-14	128

278	Performance of transient elastography for the staging of liver fibrosis: a meta-analysis. 2008 , 134, 960-74	1129
277	Magnetic resonance elastography for the noninvasive staging of liver fibrosis. 2008 , 135, 32-40	575
276	Medici3n de la elasticidad hep3tica. Utilidad potencial del FibroScan® . 2008 , 7, 27-30	
275	Noninvasive assessment of liver fibrosis by measurement of stiffness in patients with nonalcoholic fatty liver disease (NAFLD). 2008 , 40, 371-8	287
274	Liver stiffness values in apparently healthy subjects: influence of gender and metabolic syndrome. 2008 , 48, 606-13	324
273	Non-invasive evaluation of liver fibrosis using transient elastography. 2008 , 48, 835-47	964
272	Evolving trends in nonalcoholic fatty liver disease. 2008 , 19, 75-82	25
271	A treatment algorithm for the management of chronic hepatitis B virus infection in the United States: 2008 update. 2008 , 6, 1315-41; quiz 1286	367
270	Screening for liver fibrosis by using a noninvasive biomarker in patients with diabetes. 2008 , 6, 828-31	52
269	New diagnostic method for liver fibrosis and cirrhosis. 2008 , 51 Suppl 1, 11-6	41
268	Assessing liver fibrosis. 2008 , 2, 541-52	31
267	Prevalence of fibrosis and cirrhosis in chronic hepatitis B: implications for treatment and management. 2008 , 103, 1421-6	43
266	[Assessment of the solid-liquid behavior of the liver for the diagnosis of diffuse disease using magnetic resonance elastography]. 2008 , 180, 1104-9	8
265	Diagnosis of nonalcoholic fatty liver disease: invasive versus noninvasive. 2008 , 28, 386-95	205
264	In vivo visualization of abdominal malignancies with acoustic radiation force elastography. 2008 , 53, 279-93	132
263	Learning curve and interobserver reproducibility evaluation of liver stiffness measurement by transient elastography. <i>European Journal of Gastroenterology and Hepatology</i> , 2008 , 20, 693-701	2.2 79
262	Risk assessment of hepatocellular carcinoma in chronic hepatitis C patients by transient elastography. 2008 , 42, 839-43	64
261	Why is transient elastography essential in hepatology?. 2009 , 8, 148-149	0

260	Noninvasive diagnosis of nonalcoholic fatty liver disease. 2009 , 8, S25-S33	29
259	Elastography for the non-invasive assessment of liver disease: limitations and future developments. 2009 , 58, 157-60	33
258	Liver fibrosis in viral hepatitis: noninvasive assessment with acoustic radiation force impulse imaging versus transient elastography. 2009 , 252, 595-604	531
257	A proteomic strategy to identify novel serum biomarkers for liver cirrhosis and hepatocellular cancer in individuals with fatty liver disease. 2009 , 9, 271	59
256	[Non-invasive diagnosis of liver fibrosis: guidelines for the use of biomarkers and FibroScan]. 2009 , 33, 334-44	6
255	[Noninvasive tools for the diagnosis of steatosis and fibrosis in patients with NAFLD]. 2009 , 33, 405-12	2
254	Factors of accuracy of transient elastography (fibrosan) for the diagnosis of liver fibrosis in chronic hepatitis C. 2009 , 49, 1083-9	175
253	Mitteilungen der GARPS. 2009 , 4, 469-472	
252	Diabetes und Fettleber. 2009 , 5, 653-665	0
251	Transient elastography and other noninvasive tests to assess hepatic fibrosis in patients with viral hepatitis. 2009 , 16, 300-14	127
250	Noninvasive diagnosis of liver fibrosis in patients with HIV infection and HCV/HBV co-infection. 2009 , 16, 249-58	23
249	An evaluation of the potential cost-effectiveness of non-invasive testing strategies in the diagnosis of significant liver fibrosis. 2009 , 24, 786-91	25
248	Non-invasive model predicting clinically-significant portal hypertension in patients with advanced fibrosis. 2009 , 24, 1289-93	26
247	Non-invasive assessment of liver fibrosis by stiffness measurement in patients with chronic hepatitis B. 2009 , 29, 242-7	369
246	Reproducibility and limitations of transient elastography. 2009 , 29, 619-20	12
245	In vivo quantification of liver stiffness in a rat model of hepatic fibrosis with acoustic radiation force. 2009 , 35, 1709-21	45
244	[Utility of Fibroscan in the evaluation of liver fibrosis]. 2009 , 32, 415-23	6
243	[Transient elastography (FibroScan): a new tool in hepatology]. 2009 , 38, 1516-25	4

242	Early detection in routine clinical practice of cirrhosis and oesophageal varices in chronic hepatitis C: comparison of transient elastography (FibroScan) with standard laboratory tests and non-invasive scores. 2009 , 50, 59-68		289
241	[Value of transient elastography in noninvasive assessment in children's hepatic fibrosis]. 2009 , 16, 1005-10		22
240	Transient elastography for the assessment of liver fibrosis in patients with chronic viral hepatitis: the missing tool?. 2009 , 41, 863-6		23
239	Assessment of liver stiffness in patients after living donor liver transplantation by transient elastography. 2009 , 44, 1115-20		11
238	Noninvasive assessment of fibrosis and steatosis in NASH and ASH. 2009 , 33, 940-8		12
237	Liver stiffness measurement in combination with noninvasive markers for the improved diagnosis of B-viral liver cirrhosis. 2009 , 43, 267-71		63
236	Ultrasound-based hepatic elastography: origins, limitations, and applications. 2010 , 44, 637-45		34
235	Non-alcoholic fatty liver disease: the hepatic consequence of obesity and the metabolic syndrome. 2010 , 69, 211-20		156
234	How can we enhance the performance of liver stiffness measurement using FibroScan in diagnosing liver cirrhosis in patients with chronic hepatitis B?. 2010 , 44, 66-71		88
233	Noninvasive markers of fibrosis and inflammation in clinical practice: prospective comparison with liver biopsy. <i>European Journal of Gastroenterology and Hepatology</i> , 2010 , 22, 474-80	2.2	26
232	Acoustic radiation force impulse: a new ultrasonographic technology for the widespread noninvasive diagnosis of liver fibrosis. <i>European Journal of Gastroenterology and Hepatology</i> , 2010 , 22, 1074-84	2.2	113
231	Interobserver concordance in the assessment of liver fibrosis in HIV/HCV-coinfected patients using transient elastometry. <i>European Journal of Gastroenterology and Hepatology</i> , 2010 , 22, 801-7	2.2	23
230	Evaluation of sound speed for detection of liver fibrosis: prospective comparison with transient dynamic elastography and histology. 2010 , 29, 1581-8		19
229	Liver transient elastography (Fibroscan): a place in the management algorithms of chronic viral hepatitis. <i>Antiviral Therapy</i> , 2010 , 15, 1-11	1.6	25
228	Assessment of liver fibrosis and steatosis in PBC with FibroScan, MRI, MR-spectroscopy, and serum markers. 2010 , 44, 58-65		69
227	Transient elastography with a new probe for obese patients for non-invasive staging of non-alcoholic steatohepatitis. 2010 , 20, 2390-6		91
226	Chronic hepatitis B: whom to treat and for how long? Propositions, challenges, and future directions. 2010 , 4, 386-95		44
225	Assessment of liver fibrosis by transient elastography in rheumatoid arthritis patients treated with methotrexate. 2010 , 77, 588-92		26

224	Pitfalls of liver stiffness measurement: a 5-year prospective study of 13,369 examinations. 2010 , 51, 828-35	736
223	Feasibility of liver transient elastography with FibroScan using a new probe for obese patients. 2010 , 30, 1043-8	139
222	Transient elastography. 2010 , 25, 1726-31	64
221	Noninvasive evaluation of renal allograft fibrosis by transient elastography--a pilot study. 2010 , 23, 871-7	75
220	Shear wave velocity is a useful marker for managing nonalcoholic steatohepatitis. 2010 , 16, 2918-25	64
219	Liver stiffness: a novel parameter for the diagnosis of liver disease. 2010 , 2, 49-67	220
218	Transient elastography in chronic hepatitis B: an Asian perspective. 2010 , 16, 5173-80	28
217	Transient elastography for the noninvasive assessment of liver fibrosis: a multicentre Canadian study. 2010 , 24, 661-70	46
216	Liver biopsy for histological assessment: The case in favor. 2010 , 16, 133-9	9
215	Decompensated chronic heart failure: increased liver stiffness measured by means of transient elastography. 2010 , 257, 872-8	128
214	Prospective comparison of two algorithms combining non-invasive methods for staging liver fibrosis in chronic hepatitis C. 2010 , 52, 191-8	158
213	Liver stiffness is directly influenced by central venous pressure. 2010 , 52, 206-10	394
212	Ultrasonographic evaluation of liver surface and transient elastography in clinically doubtful cirrhosis. 2010 , 52, 846-53	90
211	[Retrospective study of FibroScan, APRI, FIB-4 and FORNS indexes compared with liver biopsy in the evaluation of liver fibrosis in patients with chronic hepatitis C mono-infection and HIV coinfection]. 2010 , 33, 425-32	10
210	Transient elastography for the diagnosis of liver fibrosis. 2010 , 7, 811-23	66
209	The role of transient elastography in patients with hepatitis B viral disease. 2011 , 43 Suppl 1, S25-31	13
208	Fatty liver and its clinical management in obese adolescents. 2011 , 58, 32-37	
207	[Fatty liver and its clinical management in obese adolescents]. 2011 , 58, 32-7	1

206	Drug metabolism alterations in nonalcoholic fatty liver disease. 2011 , 43, 317-34	114
205	[Non-alcoholic fatty liver disease. How and who to screen]. 2011 , 34, 278-88	
204	Comparison of NAFLD fibrosis score and BARD score in predicting fibrosis in nonalcoholic fatty liver disease. 2011 , 54, 160-3	57
203	Noninvasive evaluation of hepatic fibrosis using acoustic radiation force-based shear stiffness in patients with nonalcoholic fatty liver disease. 2011 , 55, 666-672	264
202	Noninvasive Alternatives for the Assessment of Liver Fibrosis. 2011 ,	
201	Non-Invasive Assessment of Liver Fibrosis by Vibration-Controlled Transient Elastography (Fibroscan®). 2011 ,	4
200	The usefulness of liver stiffness measurement using FibroScan in chronic hepatitis C in South Korea: a multicenter, prospective study. 2011 , 26, 171-8	33
199	Transient elastography: the bigger we are, the harder to scan. 2011 , 26, 217-9	3
198	Reproducibility of liver stiffness measurements in hepatitis C virus (HCV)-infected patients in Egypt. 2011 , 18, e358-65	12
197	Effect of detoxification on liver stiffness assessed by Fibroscan® in alcoholic patients. 2011 , 35, 566-70	50
196	Prospective evaluation of FibroScan for the diagnosis of hepatic fibrosis compared with liver biopsy/AST platelet ratio index and FIB-4 in patients with chronic HBV infection. 2011 , 56, 2742-9	71
195	The use of transient elastography in the management of chronic hepatitis B. 2011 , 5, 868-875	13
194	Multi-excitation fluorescence spectroscopy for analysis of non-alcoholic fatty liver disease. 2011 , 43, 392-400	10
193	Test-retest repeatability of MR elastography for noninvasive liver fibrosis assessment in hepatitis C. 2011 , 34, 947-55	107
192	MR elastography for noninvasive assessment of hepatic fibrosis: experience from a tertiary center in Asia. 2011 , 34, 1110-6	81
191	Liver stiffness assessment by tagged MRI of cardiac-induced liver motion. 2011 , 65, 949-55	38
190	Évaluation de la fibrose hépatique par l'élastométrie chez les patients atteints de polyarthrite rhumatoïde traités par méthotrexate. 2011 , 78, 47-51	
189	Hepatic elasticity in patients with ascites: evaluation with real-time tissue elastography. 2011 , 196, W766-71	26

188	Breast mass evaluation: factors influencing the quality of US elastography. 2011 , 259, 59-64	144
187	Cholestatic liver diseases from child to adult: the diversity of MDR3 disease. 2011 , 49, 728-36	28
186	What did we learn from the first 3,459 cases of liver stiffness measurement by transient elastography (FibroScan®)? 2011 , 32, 40-5	14
185	Accuracy of VirtualTouch Acoustic Radiation Force Impulse (ARFI) imaging for the diagnosis of cirrhosis during liver ultrasonography. 2011 , 32, 167-75	94
184	Preliminary Experiments on Magnetic Resonance Elastography. 2011 ,	
183	Evaluation of acoustic radiation force impulse imaging for determination of liver stiffness using transient elastography as a reference. 2012 , 18, 1077-84	45
182	Systemic mastocytosis: a rare case of increased liver stiffness. 2012 , 2012, 728172	9
181	[Serological markers of fibrosis]. 2012 , 35 Suppl 2, 10-6	3
180	MRI-guided biopsy to correlate tissue specimens with MR elastography stiffness readings in liver transplants. 2012 , 19, 1121-6	15
179	Comparison of acoustic radiation force impulse imaging with transient elastography for the detection of complications in patients with cirrhosis. 2012 , 32, 852-8	63
178	Discordance in fibrosis staging between liver biopsy and transient elastography using the FibroScan XL probe. 2012 , 56, 564-70	112
177	Elastographic assessment of liver fibrosis in children: A prospective single center experience. 2012 , 81, e870-4	41
176	Can NASH be diagnosed, graded, and staged noninvasively?. 2012 , 16, 567-85	53
175	Non-Invasive Evaluation of Liver Steatosis, Fibrosis and Cirrhosis in Hepatitis C Virus Infected Patients Using Unidimensional Transient Elastography (Fibroscan®). 2012 ,	1
174	Overview and developments in noninvasive diagnosis of nonalcoholic fatty liver disease. 2012 , 18, 3945-54	55
173	Transient elastography in non-alcoholic fatty liver disease. 2012 , 11, 172-178	42
172	Medical management of bleeding varices. 2012 , 1129-1134.e2	
171	Liver stiffness measurement using acoustic radiation force impulse (ARFI) elastography and effect of necroinflammation. 2012 , 57, 1682-91	73

170	Noninvasive measurement of liver fibrosis by transient elastography and influencing factors in patients with chronic hepatitis B-A single center retrospective study of 466 patients. 2012 , 32, 69-74	7
169	Beneficial effect of successful HCV treatment in patients with inherited bleeding disorders, assessed by liver stiffness measurements. 2012 , 18, e266-72	4
168	Feasibility and diagnostic performance of the FibroScan XL probe for liver stiffness measurement in overweight and obese patients. 2012 , 55, 199-208	335
167	Prospective biopsy-controlled evaluation of cell death biomarkers for prediction of liver fibrosis and nonalcoholic steatohepatitis. 2012 , 55, 455-64	126
166	Factors influencing reliability of liver stiffness measurements using transient elastography (M-probe)-monocentric experience. 2013 , 82, e313-6	27
165	Value of acoustic radiation force impulse imaging elastography for non-invasive evaluation of patients with nonalcoholic fatty liver disease. 2013 , 39, 1942-50	69
164	JSUM ultrasound elastography practice guidelines: liver. 2013 , 40, 325-57	75
163	Role of fibroscan and APRI in detection of liver fibrosis: a systematic review and meta-analysis. 2013 , 14, 44-50	28
162	Longitudinal assessment of liver stiffness in patients undergoing antiviral treatment for hepatitis C. 2013 , 45, 840-3	41
161	Influence of different frequencies and insertion depths on the diagnostic accuracy of liver elastography by acoustic radiation force impulse imaging (ARFI). 2013 , 82, 1207-12	40
160	Clinical application of liver stiffness measurement using transient elastography: a surgical perspective. 2013 , 88, 258-65	17
159	Strategies for biomarker discovery in fibrotic disease. 2013 , 1832, 1079-87	11
158	Quantification of liver viscoelasticity with acoustic radiation force: a study of hepatic fibrosis in a rat model. 2013 , 39, 2091-102	45
157	Assessment of renal allograft fibrosis by transient elastography. 2013 , 26, 545-51	45
156	Liver fibrosis: noninvasive assessment with acoustic radiation force impulse elastography--comparison with FibroScan M and XL probes and FibroTest in patients with chronic liver disease. 2013 , 269, 283-92	91
155	Utility of controlled attenuation parameter measurement for assessing liver steatosis in Japanese patients with chronic liver diseases. 2013 , 43, 1125-6	3
154	. 2013 ,	
153	Acoustic radiation force impulse of the liver. 2013 , 19, 4841-9	46

152	Direct comparison of the FibroScan XL and M probes for assessment of liver fibrosis in obese and nonobese patients. 2013 , 5, 43-52	16
151	Defining normal liver stiffness range in a normal healthy Chinese population without liver disease. 2013 , 8, e85067	29
150	Feasibility and reliability of the FibroScan S2 (pediatric) probe compared with the M probe for liver stiffness measurement in small adults with chronic liver disease. 2013 , 12, 100-107	18
149	Non-invasive assessment of liver fibrosis using ARFI with pathological correlation, a prospective study. 2013 , 12, 440-447	10
148	The clinical utility of FibroScan(®) as a noninvasive diagnostic test for liver disease. 2014 , 7, 107-114	32
147	Invasive and non-invasive diagnosis of cirrhosis and portal hypertension. 2014 , 20, 4300-15	69
146	Non-invasive diagnosis of hepatitis B virus-related cirrhosis. 2014 , 20, 445-59	22
145	Non-invasive prediction of development of hepatocellular carcinoma using transient elastography in patients with chronic liver disease. 2014 , 8, 501-11	7
144	Fibroscan. 2014 , 4, 97-101	7
143	Correlation of MR elastography with morphometric quantification of liver fibrosis (Fibro-C-Index) in chronic hepatitis B. 2014 , 72, 1123-9	30
142	Determination of normal hepatic elasticity by using real-time shear-wave elastography. 2014 , 271, 895-900	53
141	Acoustic radiation force impulse elastography for non-invasive assessment of disease stage in patients with primary biliary cirrhosis: A preliminary study. 2014 , 69, 836-40	20
140	[Using transient elastography as a screening tool for liver fibrosis in addiction service]. 2014 , 43, e17-31	3
139	MR elastography for noninvasive assessment of hepatic fibrosis: reproducibility of the examination and reproducibility and repeatability of the liver stiffness value measurement. 2014 , 39, 326-31	79
138	Utilization of FibroScan in clinical practice. 2014 , 16, 372	67
137	Use of magnetic resonance elastography to assess hepatic fibrosis in children with chronic liver disease. 2014 , 164, 186-8	102
136	Serumuma refeição ligeira fator de erro na avaliação da dureza hepática por elastografia hepática transitória? Um estudo prospectivo. 2014 , 21, 102-108	1
135	Matrix biology of idiopathic pulmonary fibrosis: a workshop report of the national heart, lung, and blood institute. 2014 , 184, 1643-51	74

134	Quantitative comparison of transient elastography (TE), shear wave elastography (SWE) and liver biopsy results of patients with chronic liver disease. 2015 , 27, 2465-8		13
133	Longer duration of transient elastography predicts unreliable liver stiffness measurements. <i>European Journal of Gastroenterology and Hepatology</i> , 2015 , 27, 655-9	2.2	1
132	Inflammation-adapted liver stiffness values for improved fibrosis staging in patients with hepatitis C virus and alcoholic liver disease. 2015 , 35, 2514-21		65
131	Patient preference and willingness to pay for transient elastography versus liver biopsy: A perspective from British Columbia. 2015 , 29, 72-6		13
130	Diagnostic imaging modalities. 2015 , 21-34		
129	Noninvasive evaluation of liver fibrosis using real-time tissue elastography and transient elastography (FibroScan). 2015 , 34, 403-10		26
128	Non-invasive detection of liver fibrosis: MR imaging features vs. MR elastography. 2015 , 40, 766-75		64
127	Performance of transient elastography assessing fibrosis of single hepatitis B virus infection: a systematic review and meta-analysis of a diagnostic test. 2015 , 9, 558-66		23
126	Efficacy and Accessibility of a Novel Liver Fibrosis Evaluation Program: Limited Abdominal Ultrasound with Concurrent Transient Elastography. 2015 , 22, 814-9		1
125	Evaluation of renal allograft fibrosis by transient elastography (Fibro Scan). 2015 , 47, 640-3		28
124	Functional Liver Recovery After Bariatric Surgery--a Prospective Cohort Study with the LiMAx Test. 2015 , 25, 2047-53		23
123	WFUMB guidelines and recommendations for clinical use of ultrasound elastography: Part 3: liver. 2015 , 41, 1161-79		390
122	Evaluation of Acoustic Radiation Force Impulse Imaging (ARFI) for the Determination of Liver Stiffness Using Transient Elastography as a Reference in Children. 2015 , 1, E2-7		15
121	Comparison of acoustic radiation force impulse imaging and transient elastography for non-invasive assessment of liver fibrosis in patients with chronic hepatitis B. 2015 , 41, 7-14		26
120	Comparison of the diagnostic accuracies of magnetic resonance elastography and transient elastography for hepatic fibrosis. 2015 , 33, 26-30		74
119	Non-invasive assessment of liver fibrosis in patients with alcoholic liver disease using acoustic radiation force impulse elastography. 2015 , 40, 723-9		33
118	Measurement of liver stiffness as a non-invasive method for diagnosis of non-alcoholic fatty liver disease. 2015 , 45, 142-51		39
117	Evaluation of the utility of serum prolidase as a marker for liver fibrosis. 2015 , 29, 208-13		6

116	Magnetic Resonance Elastography and Other Magnetic Resonance Imaging Techniques in Chronic Liver Disease: Current Status and Future Directions. 2016 , 10, 672-86	29
115	A Review of Western and Traditional Chinese Medical Approaches to Managing Nonalcoholic Fatty Liver Disease. 2016 , 2016, 6491420	8
114	FibroScan: assessing cirrhosis and advanced fibrosis with vibration-controlled transient elastography. 2016 , 14, S10-S15	
113	Automatic detection of cirrhosis in hospitalized patients: a pragmatic experience. <i>European Journal of Gastroenterology and Hepatology</i> , 2016 , 28, 74-81	2.2
112	Controlled attenuation parameter is correlated with actual hepatic fat content in patients with non-alcoholic fatty liver disease with none-to-mild obesity and liver fibrosis. 2016 , 46, 1019-27	30
111	Comparison of three dimensional strain volume reconstructions using SOUPR and wobbler based acquisitions: A phantom study. 2016 , 43, 1615	3
110	Cytokeratin 18-Aspartate396 apoptotic fragment for fibrosis detection in patients with non-alcoholic fatty liver disease and chronic viral hepatitis. 2016 , 48, 55-61	16
109	Noninvasive Markers of Fibrosis and Inflammation in Nonalcoholic Fatty Liver Disease. 2016 , 15, 86-95	16
108	Serum Fibrosis Markers for the Diagnosis of Liver Disease Among People With Chronic Hepatitis C in Chennai, India. 2016 , 3, ofw156	4
107	Decision analytic model of the diagnostic pathways for patients with suspected non-alcoholic fatty liver disease using non-invasive transient elastography and multiparametric magnetic resonance imaging. 2016 , 6, e010507	20
106	Reply. 2016 , 151, 375-6	
105	The Correlation Between Obesity-Related Diseases and Non-alcoholic Fatty Liver Disease in Women in the Pre-operative Evaluation for Bariatric Surgery Assessed by Transient Hepatic Elastography. 2016 , 26, 2089-2097	9
104	Magnetic Resonance Imaging More Accurately Classifies Steatosis and Fibrosis in Patients With Nonalcoholic Fatty Liver Disease Than Transient Elastography. 2016 , 150, 626-637.e7	438
103	Liver Steatosis Assessed by Controlled Attenuation Parameter (CAP) Measured with the XL Probe of the FibroScan: A Pilot Study Assessing Diagnostic Accuracy. 2016 , 42, 92-103	84
102	Robust sound speed estimation for ultrasound-based hepatic steatosis assessment. 2017 , 62, 3582-3598	61
101	Diagnostic Accuracy of Noninvasive Fibrosis Scores in a Population of Individuals With a Low Prevalence of Fibrosis. 2017 , 15, 1453-1460.e1	20
100	Cell death biomarker M65 is a useful indicator of liver inflammation and fibrosis in chronic hepatitis B: A cross-sectional study of diagnostic accuracy. 2017 , 96, e6807	8
99	Reproducibility of 2-Dimensional Shear Wave Elastography Assessment of the Liver: A Direct Comparison With Point Shear Wave Elastography in Healthy Volunteers. 2017 , 36, 1563-1569	25

98	Multiparametric or practical quantitative liver MRI: towards millisecond, fat fraction, kilopascal and function era. 2017 , 11, 167-182	15
97	Non-invasive Assessment of Liver Fibrosis with ElastPQ: Comparison with Transient Elastography and Serologic Fibrosis Marker Tests, and Correlation with Liver Pathology Results. 2017 , 43, 2515-2521	13
96	Central Hepatectomy Still Plays an Important Role in Treatment of Early-Stage Centrally Located Hepatocellular Carcinoma. 2017 , 41, 2830-2837	6
95	Primary Sclerosing Cholangitis. 2017 ,	
94	Strain elastography for assessment of liver fibrosis and prognosis in patients with chronic liver diseases. 2017 , 52, 724-733	8
93	Magnetic Resonance Elastography vs Transient Elastography in Detection of Fibrosis and Noninvasive Measurement of Steatosis in Patients With Biopsy-Proven Nonalcoholic Fatty Liver Disease. 2017 , 152, 598-607.e2	342
92	Liver stiffness measurement to platelet ratio index predicts the stage of liver fibrosis in non-alcoholic fatty liver disease. 2017 , 47, 721-730	11
91	Liver stiffness measurement predicts high-grade post-hepatectomy liver failure: A prospective cohort study. 2017 , 32, 506-514	26
90	Technical and clinical biases of transient elastography in liver diseases. 2017 ,	
89	Effects of lifestyle modification on liver enzyme and Fibroscan in Indian patients with non-alcoholic fatty liver disease. 2018 , 6, 49-53	2
88	Evaluating Diagnostic Accuracy of Noninvasive Tests in Assessment of Significant Liver Fibrosis in Chronic Hepatitis C Egyptian Patients. 2018 , 31, 315-320	1
87	Abbreviated mpMRI protocol for diffuse liver disease: a practical approach for evaluation and follow-up of NAFLD. 2018 , 43, 2340-2350	14
86	Quantitative estimation of progression of chronic liver disease using gadoxetate disodium-enhanced magnetic resonance imaging. 2018 , 48, 735-745	3
85	Diagnosis and Evaluation of Nonalcoholic Fatty Liver Disease/Nonalcoholic Steatohepatitis, Including Noninvasive Biomarkers and Transient Elastography. 2018 , 22, 73-92	44
84	Role of imaging-based biomarkers in NAFLD: Recent advances in clinical application and future research directions. 2018 , 68, 296-304	69
83	Optimal threshold of controlled attenuation parameter with MRI-PDFF as the gold standard for the detection of hepatic steatosis. 2018 , 67, 1348-1359	144
82	Bariatric Surgery as an Efficient Treatment for Non-Alcoholic Fatty Liver Disease in a Prospective Study with 1-Year Follow-up : BariScan Study. 2018 , 28, 1342-1350	50
81	Ultrasound diagnosis of hepaticfibrosis and steatosis. 2018 , 59, 384-392	

80	A Noninvasive Score to Predict Liver Fibrosis in HBeAg-Positive Hepatitis B Patients with Normal or Minimally Elevated Alanine Aminotransferase Levels. 2018 , 2018, 3924732	2
79	The XL probe: A luxury or a necessity? Risk stratification in an obese community cohort using transient elastography. 2018 , 6, 1372-1379	3
78	Monitoring methotrexate-induced liver fibrosis in patients with psoriasis: utility of transient elastography. 2018 , 8, 21-29	10
77	Non-invasive tests for the detection of oesophageal varices in compensated cirrhosis: systematic review and meta-analysis. 2018 , 6, 806-818	8
76	Magnetic resonance elastography increases usefulness and safety of non-invasive screening for esophageal varices. 2018 , 33, 2022-2028	15
75	. 2018 , 6, 33969-33976	2
74	Transient elastography: should XL probe be used in all overweight patients?. 2019 , 54, 1022-1026	4
73	Liver Fibrosis Assessment with Diffusion-Weighted Imaging: Value of Liver Apparent Diffusion Coefficient Normalization Using the Spleen as a Reference Organ. 2019 , 9,	8
72	Noninvasive monitoring of hepatic steatosis: controlled attenuation parameter and magnetic resonance imaging-proton density fat fraction in patients with nonalcoholic fatty liver disease. 2019 , 13, 523-530	11
71	Quick assessment with controlled attenuation parameter for hepatic steatosis in children based on MRI-PDFF as the gold standard. 2019 , 19, 112	20
70	Nonalcoholic Fatty Liver Disease in Diabetes. Part I: Epidemiology and Diagnosis. 2019 , 43, 31-45	60
69	Evaluation of hepatic fibrosis by ultrasonic acoustic structure quantification. 2019 , 98, e16533	3
68	Fibroscan and low-density lipoprotein as determinants of severe liver fibrosis in diabetic patients with nonalcoholic fatty liver disease. <i>European Journal of Gastroenterology and Hepatology</i> , 2019 , 31, 1540-1544	2.2 7
67	Comparison of Acoustic Structure Quantification, Transient Elastography (FibroScan) and Histology in Patients with Chronic Hepatitis B and without Moderate to Severe Hepatic Steatosis. 2019 , 45, 684-692	3
66	Subgroup analysis of the predictive ability of aspartate aminotransferase to platelet ratio index (APRI) and fibrosis-4 (FIB-4) for assessing hepatic fibrosis among patients with chronic hepatitis C. 2020 , 53, 542-549	6
65	Non-Alcoholic Fatty Liver Disease. 2020 ,	1
64	Diagnostic of hepatic fibrosis with the XL probe of the Fibroscan versus biopsies in patients candidates to bariatric surgery. 2020 , 37, 226-232	2
63	Total bile acid-to-cholesterol ratio as a novel noninvasive marker for significant liver fibrosis and cirrhosis in patients with non-cholestatic chronic hepatitis B virus infection. 2020 , 99, e19248	4

62	Utility and accuracy of transient elastography in determining liver fibrosis: a case-control study. 2020 , 179, 671-677	3
61	Use of Direct Liver Stiffness Measurement in Evaluating the Severity of Liver Cirrhosis in Patients with Hepatocellular Carcinoma. 2020 , 44, 2777-2783	4
60	A meta-analysis on the diagnostic performance of magnetic resonance imaging and transient elastography in nonalcoholic fatty liver disease. 2021 , 51, e13446	11
59	Liver stiffness measurement predicts short-term and long-term outcomes in patients with hepatocellular carcinoma after curative liver resection. 2021 ,	0
58	Estimation of liver elasticity using the finite element method and four-dimensional computed tomography images as a biomarker of liver fibrosis. 2021 , 48, 1286-1298	0
57	N-glycan structures of Wisteria floribunda agglutinin-positive Mac2 binding protein in the serum of patients with liver fibrosis 2021 , 31, 1268-1278	0
56	Viral Biomarkers for Hepatitis B Virus-Related Hepatocellular Carcinoma Occurrence and Recurrence. 2021 , 12, 665201	3
55	Feasibility of Fibroscan in Assessment of Hepatic Steatosis and Fibrosis in Obese Patients: Report From a General Internal Medicine Clinic. 2021 , 32, 466-472	0
54	Usefulness of US attenuation imaging for the detection and severity grading of hepatic steatosis in routine abdominal ultrasonography. 2021 , 76, 53-59	4
53	The severity of histologic liver lesions is independent of body mass index in patients with nonalcoholic fatty liver disease. 2013 , 47, 280-6	28
52	Evaluating the Significance of Viscoelasticity in Diagnosing Early-Stage Liver Fibrosis with Transient Elastography. 2017 , 12, e0170073	9
51	Principles and clinical application of ultrasound elastography for diffuse liver disease. 2014 , 33, 149-60	86
50	Clinical applications of transient elastography. 2012 , 18, 163-73	74
49	Magnetization-tagged MRI is a simple method for predicting liver fibrosis. 2016 , 22, 140-5	1
48	[Liver stiffness measurement for the diagnosis of hepatic fibrosis in patients with chronic viral hepatitis]. 2007 , 13, 521-9	8
47	[Clinical factors influencing liver stiffness as measured by transient elastography (Fibroscan) in patients with chronic liver disease]. 2010 , 16, 123-30	10
46	Hepatocellular Carcinoma and Non-Alcoholic Fatty Liver Disease: A Step Forward for Better Evaluation Using Ultrasound Elastography. 2020 , 12,	6
45	The Use of Two-Dimensional Shear Wave Elastography in People with Obesity for the Assessment of Liver Fibrosis in Non-Alcoholic Fatty Liver Disease. 2020 , 10,	5

44	Nonalcoholic fatty liver disease: an overview of current insights in pathogenesis, diagnosis and treatment. 2008 , 14, 2474-86	135
43	Increased liver stiffness in alcoholic liver disease: differentiating fibrosis from steatohepatitis. 2010 , 16, 966-72	169
42	Value of transient elastography for the prediction of variceal bleeding. 2011 , 17, 2206-10	31
41	Checkmate to liver biopsy in chronic hepatitis C?. 2012 , 18, 5514-20	13
40	Clinical application of liver stiffness measurement using transient elastography in chronic liver disease from longitudinal perspectives. 2013 , 19, 1890-900	58
39	Study of detection times for liver stiffness evaluation by shear wave elastography. 2014 , 20, 9578-84	21
38	Non-invasive assessment of liver fibrosis in chronic hepatitis B. 2014 , 20, 14568-80	39
37	Acoustic radiation force impulse imaging for assessing liver fibrosis in alcoholic liver disease. 2016 , 22, 4926-35	25
36	The Largest American Study Comparing Transient Elastography and the Aminotransferase-to-platelet Ratio Index Score, Two Noninvasive Tests for Liver Fibrosis Staging in HIV/HCV Coinfected Patients. 2019 , 11, 80-82	1
35	Clinical Evaluation of Hepatic Fibrosis in Chinese Patients with Nonalcoholic Fatty Liver Disease with FibroScan. 2015 , 06, 115-123	1
34	Impact of bariatric surgery on non-alcoholic fatty liver disease. 2017 , 89, 1-4	4
33	Endoscopic Ultrasound Real-Time Elastography in Liver Disease. 2018 , 51, 118-119	2
32	Valuer la fibrose: Pourquoi? Comment? Pour qui?. 2010 , 245-257	
31	Liver Biopsy and FibroScan to Detect Early Histopathological Changes in Chronic HBV Patients Not Candidate for Treatment. 2014 , 7, 56-63	1
30	Liver stiffness measurement by transient elastography (FibroScan). 2015 , 42, 477-485	1
29	Diagnosis of experimental steatohepatosis using ultrasound shear wave elastography. 2015 , 26, 109-113	0
28	Noninvasive Imaging of Primary Sclerosing Cholangitis: A Radiologic Perspective. 2017 , 167-179	
27	The diagnostic value of normalized ADC using spleen as reference organ in assessment liver fibrosis. 2020 , 51,	0

26	Noninvasive Diagnostic Approach to NASH: Radiological Diagnostics. 2020 , 257-269		
25	Non-biopsy methods to determine hepatic fibrosis. 2009 , 2, 401-6		4
24	NASH: The Hepatic injury of Metabolic syndrome: a brief update. 2009 , 3, 265-70		4
23	Correlation of Laparoscopic Liver Biopsy to Elasticity Measurements (FibroScan) in Patients With Chronic Liver Disease. 2008 , 4, 862-70		7
22	Assessment of noninvasive liver stiffness in inactive HBsAg carriers by transient elastography: Fibroscan in inactive HBsAg carriers. 2011 , 11, 182-5		11
21	Quantitative assessment of the elasticity values of liver with shear wave ultrasonographic elastography. 2013 , 137, 911-5		2
20	EUS Elastography in Evaluating Chronic Liver Disease. Why not from Inside?. 2009 , 35, 225-7		5
19	The Role of Elastography in Non-Alcoholic Fatty Liver Disease. 2020 , 46, 255-269		
18	The Evolving Role of Advanced Endoscopic Techniques in Hepatology. 2021 , 17, 67-72		0
17	Liver Elastography: Basic Principles, Evaluation Technique, and Confounding Factors.		
16	Noninvasive Assessment of Liver Fibrosis with ElastPQ in Patients with Chronic Viral Hepatitis: Comparison Using Histopathological Findings.. 2022 , 12,		
15	Reproducibility and diagnostic accuracy of pocket-sized ultrasound devices in ruling out compensated cirrhosis of mixed etiology.. 2022 ,		
14	HVPG10 score for prediction of clinically significant portal hypertension - ready for prime time?. <i>Hpb</i> , 2021 ,	3.8	
13	Diagnosis, Diagnostic Tests and Monitoring of Hepatitis B Virus in Monoinfected and HIV-Coinfected Patients. <i>Antiviral Therapy</i> , 2007 , 12, 25-31	1.6	11
12	Use of doppler ultrasound to predict need for transjugular intrahepatic portosystemic shunt revision. <i>World Journal of Hepatology</i> , 2022 , 14, 1200-1209	3-4	
11	Noninvasive Assessment of Liver Fibrosis by Transient Elastography(The Forefront of Noninvasive Test). <i>Ika Kikaigaku</i> , 2006 , 76, 883-887		
10	Fibroscan and Shear Wave correlated well in hepatic fibrosis evaluation of patients with chronic liver diseases in real life situation 2022, 101, e30025		0
9	Non-Invasive Imaging Modalities in Nonalcoholic Fatty Liver Disease: Where Do We Stand?. 80-85		0

- 8 Noninvasive Imaging Modalities in Nonalcoholic Fatty Liver Disease: Where Do We Stand?. 57-62 1
- 7 Non-invasive methods to evaluate liver fibrosis in patients with non-alcoholic fatty liver disease. 13, 0
- 6 Deep attenuation transducer to measure liver stiffness in obese patients with liver disease. 0
- 5 The role of transient elastography and novel serum biomarkers in assessment of liver fibrosis in patients with psoriasis and rheumatic disease. 0
- 4 The Non-Invasive Assessment of Circulating D-Loop and mt-ccf Levels Opens an Intriguing Spyhole into Novel Approaches for the Tricky Diagnosis of NASH. **2023**, 24, 2331 1
- 3 Performance of two-dimensional shear wave elastography and transient elastography compared to liver biopsy for staging of liver fibrosis. 0
- 2 Comparative Accuracy of Clinical Fibrosis Markers, Hepascore and Fibroscan[®] to Detect Advanced Fibrosis in Patients with Nonalcoholic Fatty Liver Disease. 0
- 1 Three-dimensional magnetic resonance elastography in chronic liver disease. **2023**, 2, 32-39 0