

Overweight and obesity increase the risk for liver cancer
and long-term oral supplementation with branched-chain
liver carcinogenesis in heavier patients with liver cirrhosis

Hepatology Research

35, 204-14

DOI: [10.1016/j.hepres.2006.04.007](https://doi.org/10.1016/j.hepres.2006.04.007)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Nutritional assessment in liver cirrhosis. <i>Journal of Gastroenterology</i> , 2006, 41, 511-512.	2.3	2
2	The effect of supplementation with branched-chain amino acids in patients with liver cirrhosis. <i>Hepatology Research</i> , 2007, 37, 510-516.	1.8	55
3	Oral branched-chain amino acid supplementation improves the oxidized/reduced albumin ratio in patients with liver cirrhosis. <i>Hepatology Research</i> , 2007, 37, 765-770.	1.8	55
4	Restoration of innate host defense responses by oral supplementation of branched-chain amino acids in decompensated cirrhotic patients. <i>Hepatology Research</i> , 2007, 37, 1062-1067.	1.8	41
5	The Benefit of the Supplementation of Perioperative Branched-Chain Amino Acids in Patients with Surgical Management for Hepatocellular Carcinoma: a Preliminary Study. <i>Digestive Diseases and Sciences</i> , 2008, 53, 204-209.	1.1	16
6	Inhibitory effect of branched-chain amino acid granules on progression of compensated liver cirrhosis due to hepatitis C virus. <i>Journal of Gastroenterology</i> , 2008, 43, 63-70.	2.3	37
7	Does a late evening meal reduce the risk of hepatocellular carcinoma among patients with chronic hepatitis C?. <i>Hepatology Research</i> , 2008, 38, 860-868.	1.8	2
8	Supplement improves nutrition and stresses caused by examination-associated fasting in patients with liver cirrhosis. <i>Hepatology Research</i> , 2008, 38, 1178-1185.	1.8	6
9	Long-term outcome of branched-chain amino acid treatment in patients with liver cirrhosis. <i>Hepatology Research</i> , 2008, 38, S102-6.	1.8	37
10	Obesity Is an Independent Risk Factor for Hepatocellular Carcinoma Development in Chronic Hepatitis C Patients. <i>Clinical Gastroenterology and Hepatology</i> , 2008, 6, 459-464.	2.4	149
11	Hepatocellular Carcinoma in Keio Affiliated Hospitals – Diagnosis, Treatment, and Prognosis of this Disease –. <i>Keio Journal of Medicine</i> , 2009, 58, 161-175.	0.5	9
12	Supplementation with Branched-chain Amino Acids Inhibits Azoxymethane-induced Colonic Preneoplastic Lesions in Male C57BL/KsJ- <i>db/db</i> Mice. <i>Clinical Cancer Research</i> , 2009, 15, 3068-3075.	3.2	60
13	Branched chain amino acids enhance the maturation and function of myeloid dendritic cells ex vivo in patients with advanced cirrhosis. <i>Hepatology</i> , 2009, 50, 1936-1945.	3.6	56
14	A randomized pilot trial of oral branched-chain amino acids in early cirrhosis: Validation using prognostic markers for pre-liver transplant status. <i>Liver Transplantation</i> , 2009, 15, 790-797.	1.3	50
15	Branched-chain amino acids suppress insulin-resistance-based hepatocarcinogenesis in obese diabetic rats. <i>Journal of Gastroenterology</i> , 2009, 44, 483-491.	2.3	60
16	Diabetes pattern on the 75- <i>g</i> oral glucose tolerance test is a risk factor for hepatocellular carcinoma in patients with hepatitis C virus. <i>Liver International</i> , 2009, 29, 1194-1201.	1.9	39
17	1. Treatment of Hepatic Cirrhosis.. <i>The Journal of the Japanese Society of Internal Medicine</i> , 2010, 99, 2223-2229.	0.0	0
18	Attenuation of insulin-resistance-based hepatocarcinogenesis and angiogenesis by combined treatment with branched-chain amino acids and angiotensin-converting enzyme inhibitor in obese diabetic rats. <i>Journal of Gastroenterology</i> , 2010, 45, 443-450.	2.3	33

#	ARTICLE	IF	CITATIONS
19	Dietary supplementation with branched-chain amino acids suppresses diethylnitrosamine-induced liver tumorigenesis in obese and diabetic C57BL/KsJ-db/db mice. <i>Cancer Science</i> , 2010, 101, 460-467.	1.7	82
20	Effects of branched-chain amino acid-enriched nutrient for patients with hepatocellular carcinoma following radiofrequency ablation: A one-year prospective trial. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2010, 25, 1550-1555.	1.4	44
21	Importance of hepatitis C virus-associated insulin resistance: Therapeutic strategies for insulin sensitization. <i>World Journal of Gastroenterology</i> , 2010, 16, 1943.	1.4	55
22	Isoleucine Prevents the Accumulation of Tissue Triglycerides and Upregulates the Expression of PPAR α and Uncoupling Protein in Diet-Induced Obese Mice. <i>Journal of Nutrition</i> , 2010, 140, 496-500.	1.3	95
23	Changes in liver function parameters after percutaneous radiofrequency ablation therapy in patients with hepatocellular carcinoma. <i>Hepatology Research</i> , 2010, 40, 550-554.	1.8	17
24	Insulin resistance raises the risk for recurrence of stage I hepatocellular carcinoma after curative radiofrequency ablation in hepatitis C virus-positive patients: A prospective, case series study. <i>Hepatology Research</i> , 2010, 40, 376-382.	1.8	43
25	Guidelines for the treatment of chronic hepatitis and cirrhosis due to hepatitis B virus infection for the fiscal year 2008 in Japan. <i>Hepatology Research</i> , 2010, 40, 1-7.	1.8	78
26	Guidelines for the treatment of chronic hepatitis and cirrhosis due to hepatitis C virus infection for the fiscal year 2008 in Japan. <i>Hepatology Research</i> , 2010, 40, 8-13.	1.8	92
27	Chapter 2: Diagnosis and surveillance. <i>Hepatology Research</i> , 2010, 40, 16-47.	1.8	15
28	Effects of late evening snack on diurnal plasma glucose profile in patients with chronic viral liver disease. <i>Hepatology Research</i> , 2010, 40, 887-893.	1.8	7
29	Effects of late evening snack including branched-chain amino acid on the function of hepatic parenchymal cells in patients with liver cirrhosis. <i>Hepatology Research</i> , 2011, 41, 417-422.	1.8	18
30	Sustained viral response (SVR) in type C liver cirrhosis patients undergoing peginterferon α -2b/ribavirin combination therapy after partial splenic embolization. <i>Acta Hepatologica Japonica</i> , 2011, 52, 26-35.	0.0	0
31	Increased levels of serum leptin are a risk factor for the recurrence of stage I/II hepatocellular carcinoma after curative treatment. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2011, 49, 153-158.	0.6	35
32	Pitavastatin suppresses diethylnitrosamine-induced liver preneoplasms in male C57BL/KsJ-db/db obese mice. <i>BMC Cancer</i> , 2011, 11, 281.	1.1	45
33	Cancer chemoprevention with green tea catechins by targeting receptor tyrosine kinases. <i>Molecular Nutrition and Food Research</i> , 2011, 55, 832-843.	1.5	105
34	Branched-chain amino acids as pharmacological nutrients in chronic liver disease. <i>Hepatology</i> , 2011, 54, 1063-1070.	3.6	276
35	Serum level of adiponectin and the risk of liver cancer development in chronic hepatitis C patients. <i>International Journal of Cancer</i> , 2011, 129, 2226-2235.	2.3	70
36	Preventive Effects of (â)-Epigallocatechin Gallate on Diethylnitrosamine-Induced Liver Tumorigenesis in Obese and Diabetic C57BL/KsJ-db/db Mice. <i>Cancer Prevention Research</i> , 2011, 4, 396-403.	0.7	76

#	ARTICLE	IF	CITATIONS
37	Oral Branched-Chain Amino Acid Granules Reduce the Incidence of Hepatocellular Carcinoma and Improve Event-Free Survival in Patients with Liver Cirrhosis. <i>Digestive Diseases</i> , 2011, 29, 326-332.	0.8	57
38	Acyclic retinoid in chemoprevention of hepatocellular carcinoma: Targeting phosphorylated retinoid X receptor-1 \pm for prevention of liver carcinogenesis. <i>Journal of Carcinogenesis</i> , 2012, 11, 11.	2.5	23
39	Nutraceutical Approach for Preventing Obesity-Related Colorectal and Liver Carcinogenesis. <i>International Journal of Molecular Sciences</i> , 2012, 13, 579-595.	1.8	37
40	Nutrition Therapy for Liver Diseases Based on the Status of Nutritional Intake. <i>Gastroenterology Research and Practice</i> , 2012, 2012, 1-8.	0.7	21
41	Clofibrate-Induced Reduction of Plasma Branched-Chain Amino Acid Concentrations Impairs Glucose Tolerance in Rats. <i>Journal of Parenteral and Enteral Nutrition</i> , 2012, 36, 337-343.	1.3	20
42	Valine, a Branched-Chain Amino Acid, Reduced HCV Viral Load and Led to Eradication of HCV by Interferon Therapy in a Decompensated Cirrhotic Patient. <i>Case Reports in Gastroenterology</i> , 2012, 6, 660-667.	0.3	11
43	Risk of Hepatocellular Carcinoma Development in Cases of Hepatitis C Treated by Long-Term, Low-Dose PEG-IFN1 \pm -2a. <i>Digestive Diseases</i> , 2012, 30, 561-567.	0.8	2
44	Hepatocellular carcinoma in nonalcoholic steatohepatitis: Growing evidence of an epidemic?. <i>Hepatology Research</i> , 2012, 42, 1-14.	1.8	106
45	Long-term treatment with l-isoleucine or l-leucine in AIN-93G diet has promoting effects on rat bladder carcinogenesis. <i>Food and Chemical Toxicology</i> , 2012, 50, 3934-3940.	1.8	14
46	Assessment of portal hypertension by transient elastography in patients with compensated cirrhosis and potentially resectable liver tumors. <i>Journal of Hepatology</i> , 2012, 56, 103-108.	1.8	142
47	Branched-chain amino acid deficiency stabilizes insulin-induced vascular endothelial growth factor mRNA in hepatocellular carcinoma cells. <i>Journal of Cellular Biochemistry</i> , 2012, 113, 3113-3121.	1.2	23
48	Nutritional support for liver disease. <i>The Cochrane Library</i> , 2012, , CD008344.	1.5	79
49	Combination of acyclic retinoid with branched-chain amino acids inhibits xenograft growth of human hepatoma cells in nude mice. <i>Hepatology Research</i> , 2012, 42, 1241-1247.	1.8	4
50	Early administration of branched-chain amino acid granules. <i>World Journal of Gastroenterology</i> , 2012, 18, 4486.	1.4	13
51	Clinical factors related to long-term administration of sorafenib in patients with hepatocellular carcinoma. <i>Cancer Management and Research</i> , 2012, 4, 423.	0.9	7
52	Guidelines on nutritional management in Japanese patients with liver cirrhosis from the perspective of preventing hepatocellular carcinoma. <i>Hepatology Research</i> , 2012, 42, 621-626.	1.8	37
53	Branched-chain amino acids prevent insulin-induced hepatic tumor cell proliferation by inducing apoptosis through mTORC1 and mTORC2-dependent mechanisms. <i>Journal of Cellular Physiology</i> , 2012, 227, 2097-2105.	2.0	57
54	Obesity and hepatocellular carcinoma: targeting obesity-related inflammation for chemoprevention of liver carcinogenesis. <i>Seminars in Immunopathology</i> , 2013, 35, 191-202.	2.8	48

#	ARTICLE	IF	CITATIONS
55	Oral supplementation of branched-chain amino acids reduces early recurrence after hepatic resection in patients with hepatocellular carcinoma: a prospective study. <i>Surgery Today</i> , 2013, 43, 720-726.	0.7	66
56	Nutritional assessment and treatment of patients with liver cirrhosis. <i>Nutrition</i> , 2013, 29, 1279-1285.	1.1	53
57	Long-term Branched Chain Amino Acid Supplementation Ameliorates Diethylnitrosamine-induced Liver Glutathione S-transferase-p Positivity in Zucker Fatty Rats. <i>Journal of Clinical and Experimental Hepatology</i> , 2013, 3, 192-197.	0.4	4
58	Effects of Oral Branched-Chain Amino Acids on Hepatic Encephalopathy and Outcome in Patients With Liver Cirrhosis. <i>Nutrition in Clinical Practice</i> , 2013, 28, 580-588.	1.1	58
59	Diuretics aggravate zinc deficiency in patients with liver cirrhosis by increasing zinc excretion in urine. <i>Hepatology Research</i> , 2013, 43, 365-373.	1.8	33
60	A targeted metabolomic protocol for short-chain fatty acids and branched-chain amino acids. <i>Metabolomics</i> , 2013, 9, 818-827.	1.4	212
61	DNA microarray analysis identified molecular pathways mediating the effects of supplementation of branched-chain amino acids on CCl ₄ -induced cirrhosis in rats. <i>Molecular Nutrition and Food Research</i> , 2013, 57, 291-306.	1.5	14
62	Branched-chain amino acids suppress the cumulative recurrence of hepatocellular carcinoma under conditions of insulin-resistance. <i>Oncology Reports</i> , 2013, 30, 545-552.	1.2	17
63	Incidence, Mortality, and Predictive Factors of Hepatocellular Carcinoma in Primary Biliary Cirrhosis. <i>Gastroenterology Research and Practice</i> , 2013, 2013, 1-8.	0.7	11
64	Effects of branched-chain amino acid granules on serum albumin level and prognosis are dependent on treatment adherence in patients with liver cirrhosis. <i>Hepatology Research</i> , 2013, 43, 459-466.	1.8	11
65	Nutritional status and quality of life in current patients with liver cirrhosis as assessed in 2007-2011. <i>Hepatology Research</i> , 2013, 43, 106-112.	1.8	63
66	Body Mass Index and Weight Change During Adulthood Are Associated With Increased Mortality From Liver Cancer: The JACC Study. <i>Journal of Epidemiology</i> , 2013, 23, 219-226.	1.1	13
67	The Effect of Long-term Supplementation With Branched-chain Amino Acid Granules in Patients With Hepatitis C Virus-related Hepatocellular Carcinoma After Radiofrequency Thermal Ablation. <i>Journal of Clinical Gastroenterology</i> , 2013, 47, 359-366.	1.1	36
68	Branched-chain amino acids in liver diseases. <i>World Journal of Gastroenterology</i> , 2013, 19, 7620.	1.4	152
69	Branched-Chain Amino Acid Supplementation Reduces Oxidative Stress and Prolongs Survival in Rats with Advanced Liver Cirrhosis. <i>PLoS ONE</i> , 2013, 8, e70309.	1.1	50
70	Branched-Chain Amino Acids Ameliorate Fibrosis and Suppress Tumor Growth in a Rat Model of Hepatocellular Carcinoma with Liver Cirrhosis. <i>PLoS ONE</i> , 2013, 8, e77899.	1.1	39
71	Branched-Chain Amino Acids Enhance Premature Senescence through Mammalian Target of Rapamycin Complex I-Mediated Upregulation of p21 Protein. <i>PLoS ONE</i> , 2013, 8, e80411.	1.1	32
72	Insulin resistance as early sign of hepatic dysfunction in liver cirrhosis. <i>Journal of Medical Investigation</i> , 2014, 61, 180-189.	0.2	18

#	ARTICLE	IF	CITATIONS
73	Impact of Serum Chemerin Levels on Liver Functional Reserves and Platelet Counts in Patients with Hepatocellular Carcinoma. <i>International Journal of Molecular Sciences</i> , 2014, 15, 11294-11306.	1.8	26
74	Clinical significance of therapy using branched-chain amino acid granules in patients with liver cirrhosis and hepatocellular carcinoma. <i>Hepatology Research</i> , 2014, 44, 149-158.	1.8	34
75	Oral supplementation with branched-chain amino acid granules prevents hepatocarcinogenesis in patients with hepatitis C-related cirrhosis: A propensity score analysis. <i>Hepatology Research</i> , 2014, 44, 288-295.	1.8	10
76	Serum albumin level is a notable profiling factor for non-B, non-C hepatitis virus-related hepatocellular carcinoma: A data-mining analysis. <i>Hepatology Research</i> , 2014, 44, 837-845.	1.8	26
77	Excess visceral adiposity induces alterations in mitochondrial function and energy metabolism in esophageal adenocarcinoma. <i>BMC Cancer</i> , 2014, 14, 907.	1.1	21
78	Pharmaceutical and nutraceutical approaches for preventing liver carcinogenesis: Chemoprevention of hepatocellular carcinoma using acyclic retinoid and branched-chain amino acids. <i>Molecular Nutrition and Food Research</i> , 2014, 58, 124-135.	1.5	8
79	Branched-chain amino acids in metabolic signalling and insulin resistance. <i>Nature Reviews Endocrinology</i> , 2014, 10, 723-736.	4.3	1,006
80	Insight into the impact of diabetes mellitus on the increased risk of hepatocellular carcinoma: mini-review. <i>Journal of Diabetes and Metabolic Disorders</i> , 2014, 13, 57.	0.8	25
81	Effect of zinc on liver cirrhosis with hyperammonemia: A preliminary randomized, placebo-controlled double-blind trial. <i>Nutrition</i> , 2014, 30, 1409-1414.	1.1	67
82	Branched-Chain Amino Acids Prevent Hepatocarcinogenesis and Prolong Survival of Patients With Cirrhosis. <i>Clinical Gastroenterology and Hepatology</i> , 2014, 12, 1012-1018.e1.	2.4	84
83	Efficacy and safety of oral branched-chain amino acid supplementation in patients undergoing interventions for hepatocellular carcinoma: a meta-analysis. <i>Nutrition Journal</i> , 2015, 14, 67.	1.5	37
84	Impact of the branched-chain amino acid to tyrosine ratio and branched-chain amino acid granule therapy in patients with hepatocellular carcinoma: A propensity score analysis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2015, 30, 1412-1419.	1.4	24
85	Changing epidemiology of hepatocellular carcinoma in Asia. <i>Bailliere's Best Practice and Research in Clinical Gastroenterology</i> , 2015, 29, 919-928.	1.0	121
86	Chemopreventive Potential of Green Tea Catechins in Hepatocellular Carcinoma. <i>International Journal of Molecular Sciences</i> , 2015, 16, 6124-6139.	1.8	42
87	Effect of branched-chain amino acid supplementation on functional liver regeneration in patients undergoing portal vein embolization and sequential hepatectomy: a randomized controlled trial. <i>Journal of Gastroenterology</i> , 2015, 50, 1197-1205.	2.3	38
88	Prognostic Impact of Visceral Fat Amount and Branched-Chain Amino Acids (BCAA) in Hepatocellular Carcinoma. <i>Annals of Surgical Oncology</i> , 2015, 22, 1041-1047.	0.7	21
89	Postoperative Outcomes for Patients with Non-B Non-C Hepatocellular Carcinoma: A Subgroup Analysis of Patients with a History of Hepatitis B Infection. <i>Annals of Surgical Oncology</i> , 2015, 22, 1034-1040.	0.7	26
90	Branched-chain amino acids reduce hepatic iron accumulation and oxidative stress in hepatitis C virus polyprotein-expressing mice. <i>Liver International</i> , 2015, 35, 1303-1314.	1.9	23

#	ARTICLE	IF	CITATIONS
91	Two randomized controlled studies comparing the nutritional benefits of branched-chain amino acid (BCAA) granules and a BCAA-enriched nutrient mixture for patients with esophageal varices after endoscopic treatment. <i>Journal of Gastroenterology</i> , 2015, 50, 109-118.	2.3	15
92	Body mass index and cholesterol level predict surgical outcome in patients with hepatocellular carcinoma in Taiwan - a cohort study. <i>Oncotarget</i> , 2016, 7, 22948-22959.	0.8	35
93	Impact of branched-chain amino acid supplementation on survival in patients with advanced hepatocellular carcinoma treated with sorafenib: A multicenter retrospective cohort study. <i>Hepatology Research</i> , 2016, 46, 1002-1010.	1.8	21
94	Role of Nutrition and Muscle in Cirrhosis. <i>Current Treatment Options in Gastroenterology</i> , 2016, 14, 257-273.	0.3	40
95	Tumour biology of obesity-related cancers: understanding the molecular concept for better diagnosis and treatment. <i>Tumor Biology</i> , 2016, 37, 14363-14380.	0.8	21
96	Leucine and Mammalian Target of Rapamycin-Dependent Activation of Muscle Protein Synthesis in Aging. <i>Journal of Nutrition</i> , 2016, 146, 2616S-2624S.	1.3	42
97	Impact of Branched-Chain Amino Acid-Enriched Nutrient on liver Cirrhosis with Hepatocellular Carcinoma Undergoing Transcatheter Arterial Chemoembolization in Barcelona Clinic Liver Cancer Stage B: A Prospective Study. <i>Journal of Nippon Medical School</i> , 2016, 83, 248-256.	0.3	14
98	Overweight in late adolescence predicts development of severe liver disease later in life: A 39years follow-up study. <i>Journal of Hepatology</i> , 2016, 65, 363-368.	1.8	68
99	Evidence-based clinical practice guidelines for liver cirrhosis 2015. <i>Journal of Gastroenterology</i> , 2016, 51, 629-650.	2.3	253
100	Branched-chain amino acids alleviate hepatic steatosis and liver injury in choline-deficient high-fat diet induced NASH mice. <i>Metabolism: Clinical and Experimental</i> , 2017, 69, 177-187.	1.5	80
101	Liver Transplantation and Bariatric Surgery. <i>Clinics in Liver Disease</i> , 2017, 21, 215-230.	1.0	21
102	Nutrition and the transplant candidate. <i>Liver Transplantation</i> , 2017, 23, 1451-1464.	1.3	46
103	Efficacy of branched-chain amino acid supplementation and walking exercise for preventing sarcopenia in patients with liver cirrhosis. <i>European Journal of Gastroenterology and Hepatology</i> , 2017, 29, 1416-1423.	0.8	113
104	Nutrition for the Patient with Advanced Liver Disease. <i>Current Hepatology Reports</i> , 2017, 16, 205-211.	0.4	0
105	Effects of branched-chain amino acids (BCAAs) on the progression of advanced liver disease. <i>Medicine (United States)</i> , 2017, 96, e6580.	0.4	45
106	Effects of branched-chain amino acid supplementation after radiofrequency ablation for hepatocellular carcinoma: A randomized trial. <i>Nutrition</i> , 2017, 33, 20-27.	1.1	37
107	Branched-chain amino acids differently modulate catabolic and anabolic states in mammals: a pharmacological point of view. <i>British Journal of Pharmacology</i> , 2017, 174, 1366-1377.	2.7	107
108	Does Nutrition Matter in Liver Disease?. , 2017, , 743-759.		2

#	ARTICLE	IF	CITATIONS
109	Nutritional Status in Liver Cirrhosis. , 2017, , .		1
110	Antiangiogenic Therapy for Hepatocellular Carcinoma. , 2017, , .		0
111	Epidemic of non-alcoholic fatty liver disease and hepatocellular carcinoma. World Journal of Clinical Oncology, 2017, 8, 429-436.	0.9	73
112	The profiling of plasma free amino acids and the relationship between serum albumin and plasma-branched chain amino acids in chronic liver disease: a single-center retrospective study. Journal of Gastroenterology, 2018, 53, 978-988.	2.3	17
113	The Value of the Câ€Reactive Proteinâ€toâ€Albumin Ratio is Useful for Predicting Survival of Patients with Childâ€Pugh Class A Undergoing Liver Resection for Hepatocellular Carcinoma. World Journal of Surgery, 2018, 42, 2218-2226.	0.8	21
114	Metabonomics of <sc>d</sc>-glucaro-1,4-lactone in preventing diethylnitrosamine-induced liver cancer in rats. Pharmaceutical Biology, 2018, 56, 643-648.	1.3	16
115	Is the Response of Tumours Dependent on the Dietary Input of Some Amino Acids or Ratios among Essential and Non-Essential Amino Acids? All That Glitters Is Not Gold. International Journal of Molecular Sciences, 2018, 19, 3631.	1.8	3
116	Microbially Produced Imidazole Propionate Impairs Insulin Signaling through mTORC1. Cell, 2018, 175, 947-961.e17.	13.5	517
117	Secondary Unconjugated Bile Acids Induce Hepatic Stellate Cell Activation. International Journal of Molecular Sciences, 2018, 19, 3043.	1.8	18
118	Non-invasive urinary metabolomic profiles discriminate biliary atresia from infantile hepatitis syndrome. Metabolomics, 2018, 14, 90.	1.4	10
119	Obesity and recurrenceâ€free survival in patients with hepatocellular carcinoma after achieving sustained virological response to interferon therapy for chronic hepatitis C. Annals of Gastroenterological Surgery, 2018, 2, 319-326.	1.2	10
120	Nutrition is often ignored in management of chronic liver diseases. Journal of Gastroenterology and Hepatology (Australia), 2019, 34, 1127-1128.	1.4	1
121	Obesity and Hepatocarcinogenesis. , 2019, , 87-102.		0
122	Late evening snacks with branched-chain amino acids improve the Fischer ratio with patients liver cirrhosis at fasting in the next morning. Clinical Nutrition ESPEN, 2019, 30, 138-144.	0.5	14
123	Prevalence of diabetes in liver cirrhosis: A systematic review and metaâ€analysis. Diabetes/Metabolism Research and Reviews, 2019, 35, e3157.	1.7	42
124	Impact of Branched-Chain Amino Acid Granule Therapy in Patients with Hepatocellular Carcinoma Who Have Normal Albumin Levels and Low Branched-Chain Amino Acid to Tyrosine Ratios. Nutrition and Cancer, 2019, 71, 1132-1141.	0.9	11
125	Effects of Zinc Acetate on Serum Zinc Concentrations in Chronic Liver Diseases: a Multicenter, Double-Blind, Randomized, Placebo-Controlled Trial and a Dose Adjustment Trial. Biological Trace Element Research, 2020, 195, 71-81.	1.9	14
126	Anti-obesity effects of Î±-amylase inhibitor enriched-extract from white common beans (<i>Phaseolus</i> Tj ETQq1 1 0.784314 rgBT /Over obese rats. Food and Function, 2020, 11, 1624-1634.	2.1	41

#	ARTICLE	IF	CITATIONS
127	Zinc and protein metabolism in chronic liver diseases. <i>Nutrition Research</i> , 2020, 74, 1-9.	1.3	48
128	Supplementation of branched-chain amino acids decreases fat accumulation in the liver through intestinal microbiota-mediated production of acetic acid. <i>Scientific Reports</i> , 2020, 10, 18768.	1.6	31
129	Dietary Approaches to Cancer Therapy. <i>Cancer Cell</i> , 2020, 37, 767-785.	7.7	105
130	Diet and liver cancer risk: a narrative review of epidemiological evidence. <i>British Journal of Nutrition</i> , 2020, 124, 330-340.	1.2	95
131	Current and new pharmacotherapy options for non-alcoholic steatohepatitis. <i>Expert Opinion on Pharmacotherapy</i> , 2020, 21, 953-967.	0.9	28
132	A Comprehensive Review Evaluating the Impact of Protein Source (Vegetarian vs. Meat Based) in Hepatic Encephalopathy. <i>Nutrients</i> , 2021, 13, 370.	1.7	7
133	A hexokinase isoenzyme switch in human liver cancer cells promotes lipogenesis and enhances innate immunity. <i>Communications Biology</i> , 2021, 4, 217.	2.0	21
134	Sarcopenic Obesity in Liver Cirrhosis: Possible Mechanism and Clinical Impact. <i>International Journal of Molecular Sciences</i> , 2021, 22, 1917.	1.8	25
135	Malnutrition in Patients with Liver Cirrhosis. <i>Nutrients</i> , 2021, 13, 540.	1.7	57
136	Increased Visceral Adipose Tissue and Hyperinsulinemia Raise the Risk for Recurrence of Non-B Non-C Hepatocellular Carcinoma after Curative Treatment. <i>Cancers</i> , 2021, 13, 1542.	1.7	9
137	Obesity and Liver Cancer in Japan: A Comprehensive Review. <i>Anticancer Research</i> , 2021, 41, 2227-2237.	0.5	5
138	A novel, multitargeted endogenous metabolic modulator composition impacts metabolism, inflammation, and fibrosis in nonalcoholic steatohepatitis-relevant primary human cell models. <i>Scientific Reports</i> , 2021, 11, 11861.	1.6	10
139	Superiority of CRP-albumin-lymphocyte index (CALLY index) as a non-invasive prognostic biomarker after hepatectomy for hepatocellular carcinoma. <i>Hpb</i> , 2022, 24, 101-115.	0.1	23
140	The Current View of Nonalcoholic Fatty Liver Disease-Related Hepatocellular Carcinoma. <i>Cancers</i> , 2021, 13, 516.	1.7	15
141	Assessment of Nutritional Status of Patients with Chronic Hepatitis C and HCV-Related Cirrhosis in the Compensated Stage. <i>American Journal of Internal Medicine</i> , 2016, 4, 24.	0.1	1
142	Branched Chain Amino Acid Suppresses Hepatocellular Cancer Stem Cells through the Activation of Mammalian Target of Rapamycin. <i>PLoS ONE</i> , 2013, 8, e82346.	1.1	34
143	Metformin Suppresses Diethylnitrosamine-Induced Liver Tumorigenesis in Obese and Diabetic C57BL/KsJ-+Leprdb/+Leprdb Mice. <i>PLoS ONE</i> , 2015, 10, e0124081.	1.1	22
144	Feasibility Study of NMR Based Serum Metabolomic Profiling to Animal Health Monitoring: A Case Study on Iron Storage Disease in Captive Sumatran Rhinoceros (<i>Dicerorhinus sumatrensis</i>). <i>PLoS ONE</i> , 2016, 11, e0156318.	1.1	10

#	ARTICLE	IF	CITATIONS
145	The Prevalence and Implication of Zinc Deficiency in Patients With Chronic Liver Disease. <i>Journal of Clinical Medicine Research</i> , 2018, 10, 437-444.	0.6	47
146	Branched-chain amino acids prevent hepatic fibrosis and development of hepatocellular carcinoma in a non-alcoholic steatohepatitis mouse model. <i>Oncotarget</i> , 2017, 8, 18191-18205.	0.8	59
147	Preventive effects of the sodium glucose cotransporter 2 inhibitor tofogliflozin on diethylnitrosamine-induced liver tumorigenesis in obese and diabetic mice. <i>Oncotarget</i> , 2017, 8, 58353-58363.	0.8	47
148	Increased visceral fat volume raises the risk for recurrence of hepatocellular carcinoma after curative treatment. <i>Oncotarget</i> , 2018, 9, 14058-14067.	0.8	15
149	Sodium alginate prevents progression of non-alcoholic steatohepatitis and liver carcinogenesis in obese and diabetic mice. <i>Oncotarget</i> , 2016, 7, 10448-10458.	0.8	22
150	Inflammation- and stress-related signaling pathways in hepatocarcinogenesis. <i>World Journal of Gastroenterology</i> , 2012, 18, 4071.	1.4	115
151	Nutritional status in relation to lifestyle in patients with compensated viral cirrhosis. <i>World Journal of Gastroenterology</i> , 2012, 18, 5759.	1.4	32
152	Impairment of innate immune responses in cirrhotic patients and treatment by branched-chain amino acids. <i>World Journal of Gastroenterology</i> , 2014, 20, 7298.	1.4	24
153	Chemoprevention of colorectal cancer by targeting obesity-related metabolic abnormalities. <i>World Journal of Gastroenterology</i> , 2014, 20, 8939-46.	1.4	14
154	Chemoprevention of obesity-related liver carcinogenesis by using pharmaceutical and nutraceutical agents. <i>World Journal of Gastroenterology</i> , 2016, 22, 394.	1.4	12
155	Supportive therapies for prevention of hepatocellular carcinoma recurrence and preservation of liver function. <i>World Journal of Gastroenterology</i> , 2016, 22, 7252.	1.4	15
156	Nutrition and metabolism in hepatocellular carcinoma. <i>Hepatobiliary Surgery and Nutrition</i> , 2013, 2, 89-96.	0.7	12
157	Effect of zinc supplementation in patients with type C liver cirrhosis. <i>Open Journal of Gastroenterology</i> , 2011, 01, 28-34.	0.1	10
158	Treating morbid obesity in cirrhosis: A quest of holy grail. <i>World Journal of Hepatology</i> , 2015, 7, 2819.	0.8	17
159	Hepatocellular Carcinoma in Asia: A Challenging Situation. <i>Euroasian Journal of Hepato-gastroenterology</i> , 2019, 9, 27-33.	0.1	23
160	Comparison of measured and predicted energy expenditure in patients with liver cirrhosis. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2014, 23, 197-204.	0.3	13
161	Insulin resistance as early sign of hepatic dysfunction in liver cirrhosis . <i>Journal of Medical Investigation</i> , 2000, 40, 180-189.	0.2	0
162	The effect of hepatic steatosis on Peginterferon alfa-2b and Ribavirin combination therapy for chronic hepatitis C. <i>Acta Hepatologica Japonica</i> , 2008, 49, 183-191.	0.0	2

#	ARTICLE	IF	CITATIONS
163	Effects of Branched-chain Amino Acid Supplementation in Rats Fed a Vitamin B₆-deficient Diet. Nihon Eiyō-Shokuryō Gakkai Shi = Nippon Eiyō-Shokuryō Gakkaishi = Journal of Japanese Society of Nutrition and Food Science, 2012, 65, 65-73.	0.2	1
164	æ,,éŠæŠ•ăŽă«ă,ă,ç™CEă†...ă¢—æ®—æ©Ÿæš«ă®ă^†ă³/4j. The Japanese Journal of SURGICAL METABOLISM and NUTRITION, 2013, 47,		
165	é«~é1/2¢è€...è,ă^†é™ă«ăšăă,ă“èj“æœŸæ,,éŠç®jç†ă®ă•éjCEç,1. The Japanese Journal of SURGICAL METABOLISM and NUTRITION, 2014, 48,		
166	Identification of Branched Chain Amino Acids; Underlying Molecular Pathways Using Transcriptomic Analysis: Application to Cirrhosis. , 2015, , 141-157.		0
167	Branched-chain amino acid metabolism and insulin resistance. The Japanese Journal of SURGICAL METABOLISM and NUTRITION, 2015, 49, 177-182.	0.1	0
168	Metabolic Syndrome and the Liver. , 2016, , 149-177.		0
169	Three step primary liver cancer prevention program utilizing dynamic tumor marker combination assay in high-risk patients with chronic hepatitis. MOJ Current Research & Reviews, 2018, 1, 114-117.	0.1	0
170	Nutrition in Liver Cirrhosis. , 2019, , 69-77.		0
172	Bariatric Surgery and Liver Transplantation. Gastroenterology and Hepatology, 2017, 13, 170-175.	0.2	2
173	ANXA2, PRKCE, and OXT are critical differentially genes in Nonalcoholic fatty liver disease. Gastroenterology and Hepatology From Bed To Bench, 2019, 12, 131-137.	0.6	6
174	Favorable and harmful effects of branched-chain amino acid administration in patients with liver cirrhosis: effects on hepatic encephalopathy and ammonia metabolism. Acta Hepatologica Japonica, 2022, 63, 1-8.	0.0	0
175	Therapeutic mechanisms and beneficial effects of non-antidiabetic drugs in chronic liver diseases. Clinical and Molecular Hepatology, 2022, 28, 425-472.	4.5	14
176	The Complex of Phycobiliproteins, Fucoxanthin, and Krill Oil Ameliorates Obesity through Modulation of Lipid Metabolism and Antioxidants in Obese Rats. Nutrients, 2022, 14, 4815.	1.7	1
177	Recent Progress of Nutritional Management for Liver Cirrhosis. Journal of Nutritional Science and Vitaminology, 2022, 68, S73-S75.	0.2	0
178	Metabolomics and Lipidomics Screening Reveal Reprogrammed Signaling Pathways toward Cancer Development in Non-Alcoholic Steatohepatitis. International Journal of Molecular Sciences, 2023, 24, 210.	1.8	9
179	VII. Nutrition and Exercise Therapies for Liver Cirrhosis. The Journal of the Japanese Society of Internal Medicine, 2022, 111, 50-57.	0.0	0