CITATION REPORT List of articles citing

Effects of systemic steroids in patients with severe community-acquired pneumonia

DOI: 10.1183/09031936.00027607 European Respiratory Journal, 2007, 30, 951-6.

Source: https://exaly.com/paper-pdf/41626889/citation-report.pdf

Version: 2024-04-28

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
1175	Pneumonia in elderly patients with chronic obstructive pulmonary disease. <i>Current Infectious Disease Reports</i> , 2008 , 10, 223-8	3.9	7
1174	Pneumonia in the emergency department. 2008 , 26, 389-411, ix		8
1173	The role of corticosteroids in severe community-acquired pneumonia: a systematic review. <i>Critical Care</i> , 2008 , 12, R76	10.8	54
1172	Corticosteroids in severe pneumonia. European Respiratory Journal, 2008, 32, 259-64	13.6	38
1171	Novel targets in the management of pneumonia. 2008 , 2, 387-400		6
1170	Early mortality in patients with community-acquired pneumonia: causes and risk factors. <i>European Respiratory Journal</i> , 2008 , 32, 733-9	13.6	83
1169	Bibliography. Current world literature. Infectious diseases. 2008 , 14, 266-73		
1168	Brazilian guidelines for community-acquired pneumonia in immunocompetent adults - 2009. 2009 , 35, 574-601		37
1167	Community-acquired pneumonia in elderly patients. 2009 , 5, 763-774		78
1166	Early and late treatment failure in community-acquired pneumonia. <i>Seminars in Respiratory and Critical Care Medicine</i> , 2009 , 30, 154-60	3.9	28
1165	Prolonged Physiologic Doses of Corticosteroids for the Treatment of Severe Community Acquired Pneumonia. 2009 , 5, 90-95		1
1164	Bacteraemic pneumococcal pneumonia in COPD patients: better outcomes than expected. 2009 , 28, 971-6		10
1163	Mortality in ICU patients with bacterial community-acquired pneumonia: when antibiotics are not enough. <i>Intensive Care Medicine</i> , 2009 , 35, 430-8	14.5	65
1162	Community-acquired pneumonia in patients with and without chronic obstructive pulmonary disease. <i>Journal of Infection</i> , 2009 , 58, 417-24	18.9	84
1161	Time course of dyspnea evolution in the emergency department: results from the URGENT dyspnea survey. <i>Critical Care</i> , 2009 , 13, P1	10.8	1
1160	Endotracheal intubation in the emergency room: a multicenter questionnaire. <i>Critical Care</i> , 2009 , 13, P2	10.8	78
1159	Radiological validation of endotracheal tube insertion depth in prehospital emergency patients. <i>Critical Care</i> , 2009 , 13, P3	10.8	78

[2009-2009]

1158	Does bedside chest ultrasound in the ICU improve early diagnosis and quick resolution of pleural effusion?. <i>Critical Care</i> , 2009 , 13, P4	10.8	78
1157	Portable chest radiography in mechanically ventilated ICU patients: does synchronizing with end-inspiration improve the quality of films?. <i>Critical Care</i> , 2009 , 13, P5	10.8	78
1156	Quality assurance report on the use of continuous positive airway pressure and end-tidal carbon dioxide during respiratory distress in field emergency care. <i>Critical Care</i> , 2009 , 13, P6	10.8	1
1155	Cardiogenic oscillations extracted from spontaneous breathing airway pressure and flow signal are related to chest wall motility and continuous positive airway pressure. <i>Critical Care</i> , 2009 , 13, P7	10.8	78
1154	Noninvasive positive pressure ventilation in infants with respiratory failure. <i>Critical Care</i> , 2009 , 13, P8	10.8	78
1153	Functional respiratory effects of noninvasive ventilation in acute hypercapnic patients with chronic obstructive pulmonary disease. <i>Critical Care</i> , 2009 , 13, P9	10.8	78
1152	Clinical application of noninvasive ventilation in acute respiratory failure in a general ICU. <i>Critical Care</i> , 2009 , 13, P10	10.8	78
1151	Relative effectiveness of two nasal continuous positive airway pressure devices in VLBW infants: first report from a multicenter, randomized, controlled trial. <i>Critical Care</i> , 2009 , 13, P11	10.8	78
1150	Contribution of noninvasive ventilation in the precocious extubation in the medical ICU. <i>Critical Care</i> , 2009 , 13, P12	10.8	2
1149	Survey of current intubation practices in Polish neonatal and pediatric ICUs. <i>Critical Care</i> , 2009 , 13, P13	10.8	78
1148	Incidence of post-intubation hemodynamic instability associated with emergent endotracheal intubations: a systematic review. <i>Critical Care</i> , 2009 , 13, P14	10.8	2
1147	A novel method to develop an elastic, thin-walled, leak-proof, inflatable tracheal tube cuff. <i>Critical Care</i> , 2009 , 13, P15	10.8	78
1146	Abstract withdrawn. <i>Critical Care</i> , 2009 , 13, P16	10.8	78
1145	Influence of tracheostomy on duration of weaning from mechanical ventilation. <i>Critical Care</i> , 2009 , 13, P17	10.8	78
1144	Safety of percutaneous tracheotomy in patients with cricoid cartilage not identified: report of 122 cases. <i>Critical Care</i> , 2009 , 13, P18	10.8	78
1143	Percutaneous dilational tracheostomy: early and late complications. <i>Critical Care</i> , 2009 , 13, P19	10.8	78
1142	Tracheostomy in the ICU: an analysis of 443 procedures. <i>Critical Care</i> , 2009 , 13, P20	10.8	78
1141	Percutaneous dilational tracheostomy in neurointensive care patients. <i>Critical Care</i> , 2009 , 13, P21	10.8	78

	Service evaluation of complications following tracheostomy insertion in ICU patients. <i>Critical Care</i> , 2009 , 13, P22	10.8	78
1139	Risk factors for unplanned extubation in critically ill patients. <i>Critical Care</i> , 2009 , 13, P23	10.8	O
1138	Protocol-driven weaning from mechanical ventilation: a study into adherence and outcomes. <i>Critical Care</i> , 2009 , 13, P24	10.8	1
1137	SmartCare is faster than paper-protocol weaning. <i>Critical Care</i> , 2009 , 13, P25	10.8	78
1136	Respiratory muscle oxygen saturation during weaning. Critical Care, 2009, 13, P26	10.8	78
	Early and continuous weaning from mechanical ventilation without formal protocols in a university hospital. <i>Critical Care</i> , 2009 , 13, P27	10.8	78
1134	Ventilator dependency among morbidly obese in the ICU. Critical Care, 2009, 13, P28	10.8	78
	Comparison of a novel humidifier with two conventional humidifiers during high-frequency oscillatory ventilation. <i>Critical Care</i> , 2009 , 13, P29	10.8	О
	A new in vitro model for force measurements at the isolated entire rat diaphragm. <i>Critical Care</i> , 2009 , 13, P30	10.8	78
	Adaptive support ventilation prevents ventilator-induced diaphragmatic dysfunction: an in vivo piglet study. <i>Critical Care</i> , 2009 , 13, P31	10.8	1
1130	Characterization of the mechanical ventilator adjustment process. Critical Care, 2009, 13, P32	10.8	78
	Impact of a dedicated ventilatory support team on how mechanical ventilation is employed in a tertiary-care hospital. <i>Critical Care</i> , 2009 , 13, P33	10.8	78
1128	Hyperoxia in mechanically ventilated patients. Critical Care, 2009, 13, P34	10.8	1
	Are chest X-rays necessary after chest tube insertion in trauma emergencies?. <i>Critical Care</i> , 2009 , 13, P35	10.8	1
	Bedside chest ultrasound reduces the rate of chest X-ray and CT examinations. <i>Critical Care</i> , 2009 , 13, P36	10.8	78
	Combination of variability with pressure support ventilation enhances lung protection and function in experimental acute lung injury. <i>Critical Care</i> , 2009 , 13, P37	10.8	78
	Measurement of the forearm to warrant low-tidal-volume ventilation in the acute respiratory distress syndrome. <i>Critical Care</i> , 2009 , 13, P38	10.8	78
	Effect of pleural effusion on gas exchange and response to positive end-expiratory pressure in acute lung injury/acute respiratory distress syndrome patients. <i>Critical Care</i> , 2009 , 13, P39	10.8	78

CT-scan lung morphology predicts the response to a recruitment maneuver in acute respiratory distress syndrome patients. <i>Critical Care</i> , 2009 , 13, P40	10.8	1
Alveolar wall disruption and lung inflammation associated with positive end-expiratory pressure and recruitment maneuver in pigs. <i>Critical Care</i> , 2009 , 13,	10.8	78
Ventilation with high positive end-expiratory pressure improves oxygenation after cardiac surgery independently of the mode of ventilation and of the use of nitric oxide. <i>Critical Care</i> , 2009 , 13, P42	10.8	1
Control system for automated titration of positive end-expiratory pressure and tidal volume using dynamic nonlinear compliance as the setpoint. <i>Critical Care</i> , 2009 , 13, P43	10.8	78
Effect of positive end-expiratory pressure on regional ventilation monitored by electrical impedance tomography in mechanically ventilated ICU patients. <i>Critical Care</i> , 2009 , 13, P44	10.8	78
Measurement of the end-expiratory lung volume without interruption of mechanical ventilation in pediatric patients. <i>Critical Care</i> , 2009 , 13, P45	10.8	78
Success of recruitment maneuvers during pneumoperitoneum is dependent on the intraabdominal pressure. <i>Critical Care</i> , 2009 , 13, P46	10.8	78
Correlation between lung sound distribution and functional residual capacity: preliminary findings. Critical Care, 2009 , 13, P47	10.8	1
Clinical utility of functional residual capacity measurement based on a modified nitrogen breath washout technique. <i>Critical Care</i> , 2009 , 13, P48	10.8	78
Pulmonary permeability index predicts progression to acute lung injury in patients with increased risk. <i>Critical Care</i> , 2009 , 13, P49	10.8	1
Pulmonary electrical impedance tomography changes in a model of hemorrhagic shock with endotoxemia and resuscitation. <i>Critical Care</i> , 2009 , 13,	10.8	1
1111 Determination of lung area in electrical impedance tomography images. <i>Critical Care</i> , 2009 , 13, P51	10.8	1
Development of a system for in vivo optical alveolar elastometry. <i>Critical Care</i> , 2009 , 13, P52	10.8	1
Effects of inhaled iloprost on acute respiratory distress syndrome in prone and supine positions. Critical Care, 2009 , 13, P53	10.8	78
Decreased vascular endothelial growth factor expression in lung tissue during acute respiratory distress syndrome. <i>Critical Care</i> , 2009 , 13, P54	10.8	78
The selective II nicotinic acetylcholine receptor agonist GTS-21 attenuates ventilator-induced inflammation and lung injury. <i>Critical Care</i> , 2009 , 13, P55	10.8	78
Usefulness of soluble E-selectin in the clinicopathologic assessment of acute lung injury/acute respiratory distress syndrome. <i>Critical Care</i> , 2009 , 13, P56	10.8	1
Prognostic value of epithelial neutrophil activating peptide 78 and monokine induced by IFNIIn bronchoalveolar lavage fluid in critically ill pediatric patients with acute respiratory distress syndrome. <i>Critical Care</i> , 2009 , 13, P57	10.8	78

1104	Cortisol-binding globulin cleavage at sites of inflammation in critically ill patients. <i>Critical Care</i> , 2009 , 13, P58	10.8	1
1103	Vascular connexins have differing responses to tumor necrosis factor. <i>Critical Care</i> , 2009 , 13, P59	10.8	78
1102	Prehospital intubation for out-of-hospital cardiac arrest. <i>Critical Care</i> , 2009 , 13, P60	10.8	78
1101	Focused echocardiography and capnography during resuscitation from pulseless electrical activity after out-of-hospital cardiac arrest. <i>Critical Care</i> , 2009 , 13, P61	10.8	2
1100	Patients with an automatic external defibrillator applied by a bystander in a public setting have a strikingly higher frequency of ventricular tachycardia/ventricular fibrillation than observed cardiac arrests in the home. <i>Critical Care</i> , 2009 , 13, P62	10.8	78
1099	Survival unchanged 5 months after implementing the 2005 American Heart Association cardiopulmonary resuscitation and emergency cardiac care guidelines for out-of-hospital cardiac arrest. <i>Critical Care</i> , 2009 , 13, P63	10.8	78
1098	Outcomes following admission to ICU post cardiac arrest. <i>Critical Care</i> , 2009 , 13, P64	10.8	1
1097	Clinical outcome in patients who experienced inhospital cardiac arrest by underlying disease. <i>Critical Care</i> , 2009 , 13, P65	10.8	78
1096	Good outcome in octogenarians after ventricular fibrillation out-of-hospital cardiac arrest. <i>Critical Care</i> , 2009 , 13, P66	10.8	78
1095	What is full recovery? Reconsidering outcome of cardiopulmonary cerebral resuscitation beyond Glasgow B ittsburgh Cerebral Performance Categories 1. <i>Critical Care</i> , 2009 , 13, P67	10.8	1
1094	Intracranial pressure monitoring during hypothermia after cardiopulmonary resuscitation. <i>Critical Care</i> , 2009 , 13, P68	10.8	78
1093	Serum neuron-specific enolase as early predictor of outcome after inhospital cardiac arrest. <i>Critical Care</i> , 2009 , 13, P69	10.8	78
1092	Controlled reperfusion prevents neurologic injury after global brain ischemia in a novel ischemic brain model. <i>Critical Care</i> , 2009 , 13, P70	10.8	78
1091	Effectiveness of an underbody forced warm-air blanket in preventing postoperative hypothermia after coronary artery bypass graft surgery with normothermic cardiopulmonary bypass. <i>Critical Care</i> , 2009 , 13, P71	10.8	78
1090	Good outcome in noncoronary out-of-hospital cardiac arrest treated with mild induced hypothermia. <i>Critical Care</i> , 2009 , 13, P72	10.8	78
1089	Nasopharyngeal cooling during resuscitation: randomized study. <i>Critical Care</i> , 2009 , 13, P73	10.8	78
1088	Introduction of an external noninvasive cooling device for effective implementation of Intensive Care Society standards post cardiac arrest. <i>Critical Care</i> , 2009 , 13, P74	10.8	78
1087	Therapeutic hypothermia for postcardiac arrest patients: physicians are warming up to the idea. <i>Critical Care</i> , 2009 , 13, P75	10.8	78

1086	Comparison between clinical tests and the Cerebral State Index or brain death determination. <i>Critical Care</i> , 2009 , 13, P76	10.8	78
1085	Effects of mechanical ventilation on Cushing's triad. Critical Care, 2009, 13, P77	10.8	2
1084	Losing potential organ donors in critical care units: data from the Donor Action Database. <i>Critical Care</i> , 2009 , 13, P78	10.8	78
1083	Critical care staff attitudes to organ donation impact on national donation rates: data from the Donor Action Database. <i>Critical Care</i> , 2009 , 13, P79	10.8	2
1082	Effect of endotracheal suctioning on intracranial pressure in severe head-injured patients. <i>Critical Care</i> , 2009 , 13, P80	10.8	1
1081	Elevated plasma ammonia concentration in patients with traumatic hemorrhage. <i>Critical Care</i> , 2009 , 13, P81	10.8	78
1080	Regulatory T cells are persistently enriched in the peripheral blood of head-injury patients. <i>Critical Care</i> , 2009 , 13, P82	10.8	78
1079	Alteration of cardiopulmonary function after severe head injury. Critical Care, 2009, 13, P83	10.8	78
1078	Abstract withdrawn. <i>Critical Care</i> , 2009 , 13, P84	10.8	78
1077	The RESCUEicp decompressive craniectomy trial. <i>Critical Care</i> , 2009 , 13, P85	10.8	1
1076	Drift analysis of a novel device for measurement of regional cerebral blood flow. <i>Critical Care</i> , 2009 , 13, P86	10.8	78
1075	Continuous monitoring of carbon dioxide reactivity in traumatic brain injury. <i>Critical Care</i> , 2009 , 13, P87	10.8	78
1074	Determining neurological prognosis in patients with severe traumatic brain injury: a survey of Canadian intensivists. <i>Critical Care</i> , 2009 , 13, P88	10.8	78
1073	Severe brain trauma management analysis using a high-rate recording tool: better definition allows better analysis of practice. <i>Critical Care</i> , 2009 , 13, P89	10.8	78
1072	Comparison of a new brain tissue oxygenation measuring probe with the established standard. <i>Critical Care</i> , 2009 , 13, P90	10.8	78
1071	The brain is a major source of S100 increase in porcine endotoxemic shock. <i>Critical Care</i> , 2009 , 13, P91	10.8	1
1070	Anemia is associated with brain tissue hypoxia and metabolic crisis after severe brain injury. <i>Critical Care</i> , 2009 , 13, P92	10.8	3
1069	Nonconvulsive seizures and renal failure after intracerebral hemorrhage. <i>Critical Care</i> , 2009 , 13, P93	10.8	78

1068	S100 beta is a marker of brain insult caused by systemic inflammatory response syndrome. <i>Critical Care</i> , 2009 , 13, P94	10.8	78
1067	Cerebral oxygenation monitoring in critical care patients with traumatic brain injury. <i>Critical Care</i> , 2009 , 13, P95	10.8	78
1066	Addition of new criteria to the Sequential Organ Failure Assessment for the patients with subarachnoid hemorrhage. <i>Critical Care</i> , 2009 , 13, P96	10.8	1
1065	Epidemiological analysis of patients with cerebral aneurysm submitted for an embolization at SB Jos[] do Ava[Hospital. <i>Critical Care</i> , 2009 , 13, P97	10.8	78
1064	Seizures and organ dysfunction after subarachnoid hemorrhage. Critical Care, 2009, 13, P98	10.8	78
1063	Alteration of cardiopulmonary function after subarachnoid hemorrhage. Critical Care, 2009, 13, P99	10.8	78
1062	Transfusion increases infection without affecting neurologic outcome in spontaneous subarachnoid hemorrhage. <i>Critical Care</i> , 2009 , 13, P100	10.8	1
1061	Emergency surgical management for prevention of symptomatic vasospasm and normal pressure hydrocephalus after subarachnoid hemorrhage due to ruptured cerebral aneurysm. <i>Critical Care</i> , 2009 , 13, P101	10.8	78
1060	Diagnosis and treatment of subarachnoid hemorrhage-induced vasospasm. <i>Critical Care</i> , 2009 , 13, P102	10.8	78
1059	Effects of simvastatin in prevention of vasospasm in nontraumatic subarachnoid hemorrhage: preliminary data. <i>Critical Care</i> , 2009 , 13, P103	10.8	13
1058	Magnesium use on prophylaxis of vasospasm morbidity and the mortality rate in subarachnoid hemorrhage. <i>Critical Care</i> , 2009 , 13, P104	10.8	78
1057	Outcomes of ventilator-associated pneumonia in aneurysmal subarachnoid hemorrhage patients. <i>Critical Care</i> , 2009 , 13, P105	10.8	78
1056	Outcomes from subarachnoid haemorrhage. <i>Critical Care</i> , 2009 , 13, P106	10.8	78
1055	Full Outline of Unresponsiveness compared with Glasgow coma scale assessment and outcome prediction in coma. <i>Critical Care</i> , 2009 , 13, P107	10.8	2
1054	Intravenous immunoglobulins versus plasma exchange in the treatment of Guillain-Barr syndrome. <i>Critical Care</i> , 2009 , 13, P108	10.8	78
1053	Determinants of critical illness polyneuropathy in the case of long-term ICU treatment. <i>Critical Care</i> , 2009 , 13, P109	10.8	78
1052	Neurally adjusted ventilatory assistance in patients with critical illness polyneuromyopathy. <i>Critical Care</i> , 2009 , 13, P110	10.8	78
1051	Assessment of muscle membrane properties using muscle velocity recovery cycles in patients with critical illness polyneuromyopathy. <i>Critical Care</i> , 2009 , 13, P111	10.8	78

1050	Risk factors for developing hypoglycemia in neurocritical care patients. <i>Critical Care</i> , 2009 , 13, P112	10.8	78
1049	Relationship between effective osmolality changes and neurological status during treatment for severe paediatric diabetic ketoacidosis. <i>Critical Care</i> , 2009 , 13, P113	10.8	78
1048	Continuous monitoring of blood parameters in intensive care patients. <i>Critical Care</i> , 2009 , 13, P114	10.8	О
1047	Morning glucose correlates poorly with daily mean glucose in cardiac surgical ICU patients: impact of ultradian variation of glucose. <i>Critical Care</i> , 2009 , 13, P115	10.8	78
1046	Hypoglycaemia is associated with a higher mortality in critically ill patients. <i>Critical Care</i> , 2009 , 13, P116	10.8	78
1045	Insulin pseudo-resistance from adsorption to burettes. <i>Critical Care</i> , 2009 , 13, P117	10.8	78
1044	High blood glucose variability in acute phase is one of the most important risk factors relating to the outcome in acutely ill severe patients with glucose intolerance. <i>Critical Care</i> , 2009 , 13, P118	10.8	1
1043	Effect of increasing intravenous glucose load in the presence of normoglycemia on outcome and metabolism in critically ill rabbits. <i>Critical Care</i> , 2009 , 13, P119	10.8	78
1042	Reduction of glucose and insulin concentrations during in vitro incubation of whole blood. <i>Critical Care</i> , 2009 , 13, P120	10.8	78
1041	Surprising result after evaluating a nurse-driven guideline on blood glucose management. <i>Critical Care</i> , 2009 , 13, P121	10.8	78
1040	Evaluation of nursing perceptions about three insulin protocols for blood glucose control in critical care. <i>Critical Care</i> , 2009 , 13, P122	10.8	1
1039	Impact of a simple computer alert on the quality of tight glycemic control. <i>Critical Care</i> , 2009 , 13, P123	10.8	78
1038	Which variables affect strict glycaemic control with intensive insulin therapy in postoperative/post-traumatic critically ill patients?. <i>Critical Care</i> , 2009 , 13, P124	10.8	78
1037	Comparison of conventional measures of glucose control versus the area under the curve from a continuous glucose monitoring device in critical care patients. <i>Critical Care</i> , 2009 , 13, P125	10.8	78
1036	Exogenous glucagon-like peptide 1: a potentially novel therapy for the management of hyperglycaemia in the critically ill. <i>Critical Care</i> , 2009 , 13, P126	10.8	78
1035	Tight glucose control: is there any influence on outcome? A retrospective cohort study. <i>Critical Care</i> , 2009 , 13, P127	10.8	78
1034	Should we treat children with hyperglycaemia with insulin after cardiac surgery?. <i>Critical Care</i> , 2009 , 13, P128	10.8	78
1033	Effect of glucose?insulin?potassium infusion on mortality in critically ill patients: a systematic review and meta-analysis. <i>Critical Care</i> , 2009 , 13, P129	10.8	78

1032	Incidence and risk factors of hypertriglyceridemia in the ICU. Critical Care, 2009, 13, P130	10.8	78
1031	Abstract withdrawn. <i>Critical Care</i> , 2009 , 13, P131	10.8	78
1030	Should perioperative immune-modulating nutrition therapy be the standard of care? A systematic review. <i>Critical Care</i> , 2009 , 13, P132	10.8	78
1029	Arginine metabolism in a small animal model of sepsis and after hemihepatectomy. <i>Critical Care</i> , 2009 , 13, P133	10.8	78
1028	Ginger extract dietary supplementation effects on delayed gastric emptying and ventilator-associated pneumonia in adult respiratory distress syndrome patients. <i>Critical Care</i> , 2009 , 13, P134	10.8	78
1027	Immunostimulated enterocytes activate extracellular arginase I which competes with enterocyte inducible nitric oxide synthase for arginine during inflammation. <i>Critical Care</i> , 2009 , 13, P135	10.8	78
1026	Role of parenteral glutamine supplementation on patient outcome in the surgical ICU. <i>Critical Care</i> , 2009 , 13, P136	10.8	78
1025	Plasma citrulline kinetics and prognostic value in the critically ill patient. <i>Critical Care</i> , 2009 , 13, P137	10.8	1
1024	Modulation of lipid utilisation by parenteral administration of a fish-oil-enriched new lipid formula (SMOFlipid[]) in surgical ICU patients: comparison with a lipid emulsion based on olive and soybean oil. <i>Critical Care</i> , 2009 , 13, P138	10.8	1
1023	Gastrointestinal function in critically ill trauma patients using motility capsule technology. <i>Critical Care</i> , 2009 , 13, P139	10.8	78
1022	Is intragastric administration of enteral nutrition safe in acute severe pancreatitis?. <i>Critical Care</i> , 2009 , 13, P140	10.8	78
1021	Critical illness gastrointestinal hypomobility disorder and success of enteral erythromycin. <i>Critical Care</i> , 2009 , 13, P141	10.8	78
1020	A randomised prospective trial to compare the efficacy of bolus versus continuous nasogastric feeding in paediatric intensive care. <i>Critical Care</i> , 2009 , 13, P142	10.8	О
1019	Nutrition therapy in the critical care setting: what is best achievable practice? An international quality improvement project. <i>Critical Care</i> , 2009 , 13, P143	10.8	3
1018	Feeding enterally the hemodynamically unstable critically ill patient: experience from a multicentre trial (the REDOXS' study). <i>Critical Care</i> , 2009 , 13, P144	10.8	1
1017	Molecular adsorbent recirculating system: a clinical experience in acute or acute on chronic liver failure (133 sessions). <i>Critical Care</i> , 2009 , 13, P145	10.8	78
1016	Comparison of two indwelling bowel catheters on economic impact by number of bedding and dressing changes per day. <i>Critical Care</i> , 2009 , 13, P146	10.8	78
1015	Alterations of the immune system in acute pancreatitis and systematic inflammatory response syndrome. <i>Critical Care</i> , 2009 , 13, P147	10.8	78

Hungarian perioperative selenium survey in patients with oesophageal cancer. <i>Critical Care</i> , 2009 , 13, P148	10.8	1
1013 Selenium in critically ill children with cardiac dysfunction. <i>Critical Care</i> , 2009 , 13, P149	10.8	78
Effect of high-dose selenium substitution on selected laboratory parameters and prognosis in critically ill patients. <i>Critical Care</i> , 2009 , 13, P150	10.8	1
1011 Dynamic generation of physiological model systems. <i>Critical Care</i> , 2009 , 13, P151	10.8	78
1010 Intraaortic balloon pumping: why should we hurry up?. Critical Care, 2009, 13, P152	10.8	78
1009 Study of risk factors for female patients with acute myocardial infarction. <i>Critical Care</i> , 2009 , 13, P1	53 10.8	78
1008 Influence of arterial pressure on tissue perfusion in septic shock. <i>Critical Care</i> , 2009 , 13, P154	10.8	78
Echocardiographic assessment of the effects of acute left ventricular pacing on patients with severe congestive heart failure and narrow QRS duration. <i>Critical Care</i> , 2009 , 13, P155	10.8	78
1006 Left ventricular pacing alone improves haemodynamic variables. <i>Critical Care</i> , 2009 , 13, P156	10.8	78
Preoperative left atrial dysfunction and new-onset atrial fibrillation in cardiac surgery patients. **Critical Care**, 2009 , 13, P157	10.8	78
1004 Right ventricular involvement in Takotsubo cardiomyopathy. <i>Critical Care</i> , 2009 , 13, P158	10.8	78
Possible cause of bradycardia developed due to 2-sympothomimetic infusion. <i>Critical Care</i> , 2009 , 13, P159	10.8	78
Automatic real-time detection of myocardial ischemia by epicardial accelerometer. <i>Critical Care</i> , 2009 , 13, P160	10.8	78
Prognosis of acute myocardial infarction outcomes using evaluation of cardiac power (product of cardiac output and mean arterial pressure). <i>Critical Care</i> , 2009 , 13, P161	10.8	78
Left ventricular torsion analysis using echocardiographic speckle tracking in a canine model of dyssynchrony and cardiac resynchronization therapy predicts global cardiac performance. <i>Critical Care</i> , 2009 , 13, P162	10.8	78
999 Cardiac diseases during pregnancy and periperium: 13 years in the ICU. <i>Critical Care</i> , 2009 , 13, P163	10.8	78
Endothelial nitric oxide synthase deficiency and inducible nitric oxide synthase inhibition in the setting of septic cardiomyopathy. <i>Critical Care</i> , 2009 , 13, P164	10.8	78
997 Use of levosimendan in myocardial dysfunction due to sepsis. <i>Critical Care</i> , 2009 , 13, P165	10.8	8

996	Haemodynamic effects of levosimendan following cardiac surgery. <i>Critical Care</i> , 2009 , 13, P166	10.8	78
995	Effects of levosimendan started 24 hours before cardiopulmonary bypass in patients admitted to the ICU with an ejection fraction lower than 30%. <i>Critical Care</i> , 2009 , 13, P167	10.8	78
994	Utilization of levosimendan in the cardiac ICU: case series. <i>Critical Care</i> , 2009 , 13, P168	10.8	78
993	Effects of levosimendan and inhaled nitric oxide on microcirculation in septic shock. <i>Critical Care</i> , 2009 , 13, P169	10.8	2
992	Effects of levosimendan on renal function in septic shock: a caseBontrol study. <i>Critical Care</i> , 2009 , 13, P170	10.8	3
991	Levosimendan versus dobutamine in septic shock. <i>Critical Care</i> , 2009 , 13, P171	10.8	2
990	Levosimendan therapy does not improve survival of post-resuscitation cardiogenic shock patients. <i>Critical Care</i> , 2009 , 13, P172	10.8	2
989	Study of hemodynamics in patients treated with landiolol in the ICU. <i>Critical Care</i> , 2009 , 13, P173	10.8	78
988	Effects of intravenous diltiazem on a porcine model of endotoxin-induced pulmonary hypertension. <i>Critical Care</i> , 2009 , 13, P174	10.8	78
987	Acute liver failure induced by intravenous amiodarone in the cardiac care unit: retrospective study of 3 years. <i>Critical Care</i> , 2009 , 13, P175	10.8	78
986	Prevention of postoperative arrhythmias after pulmonary resection: celiprolol versus magnesium. <i>Critical Care</i> , 2009 , 13, P176	10.8	78
985	Low-dose dopamine is not useful in kidney transplantation. <i>Critical Care</i> , 2009 , 13, P177	10.8	78
984	Low-dose nitroglycerin improves microcirculation in hospitalized patients with acute heart failure. <i>Critical Care</i> , 2009 , 13, P178	10.8	78
983	Effects of intravenous nitroglycerin and noradrenaline on gastric microvascular perfusion in an experimental model of gastric tube reconstruction. <i>Critical Care</i> , 2009 , 13, P179	10.8	1
982	Norepinephrine: more than blood pressure cosmetics?. <i>Critical Care</i> , 2009 , 13, P180	10.8	1
981	Vasopressin in pediatric vasodilatory shock: a multicentred randomized controlled trial. <i>Critical Care</i> , 2009 , 13, P181	10.8	78
980	Arginine vasopressin increases plasma levels of von Willebrand factor in sheep. <i>Critical Care</i> , 2009 , 13, P182	10.8	4
979	Selective V1-agonism and selective V2-antagonism are superior to arginine vasopressin in ovine septic shock. <i>Critical Care</i> , 2009 , 13, P183	10.8	78

978	Adjunct terlipressin effect on vital organ perfusion during advanced life support in a porcine model of ventricular fibrillation. <i>Critical Care</i> , 2009 , 13, P184	10.8	78
977	Vasopressin versus norepinephrine infusion in patients with vasoplegic shock after cardiac surgery. <i>Critical Care</i> , 2009 , 13, P185	10.8	78
976	Arterial catheter-related infection according to the catheter site. Critical Care, 2009, 13, P186	10.8	78
975	Inline filtration reduces the incidence of systemic inflammatory response syndrome in critically ill children. <i>Critical Care</i> , 2009 , 13, P187	10.8	78
974	Central venous catheter infection: incidence and risk factors. Critical Care, 2009, 13, P188	10.8	78
973	Impact of bloodstream infections on ICU mortality. <i>Critical Care</i> , 2009 , 13, P189	10.8	78
972	Reduction of the catheter-related bloodstream infections in critically ill patients. <i>Critical Care</i> , 2009 , 13, P190	10.8	78
971	Definition of catheter-related bloodstream infection as a quality improvement measure in intensive care. <i>Critical Care</i> , 2009 , 13, P191	10.8	3
970	Comparative study between conventional and antiseptic impregnated central venous catheters. <i>Critical Care</i> , 2009 , 13, P192	10.8	78
969	Impact of early catheter removal during treatment of invasive candidiasis: analysis from two phase 3 micafungin trials. <i>Critical Care</i> , 2009 , 13, P193	10.8	78
968	Cholorhexidine, octenidine or povidone iodine for catheter-related infections: a randomised controlled trial. <i>Critical Care</i> , 2009 , 13, P194	10.8	1
967	Review of central-line-related sepsis in neurointensive care patients. <i>Critical Care</i> , 2009 , 13, P195	10.8	78
966	Central venous catheterization: a randomized comparison between external and internal jugular access. <i>Critical Care</i> , 2009 , 13, P196	10.8	1
965	Ultrasound-guided positioning of totally implantable access port systems: a single-center experience. <i>Critical Care</i> , 2009 , 13, P197	10.8	78
964	Ultrasound-guided catheterization of the subclavian vein: a prospective comparison with the landmark technique in ICU patients. <i>Critical Care</i> , 2009 , 13, P198	10.8	5
963	Perioperative fluid administration in pancreatic surgery: comparison of three regimens. <i>Critical Care</i> , 2009 , 13, P199	10.8	2
962	B-type natriuretic peptide, corrected flow time and central venous pressure as predictors of fluid responsiveness in septic shock. <i>Critical Care</i> , 2009 , 13, P200	10.8	78
961	Intraoperative fluid optimization using stroke volume variation in high-risk surgical patients: preliminary results of a randomized prospective single-center study. <i>Critical Care</i> , 2009 , 13, P201	10.8	78

960	Impedance cardiography in the estimation of hemodynamic and fluid status of coma patients during continuous venovenous hemodiafiltration. <i>Critical Care</i> , 2009 , 13, P202	10.8	1
959	Pulse and systolic pressure variation assessment in partially assisted ventilatory support. <i>Critical Care</i> , 2009 , 13, P203	10.8	78
958	Does the pleth variability index improve fluid management during major abdominal surgery?. <i>Critical Care</i> , 2009 , 13, P204	10.8	3
957	Plethysmography variability index: a new fluid responsiveness parameter. <i>Critical Care</i> , 2009 , 13, P205	10.8	2
956	Calibration of pulse contour continuous cardiac output analysis. <i>Critical Care</i> , 2009 , 13, P206	10.8	78
955	Is the pulse pressure variation a good predictor of fluid responsiveness in mechanically ventilated patients with low tidal volume?. <i>Critical Care</i> , 2009 , 13, P207	10.8	78
954	Proving the effectiveness of three dynamic indices to predict fluid responsiveness in septic mechanically ventilated patients. <i>Critical Care</i> , 2009 , 13, P208	10.8	78
953	Cross-comparison of the trending accuracy of continuous cardiac output measurement devices in postoperation patients. <i>Critical Care</i> , 2009 , 13, P209	10.8	78
952	Comparison of three noninvasive cardiac output monitors in patients undergoing therapeutic hypothermia after cardiac arrest. <i>Critical Care</i> , 2009 , 13, P210	10.8	2
951	Head-up tilt and passive leg raising in healthy volunteers as a preclinical model for preload-induced stroke volume modification. <i>Critical Care</i> , 2009 , 13, P211	10.8	Ο
950	Global end-diastolic volume as a predictor of the need for massive transfusion in multiple-trauma patients with hemorrhagic shock. <i>Critical Care</i> , 2009 , 13, P212	10.8	78
949	Should pulse pressure variation be indexed to tidal volume?. <i>Critical Care</i> , 2009 , 13, P213	10.8	78
948	Fluid responsiveness in patients following major surgery. <i>Critical Care</i> , 2009 , 13, P214	10.8	78
947	Alternative echographic assessment of inferior vena cava diameter variation in mechanically ventilated patients. <i>Critical Care</i> , 2009 , 13, P215	10.8	1
946	Global end-diastolic volume and global end-diastolic volume index are age dependent in awake, spontaneous-breathing patients after elective craniotomy. <i>Critical Care</i> , 2009 , 13, P216	10.8	78
945	Recalibration of pulse contour cardiac output using the PiCCO-2 device: when to perform the next thermodilution?. <i>Critical Care</i> , 2009 , 13, P217	10.8	1
944	Transpulmonary lithium dilution technique: time to recalibration and calibration drift. <i>Critical Care</i> , 2009 , 13, P218	10.8	78
943	Individualized intraoperative patient optimization using uncalibrated arterial pressure waveform analysis in high-risk patients undergoing major abdominal surgery. <i>Critical Care</i> , 2009 , 13, P219	10.8	1

942	Evaluation of the effectiveness of different volume replacement therapies in postoperative hypovolemic patients using the PiCCO monitoring system. <i>Critical Care</i> , 2009 , 13, P220	10.8	78
941	Coupling of cardiac index and global end-diastolic volume index: is it mathematical or something else?. <i>Critical Care</i> , 2009 , 13, P221	10.8	78
940	Comparison of transpulmonary thermodilution measurements of global end-diastolic volume index, extravascular lung water index and central venous pressure with radiographic estimation of these parameters using computed tomography. <i>Critical Care</i> , 2009 , 13, P222	10.8	78
939	Correlation of clinical evaluation and invasive monitoring evaluation in critically ill patients. <i>Critical Care</i> , 2009 , 13, P223	10.8	78
938	Influence of prone positioning on measurement of extravascular lung water and pulmonary vascular permeability indexes by transpulmonary thermodilution. <i>Critical Care</i> , 2009 , 13, P224	10.8	78
937	Cardiac output by thermodilution and the MostCare system in patients on intra-aortic balloon pump. <i>Critical Care</i> , 2009 , 13, P225	10.8	1
936	Validation of the extravascular lung water by single transpulmonary thermodilution in the clinical setting. <i>Critical Care</i> , 2009 , 13, P226	10.8	78
935	Relation between mixed venous and central venous saturation in sepsis: influence of source of sepsis. <i>Critical Care</i> , 2009 , 13, P227	10.8	78
934	Intraoperative central venous oxygen saturation and patient outcome in patients undergoing major abdominal surgery. <i>Critical Care</i> , 2009 , 13, P228	10.8	78
933	Bioelectrical impedance analysis in ICU patients. <i>Critical Care</i> , 2009 , 13, P229	10.8	78
932	Estimated respiratory quotient and venoarterial pCO2 difference are outcome markers in patients with left ventricular dysfunction submitted to coronary artery bypass surgery. <i>Critical Care</i> , 2009 , 13, P230	10.8	78
931	Clinical value of noninvasive pulmonary artery systolic pressure estimates in our patients with pulmonary artery hypertension. <i>Critical Care</i> , 2009 , 13, P231	10.8	78
930	Goal-directed fluid therapy based on the continuous left ventricle end-diastolic volume improves acute mesenteric ischemia/reperfusion injury in rats. <i>Critical Care</i> , 2009 , 13, P232	10.8	78
929	Utility of brain natriuretic peptide as a marker of early cardiac risk in patients undergoing elective abdominal aortic aneurism repair. <i>Critical Care</i> , 2009 , 13, P233	10.8	78
928	Accuracy of central venous oxygen saturation with a fiberoptic catheter. <i>Critical Care</i> , 2009 , 13, P234	10.8	78
927	Venous oxygen saturation and lactate gradient from the superior vena cava to the pulmonary artery in ICU patients with septic shock. <i>Critical Care</i> , 2009 , 13, P235	10.8	1
926	Use of tissue oxygenation saturation in association with skin temperature as an indicator of the peripheral tissue perfusion in critically ill patients. <i>Critical Care</i> , 2009 , 13, P236	10.8	78
925	Effects of peripheral vasodilation induced by regional anaesthesia blocks on resting tissue oxygenation values. <i>Critical Care</i> , 2009 , 13, P237	10.8	0

924	Effect of anemia on tissue oxygenation saturation and the tissue deoxygenation rate during ischemia. <i>Critical Care</i> , 2009 , 13, P238	10.8	6
923	Near-infrared spectroscopy monitoring tissue oxygen saturation after cardiac surgery. <i>Critical Care</i> , 2009 , 13, P239	10.8	78
922	Comparison of muscle tissue oxygenation response curves to two time-based vascular occlusion tests: evidence of diminishing returns?. <i>Critical Care</i> , 2009 , 13, P240	10.8	78
921	Adaptability of muscle tissue oxygenation to repeated vascular occlusion. <i>Critical Care</i> , 2009 , 13, P241	10.8	78
920	Microcirculatory, leucocyte/endothelium interaction and survival time effects of dobutamine in nonhypotensive endotoxemia. <i>Critical Care</i> , 2009 , 13, P242	10.8	78
919	Microcirculatory, leukocyte/endothelium interaction and survival time effects of recombinant C-reactive protein in nonhypotensive endotoxemia in hamsters. <i>Critical Care</i> , 2009 , 13, P243	10.8	78
918	Assessment of tissue oxygen saturation during a vascular occlusion test using near-infrared spectroscopy: role of the probe spacing and measurement site studied in healthy volunteers. <i>Critical Care</i> , 2009 , 13, P244	10.8	0
917	Effects of perfusion pressure on microcirculatory perfusion of gastric tubes in pigs. <i>Critical Care</i> , 2009 , 13, P245	10.8	78
916	Lack of autoregulatory blood flow escape in the skin after infusion of therapeutic levels of noradrenaline through a microdialysis system in healthy volunteers. <i>Critical Care</i> , 2009 , 13, P246	10.8	78
915	Alterations of tissue-dependent microcirculation in patients after successful resuscitation. <i>Critical Care</i> , 2009 , 13, P247	10.8	78
914	Microcirculatory changes caused by magnesium sulphate infusion in severe sepsis and septic shock. <i>Critical Care</i> , 2009 , 13, P248	10.8	78
913	Sepsis induces an early impairment of endothelial glycocalyx in glomerular capillaries. <i>Critical Care</i> , 2009 , 13, P249	10.8	78
912	Electron microscopic renal tubular injury score in ovine endotoxemia. <i>Critical Care</i> , 2009 , 13, P250	10.8	78
911	Inflammation-induced renal injury subsides when endotoxin tolerance develops in humans as measured by urine proteomics. <i>Critical Care</i> , 2009 , 13, P251	10.8	78
910	Urinary glutathione S-transferase as an early marker for acute kidney injury in patients admitted to intensive care with sepsis. <i>Critical Care</i> , 2009 , 13, P252	10.8	78
909	Plasma neutrophil gelatinase-associated lipocalin is an early marker of acute kidney injury in critically ill patients: a prospective study. <i>Critical Care</i> , 2009 , 13, P253	10.8	1
908	Plasma cholinesterase activity in patients during peritoneal dialysis. <i>Critical Care</i> , 2009 , 13, P254	10.8	
907	Early predictive value of neutrophil gelatinase-associated lipocalin in adult ICU patients with acute kidney injury. <i>Critical Care</i> , 2009 , 13, P255	10.8	78

906	Novel time series analysis approach for prediction of dialysis in critically ill patients using echo-state networks. <i>Critical Care</i> , 2009 , 13, P256	10.8	78
905	Contrast-induced nephropathy: a prospective analysis of long-term outcome and persistence of renal impairment. <i>Critical Care</i> , 2009 , 13, P257	10.8	78
904	Outcome of ICU patients requiring dialysis in an African institution. Critical Care, 2009, 13, P258	10.8	78
903	Early goal-directed therapy of septic patients coming from the ward does not protect the kidneys. <i>Critical Care</i> , 2009 , 13, P259	10.8	78
902	Is goal-directed therapy useful in kidney transplantation?. Critical Care, 2009, 13, P260	10.8	78
901	Clinical characteristics and outcomes of critically ill adults with septic acute kidney injury in a general hospital in Singapore. <i>Critical Care</i> , 2009 , 13, P261	10.8	78
900	Incidence and outcome of acute renal failure necessitating renal replacement therapy after trauma. <i>Critical Care</i> , 2009 , 13, P262	10.8	78
899	Early acute kidney injury in Northern Ireland ICUs. <i>Critical Care</i> , 2009 , 13, P263	10.8	О
898	RIFLE classification can predict hospital mortality of critically ill patients. Critical Care, 2009, 13, P264	10.8	78
897	Assessment of acute kidney injury with modified RIFLE criteria in critically ill pediatric burn patients. <i>Critical Care</i> , 2009 , 13, P265	10.8	1
896	Extracorporeal renal replacement therapy in patients with heparin-induced thrombocytopenia. <i>Critical Care</i> , 2009 , 13, P266	10.8	78
895	Coagulopathy of liver disease does not increase the filter life during continuous renal replacement therapy. <i>Critical Care</i> , 2009 , 13, P267	10.8	78
894	Optimal dose of renal replacement therapy in acute kidney injury: a meta-analysis. <i>Critical Care</i> , 2009 , 13, P268	10.8	78
893	Removal of inflammatory mediators by continuous renal replacement therapy in severe sepsis. <i>Critical Care</i> , 2009 , 13, P269	10.8	1
892	Blood epuration of middle molecules in continuous venovenous hemodiafiltration with regional citrate anticoagulation versus systemic heparinization. <i>Critical Care</i> , 2009 , 13, P270	10.8	78
891	Utilization of slow low-efficiency dialysis may help to optimize the need for continuous renal replacement therapy in Indian ICUs. <i>Critical Care</i> , 2009 , 13, P271	10.8	1
890	Observational cohort of renal replacement therapy patients at a district general hospital ICU: case mix and outcomes. <i>Critical Care</i> , 2009 , 13, P272	10.8	1
889	Effects of continuous hemofiltration on organ perfusion, energy metabolism, oxidative stress, endothelial dysfunction and inflammation. <i>Critical Care</i> , 2009 , 13, P273	10.8	78

888	Influence of continuous venovenous hemofiltration on transpulmonary thermodilution-derived parameters. <i>Critical Care</i> , 2009 , 13, P274	10.8	1
887	Antibiotic dosing regimens for septic patients receiving continuous venovenous haemofiltration: do current studies supply sufficient data?. <i>Critical Care</i> , 2009 , 13, P275	10.8	2
886	Comparing adsorption of gentamicin by polyacrylonitrile and polyamide hemofiltration filter in an in vitro continuous venovenous hemofiltration model. <i>Critical Care</i> , 2009 , 13, P276	10.8	78
885	Slow continuous ultrafiltration: how many fluids must be taken out?. <i>Critical Care</i> , 2009 , 13, P277	10.8	78
884	Plasmatic cytokines and intermittent hemodialysis with polymethylmethacrylate membrane in septic shock patients. <i>Critical Care</i> , 2009 , 13, P278	10.8	78
883	Immunoadsorption in dilated cardiomyopathy. <i>Critical Care</i> , 2009 , 13, P279	10.8	78
882	Lipopolysaccharide adsorber in abdominal septic shock. <i>Critical Care</i> , 2009 , 13, P280	10.8	2
881	Experience with the use of selective sorbents in complex intensive care of sepsis in patients after cardiac surgery. <i>Critical Care</i> , 2009 , 13, P281	10.8	2
880	Best choice of acute blood purification therapy based on the severity score and blood lactic acid values in septic shock patients. <i>Critical Care</i> , 2009 , 13, P282	10.8	78
879	Clinical effects of polymyxin B immobilised fiber with direct hemoperfusion for the patients of severe sepsis or septic shock caused by intra-abdominal infections. <i>Critical Care</i> , 2009 , 13, P283	10.8	78
878	Improvement of haemodynamic and respiratory parameters during coupled plasma filtration and adsorption correlates with the clearance of inflammatory mediators. <i>Critical Care</i> , 2009 , 13, P284	10.8	2
877	Improving of APACHE II score at the early phase using CTR-001 direct hemoperfusion in patients with severe sepsis and septic shock. <i>Critical Care</i> , 2009 , 13, P285	10.8	O
876	Beneficial effects of early hemoperfusion with a polymyxin B fibre column on septic shock. <i>Critical Care</i> , 2009 , 13, P286	10.8	78
875	Observational study for direct hemoperfusion therapy with polymyxin-B immobilized fibers (PMX-DHP) in patients with septic shock in Japan: PMX-DHP study group. <i>Critical Care</i> , 2009 , 13, P287	10.8	78
874	Investigation of type II phospholipase A2 values and eicosanoid values during polymyxin-B-immobilized fiber direct hemoperfusion of septic shock patients. <i>Critical Care</i> , 2009 , 13, P288	10.8	78
873	Lipid mediator adosorption with dialyser membrane in patients with septic shock. <i>Critical Care</i> , 2009 , 13, P289	10.8	78
872	Microbiologic contamination of ultrasound transducers utilized by anesthesiologists in the operating room and ICU. <i>Critical Care</i> , 2009 , 13, P290	10.8	2
871	Assessment of the pathogen microorganisms and resistance patterns in patient with device-associated infections in the ICU. <i>Critical Care</i> , 2009 , 13, P291	10.8	78

870	Does improved oral hygiene alone prevent ventilator-associated pneumonia?. <i>Critical Care</i> , 2009 , 13, P292	10.8	1
869	Effects of body orientation on the development of ventilator-associated pneumonia in mechanically ventilated swine. <i>Critical Care</i> , 2009 , 13, P293	10.8	78
868	Gastroesophageal reflux in mechanically ventilated pediatric patients and its relation to ventilator acquired pneumonia. <i>Critical Care</i> , 2009 , 13, P294	10.8	78
867	Incidence of ventilator-associated pneumonia in patients undergoing elective tube exchange to LoTrach endotracheal tubes. <i>Critical Care</i> , 2009 , 13, P295	10.8	2
866	Reduction in ventilator-associated pneumonia following the introduction of subglottic suction endotracheal tubes. <i>Critical Care</i> , 2009 , 13, P296	10.8	78
865	Ventilator-associated pneumonia in a Greek ICU: prevalence and etiology. <i>Critical Care</i> , 2009 , 13, P297	10.8	78
864	Ventilator-associated pneumonia after procedures in cardiac surgery. Critical Care, 2009, 13, P298	10.8	78
863	Epidemiology of ventilator-associated pneumonia in ICU patients. <i>Critical Care</i> , 2009 , 13, P299	10.8	1
862	Clara cell protein in bronchoalveolar lavage fluid: a predictor of ventilator-associated pneumonia?. <i>Critical Care</i> , 2009 , 13, P300	10.8	78
861	Presence of human metapneumovirus in bronchoalveolar lavage fluid of critically ill patients. <i>Critical Care</i> , 2009 , 13, P301	10.8	78
860	Trend analysis of antibiotic resistance and minimum inhibitory concentration distribution from 1997 to 2007 among Pseudomonas aeruginosa, Escherichia coli and Klebsiella pneumoniae in European ICUs. <i>Critical Care</i> , 2009 , 13, P302	10.8	78
859	Antimicrobial resistance pattern of Pseudomonas aeruginosa in clinical isolates from ICU patients. <i>Critical Care</i> , 2009 , 13, P303	10.8	78
858	Bacteriological profile and antibiotic resistance of bacteria isolates in a burn department. <i>Critical Care</i> , 2009 , 13, P304	10.8	78
857	Antibiotic resistance of Staphylococcus aureus from ICUs in the Netherlands 1996 to 2006. <i>Critical Care</i> , 2009 , 13, P305	10.8	78
856	Clinical and microbiological efficacy of continuous versus intermittent administration of vancomycin in critical care patients. <i>Critical Care</i> , 2009 , 13, P306	10.8	78
855	Over-increased creatinine renal clearance in septic patients and implications for vancomycin optimization. <i>Critical Care</i> , 2009 , 13, P307	10.8	78
854	Vancomycin-resistant enterococci colonisation in the ICU and control measures. <i>Critical Care</i> , 2009 , 13, P308	10.8	78
853	Activities of vancomycin, teicoplanin and linezolid against bacteraemic methicillin-resistant Staphylococcus aureus strains in Gauteng, South Africa. <i>Critical Care</i> , 2009 , 13, P309	10.8	78

852	Telavancin for the treatment of hospital-acquired pneumonia in severely ill and older patients: the ATTAIN studies. <i>Critical Care</i> , 2009 , 13, P310	10.8	2
851	Piperacillin/tazobactam administered by continuous or intermittent infusion for the treatment of nosocomial pneumonia. <i>Critical Care</i> , 2009 , 13, P311	10.8	78
850	Efficiency of rifampicinthiconazole-impregnated catheters in the femoral venous site. <i>Critical Care</i> , 2009 , 13, P312	10.8	78
849	Micafungin for the treatment of pediatric invasive fungal infections. <i>Critical Care</i> , 2009 , 13, P313	10.8	78
848	Malignant Boutonneuse fever with multiple organ failure: a three-case series. <i>Critical Care</i> , 2009 , 13, P314	10.8	78
847	Impact of microbiology and antimicrobial treatment on mortality in septic shock. <i>Critical Care</i> , 2009 , 13, P315	10.8	78
846	Do changes in ICU flora and their antibiotic susceptibility parallel changes in infection control and antibiotic use?. <i>Critical Care</i> , 2009 , 13, P316	10.8	78
845	Prehospital administered intravenous antimicrobial protocol for septic shock: a prospective randomized clinical trial. <i>Critical Care</i> , 2009 , 13, P317	10.8	5
844	Clinical experiment with the use of colistin for the treatment of multiresistant Acinetobacter baumanii nosocomial infection. <i>Critical Care</i> , 2009 , 13, P318	10.8	78
843	Evaluation of gentamicin first-dose pharmacokinetics in septic critically ill patients: pilot study. <i>Critical Care</i> , 2009 , 13, P319	10.8	1
842	Does early appropriate antibiotic therapy improve the outcome of severe sepsis or septic shock?. <i>Critical Care</i> , 2009 , 13, P320	10.8	78
841	Strategy to reduce antibiotic prescription in cases of airway aspiration. <i>Critical Care</i> , 2009 , 13, P321	10.8	78
840	Effect of intravenous administration of freeze-dried sulfonated human normal immunoglobulin for septic patients. <i>Critical Care</i> , 2009 , 13, P322	10.8	78
839	Levels of soluble fibrin in severe septic patients given intravenous immunoglobulin. <i>Critical Care</i> , 2009 , 13, P323	10.8	78
838	Efficacy of corticosteroids on survival in patients with sepsis and septic shock: meta-analysis. <i>Critical Care</i> , 2009 , 13, P324	10.8	1
837	Effect of hydrocortisone therapy on outcome and the incidence of infection in patients with septic shock. <i>Critical Care</i> , 2009 , 13, P325	10.8	78
836	Effects of methylprednisolone on sepsis-induced blood B rain barrier permeability and reflex responsiveness in rats. <i>Critical Care</i> , 2009 , 13, P326	10.8	78
835	Systemic administration of E-selectin-directed dexamethasone liposomes attenuates pulmonary inflammation in a mouse model of ventilator-induced lung injury. <i>Critical Care</i> , 2009 , 13, P327	10.8	78

834	Relative adrenal insufficiency in cardiopulmonary bypass surgery patients: impact on the postoperative hemodynamic status. <i>Critical Care</i> , 2009 , 13, P328	10.8	78
833	Adrenal gland evaluation in septic shock patients: preliminary results of the first CT-scan study. <i>Critical Care</i> , 2009 , 13, P329	10.8	78
832	Effects of systemic steroid in patients with severe community-acquired pneumonia requiring mechanical ventilation. <i>Critical Care</i> , 2009 , 13, P330	10.8	78
831	Statin therapy in patients admitted to hospital with presumed infection. <i>Critical Care</i> , 2009 , 13, P331	10.8	78
830	Evolution of inflammation in non-ICU patients with infections: pilot prospective cohort study. <i>Critical Care</i> , 2009 , 13, P332	10.8	78
829	Heart rate variability in Egyptian children with acute rheumatic fever. <i>Critical Care</i> , 2009 , 13, P333	10.8	78
828	Recombinant human activated protein C reduces cardiac 3-nitrotyrosine and malondialdehyde levels in ovine acute respiratory distress syndrome and septic shock. <i>Critical Care</i> , 2009 , 13, P334	10.8	78
827	Altered plasma proteome during an early phase of an experimental model of peritonitis-induced sepsis. <i>Critical Care</i> , 2009 , 13,	10.8	78
826	Increased epithelial apoptosis and decreased oxidant injury in silymarin-treated rats with sepsis-induced acute lung injury. <i>Critical Care</i> , 2009 , 13, P336	10.8	1
825	Effect of nattokinase supplementation on plasma fibrinogen levels, whole blood viscosity and mortality in experimental sepsis in rats. <i>Critical Care</i> , 2009 , 13, P337	10.8	78
824	Physicians' perceptions of current management strategies for Gram-negative pneumonia: a multinational study. <i>Critical Care</i> , 2009 , 13, P338	10.8	78
823	Pilot proforma to aid the diagnosis of sepsis in burns patients. <i>Critical Care</i> , 2009 , 13, P339	10.8	78
822	Surviving sepsis campaign guidelines for severe sepsis and septic shock: implementation and outcome of a 3-year follow up. <i>Critical Care</i> , 2009 , 13, P340	10.8	78
821	Sepsis bundles: just think about it?. <i>Critical Care</i> , 2009 , 13, P341	10.8	78
820	Compliance with sepsis resuscitation but not management bundles improves the survival of surgical patients with septic shock. <i>Critical Care</i> , 2009 , 13, P342	10.8	78
819	Standard operating procedure in patients with severe sepsis and septic shock. <i>Critical Care</i> , 2009 , 13, P343	10.8	78
818	Epidemiology and outcome of patients with severe sepsis in six Spanish ICUs. <i>Critical Care</i> , 2009 , 13, P3	41 0.8	78
817	Epidemiology of severe sepsis in India. <i>Critical Care</i> , 2009 , 13, P345	10.8	3

816	Improving identification of severe sepsis by junior doctors: an observational study. <i>Critical Care</i> , 2009 , 13, P346	10.8	78
815	Prevalence of ICU infection in South Africa and accuracy of treating physician diagnosis and treatment. <i>Critical Care</i> , 2009 , 13, P347	10.8	2
814	Defining severe community-acquired pneumonia: a significant barrier to improving patient outcomes. <i>Critical Care</i> , 2009 , 13, P348	10.8	78
813	Outcome of severe sepsis in the ICU is independent of haemoglobin levels. <i>Critical Care</i> , 2009 , 13, P349	10.8	78
812	Abstract withdrawn. <i>Critical Care</i> , 2009 , 13, P350	10.8	78
811	TNFI+250 polymorphism and hyperdynamic state in cardiac surgery with extracorporeal circulation. <i>Critical Care</i> , 2009 , 13, P351	10.8	78
810	Role of 4G/5G plasminogen activator inhibitor-1 (PAI-1) gene polymorphism and the PAI-1 plasma levels in the outcome of patients with severe sepsis. <i>Critical Care</i> , 2009 , 13, P352	10.8	2
809	Role of monocyte apoptosis for the final outcome of the septic host with peritonitis. <i>Critical Care</i> , 2009 , 13, P353	10.8	78
808	Genomic variations within matrix metalloproteinase-9 and severe sepsis. <i>Critical Care</i> , 2009 , 13, P354	10.8	2
807	Association of IL-10 promoter polymorphism -1082 G/A with adverse outcome in severe sepsis and septic shock. <i>Critical Care</i> , 2009 , 13, P355	10.8	2
806	Activated protein C restores kidney function in endotoxin-induced acute renal failure in the rat. <i>Critical Care</i> , 2009 , 13, P356	10.8	1
805	Angiopoietin-2 is a mediator involved in the advent of hypotension following endotoxin shock. <i>Critical Care</i> , 2009 , 13, P357	10.8	78
804	Inhibition of the lectin-like oxidized low-density lipoprotein receptor-1 improves intestinal microcirculation in experimental endotoxaemia. <i>Critical Care</i> , 2009 , 13, P358	10.8	78
803	Effect of the a7nAChR agonist GTS-21 on inflammation during human endotoxemia. <i>Critical Care</i> , 2009 , 13, P359	10.8	1
802	Inflammatory response of critically ill and haemodialysis patients after whole blood stimulation by cell wall components of Gram-positive bacteria and fungi. <i>Critical Care</i> , 2009 , 13, P360	10.8	78
801	Asymmetric and symmetric dimethylarginines: metabolism and role in severe sepsis. <i>Critical Care</i> , 2009 , 13, P361	10.8	2
800	IL-2 modulates IFNImRNA gene expression in cultured peripheral blood mononuclear cells from septic patients. <i>Critical Care</i> , 2009 , 13, P362	10.8	78
799	Compliment activity patterns in patients with sepsis after human purified C1-esterase inhibitor infusion. <i>Critical Care</i> , 2009 , 13, P363	10.8	1

798	Detrimental hemodynamic and inflammatory effects of microparticles. Critical Care, 2009, 13, P364	10.8	78
797	Hydrogen sulfide: anti-inflammatory and cytoprotective effects. Critical Care, 2009, 13, P365	10.8	2
796	Dimethyl sulphoxide administration decreases renal ischemicEeperfusion injury. <i>Critical Care</i> , 2009 , 13, P366	10.8	78
795	Beneficial effects of the heme oxygenase-1/carbon monoxide system. <i>Critical Care</i> , 2009 , 13, P367	10.8	78
794	Assessment of IL-18 values in septic acute lung injury/acute respiratory distress syndrome patients. Critical Care, 2009, 13, P368	10.8	2
793	Cytochrome C is not released in the heart during sepsis-induced myocardial depression. <i>Critical Care</i> , 2009 , 13, P369	10.8	78
792	Strain-specific and pathogen-specific physiologic and genomic differences in murine inflammatory cardiac dysfunction. <i>Critical Care</i> , 2009 , 13, P370	10.8	78
791	Diastolic but not systolic dysfunction is associated with troponin and N-terminal pro-brain natriuretic peptide elevation in sepsis. <i>Critical Care</i> , 2009 , 13, P371	10.8	78
790	L-Threonine treatment enhances heat shock protein 25 expression and prevents apoptosis in heat-stressed intestinal epithelial-18 cells. <i>Critical Care</i> , 2009 , 13, P372	10.8	78
789	Repeated measurements of N-terminal pro-brain natriuretic peptide enable dynamic risk stratification in critically ill patients. <i>Critical Care</i> , 2009 , 13, P373	10.8	78
788	CT-pro-endothelin-1 and prognosis in critically ill patients with respiratory failure. <i>Critical Care</i> , 2009 , 13, P374	10.8	78
787	Serum high-mobility group box-1 protein as a specific marker of severe abdominal injury. <i>Critical Care</i> , 2009 , 13, P375	10.8	78
786	LightCycler SeptiFast in early diagnosis of sepsis: our experience. Critical Care, 2009, 13, P376	10.8	78
7 ⁸ 5	Differential diagnosis of systemic inflammatory response syndrome versus sepsis based on a multiplex quantitative PCR assay. <i>Critical Care</i> , 2009 , 13, P377	10.8	78
784	Sepsis in the emergency department: pathogen identification by blood cultures and PCR. <i>Critical Care</i> , 2009 , 13, P378	10.8	1
783	Rapid detection and identification of pathogens in critically ill and immunocompromised hosts using molecular techniques. <i>Critical Care</i> , 2009 , 13, P379	10.8	1
782	Microalbuminuria: a biomarker of sepsis. <i>Critical Care</i> , 2009 , 13, P380	10.8	1
781	Newly developed endotoxin measurement method (endotoxin activity assay) may reflect the severity of sepsis. <i>Critical Care</i> , 2009 , 13, P381	10.8	1

780	Activated protein Cprotein C inhibitor complex as a prognostic marker in sepsis. <i>Critical Care</i> , 2009 , 13, P382	10.8	78
779	Copeptin is a strong and independent predictor of outcome in cardiogenic shock. <i>Critical Care</i> , 2009 , 13, P383	10.8	4
778	Daily assay of procalcitonin, C-reactive protein and IL-6 roles in diagnosis and management of severe sepsis. <i>Critical Care</i> , 2009 , 13, P384	10.8	78
777	Effect of a clinical procalcitonin algorithm to guide antibiotic therapy in patients with lower respiratory tract infections outside study conditions: a post-study survey. <i>Critical Care</i> , 2009 , 13, P385	10.8	78
776	Effect of procalcitonin-based guidelines compared with standard guidelines on antibiotic use in lower respiratory tract infections: the randomized-controlled multicenter ProHOSP trial. <i>Critical Care</i> , 2009 , 13, P386	10.8	2
775	Procalcitonin: a prognostic marker for critically ill. <i>Critical Care</i> , 2009 , 13, P388	10.8	78
774	Abstract withdrawn. <i>Critical Care</i> , 2009 , 13, P389	10.8	78
773	Raised C-reactive protein on ICU discharge is not associated with subsequent poor outcome. <i>Critical Care</i> , 2009 , 13, P391	10.8	78
772	Influence of age on the distribution of morphine and morphine-3-glucuronide across the bloodBrain barrier in sheep. <i>Critical Care</i> , 2009 , 13, P392	10.8	78
771	Epidural anesthesia during surgery: friend or foe?. <i>Critical Care</i> , 2009 , 13, P393	10.8	78
770	Epidural analgesia compared with peripheral nerve blockade after major knee surgery. <i>Critical Care</i> , 2009 , 13, P394	10.8	2
769	Postconditioning effects following sevoflurane inhalational sedation in the ICU: a pilot study in cardiac surgery patients. <i>Critical Care</i> , 2009 , 13, P395	10.8	78
768	Multicenter randomized trial of sedation using daily wake-up calls, bispectral index or clinical sedation scores in a mixed medical Burgical ICU population. <i>Critical Care</i> , 2009 , 13, P396	10.8	3
767	Safety and length of different sedations for endoscopic procedures. <i>Critical Care</i> , 2009 , 13, P397	10.8	78
766	Novel responsiveness index measures the level of sedation in cardiac ICU patients. <i>Critical Care</i> , 2009 , 13, P398	10.8	78
765	Sedation protocols for intubated patients and noninvasive ventilation: additional concepts for a noniatrogenic intensive care. <i>Critical Care</i> , 2009 , 13, P399	10.8	2
764	Does choice of sedative agent affect duration of ICU stay, mortality or neurological outcome in patients undergoing therapeutic hypothermia?. <i>Critical Care</i> , 2009 , 13, P400	10.8	78
763	Positive inotropic effects of midazolam anaesthesia in rats during noninvasive heart studies. <i>Critical Care</i> , 2009 , 13, P401	10.8	78

762	Comparison of cerebral hemodynamic variables in hemorrhagic stroke using dexmedetomidine propofol versus dexmedetomidine bidazolam. <i>Critical Care</i> , 2009 , 13, P402	10.8	78
761	Effect of low-dose ketamine regimen with or without magnesium sulphate adjunct in tramadol patient controlled analgesia in a major abdominal surgery ICU population. <i>Critical Care</i> , 2009 , 13, P403	10.8	78
760	Impact of ketamine on dynamic compliance and airway resistance of sedated and mechanically ventilated ICU patients. <i>Critical Care</i> , 2009 , 13, P404	10.8	1
759	Etomidate versus ketamine for rapid sequence intubation in acutely ill patients: a multicenter randomized controlled trial. <i>Critical Care</i> , 2009 , 13, P405	10.8	1
758	Effect of clonidine on vomiting and diarrhea associated with ultra-rapid opioid detoxification: a randomized clinical trial. <i>Critical Care</i> , 2009 , 13, P406	10.8	1
757	Efficacy and pharmacokinetics of intravenous paracetamol in the critically ill patient. <i>Critical Care</i> , 2009 , 13, P407	10.8	1
756	Abstract withdrawn. <i>Critical Care</i> , 2009 , 13, P408	10.8	78
755	Predictive value of a preoperative respiratory failure index on the development of postoperative delirium. <i>Critical Care</i> , 2009 , 13, P409	10.8	78
754	Risk factors for intensive care delirium. <i>Critical Care</i> , 2009 , 13, P410	10.8	1
753	Delirium assessment in ICU patients: a comparison study. <i>Critical Care</i> , 2009 , 13, P411	10.8	1
752	Effect of magnesium on the incidence of delirium occurrence in the ICU. Critical Care, 2009, 13, P412	10.8	78
751	Memory dysfunction after brain traumatic injury depends on the Marshall score but not on the duration of sedation: preliminary findings. <i>Critical Care</i> , 2009 , 13, P413	10.8	78
750	COX-2 and E-selectin expression evaluation after acute normovolemic hemodilution. <i>Critical Care</i> , 2009 , 13,	10.8	78
749	Influence of hematocrit detection methodology on transfusion practice. Critical Care, 2009, 13, P415	10.8	78
			2
748	Effect of correction of anemia in severe congestive heart failure. <i>Critical Care</i> , 2009 , 13, P416	10.8	
748	Effect of correction of anemia in severe congestive heart failure. <i>Critical Care</i> , 2009 , 13, P416 Effect of length of storage of red blood cell units on outcome in critically ill children. <i>Critical Care</i> , 2009 , 13, P417	10.8	
	Effect of length of storage of red blood cell units on outcome in critically ill children. <i>Critical Care</i> ,		78

744	CT scans at outlying hospitals improve understanding of advanced trauma life support noncompliance. <i>Critical Care</i> , 2009 , 13, P420	10.8	78
743	Blunt abdominal trauma in children: a score to predict the absence of organ injury. <i>Critical Care</i> , 2009 , 13, P421	10.8	78
742	Risk of infection to packed gauze in damage control surgery for patients with hemorrhagic shock: safe limit of duration of packing. <i>Critical Care</i> , 2009 , 13, P422	10.8	2
74 ¹	Comparison between open and open-vacuum-assisted closure treatment in the adjuvant treatment of the abdominal compartment syndrome. <i>Critical Care</i> , 2009 , 13, P423	10.8	78
740	Upper gastrointestinal tract bleeding in the ICU. <i>Critical Care</i> , 2009 , 13, P424	10.8	78
739	A circulating factor modulates apoptosis of lymphocytes and monocytes after multiple trauma. <i>Critical Care</i> , 2009 , 13, P425	10.8	78
738	Abstract withdrawn. <i>Critical Care</i> , 2009 , 13, P426	10.8	78
737	Treatment with p-hydroxyphenylpyruvate in the absence of resuscitation extends survival in rats subjected to lethal blood loss. <i>Critical Care</i> , 2009 , 13, P427	10.8	78
736	Guidelines-associated decrease in mortality with recombinant activated factor VII for refractory hemorrhage. <i>Critical Care</i> , 2009 , 13, P428	10.8	78
735	Recombinant activated factor VII: does it really save lives?. <i>Critical Care</i> , 2009 , 13, P429	10.8	78
734	Recombinant activated factor VII for the treatment of bleeding in abdominal surgery and cardiac surgery. <i>Critical Care</i> , 2009 , 13, P430	10.8	78
733	Safety of activated recombinant factor VII in patients with circulatory assist devices. <i>Critical Care</i> , 2009 , 13, P431	10.8	78
732	Analysis of the off-licence use of recombinant activated factor VII for patients with uncontrolled haemorrhage in a UK tertiary referral hospital. <i>Critical Care</i> , 2009 , 13, P432	10.8	78
731	Treatment of obstetric hemorrhage with recombinant activated factor VII. <i>Critical Care</i> , 2009 , 13, P433	10.8	78
730	Thromboelastography versus viscoelastography for evaluation of coagulopathies in patients after abdominal surgery for cancer. <i>Critical Care</i> , 2009 , 13, P434	10.8	78
729	Prediction of long-term survival and coagulation factor VIII levels: interactions with gender in a large hospital-based cohort. <i>Critical Care</i> , 2009 , 13, P435	10.8	78
728	Gel point and fractal microstructure of incipient blood clots are significant new markers of haemostasis. <i>Critical Care</i> , 2009 , 13, P436	10.8	78
727	Acquired bleeding model induced by dilutional coagulopathy in the rabbit. <i>Critical Care</i> , 2009 , 13, P437	10.8	78

726	Impact of nebulized unfractionated heparin and N-acetylcysteine in management of smoke inhalation injury. <i>Critical Care</i> , 2009 , 13, P438	10.8	2
725	Anti-activated factor (response to enoxaparin in critically ill patients. Critical Care, 2009, 13, P439	10.8	78
724	Effect of amniotic fluid on blood coagulation. <i>Critical Care</i> , 2009 , 13, P440	10.8	78
723	Is it safe to thrombolyse patients with pulmonary embolism and right heart mobile thrombi?. <i>Critical Care</i> , 2009 , 13, P441	10.8	78
722	Argatroban for heparin-induced thrombocytopenia therapy in ICU patients with sepsis or multiple organ dysfunction syndrome. <i>Critical Care</i> , 2009 , 13, P442	10.8	78
721	Compression ultrasonography training program for ICU physicians does improve deep vein thrombosis risk management. <i>Critical Care</i> , 2009 , 13, P443	10.8	78
720	Prothrombotic state at hospital discharge increases 1-year mortality in pneumonia and sepsis survivors. <i>Critical Care</i> , 2009 , 13, P444	10.8	78
719	Thromboembolic risk factors and thromboprophylaxis in a medicalBurgical ICU in Mexico. <i>Critical Care</i> , 2009 , 13, P445	10.8	78
718	Hydroxyethyl starch or gelatin impairs, but Ringer's acetate enhances, coagulation capacity dose dependently after cardiac surgery. <i>Critical Care</i> , 2009 , 13, P446	10.8	78
717	Effects of Ringer's lactate or Ringerfundin resuscitation on the acid B ase status and serum electrolytes in septic oncologic patients. <i>Critical Care</i> , 2009 , 13, P447	10.8	1
716	Is Stewart's approach useful in evaluation of acidBase disorders in septic shock patients?. <i>Critical Care</i> , 2009 , 13, P448	10.8	78
715	Comparative study of factors that contribute to the severity of metabolic alkalosis in a surgical ICU. <i>Critical Care</i> , 2009 , 13, P450	10.8	78
714	AcidBase disorders evaluation in critically ill patients: hyperchloremia is associated with mortality. <i>Critical Care</i> , 2009 , 13, P451	10.8	78
713	Buffer therapy in severe metabolic or mixed acidosis. <i>Critical Care</i> , 2009 , 13, P452	10.8	78
712	Incidence, severity and timing of hypophosphataemia in Glasgow Royal Infirmary ICU. <i>Critical Care</i> , 2009 , 13, P453	10.8	78
711	Hypernatraemia in the neurointensive care unit: central diabetes insipidus and noncentral diabetes insipidus. <i>Critical Care</i> , 2009 , 13, P454	10.8	78
710	Impact of hypernatremia, hyponatremia and their comparison on ICU mortality. <i>Critical Care</i> , 2009 , 13, P455	10.8	1
709	Hyponatremia is associated with increased mortality in patients with acute decompensated hear failure. <i>Critical Care</i> , 2009 , 13, P456	10.8	78

708	Abnormal liver function tests are associated with increased mortality in both cardiothoracic and general intensive care. <i>Critical Care</i> , 2009 , 13, P457	10.8	1
707	Sexual hormone pattern in chronic critically ill patients. <i>Critical Care</i> , 2009 , 13, P458	10.8	78
706	Economic comparison of the traditional bathing method with the basinless bathing method in coma patients. <i>Critical Care</i> , 2009 , 13, P459	10.8	78
705	In critically ill patients, how often is their weight estimated and how accurate is that estimate?. <i>Critical Care</i> , 2009 , 13, P460	10.8	1
704	Alcohol misuse requiring admission to a tertiary referral ICU: resource implications. <i>Critical Care</i> , 2009 , 13, P461	10.8	2
703	Systemic rheumatic diseases in the ICU. <i>Critical Care</i> , 2009 , 13, P462	10.8	78
702	Distances covered by ICU nurses during their shifts. <i>Critical Care</i> , 2009 , 13, P463	10.8	78
701	ICU triage in terror-related mass casualty events. <i>Critical Care</i> , 2009 , 13, P464	10.8	1
700	Outcome of weekend and afterhour admissions compared with weekday and daytime admissions in the ICU with onsite intensivist coverage. <i>Critical Care</i> , 2009 , 13, P465	10.8	78
699	Scottish audit of interhospital transfers of acutely ill adults. <i>Critical Care</i> , 2009 , 13, P466	10.8	78
698	Outcome and resource utilization following interhospital transfer of critically ill patients. <i>Critical Care</i> , 2009 , 13, P467	10.8	78
697	Patient monitor and ventilation workstation alarming patterns during cardiac surgery. <i>Critical Care</i> , 2009 , 13, P468	10.8	78
696	Second-day discharge from the ICU after non-emergency cardiac surgery. <i>Critical Care</i> , 2009 , 13, P469	10.8	78
695	Audit on compliance with NICE Clinical Guideline 50 (NCG 50): 'Recognition of and response to acute illness in adults in hospital'. <i>Critical Care</i> , 2009 , 13, P470	10.8	78
694	Unique rapid response system with emergency call using an inhospital whole paging system. <i>Critical Care</i> , 2009 , 13, P471	10.8	78
693	Influence of medical emergency team call system on critically ill patients. <i>Critical Care</i> , 2009 , 13, P472	10.8	78
692	Impact of critical care outreach in blood and marrow transplantation. Critical Care, 2009, 13, P473	10.8	78
691	Audit of care provided prior to ICU admission. <i>Critical Care</i> , 2009 , 13, P474	10.8	78

690	Covering the cardiac ICU at night: a national survey of existing levels of cover and the potential for change with implementation of the European working time directive. <i>Critical Care</i> , 2009 , 13, P475	10.8	78
689	Timing of first review of new ICU admissions by consultant intensivists in a UK district general hospital. <i>Critical Care</i> , 2009 , 13, P476	10.8	78
688	Effects of establishing an intermediate care unit: changes in case mix at a cardiologic-pulmonary ICU. <i>Critical Care</i> , 2009 , 13, P477	10.8	78
687	Improvement in clinical outcomes following institution of dedicated critical care support in the high-dependency unit. <i>Critical Care</i> , 2009 , 13, P478	10.8	78
686	Effects of a closed ICU model on a general surgical ICU. <i>Critical Care</i> , 2009 , 13, P479	10.8	1
685	Early transfer to a high-volume ICU (upgrading) reduces mortality. <i>Critical Care</i> , 2009 , 13, P480	10.8	78
684	Teenagers: paediatric or adult ICU?. <i>Critical Care</i> , 2009 , 13, P481	10.8	78
683	Verbal orders drug therapy and prescription in a UK ICU setting. <i>Critical Care</i> , 2009 , 13, P482	10.8	O
682	Effects of organizational change in the ICU on high-risk surgical patients: a comparison of open and closed formats. <i>Critical Care</i> , 2009 , 13, P483	10.8	78
681	Study of the ability of ICU staff to set a bed to the semi-recumbent position. <i>Critical Care</i> , 2009 , 13, P48	4 10.8	78
680	Diagnosis-related group-based reimbursement is unrealistic for ICUs. <i>Critical Care</i> , 2009 , 13, P485	10.8	78
679	Waiting rooms and facilities available to patients' families and visitors in Italian pediatric ICUs: a national survey. <i>Critical Care</i> , 2009 , 13, P486	10.8	78
678	Patient satisfaction in the ICU: level of satisfaction and influencing factors. Critical Care, 2009, 13, P487	10.8	4
677	Use of deferred proxy consent in two emergency critical care trials. <i>Critical Care</i> , 2009 , 13, P488	10.8	78
676	Outcome and prognostic factors in critically ill cancer patients with diagnosis of sepsis in the ICU. <i>Critical Care</i> , 2009 , 13, P489	10.8	78
675	Toxic epidermal necrolysis: review of 15 cases. <i>Critical Care</i> , 2009 , 13, P490	10.8	78
674	Pesticide poisoning in the region of Rodopi, Greece. <i>Critical Care</i> , 2009 , 13, P491	10.8	78
673	Characterization of organophosphate poisoning patients in the ICU: a 4-year review. <i>Critical Care</i> , 2009 , 13, P492	10.8	78

672	Metformin-associated lactic acidosis: three patients surviving after continuous venovenous haemofiltration. <i>Critical Care</i> , 2009 , 13, P493	10.8	2
671	Epidemiology and nature of adverse events in the ICU. <i>Critical Care</i> , 2009 , 13, P494	10.8	78
670	Drug-drug interactions in the ICU. Critical Care, 2009, 13, P495	10.8	1
669	Outcome of patients with metastatic cancer or haematological malignancy in intensive care. <i>Critical Care</i> , 2009 , 13, P496	10.8	78
668	Mortality rate of vasopressor-dependent septic shock from 1997 through 2006. <i>Critical Care</i> , 2009 , 13, P497	10.8	78
66 7	Patients with severe sepsis and solid tumours: outcome and prognostic factors. <i>Critical Care</i> , 2009 , 13, P498	10.8	78
666	Evolution of acute lung injury/acute respiratory distress syndrome mortality in an ICU in Southern Brazil. <i>Critical Care</i> , 2009 , 13, P499	10.8	78
665	Demographic changes over a 12-year period in intensive care. <i>Critical Care</i> , 2009 , 13, P500	10.8	78
664	ICU admittance prediction of the seventh-day organ dysfunction clustering. <i>Critical Care</i> , 2009 , 13, P502	210.8	78
663	Serum lactic acid in the evaluation of outcome in the ICU. <i>Critical Care</i> , 2009 , 13, P503	10.8	78
662	APACHE II score may predict mortality in burns patients. <i>Critical Care</i> , 2009 , 13, P504	10.8	78
661	Scoring systems (APACHE II and Simplified Acute Physiology Score II) for predicting outcome in status asthmaticus. <i>Critical Care</i> , 2009 , 13, P505	10.8	78
660	Is Simplified Acute Physiology Score 3 better than APACHE II to predict mortality in transplanted critical patients?. <i>Critical Care</i> , 2009 , 13, P506	10.8	78
659	Inter-rater reliability of APACHE II scores in the medical ICU. Critical Care, 2009, 13, P507	10.8	78
658	Validation of Simplified Acute Physiology Score II and Simplified Acute Physiology Score III as mortality and morbidity risk models. <i>Critical Care</i> , 2009 , 13, P508	10.8	1
657	Performance of the APACHE IV system in patients with acute renal failure. <i>Critical Care</i> , 2009 , 13, P509	10.8	78
656	APACHE IV: benchmarking in an Indian ICU. Critical Care, 2009, 13, P510	10.8	3
655	Validation of the APACHE IV scoring system in patients with severe sepsis and comparison with the APACHE II system. <i>Critical Care</i> , 2009 , 13, P511	10.8	8

654	Prolonged intensive care stay and subsequent psychological distress: a study in cardiac patients. <i>Critical Care</i> , 2009 , 13, P512	10.8	78
653	Prediction of hospital mortality by classification trees and smoothing splines in patients with haematological malignancies admitted to the ICU. <i>Critical Care</i> , 2009 , 13, P513	10.8	78
652	Health-related quality of life is related to the lactate level at ICU admission in brain traumatic injury and major trauma patients. <i>Critical Care</i> , 2009 , 13, P514	10.8	78
651	Long-term survival after surgical intensive care admission: 50% die within 10 years. <i>Critical Care</i> , 2009 , 13, P515	10.8	78
650	Quality of life 1 year after intensive care. <i>Critical Care</i> , 2009 , 13, P516	10.8	78
649	Health-related quality of life in critically ill octogenarians: a long-term follow-up study. <i>Critical Care</i> , 2009 , 13, P517	10.8	78
648	Assessment and health-related quality of life evaluation of critical care patients, 1 year after intensive care discharge. <i>Critical Care</i> , 2009 , 13, P518	10.8	78
647	Clinical characteristics and prognostic factors of severe acute pancreatitis. <i>Critical Care</i> , 2009 , 13, P501	10.8	78
646	Abstract withdrawn. <i>Critical Care</i> , 2009 , 13, P390	10.8	78
645	Optimization of antibiotic therapy by procalcitonin. <i>Critical Care</i> , 2009 , 13, P387	10.8	78
644	Acidbase.org: online decision support in complex acidBase disorders using the Stewart approach. <i>Critical Care</i> , 2009 , 13, P449	10.8	1
643	Inhaled corticosteroids in COPD and the risk of pneumonia. <i>Lancet, The</i> , 2009 , 374, 668-70	40	21
642	Time to define the disorders of the syndrome of COPD. <i>Lancet, The</i> , 2009 , 374, 670-2	40	38
641	Severe community-acquired pneumonia. <i>Infectious Disease Clinics of North America</i> , 2009 , 23, 503-20	6.5	55
640	Actualit□ s dans la pneumopathie communautaire. 2009 , 1, 545-547		
639	Pneumonia in the elderly. 2009 , 15, 223-9		25
638	Current world literature. 2009 , 22, 198-213		
637	Community-acquired pneumonia. 2009 , 15, 261-73		19

636 Corticosteroids for pneumonia. 2009,

625	Statins for community-acquired pneumonia: current state of the science. 2010 , 29, 143-52		27
635	Statins for community acquired phedmonia. current state of the science. 2010, 25, 145-52		37
634	[New challenges in community-acquired pneumonia]. 2010 , 46 Suppl 6, 22-6		1
633	Dysphagia and aspiration pneumonia in older adults. 2010 , 22, 17-22		65
632	The influence of COPD on mortality and severity scoring in community-acquired pneumonia. 2010 , 79, 46-53		31
631	Infection as a comorbidity of COPD. European Respiratory Journal, 2010, 35, 1209-15	13.6	160
630	Efficacy of corticosteroids in community-acquired pneumonia: a randomized double-blinded clinical trial. 2010 , 181, 975-82		199
629	Emerging drugs for pneumococcal pneumonia. 2011 , 16, 459-77		3
628	Arguments against corticosteroids in community acquired pneumonia. 2011 , 47, 219-21		
627	Arguments in Favor of Corticosteroids in Pneumonia. 2011 , 47, 222-223		
626	Bacteraemic pneumococcal pneumonia: current therapeutic options. 2011 , 71, 131-53		30
625	Corticosteroids for pneumonia. 2011 , CD007720		27
624	Effect of corticosteroids on the clinical course of community-acquired pneumonia: a randomized controlled trial. <i>Critical Care</i> , 2011 , 15, R96	10.8	84
623	Dexamethasone and length of hospital stay in patients with community-acquired pneumonia: a randomised, double-blind, placebo-controlled trial. <i>Lancet, The</i> , 2011 , 377, 2023-30	40	225
622	Impact of systemic corticosteroids on the clinical course and outcomes of patients with severe community-acquired pneumonia: a cohort study. <i>Journal of Critical Care</i> , 2011 , 26, 193-200	4	40
621	Corticoides en la neumon adquirida en la comunidad. Argumentos en contra. 2011 , 47, 219-221		
620	Arguments in favor of corticosteroids in pneumonia. 2011 , 47, 222-3		
619	Analysis of systemic corticosteroid usage and survival in patients requiring mechanical ventilation for severe community-acquired pneumonia. 2011 , 17, 449-55		10

618	Use of early corticosteroid therapy on ICU admission in patients affected by severe pandemic (H1N1)v influenza A infection. <i>Intensive Care Medicine</i> , 2011 , 37, 272-83	14.5	164
617	Adjunctive therapies in severe pneumonia in critical care patients. 2011 , 11, 395-400		3
616	Adjunct corticosteroids in children hospitalized with community-acquired pneumonia. 2011 , 127, e255-	63	39
615	How Does One Diagnose and Manage Severe Community-Acquired Pneumonia?. 2011 , 253-262		
614	What is the role of steroids in pneumonia therapy?. 2012 , 25, 199-204		15
613	Adjunctive therapy in community-acquired pneumonia. <i>Seminars in Respiratory and Critical Care Medicine</i> , 2012 , 33, 311-8	3.9	19
612	Severity and outcomes of hospitalised community-acquired pneumonia in COPD patients. <i>European Respiratory Journal</i> , 2012 , 39, 855-61	13.6	69
611	Guidelines for diagnosis and management of community- and hospital-acquired pneumonia in adults: Joint ICS/NCCP(I) recommendations. 2012 , 29, S27-62		71
610	Corticosteroids in the treatment of community-acquired pneumonia in adults: a meta-analysis. <i>PLoS ONE</i> , 2012 , 7, e47926	3.7	73
609	[Influence of systemic corticosteroid administration in the prognosis of patients with community-acquired pneumonia]. 2012 , 212, 337-43		1
608	Techniques to measure cardiac output: minimally invasive method versus thermodilution. <i>Critical Care</i> , 2012 , 16,	10.8	78
607	Effects of cardiac output levels on the measurement of transpulmonary thermodilution cardiac output in patients with acute lung injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
606	Comparison of bioimpedance and oesophageal Doppler cardiac output monitoring during abdominal aortic surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
605	Cardiac output monitoring in cirrhotic patients: EV1000 versus pulmonary artery catheter - preliminary data. <i>Critical Care</i> , 2012 , 16,	10.8	78
604	Pulse contour cardiac output monitoring is less reliable in critically ill children. <i>Critical Care</i> , 2012 , 16,	10.8	78
603	Impact of arterial catheter location on the accuracy of cardiac output provided by an endotracheal bioimpedance device. <i>Critical Care</i> , 2012 , 16,	10.8	78
602	Accuracy of the PiCCO2-derived pulse contour cardiac index (CIpc): development and validation of a calibration index in two independent collectives. <i>Critical Care</i> , 2012 , 16,	10.8	78
601	Cardiac output monitoring using the LiDCOplusImonitor in abdominal aortic surgery: changes in calibration factor in aortic aneurysm disease versus aortic occlusive disease. <i>Critical Care</i> , 2012 , 16,	10.8	1

600	Haemodynamic changes during the peri-extubation period using bioreactance flow monitoring. <i>Critical Care</i> , 2012 , 16,	10.8	78
599	Left ventricular stroke volume measurement by impedance cardiography correlates with echocardiography in neonates. <i>Critical Care</i> , 2012 , 16,	10.8	2
598	Validation of less-invasive hemodynamic monitoring with Pulsioflex in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	1
597	A preliminary study on the use of noninvasive hemodynamic monitoring with the Nexfin monitor in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
596	Computer-based monitoring of global cardiovascular dynamics during acute pulmonary embolism and septic shock in swine. <i>Critical Care</i> , 2012 , 16,	10.8	78
595	Homeodynamic complexity: multifractal analysis of physiological instability. <i>Critical Care</i> , 2012 , 16,	10.8	78
594	Accuracy of conventional urinary output monitoring in the ICU. Critical Care, 2012, 16,	10.8	2
593	Changes in B-type natriuretic peptide and related hemodynamic parameters following a fluid challenge in patients with severe sepsis or septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
592	The Brody effect to detect hypovolemia in clinical practice. <i>Critical Care</i> , 2012 , 16,	10.8	1
591	Stroke volume variation guided fluid therapy in septic shock with ARDS. <i>Critical Care</i> , 2012 , 16,	10.8	78
590	Assessment fluid responsiveness in septic shock patients: a comparison of automated pulse pressure variation and manually calculated pulse pressure variation. <i>Critical Care</i> , 2012 , 16,	10.8	78
589	Applicability of methods for fluid responsiveness prediction in the ICU. Critical Care, 2012, 16,	10.8	78
588	Fluid responsiveness during weaning from mechanical ventilation. Critical Care, 2012, 16,	10.8	78
587	Microcirculatory blood flow is related to clinical signs of impaired organ perfusion, and its dynamics to the macrohemodynamic concept of fluid responsiveness. <i>Critical Care</i> , 2012 , 16,	10.8	1
586	Frank-Starling and Guyton together at bedside during a fluid challenge. Critical Care, 2012, 16,	10.8	78
585	Prediction of fluid responsiveness in intensive care (PREFERENCE study): fluid challenge versus passive leg raising in high-risk surgical patients. <i>Critical Care</i> , 2012 , 16,	10.8	5
584	Tight control of fluid balance may reduce incidence of intra-abdominal hypertension in patients after major abdominal surgery and trauma: a pilot study. <i>Critical Care</i> , 2012 , 16,	10.8	78
583	Negative fluid balance 48 hours after admission improves survival at 28 days in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78

(2012-2012)

582	Goal-directed fluid and hemodynamic therapy in major colon surgery with the pressure recording analytical method cardiac output monitor (MostCare[] -PRAM[]): prospective analysis of 58 patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
581	Implementation of an optimal fluid management protocol using the PiCCO system delays development of ARDS secondary to severe sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	1
580	Confrontation of the increase in ELWI rate regarding the septic polytraumatised patient administering furosemide: is it effective?. <i>Critical Care</i> , 2012 , 16,	10.8	78
579	Indexation of extravascular lung water in unselected adult patients with and without mechanical ventilation: a prospective study in 50 patients with 843 transpulmonary thermodilutions. <i>Critical Care</i> , 2012 , 16,	10.8	78
578	How to perform indexing of extravascular lung water data. Critical Care, 2012, 16,	10.8	78
577	Near-normal values of extravascular lung water in children. <i>Critical Care</i> , 2012 , 16,	10.8	78
576	Transthoracic ultrasound assessment of B-lines for identifying the increment of extravascular lung water in shock patients requiring fluid resuscitation. <i>Critical Care</i> , 2012 , 16,	10.8	78
575	Fluid therapy tactics in patients with polytrauma during interhospital transportation. <i>Critical Care</i> , 2012 , 16,	10.8	78
574	A prospective, randomized, clinical trial comparing the hemodynamics, efficacy, and safety of 6% hydroxyethyl starch 130/0.4 compared to albumin in postoperative patients undergoing pancreaticoduodenectomy. <i>Critical Care</i> , 2012 , 16,	10.8	78
573	Effect of balanced versus unbalanced HES solution on cytokine response in a rat model of peritonitis. <i>Critical Care</i> , 2012 , 16,	10.8	78
572	Evaluation of effectiveness and safety of hydroxyethyl starch (HES 130 kDa/0.4) in burn resuscitation. <i>Critical Care</i> , 2012 , 16,	10.8	1
571	Normal saline resuscitation worsens lactic acidosis in experimental sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	4
57°	Albumin in early septic shock resuscitation: examination of plasma and urine inflammatory markers. <i>Critical Care</i> , 2012 , 16,	10.8	78
569	Study of the correlation between central venous oxygen saturation and venous saturation from the antecubital vein in severe sepsis/septic shock patients. <i>Critical Care</i> , 2012 , 16,	10.8	1
568	Central venous hyperoxia is related to changes in tissue perfusion and morbi-mortality of patients in shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
567	Curve analysis of tissue oxygen desaturation after a venous occlusion test does not identify the central venous hemoglobin oxygen saturation. <i>Critical Care</i> , 2012 , 16,	10.8	78
566	Lactate in burn patients: biomarker of sepsis and mortality. <i>Critical Care</i> , 2012 , 16,	10.8	1
565	Can we predict arterial lactate from venous lactate in the emergency department?. <i>Critical Care</i> , 2012 , 16,	10.8	2

564	Cross-correlation analysis of blood and microdialysis-assessed tissue lactate monitoring: a study in critically ill septic patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
563	Admission lactate and outcome after high-risk surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
562	Effects of induced relative hypoxia during the postoperative period of abdominal oncologic surgery, on hemoglobin and reticulocyte levels: a prospective, randomized controlled clinical trial. <i>Critical Care</i> , 2012 , 16,	10.8	1
561	Pre-emptive hypothermia during resuscitated porcine hemorrhagic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
560	Carbon monoxide therapy protects against hepatic microvascular injury in a mouse model of murine hemorrhagic shock and resuscitation. <i>Critical Care</i> , 2012 , 16,	10.8	78
559	Customized modeling to predict the use of vasopressors in ICUs. Critical Care, 2012, 16,	10.8	78
558	Implementation of the fifth link of the Chain of Survival concept for out-of-hospital cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
557	Examination of out-of-hospital cardiac arrest patients with the Utstein style in Saga prefecture, Japan. <i>Critical Care</i> , 2012 , 16,	10.8	1
556	Effectiveness and limitations of learning cardiopulmonary resuscitation with an automated external defibrillator in the curriculum of First Aid courses among lay people. <i>Critical Care</i> , 2012 , 16,	10.8	78
555	Survival after out-of-hospital cardiac arrest during nights and weekends. <i>Critical Care</i> , 2012 , 16,	10.8	78
554	CPR initiated after telephone-assisted instruction produces a better outcome of bystander-witnessed out-of-hospital cardiac arrests than no bystander CPR but is less effective than CPR on the bystander's own initiative. <i>Critical Care</i> , 2012 , 16,	10.8	78
553	Critical times in pediatric out-of-hospital cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	1
552	Don't stop your heart in front of your family: family as a bystander is associated with poor outcome of bystander-witnessed, bystander-CPR-performed out-of-hospital cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
551	Coronary perfusion pressure in a pig model of prolonged cardiac arrest treated by different modes of venoarterial extracorporeal membrane oxygenation and intraaortic balloon counterpulsation. <i>Critical Care</i> , 2012 , 16,	10.8	3
550	Capnometry successfully predicts outcome and determination of the cessation of cardiopulmonary resuscitation efforts. <i>Critical Care</i> , 2012 , 16,	10.8	1
549	Modified clinical decision rule for termination-of-resuscitation in cases of refractory out-of-hospital cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
548	Survival benefit for patients receiving antibiotics following out-of-hospital cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
547	Correlation between IL-6 and S-100B blood levels and outcome of post-cardiac arrest syndrome and influence of therapeutic hypothermia on these mediator blood levels. <i>Critical Care</i> , 2012 , 16,	10.8	78

546	Early neurological outcome prediction model after bystander-witnessed out-of-hospital cardiac arrest: a nationwide population-based study. <i>Critical Care</i> , 2012 , 16,	10.8	78
545	Helium ventilation is safe and feasible in ICU patients admitted after cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	2
544	Therapeutic hypothermia in an out-of-hospital arrest population: are we selecting appropriately?. <i>Critical Care</i> , 2012 , 16,	10.8	78
543	Therapeutic hypothermia for nonventricular fibrillation/ventricular tachycardia cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
542	Comparison of cold crystalloid and colloid infusions for induction of therapeutic hypothermia. <i>Critical Care</i> , 2012 , 16,	10.8	78
541	Cerebral oxygenation during induction of therapeutic hypothermia after cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
540	Survey on the management of patients treated with therapeutic hypothermia post cardiac arrest in London hospitals. <i>Critical Care</i> , 2012 , 16,	10.8	78
539	Simplified EEG/aEEG to monitor the injured brain after cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
538	Usefulness of a Bispectral index oriented sedative method without neuromuscular blocker for therapeutic hypothermia after cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
537	Predictive factors of neurologic outcome in therapeutic hypothermia after prehospital return of spontaneous circulation. <i>Critical Care</i> , 2012 , 16,	10.8	78
536	Employment status 1 year after out-of-hospital cardiac arrest in comatose patients treated with therapeutic hypothermia. <i>Critical Care</i> , 2012 , 16,	10.8	78
535	Changes in cerebrospinal fluid and serum cytokine levels in severe traumatic brain injury patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
534	Noninvasive cerebral oxygenation monitoring during rapid ventricular pacing in transcutaneous aortic valve implant. <i>Critical Care</i> , 2012 , 16,	10.8	78
533	Novel models to predict elevated intracranial pressure during intensive care and long-term neurological outcome after TBI. <i>Critical Care</i> , 2012 , 16,	10.8	78
532	Transcranial Doppler pulsatility index is a poor predictor of hydrocephalus in patients with aneurysmal subarachnoid haemorrhage. <i>Critical Care</i> , 2012 , 16,	10.8	78
531	Transcranial cerebral oximetry in newborn infants on mechanical ventilation as a method for prevention of hyperoxia and oxidative stress. <i>Critical Care</i> , 2012 , 16,	10.8	78
530	Cerebral oximetry and brain death in the ICU: data from seven cases. Critical Care, 2012, 16,	10.8	2
529	Deoxyhaemoglobin as a biomarker of cerebral autoregulation. <i>Critical Care</i> , 2012 , 16,	10.8	3

528	Study of the acoustic stem evoked potentials in blood circulation disorder in the vertebral basilar basin. <i>Critical Care</i> , 2012 , 16,	10.8	78
527	Examination of the autonomic nervous system in the ICU: a pilot study. Critical Care, 2012, 16,	10.8	78
526	Predictive value of glial fibrillary acidic protein for prognosis in patients with moderate and severe traumatic brain injury: a systematic review and meta-analysis. <i>Critical Care</i> , 2012 , 16,	10.8	3
525	Prevalence of pituitary disorders associated with traumatic brain injury: a systematic review. <i>Critical Care</i> , 2012 , 16,	10.8	2
524	Mannose binding lectin deficiency attenuates neurobehavioral deficits following experimental traumatic brain injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
523	Azathioprine and aspirin in treatment of childhood primary arterial stroke: therapeutic benefits and side effects. <i>Critical Care</i> , 2012 , 16,	10.8	78
522	Changes of ribosomal protein S3 immunoreactivity and its new expression in microglia in the mice hippocampus after lipopolysaccharide treatment. <i>Critical Care</i> , 2012 , 16,	10.8	78
521	Neuronal damage using Fluoro-Jade B histofluorescence and gliosis in the striatum after various durations of transient cerebral ischemia in gerbils. <i>Critical Care</i> , 2012 , 16,	10.8	3
520	Molecular, histological and microcirculatory modeling of cerebral ischemia in pigs. <i>Critical Care</i> , 2012 , 16,	10.8	78
519	Delayed post-ischaemic administration of xenon reduces brain damage in a rat model of global ischaemia. <i>Critical Care</i> , 2012 , 16,	10.8	78
518	Seizures in the respiratory ICU: single-center study of patients with new-onset seizures. <i>Critical Care</i> , 2012 , 16,	10.8	1
517	Early treatment with intravenous immunoglobulins in patients with critical illness polyneuropathy: a randomized controlled, double-blinded study. <i>Critical Care</i> , 2012 , 16,	10.8	78
516	Intracranial pressure monitoring in acute liver failure: a retrospective cohort study. <i>Critical Care</i> , 2012 , 16,	10.8	2
515	Retrospective observation of 6-month survival following decompressive craniectomy in a London major trauma and stroke centre. <i>Critical Care</i> , 2012 , 16,	10.8	1
514	Feasibility of a multicenter prospective cohort study on the evaluation of prognosis in severe traumatic brain injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
513	Predictive value of neuron-specific enolase following moderate and severe traumatic brain injury: a systematic review and meta-analysis. <i>Critical Care</i> , 2012 , 16,	10.8	78
512	Blood-brain barrier permeability following traumatic brain injury. Critical Care, 2012, 16,	10.8	1
511	Can urinary 8-OHdG be a good indicator of vasospasm occurrence following subarachnoid hemorrhage?. <i>Critical Care</i> , 2012 , 16,	10.8	8

510	Cortical capillary recruitment by rosuvastatin in experimental brain trauma is associated with increased NO production. <i>Critical Care</i> , 2012 , 16,	10.8	78
509	Effects of sinvastatin in prevention of vasospasm in nontraumatic subarachnoid hemorrhage: preliminary data. <i>Critical Care</i> , 2012 , 16,	10.8	1
508	Evaluation of arterial and venous ophthalmic hemodynamics in preeclamptic pregnant women. <i>Critical Care</i> , 2012 , 16,	10.8	78
507	Data classification of magnetic resonance tomography and computer tomography images of brain in parturients with neurological complications of eclampsia. <i>Critical Care</i> , 2012 , 16,	10.8	78
506	Eleven years of critical obstetric pathology: epidemiologic study. Critical Care, 2012, 16,	10.8	78
505	Clinical outcomes in neonates following maternal magnesium sulfate therapy in preeclampsia/eclampsia. <i>Critical Care</i> , 2012 , 16,	10.8	78
504	Sleep monitoring by actigraphy in short-stay ICU patients. Critical Care, 2012, 16,	10.8	1
503	Quality and quantity of sleep in multipatient versus single-room ICUs. Critical Care, 2012, 16,	10.8	78
502	Oral melatonin in high-risk critically ill patients: quality of sedative effect. Critical Care, 2012, 16,	10.8	1
501	Sedation depth and mortality in mechanically ventilated critically ill adults. Critical Care, 2012, 16,	10.8	1
500	Sedation in the ICU: nurses' perceptions of practices and influencing factors. <i>Critical Care</i> , 2012 , 16,	10.8	78
499	Implementation of a national guideline for analgesia and sedation: how often can a RASS of 0 to -2 be achieved?. <i>Critical Care</i> , 2012 , 16,	10.8	78
498	Comparison of the RAMSAY score and the Richmond Agitation Sedation Score for the measurement of sedation depth. <i>Critical Care</i> , 2012 , 16,	10.8	12
497	Dexmedetomidine is associated with better outcomes in patients undergoing cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
496	Cerebral ischemia-reperfusion model in rabbits: relationship between dexmedetomidine and biochemical parameters in lowering intraparenchymal pressure. <i>Critical Care</i> , 2012 , 16,	10.8	78
495	Evaluation of sedation using pupilometry in ICUs: a pilot study. <i>Critical Care</i> , 2012 , 16,	10.8	78
494	Effect of critical illness on the pharmacokinetics and dose-response relationship of midazolam. <i>Critical Care</i> , 2012 , 16,	10.8	1
493	Effect of propofol and midazolan on microcirculation of septic shock patients. <i>Critical Care</i> , 2012 , 16,	10.8	78

492	Current use of pain scores in Dutch ICUs: a postal survey in the Netherlands. Critical Care, 2012, 16,	10.8	78
491	Efficiency estimation of intrapleural and thoracic paravertebral block in combination with general anesthesia at thoracoscopic interventions. <i>Critical Care</i> , 2012 , 16,	10.8	78
490	Preoperative diclofenc reduces postcraniotomy headache: a randomized, placebo-controlled trial. <i>Critical Care</i> , 2012 , 16,	10.8	78
489	Long-term adverse neuropsychological functioning in children who survived meningococcal septic shock: is there a relationship with sedation and analgesia during paediatric ICU admission?. <i>Critical Care</i> , 2012 , 16,	10.8	78
488	Delirium could be an indicator of sepsis in patients under 65 years old with urinary tract infections. <i>Critical Care</i> , 2012 , 16,	10.8	78
487	Delirium screening in critically ill patients: a systematic review and meta-analysis. <i>Critical Care</i> , 2012 , 16,	10.8	3
486	Electroencephalography-based monitoring of delirium in the ICU: what are the opportunities?. <i>Critical Care</i> , 2012 , 16,	10.8	0
485	Performance of SAPS 3 in predicting delirium among critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
484	Incidence of delirium and inadequacy of the clinical diagnosis in patients in intensive care. <i>Critical Care</i> , 2012 , 16,	10.8	78
483	Investigation into detection and treatment rates of hyperactive and hypoactive delirium in the ICU setting. <i>Critical Care</i> , 2012 , 16,	10.8	78
482	Memories and post-traumatic stress-related symptoms in older, post-cardiac surgery patients: substudy of an RCT. <i>Critical Care</i> , 2012 , 16,	10.8	78
481	Using tramadol to monitor hepatic drug metabolism in the critically ill. Critical Care, 2012, 16,	10.8	78
480	Data mining techniques for predicting acute kidney injury after elective cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	3
479	Acute kidney injury in critically ill patients with A/H1N1 pneumonitis in 2010/11. <i>Critical Care</i> , 2012 , 16,	10.8	78
478	A RIFLE score-based trigger for renal replacement therapy and survival after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
477	Effect of off-pump versus on-pump coronary artery bypass grafting in patients with chronic kidney disease. <i>Critical Care</i> , 2012 , 16,	10.8	1
476	Effects of renal-dose dopamine on renal tubular functions following coronary artery bypass grafting surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
475	Nurses' knowledge regarding the early identification of acute kidney injury. <i>Critical Care</i> , 2012 , 16,	10.8	78

474	Neutrophil gelatinase-associated lipocalin predicts postoperative fluid overload, a potentially modifiable risk factor for mortality after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	1
473	Plasma and urine neutrophil gelatinase-associated lipocalin as markers of acute kidney injury in critically ill adults. <i>Critical Care</i> , 2012 , 16,	10.8	78
472	Plasma and urine neutrophil gelatinase-associated lipocalin in septic and nonseptic ICU patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
471	Urinary neutrophil gelatinase-associated lipocalin as an early marker of acute kidney injury complicating circulatory shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
470	Additive value to clinical judgement of blood neutrophil gelatinase-associated lipocalin in diagnosis of acute kidney injury and prediction of mortality in patients hospitalized from the emergency department. <i>Critical Care</i> , 2012 , 16,	10.8	2
469	Is cystatin C reliable in the anesthetized pig? An experimental study with special reference to septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	1
468	Are serum cystatin-C-based estimates better than those derived from serum creatinine in critically ill patients?. <i>Critical Care</i> , 2012 , 16,	10.8	2
467	Assessment of glomerular filtration rate in trauma patients in early resuscitation phase. <i>Critical Care</i> , 2012 , 16,	10.8	1
466	Validation of a continuous low-dose iohexol infusion to measure the glomerular filtration rate. <i>Critical Care</i> , 2012 , 16,	10.8	78
465	Investigation into the effects of commencing haemodialysis in the critically ill. <i>Critical Care</i> , 2012 , 16,	10.8	78
464	Is the AKIN score useful as an indicator of the optimum time for intervention with renal replacement therapy in critically ill patients?. <i>Critical Care</i> , 2012 , 16,	10.8	78
463	Timing for initiation of continuous renal replacement therapy in patients with septic shock and acute kidney injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
462	Early application of CVVH In the complex treatment of patients with early severe acute pancreatitis. <i>Critical Care</i> , 2012 , 16,	10.8	1
461	Timing of the initiation of continuous renal replacement therapy and clinical outcome in patients with severe sepsis and septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
460	Amino acid concentrations in serum, urine and dialysate/ultrafiltrate solutions of continuous venovenous hemodiafiltration patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
459	Evaluation of the potential adverse effects associated with calcium carbonate precipitate during continuous venovenous hemofiltration. <i>Critical Care</i> , 2012 , 16,	10.8	78
458	Regional citrate anticoagulation in CVVH: a new protocol combining citrate solution with a phosphate-containing replacement fluid. <i>Critical Care</i> , 2012 , 16,	10.8	78
457	Regional citrate anticoagulation with a low-concentration solution in predilution-postdilution CVVH. <i>Critical Care</i> , 2012 , 16,	10.8	78

456	Exposure to intermittent hemodialysis and renal recovery after acute kidney injury: a systematic review. <i>Critical Care</i> , 2012 , 16,	10.8	2
455	Sustained low-efficiency dialysis for renal replacement therapy in the ICU: a cost-benefit analysis of the years 2006 to 2010. <i>Critical Care</i> , 2012 , 16,	10.8	78
454	The new dialysis method Mini-SLED is useful for dialyzing acute brain stroke patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
453	Investigation into haemodynamic stability during intermittent haemodialysis in the critically ill. <i>Critical Care</i> , 2012 , 16,	10.8	78
452	Evaluation of microcirculation before and during continuous renal replacement therapy and the impact of dose prescription. <i>Critical Care</i> , 2012 , 16,	10.8	78
451	Ultrafiltration during continuous hemofiltration in stabilized ICU patients is not associated with microcirculatory perfusion derangements. <i>Critical Care</i> , 2012 , 16,	10.8	78
450	Plasmapheresis without apparatus in complex care of victims with crush syndrome during the first hours after extrication in a field hospital of EMERCOM of Russia in emergency areas. <i>Critical Care</i> , 2012 , 16,	10.8	78
449	Degree of impaired kidney function at hospital discharge has a major impact on long-term survival of critically ill patients recovered from renal failure. <i>Critical Care</i> , 2012 , 16,	10.8	78
448	Long-term survival for ICU patients with acute kidney injury. Critical Care, 2012, 16,	10.8	78
447	Super high-flux continuous hemodialysis: an efficient compromise for blood purification in sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	78
446	Efficacy of continuous haemodiafiltration using a polymethylmethacrylate membrane haemofilter in the treatment of sepsis and acute respiratory distress syndrome. <i>Critical Care</i> , 2012 , 16,	10.8	1
445	Possible adsorption mechanism of high mobility group box 1 protein on a polyacrylonitrile (AN69ST) membrane filter. <i>Critical Care</i> , 2012 , 16,	10.8	78
444	High mobility group box 1 levels in septic disseminated intravascular coagulation patients undergoing Polymixin-B immobilized fiber-direct hemoperfusion. <i>Critical Care</i> , 2012 , 16,	10.8	78
443	Polymyxin B-immobilized fiber column hemoperfusion has the ability of endotoxin removal during 24 hours. <i>Critical Care</i> , 2012 , 16,	10.8	78
442	Polymyxin B-direct hemoperfusion therapy could contribute to hemodynamics and outcomes in emergency surgical patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
441	Clinical impact of enhanced cytokine clearance with sustained high-efficiency daily diafiltration using a mediator-adsorbing membrane (SHEDD-fA) in patients with severe sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	78
440	Mortality and priority level for ICU admission in the setting of limited critical care beds in El Salvador. <i>Critical Care</i> , 2012 , 16,	10.8	1
439	Mainz Emergency Evaluation Scoring in combination with capnometry predicts outcome in trauma patients. <i>Critical Care</i> , 2012 , 16,	10.8	78

438	Predicting outcome in the ICU: comparison of Ranson criteria and Ranson + CRP levels in acute pancreatitis. <i>Critical Care</i> , 2012 , 16,	10.8	1
437	Number of failed organs and response to therapy determine outcome in patients with acute pancreatitis requiring level 1 organ support. <i>Critical Care</i> , 2012 , 16,	10.8	78
436	Mortality predictors in acute pancreatitis admitted to the ICU. Critical Care, 2012, 16,	10.8	78
435	System biology prediction model based on clinical data: highly accurate outcome prediction in patients with acute-on-chronic liver failure. <i>Critical Care</i> , 2012 , 16,	10.8	78
434	Prognostic relevance of arterial ammonia levels in different acute and acute-on-chronic liver diseases. <i>Critical Care</i> , 2012 , 16,	10.8	1
433	Liver failure secondary to alcoholic liver disease carries a worse prognosis than other aetiologies of liver failure: retrospective analysis of routine biochemical markers in critically ill patients with liver failure. <i>Critical Care</i> , 2012 , 16,	10.8	1
432	Incidence, morbidity and mortality of admissions related to alcohol consumption on critical care: a single-centre experience. <i>Critical Care</i> , 2012 , 16,	10.8	1
431	Changing outcomes in patients with chronic liver disease in intensive care: a decade of experience. <i>Critical Care</i> , 2012 , 16,	10.8	2
430	Multivariate regression analysis of outcomes following orthotopic liver transplantation in decompensated cirrhotics transplanted from the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
429	Liver transplantation in the critically ill: a Canadian collaboration. <i>Critical Care</i> , 2012 , 16,	10.8	78
428	Acute respiratory distress syndrome: analysis of incidence and mortality in a university hospital critical care unit. <i>Critical Care</i> , 2012 , 16,	10.8	2
427	Epidemiology and outcome of sepsis syndromes in Italian ICUs: a regional multicenter observational cohort. <i>Critical Care</i> , 2012 , 16,	10.8	2
426	Outcome of faecal peritonitis in the ICU. Critical Care, 2012, 16,	10.8	2
425	HIV patients in the ICU: our experience. <i>Critical Care</i> , 2012 , 16,	10.8	78
424	Impact of congestive heart failure on severe sepsis and septic shock survivors: outcomes and performance status after 1-year hospital discharge. <i>Critical Care</i> , 2012 , 16,	10.8	3
423	Predictive value of N-terminal pro-brain natriuretic peptide among critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
422	Low preoperative hepcidin concentration is a risk factor for mortality but not for acute kidney injury after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
421	Outcomes and resource use for over 80 year olds admitted to a UK critical care unit after an emergency laparotomy over a 3-year period. <i>Critical Care</i> , 2012 , 16,	10.8	78

420	Correlation between APACHE II score and quality of life among patients discharged from the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
419	Parameters that affect outcome in surgical ICU patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
418	Relationship between illness severity scores in the ICU. Critical Care, 2012, 16,	10.8	78
417	Predicting hospital mortality: comparing accuracy of SAPS II and clinical staff prognosis. <i>Critical Care</i> , 2012 , 16,	10.8	78
416	Predictors of mortality in patients from a hematological ICU in Brazil. Critical Care, 2012, 16,	10.8	78
415	Retrospective study of the outcomes of patients admitted to the ICU with a hematological malignancy. <i>Critical Care</i> , 2012 , 16,	10.8	78
414	Six-month survival of patients with lung cancer admitted to a medical ICU: a retrospective study. <i>Critical Care</i> , 2012 , 16,	10.8	78
413	Health-related quality of life and survival of cancer patients admitted to ICUs: Results of the QALY study. <i>Critical Care</i> , 2012 , 16,	10.8	1
412	Characteristics, resource consumption and outcome of cancer patients admitted to ICUs. <i>Critical Care</i> , 2012 , 16,	10.8	78
411	Managing critically ill oncological patients in hospital: a survey across all ICUs in the UK. <i>Critical Care</i> , 2012 , 16,	10.8	78
410	Role of illness severity scores in predicting mortality in the coronary care unit. <i>Critical Care</i> , 2012 , 16,	10.8	78
409	New severity score of acute respiratory failure. <i>Critical Care</i> , 2012 , 16,	10.8	78
408	Validity of six prognostic scoring systems for septic shock patients admitted to a medical ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
407	Risk factors of venous thrombosis in knee joint endoprosthesis. <i>Critical Care</i> , 2012 , 16,	10.8	78
406	Proximal and distal deep venous thrombosis in critically ill patients: incidence and prevalence. <i>Critical Care</i> , 2012 , 16,	10.8	78
405	Saddle embolism is associated with the major adverse events in patients with nonhigh-risk pulmonary embolism. <i>Critical Care</i> , 2012 , 16,	10.8	1
404	Efficacy and safety of enoxaparin as deep vein thrombosis prophylaxis in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
403	Reducing the level of postoperative thrombotic complications by the combination of low molecular weight heparin and epidural anesthesia at the patients after total hysterectomy. <i>Critical Care</i> , 2012 , 16,	10.8	78

402	Cost-effectiveness analysis of two thromboprophylactic strategies following major surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
401	Consequences of suspected heparin-induced thrombocytopenia in the ICU. Critical Care, 2012, 16,	10.8	78
400	Evaluation of iron, transferrin and ferritin serum levels in patients with severe sepsis and septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	2
399	Using angiogenic factors and their soluble receptors to predict organ dysfunction in patients with disseminated intravascular coagulation associated with severe trauma. <i>Critical Care</i> , 2012 , 16,	10.8	78
398	A simple blood-saving bundle reduces diagnostic blood loss in mechanically ventilated patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
397	Comparative assessment of invasive and noninvasive Methods for detection of total hemoglobin in gynecological patients' blood. <i>Critical Care</i> , 2012 , 16,	10.8	4
396	Use of coagulation screening in the critical care unit. Critical Care, 2012, 16,	10.8	О
395	Templating effect of clot structure can predict clot development and outcome in diluted blood: a comparison with thromboelastography. <i>Critical Care</i> , 2012 , 16,	10.8	5
394	Fractal analysis: a new biomarker for determining clot characteristics in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	1
393	Fractal dimension: a biomarker for detecting acute thromboembolic disease. <i>Critical Care</i> , 2012 , 16,	10.8	1
392	Thromboelastography (platelet contribution to clot strength) for the assessment of platelet residual function. <i>Critical Care</i> , 2012 , 16,	10.8	78
391	Measurement of hemoglobin in the operating room: what Methods can we trust?. <i>Critical Care</i> , 2012 , 16,	10.8	78
390	Retrospective comparison study of warfarinised trauma patients and an age-matched control group of nonwarfarinised patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
389	In medical-surgical ICU patients, major bleeding is common but independent of heparin prophylaxis. <i>Critical Care</i> , 2012 , 16,	10.8	2
388	Reducing ICU blood draws with artificial intelligence. Critical Care, 2012, 16,	10.8	78
387	Hemostasis system condition in infectious complication development in severe burned patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
386	Randomized comparison of fibrinogen concentrate versus cryoprecipitate for bleeding control in pediatric cardiac surgery (FICCS study). <i>Critical Care</i> , 2012 , 16,	10.8	5
385	Efficacy of tranexamic acid in decreasing blood loss during cesarean section. <i>Critical Care</i> , 2012 , 16,	10.8	2

384	Reduced EPO receptor expression may contribute to limited pleiotropic effects of EPO during critical illness. <i>Critical Care</i> , 2012 , 16,	10.8	1
383	Recognition and management of haemophagocytic lymphohistiocytosis on the ICU: a case series. <i>Critical Care</i> , 2012 , 16,	10.8	78
382	Blood transfusion is an independent predicting factor for poor outcome after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
381	Red blood transfusion is a predictor of poor outcome in pediatric cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	1
380	No impact of a massive transfusion protocol on coagulopathy and mortality at a level 1 trauma center: why?. <i>Critical Care</i> , 2012 , 16,	10.8	78
379	Massive transfusion practice. <i>Critical Care</i> , 2012 , 16,	10.8	78
378	Red blood cell transfusion improves microdialysis-assessed interstitial lactate/pyruvate ratio in critically ill septic patients. <i>Critical Care</i> , 2012 , 16,	10.8	1
377	Blood transfusion after cardiac surgery increases the hospital length of stay in adult patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
376	Transfusion of blood stored for longer periods of time does not alter the reactive hyperemia index in healthy volunteers. <i>Critical Care</i> , 2012 , 16,	10.8	О
375	Liberal use of platelet transfusions in the acute phase of trauma resuscitation: a systematic review. <i>Critical Care</i> , 2012 , 16,	10.8	78
374	Impact on early trauma mortality of the adoption of the Updated European Guidelines on the management of bleeding. <i>Critical Care</i> , 2012 , 16,	10.8	78
373	Hemodynamics in the severely injured patient with significant hemorrhage. Critical Care, 2012, 16,	10.8	78
372	Critical older trauma patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
371	Outcomes in older blunt chest wall trauma patients: a retrospective study. Critical Care, 2012, 16,	10.8	2
370	Mortality trend alteration of thoracic injury after rapid response trauma team establishment. <i>Critical Care</i> , 2012 , 16,	10.8	78
369	Trauma patients and cervical spine protection in critical care: the impact of a spinal checklist on clinical care and documentation. <i>Critical Care</i> , 2012 , 16,	10.8	78
368	Effect of instrumented spinal fixation on outcome in polytrauma patients in the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
367	Whole body computed tomography scanning for severe blunt polytrauma: analysis of Trauma Audit and Research Network database 2005 to 2010. <i>Critical Care</i> , 2012 , 16,	10.8	1

366	Post-traumatic rhabdomyolysis: an observational study in seven patients. Critical Care, 2012, 16,	10.8	78
365	Exertional rhabdomyolysis in female amateur triathletes. <i>Critical Care</i> , 2012 , 16,	10.8	1
364	Controlled mechanical ventilation tactics in patients with polytrauma during interhospital transportation to the specialized center. <i>Critical Care</i> , 2012 , 16,	10.8	11
363	Impact of fluid resuscitation volume on the severity of organ failures in severely burned patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
362	Organ dysfunction in the resuscitation phase of critical burn patients. Critical Care, 2012, 16,	10.8	78
361	Epidemiological study of critical burn patients in an ICU. Critical Care, 2012, 16,	10.8	1
360	Fluid creep in burn resuscitation: the tide has not yet turned. Critical Care, 2012, 16,	10.8	5
359	Early administration of parenteral estrogen suppresses the deleterious local and systemic inflammatory response in severe burns. <i>Critical Care</i> , 2012 , 16,	10.8	78
358	Reducing the indication of ventilatory support in the severely burnt patient and improving outcomes: Results of a new protocol approach within a regional burns centre. <i>Critical Care</i> , 2012 , 16,	10.8	78
357	Cardiopulmonary exercise testing and elective open abdominal aortic aneurysm surgery over a 6-year period in a UK teaching hospital. <i>Critical Care</i> , 2012 , 16,	10.8	1
356	Perioperative evaluation of elective surgical patients: is it possible to plan ICU admission?. <i>Critical Care</i> , 2012 , 16,	10.8	78
355	Cardiac-specific biomarkers and life-threatening complications of off-pump versus on-pump coronary bypass surgery in Egyptian patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
354	Aortic aneurysm disease versus aortic occlusive disease: differences in postoperative ICU requirements after open elective abdominal aortic surgery. <i>Critical Care</i> , 2012 , 16,	10.8	1
353	High postoperative blood levels of macrophage migration inhibitory factor are associated with less organ dysfunction in patients after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
352	Predictors of prolonged mechanical ventilation after heart transplantation. <i>Critical Care</i> , 2012 , 16,	10.8	78
351	Atrial fibrillation following major noncardiac thoracic surgery: significance and impact on morbidity. <i>Critical Care</i> , 2012 , 16,	10.8	1
350	Oxygen delivery index during goal-directed therapy predicts complications and hospital length of stay in patients undergoing high-risk surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
349	Transfer delays in patients referred for neurosurgical intervention with traumatic brain injury. Critical Care, 2012 , 16,	10.8	1

348	Performances of ventilator at simulated altitude. Critical Care, 2012, 16,	10.8	78
347	Impact of H1N1 influenza on critical care and dependent services in Wales during winter 2010/2011. <i>Critical Care</i> , 2012 , 16,	10.8	78
346	Effects of levels of clinical supervision during simulated ICU scenarios on resident learning and patient care: a qualitative study. <i>Critical Care</i> , 2012 , 16,	10.8	78
345	Virtual reality and live scenario simulation: options for training medical students in mass casualty incident triage. <i>Critical Care</i> , 2012 , 16,	10.8	3
344	Utilization of iPad in the system of emergency demand and acceptance. Critical Care, 2012, 16,	10.8	78
343	Mass evacuation of victims from emergency areas by medical modules aboard the aircraft of EMERCOM of Russia. <i>Critical Care</i> , 2012 , 16,	10.8	78
342	Reliability and validity of an Italian four-level emergency triage system. Critical Care, 2012, 16,	10.8	78
341	Coordination of emergency resources after Lorca's earthquakes. Critical Care, 2012, 16,	10.8	78
340	Lightning injuries in a lightning city: a district hospital experience in Singapore. <i>Critical Care</i> , 2012 , 16,	10.8	78
339	Satisfaction survey among medical staff involved in relief operations following the Great East Japan Earthquake and Tsunami. <i>Critical Care</i> , 2012 , 16,	10.8	78
338	Nuclear disaster and the medical problems during the earthquake in Japan, 2011. <i>Critical Care</i> , 2012 , 16,	10.8	78
337	Stressors in the ICU: different perceptions of patients, relatives and staff members. <i>Critical Care</i> , 2012 , 16,	10.8	78
336	Role of ICU nurses in the confrontation of post-traumatic stress disorder in relatives of ICU patients in a general hospital of Athens, Greece. <i>Critical Care</i> , 2012 , 16,	10.8	78
335	Family meetings and end-of-life decision-making in Thai critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
334	Influence of burnout on attitudes of ICU doctors and nurses towards liberalization of visiting polices. <i>Critical Care</i> , 2012 , 16,	10.8	78
333	Prevalence, risk factors and impact of severe burnout syndrome in 12 Uruguayan ICUs. <i>Critical Care</i> , 2012 , 16,	10.8	78
332	Opening the ICU: views of ICU doctors and nurses before and after liberalization of visiting policies. <i>Critical Care</i> , 2012 , 16,	10.8	3
331	A family-based satisfaction survey on the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	1

330	Immediate needs and level of anxiety of families with traumatic brain injury patients admitted to ICUs. <i>Critical Care</i> , 2012 , 16,	10.8	78
329	Families: the newest members of the ICU multidisciplinary team. Critical Care, 2012, 16,	10.8	78
328	Family satisfaction in an interdisciplinary ICU: a quality audit. Critical Care, 2012, 16,	10.8	78
327	Incidence of post-traumatic stress, anxiety and depression symptoms in patients and relatives during the ICU stay and after discharge. <i>Critical Care</i> , 2012 , 16,	10.8	3
326	Application of a new German law as a basis for end-of life decisions in a medical ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
325	Effect of a full moon on mortality of patients admitted to the ICU. Critical Care, 2012, 16,	10.8	78
324	Potential association of gender with mortality and withdrawal of life-sustaining therapies in patients with severe TBI: a Canadian multicentre cohort study. <i>Critical Care</i> , 2012 , 16,	10.8	78
323	Making drug delivery in the ICU safer: the implementation of advanced computerised intravenous infusion pumps. <i>Critical Care</i> , 2012 , 16,	10.8	1
322	Growing a positive culture in an ICU antimicrobial stewardship program. Critical Care, 2012, 16,	10.8	78
321	Injectional anthrax: the inflammatory response. Critical Care, 2012, 16,	10.8	78
320	Multicenter consensus development of a checklist for lung injury prevention. Critical Care, 2012, 16,	10.8	1
319	Impact of the Paediatric Intensive Care Outreach Network service on mortality within intensive care. <i>Critical Care</i> , 2012 , 16,	10.8	78
318	In-hospital rapid response system: effects on outcome and workload. Critical Care, 2012, 16,	10.8	1
317	Medical emergency team admittance to intensive care versus conventional admittance: characteristics and outcome. <i>Critical Care</i> , 2012 , 16,	10.8	78
316	Factors affecting critical care admission to a UK university hospital. Critical Care, 2012, 16,	10.8	1
315	Intensive care services in Hungary 2000 to 2010: an analysis of bed numbers, occupancy rates, case mix and economics. <i>Critical Care</i> , 2012 , 16,	10.8	78
314	Data acquisition for the UK Critical Care Minimum Data Set: validation of a computer model for automatic calculation from an electronic patient record. <i>Critical Care</i> , 2012 , 16,	10.8	78

312	Out-of-hours discharge from the ICU: defining the out-of-hours period and its effect on mortality. <i>Critical Care</i> , 2012 , 16,	10.8	3
311	Delayed discharges revisited: impact of a liaison post on patients' transition from ICU to ward care. <i>Critical Care</i> , 2012 , 16,	10.8	78
310	Assessing demand for intensive care services: the role of readmission rates. Critical Care, 2012, 16,	10.8	78
309	Management of cardiac drugs in a critical care setting. <i>Critical Care</i> , 2012 , 16,	10.8	78
308	Pharmacists and fastidiousness improve compliance with guidelines for stress ulcer prophylaxis. <i>Critical Care</i> , 2012 , 16,	10.8	78
307	Healthcare workers' experience when using an electronic medical order entry and bar-code technology in an ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
306	Safer ICU trainee handover: a service improvement project. Critical Care, 2012, 16,	10.8	1
305	ICU handover: are we forgetting something? A preliminary study. Critical Care, 2012, 16,	10.8	1
304	Quality and value of intensive care discharge summaries for general practitioners. <i>Critical Care</i> , 2012 , 16,	10.8	3
303	Volume-outcome relationship in critical care: a systematic review. <i>Critical Care</i> , 2012 , 16,	10.8	78
302	Radiation doses in young ICU patients: a cause for concern?. Critical Care, 2012, 16,	10.8	1
301	Accuracy of height and weight estimation by critical care staff. Critical Care, 2012, 16,	10.8	78
300	Implementation of evidence-based care bundles in the ICU. Critical Care, 2012, 16,	10.8	1
299	Awareness of difficult airway equipment on the ICU. Critical Care, 2012, 16,	10.8	78
298	A new patient mobilization scoring system in the ICU: what is the degree of similarity in scores between assessors in daily use?. <i>Critical Care</i> , 2012 , 16,	10.8	78
297	Motor and respiratory intensive rehabilitation in bedridden patients. Critical Care, 2012, 16,	10.8	1
296	Severity of electrophysiological alterations correlates with severity of illness in the early phase of critical illness polyneuropathy. <i>Critical Care</i> , 2012 , 16,	10.8	78
295	Muscle strength assessment of critically ill patients is associated with functional ability and quality of life at hospital discharge. <i>Critical Care</i> , 2012 , 16,	10.8	1

294	Functional dependency in the direct post-ICU phase in patients with prolonged mechanical ventilation. <i>Critical Care</i> , 2012 , 16,	10.8	78
293	IL-17A rs1974226 GG genotype is associated with increased susceptibility to Gram-positive infection and mortality of severe sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	78
292	Prevalence of TLR4 single nucleotide polymorphisms (ASP299GLY, THR399ILE) in healthy subjects and septic patients, and association with outcome. <i>Critical Care</i> , 2012 , 16,	10.8	78
291	Modelling immune responses in sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	78
29 0	Decreased peripheral CD4+/CD8+ lymphocytes and poor prognosis in aged sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	78
289	Homeostatic pulmonary microenvironment is responsible for alveolar macrophages resistance to endotoxin tolerance. <i>Critical Care</i> , 2012 , 16,	10.8	78
288	In vivo natural killer and natural killer T-cell depletion affects mortality in a murine pneumococcal pneumonia sepsis model. <i>Critical Care</i> , 2012 , 16,	10.8	78
287	Mobilization of hematopoietic and nonhematopoietic stem cell subpopulations in sepsis: a preliminary report. <i>Critical Care</i> , 2012 , 16,	10.8	1
286	Blunted IL-17 responses early after advent of multiple injuries. <i>Critical Care</i> , 2012 , 16,	10.8	78
285	Apoptosis of neutrophils, expression of TREM-1 on neutrophils and IL-17 responses in experimental burn in injury are related to the type and time of burn exposure. <i>Critical Care</i> , 2012 , 16,	10.8	78
284	Insufficient autophagy relates to mitochondrial dysfunction, organ failure and adverse outcome in an animal model of critical illness. <i>Critical Care</i> , 2012 , 16,	10.8	1
283	Modulation of mediators derived from whole blood or monocytic cells stimulated with lipopolysaccharide reduces endothelial cell activation. <i>Critical Care</i> , 2012 , 16,	10.8	78
282	A/H1N1 infection: immunological parameters in ICU patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
281	Time of course CD64, a leukocyte activation marker, during extracorporeal circulation. <i>Critical Care</i> , 2012 , 16,	10.8	78
280	Oral neutrophil quantitation in patients undergoing elective cardiopulmonary bypass. <i>Critical Care</i> , 2012 , 16,	10.8	78
279	C13-pyruvate administration revealed differential metabolism between heart, liver and red blood cells and improved heart function during endotoxemia. <i>Critical Care</i> , 2012 , 16,	10.8	78
278	AMP-activated protein kinase controls liposaccharide-induced hyperpermeability. <i>Critical Care</i> , 2012 , 16,	10.8	2
277	Reduced expression of PPAR-Mimits the potential beneficial effects of GW0742 during septic shock in atherosclerotic swine. <i>Critical Care</i> , 2012 , 16,	10.8	78

276	Effects of hexafluoro-2-propanol on inflammatory and hemodynamic responses in a rat model of endotoxic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
275	Effects of noradrenaline and lipopolysaccharide exposure on mitochondrial respiration in alveolar macrophages. <i>Critical Care</i> , 2012 , 16,	10.8	78
274	Effects of the anti-diabetic imeglimin in hyperglycemic mice with septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	1
273	Adrenomedullin blockade improves catecholamine responsiveness and kidney function in resuscitated murine septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
272	Activated protein C, severe sepsis and 28-day mortality. <i>Critical Care</i> , 2012 , 16,	10.8	78
271	Soluble usokinase plasminogen activator receptor as a useful biomarker to define advent of sepsis in patients with multiple injuries. <i>Critical Care</i> , 2012 , 16,	10.8	78
270	Role of serum biomarkers in the diagnosis of infection in patients undergoing extracorporeal membrane oxygenation. <i>Critical Care</i> , 2012 , 16,	10.8	1
269	Correlation of VAP diagnosis with parameters of critically ill patients in a general ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
268	Usefulness of daily monitoring of procalcitonin and C-reactive protein in the early diagnosis of infection after elective colonic surgery. <i>Critical Care</i> , 2012 , 16,	10.8	1
267	Procalcitonin as a predictive marker for PCR test and blood culture results in suspected invasive candidemia. <i>Critical Care</i> , 2012 , 16,	10.8	78
266	Would procalcitonin measurement aid antimicrobial stewardship in a UK district general hospital mixed adult critical care population?. <i>Critical Care</i> , 2012 , 16,	10.8	78
265	Procalcitonin has a poor prognosis value in critically ill patients with candidemia. <i>Critical Care</i> , 2012 , 16,	10.8	78
264	Assessment of the usefulness of presepsin (soluble CD14 subtype) in septic patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
263	Circulating cell-free DNA levels measured by a novel simple fluorescent assay are predictive for outcome of severe sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	78
262	Clinical usefulness of measuring endotoxin activity on ICU admission. Critical Care, 2012, 16,	10.8	78
261	Prognostic value of serum galactomannan in mixed ICU patients: a retrospective observational study. <i>Critical Care</i> , 2012 , 16,	10.8	78
260	Analysis of (1-ȝ) 원-glucan as a diagnostic adjunct for invasive fungal infections in the ICU setting. <i>Critical Care</i> , 2012 , 16,	10.8	78
259	Impaired heart rate variability predicts clinical deterioration and progressive organ failure in emergency department sepsis patients. <i>Critical Care</i> , 2012 , 16,	10.8	4

258	Severe community-acquired pneumonia: risk factors for in-hospital mortality. Critical Care, 2012, 16,	10.8	1
257	Systemic corticosteroids for community-acquired pneumonia in adults. <i>Critical Care</i> , 2012 , 16,	10.8	78
256	Characteristics of leptospirosis patients admitted to a tropical university hospital during the 2000 to 2010 period. <i>Critical Care</i> , 2012 , 16,	10.8	1
255	Prognostic impact of imported and newly-isolated methicillin-resistant Staphylococcus aureus in the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
254	Necrotizing pneumonia due to methicillin-sensitive Staphylococcus aureus secreting Panton-Valentine leukocidin: a review of case reports. <i>Critical Care</i> , 2012 , 16,	10.8	78
253	Predicting methicillin-resistant Staphylococcus aureus in critically ill patients with pneumonia presenting to the hospital. <i>Critical Care</i> , 2012 , 16,	10.8	78
252	Predictors of multidrug-resistant Acinetobacter baumannii infections: a retrospective analysis in surgical ICU patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
251	Risk factors for bronchial acquisition of resistant Gram-negative bacteria in critically ill patients and outcome. <i>Critical Care</i> , 2012 , 16,	10.8	78
250	Improved antibiotic stewardship resulting from a multifaceted strategy implemented after an outbreak of multiresistant Acinetobacter baumannii in a university ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
249	Empirical antifungal treatment in the critically ill patients: how does it impact on the outcome?. <i>Critical Care</i> , 2012 , 16,	10.8	78
248	Relationship between polyclonal immunoglobulin therapy and colonization by Candida spp <i>Critical Care</i> , 2012 , 16,	10.8	78
247	Predictive and prognostic factors of septic shock of nosocomial origin. Critical Care, 2012, 16,	10.8	1
246	Catheter-related bloodstream infection: factors affecting incidence. Critical Care, 2012, 16,	10.8	78
245	Prognostic factors of septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
244	Severe sepsis in the United Sates: a 5-year analysis. <i>Critical Care</i> , 2012 , 16,	10.8	78
243	District hospital experience of organ support requirements for H1N1-associated pneumonia. <i>Critical Care</i> , 2012 , 16,	10.8	78
242	Compliance with the sepsis resuscitation bundle in patients with severe sepsis and septic shock admitted to Scottish ICUs. <i>Critical Care</i> , 2012 , 16,	10.8	78
241	Improving early administration of antibiotics: a 'Plan Do Study Act' approach. <i>Critical Care</i> , 2012 , 16,	10.8	78

240	Source-directed antimicrobials: a shot in the dark?. Critical Care, 2012, 16,	10.8	78
239	Relation between temperature in the initial 24 hours in patients with severe sepsis or septic shock with mortality and length of stay in the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
238	Temperature management for patients without brain injury in Australia and New Zealand ICUs: a point prevalence study. <i>Critical Care</i> , 2012 , 16,	10.8	78
237	Impact of antifungal treatment in ICU patients with Candida colonization: analysis of the EPIC II study population. <i>Critical Care</i> , 2012 , 16,	10.8	78
236	Pharmacokinetics of micafungin in patients with severe burn injuries. Critical Care, 2012, 16,	10.8	78
235	Bacteremia affects the mortality of septic patients with high serum procalcitonin level in the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
234	Cases of tetanus after the Japan crisis 2011. <i>Critical Care</i> , 2012 , 16,	10.8	1
233	Does the day of the week predict the presence of microbiologically confirmed ventilator-associated pneumonia?. <i>Critical Care</i> , 2012 , 16,	10.8	78
232	Assessing perforation of acute appendicitis using the delta neutrophil index reflecting the peripheral immature granulocyte count. <i>Critical Care</i> , 2012 , 16,	10.8	78
231	Effects of a multifaceted quality improvement intervention in reducing mortality and bloodstream infection in ICUs: insights from the QUALITI initiative. <i>Critical Care</i> , 2012 , 16,	10.8	78
230	Ceftazidime dosage regimen recommendations in burn patients based on a Monolix population pharmacokinetic study. <i>Critical Care</i> , 2012 , 16,	10.8	78
229	Continuous versus intermittent vancomycin in children after cardiac surgery with delayed sternal closure. <i>Critical Care</i> , 2012 , 16,	10.8	1
228	Elimination of linezolid in patients undergoing low-flow continuous venovenous haemodiafiltration. <i>Critical Care</i> , 2012 , 16,	10.8	1
227	A post-authorisation survey to analyse the perioperative teicoplanin plasma concentrations in adult patients with chronic bone sepsis, who received loading doses of 12 mg/kg 12-hourly for 48 hours followed by 12 mg/kg once daily. <i>Critical Care</i> , 2012 , 16,	10.8	1
226	Pharmacokinetics of inhaled colistin in critically ill patients with ventilator-associated tracheobronchitis. <i>Critical Care</i> , 2012 , 16,	10.8	78
225	Efficacy of inhaled tobramycin in severe nosocomial pneumonia. Critical Care, 2012, 16,	10.8	1
224	Comparison of a bronchoscopic microsample probe with bronchoalveolar lavage to measure cytokine levels in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
223	Clinical and epidemiological risk factors for ventilator-associated pneumonia in a cohort of critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	1

222	Use of a ventilator-associated pneumonia (VAP) bundle to decrease the VAP rate in Syria. <i>Critical Care</i> , 2012 , 16,	10.8	78
221	A strategy for prevention and control of catheter-related bloodstream infection of ICU patients in China (Prevent CRBSI): a prospective, multicenter, controlled study. <i>Critical Care</i> , 2012 , 16,	10.8	1
220	Wash your hands: simple measures save lives. <i>Critical Care</i> , 2012 , 16,	10.8	78
219	Comparison of hand hygiene in single-room versus open-plan ICUs. Critical Care, 2012, 16,	10.8	78
218	Compliance for decontamination of bedside computer keyboards on an ICU. Critical Care, 2012, 16,	10.8	78
217	Reduced air contamination in an ICU environment with a portable air purification system. <i>Critical Care</i> , 2012 , 16,	10.8	78
216	Effectiveness of an innovative system for the bio-decontamination of the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
215	Massive hemoptysis in a respiratory ICU: causes, interventions and outcomes - Indian study. <i>Critical Care</i> , 2012 , 16,	10.8	1
214	Capnography use in Scottish ICUs. <i>Critical Care</i> , 2012 , 16,	10.8	78
213	Digitalized acoustic monitoring of lung congestion. <i>Critical Care</i> , 2012 , 16,	10.8	78
212	Usefulness of electrical activity of the diaphragm to detect intrinsic positive end-expiratory pressure during pressure support ventilation. <i>Critical Care</i> , 2012 , 16,	10.8	78
211	Adequate lung sliding identification is not influenced by the level of academic or ultrasound training. <i>Critical Care</i> , 2012 , 16,	10.8	78
210	Lung ultrasound can differentiate Pneumocystis jiroveci versus other etiologies among critically ill AIDS patients with pneumonia. <i>Critical Care</i> , 2012 , 16,	10.8	5
209	Difference in accuracy of lung sliding identification between the right and left hemithorax. <i>Critical Care</i> , 2012 , 16,	10.8	78
208	Trans-thoracic echo evaluation before and during noninvasive ventilation. Critical Care, 2012, 16,	10.8	1
207	Listen to PaO2/FiO2 ratios: they tell us about length of stay. <i>Critical Care</i> , 2012 , 16,	10.8	78
206	Worst Oxygenation Index during the first 24 hours of ventilation predicts mortality. <i>Critical Care</i> , 2012 , 16,	10.8	78
	The Oxygenation Index compared with the P/F ratio in ALI/ARDS. <i>Critical Care</i> , 2012 , 16,	10.8	

204	Do actual tidal volumes differ from prescribed tidal volumes?. Critical Care, 2012, 16,	10.8	78
203	Intratracheal administration of siRNA targeting FAS reduces ischemia-reperfusion-induced lung injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
202	Acute lung injury in mice associates with p44/42 and c-Jun N-terminal kinase activation and requires the function of TNFI eceptor I. <i>Critical Care</i> , 2012 , 16,	10.8	78
201	Retrieval of patients with severe respiratory failure on venovenous extracorporeal membrane oxygenation: an intensivist-led model. <i>Critical Care</i> , 2012 , 16,	10.8	78
200	A new miniaturized extracorporeal membrane oxygenator with integrated rotary blood pump (Ilias): first results in a porcine model of lung injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
199	ECMO in nonintubated patients as a bridge to lung transplant: our experience. <i>Critical Care</i> , 2012 , 16,	10.8	78
198	Extracorporeal life support in major trauma: case series from a tertiary referral trauma. <i>Critical Care</i> , 2012 , 16,	10.8	78
197	Two years' experience with bicaval dual lumen cannula for venovenous extracorporeal membrane oxygenation in adult refractory acute respiratory distress syndrome. <i>Critical Care</i> , 2012 , 16,	10.8	1
196	Resolution of organ functional scores to predict outcomes in severe acute respiratory distress syndrome patients receiving extracorporeal membrane oxygenation. <i>Critical Care</i> , 2012 , 16,	10.8	1
195	Corticosteroids for critically ill patients: an international survey of intensivists. <i>Critical Care</i> , 2012 , 16,	10.8	78
194	Effects of salbutamol on airway characteristics in mechanically ventilated adults without COPD. <i>Critical Care</i> , 2012 , 16,	10.8	78
193	Respiratory system elastance monitoring during PEEP titration. Critical Care, 2012, 16,	10.8	2
192	Accuracy of the pressure-volume curve method compared to quantitative lung CT scan to assess the recruitable lung in patients with acute respiratory failure. <i>Critical Care</i> , 2012 , 16,	10.8	78
191	Flow-controlled expiration discloses PEEP-dependent dynamic hysteresis of the pressure-volume loop. <i>Critical Care</i> , 2012 , 16,	10.8	78
190	Feasibility of early spontaneous breathing in acute respiratory distress syndrome. <i>Critical Care</i> , 2012 , 16,	10.8	78
189	A device for ventilation-analogue mechanostimulation in vitro. Critical Care, 2012, 16,	10.8	78
188	Hemodynamics effects of recruitment maneuver. Critical Care, 2012, 16,	10.8	2
187	Differential pulmonary and circulatory effects of preventive lung protective ventilation in an experimental postoperative sepsis model. <i>Critical Care</i> , 2012 , 16,	10.8	78

186	High NT-proBNP level is correlated with high PEEP, low PH and low PaO2/FiO2 in ARDS. <i>Critical Care</i> , 2012 , 16,	10.8	78
185	Early application of high-frequency oscillatory ventilation in H1N1 influenza-related severe ARDS is associated with better outcome. <i>Critical Care</i> , 2012 , 16,	10.8	78
184	Outcomes of early delivery in pregnant patients with acute respiratory distress syndrome. <i>Critical Care</i> , 2012 , 16,	10.8	3
183	Mechanical ventilation demographics between 1999 and 2009. Critical Care, 2012, 16,	10.8	78
182	Mechanical ventilation in intensive and critical care units of Russia: RuVent national epidemiologic study. <i>Critical Care</i> , 2012 , 16,	10.8	3
181	Use of a fully closed-loop ventilation mode in long-term ventilated ICU patients: a prospective study. <i>Critical Care</i> , 2012 , 16,	10.8	78
180	Effects of low and high tidal volume and pentoxifylline on intestinal blood flow and leukocyte-endothelial interactions in mechanically ventilated rats. <i>Critical Care</i> , 2012 , 16,	10.8	78
179	A method for continuous noninvasive assessment of respiratory mechanics during spontaneous breathing. <i>Critical Care</i> , 2012 , 16,	10.8	78
178	Influence of catheter diameter, endotracheal tube diameter, suction pressure, and PEEP on the tracheal pressure and lung volume during endotracheal suctioning using a lung model. <i>Critical Care</i> , 2012 , 16,	10.8	78
177	Risk factors of mortality in severe cutaneous adverse reactions patients with pulmonary involvement. <i>Critical Care</i> , 2012 , 16,	10.8	78
176	During spontaneous breathing cardiac output lacks major effect on pulmonary shunting in porcine lungs with partial collapse. <i>Critical Care</i> , 2012 , 16,	10.8	78
175	Oxygenation correlates with lung aeration during unsupported spontaneous breathing in porcine lung collapse model. <i>Critical Care</i> , 2012 , 16,	10.8	78
174	Safety and effect of intermittent intrapulmonary percussive ventilation on oxygen saturation and hemodynamic functions. <i>Critical Care</i> , 2012 , 16,	10.8	78
173	Impact of an open lung approach on hemodynamic parameters after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
172	A protective-ventilation strategy reduces pulmonary complications after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
171	A rule for predicting the new equilibrated carbon dioxide partial pressure after changes in the ventilation frequency. <i>Critical Care</i> , 2012 , 16,	10.8	78
170	Patient-ventilator asynchrony during conventional or automated pressure support ventilation in difficult-to-wean patients. <i>Critical Care</i> , 2012 , 16,	10.8	2
169	High levels of B-type natriuretic peptide predict weaning failure from mechanical ventilation in adult patients after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78

168	Case-control study of failed extubation. <i>Critical Care</i> , 2012 , 16,	10.8	2
167	Out-of-bed extubation: changing paradigms. <i>Critical Care</i> , 2012 , 16,	10.8	78
166	Prediction of post-extubation failure by portable ICU ultrasound. Critical Care, 2012, 16,	10.8	78
165	Intermittent aspiration of pharyngeal secretion for re-intubation prevention. Critical Care, 2012, 16,	10.8	78
164	Efficacy of biphasic cuirass ventilation in the critical care department. Critical Care, 2012, 16,	10.8	1
163	Differences in neurophysiologic effects between CPAP and a novel high-flow therapy system. <i>Critical Care</i> , 2012 , 16,	10.8	78
162	Management of acute bronchospasm respiratory distress with CPAP ventilation associated with nebulization in the prehospital emergency setting. <i>Critical Care</i> , 2012 , 16,	10.8	78
161	Difference between continuous positive airway pressure via mask therapy plus chest physiotherapy (CPT) and incentive spirometry therapy plus CPT to treat or prevent acute atelectasis after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
160	Nasal high-flow oxygen in patients with hypoxic respiratory failure: effect on functional and subjective respiratory parameters compared to conventional oxygen therapy and noninvasive ventilation. <i>Critical Care</i> , 2012 , 16,	10.8	2
159	Short-term effect of humidified high nasal flow oxygen in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
158	Good response on high nasal oxygen flow reduces the need for intubation in adult respiratory failure. <i>Critical Care</i> , 2012 , 16,	10.8	78
157	An audit of airway complications in a district general hospital ICU. Critical Care, 2012, 16,	10.8	78
156	Multidisciplinary care for patients with tracheostomy shortened time to decannulation. <i>Critical Care</i> , 2012 , 16,	10.8	78
155	Influence of percutaneous tracheostomy on gas exchange in mechanically ventilated patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
154	Bronchoscope-guided percutaneous dilatational tracheostomy performed by an experienced intensivist: a 26-month experience at a tertiary care center in United Arab Emirates. <i>Critical Care</i> , 2012 , 16,	10.8	78
153	Risk factors for poor outcome in patients with osmotic demyelination syndrome. <i>Critical Care</i> , 2012 , 16,	10.8	3
152	Is inappropriate secretion of anti-diuretic hormone (SIADH) the cause of hyponatremia in Legionella pneumonia?. <i>Critical Care</i> , 2012 , 16,	10.8	78
151	Fluctuations in serum sodium level are associated with an increased risk of death in surgical ICU patients. <i>Critical Care</i> , 2012 , 16,	10.8	78

150	Impact of ketogenesis and strong ion difference on acid-base in our CICU. Critical Care, 2012, 16,	10.8	78
149	Buffer therapy in metabolic acidosis after surgery-associated hemorrhage in pediatric oncology. <i>Critical Care</i> , 2012 , 16,	10.8	78
148	An unusual cause of high anion gap metabolic acidosis: pyroglutamic acidosis. <i>Critical Care</i> , 2012 , 16,	10.8	0
147	Metabolic acid-base disturbances in patients in the emergency department. Critical Care, 2012, 16,	10.8	78
146	Aberrant bone metabolism in critical illness. <i>Critical Care</i> , 2012 , 16,	10.8	78
145	Low serum 25-hydroxyvitamin D levels and acute kidney injury in the critically ill. <i>Critical Care</i> , 2012 , 16,	10.8	78
144	Concentration of major vitamins in critically ill patients. Critical Care, 2012, 16,	10.8	78
143	No association between vitamin D deficiency at admission and outcome in a medical ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
142	Plasma levels of Coenzyme Q10 are reduced in critically ill patients as compared to healthy volunteers and correlate with age. <i>Critical Care</i> , 2012 , 16,	10.8	3
141	Reduced cortisol metabolism drives hypercortisolism in critical illness. <i>Critical Care</i> , 2012 , 16,	10.8	78
140	Effect of low-dose hydrocortisone on the expression of glucocorticoid receptor alpha of the septic kidney in rats and its protective effect on kidney injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
139	Hydrocortisone increases the risk of dysglycemia in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
138	Nutritional status of patients occupying ICUs in the state of Rio de Janeiro. Critical Care, 2012, 16,	10.8	78
137	Investigating diarrhoea on the ICU: a retrospective study. Critical Care, 2012, 16,	10.8	78
136	Preliminary report of surface electrogastrography in critically ill septic patients after resuscitation. <i>Critical Care</i> , 2012 , 16,	10.8	78
135	Frequency, determinants and impact of feed intolerance amongst the critically ill. <i>Critical Care</i> , 2012 , 16,	10.8	78
134	Gastric emptying assessment in critically ill patients with feed intolerance; comparison of 13C octanoic acid, paracetamol and 3-O-methylglucose absorption tests. <i>Critical Care</i> , 2012 , 16,	10.8	78
133	Early enteral nutrition in the critically ill: a single-centre study. Critical Care, 2012, 16,	10.8	78

132	Preliminary experience with ketone-targeted treatment of diabetic ketoacidosis. <i>Critical Care</i> , 2012 , 16,	10.8	78
131	Impact of blood glucose on blood lactate levels in a medical ICU: a retrospective cohort study. <i>Critical Care</i> , 2012 , 16,	10.8	5
130	Relationship between glycemic Lability Index, infections and outcome in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	Ο
129	Variability of insulin sensitivity during the first 4 days of critical illness. Critical Care, 2012, 16,	10.8	4
128	Endogenous insulin secretion in critically ill patients. Critical Care, 2012, 16,	10.8	78
127	Impact of the type of glucose monitoring on the assessment of glycemic variability in critical care patients. <i>Critical Care</i> , 2012 , 16,	10.8	2
126	Evaluation of a continuous blood glucose monitoring system using a central venous catheter with an integrated microdialysis function. <i>Critical Care</i> , 2012 , 16,	10.8	78
125	Variable and maximum blood glucose levels during the first week after ICU admission are related to the severity of the patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
124	Pilot trial of STAR in the medical ICU. Critical Care, 2012, 16,	10.8	78
123	Glucometer accuracy and implications for clinical studies. Critical Care, 2012, 16,	10.8	1
122	Initial experience with continuous intra-arterial fluorescent glucose monitoring in patients in the ICU following cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
121	Preliminary ICU experience of a novel intravascular blood glucose sensor. Critical Care, 2012, 16,	10.8	3
120	Does tight glycemic control positively impact on patient mortality?. Critical Care, 2012, 16,	10.8	78
119	Glycaemic control in ICUs in large English hospitals: a follow-up telephone survey. <i>Critical Care</i> , 2012 , 16,	10.8	78
118	Model-based regulation of glucose in critical care. <i>Critical Care</i> , 2012 , 16,	10.8	78
117	Perioperative glycemic control with a computerized algorithm versus conventional glycemic control. <i>Critical Care</i> , 2012 , 16,	10.8	78
116	Efficacy of the novel heart attack centre extension pathway: a pilot study. Critical Care, 2012, 16,	10.8	78
115	Integral assistance process implantation for ST-elevated acute coronary syndrome. <i>Critical Care</i> , 2012 , 16,	10.8	78

114	Prognostic value of Killip classification in terms of health-related quality of life. <i>Critical Care</i> , 2012 , 16,	10.8	78
113	Modes of arrival, door to balloon time and its impact on morbidity and mortality for ST elevation myocardial infarct. <i>Critical Care</i> , 2012 , 16,	10.8	3
112	Next-generation, fast and accurate point-of-care test for NT-proBNP based on Magnotech technology. <i>Critical Care</i> , 2012 , 16,	10.8	78
111	Right ventricular apical versus septal pacing: impact on left ventricular synchrony and function. <i>Critical Care</i> , 2012 , 16,	10.8	78
110	Consecutive case series of Takotsubo cardiomyopathy: a disease potentially triggered by the Great East Japan Earthquake. <i>Critical Care</i> , 2012 , 16,	10.8	78
109	Stress cardiomyopathy after live donor liver transplantation: incidence, risk factors and mortality. <i>Critical Care</i> , 2012 , 16,	10.8	78
108	Short-term hemodynamic effects of nebivolol in acute decompensated heart failure: a randomized clinical trial. <i>Critical Care</i> , 2012 , 16,	10.8	78
107	Patients with infective endocarditis patients in the ICU: how are they?. Critical Care, 2012, 16,	10.8	78
106	Malperfusion and branch compromise in acute type A aortic syndrome. Critical Care, 2012, 16,	10.8	78
105	Prognostic value of the echocardiographic-derived calcium index in coronary artery disease. <i>Critical Care</i> , 2012 , 16,	10.8	78
104	Echocardiography in the ICU: an audit of 3 years practice. Critical Care, 2012, 16,	10.8	1
103	Left ventriculum diastolic dysfunction in pediatric septic shock. Critical Care, 2012, 16,	10.8	78
102	Existence of interference between the heart and respiratory sounds: preliminary report. <i>Critical Care</i> , 2012 , 16,	10.8	78
101	Effects of the intravenous administration of purine nucleosides guanosine or inosine against hemorrhagic shock in pigs. <i>Critical Care</i> , 2012 , 16,	10.8	78
100	Norepinephrine versus angiotensin II in septic shock: effects on isolated kidney, heart and liver mitochondrial respiration. <i>Critical Care</i> , 2012 , 16,	10.8	78
99	Goal-directed hemodynamic resuscitation in high-risk patients undergoing cardiac surgery: a randomized controlled trial - preliminary data (GRICCS STUDY). <i>Critical Care</i> , 2012 , 16,	10.8	1
98	Economic evaluation of early-goal directed therapy for high-risk surgical patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
97	What matters during a hypotensive episode: fluids, vasopressors, or both?. <i>Critical Care</i> , 2012 , 16,	10.8	78

96	Elevated central venous pressure in septic patients is associated with impairment of microcirculatory blood flow. <i>Critical Care</i> , 2012 , 16,	10.8	78
95	Human protein C concentrate to restore physiological values in adult septic shock patients: effects on microcirculation. <i>Critical Care</i> , 2012 , 16,	10.8	78
94	Heart rate reduction with esmolol in septic shock: effects on microcirculation. <i>Critical Care</i> , 2012 , 16,	10.8	78
93	Microcirculation and blood transfusion: effects of three different types of concentrated red blood cells - preliminary results. <i>Critical Care</i> , 2012 , 16,	10.8	2
92	Peripheral perfusion is correlated to metabolic perfusion parameters and microvascular reactivity but not with hepatosplanchnic or microcirculatory flow parameters in hyperdynamic septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
91	Hyperoxia affects peripheral tissue microcirculation in patients with pulmonary arterial hypertension. <i>Critical Care</i> , 2012 , 16,	10.8	2
90	Supraclavicular ultrasound-guided subclavian vein cannulation in infants under 5 kg. <i>Critical Care</i> , 2012 , 16,	10.8	78
89	Central venous catheter placement: where is the end of the line?. Critical Care, 2012, 16,	10.8	78
88	Power-injectable peripherally inserted central catheters in intensive care patients. <i>Critical Care</i> , 2012 , 16,	10.8	1
87	Comparison of internal jugular and subclavian access for central venous catheterization in pediatric cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	1
86	Ultrasound-guided central venous line placement in critically ill patients: is chest X-ray needed to assess post-insertion pneumothorax?. <i>Critical Care</i> , 2012 , 16,	10.8	78
85	Errors in the arterial blood pressure measurement. <i>Critical Care</i> , 2012 , 16,	10.8	1
84	The T-Line TL-200 system for continuous noninvasive blood pressure measurement in medical ICU patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
83	Reliability of radial arterial pressure monitoring after cardiac surgery. Critical Care, 2012, 16,	10.8	1
82	Right/left ventricular area ratio does not correlate with right ventricular impedance. <i>Critical Care</i> , 2012 , 16,	10.8	1
81	Impact of prior systemic corticosteroid use in patients admitted with community-acquired pneumonia. 2012 , 6, 323-30		1
80	Role of mannose-binding lectin on pneumococcal infections. Critical Care, 2012, 16,	10.8	78
79	Severe sepsis in community-acquired pneumoniaearly recognition and treatment. 2012 , 23, 412-9		19

78	Abstracts of the 32nd International Symposium on Intensive Care and Emergency Medicine. March 20-23, 2012. Brussels, Belgium. <i>Critical Care</i> , 2012 , 16 Suppl 1, P1-520	10.8	1
77	Relation between chronic obstructive pulmonary disease and antibiotics. <i>Current Infectious Disease Reports</i> , 2012 , 14, 300-7	3.9	6
76	Treatment with anti-inflammatory drugs in community-acquired pneumonia. 2012, 272, 25-35		24
75	Corticosteroid therapy in patients with primary viral pneumonia due to pandemic (H1N1) 2009 influenza. <i>Journal of Infection</i> , 2012 , 64, 311-8	18.9	50
74	Efficacy of methylprednisolone in children with severe community acquired pneumonia. <i>Pediatric Pulmonology</i> , 2013 , 48, 168-75	3.5	20
73	Adjuvant role of corticosteroids in the treatment of community-acquired pneumonia. <i>The Egyptian Journal of Chest Diseases and Tuberculosis</i> , 2013 , 62, 439-445	0.1	38
72	Community-acquired pneumonia in patients with chronic obstructive pulmonary disease requiring admission to the intensive care unit: risk factors for mortality. <i>Journal of Critical Care</i> , 2013 , 28, 975-9	4	21
71	Adjuvant steroid therapy in community-acquired pneumonia: a systematic review and meta-analysis. <i>Journal of Hospital Medicine</i> , 2013 , 8, 68-75	2.7	23
70	What is the best antimicrobial treatment for severe community-acquired pneumonia (including the role of steroids and statins and other immunomodulatory agents). <i>Infectious Disease Clinics of North America</i> , 2013 , 27, 133-47	6.5	13
69	Corticosteroids for pneumonia: are we there yet?. <i>Respirology</i> , 2013 , 18, 199-200	3.6	1
68	Systemic corticosteroids for community-acquired pneumonia: reasons for use and lack of benefit on outcome. <i>Respirology</i> , 2013 , 18, 263-71	3.6	24
67	Severe community-acquired pneumonia. <i>Critical Care Clinics</i> , 2013 , 29, 563-601	4.5	47
66	Glucocorticosteroid in treatment of severe pneumonia. <i>Mediators of Inflammation</i> , 2013 , 2013, 865635	4.3	9
65	Overview of community-acquired pneumonia and the role of inflammatory mechanisms in the immunopathogenesis of severe pneumococcal disease. <i>Mediators of Inflammation</i> , 2013 , 2013, 490346	4.3	63
64	Impact and indication of early systemic corticosteroids for very severe community-acquired pneumonia. <i>International Journal of General Medicine</i> , 2013 , 6, 693-701	2.3	3
63	Recent advances in our understanding of Streptococcus pneumoniae infection. <i>F1000prime Reports</i> , 2014 , 6, 82		35
62	Inhaled corticosteroids and systemic inflammatory response in community-acquired pneumonia: a prospective clinical study. <i>Respirology</i> , 2014 , 19, 929-35	3.6	14
61	Corticosteroid treatment for community-acquired pneumoniathe STEP trial: study protocol for a randomized controlled trial. <i>Trials</i> , 2014 , 15, 257	2.8	14

60	Annual Update in Intensive Care and Emergency Medicine 2014. <i>Annual Update in Intensive Care and Emergency Medicine</i> , 2014 ,	0.2	3
59	Corticosteroid therapy for severe community-acquired pneumonia: a meta-analysis. <i>Respiratory Care</i> , 2014 , 59, 557-63	2.1	29
58	Polmoniti comunitarie gravi delladulto immunocompetente. <i>EMC - Anestesia-Rianimazione</i> , 2015 , 20, 1-10	О	
57	Effect of 23-Valent Pneumococcal Polysaccharide Vaccine Inoculated During Anti-Cancer Treatment Period in Elderly Lung Cancer Patients on Community-Acquired Pneumonia Hospitalization: A Nationwide Population-Based Cohort Study. <i>Medicine (United States)</i> , 2015 , 94, e1022	1.8 2	18
56	Adjunctive Systemic Corticosteroids for Hospitalized Community-Acquired Pneumonia: Systematic Review and Meta-Analysis 2015 Update. <i>Scientific Reports</i> , 2015 , 5, 14061	4.9	52
55	Corticosteroids for all adult patients with community-acquired pneumonia?. <i>Pneumonia (Nathan Qld)</i> , 2015 , 6, 44-47	2.8	
54	Steroids for Community-Acquired Pneumonia. <i>Infectious Diseases in Clinical Practice</i> , 2015 , 23, 279-280	0.2	
53	The Addition of Prednisone to Antibiotics for the Treatment of Community-acquired Pneumonia May Shorten the Time to Clinical Stability. <i>Clinical Pulmonary Medicine</i> , 2015 , 22, 323-324	0.3	
52	To use or not to use corticosteroids for pneumonia? A clinician's perspective. <i>Monaldi Archives for Chest Disease</i> , 2012 , 77, 94-101	2.7	9
51	Etiological Diagnosis of Community-Acquired Pneumonia in Adult Patients: A Prospective Hospital-Based Study in Mashhad, Iran. <i>Jundishapur Journal of Microbiology</i> , 2015 , 8, e22780	1.2	9
50	Systemic steroid treatment for severe expanding pneumococcal pneumonia. <i>Case Reports in Pediatrics</i> , 2015 , 2015, 186302	0.7	4
49	Neumonਬs comunitarias graves del adulto inmunocompetente. <i>EMC - Anestesia-Reanimac</i> 內 , 2015 , 41, 1-11	0.1	
48	Adjunct prednisone therapy for patients with community-acquired pneumonia: a multicentre, double-blind, randomised, placebo-controlled trial. <i>Lancet, The</i> , 2015 , 385, 1511-8	40	244
47	Effect of corticosteroids on treatment failure among hospitalized patients with severe community-acquired pneumonia and high inflammatory response: a randomized clinical trial. <i>JAMA</i> - Journal of the American Medical Association, 2015, 313, 677-86	27.4	306
46	Defining low-dose corticosteroid: the pendulum still oscillates. <i>European Respiratory Journal</i> , 2015 , 46, 574-6	13.6	0
45	Therapeutic strategies in pneumonia: going beyond antibiotics. <i>European Respiratory Review</i> , 2015 , 24, 516-24	9.8	11
44	Community-acquired pneumonia in patients with chronic obstructive pulmonary disease. <i>Current Infectious Disease Reports</i> , 2015 , 17, 444	3.9	10
43	Low-dose corticosteroid use and mortality in severe community-acquired pneumonia patients. <i>European Respiratory Journal</i> , 2015 , 45, 463-72	13.6	64

42	Nonantibiotic Adjunctive Therapies for Community-Acquired Pneumonia (Corticosteroids and Beyond): Where Are We with Them?. <i>Seminars in Respiratory and Critical Care Medicine</i> , 2016 , 37, 913-92	23.9	5
41	[New Pathogenetic Concepts and Pharmacological Studies on Adjuvant Therapy in Severe Pneumonia]. <i>Pneumologie</i> , 2016 , 70, 372-8	0.5	1
40	Bacterial Pneumonia in Older Adults. <i>Clinics in Geriatric Medicine</i> , 2016 , 32, 459-77	3.8	13
39	New aspects in the management of pneumonia. <i>Critical Care</i> , 2016 , 20, 267	10.8	16
38	Pathogen- and antibiotic-specific effects of prednisone in community-acquired pneumonia. European Respiratory Journal, 2016 , 48, 1150-1159	13.6	26
37	Benefit of adjunct corticosteroids for community-acquired pneumonia in diabetic patients. <i>Diabetologia</i> , 2016 , 59, 2552-2560	10.3	12
36	Precision medicine for the treatment of severe pneumonia in intensive care. <i>Expert Review of Respiratory Medicine</i> , 2016 , 10, 297-316	3.8	20
35	Efficacy and Safety of Corticosteroids for Community-Acquired Pneumonia: A Systematic Review and Meta-Analysis. <i>Chest</i> , 2016 , 149, 209-19	5.3	104
34	Has Mortality of Community-Acquired Pneumonia Really Reduced?. <i>Clinical Pulmonary Medicine</i> , 2017 , 24, 258-262	0.3	1
33	Bacteriological incidence in pneumonia patients with pulmonary emphysema: a bacterial floral analysis using the 16S ribosomal RNA gene in bronchoalveolar lavage fluid. <i>International Journal of COPD</i> , 2017 , 12, 2111-2120	3	10
32	Adjuvant therapies in critical care: steroids to treat infectious diseases. <i>Intensive Care Medicine</i> , 2018 , 44, 1306-1309	14.5	4
31	Bronchoalveolar carcinoma as an unsuspected cause for worsening shortness of breath in a patient with metastatic breast cancer. <i>BMJ Case Reports</i> , 2018 , 2018,	0.9	
30	Adjunctive Therapies for Community-Acquired Pneumonia. <i>Clinics in Chest Medicine</i> , 2018 , 39, 753-764	5.3	7
29	Effectiveness of 23-valent pneumococcal polysaccharide vaccine on elderly long-term cancer survivors: a population-based propensity score matched cohort study. <i>BMJ Open</i> , 2018 , 8, e019364	3	7
28	Effects and safety of triamcinolone acetonide-controlled common therapy in keloid treatment: a Bayesian network meta-analysis. <i>Therapeutics and Clinical Risk Management</i> , 2018 , 14, 973-980	2.9	5
27	Community-acquired Pneumonia and Acute Respiratory Distress Syndrome: Prevalence, Risk, and Prognosis. <i>Clinical Pulmonary Medicine</i> , 2018 , 25, 100-106	0.3	1
26	Severe community-acquired pneumonia: current management and future therapeutic alternatives. <i>Expert Review of Anti-Infective Therapy</i> , 2018 , 16, 667-677	5.5	28
25	Corticosteroids in Acute Lung Injury: The Dilemma Continues. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	49

24	Why the Adjunctive Corticosteroid Treatment in Critically Ill Patients With Septic Shock (ADRENAL) Trial Did Not Show a Difference in Mortality. <i>Critical Care Medicine</i> , 2019 , 47, 1785-1788	1.4	11
23	Use of glucocorticoids in the critical care setting: Science and clinical evidence. <i>Pharmacology & Therapeutics</i> , 2020 , 206, 107428	13.9	14
22	Comparative Efficacy and Safety of Common Therapies in Keloids and Hypertrophic Scars: A Systematic Review and Meta-analysis. <i>Aesthetic Plastic Surgery</i> , 2020 , 44, 207-218	2	24
21	Low-Dose Corticosteroid Treatment in Children With Pneumonia: A Retrospective Cohort Study. <i>Frontiers in Pediatrics</i> , 2020 , 8, 566371	3.4	
20	Kortikosteroidi kod telle pneumonije. <i>Infektoloski Glasnik</i> , 2020 , 39, 119-123	0.1	
19	Granulocyte-targeted therapies for airway diseases. <i>Pharmacological Research</i> , 2020 , 157, 104881	10.2	8
18	The role of steroids in severe CAP. Hospital Practice (1995), 2020, 48, 12-22	2.2	1
17	The REMAP-CAP (Randomized Embedded Multifactorial Adaptive Platform for Community-acquired Pneumonia) Study. Rationale and Design. <i>Annals of the American Thoracic Society</i> , 2020 , 17, 879-891	4.7	128
16	Impact of 23-valent pneumococcal polysaccharide vaccination on the frequency of pneumonia-related hospitalization and survival in elderly patients with prostate cancer: A seven-year nationwide matched cohort study. <i>Cancer</i> , 2020 , 127, 124-136	6.4	1
15	Burden of pneumococcal disease among adults in Southern Europe (Spain, Portugal, Italy, and Greece): a systematic review and meta-analysis. <i>Human Vaccines and Immunotherapeutics</i> , 2021 , 17, 36	70 ¹ 368	6 ^O
14	Non-antibiotic therapies for CAP. 2014 , 219-233		1
13	Clinical features, etiology and outcomes of community-acquired pneumonia in patients with chronic obstructive pulmonary disease. <i>PLoS ONE</i> , 2014 , 9, e105854	3.7	33
12	Epidemiology, Seasonality and Treatment of Hospitalized Adults and Adolescents with Influenza in Jingzhou, China, 2010-2012. <i>PLoS ONE</i> , 2016 , 11, e0150713	3.7	10
11	Advances in the prevention, management, and treatment of community-acquired pneumonia. <i>F1000 Medicine Reports</i> , 2010 , 2, 53		4
10	Corticosteroids and ICU Course of Community Acquired Pneumonia in Egyptian Settings. <i>Pharmacology & Pharmacy</i> , 2011 , 02, 73-81	0.3	57
9	Acute lung injury (ALI) and acute respiratory distress syndrome (ARDS). <i>Nihon Kyukyu Igakukai Zasshi</i> , 2010 , 21, 819-827	O	
8	Systemic Corticosteroid Treatment in Severe Community-Acquired Pneumonia Requiring Mechanical Ventilation: Impact on Outcomes and Complications. <i>Tuberculosis and Respiratory Diseases</i> , 2012 , 72, 149	3.2	
7	Breathing Easier: Respiratory Disease in the Older Adult. 43-57		

CITATION REPORT

6 Corticosteroids as Adjunctive Treatment in Community-Acquired Pneumonia. **2014**, 53-63

5	Community Acquired Pneumonia. 2017 , 181-188		
4	Ribociclib-associated organising pneumonia BMJ Case Reports, 2021, 14,	0.9	1
3	Data_Sheet_1.docx. 2020 ,		
2	Exploring the role of antibiotics and steroids in managing respiratory diseases. <i>Journal of Biochemical and Molecular Toxicology</i> ,	3.4	1
1	The effects of systemic corticosteroid on pediatric community-acquired pneumonia: comprehensive review. 2,		O