

Evaluation of protease-activated receptor 2 in murine n

Arthritis and Rheumatism

56, 101-107

DOI: [10.1002/art.22312](https://doi.org/10.1002/art.22312)

Citation Report

#	ARTICLE	IF	CITATIONS
2	The therapeutic potential of proteinase-activated receptors in arthritis. <i>Current Opinion in Pharmacology</i> , 2007, 7, 334-338.	1.7	18
3	Activation of proteinase-activated receptor 2 in human osteoarthritic cartilage upregulates catabolic and proinflammatory pathways capable of inducing cartilage degradation: a basic science study. <i>Arthritis Research and Therapy</i> , 2007, 9, R121.	1.6	61
4	Development of adjuvant monoarthritis is critically dependent on the induction protocol: Comment on the article by Busso et al. <i>Arthritis and Rheumatism</i> , 2007, 56, 3510-3510.	6.7	2
5	Role of protease-activated receptor 2 in joint inflammation. <i>Arthritis and Rheumatism</i> , 2007, 56, 3514-3517.	6.7	7
6	Protease-activated receptor 2 mediates the proinflammatory effects of synovial mast cells. <i>Arthritis and Rheumatism</i> , 2007, 56, 3532-3540.	6.7	51
8	Strain dependence in murine models of monoarthritis. <i>Inflammation Research</i> , 2007, 56, 511-514.	1.6	5
9	Emerging roles of serine proteinases in tissue turnover in arthritis. <i>Arthritis and Rheumatism</i> , 2008, 58, 3644-3656.	6.7	57
10	Role of protease-activated receptors in inflammatory responses, innate and adaptive immunity. <i>Journal of Leukocyte Biology</i> , 2008, 83, 1309-1322.	1.5	166
11	Advances in multiple Sclerosis and Experimental Demyelinating Diseases. <i>Current Topics in Microbiology and Immunology</i> , 2008, , .	0.7	4
12	The Multiple Sclerosis Degradome: Enzymatic Cascades in Development and Progression of Central Nervous System Inflammatory Disease. <i>Current Topics in Microbiology and Immunology</i> , 2008, 318, 133-175.	0.7	36
13	Essential role of platelet activation via protease activated receptor 4 in tissue factor-initiated inflammation. <i>Arthritis Research and Therapy</i> , 2008, 10, R42.	1.6	35
15	Mast Cells Contribute to Autoimmune Inflammatory Arthritis via Their Tryptase/Heparin Complexes. <i>Journal of Immunology</i> , 2009, 182, 647-656.	0.4	153
16	Modulation of OPG, RANK and RANKL by human chondrocytes and their implication during osteoarthritis. <i>Rheumatology</i> , 2009, 48, 1482-1490.	0.9	99
17	Protease-activated receptors as drug targets in inflammation and pain. , 2009, 123, 292-309.		113
18	Triggering of proteinase-activated receptor 4 leads to joint pain and inflammation in mice. <i>Arthritis and Rheumatism</i> , 2009, 60, 728-737.	6.7	66
19	Activation of cartilage matrix metalloproteinases by activated protein C. <i>Arthritis and Rheumatism</i> , 2009, 60, 780-791.	6.7	44
20	Proteinase activated receptor (PAR) involvement in mediating arthritis pain and inflammation. <i>Inflammation Research</i> , 2009, 58, 119-126.	1.6	46
21	Treatment with ephrin B2 positively impacts the abnormal metabolism of human osteoarthritic chondrocytes. <i>Arthritis Research and Therapy</i> , 2009, 11, R119.	1.6	32

#	ARTICLE	IF	CITATIONS
22	Proteinase-activated receptor (PAR)-2 activation impacts bone resorptive properties of human osteoarthritic subchondral bone osteoblasts. <i>Bone</i> , 2009, 44, 1143-1150.	1.4	34
23	Expression of Proteinase-Activated Receptor-2 in the Intervertebral Disc. <i>Spine</i> , 2009, 34, 470-478.	1.0	13
24	Involvement of transient receptor potential vanilloid 1 receptors in protease-activated receptor-2-induced joint inflammation and nociception. <i>European Journal of Pain</i> , 2010, 14, 351-358.	1.4	34
25	Protease-activated receptor 2 signalling promotes dendritic cell antigen transport and T cell activation <i>in vivo</i> . <i>Immunology</i> , 2010, 129, 20-27.	2.0	37
26	Expression and function of the NALP3 inflammasome in rheumatoid synovium. <i>Immunology</i> , 2010, 129, 178-185.	2.0	85
27	Protease-activated Receptor-2 (par ₂) in Human Periodontitis. <i>Journal of Dental Research</i> , 2010, 89, 948-953.	2.5	39
28	Oral treatment with a <i>Brachytemma calycinum</i> D don plant extract reduces disease symptoms and the development of cartilage lesions in experimental dog osteoarthritis: inhibition of protease-activated receptor 2. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, 1179-1184.	0.5	13
29	<i>Porphyrromonas Gingivalis</i> is Associated With Protease-Activated Receptor-2 Upregulation in Chronic Periodontitis. <i>Journal of Periodontology</i> , 2011, 82, 1596-1601.	1.7	27
30	The Role of Coagulation in Chronic Inflammatory Disorders: A Jack of All Trades. <i>Current Pharmaceutical Design</i> , 2011, 17, 9-16.	0.9	20
31	Structure, function and pathophysiology of protease activated receptors. , 2011, 130, 248-282.		315
32	Anti-Inflammatory mechanisms of the proteinase-activated receptor 2-inhibiting peptide in human synovial cells. <i>Journal of Biomedical Science</i> , 2011, 18, 43.	2.6	9
33	Proteinase-Activated Receptors and Arthritis. , 2011, , 217-242.		0
34	Immunomodulatory role of proteinase-activated receptor-2. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 1559-1566.	0.5	21
35	An antagonist of human protease activated receptor-2 attenuates PAR2 signaling, macrophage activation, mast cell degranulation, and collagen-induced arthritis in rats. <i>FASEB Journal</i> , 2012, 26, 2877-2887.	0.2	91
36	15 Kallikrein-related Peptidases (KLKs), Proteinase-mediated Signaling and Proteinase-activated Receptors (PARs). , 2012, , 373-398.		1
37	PAR2 expression in peripheral blood monocytes of patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 1049-1054.	0.5	22
38	Proteinase-activated receptor-2 is required for normal osteoblast and osteoclast differentiation during skeletal growth and repair. <i>Bone</i> , 2012, 50, 704-712.	1.4	25
39	Activation of PAR ₂ receptors sensitizes primary afferents and causes leukocyte rolling and adherence in the rat knee joint. <i>British Journal of Pharmacology</i> , 2012, 167, 1665-1678.	2.7	25

#	ARTICLE	IF	CITATIONS
40	Correlation of protease-activated receptor-2 expression and synovitis in rheumatoid and osteoarthritis. <i>Rheumatology International</i> , 2012, 32, 3077-3086.	1.5	21
41	Interleukin-10 produced by B cells is crucial for the suppression of Th17/Th1 responses, induction of T regulatory type 1 cells and reduction of collagen-induced arthritis. <i>Arthritis Research and Therapy</i> , 2012, 14, R32.	1.6	236
42	Role of transient receptor potential vanilloid 4 in rat joint inflammation. <i>Arthritis and Rheumatism</i> , 2012, 64, 1848-1858.	6.7	39
43	Protease-activated receptor 2, rather than protease-activated receptor 1, contributes to the aggressive properties of synovial fibroblasts in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2012, 64, 88-98.	6.7	32
44	Toward Drugs for Protease-Activated Receptor 2 (PAR2). <i>Journal of Medicinal Chemistry</i> , 2013, 56, 7477-7497.	2.9	68
45	Evolutionary Rate Covariation in Meiotic Proteins Results from Fluctuating Evolutionary Pressure in Yeasts and Mammals. <i>Genetics</i> , 2013, 193, 529-538.	1.2	34
46	Depletion of Protease-Activated Receptor 2 but Not Protease-Activated Receptor 1 May Confer Protection Against Osteoarthritis in Mice Through Extracartilaginous Mechanisms. <i>Arthritis and Rheumatology</i> , 2014, 66, 3337-3348.	2.9	95
47	B Cells Regulate CD4+ T Cell Responses to Papain following B Cell Receptor-Independent Papain Uptake. <i>Journal of Immunology</i> , 2014, 193, 529-539.	0.4	11
48	TRPV4 as a therapeutic target for joint diseases. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2015, 388, 437-450.	1.4	78
49	The Role of Proteases in Pain. <i>Handbook of Experimental Pharmacology</i> , 2015, 227, 239-260.	0.9	15
51	Silencing of protease-activated receptors attenuates synovitis and cartilage damage following a joint bleed in haemophilic mice. <i>Haemophilia</i> , 2016, 22, 152-159.	1.0	7
52	Neutrophil elastase induces inflammation and pain in mouse knee joints via activation of proteinase-activated receptor-2. <i>British Journal of Pharmacology</i> , 2016, 173, 766-777.	2.7	57
53	Proteinase-activated receptor 2 expression on peripheral blood monocytes and T-cells in patients with rheumatoid arthritis. <i>Egyptian Rheumatologist</i> , 2016, 38, 91-98.	0.5	3
54	Proteinases and their receptors in inflammatory arthritis: an overview. <i>Nature Reviews Rheumatology</i> , 2018, 14, 170-180.	3.5	45
55	Attenuation of nociceptive pain and inflammatory disorders by total steroid and terpenoid fraction of <i>Euphorbia tirucalli</i> Linn root in experimental in vitro and in vivo model. <i>Inflammopharmacology</i> , 2018, 26, 235-250.	1.9	12
56	Neutrophils: Novel key players in Rheumatoid Arthritis. Current and future therapeutic targets. <i>Autoimmunity Reviews</i> , 2018, 17, 1138-1149.	2.5	88
57	The phytosphingosine-CD300b interaction promotes zymosan-induced, nitric oxide-dependent neutrophil recruitment. <i>Science Signaling</i> , 2019, 12, .	1.6	12
58	Proteinase-Mediated Macrophage Signaling in Psoriatic Arthritis. <i>Frontiers in Immunology</i> , 2020, 11, 629726.	2.2	8

#	ARTICLE	IF	CITATIONS
59	Protease Activated Receptors and Arthritis. International Journal of Molecular Sciences, 2021, 22, 9352.	1.8	9
60	Down-regulation of protease-activated receptor 2 ameliorated osteoarthritis in rats through regulation of MAPK/NF- κ B signaling pathway <i>in vivo</i> and <i>in vitro</i> . Bioscience Reports, 2020, 40, .	1.1	14
61	Role of protease activated receptor-2 in rheumatoid arthritis: recent progress. Academic Journal of Second Military Medical University, 2010, 29, 725-726.	0.0	0
62	Deficiency of protease-activated receptor (PAR) 1 and PAR2 exacerbates collagen-induced arthritis in mice via differing mechanisms. Rheumatology, 2021, 60, 2990-3003.	0.9	5