

# Barriers to Propofol Use in Emergency Medicine

Annals of Emergency Medicine

52, 392-398

DOI: [10.1016/j.annemergmed.2007.12.002](https://doi.org/10.1016/j.annemergmed.2007.12.002)

Citation Report

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Emergency department procedural sedation practice in Cape Town, South Africa. <i>International Journal of Emergency Medicine</i> , 2009, 2, 91-97.   | 1.6 | 6         |
| 2  | Ketamine and Neurotoxicity: Clinical Perspectives and Implications for Emergency Medicine. <i>Annals of Emergency Medicine</i> , 2009, 54, 181-190.  | 0.6 | 64        |
| 3  | Is it safe to use propofol in the emergency department? A randomized controlled trial to compare propofol and midazolam. <i>International Journal of Emergency Medicine</i> , 2010, 3, 105-113.  | 1.6 | 18        |
| 4  | Professional Skills and Competence for Safe and Effective Procedural Sedation in Children: Recommendations Based on a Systematic Review of the Literature. <i>International Journal of Pediatrics (United Kingdom)</i> , 2010, 2010, 1-16. | 0.8 | 33        |
| 5  | Paediatric procedural sedation based on nitrous oxide and ketamine: sedation registry data from Australia. <i>Emergency Medicine Journal</i> , 2010, 27, 607-612.  | 1.0 | 33        |
| 6  | Reformulation of the Sedation Continuum. <i>JAMA - Journal of the American Medical Association</i> , 2010, 303, 876.   | 7.4 | 34        |
| 7  | Stratification of sedation risk – a challenge to the sedation continuum. <i>Paediatric Anaesthesia</i> , 2011, 21, 924-931.  | 1.1 | 28        |
| 8  | Procedural Sedation and Analgesia in the Emergency Department: Recommendations for Physician Credentialing, Privileging, and Practice. <i>Annals of Emergency Medicine</i> , 2011, 58, 365-370.  | 0.6 | 59        |
| 10 | Who Owns Deep Sedation?. <i>Annals of Emergency Medicine</i> , 2011, 57, 470-474.  | 0.6 | 16        |
| 11 | Safe and Effective Procedural Sedation for Gastrointestinal Endoscopy in Children. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2012, 54, 171-185.   | 1.8 | 94        |
| 12 | Quality of care for joint dislocation in 47 US EDs. <i>American Journal of Emergency Medicine</i> , 2012, 30, 1105-1113.   | 1.6 | 2         |
| 13 | Pediatric Sedation Outside of the Operating Room. , 2012, , .  |     | 10        |
| 14 | Pediatric Sedation with Propofol – Continuing Evolution of Procedural Sedation Practice. <i>Journal of Pediatrics</i> , 2012, 160, 714-716.  | 1.8 | 16        |
| 15 | A window into diaphragm kinetics: feasibility, precision, and physiological meaning of ultrasound measurements of diaphragm thickness. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 3         |
| 17 | Reversible depressive effect of TNF $\alpha$ on a model of isolated perfused rat heart. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 18 | Effect of cdp-choline on microcirculatory alterations during endotoxemia. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 19 | Immune response after stimulation with wall components of Gram-positive bacteria. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 20 | Correlation of the oxygen radical activity and antioxidants and severity in critically ill surgical patients: preliminary report. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 21 | Simultaneous analysis of the expression of CD64 and HLA-DR in the peripheral blood and bronchoalveolar lavage fluid in sepsis. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 22 | Anti-inflammatory effects of Kupffer cells through $\alpha 7$ -nicotinic acetylcholine receptors. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 23 | Prevention of sepsis by correcting apoptosis. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 24 | Immune paralysis in trauma patients; implications for prehospital intervention. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 25 | Is hemoglobin concentration affected by sepsis in the acute phase?. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 4         |
| 26 | Do changes in red blood cell deformability in patients with septic shock correlate with changes in SOFA scores?. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 27 | Do erythrocytes subjected to cardiopulmonary bypass exhibit changes in their membrane mechanical properties?. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 28 | Platelet-associated oxidative stress and ADAMTS-13 levels are inversely associated with a poor prognosis in septic shock. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 29 | Neutrophil gelatinase-associated lipocalin/lipocalin2, derived from gut crypt cells, exerts intestinal antimicrobial effect via bacterial stimulation of Toll-like receptor 4 and 9. <i>Critical Care</i> , 2013, 17, .                                 | 5.8 | 0         |
| 30 | Lethal influenza virus A H1N1 infection in two relatives with autosomal dominant GATA-2 deficiency. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 31 | Bacterial translocation primes proinflammatory responses and is connected to early death in an experimental model of lethal injury. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 32 | Effects of the common 34C>T variant of the AMPD1 gene on immune function, multiorgan dysfunction and mortality in patients with sepsis. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 33 | Exploring the translational disconnect between the murine and human inflammatory response: in vitro analysis of the dose-response relationship of LPS and NF $\kappa$ B activation in murine and human immune cells. <i>Critical Care</i> , 2013, 17, . | 5.8 | 0         |
| 34 | Hypogammaglobulinemia in sepsis is not correlated to high circulating angiotensin-2 levels. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 35 | TREM-1 levels are elevated in patients with liver cirrhosis. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 36 | Effects of different doses and serotypes of LPS on blood-brain barrier permeability in Sprague-Dawley rats. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 37 | Bioenergetic imbalance and oxidative stress in the pathophysiology of septic encephalopathy. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 38 | Pathophysiology of sepsis-associated brain dysfunction: an experimental study of cerebral microdialysis and mitochondrial function. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 39 | Cholinergic modulation of hippocampal activity during septic encephalopathy. Critical Care, 2013, 17, .  | 5.8 | 1         |
| 40 | Neutrophil-lymphocyte count ratio as a biomarker of severe sepsis in Escherichia coli infections in adults. Critical Care, 2013, 17, .   | 5.8 | 3         |
| 41 | Obesity and inflammatory markers in severe sepsis. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 42 | Cytokine gene expression can predict infectious complications following severe trauma. Critical Care, 2013, 17, .  | 5.8 | 3         |
| 43 | A cohort study of routinely used sepsis biomarkers and 28-day mortality. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 44 | Procalcitonin as prognostic marker of mortality. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 45 | Changes in circulating procalcitonin versus C-reactive protein in predicting evolution of infectious disease in febrile, critically ill patients. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 46 | Procalcitonin-guided antibiotic therapy in patients with congestive heart failure and suspicion of lower respiratory tract infection: results from a randomized trial. Critical Care, 2013, 17, .                                | 5.8 | 2         |
| 47 | Pro-adrenomedullin as a clinical predictor after cardiac surgery. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 48 | Pro-adrenomedullin as prognostic biomarker in the sepsis. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 49 | Early IL-6 response in sepsis is correlated with mortality and severity score. Critical Care, 2013, 17, .  | 5.8 | 17        |
| 50 | Compared values of presepsin (sCD14-ST) and procalcitonin as early markers of outcome in severe sepsis and septic shock: a preliminary report from the Albumin Italian Outcome Sepsis (ALBIOS) study. Critical Care, 2013, 17, . | 5.8 | 0         |
| 51 | Usefulness of presepsin in the diagnosis of sepsis in acute kidney injury patients. Critical Care, 2013, 17, .   | 5.8 | 1         |
| 52 | Serum levels of presepsin reflects the APACHE II and SOFA scores in patients with sepsis. Critical Care, 2013, 17, .   | 5.8 | 1         |
| 53 | Relationship between serum nitrite/nitrate levels in the early phase of septic acute lung injury and prognosis. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 54 | Simple and fast prediction of Legionella sp. in community-acquired pneumonia: validation of a prediction rule. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 55 | Host genetic variants associated with community-acquired pneumonia. Critical Care, 2013, 17, .   | 5.8 | 1         |
| 56 | Outcome of severe community-acquired pneumonia: the impact of comorbidities. Critical Care, 2013, 17, .  | 5.8 | 2         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 57 | Prognostic scores and infective endocarditis. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 58 | Effectiveness of early ureteric stenting for urosepsis associated with urinary tract calculi. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 59 | Outcomes in urinary sepsis. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 60 | Nosocomial pneumonia in the postoperative period after heart transplantation. Critical Care, 2013, 17, .  | 5.8 | 2         |
| 61 | Abdominal infection plays a role in the incidence of ventilator-associated pneumonia. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 62 | Incidence and risk factors for ventilator-associated pneumonia in Siriraj Hospital. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 63 | Hospital-acquired bloodstream infection: Indian perspective. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 64 | Risk factors for catheter-related bloodstream infections in a surgical ICU. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 65 | Analysis of risk factors for catheter-related bloodstream infection in a parenteral nutrition population. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 66 | Can we predict the postoperative infections?. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 67 | Results for introduction of a new hand hygiene program in the ICU. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 68 | Improving hand hygiene performance in a surgical ICU: a novel method for data collection to change physician behavior. Critical Care, 2013, 17, .                           | 5.8 | 0         |
| 69 | Compliance with the implementation of an ICU cluster-randomized trial assessing the benefits and potential harms of universal glove and gowning. Critical Care, 2013, 17, . | 5.8 | 1         |
| 70 | Fighting hospital sepsis. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 71 | Sinks as a correctable source of ESBL contamination for patients in the ICU. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 72 | Hemodynamic and mitochondrial effects of enalapril in experimental sepsis. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 73 | Effects of intraperitoneal immunoglobulin therapy on behavior and cognitive functions in CLP-induced sepsis model in rats. Critical Care, 2013, 17, .                       | 5.8 | 0         |
| 74 | Clinical efficacy of recombinant human soluble thrombomodulin in patients with septic disseminated intravascular coagulation. Critical Care, 2013, 17, .                    | 5.8 | 0         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 75 | Antithrombin III concentrate therapy may have an efficacy in sepsis. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 76 | A clinical randomized trial on the use of atorvastatin in patients with sepsis or septic shock: effects on endothelial function. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 3         |
| 77 | A multicenter randomized controlled study of an extracorporeal cytokine hemoabsorption device in septic patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 35        |
| 78 | Continuous renal replacement therapy with the adsorbent membrane oXiris in septic patients: a clinical experience. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 9         |
| 79 | Combined use of pumpless extracorporeal lung assist system and continuous renal replacement therapy with citrate anticoagulation in polytrauma patients. <i>Critical Care</i> , 2013, 17, .                               | 5.8 | 0         |
| 81 | Use of extracorporeal endotoxin elimination therapy for septic shock. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 5         |
| 82 | Selective adsorption of lipopolysaccharide in the complex treatment of patients with severe sepsis. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 83 | Alteco endotoxin hemoabsorption in Gram-negative septic shock patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 84 | Steroid therapy in septic shock: survey of practice amongst UK critical care physicians. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 85 | Need for restoration of cortisol serum levels for successful antimicrobial therapy in experimental sepsis. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 86 | Impact of time to initiation of appropriate antibiotic therapy on early mortality of patients with septic shock. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 87 | Comparison of survivors and nonsurvivors in 27 confirmed injectional anthrax cases from the 2009 outbreak in Scotland. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 3         |
| 88 | Efficacy and utility of preemptive anti-Gr(+) coverage in selected critically ill patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 89 | Gram-negative resistance and need for ICU among urinary tract infections in the United States. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 90 | Multidrug resistance among <i>P. aeruginosa</i> and Enterobacteriaceae in US hospitals, 2000 to 2009. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 91 | Prevalence of multidrug resistance, extensive drug resistance and pandrug resistance among multiple Gram-negative isolates: experience in a tertiary-care hospital ICU in North India. <i>Critical Care</i> , 2013, 17, . | 5.8 | 6         |
| 92 | Risk factors associated with <i>Acinetobacter baumannii</i> septicemia and its mortality rates in critically ill patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 93 | Prevalence of colonization by multiresistant bacteria on admission to the ICU in the French military hospital in Kabul, Afghanistan. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 94  | Extended-spectrum $\beta$ -lactamase <i>Providencia stuartii</i> in a general ICU. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 95  | High-dose tigecycline use in severe infections. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 96  | In vitro efficiency of Amikacin Inhale, a novel drug-device delivery system. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 6         |
| 97  | Analysis of parameters in critically ill patients influencing amikacin peak levels. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 98  | Cost implication of antibiotic-associated diarrhoea and the financial impact of probiotic use in its prevention. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 99  | Empirical versus preemptive antimycotic therapy in terms of outcome benefit. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 100 | Impact of echinocandins as first-choice therapy in cardiothoracic high-risk patients with candidemia. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 101 | Fungal ventricular assist device infections occur in colonized patients and are associated with high mortality rate. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 102 | Cost-effectiveness analysis of anidulafungin in the treatment of candidaemia. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 103 | Prevalence and impact of invasive fungal infections in intensive care. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 104 | Beliefs and actual practice of oxygen therapy in the ICU. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 105 | FiO <sub>2</sub> /PEEP index: a simple tool for optimizing ventilator settings. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 106 | Use of high-flow nasal canulae: effect on alveolar pressure and its limitation. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 4         |
| 107 | Relation between PaO <sub>2</sub> /FiO <sub>2</sub> ratio, SpO <sub>2</sub> /FiO <sub>2</sub> ratio, oxygenation index and ventilation ratio in critically ill patients. <i>Critical Care</i> , 2013, 17, .         | 5.8 | 0         |
| 108 | Decreased respiratory complexity during apnea in acute respiratory distress syndrome. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 109 | Acute respiratory distress syndrome - the Berlin definition: impact on an ICU of a university hospital. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 110 | Extravascular lung water, B-type natriuretic peptide and blood volume contraction for diagnosing weaning-induced pulmonary edema. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 111 | Pressure support ventilation with minimal sedation as the main ventilatory mode in critically ill patients with lung injury: effect on mortality and incidence of complications. <i>Critical Care</i> , 2013, 17, . | 5.8 | 1         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 112 | Comparison of a fully automatic ARDSNet protocol and a feedback-controlled open lung management concept. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 113 | Comparison of an entirely automated ventilation mode, Intellivent-ASV, with conventional ventilation in ARDS patients: a 48-hour study. <i>Critical Care</i> , 2013, 17, .                       | 5.8 | 3         |
| 114 | Difference in pulmonary permeability between indirect and direct acute respiratory distress syndrome assessed by the transpulmonary thermodilution technique. <i>Critical Care</i> , 2013, 17, . | 5.8 | 0         |
| 115 | Biomarkers for early stage of acute respiratory distress syndrome in septic patients: surfactant protein D and Clara cell protein. <i>Critical Care</i> , 2013, 17, .                            | 5.8 | 1         |
| 116 | Intravenous perfluorocarbons for prevention of ventilator-associated ARDS. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 117 | Effect of angiotensin-converting enzyme gene I/D polymorphism and its expression on clinical outcome in acute respiratory distress syndrome. <i>Critical Care</i> , 2013, 17, .                  | 5.8 | 0         |
| 118 | End-expiratory esophageal pressure versus lower inflection point in acute lung injury. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 119 | Lung recruitment induced by Sigh in hypoxemic intubated critically ill patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 120 | Quantification of stress raisers in ARDS. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 5         |
| 121 | Recruited lung tissue does not resume normal mechanical properties. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 2         |
| 122 | Intraoperative management of hypoxemia with recruitment maneuvers: are the benefits worth the costs?. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 123 | Haemodynamic changes during alveolar recruitment manoeuvre in patients with diastolic dysfunction. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 124 | Alveolar Recruitment for ARDS Trial: preliminary results. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 125 | Reliability of pressure measurements via balloon catheters is high: an evaluation of six esophageal catheters. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 126 | Effect of protective ventilation on organ-specific cytokine production in an experimental postoperative sepsis model. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 127 | Time required for gas exchange equilibration after a change of positive end-expiratory pressure in acute respiratory distress syndrome. <i>Critical Care</i> , 2013, 17, .                       | 5.8 | 0         |
| 128 | Comparison between the standard and low-dose chest CT scans on the lung quantitative analysis in critically ill patients. <i>Critical Care</i> , 2013, 17, .                                     | 5.8 | 0         |
| 129 | Preliminary experience with airway pressure release ventilation on hemodynamics in patients with septic shock in a medical/surgical ICU. <i>Critical Care</i> , 2013, 17, .                      | 5.8 | 0         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 130 | Feasibility and effectiveness of prone position in morbidly obese ARDS patients: a case-control clinical study. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 2         |
| 131 | Provision of mechanical ventilation to pregnant/postpartum women with H1N1 influenza: a case-control study. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 132 | Inhalation injury in critical burn patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 133 | Beneficial effects of intrapulmonary percussive ventilation in patients with respiratory insufficiency in the ICU. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 134 | High-frequency ventilation for acute traumatic and nontraumatic lung injury. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 135 | Improvement in oxygenation with high-frequency oscillatory ventilation combined with tracheal gas insufflation is correlated to the extravascular lung water index. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 136 | Physiologic effects and regional ventilation of high-frequency positive-pressure ventilation using a conventional ventilator in a severe ARDS animal model associated with an inspiratory pause or recruitment maneuvers. <i>Critical Care</i> , 2013, 17, . | 5.8 | 0         |
| 137 | High-frequency percussive ventilation at altitude: study in a hypobaric chamber with a mechanical test lung. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 138 | Recruitable volume is comparable in acute respiratory distress syndrome and in healthy lungs. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 139 | Nosocomial bloodstream infection and venovenous extracorporeal membrane oxygenation: a retrospective cohort study. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 140 | Monitoring anticoagulation during extracorporeal membrane oxygenation in patients with acute respiratory failure. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 5         |
| 141 | Extracorporeal membrane oxygenation: the MOTOR of cytokine production?. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 142 | Effect of extracorporeal CO <sub>2</sub> removal on respiratory rate in spontaneously breathing patients with chronic obstructive pulmonary disease exacerbation. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 8         |
| 143 | ECCO <sub>2</sub> removal with a phosphorylcholine-coated membrane oxygenator in difficult respiratory weaning patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 144 | Lung elastance can be determined without esophageal pressure measurements. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 145 | Feasibility of minimally invasive stimulation of the phrenic nerves for supporting ventilation in fully anesthetized swine. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 146 | Prehospital management of COPD patients in respiratory failure and short-term outcome. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 147 | N-terminal pro-brain natriuretic peptide as a prognostic marker in patients with exacerbation of chronic obstructive pulmonary disease. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 148 | Novel biomarkers for prediction of mortality after acute exacerbation of chronic obstructive pulmonary disease. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 149 | High incidence of respiratory viruses in critically ill adult patients with respiratory failure. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 150 | Role of bronchoscopy during non-invasive ventilation in hypercapnic respiratory failure. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 151 | SMART-COP score for patients admitted with community-acquired pneumonia (CAP) to an ICU in a district general hospital: a smarter way of identifying patients with severe CAP?. Critical Care, 2013, 17, .         | 5.8 | 0         |
| 152 | Effects of steroid therapies to severe streptococcal pneumoniae that required mechanical ventilation. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 153 | Risk factors for ventilator-associated pneumonia in a surgical ICU. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 154 | Ventilation distribution measured with electrical impedance tomography at varying levels of assist during pressure support versus neurally adjusted ventilatory assist in ICU patients. Critical Care, 2013, 17, . | 5.8 | 0         |
| 155 | Analysis of bilevel positive airway pressure therapy in children based on weight. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 156 | Determinants of non-invasive ventilation success or failure in morbidly obese patients in acute respiratory failure. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 157 | Non-invasive ventilation in patients with a clinical diagnosis of pneumonia: a clinical audit. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 158 | Analysis of management of non-invasive ventilation support in prehospital care for COPD patients and short-term outcome. Critical Care, 2013, 17, .  | 5.8 | 1         |
| 159 | Retrospective review of utilisation and outcomes of diaphragmatic EMG monitoring and neurally adjusted ventilatory assist in a central London teaching hospital over a 3-year period. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 160 | Validation of a lung ultrasound protocol in acute respiratory failure: preliminary results. Critical Care, 2013, 17, .   | 5.8 | 2         |
| 161 | Risk of weaning failure with PAOP $\leq 15$ versus PAOP $\leq 18$ . Critical Care, 2013, 17, .   | 5.8 | 1         |
| 162 | Impact of continuous positive airway pressure on the treatment of acute exacerbation of asthma: a randomised controlled trial. Critical Care, 2013, 17, .  | 5.8 | 2         |
| 163 | Novel method to confirm tracheal intubation based on airway pressures: validation in patients with lung disease. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 164 | Main indications, diagnostic and therapeutic yield of bronchoscopy in the ICU. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 165 | Benchtop study comparing leakages across cuffs of three endotracheal tubes. Critical Care, 2013, 17, .   | 5.8 | 1         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 166 | Endotracheal tube with tapered-type cuff for preventing ventilator-associated pneumonia: a randomized clinical trial. <i>Critical Care</i> , 2013, 17, .                                       | 5.8 | 1         |
| 167 | Effect of body position changes on endotracheal tube cuff pressure in ventilated patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 168 | Towards safer airway management in the critically ill: lessons from National Audit Project 4. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 169 | Quantitative assessment of endotracheal tube (ETT) biofilm by micro-CT scan: evaluation of the effectiveness of ETT cleaning devices. <i>Critical Care</i> , 2013, 17, .                       | 5.8 | 0         |
| 170 | Early identification of patients at risk of difficult intubation in the ICU: development and validation of the MACOCHA score in a multicenter cohort study. <i>Critical Care</i> , 2013, 17, . | 5.8 | 4         |
| 171 | Incidence of major complications related to endotracheal intubation in ICUs of nonacademic Brazilian public hospitals. <i>Critical Care</i> , 2013, 17, .                                      | 5.8 | 0         |
| 172 | Tracheal intubation for a difficult airway using Airway scopeÂ®, KingVisionÂ® and McGRATHÂ®: a comparative manikin study of inexperienced personnel. <i>Critical Care</i> , 2013, 17, .        | 5.8 | 5         |
| 173 | Resistance of neonatal endotracheal tubes: a comparison of four commercially available types. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 2         |
| 174 | NT-proBNP and cardiac cycle efficiency changes during extubation process in critically ill patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 175 | Comparison of outcomes between early and late tracheostomy for critically ill patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 3         |
| 176 | Ultrasound scanning for percutaneous dilatational tracheostomy: a systematic review. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 177 | Evaluation of a new device for emergency transcrioid ventilation in a manikin model. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 178 | Device for measuring bronchodilator delivery and response in resource-limited settings. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 179 | Evidence-based guidelines and protocols for the management of adult patients with a tracheostomy: a systematic review. <i>Critical Care</i> , 2013, 17, .                                      | 5.8 | 0         |
| 180 | Survey of emergency tracheostomy management in the East of England region. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 181 | Can we keep neck breathers safe? A survey on training adequacy of medical staff caring for tracheostomy/laryngectomy patients. <i>Critical Care</i> , 2013, 17, .                              | 5.8 | 0         |
| 182 | Laryngeal mask as a safe and effective ventilatory device during Blue Dolphin tracheostomy in the ICU. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 183 | Real-time ultrasound-guided balloon dilatational percutaneous tracheostomy is a safe procedure in critically ill patients: an evaluation study. <i>Critical Care</i> , 2013, 17, .             | 5.8 | 20        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 184 | Right ventricular restriction in interventional lung assist for acute respiratory distress syndrome. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 185 | Left ventricular electromechanical dyssynchrony and mortality in cardiothoracic intensive care. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 186 | Safety of ultrasound-guided central venous access in critically ill patients with uncorrected coagulopathy. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 187 | Cyanoacrylate glue prevents early bleeding of the exit site after CVC or PICC placement. Critical Care, 2013, 17, .   | 5.8 | 10        |
| 188 | Intracardiac ECG for confirmation of correct positioning of central venous catheters is safe and cost-effective. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 189 | Is automated brachial cuff measurement of arterial pressure less accurate in cases of arrhythmia?. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 190 | Cardiac output monitoring in brain-stem-dead potential organ donors: an audit of current UK practice. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 191 | Indocyanine green plasma disappearance rate for assessment of liver function: re-evaluation of normal ranges and impact of biometric data. Critical Care, 2013, 17, .   | 5.8 | 2         |
| 192 | Prediction of 28-day mortality by indocyanine green disappearance rate, other markers of hepatic function and transpulmonary thermodilution parameters: a prospective study in 154 patients. Critical Care, 2013, 17, . | 5.8 | 0         |
| 193 | Mixed and central venous oxygen saturation are not interchangeable in patients with cardiogenic shock after cardiac surgery. Critical Care, 2013, 17, .   | 5.8 | 1         |
| 194 | Pulmonary hemodynamic disorder in pediatric sepsis and their correction with L-arginine infusion. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 195 | Utility of transesophageal echocardiography in the ICU: a preliminary US perspective. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 196 | Cardiopulmonary monitoring in Thai ICUs: results of ICU-RESOURCE I surveys. Critical Care, 2013, 17, .  | 5.8 | 1         |
| 197 | Establishing a critical care echocardiography laboratory. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 198 | Contrast-enhanced ultrasonography in the ICU: promising tool or exciting toy?. Critical Care, 2013, 17, .   | 5.8 | 1         |
| 199 | Hemodynamic disclosure of septic shock patients by intensive care ultrasound. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 200 | Influence of positive end-expiratory pressure on four-chamber longitudinal strain analysis by speckle tracking echocardiography. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 201 | Evaluation of a new calibration index suggesting recalibration of the pulse contour cardiac index by transpulmonary thermodilution: a prospective study. Critical Care, 2013, 17, .                                     | 5.8 | 0         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 202 | Validation of less-invasive hemodynamic monitoring with Pulsioflex in critically ill patients: interim results of a multicentre study. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 203 | Interim results of an ongoing study on the use of non-invasive hemodynamic monitoring with Nexfin in critically ill patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 204 | Cost-effectiveness analysis of stroke volume variation guided perioperative hemodynamic optimization. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 205 | EV1000/VolumeView: a new device for a more reliable measurement of extravascular lung water index in patients with lung resections. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 206 | Goal-directed resuscitation therapy in high-risk patients undergoing cardiac surgery (GRICS study): a randomized controlled trial - preliminary results. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 207 | Perioperative volume management for esophageal cancer surgery. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 208 | Treatment-related cardiac complications associated with goal-directed therapy in high-risk surgical patients: a meta-analysis. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 209 | Estimation of potential cost-savings related to the implementation of perioperative hemodynamic goal-directed therapy. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 210 | Cardiac output estimation: Vigileo and Mostcare versus echocardiography. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 211 | Brachial pulse waveform characteristics predict development of organ failure in septic patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 212 | Computerized decision support system improves consistency of haemodynamic assessment amongst ICU team members. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 213 | Volume quantification by contrast-enhanced ultrasound and thermodilution: an in vitro comparison. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 214 | Successful introduction of cardiac index, fluid responsiveness and oxygen delivery data into the primary survey at a central London major trauma centre and impact on time to CT, fluid resuscitation and disposal. <i>Critical Care</i> , 2013, 17, . | 5.8 | 0         |
| 215 | Fluid responsiveness in pressure support ventilation: role of asynchrony. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 216 | Comparison of pulse pressure variation with radial arterial systolic, diastolic and pulse transit time interval variation in pediatric patients undergoing liver transplantation. <i>Critical Care</i> , 2013, 17, .                                   | 5.8 | 0         |
| 217 | Mini-fluid challenge: how much fluid and what parameter to use?. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 218 | Looking for the inflexion point of the Frank-Starling curve. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 219 | Prediction of stroke volume response to fluid bolus in 100 children. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 220 | Microcirculatory changes during goal-directed or mean arterial pressure-guided fluid therapy in abdominal surgery. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 221 | Postoperative imaging of the intestinal microcirculation. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 222 | Monitoring microcirculatory blood flow with a new sublingual tonometer in a porcine model of haemorrhagic shock. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 223 | Sublingual microcirculatory changes during transient intra-abdominal hypertension: a study in laparoscopic surgery patients. <i>Critical Care</i> , 2013, 17, .                                   | 5.8 | 0         |
| 224 | Changes in microcirculatory perfusion during cardiac surgery are paralleled by alterations in glycocalyx integrity. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 8         |
| 225 | Microvascular response to cold stress in healthy humans. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 226 | Microcirculatory response to experimentally-induced whole body heat stress in healthy humans. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 227 | Observational study of the effects of age, diabetes mellitus, cirrhosis and chronic kidney disease on sublingual microvascular flow. <i>Critical Care</i> , 2013, 17, .                           | 5.8 | 0         |
| 228 | Lactate clearance as a determinant of mortality in surgical patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 229 | Assessment of the mottling score as a mortality predictor in critically ill patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 4         |
| 230 | Impact of different haemodynamic resuscitation strategies on brain perfusion and tissue oedema markers in a model of severe haemorrhagic shock. <i>Critical Care</i> , 2013, 17, .                | 5.8 | 0         |
| 231 | Accuracy of the shock index and various modified shock indexes to predict early mortality in patients suffering from gastrointestinal haemorrhage. <i>Critical Care</i> , 2013, 17, .             | 5.8 | 4         |
| 232 | Inhaled carbon monoxide or nebulized sodium nitrite protect against hemorrhagic shock-induced mitochondrial dysfunction. <i>Critical Care</i> , 2013, 17, .                                       | 5.8 | 0         |
| 233 | Is norepinephrine more effective than other vasopressors for septic shock? A systematic review and meta-analysis. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 234 | Vasopressin Versus Norepinephrine for the Management of Shock After Cardiac Surgery (VaNCS) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 1  | 5.8 | 1         |
| 235 | Intraaortic counterpulsation in a second-level institution: indications, management and outcome. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 236 | Impact of left ventricular ejection fraction and elevated filling pressures on mortality in mechanically ventilated patients in severe sepsis or septic shock. <i>Critical Care</i> , 2013, 17, . | 5.8 | 0         |
| 237 | Effect of $\beta$ -blockers and calcium channel blockers on shock index predictability in patients suffering from urosepsis. <i>Critical Care</i> , 2013, 17, .                                   | 5.8 | 0         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 238 | Levosimendan in critically ill adults: a utilisation review. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 239 | Levosimendan versus dobutamine in Tako-tsubo cardiomyopathy. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 2         |
| 240 | Combined effects of ivabradine with dobutamine or levosimendan in isolated perfused hearts. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 241 | Influence of temperature on the cardiac action of dobutamine and levosimendan in isolated perfused hearts. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 242 | Conflicting results between V/Q SPECT and angioscan in pulmonary embolism: what to do?. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 243 | Intraabdominal hypertension and abdominal compartment syndrome in patients admitted to a medical and surgical ICU at a third referral university hospital. <i>Critical Care</i> , 2013, 17, . | 5.8 | 2         |
| 244 | Red cell distribution width predicts cardiovascular complications after high-risk surgery. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 245 | Consequences of obesity in outcomes after cardiac surgery: analysis of the ARIAM registry. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 246 | Myocardial injury in critically ill patients admitted with noncardiac diagnoses. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 247 | Validation of the GRACE score risk in our population. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 248 | Outcomes of incomplete and complete revascularization in ACS. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 249 | Non-invasive valve implantation with TAVI versus conventional aortic valve replacement. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 250 | Time to recover from shock is determinant of a positive fluid balance in septic shock. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 251 | Lactate clearance is a predictor of sustained bleeding in emergency room patients with moderate upper gastrointestinal bleeding. <i>Critical Care</i> , 2013, 17, .                           | 5.8 | 0         |
| 252 | Muscular glucose assessed by microdialysis and blood glucose can predict mortality in septic shock. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 253 | Constipation in critically ill patients and its relationship to feeding and weaning from respiratory support. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 7         |
| 254 | Nasogastric tube dislodgement: a problem on our ICU. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 255 | Handling of dietary protein in critically ill patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 256 | Confirmation of nasogastric tube placement in critical care. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 257 | pH testing to confirm nasogastric tube position on the ICU: are we wasting our time?. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 258 | Effect of probiotic containing lactobacillus, bifidobacterium and streptococcus thermophilus in critically ill patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 259 | Selenium levels in patients with different sources of sepsis. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 260 | Is a combined i.v. and enteral glutamine regimen superior to a single i.v. glutamine regimen in severe thoracic trauma?. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 261 | Clinical effect of altered lipid emulsion composition containing fish oil in postoperative patients following abdominal surgery: a prospective, randomized, open-label, comparative, multicenter phase 3 clinical study. <i>Critical Care</i> , 2013, 17, . | 5.8 | 1         |
| 262 | Influence of fat-based versus glucose-based enteral nutrition formulas on glucose homeostasis. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 3         |
| 263 | Effect of hypocaloric versus normocaloric parenteral nutrition on whole body protein kinetics in critically ill neurosurgical patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 264 | Withholding parenteral nutrition for 1 week reduces ICU-acquired weakness. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 2         |
| 265 | Impact of early parenteral nutrition on muscle and adipose tissue compartments during critical illness. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 10        |
| 266 | Withholding parenteral nutrition during the first week of critical illness increases plasma bilirubin but lowers the incidence of cholestasis and gallbladder sludge. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 267 | Impact of early parenteral nutrition on catabolism. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 268 | Impact of early versus late parenteral nutrition on morphological and molecular markers of atrophy and autophagy in skeletal muscle of critically ill patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 269 | Impact of closing an emergency department on a neighbouring teaching hospital: the concentrate effect. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 270 | Overcrowding estimation in the emergency department: is the simplest score the best?. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 271 | GP-led walk-in centre in the UK: another way for urgent healthcare provision. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 272 | Establishing a new emergency department: effects on patient flow. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 273 | Prehospital EKG evaluation in Rio de Janeiro ambulances. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 274 | Technique of ultrasound-guided peripheral venous access in the emergency room. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 275 | FIB the fractured femur. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 276 | Abdominal pain in adolescent females: a single-centre audit and review of management. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 277 | Multidisciplinary approach to improving documentation of visual acuity in patients presenting with ocular trauma. Critical Care, 2013, 17, .              | 5.8 | 0         |
| 278 | BiPAP for treating moderate and severe asthma exacerbations in a PED. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 279 | Report of an outbreak of toxicity from a novel drug of abuse in the UK: Eric-3. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 281 | Thallium group poisoning incident in Japan 2011. Critical Care, 2013, 17, .   | 5.8 | 1         |
| 282 | Heart rate variability in children with tricyclic antidepressant intoxication. Critical Care, 2013, 17, .   | 5.8 | 2         |
| 283 | Diagnosing anisakiasis in the emergency department. Critical Care, 2013, 17, .  | 5.8 | 1         |
| 284 | Implementation of a sepsis protocol in a community hospital using a Telemedicine Program. Critical Care, 2013, 17, .                                      | 5.8 | 1         |
| 285 | Initial Brazilian experience of Telestroke for thrombolysis in a community hospital. Critical Care, 2013, 17, .   | 5.8 | 1         |
| 286 | Analysis of emergency calls achieved in a French emergency dispatching centre: what resources for which patients. Critical Care, 2013, 17, .              | 5.8 | 0         |
| 287 | Reform of emergency services: immediate effects on cardiac care unit and ICU patient intake. Critical Care, 2013, 17, .                                   | 5.8 | 0         |
| 288 | Overtriage and undertriage in a prehospital system over 7 years. Critical Care, 2013, 17, .   | 5.8 | 1         |
| 289 | Lorca's hospital evacuation and medical attention by the emergency department. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 290 | War injuries and civilian accidents in Afghan children. Critical Care, 2013, 17, .  | 5.8 | 2         |
| 291 | Intensive care and disaster medicine: the role of a compendium. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 292 | Factors involved in ICU mortality following medical retrieval: identifying differences between ICU survivors and nonsurvivors. Critical Care, 2013, 17, . | 5.8 | 0         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 293 | Algorithm for the resuscitation of traumatic cardiac arrest patients in a physician-staffed helicopter emergency medical service. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 294 | Are physicians required during HEMS winch rescue missions?. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 295 | Improving maintenance of critical care land and aeromedical transfer equipment. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 296 | Safety standards for intrahospital transfer of critical care patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 297 | Prehospital hyperoxemia does not influence the functional neurological outcome in polytraumatized patients with traumatic head injury. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 298 | Innate immune response-mediated late increase in SuPAR in multi-trauma patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 299 | Single-dose estrogen infusion can amplify brain levels of Sonic hedgehog, a signal protein for neuro stem cells and repair following the indirect brain injury resulting after severe torso burns. <i>Critical Care</i> , 2013, 17, .               | 5.8 | 3         |
| 300 | Feasibility and experience of using exception from informed consent in a pilot study of immediate estrogen infusion for hypotensive trauma patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 301 | Early administration of a single dose of parenteral estrogen decreases inflammation in the heart for 45 days after severe burns. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 302 | Accuracy of FAST scan in blunt abdominal trauma in a major London trauma centre. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 6         |
| 303 | Level of lipopolysaccharide-binding protein and microbial landscape with account of severity of sepsis syndromes in polytrauma. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 304 | Risk factors for the development of complications following blunt chest trauma: a new risk stratification tool. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 305 | Guidelines for chest drain insertion do not protect relevant anatomical structures. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 306 | Prediction score of indications for whole body computed tomography in blunt trauma patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 307 | Prehospital blood transfusion: 5-year experience of an Australian helicopter emergency medical service. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 17        |
| 308 | Severe hyperkalemia is prevalent and can be detected by quantitative electrocardiographic findings and medication history of calcium channel blocker among the patients with symptomatic or extreme bradycardia. <i>Critical Care</i> , 2013, 17, . | 5.8 | 1         |
| 309 | Prehospital risk stratification following out-of-hospital cardiac arrest. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 310 | Multidisciplinary Medical Emergency Team System increases survival in cancer patients undergoing cardiopulmonary resuscitation. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 311 | Focused cardiac ultrasound in resuscitation. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 312 | Acute intrathoracic gastric herniation as a rare cause of cardiac arrest. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 313 | Implementation of dispatcher-assisted cardiopulmonary instructions using the ALERT protocol: preliminary results in Belgium. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 314 | Prehospital epinephrine administration and survival among patients with unshockable initial rhythm after out-of-hospital cardiac arrest. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 315 | Impact of termination of resuscitation for out-of-hospital cardiopulmonary arrest in Japan. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 316 | Survival from cardiopulmonary arrest after regular working hours in a tertiary-care hospital: retrospective study. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 317 | Cognitive performance and capacity to return home following out-of-hospital cardiac arrest. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 318 | Start value of cerebral saturation in prehospital cardiac arrest patients: does it mean something?. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 319 | Intensive care admission after cardiac arrest: cardiac versus noncardiac causes and consequences for treatment. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 320 | Comparison of 1-year neurological outcome between intra-hospital and extra-hospital cardiac arrest survivors submitted to mild therapeutic hypothermia in a community-based setting in Brazil. <i>Critical Care</i> , 2013, 17, . | 5.8 | 0         |
| 321 | Primary percutaneous coronary angioplasty and therapeutic hypothermia in out-of-hospital cardiac arrest. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 322 | Infrared pupillometry for outcome prediction after cardiac arrest and therapeutic hypothermia. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 323 | Effect of moderate hyperventilation and induced hypertension on cerebral tissue oxygen saturation in comatose post-cardiac arrest patients treated with hypothermia. <i>Critical Care</i> , 2013, 17, .                           | 5.8 | 1         |
| 324 | Effects of viscosity on cerebral blood flow after cardiac arrest. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 325 | Therapeutic mild hypothermia after cardiac arrest in shockable and nonshockable rhythms: does it improve both survival and neurological outcome?. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 326 | Agreement between temperature measurements during cooling and warming through the esophagus. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 327 | Association between oxygenation and 6-month mortality during post-cardiac arrest care. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 328 | Neuron-specific enolase and bispectral index/suppression ratio for prognostication after cardiac arrest. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 329 | Motor scores, therapeutic hypothermia and neurological outcome after cardiac arrest. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 330 | Differences of calcium binding protein immunoreactivities in the young hippocampal CA1 region from the adult following transient ischemic damage. <i>Critical Care</i> , 2013, 17, . | 5.8 | 0         |
| 331 | Early hypothermia improves survival and reduces the rise of serum biomarkers after traumatic brain injury in swine. <i>Critical Care</i> , 2013, 17, .                               | 5.8 | 0         |
| 332 | Epidemiology of moderate and severe traumatic brain injury in Cairo University Hospital in 2010. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 333 | Alcohol intoxication impedes the recognition of traumatic brain injury in the prehospital setting and may worsen 6-month outcome. <i>Critical Care</i> , 2013, 17, .                 | 5.8 | 0         |
| 334 | Predictive ability of the IMPACT-TBI prognostic model is improved by adding markers of coagulation. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 335 | Nonischemic endogenous lactate production in humans with severe traumatic brain injury. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 336 | Japan Coma Scale used in the prehospital setting can predict clinical outcome in severe pediatric trauma. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 10        |
| 337 | Effect of positive end-expiratory pressure on ultrasound measurement of optical sheath nerve diameter: preliminary study. <i>Critical Care</i> , 2013, 17, .                         | 5.8 | 0         |
| 338 | National survey of current protocols and management of the traumatic brain injury patients in UK ICUs. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 339 | Low-frequency autoregulation index for calculation of optimal cerebral perfusion pressure in severe traumatic brain injury. <i>Critical Care</i> , 2013, 17, .                       | 5.8 | 1         |
| 340 | Cerebral rSO <sub>2</sub> monitoring in pediatric altered mental status patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 341 | Intravenous paracetamol for fever control in acute brain-injured patients: cerebral and hemodynamic effects. <i>Critical Care</i> , 2013, 17, .                                      | 5.8 | 2         |
| 342 | Retrospective analysis of the hemodynamic effects of induction of barbiturate coma in patients with refractory elevated intracranial pressure. <i>Critical Care</i> , 2013, 17, .    | 5.8 | 0         |
| 343 | Hospital readmissions following traumatic brain injury. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 6         |
| 344 | Risk factors of pituitary disorders following traumatic brain injury. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 345 | Vasopressin reduces the cumulative epinephrine dose in hypothermic swine with traumatic brain injury. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 346 | Long-term functional outcome in adults with severe TBI: a meta-analysis. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 347 | Monitoring of severe traumatic brain injury patients in UK ICUs: a national survey. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 348 | Survival analysis after head injury: is a normal INR reassuring?. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 349 | One-year extended Glasgow Outcome Scale and hospital mortality predictors in patients with severe traumatic brain injury in Brazil. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 350 | Early respiratory deterioration predicts poor outcome after aneurysmal subarachnoid hemorrhage. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 351 | Evaluation of proximal-distal velocity gradient in spastic middle cerebral artery after aneurysmal subarachnoid hemorrhage. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 352 | Microparticles in aneurysmal subarachnoid hemorrhage: role in acute and delayed cerebral ischemia. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 353 | Intracranial pressure after subarachnoid hemorrhage. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 354 | Prevalence of pituitary disorders associated with aneurysmal subarachnoid hemorrhage. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 355 | Correlation between the timing of endovascular coiling or microsurgical clipping and long-term outcomes of patients after aneurysmal subarachnoid haemorrhage. <i>Critical Care</i> , 2013, 17, .                                  | 5.8 | 0         |
| 356 | Time course of neuropsychological functioning after aneurysmal subarachnoid hemorrhage. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 357 | Neuroprotective effect of a new synthetic aspirin-decurisinol adduct in a rat model of ischemic stroke. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 358 | Prevalence of acute critical neurological disease in children: a global epidemiological assessment (PANGEA). <i>Critical Care</i> , 2013, 17, .  | 5.8 | 4         |
| 359 | Survival analysis of patients taking dabigatran who consulted an emergency department for acute bleeding: a first alarm. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 360 | Bedside anti-Xa measurement for therapeutic assessment of a prophylactic anticoagulation regimen. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 361 | Effective dosage of enoxaparin for intensive care patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 362 | Anti-Xa activities following 5,000 IU and 7,500 IU of s.c. dalteparin in critically ill patients and 5,000 IU in medical patients: a prospective randomized study. <i>Critical Care</i> , 2013, 17, .                              | 5.8 | 0         |
| 363 | Randomized double-blind placebo-controlled PK/PD study on the effects of a single intravenous dose of the anti-hepcidin Spiegelmer NOX-H94 on serum iron during experimental human endotoxemia. <i>Critical Care</i> , 2013, 17, . | 5.8 | 7         |
| 364 | Haemoglobin concentration on admission to intensive care influences hospital mortality rates and length of stay: a retrospective study. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 365 | Influence of epidural anesthesia on the hemocoagulation disorders and quantity of septic complications in patients with acute necrotizing pancreatitis. Critical Care, 2013, 17, .               | 5.8 | 1         |
| 366 | Hemoviscoelastography: a new method of hemostasis monitoring. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 367 | Thromboelastographic monitoring of fondaparinux in surgical patients. Critical Care, 2013, 17, .   | 5.8 | 2         |
| 368 | Urokinase-induced fibrinolysis in thromboelastography (UKIF-TEG) to assess fibrinolysis in critically ill patients. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 369 | Bleeding complications after central line insertions and the relevance of pre-procedure coagulation tests and blood component therapy. Critical Care, 2013, 17, .                                | 5.8 | 4         |
| 370 | Efficacy of tranexamic acid in decreasing blood loss during cesarean section. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 371 | Dose-response of fibrinogen and factor XIII concentrate for correcting albumin-induced coagulopathy. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 372 | Intraoperative cell savage for life-threatening haemorrhagic caesarean section. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 373 | Does protamine infusion reduce postoperative blood loss and heparin rebound in cardiac surgery patients? A retrospective analysis of 240 patients on the cardiac ICU. Critical Care, 2013, 17, . | 5.8 | 0         |
| 374 | Transfusion Requirements in Surgical Oncology Patients (TRISOP): a randomized, controlled trial. Critical Care, 2013, 17, .  | 5.8 | 3         |
| 375 | Red blood cell transfusion is a predictor of mortality and morbidity in children undergoing cardiac surgery. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 376 | Red blood cell transfusion results in worse outcomes in patients with left ventricular dysfunction undergoing on-pump cardiac surgery. Critical Care, 2013, 17, .                                | 5.8 | 0         |
| 377 | Defining the need for a massive transfusion in severe blunt traumatic patients. Critical Care, 2013, 17, .   | 5.8 | 1         |
| 378 | Association between red blood cell transfusions and clinical outcome in critically ill children. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 379 | Microcirculation and blood transfusions during sepsis: leukoreduced (LR) versus non-LR red blood cells. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 380 | Massive blood transfusion for obstetric haemorrhage. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 381 | Blood transfusion practices in the ICU of a level 1 trauma centre and tertiary cardiac unit. Critical Care, 2013, 17, .  | 5.8 | 1         |
| 382 | C1 inhibitor attenuates pulmonary inflammation in an in vivo model of transfusion-related acute lung injury. Critical Care, 2013, 17, .  | 5.8 | 0         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 383 | Incidence, risk factors and outcome of TRALI after cardiac surgery. Critical Care, 2013, 17, .   | 5.8 | 2         |
| 384 | Albumin-induced coagulopathy is less severe and more effectively reversed with fibrinogen concentrate than is synthetic colloid-induced coagulopathy. Critical Care, 2013, 17, .                 | 5.8 | 0         |
| 385 | Network meta-analysis of clinical trials of fluid treatments in sepsis demonstrates improved survival with albumin compared with crystalloid and hydroxyethyl starch. Critical Care, 2013, 17, . | 5.8 | 0         |
| 386 | Assessing the economic impact of catheter-related bloodstream infections when switching from open to closed system delivery of human albumin solutions in ICUs. Critical Care, 2013, 17, .       | 5.8 | 0         |
| 387 | Comparing the hemodynamic effects of six different colloids. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 388 | Strong ion difference and arterial bicarbonate concentration as cornerstones of the impact of fluid therapy on acid-base balance. Critical Care, 2013, 17, .                                     | 5.8 | 1         |
| 389 | Microcirculatory response to fluid challenge: should we prefer balanced colloids to rebalance tissue perfusion?. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 390 | Procedural sedation in the emergency department. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 391 | Impact of an age, kidney and liver function adjusted sedation protocol in critically ill patients. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 392 | Risk and benefit of protocol sedation and daily interruption of sedation for mechanically ventilated patients: a systematic review. Critical Care, 2013, 17, .                                   | 5.8 | 0         |
| 393 | Chemical and physical compatibility of continuous intravenous drug infusion combinations used in paediatric intensive care. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 395 | Benzodiazepines induce hyperglycemia in rats by affecting peripheral disposal of glucose. Critical Care, 2013, 17, .   | 5.8 | 1         |
| 396 | Prolonged sedation in ARDS patients with inhaled anesthetics: our experience. Critical Care, 2013, 17, .   | 5.8 | 3         |
| 397 | Involvement of mitochondrial ATP-sensitive K <sup>+</sup> channels in fentanyl-induced mitochondrial dysfunction of cultured human hepatocytes. Critical Care, 2013, 17, .                       | 5.8 | 0         |
| 398 | Effects of clonidine on microcirculatory alterations during endotoxemia. Critical Care, 2013, 17, .  | 5.8 | 1         |
| 399 | Conjunctival administration of S+ketamine in rabbits. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 400 | Use of propofol for procedural sedation reduces length of stay in the emergency department. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 401 | Pre-PRIS? Prospective monitoring for early markers of propofol infusion syndrome. Critical Care, 2013, 17, .   | 5.8 | 1         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 402 | Safety profile of 1,008 cases of propofol sedation in the emergency department. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 403 | Use of physical restraint in Dutch ICUs: prevalence and motives. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 404 | Factors influencing the use of physical restraints in Canadian ICUs. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 405 | Current practices in ICU delirium management: a prospective multicenter study in the Netherlands. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 406 | Intraoperative dexamethasone and delirium after cardiac surgery: a randomized clinical trial. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 407 | Current perspectives, beliefs and practices concerning delirium in critically ill patients: a multicenter survey among Dutch healthcare professionals. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 408 | International validation of the delirium prediction model for ICU patients (PREDELIRIC): a multicenter observational study. <i>Critical Care</i> , 2013, 17, .                             | 5.8 | 1         |
| 409 | Delirium in ICU patients is associated with increased temperature variability. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 410 | Long-term outcome of delirium in critically ill patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 411 | Gastrointestinal symptoms in the ICU: a comparison of primary and secondary acute gastrointestinal injury. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 412 | Effects of proton pump inhibitor versus H2-receptor antagonist stress ulcer prophylaxis on ventilator-associated pneumonia: a pilot study. <i>Critical Care</i> , 2013, 17, .              | 5.8 | 6         |
| 413 | Early diagnosis of liver failure in septic patients using the maximal liver function capacity test (LiMAx) Tj ETQq1 1 0.784314 rgBT /Ove lo  | 5.8 | 1         |
| 414 | Impact of liver resection on portal venous pressure and renal function. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 415 | Value of the preoperative Model for End-stage Liver Disease score in predicting postoperative outcome in living donor liver transplantation recipients. <i>Critical Care</i> , 2013, 17, . | 5.8 | 0         |
| 416 | Does just fungal evidence increase the mortality of critically ill patients with liver cirrhosis? A retrospective analysis. <i>Critical Care</i> , 2013, 17, .                             | 5.8 | 0         |
| 417 | Predictors of persistent organ failure in ICU patients with severe acute pancreatitis. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 418 | Increased plasma levels of thrombopoietin in patients with severe acute pancreatitis. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 419 | Severe renal dysfunction measured by pRIFLE in children is an independent factor associated with death and not predicted by PIM2. <i>Critical Care</i> , 2013, 17, .                       | 5.8 | 0         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 420 | Thorough evaluation for the new acute kidney injury criteria by Kidney Disease Improving Global Outcomes. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 421 | Urinary neutrophil gelatinase-associated lipocalin as an early predictor of acute kidney injury in cardiac surgery patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 422 | Atrial natriuretic peptide attenuates metabolic acidosis and inflammation of the kidney, lung and heart in a rat model of renal ischemia-reperfusion injury. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 423 | Perioperative sodium bicarbonate to prevent acute kidney injury after cardiac surgery: a multicenter double-blind randomized controlled trial. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 424 | Tubular damage biomarkers linked to inflammation or iron metabolism predict acute kidney injury. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 425 | Renal response and acid-base balance alterations during furosemide administration. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 2         |
| 426 | Red cell distribution width is not a predictor of mortality in acute kidney injury. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 427 | Value of Tc-99m DMSA renal scan in assessing the prognosis and outcome of acute renal failure due to sepsis. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 428 | Estimated glomerular filtration rate based on hospital discharge creatinine may significantly overestimate renal function and underestimate chronic kidney disease in survivors of critical illness. <i>Critical Care</i> , 2013, 17, . | 5.8 | 1         |
| 429 | Varying models of intra-abdominal hypertension and their effect on renal function in a porcine model. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 430 | Renal resistive index can predict response to furosemide in septic patients with oliguria. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 431 | Urinary chitinase 3-like 1, a novel biomarker for acute kidney injury in adult cardiac intensive care patients: a pilot study. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 432 | Use, type and duration of cardiopulmonary bypass are related to acute kidney injury occurrence and urinary NGAL concentrations. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 433 | Symmetrical dimethylarginine is a more sensitive biomarker of renal dysfunction than creatinine. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 11        |
| 434 | Prognostic value of neutrophil gelatinase-associated lipocalin and transpulmonary thermodilution-derived parameters within 48 hours after admission. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 435 | Assessment of neutrophil gelatinase-associated lipocalin as a biomarker of acute kidney injury in acute heart failure patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 436 | Neutrophil gelatinase-associated lipocalin predicts postoperative fluid overload after cardiac surgery. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 437 | Survivors of acute kidney injury requiring renal replacement therapy rarely receive follow-up: identification of an unmet need. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 438 | Aetiology and outcomes for dialysis-dependent acute kidney injury patients on the ICU. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 439 | Efficiency of substitutive renal therapy in the complex intensive care of multiple organ failure in patients with polytrauma. <i>Critical Care</i> , 2013, 17, .                         | 5.8 | 0         |
| 440 | Can treatment with the molecular adsorbent recirculation system be the solution for type-1 hepatorenal syndrome?. <i>Critical Care</i> , 2013, 17, .                                     | 5.8 | 0         |
| 441 | Effect of continuous high-volume hemofiltration on patients with acute respiratory distress syndrome. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 2         |
| 442 | Dialyzer clearance of myoglobin with middle molecule filter and low blood flow CVHD in patients with rhabdomyolysis-associated acute kidney injury. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 443 | Daily fluid balance predicts hospital mortality in critically ill patients receiving continuous renal replacement therapy. <i>Critical Care</i> , 2013, 17, .                            | 5.8 | 0         |
| 444 | Fluid balance and renal outcomes in patients requiring renal replacement therapy in the ICU. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 445 | Pharmacokinetics and pharmacodynamics of esomeprazole and ranitidine during continuous venovenous hemodiafiltration. <i>Critical Care</i> , 2013, 17, .                                  | 5.8 | 0         |
| 446 | Pharmacokinetics of gentamicin and vancomycin during continuous venovenous hemofiltration in critically ill septic patients with acute kidney injury. <i>Critical Care</i> , 2013, 17, . | 5.8 | 1         |
| 447 | Effects of two different regional citrate anticoagulation CVVH protocols on acid-base status and phosphate supplementation. <i>Critical Care</i> , 2013, 17, .                           | 5.8 | 0         |
| 449 | Muscle wasting in the ICU can be reliably monitored using ultrasound. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 450 | Critical illness induces nutrient-independent adipogenesis and accumulation of alternatively activated tissue macrophages. <i>Critical Care</i> , 2013, 17, .                            | 5.8 | 0         |
| 452 | Hypocalcemia during magnesium therapy for obstetric ICU admissions. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 453 | Hyponatremia and intensive care outcome. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 454 | Strong ion gap can be accurately estimated with a simple bedside equation. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 455 | Relative adrenal insufficiency in burns. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 456 | Albumin-adjusted calcium concentration should not be used to identify hypocalcaemia in critical illness. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 457 | Vitamin D status and overall and sepsis mortality in critically ill patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 2         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 458 | Etiology of low testosterone levels in male patients with severe sepsis requiring mechanical ventilation. <i>Critical Care</i> , 2013, 17, .                           | 5.8 | 5         |
| 459 | Efficacy of calcium replacement in hypocalcaemia. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 460 | Endogenous melatonin in critically ill patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 461 | Succinate ameliorates mitochondrial oxygen consumption of metformin-intoxicated human platelets. <i>Critical Care</i> , 2013, 17, .                                    | 5.8 | 0         |
| 462 | Does chronic metformin therapy offer cardiomyoprotection in coronary artery bypass grafting surgery patients? A case-control study. <i>Critical Care</i> , 2013, 17, . | 5.8 | 0         |
| 463 | Metformin-associated lactic acidosis: risk factors and prognostic factors. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 464 | Multiscale modeling of acute insulin resistance in critical care. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 465 | Impact of a real-time electronic persistent hyperglycemia alert on glucose control in the ICU. <i>Critical Care</i> , 2013, 17, .                                      | 5.8 | 1         |
| 466 | Individualized targeted glucose control to avoid hypoglycemia. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 467 | Hyperglycemia in intensive care identifies patients at high risk of developing diabetes: preliminary data of a multicentre study. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 468 | Association between average glucose levels and hospital mortality among critically ill patients. <i>Critical Care</i> , 2013, 17, .                                    | 5.8 | 2         |
| 469 | Continuous glucose monitoring in critically ill adults: comparison of two different calibration protocols. <i>Critical Care</i> , 2013, 17, .                          | 5.8 | 1         |
| 470 | Root cause analysis of hypoglycemic events in critically ill patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 471 | Continuous arterial and venous glucose monitoring by quenched chemical fluorescence in ICU patients after cardiac surgery. <i>Critical Care</i> , 2013, 17, .          | 5.8 | 2         |
| 472 | Performance of the Medtronic SentrinoÂ® continuous glucose management system in the cardiac ICU. <i>Critical Care</i> , 2013, 17, .                                    | 5.8 | 2         |
| 473 | Intensive insulin therapy in critically ill children: impact on blood glucose dynamics and its relation with mortality. <i>Critical Care</i> , 2013, 17, .             | 5.8 | 0         |
| 474 | Implementation of evidence-based protocols improves survival: a 15-year surgical ICU experience in 10,172 patients. <i>Critical Care</i> , 2013, 17, .                 | 5.8 | 0         |
| 475 | Decubitus ulcers are associated with prolonged length of stay in critically ill patients. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 476 | HIV patients admitted to an ICU of a university hospital - experience of 15 years: 1995 to 2009. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 477 | Prevalence and prognosis of organ dysfunctions of AIDS critically ill patients. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 478 | Mortality predictors in septic HIV patients in the ICU. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 479 | Outcome of intensive care treatment for lung cancer patients. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 480 | Outcome of older patients with cancer requiring ICU admission. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 481 | Identifying risk factors for readmission to the ICU: a qualitative approach. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 482 | Clinical and laboratory postoperative impact on outcome of living donor liver transplantation recipients. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 483 | Are APACHE II scores better predictors of mortality than routine laboratory values?. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 484 | Complementarity of prognosis models SAPS 3 and EuroSCORE in cardiac surgery. Critical Care, 2013, 17, .  | 5.8 | 3         |
| 485 | Comparing intensive care treated primary, secondary and tertiary sepsis with respect to length of stay, organ failure and mortality: a pilot study. Critical Care, 2013, 17, . | 5.8 | 0         |
| 486 | Survival of older patients after critical care. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 487 | Eldicus scores for admission to our ICU: a retrospective study. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 488 | Developing a screening instrument for predicting psychological morbidity after critical illness. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 489 | Prediction of sequences of organ dysfunction in critical patients studied with Bayesian analysis. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 490 | Comparison of IGS 2 score and a new neural network biomarker score performance in predicting 1-year mortality. Critical Care, 2013, 17, .                                      | 5.8 | 0         |
| 491 | Significance of preadmission quality of life for mortality in the ICU: a prospective cohort study. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 492 | Predicting physical function and mental health in trauma intensive care patients 2 years after hospitalisation. Critical Care, 2013, 17, .                                     | 5.8 | 0         |
| 493 | Do ICU patients' discharge characteristics change with Swiss Diagnosis Related Groups?. Critical Care, 2013, 17, .   | 5.8 | 0         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 494 | Do Diagnosis Related Groups change admission practice to a large Swiss ICU?. Critical Care, 2013, 17, .   | 5.8 | 1         |
| 495 | Variable cost of a Greek ICU. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 496 | Audit of the medical staffing of and referrals to a mixed adult ICU in a teaching hospital. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 497 | Impact of the time elapsed between ICU request and actual admission on mortality and length of stay. Critical Care, 2013, 17, .                                       | 5.8 | 0         |
| 498 | Medical problems: review from the major nuclear incidents. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 499 | Variation in acute care burden and supply across diverse urban settings. Critical Care, 2013, 17, .   | 5.8 | 3         |
| 500 | Differences in severity and resource utilization for medical and surgical ICU patients. Critical Care, 2013, 17, .  | 5.8 | 1         |
| 501 | Educational effectiveness of introductory training for the Rapid Response System in Japan. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 502 | Standards of referrals and admissions of critically ill patients to the ICU. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 503 | Knowledge of confidentiality, consent and information disclosure is not affected by position or experience in adult critical care. Critical Care, 2013, 17, .         | 5.8 | 0         |
| 504 | Survey of alcohol-related admissions to critical care and the resource and financial implications. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 505 | Association of sleep disordered breathing with clinical trajectories in patients undergoing cardiac surgery. Critical Care, 2013, 17, .                               | 5.8 | 0         |
| 506 | Can massage therapy help cardiac patients? A literature review plus a case study. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 507 | Ischemic stroke in different ICU types. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 508 | Development of a severe influenza critical care curriculum and training materials for resource-limited settings. Critical Care, 2013, 17, .                           | 5.8 | 3         |
| 509 | Quality assurance in severe sepsis: an individualised audit/feedback system results in substantial improvements at a UK teaching hospital. Critical Care, 2013, 17, . | 5.8 | 1         |
| 510 | Safety climate perception for staff nurses in an ICU. Critical Care, 2013, 17, .  | 5.8 | 1         |
| 511 | Physico-technical medicine: teaching technical skills to the intensivist. Critical Care, 2013, 17, .  | 5.8 | 0         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 512 | First-hour protocol clarifies resource management and nursing staff education in the ICU. Critical Care, 2013, 17, .                                      | 5.8 | 0         |
| 513 | Influence of critical care nurse education and work environment on outcomes of mechanically ventilated older adults. Critical Care, 2013, 17, .           | 5.8 | 0         |
| 514 | Activity and case-mix changes in a medical ICU after the geographical transfer of a third-level university hospital. Critical Care, 2013, 17, .           | 5.8 | 0         |
| 515 | Are general wards sufficiently staffed to care for level 1 patients?. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 516 | Sleep quality among ICU workers. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 517 | Work-related stress amongst doctors in intensive care, anaesthetics, accident and emergency and general medicine. Critical Care, 2013, 17, .              | 5.8 | 0         |
| 518 | Audit of obstetric critical care admissions in a high-risk population. Critical Care, 2013, 17, .   | 5.8 | 0         |
| 519 | Influence of adverse drug events on the mortality and the length of hospital stay in ICUs in Japan: the JADE Study. Critical Care, 2013, 17, .            | 5.8 | 0         |
| 520 | Attitude and knowledge of intensive care staff concerning donation in Hungary: it is the first step to change. Critical Care, 2013, 17, .                 | 5.8 | 0         |
| 521 | Specialist nurse for organ donation in an emergency department will increase organ donation. Critical Care, 2013, 17, .                                   | 5.8 | 0         |
| 522 | Sepsis in HIV patients admitted to the ICU. Critical Care, 2013, 17, .  | 5.8 | 0         |
| 523 | Mortality related to respiratory failure among pediatric hematology patients requiring intensive care. Critical Care, 2013, 17, .                         | 5.8 | 0         |
| 524 | Survival of critically ill patients with haematological malignancies compared with patients without haematological malignancy. Critical Care, 2013, 17, . | 5.8 | 0         |
| 525 | Outcome of critically ill patients with haematological malignancy admitted to the ICU as an emergency. Critical Care, 2013, 17, .                         | 5.8 | 0         |
| 526 | Effect of organizational structure of the ICU on the prognosis: open format versus semi-closed format. Critical Care, 2013, 17, .                         | 5.8 | 2         |
| 527 | Use of defaults on an electronic prescribing tool influences the type of fluid received by patients. Critical Care, 2013, 17, .                           | 5.8 | 0         |
| 528 | Automated, electronic monitoring for early detection of sepsis. Critical Care, 2013, 17, .  | 5.8 | 2         |
| 529 | Admission to the ICU overnight: is it really a bad thing?. Critical Care, 2013, 17, .   | 5.8 | 0         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 530 | Assessing the quality of interdisciplinary rounds. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 2         |
| 531 | Leadership training and quality improvement of interdisciplinary rounds in the ICU. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 1         |
| 532 | Let's go round again! Quality improvement through intentional rounding. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 533 | Prospective observational study of handover in a medical ICU. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 534 | Impact of knowledge transfer through the implementation of a telemedicine program in a community hospital in Brazil. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 535 | Utilization of telemedicine to improve burn care in a developing country. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 536 | Different aspects of therapy limitation: a comparative study of the nurse's view. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 537 | Alternative to improve palliative care for all patients and families in critical care units: development and preliminary evaluation following MRC guidance of the King's Psychosocial, Assessment and Care tool. <i>Critical Care</i> , 2013, 17, . | 5.8 | 0         |
| 538 | End-of life thoughts in the ICU: results of a survey. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 539 | End-of-life decisions in Slovenian ICUs: a cross-sectional survey. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 540 | Attitudes of intensivists in the UK to withdrawal of futile therapy. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 541 | Parent's perception on end-of-life care in Brazil. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 0         |
| 542 | Getting critical care patients home for end-of-life care. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 543 | Project PIX (Post Intensive care eXercise): impact on physical fitness and focus group analysis of quality of life following exercise rehabilitation. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 544 | Feasibility of neuromuscular electrical stimulation in critically ill patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 2         |
| 545 | Follow-up and counseling service in trauma patients: needs and goals - a preliminary study. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 546 | Post-traumatic stress disorder - are we doing all we can? A UK national-based survey. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 547 | Implementation of an ICU follow-up clinic: outcomes and patient satisfaction after 1 year. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 2         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 548 | Acute exercise response in the critically ill. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 549 | Randomized, controlled pilot study of early rehabilitation strategies in acute respiratory failure. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 6         |
| 550 | Early results of a 6-week exercise programme in post-ICU patients. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 4         |
| 551 | Recognition of the primary stressors affecting intensive care patients: a systematic review. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 552 | Quality of care in the ICU from the perspective of relatives. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 553 | Family satisfaction in the ICU: enhancing patient experience. <i>Critical Care</i> , 2013, 17, .   | 5.8 | 1         |
| 554 | Family satisfaction with patient care in ICUs of nonacademic Brazilian public hospitals. <i>Critical Care</i> , 2013, 17, .  | 5.8 | 0         |
| 555 | Propofol for Procedural Sedation and Analgesia Reduced Dedicated Emergency Nursing Time While Maintaining Safety in a Community Emergency Department. <i>Journal of Emergency Nursing</i> , 2013, 39, 502-507. | 1.0 | 8         |
| 557 | Propofol for adult procedural sedation in a UK emergency department: safety profile in 1008 cases. <i>British Journal of Anaesthesia</i> , 2013, 111, 651-655.   | 3.4 | 45        |
| 558 | An update on pediatric hospital-based sedation. <i>Current Opinion in Pediatrics</i> , 2013, 25, 310-316.  | 2.0 | 21        |
| 559 | In vitro correction of hypothermic and dilutive crystalloid and colloid rotational thromboelastography-monitored coagulopathy with fibrinogen and factor XIII. <i>Critical Care</i> , 2013, 17, .              | 5.8 | 0         |
| 563 | Soft Tissue Trauma and Scar Revision. <i>Facial Plastic Surgery Clinics of North America</i> , 2014, 22, 639-651.  | 1.5 | 16        |
| 564 | Provision of deep procedural sedation by a pediatric sedation team at a freestanding imaging center. <i>Pediatric Radiology</i> , 2014, 44, 1020-1025.   | 2.0 | 22        |
| 565 | The Anesthesia-Directed Sedation Service: Models, Protocols, and Challenges. , 2015, , 227-241.  |     | 0         |
| 566 | Complications of IV sedation for dental treatment in individuals with intellectual disability. <i>Egyptian Journal of Anaesthesia</i> , 2015, 31, 143-148.   | 0.5 | 2         |
| 567 | Propofol Use in Israeli PICUs*. <i>Pediatric Critical Care Medicine</i> , 2016, 17, e117-e120.   | 0.5 | 10        |
| 568 | Paediatric procedural sedation within the emergency department. <i>Journal of Paediatrics and Child Health</i> , 2016, 52, 197-203.  | 0.8 | 9         |
| 569 | Procedural sedation and its place in paediatric emergency medicine. <i>Emergency Medicine Journal</i> , 2016, 33, 525-526.   | 1.0 | 5         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 570 | Anesthesia for Liposuction. , 2016, , 3-49.  |     | 1         |
| 571 | Ketamine and propofol sedation by emergency medicine specialists: mainstream or menace?. British Journal of Anaesthesia, 2016, 116, 449-451.   | 3.4 | 6         |
| 572 | Adverse Events With Ketamine Versus Ketofol for Procedural Sedation on Adults: A Double-blind, Randomized Controlled Trial. Academic Emergency Medicine, 2017, 24, 1441-1449.          | 1.8 | 16        |
| 573 | The Newest Threat to Emergency Department Procedural Sedation. Annals of Emergency Medicine, 2018, 72, 115-119.  | 0.6 | 7         |
| 574 | Unscheduled Procedural Sedation: A Multidisciplinary Consensus Practice Guideline. Annals of Emergency Medicine, 2019, 73, e51-e65.  | 0.6 | 45        |
| 575 | Emergency Department Procedural Sedation Practice Limitations: A Statewide California American College of Emergency Physicians Survey. Academic Emergency Medicine, 2019, 26, 539-548. | 1.8 | 3         |
| 576 | Pediatric Sedation: The European Experience and Approach. , 2021, , 561-586.   |     | 0         |
| 577 | The Anesthesia Directed Sedation Service: Models, Protocols, and Challenges. , 2012, , 179-197.  |     | 0         |
| 578 | Pediatric Sedation: The European Experience and Approach. , 2012, , 351-367.   |     | 0         |
| 579 | Impact of the United States propofol ban on emergency providers' procedural sedation agent choice and patient length of stay. World Journal of Emergency Medicine, 2012, 3, 177.       | 1.0 | 1         |
| 580 | Anesthesia for Aesthetic Surgery. , 2013, , 105-144.   |     | 1         |
| 581 | Pediatric Sedation: The European Experience and Approach. , 2015, , 461-483.   |     | 0         |
| 582 | Pain management mini-series part III. Procedural sedation for the non-anesthesia provider. Missouri Medicine, 2013, 110, 355-60.   | 0.3 | 1         |