

Human umbilical cord blood-derived non-hematopoietic proliferation and CD4, CD8 expression

Journal of Neuroimmunology

197, 99-109

DOI: [10.1016/j.jneuroim.2008.04.013](https://doi.org/10.1016/j.jneuroim.2008.04.013)

Citation Report

#	ARTICLE	IF	CITATIONS
1	The homing of human cord blood stem cells to sites of inflammation. <i>Cell Cycle</i> , 2012, 11, 2303-2313.	1.3	10
2	Pleiotrophin is involved in the amniotic epithelial cell-induced differentiation of human umbilical cord blood-derived mesenchymal stem cells into dopaminergic neuron-like cells. <i>Neuroscience Letters</i> , 2013, 539, 86-91.	1.0	16
4	Effect of Panax ginseng combined with Angelica sinensis on the dissolution of ginsenosides and in chemotherapy mice hematopoietic function. <i>Bioorganic and Medicinal Chemistry</i> , 2019, 27, 4211-4218.	1.4	7
5	An Analysis of Structure-Function Co-relation between GLI Oncoprotein and HLA Immune-gene Transcriptional Regulation through Molecular Docking. <i>Current Cancer Therapy Reviews</i> , 2021, 17, .	0.2	0
6	Possible Therapeutic Effect of Stem Cell in Atherosclerosis in Albino Rats. A Histological and Immunohistochemical Study. <i>International Journal of Stem Cells</i> , 2015, 8, 200-208.	0.8	12