

Detection of PhIP in Grilled Chicken Entrées at Popular Restaurants in California

Nutrition and Cancer

60, 592-602

DOI: 10.1080/01635580801956519

Citation Report

#	ARTICLE	IF	CITATIONS
1	Uptake of the colon carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine by different segments of the rat gastrointestinal tract: Its implication in colorectal carcinogenesis. <i>Toxicology Letters</i> , 2010, 196, 60-66.	0.8	12
2	Factors That Affect the Content of Heterocyclic Aromatic Amines in Foods. <i>Comprehensive Reviews in Food Science and Food Safety</i> , 2011, 10, 52-108.	11.7	215
3	Paper-Based Electrochemiluminescent Screening for Genotoxic Activity in the Environment. <i>Environmental Science & Technology</i> , 2013, 47, 1937-1944.	10.0	74
4	Impact of dextran sulphate sodium-induced colitis on the intestinal transport of the colon carcinogen PhIP. <i>Archives of Toxicology</i> , 2016, 90, 1093-1102.	4.2	6
5	Presence of heterocyclic amine carcinogens in home-cooked and fast-food camel meat burgers commonly consumed in Saudi Arabia. <i>Scientific Reports</i> , 2017, 7, 1707.	3.3	11
6	Reduced cancer risk in vegetarians: an analysis of recent reports. <i>Cancer Management and Research</i> , 2010, 3, 1.	1.9	26