CITATION REPORT List of articles citing

Chronic liver disease--an increasing problem: a study of hospital admission and mortality rates in England, 1979-2005, with particular reference to alcoholic liver disease

DOI: 10.1093/alcalc/agn020 Alcohol and Alcoholism, 2008, 43, 416-22.

Source: https://exaly.com/paper-pdf/44616180/citation-report.pdf

Version: 2024-04-19

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
592	The increasing hospital disease burden of haemochromatosis in England. <i>Alimentary Pharmacology and Therapeutics</i> , 2010 , 31, 247-52	6.1	1
591	Outcomes of critically ill patients with cirrhosis admitted to intensive care: an important perspective from the non-transplant setting. <i>Alimentary Pharmacology and Therapeutics</i> , 2010 , 32, 233-4	43 ^{.1}	41
590	Hospital-related morbidity in people notified with hepatitis C: a population-based record linkage study in New South Wales, Australia. <i>Journal of Hepatology</i> , 2010 , 53, 43-9	13.4	13
589	The epidemiological pattern of chronic liver diseases in a community undergoing voluntary screening for hepatitis B and C. 2011 , 43, 653-8		12
588	Under-reporting of foetal alcohol spectrum disorders: an analysis of hospital episode statistics. 2011 , 11, 14		33
587	Trends in all cause and viral liver disease-related hospitalizations in people with hepatitis B or C: a population-based linkage study. <i>BMC Public Health</i> , 2011 , 11, 52	4.1	15
586	Hepatology training in the UK. Frontline Gastroenterology, 2012 , 3, 52-56	2.6	2
585	Popular intoxicants: what lessons can be learned from the last 40 years of alcohol and cannabis regulation?. 2012 , 26, 213-20		11
5 ⁸ 4	EASL clinical practical guidelines: management of alcoholic liver disease. <i>Journal of Hepatology</i> , 2012 , 57, 399-420	13.4	489
583	Techniques to measure cardiac output: minimally invasive method versus thermodilution. <i>Critical Care</i> , 2012 , 16,	10.8	78
582	Effects of cardiac output levels on the measurement of transpulmonary thermodilution cardiac output in patients with acute lung injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
581	Comparison of bioimpedance and oesophageal Doppler cardiac output monitoring during abdominal aortic surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
580	Cardiac output monitoring in cirrhotic patients: EV1000 versus pulmonary artery catheter - preliminary data. <i>Critical Care</i> , 2012 , 16,	10.8	78
579	Pulse contour cardiac output monitoring is less reliable in critically ill children. <i>Critical Care</i> , 2012 , 16,	10.8	78
578	Impact of arterial catheter location on the accuracy of cardiac output provided by an endotracheal bioimpedance device. <i>Critical Care</i> , 2012 , 16,	10.8	78
577	Accuracy of the PiCCO2-derived pulse contour cardiac index (CIpc): development and validation of a calibration index in two independent collectives. <i>Critical Care</i> , 2012 , 16,	10.8	78
576	Cardiac output monitoring using the LiDCOplusImonitor in abdominal aortic surgery: changes in calibration factor in aortic aneurysm disease versus aortic occlusive disease. <i>Critical Care</i> , 2012 , 16,	10.8	1

575	Haemodynamic changes during the peri-extubation period using bioreactance flow monitoring. <i>Critical Care</i> , 2012 , 16,	10.8	78
574	Left ventricular stroke volume measurement by impedance cardiography correlates with echocardiography in neonates. <i>Critical Care</i> , 2012 , 16,	10.8	2
573	Validation of less-invasive hemodynamic monitoring with Pulsioflex in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	1
572	A preliminary study on the use of noninvasive hemodynamic monitoring with the Nexfin monitor in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
571	Computer-based monitoring of global cardiovascular dynamics during acute pulmonary embolism and septic shock in swine. <i>Critical Care</i> , 2012 , 16,	10.8	78
570	Homeodynamic complexity: multifractal analysis of physiological instability. <i>Critical Care</i> , 2012 , 16,	10.8	78
569	Accuracy of conventional urinary output monitoring in the ICU. Critical Care, 2012, 16,	10.8	2
568	Changes in B-type natriuretic peptide and related hemodynamic parameters following a fluid challenge in patients with severe sepsis or septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
567	The Brody effect to detect hypovolemia in clinical practice. <i>Critical Care</i> , 2012 , 16,	10.8	1
566	Stroke volume variation guided fluid therapy in septic shock with ARDS. <i>Critical Care</i> , 2012 , 16,	10.8	78
565	Assessment fluid responsiveness in septic shock patients: a comparison of automated pulse pressure variation and manually calculated pulse pressure variation. <i>Critical Care</i> , 2012 , 16,	10.8	78
564	Applicability of methods for fluid responsiveness prediction in the ICU. Critical Care, 2012, 16,	10.8	78
563	Fluid responsiveness during weaning from mechanical ventilation. Critical Care, 2012, 16,	10.8	78
562	Microcirculatory blood flow is related to clinical signs of impaired organ perfusion, and its dynamics to the macrohemodynamic concept of fluid responsiveness. <i>Critical Care</i> , 2012 , 16,	10.8	1
561	Frank-Starling and Guyton together at bedside during a fluid challenge. Critical Care, 2012, 16,	10.8	78
560	Prediction of fluid responsiveness in intensive care (PREFERENCE study): fluid challenge versus passive leg raising in high-risk surgical patients. <i>Critical Care</i> , 2012 , 16,	10.8	5
559	Tight control of fluid balance may reduce incidence of intra-abdominal hypertension in patients after major abdominal surgery and trauma: a pilot study. <i>Critical Care</i> , 2012 , 16,	10.8	78
558	Negative fluid balance 48 hours after admission improves survival at 28 days in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78

557	Goal-directed fluid and hemodynamic therapy in major colon surgery with the pressure recording analytical method cardiac output monitor (MostCare[] -PRAM[]): prospective analysis of 58 patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
556	Implementation of an optimal fluid management protocol using the PiCCO system delays development of ARDS secondary to severe sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	1
555	Confrontation of the increase in ELWI rate regarding the septic polytraumatised patient administering furosemide: is it effective?. <i>Critical Care</i> , 2012 , 16,	10.8	78
554	Indexation of extravascular lung water in unselected adult patients with and without mechanical ventilation: a prospective study in 50 patients with 843 transpulmonary thermodilutions. <i>Critical Care</i> , 2012 , 16,	10.8	78
553	How to perform indexing of extravascular lung water data. Critical Care, 2012, 16,	10.8	78
552	Near-normal values of extravascular lung water in children. <i>Critical Care</i> , 2012 , 16,	10.8	78
551	Transthoracic ultrasound assessment of B-lines for identifying the increment of extravascular lung water in shock patients requiring fluid resuscitation. <i>Critical Care</i> , 2012 , 16,	10.8	78
550	Fluid therapy tactics in patients with polytrauma during interhospital transportation. <i>Critical Care</i> , 2012 , 16,	10.8	78
549	A prospective, randomized, clinical trial comparing the hemodynamics, efficacy, and safety of 6% hydroxyethyl starch 130/0.4 compared to albumin in postoperative patients undergoing pancreaticoduodenectomy. <i>Critical Care</i> , 2012 , 16,	10.8	78
548	Effect of balanced versus unbalanced HES solution on cytokine response in a rat model of peritonitis. <i>Critical Care</i> , 2012 , 16,	10.8	78
547	Evaluation of effectiveness and safety of hydroxyethyl starch (HES 130 kDa/0.4) in burn resuscitation. <i>Critical Care</i> , 2012 , 16,	10.8	1
546	Normal saline resuscitation worsens lactic acidosis in experimental sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	4
545	Albumin in early septic shock resuscitation: examination of plasma and urine inflammatory markers. <i>Critical Care</i> , 2012 , 16,	10.8	78
544	Study of the correlation between central venous oxygen saturation and venous saturation from the antecubital vein in severe sepsis/septic shock patients. <i>Critical Care</i> , 2012 , 16,	10.8	1
543	Central venous hyperoxia is related to changes in tissue perfusion and morbi-mortality of patients in shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
542	Curve analysis of tissue oxygen desaturation after a venous occlusion test does not identify the central venous hemoglobin oxygen saturation. <i>Critical Care</i> , 2012 , 16,	10.8	78
541	Lactate in burn patients: biomarker of sepsis and mortality. Critical Care, 2012, 16,	10.8	1
540	Can we predict arterial lactate from venous lactate in the emergency department?. <i>Critical Care</i> , 2012 , 16,	10.8	2

539	Cross-correlation analysis of blood and microdialysis-assessed tissue lactate monitoring: a study in critically ill septic patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
538	Admission lactate and outcome after high-risk surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
537	Effects of induced relative hypoxia during the postoperative period of abdominal oncologic surgery, on hemoglobin and reticulocyte levels: a prospective, randomized controlled clinical trial. <i>Critical Care</i> , 2012 , 16,	10.8	1
536	Pre-emptive hypothermia during resuscitated porcine hemorrhagic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
535	Carbon monoxide therapy protects against hepatic microvascular injury in a mouse model of murine hemorrhagic shock and resuscitation. <i>Critical Care</i> , 2012 , 16,	10.8	78
534	Customized modeling to predict the use of vasopressors in ICUs. Critical Care, 2012, 16,	10.8	78
533	Implementation of the fifth link of the Chain of Survival concept for out-of-hospital cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
532	Examination of out-of-hospital cardiac arrest patients with the Utstein style in Saga prefecture, Japan. <i>Critical Care</i> , 2012 , 16,	10.8	1
531	Effectiveness and limitations of learning cardiopulmonary resuscitation with an automated external defibrillator in the curriculum of First Aid courses among lay people. <i>Critical Care</i> , 2012 , 16,	10.8	78
530	Survival after out-of-hospital cardiac arrest during nights and weekends. <i>Critical Care</i> , 2012 , 16,	10.8	78
529	CPR initiated after telephone-assisted instruction produces a better outcome of bystander-witnessed out-of-hospital cardiac arrests than no bystander CPR but is less effective than CPR on the bystanderß own initiative. <i>Critical Care</i> , 2012 , 16,	10.8	78
528	Critical times in pediatric out-of-hospital cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	1
527	DonR stop your heart in front of your family: family as a bystander is associated with poor outcome of bystander-witnessed, bystander-CPR-performed out-of-hospital cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
526	Coronary perfusion pressure in a pig model of prolonged cardiac arrest treated by different modes of venoarterial extracorporeal membrane oxygenation and intraaortic balloon counterpulsation. <i>Critical Care</i> , 2012 , 16,	10.8	3
525	Capnometry successfully predicts outcome and determination of the cessation of cardiopulmonary resuscitation efforts. <i>Critical Care</i> , 2012 , 16,	10.8	1
524	Modified clinical decision rule for termination-of-resuscitation in cases of refractory out-of-hospital cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
523	Survival benefit for patients receiving antibiotics following out-of-hospital cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
522	Correlation between IL-6 and S-100B blood levels and outcome of post-cardiac arrest syndrome and influence of therapeutic hypothermia on these mediator blood levels. <i>Critical Care</i> , 2012 , 16,	10.8	78

521	Early neurological outcome prediction model after bystander-witnessed out-of-hospital cardiac arrest: a nationwide population-based study. <i>Critical Care</i> , 2012 , 16,	10.8	78
520	Helium ventilation is safe and feasible in ICU patients admitted after cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	2
519	Therapeutic hypothermia in an out-of-hospital arrest population: are we selecting appropriately?. <i>Critical Care</i> , 2012 , 16,	10.8	78
518	Therapeutic hypothermia for nonventricular fibrillation/ventricular tachycardia cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
517	Comparison of cold crystalloid and colloid infusions for induction of therapeutic hypothermia. <i>Critical Care</i> , 2012 , 16,	10.8	78
516	Cerebral oxygenation during induction of therapeutic hypothermia after cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
515	Survey on the management of patients treated with therapeutic hypothermia post cardiac arrest in London hospitals. <i>Critical Care</i> , 2012 , 16,	10.8	78
514	Simplified EEG/aEEG to monitor the injured brain after cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
513	Usefulness of a Bispectral index oriented sedative method without neuromuscular blocker for therapeutic hypothermia after cardiac arrest. <i>Critical Care</i> , 2012 , 16,	10.8	78
512	Predictive factors of neurologic outcome in therapeutic hypothermia after prehospital return of spontaneous circulation. <i>Critical Care</i> , 2012 , 16,	10.8	78
511	Employment status 1 year after out-of-hospital cardiac arrest in comatose patients treated with therapeutic hypothermia. <i>Critical Care</i> , 2012 , 16,	10.8	78
510	Changes in cerebrospinal fluid and serum cytokine levels in severe traumatic brain injury patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
509	Noninvasive cerebral oxygenation monitoring during rapid ventricular pacing in transcutaneous aortic valve implant. <i>Critical Care</i> , 2012 , 16,	10.8	78
508	Novel models to predict elevated intracranial pressure during intensive care and long-term neurological outcome after TBI. <i>Critical Care</i> , 2012 , 16,	10.8	78
507	Transcranial Doppler pulsatility index is a poor predictor of hydrocephalus in patients with aneurysmal subarachnoid haemorrhage. <i>Critical Care</i> , 2012 , 16,	10.8	78
506	Transcranial cerebral oximetry in newborn infants on mechanical ventilation as a method for prevention of hyperoxia and oxidative stress. <i>Critical Care</i> , 2012 , 16,	10.8	78
505	Cerebral oximetry and brain death in the ICU: data from seven cases. Critical Care, 2012, 16,	10.8	2
504	Deoxyhaemoglobin as a biomarker of cerebral autoregulation. <i>Critical Care</i> , 2012 , 16,	10.8	3

503	Study of the acoustic stem evoked potentials in blood circulation disorder in the vertebral basilar basin. <i>Critical Care</i> , 2012 , 16,	10.8	78
502	Examination of the autonomic nervous system in the ICU: a pilot study. Critical Care, 2012, 16,	10.8	78
501	Predictive value of glial fibrillary acidic protein for prognosis in patients with moderate and severe traumatic brain injury: a systematic review and meta-analysis. <i>Critical Care</i> , 2012 , 16,	10.8	3
500	Prevalence of pituitary disorders associated with traumatic brain injury: a systematic review. <i>Critical Care</i> , 2012 , 16,	10.8	2
499	Mannose binding lectin deficiency attenuates neurobehavioral deficits following experimental traumatic brain injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
498	Azathioprine and aspirin in treatment of childhood primary arterial stroke: therapeutic benefits and side effects. <i>Critical Care</i> , 2012 , 16,	10.8	78
497	Changes of ribosomal protein S3 immunoreactivity and its new expression in microglia in the mice hippocampus after lipopolysaccharide treatment. <i>Critical Care</i> , 2012 , 16,	10.8	78
496	Neuronal damage using Fluoro-Jade B histofluorescence and gliosis in the striatum after various durations of transient cerebral ischemia in gerbils. <i>Critical Care</i> , 2012 , 16,	10.8	3
495	Molecular, histological and microcirculatory modeling of cerebral ischemia in pigs. <i>Critical Care</i> , 2012 , 16,	10.8	78
494	Delayed post-ischaemic administration of xenon reduces brain damage in a rat model of global ischaemia. <i>Critical Care</i> , 2012 , 16,	10.8	78
493	Seizures in the respiratory ICU: single-center study of patients with new-onset seizures. <i>Critical Care</i> , 2012 , 16,	10.8	1
492	Early treatment with intravenous immunoglobulins in patients with critical illness polyneuropathy: a randomized controlled, double-blinded study. <i>Critical Care</i> , 2012 , 16,	10.8	78
491	Intracranial pressure monitoring in acute liver failure: a retrospective cohort study. <i>Critical Care</i> , 2012 , 16,	10.8	2
490	Retrospective observation of 6-month survival following decompressive craniectomy in a London major trauma and stroke centre. <i>Critical Care</i> , 2012 , 16,	10.8	1
489	Feasibility of a multicenter prospective cohort study on the evaluation of prognosis in severe traumatic brain injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
488	Predictive value of neuron-specific enolase following moderate and severe traumatic brain injury: a systematic review and meta-analysis. <i>Critical Care</i> , 2012 , 16,	10.8	78
487	Blood-brain barrier permeability following traumatic brain injury. Critical Care, 2012, 16,	10.8	1
486	Can urinary 8-OHdG be a good indicator of vasospasm occurrence following subarachnoid hemorrhage?. <i>Critical Care</i> , 2012 , 16,	10.8	8

485	Cortical capillary recruitment by rosuvastatin in experimental brain trauma is associated with increased NO production. <i>Critical Care</i> , 2012 , 16,	10.8	78
484	Effects of sinvastatin in prevention of vasospasm in nontraumatic subarachnoid hemorrhage: preliminary data. <i>Critical Care</i> , 2012 , 16,	10.8	1
483	Evaluation of arterial and venous ophthalmic hemodynamics in preeclamptic pregnant women. <i>Critical Care</i> , 2012 , 16,	10.8	78
482	Data classification of magnetic resonance tomography and computer tomography images of brain in parturients with neurological complications of eclampsia. <i>Critical Care</i> , 2012 , 16,	10.8	78
481	Eleven years of critical obstetric pathology: epidemiologic study. Critical Care, 2012, 16,	10.8	78
480	Clinical outcomes in neonates following maternal magnesium sulfate therapy in preeclampsia/eclampsia. <i>Critical Care</i> , 2012 , 16,	10.8	78
479	Sleep monitoring by actigraphy in short-stay ICU patients. Critical Care, 2012, 16,	10.8	1
478	Quality and quantity of sleep in multipatient versus single-room ICUs. Critical Care, 2012, 16,	10.8	78
477	Oral melatonin in high-risk critically ill patients: quality of sedative effect. Critical Care, 2012, 16,	10.8	1
476	Sedation depth and mortality in mechanically ventilated critically ill adults. Critical Care, 2012, 16,	10.8	1
475	Sedation in the ICU: nursesPperceptions of practices and influencing factors. <i>Critical Care</i> , 2012 , 16,	10.8	78
474	Implementation of a national guideline for analgesia and sedation: how often can a RASS of 0 to -2 be achieved?. <i>Critical Care</i> , 2012 , 16,	10.8	78
473	Comparison of the RAMSAY score and the Richmond Agitation Sedation Score for the measurement of sedation depth. <i>Critical Care</i> , 2012 , 16,	10.8	12
472	Dexmedetomidine is associated with better outcomes in patients undergoing cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
471	Cerebral ischemia-reperfusion model in rabbits: relationship between dexmedetomidine and biochemical parameters in lowering intraparenchymal pressure. <i>Critical Care</i> , 2012 , 16,	10.8	78
470	Evaluation of sedation using pupilometry in ICUs: a pilot study. Critical Care, 2012, 16,	10.8	78
469	Effect of critical illness on the pharmacokinetics and dose-response relationship of midazolam. <i>Critical Care</i> , 2012 , 16,	10.8	1
468	Effect of propofol and midazolan on microcirculation of septic shock patients. <i>Critical Care</i> , 2012 , 16,	10.8	78

467	Current use of pain scores in Dutch ICUs: a postal survey in the Netherlands. Critical Care, 2012, 16,	10.8	78
466	Efficiency estimation of intrapleural and thoracic paravertebral block in combination with general anesthesia at thoracoscopic interventions. <i>Critical Care</i> , 2012 , 16,	10.8	78
465	Preoperative diclofenc reduces postcraniotomy headache: a randomized, placebo-controlled trial. <i>Critical Care</i> , 2012 , 16,	10.8	78
464	Long-term adverse neuropsychological functioning in children who survived meningococcal septic shock: is there a relationship with sedation and analgesia during paediatric ICU admission?. <i>Critical Care</i> , 2012 , 16,	10.8	78
463	Delirium could be an indicator of sepsis in patients under 65 years old with urinary tract infections. <i>Critical Care</i> , 2012 , 16,	10.8	78
462	Delirium screening in critically ill patients: a systematic review and meta-analysis. <i>Critical Care</i> , 2012 , 16,	10.8	3
461	Electroencephalography-based monitoring of delirium in the ICU: what are the opportunities?. <i>Critical Care</i> , 2012 , 16,	10.8	0
460	Performance of SAPS 3 in predicting delirium among critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
459	Incidence of delirium and inadequacy of the clinical diagnosis in patients in intensive care. <i>Critical Care</i> , 2012 , 16,	10.8	78
458	Investigation into detection and treatment rates of hyperactive and hypoactive delirium in the ICU setting. <i>Critical Care</i> , 2012 , 16,	10.8	78
457	Memories and post-traumatic stress-related symptoms in older, post-cardiac surgery patients: substudy of an RCT. <i>Critical Care</i> , 2012 , 16,	10.8	78
456	Using tramadol to monitor hepatic drug metabolism in the critically ill. Critical Care, 2012, 16,	10.8	78
455	Data mining techniques for predicting acute kidney injury after elective cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	3
454	Acute kidney injury in critically ill patients with A/H1N1 pneumonitis in 2010/11. <i>Critical Care</i> , 2012 , 16,	10.8	78
453	A RIFLE score-based trigger for renal replacement therapy and survival after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
452	Effect of off-pump versus on-pump coronary artery bypass grafting in patients with chronic kidney disease. <i>Critical Care</i> , 2012 , 16,	10.8	1
451	Effects of renal-dose dopamine on renal tubular functions following coronary artery bypass grafting surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
450	NursesPknowledge regarding the early identification of acute kidney injury. Critical Care, 2012, 16,	10.8	78

449	Neutrophil gelatinase-associated lipocalin predicts postoperative fluid overload, a potentially modifiable risk factor for mortality after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	1
448	Plasma and urine neutrophil gelatinase-associated lipocalin as markers of acute kidney injury in critically ill adults. <i>Critical Care</i> , 2012 , 16,	10.8	78
447	Plasma and urine neutrophil gelatinase-associated lipocalin in septic and nonseptic ICU patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
446	Urinary neutrophil gelatinase-associated lipocalin as an early marker of acute kidney injury complicating circulatory shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
445	Additive value to clinical judgement of blood neutrophil gelatinase-associated lipocalin in diagnosis of acute kidney injury and prediction of mortality in patients hospitalized from the emergency department. <i>Critical Care</i> , 2012 , 16,	10.8	2
444	Is cystatin C reliable in the anesthetized pig? An experimental study with special reference to septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	1
443	Are serum cystatin-C-based estimates better than those derived from serum creatinine in critically ill patients?. <i>Critical Care</i> , 2012 , 16,	10.8	2
442	Assessment of glomerular filtration rate in trauma patients in early resuscitation phase. <i>Critical Care</i> , 2012 , 16,	10.8	1
441	Validation of a continuous low-dose iohexol infusion to measure the glomerular filtration rate. <i>Critical Care</i> , 2012 , 16,	10.8	78
440	Investigation into the effects of commencing haemodialysis in the critically ill. <i>Critical Care</i> , 2012 , 16,	10.8	78
439	Is the AKIN score useful as an indicator of the optimum time for intervention with renal replacement therapy in critically ill patients?. <i>Critical Care</i> , 2012 , 16,	10.8	78
438	Timing for initiation of continuous renal replacement therapy in patients with septic shock and acute kidney injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
437	Early application of CVVH In the complex treatment of patients with early severe acute pancreatitis. <i>Critical Care</i> , 2012 , 16,	10.8	1
436	Timing of the initiation of continuous renal replacement therapy and clinical outcome in patients with severe sepsis and septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
435	Amino acid concentrations in serum, urine and dialysate/ultrafiltrate solutions of continuous venovenous hemodiafiltration patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
434	Evaluation of the potential adverse effects associated with calcium carbonate precipitate during continuous venovenous hemofiltration. <i>Critical Care</i> , 2012 , 16,	10.8	78
433	Regional citrate anticoagulation in CVVH: a new protocol combining citrate solution with a phosphate-containing replacement fluid. <i>Critical Care</i> , 2012 , 16,	10.8	78
432	Regional citrate anticoagulation with a low-concentration solution in predilution-postdilution CVVH. <i>Critical Care</i> , 2012 , 16,	10.8	78

431	Exposure to intermittent hemodialysis and renal recovery after acute kidney injury: a systematic review. <i>Critical Care</i> , 2012 , 16,	10.8	2
430	Sustained low-efficiency dialysis for renal replacement therapy in the ICU: a cost-benefit analysis of the years 2006 to 2010. <i>Critical Care</i> , 2012 , 16,	10.8	78
429	The new dialysis method Mini-SLED is useful for dialyzing acute brain stroke patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
428	Investigation into haemodynamic stability during intermittent haemodialysis in the critically ill. <i>Critical Care</i> , 2012 , 16,	10.8	78
427	Evaluation of microcirculation before and during continuous renal replacement therapy and the impact of dose prescription. <i>Critical Care</i> , 2012 , 16,	10.8	78
426	Ultrafiltration during continuous hemofiltration in stabilized ICU patients is not associated with microcirculatory perfusion derangements. <i>Critical Care</i> , 2012 , 16,	10.8	78
425	Plasmapheresis without apparatus in complex care of victims with crush syndrome during the first hours after extrication in a field hospital of EMERCOM of Russia in emergency areas. <i>Critical Care</i> , 2012 , 16,	10.8	78
424	Degree of impaired kidney function at hospital discharge has a major impact on long-term survival of critically ill patients recovered from renal failure. <i>Critical Care</i> , 2012 , 16,	10.8	78
423	Long-term survival for ICU patients with acute kidney injury. Critical Care, 2012, 16,	10.8	78
422	Super high-flux continuous hemodialysis: an efficient compromise for blood purification in sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	78
421	Efficacy of continuous haemodiafiltration using a polymethylmethacrylate membrane haemofilter in the treatment of sepsis and acute respiratory distress syndrome. <i>Critical Care</i> , 2012 , 16,	10.8	1
420	Possible adsorption mechanism of high mobility group box 1 protein on a polyacrylonitrile (AN69ST) membrane filter. <i>Critical Care</i> , 2012 , 16,	10.8	78
419	High mobility group box 1 levels in septic disseminated intravascular coagulation patients undergoing Polymixin-B immobilized fiber-direct hemoperfusion. <i>Critical Care</i> , 2012 , 16,	10.8	78
418	Polymyxin B-immobilized fiber column hemoperfusion has the ability of endotoxin removal during 24 hours. <i>Critical Care</i> , 2012 , 16,	10.8	78
417	Polymyxin B-direct hemoperfusion therapy could contribute to hemodynamics and outcomes in emergency surgical patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
416	Clinical impact of enhanced cytokine clearance with sustained high-efficiency daily diafiltration using a mediator-adsorbing membrane (SHEDD-fA) in patients with severe sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	78
415	Mortality and priority level for ICU admission in the setting of limited critical care beds in El Salvador. <i>Critical Care</i> , 2012 , 16,	10.8	1
414	Mainz Emergency Evaluation Scoring in combination with capnometry predicts outcome in trauma patients. <i>Critical Care</i> , 2012 , 16,	10.8	78

413	Predicting outcome in the ICU: comparison of Ranson criteria and Ranson + CRP levels in acute pancreatitis. <i>Critical Care</i> , 2012 , 16,	10.8	1
412	Number of failed organs and response to therapy determine outcome in patients with acute pancreatitis requiring level 1 organ support. <i>Critical Care</i> , 2012 , 16,	10.8	78
411	Mortality predictors in acute pancreatitis admitted to the ICU. Critical Care, 2012, 16,	10.8	78
410	System biology prediction model based on clinical data: highly accurate outcome prediction in patients with acute-on-chronic liver failure. <i>Critical Care</i> , 2012 , 16,	10.8	78
409	Prognostic relevance of arterial ammonia levels in different acute and acute-on-chronic liver diseases. <i>Critical Care</i> , 2012 , 16,	10.8	1
408	Liver failure secondary to alcoholic liver disease carries a worse prognosis than other aetiologies of liver failure: retrospective analysis of routine biochemical markers in critically ill patients with liver failure. <i>Critical Care</i> , 2012 , 16,	10.8	1
407	Incidence, morbidity and mortality of admissions related to alcohol consumption on critical care: a single-centre experience. <i>Critical Care</i> , 2012 , 16,	10.8	1
406	Changing outcomes in patients with chronic liver disease in intensive care: a decade of experience. <i>Critical Care</i> , 2012 , 16,	10.8	2
405	Multivariate regression analysis of outcomes following orthotopic liver transplantation in decompensated cirrhotics transplanted from the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
404	Liver transplantation in the critically ill: a Canadian collaboration. <i>Critical Care</i> , 2012 , 16,	10.8	78
403	Acute respiratory distress syndrome: analysis of incidence and mortality in a university hospital critical care unit. <i>Critical Care</i> , 2012 , 16,	10.8	2
402	Epidemiology and outcome of sepsis syndromes in Italian ICUs: a regional multicenter observational cohort. <i>Critical Care</i> , 2012 , 16,	10.8	2
401	Outcome of faecal peritonitis in the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	2
400	HIV patients in the ICU: our experience. <i>Critical Care</i> , 2012 , 16,	10.8	78
399	Impact of congestive heart failure on severe sepsis and septic shock survivors: outcomes and performance status after 1-year hospital discharge. <i>Critical Care</i> , 2012 , 16,	10.8	3
398	Predictive value of N-terminal pro-brain natriuretic peptide among critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
397	Low preoperative hepcidin concentration is a risk factor for mortality but not for acute kidney injury after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
396	Outcomes and resource use for over 80 year olds admitted to a UK critical care unit after an emergency laparotomy over a 3-year period. <i>Critical Care</i> , 2012 , 16,	10.8	78

395	Correlation between APACHE II score and quality of life among patients discharged from the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
394	Parameters that affect outcome in surgical ICU patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
393	Relationship between illness severity scores in the ICU. Critical Care, 2012, 16,	10.8	78
392	Predicting hospital mortality: comparing accuracy of SAPS II and clinical staff prognosis. <i>Critical Care</i> , 2012 , 16,	10.8	78
391	Predictors of mortality in patients from a hematological ICU in Brazil. Critical Care, 2012, 16,	10.8	78
390	Retrospective study of the outcomes of patients admitted to the ICU with a hematological malignancy. <i>Critical Care</i> , 2012 , 16,	10.8	78
389	Six-month survival of patients with lung cancer admitted to a medical ICU: a retrospective study. <i>Critical Care</i> , 2012 , 16,	10.8	78
388	Health-related quality of life and survival of cancer patients admitted to ICUs: Results of the QALY study. <i>Critical Care</i> , 2012 , 16,	10.8	1
387	Characteristics, resource consumption and outcome of cancer patients admitted to ICUs. <i>Critical Care</i> , 2012 , 16,	10.8	78
386	Managing critically ill oncological patients in hospital: a survey across all ICUs in the UK. <i>Critical Care</i> , 2012 , 16,	10.8	78
385	Role of illness severity scores in predicting mortality in the coronary care unit. <i>Critical Care</i> , 2012 , 16,	10.8	78
384	New severity score of acute respiratory failure. <i>Critical Care</i> , 2012 , 16,	10.8	78
383	Validity of six prognostic scoring systems for septic shock patients admitted to a medical ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
382	Risk factors of venous thrombosis in knee joint endoprosthesis. <i>Critical Care</i> , 2012 , 16,	10.8	78
381	Proximal and distal deep venous thrombosis in critically ill patients: incidence and prevalence. <i>Critical Care</i> , 2012 , 16,	10.8	78
380	Saddle embolism is associated with the major adverse events in patients with nonhigh-risk pulmonary embolism. <i>Critical Care</i> , 2012 , 16,	10.8	1
379	Efficacy and safety of enoxaparin as deep vein thrombosis prophylaxis in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
378	Reducing the level of postoperative thrombotic complications by the combination of low molecular weight heparin and epidural anesthesia at the patients after total hysterectomy. <i>Critical Care</i> , 2012 , 16,	10.8	78

377	Cost-effectiveness analysis of two thromboprophylactic strategies following major surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
376	Consequences of suspected heparin-induced thrombocytopenia in the ICU. Critical Care, 2012, 16,	10.8	78
375	Evaluation of iron, transferrin and ferritin serum levels in patients with severe sepsis and septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	2
374	Using angiogenic factors and their soluble receptors to predict organ dysfunction in patients with disseminated intravascular coagulation associated with severe trauma. <i>Critical Care</i> , 2012 , 16,	10.8	78
373	A simple blood-saving bundle reduces diagnostic blood loss in mechanically ventilated patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
372	Comparative assessment of invasive and noninvasive Methods for detection of total hemoglobin in gynecological patients Polood. <i>Critical Care</i> , 2012 , 16,	10.8	4
371	Use of coagulation screening in the critical care unit. Critical Care, 2012, 16,	10.8	0
370	Templating effect of clot structure can predict clot development and outcome in diluted blood: a comparison with thromboelastography. <i>Critical Care</i> , 2012 , 16,	10.8	5
369	Fractal analysis: a new biomarker for determining clot characteristics in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	1
368	Fractal dimension: a biomarker for detecting acute thromboembolic disease. <i>Critical Care</i> , 2012 , 16,	10.8	1
367	Thromboelastography (platelet contribution to clot strength) for the assessment of platelet residual function. <i>Critical Care</i> , 2012 , 16,	10.8	78
366	Measurement of hemoglobin in the operating room: what Methods can we trust?. <i>Critical Care</i> , 2012 , 16,	10.8	78
365	Retrospective comparison study of warfarinised trauma patients and an age-matched control group of nonwarfarinised patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
364	In medical-surgical ICU patients, major bleeding is common but independent of heparin prophylaxis. <i>Critical Care</i> , 2012 , 16,	10.8	2
363	Reducing ICU blood draws with artificial intelligence. <i>Critical Care</i> , 2012 , 16,	10.8	78
362	Hemostasis system condition in infectious complication development in severe burned patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
361	Randomized comparison of fibrinogen concentrate versus cryoprecipitate for bleeding control in pediatric cardiac surgery (FICCS study). <i>Critical Care</i> , 2012 , 16,	10.8	5
360	Efficacy of tranexamic acid in decreasing blood loss during cesarean section. <i>Critical Care</i> , 2012 , 16,	10.8	2

359	Reduced EPO receptor expression may contribute to limited pleiotropic effects of EPO during critical illness. <i>Critical Care</i> , 2012 , 16,	10.8	1
358	Recognition and management of haemophagocytic lymphohistiocytosis on the ICU: a case series. <i>Critical Care</i> , 2012 , 16,	10.8	78
357	Blood transfusion is an independent predicting factor for poor outcome after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
356	Red blood transfusion is a predictor of poor outcome in pediatric cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	1
355	No impact of a massive transfusion protocol on coagulopathy and mortality at a level 1 trauma center: why?. <i>Critical Care</i> , 2012 , 16,	10.8	78
354	Massive transfusion practice. <i>Critical Care</i> , 2012 , 16,	10.8	78
353	Red blood cell transfusion improves microdialysis-assessed interstitial lactate/pyruvate ratio in critically ill septic patients. <i>Critical Care</i> , 2012 , 16,	10.8	1
352	Blood transfusion after cardiac surgery increases the hospital length of stay in adult patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
351	Transfusion of blood stored for longer periods of time does not alter the reactive hyperemia index in healthy volunteers. <i>Critical Care</i> , 2012 , 16,	10.8	0
350	Liberal use of platelet transfusions in the acute phase of trauma resuscitation: a systematic review. <i>Critical Care</i> , 2012 , 16,	10.8	78
349	Impact on early trauma mortality of the adoption of the Updated European Guidelines on the management of bleeding. <i>Critical Care</i> , 2012 , 16,	10.8	78
348	Hemodynamics in the severely injured patient with significant hemorrhage. Critical Care, 2012, 16,	10.8	78
347	Critical older trauma patients. Critical Care, 2012, 16,	10.8	78
346	Outcomes in older blunt chest wall trauma patients: a retrospective study. <i>Critical Care</i> , 2012 , 16,	10.8	2
345	Mortality trend alteration of thoracic injury after rapid response trauma team establishment. <i>Critical Care</i> , 2012 , 16,	10.8	78
344	Trauma patients and cervical spine protection in critical care: the impact of a spinal checklist on clinical care and documentation. <i>Critical Care</i> , 2012 , 16,	10.8	78
343	Effect of instrumented spinal fixation on outcome in polytrauma patients in the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
342	Whole body computed tomography scanning for severe blunt polytrauma: analysis of Trauma Audit and Research Network database 2005 to 2010. <i>Critical Care</i> , 2012 , 16,	10.8	1

341	Post-traumatic rhabdomyolysis: an observational study in seven patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
340	Exertional rhabdomyolysis in female amateur triathletes. <i>Critical Care</i> , 2012 , 16,	10.8	1
339	Controlled mechanical ventilation tactics in patients with polytrauma during interhospital transportation to the specialized center. <i>Critical Care</i> , 2012 , 16,	10.8	11
338	Impact of fluid resuscitation volume on the severity of organ failures in severely burned patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
337	Organ dysfunction in the resuscitation phase of critical burn patients. Critical Care, 2012, 16,	10.8	78
336	Epidemiological study of critical burn patients in an ICU. Critical Care, 2012, 16,	10.8	1
335	Fluid creep in burn resuscitation: the tide has not yet turned. Critical Care, 2012, 16,	10.8	5
334	Early administration of parenteral estrogen suppresses the deleterious local and systemic inflammatory response in severe burns. <i>Critical Care</i> , 2012 , 16,	10.8	78
333	Reducing the indication of ventilatory support in the severely burnt patient and improving outcomes: Results of a new protocol approach within a regional burns centre. <i>Critical Care</i> , 2012 , 16,	10.8	78
332	Cardiopulmonary exercise testing and elective open abdominal aortic aneurysm surgery over a 6-year period in a UK teaching hospital. <i>Critical Care</i> , 2012 , 16,	10.8	1
331	Perioperative evaluation of elective surgical patients: is it possible to plan ICU admission?. <i>Critical Care</i> , 2012 , 16,	10.8	78
330	Cardiac-specific biomarkers and life-threatening complications of off-pump versus on-pump coronary bypass surgery in Egyptian patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
329	Aortic aneurysm disease versus aortic occlusive disease: differences in postoperative ICU requirements after open elective abdominal aortic surgery. <i>Critical Care</i> , 2012 , 16,	10.8	1
328	High postoperative blood levels of macrophage migration inhibitory factor are associated with less organ dysfunction in patients after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
327	Predictors of prolonged mechanical ventilation after heart transplantation. Critical Care, 2012, 16,	10.8	78
326	Atrial fibrillation following major noncardiac thoracic surgery: significance and impact on morbidity. <i>Critical Care</i> , 2012 , 16,	10.8	1
325	Oxygen delivery index during goal-directed therapy predicts complications and hospital length of stay in patients undergoing high-risk surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
324	Transfer delays in patients referred for neurosurgical intervention with traumatic brain injury. <i>Critical Care</i> , 2012 , 16,	10.8	1

323	Performances of ventilator at simulated altitude. Critical Care, 2012, 16,	10.8	78
322	Impact of H1N1 influenza on critical care and dependent services in Wales during winter 2010/2011. <i>Critical Care</i> , 2012 , 16,	10.8	78
321	Effects of levels of clinical supervision during simulated ICU scenarios on resident learning and patient care: a qualitative study. <i>Critical Care</i> , 2012 , 16,	10.8	78
320	Virtual reality and live scenario simulation: options for training medical students in mass casualty incident triage. <i>Critical Care</i> , 2012 , 16,	10.8	3
319	Utilization of iPad in the system of emergency demand and acceptance. Critical Care, 2012, 16,	10.8	78
318	Mass evacuation of victims from emergency areas by medical modules aboard the aircraft of EMERCOM of Russia. <i>Critical Care</i> , 2012 , 16,	10.8	78
317	Reliability and validity of an Italian four-level emergency triage system. Critical Care, 2012, 16,	10.8	78
316	Coordination of emergency resources after Lorcaß earthquakes. Critical Care, 2012, 16,	10.8	78
315	Lightning injuries in a lightning city: a district hospital experience in Singapore. <i>Critical Care</i> , 2012 , 16,	10.8	78
314	Satisfaction survey among medical staff involved in relief operations following the Great East Japan Earthquake and Tsunami. <i>Critical Care</i> , 2012 , 16,	10.8	78
313	Nuclear disaster and the medical problems during the earthquake in Japan, 2011. <i>Critical Care</i> , 2012 , 16,	10.8	78
312	Stressors in the ICU: different perceptions of patients, relatives and staff members. <i>Critical Care</i> , 2012 , 16,	10.8	78
311	Role of ICU nurses in the confrontation of post-traumatic stress disorder in relatives of ICU patients in a general hospital of Athens, Greece. <i>Critical Care</i> , 2012 , 16,	10.8	78
310	Family meetings and end-of-life decision-making in Thai critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
309	Influence of burnout on attitudes of ICU doctors and nurses towards liberalization of visiting polices. <i>Critical Care</i> , 2012 , 16,	10.8	78
308	Prevalence, risk factors and impact of severe burnout syndrome in 12 Uruguayan ICUs. <i>Critical Care</i> , 2012 , 16,	10.8	78
307	Opening the ICU: views of ICU doctors and nurses before and after liberalization of visiting policies. Critical Care, 2012 , 16,	10.8	3
306	A family-based satisfaction survey on the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	1

305	Immediate needs and level of anxiety of families with traumatic brain injury patients admitted to ICUs. <i>Critical Care</i> , 2012 , 16,	10.8	78
304	Families: the newest members of the ICU multidisciplinary team. Critical Care, 2012, 16,	10.8	78
303	Family satisfaction in an interdisciplinary ICU: a quality audit. Critical Care, 2012, 16,	10.8	78
302	Incidence of post-traumatic stress, anxiety and depression symptoms in patients and relatives during the ICU stay and after discharge. <i>Critical Care</i> , 2012 , 16,	10.8	3
301	Application of a new German law as a basis for end-of life decisions in a medical ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
300	Effect of a full moon on mortality of patients admitted to the ICU. Critical Care, 2012, 16,	10.8	78
299	Potential association of gender with mortality and withdrawal of life-sustaining therapies in patients with severe TBI: a Canadian multicentre cohort study. <i>Critical Care</i> , 2012 , 16,	10.8	78
298	Making drug delivery in the ICU safer: the implementation of advanced computerised intravenous infusion pumps. <i>Critical Care</i> , 2012 , 16,	10.8	1
297	Growing a positive culture in an ICU antimicrobial stewardship program. Critical Care, 2012, 16,	10.8	78
296	Injectional anthrax: the inflammatory response. Critical Care, 2012, 16,	10.8	78
295	Multicenter consensus development of a checklist for lung injury prevention. Critical Care, 2012, 16,	10.8	1
294	Impact of the Paediatric Intensive Care Outreach Network service on mortality within intensive care. <i>Critical Care</i> , 2012 , 16,	10.8	78
293	In-hospital rapid response system: effects on outcome and workload. Critical Care, 2012, 16,	10.8	1
292	Medical emergency team admittance to intensive care versus conventional admittance: characteristics and outcome. <i>Critical Care</i> , 2012 , 16,	10.8	78
291	Factors affecting critical care admission to a UK university hospital. Critical Care, 2012, 16,	10.8	1
290	Intensive care services in Hungary 2000 to 2010: an analysis of bed numbers, occupancy rates, case mix and economics. <i>Critical Care</i> , 2012 , 16,	10.8	78
289	Data acquisition for the UK Critical Care Minimum Data Set: validation of a computer model for automatic calculation from an electronic patient record. <i>Critical Care</i> , 2012 , 16,	10.8	78
288	To admit or not to admit? The suitability of critical care admission criteria. <i>Critical Care</i> , 2012 , 16,	10.8	2

287	Out-of-hours discharge from the ICU: defining the out-of-hours period and its effect on mortality. <i>Critical Care</i> , 2012 , 16,	10.8	3
286	Delayed discharges revisited: impact of a liaison post on patientsPtransition from ICU to ward care. <i>Critical Care</i> , 2012 , 16,	10.8	78
285	Assessing demand for intensive care services: the role of readmission rates. Critical Care, 2012, 16,	10.8	78
284	Management of cardiac drugs in a critical care setting. <i>Critical Care</i> , 2012 , 16,	10.8	78
283	Pharmacists and fastidiousness improve compliance with guidelines for stress ulcer prophylaxis. <i>Critical Care</i> , 2012 , 16,	10.8	78
282	Healthcare workersPexperience when using an electronic medical order entry and bar-code technology in an ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
281	Safer ICU trainee handover: a service improvement project. Critical Care, 2012, 16,	10.8	1
280	ICU handover: are we forgetting something? A preliminary study. Critical Care, 2012, 16,	10.8	1
279	Quality and value of intensive care discharge summaries for general practitioners. <i>Critical Care</i> , 2012 , 16,	10.8	3
278	Volume-outcome relationship in critical care: a systematic review. <i>Critical Care</i> , 2012 , 16,	10.8	78
277	Radiation doses in young ICU patients: a cause for concern?. Critical Care, 2012, 16,	10.8	1
276	Accuracy of height and weight estimation by critical care staff. Critical Care, 2012, 16,	10.8	78
275	Implementation of evidence-based care bundles in the ICU. Critical Care, 2012, 16,	10.8	1
274	Awareness of difficult airway equipment on the ICU. Critical Care, 2012, 16,	10.8	78
273	A new patient mobilization scoring system in the ICU: what is the degree of similarity in scores between assessors in daily use?. <i>Critical Care</i> , 2012 , 16,	10.8	78
272	Motor and respiratory intensive rehabilitation in bedridden patients. Critical Care, 2012, 16,	10.8	1
271	Severity of electrophysiological alterations correlates with severity of illness in the early phase of critical illness polyneuropathy. <i>Critical Care</i> , 2012 , 16,	10.8	78
270	Muscle strength assessment of critically ill patients is associated with functional ability and quality of life at hospital discharge. <i>Critical Care</i> , 2012 , 16,	10.8	1

269	Functional dependency in the direct post-ICU phase in patients with prolonged mechanical ventilation. <i>Critical Care</i> , 2012 , 16,	10.8	78
268	IL-17A rs1974226 GG genotype is associated with increased susceptibility to Gram-positive infection and mortality of severe sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	78
267	Prevalence of TLR4 single nucleotide polymorphisms (ASP299GLY, THR399ILE) in healthy subjects and septic patients, and association with outcome. <i>Critical Care</i> , 2012 , 16,	10.8	78
266	Modelling immune responses in sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	78
265	Decreased peripheral CD4+/CD8+ lymphocytes and poor prognosis in aged sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	78
264	Homeostatic pulmonary microenvironment is responsible for alveolar macrophages resistance to endotoxin tolerance. <i>Critical Care</i> , 2012 , 16,	10.8	78
263	In vivo natural killer and natural killer T-cell depletion affects mortality in a murine pneumococcal pneumonia sepsis model. <i>Critical Care</i> , 2012 , 16,	10.8	78
262	Mobilization of hematopoietic and nonhematopoietic stem cell subpopulations in sepsis: a preliminary report. <i>Critical Care</i> , 2012 , 16,	10.8	1
261	Blunted IL-17 responses early after advent of multiple injuries. Critical Care, 2012, 16,	10.8	78
260	Apoptosis of neutrophils, expression of TREM-1 on neutrophils and IL-17 responses in experimental burn in injury are related to the type and time of burn exposure. <i>Critical Care</i> , 2012 , 16,	10.8	78
259	Insufficient autophagy relates to mitochondrial dysfunction, organ failure and adverse outcome in an animal model of critical illness. <i>Critical Care</i> , 2012 , 16,	10.8	1
258	Modulation of mediators derived from whole blood or monocytic cells stimulated with lipopolysaccharide reduces endothelial cell activation. <i>Critical Care</i> , 2012 , 16,	10.8	78
257	A/H1N1 infection: immunological parameters in ICU patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
256	Time of course CD64, a leukocyte activation marker, during extracorporeal circulation. <i>Critical Care</i> , 2012 , 16,	10.8	78
255	Oral neutrophil quantitation in patients undergoing elective cardiopulmonary bypass. <i>Critical Care</i> , 2012 , 16,	10.8	78
254	C13-pyruvate administration revealed differential metabolism between heart, liver and red blood cells and improved heart function during endotoxemia. <i>Critical Care</i> , 2012 , 16,	10.8	78
253	AMP-activated protein kinase controls liposaccharide-induced hyperpermeability. <i>Critical Care</i> , 2012 , 16,	10.8	2
252	Reduced expression of PPAR- Mimits the potential beneficial effects of GW0742 during septic shock in atherosclerotic swine. <i>Critical Care</i> , 2012 , 16,	10.8	78

251	Effects of hexafluoro-2-propanol on inflammatory and hemodynamic responses in a rat model of endotoxic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
250	Effects of noradrenaline and lipopolysaccharide exposure on mitochondrial respiration in alveolar macrophages. <i>Critical Care</i> , 2012 , 16,	10.8	78
249	Effects of the anti-diabetic imeglimin in hyperglycemic mice with septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	1
248	Adrenomedullin blockade improves catecholamine responsiveness and kidney function in resuscitated murine septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
247	Activated protein C, severe sepsis and 28-day mortality. <i>Critical Care</i> , 2012 , 16,	10.8	78
246	Soluble usokinase plasminogen activator receptor as a useful biomarker to define advent of sepsis in patients with multiple injuries. <i>Critical Care</i> , 2012 , 16,	10.8	78
245	Role of serum biomarkers in the diagnosis of infection in patients undergoing extracorporeal membrane oxygenation. <i>Critical Care</i> , 2012 , 16,	10.8	1
244	Correlation of VAP diagnosis with parameters of critically ill patients in a general ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
243	Usefulness of daily monitoring of procalcitonin and C-reactive protein in the early diagnosis of infection after elective colonic surgery. <i>Critical Care</i> , 2012 , 16,	10.8	1
242	Procalcitonin as a predictive marker for PCR test and blood culture results in suspected invasive candidemia. <i>Critical Care</i> , 2012 , 16,	10.8	78
241	Would procalcitonin measurement aid antimicrobial stewardship in a UK district general hospital mixed adult critical care population?. <i>Critical Care</i> , 2012 , 16,	10.8	78
240	Procalcitonin has a poor prognosis value in critically ill patients with candidemia. <i>Critical Care</i> , 2012 , 16,	10.8	78
239	Assessment of the usefulness of presepsin (soluble CD14 subtype) in septic patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
238	Circulating cell-free DNA levels measured by a novel simple fluorescent assay are predictive for outcome of severe sepsis. <i>Critical Care</i> , 2012 , 16,	10.8	78
237	Clinical usefulness of measuring endotoxin activity on ICU admission. <i>Critical Care</i> , 2012 , 16,	10.8	78
236	Prognostic value of serum galactomannan in mixed ICU patients: a retrospective observational study. <i>Critical Care</i> , 2012 , 16,	10.8	78
235	Analysis of (1-隂) 턴-glucan as a diagnostic adjunct for invasive fungal infections in the ICU setting. <i>Critical Care</i> , 2012 , 16,	10.8	78
234	Impaired heart rate variability predicts clinical deterioration and progressive organ failure in emergency department sepsis patients. <i>Critical Care</i> , 2012 , 16,	10.8	4

233	Severe community-acquired pneumonia: risk factors for in-hospital mortality. Critical Care, 2012, 16,	10.8	1
232	Systemic corticosteroids for community-acquired pneumonia in adults. <i>Critical Care</i> , 2012 , 16,	10.8	78
231	Characteristics of leptospirosis patients admitted to a tropical university hospital during the 2000 to 2010 period. <i>Critical Care</i> , 2012 , 16,	10.8	1
230	Prognostic impact of imported and newly-isolated methicillin-resistant Staphylococcus aureus in the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
229	Necrotizing pneumonia due to methicillin-sensitive Staphylococcus aureus secreting Panton-Valentine leukocidin: a review of case reports. <i>Critical Care</i> , 2012 , 16,	10.8	78
228	Predicting methicillin-resistant Staphylococcus aureus in critically ill patients with pneumonia presenting to the hospital. <i>Critical Care</i> , 2012 , 16,	10.8	78
227	Predictors of multidrug-resistant Acinetobacter baumannii infections: a retrospective analysis in surgical ICU patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
226	Risk factors for bronchial acquisition of resistant Gram-negative bacteria in critically ill patients and outcome. <i>Critical Care</i> , 2012 , 16,	10.8	78
225	Improved antibiotic stewardship resulting from a multifaceted strategy implemented after an outbreak of multiresistant Acinetobacter baumannii in a university ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
224	Empirical antifungal treatment in the critically ill patients: how does it impact on the outcome?. <i>Critical Care</i> , 2012 , 16,	10.8	78
223	Relationship between polyclonal immunoglobulin therapy and colonization by Candida spp <i>Critical Care</i> , 2012 , 16,	10.8	78
222	Predictive and prognostic factors of septic shock of nosocomial origin. Critical Care, 2012, 16,	10.8	1
221	Catheter-related bloodstream infection: factors affecting incidence. Critical Care, 2012, 16,	10.8	78
220	Prognostic factors of septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
219	Severe sepsis in the United Sates: a 5-year analysis. <i>Critical Care</i> , 2012 , 16,	10.8	78
218	District hospital experience of organ support requirements for H1N1-associated pneumonia. <i>Critical Care</i> , 2012 , 16,	10.8	78
217	Compliance with the sepsis resuscitation bundle in patients with severe sepsis and septic shock admitted to Scottish ICUs. <i>Critical Care</i> , 2012 , 16,	10.8	78
216	Improving early administration of antibiotics: a PPlan Do Study ActPapproach. Critical Care, 2012, 16,	10.8	78

215	Source-directed antimicrobials: a shot in the dark?. Critical Care, 2012, 16,	10.8	78
214	Relation between temperature in the initial 24 hours in patients with severe sepsis or septic shock with mortality and length of stay in the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
213	Temperature management for patients without brain injury in Australia and New Zealand ICUs: a point prevalence study. <i>Critical Care</i> , 2012 , 16,	10.8	78
212	Impact of antifungal treatment in ICU patients with Candida colonization: analysis of the EPIC II study population. <i>Critical Care</i> , 2012 , 16,	10.8	78
211	Pharmacokinetics of micafungin in patients with severe burn injuries. <i>Critical Care</i> , 2012 , 16,	10.8	78
2 10	Bacteremia affects the mortality of septic patients with high serum procalcitonin level in the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
209	Cases of tetanus after the Japan crisis 2011. <i>Critical Care</i> , 2012 , 16,	10.8	1
208	Does the day of the week predict the presence of microbiologically confirmed ventilator-associated pneumonia?. <i>Critical Care</i> , 2012 , 16,	10.8	78
207	Assessing perforation of acute appendicitis using the delta neutrophil index reflecting the peripheral immature granulocyte count. <i>Critical Care</i> , 2012 , 16,	10.8	78
206	Effects of a multifaceted quality improvement intervention in reducing mortality and bloodstream infection in ICUs: insights from the QUALITI initiative. <i>Critical Care</i> , 2012 , 16,	10.8	78
205	Ceftazidime dosage regimen recommendations in burn patients based on a Monolix population pharmacokinetic study. <i>Critical Care</i> , 2012 , 16,	10.8	78
204	Continuous versus intermittent vancomycin in children after cardiac surgery with delayed sternal closure. <i>Critical Care</i> , 2012 , 16,	10.8	1
203	Elimination of linezolid in patients undergoing low-flow continuous venovenous haemodiafiltration. <i>Critical Care</i> , 2012 , 16,	10.8	1
202	A post-authorisation survey to analyse the perioperative teicoplanin plasma concentrations in adult patients with chronic bone sepsis, who received loading doses of 12 mg/kg 12-hourly for 48 hours followed by 12 mg/kg once daily. <i>Critical Care</i> , 2012 , 16,	10.8	1
201	Pharmacokinetics of inhaled colistin in critically ill patients with ventilator-associated tracheobronchitis. <i>Critical Care</i> , 2012 , 16,	10.8	78
200	Efficacy of inhaled tobramycin in severe nosocomial pneumonia. Critical Care, 2012, 16,	10.8	1
199	Comparison of a bronchoscopic microsample probe with bronchoalveolar lavage to measure cytokine levels in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
198	Clinical and epidemiological risk factors for ventilator-associated pneumonia in a cohort of critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	1

197	Use of a ventilator-associated pneumonia (VAP) bundle to decrease the VAP rate in Syria. <i>Critical Care</i> , 2012 , 16,	10.8	78
196	A strategy for prevention and control of catheter-related bloodstream infection of ICU patients in China (Prevent CRBSI): a prospective, multicenter, controlled study. <i>Critical Care</i> , 2012 , 16,	10.8	1
195	Wash your hands: simple measures save lives. <i>Critical Care</i> , 2012 , 16,	10.8	78
194	Comparison of hand hygiene in single-room versus open-plan ICUs. <i>Critical Care</i> , 2012 , 16,	10.8	78
193	Compliance for decontamination of bedside computer keyboards on an ICU. Critical Care, 2012, 16,	10.8	78
192	Reduced air contamination in an ICU environment with a portable air purification system. <i>Critical Care</i> , 2012 , 16,	10.8	78
191	Effectiveness of an innovative system for the bio-decontamination of the ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
190	Massive hemoptysis in a respiratory ICU: causes, interventions and outcomes - Indian study. <i>Critical Care</i> , 2012 , 16,	10.8	1
189	Capnography use in Scottish ICUs. <i>Critical Care</i> , 2012 , 16,	10.8	78
188	Digitalized acoustic monitoring of lung congestion. <i>Critical Care</i> , 2012 , 16,	10.8	78
187	Usefulness of electrical activity of the diaphragm to detect intrinsic positive end-expiratory pressure during pressure support ventilation. <i>Critical Care</i> , 2012 , 16,	10.8	78
186	Adequate lung sliding identification is not influenced by the level of academic or ultrasound training. <i>Critical Care</i> , 2012 , 16,	10.8	78
185	Lung ultrasound can differentiate Pneumocystis jiroveci versus other etiologies among critically ill AIDS patients with pneumonia. <i>Critical Care</i> , 2012 , 16,	10.8	5
184	Difference in accuracy of lung sliding identification between the right and left hemithorax. <i>Critical Care</i> , 2012 , 16,	10.8	78
183	Trans-thoracic echo evaluation before and during noninvasive ventilation. <i>Critical Care</i> , 2012 , 16,	10.8	1
10)	Trans choracte cens evaluation service and daring noninvasive ventilation. Critical care, 2012, 10,	10.0	
182	Listen to PaO2/FiO2 ratios: they tell us about length of stay. <i>Critical Care</i> , 2012 , 16,	10.8	78
			78 78

179	Do actual tidal volumes differ from prescribed tidal volumes?. Critical Care, 2012, 16,	10.8	78
178	Intratracheal administration of siRNA targeting FAS reduces ischemia-reperfusion-induced lung injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
177	Acute lung injury in mice associates with p44/42 and c-Jun N-terminal kinase activation and requires the function of TNFI receptor I. <i>Critical Care</i> , 2012 , 16,	10.8	78
176	Retrieval of patients with severe respiratory failure on venovenous extracorporeal membrane oxygenation: an intensivist-led model. <i>Critical Care</i> , 2012 , 16,	10.8	78
175	A new miniaturized extracorporeal membrane oxygenator with integrated rotary blood pump (Ilias): first results in a porcine model of lung injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
174	ECMO in nonintubated patients as a bridge to lung transplant: our experience. <i>Critical Care</i> , 2012 , 16,	10.8	78
173	Extracorporeal life support in major trauma: case series from a tertiary referral trauma. <i>Critical Care</i> , 2012 , 16,	10.8	78
172	Two yearsPexperience with bicaval dual lumen cannula for venovenous extracorporeal membrane oxygenation in adult refractory acute respiratory distress syndrome. <i>Critical Care</i> , 2012 , 16,	10.8	1
171	Resolution of organ functional scores to predict outcomes in severe acute respiratory distress syndrome patients receiving extracorporeal membrane oxygenation. <i>Critical Care</i> , 2012 , 16,	10.8	1
170	Corticosteroids for critically ill patients: an international survey of intensivists. <i>Critical Care</i> , 2012 , 16,	10.8	78
169	Effects of salbutamol on airway characteristics in mechanically ventilated adults without COPD. <i>Critical Care</i> , 2012 , 16,	10.8	78
168	Respiratory system elastance monitoring during PEEP titration. Critical Care, 2012, 16,	10.8	2
167	Accuracy of the pressure-volume curve method compared to quantitative lung CT scan to assess the recruitable lung in patients with acute respiratory failure. <i>Critical Care</i> , 2012 , 16,	10.8	78
166	Flow-controlled expiration discloses PEEP-dependent dynamic hysteresis of the pressure-volume loop. <i>Critical Care</i> , 2012 , 16,	10.8	78
165	Feasibility of early spontaneous breathing in acute respiratory distress syndrome. <i>Critical Care</i> , 2012 , 16,	10.8	78
164	A device for ventilation-analogue mechanostimulation in vitro. <i>Critical Care</i> , 2012 , 16,	10.8	78
163	Hemodynamics effects of recruitment maneuver. Critical Care, 2012, 16,	10.8	2
162	Differential pulmonary and circulatory effects of preventive lung protective ventilation in an experimental postoperative sepsis model. <i>Critical Care</i> , 2012 , 16,	10.8	78

161	High NT-proBNP level is correlated with high PEEP, low PH and low PaO2/FiO2 in ARDS. <i>Critical Care</i> , 2012 , 16,	10.8	78
160	Early application of high-frequency oscillatory ventilation in H1N1 influenza-related severe ARDS is associated with better outcome. <i>Critical Care</i> , 2012 , 16,	10.8	78
159	Outcomes of early delivery in pregnant patients with acute respiratory distress syndrome. <i>Critical Care</i> , 2012 , 16,	10.8	3
158	Mechanical ventilation demographics between 1999 and 2009. Critical Care, 2012, 16,	10.8	78
157	Mechanical ventilation in intensive and critical care units of Russia: RuVent national epidemiologic study. <i>Critical Care</i> , 2012 , 16,	10.8	3
156	Use of a fully closed-loop ventilation mode in long-term ventilated ICU patients: a prospective study. <i>Critical Care</i> , 2012 , 16,	10.8	78
155	Effects of low and high tidal volume and pentoxifylline on intestinal blood flow and leukocyte-endothelial interactions in mechanically ventilated rats. <i>Critical Care</i> , 2012 , 16,	10.8	78
154	A method for continuous noninvasive assessment of respiratory mechanics during spontaneous breathing. <i>Critical Care</i> , 2012 , 16,	10.8	78
153	Influence of catheter diameter, endotracheal tube diameter, suction pressure, and PEEP on the tracheal pressure and lung volume during endotracheal suctioning using a lung model. <i>Critical Care</i> , 2012 , 16,	10.8	78
152	Risk factors of mortality in severe cutaneous adverse reactions patients with pulmonary involvement. <i>Critical Care</i> , 2012 , 16,	10.8	78
151	During spontaneous breathing cardiac output lacks major effect on pulmonary shunting in porcine lungs with partial collapse. <i>Critical Care</i> , 2012 , 16,	10.8	78
150	Oxygenation correlates with lung aeration during unsupported spontaneous breathing in porcine lung collapse model. <i>Critical Care</i> , 2012 , 16,	10.8	78
149	Safety and effect of intermittent intrapulmonary percussive ventilation on oxygen saturation and hemodynamic functions. <i>Critical Care</i> , 2012 , 16,	10.8	78
148	Impact of an open lung approach on hemodynamic parameters after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
147	A protective-ventilation strategy reduces pulmonary complications after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
146	A rule for predicting the new equilibrated carbon dioxide partial pressure after changes in the ventilation frequency. <i>Critical Care</i> , 2012 , 16,	10.8	78
145	Patient-ventilator asynchrony during conventional or automated pressure support ventilation in difficult-to-wean patients. <i>Critical Care</i> , 2012 , 16,	10.8	2
144	High levels of B-type natriuretic peptide predict weaning failure from mechanical ventilation in adult patients after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78

143	Case-control study of failed extubation. <i>Critical Care</i> , 2012 , 16,	10.8	2
142	Out-of-bed extubation: changing paradigms. <i>Critical Care</i> , 2012 , 16,	10.8	78
141	Prediction of post-extubation failure by portable ICU ultrasound. Critical Care, 2012, 16,	10.8	78
140	Intermittent aspiration of pharyngeal secretion for re-intubation prevention. Critical Care, 2012, 16,	10.8	78
139	Efficacy of biphasic cuirass ventilation in the critical care department. Critical Care, 2012, 16,	10.8	1
138	Differences in neurophysiologic effects between CPAP and a novel high-flow therapy system. <i>Critical Care</i> , 2012 , 16,	10.8	78
137	Management of acute bronchospasm respiratory distress with CPAP ventilation associated with nebulization in the prehospital emergency setting. <i>Critical Care</i> , 2012 , 16,	10.8	78
136	Difference between continuous positive airway pressure via mask therapy plus chest physiotherapy (CPT) and incentive spirometry therapy plus CPT to treat or prevent acute atelectasis after cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
135	Nasal high-flow oxygen in patients with hypoxic respiratory failure: effect on functional and subjective respiratory parameters compared to conventional oxygen therapy and noninvasive ventilation. <i>Critical Care</i> , 2012 , 16,	10.8	2
134	Short-term effect of humidified high nasal flow oxygen in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
133	Good response on high nasal oxygen flow reduces the need for intubation in adult respiratory failure. <i>Critical Care</i> , 2012 , 16,	10.8	78
132	An audit of airway complications in a district general hospital ICU. Critical Care, 2012, 16,	10.8	78
131	Multidisciplinary care for patients with tracheostomy shortened time to decannulation. <i>Critical Care</i> , 2012 , 16,	10.8	78
130	Influence of percutaneous tracheostomy on gas exchange in mechanically ventilated patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
129	Bronchoscope-guided percutaneous dilatational tracheostomy performed by an experienced intensivist: a 26-month experience at a tertiary care center in United Arab Emirates. <i>Critical Care</i> , 2012 , 16,	10.8	78
128	Risk factors for poor outcome in patients with osmotic demyelination syndrome. <i>Critical Care</i> , 2012 , 16,	10.8	3
127	Is inappropriate secretion of anti-diuretic hormone (SIADH) the cause of hyponatremia in Legionella pneumonia?. <i>Critical Care</i> , 2012 , 16,	10.8	78
126	Fluctuations in serum sodium level are associated with an increased risk of death in surgical ICU patients. <i>Critical Care</i> , 2012 , 16,	10.8	78

125	Impact of ketogenesis and strong ion difference on acid-base in our CICU. Critical Care, 2012, 16,	10.8	78
124	Buffer therapy in metabolic acidosis after surgery-associated hemorrhage in pediatric oncology. <i>Critical Care</i> , 2012 , 16,	10.8	78
123	An unusual cause of high anion gap metabolic acidosis: pyroglutamic acidosis. <i>Critical Care</i> , 2012 , 16,	10.8	О
122	Metabolic acid-base disturbances in patients in the emergency department. Critical Care, 2012, 16,	10.8	78
121	Aberrant bone metabolism in critical illness. <i>Critical Care</i> , 2012 , 16,	10.8	78
120	Low serum 25-hydroxyvitamin D levels and acute kidney injury in the critically ill. <i>Critical Care</i> , 2012 , 16,	10.8	78
119	Concentration of major vitamins in critically ill patients. Critical Care, 2012, 16,	10.8	78
118	No association between vitamin D deficiency at admission and outcome in a medical ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
117	Plasma levels of Coenzyme Q10 are reduced in critically ill patients as compared to healthy volunteers and correlate with age. <i>Critical Care</i> , 2012 , 16,	10.8	3
116	Reduced cortisol metabolism drives hypercortisolism in critical illness. <i>Critical Care</i> , 2012 , 16,	10.8	78
115	Effect of low-dose hydrocortisone on the expression of glucocorticoid receptor alpha of the septic kidney in rats and its protective effect on kidney injury. <i>Critical Care</i> , 2012 , 16,	10.8	78
114	Hydrocortisone increases the risk of dysglycemia in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
113	Nutritional status of patients occupying ICUs in the state of Rio de Janeiro. Critical Care, 2012, 16,	10.8	78
112	Investigating diarrhoea on the ICU: a retrospective study. Critical Care, 2012, 16,	10.8	78
111	Preliminary report of surface electrogastrography in critically ill septic patients after resuscitation. <i>Critical Care</i> , 2012 , 16,	10.8	78
110	Frequency, determinants and impact of feed intolerance amongst the critically ill. <i>Critical Care</i> , 2012 , 16,	10.8	78
109	Gastric emptying assessment in critically ill patients with feed intolerance; comparison of 13C octanoic acid, paracetamol and 3-O-methylglucose absorption tests. <i>Critical Care</i> , 2012 , 16,	10.8	78
108	Early enteral nutrition in the critically ill: a single-centre study. Critical Care, 2012, 16,	10.8	78

107	Preliminary experience with ketone-targeted treatment of diabetic ketoacidosis. <i>Critical Care</i> , 2012 , 16,	10.8	78
106	Impact of blood glucose on blood lactate levels in a medical ICU: a retrospective cohort study. <i>Critical Care</i> , 2012 , 16,	10.8	5
105	Relationship between glycemic Lability Index, infections and outcome in critically ill patients. <i>Critical Care</i> , 2012 , 16,	10.8	О
104	Variability of insulin sensitivity during the first 4 days of critical illness. Critical Care, 2012, 16,	10.8	4
103	Endogenous insulin secretion in critically ill patients. Critical Care, 2012, 16,	10.8	78
102	Impact of the type of glucose monitoring on the assessment of glycemic variability in critical care patients. <i>Critical Care</i> , 2012 , 16,	10.8	2
101	Evaluation of a continuous blood glucose monitoring system using a central venous catheter with an integrated microdialysis function. <i>Critical Care</i> , 2012 , 16,	10.8	78
100	Variable and maximum blood glucose levels during the first week after ICU admission are related to the severity of the patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
99	Pilot trial of STAR in the medical ICU. <i>Critical Care</i> , 2012 , 16,	10.8	78
98	Glucometer accuracy and implications for clinical studies. Critical Care, 2012, 16,	10.8	1
97	Initial experience with continuous intra-arterial fluorescent glucose monitoring in patients in the ICU following cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	78
96	Preliminary ICU experience of a novel intravascular blood glucose sensor. Critical Care, 2012, 16,	10.8	3
95	Does tight glycemic control positively impact on patient mortality?. Critical Care, 2012, 16,	10.8	78
94	Glycaemic control in ICUs in large English hospitals: a follow-up telephone survey. <i>Critical Care</i> , 2012 , 16,	10.8	78
93	Model-based regulation of glucose in critical care. Critical Care, 2012, 16,	10.8	78
92	Perioperative glycemic control with a computerized algorithm versus conventional glycemic control. <i>Critical Care</i> , 2012 , 16,	10.8	78
91	Efficacy of the novel heart attack centre extension pathway: a pilot study. Critical Care, 2012, 16,	10.8	78
90	Integral assistance process implantation for ST-elevated acute coronary syndrome. <i>Critical Care</i> , 2012 , 16,	10.8	78

89	Prognostic value of Killip classification in terms of health-related quality of life. <i>Critical Care</i> , 2012 , 16,	10.8	78
88	Modes of arrival, door to balloon time and its impact on morbidity and mortality for ST elevation myocardial infarct. <i>Critical Care</i> , 2012 , 16,	10.8	3
87	Next-generation, fast and accurate point-of-care test for NT-proBNP based on Magnotech technology. <i>Critical Care</i> , 2012 , 16,	10.8	78
86	Right ventricular apical versus septal pacing: impact on left ventricular synchrony and function. <i>Critical Care</i> , 2012 , 16,	10.8	78
85	Consecutive case series of Takotsubo cardiomyopathy: a disease potentially triggered by the Great East Japan Earthquake. <i>Critical Care</i> , 2012 , 16,	10.8	78
84	Stress cardiomyopathy after live donor liver transplantation: incidence, risk factors and mortality. <i>Critical Care</i> , 2012 , 16,	10.8	78
83	Short-term hemodynamic effects of nebivolol in acute decompensated heart failure: a randomized clinical trial. <i>Critical Care</i> , 2012 , 16,	10.8	78
82	Patients with infective endocarditis patients in the ICU: how are they?. Critical Care, 2012, 16,	10.8	78
81	Malperfusion and branch compromise in acute type A aortic syndrome. Critical Care, 2012, 16,	10.8	78
80	Prognostic value of the echocardiographic-derived calcium index in coronary artery disease. <i>Critical Care</i> , 2012 , 16,	10.8	78
79	Echocardiography in the ICU: an audit of 3 years practice. Critical Care, 2012, 16,	10.8	1
78	Left ventriculum diastolic dysfunction in pediatric septic shock. Critical Care, 2012, 16,	10.8	78
77	Existence of interference between the heart and respiratory sounds: preliminary report. <i>Critical Care</i> , 2012 , 16,	10.8	78
76	Effects of the intravenous administration of purine nucleosides guanosine or inosine against hemorrhagic shock in pigs. <i>Critical Care</i> , 2012 , 16,	10.8	78
75	Norepinephrine versus angiotensin II in septic shock: effects on isolated kidney, heart and liver mitochondrial respiration. <i>Critical Care</i> , 2012 , 16,	10.8	78
74	Goal-directed hemodynamic resuscitation in high-risk patients undergoing cardiac surgery: a randomized controlled trial - preliminary data (GRICCS STUDY). <i>Critical Care</i> , 2012 , 16,	10.8	1
73	Economic evaluation of early-goal directed therapy for high-risk surgical patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
72	What matters during a hypotensive episode: fluids, vasopressors, or both?. <i>Critical Care</i> , 2012 , 16,	10.8	78

71	Elevated central venous pressure in septic patients is associated with impairment of microcirculatory blood flow. <i>Critical Care</i> , 2012 , 16,	10.8	78
70	Human protein C concentrate to restore physiological values in adult septic shock patients: effects on microcirculation. <i>Critical Care</i> , 2012 , 16,	10.8	78
69	Heart rate reduction with esmolol in septic shock: effects on microcirculation. <i>Critical Care</i> , 2012 , 16,	10.8	78
68	Microcirculation and blood transfusion: effects of three different types of concentrated red blood cells - preliminary results. <i>Critical Care</i> , 2012 , 16,	10.8	2
67	Peripheral perfusion is correlated to metabolic perfusion parameters and microvascular reactivity but not with hepatosplanchnic or microcirculatory flow parameters in hyperdynamic septic shock. <i>Critical Care</i> , 2012 , 16,	10.8	78
66	Hyperoxia affects peripheral tissue microcirculation in patients with pulmonary arterial hypertension. <i>Critical Care</i> , 2012 , 16,	10.8	2
65	Supraclavicular ultrasound-guided subclavian vein cannulation in infants under 5 kg. <i>Critical Care</i> , 2012 , 16,	10.8	78
64	Central venous catheter placement: where is the end of the line?. Critical Care, 2012, 16,	10.8	78
63	Power-injectable peripherally inserted central catheters in intensive care patients. <i>Critical Care</i> , 2012 , 16,	10.8	1
62	Comparison of internal jugular and subclavian access for central venous catheterization in pediatric cardiac surgery. <i>Critical Care</i> , 2012 , 16,	10.8	1
61	Ultrasound-guided central venous line placement in critically ill patients: is chest X-ray needed to assess post-insertion pneumothorax?. <i>Critical Care</i> , 2012 , 16,	10.8	78
60	Errors in the arterial blood pressure measurement. <i>Critical Care</i> , 2012 , 16,	10.8	1
59	The T-Line TL-200 system for continuous noninvasive blood pressure measurement in medical ICU patients. <i>Critical Care</i> , 2012 , 16,	10.8	78
58	Reliability of radial arterial pressure monitoring after cardiac surgery. Critical Care, 2012, 16,	10.8	1
57	Right/left ventricular area ratio does not correlate with right ventricular impedance. <i>Critical Care</i> , 2012 , 16,	10.8	1
56	Role of mannose-binding lectin on pneumococcal infections. Critical Care, 2012, 16,	10.8	78
55	The impact of organ dysfunction in cirrhosis: survival at a cost?. <i>Journal of Hepatology</i> , 2012 , 57, 707-8; author reply 709	13.4	2
54	Abstracts of the 32nd International Symposium on Intensive Care and Emergency Medicine. March 20-23, 2012. Brussels, Belgium. <i>Critical Care</i> , 2012 , 16 Suppl 1, P1-520	10.8	1

53	Opportunities for prevention of alcohol-related death in primary care: results from a population-based cross-sectional study. <i>Alcohol</i> , 2012 , 46, 703-7	2.7	9
52	Descriptive epidemiology of chronic liver disease in northeastern Italy: an analysis of multiple causes of death. <i>Population Health Metrics</i> , 2013 , 11, 20	3	8
51	Review article: towards a considered and ethical approach to organ support in critically-ill patients with cirrhosis. <i>Alimentary Pharmacology and Therapeutics</i> , 2013 , 37, 174-82	6.1	25
50	Unplanned alcohol withdrawal: a survey of consecutive admissions to an acute medical unit in 2010 and 2011. <i>QJM - Monthly Journal of the Association of Physicians</i> , 2013 , 106, 43-9	2.7	1
49	An identification and brief advice programme for low-risk alcohol consumption in an acute medical setting: an implementation study. <i>JRSM Short Reports</i> , 2013 , 4, 2042533313476696		2
48	Alcohol-induced Hyperlipidemia Is Ameliorated by Orally Administered DWP208, a Sodium Succinate Form of ZYM201. <i>Korean Journal of Physiology and Pharmacology</i> , 2014 , 18, 469-74	1.8	7
47	Effect of financial incentives on delivery of alcohol screening and brief intervention (ASBI) in primary care: longitudinal study. <i>Journal of Public Health</i> , 2014 , 36, 450-9	3.5	21
46	Alcohol withdrawal recognition and screening in the acute medical setting. <i>European Journal of Internal Medicine</i> , 2014 , 25, e87	3.9	
45	Alcohol age of initiation and long-term impact: a cross sectional survey of adults in England. <i>Journal of Substance Use</i> , 2014 , 1-7	0.8	2
44	New real-time bowel sound analysis may predict disease severity in septic patients. <i>Critical Care</i> , 2015 , 19, P2	10.8	78
43	Patient epidemiology in a level II hospital ICU and how main nosocomial infections affect morbidity and mortality. <i>Critical Care</i> , 2015 , 19, P94	10.8	78
42	Emergence of isolates that are intrinsically resistant to colistin in critically ill patients: are we selecting them out?. <i>Critical Care</i> , 2015 , 19, P95	10.8	1
41	Risk factors for severe vasodilatory shock after cardiac surgery. <i>Critical Care</i> , 2015 , 19, P149	10.8	1
40	Preoperative treatment with levosimendan helps to evaluate myocardial reserves in cardiosurgical patients with chronic heart failure. <i>Critical Care</i> , 2015 , 19, P150	10.8	1
39	Ultrasound assessment for extravascular lung water in patients with septic shock. <i>Critical Care</i> , 2015 , 19, P223	10.8	78
38	Quantitative ultrasonography for pneumonia. <i>Critical Care</i> , 2015 , 19, P225	10.8	78
37	Mesenchymal stem cell and endothelial cell interaction restores endothelial permeability via paracrine hepatocyte growth factor in vitro. <i>Critical Care</i> , 2015 , 19, P236	10.8	3
36	Does it make a difference to add automatic EPAP titration to the volume-targeted pressure support mode in noninvasive ventilation of hypercapnic ICU patients?. <i>Critical Care</i> , 2015 , 19, P251	10.8	78

35	Effects of iron deficiency on transfusion requirements in critically ill patients: a preliminary observational study. <i>Critical Care</i> , 2015 , 19, P339	10.8	78
34	Intensive care referral and admission: do the criteria for liver disease match?. <i>Critical Care</i> , 2015 , 19, P3	84 0.8	78
33	Evaluation of emergency call Code Blue over a 5-year period. <i>Critical Care</i> , 2015 , 19, P411	10.8	78
32	Utilisation and prognostic impact of angiography and primary percutaneous coronary intervention prior to intensive care admission for patients following out-of-hospital cardiac arrest. <i>Critical Care</i> , 2015 , 19, P423	10.8	78
31	Ten years of hospital admissions for liver cirrhosis in Portugal. <i>European Journal of Gastroenterology and Hepatology</i> , 2015 , 27, 1320-6	2.2	9
30	Increased Survival for Patients With Cirrhosis and Organ Failure in Liver Intensive Care and Validation of the Chronic Liver Failure-Sequential Organ Failure Scoring System. <i>Clinical Gastroenterology and Hepatology</i> , 2015 , 13, 1353-1360.e8	6.9	65
29	35th international symposium on intensive care and emergency medicine. <i>Critical Care</i> , 2015 , 19 Suppl 1, P1-P578	10.8	4
28	Evaluation of work-based screening for early signs of alcohol-related liver disease in hazardous and harmful drinkers: the PrevAIL study. <i>BMC Public Health</i> , 2015 , 15, 532	4.1	6
27	The burden of alcoholism in fifteen years of cirrhosis hospital admissions in Portugal. <i>Liver International</i> , 2015 , 35, 746-55	7.9	15
26	Angelica keiskei Koidzumi extracts improve some markers of liver function in habitual alcohol drinkers: a randomized double-blind clinical trial. <i>Journal of Medicinal Food</i> , 2015 , 18, 166-72	2.8	9
25	Assessment of renal functions in patients of chronic liver disease. <i>Renal Failure</i> , 2015 , 37, 1457-63	2.9	2
24	Implementation of a Rare bundlePimproves the management of patients admitted to hospital with decompensated cirrhosis. <i>Alimentary Pharmacology and Therapeutics</i> , 2016 , 44, 1030-1038	6.1	14
23	Alcoholic Liver Disease in Asia, Europe, and North America. <i>Gastroenterology</i> , 2016 , 150, 1786-97	13.3	94
22	Patients with alcohol-related liver diseasebeliefs about their illness and factors that influence their self-management. <i>Journal of Advanced Nursing</i> , 2016 , 72, 173-85	3.1	14
21	Response to the NCEPOD report: development of a care bundle for patients admitted with decompensated cirrhosis-the first 24 h. <i>Frontline Gastroenterology</i> , 2016 , 7, 16-23	2.6	16
20	Trends in alcohol-related admissions to hospital by age, sex and socioeconomic deprivation in England, 2002/03 to 2013/14. <i>BMC Public Health</i> , 2017 , 17, 412	4.1	20
19	Disease spectrum of alcoholic liver disease in Beijing 302 Hospital from 2002 to 2013: A large tertiary referral hospital experience from 7422 patients. <i>Medicine (United States)</i> , 2017 , 96, e6163	1.8	21
18	Educating students with FASD: linking policy, research and practice. <i>Journal of Research in Special Educational Needs</i> , 2017 , 17, 3-17	1.7	13

17	National Early Warning Score Accurately Discriminates the Risk of Serious Adverse Events in Patients With Liver Disease. <i>Clinical Gastroenterology and Hepatology</i> , 2018 , 16, 1657-1666.e10	6.9	6
16	Incidence and Outcomes for Patients With Cirrhosis Admitted to the United Kingdom Critical Care Units. <i>Critical Care Medicine</i> , 2018 , 46, 705-712	1.4	31
15	Comparing assessment tools for detecting undernutrition in patients with liver cirrhosis. <i>Clinical Nutrition ESPEN</i> , 2018 , 23, 156-161	1.3	12
14	NCEPOD and alcohol-related liver disease, what are the views of those who deliver the service? A survey of consultants and trainees in North Eastern England. <i>Journal of the Royal College of Physicians of Edinburgh, The</i> , 2018 , 48, 293-298	0.9	1
13	The changing aetiology of head and neck squamous cell cancer: A tale of three cancers?. <i>Clinical Otolaryngology</i> , 2018 , 43, 999-1003	1.8	10
12	Understanding alcohol-related liver disease patients Pillness beliefs and views about their medicine. <i>British Journal of Nursing</i> , 2018 , 27, 730-736	0.7	2
11	Place of death and factors associated with hospital death in patients who have died from liver disease in England: a national population-based study. <i>The Lancet Gastroenterology and Hepatology</i> , 2019 , 4, 52-62	18.8	6
10	Risk factors for intrahepatic and extrahepatic cholangiocarcinoma: A systematic review and meta-analysis. <i>Journal of Hepatology</i> , 2020 , 72, 95-103	13.4	102
9	Incidence and mortality of primary liver cancer in England and Wales: changing patterns and ethnic variations. <i>World Journal of Gastroenterology</i> , 2014 , 20, 1544-53	5.6	27
8	A Study on the Temporal Trends in the Etiology of Cirrhosis of Liver in Coastal Eastern Odisha. <i>Euroasian Journal of Hepato-gastroenterology</i> , 2020 , 10, 1-6	1.6	3
7	[Investigation of the frequency of alcohol related problems in J团sefvflos, Budapest. Data and conclusions of three surveys conducted among doctors and outpatients]. <i>Orvosi Hetilap</i> , 2012 , 153, 12	63 ⁻⁸⁰	
6	How to Find and Apply Large Databases for Epidemiologic Research. 83-97		
5	Hepatitis C in the three main health institutions in Mexico: a 13-year mortality and hospitalization analysis. <i>Revista Do Instituto De Medicina Tropical De Sao Paulo</i> , 2020 , 62, e11	2.2	0
4	Clinical, Epidemiological and Histopathological Aspects in Patients with Hepatocellular Carcinoma Undergoing Liver Transplantation. <i>Asian Pacific Journal of Cancer Prevention</i> , 2018 , 19, 2795-2802	1.7	5
3	A decade of rising alcoholic liver disease hospital admissions and deaths in Irish hospitals, 2007-2016: a retrospective cross-sectional analysis <i>European Journal of Gastroenterology and Hepatology</i> , 2022 ,	2.2	0
2	Risk factors of chronic liver disease among adult patients in tertiary hospitals, Northern Ethiopia: An unmatched case-control study.		
1	Behandlung von riskantem, schälichem und abhligigem Alkoholgebrauch. 2022 , 51-356		O